

ROBUST PARAMETER ESTIMATION FOR PARTIALLY OBSERVED SECOND-ORDER DIFFUSION PROCESSES

JAN ALBRECHT AND SEBASTIAN REICH

ABSTRACT. Estimating parameters of a diffusion process given continuous-time observations of the process via maximum likelihood approaches or, online, via stochastic gradient descent or Kalman filter formulations constitutes a well-established research area. It has also been established previously that these techniques are, in general, not robust to perturbations in the data in the form of temporal correlations. While the subject is relatively well understood and appropriate modifications have been suggested in the context of multi-scale diffusion processes and their reduced model equations, we consider here an alternative setting where a second-order diffusion process in positions and velocities is only observed via its positions. In this note, we propose a simple modification to standard stochastic gradient descent and Kalman filter formulations, which eliminates the arising systematic estimation biases. The modification can be extended to standard maximum likelihood approaches and avoids computation of previously proposed correction terms.

1. INTRODUCTION

We consider the problem of online learning parameters $\theta \in \mathbb{R}^{d_\theta}$ of second-order stochastic differential equations (SDEs)

$$(1a) \quad dU_t = f(X_t, U_t)dt + F(X_t, U_t)\theta dt + \sqrt{\sigma}dW_t,$$

$$(1b) \quad dX_t = U_t dt$$

from incoming observations $X_t^\dagger \in \mathbb{R}^d$, $t \geq 0$, of an underlying reference process. Here $f : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $F : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d_\theta}$ are given functions, $\sigma > 0$ is the diffusion constant, and W_t denotes d -dimensional Brownian motion.

The data X_t^\dagger is being generated from a second-order SDE of the same type, i.e.,

$$(2a) \quad dU_t^\dagger = g(X_t^\dagger, U_t^\dagger)dt + \sqrt{\sigma}dW_t^\dagger,$$

$$(2b) \quad dX_t^\dagger = U_t^\dagger dt$$

with the drift $g(x, u)$ being unknown. We assume that there exists a unique θ_* such that

$$(3) \quad g(x, u) = f(x, u) + F(x, u)\theta_*.$$

While standard online estimation techniques, such as stochastic gradient descent (SGD) [17] and Kalman filtering [11, 16], can be applied in case both $(X_t^\dagger, U_t^\dagger)$ from (2) are being

Date: June 24, 2024.

observed, the estimation problem becomes more challenging in case only positions X_t^\dagger are being observed and velocities need to be estimated using finite differencing [15, 5], i.e.,

$$(4) \quad \tilde{U}_{n+1/2} := \frac{X_{t_{n+1}}^\dagger - X_{t_n}^\dagger}{\tau}$$

with $\tau > 0$ denoting the sampling interval, $t_n = \tau n$, for all $n \geq 0$.

As demonstrated previously [9], naive application of standard maximum likelihood-based (MLE) estimators [10, 13] leads to a systematic bias. In order to correct for the arising estimation bias, appropriate correction terms to the MLE estimator have been proposed in [9, 5, 3]. Related phenomena have been investigated in the context of multi-scale diffusion [2, 12, 4], which again can be corrected via an appropriately modified MLE estimator [8].

While these correction terms can be applied to online learning, we follow an alternative approach in this note, which has been motivated by the related work [1, 14]. Our approach can easily be implemented both for SGD [17] as well as Kalman filter [11, 16] formulations.

As demonstrated in the next section by means of a simple example, a discretised SGD of the form

$$(5a) \quad \theta_{n+1} := \theta_n + \alpha_t F(X_{t_n}^\dagger, \tilde{U}_n)^\top \Delta I_n(\theta_n),$$

$$(5b) \quad \Delta I_n(\theta) := (\tilde{U}_{n+1/2} - \tilde{U}_{n-1/2}) - \left(f(X_{t_n}^\dagger, \tilde{U}_n) + F(X_{t_n}^\dagger, \tilde{U}_n) \theta \right) \tau,$$

with $U_{t_{n+1/2}}$ being replaced by its estimated $\tilde{U}_{n+1/2}$ etc. and the symmetric approximation

$$(6) \quad \tilde{U}_n := \frac{1}{2} \left(\tilde{U}_{n+1/2} + \tilde{U}_{n-1/2} \right) = \frac{X_{t_{n+1}}^\dagger - X_{t_{n-1}}^\dagger}{2\tau}$$

will fail even in the limit $\tau \rightarrow 0$. Here

$$(7) \quad \alpha_t = c_1 / (c_2 + t)$$

denotes the learning rate for appropriate constants $c_i > 0$, $i = 1, 2$.

The discussion from the following section will lead to an appropriate modification of (5), which removes the undesirable bias in the SGD estimator (5). See Section 3.

We extend the proposed methodology to Kalman filter based estimation in Section 4. The unbiasedness of the corrected SGD approach is demonstrated for a nonlinear second-order diffusion process in Section 5. The note closes with comments on further directions for research.

2. MOTIVATING EXAMPLE

In order to demonstrate the difficulties the standard SGD implementation (5) encounters, we consider velocity data U_t^\dagger being generated by the linear OU process

$$(8) \quad dU_t^\dagger = -\theta_* U_t^\dagger dt + \sqrt{\sigma} dW_t^\dagger$$

for given diffusion constant $\sigma > 0$ and unknown drift parameter $\theta_* \geq 0$.

We will set $\theta_* = 0$ for some of the analysis to be conducted later in this section, which implies that solutions of (8) satisfy

$$(9) \quad U_t^\dagger = U_s^\dagger + \sqrt{\sigma} W_{t-s}^\dagger$$

for any $t \geq s$ and $s \geq 0$ and, consequently,

$$(10) \quad X_t^\dagger = X_s^\dagger + (t-s)U_s^\dagger + \sqrt{\sigma} \int_s^t W_{r-s}^\dagger dr.$$

Here we have introduced the shorthand $W_{t-s}^\dagger = W_t^\dagger - W_s^\dagger$.

We first recall some of the properties of SGD when a solution of (8) has in fact been fully observed. An application of SGD [17] to this problem yields

$$(11a) \quad d\theta_t = -\alpha_t U_t^\dagger dI_t = -\alpha_t U_t^\dagger \left(dU_t^\dagger + \theta_t U_t^\dagger dt \right)$$

$$(11b) \quad = -\alpha_t U_t^\dagger \left((\theta_t - \theta_*) U_t^\dagger dt + \sqrt{\sigma} dW_t^\dagger \right).$$

We call

$$(12) \quad I_t = \int_0^t \left(dU_s^\dagger + \theta_s U_s^\dagger ds \right) = \int_0^t \left((\theta_s - \theta_*) U_s^\dagger dt + \sqrt{\sigma} dW_s^\dagger \right)$$

the innovation and $\alpha_t U_t^\dagger$ the gain. It is obvious that I_t becomes a martingale at the optimal parameter choice $\theta_s = \theta_*$, $s \in [0, t]$.

It has been shown that $\theta_t \rightarrow \theta_*$ as $t \rightarrow \infty$ [17]. In our context, a key observation is that the integral

$$(13) \quad \int_0^t \alpha_s U_s^\dagger \left(dU_s^\dagger + \theta_s U_s^\dagger ds \right)$$

also becomes a martingale for any $t > 0$ if and only if $\theta_s = \theta_*$ for all $s \in [0, T]$.

In terms of numerical implementations of the SGD formulation (11), we consider the discrete time SGD method

$$(14) \quad \theta_{n+1} := \theta_n - \alpha_{t_n} U_{t_n}^\dagger \left(U_{t_{n+1}}^\dagger - U_{t_n}^\dagger + \frac{\theta_n}{2} \left(U_{t_{n+1}}^\dagger + U_{t_n}^\dagger \right) \tau \right).$$

This discrete-time SGD method still implies that θ_n converges to θ_* as $n \rightarrow \infty$ and $\tau \rightarrow 0$. More precisely, any finite sum

$$(15) \quad \sum_{n=0}^{N-1} U_{t_n}^\dagger \left(U_{t_{n+1}}^\dagger - U_{t_n}^\dagger + \frac{\theta_n}{2} \left(U_{t_{n+1}}^\dagger + U_{t_n}^\dagger \right) \tau \right)$$

becomes a martingale as $\tau \rightarrow 0$ for $T = N\tau$ fixed if and only if $\theta_n = \theta_*$.

However, the focus of this note is on the case when only positions X_t^\dagger , $t \geq 0$, can be observed and we recall the velocity approximations (4), where the sampling interval $\tau > 0$

can be made arbitrarily small. We note that (4) can also be viewed as a local time average, i.e.,

$$(16) \quad \tilde{U}_{n+1/2} = \frac{1}{\tau} \int_{t_n}^{t_{n+1}} U_t^\dagger dt,$$

which provides a link to the related work [9, 7].

Using (6) instead of the true U_{t_n} etc. lead to the previously stated (5), which reduces here to

$$(17a) \quad \theta_{n+1} := \theta_n - \alpha_{t_n} \tilde{U}_n \left(\tilde{U}_{n+1/2} - \tilde{U}_{n-1/2} + \theta_n \tilde{U}_n \tau \right)$$

$$(17b) \quad = \theta_n - \frac{\alpha_{t_n}}{2} \left(X_{t_{n+1}}^\dagger - X_{t_{n-1}}^\dagger \right) \left(\frac{X_{t_{n+1}}^\dagger - 2X_{t_n}^\dagger + X_{t_{n-1}}^\dagger}{\tau^2} + \frac{\theta_n}{2} \frac{X_{t_{n+1}}^\dagger - X_{t_{n-1}}^\dagger}{\tau} \right).$$

We note that, while the discrete innovations

$$(18) \quad I_N = \sum_{n=1}^{N-1} \left(\frac{X_{t_{n+1}}^\dagger - 2X_{t_n}^\dagger + X_{t_{n-1}}^\dagger}{\tau^2} + \frac{\theta_n}{2} \frac{X_{t_{n+1}}^\dagger - X_{t_{n-1}}^\dagger}{\tau} \right) \tau$$

still converge to a martingale as $\tau \rightarrow 0$ for $T = N\tau$ fixed provided $\theta_n = \theta_*$ for all $n \geq 0$, this is no longer true for

$$(19) \quad \frac{1}{2} \sum_{n=1}^{N-1} \left(X_{t_{n+1}}^\dagger - X_{t_{n-1}}^\dagger \right) \left(\frac{X_{t_{n+1}}^\dagger - 2X_{t_n}^\dagger + X_{t_{n-1}}^\dagger}{\tau^2} + \frac{\theta_n}{2} \frac{X_{t_{n+1}}^\dagger - X_{t_{n-1}}^\dagger}{\tau} \right).$$

This violation of the martingale property is perhaps not surprising since (19) resembles a discretised Stratonovitch integral [13]. We note that such an interpretation as a discretised stochastic integral concerns only the second-order differencing term in (19) and we may therefore set $\theta = \theta_* = 0$ in the subsequent derivations. Recall that

$$(20) \quad \frac{X_{t_{n+1}} - X_{t_n}}{\tau} = \tilde{U}_{n+1/2} = U_{t_n}^\dagger + \frac{\sqrt{\sigma}}{\tau} \int_{t_n}^{t_{n+1}} W_{t-t_n}^\dagger dt$$

and we therefore obtain

$$(21a) \quad \frac{1}{2} \mathbb{E} \left[\left(\tilde{U}_{n+1/2} + \tilde{U}_{n-1/2} \right) \left(\tilde{U}_{n+1/2} - \tilde{U}_{n-1/2} \right) \right] = \frac{1}{2} \mathbb{E} \left[\tilde{U}_{n+1/2}^2 - \tilde{U}_{n-1/2}^2 \right]$$

$$(21b) \quad = \frac{1}{2} \mathbb{E} \left[(U_{t_n}^\dagger)^2 - (U_{t_{n-1}}^\dagger)^2 \right] = \frac{\tau\sigma}{2},$$

which implies an estimation bias induced by the discrete sum (19) even in the limit $\tau \rightarrow 0$.

In order to remove the resulting bias in the SGD formulation (17), we propose to simply shift the innovation term forward in time such that it does no longer overlap with the gain; thus regaining an Itô-type stochastic integral approximation. A related idea has been proposed and investigated in [14] in the context of multi-scale diffusion. This modification leads to the following implementation:

$$(22) \quad \theta_{n+1} := \theta_n + \alpha_{t_n} \tilde{U}_n \left(\tilde{U}_{n+5/2} - \tilde{U}_{n+3/2} - \theta_n \tilde{U}_{n+2\tau} \right).$$

Let us introduce the random variables

$$(23) \quad \tilde{\Xi}_n := \frac{1}{\sqrt{\sigma\tau}} \left(\tilde{U}_{n+1/2} - \tilde{U}_{n-1/2} \right)$$

for $n \geq 1$, which is a rescaling of the innovation $\Delta I_n(\theta_*)$, as defined in (5b), with θ_* again set to $\theta_* = 0$. It is crucial to observe that $\tilde{\Xi}_{n+2}$ is correlated with both $\tilde{U}_{n+5/2}$ and $\tilde{U}_{n+3/2}$, while being independent of \tilde{U}_n . Hence, provided $\tilde{\Xi}_n$ is a martingale for all $n \geq 1$, the SGD scheme (22) will become unbiased in the limit $\tau \rightarrow 0$.

Hence let us analyse the statistical properties of $\tilde{\Xi}_n$. We find that

$$(24a) \quad \tilde{\Xi}_n = \frac{1}{\sqrt{\sigma\tau}} \left(\tilde{U}_{n+1/2} - \tilde{U}_{n-1/2} \right)$$

$$(24b) \quad = \frac{1}{\sqrt{\sigma\tau}} \left(U_{t_n}^\dagger - U_{t_{n-1}}^\dagger \right) + \frac{1}{\sqrt{\tau}} \left(\frac{\int_{t_n}^{t_{n+1}} W_{t-t_n}^\dagger dt - \int_{t_{n-1}}^{t_n} W_{t-t_{n-1}}^\dagger dt}{\tau} \right)$$

$$(24c) \quad = \frac{W_{t_n-t_{n-1}}^\dagger}{\sqrt{\tau}} + \frac{1}{\sqrt{\tau}} \left(\frac{\int_{t_n}^{t_{n+1}} W_{t-t_n}^\dagger dt - \int_{t_{n-1}}^{t_n} W_{t-t_{n-1}}^\dagger dt}{\tau} \right).$$

It follows that $\tilde{\Xi}_n$ is Gaussian with mean zero and that $\tilde{\Xi}_{n+2}$ is indeed independent of \tilde{U}_n .

We recall the following two properties of integrated Brownian motion:

$$(25) \quad \text{var} \left(\int_{t_n}^{t_{n+1}} W_{t-t_n}^\dagger dt \right) = \frac{1}{3} \tau^3.$$

and

$$(26) \quad \mathbb{E} \left[W_{t_n-t_{n-1}}^\dagger \left(\int_{t_{n-1}}^{t_n} W_{t-t_{n-1}}^\dagger dt \right) \right] = \int_{t_{n-1}}^{t_n} \mathbb{E} \left[(W_{t-t_{n-1}}^\dagger)^2 \right] dt = \frac{1}{2} \tau^2.$$

Upon applying these results to (24c) when computing the variance of $\tilde{\Xi}_n$, it is found that

$$(27) \quad \text{var}(\tilde{\Xi}_n) = \frac{2}{3}.$$

In other words, we obtain that $\tilde{\Xi}_n \sim \text{N}(0, 2/3)$ in agreement with results from [9, 6]. This result carries over to $\theta_* \neq 0$ since the arising additional terms vanish as $\tau \rightarrow 0$.

In line with the previous result [9], our calculations imply the following estimator for the diffusion constant σ :

$$(28) \quad \sigma_N := \frac{3}{2\tau N} \sum_{n=1}^N \|\tilde{U}_{n+1/2} - \tilde{U}_{n-1/2}\|^2.$$

We finally formally investigate the time-continuous limit of the modified discrete-time SGD formulation (22). We first note that

$$(29) \quad \mathbb{E}[\tilde{\Xi}_n \tilde{\Xi}_{n+k}] = \begin{cases} \frac{1}{6} & |k| = 1 \\ 0 & |k| > 1 \end{cases},$$

which implies that $\sqrt{\tau}\tilde{\Xi}_n$ converges to standard Brownian motion in the limit $\tau \rightarrow 0$. When using this result in the SGD formulation (22) for general θ_* , we obtain

$$(30a) \quad \theta_{n+1} \approx \theta_n - \alpha_{t_n} \tilde{U}_n \left((\theta_n - \theta_*) \tilde{U}_{n+2} \tau + \sqrt{\sigma\tau} \tilde{\Xi}_{n+2} \right)$$

$$(30b) \quad \approx \theta_n - \alpha_{t_n} \tilde{U}_n \left((\theta_n - \theta_*) \tilde{U}_n \tau + \sqrt{\sigma\tau} \tilde{\Xi}_{n+2} \right)$$

and the limit $\tau \rightarrow 0$ leads formally to the time-continuous formulation

$$(31) \quad \dot{\theta}_t = -\alpha_t U_t^\dagger \left((\theta - \theta_*) U_t^\dagger + \sqrt{\sigma} \dot{W}_t^\dagger \right),$$

which coincides with (11b).

3. BIAS-CORRECTED SGD

The analysis from Section 2 leads us to the following modification of the discrete-time SGD implementation (5):

$$(32) \quad \theta_{n+1} = \theta_n + \alpha_{t_n} F(X_{t_n}^\dagger, \tilde{U}_n)^\top \Delta I_{n+2}(\theta_n),$$

for $n \geq 1$ with the innovations $\Delta I_n(\theta)$ defined by (5b) and the learning rate α_{t_n} by (7) with $t = t_n = n\tau$. We recall that the key point of shifting the innovation forward in time is to retain an Itô-type approximation in the gain times innovation stochastic integral approximation. This is the property that maintains the desired unbiasedness of the resulting sequential learning scheme in the limit $\tau \rightarrow 0$.

We note that unbiased formulations are not unique. For example, (32) could be replaced by

$$(33) \quad \theta_{n+1} = \theta_n + \alpha_{t_n} F(X_{t_n}^\dagger, \tilde{U}_n)^\top \left(\tilde{U}_{n+5/2} - \tilde{U}_{n+3/2} - \left(f(X_{t_n}^\dagger, \tilde{U}_n) + F(X_{t_n}^\dagger, \tilde{U}_n) \theta_n \right) \tau \right).$$

This freedom is also utilised when formulating the modified MLE estimator (37) further below in Section 6.

4. BIAS-CORRECTED KALMAN FILTER

The SGD method (32) provides an online point estimator for the parameter θ . However, it is also possible to implement an online Bayesian estimator using the Kalman filter [11, 16]. The key idea is to treat the unknown parameter as a Gaussian random variable Θ_n with mean m_n and covariance Σ_n , $n \geq 0$. The evolution equation for the mean and the covariance follow the standard Kalman update formulas:

$$(34a) \quad m_{n+1} := m_n + K_n \frac{\Delta I_{n+2}(m_n)}{\tau},$$

$$(34b) \quad \Sigma_{n+1} := \Sigma_n - K_n F(X_{t_n}^\dagger, \tilde{U}_n) \Sigma_n$$

with Kalman gain

$$(35) \quad K_n := \Sigma_n F(X_{t_n}^\dagger, \tilde{U}_n)^\top \left(F(X_{t_n}^\dagger, \tilde{U}_n)^\top \Sigma_n F(X_{t_n}^\dagger, \tilde{U}_n) + \frac{\sigma}{\tau} I \right)^{-1}$$

for $n \geq 1$. The iteration is initialised from the Gaussian prior $\Theta_1 \sim \mathcal{N}(m_{\text{prior}}, \Sigma_{\text{prior}})$.

In contrast to the SGD formulation (32), the learning rate α_{t_n} no longer appears and has instead been submerged implicitly into the Kalman gain K_n . Indeed, the singular values of the covariance matrix Σ_n decay with rate $1/t_n$ under suitable ergodicity assumptions on (2). If the required diffusion constant σ is unknown, it can either be estimated via (28) or, alternatively, be treated as part of the prior covariance matrix Σ_{prior} and then set to $\sigma = 1$ in the Kalman gain (35).

5. NUMERICAL EXAMPLE

We consider the nonlinear second-order SDEs

$$(36a) \quad dU_t = -X_t dt + U_t dt - \theta U_t^3 dt + \sqrt{2} dW_t,$$

$$(36b) \quad dX_t = U_t dt$$

with unknown parameter $\theta \in \mathbb{R}$. The data X_t^\dagger is generated with $\theta_* = 1$, while the SGD iteration starts from $\theta_0 = 2$. The sample interval is set to $\tau = 0.025$. The data is generated with a much smaller step-size of $h = \tau/100$. A total of $N = 10^5$ positions X_{t_n} , $n = 1, \dots, N$ have been generated. The learning rate is given by $\alpha_{t_n} = 0.2/n$. See Figure 5 for the time evolved estimates θ_n , which clearly converge to θ_* . We also display results from the standard SGD in Figure 5. While the standard SGD converges slightly faster, it converges to a wrong parameter value.




FIGURE 1. Left panel: Estimated parameter value from standard SGD as a function of the iteration index. The iteration starts at $\theta_0 = 2$ and the reference value is $\theta_* = 1$. A systematic bias can clearly be seen. Right panel: Same data is used in the modified SGD. The modified SGD algorithm converges to the correct reference value.

6. CONCLUSIONS

In this note, we have proposed a simple-to-implement modification of standard SGD and Kalman filter formulations for online parameter estimation from position data only. The essential idea is to establish an Itô-like modification that re-establishes the martingale property in the presence of temporal correlations. The approach has been inspired by related work [1, 14] on multi-scale diffusion and parameter estimation for the arising reduced equations. We finally note that a corrected MLE estimator can be based on the negative log-likelihood function

$$(37a) \quad \mathcal{L}(\theta, \sigma) = \frac{1}{2\sigma} \sum_{n=1}^{N-3} \theta^T F(X_{t_n}^\dagger, \tilde{U}_n)^T F(X_{t_n}^\dagger, \tilde{U}_n) \theta \tau$$

$$(37b) \quad - \frac{1}{\sigma} \sum_{n=1}^{N-3} \theta^T F(X_{t_n}^\dagger, \tilde{U}_n)^T \left(\tilde{U}_{n+5/2} - \tilde{U}_{n+3/2} - f(X_{t_n}^\dagger, \tilde{U}_n) \tau \right)$$

$$(37c) \quad + \frac{3}{4\tau\sigma} \sum_{n=1}^{N-3} \|\tilde{U}_{n+5/2} - \tilde{U}_{n+3/2}\|^2 + \frac{N-3}{2} \log \sigma.$$

Future directions include the investigation of the proposed estimators in the context multi-scale diffusion when (1) arises from homogenisation of an underlying multi-scale diffusion process and the inclusion of observation errors in the positions X_t . In the former case, a combination of the approaches taken here and in [14] emerges as a feasible starting point.

Acknowledgements. This work has been partially funded by Deutsche Forschungsgemeinschaft (DFG) - Project-ID 318763901 - SFB1294.

REFERENCES

- [1] Assyr Abdulle, Giacomo Garegnani, Grigorios Pavliotis, Andrew Stuart, and Andrea Zanoni. Drift estimation of multiscale diffusions based on filtered data. *Foundations of Computational Mathematics*, 23:33–84, 2023. doi:10.1007/s10208-021-09541-9.
- [2] Yacine Ait-Sahalia, Per A Mykland, and Lan Zhang. How often to sample a continuous-time process in the presence of market microstructure noise. *The Review of Financial Studies*, 18:351–416, 2005.
- [3] Jan Albrecht, Carsten Beta, Robert Großmann, and Manfred Opper. Heterogeneity inference for second order Langevin models, 2024. in preparation.
- [4] S. Bo and A. Celani. White-noise limit of nonwhite nonequilibrium processes. *Physical Review E*, 88:062150, 2013. doi:10.1103/PhysRevE.88.062150.
- [5] D.B. Brückner, P. Ronceray, and C.P. Broedersz. Inferring the dynamics of underdamped stochastic systems. *Phy. Rev. Lett.*, 125:28005, 2010.
- [6] I. Cialenco, H.-J. Kim, and G. Pasemann. Statistical analysis of discretely sampled semilinear SPDEs: A power variation approach. *Stochastics and Partial Differential Equations: Analysis and Computations*, 12:326–351, 2024.

- [7] Igor Cialenco and Y. Huang. A note on parameter estimation for discretely sampled SPDEs. *Stoch. Dyn.*, 20:2050016, 2020.
- [8] J. Diehl, P. Friz, and H. Mai. Pathwise stability of likelihood estimators for diffusion via rough paths. *The Annals of Applied Probability*, 26:2169–2192, 2016. doi:10.1214/15-AAP1143.
- [9] Arnaud Gloter. Parameter estimation for a discretely observed integrated diffusion process. *Scandinavian Journal of Statistics*, 33:83–104, 2006.
- [10] Yury A Kutoyants. *Statistical inference for ergodic diffusion processes*. Springer Science & Business Media, 2013.
- [11] Nikolas Nüsken, Sebastian Reich, and Paul J Rozdeba. State and parameter estimation from observed signal increments. *Entropy*, 21(5):505, 2019. doi:10.3390/e21050505.
- [12] A. Papavasiliou, G. Pavliotis, and A.M. Stuart. Maximum likelihood estimation for multiscale diffusions. *Stochastic Processes and their Applications*, 19:3173–3210, 2009.
- [13] G.A. Pavliotis. *Stochastic Processes and Applications*. Springer Verlag, New York, 2016.
- [14] Grigorios A Pavliotis, Sebastian Reich, and Andrea Zanoni. Filtered data based estimators for stochastic processes driven by colored noise, 2023. arXiv:2312.15975.
- [15] Jonas N. Pedersen, Liang Li, Cristian Gradinaru, Robert H. Austin, Edward C. Cox, , and Henrik Flyvbjerg. How to connect time-lapse recorded trajectories of motile microorganisms with dynamical models in continuous time. *Physical Review E*, 94:062401, 2016.
- [16] Sebastian Reich. Frequentist perspective on robust parameter estimation using the ensemble Kalman filter. In B. Chapron, D. Crisan, D. Holm, E. Mémin, and A. Radomska, editors, *Stochastic Transport in Upper Ocean Dynamics. STUOD 2021. Mathematics of Planet Earth*, volume 10, pages 237–258, Cham., 2023. Springer. doi:10.1007/978-3-031-18988-3_15.
- [17] J. Sirignano and K. Spiliopoulos. Stochastic gradient descent in continuous time. *SIAM J. Financial Math.*, 8:933–961, 2017. doi:10.1137/17M1126825.