

THE DISCRETE DISLOCATION DYNAMICS OF MULTIPLE DISLOCATION LOOPS

STEFANIA PATRIZI AND MARY VAUGHAN

ABSTRACT. We consider a nonlocal reaction-diffusion equation that physically arises from the classical Peierls–Nabarro model for dislocations in crystalline structures. Our initial configuration corresponds to multiple slip loop dislocations in \mathbb{R}^n , $n \geq 2$. After suitably rescaling the equation with a small phase parameter $\varepsilon > 0$, the rescaled solution solves a fractional Allen–Cahn equation. We show that, as $\varepsilon \rightarrow 0$, the limiting solution exhibits multiple interfaces evolving independently and according to their mean curvature.

1. INTRODUCTION

We study the fractional Allen–Cahn equation

$$(1.1) \quad \varepsilon \partial_t u^\varepsilon = \frac{1}{\varepsilon |\ln \varepsilon|} (\varepsilon \mathcal{I}_n[u^\varepsilon] - W'(u^\varepsilon)) \quad \text{in } (0, \infty) \times \mathbb{R}^n, \quad n \geq 2,$$

where $\varepsilon > 0$ is a small parameter, $\mathcal{I}_n = -c_n(-\Delta)^{\frac{1}{2}}$ denotes up to a constant the square root of the Laplacian in \mathbb{R}^n , and W is a multi-well potential, see (1.2) and (1.3) respectively.

Equation (1.1) is a rescaled version of the evolutionary Peierls–Nabarro model for atomic dislocations in crystalline structures. In this article, we initiate the mathematical analysis of the evolution of multiple slip dislocations according to (1.1). Our initial configuration corresponds to multiple slip loops, namely simple closed curves embedded in the same plane, with each loop enclosing the next within its confines. We show that, as $\varepsilon \rightarrow 0$, the curves move independently and according to their mean curvature. To the best of our knowledge, we are the first to consider the movement of multiple fronts in \mathbb{R}^n , $n \geq 2$, (the physical dimension being $n = 2$) in this setting. Furthermore, our results are new even for the case of a single curve and completes the work of Imbert and Souganidis in [29].

Before further discussing the Peierls–Nabarro model for slip loop dislocations, let us formalize our problem mathematically.

1.1. Setting of the problem and main result. The operator \mathcal{I}_n is a nonlocal integro-differential operator of order 1 and is given by

$$(1.2) \quad \mathcal{I}_n u(x) = \text{P. V.} \int_{\mathbb{R}^n} (u(x+y) - u(x)) \frac{dy}{|y|^{n+1}}, \quad x \in \mathbb{R}^n, \quad n \in \mathbb{N},$$

2010 *Mathematics Subject Classification.* Primary: 82D25, 35R09, 35R11. Secondary: 74E15, 47G20.

Key words and phrases. Peierls–Nabarro model, nonlocal integro-differential equations, dislocation dynamics, fractional Allen–Cahn, phase transitions.

where P.V. indicates that the integral is taken in the principal value sense. For further background on fractional Laplacians, see for example [14, 49]. The potential W satisfies

$$(1.3) \quad \begin{cases} W \in C^{4,\beta}(\mathbb{R}) & \text{for some } 0 < \beta < 1 \\ W(u+1) = W(u) & \text{for any } u \in \mathbb{R} \\ W = 0 & \text{on } \mathbb{Z} \\ W > 0 & \text{on } \mathbb{R} \setminus \mathbb{Z} \\ W''(0) > 0. \end{cases}$$

We let u^ε be the solution to (1.1) when the initial condition u_0^ε is a superposition of layer solutions. The layer solution (also called the phase transition) $\phi : \mathbb{R} \rightarrow (0, 1)$ is the unique solution to

$$(1.4) \quad \begin{cases} C_n \mathcal{I}_1[\phi] = W'(\phi) & \text{in } \mathbb{R} \\ \dot{\phi} > 0 & \text{in } \mathbb{R} \\ \phi(-\infty) = 0, \quad \phi(+\infty) = 1, \quad \phi(0) = \frac{1}{2}, \end{cases}$$

where \mathcal{I}_1 is the fractional operator in (1.2) on \mathbb{R} and the constant $C_n > 0$ (given explicitly in (3.1)) depends only on $n \geq 2$. Further discussion on ϕ will be presented in Section 4.

For a fixed $N \in \mathbb{N}$, let $(\Omega_0^i)_{i=1}^N$ be a finite sequence of open subsets of \mathbb{R}^n that are bounded, satisfy $\Omega_0^{i+1} \subset \subset \Omega_0^i$ and have smooth boundaries $\Gamma_0^i = \partial\Omega_0^i$. When $n = 2$, the curves Γ_0^i can be understood as the initial dislocation loops in the crystal. Let $d_i^0(x)$ be the signed distance function associated to Ω_0^i , $i = 1, \dots, N$, given by

$$(1.5) \quad d_i^0(x) = \begin{cases} d(x, \Gamma_0^i) & \text{if } x \in \Omega_0^i \\ -d(x, \Gamma_0^i) & \text{otherwise.} \end{cases}$$

For our initial condition to be well-prepared, we let u_0^ε be the N -fold sum of the layer solutions $\phi(d_i^0(x)/\varepsilon)$, see (1.7) and Figure 1.




FIGURE 1. Initial configuration for $N = 3$ in dimension $n = 2$

We show that the evolution of the surfaces Γ_0^i , denoted by $(\Gamma_t^i)_{t \geq 0}$, corresponds to interfaces moving independently by mean curvature. Specifically, $(\Gamma_t^i)_{t \geq 0}$ move in the direction of the interior normal vector with scalar velocity

$$(1.6) \quad v = \mu \sum_{j=1}^{n-1} \kappa_j, \quad \mu := \frac{c_0 |S^{n-2}|}{2(n-1)} > 0,$$

where κ_j are the principal curvatures and $c_0 > 0$ is explicit (see (4.1)). In dimension 2, Γ_t^i can be interpreted as the dislocation loop at time $t > 0$ evolved from Γ_0^i . To handle possible

singularities for large times $t > 0$, we use the level set approach. Roughly speaking, consider a solution $u^i(t, x)$ to the mean curvature equation (see (2.2) with μ as in (1.6)) whose positive, zero and negative sets at time $t = 0$ respectively are Ω_0^i, Γ_0^i and $(\overline{\Omega_0^i})^c$. If ${}^+\Omega_t^i, \Gamma_t^i$, and ${}^-\Omega_t^i$ respectively are the positive, zero, and negative sets of $u^i(t, \cdot)$ at time $t > 0$, then we say that the triplet $({}^+\Omega_t^i, \Gamma_t^i, {}^-\Omega_t^i)_{t \geq 0}$ denotes the level set evolution of $(\Omega_0^i, \Gamma_0^i, (\overline{\Omega_0^i})^c)$. Since $\Omega_0^{i+1} \subset\subset \Omega_0^i$, one has that ${}^+\Omega_t^{i+1} \subset\subset {}^+\Omega_t^i$ and ${}^-\Omega_t^i \subset\subset {}^-\Omega_t^{i+1}$, for all times $t > 0$ such that ${}^+\Omega_t^i \neq \emptyset$. See Section 2 for definitions and details on the level set approach to motion by mean curvature.

We now present the main result of our paper.

Theorem 1.1. *Let $u^\varepsilon = u^\varepsilon(t, x)$ be the unique solution of the reaction-diffusion equation (1.1) with the initial datum $u_0^\varepsilon : \mathbb{R}^n \rightarrow (0, N)$ defined by*

$$(1.7) \quad u_0^\varepsilon(x) = \sum_{i=1}^N \phi\left(\frac{d_i^0(x)}{\varepsilon}\right),$$

where ϕ solves (1.4), d_i^0 are given in (1.5), and $N \geq 1$. Then, as $\varepsilon \rightarrow 0$, the solutions u^ε satisfy

$$\begin{cases} u^\varepsilon \rightarrow N & \text{in } {}^+\Omega_t^N, \\ u^\varepsilon \rightarrow i & \text{in } {}^+\Omega_t^i \cap {}^-\Omega_t^{i+1}, \quad i = 1, \dots, N-1, \\ u^\varepsilon \rightarrow 0 & \text{in } {}^-\Omega_t^1, \end{cases}$$

where $({}^+\Omega_t^i, \Gamma_t^i, {}^-\Omega_t^i)$ denotes the level set evolution of $(\Omega_0^i, \Gamma_0^i, (\overline{\Omega_0^i})^c)$ with velocity (1.6).




FIGURE 2. Convergence result for $N = 3$ in dimension $n = 2$

As illustrated in Figure 2, Theorem 1.1 says that the solutions u^ε converge to integers “between” the interfaces Γ_t^i . For the case in which the sets Γ_t^i do not develop interior, i.e. $\Gamma_t^i = \partial({}^+\Omega_t^i) = \partial({}^-\Omega_t^i)$, the limiting function in Theorem 1.1 makes integer jumps on the surfaces Γ_t^i and satisfies

$$\lim_{\varepsilon \rightarrow 0} u^\varepsilon = \frac{N}{2} + \frac{1}{2} \sum_{i=1}^N \left(\mathbb{1}_{{}^+\Omega_t^i} - \mathbb{1}_{(\overline{{}^+\Omega_t^i})^c} \right) \quad \text{in } ((0, \infty) \times \mathbb{R}^n) \setminus \bigcup_{i=1}^N \left(\bigcup_{t>0} \{t\} \times \Gamma_t^i \right)$$

where $\mathbb{1}_\Omega$ denotes the characteristic function of the set $\Omega \subset \mathbb{R}^n$. However, it is well known that Γ_t^i may develop interior in finite time, even if Γ_0^i has none. In this situation, the discontinuity

set at time t of the limiting function is contained in the sets Γ_t^i , but we cannot say exactly where the jumps occur within these sets.

Theorem 1.1 for $N = 1$ has been addressed in the literature in the local case when W is instead a double-well potential. For instance, the classical Allen–Cahn equation for which (1.1) is instead driven by the usual Laplacian Δ was studied famously by Modica–Mortola [32] for the stationary case. Chen studied the corresponding evolutionary Allen–Cahn problem and proved that the solution exhibits an interface moving by mean curvature [9]. See Section 2 for more on the phase field theory. In the fractional setting and again for a double-well potential, the stationary case was studied by Ambrosio–De Philippis–Martinazzi [1] and Savin–Valdinoci [47] and the evolution problem was considered by Imbert–Souganidis in the preprint [29]. Our result is not included in and does not follow from [29, 47]. Indeed, our problem deals with $N \geq 1$ interfaces and a multi-well potential.

The proof of Theorem 1.1 relies on the abstract method introduced in [4, 6] for the study of front propagation. For this, we construct new barriers that are strict sub/super solutions to (1.1), see Section 6. We found that a superposition of the corresponding barriers in [29] cannot be applied as one cannot control the errors associated with the nonlinear potential. Loosely speaking, our barriers are a superposition of layer solutions $\phi(d_i(t, x)/\varepsilon)$ where d_i is the signed distance function associated to the front Γ_t^i . Since the equation is nonlocal, an added difficulty arises in dealing with $d_i(t, x)$ when (t, x) is far from the front since d_i may not be smooth at such points.

In the actual construction of the barriers, we add lower-order correctors to control the error as $\varepsilon \rightarrow 0$. We use a superposition of solutions $\psi = \psi_\varepsilon^i$ to the linearized equation

$$(1.8) \quad -C_n \mathcal{I}_1[\psi] + W''(\phi)\psi = g$$

for some right-hand side $g = g_\varepsilon^i$ depending on $\varepsilon > 0$ and on the signed distance function d_i . The explicit form of g is somewhat technical and is given in Section 4. In [29], they *assume* the existence and uniqueness of correctors with derivatives bounded independently of ε . It was left as an open question whether such correctors exist. In this paper, we prove both existence and uniqueness of the solution ψ . In addition, we have found, via delicate analysis, that $\psi = \psi_\varepsilon^i$ can only be bounded independently of $\varepsilon > 0$ near the front Γ_t^i . In general, it holds that $\psi = O(\varepsilon^{-\frac{1}{2}} |\ln \varepsilon|^{-1})$. See Theorem 4.8 for detailed properties of ψ which are sufficient to prove Theorem 1.1.

A further novelty of our work is the addition of two heuristical derivations in Section 5. First, we provide heuristics for the evolution of the fronts Γ_t^i by mean curvature in Theorem 1.1 in order to give an intuitive explanation of the behavior of the fronts. Second, we write a heuristical derivation for the equation of the corrector (1.8) which clarifies the choice of g .

1.2. The Peierls–Nabarro model for slip loop dislocations. We now describe how (1.1) with initial data (1.7) arises from the Peierls–Nabarro model for slip loop dislocations.

Dislocations are line defects in crystalline materials whose motion is responsible for the plastic behavior of metals (see [25, 26] for more details). The dynamics of an *ensemble of dislocation lines all contained in the same plane* (the slip plane) can be described at different scales by different models:

- atomic scale (Frenkel–Kontorova model),
- microscopic scale (Peierls–Nabarro model),
- mesoscopic scale (Discrete dislocation dynamics),
- macroscopic scale (Elasto-visco-plasticity with density of dislocation).

Several changes of scales exist in the literature. The passage from the Frenkel–Kontorova model to the Peierls–Nabarro model is performed in [17]. In the evolutive models, further changes of scales are only known when the dislocations are *straight*, parallel lines (i.e. one-dimensional models): Gonzalez–Monneau [23] passed from the Peierls–Nabarro model to the discrete dislocation dynamics (see also [31, 41, 43, 44] for the case of dislocations with different orientation); the passage from the discrete dislocation dynamics to the elasto-visco-plasticity with density of dislocation is done in [18, 30]; finally, a direct passage from the Peierls–Nabarro model to the elasto-visco-plasticity with density of dislocation is performed in [39, 40]. We reference the reader to [15, 24] and the references therein for more on the dynamics of edge dislocations in one-dimension.

For different models for straight dislocation lines, we refer to [2, 3, 19, 20] and the references therein.

In reality, dislocation lines are rarely straight and often form loops, see [48]. The dynamics of an ensemble of *curved* dislocation lines lying in the same slip plane is described by the above models in dimension $n = 2$. Our main result, Theorem 1.1, is a passage from the Peierls–Nabarro model to the discrete dislocation dynamics in any dimension $n \geq 2$. Differently from the one-dimensional case where lines move according to an interaction potential (see [23]), here the line tension effect is stronger than the interaction between the curves, and we have a movement by mean curvature as established in Theorem 1.1. See also the work of Garroni–Müller [21] and Garroni–Conti–Muller [11] for variational Peierls–Nabarro models in dimension two. We also mention that Monneau and the first author in [35] studied homogenization of the Peierls–Nabarro model in any dimension to find an effective plastic law for density of dislocations at the macroscopic scale. See [28, 42] for related results.

We next briefly review the Peierls–Nabarro model for a dislocation loop. For the original model, we refer to [36, 37, 46]. A slip dislocation occurs when a portion of atoms slides over another along a particular plane, called the slip plane, and is the curve created by the boundary between the shifted and unshifted regions. In Cartesian coordinates $x_1x_2x_3$, we assume that the slip plane is the x_1x_2 -plane. The movement of the dislocation is determined by the so-called Burgers’ vector b . For slip dislocations, the Burgers’ vector is contained within the slip plane, so we assume it to be in the direction of the x_1 -axis, say $b = e_1$.

There are two types of slip dislocations that correspond to straight lines in \mathbb{R}^2 : edge and screw. An edge dislocation is formed when an extra half-plane of atoms is included in the crystal. In this case, the Burgers’ vector is perpendicular to the dislocation line and the motion of the dislocation line is in the direction of b . A screw dislocation instead creates a spiral or helical path around the core. Here, the Burgers’ vector is parallel to the dislocation curve which means that the dislocation line moves parallel to the direction of b . It is probable that most slip dislocations are not straight lines and thus exhibit components of both edge and screw dislocations. These are called mixed dislocations. *Slip loops* (or loop dislocations) are slip dislocations in the form of simple closed curves. Since the Burgers’ vector is fixed for a given slip loop, the type of dislocation changes from point to point along the dislocation curve, see Figure 3. Depending on the orientation of the curve with respect to b , the slip loop will either shrink or expand.

The Peierls–Nabarro model is a phase field model for dislocation dynamics which incorporates the atomic features of a crystalline structure into continuum framework. In the phase field approach, dislocations are interfaces represented by a transition of a continuous field: a phase parameter $\Phi(x_1, x_2)$ between 0 and 1 is used to capture the disregistry of the upper half crystal $\{x_3 > 0\}$ from the lower half crystal $\{x_3 < 0\}$. In particular, the dislocation loop




FIGURE 3. Dislocation types in a slip loop dislocation with fixed Burgers' vector

is the set $\Gamma_0 := \{\Phi = 1/2\} \subset \mathbb{R}^2$, and the phase parameter satisfies $\Phi \simeq 1$ inside the loop and $\Phi \simeq 0$ outside the loop (as indicated in Figure 3).

Next, let $U = U(x_1, x_2, x_3)$ be the distance between an atom at location (x_1, x_2, x_3) in the upper half crystal and its rest position. Let $\Phi(x_1, x_2) = U(x_1, x_2, 0)$ be the displacement on the slip plane. Most of the mismatch of atoms from their perfect lattice structure occurs within the slip plane. To quantify this, we use a multi-well potential W satisfying (1.3). Here, the periodicity of W reflects the periodicity of the crystal while the minimum property on \mathbb{Z} captures how the energy penalizes atoms which are not aligned with the lattice structure. In the Peierls–Nabarro model, the total energy is the elastic energy for bonds between atoms plus the so-called misfit energy due to the nonlinear atomic interaction across the slip plane:

$$\mathcal{E}(U) = \frac{1}{2} \int_{\mathbb{R}^2 \times \mathbb{R}^+} |U(x_1, x_2, x_3)|^2 dx_1 dx_2 dx_3 + \int_{\mathbb{R}^2} W(\Phi(x_1, x_2)) dx_1 dx_2.$$

The equilibrium configuration is obtained by minimizing the energy \mathcal{E} with respect to U under the constraint that $\Phi \simeq 1$ in the loop and $\Phi \simeq 0$ outside the loop. The corresponding Euler-Lagrange equation is

$$\mathcal{I}_2[\Phi] = W'(\Phi) \quad \text{in } \mathbb{R}^2$$

where \mathcal{I}_2 is the fractional Laplacian of order 1 in \mathbb{R}^2 .

We are interested in the evolution of multiple loop dislocations in the same slip plane, with the same Burgers' vector. For this, we use a single parameter $u(t, x)$ defined for x in the slip plane \mathbb{R}^n , $n \geq 2$, the physical dimension being $n = 2$. The dislocation dynamics are then captured by the evolutionary Peierls–Nabarro model:

$$\partial_t u = \mathcal{I}_n[u] - W'(u) \quad \text{in } (0, \infty) \times \mathbb{R}^n.$$

At the microscopic scale, we assume that the dislocations curves are at a distance of order $1/\varepsilon$ from each other. This can be represented by the initial condition

$$u(0, x) = \sum_{i=1}^N \phi \left(\frac{d_i^0(\varepsilon x)}{\varepsilon} \right) \quad \text{for } x \in \mathbb{R}^n, \varepsilon > 0,$$

where ϕ solves (1.4) in \mathbb{R} and d_i^0 is the signed distance function associated to the loop Γ_0^i .

In order to understand the movement of the dislocation curves at a larger (mesoscopic) scale, we consider the rescaling

$$u^\varepsilon(t, x) = u\left(\frac{t}{\varepsilon^2 |\ln \varepsilon|}, \frac{x}{\varepsilon}\right), \quad (t, x) \in [0, \infty) \times \mathbb{R}^n, \quad \varepsilon > 0.$$

Here, $\varepsilon > 0$ represents the scaling between the microscopic scale and the mesoscopic scale. The presence of the factor $|\ln \varepsilon|$ is well-known in physics, see [7, 13, 25]. Roughly speaking, it arises from an integrability condition for the kernel of \mathcal{I}_n in (1.2). See also [47]. One can easily check that u^ε solves (1.1) with the initial condition (1.7), so that Theorem 1.1 indeed describes the evolution of the dislocation curves $(\Gamma_t^i)_{t \geq 0}$.

1.3. Organization of the paper. The rest of the paper is organized as follows. First, in Section 2, we provide the necessary background pertaining to motion by mean curvature and the level set approach. Section 3 contains preliminary results on fractional Laplacians. Then, we establish preliminary results pertaining to the phase transition ϕ , the corrector ψ , and other auxiliary functions needed for the rest of the paper in Section 4. Section 5 contains heuristics for the evolution of fronts in Theorem 1.1 and heuristics for the equation for the corrector. The construction of barriers is presented in Section 6. Section 7 contains the proof of Theorem 1.1. Lastly, since the proofs of some of auxiliary results in Section 4 are rather technical, we write them in Sections 8, 9, 10, 11.

1.4. Notations. In the paper, we will denote by $C > 0$ any universal constant depending only on the dimension n and W .

We let $B(x_0, r)$ and $\bar{B}(x_0, r)$ denote respectively the ball of radius $r > 0$ centered at $x_0 \in \mathbb{R}^n$ and its closure, and let S^n denote the unit sphere in \mathbb{R}^{n+1} .

For $\beta \in (0, 1]$ and $k \in \mathbb{N} \cup \{0\}$, we denote by $C^{k, \beta}(\mathbb{R})$ the usual class of functions with bounded $C^{k, \beta}$ norm over \mathbb{R} . The class $H^{\frac{1}{2}}(\mathbb{R})$ is the set of functions $g \in L^2(\mathbb{R})$ such that

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|g(x) - g(y)|^2}{|x - y|^3} dy dx < \infty.$$

For multi-variable functions, we let $C_\xi^{k, \beta}(\mathbb{R})$ and $H_\xi^{\frac{1}{2}}(\mathbb{R})$ denote the classes $C^{k, \beta}(\mathbb{R})$ and $H^{\frac{1}{2}}(\mathbb{R})$ in the variable $\xi \in \mathbb{R}$, respectively.

Given a function $\eta = \eta(t, x)$, defined on a set A , we write $\eta = O(\varepsilon)$ if there is $C > 0$ such that $|\eta(t, x)| \leq C\varepsilon$ for all $(t, x) \in A$, and we write $\eta = o_\varepsilon(1)$ if $\lim_{\varepsilon \rightarrow 0} \eta(t, x) = 0$, uniformly in $(t, x) \in A$. Given a sequence of functions $u^\varepsilon(t, x)$, we write

$$\liminf_{\varepsilon \rightarrow 0} *u^\varepsilon(t, x) = \inf \left\{ \liminf_{\varepsilon \rightarrow 0} u^\varepsilon(t_\varepsilon, x_\varepsilon) : (t_\varepsilon, x_\varepsilon) \rightarrow (t, x) \right\}$$

and

$$\limsup_{\varepsilon \rightarrow 0} *u^\varepsilon(t, x) = \sup \left\{ \limsup_{\varepsilon \rightarrow 0} u^\varepsilon(t_\varepsilon, x_\varepsilon) : (t_\varepsilon, x_\varepsilon) \rightarrow (t, x) \right\}.$$

Given matrices M, N , we denote by $M \otimes N$ the usual tensor product of M and N . For a vector $p \in \mathbb{R}^n$, we write $\hat{p} = p/|p|$.

For a set A , we denote by $\mathbb{1}_A$ the characteristic function of the set A .

2. MOTION BY MEAN CURVATURE

In this section, we present the necessary background and preliminaries regarding the mean curvature of an evolving front.

2.1. The level set approach. Here, we review the level set approach for the geometric motions of the fronts. For a smooth function $u = u(t, x)$, consider a level set $\Gamma_t = \{x \in \mathbb{R}^n : u(t, x) = \ell\}$ of $u(t, \cdot)$ at level $\ell \in \mathbb{R}$ and assume that Γ_t is bounded and ∇u does not vanish on Γ_t . Let $d(t, x)$ denote the signed distance function to $\{u > \ell\}$:

$$d(t, x) = \begin{cases} d(x, \Gamma_t) & \text{for } u(t, x) \geq \ell \\ -d(x, \Gamma_t) & \text{for } u(t, x) < \ell. \end{cases}$$

Note that $n(t, x) := \nabla d(t, x)$ is normal to Γ_t and satisfies $|\nabla d| = 1$ in a time-space neighborhood \mathcal{N} of Γ_t . Then, as theorized by Osher–Sethian [38] and justified by Evans–Spruck in [16], the level set Γ_t moves in the normal direction with velocity $v(t, x)$ proportional to its mean curvature, i.e.,

$$(2.1) \quad \begin{aligned} v(t, x) &= -\mu \operatorname{div}(n(t, x))n(t, x) \\ &= -\mu \Delta d(t, x) \nabla d(t, x) \quad \text{in } \mathcal{N}, \end{aligned}$$

where $\mu > 0$ is a constant, if and only if u is a solution in \mathcal{N} to the *mean curvature equation*

$$(2.2) \quad \partial_t u = \mu \operatorname{tr} \left((I - \widehat{\nabla} u \otimes \widehat{\nabla} u) D^2 u \right).$$

Consequently, if u is smooth and $\nabla u \neq 0$ in an open region $O \subset (0, \infty) \times \mathbb{R}^n$, then u is a solution to (2.2) in O if and only if all its level sets move according to their mean curvature (2.1) in O .

However, level sets that move by mean curvature may develop singularities in finite time. Indeed, observe that equation (2.2) is degenerate, and it is not well-defined when $\nabla u = 0$. To overcome these difficulties and develop a weak notion of evolving fronts, they use in [16] the theory of viscosity solutions (see [12] for more on viscosity solutions). More precisely, for a bounded, open set $\Omega_0 \subset \mathbb{R}^n$, set $\Gamma_0 = \partial\Omega_0$ and consider the initial triplet $(\Omega_0, \Gamma_0, (\overline{\Omega_0})^c)$. Focusing on the zero level set of the solution, let $u_0(x)$ be a continuous function such that

$$\Omega_0 = \{x : u_0(x) > 0\}, \quad \Gamma_0 = \{x : u_0(x) = 0\}, \quad \text{and} \quad (\overline{\Omega_0})^c = \{x : u_0(x) < 0\},$$

and u is a constant outside a ball. Then, there exists a unique continuous viscosity solution u to (2.2) in $(0, \infty) \times \mathbb{R}^n$ with initial datum $u(0, x) = u_0(x)$, and such that u is constant outside a ball of $(0, \infty) \times \mathbb{R}^n$. By our previous discussion, the zero level sets of u move according to their mean curvature, at least in the region where u is smooth and $\nabla u \neq 0$. Set

$$(2.3) \quad {}^+\Omega_t = \{x : u(t, x) > 0\}, \quad \Gamma_t = \{x : u(t, x) = 0\}, \quad \text{and} \quad {}^-\Omega_t = \{x : u(t, x) < 0\}.$$

The triplet $({}^+\Omega_t, \Gamma_t, {}^-\Omega_t)_{t \geq 0}$ is called the *level set evolution* of $(\Omega_0, \Gamma_0, (\overline{\Omega_0})^c)$. As shown in [16], Γ_t does not depend on the particular choice of u_0 but only on Γ_0 .

If $({}^+\tilde{\Omega}_t, \tilde{\Gamma}_t, {}^-\tilde{\Omega}_t)_{t \geq 0}$ denotes the level set evolution of $(\tilde{\Omega}_0, \tilde{\Gamma}_0, (\overline{\tilde{\Omega}_0})^c)$ with $\tilde{\Gamma}_0 = \partial\tilde{\Omega}_0$, and $\Omega_0 \subset\subset \tilde{\Omega}_0$, then ${}^+\Omega_t \subset\subset {}^+\tilde{\Omega}_t$ and ${}^-\tilde{\Omega}_t \subset\subset {}^-\Omega_t$ for all times $t > 0$ such that ${}^+\tilde{\Omega}_t \neq \emptyset$, see [16, Theorem 7.3].

In the special setting in which Γ_t is smooth for $t \in [t_0, t_0 + h]$, the signed distance function d is a solution to (2.2) in $\{(t, x) : t \in [t_0, t_0 + h], x \in \Gamma_t\}$. More generally, if u solves

$$\partial_t u = \mu \operatorname{tr} \left((I - \widehat{\nabla} u \otimes \widehat{\nabla} u) D^2 u \right) + \sigma$$

in a neighborhood of Γ_t for some $\sigma = \sigma(t, x)$, then d solves

$$(2.4) \quad \partial_t d = \mu \operatorname{tr} \left((I - \widehat{\nabla} d \otimes \widehat{\nabla} d) D^2 d \right) + \frac{\sigma}{|\nabla u|} = \mu \Delta d + \frac{\sigma}{|\nabla u|} \quad \text{in} \quad \bigcup_{t \in [t_0, t_0 + h]} \{t\} \times \Gamma_t,$$

where the last equality follows from $|\nabla d| = 1$ in $\{t\} \times \Gamma_t$.

We remark that around the same time as [16], Chen–Giga–Goto independently established a level set approach for degenerate parabolic PDEs [10]. Shortly after, Barles–Soner–Souganidis rigorously connected the level set approach and the phase field theory for reaction-diffusion equations [5].

2.2. Generalized flows. A different approach for defining the weak evolutions of fronts was considered in [6] and [4]. We present a slight variation of the definition in [4] (see also [27]).

Let $F(p, X)$ be given by

$$(2.5) \quad F(p, X) = -\mu \operatorname{tr}((I - \widehat{p} \otimes \widehat{p})X)$$

and the lower and upper semi-continuous envelopes of F be denoted by F_* and F^* respectively.

Definition 2.1. A family $(D_t)_{t>0}$ (resp., $(E_t)_{t>0}$) of open (resp., closed) subsets of \mathbb{R}^n is a *generalized super-flow* (resp., *sub-flow*) of the mean curvature equation (2.2) if for all $(t_0, x_0) \in (0, \infty) \times \mathbb{R}^n$, $h, r > 0$, and for all smooth functions $\varphi : (0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that

(i) (Boundedness) for all $t \in [t_0, t_0 + h]$, the set

$$\{x \in \mathbb{R}^n : \varphi(t, x) > 0\} \quad (\text{resp. } \{x \in \mathbb{R}^n : \varphi(t, x) < 0\})$$

is bounded and

$$\{x \in \overline{B}(x_0, r) : \varphi(t, x) > 0\} \quad (\text{resp. } \{x \in \overline{B}(x_0, r) : \varphi(t, x) < 0\})$$

is non-empty,

(ii) (Speed) there exists $\delta = \delta(\varphi) > 0$ such that

$$\begin{aligned} \partial_t \varphi + F^*(\nabla \varphi, D^2 \varphi) &\leq -\delta \quad \text{in } [t_0, t_0 + h] \times \overline{B}(x_0, r), \\ (\text{resp.}, \partial_t \varphi + F_*(\nabla \varphi, D^2 \varphi) &\geq \delta) \end{aligned}$$

(iii) (Non-degeneracy)

$$\nabla \varphi \neq 0 \quad \text{on } \{(t, x) \in [t_0, t_0 + h] \times \overline{B}(x_0, r) : \varphi(t, x) = 0\},$$

(iv) (Initial condition)

$$\begin{aligned} \{x \in \mathbb{R}^n : \varphi(t_0, x) \geq 0\} &\subset D_{t_0} \\ (\text{resp.}, \{x \in \mathbb{R}^n : \varphi(t_0, x) \leq 0\} &\subset \mathbb{R}^n \setminus E_{t_0}), \end{aligned}$$

(v) (Boundary condition) for all $t \in [t_0, t_0 + h]$,

$$\begin{aligned} \{x \in \mathbb{R}^n \setminus B(x_0, r) : \varphi(t, x) \geq 0\} &\subset D_t \\ (\text{resp.}, \{x \in \mathbb{R}^n \setminus B(x_0, r) : \varphi(t, x) \leq 0\} &\subset \mathbb{R}^n \setminus E_t), \end{aligned}$$

then

$$\begin{aligned} \{x \in B(x_0, r) : \varphi(t_0 + h, x) > 0\} &\subset D_{t_0+h} \\ (\text{resp.}, \{x \in B(x_0, r) : \varphi(t_0 + h, x) < 0\} &\subset \mathbb{R}^n \setminus E_{t_0+h}). \end{aligned}$$

In this paper, we will implement the abstract method developed in [4]-[6]. Precisely, for $i = 1, \dots, N$, let Ω_0^i be the open sets defined in (1.5). We will show that there exist families of open sets $(D_t^i)_{t \geq 0}$ and $(E_t^i)_{t \geq 0}$, $i = 1, \dots, N$, such that the following hold: $(D_t^i)_{t \geq 0}$ and $((E_t^i)^c)_{t \geq 0}$ are respectively a generalized super and sub-flow of the mean curvature equation with μ as in (1.6), $\Omega_0^i \subset D_0^i$, $(\overline{\Omega}_0^i)^c \subset E_0^i$, and

$$u^\varepsilon(t, x) \rightarrow i \quad \text{if } x \in D_t^i \cap E_t^{i+1}.$$

Therefore, if $({}^+\Omega_t^i, \Gamma_t^i, -\Omega_t^i)_{t \geq 0}$ denotes the level set evolution of $(\Omega_0^i, \Gamma_0^i, (\overline{\Omega_0^i})^c)$, by [4, Theorem 1.1, Theorem 1.2] (see also [6, Proposition 2.1, Theorem 2.4]),

$${}^+\Omega_t^i \subset D_t^i \subset {}^+\Omega_t^i \cup \Gamma_t^i \quad \text{and} \quad -\Omega_t^i \subset E_t^i \subset -\Omega_t^i \cup \Gamma_t^i.$$

In particular, if the sets $(\Gamma_t^i)_{t \geq 0}$ do not develop interior, then

$${}^+\Omega_t^i = D_t^i \quad \text{and} \quad -\Omega_t^i = E_t^i.$$

Our main result, Theorem 1.1, immediately follows.

2.3. Extension of the signed distance function. Recall that the signed distance function $d = d(t, x)$ associated to the front Γ_t in (2.3) is smooth in some neighborhood $Q_\rho = \{|d| < \rho\}$ of the front, provided Γ_t is smooth. However, in general, d is not smooth away from the front. Throughout the paper, we will use the following smooth extension of the distance function away from Γ_t .

Definition 2.2 (Extension of the signed distance function). For $t \in [t_0, t_0 + h]$, let \tilde{d} be the signed distance function from a bounded domain Ω_t with boundary Γ_t and let $\rho > 0$ be such that \tilde{d} is smooth in

$$Q_{2\rho} = \{(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n : |\tilde{d}(t, x)| \leq 2\rho\}.$$

Let $\eta(t, x)$ be a smooth, bounded function such that

$$\eta = 1 \text{ in } \{|\tilde{d}| \leq \rho\}, \quad \eta = 0 \text{ in } \{|\tilde{d}| \geq 2\rho\}, \quad 0 \leq \eta \leq 1.$$

We extend $\tilde{d}(t, x)$ in the set $\{(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n : |\tilde{d}(t, x)| > \rho\}$ with the smooth bounded function $d(t, x)$ given by

$$d(t, x) = \begin{cases} \tilde{d}(t, x) & \text{in } Q_\rho = \{|\tilde{d}(t, x)| \leq \rho\} \\ \tilde{d}(t, x)\eta(t, x) + 2\rho(1 - \eta(t, x)) & \text{in } \{\rho < \tilde{d}(t, x) < 2\rho\} \\ \tilde{d}(t, x)\eta(t, x) - 2\rho(1 - \eta(t, x)) & \text{in } \{-2\rho < \tilde{d}(t, x) < -\rho\} \\ 2\rho & \text{in } \{\tilde{d}(t, x) \geq 2\rho\} \\ -2\rho & \text{in } \{\tilde{d}(t, x) \leq -2\rho\}. \end{cases}$$

Notice that, in $\{\rho < \tilde{d} < 2\rho\}$, the function d satisfies

$$d = 2\rho + (\tilde{d} - 2\rho)\eta > 2\rho - \rho\eta \geq \rho,$$

and, in $\{-2\rho < \tilde{d} < -\rho\}$, the function d satisfies

$$d = -2\rho + (\tilde{d} + 2\rho)\eta < -2\rho + \rho\eta \leq -\rho.$$

3. PRELIMINARY RESULTS ON THE FRACTIONAL LAPLACIAN

In this section, we recall a few basic properties of the operator \mathcal{I}_n that will be used later on in the paper. First of all, by using that

$$\text{P. V.} \int_{\{|y| < R\}} \nabla u(x) \cdot y \frac{dy}{|y|^{n+1}} = 0,$$

for any $R > 0$, we can write

$$\mathcal{I}_n u(x) = \int_{\{|y| < R\}} (u(x+y) - u(x) - \nabla u(x) \cdot y) \frac{dy}{|y|^{n+1}} + \int_{\{|y| > R\}} (u(x+y) - u(x)) \frac{dy}{|y|^{n+1}}.$$

In particular if u is smooth and bounded, both integrals above are finite and we can bound $\mathcal{I}_n u$ as follows,

$$|\mathcal{I}_n u(x)| \leq C \left(\|D^2 u\|_\infty R + \frac{\|u\|_\infty}{R} \right),$$

where we used the following lemma whose proof is a direct computation in polar coordinates.

Lemma 3.1. *There exists $C_1, C_2 > 0$ such that for any $R > 0$,*

$$\int_{\{|z| < R\}} \frac{dz}{|z|^{n-1}} = C_1 R \quad \text{and} \quad \int_{\{|z| > R\}} \frac{dz}{|z|^{n+1}} = \frac{C_2}{R}.$$

We will frequently use Lemma 3.1 throughout the paper without reference.

The next is an auxiliary lemma that allows us to view one-dimensional fractional Laplacians of functions defined over \mathbb{R} equivalently as n -dimensional fractional Laplacians.

Lemma 3.2. [45, Lemma 3.2] *For a vector $e \in \mathbb{R}^n$ and a function $v \in C^{1,1}(\mathbb{R})$, let $v_e(x) = v(e \cdot x) : \mathbb{R}^n \rightarrow \mathbb{R}$. Then,*

$$\mathcal{I}_n[v_e](x) = |e| C_n \mathcal{I}_1[v](e \cdot x)$$

where

$$(3.1) \quad C_n = \int_{\mathbb{R}^{n-1}} \frac{1}{(|y|^2 + 1)^{\frac{n+1}{2}}} dy.$$

Consequently,

$$|e| C_n \mathcal{I}_1[v](\xi) = \text{P. V.} \int_{\mathbb{R}^n} (v(\xi + e \cdot z) - v(\xi)) \frac{dz}{|z|^{n+1}}, \quad \xi \in \mathbb{R}.$$

4. THE PHASE TRANSITION, THE CORRECTOR, AND THE AUXILIARY FUNCTIONS

In this section, we will introduce the phase transition ϕ and the corrector ψ . Along the way, we will also define the auxiliary functions a_ε and \bar{a}_ε and exhibit their relationship with fractional Laplacians and the mean curvature equation, respectively.

4.1. The phase transition ϕ . Let ϕ be the solution to (1.4). For convenience in the notation, let c_0 and α be given respectively by

$$(4.1) \quad c_0^{-1} = \int_{\mathbb{R}} [\dot{\phi}(\xi)]^2 d\xi \quad \text{and} \quad \alpha = \frac{W''(0)}{C_n}$$

where C_n is defined in (3.1), and let $H(\xi)$ be the Heaviside function.

Lemma 4.1. *There is a unique solution $\phi \in C^{4,\beta}(\mathbb{R})$ of (1.4), with β as in (1.3). Moreover, there exists a constant $C > 0$ such that*

$$(4.2) \quad \left| \phi(\xi) - H(\xi) + \frac{1}{\alpha \xi} \right| \leq \frac{C}{|\xi|^2}, \quad \text{for } |\xi| \geq 1$$

and

$$(4.3) \quad \frac{1}{C|\xi|^2} \leq \dot{\phi}(\xi) \leq \frac{C}{|\xi|^2}, \quad |\ddot{\phi}(\xi)| \leq \frac{C}{|\xi|^2}, \quad |\dddot{\phi}(\xi)| \leq \frac{C}{|\xi|^2}, \quad \text{for } |\xi| \geq 1.$$

Proof. Existence of a unique solution $\phi \in C^{2,\beta}(\mathbb{R})$ of (1.4) is proven in [8], see Theorem 1.2 and Lemma 2.3. The regularity of W in (1.3) implies that ϕ is actually in $C^{4,\beta}(\mathbb{R})$, see [8, Lemma 2.3]. Estimate (4.2) and the estimate on $\dot{\phi}$ in (4.3) are proven in [23, Theorem 3.1]. Finally, the estimates on $\ddot{\phi}$ and $\dddot{\phi}$ in (4.3) are proven in [34, Lemma 3.1]. \square

4.2. The auxiliary functions a_ε and \bar{a}_ε . Now, we will introduce two auxiliary functions that are necessary for our analysis. Let Ω_t be a bounded domain with smooth boundary Γ_t , for $t \in [t_0, t_0 + h]$. Throughout this section, let $d = d(t, x)$ and ρ be such that d is the smooth extension of the signed distance function \tilde{d} to Γ_t outside of Q_ρ as given in Definition 2.2. We also introduce a new parameter $0 < \gamma < 1$.

Define the function $a_\varepsilon = a_\varepsilon(\xi; t, x)$ by

$$(4.4) \quad a_\varepsilon = \int_{\{|z| < \frac{\gamma}{\varepsilon}\}} \left(\phi \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \phi(\xi + \nabla d(t, x) \cdot z) \right) \frac{dz}{|z|^{n+1}},$$

where $(\xi, t, x) \in \mathbb{R} \times [t_0, t_0 + h] \times \mathbb{R}^n$. By Lemma 3.1 and by the regularity of ϕ and d , the integral in (4.4) is well defined. We note that the integral is taken over $\{|z| < \frac{\gamma}{\varepsilon}\}$ instead of \mathbb{R}^n so that a_ε is sufficiently small for (t, x) close to Γ_t and large ξ (see (4.14)) and its derivatives in t and x remain bounded (see the proof of (4.7)).

The corresponding function $\bar{a}_\varepsilon = \bar{a}_\varepsilon(t, x)$ is given by

$$(4.5) \quad \bar{a}_\varepsilon(t, x) = \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} a_\varepsilon(\xi; t, x) \dot{\phi}(\xi) d\xi.$$

One of the main observations in [29] is that $\bar{a}_\varepsilon(t, x)$ converges, up to constants, to $\Delta d(t, x)$ in the neighborhood Q_ρ , and therefore to the mean curvature of Γ_t at points $x \in \Gamma_t$, see [29, Lemma 4]. In our setting, we use the recent result in [45].

Theorem 4.2. *For $t \in [t_0, t_0 + h]$, let Ω_t be a bounded domain with smooth boundary Γ_t . Let d be as in Definition 2.2. Then,*

$$\lim_{\varepsilon \rightarrow 0} \bar{a}_\varepsilon(t, x) = \frac{1}{2} \frac{|S^{n-2}|}{n-1} \Delta d(t, x) = \mu c_0^{-1} \operatorname{tr} \left((I - \widehat{\nabla} d \otimes \widehat{\nabla} d) D^2 d \right)$$

uniformly in $(t, x) \in Q_\rho$ where $\mu > 0$ is in (1.6).

Proof. This is [45, Theorem 1.3] (see also Remark 1.1 there) when $\gamma = +\infty$. For $0 < \gamma < 1$, the proof is the same after making a minor adaptation to [45, Lemma 4.1]. \square

We will also need several estimates on $a_\varepsilon, \bar{a}_\varepsilon$ and their derivatives, none of which are proven in [29]. First, we have the following general estimate. The proof is delayed until Section 8.

Lemma 4.3. *There exists $C > 0$ such that, for all $(\xi, t, x) \in \mathbb{R} \times [t_0, t_0 + h] \times \mathbb{R}^n$,*

$$(4.6) \quad \|a_\varepsilon\|_{C_\xi^3(\mathbb{R})} \leq C \varepsilon^{\frac{1}{2}},$$

$$(4.7) \quad \|\partial_t a_\varepsilon\|_{C_\xi^3(\mathbb{R})}, \|\nabla_x a_\varepsilon\|_{C_\xi^3(\mathbb{R})} \leq C |\ln \varepsilon|, \quad \|D_x^2 a_\varepsilon\|_{C_\xi^3(\mathbb{R})} \leq \frac{C}{\varepsilon},$$

$$(4.8) \quad |a_\varepsilon(\xi; t, x)|, |\dot{a}_\varepsilon(\xi; t, x)| \leq \frac{C}{1 + |\xi|},$$

$$(4.9) \quad \begin{aligned} |\partial_t a_\varepsilon(\xi; t, x)|, |\nabla_x a_\varepsilon(\xi; t, x)| &\leq \frac{C}{\varepsilon(1 + |\xi|)}, & |D_x^2 a_\varepsilon(\xi; t, x)| &\leq \frac{C}{\varepsilon^2(1 + |\xi|)}, \\ |\partial_t \dot{a}_\varepsilon(\xi; t, x)|, |\nabla_x \dot{a}_\varepsilon(\xi; t, x)| &\leq \frac{C}{\varepsilon(1 + |\xi|)}, & |D_x^2 \dot{a}_\varepsilon(\xi; t, x)| &\leq \frac{C}{\varepsilon^2(1 + |\xi|)}. \end{aligned}$$

Consequently, for all $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$,

$$(4.10) \quad |\bar{a}_\varepsilon(t, x)| \leq \frac{C}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|},$$

and

$$(4.11) \quad |\partial_t \bar{a}_\varepsilon(t, x)|, |\nabla_x \bar{a}_\varepsilon(t, x)| \leq \frac{C}{\varepsilon}, \quad |D_x^2 \bar{a}_\varepsilon(t, x)| \leq \frac{C}{\varepsilon^2 |\ln \varepsilon|}.$$

We also have the following refined estimates near the front Γ_t . The proof is in Section 9.

Lemma 4.4. *There exists $C > 0$ such that, for $(\xi, t, x) \in \mathbb{R} \times [t_0, t_0 + h] \times \mathbb{R}^n$, if $|\nabla d(t, x)| \neq 0$, then*

$$(4.12) \quad \begin{aligned} |a_\varepsilon(\xi; t, x)|, |\dot{a}_\varepsilon(\xi; t, x)|, |\ddot{a}_\varepsilon(\xi; t, x)| &\leq \frac{C\varepsilon |\ln \varepsilon|}{|\nabla d(t, x)|}, \\ |\bar{a}_\varepsilon(t, x)| &\leq \frac{C}{|\nabla d(t, x)|}. \end{aligned}$$

Moreover, if $(t, x) \in Q_\rho$, then

$$(4.13) \quad \begin{aligned} |\partial_t a_\varepsilon(\xi; t, x)|, |\nabla_x a_\varepsilon(\xi; t, x)|, |\nabla_x \dot{a}_\varepsilon(\xi; t, x)| &\leq C, \\ |\partial_t \bar{a}_\varepsilon(t, x)|, |\nabla_x \bar{a}_\varepsilon(t, x)| &\leq \frac{C}{\varepsilon |\ln \varepsilon|}, \end{aligned}$$

and, there exists $\varepsilon_0 = \varepsilon_0(\gamma)$, with γ as in (4.4), such that for $0 < \varepsilon < \varepsilon_0$,

$$(4.14) \quad |a_\varepsilon(\xi; t, x)| \leq \frac{C\gamma}{1 + |\xi|}.$$

It is also important to notice that when $\xi = d(t, x)/\varepsilon$, morally, $a_\varepsilon(d/\varepsilon)$ is the difference between an n -dimensional and a 1-dimensional fractional Laplacian of $\phi(d/\varepsilon)$. This is made precise in the following lemma whose proof is delayed until Section 10.

Lemma 4.5. *There is $C = C(\gamma, \rho) > 0$ such that*

$$\left| a_\varepsilon \left(\frac{d(t, x)}{\varepsilon}; t, x \right) - \left(\varepsilon \mathcal{I}_n \left[\phi \left(\frac{d(t, \cdot)}{\varepsilon} \right) \right] (x) - C_n \mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) \right) \right| \leq C\varepsilon$$

for any $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$.

4.3. The corrector ψ . The linearized operator \mathcal{L} associated to (1.4) around ϕ is given by

$$(4.15) \quad \mathcal{L}[\psi] = -C_n \mathcal{I}_1[\psi] + W''(\phi)\psi.$$

In the constructions of barriers, we will need the corrector $\psi = \psi(\xi; t, x)$ that solves

$$(4.16) \quad \begin{cases} \mathcal{L}[\psi] = \frac{a_\varepsilon(\xi; t, x)}{\varepsilon |\ln \varepsilon|} + c_0 \dot{\phi}(\xi) (\sigma - \bar{a}_\varepsilon(t, x)) + \tilde{\sigma} (W''(\phi(\xi)) - W''(0)), & \xi \in \mathbb{R} \\ \psi(\pm\infty; t, x) = 0, \end{cases}$$

where $\sigma \in \mathbb{R}$ is a small constant and $\tilde{\sigma}$ is such that $\sigma = W''(0)\tilde{\sigma}$. See Section 5 for a formal derivation of (4.16).

To prove existence of the solution of (4.16), we will use the following result from [23].

Lemma 4.6. *Let $g \in H^{\frac{1}{2}}(\mathbb{R})$ be such that*

$$\int_{\mathbb{R}} g(\xi) \dot{\phi}(\xi) d\xi = 0.$$

Then, there exists a unique solution $\psi \in H^{\frac{1}{2}}(\mathbb{R})$ such that $\int_{\mathbb{R}} \psi(\xi) \dot{\phi}(\xi) d\xi = 0$ of

$$\mathcal{L}[\psi] = g \quad \text{in } \mathbb{R}.$$

Moreover, if $g \in C^{2,\beta}(\mathbb{R})$ with β as in (1.3), then $\psi \in C^{3,\beta}(\mathbb{R})$, and

$$(4.17) \quad \|\psi\|_{C^{3,\beta}(\mathbb{R})} \leq C(\|g\|_{C^{2,\beta}(\mathbb{R})} + \|\psi\|_{L^\infty(\mathbb{R})}).$$

Proof. The proof of existence of a unique solution $\psi \in H^{\frac{1}{2}}(\mathbb{R})$ satisfying $\int_{\mathbb{R}} \psi(\xi) \dot{\phi}(\xi) d\xi = 0$ is contained in the proof of [23, Theorem 3.2]. If $g \in C^{2,\beta}(\mathbb{R}) \cap H^{\frac{1}{2}}(\mathbb{R})$, then for all $p \geq 2$,

$$\|g\|_{L^p(\mathbb{R})} \leq C$$

and the boundedness of ψ follows from [23, Corollary 5.16]. The $C^{3,\beta}$ regularity of ψ and (4.17) are a consequence of the regularity of ϕ given by Lemma 4.1 and [8, Lemma 2.3]. \square

Lemma 4.7. *Let*

$$(4.18) \quad g(\xi; t, x) := \frac{a_\varepsilon(\xi; t, x)}{\varepsilon |\ln \varepsilon|} + c_0 \dot{\phi}(\xi) (\sigma - \bar{a}_\varepsilon(t, x)) + \tilde{\sigma} (W''(\phi(\xi)) - W''(0)).$$

Then,

$$(4.19) \quad \int_{\mathbb{R}} g(\xi; t, x) \dot{\phi}(\xi) d\xi = 0.$$

Moreover, $g \in H_\xi^{\frac{1}{2}}(\mathbb{R}) \cap C_\xi^{2,\beta}(\mathbb{R})$ with β as in (1.3), uniformly in $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$.

Proof. Recalling that $\sigma = W''(0)\tilde{\sigma}$, we compute

$$\begin{aligned} \int_{\mathbb{R}} g(\xi; t, x) \dot{\phi}(\xi) d\xi &= \int_{\mathbb{R}} \left(\frac{1}{\varepsilon |\ln \varepsilon|} a_\varepsilon(\xi; t, x) - \dot{\phi}(\xi) c_0 \bar{a}_\varepsilon(t, x) \right) \dot{\phi}(\xi) d\xi \\ &\quad + \int_{\mathbb{R}} \left(c_0 \sigma \dot{\phi}(\xi) + \tilde{\sigma} W''(\phi(\xi)) - \sigma \right) \dot{\phi}(\xi) d\xi. \end{aligned}$$

Using the definitions of c_0 and \bar{a}_ε in (4.1) and (4.5), respectively, we get

$$\int_{\mathbb{R}} \left(\frac{1}{\varepsilon |\ln \varepsilon|} a_\varepsilon(\xi; t, x) - \dot{\phi}(\xi) c_0 \bar{a}_\varepsilon(t, x) \right) \dot{\phi}(\xi) d\xi = \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} a_\varepsilon(\xi; t, x) \dot{\phi}(\xi) d\xi - \bar{a}_\varepsilon(t, x) = 0.$$

Then, we use that W' is periodic, $\phi(\infty) = 1$ and $\phi(-\infty) = 0$ to find that

$$\int_{\mathbb{R}} \tilde{\sigma} W''(\phi(\xi)) \dot{\phi}(\xi) d\xi = \tilde{\sigma} \int_{\mathbb{R}} \frac{d}{d\xi} [W'(\phi(\xi))] d\xi = \tilde{\sigma} [W'(1) - W'(0)] = 0$$

and again the definition of c_0 to see that

$$\int_{\mathbb{R}} \left(c_0 \sigma \dot{\phi}(\xi) - \sigma \right) \dot{\phi}(\xi) d\xi = c_0 \sigma \int_{\mathbb{R}} [\dot{\phi}(\xi)]^2 d\xi - \sigma = 0,$$

as desired. This proves (4.19).

Next, from (4.2), (4.3) and (4.8) we have that

$$W''(\phi(\xi)) - W''(0) = O(\phi(\xi)), \dot{\phi}(\xi), a_\varepsilon(\xi; t, x) \in H_\xi^1(\mathbb{R}),$$

which implies that $g \in H_\xi^{\frac{1}{2}}(\mathbb{R})$, with $\|g\|_{H_\xi^{\frac{1}{2}}(\mathbb{R})} \leq C_\varepsilon$ for all $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$. Moreover,

from the regularity of ϕ and W and (4.6), it follows that $g \in C_\xi^{2,\beta}(\mathbb{R})$ with $\|g\|_{C_\xi^{2,\beta}(\mathbb{R})} \leq C_\varepsilon$, for all $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$. \square

Theorem 4.8. *There is a unique solution $\psi = \psi(\xi; t, x) \in C^{3,\beta}(\mathbb{R})$ to (4.16) with β as in (1.3), and $C > 0$ such that, for all $(\xi, t, x) \in \mathbb{R} \times [t_0, t_0 + h] \times \mathbb{R}^n$*

$$(4.20) \quad \begin{aligned} |\psi(\xi; t, x)|, |\dot{\psi}(\xi; t, x)|, |\ddot{\psi}(\xi; t, x)| &\leq \frac{C}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|}, \\ |\partial_t \psi(\xi; t, x)|, |\nabla_x \psi(\xi; t, x)|, |\nabla_x \dot{\psi}(\xi; t, x)| &\leq \frac{C}{\varepsilon}, \\ |D_x^2 \psi(\xi; t, x)| &\leq \frac{C}{\varepsilon^2 |\ln \varepsilon|}, \end{aligned}$$

and

$$(4.21) \quad |\psi(\xi; t, x)| \leq \frac{C}{\varepsilon |\ln \varepsilon| (1 + |\xi|)}.$$

Moreover, if $(t, x) \in Q_\rho$, then

$$(4.22) \quad \begin{aligned} |\psi(\xi; t, x)|, |\dot{\psi}(\xi; t, x)|, |\ddot{\psi}(\xi; t, x)| &\leq C, \\ |\partial_t \psi(\xi; t, x)|, |\nabla_x \psi(\xi; t, x)|, |\nabla_x \dot{\psi}(\xi; t, x)| &\leq \frac{C}{\varepsilon |\ln \varepsilon|} \end{aligned}$$

and, there exists $\varepsilon_0 = \varepsilon_0(\gamma)$, with γ as in (4.4), such that for $\varepsilon < \varepsilon_0$,

$$(4.23) \quad |\psi(\xi; t, x)| \leq \frac{C\gamma}{\varepsilon |\ln \varepsilon| (1 + |\xi|)},$$

with γ as in (4.4).

Proof. First, the existence of a unique solution $\psi \in H_\xi^{\frac{1}{2}}(\mathbb{R}) \cap C_\xi^{3,\beta}(\mathbb{R})$ to (4.16) satisfying $\int_{\mathbb{R}} \psi(\xi; t, x) \dot{\phi}(\xi) d\xi = 0$ follows from Lemmas 4.6 and 4.7.

Let us show (4.20) for ψ . Let g be defined as in (4.18). We first notice that by (4.6) and (4.10), we have that, for all $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$

$$(4.24) \quad \|g(\cdot; t, x)\|_{C^{2,\beta}(\mathbb{R})} \leq \frac{C}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|}.$$

Suppose by contradiction that there is a sequence $\varepsilon_k \rightarrow 0$ as $k \rightarrow \infty$ such that if ψ_k is solution to (4.16) with $\varepsilon = \varepsilon_k$, then for some $(t_k, x_k) \in [t_0, t_0 + h] \times \mathbb{R}^n$,

$$(4.25) \quad \|\psi_k(\cdot; t_k, x_k)\|_{L^\infty(\mathbb{R})} \geq \frac{1}{\varepsilon_k^{\frac{1}{2}} |\ln \varepsilon_k| b_k}$$

with $0 < b_k \rightarrow 0$ as $k \rightarrow \infty$. Define the function $\tilde{\psi}_k(\xi) := \psi_k(\cdot; t_k, x_k) / \|\psi_k(\cdot; t_k, x_k)\|_{L^\infty(\mathbb{R})}$. Clearly, $\|\tilde{\psi}_k\|_{L^\infty(\mathbb{R})} = 1$, $\int_{\mathbb{R}} \tilde{\psi}_k(\xi) \dot{\phi}(\xi) d\xi = 0$ and $\tilde{\psi}_k$ solves

$$\mathcal{L}[\tilde{\psi}_k] = \tilde{g}_k \quad \text{in } \mathbb{R},$$

with

$$\tilde{g}_k(\xi) = \frac{g(\xi; t, x)}{\|\psi_k(\cdot; t_k, x_k)\|_{L^\infty(\mathbb{R})}}.$$

Notice that by (4.24) and (4.25), $\tilde{g}_k \rightarrow 0$ in $C^{2,\beta}(\mathbb{R})$ as $k \rightarrow \infty$. By (4.17) the sequence of functions $\tilde{\psi}_k$ is uniformly bounded in $C^{3,\beta}(\mathbb{R})$. Therefore, up to a subsequence, it converges in $C^3(\mathbb{R})$ to a function $\tilde{\psi}_\infty \in H^{\frac{1}{2}}(\mathbb{R}) \cap C^{3,\beta}(\mathbb{R})$ which is solution to

$$\mathcal{L}[\tilde{\psi}_\infty] = 0 \quad \text{in } \mathbb{R}.$$

Moreover, $\int_{\mathbb{R}} \tilde{\psi}_{\infty}(\xi) \dot{\phi}(\xi) d\xi = 0$. Indeed, by the uniform convergence of $\tilde{\psi}_k$ to $\tilde{\psi}_{\infty}$ and using that $\phi(-\infty) = 0$ and $\phi(\infty) = 1$, as $k \rightarrow \infty$,

$$\begin{aligned} \left| \int_{\mathbb{R}} \tilde{\psi}_{\infty}(\xi) \dot{\phi}(\xi) d\xi \right| &= \left| \int_{\mathbb{R}} \tilde{\psi}_k(\xi) \dot{\phi}(\xi) d\xi - \int_{\mathbb{R}} \tilde{\psi}_{\infty}(\xi) \dot{\phi}(\xi) d\xi \right| \\ &\leq \|\tilde{\psi}_k - \psi_{\infty}\|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} \dot{\phi}(\xi) d\xi \\ &= \|\tilde{\psi}_k - \psi_{\infty}\|_{L^{\infty}(\mathbb{R})} \rightarrow 0. \end{aligned}$$

The uniqueness of the solution guaranteed by Lemma 4.6 implies that $\tilde{\psi}_{\infty} \equiv 0$ which is in contradiction with $\|\tilde{\psi}_{\infty}\|_{L^{\infty}(\mathbb{R})} = 1$. This proves (4.20) for ψ . The estimates for $\dot{\psi}$ and $\ddot{\psi}$ in (4.20) then follow from (4.24), (4.17) and (4.20) for ψ , just proven.

Next, we establish (4.20) for the time/space derivatives of ψ . From (4.19) it follows that, for $i, j = 1, \dots, n$,

$$\int_{\mathbb{R}} \partial_t g(\xi; t, x) \dot{\phi}(\xi) d\xi = 0, \quad \int_{\mathbb{R}} \partial_{x_i} g(\xi; t, x) \dot{\phi}(\xi) d\xi = 0, \quad \int_{\mathbb{R}} \partial_{x_i x_j}^2 g(\xi; t, x) \dot{\phi}(\xi) d\xi = 0.$$

Moreover, from (4.3), (4.7), (4.9) and (4.11), the functions $\partial_t g$, $\partial_{x_i} g$ and $\partial_{x_i x_j}^2 g$ belong to the space $H_{\xi}^{\frac{1}{2}}(\mathbb{R}) \cap C_{\xi}^{2,\beta}(\mathbb{R})$ uniformly in (t, x) . Thus, it is easy to see that $\partial_t \psi$, $\partial_{x_i} \psi$ and $\partial_{x_i x_j}^2 \psi$ exist and are the unique solutions in $H_{\xi}^{\frac{1}{2}}(\mathbb{R})$ such that $\int_{\mathbb{R}} \partial_t \psi(\xi; t, x) \dot{\phi}(\xi) d\xi = 0$, $\int_{\mathbb{R}} \partial_{x_i} \psi(\xi; t, x) \dot{\phi}(\xi) d\xi = 0$ and $\int_{\mathbb{R}} \partial_{x_i x_j}^2 \psi(\xi; t, x) \dot{\phi}(\xi) d\xi = 0$, to

$$\mathcal{L}[\partial_t \psi] = \partial_t g, \quad \mathcal{L}[\partial_{x_i} \psi] = \partial_{x_i}^2 g, \quad \mathcal{L}[\partial_{x_i x_j}^2 \psi] = \partial_{x_i x_j}^2 g \quad \text{in } \mathbb{R},$$

respectively. Therefore, as above, from (4.7), the estimates in (4.20) for $\partial_t \psi$, $\nabla_x \psi$, $\nabla_x \dot{\psi}$ and $D_x^2 \psi$ follow.

Now, we prove estimate (4.21). By (4.2), (4.3), (4.8) and (4.10), for $|\xi| \geq 1$,

$$(4.26) \quad |g(\xi; t, x)| \leq \frac{C}{\varepsilon |\ln \varepsilon| |\xi|}.$$

For $a > 0$ let us denote $\phi_a(\xi) := \phi\left(\frac{\xi}{a}\right)$. Then, ϕ_a solves

$$C_n \mathcal{I}_1[\phi_a] = \frac{1}{a} W'(\phi_a) \quad \text{in } \mathbb{R}.$$

Therefore, recalling (4.1) and that $W'(0) = 0$, for $\xi \leq -1$, by (4.2) (note that $H(\xi) = 0$),

$$\begin{aligned} \mathcal{L}[\phi_a](\xi) &= W''(\phi(\xi)) \phi_a(\xi) - \frac{1}{a} W'(\phi_a(\xi)) \\ &= W''(0) \phi_a(\xi) - \frac{W''(0)}{a} \phi_a(\xi) + O(\phi \phi_a) + O(\phi_a^2) \\ &= \alpha C_n \left(\phi_a(\xi) - \frac{1}{a} \phi_a(\xi) \right) + O\left(\frac{1}{\xi^2}\right) \\ &= -C_n \frac{a-1}{\xi} + O\left(\frac{1}{\xi^2}\right). \end{aligned}$$

Choose $a = 3$. Then, there exists $R_0 > 0$ such that,

$$(4.27) \quad \mathcal{L}[\phi_a](\xi) = -\frac{2C_n}{\xi} + O\left(\frac{1}{\xi^2}\right) = \frac{2C_n}{|\xi|} + O\left(\frac{1}{\xi^2}\right) \geq \frac{C_n}{|\xi|}, \quad \text{for } \xi < -R_0.$$

Choose R_0 so large that for $\xi < -R_0$, by (4.2),

$$W''(\phi(\xi)) \geq W''(0) - C\phi(\xi) \geq \alpha C_n - \frac{C}{|\xi|} \geq \frac{\alpha C_n}{2} > 0.$$

Then, the operator \mathcal{L} satisfies the maximum principle in $(-\infty, -R_0)$. By (4.20) and the monotonicity of ϕ ,

$$(4.28) \quad \psi(\xi; t, x) \leq \frac{C\varepsilon^{\frac{1}{2}}}{\varepsilon|\ln\varepsilon|} \leq \frac{\tilde{C}}{\varepsilon|\ln\varepsilon|} \phi\left(-\frac{R_0}{3}\right) \leq \frac{\tilde{C}\phi_a(\xi)}{\varepsilon|\ln\varepsilon|}, \quad \text{for } \xi \geq -R_0.$$

Choose $K \geq \tilde{C}$, with \tilde{C} as in (4.28), such that, if we denote $\hat{\phi}(\xi) = \frac{K\phi_a(\xi)}{\varepsilon|\ln\varepsilon|}$, then by (4.26) and (4.27),

$$\mathcal{L}[\hat{\phi}](\xi) \geq g(\xi; t, x) \quad \text{for } \xi < -R_0.$$

Since in addition by (4.28),

$$\psi(\xi; t, x) \leq \hat{\phi}(\xi) \quad \text{for } \xi \geq -R_0,$$

the maximum principle implies that

$$\psi(\xi; t, x) \leq \hat{\phi}(\xi) \leq \frac{C}{\varepsilon|\ln\varepsilon||\xi|} \quad \text{for } \xi < -R_0.$$

Comparing ψ with $-\hat{\phi}$ we also get

$$\psi(\xi; t, x) \geq -\frac{C}{\varepsilon|\ln\varepsilon||\xi|} \quad \text{for } \xi < -R_0.$$

Similarly, choosing $a < 0$ and using the periodicity of W , one can prove that

$$|\psi(\xi; t, x)| \leq \frac{C}{\varepsilon|\ln\varepsilon||\xi|} \quad \text{for } \xi > R_0.$$

Estimate (4.21) then follows.

Next, we prove (4.22). For $(t, x) \in Q_\rho = \{|d(t, x)| < \rho\}$, recall that $|\nabla d(t, x)| = 1$. By (4.2), (4.3), and (4.12), we have

$$|g(\xi; t, x)| \leq C \quad \text{for all } \xi \in \mathbb{R} \text{ and } (t, x) \in Q_\rho.$$

By the above arguments for the proof of (4.20), we can show (4.22) for ψ . The estimates for $\dot{\psi}$ and $\ddot{\psi}$ then follow. With (4.2), (4.3), and (4.13), we find (4.22) for $\partial_t \psi, \nabla_x \psi, \nabla_x \dot{\psi}$ in the same way as above for (4.20).

Lastly, we check (4.23). By (4.2), (4.3), (4.14), and (4.12), for $|\xi| \geq 1$ and $(t, x) \in Q_\rho$,

$$|g(\xi; t, x)| \leq \frac{C\gamma}{\varepsilon|\ln\varepsilon||\xi|}.$$

Then, following the above arguments for the proof of (4.21), we obtain (4.23). \square

We conclude this section by stating the following estimate for the n - and 1-dimensional fractional Laplacians of ψ . The proof is in Section 11.

Lemma 4.9. *There is $C = C(\rho) > 0$ such that*

$$\left| \varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(t, \cdot)}{\varepsilon}; t, \cdot \right) \right] (x) - C_n \mathcal{I}_1 [\psi(\cdot; t, x)] \left(\frac{d(t, x)}{\varepsilon} \right) \right| \leq \frac{C}{|\ln\varepsilon|^{\frac{1}{2}}}$$

for any $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$.

5. HEURISTICS

5.1. Derivation of the mean curvature equation. We believe it is helpful to view the heuristical derivation of the evolution of the fronts Γ_t^i by mean curvature in Theorem 1.1. For simplicity, we present the case $N = 2$.

For the following formal computations, assume that the signed distance function $d_i(t, x)$ associated to Γ_t^i is smooth and that $|\nabla d_i| = 1$. Moreover, we assume that there is a positive, uniform distance ρ between Γ_t^1 and Γ_t^2 .

Consider the following ansatz for the solution of (1.1)-(1.7)

$$(5.1) \quad u^\varepsilon(t, x) \simeq \phi\left(\frac{d_1(t, x)}{\varepsilon}\right) + \phi\left(\frac{d_2(t, x)}{\varepsilon}\right),$$

with ϕ the solution of (1.4). Plugging the ansatz into (1.1), the left-hand side gives

$$(5.2) \quad \varepsilon \partial_t u^\varepsilon \simeq \dot{\phi}\left(\frac{d_1}{\varepsilon}\right) \partial_t d_1 + \dot{\phi}\left(\frac{d_2}{\varepsilon}\right) \partial_t d_2.$$

On the other hand, we use the equation for ϕ and estimates on a_ε (see Lemma 4.5) to write the fractional Laplacian of the ansatz as

$$(5.3) \quad \begin{aligned} \varepsilon \mathcal{I}_n[u^\varepsilon] &\simeq \varepsilon \mathcal{I}_n\left[\phi\left(\frac{d_1}{\varepsilon}\right)\right] + \varepsilon \mathcal{I}_n\left[\phi\left(\frac{d_2}{\varepsilon}\right)\right] \\ &= \left(\varepsilon \mathcal{I}_n\left[\phi\left(\frac{d_1}{\varepsilon}\right)\right] - C_n \mathcal{I}_1[\phi]\left(\frac{d_1}{\varepsilon}\right)\right) + \left(\varepsilon \mathcal{I}_n\left[\phi\left(\frac{d_2}{\varepsilon}\right)\right] - C_n \mathcal{I}_1[\phi]\left(\frac{d_2}{\varepsilon}\right)\right) \\ &\quad + C_n \mathcal{I}_1[\phi]\left(\frac{d_1}{\varepsilon}\right) + C_n \mathcal{I}_1[\phi]\left(\frac{d_2}{\varepsilon}\right) \\ &\simeq a_\varepsilon\left(\frac{d_1}{\varepsilon}\right) + a_\varepsilon\left(\frac{d_2}{\varepsilon}\right) + W'\left(\phi\left(\frac{d_1}{\varepsilon}\right)\right) + W'\left(\phi\left(\frac{d_2}{\varepsilon}\right)\right). \end{aligned}$$

Freeze a point (t, x) near the front Γ_t^1 . Let $\xi = d_1(t, x)/\varepsilon$ and assume separation of scales. That is, assume that ξ and (t, x) are unrelated. In this regard, let $\eta := d_2(t, x)$ with $|\eta| \geq \rho$. Since the ansatz u^ε is a solution to (1.1), we can multiply the equation by $\dot{\phi}(\xi)$ and integrate over $\xi \in \mathbb{R}$ to write

$$(5.4) \quad \int_{\mathbb{R}} \varepsilon \partial_t u^\varepsilon \dot{\phi}(\xi) d\xi \simeq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} (\varepsilon \mathcal{I}_n[u^\varepsilon] - W'(u^\varepsilon)) \dot{\phi}(\xi) d\xi.$$

For convenience, we will consider the left and right-hand sides separately again.

First, the left-hand side of (5.4) with (5.2) gives

$$\begin{aligned} \int_{\mathbb{R}} \varepsilon \partial_t u^\varepsilon \dot{\phi}(\xi) d\xi &\simeq \partial_t d_1(t, x) \int_{\mathbb{R}} [\dot{\phi}(\xi)]^2 d\xi + \partial_t d_2(t, x) \int_{\mathbb{R}} \dot{\phi}\left(\frac{\eta}{\varepsilon}\right) \dot{\phi}(\xi) d\xi \\ &\simeq c_0^{-1} \partial_t d_1(t, x) + \partial_t d_2(t, x) O\left(\frac{\varepsilon^2}{\rho^2}\right) \int_{\mathbb{R}} \dot{\phi}(\xi) d\xi \\ &\simeq c_0^{-1} \partial_t d_1(t, x), \end{aligned}$$

where we used (4.1), (1.4), and the asymptotics on $\dot{\phi}$ in (4.3).

Then, we look at the right-hand side of (5.4) with (5.3) and write

$$\begin{aligned} \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} [\varepsilon \mathcal{I}_n[u^\varepsilon] - W'(u^\varepsilon)] \dot{\phi}(\xi) d\xi \\ \simeq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} \left[a_\varepsilon(\xi) + a_\varepsilon\left(\frac{\eta}{\varepsilon}\right) + W'(\phi(\xi)) + W'\left(\phi\left(\frac{\eta}{\varepsilon}\right)\right) \right] \dot{\phi}(\xi) d\xi \end{aligned}$$

$$- \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W' \left(\phi(\xi) + \phi \left(\frac{\eta}{\varepsilon} \right) \right) \dot{\phi}(\xi) d\xi.$$

Using that $\phi(-\infty) = 0$, $\phi(\infty) = 1$ and that W is periodic, we have

$$\frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W'(\phi(\xi)) \dot{\phi}(\xi) d\xi = \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} \frac{d}{d\xi} [W(\phi(\xi))] d\xi = 0.$$

Next, we use again (1.4), the asymptotics on ϕ in (4.2), and that $W'(0) = 0$ to expand W' around the origin and estimate

$$\begin{aligned} \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W' \left(\phi \left(\frac{\eta}{\varepsilon} \right) \right) \dot{\phi}(\xi) d\xi &= \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W' \left(\phi \left(\frac{\eta}{\varepsilon} \right) - H \left(\frac{\eta}{\varepsilon} \right) \right) \dot{\phi}(\xi) d\xi \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} \left[W'(0) + W''(0) \left(\phi \left(\frac{\eta}{\varepsilon} \right) - H \left(\frac{\eta}{\varepsilon} \right) \right) \right] \dot{\phi}(\xi) d\xi \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} O \left(\frac{\varepsilon}{\rho} \right) \int_{\mathbb{R}} \dot{\phi}(\xi) d\xi \simeq 0. \end{aligned}$$

For the remaining W' term, we expand around $\phi(\xi)$ and use similar estimates to obtain

$$\begin{aligned} \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W' \left(\phi(\xi) + \phi \left(\frac{\eta}{\varepsilon} \right) \right) \dot{\phi}(\xi) d\xi &= \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W' \left(\phi(\xi) + \phi \left(\frac{\eta}{\varepsilon} \right) - H \left(\frac{\eta}{\varepsilon} \right) \right) \dot{\phi}(\xi) d\xi \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} \left[W'(\phi(\xi)) + W''(\phi(\xi)) \left(\phi \left(\frac{\eta}{\varepsilon} \right) - H \left(\frac{\eta}{\varepsilon} \right) \right) \right] \dot{\phi}(\xi) d\xi \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} W'(\phi(\xi)) \dot{\phi}(\xi) d\xi + \frac{1}{\varepsilon |\ln \varepsilon|} O \left(\frac{\varepsilon}{\rho} \right) \int_{\mathbb{R}} W''(\phi(\xi)) \dot{\phi}(\xi) d\xi \\ &= 0. \end{aligned}$$

Lastly, for the nonlocal terms, by (4.8),

$$\frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} a_\varepsilon \left(\frac{\eta}{\varepsilon} \right) \dot{\phi}(\xi) d\xi \simeq \frac{1}{\varepsilon |\ln \varepsilon|} O \left(\frac{\varepsilon}{\rho} \right) \int_{\mathbb{R}} \dot{\phi}(\xi) d\xi \simeq 0$$

and by Theorem 4.2,

$$\begin{aligned} \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} a_\varepsilon(\xi) \dot{\phi}(\xi) d\xi &= \bar{a}_\varepsilon(t, x) \\ &\simeq c_0^{-1} \mu \operatorname{tr} \left((I - \widehat{\nabla d_1(t, x)}) \otimes \widehat{\nabla d_1(t, x)} \right) D^2 d_1(t, x). \end{aligned}$$

Combing all these pieces, (5.4) for the ansatz gives

$$c_0^{-1} \partial_t d_1(t, x) \simeq \mu c_0^{-1} \operatorname{tr} \left((I - \widehat{\nabla d_1(t, x)}) \otimes \widehat{\nabla d_1(t, x)} \right) D^2 d_1(t, x).$$

The computation for (t, x) near Γ_t^2 is similar. We conclude that the fronts move according to their mean curvature:

$$\begin{cases} \partial_t d_1(t, x) \simeq \mu \operatorname{tr} \left((I - \widehat{\nabla d_1(t, x)}) \otimes \widehat{\nabla d_1(t, x)} \right) D^2 d_1(t, x) & \text{near } \Gamma_t^1 \\ \partial_t d_2(t, x) \simeq \mu \operatorname{tr} \left((I - \widehat{\nabla d_2(t, x)}) \otimes \widehat{\nabla d_2(t, x)} \right) D^2 d_2(t, x) & \text{near } \Gamma_t^2. \end{cases}$$

5.2. Derivation of (4.16). It is actually necessary to add a lower order correction to (5.1) for the ansatz to solve the fractional Allen-Chan equation (1.1). This was already observed in the one-dimensional case in [23, Section 3.1]. In order to showcase the equation for the corrector, for $\sigma \in \mathbb{R}$, let v^ε be the solution to

$$(5.5) \quad \varepsilon \partial_t v^\varepsilon = \frac{1}{\varepsilon |\ln \varepsilon|} (\varepsilon \mathcal{I}_n v^\varepsilon - W'(v^\varepsilon)) - \sigma,$$

with initial condition (1.7). Consider the simplest case in which $N = 1$ and assume that $d(t, x) = d_1(t, x)$ is smooth with $|\nabla d| = 1$ and satisfies

$$(5.6) \quad \partial_t d = \mu \Delta d - c_0 \sigma \simeq c_0 \bar{a}_\varepsilon(t, x) - c_0 \sigma,$$

in a neighborhood of Γ_t^1 . Consider the new ansatz

$$v^\varepsilon(t, x) \simeq \phi\left(\frac{d(t, x)}{\varepsilon}\right) + \varepsilon |\ln \varepsilon| \psi\left(\frac{d(t, x)}{\varepsilon}\right) - \varepsilon |\ln \varepsilon| \bar{\sigma},$$

where the function ψ and the constant $\bar{\sigma}$ are to be determined. Assume that ψ is smooth and bounded with bounded derivatives. This in particular implies that the estimate in Lemma 4.9 holds true.

Plugging the ansatz into (5.5), the left-hand side gives

$$(5.7) \quad \varepsilon \partial_t v^\varepsilon \simeq \dot{\phi}\left(\frac{d}{\varepsilon}\right) \partial_t d + \varepsilon |\ln \varepsilon| \dot{\psi}\left(\frac{d}{\varepsilon}\right) \partial_t d \simeq \dot{\phi}\left(\frac{d}{\varepsilon}\right) \partial_t d,$$

where we used that $\dot{\psi}$ and $\partial_t d$ are bounded. Next, we look at the right-hand side of (5.5) for the ansatz. First, we use the equation for ϕ (see (1.4)) and Lemmas 4.5 and 4.9 to find that

$$(5.8) \quad \begin{aligned} \frac{\varepsilon}{\varepsilon |\ln \varepsilon|} \mathcal{I}_n[v^\varepsilon] &\simeq \frac{\varepsilon}{\varepsilon |\ln \varepsilon|} \mathcal{I}_n \left[\phi\left(\frac{d}{\varepsilon}\right) \right] + \varepsilon \mathcal{I}_n \left[\psi\left(\frac{d}{\varepsilon}\right) \right] \\ &= \frac{1}{\varepsilon |\ln \varepsilon|} \left(\varepsilon \mathcal{I}_n \left[\phi\left(\frac{d}{\varepsilon}\right) \right] - C_n \mathcal{I}_1[\phi]\left(\frac{d}{\varepsilon}\right) \right) + \frac{1}{\varepsilon |\ln \varepsilon|} C_n \mathcal{I}_1[\psi]\left(\frac{d}{\varepsilon}\right) \\ &\quad + \left(\varepsilon \mathcal{I}_n \left[\psi\left(\frac{d}{\varepsilon}\right) \right] - C_n \mathcal{I}_1[\psi]\left(\frac{d}{\varepsilon}\right) \right) + C_n \mathcal{I}_1[\psi]\left(\frac{d}{\varepsilon}\right) \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} a_\varepsilon\left(\frac{d}{\varepsilon}\right) + \frac{1}{\varepsilon |\ln \varepsilon|} W'\left(\phi\left(\frac{d}{\varepsilon}\right)\right) + C_n \mathcal{I}_1[\psi]\left(\frac{d}{\varepsilon}\right). \end{aligned}$$

On the other hand, we do a Taylor expansion for W' around $\phi(d/\varepsilon)$ to estimate

$$(5.9) \quad \begin{aligned} \frac{1}{\varepsilon |\ln \varepsilon|} W'(v^\varepsilon) &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} \left[W'\left(\phi\left(\frac{d}{\varepsilon}\right)\right) + W''\left(\phi\left(\frac{d}{\varepsilon}\right)\right) \left(v^\varepsilon - \phi\left(\frac{d}{\varepsilon}\right)\right) \right] \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} \left[W'\left(\phi\left(\frac{d}{\varepsilon}\right)\right) + W''\left(\phi\left(\frac{d}{\varepsilon}\right)\right) \left(\varepsilon |\ln \varepsilon| \psi\left(\frac{d}{\varepsilon}\right) - \varepsilon |\ln \varepsilon| \bar{\sigma}\right) \right]. \end{aligned}$$

Equating (5.7) with (5.8) and (5.9), the equation for the ansatz gives

$$(5.10) \quad \begin{aligned} \dot{\phi}\left(\frac{d}{\varepsilon}\right) \partial_t d &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} a_\varepsilon\left(\frac{d}{\varepsilon}\right) + C_n \mathcal{I}_1[\psi]\left(\frac{d}{\varepsilon}\right) \\ &\quad - W''\left(\phi\left(\frac{d}{\varepsilon}\right)\right) \psi\left(\frac{d}{\varepsilon}\right) + \bar{\sigma} W''\left(\phi\left(\frac{d}{\varepsilon}\right)\right) - \sigma. \end{aligned}$$

Rearranging and using (5.6), we see that ψ satisfies

$$-C_n \mathcal{I}_1[\psi]\left(\frac{d}{\varepsilon}\right) + W''\left(\phi\left(\frac{d}{\varepsilon}\right)\right) \psi\left(\frac{d}{\varepsilon}\right)$$

$$\begin{aligned} &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} a_\varepsilon \left(\frac{d}{\varepsilon} \right) - \dot{\phi} \left(\frac{d}{\varepsilon} \right) \partial_t d + \tilde{\sigma} W'' \left(\phi \left(\frac{d}{\varepsilon} \right) \right) - \sigma \\ &\simeq \frac{1}{\varepsilon |\ln \varepsilon|} a_\varepsilon \left(\frac{d}{\varepsilon} \right) - \dot{\phi} \left(\frac{d}{\varepsilon} \right) c_0 \bar{a}_\varepsilon + c_0 \sigma \dot{\phi} \left(\frac{d}{\varepsilon} \right) + \tilde{\sigma} W'' \left(\phi \left(\frac{d}{\varepsilon} \right) \right) - \sigma. \end{aligned}$$

Evaluating the equation when $|d(t, x)| \gg \varepsilon$, from (4.2), (4.3), (4.8) and (4.10), if $\psi(\pm\infty) = 0$, we see that $\tilde{\sigma}$ must satisfy

$$\tilde{\sigma} W''(0) = \sigma.$$

This shows that the corrector $\psi = \psi(\xi; t, x)$ is solution to (4.16).

Note that for equation (4.16) to be solvable, the right hand-side $g = g(\xi; t, x)$ must satisfy the compatibility condition (4.19) guaranteed by Lemma 4.7. Indeed, multiply both sides of (4.16) by $\dot{\phi}(\xi)$ and integrate over \mathbb{R} to write

$$(5.11) \quad \int_{\mathbb{R}} \mathcal{L}[\psi](\xi) \dot{\phi}(\xi) d\xi = \int_{\mathbb{R}} g(\xi; t, x) \dot{\phi}(\xi) d\xi.$$

Since \mathcal{I}_1 is self-adjoint and ϕ satisfies (1.4), the left-hand side of the equation gives

$$(5.12) \quad \begin{aligned} \int_{\mathbb{R}} \mathcal{L}[\psi] \dot{\phi} d\xi &= \int_{\mathbb{R}} \left(-C_n \mathcal{I}_1[\dot{\phi}] + W''(\phi) \dot{\phi} \right) \psi d\xi \\ &= \int_{\mathbb{R}} \frac{d}{d\xi} \left(-C_n \mathcal{I}_1[\phi] + W'(\phi) \right) \psi d\xi = 0. \end{aligned}$$

Finally, note that in view of (5.11) and the above computations, the addition of the corrector ψ in the ansatz for the solution of (1.1)-(1.7) does not give any additional contributions in the derivation of the mean curvature equation obtained in Section 5.1.

Remark 5.1. Notice that ψ depends on the signed distance function $d(t, x)$ through a_ε and \bar{a}_ε . Hence, when $N > 1$, we have a finite sequence of correctors, denoted by ψ_1, \dots, ψ_N , depending on the signed distance function $d_i(t, x)$ to the front Γ_t^i , $i = 1, \dots, N$.

6. CONSTRUCTION OF BARRIERS

We now construct local and global strict subsolutions (supersolutions) to (1.1) needed for the proof of Theorem 1.1. We will focus on the construction of subsolutions as the construction of supersolutions is analogous. We will start with the global ones.

6.1. Global subsolutions. Fix $t_0 \in [0, \infty)$ and $h > 0$. For $i = 1, \dots, N$ and $t \in [t_0, t_0 + h]$, let Ω_t^i be bounded open sets with boundaries $\Gamma_t^i := \partial\Omega_t^i$ such that

$$(6.1) \quad \Omega_t^{i+1} \subset\subset \Omega_t^i \quad \text{for } i = 1, \dots, N-1.$$

Let $\tilde{d}_i(t, x)$ be the signed distance function associated to the set Ω_t^i , then $\Gamma_t^i = \{x \in \mathbb{R}^n : \tilde{d}_i(t, x) = 0\}$. Assume that there exists a $\rho > 0$ such that, for all $1 \leq i \leq N$, $\tilde{d}_i(t, x)$ is smooth in the set

$$(6.2) \quad Q_{2\rho}^i = \{(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n : |\tilde{d}_i(t, x)| \leq 2\rho\},$$

and let d_i be the smooth, bounded extension of \tilde{d}_i outside of Q_ρ^i as defined in Definition 2.2. Assume in addition that there exists $\sigma > 0$ such that

$$(6.3) \quad \partial_t d_i \leq \mu \Delta d_i - c_0 \sigma \quad \text{in } Q_\rho^i.$$

Let $\tilde{\sigma} > 0$ be such that $\sigma = W''(0)\tilde{\sigma}$. By (6.1), and perhaps making σ smaller, we can assume that, for all $t \in [t_0, t_0 + h]$,

$$(6.4) \quad \{x \in \mathbb{R}^n : |d_i(t, x)| < \tilde{\sigma}\} \cap \{x \in \mathbb{R}^n : |d_j(t, x)| < \tilde{\sigma}\} = \emptyset \quad i \neq j.$$

We define the smooth barrier $v^\varepsilon(t, x)$ by

$$(6.5) \quad v^\varepsilon(t, x) = \sum_{i=1}^N \phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon}; t, x \right) - \tilde{\sigma} \varepsilon |\ln \varepsilon|.$$

Lemma 6.1 (Global subsolutions to (1.1)). *Let ρ and γ be respectively as in Definitions 2.2 and 4.4. Assume (6.3) with μ as in (1.6) and (6.4), and let v^ε be defined as in (6.5). Then there exists $\varepsilon_0 = \varepsilon_0(\rho, \gamma, \sigma) > 0$ such that for any $0 < \varepsilon < \varepsilon_0$, v^ε is a solution to*

$$(6.6) \quad \varepsilon \partial_t v^\varepsilon - \frac{1}{\varepsilon |\ln \varepsilon|} (\varepsilon \mathcal{I}_n[v^\varepsilon] - W'(v^\varepsilon)) \leq -\frac{\sigma}{2} \quad \text{in } [t_0, t_0 + h] \times \mathbb{R}^n.$$

Moreover, there is a constant $C > 0$ such that, for $\varepsilon < \varepsilon_0$,

$$(6.7) \quad v^\varepsilon(t, x) \geq N - 2\tilde{\sigma}\varepsilon |\ln \varepsilon| \quad \text{in } \left\{ (t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n : d_N(t, x) - \tilde{\sigma} \geq \frac{C}{\tilde{\sigma} |\ln \varepsilon|} \right\}.$$

Proof. We will break the proof into four main steps. First, we estimate the equation for $v^\varepsilon(t, x)$ for any (t, x) . Then, we will show that $v^\varepsilon(t, x)$ satisfies (6.6) when (t, x) is near a single front $\Gamma_t^{i_0}$ and then when (t, x) is far from all fronts Γ_t^i , $i = 1, \dots, N$. Lastly, we prove the estimate in (6.7). For convenience, we use the following notation throughout the proof:

$$(6.8) \quad \begin{aligned} \phi_i &:= \phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) \\ \psi_i &:= \psi_i \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon}; t, x \right) \\ \tilde{\phi}_i &:= \phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) - H \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) \\ a_\varepsilon^i &:= a_\varepsilon \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon}; t, x \right) \\ \bar{a}_\varepsilon^i &:= \bar{a}_\varepsilon(t, x) \quad \text{corresponding to } a_\varepsilon^i. \end{aligned}$$

We note that it will be important for the reader to remember the dependence of ψ_i and a_ε on the variables t, x and $\xi = d_i(t, x)/\varepsilon$ when taking derivatives in t, x .

Step 1. Computation for $v^\varepsilon(t, x)$ in (1.1) for an arbitrary $(t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n$.

First, the time derivative of v^ε at (t, x) is given by

$$\varepsilon \partial_t v^\varepsilon(t, x) = \sum_{i=1}^N \dot{\phi}_i \partial_t d_i(t, x) + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \left[\dot{\psi}_i \partial_t d_i(t, x) + \varepsilon \partial_t \psi_i \right].$$

By (4.20) for $\dot{\psi}_i$ and $\partial_t \psi_i$,

$$(6.9) \quad \varepsilon \partial_t v^\varepsilon = \sum_{i=1}^N \dot{\phi}_i \partial_t d_i(t, x) + O(\varepsilon^{\frac{1}{2}}) + O(\varepsilon |\ln \varepsilon|) = \sum_{i=1}^N \dot{\phi}_i \partial_t d_i(t, x) + O(\varepsilon^{\frac{1}{2}}).$$

Next, we consider the nonlocal term. For each $i = 1, \dots, N$, we use that ϕ satisfies (1.4) and apply Lemma 4.5 to find

$$\begin{aligned} \varepsilon \mathcal{I}_n[\phi_i](x) &= \varepsilon \mathcal{I}_n[\phi_i](x) - C_n \mathcal{I}_1[\phi] \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) + W'(\phi_i) \\ &= a_\varepsilon^i + O(\varepsilon) + W'(\phi_i). \end{aligned}$$

Also, using that ψ satisfies (4.16) and Lemma 4.9, we find that

$$\begin{aligned}\varepsilon \mathcal{I}_n[\psi_i](x) &= \varepsilon \mathcal{I}_n[\psi_i](x) - C_n \mathcal{I}_1[\psi_i] \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) - \mathcal{L}[\psi] \left(\frac{d_i(t, \cdot) - \tilde{\sigma}}{\varepsilon} \right) + W''(\phi_i) \psi_i \\ &= O(|\ln \varepsilon|^{-\frac{1}{2}}) - \frac{a_\varepsilon^i}{\varepsilon |\ln \varepsilon|} + \dot{\phi}_i c_0 (\bar{a}_\varepsilon^i - \sigma) - \tilde{\sigma} (W''(\phi_i) - W''(0)) + W''(\phi_i) \psi_i.\end{aligned}$$

Therefore, the fractional Laplacian of v^ε can be written as

$$\begin{aligned}\varepsilon \mathcal{I}_n[v^\varepsilon](x) &= \sum_{i=1}^N [a_\varepsilon^i + O(\varepsilon) + W'(\phi_i)] \\ &\quad + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \left[O(|\ln \varepsilon|^{-\frac{1}{2}}) - \frac{a_\varepsilon^i}{\varepsilon |\ln \varepsilon|} + \dot{\phi}_i c_0 (\bar{a}_\varepsilon^i - \sigma) \right. \\ &\quad \left. - \tilde{\sigma} (W''(\phi_i) - W''(0)) + W''(\phi_i) \psi_i \right].\end{aligned}$$

Recall the definition of $\tilde{\phi}_i$ introduced in (6.8). Since W is periodic, we have that $W'(\phi_i) = W'(\tilde{\phi}_i)$ and similarly $W''(\phi_i) = W''(\tilde{\phi}_i)$. Using this and rearranging, we equivalently write

$$\begin{aligned}(6.10) \quad \varepsilon \mathcal{I}_n[v^\varepsilon](x) &= \sum_{i=1}^N W'(\tilde{\phi}_i) + O(\varepsilon) + O(\varepsilon |\ln \varepsilon|^{\frac{1}{2}}) \\ &\quad + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \left[W''(\tilde{\phi}_i) \psi_i + \dot{\phi}_i c_0 (\bar{a}_\varepsilon^i - \sigma) - \tilde{\sigma} (W''(\tilde{\phi}_i) - W''(0)) \right].\end{aligned}$$

Then, with (6.9) and (6.10), the equation for v^ε at (t, x) can be written as

$$\begin{aligned}\text{Eqn}(v^\varepsilon) &:= \varepsilon \partial_t v^\varepsilon(t, x) - \frac{1}{\varepsilon |\ln \varepsilon|} (\varepsilon \mathcal{I}_n[v^\varepsilon(t, \cdot)](x) - W'(v^\varepsilon(t, x))) \\ &= O(\varepsilon^{\frac{1}{2}}) + \sum_{i=1}^N \dot{\phi}_i \partial_t d_i(t, x) \\ &\quad - \frac{1}{\varepsilon |\ln \varepsilon|} \left\{ \sum_{i=1}^N W'(\tilde{\phi}_i) + O(\varepsilon) + O(\varepsilon |\ln \varepsilon|^{\frac{1}{2}}) \right. \\ &\quad \left. + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \left[W''(\tilde{\phi}_i) \psi_i + \dot{\phi}_i c_0 (\bar{a}_\varepsilon^i - \sigma) - \tilde{\sigma} (W''(\tilde{\phi}_i) - W''(0)) \right] \right. \\ &\quad \left. - W' \left(\sum_{i=1}^N \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right) \right\}.\end{aligned}$$

Grouping the error terms, the $\dot{\phi}_i$ terms, and the nonlinear terms together, we have

$$(6.11) \quad \begin{aligned} \text{Eqn}(v^\varepsilon) &= O(|\ln \varepsilon|^{-\frac{1}{2}}) + \sum_{i=1}^N \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] \\ &+ \frac{1}{\varepsilon |\ln \varepsilon|} \left\{ W' \left(\sum_{i=1}^N \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right) \right. \\ &\quad \left. - \sum_{i=1}^N \left(W'(\tilde{\phi}_i) + \varepsilon |\ln \varepsilon| \left[W''(\tilde{\phi}_i) \psi_i - \tilde{\sigma} (W''(\tilde{\phi}_i) - W''(0)) \right] \right) \right\}. \end{aligned}$$

Fix an index $i_0 \in \{1, \dots, N\}$. For the remainder of Step 1, we will conveniently isolate every term indexed with i_0 to help with Step 2. First, we do a Taylor expansion for W' around $\tilde{\phi}_{i_0}$ to obtain

$$(6.12) \quad \begin{aligned} &W' \left(\sum_{i=1}^N \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right) \\ &= W'(\tilde{\phi}_{i_0}) + W''(\tilde{\phi}_{i_0}) \left(\sum_{i \neq i_0} \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right) \\ &\quad + O \left(\left(\sum_{i \neq i_0} \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right)^2 \right). \end{aligned}$$

By (4.20) for ψ_i , we have that

$$\begin{aligned} &\frac{1}{\varepsilon |\ln \varepsilon|} O \left(\left(\sum_{i \neq i_0} \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right)^2 \right) \\ &= \sum_{i \neq i_0} O \left(\frac{(\tilde{\phi}_i)^2}{\varepsilon |\ln \varepsilon|} \right) + O(|\ln \varepsilon|^{-1}) + O(\varepsilon |\ln \varepsilon|). \end{aligned}$$

Hence, with (6.12) and grouping error terms, we can write (6.11) as

$$\begin{aligned} \text{Eqn}(v^\varepsilon) &= O(|\ln \varepsilon|^{-\frac{1}{2}}) + \sum_{i \neq i_0} O \left(\frac{(\tilde{\phi}_i)^2}{\varepsilon |\ln \varepsilon|} \right) + \sum_{i=1}^N \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] \\ &+ \frac{1}{\varepsilon |\ln \varepsilon|} \left\{ W'(\tilde{\phi}_{i_0}) + W''(\tilde{\phi}_{i_0}) \left(\sum_{i \neq i_0} \tilde{\phi}_i + \varepsilon |\ln \varepsilon| \sum_{i=1}^N \psi_i - \tilde{\sigma} \varepsilon |\ln \varepsilon| \right) \right. \\ &\quad - \left(W'(\tilde{\phi}_{i_0}) + \varepsilon |\ln \varepsilon| \left[W''(\tilde{\phi}_{i_0}) \psi_{i_0} - \tilde{\sigma} (W''(\tilde{\phi}_{i_0}) - W''(0)) \right] \right) \\ &\quad \left. - \sum_{i \neq i_0} \left(W'(\tilde{\phi}_i) + \varepsilon |\ln \varepsilon| \left[W''(\tilde{\phi}_i) \psi_i - \tilde{\sigma} (W''(\tilde{\phi}_i) - W''(0)) \right] \right) \right\} \end{aligned}$$

where in the last two lines we extracted the i_0 term. Cancelling the $W'(\tilde{\phi}_{i_0})$ and $W''(\tilde{\phi}_{i_0})\psi_{i_0}$ terms then distributing $1/(\varepsilon |\ln \varepsilon|)$, we simplify to

$$\begin{aligned} \text{Eqn}(v^\varepsilon) &= O(|\ln \varepsilon|^{-\frac{1}{2}}) + \sum_{i \neq i_0} O\left(\frac{(\tilde{\phi}_i)^2}{\varepsilon |\ln \varepsilon|}\right) + \sum_{i=1}^N \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] \\ &\quad + W''(\tilde{\phi}_{i_0}) \left(\sum_{i \neq i_0} \frac{\tilde{\phi}_i}{\varepsilon |\ln \varepsilon|} + \sum_{i \neq i_0} \psi_i - \tilde{\sigma} \right) + \tilde{\sigma} \left(W''(\tilde{\phi}_{i_0}) - W''(0) \right) \\ &\quad - \sum_{i \neq i_0} \left[\frac{W'(\tilde{\phi}_i)}{\varepsilon |\ln \varepsilon|} + W''(\tilde{\phi}_i)\psi_i - \tilde{\sigma} \left(W''(\tilde{\phi}_i) - W''(0) \right) \right]. \end{aligned}$$

Next, we do a Taylor expansion for W' around 0 and recall that $W'(0) = 0$ to write

$$W'(\tilde{\phi}_i) = W'(0) + W''(0)\tilde{\phi}_i + O((\tilde{\phi}_i)^2) = W''(0)\tilde{\phi}_i + O((\tilde{\phi}_i)^2).$$

With this, we now have that

$$\begin{aligned} \text{Eqn}(v^\varepsilon) &= O(|\ln \varepsilon|^{-\frac{1}{2}}) + \sum_{i \neq i_0} O\left(\frac{(\tilde{\phi}_i)^2}{\varepsilon |\ln \varepsilon|}\right) + \sum_{i=1}^N \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] \\ &\quad + W''(\tilde{\phi}_{i_0}) \left(\sum_{i \neq i_0} \frac{\tilde{\phi}_i}{\varepsilon |\ln \varepsilon|} + \sum_{i \neq i_0} \psi_i - \tilde{\sigma} \right) + \tilde{\sigma} \left(W''(\tilde{\phi}_{i_0}) - W''(0) \right) \\ &\quad - \sum_{i \neq i_0} \left[\frac{W''(0)\tilde{\phi}_i}{\varepsilon |\ln \varepsilon|} + W''(\tilde{\phi}_i)\psi_i - \tilde{\sigma} \left(W''(\tilde{\phi}_i) - W''(0) \right) \right]. \end{aligned}$$

We rearrange to group the terms with $W''(\tilde{\phi}_{i_0}) - W''(0)$ together

$$\begin{aligned} \text{Eqn}(v^\varepsilon) &= O(|\ln \varepsilon|^{-\frac{1}{2}}) + \sum_{i \neq i_0} O\left(\frac{(\tilde{\phi}_i)^2}{\varepsilon |\ln \varepsilon|}\right) + \sum_{i=1}^N \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] \\ (6.13) \quad &\quad + \left(W''(\tilde{\phi}_{i_0}) - W''(0) \right) \sum_{i \neq i_0} \frac{\tilde{\phi}_i}{\varepsilon |\ln \varepsilon|} - \tilde{\sigma} W''(0) \\ &\quad + \sum_{i \neq i_0} \left[\left(W''(\tilde{\phi}_{i_0}) - W''(\tilde{\phi}_i) \right) \psi_i + \tilde{\sigma} \left(W''(\tilde{\phi}_i) - W''(0) \right) \right]. \end{aligned}$$

Looking at the last two lines for $i \neq i_0$, note first that $(W''(\tilde{\phi}_{i_0}) - W''(0))\tilde{\phi}_i = O(\tilde{\phi}_i)$ and then Taylor expand W'' around 0 to find

$$\begin{aligned} (W''(\tilde{\phi}_{i_0}) - W''(\tilde{\phi}_i))\psi_i + \tilde{\sigma} \left(W''(\tilde{\phi}_i) - W''(0) \right) &= O(\psi_i) - \tilde{\sigma} \left(W'''(0)\tilde{\phi}_i + O(\tilde{\phi}_i^2) \right) \\ &= O(\psi_i) + O(\tilde{\phi}_i). \end{aligned}$$

Therefore, (6.13) can be written as

$$\begin{aligned} \text{Eqn}(v^\varepsilon) &= O(|\ln \varepsilon|^{-\frac{1}{2}}) + \sum_{i \neq i_0} \left[O\left(\frac{\tilde{\phi}_i}{\varepsilon |\ln \varepsilon|}\right) + O(\psi_i) \right] \\ (6.14) \quad &\quad + \sum_{i=1}^N \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] - \sigma. \end{aligned}$$

where we used that $\sigma = W''(0)\tilde{\sigma}$.

Step 2. $v^\varepsilon(t, x)$ satisfies (6.6) when (t, x) is near the front $\Gamma_t^{i_0}$.

Assume that $|d_{i_0}(t, x) - \tilde{\sigma}| \leq |\ln \varepsilon|^{-1/2}$ for some index $1 \leq i_0 \leq N$. Then, by (6.4), for ε sufficiently small,

$$|d_i(t, x) - \tilde{\sigma}| \geq |\ln \varepsilon|^{-\frac{1}{2}} \quad \text{for all } i \neq i_0.$$

We begin by estimating the error terms in (6.14) for $i \neq i_0$. First, we use (4.2) to estimate

$$|\tilde{\phi}_i| = \left| \phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) - H \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) \right| \leq C \frac{\varepsilon}{|d_i(t, x) - \tilde{\sigma}|} = O(\varepsilon |\ln \varepsilon|^{\frac{1}{2}}),$$

from which it follows that

$$\frac{|\tilde{\phi}_i|}{\varepsilon |\ln \varepsilon|} = O\left(|\ln \varepsilon|^{-\frac{1}{2}}\right).$$

Similarly, from (4.21), for $i \neq i_0$,

$$|\psi_i| \leq \frac{C}{\varepsilon |\ln \varepsilon|} \frac{\varepsilon}{|d_i(t, x) - \tilde{\sigma}|} = O\left(|\ln \varepsilon|^{-\frac{1}{2}}\right).$$

Combining the above estimates in view of (6.14), we have

$$\sum_{i \neq i_0} \left(O\left(\frac{\tilde{\phi}_i}{\varepsilon |\ln \varepsilon|}\right) + O(\psi_i) \right) = O\left(|\ln \varepsilon|^{-\frac{1}{2}}\right).$$

Next, we check the terms with the auxiliary functions \bar{a}_ε^i and a_ε^i . For $i \neq i_0$, we use that d_i is smooth, (4.3) and (4.10) to obtain

$$(6.15) \quad \left| \sum_{i \neq i_0} \dot{\phi}_i [\partial_t d_i(t, x) - c_0 (\bar{a}_\varepsilon^i - \sigma)] \right| \leq \sum_{i \neq i_0} \left(\frac{\varepsilon}{|d_i(t, x) - \tilde{\sigma}|} \right)^2 \frac{C}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|} = O\left(\varepsilon^{\frac{3}{2}}\right).$$

For $i = i_0$, we use that $\dot{\phi}_{i_0} \geq 0$, (6.3), and Theorem 4.2 to estimate

$$\begin{aligned} \dot{\phi}_{i_0} [\partial_t d_{i_0}(t, x) - c_0 (\bar{a}_\varepsilon^{i_0} - \sigma)] &= \dot{\phi}_{i_0} ([\partial_t d_{i_0}(t, x) - \mu \Delta d_{i_0}(t, x) + c_0 \sigma] + [\mu \Delta d_{i_0}(t, x) - c_0 \bar{a}_\varepsilon^{i_0}]) \\ &\leq \dot{\phi}_{i_0} (0 + o_\varepsilon(1)) = o_\varepsilon(1). \end{aligned}$$

Consequently, in (6.14), we have that

$$\text{Eqn}(v^\varepsilon) \leq o_\varepsilon(1) - \sigma.$$

Taking ε sufficiently small, (6.6) holds.

Step 3. $v^\varepsilon(t, x)$ satisfies (6.6) when (t, x) is away from all fronts Γ_t^i .

Assume that for all $i = 1, \dots, N$,

$$|d_i(t, x) - \tilde{\sigma}| \geq |\ln \varepsilon|^{-\frac{1}{2}}.$$

Then, we estimate exactly as in Step 2 but we include $i = i_0$ in (6.15) to obtain

$$\text{Eqn}(v^\varepsilon) \leq o_\varepsilon(1) - \sigma.$$

Taking ε sufficiently small, (6.6) holds.

Step 4. $v^\varepsilon(t, x)$ satisfies (6.7).

Let $\tilde{\rho} := \frac{C}{\tilde{\sigma}|\ln \varepsilon|}$ for $C > 0$ to be determined and $\varepsilon > 0$ sufficiently small. Fix (t, x) such that $d_N(t, x) - \tilde{\sigma} \geq \tilde{\rho}$. Then,

$$d_i(t, x) - \tilde{\sigma} \geq \tilde{\rho} \quad \text{for all } 1 \leq i \leq N,$$

and by (4.2) and (4.21),

$$\begin{aligned} v^\varepsilon(t, x) &\geq N + \sum_{i=1}^N \left(\phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) - H \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) \right) - C \sum_{i=1}^N \frac{\varepsilon}{d_i(t, x) - \tilde{\sigma}} - \tilde{\sigma} \varepsilon |\ln \varepsilon| \\ &\geq N - \sum_{i=1}^N \left(\frac{\varepsilon}{\alpha(d_i(t, x) - \tilde{\sigma})} + \frac{C\varepsilon^2}{(d_i(t, x) - \tilde{\sigma})^2} \right) - C \sum_{i=1}^N \frac{\varepsilon}{d_i(t, x) - \tilde{\sigma}} - \tilde{\sigma} \varepsilon |\ln \varepsilon| \\ &\geq N - \frac{C\varepsilon}{\rho} - \frac{C\varepsilon^2}{\rho^2} - \tilde{\sigma} \varepsilon |\ln \varepsilon| \\ &\geq N - 2\tilde{\sigma} \varepsilon |\ln \varepsilon| \end{aligned}$$

for C sufficiently large in the choice of $\tilde{\rho}$. This proves (6.7). \square

6.2. Local subsolutions. We now construct local subsolutions to (1.1). For $i = 1, \dots, N$ and $t \in [t_0, t_0 + h]$, let Ω_t^i be bounded open sets satisfying (6.1). Assume that there exists R, ρ and $x_0 \in \mathbb{R}^n$ such that the signed distance function $\tilde{d}_i(t, x)$ associated to the set Ω_t^i is smooth in the set $Q_{2\rho}^i \cap ([t_0, t_0 + h] \times B(x_0, R))$ (recall (6.2)). Let d_i be the bounded extension of \tilde{d}_i outside of Q_ρ^i as defined in Definition 2.2, then d_i is smooth in $[t_0, t_0 + h] \times B(x_0, R)$. Assume that there exist $\sigma > 0$ and $0 < r < R$ such that,

$$(6.16) \quad \partial_t d_i \leq \mu \Delta d_i - c_0 \sigma \quad \text{in } Q_\rho \cap ([t_0, t_0 + h] \times B(x_0, r)).$$

As before, let $\tilde{\sigma} > 0$ be such that $\sigma = W''(0)\tilde{\sigma}$, then, by eventually making σ smaller, we can assume that, for all $t \in [t_0, t_0 + h]$, (6.4) holds true.

The following lemma is the local version of Lemma 6.1.

Lemma 6.2 (Local subsolutions to (1.1)). *For $0 < r < R$, let ρ and γ be respectively as in Definitions 2.2 and 4.4 be such that $0 < \rho, \gamma < R - r$. Assume that d_i is smooth in $[t_0, t_0 + h] \times B(x_0, R)$ and that (6.16) with μ as in (1.6) and (6.4) hold. Let v^ε be defined as in (6.5). Then there exists $\varepsilon_0 = \varepsilon_0(\rho, \gamma, \sigma) > 0$ such that for any $0 < \varepsilon < \varepsilon_0$, v^ε is a solution to*

$$\varepsilon \partial_t v^\varepsilon - \frac{1}{\varepsilon |\ln \varepsilon|} (\varepsilon \mathcal{I}_n[v^\varepsilon] - W'(v^\varepsilon)) \leq -\frac{\sigma}{2} \quad \text{in } [t_0, t_0 + h] \times B(x_0, r).$$

Moreover, there is a constant $C > 0$ such that, for $\varepsilon < \varepsilon_0$,

$$(6.17) \quad v^\varepsilon(t, x) \geq N - 2\tilde{\sigma} \varepsilon |\ln \varepsilon| \quad \text{in } \left\{ (t, x) \in [t_0, t_0 + h] \times \mathbb{R}^n : d_N(t, x) - \tilde{\sigma} \geq \frac{C}{\tilde{\sigma} |\ln \varepsilon|} \right\}.$$

Proof. The proof of the lemma is exactly as the proof of Lemma 6.1 except that we now only consider points $(t, x) \in [t_0, t_0 + h] \times B(x_0, r)$. Note that if $0 < \gamma < R - r$ and $x \in B(x_0, r)$, then $d(t, x + \varepsilon(\cdot))$ is smooth at all points in the domain of integration of a_ε and all estimates in Lemma 4.3 and Theorem 4.8 hold true. In addition, the smoothness of $d(t, \cdot)$ in $B(x_0, R)$ and its boundedness in the whole \mathbb{R}^n are sufficient to guarantee that for $0 < \rho < R - r$ and $x \in B(x_0, r)$, Lemmas 4.5 and 4.9 hold true, see Sections 10 and 11. \square

7. PROOF OF THEOREM 1.1

Proof. We apply an adaptation of the abstract method in [4, 6] as described in Section 2.2.

Begin by defining the families of open sets $(D^i)_{i=1}^N$ and $(E^i)_{i=1}^N$ by

$$D^i = \text{Int} \left\{ (t, x) \in (0, \infty) \times \mathbb{R}^n : \liminf_{\varepsilon \rightarrow 0} * \frac{u^\varepsilon(t, x) - i}{\varepsilon |\ln \varepsilon|} \geq 0 \right\} \subset (0, \infty) \times \mathbb{R}^n$$

$$E^i = \text{Int} \left\{ (t, x) \in (0, \infty) \times \mathbb{R}^n : \limsup_{\varepsilon \rightarrow 0} * \frac{u^\varepsilon(t, x) - (i - 1)}{\varepsilon |\ln \varepsilon|} \leq 0 \right\} \subset (0, \infty) \times \mathbb{R}^n.$$

To define the traces of D^i and E^i , we first define the functions $\underline{\chi}^i, \bar{\chi}^i : (0, \infty) \times \mathbb{R}^n \rightarrow \{-1, 1\}$, respectively, by

$$\underline{\chi}^i = \mathbb{1}_{D^i} - \mathbb{1}_{(D^i)^c} \quad \text{and} \quad \bar{\chi}^i = \mathbb{1}_{(E^i)^c} - \mathbb{1}_{E^i}.$$

Since D^i is open, $\underline{\chi}^i$ is lower semicontinuous, and since $(E^i)^c$ is closed, $\bar{\chi}^i$ is upper semicontinuous. To ensure that $\bar{\chi}^i$ and $\underline{\chi}^i$ remain lower and upper semicontinuous, respectively, at $t = 0$, we set

$$\underline{\chi}^i(0, x) = \liminf_{t \rightarrow 0, y \rightarrow x} \underline{\chi}^i(t, y) \quad \text{and} \quad \bar{\chi}^i(0, x) = \limsup_{t \rightarrow 0, y \rightarrow x} \bar{\chi}^i(t, y).$$

Define the traces D_0^i and E_0^i by

$$D_0^i = \{x \in \mathbb{R}^n : \underline{\chi}^i(0, x) = 1\} \quad \text{and} \quad E_0^i = \{x \in \mathbb{R}^n : \bar{\chi}^i(0, x) = -1\}.$$

Note that D_0^i and E_0^i are open sets. For $t > 0$, define the sets D_t^i and E_t^i by

$$D_t^i = \{x \in \mathbb{R}^n : (t, x) \in D^i\} \quad \text{and} \quad E_t^i = \{x \in \mathbb{R}^n : (t, x) \in E^i\}.$$

We need the following propositions for the abstract method. Their proofs are delayed until the end of the section.

Proposition 7.1 (Initialization). *For each $i = 1, \dots, N$,*

$$\Omega_0^i \subset D_0^i \quad \text{and} \quad (\bar{\Omega}_0^i)^c \subset E_0^i.$$

Proposition 7.2 (Propagation). *For each $i = 1, \dots, N$, $(D_t^i)_{t>0}$ is a generalized super-flow, and $((E_t^i)^c)_{t>0}$ is a generalized sub-flow, according to Definition 2.1.*

By [4, Theorem 1.1, Theorem 1.2], it follows from Propositions 7.1 and 7.2 that

$${}^+\Omega_t^i \subset D_t^i \subset {}^+\Omega_t^i \cup \Gamma_t^i \quad \text{and} \quad {}^-\Omega_t^i \subset E_t^i \subset {}^-\Omega_t^i \cup \Gamma_t^i.$$

The conclusion readily follows; we write the details for completeness.

First, since ${}^+\Omega_t^i \subset D_t^i$, we use the definition of D_t^i to see that

$$(7.1) \quad \liminf_{\varepsilon \rightarrow 0} * u^\varepsilon(t, x) \geq i \quad \text{for } x \in {}^+\Omega_t^i.$$

Using that ${}^-\Omega_t^{i+1} \subset E_t^{i+1}$, we similarly get

$$(7.2) \quad \limsup_{\varepsilon \rightarrow 0} * u^\varepsilon(t, x) \leq (i + 1) - 1 = i \quad \text{for } x \in {}^-\Omega_t^{i+1}.$$

Therefore, for $i = 1, \dots, N - 1$,

$$\lim_{\varepsilon \rightarrow 0} u^\varepsilon(t, x) = i \quad \text{in } {}^+\Omega_t^i \cap {}^-\Omega_t^{i+1}.$$

Now, since constant functions solve (1.1) and $0 \leq u_0^\varepsilon \leq N$, by the comparison principle, we know $0 \leq u^\varepsilon \leq N$. In particular,

$$0 \leq \liminf_{\varepsilon \rightarrow 0} * u^\varepsilon \quad \text{and} \quad \limsup_{\varepsilon \rightarrow 0} * u^\varepsilon \leq N.$$

Together with (7.2) and respectively (7.1) we have

$$\lim_{\varepsilon \rightarrow 0} u^\varepsilon(t, x) = 0 \quad \text{in } {}^{-}\Omega_t^1 \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} u^\varepsilon(t, x) = N \quad \text{in } {}^{+}\Omega_t^N.$$

□

It remains to prove Propositions 7.1 and 7.2.

7.1. Proof of Proposition 7.1.

Proof. We will prove that $\Omega_0^{i_0} \subset D_0^{i_0}$ for all $1 \leq i_0 \leq N$. The proof of $(\overline{\Omega}_0^{i_0})^c \subset E_0^{i_0}$ is similar.

Fix i_0 and a point $x_0 \in \Omega_0^{i_0}$. To prove that $x_0 \in D_0^{i_0}$, it is enough to show that, for all (t, x) in a neighborhood of $(0, x_0)$ in $[0, \infty) \times \mathbb{R}^n$,

$$\liminf_{\varepsilon \rightarrow 0} * \frac{u^\varepsilon(t, x) - i_0}{\varepsilon |\ln \varepsilon|} \geq 0.$$

For this, we will use (6.5) to construct a suitable (global in space) subsolution $v^\varepsilon \leq u^\varepsilon$, depending on a small constant $\tilde{\sigma} > 0$.

For $0 < \tilde{\sigma} < d_i^0(x_0)$, where d_i^0 is defined in (1.5), let $r_i > 0$ be given by

$$(7.3) \quad r_i = d_i^0(x_0) - \tilde{\sigma}, \quad i = 1, \dots, i_0.$$

Note that $B(x_0, r_i) \subset \subset \Omega_0^i$, and since the sets Γ_0^i are separated, for $\tilde{\sigma}$ small enough,

$$(7.4) \quad r_i - r_{i+1} = d_i^0(x_0) - d_{i+1}^0(x_0) \geq d(\Gamma_0^i, \Gamma_0^{i+1}) \geq 4\tilde{\sigma}.$$

For a constant $C > 0$, to be determined and for $t \leq r_{i_0}/(2C) \leq r_i/(2C)$, let $\tilde{d}_i(t, x)$ be the signed distance function associated to the ball $B(x_0, r_i - Ct)$, namely

$$\tilde{d}_i(t, x) = r_i - Ct - |x - x_0|, \quad i = 1, \dots, i_0.$$

Then, by (7.3) and (7.4),

$$(7.5) \quad d_i^0(x) - \tilde{\sigma} \geq \tilde{d}_i(0, x) \geq \tilde{d}_{i+1}(0, x) + 4\tilde{\sigma}.$$

For $0 < \rho < r_{i_0}/4$, let d_i be the smooth, bounded extension of \tilde{d}_i outside of

$$Q_\rho^i = \left\{ (t, x) \in \left[0, \frac{r_{i_0}}{2C}\right] \times \mathbb{R}^n : |\tilde{d}_i(t, x)| \leq \rho \right\},$$

as in Definition 2.2. Then, for $(t, x) \in Q_\rho^i$, we have that $d_i = \tilde{d}_i$, $|x - x_0| \geq r_{i_0}/4$ and

$$\partial_t d_i(t, x) = -C, \quad D^2 d_i(t, x) = -\frac{1}{|x - x_0|} \left(I - \frac{(x - x_0) \otimes (x - x_0)}{|x - x_0|^2} \right)$$

from which we find that

$$(7.6) \quad \partial_t d_i(t, x) - \mu \Delta d_i(t, x) = -C + \mu \frac{n-1}{|x - x_0|} \leq -C + 4\mu \frac{n-1}{r_{i_0}} \leq -c_0 \sigma$$

for $C > 0$ sufficiently large, with $\sigma = W''(0)\tilde{\sigma}$ and μ as in (1.6). Moreover, we can assume, by eventually making $\tilde{\sigma}$ smaller, that $\rho > 2\tilde{\sigma}$.

Let $v^\varepsilon = v^\varepsilon(t, x)$ be given by

$$v^\varepsilon(t, x) = \sum_{i=1}^{i_0} \phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) + \varepsilon |\ln \varepsilon| \sum_{i=1}^{i_0} \psi_i \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon}; t, x \right) - \tilde{\sigma} \varepsilon |\ln \varepsilon|.$$

Note that v^ε is the same as (6.5) except we only sum over $1 \leq i \leq i_0$. By (7.5) and the fact that the balls $B(x_0, r_i - Ct)$ shrink with the same constant velocity, condition (6.4) is satisfied in the time interval $[0, r_{i_0}/(2C)]$. Since in addition (7.6) holds true, we are in position to apply Lemma 6.1 (with $N = i_0$) to conclude that v^ε is a solution to (6.6) in

$[0, r_{i_0}/(2C)] \times \mathbb{R}^n$ for ε small enough. We claim that choosing γ in (4.4) sufficiently small (but independent of ε), and ε sufficiently small,

$$(7.7) \quad v^\varepsilon(0, x) \leq u^\varepsilon(0, x) = u_0^\varepsilon(x) \quad \text{for all } x \in \mathbb{R}^n,$$

with u_0^ε as in (1.7). We break into two cases: we first consider the case when x is near the boundary of one of the balls, say $\partial B(x_0, r_j)$ for some j , and then when x is far from the boundaries of all balls $\partial B(x_0, r_i)$, $i = 1, \dots, i_0$.

Case 1: *There is a $j \in \{1, \dots, i_0\}$ such that $|\tilde{d}_j(0, x) - \tilde{\sigma}| \leq \frac{C_0}{|\ln \varepsilon|}$ for $C_0 > 0$ to be determined.*

For ε small enough, $\frac{C_0}{|\ln \varepsilon|} + \tilde{\sigma} \leq 2\tilde{\sigma} < \rho$, so that $0 \leq \tilde{d}_j(0, x) \leq 2\tilde{\sigma} < \rho$. Hence, recalling the Definition 2.2 of d_i and by the second inequality in (7.5), we have that

$$\begin{aligned} d_j(0, x) &= \tilde{d}_j(0, x), \\ d_i(0, x) - \tilde{\sigma} &\geq \tilde{\sigma} \quad \text{for } i = 1, \dots, j-1, \\ d_i(0, x) - \tilde{\sigma} &\leq -\tilde{\sigma} \quad \text{for } i = j+1, \dots, i_0. \end{aligned}$$

Therefore, by (4.2) and (4.23),

$$\phi\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}\right) + \varepsilon |\ln \varepsilon| \psi_i\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}; 0, x\right) \leq \begin{cases} 1 + \frac{C\varepsilon}{\tilde{\sigma}} & \text{for } i = 1, \dots, j-1 \\ \frac{C\varepsilon}{\tilde{\sigma}} & \text{for } i = j+1, \dots, i_0, \end{cases}$$

and by the monotonicity of ϕ , (4.2) and (4.22),

$$\begin{aligned} \phi\left(\frac{d_j(0, x) - \tilde{\sigma}}{\varepsilon}\right) + \varepsilon |\ln \varepsilon| \psi_j\left(\frac{d_j(0, x) - \tilde{\sigma}}{\varepsilon}; 0, x\right) &\leq \phi\left(\frac{C_0}{\varepsilon |\ln \varepsilon|}\right) + C\varepsilon |\ln \varepsilon| \\ &\leq 1 - \frac{C\varepsilon |\ln \varepsilon|}{C_0} + C\varepsilon |\ln \varepsilon|. \end{aligned}$$

From the last two estimates, we deduce that, for ε small enough,

$$v^\varepsilon(0, x) \leq j - \frac{C\varepsilon |\ln \varepsilon|}{C_0} + C\varepsilon |\ln \varepsilon| - \tilde{\sigma}\varepsilon |\ln \varepsilon| \leq j - C\varepsilon |\ln \varepsilon|,$$

if C_0 is sufficiently small.

Similarly, by the first inequality in (7.5), (4.2), and using that $\phi > 0$,

$$\phi\left(\frac{d_i^0(x)}{\varepsilon}\right) \geq \begin{cases} 1 - \frac{C\varepsilon}{\tilde{\sigma}} & \text{for } i = 1, \dots, j-1 \\ 0 & \text{for } i = j+1, \dots, i_0, \end{cases}$$

and since $d_j^0(x) \geq \tilde{d}_j(0, x) + \tilde{\sigma} \geq \tilde{\sigma}$,

$$\phi\left(\frac{d_j^0(x)}{\varepsilon}\right) \geq 1 - \frac{C\varepsilon}{\tilde{\sigma}}.$$

Putting it all together, we infer that, for ε small enough,

$$v^\varepsilon(0, x) \leq j - C\varepsilon |\ln \varepsilon| \leq j - \frac{C\varepsilon}{\tilde{\sigma}} \leq \sum_{i=1}^N \phi\left(\frac{d_i^0(x)}{\varepsilon}\right) = u_0(x),$$

which proves (7.7) in Case 1.

Case 2: $|\tilde{d}_i(0, x) - \tilde{\sigma}| \geq \frac{C_0}{|\ln \varepsilon|}$ for all $i = 1, \dots, i_0$.

Recalling the Definition 2.2 of d_i , if $|\tilde{d}_i(0, x)| < \rho$, then $d_i(0, x) = \tilde{d}_i(0, x)$. In this case, by the first inequality in (7.5) and the monotonicity of ϕ , we have that

$$\phi\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}\right) \leq \phi\left(\frac{d_i^0(x)}{\varepsilon}\right),$$

and by (4.23),

$$\varepsilon |\ln \varepsilon| \left| \psi_i\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}; 0, x\right) \right| \leq C\gamma\varepsilon |\ln \varepsilon|.$$

If instead, $\tilde{d}_i(0, x) \geq \rho$, then $d_i(0, x) \geq \rho$ and $d_i^0(x) \geq \tilde{d}_i(0, x) \geq \rho$, and by (4.2),

$$\phi\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}\right) \leq 1 \leq \phi\left(\frac{d_i^0(x)}{\varepsilon}\right) + \frac{C\varepsilon}{\rho}.$$

Finally, if $\tilde{d}_i(0, x) \leq -\rho$, then $d_i(0, x) \leq -\rho$ and

$$\phi\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}\right) \leq \frac{C\varepsilon}{\rho} \leq \phi\left(\frac{d_i^0(x)}{\varepsilon}\right) + \frac{C\varepsilon}{\rho}.$$

For $|\tilde{d}_i(0, x)| \geq \rho$, by (4.21) we also have

$$\varepsilon |\ln \varepsilon| \left| \psi_i\left(\frac{d_i(0, x) - \tilde{\sigma}}{\varepsilon}; 0, x\right) \right| \leq \frac{C\varepsilon}{\rho}.$$

Putting it all together, we infer that, for ε small enough,

$$v^\varepsilon(0, x) \leq u_0(x) + C\gamma\varepsilon |\ln \varepsilon| - \tilde{\sigma}\varepsilon |\ln \varepsilon| \leq u_0(x),$$

choosing $\gamma \leq \tilde{\sigma}/C$. This proves (7.7) in Case 2.

By (7.7) and the comparison principle,

$$u^\varepsilon(t, x) \geq v^\varepsilon(t, x) \quad \text{for all } (t, x) \in \left[0, \frac{r_{i_0}}{2C}\right] \times \mathbb{R}^n.$$

Since $2\sigma < \rho$, note that $\{x : d_{i_0}(t, x) \geq 2\tilde{\sigma}\} = \{x : \tilde{d}_{i_0}(t, x) \geq 2\tilde{\sigma}\}$. Consequently, by (6.7) (with $N = i_0$) and the inequality above, for $t \in [0, r_{i_0}/(2C)]$ and

$$x \in \{x : \tilde{d}_{i_0}(t, x) \geq 2\tilde{\sigma}\} = \{|x - x_0| \leq r_{i_0} - Ct - 2\tilde{\sigma}\} \supset \left\{|x - x_0| \leq \frac{r_{i_0}}{2} - 2\tilde{\sigma}\right\},$$

we have

$$\liminf_{\varepsilon \rightarrow 0} \frac{u^\varepsilon(t, x) - i_0}{\varepsilon |\ln \varepsilon|} \geq \liminf_{\varepsilon \rightarrow 0} \frac{v^\varepsilon(t, x) - i_0}{\varepsilon |\ln \varepsilon|} \geq -2\tilde{\sigma}.$$

Letting $\tilde{\sigma} \rightarrow 0$, the result follows. \square

7.2. Proof of Proposition 7.2.

Proof. Fix $1 \leq i_0 \leq N$. We will show that $(D_t^{i_0})_{t>0}$ is a generalized super-flow. The proof that $((E_t^{i_0})^c)_{t>0}$ is a generalized sub-flow is similar.

Let $(t_0, x_0) \in (0, \infty) \times \mathbb{R}^n$, $h, r > 0$, and $\varphi_{i_0} : (0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function satisfying (i)-(v) in Definition 2.1 in $[t_0, t_0 + h]$ with F given in (2.5) and μ in (1.6). Then, there exists $h' > h$ such that φ_{i_0} satisfies (i)-(v) in $[t_0, t_0 + h']$, with an eventually smaller δ in (ii). For $t \in [t_0, t_0 + h']$, let us denote

$$\Omega_t^{i_0} = \{x \in \mathbb{R}^n : \varphi_{i_0}(t, x) > 0\} \quad \text{and} \quad \Gamma_t^{i_0} = \partial\Omega_t^{i_0}.$$

By (iii) there exists $R' > r$ such that $\nabla\varphi_{i_0} \neq 0$ on $\Gamma_t^{i_0} \cap \overline{B}(x_0, R')$ which is therefore a smooth (and by (i) a non-empty) set. Let $\tilde{d}_{i_0}(t, x)$ be the signed distance function associated to $\Omega_t^{i_0}$,

and let $Q_\rho^{i_0} = \{(t, x) \in [t_0, t_0 + h'] \times \mathbb{R}^n : |\tilde{d}_{i_0}(t, x)| \leq \rho\}$ for $\rho > 0$. Then, there exist $\rho_0 > 0$ and $r' > r$ such that \tilde{d}_{i_0} is smooth in $Q_{\rho_0}^{i_0} \cap ([t_0, t_0 + h'] \times B(x_0, R'))$ and by (ii) (recall (2.4)),

$$(7.8) \quad \partial_t \tilde{d}_{i_0} \leq \mu \Delta \tilde{d}_{i_0} - \frac{\delta}{2} \quad \text{in } Q_{\rho_0}^{i_0} \cap ([t_0, t_0 + h'] \times B(x_0, r')).$$

Next, for $i = 1, \dots, i_0 - 1$, and $\tilde{\sigma} > 0$, define

$$\Omega_t^i = \{x \in \mathbb{R}^n : \tilde{d}_{i_0}(t, x) \geq -4(i_0 - i)\tilde{\sigma}\} \quad \text{and} \quad \Gamma_t^i = \partial\Omega_t^i.$$

Clearly, $\Omega_t^{i+1} \subset \Omega_t^i$, and if $\tilde{d}_i(t, x)$ is the signed distance function associated to Ω_t^i , then

$$(7.9) \quad \tilde{d}_i(t, x) \geq \tilde{d}_{i+1}(t, x) + 4\tilde{\sigma} \quad \text{for all } i = 1, \dots, i_0 - 1.$$

Moreover, by (7.8) there exist $R, \rho, \tilde{\sigma} > 0$ such that if $d_i(t, x)$ is the bounded extension of $\tilde{d}_i(t, x)$ outside of Q_ρ^i as in Definition 2.2, then d_i is smooth in $[t_0, t_0 + h'] \times B(x_0, R)$ and

$$\partial_t d_i \leq \mu \Delta d_i - c_0 \sigma \quad \text{in } Q_\rho^i \cap ([t_0, t_0 + h'] \times B(x_0, r)) \quad \text{for all } i = 1, \dots, i_0,$$

where $\sigma = W''(0)\tilde{\sigma}$. Then, after taking ρ perhaps smaller and γ in (4.4) such that $0 < \rho, \gamma < R - r$, Lemma 6.2 (with $N = i_0$) implies that the function

$$v^\varepsilon(t, x) = \sum_{i=1}^{i_0} \phi \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon} \right) + \varepsilon |\ln \varepsilon| \sum_{i=1}^{i_0} \psi_i \left(\frac{d_i(t, x) - \tilde{\sigma}}{\varepsilon}; t, x \right) - \tilde{\sigma} \varepsilon |\ln \varepsilon|$$

is a solution to (6.6) in $[t_0, t_0 + h'] \times B(x_0, r)$. We now show that for $\tilde{\sigma} < \rho/2$ and ε, γ small enough,

$$(7.10) \quad v^\varepsilon(t_0, x) \leq u^\varepsilon(t_0, x) \quad \text{for all } x \in \mathbb{R}^n$$

and

$$(7.11) \quad v^\varepsilon(t, x) \leq u^\varepsilon(t, x) \quad \text{for all } (t, x) \in [t_0, t_0 + h'] \times (\mathbb{R}^n \setminus B(x_0, r)).$$

We start with (7.10). Since φ_{i_0} satisfies (i) and (iv) in Definition 2.1, for $\tilde{\sigma}$ small enough we have that $\Omega_{t_0}^i \subset \subset D_{t_0}^{i_0}$ for all $i = 1, \dots, i_0$, and there is a compact set $K \subset D_{t_0}^{i_0}$ such that

$$(7.12) \quad d_K(x) - 2\tilde{\sigma} \geq \tilde{d}_i(t_0, x) \quad \text{for all } i = 1, \dots, i_0$$

where d_K denotes the signed distance function from K .

The proof of (7.10) is broken into three cases: we first consider the case when x is close to one of the sets $\Gamma_{t_0}^j$, then when x is in K but far from all sets $\Gamma_{t_0}^i$, and finally when x is not in K .

Case 1: *There is a $j \in \{1, \dots, i_0\}$ such that $|\tilde{d}_j(t_0, x) - \tilde{\sigma}| \leq \frac{C_0}{|\ln \varepsilon|}$ for $C_0 > 0$ to be determined.*

Like in Case 1 in the proof of Proposition 7.1, we can show, for $\varepsilon, C_0 > 0$ small enough

$$v^\varepsilon(t_0, x) \leq j - C\varepsilon |\ln \varepsilon|.$$

Since $\tilde{d}_j(t_0, x) \geq 0$ for ε small enough, and by (7.12), we know that $x \in K \subset D_{t_0}^{i_0}$. Since K is compact and by the definition of $D_{t_0}^{i_0}$, given $\delta_0 > 0$, for ε small enough and $y \in K$,

$$(7.13) \quad \frac{u^\varepsilon(t_0, y) - i_0}{\varepsilon |\ln \varepsilon|} \geq -\delta_0.$$

In particular, for $\delta_0 \leq C$,

$$u^\varepsilon(t_0, x) \geq i_0 - \delta_0 \varepsilon |\ln \varepsilon| \geq j - C\varepsilon |\ln \varepsilon| \geq v^\varepsilon(t_0, x).$$

Therefore, (7.10) holds for Case 1 for sufficiently small $\varepsilon > 0$.

Case 2: *$x \in K$ and $|\tilde{d}_i(t_0, x) - \tilde{\sigma}| \geq \frac{C_0}{|\ln \varepsilon|}$ for all $i = 1, \dots, i_0$.*

We note that

$$\phi\left(\frac{d_i(t_0, x) - \tilde{\sigma}}{\varepsilon}\right) \leq 1$$

and then estimate ψ_i as in Case 2 in the proof of Proposition 7.1, to find

$$v^\varepsilon(t_0, x) \leq i_0 + C\gamma\varepsilon|\ln\varepsilon| - \tilde{\sigma}\varepsilon|\ln\varepsilon|.$$

On the other hand, since $x \in K$, we know that (7.13) holds for given $\delta_0 > 0$ and ε small enough. Choosing $\gamma \leq \tilde{\sigma}/(2C)$ and $\delta_0 = \tilde{\sigma}/2$,

$$u^\varepsilon(t_0, x) \geq i_0 - \delta_0\varepsilon|\ln\varepsilon| \geq v^\varepsilon(t_0, x)$$

for $\varepsilon > 0$ small. We now have (7.10) in Case 2.

Case 3: $x \notin K$.

Since $d_K(t_0, x) \leq 0$, by (7.12), we have that $\tilde{d}_i(t_0, x) \leq -2\tilde{\sigma}$, for all $i = 1, \dots, i_0$. If $-\rho < \tilde{d}_i(t_0, x) < -2\tilde{\sigma}$, then $\tilde{d}_i(t_0, x) = d_i(t_0, x)$. By (4.2) and (4.21),

$$\phi\left(\frac{d_i(t_0, x) - \tilde{\sigma}}{\varepsilon}\right) + \varepsilon|\ln\varepsilon|\psi\left(\frac{d_i(t_0, x) - \tilde{\sigma}}{\varepsilon}; t_0, x\right) - \tilde{\sigma}\varepsilon|\ln\varepsilon| \leq \frac{C\varepsilon}{\tilde{\sigma}} + \frac{C\varepsilon}{\tilde{\sigma}} - \tilde{\sigma}\varepsilon|\ln\varepsilon| \leq 0$$

for $\varepsilon > 0$ sufficiently small. If $\tilde{d}_i(t_0, x) < -\rho$, then $d_i(t_0, x) < -\rho$ and we similarly find

$$\phi\left(\frac{d_i(t_0, x) - \tilde{\sigma}}{\varepsilon}\right) + \varepsilon|\ln\varepsilon|\psi\left(\frac{d_i(t_0, x) - \tilde{\sigma}}{\varepsilon}; t_0, x\right) - \tilde{\sigma}\varepsilon|\ln\varepsilon| \leq \frac{C\varepsilon}{\rho} + \frac{C\varepsilon}{\rho} - \tilde{\sigma}\varepsilon|\ln\varepsilon| \leq 0$$

for small $\varepsilon > 0$. In particular, we have shown $v^\varepsilon(t_0, x) \leq 0$. Now, since the zero function is a solution to (1.1) and $u_0^\varepsilon \geq 0$, the comparison principle implies $u^\varepsilon(t_0, x) \geq 0$. Therefore,

$$u^\varepsilon(t_0, x) \geq 0 \geq v^\varepsilon(t_0, x),$$

and (7.10) holds for Case 3.

This proves (7.10). Inequality (7.11) follows with a similar argument using that φ_{i_0} satisfies (i) and (v) in Definition 2.1. With (7.10) and (7.11), the comparison principle then implies

$$(7.14) \quad u^\varepsilon(t, x) \geq v^\varepsilon(t, x) \quad \text{for all } (t, x) \in [t_0, t_0 + h'] \times \mathbb{R}^n.$$

By (6.7) (with $N = i_0$), we have that, for all $t \in [t_0, t_0 + h']$,

$$\frac{u^\varepsilon(t, x) - i_0}{\varepsilon|\ln\varepsilon|} \geq \frac{v^\varepsilon(t, x) - i_0}{\varepsilon|\ln\varepsilon|} \geq -2\tilde{\sigma} \quad \text{in } \left\{x \in \mathbb{R}^n : d_{i_0}(t, x) - \tilde{\sigma} \geq \frac{C}{\tilde{\sigma}|\ln\varepsilon|}\right\}.$$

Letting $\varepsilon \rightarrow 0$, it follows that, for all $t \in [t_0, t_0 + h']$,

$$(7.15) \quad \{x \in \mathbb{R}^n : d_{i_0}(t, x) - \tilde{\sigma} \geq 0\} \subset \left\{x \in \mathbb{R}^n : \liminf_{\varepsilon \rightarrow 0} \frac{u^\varepsilon(t, x) - i_0}{\varepsilon|\ln\varepsilon|} \geq -2\tilde{\sigma}\right\}.$$

Now, let $x_1 \in \{x \in B(x_0, r) : \varphi_{i_0}(t_0 + h, x) > 0\}$, so that $d_{i_0}(t_0 + h, x_1) > 0$. Then, there exist $r_1 > 0$ and $0 < \tau < h' - h$ such that for $|t - (t_0 + h)| < \tau$, it holds that $B(x_1, r_1) \subset \{x \in B(x_0, r) : d_{i_0}(t, x) > 0\}$ and by (7.15), for $\tilde{\sigma} < r_1/2$,

$$[t_0 + h - \tau, t_0 + h + \tau] \times B\left(x_1, \frac{r_1}{2}\right) \subset \left\{(t, x) : \liminf_{\varepsilon \rightarrow 0} \frac{u^\varepsilon(t, x) - i_0}{\varepsilon|\ln\varepsilon|} \geq -2\tilde{\sigma}\right\}.$$

Taking $\tilde{\sigma} \rightarrow 0$, we see that $(t_0 + h, x_1)$ is an interior point of the set

$$\left\{(t, x) \in (0, \infty) \times \mathbb{R}^n : \liminf_{\varepsilon \rightarrow 0} \frac{u^\varepsilon(t, x) - i_0}{\varepsilon|\ln\varepsilon|} \geq 0\right\},$$

namely, it belongs to D^{i_0} . This proves the desired inclusion

$$\{x \in B(x_0, r) : \varphi_{i_0}(t_0 + h, x) > 0\} = \{x \in B(x_0, r) : d_{i_0}(t_0 + h, x) > 0\} \subset D_{t_0+h}^{i_0}.$$

□

8. PROOF OF LEMMA 4.3

First, note that by the regularity of ϕ and d , there is some $\tau \in (0, 1)$ and $C > 0$ such that

$$(8.1) \quad \begin{aligned} & \left| \phi \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \phi(\xi + \nabla d(t, x) \cdot z) \right| \\ & \leq \dot{\phi} \left(\xi + (1 - \tau) \nabla d(t, x) \cdot z + \tau \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) C \varepsilon |z|^2. \end{aligned}$$

8.1. Proof of (4.6) and (4.10). To prove (4.6), we will show that for all $(\xi, t, x) \in \mathbb{R} \times [t_0, t_0 + h] \times \mathbb{R}^n$,

$$(8.2) \quad |a_\varepsilon(\xi; t, x)| \leq C \varepsilon^{\frac{1}{2}}.$$

The same estimate for \dot{a}_ε , \ddot{a}_ε and $\ddot{\ddot{a}}_\varepsilon$ will follow similarly by replacing ϕ in expression (4.4) by $\dot{\phi}$, $\ddot{\phi}$ and $\ddot{\ddot{\phi}}$, respectively.

Begin by writing

$$\begin{aligned} a_\varepsilon(\xi; t, x) &= \int_{|z| < \varepsilon^{-\frac{1}{2}}} \left(\phi \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \phi(\xi + \nabla d(t, x) \cdot z) \right) \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{\varepsilon^{-\frac{1}{2}} < |z| < \gamma \varepsilon^{-1}} \left(\phi \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \phi(\xi + \nabla d(t, x) \cdot z) \right) \frac{dz}{|z|^{n+1}} \\ &=: I + II. \end{aligned}$$

For the long-range interactions,

$$|II| \leq 2 \int_{|z| > \varepsilon^{-\frac{1}{2}}} \frac{dz}{|z|^{n+1}} = C \varepsilon^{\frac{1}{2}}.$$

For the short range interactions, we use (8.1) to estimate

$$|I| \leq C \int_{|z| < \varepsilon^{-\frac{1}{2}}} \varepsilon |z|^2 \frac{dz}{|z|^{n+1}} = C \varepsilon^{\frac{1}{2}}.$$

Estimate (8.2) follows.

Consequently, using the behavior of ϕ at $\pm\infty$ in (1.4), we estimate

$$(8.3) \quad |\bar{a}_\varepsilon(t, x)| \leq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} |a_\varepsilon(\xi; t, x)| \dot{\phi}(\xi) d\xi \leq \frac{C \varepsilon^{\frac{1}{2}}}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} \dot{\phi}(\xi) d\xi = \frac{C}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|},$$

which gives (4.10).

8.2. Proof of (4.7) and (4.11). We first show the estimate for $\partial_t a_\varepsilon$ in (4.7). For this, we write

$$\begin{aligned} \partial_t a_\varepsilon(\xi; t, x) &= \int_{|z| < \frac{\gamma}{\varepsilon}} \left\{ \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) \frac{d_t(t, x + \varepsilon z) - d_t(t, x)}{\varepsilon} \right. \\ &\quad \left. - \dot{\phi}(\xi + \nabla d(t, x) \cdot z) \nabla d_t(t, x) \cdot z \right\} \frac{dz}{|z|^{n+1}} \\ &= \int_{|z| < \gamma} (\dots) + \int_{\gamma < |z| < \frac{\gamma}{\varepsilon}} (\dots) =: I + II. \end{aligned}$$

Beginning with I , we write

$$I = \int_{|z| < \gamma} \left\{ \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) \frac{d_t(t, x + \varepsilon z) - d_t(t, x) - \nabla d_t(t, x) \cdot (\varepsilon z)}{\varepsilon} \right. \\ \left. + \left(\dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \dot{\phi}(\xi + \nabla d(t, x) \cdot z) \right) \nabla d_t(t, x) \cdot z \right\} \frac{dz}{|z|^{n+1}}.$$

With the regularity of ϕ and d , we have

$$|I| \leq C \int_{|z| < \gamma} \left\{ \frac{|d_t(t, x + \varepsilon z) - d_t(t, x) - \nabla d_t(t, x) \cdot (\varepsilon z)|}{\varepsilon} \right. \\ \left. + \frac{|d_t(t, x + \varepsilon z) - d_t(t, x) - \nabla d_t(t, x) \cdot (\varepsilon z)|}{\varepsilon} |\nabla d_t(t, x) \cdot z| \right\} \frac{dz}{|z|^{n+1}} \\ \leq C \int_{|z| < \gamma} (\varepsilon |z|^2 + \varepsilon |z|^3) \frac{dz}{|z|^{n+1}} \leq C \gamma \varepsilon \leq C |\ln \varepsilon|.$$

Next, we estimate II . By the regularity of ϕ and d , we have

$$|II| \leq C \int_{\gamma < |z| < \frac{\gamma}{\varepsilon}} |z| \frac{dz}{|z|^{n+1}} = C |\ln \varepsilon|.$$

Combining the estimates for I and II , we have (4.7) for $\partial_t a_\varepsilon$. The estimate for $\nabla_x a_\varepsilon$ in (4.7) follows along the same lines.

We now check (4.7) for $D_x^2 a_\varepsilon$. For ease, we drop the notation in t and write

$$D_x^2 a_\varepsilon(\xi; x) \\ = \int_{|z| < \frac{\gamma}{\varepsilon}} \left\{ \ddot{\phi} \left(\xi + \frac{d(x + \varepsilon z) - d(x)}{\varepsilon} \right) \frac{(\nabla d(x + \varepsilon z) - \nabla d(x)) \otimes (\nabla d(x + \varepsilon z) - \nabla d(x))}{\varepsilon^2} \right. \\ - \ddot{\phi}(\xi + \nabla d(x) \cdot z) D^2 d(x) z \otimes D^2 d(x) z \\ + \dot{\phi} \left(\xi + \frac{d(x + \varepsilon z) - d(x)}{\varepsilon} \right) \frac{D^2 d(x + \varepsilon z) - D^2 d(x)}{\varepsilon} \\ \left. - \dot{\phi}(\xi + \nabla d(x) \cdot z) D^3 d(x) z. \right\} \frac{dz}{|z|^{n+1}}.$$

Using the regularity of ϕ and d , we estimate to find

$$\begin{aligned}
(8.4) \quad & |D_x^2 a_\varepsilon(\xi; x)| \\
& \leq \int_{|z| < \frac{\gamma}{\varepsilon}} \left\{ \left| \ddot{\phi} \left(\xi + \frac{d(x + \varepsilon z) - d(x)}{\varepsilon} \right) \frac{(\nabla d(x + \varepsilon z) - \nabla d(x)) \otimes (\nabla d(x + \varepsilon z) - \nabla d(x))}{\varepsilon^2} \right| \right. \\
& \quad + \left| \ddot{\phi}(\xi + \nabla d(x) \cdot z) D^2 d(x) z \otimes D^2 d(x) z \right| \\
& \quad + \left| \left(\dot{\phi} \left(\xi + \frac{d(x + \varepsilon z) - d(x)}{\varepsilon} \right) - \dot{\phi}(\xi) \right) \frac{D^2 d(x + \varepsilon z) - D^2 d(x)}{\varepsilon} \right| \\
& \quad + \left| \left(\dot{\phi}(\xi + \nabla d(x) \cdot z) - \dot{\phi}(\xi) \right) D^3 d(x) z \right| \\
& \quad \left. + \left| \dot{\phi}(\xi) \left(\frac{D^2 d(x + \varepsilon z) - D^2 d(x)}{\varepsilon} - D^3 d(x) z \right) \right| \right\} \frac{dz}{|z|^{n+1}} \\
& \leq C \int_{|z| < \frac{\gamma}{\varepsilon}} \left\{ \frac{|\nabla d(x + \varepsilon z) - \nabla d(x)|^2}{\varepsilon^2} + |D^2 d(x) z \otimes D^2 d(x) z| \right. \\
& \quad + \frac{|d(x + \varepsilon z) - d(x)|}{\varepsilon} \frac{|D^2 d(x + \varepsilon z) - D^2 d(x)|}{\varepsilon} + |\nabla d(x) \cdot z| |D^3 d(x) z| \\
& \quad \left. + \left| \frac{D^2 d(x + \varepsilon z) - D^2 d(x)}{\varepsilon} - D^3 d(x) z \right| \right\} \frac{dz}{|z|^{n+1}} \\
& \leq C \int_{|z| < \frac{\gamma}{\varepsilon}} |z|^2 \frac{dz}{|z|^{n+1}} \leq \frac{C}{\varepsilon}.
\end{aligned}$$

With (4.7), we estimate as in (8.3) to obtain

$$|\partial_t \bar{a}_\varepsilon(t, x)| \leq \frac{1}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} |\partial_t a_\varepsilon(\xi; t, x)| \dot{\phi}(\xi) d\xi \leq \frac{C |\ln \varepsilon|}{\varepsilon |\ln \varepsilon|} \int_{\mathbb{R}} \dot{\phi}(\xi) d\xi = \frac{C}{\varepsilon}$$

and similarly for $\nabla_x a_\varepsilon$ and $D_x^2 a_\varepsilon$. This gives (4.11).

8.3. Proof of (4.8). By (4.6), we can assume that $|\xi| > 1$.

Choose $\kappa > 0$ such that

$$(8.5) \quad \|\nabla d\|_\infty \kappa < \frac{1}{4}.$$

Then, for $|z| < \kappa|\xi|$, we have that $|\nabla d(t, x) \cdot z| \leq |\xi|/4$ and $\frac{|d(t, x + \varepsilon z) - d(t, x)|}{\varepsilon} \leq \|\nabla d\|_\infty |z| \leq |\xi|/4$. In particular, by (8.1) and the estimate (4.3) for $\dot{\phi}$,

$$(8.6) \quad \left| \phi \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \phi(\xi + \nabla d(t, x) \cdot z) \right| \leq \frac{C \varepsilon |z|^2}{|\xi|^2}.$$

Now, we write

$$\begin{aligned}
a_\varepsilon(\xi; t, x) &= \int_{\{|z| < \frac{\gamma}{\varepsilon}\}} \left(\phi \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \phi(\xi + \nabla d(t, x) \cdot z) \right) \frac{dz}{|z|^{n+1}} \\
&= \int_{\{|z| < \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} (\dots) + \int_{\{|z| > \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} (\dots) \\
&=: I + II.
\end{aligned}$$

By (8.6),

$$|I| \leq \frac{C\varepsilon}{|\xi|^2} \int_{\{|z| < \kappa|\xi|\}} \frac{dz}{|z|^{n-1}} = \frac{C\varepsilon}{|\xi|}.$$

Next, we have

$$|II| \leq 2 \int_{\{|z| > \kappa|\xi|\}} \frac{dz}{|z|^{n+1}} \leq \frac{C}{|\xi|}.$$

Estimate (4.8) for a_ε then follows.

Estimate (4.8) for \dot{a}_ε is proven in the same way, just replacing ϕ by $\dot{\phi}$.

8.4. Proof of (4.9). By (4.7), we can assume that $|\xi| > 1$.

Choose $\kappa > 0$ satisfying (8.5). We split

$$\begin{aligned} \partial_t a_\varepsilon(\xi; t, x) &= \int_{\{|z| < \frac{\gamma}{\varepsilon}\}} \left\{ \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) \frac{d_t(t, x + \varepsilon z) - d_t(t, x)}{\varepsilon} \right. \\ &\quad \left. - \dot{\phi}(\xi + \nabla d(t, x) \cdot z) \nabla d_t(t, x) \cdot z \right\} \frac{dz}{|z|^{n+1}} \\ &= \int_{\{|z| < \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} (\dots) + \int_{\{|z| > \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} (\dots) =: I + II. \end{aligned}$$

Beginning with I , we write

$$\begin{aligned} I &= \int_{\{|z| < \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} \left\{ \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) \frac{d_t(t, x + \varepsilon z) - d_t(t, x) - \nabla d_t(t, x) \cdot (\varepsilon z)}{\varepsilon} \right. \\ &\quad \left. + \left(\dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \dot{\phi}(\xi + \nabla d(t, x) \cdot z) \right) \nabla d_t(t, x) \cdot z \right\} \frac{dz}{|z|^{n+1}}. \end{aligned}$$

As in the proof of (4.8), by the estimates (4.3) for $\dot{\phi}$ and $\ddot{\phi}$, if $|z| < \kappa|\xi|$ and $|z| < \frac{\gamma}{\varepsilon}$, then

$$0 < \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) \leq \frac{C}{|\xi|^2}$$

and

$$\left| \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \dot{\phi}(\xi + \nabla d(t, x) \cdot z) \right| \leq \frac{C\varepsilon}{|\xi|^2} |z|^2 \leq \frac{C}{|\xi|^2} |z|.$$

Therefore,

$$|I| \leq \frac{C}{|\xi|^2} \int_{\{|z| < \kappa|\xi|\}} (\varepsilon|z|^2 + |z|^2) \frac{dz}{|z|^{n+1}} \leq \frac{C}{|\xi|}.$$

Regarding II , we have

$$|II| \leq C \int_{\{|z| > \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} \left(\frac{1}{\varepsilon} + |z| \right) \frac{dz}{|z|^{n+1}} \leq \frac{C}{\varepsilon} \int_{\{|z| > \kappa|\xi|\}} \frac{dz}{|z|^{n+1}} \leq \frac{C}{\varepsilon|\xi|}.$$

From the estimates of I and II , (4.9) for $\partial_t a_\varepsilon$ follows.

The estimate for $\nabla_x a_\varepsilon$ can be proven in the same way.

We next check (4.9) for $D_x^2 a_\varepsilon$. As before, if κ satisfies (8.5) and if $|z| < \kappa|\xi|$, then, by (4.3) for $\dot{\phi}$ and $\ddot{\phi}$,

$$\begin{aligned} &\left| \ddot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) \right|, \left| \ddot{\phi}(\xi + \nabla d(t, x) \cdot z) \right|, \dot{\phi}(\xi) \leq \frac{C}{|\xi|^2}, \\ &\left| \dot{\phi}(\xi + \nabla d(t, x) \cdot z) - \dot{\phi}(\xi) \right|, \left| \dot{\phi} \left(\xi + \frac{d(t, x + \varepsilon z) - d(t, x)}{\varepsilon} \right) - \dot{\phi}(\xi) \right| \leq \frac{C|z|}{|\xi|^2}. \end{aligned}$$

Therefore, recalling (8.4),

$$\begin{aligned} |D_x^2 a_\varepsilon(\xi; t, x)| &\leq \frac{C}{|\xi|^2} \int_{\{|z| < \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} |z|^2 \frac{dz}{|z|^{n+1}} + C \int_{\{|z| > \kappa|\xi|\} \cap \{|z| < \frac{\gamma}{\varepsilon}\}} |z|^2 \frac{dz}{|z|^{n+1}} \\ &\leq \frac{C}{|\xi|^2} \int_{\{|z| < \kappa|\xi|\}} \frac{dz}{|z|^{n-1}} + \frac{C}{\varepsilon^2} \int_{\{|z| > \kappa|\xi|\}} \frac{dz}{|z|^{n+1}} \leq \frac{C}{\varepsilon^2 |\xi|}. \end{aligned}$$

The proof of the second line in (4.9) is proven the same way, just replacing ϕ with $\dot{\phi}$.

The proof of Lemma 4.3 is then complete. \square

9. PROOF OF LEMMA 4.4

9.1. **Proof of (4.12).** For ease, we drop the notation in t . Note that

$$\frac{d(x + \varepsilon z) - d(x)}{\varepsilon} = \nabla d(x) \cdot z + \varepsilon O(|z|^2).$$

If $|\nabla d(x)| \leq C_0 \varepsilon$, with $C_0 > 0$ to be determined, then

$$|a_\varepsilon(\xi; x)| \leq C \varepsilon \int_{\{|z| < \frac{\gamma}{\varepsilon}\}} \frac{dz}{|z|^{n-1}} \leq C \leq \frac{C \varepsilon |\ln \varepsilon|}{|\nabla d(x)|},$$

as desired.

Next, let us assume

$$(9.1) \quad c_1 := |\nabla d(x)| > C_0 \varepsilon.$$

Let T be an orthogonal matrix such that if $z = Ty$, then

$$(9.2) \quad \nabla d(x) \cdot z = \nabla d(x) \cdot Ty = c_1 y_n.$$

With the change of variables $\varepsilon z = Ty$ and using the monotonicity of ϕ , we obtain

$$\begin{aligned} a_\varepsilon(\xi; x) &= \varepsilon \int_{\{|y| < \gamma\}} \left\{ \phi \left(\xi + \frac{c_1 y_n}{\varepsilon} + \frac{O(|y|^2)}{\varepsilon} \right) - \phi \left(\xi + \frac{c_1 y_n}{\varepsilon} \right) \right\} \frac{dy}{|y|^{n+1}} \\ &\leq \varepsilon \int_{\mathbb{R}^n} \left\{ \phi \left(\xi + \frac{c_1 y_n}{\varepsilon} + \frac{C|y|^2}{\varepsilon} \right) - \phi \left(\xi + \frac{c_1 y_n}{\varepsilon} \right) \right\} \frac{dy}{|y|^{n+1}}. \end{aligned}$$

Fix $r > \varepsilon$, to be chosen. For $y = (y', y_n)$, we write

$$\begin{aligned} \frac{a_\varepsilon(\xi; x)}{\varepsilon} &\leq \int_{\{|y'| > r\} \cup \{|y_n| > r\}} (\dots) + \int_{\{|y'|, |y_n| < \varepsilon\}} (\dots) \\ &\quad + \underbrace{\int_{\{|y_n| < r\} \cap \{\varepsilon < |y'| < r\}} (\dots)}_{:=I} + \underbrace{\int_{\{|y'| < \varepsilon\} \cap \{\varepsilon < |y_n| < r\}} (\dots)}_{:=II}. \end{aligned}$$

Since

$$\int_{\{|y'| > r\} \cup \{|y_n| > r\}} (\dots) \frac{dy}{|y|^{n+1}} \leq 2 \int_{\{|y'| > r\}} \frac{dy}{|y|^{n+1}} \leq \frac{C}{r}$$

and

$$\int_{\{|y'|, |y_n| < \varepsilon\}} (\dots) \frac{dy}{|y|^{n+1}} \leq \frac{C}{\varepsilon} \int_{\{|y| < \sqrt{2}\varepsilon\}} |y|^2 \frac{dy}{|y|^{n+1}} \leq C,$$

we have

$$(9.3) \quad \frac{a_\varepsilon(\xi; x)}{\varepsilon} \leq O(1) + O\left(\frac{1}{r}\right) + I + II.$$

Next, we will show that

$$(9.4) \quad I \leq \frac{C}{c_1} |\ln \varepsilon|.$$

For this, we will make the change of variables

$$(9.5) \quad t = \frac{y_n}{|y'|} \quad \text{and} \quad y' = \theta p, \quad \theta \in S^{n-2}.$$

Indeed, changing variables first in t for a fixed y' , we have

$$\begin{aligned} I &= \int_{\{\varepsilon < |y'| < r\}} dy' \int_{\{|y_n| < r\}} \frac{dy_n}{(|y'|^2 + |y_n|^2)^{\frac{n+1}{2}}} \\ &\quad \left\{ \phi\left(\xi + \frac{c_1 y_n}{\varepsilon} + \frac{C}{\varepsilon}(|y'|^2 + |y_n|^2)\right) - \phi\left(\xi + \frac{c_1 y_n}{\varepsilon}\right) \right\} \\ &= \int_{\{\varepsilon < |y'| < r\}} \frac{dy'}{|y'|^n} \int_{\{|t| < \frac{r}{|y'|}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \\ &\quad \left\{ \phi\left(\xi + \frac{|y'|c_1}{\varepsilon}t + C\frac{|y'|^2}{\varepsilon}(1+t^2)\right) - \phi\left(\xi + \frac{|y'|c_1}{\varepsilon}t\right) \right\}. \end{aligned}$$

Then, we change to polar coordinates in y' to write

$$\begin{aligned} I &= \int_{S^{n-2}} d\theta \int_\varepsilon^r \frac{dp}{p^2} \int_{\{|t| < \frac{r}{p}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \\ &\quad \left\{ \phi\left(\xi + \frac{pc_1}{\varepsilon}t + C\frac{p^2}{\varepsilon}(1+t^2)\right) - \phi\left(\xi + \frac{pc_1}{\varepsilon}t\right) \right\}. \end{aligned}$$

We split

$$I = \int_{S^{n-2}} d\theta \int_\varepsilon^r \frac{dp}{p^2} \int_{\{|t| < 1\}} dt(\dots) + \int_{S^{n-2}} d\theta \int_\varepsilon^r \frac{dp}{p^2} \int_{\{1 < |t| < \frac{r}{p}\}} dt(\dots) =: I_1 + I_2.$$

If $|t| < 1$, then $1 + t^2 \leq C$. With this and the monotonicity of ϕ ,

$$\begin{aligned} I_1 &\leq C \int_\varepsilon^r \frac{dp}{p^2} \int_{\{|t| < 1\}} \left\{ \phi\left(\xi + \frac{pc_1}{\varepsilon}t + C\frac{p^2}{\varepsilon}\right) - \phi\left(\xi + \frac{pc_1}{\varepsilon}t\right) \right\} dt \\ &= C \int_\varepsilon^r \frac{dp}{p^2} \int_{\{|t| < 1\}} dt \int_0^1 \dot{\phi}\left(\xi + \frac{pc_1}{\varepsilon}t + C\frac{p^2\tau}{\varepsilon}\right) \frac{p^2}{\varepsilon} d\tau \\ &= \frac{C}{\varepsilon} \int_\varepsilon^r dp \int_0^1 d\tau \int_{\{|t| < 1\}} \partial_t \left[\phi\left(\xi + \frac{pc_1}{\varepsilon}t + C\frac{p^2\tau}{\varepsilon}\right) \right] \frac{\varepsilon}{pc_1} dt \\ &= \frac{C}{c_1} \int_\varepsilon^r \frac{dp}{p} \int_0^1 d\tau \left\{ \phi\left(\xi + \frac{pc_1}{\varepsilon} + C\frac{p^2\tau}{\varepsilon}\right) - \phi\left(\xi - \frac{pc_1}{\varepsilon} + C\frac{p^2\tau}{\varepsilon}\right) \right\} \\ &\leq \frac{C |\ln \varepsilon|}{c_1}. \end{aligned}$$

This shows that

$$(9.6) \quad I_1 \leq \frac{C |\ln \varepsilon|}{c_1}.$$

Next, estimating I_2 , we use that the integrand function is positive to find

$$I_2 \leq C \int_{\varepsilon}^r \frac{dp}{p^2} \int_{\{|t|>1\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \left\{ \phi \left(\xi + \frac{pc_1}{\varepsilon}t + \frac{Cp^2}{\varepsilon}(1+t^2) \right) - \phi \left(\xi + \frac{pc_1}{\varepsilon}t \right) \right\}.$$

Let r satisfy

$$(9.7) \quad c_1 \geq r,$$

so that $c_1/p > 1$ for $0 < p < r$. Then we can split

$$\begin{aligned} I_2 &\leq C \int_{\varepsilon}^r \frac{dp}{p^2} \int_{\{|t|>\sqrt{\frac{c_1}{p}}\}} dt(\dots) + C \int_{\varepsilon}^r \frac{dp}{p^2} \int_{\{1<|t|<\sqrt{\frac{c_1}{p}}\}} dt(\dots) \\ &=: J_1 + J_2. \end{aligned}$$

We will show that

$$(9.8) \quad J_1, J_2 \leq \frac{C|\ln \varepsilon|}{c_1}.$$

For J_1 , we estimate

$$J_1 \leq C \int_{\varepsilon}^r \frac{dp}{p^2} \int_{\{|t|>\sqrt{\frac{c_1}{p}}\}} \frac{dt}{|t|^{n+1}} = \frac{C}{c_1^{\frac{n}{2}}} \int_{\varepsilon}^r p^{\frac{n}{2}-2} dp \leq \frac{C}{c_1} \int_{\varepsilon}^r \frac{dp}{p} \leq \frac{C}{c_1} |\ln \varepsilon|.$$

Estimating J_2 , we find

$$\begin{aligned} J_2 &= C \int_{\varepsilon}^r \frac{dp}{p^2} \int_{\{1<|t|<\sqrt{\frac{c_1}{p}}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \int_0^1 \dot{\phi} \left(\xi + \frac{pc_1}{\varepsilon}t + \frac{Cp^2\tau}{\varepsilon}(1+t^2) \right) \frac{p^2}{\varepsilon}(1+t^2) d\tau \\ &= C \int_{\varepsilon}^r \frac{dp}{p} \int_0^1 d\tau \int_{\{1<|t|<\sqrt{\frac{c_1}{p}}\}} \frac{dt}{(1+t^2)^{\frac{n-1}{2}}} \partial_t \left[\phi \left(\xi + \frac{pc_1}{\varepsilon}t + \frac{Cp^2\tau}{\varepsilon}(1+t^2) \right) \right] \frac{1}{c_1 + 2Cp\tau t}. \end{aligned}$$

Now, if

$$\varepsilon < p < r, \quad 0 < \tau < 1 \quad \text{and} \quad 1 \leq |t| \leq \sqrt{\frac{c_1}{p}},$$

then

$$|2Cp\tau t| \leq 2Cp\sqrt{\frac{c_1}{p}} = 2C\sqrt{c_1 p} \leq 2C\sqrt{c_1 r},$$

where we can assume $C > 1$ in the last inequality above. Choose

$$(9.9) \quad r = \frac{c_1}{16C^2},$$

then (9.7) is satisfied, and choosing $C_0 = 16C^2$ in (9.1) we also have that $r > \varepsilon$. Moreover,

$$c_1 + 2Cp\tau t \geq c_1 - 2C\sqrt{c_1 r} \geq \frac{c_1}{2}.$$

Then, integrating by parts, we get

$$\begin{aligned} J_2 &\leq \frac{C}{c_1} \int_{\varepsilon}^r \frac{dp}{p} \int_0^1 d\tau \int_{\{1<|t|<\sqrt{\frac{c_1}{p}}\}} \partial_t \left[\phi \left(\xi + \frac{pc_1}{\varepsilon}t + \frac{Cp^2\tau}{\varepsilon}(1+t^2) \right) \right] \frac{dt}{|t|^{n-1}} \\ &= \frac{C}{c_1} \int_{\varepsilon}^r \frac{dp}{p} \int_0^1 d\tau \left[(n-1) \int_{\{1<|t|<\sqrt{\frac{c_1}{p}}\}} \phi \left(\xi + \frac{pc_1}{\varepsilon}t + \frac{Cp^2\tau}{\varepsilon}(1+t^2) \right) \frac{t}{|t|} \frac{dt}{|t|^n} \right. \\ &\quad \left. + \phi \left(\xi + \frac{pc_1}{\varepsilon}t + \frac{Cp^2\tau}{\varepsilon}(1+t^2) \right) \frac{1}{|t|^{n-1}} \Big|_{-\sqrt{\frac{c_1}{p}}}^{-1} \right] \end{aligned}$$

$$\begin{aligned}
& + \phi \left(\xi + \frac{pc_1}{\varepsilon} t + \frac{Cp^2\tau}{\varepsilon} (1+t^2) \right) \frac{1}{|t|^{n-1}} \Big|_1^{\sqrt{\frac{c_1}{p}}} \\
& \leq \frac{C}{c_1} \int_{\varepsilon}^r \frac{dp}{p} \int_0^1 d\tau \left[\int_{|t|>1} \frac{dt}{|t|^n} + 1 \right] \\
& = \frac{C|\ln \varepsilon|}{c_1}.
\end{aligned}$$

Hence, we obtain (9.8). The desired estimate (9.4) follows from (9.6) and (9.8).

It remains to estimate II . First, we make the same change of variables as in I to write

$$\begin{aligned}
II &= \int_{S^{n-2}} d\theta \int_0^{\varepsilon} \frac{dp}{p^2} \int_{\{\frac{\varepsilon}{p} < |t| < \frac{r}{p}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \\
& \quad \left\{ \phi \left(\xi + \frac{pc_1}{\varepsilon} t + C \frac{p^2}{\varepsilon} (1+t^2) \right) - \phi \left(\xi + \frac{pc_1}{\varepsilon} t \right) \right\}.
\end{aligned}$$

With the regularity of ϕ and d , we have

$$\begin{aligned}
II &\leq C \int_0^{\varepsilon} \frac{dp}{p^2} \int_{\{\frac{\varepsilon}{p} < |t| < \frac{r}{p}\}} \frac{p^2}{\varepsilon} \frac{dt}{(1+t^2)^{\frac{n-1}{2}}} \\
&\leq \frac{C}{\varepsilon} \int_0^{\varepsilon} dp \int_{\{1 < |t| < \frac{r}{p}\}} \frac{dt}{|t|} \\
&\leq \frac{C}{\varepsilon} \int_0^{\varepsilon} (\ln r - \ln p) dp \leq C|\ln \varepsilon|.
\end{aligned}$$

Combining the previous estimate for II with (9.3), (9.4) and (9.9) we get

$$\frac{a_{\varepsilon}(\xi; x)}{\varepsilon} \leq \frac{C}{r} + \frac{C|\ln \varepsilon|}{c_1} \leq \frac{C|\ln \varepsilon|}{c_1}.$$

Similarly, one can show

$$\frac{a_{\varepsilon}(\xi; x)}{\varepsilon} \geq -\frac{C|\ln \varepsilon|}{c_1}.$$

This proves (4.12) for a_{ε} . The estimates for \dot{a}_{ε} and \ddot{a}_{ε} are proved in the same way, just replacing ϕ by $\dot{\phi}$ and $\ddot{\phi}$, respectively. Lastly, (4.12) for \bar{a}_{ε} follows from the estimate for a_{ε} .

9.2. Proof of (4.13). We will only prove the estimate for $\nabla_x a_{\varepsilon}$, the estimate for $\partial_t a_{\varepsilon}$ and $\nabla_x \dot{a}_{\varepsilon}$ will follow similarly. Moreover, (4.13) for the derivatives of \bar{a}_{ε} follows from the estimates for the derivatives of a_{ε} . For ease, we continue to drop the notation in t .

We perform the same Taylor expansion as above but we make the error term explicit,

$$\frac{d(x + \varepsilon z) - d(x)}{\varepsilon} = \nabla d(x) \cdot z + \varepsilon \int_0^1 D^2 d(x + \tau \varepsilon z) (1 - \tau) d\tau z \cdot z.$$

Recall that by Definition 2.2, d is the smooth signed distance function from the front Γ_t in the set $\{|d| < \rho\}$. Therefore, $|\nabla d(x)| = 1$ for all x in that set. As in (9.2) (with $c_1 = 1$), let T be an orthogonal matrix such that if $z = Ty$, then

$$(9.10) \quad \nabla d(x) \cdot z = \nabla d(x) \cdot Ty = y_n.$$

With the change of variables $z = Ty$, we obtain, for all $x \in \{|d| < \rho\}$,

$$a_{\varepsilon}(\xi; x) = \int_{\{|y| < \frac{\rho}{\varepsilon}\}} \frac{dy}{|y|^{n+1}} \left\{ \phi \left(\xi + y_n + \varepsilon \int_0^1 D^2 d(x + \tau \varepsilon Ty) (1 - \tau) d\tau Ty \cdot Ty \right) - \phi(\xi + y_n) \right\}.$$

Therefore, for $i = 1, \dots, n$,

$$\begin{aligned} \partial_{x_i} a_\varepsilon(\xi; x) &= \int_{\{|y| < \frac{\gamma}{\varepsilon}\}} \frac{dy}{|y|^{n+1}} \dot{\phi} \left(\xi + y_n + \varepsilon \int_0^1 D^2 d(x + \tau \varepsilon T y) (1 - \tau) d\tau T y \cdot T y \right) \\ &\quad \cdot \varepsilon \int_0^1 D^2 d_{x_i}(x + \tau \varepsilon T y) (1 - \tau) d\tau T y \cdot T y, \end{aligned}$$

from which we deduce that

$$|\partial_{x_i} a_\varepsilon(\xi; x)| \leq \varepsilon \int_{\{|y| < \frac{\gamma}{\varepsilon}\}} \frac{dy}{|y|^{n-1}} \leq C,$$

as desired.

9.3. Proof of (4.14). For ease, we continue to drop the notation in t .

If $x \in Q_\rho$, then, recalling Definition 2.2, $|\nabla d(x)| = 1$. Estimate (4.14) is then a consequence of estimate (4.12) for a_ε when $|\xi| \leq \gamma^{-2}$ for ε small enough. Therefore, we assume $|\xi| > \gamma^{-2}$.

We will show that

$$(9.11) \quad a_\varepsilon(\xi; x) \leq \frac{C\gamma}{|\xi|}.$$

Performing first the change of variable $\varepsilon z = Tz$ with T as in (9.10) and then the change of variable in (9.5), we get

$$\begin{aligned} a_\varepsilon(\xi; x) &= \varepsilon \int_{S^{n-2}} d\theta \int_0^\gamma \frac{dp}{p^2} \int_{\{|t| < \sqrt{\frac{\gamma^2}{p^2} - 1}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \\ &\quad \left\{ \phi \left(\xi + \frac{p}{\varepsilon} t + \frac{p^2}{\varepsilon} O(1+t^2) \right) - \phi \left(\xi + \frac{p}{\varepsilon} t \right) \right\}. \end{aligned}$$

By the monotonicity of ϕ , for some $C_0 > 0$,

$$a_\varepsilon(\xi; x) \leq C\varepsilon \int_0^\gamma \frac{dp}{p^2} \int_{\{|t| < \frac{\gamma}{p}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \left\{ \phi \left(\xi + \frac{p}{\varepsilon} t + \frac{C_0 p^2}{\varepsilon} (1+t^2) \right) - \phi \left(\xi + \frac{p}{\varepsilon} t \right) \right\}.$$

If $|\xi| \geq 2\gamma\varepsilon^{-1}(1+2C_0)$, then for $0 < p < \gamma < 1$ and $|t| < \frac{\gamma}{p}$,

$$\left| \xi + \frac{p}{\varepsilon} t + \frac{C_0 p^2}{\varepsilon} (1+t^2) \right| \geq |\xi| - \frac{|pt|}{\varepsilon} - \frac{C_0 p^2}{\varepsilon} - \frac{C_0 |pt|}{\varepsilon} \geq |\xi| - \frac{\gamma}{\varepsilon} (1+2C_0) \geq \frac{|\xi|}{2}$$

and by (4.3), for some $\tau \in (0, 1)$,

$$\begin{aligned} \phi \left(\xi + \frac{p}{\varepsilon} t + \frac{C_0 p^2}{\varepsilon} (1+t^2) \right) - \phi \left(\xi + \frac{p}{\varepsilon} t \right) &= \dot{\phi} \left(\xi + \frac{p}{\varepsilon} t + \tau \frac{C_0 p^2}{\varepsilon} (1+t^2) \right) \frac{C_0 p^2}{\varepsilon} (1+t^2) \\ &\leq \frac{C}{\left| \xi + \frac{p}{\varepsilon} t + \frac{C_0 p^2}{\varepsilon} (1+t^2) \right|^2} \frac{C_0 p^2}{\varepsilon} (1+t^2) \\ &\leq \frac{C p^2}{\varepsilon |\xi|^2} (1+t^2). \end{aligned}$$

Therefore,

$$a_\varepsilon(\xi; x) \leq \frac{C}{|\xi|^2} \int_0^\gamma dp \int_{\{|t| < \frac{\gamma}{p}\}} \frac{dt}{(1+t^2)^{\frac{n-1}{2}}} \leq \frac{C}{|\xi|^2} \left(\gamma + \int_0^\gamma dp \int_{\{1 < |t| < \frac{\gamma}{p}\}} \frac{dt}{|t|} \right)$$

$$\leq \frac{C}{|\xi|^2} \left(\gamma + \int_0^\gamma (\ln \gamma - \ln p) dp \right) \leq \frac{C\gamma}{|\xi|^2} \leq \frac{C\varepsilon}{|\xi|},$$

where we used that $|\xi| \geq 2\gamma\varepsilon^{-1}(1+2C_0)$ in the last inequality. This implies (9.11).

Next, let us assume $\gamma^{-2} < |\xi| < 2\gamma\varepsilon^{-1}(1+2C_0)$. We split,

$$\begin{aligned} (9.12) \quad \frac{a_\varepsilon(\xi; x)}{\varepsilon} &\leq C \int_0^\gamma \frac{dp}{p^2} \int_{\{|t| < \frac{\gamma}{p}\}} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \left\{ \phi \left(\xi + \frac{p}{\varepsilon}t + \frac{C_0 p^2}{\varepsilon}(1+t^2) \right) - \phi \left(\xi + \frac{p}{\varepsilon}t \right) \right\} \\ &= C \int_0^\gamma \frac{dp}{p^2} \int_{\{|t| < 1\}} (\dots) + C \int_0^\gamma \frac{dp}{p^2} \int_{\{1 < |t| < \frac{\gamma}{p}\}} (\dots) \\ &=: I + II. \end{aligned}$$

As in the proof of (4.12), for $C_1 = 2C_0$,

$$(9.13) \quad I \leq C \int_0^\gamma \frac{dp}{p} \int_0^1 \left(\phi \left(\xi + \frac{p}{\varepsilon} + \frac{C_1 p^2 \tau}{\varepsilon} \right) - \phi \left(\xi - \frac{p}{\varepsilon} + \frac{C_1 p^2 \tau}{\varepsilon} \right) \right) d\tau.$$

We next observe that

$$(9.14) \quad \frac{p}{\varepsilon} + \frac{C_1 p^2 \tau}{\varepsilon} < \frac{|\xi|}{2} \quad \text{if} \quad 0 < p < p_1 := \frac{\varepsilon|\xi|}{1 + \sqrt{1 + 2C_1 \tau \varepsilon |\xi|}},$$

and since $|\xi| < 2\gamma\varepsilon^{-1}(1+2C_0)$,

$$(9.15) \quad c\varepsilon|\xi| \leq p_1 \leq C\varepsilon|\xi|.$$

Since the integrand function in (9.13) is positive, we can write

$$I \leq C \int_0^1 d\tau \int_0^{p_1} dp (\dots) + C \int_0^1 d\tau \int_{p_1}^{\max\{\gamma, p_1\}} dp (\dots) =: J_1 + J_2.$$

By (9.14) and (4.3), if $0 < p < p_1$,

$$\phi \left(\xi + \frac{p}{\varepsilon} + \frac{C_1 p^2 \tau}{\varepsilon} \right) - \phi \left(\xi - \frac{p}{\varepsilon} + \frac{C_1 p^2 \tau}{\varepsilon} \right) \leq \frac{C}{\left(|\xi| - \frac{p}{\varepsilon} - \frac{C_1 p^2 \tau}{\varepsilon} \right)^2} \frac{p}{\varepsilon} \leq \frac{Cp}{\varepsilon|\xi|^2}.$$

Therefore, using (9.15) and that $|\xi| < 2\gamma\varepsilon^{-1}(1+2C_0)$,

$$J_1 \leq \frac{C}{\varepsilon|\xi|^2} \int_0^{p_1} p dp \leq \frac{Cp_1^2}{\varepsilon|\xi|^2} \leq C\varepsilon \leq \frac{C\gamma}{|\xi|}.$$

Next, again by (9.15),

$$J_2 \leq C \int_{p_1}^{\max\{\gamma, p_1\}} \frac{dp}{p} \leq C \frac{\max\{\gamma, p_1\} - p_1}{p_1} \leq \frac{C\gamma}{\varepsilon|\xi|}.$$

Putting it all together, we obtain

$$(9.16) \quad \varepsilon I \leq \frac{C\gamma}{|\xi|}.$$

Let us now estimate II in (9.12). By the monotonicity of ϕ ,

$$II \leq \int_0^\gamma \frac{dp}{p^2} \int_{\{1 < |t| < \frac{\gamma}{p}\}} \frac{dt}{|t|^{n+1}} \left\{ \phi \left(\xi + \frac{pt}{\varepsilon} + \frac{C_1 p^2 t^2}{\varepsilon} \right) - \phi \left(\xi + \frac{pt}{\varepsilon} \right) \right\}$$

$$\begin{aligned}
&= \int_0^\gamma \frac{dp}{p^2} \int_{\{1 < t < \frac{\gamma}{p}\}} (\dots) + \int_0^\gamma \frac{dp}{p^2} \int_{\{-\frac{\gamma}{p} < t < -1\}} (\dots) \\
&=: II_1 + II_2.
\end{aligned}$$

We are only going to estimate II_1 since the estimate for II_2 follows with a similar argument. Thus, we assume below that $t > 1$. Observe that

$$(9.17) \quad \frac{pt}{\varepsilon} + \frac{C_1 p^2 t^2}{\varepsilon} < |\xi| - \frac{\gamma}{2} |\xi| \quad \text{if} \quad 0 < tp < p_1 := \frac{2\varepsilon (|\xi| - \frac{\gamma}{2} |\xi|)}{1 + \sqrt{1 + 4C_1 \varepsilon (|\xi| - \frac{\gamma}{2} |\xi|)}}$$

and

$$(9.18) \quad \frac{pt}{\varepsilon} > |\xi| + \gamma |\xi| \quad \text{if} \quad tp > p_2 := \varepsilon (|\xi| + \gamma |\xi|).$$

Notice that for $|\xi| < 2\gamma\varepsilon^{-1}(1 + 2C_0)$ and $0 < \gamma < 1$,

$$(9.19) \quad c\varepsilon|\xi| \leq p_1, \quad p_2 \leq C\varepsilon|\xi|$$

and

$$(9.20) \quad 0 \leq p_2 - p_1 \leq \varepsilon|\xi| \left(1 - \frac{2}{1 + \sqrt{1 + O(\gamma)}} \right) + C\varepsilon\gamma|\xi| \leq C\gamma\varepsilon|\xi|.$$

Therefore, if $0 < tp < p_1$, by (4.3) and (9.17),

$$(9.21) \quad \phi \left(\xi + \frac{pt}{\varepsilon} + \frac{C_1 p^2 t^2}{\varepsilon} \right) - \phi \left(\xi + \frac{pt}{\varepsilon} \right) \leq \frac{C}{\left(|\xi| - \frac{pt}{\varepsilon} - \frac{C_1 p^2 t^2}{\varepsilon} \right)^2} \frac{p^2 t^2}{\varepsilon} \leq \frac{C p^2 t^2}{\varepsilon \gamma^2 |\xi|^2},$$

and if $tp > p_2$, by (4.2) and (9.18),

$$(9.22) \quad \phi \left(\xi + \frac{pt}{\varepsilon} + \frac{C_1 p^2 t^2}{\varepsilon} \right) - \phi \left(\xi + \frac{pt}{\varepsilon} \right) \leq 1 - \left(1 - \frac{C}{\xi + \frac{pt}{\varepsilon}} \right) \leq \frac{C}{\gamma |\xi|}.$$

Assume $\gamma > p_2$. The cases $\gamma < p_1$ and $p_1 \leq \gamma \leq p_2$ can be treated similarly. We split

$$II_1 \leq \int_0^{p_1} (\dots) + \int_{p_1}^{p_2} (\dots) + \int_{p_2}^\gamma (\dots) =: J_1 + J_2 + J_3.$$

We further split

$$J_1 = \int_0^{p_1} \frac{dp}{p^2} \int_1^{\frac{p_1}{p}} dt (\dots) + \int_0^{p_1} \frac{dp}{p^2} \int_{\frac{p_1}{p}}^{\frac{p_2}{p}} dt (\dots) + \int_0^{p_1} \frac{dp}{p^2} \int_{\frac{p_2}{p}}^{\frac{\gamma}{p}} dt (\dots) =: J_1^1 + J_1^2 + J_1^3.$$

By (9.19), (9.21) and for $\varepsilon < \gamma^3$,

$$\begin{aligned}
J_1^1 &\leq \frac{C}{\varepsilon \gamma^2 |\xi|^2} \int_0^{p_1} dp \int_1^{\frac{p_1}{p}} \frac{dt}{t^{n-1}} \leq \frac{C}{\varepsilon \gamma^2 |\xi|^2} \int_0^{p_1} dp \int_1^{\frac{p_1}{p}} \frac{dt}{t} = \frac{C}{\varepsilon \gamma^2 |\xi|^2} \int_0^{p_1} (\ln p_1 - \ln p) dp \\
&= \frac{C p_1}{\varepsilon \gamma^2 |\xi|^2} \leq \frac{C}{\gamma^2 |\xi|} \leq \frac{C \gamma}{\varepsilon |\xi|}.
\end{aligned}$$

By (9.19) and (9.20),

$$\begin{aligned}
J_1^2 &\leq 2 \int_0^{p_1} \frac{dp}{p^2} \int_{\frac{p_1}{p}}^{\frac{p_2}{p}} \frac{dt}{t^{n+1}} \leq 2 \int_0^{p_1} \frac{dp}{p^2} \int_{\frac{p_1}{p}}^{\frac{p_2}{p}} \frac{dt}{t^3} = \int_0^{p_1} \left(\frac{1}{p_1^2} - \frac{1}{p_2^2} \right) dp = \frac{(p_2 - p_1)(p_1 + p_2)}{p_1 p_2^2} \\
&\leq \frac{C \gamma}{\varepsilon |\xi|}.
\end{aligned}$$

By (9.19), (9.22) and since $|\xi| > \gamma^{-2}$,

$$J_1^3 \leq \frac{C}{\gamma|\xi|} \int_0^{p_1} \frac{dp}{p^2} \int_{\frac{p_2}{p}}^{\frac{\gamma}{p}} \frac{dt}{t^{n+1}} \leq \frac{C}{\gamma|\xi|} \int_0^{p_1} \frac{dp}{p^2} \int_{\frac{p_2}{p}}^{\infty} \frac{dt}{t^3} = \frac{C}{\gamma|\xi|} \int_0^{p_1} \frac{dp}{p^2} = \frac{Cp_1}{\gamma|\xi|p_2^2} \leq \frac{C}{\varepsilon\gamma|\xi|^2} \leq \frac{C\gamma}{\varepsilon|\xi|}.$$

Therefore,

$$J_1 \leq \frac{C\gamma}{\varepsilon|\xi|}.$$

Next, by (9.19) and (9.20),

$$J_2 \leq 2 \int_{p_1}^{p_2} \frac{dp}{p^2} \int_1^{\infty} \frac{dt}{t^{n+1}} \leq C \int_{p_1}^{p_2} \frac{dp}{p^2} = C \frac{p_2 - p_1}{p_1 p_2} \leq \frac{C\gamma}{\varepsilon|\xi|}.$$

Finally, if $p_2 < p < \gamma$ and $t > 1$, then $tp > p_2$. Hence, by (9.19), (9.22) and since $|\xi| > \gamma^{-2}$,

$$J_3 \leq \frac{C}{\gamma|\xi|} \int_{p_2}^{\gamma} \frac{dp}{p^2} \int_1^{\infty} \frac{dt}{t^{n+1}} \leq \frac{C}{\gamma|\xi|} \int_{p_2}^{\gamma} \frac{dp}{p^2} \leq \frac{C}{\gamma|\xi|p_2} \leq \frac{C}{\varepsilon\gamma|\xi|^2} \leq \frac{C\gamma}{\varepsilon|\xi|}.$$

From the last three estimates, we infer that

$$\varepsilon II_1 \leq \frac{C\gamma}{|\xi|}.$$

Similarly, one gets the same upper bound for εII_2 and thus

$$\varepsilon II \leq \frac{C\gamma}{|\xi|},$$

which together with (9.12) and (9.16) gives (9.11). The lower bound for $a_\varepsilon(\xi; x)$ is obtained with a similar argument. Estimate (4.14) follows.

This completes the proof of Lemma 4.4. \square

10. PROOF OF LEMMA 4.5

Lemma 4.5 follows from the next four lemmas.

Lemma 10.1. *Let d be as in Definition 2.2. If $|d(t, x)| < \rho$, then there is $C > 0$ such that*

$$\left| a_\varepsilon \left(\frac{d(t, x)}{\varepsilon}; t, x \right) - \left(\varepsilon \mathcal{I}_n \left[\phi \left(\frac{d(t, \cdot)}{\varepsilon} \right) \right] (x) - C_n \mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) \right) \right| \leq \frac{C\varepsilon}{\gamma}.$$

Lemma 10.2. *Let d be as in Definition 2.2. If $|d(t, x)| > \rho$, then there is $C > 0$ such that*

$$\left| a_\varepsilon \left(\frac{d(t, x)}{\varepsilon}; t, x \right) \right| \leq \frac{C\varepsilon}{\rho}.$$

Lemma 10.3. *Let d be as in Definition 2.2. If $|d(t, x)| > \rho$, then there is $C > 0$ such that*

$$\left| \mathcal{I}_n \left[\phi \left(\frac{d(t, \cdot)}{\varepsilon} \right) \right] (x) \right| \leq \frac{C}{\rho}.$$

Lemma 10.4. *Let d be as in Definition 2.2. If $|d(t, x)| > \rho$, then there is $C > 0$ such that*

$$\left| \mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) \right| \leq \frac{C\varepsilon}{\rho}.$$

10.1. **Proof of Lemma 10.1.** First, we write $a_\varepsilon = a_\varepsilon(d(t, x)/\varepsilon; t, x)$ as

$$(10.1) \quad \begin{aligned} a_\varepsilon &= \text{P. V.} \int_{\mathbb{R}^n} \left(\phi \left(\frac{d(t, x + \varepsilon z)}{\varepsilon} \right) - \phi \left(\frac{d(t, x)}{\varepsilon} \right) \right) \frac{dz}{|z|^{n+1}} \\ &\quad - \text{P. V.} \int_{\mathbb{R}^n} \left(\phi \left(\frac{d(t, x)}{\varepsilon} + \nabla d(t, x) \cdot z \right) - \phi \left(\frac{d(t, x)}{\varepsilon} \right) \right) \frac{dz}{|z|^{n+1}} \\ &\quad - \int_{|z| > \frac{\gamma}{\varepsilon}} \left(\phi \left(\frac{d(t, x + \varepsilon z)}{\varepsilon} \right) - \phi \left(\frac{d(t, x)}{\varepsilon} + \nabla d(t, x) \cdot z \right) \right) \frac{dz}{|z|^{n+1}}. \end{aligned}$$

Recalling that $0 < \phi < 1$, we first estimate

$$\int_{|z| > \frac{\gamma}{\varepsilon}} \left| \phi \left(\frac{d(t, x + \varepsilon z)}{\varepsilon} \right) - \phi \left(\frac{d(t, x)}{\varepsilon} + \nabla d(t, x) \cdot z \right) \right| \frac{dz}{|z|^{n+1}} \leq 2 \int_{|z| > \frac{\gamma}{\varepsilon}} \frac{dz}{|z|^{n+1}} = \frac{C\varepsilon}{\gamma}.$$

Since $e = \nabla d(t, x)$ is a unit vector when $|d(t, x)| \leq \rho$, by applying Lemma 3.2 to the second integral in the right-hand side of (10.1) and a change of variable in the first integral, we obtain

$$\begin{aligned} a_\varepsilon &= \varepsilon \text{P. V.} \int_{\mathbb{R}^n} \left(\phi \left(\frac{d(t, x + z)}{\varepsilon} \right) - \phi \left(\frac{d(t, x)}{\varepsilon} \right) \right) \frac{dz}{|z|^{n+1}} - C_n \mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) + O \left(\frac{\varepsilon}{\gamma} \right) \\ &= \varepsilon \mathcal{I}_n \left[\phi \left(\frac{d(t, \cdot)}{\varepsilon} \right) \right] (x) - C_n \mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) + O \left(\frac{\varepsilon}{\gamma} \right). \end{aligned}$$

□

10.2. **Proof of Lemma 10.2.** The lemma is an immediate consequence of (4.8).

10.3. **Proof of Lemma 10.3.** We drop the notation in t . For $c = \|\nabla d\|_\infty$, we write

$$\begin{aligned} \mathcal{I}_n \left[\phi \left(\frac{d(\cdot)}{\varepsilon} \right) \right] (x) &= \text{P. V.} \int_{\mathbb{R}^n} \left(\phi \left(\frac{d(x + z)}{\varepsilon} \right) - \phi \left(\frac{d(x)}{\varepsilon} \right) \right) \frac{dz}{|z|^{n+1}} \\ &= \int_{|z| < \frac{\rho}{2c}} \left(\phi \left(\frac{d(x + z)}{\varepsilon} \right) - \phi \left(\frac{d(x)}{\varepsilon} \right) - \dot{\phi} \left(\frac{d(x)}{\varepsilon} \right) \frac{\nabla d(x)}{\varepsilon} \cdot z \right) \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{|z| > \frac{\rho}{2c}} \left(\phi \left(\frac{d(x + z)}{\varepsilon} \right) - \phi \left(\frac{d(x)}{\varepsilon} \right) \right) \frac{dz}{|z|^{n+1}} \\ &=: I + II. \end{aligned}$$

Looking first at the long-range interactions, we use $0 < \phi < 1$ to estimate

$$|II| \leq 2 \int_{|z| > \frac{\rho}{2c}} \frac{dz}{|z|^{n+1}} = \frac{C}{\rho}.$$

For the short-range interactions, fix z such that $|z| < \rho/(2c)$. Then, by Taylor's theorem,

$$\begin{aligned} &\phi \left(\frac{d(x + z)}{\varepsilon} \right) - \phi \left(\frac{d(x)}{\varepsilon} \right) - \dot{\phi} \left(\frac{d(x)}{\varepsilon} \right) \frac{\nabla d(x)}{\varepsilon} \cdot z \\ &= \frac{1}{2} \left(\ddot{\phi} \left(\frac{d(x + \tau z)}{\varepsilon} \right) \frac{\nabla d(x + \tau z) \otimes \nabla d(x + \tau z)}{\varepsilon^2} + \dot{\phi} \left(\frac{d(x + \tau z)}{\varepsilon} \right) \frac{D^2 d(x + \tau z)}{\varepsilon} \right) z \cdot z \end{aligned}$$

for some $0 \leq \tau \leq 1$. Since

$$\left| \frac{d(x + \tau z)}{\varepsilon} \right| \geq \frac{|d(x)|}{\varepsilon} - \frac{c|z|}{\varepsilon} > \frac{\rho}{\varepsilon} - \frac{\rho}{2\varepsilon} = \frac{\rho}{2\varepsilon},$$

we can apply (4.3) to estimate

$$\left| \dot{\phi} \left(\frac{d(x + \tau z)}{\varepsilon} \right) \right| \leq \frac{C\varepsilon^2}{\rho^2} \quad \text{and} \quad \left| \ddot{\phi} \left(\frac{d(x + \tau z)}{\varepsilon} \right) \right| \leq \frac{C\varepsilon^2}{\rho^2}.$$

Therefore, with the regularity of d ,

$$\left| \phi \left(\frac{d(x + z)}{\varepsilon} \right) - \phi \left(\frac{d(x)}{\varepsilon} \right) - \dot{\phi} \left(\frac{d(x)}{\varepsilon} \right) \frac{\nabla d(x)}{\varepsilon} \cdot z \right| \leq \frac{C}{\rho^2} |z|^2.$$

Thus, we have that

$$|I| \leq \frac{C}{\rho^2} \int_{|z| < \frac{\rho}{2c}} |z|^2 \frac{dz}{|z|^{n+1}} = \frac{C}{\rho}.$$

The conclusion follows by combining the estimates for I and II .

10.4. Proof of Lemma 10.4. From (1.4), the periodicity of W and $W'(0) = 0$, we see that

$$\mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) = \frac{1}{C_n} W' \left(\phi \left(\frac{d(t, x)}{\varepsilon} \right) \right) = \frac{1}{C_n} W' \left(\tilde{\phi} \left(\frac{d(t, x)}{\varepsilon} \right) \right) = O \left(\tilde{\phi} \left(\frac{d(t, x)}{\varepsilon} \right) \right),$$

where, as usual, $\tilde{\phi}(\xi) = \phi(\xi) - H(\xi)$ and H is the Heaviside function. Therefore, by (4.2),

$$\left| \mathcal{I}_1[\phi] \left(\frac{d(t, x)}{\varepsilon} \right) \right| \leq \frac{C\varepsilon}{|d(t, x)|} \leq \frac{C\varepsilon}{\rho}.$$

The proof of Lemma 4.5 is then completed. \square

11. PROOF OF LEMMA 4.9

Lemma 4.9 is a consequence of the following three lemmas.

Lemma 11.1. *Let d be as in Definition 2.2. If $|d(t, x)| < \rho$, then there is $C > 0$ such that*

$$\left| \varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(t, \cdot)}{\varepsilon}; t, \cdot \right) \right] (x) - C_n \mathcal{I}_1[\psi(\cdot; t, x)] \left(\frac{d(t, x)}{\varepsilon} \right) \right| \leq \frac{C}{|\ln \varepsilon|^{\frac{1}{2}}}.$$

Lemma 11.2. *Let d be as in Definition 2.2. If $|d(t, x)| > \rho$, then there is $C > 0$ such that*

$$\left| \mathcal{I}_1[\psi(\cdot; t, x)] \left(\frac{d(t, x)}{\varepsilon} \right) \right| \leq \frac{C}{|\ln \varepsilon| \rho}.$$

Lemma 11.3. *Let d be as in Definition 2.2. If $|d(t, x)| > \rho$, then there is $C > 0$ such that*

$$\left| \varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(t, \cdot)}{\varepsilon}; t, \cdot \right) \right] (x) \right| \leq \frac{C}{|\ln \varepsilon|^{\frac{1}{2}} \rho}.$$

11.1. Proof of Lemma 11.1. For simplicity in notation, we drop the dependence on t . By Lemma 3.2 and the fact that $|\nabla d(x)| = 1$ if $|d(x)| < \rho$, we have

$$\begin{aligned} & \varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(\cdot)}{\varepsilon}; \cdot \right) \right] (x) - C_n \mathcal{I}_1[\psi(\cdot; x)] \left(\frac{d(x)}{\varepsilon} \right) \\ &= \text{P. V.} \int_{\mathbb{R}^n} \left(\psi \left(\frac{d(x + \varepsilon z)}{\varepsilon}; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right) \frac{dz}{|z|^{n+1}} \\ &= \text{P. V.} \int_{\mathbb{R}^n} \left(\psi \left(\frac{d(x + \varepsilon z)}{\varepsilon}; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) \right) \frac{dz}{|z|^{n+1}} \\ &\quad + \text{P. V.} \int_{\mathbb{R}^n} \left(\psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right) \frac{dz}{|z|^{n+1}} \\ &=: I + II. \end{aligned}$$

First, let us estimate I . We will show that

$$(11.1) \quad |I| \leq \frac{C}{|\ln \varepsilon|}.$$

Notice that

$$(11.2) \quad \left| \psi \left(\frac{d(x + \varepsilon z)}{\varepsilon}; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) \right| \leq C \|\dot{\psi}\|_{\infty} \varepsilon |z|^2,$$

so we can actually write

$$I = \int_{\mathbb{R}^n} \left(\psi \left(\frac{d(x + \varepsilon z)}{\varepsilon}; x \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right) \frac{dz}{|z|^{n+1}}.$$

From here, we split

$$I = \int_{\{|z| < \varepsilon^{-\frac{1}{2}}\}} (\dots) + \int_{\{|z| > \varepsilon^{-\frac{1}{2}}\}} (\dots) =: I_1 + I_2.$$

By (11.2) and (4.20) for $\dot{\psi}$

$$|I_1| \leq C \varepsilon \|\dot{\psi}\|_{\infty} \int_{\{|z| < \varepsilon^{-\frac{1}{2}}\}} \frac{dz}{|z|^{n-1}} \leq \frac{C \varepsilon}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|} \varepsilon^{-\frac{1}{2}} = \frac{C}{|\ln \varepsilon|},$$

and from (4.20) for ψ ,

$$|I_2| \leq 2 \|\psi\|_{\infty} \int_{\{|z| > \varepsilon^{-\frac{1}{2}}\}} \frac{dz}{|z|^{n+1}} \leq C \frac{\varepsilon^{\frac{1}{2}}}{\varepsilon^{\frac{1}{2}} |\ln \varepsilon|} = \frac{C}{|\ln \varepsilon|}.$$

This proves (11.1).

Next, using that P. V. $\int_{\{|z| < |\ln \varepsilon|^{\frac{1}{2}}\}} \nabla_x \psi \left(\frac{d(x)}{\varepsilon}; x \right) \cdot z \frac{dz}{|z|^{n+1}} = 0$, we write

$$\begin{aligned} II &= \int_{\{|z| < |\ln \varepsilon|^{\frac{1}{2}}\}} \left\{ \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right. \\ &\quad \left. - \nabla_x \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \cdot (\varepsilon z) \right\} \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{\{|z| < |\ln \varepsilon|^{\frac{1}{2}}\}} \left\{ \nabla_x \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) - \nabla_x \psi \left(\frac{d(x)}{\varepsilon}; x \right) \right\} \cdot (\varepsilon z) \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{\{|z| > |\ln \varepsilon|^{\frac{1}{2}}\}} \left\{ \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right\} \frac{dz}{|z|^{n+1}} \\ &=: II_1 + II_2 + II_3. \end{aligned}$$

By (4.20) for $D_x^2 \psi$,

$$|II_1| \leq C \|D_x^2 \psi\|_{\infty} \varepsilon^2 \int_{\{|z| < |\ln \varepsilon|^{\frac{1}{2}}\}} \frac{dz}{|z|^{n-1}} \leq \frac{C}{\varepsilon^2 |\ln \varepsilon|} \varepsilon^2 |\ln \varepsilon|^{-\frac{1}{2}} = \frac{C}{|\ln \varepsilon|^{\frac{1}{2}}}.$$

By (4.22) for $\nabla_x \dot{\psi}$,

$$|II_2| \leq \|\nabla_x \dot{\psi}(\cdot; x)\|_{\infty} \varepsilon \int_{\{|z| < |\ln \varepsilon|^{\frac{1}{2}}\}} \frac{dz}{|z|^{n-1}} \leq \frac{C}{\varepsilon |\ln \varepsilon|} \varepsilon |\ln \varepsilon|^{\frac{1}{2}} = \frac{C}{|\ln \varepsilon|^{\frac{1}{2}}}.$$

Finally, from (4.22) for ψ ,

$$|II_3| \leq 2 \|\psi\|_{\infty} \int_{\{|z| > |\ln \varepsilon|^{\frac{1}{2}}\}} \frac{dz}{|z|^{n+1}} \leq \frac{C}{|\ln \varepsilon|^{\frac{1}{2}}}.$$

This proves that

$$(11.3) \quad |II| \leq \frac{C}{|\ln \varepsilon|^{\frac{1}{2}}}.$$

From (11.1) and (11.3), we have Lemma 11.1.

11.2. Proof of Lemma 11.2. From (4.16), (4.2), (4.3), (4.8), (4.10) and (4.21), for $|\xi| \geq 1$,

$$\begin{aligned} |\mathcal{I}_1[\psi(\cdot; t, x)](\xi)| &\leq C |\psi(\xi)| + \frac{|a_\varepsilon(\xi; t, x)|}{\varepsilon |\ln \varepsilon|} + C |1 + \bar{a}_\varepsilon(t, x)| \dot{\phi}(\xi) + C |\phi(\xi)| \\ &\leq \frac{C}{\varepsilon |\ln \varepsilon| |\xi|}. \end{aligned}$$

In particular,

$$\left| \mathcal{I}_1[\psi(\cdot; t, x)] \left(\frac{d(t, x)}{\varepsilon} \right) \right| \leq \frac{C}{|\ln \varepsilon| |d(t, x)|} \leq \frac{C}{|\ln \varepsilon| \rho}.$$

11.3. Proof of Lemma 11.3. For convenience, we drop the dependence on t and begin by writing

$$(11.4) \quad \begin{aligned} \varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(\cdot)}{\varepsilon}; \cdot \right) \right] (x) &= \varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(\cdot)}{\varepsilon}; \cdot \right) \right] (x) - |\nabla d(x)| C_n \mathcal{I}_1[\psi(\cdot; x)] \left(\frac{d(x)}{\varepsilon} \right) \\ &\quad + |\nabla d(x)| C_n \mathcal{I}_1[\psi(\cdot; x)] \left(\frac{d(x)}{\varepsilon} \right). \end{aligned}$$

With Lemma 3.2 and as in the proof of Lemma 11.1, we obtain

$$\begin{aligned} &\varepsilon \mathcal{I}_n \left[\psi \left(\frac{d(\cdot)}{\varepsilon}; \cdot \right) \right] (x) - |\nabla d(x)| C_n \mathcal{I}_1[\psi(\cdot; x)] \left(\frac{d(x)}{\varepsilon} \right) \\ &= \text{P. V.} \int_{\mathbb{R}^n} \left(\psi \left(\frac{d(x + \varepsilon z)}{\varepsilon}; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) \right) \frac{dz}{|z|^{n+1}} \\ &\quad + \text{P. V.} \int_{\mathbb{R}^n} \left(\psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right) \frac{dz}{|z|^{n+1}} \\ &=: I + II, \end{aligned}$$

with

$$|I| \leq \frac{C}{|\ln \varepsilon|}.$$

Next, using that $\text{P. V.} \int_{\{|z| < |\ln \varepsilon|^{-\frac{1}{2}}\}} \nabla_x \psi \left(\frac{d(x)}{\varepsilon}; x \right) \cdot z \frac{dz}{|z|^{n+1}} = 0$, we write, for $c = \|\nabla d\|_\infty$,

$$\begin{aligned} II &= \int_{\{|z| < |\ln \varepsilon|^{-\frac{1}{2}}\}} \left\{ \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right. \\ &\quad \left. - \nabla_x \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \cdot (\varepsilon z) \right\} \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{\{|z| < |\ln \varepsilon|^{-\frac{1}{2}}\}} \left\{ \nabla_x \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) - \nabla_x \psi \left(\frac{d(x)}{\varepsilon}; x \right) \right\} \cdot (\varepsilon z) \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{\{|\ln \varepsilon|^{-\frac{1}{2}} < |z| < \frac{\rho}{2c\varepsilon}\}} \left\{ \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right\} \frac{dz}{|z|^{n+1}} \\ &\quad + \int_{\{|z| > \frac{\rho}{2c\varepsilon}\}} \left\{ \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x + \varepsilon z \right) - \psi \left(\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z; x \right) \right\} \frac{dz}{|z|^{n+1}} \end{aligned}$$

$$=: II_1 + II_2 + II_3 + II_4.$$

By (4.20) for $D_x^2\psi$,

$$|II_1| \leq C \|D_x^2\psi\|_\infty \varepsilon^2 \int_{\{|z| < |\ln \varepsilon|^{-\frac{1}{2}}\}} \frac{dz}{|z|^{n-1}} \leq \frac{C}{\varepsilon^2 |\ln \varepsilon|} \varepsilon^2 |\ln \varepsilon|^{-\frac{1}{2}} = \frac{C}{|\ln \varepsilon|^{\frac{3}{2}}}.$$

By (4.20) for $\nabla_x \dot{\psi}$,

$$|II_2| \leq \|\nabla_x \dot{\psi}(\cdot; x)\|_\infty \varepsilon \int_{\{|z| < |\ln \varepsilon|^{-\frac{1}{2}}\}} \frac{dz}{|z|^{n-1}} \leq \frac{C}{\varepsilon} \varepsilon |\ln \varepsilon|^{-\frac{1}{2}} = \frac{C}{|\ln \varepsilon|^{\frac{1}{2}}}.$$

Next, note that if $|d(x)| > \rho$ and $|z| < \frac{\rho}{2c\varepsilon}$, then $|\frac{d(x)}{\varepsilon} + \nabla d(x) \cdot z| > \frac{\rho}{2\varepsilon}$. Therefore, by (4.21),

$$|II_3| \leq \frac{C}{\varepsilon |\ln \varepsilon|} \frac{\varepsilon}{\rho} \int_{\{|z| > |\ln \varepsilon|^{-\frac{1}{2}}\}} \frac{dz}{|z|^{n+1}} = \frac{C}{|\ln \varepsilon|^{\frac{1}{2}} \rho}.$$

Finally, from (4.20) for ψ ,

$$|II_4| \leq 2\|\psi\|_\infty \int_{\{|z| > \frac{\rho}{2c\varepsilon}\}} \frac{dz}{|z|^{n+1}} \leq \frac{C\varepsilon^{\frac{1}{2}}}{|\ln \varepsilon| \rho}.$$

We conclude that

$$\left| \mathcal{I}_n \left[\psi \left(\frac{d(\cdot)}{\varepsilon}; \cdot \right) \right] (x) - |\nabla d(x)| C_n \mathcal{I}_1[\psi(\cdot; x)] \right| \leq \frac{C}{|\ln \varepsilon|^{\frac{1}{2}} \rho}.$$

Moreover, by Lemma 11.2,

$$|\nabla d(x)| C_n \mathcal{I}_1[\psi(\cdot; x)] \leq \frac{C}{|\ln \varepsilon| \rho}.$$

Recalling (11.4), the lemma then follows.

The proof of Lemma 4.9 is then completed. \square

ACKNOWLEDGEMENTS

The first author has been supported by the NSF Grant DMS-2155156 “Nonlinear PDE methods in the study of interphases.” The second author has been supported by the Australian Laureate Fellowship FL190100081 “Minimal surfaces, free boundaries and partial differential equations.” Both authors acknowledge the support of NSF Grant DMS RTG 18403.

REFERENCES

- [1] L. AMBROSIO, G. DE PHILIPPIS AND L. MARTINAZZI, Gamma-convergence of nonlocal perimeter functionals, *Manuscripta Math.* **134** (2011), no 3-4, 377–403.
- [2] R. ALICANDRO, L. DE LUCA, A. GARRONI AND M. PONSIGLIONE, Metastability and dynamics of discrete topological singularities in two dimensions: a Γ -convergence approach, *Arch. Rational Mech. Anal.* **214** (2014), 269–330.
- [3] R. ALICANDRO, L. DE LUCA, A. GARRONI AND M. PONSIGLIONE, Dynamics of discrete screw dislocations on glide directions, *J. Mech. Phys. Solids*, **92** (2016), 87–104.
- [4] G. BARLES AND F. DA LIO, A geometrical approach to front propagation problems in bounded domains with Neumann-type boundary conditions, *Interfaces Free Bound.* **5** (2003), 239–274.
- [5] G. BARLES, H. M. SONER AND P. E. SOUGANIDIS, Front propagation and phase field theory, *SIAM J. Control Optim.* **31** (1993), 439–469.
- [6] G. BARLES AND P. E. SOUGANIDIS, A new approach to front propagation problems: theory and applications, *Arch. Rational Mech. Anal.* **141** (1998), 237–296.
- [7] L. BROWN, The self-stress of dislocations and the shape of extended nodes, *Phil. Mag.* **10** (1964), p. 441.

- [8] X. CABRÉ AND J. SOLÀ-MORALES, Layer solutions in a half-space for boundary reactions, *Comm. Pure Appl. Math.*, **58** (2005), 911–941.
- [9] X. CHEN, Generation and propagation of interfaces for reaction-diffusion equations, *J. Differential Equations* **96** (1992), 116–141.
- [10] Y. G. CHEN, Y. GIGA AND S. GOTO, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Differential Geom.* **33** (1991), 749–786.
- [11] S. CONTI, A. GARRONI AND S. MÜLLER, Derivation of strain-gradient plasticity from a generalized Peierls-Nabarro model, *J. Eur. Math. Soc.* **25** (2023), 2487–2524.
- [12] M. G. CRANDALL, H. ISHII AND P.-L. LIONS, User’s guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc.* **27** (1992), 1–67.
- [13] F. DA LIO, N. FORCADEL AND R. MONNEAU, Convergence of a non-local eikonal equation to anisotropic mean curvature motion. Application to dislocation dynamics, *J. Eur. Math. Soc.* **10** (2008), 1061–1104.
- [14] E. DI NEZZA, G. PALATUCCI AND E. VALDINOCI, Hitchhiker’s guide to the fractional Sobolev spaces, *Bull. Sci. Math.* **136** (2012), 521–573.
- [15] S. DIPIERRO, S. PATRIZI AND E. VALDINOCI, A fractional glance to the theory of edge dislocations, In *Geometric and Functional Inequalities and Recent Topics in Nonlinear PDE’s*, *Contemp. Math.* **781**, 103–135. American Mathematical Society, Providence, RI, 2023
- [16] L. C. EVANS AND J. SPRUCK, Motion of level sets by mean curvature. I. *J. Differential Geom.* **33** (1991), 635–681.
- [17] A. Z. FINO, H. IBRAHIM AND R. MONNEAU, The Peierls-Nabarro model as a limit of a Frenkel-Kontorova model, *J. Differential Equations* **252** (2012), 258–293.
- [18] N. FORCADEL, C. IMBERT AND R. MONNEAU, Homogenization of some particle systems with two-body interactions and of the dislocation dynamics, *Discrete Contin. Dyn. Syst.* **23** (2009), 785–826.
- [19] A. GARRONI, G. LEONI AND M. PONSIGLIONE, Gradient theory for plasticity via homogenization of discrete dislocations, *J. Eur. Math. Soc.*, **12** (2010), 1231–1266.
- [20] A. GARRONI, P. VAN MEURS, M. PELETIER AND L. SCARDIA, Convergence and non-convergence of many-particle evolutions with multiple signs, *Arch. Ration. Mech. Anal.*, **235** (2020), 3–49.
- [21] A. GARRONI AND S. MÜLLER, A variational model for dislocations in the line tension limit, *Arch. Ration. Mech. Anal.* **181** (2006), 535–578.
- [22] D. GILBARG AND N. S. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin Heidelberg, 2001.
- [23] M. GONZÁLEZ AND R. MONNEAU, Slow motion of particle systems as a limit of a reaction-diffusion equation with half-Laplacian in dimension one, *Discrete Contin. Dyn. Syst.* **32** (2012), 1255–1286.
- [24] A. EL HAJJ, H. IBRAHIM AND R. MONNEAU, Dislocation dynamics: from microscopic models to macroscopic crystal plasticity, *Contin. Mech. Thermodyn.* **21** (2009), 109–123.
- [25] J. R. HIRTH AND L. LOTHE, *Theory of Dislocations*, Second Edition. Malabar, Florida: Krieger, 1992.
- [26] D. HULL AND D. J. BACON, *Introduction to Dislocations*, ButterworthHeinemann, Oxford, 2011.
- [27] C. IMBERT, Level set approach for fractional mean curvature flows, *Interfaces Free Bound.* **11** (2009), 153–176.
- [28] C. IMBERT, R. MONNEAU AND E. ROUY, Homogenization of first order equations with u/ϵ -periodic Hamiltonians. Part II: application to dislocations dynamics, *Communications in Partial Differential Equations* **33** (2008), 479–516.
- [29] C. IMBERT AND P. E. SOUGANIDIS, Phasefield theory for fractional diffusion-reaction equations and applications, preprint (2009), pp 41.
- [30] P. VAN MEURS, M. A. PELETIER AND N. POŽÁR, , Discrete-to-continuum convergence of charged particles in 1D with annihilation, *Arch. Ration. Mech. Anal.*, **246** (2022), 241–297.
- [31] P. VAN MEURS AND S. PATRIZI, Discrete dislocation dynamics with annihilation as the limit of the Peierls-Nabarro model in one dimension, *SIAM J. Math. Anal.* **56** (2024), 197–233.
- [32] L. MODICA AND S. MORTOLA, Un esempio di Γ^- -convergenza, *Boll. Un. Mat. Ital. B (5)* **14** (1977), 285–299.
- [33] C. W. PASSCHIER AND R. A. J. TROUW, *Microtectonics*. Springer, 1998.
- [34] R. MONNEAU AND S. PATRIZI, Derivation of the Orowan’s law from the Peierls–Nabarro model, *Commun. Partial. Differ. Equ.* **37** (2012), 1887–1911.
- [35] R. MONNEAU AND S. PATRIZI, Homogenization of the Peierls-Nabarro model for dislocation dynamics, *J. Differential Equations* **253** (2012), 2064–2105.
- [36] F. R. N. NABARRO, Dislocations in a simple cubic lattice, *Proc. Phys. Soc.* **59** (1947), 256–272.

- [37] F. R. N. NABARRO, Fifty-year study of the Peierls–Nabarro stress, *Mater. Sci. Engrg. A* **234–236** (1997), 67–76.
- [38] S. OSHER AND J. A. SETHIAN, Fronts propagating with curvature dependent speed: algorithms based on Hamilton–Jacobi formulations, *J. Computational Phys.* **79** (1988), 12–49.
- [39] S. PATRIZI AND T. SANGSAWANG, From the Peierls–Nabarro model to the equation of motion of the dislocation continuum, *Nonlinear Anal.* **202** (2021), 112096.
- [40] S. PATRIZI AND T. SANGSAWANG, Derivation of the 1-D Groma–Balogh equations from the Peierls–Nabarro model, *Calc. Var. Partial Differential Equations* **62** (2023), 55 pp.
- [41] S. PATRIZI AND E. VALDINOCI, Crystal dislocations with different orientations and collisions, *Arch. Rational Mech. Anal.* **217** (2015), 231–261.
- [42] S. PATRIZI AND E. VALDINOCI, Homogenization and Orowan’s law for anisotropic fractional operators of any order, *Nonlinear Anal.* **119** (2015), 3–36.
- [43] S. PATRIZI AND E. VALDINOCI, Long-time behavior for crystal dislocation dynamics, *Math. Models Methods Appl. Sci.* **27** (2017), 2185–2228.
- [44] S. PATRIZI AND E. VALDINOCI, Relaxation times for atom dislocations in crystals, *Calc. Var. Partial Differential Equations* **55** (2016), 1–44.
- [45] S. PATRIZI AND M. VAUGHAN, A convergence result for the derivation of front propagation in nonlocal phase field models, preprint (2024), pp. 39.
- [46] R. PEIERLS, The size of a dislocation, *Proc. Phys. Soc.* **52** (1940), 34–37.
- [47] O. SAVIN AND E. VALDINOCI, Γ -convergence for nonlocal phase transitions *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **29** (2012), 479–500.
- [48] R. SEDLACEK, J. KRATOCHVI AND E. WERNER, The importance of being curved: bowing dislocations in a continuum description, *Philosophical Magazine* **83** (2003), 3735–3752.
- [49] P. R. STINGA, User’s guide to fractional Laplacians and the method of semigroups, In Anatoly Kochubei, Yuri Luchko (Eds.), *Fractional Differential Equations*, 235–266, Berlin, Boston: De Gruyter, 2019.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, 2515 SPEEDWAY, AUSTIN, TX 78712, UNITED STATES OF AMERICA

Email address: `spatrizi@math.utexas.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF WESTERN AUSTRALIA, 35 STIRLING HWY, CRAWLEY WA 6009, AUSTRALIA

Email address: `mary.vaughan@uwa.edu.au`