

Probabilistic Emulation of a Global Climate Model with Spherical Dyffusion

Salva Rühling Cachay
UC San Diego

Brian Henn
Allen Institute for AI

Oliver Watt-Meyer
Allen Institute for AI

Christopher S. Bretherton
Allen Institute for AI

Rose Yu
UC San Diego

Abstract

Data-driven deep learning models are on the verge of transforming global weather forecasting. It is an open question if this success can extend to climate modeling, where long inference rollouts and data complexity pose significant challenges. Here, we present the first conditional generative model able to produce global climate ensemble simulations that are accurate and physically consistent. Our model runs at 6-hourly time steps and is shown to be stable for 10-year-long simulations. Our approach beats relevant baselines and nearly reaches a gold standard for successful climate model emulation. We discuss the key design choices behind our dynamics-informed diffusion model-based approach which enables this significant step towards efficient, data-driven climate simulations that can help us better understand the Earth and adapt to a changing climate.

1 Introduction

Climate models are foundational tools used to understand how the Earth system evolves over long time periods and how it may change as a response to possible greenhouse gas emission scenarios. Such climate simulations are currently very expensive to generate due to the computational complexity of the underlying physics-based climate models, which must be run on supercomputers. As a result, scientists and policymakers are limited to exploring only a small subset of possibilities for different mitigation and adaptation strategies [38].

Training relatively cheap-to-run data-driven surrogates to emulate global climate models could provide a compelling alternative. Although recent deep learning models are on the verge of transforming the conceptually similar field of medium-range weather forecasting [4, 31, 8, 41], these advances do not directly transfer to long-term climate projections. Indeed, most such models only report forecasts up to two weeks into the future and may diverge or become physically inconsistent over longer simulations. In contrast, climate projections demand accurate and stable simulations of the global Earth system spanning decades or centuries, requiring reliable reproduction of long-term statistics.

In Figure 1 we quantitatively show this divergence between the medium-range weather forecasting skill of ML models (measured as the average RMSE on 5-day forecasts) and their performance on longer




Figure 1: Weather performance (x-axis) is not a strong indicator of climate performance (y-axis). Each dot corresponds to a distinct sample or checkpoint epoch.

climate time scales (measured as the RMSE of the 10-year time-mean). We have verified that this finding holds regardless of the analyzed variable and the proxy used for weather performance, which we discuss in more detail in Appendix D.4. Heuristically, optimizing weather skill ensures that a climate model takes a locally accurate path around the climate 'attractor', but it does not guarantee that small but systematic errors may not build up to distort that simulated attractor to have biased long-term climate statistics. While this is a little-discussed observation in the ML community, the climate modeling community has documented it for physics-based models [13, 45].

A recent breakthrough is a deterministic surrogate called ACE (AI2 Climate Emulator) [58], which remains remarkably stable and physically consistent over 10-year simulations at 6-hourly time steps, forced by time-varying specified sea-surface temperature and sea-ice. Its success can be attributed to careful data processing, problem design, and the Spherical Fourier Neural Operator (SFNO) [5] architecture. ACE is trained to emulate the United States' primary operational global forecast model, the physics-based FV3GFS [63], which is operationally used at the US National Weather Service and US National Centers for Environmental Prediction. ACE produces encouragingly small ten-year mean climate biases (i.e. biased long-term averages), but they are still significantly larger than the theoretical minimum imposed by internal variability of the reference physics-based model.

ACE's deterministic nature restricts its ability to model the full distribution of climate states or to facilitate ensemble simulations, which involve drawing multiple samples from the same model. These capabilities are crucial for climate modeling, as they enable better uncertainty quantification, more robust and physically consistent predictions, and a deeper understanding of potential future climate scenarios and associated risks [26]. While it is possible to ensemble a deterministic model by perturbing its inputs, this approach often leads to under-dispersed (i.e. overly confident) ensembles compared to generative or physics-based approaches [48]. Even then, the problem remains that due to optimizing them on MSE-based loss functions, the deterministic predictions may degrade to a mean prediction for longer forecast time scales and underestimate unlikely events [6].

A generative modeling approach, particularly the use of diffusion models [50, 20], appears to be a promising solution to these challenges. However, standard diffusion models are computationally intensive to train and sample from. This complexity poses significant problems for climate modeling because: 1) atmospheric data is extremely high-dimensional, making the use of video diffusion models [54, 22, 60, 49, 21, 18] prohibitive, even more so as this class of models still struggle with videos longer than a few seconds; and 2) the sampling speed of standard diffusion models is particularly problematic for long, sequential inference rollouts. For instance, generating a single 10-year-long simulation, as in our experiments, with a standard autoregressive diffusion model [29, 41] that uses N diffusion steps would require $14600 \times N$ neural network forward passes. If a second-order solver for sampling is used [25, 41], this number doubles. Even with N as small as 30, this results in half a million forward passes to generate a single sample trajectory, severely limiting the potential of data-driven models to serve as fast surrogates for expensive physics-based models.

As a solution to this computational problem, we build upon the dynamics-informed diffusion model framework, DYffusion, from Rühling Cachay et al. [48], which caps the computational overhead at inference time (as measured by the number of neural net forward passes) to less than $3 \times$ as much as for a deterministic next-step forecasting model such as SFNO or ACE. Unfortunately, the original DYffusion method relies on an UNet-based architecture designed for Euclidean data rather than physical fields on a sphere. As we show in Figure 1, this mismatch of inductive biases becomes more problematic at the long climate time scales that we focus on in this paper.

We address these limitations by carefully integrating the DYffusion framework with the SFNO architecture from Bonev et al. [5], and the data and evaluation procedure from Watt-Meyer et al. [58]. To achieve this integration, we extend SFNO with time conditioning and inference stochasticity modules. Our proposed framework, Spherical DYffusion, achieves strong results: On average, across all 34 predicted fields, our model reduces climate biases to within 50% of the reference model, which is more than $2 \times$ and $4 \times$ lower than the best baselines. For critical fields, such as the derived total water path quantity, our method achieves results within 20% of the reference model, representing a $5 \times$ improvement over the next best baseline (see Fig. 2). Additionally, our method proves effective for ensemble climate simulations, reproducing climate variability consistent with the reference model and further reducing climate biases towards the theoretical minimum through ensemble-averaging.

Our generative model is a leap forward toward purely ML-based climate projections that are both efficient and accurate. Our main contributions are:




Figure 2: RMSE of 10-year time-means for a subset of important fields. The leftmost bar in the first two subplots shows the reference noise floor, determined by comparing ten independent 10-year reference FV3GFS simulations with the validation simulation. The scores computed using the mean over these ten simulations (a proxy for an "ensemble prediction") are shown in light shade. The subsequent bars show the corresponding scores for our method and the deep-learning baselines, using a 25-member ensemble for the probabilistic methods (all except ACE, which only reports scores for its single deterministic prediction). Scores computed using the ensemble-mean prediction are shown in light shade. The dark shaded bar on top indicates the performance drop when using a single member's prediction only, with error bars representing the standard deviation over the 25 different member choices. The rightmost subplot displays the average time-mean RMSE of the ML-based emulators relative to the reference across all 34 variables. On average, our method's time-mean RMSEs are 50% higher than the noise floor, which is less than half the average RMSE of the next best method, ACE. When using the 25-member ensemble mean prediction, this reduces to 29.28%.

1. We present the first conditional generative model for probabilistic emulation of a realistic climate model, with minimal computational overhead over deterministic baselines.
2. We carefully integrate two distinct frameworks, ACE and Dyffusion, including additional modifications to the SFNO architecture such as time-conditioning modules.
3. We show that our integrated method performs considerably better than relevant baselines in terms of reduced climate biases, ensemble-based climate modeling, and consistent variability of the climate predictions.
4. We show that weather forecasting performance is not a strong indicator of long-term climate biases.

2 Related Work

ML for weather and climate modeling. There are fundamental differences in weather and climate modeling. Climate refers to the average weather over long periods of time¹. While weather forecasting focuses on short time scales in the order of days or weeks, climate modeling simulates longer periods of decades to centuries. Weather forecasting is primarily an initial-value problem, for which it is important to analyze short-term time-specific predictions. Climate modeling is primarily a boundary-condition (or forcing-driven) problem [56], characterized by long-term averages and distributions.

Deep learning-based models have emerged as a much more computationally efficient alternative to traditional physics-based numerical weather prediction (NWP) models, showing impressive skill for deterministic medium-range weather forecasting [39, 27, 4, 5, 7, 37, 31]. This success has been more recently extended to ensemble-based probabilistic weather forecasting [8, 41]. An alternative approach is hybrid modeling, where a physics-based component is complemented by ML-based parameterizations or corrections [42, 62, 47, 1, 30, 61, 28]. At longer lead times, when weather becomes chaotic and less predictable, the ensemble mean prediction of a physics-based or probabilistic ML-based ensemble improves deterministic metrics such as root mean squared error (RMSE) over non-ensembled methods [41, 28, 43].

However, advances in weather forecasting hardly transfer to long-term climate projections. Fully data-driven models fail to maintain stability beyond two-week-ahead forecasts, as errors accumulate over their autoregressive rollouts. Weyn et al. [59] and Bonev et al. [5] showed stable forecasts for horizons of up to six weeks and one year, respectively. Only recently, Watt-Meyer et al. [58] notably achieved stable and accurate 10-year simulations, followed by another deterministic SFNO-based climate emulator showing promising results using four prognostic variables [17]. Easier, but less flexible and informative, alternatives to full-scale temporal modeling of atmospheric dynamics, include emulation of annual means given an emission scenario [57, 36] or temporal super-resolution of monthly means [3].

¹For example, see https://oceanservice.noaa.gov/facts/weather_climate

Diffusion models. Diffusion models [20, 50–52] have demonstrated significant success in generating data such as natural images and videos. Their direct applications to autoregressive forecasting [29, 41] and downscaling [55, 34] of physical data have shown promising results. However, these approaches inherit the computational complexity associated with training and sampling from standard diffusion models. This is particularly prohibitive for autoregressive predictions on climate time scales, as the total number of neural network forward passes increases proportionally with the number of sampling steps, typically ranging from 20 to 1000. Consequently, recent research that leverages insights from diffusion models to balance predictive performance and sampling speed appears more promising for assessing their viability in climate simulations [48, 33]. Diffusion models have been largely used in tandem with a U-Net [46, 11] backbone architecture but recent work reveals promising results for using a vision transformer architecture for natural image synthesis tasks [40, 24, 19]. Our work explores a different, neural operator-based, architecture for Earth data.

3 Background

We first define the problem and then introduce the key components in our framework, namely DYffusion and SFNO. We abbreviate a time series of tensors $\mathbf{y}_0, \dots, \mathbf{y}_t$ with $\mathbf{y}_{0:t}$.

3.1 Problem Setting

Our goal is to learn the probability distribution $P(\mathbf{x}_{1:H} | \mathbf{x}_0, \mathbf{f}_{0:H})$ over a horizon of H time steps, conditional on initial conditions \mathbf{x}_0 and a scenario of forcing variables $\mathbf{f}_{0:H}$ (i.e. time-varying boundary conditions). In our paper, these forcings correspond to prescribed sea surface temperatures and incoming solar radiation (see Section 5.1), leaving it to future work to force based on greenhouse gas emission scenarios explicitly. Each $\mathbf{x}_t \in \mathbb{R}^D$ represents the state of the atmosphere at a given timestep, t , consisting of two- and three-dimensional surface and atmospheric variables across a latitude-longitude grid. These variables, which serve as both input and output, are referred to as *prognostic* variables. We assume a constant time interval between successive time steps t and $t + 1$. To make training feasible, it is necessary to train on a much shorter horizon h , i.e. learn the distribution $P(\mathbf{x}_{t+1:t+h} | \mathbf{x}_t, \mathbf{f}_{t:t+h})$, and apply the model autoregressively. This process begins with $P(\mathbf{x}_{1:h} | \mathbf{x}_0, \mathbf{f}_{0:h})$ and continues until reaching time step H at inference time.

3.2 Diffusion Models and DYffusion

Diffusion models can be seen as a general paradigm to learn the target distribution $p(\mathbf{s}^{(0)})$, by iterating over N diffusion steps of a forward or reverse process. We denote the states of each diffusion step with $\mathbf{s}^{(n)}$, using a superscript n to clearly distinguish them from the physical time steps of the data \mathbf{x}_t . Standard diffusion models [50, 20, 25], initialize the reverse process from a simple isotropic Gaussian distribution $\mathbf{s}^{(N)} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ so that as $n \rightarrow 0$ the intermediate states $\mathbf{s}^{(n)}$ are gradually denoised towards a real data sample $\mathbf{s}^{(0)}$.

In Cold Diffusion [2], this paradigm is extended to more general data corruption processes such as blurring. Rühling Cachay et al. [48] propose DYffusion, by adapting cold diffusion models to forecasting problems. The key idea is to make the forward and reverse processes dynamics-informed by directly coupling them to the physical time steps of the data. That is, the reverse process is initialized with $\mathbf{s}^{(N)} = \mathbf{x}_0$ and iteratively evolves jointly with the dynamics of the data $\mathbf{x}_1, \dots, \mathbf{x}_{h-1}$ to reach the data at some target time step, $\mathbf{s}^{(0)} = \mathbf{x}_h$.

In DYffusion, the forward and reverse processes are informed by temporal dynamics in the data and do not rely on data corruption. Their only source of stochasticity comes from using a stochastic neural network as an operator for the forward process and is implemented by using Monte Carlo (MC) dropout [15]. This forward process essentially corresponds to a temporal interpolator network, while the reverse process is represented by a multi-step forecasting network. Thus, compared to standard diffusion models, DYffusion requires training one more neural network, which they propose doing in separate stages, beginning with the interpolator model. Due to its dynamics-informed nature, DYffusion was shown to be faster at sampling time and more memory-efficient than standard diffusion models, while matching or outperforming their accuracy.




Figure 3: The diagram shows how our proposed approach functions at inference time. Given an initial condition \mathbf{x}_t and forcings $\mathbf{f}_{t:t+h}$, our method uses the DYffusion framework, integrated with two SFNO backbone networks, to generate predictions for the next h time steps based on an alternation of direct multi-step forecasts and temporal interpolations. To simplify the visualization, we exclude the facts that the interpolator network, SFNO_ϕ , is conditioned on \mathbf{x}_t and \mathbf{f}_t in addition to an estimate of \mathbf{x}_{t+h} . We also exclude the time-conditioning of both networks. To forecast more time steps beyond $t+h$, our method is applied autoregressively.




Figure 4: Diagram of one of the blocks of the modified SFNO architecture for our proposed method. The full architecture consists of a sequence of 8 such blocks. Our newly introduced time-conditioning modules correspond to the Time Embedding, followed by the MLP on the right, and the scale-shift operation. Our method relies on dropout, which is part of the two-layer MLP on the top. SFNO-based baselines use the same architecture and hyperparameters without the time embedding module.

3.3 Spherical Fourier Neural Operator (SFNO)

The SFNO architecture [5] extends the FNO framework from Li et al. [32] to spherical data and symmetries such as the Earth. FNOs efficiently model long-range interactions in the Fourier space, but because the underlying Fast Fourier Transform is defined on a Euclidean domain, this can lead to modeling artifacts. SFNOs overcome this issue by using the spherical harmonic transform (SHT) [12], a generalization of the Fourier transform, instead. The SFNO model achieves higher long-term stability of autoregressive rollouts than the FNO model, showing stable forecasts of Earth’s atmospheric dynamics for up to 1-year-long rollouts at six-hourly time steps. The ACE model from Watt-Meyer et al. [58] is based on the SFNO architecture, modifying some of the hyperparameters and the grid used for the first and last SHT of the SFNO. We use the SFNO configuration from ACE in our experiments.

4 Spherical DYffusion

SFNO and ACE are deterministic models that cannot be readily used for uncertainty quantification or ensemble-based climate modeling. DYffusion introduces an efficient diffusion-based approach specifically for forecasting problems but only for Euclidean data. Thus, we propose Spherical DYffusion, a deep generative model for data-driven probabilistic climate simulations that carefully integrates SFNO and DYffusion into an unified framework.

DYffusion requires two neural networks that are used for temporal interpolation and direct multi-step forecasts. In the original framework, these are UNet-like networks. For our approach, we propose to replace them with modified versions of the SFNO architecture, which we denote by SFNO_ϕ and SFNO_θ , respectively.

Training. We follow the original training procedure from DYffusion, complementing it with the use of the input-only forcing variables. That is, for a specified training horizon h , these networks are trained in two stages such that for sequences of prognostic data $\mathbf{x}_{t:t+h}$ and forcings $\mathbf{f}_{t:t+h}$

$$\begin{aligned} \text{SFNO}_\phi(\mathbf{x}_t, \mathbf{x}_{t+h}, \mathbf{f}_t, i | \xi) &\approx \mathbf{x}_{t+i} \\ \text{SFNO}_\theta(\text{SFNO}_\phi(\mathbf{x}_t, \mathbf{x}_{t+h}, \mathbf{f}_t, j | \xi), \mathbf{f}_{t+j}, j) &\approx \mathbf{x}_{t+h}, \end{aligned}$$

where $i \in \{1, \dots, h-1\}$ and we use $j \in \{0, 1, \dots, h-1\}$, defining $\text{SFNO}_\phi(\mathbf{x}_t, \cdot, \cdot, 0 | \xi) = \mathbf{x}_t$. In our experiments, we use $h = 6$. Here, ξ refers to the random variable representing the interpolator network’s inference stochasticity. We discuss its implementation further below. The forecaster network, SFNO_θ , is deterministic. The full training scheme is defined in Algorithm 1.

Inference. At inference time, we follow the DYffusion sampling scheme based on cold sampling [2]. Essentially, we start with the initial conditions \mathbf{x}_0 to generate a first forecast of time step h through a forward pass of the forecaster network, i.e. $\hat{\mathbf{x}}_h = \text{SFNO}_\theta(\mathbf{x}_0, \mathbf{f}_0, 0)$. Given this prediction, we can now use the interpolator network to interpolate $\hat{\mathbf{x}}_1 = \text{SFNO}_\phi(\mathbf{x}_0, \hat{\mathbf{x}}_h, \mathbf{f}_0, 1 | \xi)$. In practice, cold sampling applies a correction term to this estimate. The prior forecast of \mathbf{x}_h can now be refined with $\hat{\mathbf{x}}_h = \text{SFNO}_\theta(\hat{\mathbf{x}}_1, \mathbf{f}_1, 1)$. The alternation between forecasting and interpolation continues until SFNO_ϕ predicts $\hat{\mathbf{x}}_{h-1}$ and the forecaster network performs a last refinement forecast of time step h , conditioned on the time $j = h-1$ and interpolated sample $\hat{\mathbf{x}}_{h-1}$. After this final forecast of \mathbf{x}_h , the process is repeated autoregressively, starting with \mathbf{x}_h as the new initial condition. This slightly simplified sampling process is illustrated in Figure 3 and fully described in Algorithm 2. Repeating this sampling process multiple times using the same initial conditions will lead to an ensemble of samples, thanks to the interpolator network being stochastic.

SFNO time-conditioning. To use SFNO as described above, it is necessary to implement time-conditioning modules that allow the interpolator and forecaster networks to be conditioned on the time i and j , respectively, given that the original SFNO architecture does not support this. We follow the same approach taken by standard diffusion models [11], which consists of transforming the time condition into a vector of sine/cosine Fourier features at 32 frequencies with base period 16, then pass them through a 2-layer MLP to obtain 128-dimensional time encodings that are mapped by a linear layer into the learnable scale and offset parameters. We scale and shift the neural representations of every SFNO block directly following the normalization layer and preceding the application of the SFNO spectral filter, as shown in Figure 4.

SFNO inference stochasticity. A stochastic interpolator network, made explicitly through the random variable ξ above, was shown to be a key design choice in the original DYffusion framework. However, to the best of our knowledge, the SFNO model has been only used for deterministic modeling. We overcome this issue through MC dropout [15], i.e. enabling dropout modules [53] at inference time. Following the original SFNO implementation (of training-time-only dropout), we propose to use a dropout module inside the MLP of each SFNO block. In addition, we enable stochastic depth [23]—also known as drop path—at inference time at a rate of 0.1. Stochastic depth randomly skips a whole SFNO block. When this happens the whole block reduces to the identity function, since only the residual connection is enabled. To the best of our knowledge, this has not been explored before as a source of inference stochasticity.

5 Experiments

5.1 Dataset and Experimental Setup

To compare our proposed method against ACE [58], we use the same dataset, training and evaluation setup. The dataset consists of 11 distinct 10-year-long simulations from the state-of-the-art global atmospheric model FV3GFS [63], saved every 6 hours. The forcings consist of annually repeating climatological sea surface temperature (1982-2012 average) and incoming solar radiation. Greenhouse gas and aerosol concentrations are kept fixed. The data was regridded conservatively from the cubed-sphere geometry of FV3GFS to a 1° Gaussian grid, and filtered with a spherical harmonic transform round-trip to remove artifacts in the high latitudes. We train on 100 years of simulated data from FV3GFS, and evaluate the models on how well they can emulate a distinct 10-year-long validation simulation (i.e. $H = 14600 = 10 \times 365 \times 4$). The 11 simulations form an initial-condition ensemble, where each simulation is independent of the other—after some discarded spinup time—due to the chaoticity of the atmosphere [26]. For more details, see Appendix B.

5.2 Baselines

We compare with the following baselines for climate projection.

- ACE [58] applied the SFNO architecture to the FV3GFS dataset described above.

- **ACE-STO**: We re-train ACE but use MC dropout, in the same way how it is applied in SFNO $_{\phi}$ for our method, to generate stochastic predictions.
- **DYffusion** [48]: We train DYffusion using the original UNet-based architecture as its interpolator and forecaster neural networks.
- **Reference** [63]: physics-based FV3GFS climate model simulations. We use the ten training simulations to create a 10-member reference ensemble that we use to more robustly estimate the ‘noise floor’ introduced in [58] and to compare the variability of the reference ensemble with sample simulations from our method. Note that this reference is *not* appropriate for weather forecasts given that it is initialized from different initial conditions.

It is worth noting that ACE also compared their results against a physics-based baseline called C48, which corresponds to running FV3GFS at half the original spatial resolution. This makes C48 around $8\times$ less computationally costly to run compared to the reference simulations but was shown to underperform ACE, which our method is shown to outperform in the experiments below.

For ACE, we directly use the pre-trained model from the original paper. ACE was trained on a next-step forecasting objective based on a MSE loss. For ACE-STO, we re-train ACE from scratch with the only difference being that we use a dropout rate of 10% for the MLP in the SFNO architecture. We use the same dropout rate for the interpolator model, SFNO $_{\phi}$, in our method. For both DYffusion and our approach, we choose $h = 6$. That is, these models are trained to forecast up to 36 hours into the future. We use the same training and sampling procedures for both, the only difference being the underlying neural architectures.

Runtime analysis. In Table 1, we report the computational complexity in terms of the number of neural function evaluations (NFEs) needed to predict h time steps, and the wall clock runtime for simulating one complete validation trajectory of 10 years. For our method, NFEs is not $3h$ because in the first and last iteration we do not need to actually run line 8 and lines 7 & 8 in Algorithm 2, respectively. Our runtime analysis confirms that the computational overhead at inference time for using our method, is less than $3\times$ as much as for a deterministic next-step forecasting model such as SFNO or ACE. This enables our method to provide significant $25\times$ speed-ups and associated energy savings over using the emulated physics-based model, FV3GFS.

Table 1: Computational complexity of the different deep learning methods in terms of: 1) the number of neural function evaluations (NFEs) needed to predict h time steps. and 2) Total inference runtime (simulating 10 years), including the time needed to compute metrics (in hours:minutes). N refers to the number of diffusion steps which usually ranges between 20 to 1000.

Method	NFE	Runtime
ACE / SFNO	h	01:08
Standard diffusion	Nh	N/A
Ours	$3(h - 1)$	02:56
Physics-based FV3GFS	N/A	78:04
FV3GFS ($2\times$ coarser)	N/A	45:38

All models were trained on A6000 GPUs using distributed training on 2 up to 8 GPUs, ensuring that the effective batch size (number of GPUs \times number of gradient accumulation steps \times data loader batch size) remains the same (see Table 5). For a fair inference runtime comparison measuring the wall clock time needed to simulate 10 years (i.e. one full validation rollout), we run all deep-learning baselines on one A100 GPU. We also include the runtime for the physics-based FV3GFS climate model which was run on 96 cores of AMD EPYC 7H12 processors. The deep learning methods are not only much faster, but also much more energy-efficient than FV3GFS.

For illustrative purposes, we also report the complexity of a standard autoregressive diffusion model [20, 29, 41] approach in terms of the number of neural function evaluations (NFEs) needed to predict h time steps, totaling to Nh where N is the number of sampling steps required to reverse the diffusion process. N usually ranges from between 20 to 1000. This makes the use of such an approach unattractive for climate emulation since the resulting inference runtime would not offer as significant speed-ups over the physics-based reference model.

5.3 Climate Biases

Metrics. The most crucial quality of an ML-based climate model is its ability to reproduce the climatology of the emulated reference system, i.e. the long-term average (“time-mean”) of weather




Figure 5: Global maps of the 10-year time-mean biases of a single sample from the reference noise floor simulation, our model, and the ACE baseline for the total water path field. Each subplot reports the global mean RMSE and bias of the respective bias map. Our model reproduces biases of similar location and magnitude to the reference noise floor, suggesting they are mainly due to internal climate variability rather than model bias, while the baseline exhibits larger climate biases.

states. The time-mean of the validation simulation is defined as $\frac{1}{H} \sum_{t=1}^H \mathbf{x}_t$. The time-mean for each model is defined as $\frac{1}{H} \sum_{t=1}^H \hat{\mathbf{x}}_t$, where $\hat{\mathbf{x}}_t$ is the model’s prediction for time step t . These two quantities are then compared against each other using the bias, i.e. prediction - target, and root mean squared error (RMSE) as key metrics of interest for analyzing climate biases. For the probabilistic methods, i.e. ours, DYffusion, and ACE-STO, we generate simulation ensembles by sampling from the model multiple times using the same initial conditions. Unless specified otherwise, all ensemble results are based on $M = 25$ ensemble members. We evaluate these ensembles by computing the RMSE on the ensemble-mean prediction $\frac{1}{MH} \sum_{m=1}^M \sum_{t=1}^H \hat{\mathbf{x}}_{t,m}$ as well as on member-wise time-means $\frac{1}{H} \sum_{t=1}^H \hat{\mathbf{x}}_{t,m}$, where m indexes the ensemble member dimension of the predictions. Standard deviations are computed over the respective member-wise errors. The corresponding “optimal noise floor” for the ML-based emulators is estimated by comparing the validation simulation with the 10-member reference ensemble. All metrics are area-weighted according to the size of the grid cell.

Quantitative analysis. Our method and all baselines consistently produce stable long-term climate simulations without diverging. In Figure 2, we compare the RMSE of the time-means of the reference, our method, and all baselines.

Our method significantly reduces climate biases compared to baseline methods across most fields, with errors often closer to the reference simulation’s noise floor than to the next best baseline. The performance of ACE is notably degraded when made stochastic through MC dropout. Similarly, a direct application of DYffusion fails to accurately reproduce long-term climate statistics. Both these baselines are unable to outperform or even match the scores of the deterministic ACE baseline. Only our proposed careful integration of these two paradigms leads to a skillful climate model emulator: On average, our method’s time-mean RMSEs are only 49.36% higher than the noise floor, which is less than half the average RMSE (110.47%) of the next best method, ACE.

Our method can be effectively ensembled to further reduce climate biases to 29.28% on average across all variables. The ensemble-mean prediction (light shade) consistently improves time-mean RMSEs across all fields compared to using a single member’s prediction (dark shade in Fig. 2). In contrast, reducing climate biases through ensemble-averaging is challenging for ACE and ineffective for DYffusion and ACE-STO. More detailed results across additional fields are presented in Figure 6.

In Appendix D.2.3 and Figure 10, we show that ensembling our method not only excels in absolute terms but that it also reduces errors more dramatically than the probabilistic baselines as a function of the ensemble size relative to single-member performance. That is, our method generates more effective ensemble climate simulations. Still, there is some gap compared to the reference ensemble for some variables such as total water path.

Qualitative analysis. In Figure 5 we show the corresponding global maps of the time-mean biases for the total water path (TWP) field. Our model reproduces small biases of remarkably similar location and magnitude to the “perfect-model” reference simulation, with spatial pattern RMSEs of approximately 1% of the global-time-mean TWP. The perfect-model bias is due to unforced random decadal variability in the mean climate of the reference model - each 10-year period has randomly different weather, leading to a slight difference in 10-year time-mean averages across this weather. The reference bias is due to comparing one such decade simulated with the reference model with other simulated decades; its spatial pattern depends strongly on which decade is used for computing the

reference model climatology. That our model (trained on 100 years of output) reproduces this pattern suggests that it emulates the long-term (e.g. century-long) time-mean statistics of the reference model even more accurately than a 10-year-mean RMSE can reliably resolve. On the other hand, the baseline ACE model exhibits somewhat larger climate biases, indicative of an actual, albeit small model deficiency that is already evident with a single 10-year estimate of climatology.

Climate variability. Above, we have verified that sampling 10-year-long trajectories from our model produces encouragingly low ensemble mean and member-wise time-mean biases. An important feature of climate is its natural variability on time scales of years, decades, or even centuries even when external forcings (e.g. sunlight or greenhouse gas concentrations) remain unchanged. For instance, multi-decadal periods of relative drought have stressed many past human civilizations. The present simulations are more constrained than natural climate variability because they employ a repeating cycle of sea-surface temperature and thus do not allow for feedbacks between the atmosphere, ocean, vegetation, and cryosphere. Nevertheless, an important quality of an ML emulator of the global atmosphere suitable for climate studies is that it simulates a similar level of low-frequency climate variability as the reference model.

Table 2: Global area-weighted mean of the spread of an ensemble of 10-yr time-mean’s for surface pressure, total water path, air temperature, zonal wind, and meridional wind (the last three at the near-surface level). The climate variability of our method is consistent with the reference model.

Model	p_s	TWP	T_7	u_7	v_7
Reference	19.961	0.199	0.090	0.142	0.110
Ours	23.517	0.214	0.094	0.167	0.121

Here, we verify that our time-mean ensemble passes this challenging test, measured using the intra-ensemble variability of time-mean averages of a few important climate statistics simulated by ensembles consisting of 25 samples of our model vs. the ten reference simulations. We measure this variability by computing the standard deviation across the ensemble of time-means for each latitude-longitude grid point and then average it over all grid points. In Table 2 we show that the resulting global mean variability of the ensemble of time-means of our method is within 10-20% of those of the reference simulations for all tabulated variables (and other predicted fields). In Appendix D.2.2 we show that the corresponding global maps of the time-mean variability reveal similar spatial patterns. That is, our method generates ensemble climate simulations with decadal variability consistent with the underlying climate model.

6 Conclusion

We introduced an efficient diffusion model-based approach, complemented by a spherical-aware neural architecture for fully data-driven ensemble climate simulations. Our model generates stable, 10-year-long probabilistic predictions, consisting of 14600 time steps, with lower climate biases than relevant deterministic and probabilistic baselines. Our method’s climate biases nearly reach a gold standard for successful climate model emulation, marking a significant leap towards efficient, data-driven climate simulations that can help society better understand the Earth and adapt to a changing climate.

Limitations. To achieve real-world impact, the dataset will need to be expanded so that ML emulators can be evaluated (and trained) on climate change scenarios/simulations. This will require using time-varying climate change forcings such as greenhouse gas and aerosol concentrations. Thanks to the use of a state-of-the-art atmospheric model, FV3GFS, such data can be generated. Additionally, we only considered emulating the atmosphere, but to achieve a full Earth System Model (ESM) we also need to emulate (or couple to a physics-based model of) other components such as ocean, land, sea-ice, etc. Additionally, we have ignored output-only variables such as precipitation. Training separate prediction heads for these variables or extending the DYffusion framework to support them is an interesting direction for future work. It is important to stress that while our method is more than $25\times$ faster than the reference physics-based climate model, it is still slower than deterministic models such as ACE. Physics-based models typically generate ensembles by perturbing both their inputs as well as internal model parameters. In this work, we focus on the latter point of model uncertainty by emulating a climate model with a diffusion model-based approach. Examining initial condition uncertainty is complementary to this approach and is left for future work.

Acknowledgements

This work was supported in part by the U.S. Army Research Office under Army-ECASE award W911NF-23-1-0231, the U.S. Department Of Energy, Office of Science, IARPA HAYSTAC Program, CDC-RFA-FT-23-0069, DARPA AIE FoundSci, NSF Grants #2205093, #2146343, and #2134274.

S.R.C. acknowledges generous support from a summer internship and subsequent collaboration with the Allen Institute for AI (AI2), which is primarily funded by the estate of Paul G. Allen. We thank Zihao Zhou, Gideon Dresdner, and Peter Eckmann for providing helpful feedback on our work.

References

- [1] Troy Arcomano, Istvan Szunyogh, Alexander Wikner, Brian R Hunt, and Edward Ott. A hybrid atmospheric model incorporating machine learning can capture dynamical processes not captured by its physics-based component. *Geophysical Research Letters*, 50(8), April 2023. doi:[10.1029/2022gl102649](https://doi.org/10.1029/2022gl102649).
- [2] Arpit Bansal, Eitan Borgnia, Hong-Min Chu, Jie S. Li, Hamid Kazemi, Furong Huang, Micah Goldblum, Jonas Geiping, and Tom Goldstein. Cold diffusion: Inverting arbitrary image transforms without noise. *Advances in Neural Information Processing Systems*, 2023. doi:[10.48550/arxiv.2208.09392](https://doi.org/10.48550/arxiv.2208.09392).
- [3] Seth Basetti, Brian Hutchinson, Claudia Tebaldi, and Ben Kravitz. DiffESM: Conditional emulation of earth system models with diffusion models. *ICLR Workshop on Tackling Climate Change with Machine Learning*, 2023. doi:[10.48550/arxiv.2304.11699](https://doi.org/10.48550/arxiv.2304.11699).
- [4] Kaifeng Bi, Lingxi Xie, Hengheng Zhang, Xin Chen, Xiaotao Gu, and Qi Tian. Accurate medium-range global weather forecasting with 3D neural networks. *Nature*, 619(7970):533–538, 2023. doi:[10.1038/s41586-023-06185-3](https://doi.org/10.1038/s41586-023-06185-3).
- [5] Boris Bonev, Thorsten Kurth, Christian Hundt, Jaideep Pathak, Maximilian Baust, Karthik Kashinath, and Anima Anandkumar. Spherical fourier neural operators: Learning stable dynamics on the sphere. *International Conference on Machine Learning*, 2023. doi:[10.48550/arxiv.2306.03838](https://doi.org/10.48550/arxiv.2306.03838).
- [6] Noah D. Brenowitz, Yair Cohen, Jaideep Pathak, Ankur Mahesh, Boris Bonev, Thorsten Kurth, Dale R. Durran, Peter Harrington, and Michael S. Pritchard. A practical probabilistic benchmark for AI weather models. *arXiv*, 2024. doi:[10.48550/arxiv.2401.15305](https://doi.org/10.48550/arxiv.2401.15305).
- [7] Kang Chen, Tao Han, Junchao Gong, Lei Bai, Fenghua Ling, Jing-Jia Luo, Xi Chen, Leiming Ma, Tianning Zhang, Rui Su, et al. FengWu: Pushing the skillful global medium-range weather forecast beyond 10 days lead. *arXiv*, 2023. doi:[10.48550/arxiv.2304.02948](https://doi.org/10.48550/arxiv.2304.02948).
- [8] Lei Chen, Xiaohui Zhong, Feng Zhang, Yuan Cheng, Yinghui Xu, Yuan Qi, and Hao Li. FuXi: a cascade machine learning forecasting system for 15-day global weather forecast. *npj Climate and Atmospheric Science*, 6(1), November 2023. ISSN 2397-3722. doi:[10.1038/s41612-023-00512-1](https://doi.org/10.1038/s41612-023-00512-1).
- [9] Kai-Yuan Cheng, Lucas Harris, Christopher Bretherton, Timothy M. Merlis, Maximilien Bolot, Linjiong Zhou, Alex Kaltenbaugh, Spencer Clark, and Stephan Fueglistaler. Impact of warmer sea surface temperature on the global pattern of intense convection: Insights from a global storm resolving model. *Geophysical Research Letters*, 49(16), 2022. doi:[10.1029/2022gl099796](https://doi.org/10.1029/2022gl099796).
- [10] Emmanuel de Bézenac, Syama Sundar Rangapuram, Konstantinos Benidis, Michael Bohlke-Schneider, Richard Kurle, Lorenzo Stella, Hilaf Hasson, Patrick Gallinari, and Tim Januschowski. Normalizing kalman filters for multivariate time series analysis. *Advances in Neural Information Processing Systems*, 2020.
- [11] Prafulla Dhariwal and Alexander Quinn Nichol. Diffusion models beat GANs on image synthesis. *Advances in Neural Information Processing Systems*, 2021. doi:[10.48550/arxiv.2105.05233](https://doi.org/10.48550/arxiv.2105.05233).

- [12] J.R. Driscoll and D.M. Healy. Computing fourier transforms and convolutions on the 2-sphere. *Advances in Applied Mathematics*, 15:202–250, 6 1994. ISSN 01968858. doi:10.1006/aama.1994.1008.
- [13] J. K. Fletcher, C. S. Bretherton, H. Xiao, R. Sun, and J. Han. Improving subtropical boundary layer cloudiness in the 2011 NCEP GFS. *Geoscientific Model Development*, 7(5):2107–2120, 2014. doi:10.5194/gmd-7-2107-2014.
- [14] V. Fortin, M. Abaza, F. Anctil, and R. Turcotte. Why should ensemble spread match the rmse of the ensemble mean? *Journal of Hydrometeorology*, 15(4):1708 – 1713, 2014. doi:https://doi.org/10.1175/JHM-D-14-0008.1.
- [15] Yarin Gal and Zoubin Ghahramani. Dropout as a bayesian approximation: Representing model uncertainty in deep learning. *International Conference on Machine Learning*, 2016. doi:10.48550/arxiv.1506.02142.
- [16] Tilmann Gneiting and Matthias Katzfuss. Probabilistic forecasting. *Annual Review of Statistics and Its Application*, 1(1):125–151, January 2014. ISSN 2326-831X. doi:10.1146/annurev-statistics-062713-085831.
- [17] Haiwen Guan, Troy Arcomano, Ashesh Chattopadhyay, and Romit Maulik. Lucie: A lightweight uncoupled climate emulator with long-term stability and physical consistency for O(1000)-member ensembles. *arXiv*, 2024. doi:10.48550/arxiv.2405.16297.
- [18] William Harvey, Saeid Naderiparizi, Vaden Masrani, Christian Weilbach, and Frank Wood. Flexible diffusion modeling of long videos. *Advances in Neural Information Processing Systems*, 2022. doi:10.48550/arxiv.2205.11495.
- [19] Ali Hatamizadeh, Jiaming Song, Guilin Liu, Jan Kautz, and Arash Vahdat. DiffiT: Diffusion vision transformers for image generation. *arXiv*, 2023. doi:10.48550/arxiv.2312.02139.
- [20] Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*, 2020. doi:10.48550/arxiv.2006.11239.
- [21] Jonathan Ho, William Chan, Chitwan Saharia, Jay Whang, Ruiqi Gao, Alexey Gritsenko, Diederik P. Kingma, Ben Poole, Mohammad Norouzi, David J. Fleet, and Tim Salimans. Imagen video: High definition video generation with diffusion models. *arXiv*, 2022. doi:10.48550/arxiv.2210.02303.
- [22] Jonathan Ho, Tim Salimans, Alexey Gritsenko, William Chan, Mohammad Norouzi, and David J Fleet. Video diffusion models. *Advances in Neural Information Processing Systems*, 2022. doi:10.48550/arxiv.2204.03458.
- [23] Gao Huang, Yu Sun, Zhuang Liu, Daniel Sedra, and Kilian Q. Weinberger. Deep networks with stochastic depth. In *Computer Vision – ECCV 2016*, page 646–661. Springer International Publishing, 2016. ISBN 9783319464930. doi:10.1007/978-3-319-46493-0_39.
- [24] Allan Jabri, David Fleet, and Ting Chen. Scalable adaptive computation for iterative generation. *International Conference on Machine Learning*, 2023. doi:10.48550/arxiv.2212.11972.
- [25] Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. *Advances in Neural Information Processing Systems*, 2022. doi:10.48550/arxiv.2206.00364.
- [26] J. E. Kay, C. Deser, A. Phillips, A. Mai, C. Hannay, G. Strand, J. M. Arblaster, S. C. Bates, G. Danabasoglu, J. Edwards, M. Holland, P. Kushner, J.-F. Lamarque, D. Lawrence, K. Lindsay, A. Middleton, E. Munoz, R. Neale, K. Oleson, L. Polvani, and M. Vertenstein. The community earth system model (cesm) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society*, 96(8):1333 – 1349, 2015. doi:10.1175/BAMS-D-13-00255.1.
- [27] Ryan Keisler. Forecasting global weather with graph neural networks. *arXiv*, 2022. doi:10.48550/arxiv.2202.07575.

- [28] Dmitrii Kochkov, Janni Yuval, Ian Langmore, Peter Norgaard, Jamie Smith, Griffin Mooers, James Lottes, Stephan Rasp, Peter Düben, Milan Klöwer, et al. Neural general circulation models. *arXiv*, 2023. doi:[10.48550/arxiv.2311.07222](https://doi.org/10.48550/arxiv.2311.07222).
- [29] Georg Kohl, Li-Wei Chen, and Nils Thuerey. Benchmarking autoregressive conditional diffusion models for turbulent flow simulation. *arXiv*, 2023. doi:[10.48550/arxiv.2309.01745](https://doi.org/10.48550/arxiv.2309.01745).
- [30] Anna Kwa, Spencer K. Clark, Brian Henn, Noah D. Brenowitz, Jeremy McGibbon, Oliver Watt-Meyer, W. Andre Perkins, Lucas Harris, and Christopher S. Bretherton. Machine-learned climate model corrections from a global storm-resolving model: Performance across the annual cycle. *Journal of Advances in Modeling Earth Systems*, 15(5), 2023. doi:[10.1029/2022ms003400](https://doi.org/10.1029/2022ms003400).
- [31] Remi Lam, Alvaro Sanchez-Gonzalez, Matthew Willson, Peter Wirsberger, Meire Fortunato, Ferran Alet, Suman Ravuri, Timo Ewalds, Zach Eaton-Rosen, Weihua Hu, et al. Learning skillful medium-range global weather forecasting. *Science*, 382(6677):1416–1421, 2023. ISSN 1095-9203. doi:[10.1126/science.adi2336](https://doi.org/10.1126/science.adi2336).
- [32] Zongyi Li, Nikola Borislavov Kovachki, Kamyar Azizzadenesheli, Burigede liu, Kaushik Bhattacharya, Andrew Stuart, and Anima Anandkumar. Fourier neural operator for parametric partial differential equations. *International Conference on Learning Representations*, 2021.
- [33] Phillip Lippe, Bastiaan S. Veeling, Paris Perdikaris, Richard E Turner, and Johannes Brandstetter. PDE-Refiner: Achieving accurate long rollouts with temporal neural pde solvers. *Advances in Neural Information Processing Systems*, 2023. doi:[10.48550/arxiv.2308.05732](https://doi.org/10.48550/arxiv.2308.05732).
- [34] Morteza Mardani, Noah Brenowitz, Yair Cohen, Jaideep Pathak, Chieh-Yu Chen, Cheng-Chin Liu, Arash Vahdat, Karthik Kashinath, Jan Kautz, and Mike Pritchard. Residual diffusion modeling for km-scale atmospheric downscaling. *arXiv*, 2023. doi:[10.48550/arxiv.2309.15214](https://doi.org/10.48550/arxiv.2309.15214).
- [35] James E. Matheson and Robert L. Winkler. Scoring rules for continuous probability distributions. *Management Science*, 22(10):1087–1096, 1976.
- [36] Tung Nguyen, Johannes Brandstetter, Ashish Kapoor, Jayesh K Gupta, and Aditya Grover. ClimaX: A foundation model for weather and climate. *International Conference on Machine Learning*, 2023. doi:[10.48550/arxiv.2301.10343](https://doi.org/10.48550/arxiv.2301.10343).
- [37] Tung Nguyen, Rohan Shah, Hritik Bansal, Troy Arcomano, Sandeep Madireddy, Romit Maulik, Veerabhadra Kotamarthi, Ian Foster, and Aditya Grover. Scaling transformer neural networks for skillful and reliable medium-range weather forecasting. *arXiv*, 2023. doi:[10.48550/arxiv.2312.03876](https://doi.org/10.48550/arxiv.2312.03876).
- [38] Brian C. O’Neill, Claudia Tebaldi, Detlef P. van Vuuren, Veronika Eyring, Pierre Friedlingstein, George Hurtt, Reto Knutti, Elmar Kriegler, Jean-Francois Lamarque, Jason Lowe, Gerald A. Meehl, Richard Moss, Keywan Riahi, and Benjamin M. Sanderson. The scenario model intercomparison project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*, 9(9): 3461–3482, September 2016. ISSN 1991-9603. doi:[10.5194/gmd-9-3461-2016](https://doi.org/10.5194/gmd-9-3461-2016).
- [39] Jaideep Pathak, Shashank Subramanian, Peter Harrington, Sanjeev Raja, Ashesh Chattopadhyay, Morteza Mardani, Thorsten Kurth, David Hall, Zongyi Li, Kamyar Azizzadenesheli, Pedram Hassanzadeh, Karthik Kashinath, and Animashree Anandkumar. FourCastNet: A global data-driven high-resolution weather model using adaptive fourier neural operators. *arXiv*, 2022. doi:[10.48550/arxiv.2202.11214](https://doi.org/10.48550/arxiv.2202.11214).
- [40] William Peebles and Saining Xie. Scalable diffusion models with transformers. In *IEEE/CVF International Conference on Computer Vision (ICCV)*. IEEE, October 2023. doi:[10.1109/iccv51070.2023.00387](https://doi.org/10.1109/iccv51070.2023.00387).
- [41] Ilan Price, Alvaro Sanchez-Gonzalez, Ferran Alet, Timo Ewalds, Andrew El-Kadi, Jacklynn Stott, Shakir Mohamed, Peter Battaglia, Remi Lam, and Matthew Willson. Gen-Cast: Diffusion-based ensemble forecasting for medium-range weather. *arXiv*, 2023. doi:[10.48550/arxiv.2312.15796](https://doi.org/10.48550/arxiv.2312.15796).

- [42] Stephan Rasp, Michael S. Pritchard, and Pierre Gentine. Deep learning to represent subgrid processes in climate models. *Proceedings of the National Academy of Sciences*, 115(39): 9684–9689, September 2018. ISSN 1091-6490. doi:[10.1073/pnas.1810286115](https://doi.org/10.1073/pnas.1810286115).
- [43] Stephan Rasp, Stephan Hoyer, Alexander Merose, Ian Langmore, Peter Battaglia, Tyler Russell, Alvaro Sanchez-Gonzalez, Vivian Yang, Rob Carver, Shreya Agrawal, Matthew Chantry, Zied Ben Bouallegue, Peter Dueben, Carla Bromberg, Jared Sisk, Luke Barrington, Aaron Bell, and Fei Sha. WeatherBench 2: A benchmark for the next generation of data-driven global weather models. *Journal of Advances in Modeling Earth Systems*, 16(6), June 2024. ISSN 1942-2466. doi:[10.1029/2023ms004019](https://doi.org/10.1029/2023ms004019).
- [44] Kashif Rasul, Calvin Seward, Ingmar Schuster, and Roland Vollgraf. Autoregressive denoising diffusion models for multivariate probabilistic time series forecasting. *International Conference on Machine Learning*, 2021. doi:[10.48550/arxiv.2101.12072](https://doi.org/10.48550/arxiv.2101.12072).
- [45] M. J. Rodwell and T. N. Palmer. Using numerical weather prediction to assess climate models. *Quarterly Journal of the Royal Meteorological Society*, 133(622):129–146, 2007. doi:<https://doi.org/10.1002/qj.23>.
- [46] Olaf Ronneberger, Philipp Fischer, and Thomas Brox. U-Net: Convolutional networks for biomedical image segmentation. *International Conference on Medical image computing and computer-assisted intervention*, pages 234–241, 2015. doi:[10.1007/978-3-319-24574-4_28](https://doi.org/10.1007/978-3-319-24574-4_28).
- [47] Salva Rühling Cachay, Venkatesh Ramesh, Jason N. S. Cole, Howard Barker, and David Rolnick. ClimART: A benchmark dataset for emulating atmospheric radiative transfer in weather and climate models. *Advances in Neural Information Processing Systems Track on Datasets and Benchmarks*, 2021. doi:[10.48550/arxiv.2111.14671](https://doi.org/10.48550/arxiv.2111.14671).
- [48] Salva Rühling Cachay, Bo Zhao, Hailey Joren, and Rose Yu. DYffusion: A dynamics-informed diffusion model for spatiotemporal forecasting. *Advances in Neural Information Processing Systems*, 2023. doi:[10.48550/arxiv.2306.01984](https://doi.org/10.48550/arxiv.2306.01984).
- [49] Uriel Singer, Adam Polyak, Thomas Hayes, Xi Yin, Jie An, Songyang Zhang, Qiyuan Hu, Harry Yang, Oron Ashual, Oran Gafni, Devi Parikh, Sonal Gupta, and Yaniv Taigman. Make-a-video: Text-to-video generation without text-video data. *arXiv*, 2022. doi:[10.48550/arxiv.2209.14792](https://doi.org/10.48550/arxiv.2209.14792).
- [50] Jascha Sohl-Dickstein, Eric Weiss, Niru Maheswaranathan, and Surya Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. *International Conference on Machine Learning*, 2015. doi:[10.48550/arxiv.1503.03585](https://doi.org/10.48550/arxiv.1503.03585).
- [51] Yang Song and Stefano Ermon. Generative modeling by estimating gradients of the data distribution. *Advances in Neural Information Processing Systems*, 32, 2019. doi:[10.48550/arxiv.1907.05600](https://doi.org/10.48550/arxiv.1907.05600).
- [52] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. *International Conference on Learning Representations*, 2020. doi:[10.48550/arxiv.2011.13456](https://doi.org/10.48550/arxiv.2011.13456).
- [53] Nitish Srivastava, Geoffrey Hinton, Alex Krizhevsky, Ilya Sutskever, and Ruslan Salakhutdinov. Dropout: A simple way to prevent neural networks from overfitting. *Journal of Machine Learning Research*, 15(56):1929–1958, 2014.
- [54] Vikram Voleti, Alexia Jolicoeur-Martineau, and Christopher Pal. MCVD: Masked conditional video diffusion for prediction, generation, and interpolation. *Advances in Neural Information Processing Systems*, 2022. doi:[10.48550/arxiv.2205.09853](https://doi.org/10.48550/arxiv.2205.09853).
- [55] Zhong Yi Wan, Ricardo Baptista, Anudhyan Boral, Yi-Fan Chen, John Anderson, Fei Sha, and Leonardo Zepeda-Nunez. Debias coarsely, sample conditionally: Statistical downscaling through optimal transport and probabilistic diffusion models. *Advances in Neural Information Processing Systems*, 2023. doi:[10.48550/arxiv.2305.15618](https://doi.org/10.48550/arxiv.2305.15618).
- [56] D. Watson-Parris. Machine learning for weather and climate are worlds apart. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 379(2194):20200098, 2021. doi:[10.1098/rsta.2020.0098](https://doi.org/10.1098/rsta.2020.0098).

- [57] D. Watson-Parris, Y. Rao, D. Olivié, Ø. Seland, P. Nowack, G. Camps-Valls, P. Stier, S. Bouabid, M. Dewey, E. Fons, J. Gonzalez, P. Harder, K. Jeggle, J. Lenhardt, P. Manshausen, M. Novitasari, L. Ricard, and C. Roesch. ClimateBench v1.0: A benchmark for data-driven climate projections. *Journal of Advances in Modeling Earth Systems*, 14(10), October 2022. doi:[10.1029/2021ms002954](https://doi.org/10.1029/2021ms002954).
- [58] Oliver Watt-Meyer, Gideon Dresdner, Jeremy McGibbon, Spencer K Clark, James Duncan, Brian Henn, Matthew Peters, Noah D Brenowitz, Karthik Kashinath, Mike Pritchard, Boris Bonev, and Christopher Bretherton. ACE: A fast, skillful learned global atmospheric model for climate prediction. *NeurIPS 2023 Workshop on Tackling Climate Change with Machine Learning*, 2023. doi:[10.48550/arxiv.2310.02074](https://doi.org/10.48550/arxiv.2310.02074).
- [59] Jonathan A. Weyn, Dale R. Durran, Rich Caruana, and Nathaniel Cresswell-Clay. Sub-seasonal forecasting with a large ensemble of deep-learning weather prediction models. *Journal of Advances in Modeling Earth Systems*, 13(7), July 2021. ISSN 1942-2466. doi:[10.1029/2021ms002502](https://doi.org/10.1029/2021ms002502).
- [60] Ruihan Yang, Prakhar Srivastava, and Stephan Mandt. Diffusion probabilistic modeling for video generation. *Entropy*, 25(10):1469, October 2023. ISSN 1099-4300. doi:[10.3390/e25101469](https://doi.org/10.3390/e25101469).
- [61] Sungduk Yu, Walter Hannah, Liran Peng, Jerry Lin, Mohamed Aziz Bhouri, Ritwik Gupta, Björn Lütjens, Justus Christopher Will, Gunnar Behrens, Julius Busecke, Nora Loose, Charles I Stern, Tom Beucler, Bryce Harrop, Benjamin R Hillman, Andrea Jenney, Savannah Ferretti, Nana Liu, Anima Anandkumar, Noah D Brenowitz, Veronika Eyring, Nicholas Geneva, Pierre Gentine, Stephan Mandt, Jaideep Pathak, Akshay Subramaniam, Carl Vondrick, Rose Yu, Laure Zanna, Tian Zheng, Ryan Abernathey, Fiaz Ahmed, David C Bader, Pierre Baldi, Elizabeth Barnes, Christopher Bretherton, Peter Caldwell, Wayne Chuang, Yilun Han, YU HUANG, Fernando Iglesias-Suarez, Sanket Jantre, Karthik Kashinath, Marat Khairoutdinov, Thorsten Kurth, Nicholas Lutsko, Po-Lun Ma, Griffin Mooers, J. David Neelin, David Randall, Sara Shamekh, Mark A Taylor, Nathan Urban, Janni Yuval, Guang Zhang, and Michael Pritchard. ClimSim: A large multi-scale dataset for hybrid physics-ML climate emulation. *Advances in Neural Information Processing Systems Track on Datasets and Benchmarks*, 2023.
- [62] Janni Yuval and Paul A. O’Gorman. Stable machine-learning parameterization of subgrid processes for climate modeling at a range of resolutions. *Nature Communications*, 11(1), July 2020. ISSN 2041-1723. doi:[10.1038/s41467-020-17142-3](https://doi.org/10.1038/s41467-020-17142-3).
- [63] Linjiong Zhou, Shian-Jiann Lin, Jan-Huey Chen, Lucas M. Harris, Xi Chen, and Shannon L. Rees. Toward convective-scale prediction within the next generation global prediction system. *Bulletin of the American Meteorological Society*, 100(7):1225 – 1243, 2019. doi:<https://doi.org/10.1175/BAMS-D-17-0246.1>.

Appendix

A Broader Impact

The goal of this work is to advance the application of machine learning to climate modeling, specifically for generating fast and cheap ML-based climate simulations. This could significantly democratize climate modeling, improve scientific understanding of the earth system, and enhance decision- and policy-making in a changing climate. However, to realize this goal, the reliability and limitations of such ML models will need to be much better understood.

B Dataset

In the subsections below, we elaborate on the dataset and variables that we use, including background information on FV3GFS and how it was configured in order to generate the training and validation data. Any listed data preprocessing steps below, are also described in appendix A from [58]. The final training and validation data can be downloaded from Google Cloud Storage following the instructions of the ACE paper at <https://github.com/ai2cm/ace>. The data are licensed under Creative Commons Attribution 4.0 International.

B.1 Input, output and forcing variables

Table 3: Input and output variables used in this work. The table was adapted based on Table 1 of [58]. The k subscript refers to a vertical layer index and ranges from 0 to 7 starting at the top of the atmosphere and increasing towards the surface. The two prognostic surface variables, T_s and p_s , do not have this additional vertical dimension. Each of their snapshots is a 2D latitude-longitude matrix. The Time column indicates whether a variable represents the value at a particular time step (“Snapshot”), the average across the 6-hour time step (“Mean”), or a quantity that does not depend on time (“Invariant”). “TOA” denotes “Top Of Atmosphere”, the climate model’s upper boundary.

Prognostic variables (input and output)			
Symbol	Description	Units	Time
T_k	Air temperature	K	Snapshot
q_k^T	Specific total water (vapor + condensates)	kg/kg	Snapshot
u_k	Wind speed in eastward direction	m/s	Snapshot
v_k	Wind speed in northward direction	m/s	Snapshot
T_s	Skin temperature of land or sea-ice	K	Snapshot
p_s	Atmospheric pressure at surface	Pa	Snapshot
Forcing variables (input-only)			
Symbol	Description	Units	Time
$DSWRF_{TOA}$	Downward shortwave radiative flux at TOA	W/m ²	Mean
T_s	Skin temperature of open ocean	K	Snapshot
Additional input-only variables			
Symbol	Description	Units	Time
z_s	Surface height of topography	m	Invariant
f_l	Land grid cell fraction	—	Invariant
f_o	Ocean grid cell fraction	—	Snapshot
f_{si}	Sea-ice grid cell fraction	—	Snapshot

The complete list of input, output, and forcing variables used in this work is given in Table 3. The only difference to the work from [58] is that we do not consider diagnostic (output only) variables. The forcings consist of annually repeating climatological sea surface temperature (1982-2012 average), T_s , and incoming solar radiation, $DSWRF_{TOA}$. Prescribed sea surface temperatures are simply “overwritten” on the skin temperature predictions of the ML models over all open ocean locations (when rolling out the ML-based simulation). The other forcing or input-only variables are added as

an additional channel dimension. For evaluation, we also consider the derived variable called total water path which is computed as $TWP = \frac{1}{g} \sum_k q_k^T dp_k$, i.e. as a function of surface pressure and the profile of specific total water.

B.2 Background on FV3GFS

Our dataset and physics-based baselines (including our “noise-floor” reference baseline) are based on simulations from a comprehensive global atmospheric model called Finite-Volume on a Cubed-Sphere Global Forecasting System (FV3GFS) [63]. It was developed by the National Oceanic and Atmospheric Administration (NOAA) Geophysical Fluid Dynamics Laboratory (GFDL)². A very similar model version is operationally used by the US National Centers for Environmental Prediction (NCEP) and the US weather forecasting service³. Its scalability to horizontal grid spacings as fine as 3 km [9] makes it an excellent candidate for generating training data for future ML-based climate model emulators, including out-of-distribution climate change simulations that may be necessary to train ML emulators on so that they can generalize.

B.3 FV3GFS configuration for data generation

In the following we summarize the reference data for this study, as also discussed in Section 2.1 of [58]. The training and validation data is generated by running an ensemble of 11 10-year (after discarding a 3-month spinup period) FV3GFS simulations on a C96 cubed-sphere grid (approximately 100 km horizontal grid spacing) with 63 vertical levels. The simulations are an initial-condition ensemble. That is, they are identical except for using different initial atmospheric states. Initial-condition ensembles are a popular tool in climate modeling [26]. Discarding a 3-month spinup period ensures that each simulation is independent of each other due to the chaoticity of the atmosphere⁴. Each simulation is forced by repeating annual cycles of sea-surface temperature and insolation. The temperature, humidity, two wind components at each grid point, and selected vertical fluxes at the surface and top of the atmosphere in each grid column are saved every six hours.

For ML training, the temperature, humidity, and two wind components are averaged along FV3GFS’s 63 levels to 8 vertical layers, and the data are interpolated to a latitude-longitude grid of 180×360 dimensions.

C Implementation details

All methods and baselines are conditioned on the forcings, f_t , by simple concatenation of the forcings with the remaining input variables across the channel dimension.

C.1 Training and inference pseudocode

In Algorithm 1 we provide the procedure used to train our proposed method.

²<https://www.gfdl.noaa.gov/fv3/>

³<https://www.weather.gov/news/fv3>

⁴E.g. see Kay et al. [26], who note that: “After initial condition memory is lost, which occurs within weeks in the atmosphere, each ensemble member evolves chaotically, affected by atmospheric circulation fluctuations characteristic of a random, stochastic process (e.g., Lorenz 1963; Deser et al. 2012b)”.

Algorithm 1 Spherical DYffusion, Training

- Input:** networks SFNO_ϕ , SFNO_θ , norm $\|\cdot\|$, horizon $h = 6$
- Stage 1:* Train interpolator network, SFNO_ϕ
1. Sample $i \sim \text{Uniform}(\{1, \dots, h-1\})$
 2. Sample $\mathbf{x}_t, \mathbf{x}_{t+i}, \mathbf{x}_{t+h} \sim \mathbb{R}^D$, and corresponding forcing \mathbf{f}_t
 3. Sample network stochasticity (dropout), ξ
 4. Optimize $\min_\phi \|\text{SFNO}_\phi(\mathbf{x}_t, \mathbf{x}_{t+h}, \mathbf{f}_t, i | \xi) - \mathbf{x}_{t+i}\|^2$
- Stage 2:* Train forecaster network, SFNO_θ
1. Freeze SFNO_ϕ and enable its inference stochasticity ξ
 2. Sample $j \sim \text{Uniform}(\{0, \dots, h-1\})$ and $\mathbf{x}_t, \mathbf{x}_{t+h} \sim \mathbb{R}^D$
 3. Retrieve corresponding forcings $\mathbf{f}_t, \mathbf{f}_{t+j}$
 4. $\hat{\mathbf{x}}_{t+j} \leftarrow \text{SFNO}_\phi(\mathbf{x}_t, \mathbf{x}_{t+h}, \mathbf{f}_t, j | \xi)$ # with $\hat{\mathbf{x}}_{t+j} := \mathbf{x}_t$ for $j = 0$
 5. Optimize $\min_\theta \|\text{SFNO}_\theta(\hat{\mathbf{x}}_{t+j}, \mathbf{f}_{t+j}, j) - \mathbf{x}_{t+h}\|^2$
-

In Algorithm 2 we provide the procedure used to sample with our proposed method.

Algorithm 2 Spherical DYffusion, Inference

- 1: **Input:** Initial conditions $\hat{\mathbf{x}}_0 := \mathbf{x}_0$, training and inference horizon h and $H = 14600$, forcings $\mathbf{f}_{0:H}$
 - 2: # Autoregressive loop:
 - 3: **for** $t = 0, h, 2 \cdot h, \dots, (\lceil H/h \rceil - 1) \cdot h$ **do**
 - 4: # Sampling loop for time steps $t + 1, \dots, t + h$:
 - 5: **for** $j = 0, 1, \dots, h - 1$ **do**
 - 6: $\hat{\mathbf{x}}_{t+h} \leftarrow \text{SFNO}_\theta(\hat{\mathbf{x}}_{t+j}, \mathbf{f}_{t+j}, j)$ # (Refine) forecast
 - 7: $\tilde{\mathbf{x}}_{t+j+1} \leftarrow \text{SFNO}_\phi(\hat{\mathbf{x}}_t, \hat{\mathbf{x}}_{t+h}, \mathbf{f}_t, j + 1 | \xi)$ # Interpolate
 - 8: $\hat{\mathbf{x}}_{t+j+1} = \tilde{\mathbf{x}}_{t+j+1} + \hat{\mathbf{x}}_{t+j} - \text{SFNO}_\phi(\hat{\mathbf{x}}_t, \hat{\mathbf{x}}_{t+h}, \mathbf{f}_t, j | \xi')$ # Cold sampling
 - 9: **end for**
 - 10: **end for**
 - 11: **Return:** $\hat{\mathbf{x}}_{1:H}$
-

C.2 Hyperparameters

Architectural hyperparameters. To fairly compare against the deterministic SFNO model from [58], we use exactly the same hyperparameters for training the interpolator and forecasting networks for our method, as described in Table 4⁵. For the stochastic version of ACE, ACE-STO, we re-train ACE from scratch with the only difference being that we use a dropout rate of 10% for the MLP in the SFNO architecture. We train the stochastic interpolator model, SFNO_ϕ , in our method using the same dropout rate. Both of these stochastic models are run using MC dropout (i.e. enabling the dropout layers at inference time).

Table 4: Table is directly taken from [58], and reports the SFNO hyperparameters used for ACE as well as the interpolator and forecasting networks of our method.

Name	Value
embed_dim	256
filter_type	linear
num_layers	8
operator_type	dhconv
scale_factor	1
spectral_layers	3

Optimization hyperparameters. We use the PyTorch Lightning framework to train our models. We train the interpolator networks for DYffusion and our method on the same relative L2 loss function used for the baseline from [58], and the corresponding forecaster networks on the L1 loss. The models that we train on our own, i.e. the

⁵Names correspond to the definition of the SphericalFourierNeuralOperatorNet class found here: <https://github.com/ai2cm/modulus/blob/94f62e1ce2083640829ec12d80b00619c40a47f8/modulus/models/sfno/sfnonet.py#L292>. All configuration options not listed here are set to the defaults at the linked code.

Table 5: Optimization hyperparameters. The effective batch size is calculated as batch size \times number of GPUs \times accumulate_grad_batches, and is ensured to be the same for all our trained models regardless of the number of GPUs used. Gradient accumulation refers to the number of forward passes that are aggregated before a gradient step is taken by the optimizer.

Name	Value
Optimizer	AdamW
Initial learning rate	4×10^{-4}
Weight decay	5×10^{-3}
Learning rate schedule	Cosine annealing (single cycle)
Number of epochs	60
Effective batch size	72
Exponential moving average decay rate	0.9999
Gradient clipping	0.5

interpolation and forecasting networks of DYffusion and our method are trained with mixed precision. Inference is always run at full precision. More optimization-related hyperparameters are discussed in Table 5.

D Additional results and figures

D.1 Additional ensemble metrics

Unless specified otherwise, all ensemble results are based on 25 ensemble members. We evaluate these ensembles with the ensemble-mean RMSE (eRMSE), the Continuous Ranked Probability Score (CRPS) [35], and the spread-skill ratio (spread / RMSE). The CRPS is a proper scoring rule and a popular metric in the probabilistic forecasting literature [16, 10, 44]. The spread-skill ratio is defined as the ratio of the square root of the ensemble variance to the corresponding ensemble-mean RMSE. It serves as a measure of the reliability of the ensemble, where values smaller than 1 indicate underdispersion (i.e. the probabilistic forecast is overconfident in its forecasts), and larger values overdispersion [14, 43]. All metrics are area-weighted according to the size of the grid cell. For the CRPS and RMSE lower is better, for spread / RMSE closer to 1 is better and indicates a well-dispersed ensemble.

D.2 Climate Biases

We quantitatively analyze the 10-year time mean biases of our model and the baselines in terms of the global mean errors in Figure 6. The time-mean prediction is the average over the 14,600 predicted snapshots during the 10 years. Our method significantly reduces climate biases over the baseline methods across most fields. Notably, the errors of our method are often closer to the noise floor of the reference simulation than to the next best baseline. We also show that our method can be effectively ensembled to further reduce climate biases. The ensemble-mean prediction reliably improves time-mean scores across all fields. Interestingly, the stochastic version of ACE, ACE-STO, significantly underperforms the deterministic version. Similarly, the direct application DYffusion fails to match the deterministic ACE baseline, even after ensemble-averaging. This shows that MC dropout and DYffusion alone are not the reason for the encouraging performance of our method, but rather the holistic integration of all components, including MC dropout in our SFNO-based interpolator network.

D.2.1 Zonal time-means

In this section, we analyze the absolute magnitudes of the simulated time-means by examining their zonal averages, where zonal refers to aggregation over the longitude dimension. We also report the standard deviation of the respective ensembles of time- and zonal-means for the reference and stochastic methods. We visualize these in Figure 7 and Figure 8. For several fields, including surface pressure, total water path (not shown), and near-surface temperature (top left subplot in Fig. 7), differences between the simulations are not visually noticeable, except for some biases near the




Figure 6: Same as Figure 2 but showing more fields for the RMSE of 10-year time-mean’s. Bars (left to right) show 1) the noise floor calculated from the pairwise differences of ten independent 10-year reference model simulations with respect to the validation simulation. In light shade we report the score computed using the mean over the ten reference simulations as ”prediction”, 2) the member-wise scores of an 25-member ensemble of our method in dark shade, and the corresponding ensemble mean score in light shade, 3) the score of the deterministic ACE baseline, 4) the member-wise scores of an 25-member ensemble of the DYffusion baseline in dark shade, and the corresponding ensemble mean score in light shade. The standard deviation error bar is computed over the set of pairwise (member-wise) time-mean RMSEs for the reference (Ours and DYffusion). ACE does not have a standard deviation since it is a deterministic model. Turning ACE stochastic through MC dropout (ACE-STO) degrades its performance. Our method significantly reduces climate biases over the baseline methods and can be effectively ensembled to reduce its climate biases further, approaching the theoretical lower limit imposed by the noise floor of the reference simulation.

poles by the baseline methods. Differences become more apparent as we analyze fields at higher altitude and wind fields. Generally, our method provides the closest emulation to the reference model. Although near-surface fields are the most relevant for society and decision-making, the clear biases of the baseline method at high-altitude levels might contribute to long-term biases, especially in longer simulations, due to the interactions of atmospheric dynamics across all levels. This observation may partly explain why our method achieves the lowest time-mean biases and RMSEs, as discussed in Appendix D.2.




Figure 7: Zonal means of the simulated 10-year time-mean climatologies for a representative subset of four temperature fields. Level 7 represents near-surface conditions, while Level 0 corresponds to the highest altitude. Our method generally provides the closest emulation to the reference data. The most notable biases in the emulations occur at Levels 2 and 0, indicating greater discrepancies at higher altitudes. Emulation challenges are also significant near the poles, including at near-surface levels, particularly for DYffusion.




Figure 8: Zonal-means of the simulated 10-year time-mean climatologies for a representative subset of four northward (meridional) wind fields. Our method generally provides the closest emulation to the reference data, except the level-0 polar latitudes.

D.2.2 Climate variability

In Fig. 9 we show the global maps corresponding to the global means of Table 2. Our method shows a consistent ensemble variability in terms of the simulated climate that also largely reflects the spatial patterns and magnitudes of the reference ensemble.

D.2.3 The impact of the ensemble size

In addition to the ensemble mean climate bias being generally significantly lower, in absolute terms than all the baselines, we find that our method’s ensemble performance also improves more dramatically than that of the probabilistic baselines as a function of the ensemble size relative to single-member performance. In Figure 10, we visualize the relative improvement in terms of ensemble-mean RMSE and CRPS, as well as the spread skill ratio as a function of the ensemble size. The relative boost in performance in terms of ensemble-mean RMSE and CRPS is generally considerably larger for our method than for the probabilistic baselines. The effectiveness of our method’s ensemble follows closely the reference 10-member ensemble for surface pressure, but reveals a significant gap for the total water path field. Our method produces better dispersed time-mean ensembles compared to the baselines as evidenced by its spread skill ratio scores, which are closer to one, but still manifest a certain amount of underdispersiveness.

D.3 Weather forecasting

While we focus on climate time scales in this work, climate is formed by the statistics of weather, so it is important to verify that our method also generates reasonable forecasts of the weather simulated by the reference model. In Figure 11, we analyze the medium-range forecasting skill of our method and the baselines for lead times up to two weeks. Interestingly, ACE and DYffusion show persistent biases for the surface pressure field that are clearly visible from the first few days of forecasts already but do not seem to reflect on the RMSEs at weather time scales. Such persistent biases, however, may be magnified over longer simulations and could explain why the baselines have problems reproducing accurate long-term climate statistics. In terms of RMSE, the deterministic model ACE generally has a slight edge over our method and DYffusion, especially on lead times of less than a week. After that, the ensembles of our method and DYffusion perform best in terms of ensemble-mean RMSE. As expected, the ensemble mean significantly reduces the RMSE compared to using a single sample from our method or DYffusion, especially at longer lead times. The probabilistic metrics, CRPS, and spread / RMSE ratio show that our method’s and DYffusion’s ensemble perform quite similarly, even though they are based on completely different ML architectures. Both ensembles tend to be underdispersed ($\text{Spread} / \text{RMSE} < 1$) on short time scales but quickly converge to a well-dispersed ensemble at longer lead times which persists for the whole 10-year climate simulations (not shown).




Figure 9: Global maps of the standard deviation of the 10-year time-mean of the reference ensemble and a 25-member ensemble of our method. The climate variability of our method is consistent with the reference model, and largely follows similar spatial patterns with adequate magnitudes. The global mean standard deviation is reported in Table 2.

D.4 Weather vs. climate performance

In Figure 12, we illustrate that weather performance does not correlate with the climate biases of the same model. We plot the average RMSE over the first 5 days of simulation (here, using the near-surface temperature field) against the 10-year time-mean RMSE of various fields, and do not observe any correlation between the two metrics. We have verified that this observation holds independently of the analyzed field. This is a little-discussed observation that has important implications for ML practitioners since it implies that optimizing for short-term forecasts alone – as is current practice for most ML-based weather forecasting models – may be suboptimal for attaining accurate climate simulations. Heuristically, optimizing weather skill ensures that a climate model takes a locally accurate path around the climate ‘attractor’, but it does not guarantee that small but systematic errors may not build up to distort that simulated attractor to have biased time-mean statistics. This observation has been documented for the case of physics-based climate models [13, 45].




Figure 10: We visualize the ensemble-mean RMSE, CRPS, and spread skill ratio as a function of the size, m , of the time-mean ensemble. For RMSE and CRPS we plot it relative to using a single time-mean prediction, i.e. $m = 1$, of the same model. Error bars are computed by repeatedly subsampling different combinations of m -member ensembles given a superset of $M = 25$ predictions.




Figure 11: Our method generates competitive probabilistic ensemble weather forecasts, a necessary but not sufficient prerequisite for achieving good climate simulations. We evaluate a 25-member ensemble as well as a single forecast for our method and DYffusion. We also report the deterministic forecasts of ACE.




Figure 12: We visualize the performances of DYffusion and Spherical DYffusion (in different marker colors) at multiple checkpoint epochs and for multiple generated samples. We plot the 10-year time-mean RMSE of three example fields versus the time-step-wise near-surface temperature RMSE averaged out over the first 20 forecasts (5 days). The 5-day weather forecast performance shows no correlation with the long-term climate biases (indeed, there seems to exist an inverse correlation). This has important implications for practitioners, implying that optimizing for short-term forecasts alone – as is current practice for most ML-based weather forecasting models – may be suboptimal for attaining accurate climate simulations. We have verified that the behavior shown above holds for fields other than near-surface temperature too (not shown).