

Replica bound for Ising spin glass models in one dimension

Manaka Okuyama¹ and Masayuki Ohzeki^{1,2,3}

¹*Graduate School of Information Sciences,*

Tohoku University, Sendai 980-8579, Japan

²*Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan and*

³*Sigma-i Co., Ltd., Tokyo 108-0075, Japan*

(Dated: June 24, 2024)

Abstract

The interpolation method is a powerful tool for rigorous analysis of mean-field spin glass models, both with and without dilution. In this study, we show that the interpolation method can be applied to Ising spin glass models in one dimension, such as a one-dimensional chain and a two-leg ladder. In one dimension, the replica symmetric (RS) cavity method is naturally expected to be rigorous for Ising spin glass models. Using the interpolation method, we rigorously prove that the RS cavity method provides lower bounds on the quenched free energies of Ising spin glass models in one dimension at any finite temperature in the thermodynamic limit.

I. INTRODUCTION

Rigorous analysis of mean-field spin glass models has made remarkable progress through the interpolation method [1]. The full-step replica symmetry breaking solution [2] has been proven to be an exact solution for the free energy of the Sherrington-Kirkpatrick model [3–5]. The interpolation method is also a powerful tool for diluted mean-field spin glass models, demonstrating that the replica symmetric (RS) and one-step replica symmetry breaking cavity methods [6] provide lower bounds on the free energies of these models [7–9]. Efforts to obtain inverse bounds for diluted mean-field spin glass models are ongoing [10–17].

Unfortunately, the interpolation method does not work well in finite dimensions. In one dimension, the Ising spin glass models have been analyzed using the replicated transfer matrix method [18–22]. Notably, a recent study [22] has proved that the RS cavity method is rigorous for the Ising spin glass model on a one-dimensional chain with free boundary condition (by computing the n -replicated partition function for any real number n and then rigorously taking the limit $n \rightarrow 0$). However, to the best of our knowledge, the interpolation method has not been applied in one dimension.

In the present study, we used the interpolation method to analyze the Ising spin glass models in one dimension, such as a one-dimensional chain and a two-leg ladder. The key element is the use of dilution variables introduced in the analysis of diluted mean-field models [7]. However, our interpolation method employs a slightly different parameter setup compared to that of a previous study [7]. We rigorously prove that the RS cavity method provides lower bounds on the free energies of the Ising spin glass models in one dimension.

The rest of this paper is organized as follows. In Sec. II, we define the Ising spin glass models in one dimension and state the main results. In Sec. III, we use the interpolation method to prove that the RS cavity method provides lower bounds on the free energies of these models. Our results are discussed in Sec. IV. For completeness, the derivation of the RS cavity formula is provided in Appendices A and B.

II. DEFINITIONS AND RESULTS

A. One-dimensional chain

Let us consider the Ising spin glass model on a one-dimensional chain with periodic boundary conditions:

$$H_N^{\text{chain}} = - \sum_{i=1}^N J_i \sigma_i \sigma_{i+1} - \sum_{i=1}^N H_i \sigma_i, \quad (1)$$

where $\sigma_i = \pm 1$, J_i are i.i.d. random variables following any symmetric distribution $P(J_i) = P(-J_i)$, and H_i are i.i.d. random variables following any distribution (the probability distribution of H_i is allowed to present no randomness, for example, a uniform magnetic field). We assume that these distributions are regular enough to guarantee that all expressions below are well-defined.

The partition function is defined as

$$Z_N^{\text{chain}} = \text{Tr}(e^{-\beta H_N^{\text{chain}}}), \quad (2)$$

where $\text{Tr}(\dots)$ denotes the summation over all spin variables, and β is the inverse temperature. The thermal average $\langle \dots \rangle$ is given by

$$\langle g(\sigma) \rangle = \frac{\text{Tr}(g(\sigma) e^{-\beta H_N^{\text{chain}}})}{Z_N^{\text{chain}}}. \quad (3)$$

The quenched free energy is

$$f_N^{\text{chain}} = -\frac{1}{\beta N} \mathbb{E} \left[\log Z_N^{\text{chain}} \right], \quad (4)$$

where $\mathbb{E}[\dots]$ denotes the expectation with respect to all random variables. The thermodynamic limit of the quenched free energy is defined as

$$f^{\text{chain}} = \lim_{N \rightarrow \infty} f_N^{\text{chain}}. \quad (5)$$

The existence of the thermodynamic limit of the quenched free energy is proved through the conventional approach [23, 24].

By applying the RS cavity method [25] to the Ising spin glass model on a one-dimensional chain, we obtain the RS cavity formula (see Appendix A for the derivation)

$$f_{\text{RS}}^{\text{chain}} = -\frac{1}{\beta} \mathbb{E} \left[\log \frac{2 \cosh \beta J \cosh \beta(\hat{h} + H)}{\cosh \beta \hat{h}} \right], \quad (6)$$

where J is an independent copy of the coupling J_i , H is an independent copy of the magnetic field H_i , and the cavity field \hat{h} satisfies

$$\tanh \beta \hat{h} \stackrel{d}{=} \tanh \beta J \tanh \beta (H + \hat{h}'), \quad (7)$$

where the equality holds in distribution and \hat{h}' is an independent copy of the cavity field \hat{h} . This equation can be rewritten by the distribution $P(\hat{h})$ as

$$P(\hat{h}) = \mathbb{E}_{J,H} \left[\int d\hat{h}' P(\hat{h}') \delta(\tanh \beta \hat{h} - \tanh \beta J \tanh \beta (H + \hat{h}')) \right], \quad (8)$$

where $\mathbb{E}_{J,H}[\dots]$ denotes the expectation with respect to J and H .

Then, our first main result is as follows:

Theorem 1. *In the thermodynamic limit, the quenched free energy of the Ising spin glass model on a one-dimensional chain is bounded below by the RS cavity formula*

$$f_{\text{RS}}^{\text{chain}} \leq f^{\text{chain}}. \quad (9)$$

B. Two-leg ladder

Next, we consider the Ising spin glass model on a two-leg ladder with periodic boundary conditions:

$$H_N^{\text{ladder}} = - \sum_{i=1}^N L_i \sigma_{i,1} \sigma_{i,2} - \sum_{i=1}^N (K_{i,1} \sigma_{i,1} \sigma_{i+1,1} + K_{i,2} \sigma_{i,2} \sigma_{i+1,2}), \quad (10)$$

where $\sigma_{i,1} = \pm 1$, $\sigma_{i,2} = \pm 1$, $K_{i,1}$ are i.i.d. random variables following any symmetric distribution $P(K_{i,1}) = P(-K_{i,1})$, $K_{i,2}$ are i.i.d. random variables following any symmetric distribution $P(K_{i,2}) = P(-K_{i,2})$, and L_i are i.i.d. random variables following any distribution. The probability distribution of L_i is allowed to present no randomness.

The partition function is given by

$$Z_N^{\text{ladder}} = \text{Tr} \left(e^{-\beta H_N^{\text{ladder}}} \right). \quad (11)$$

The quenched free energy and its thermodynamic limit are defined as

$$f_N^{\text{ladder}} = -\frac{1}{\beta N} \mathbb{E} \left[\log Z_N^{\text{ladder}} \right], \quad (12)$$

$$f^{\text{ladder}} = \lim_{N \rightarrow \infty} f_N^{\text{ladder}}. \quad (13)$$

By applying the RS cavity method to the Ising spin glass model on a two-leg ladder, we obtain the RS cavity formula (see Appendix B for the derivation)

$$f_{\text{RS}}^{\text{ladder}} = -\frac{1}{\beta} \mathbb{E} \left[\log \frac{4 \cosh \beta K_1 \cosh \beta K_2 \cosh \beta(L + \hat{l})}{\cosh \beta \hat{l}} \right], \quad (14)$$

where K_1 is an independent copy of the coupling $K_{i,1}$, K_2 is an independent copy of the coupling $K_{i,2}$, L is an independent copy of the coupling L_i , and the cavity coupling \hat{l} satisfies

$$\tanh \beta \hat{l} \stackrel{d}{=} \tanh \beta K_1 \tanh \beta K_2 \tanh \beta(L + \hat{l}'), \quad (15)$$

where the equality holds in distribution, and \hat{l}' is an independent copy of the cavity coupling \hat{l} . This equation can be rewritten using the distribution $P(\hat{l})$ as

$$P(\hat{l}) = \mathbb{E}_{L, K_1, K_2} \left[\int d\hat{l}' P(\hat{l}') \delta(\tanh \beta \hat{l} - \tanh \beta K_1 \tanh \beta K_2 \tanh \beta(L + \hat{l}')) \right], \quad (16)$$

where $\mathbb{E}_{L, K_1, K_2} [\dots]$ denotes the expectation with respect to L , K_1 , and K_2 .

The second main result is as follows:

Theorem 2. *The thermodynamic limit of the quenched free energy of the Ising spin glass model on the two-leg ladder is bounded by the RS cavity formula*

$$f_{\text{RS}}^{\text{ladder}} \leq f^{\text{ladder}}. \quad (17)$$

III. PROOF

A. Proof of Theorem 1

We define the interpolating Hamiltonian as

$$H_N^{\text{chain}}(t) = -\sum_{i=1}^N \left(J_i M_i(t) \sigma_i \sigma_{i+1} + (1 - M_i(t)) (\hat{h}_{i,1} \sigma_i + \hat{h}_{i,2} \sigma_{i+1}) \right) - \sum_{i=1}^N H_i \sigma_i, \quad (18)$$

$$H_N^{\text{chain}}(1) = H_N^{\text{chain}}, \quad (19)$$

$$H_N^{\text{chain}}(0) = -\sum_{i=1}^N (H_i + \hat{h}_{i,1} + \hat{h}_{i-1,2}) \sigma_i, \quad (20)$$

where $M_i(t)$ are i.i.d. Bernoulli variables with average t , and $\hat{h}_{i,1}$ and $\hat{h}_{i,2}$ are independent copies of the cavity field \hat{h} (see Eq. (7)). We define the interpolating quenched free energy as

$$f_N^{\text{chain}}(t) = -\frac{1}{\beta N} \mathbb{E} \left[\log Z_N^{\text{chain}}(t) \right]. \quad (21)$$

At $t = 1$, we recover the free energy of the Ising spin glass model on the one-dimensional chain

$$f_N^{\text{chain}}(1) = f_N^{\text{chain}}. \quad (22)$$

For any function $g(x)$, the following identity holds:

$$\frac{d}{dt} \mathbb{E}[g(M_i(t))] = g(1) - g(0). \quad (23)$$

Using this identity, we have

$$\begin{aligned} \frac{d}{dt} f_N^{\text{chain}}(t) &= -\frac{1}{\beta N} \sum_{i=1}^N \mathbb{E} \left[\log \frac{\langle e^{\beta J_i \sigma_i \sigma_{i+1}} \rangle_{H_{\setminus i}^{\text{chain}}(t)}}{\langle e^{\beta(\hat{h}_{i,1} \sigma_i + \hat{h}_{i,2} \sigma_{i+1})} \rangle_{H_{\setminus i}^{\text{chain}}(t)}} \right] \\ &= -\frac{1}{\beta} \mathbb{E} \left[\log \frac{\langle e^{\beta J_i \sigma_i \sigma_{i+1}} \rangle_{H_{\setminus i}^{\text{chain}}(t)}}{\langle e^{\beta(\hat{h}_{i,1} \sigma_i + \hat{h}_{i,2} \sigma_{i+1})} \rangle_{H_{\setminus i}^{\text{chain}}(t)}} \right], \end{aligned} \quad (24)$$

where we have applied the translation invariance property, and $\langle \cdots \rangle_{H_{\setminus i}^{\text{chain}}(t)}$ denotes the thermal average with respect to $H_{\setminus i}^{\text{chain}}(t)$. Here, $H_{\setminus i}^{\text{chain}}(t)$ is defined as

$$H_{\setminus i}^{\text{chain}}(t) = H_N^{\text{chain}}(t) + J_i M_i(t) \sigma_i \sigma_{i+1} + (1 - M_i(t)) (\hat{h}_{i,1} \sigma_i + \hat{h}_{i,2} \sigma_{i+1}). \quad (25)$$

Note that $H_{\setminus i}^{\text{chain}}(t)$ does not contain J_i , $\hat{h}_{i,1}$, and $\hat{h}_{i,2}$. The lattice structure of $H_{\setminus i}^{\text{chain}}(t)$ is that of a one-dimensional chain with free boundary conditions instead of periodic boundary conditions.

By using the identity $\exp(\beta J_i \sigma_i \sigma_{i+1}) = \cosh \beta J_i (1 + \sigma_i \sigma_{i+1} \tanh \beta J_i)$ and Eq. (7), we obtain

$$f_N^{\text{chain}}(0) + \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} = f_{\text{RS}}^{\text{chain}}. \quad (26)$$

This equation implies that the information at $t = 0$ coincides with the result of the RS cavity method. This structure is also common in the analysis of mean-field models using the interpolation method [3, 7]. By using the fundamental theorem of calculus and Eq. (26), we have

$$\begin{aligned} f_N^{\text{chain}}(1) &= f_N^{\text{chain}}(0) + \int_0^1 dt \frac{d}{dt} f_N^{\text{chain}}(t) \\ &= f_{\text{RS}}^{\text{chain}} + \int_0^1 dt \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right). \end{aligned} \quad (27)$$

Thus, to obtain Theorem 1, it is sufficient to prove

$$\int_0^1 dt \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right) \geq O(N^{-1}). \quad (28)$$

We obtain

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right) \\
&= \mathbb{E} \left[\log \left(1 + \tanh \beta J_i \langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right) \right] - \mathbb{E} \left[\log \left((1 + \tanh \beta \hat{h}_{i,1} \sigma_i) (1 + \tanh \beta \hat{h}_{i,2} \sigma_{i+1}) \right)_{H_{\hat{v}_i}^{\text{chain}}(t)} \right] \\
& \quad + \mathbb{E} \left[\log \left(1 + \tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \tanh \beta (H_{i+1} + \hat{h}_{i+1,1}) \right) \right]. \tag{29}
\end{aligned}$$

Here, we recall that, in the thermal average $\langle \cdots \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}$, sites i and $i+1$ are both ends of a one-dimensional chain with free boundary conditions. This fact implies that sites i and $i+1$ are connected with probability t^{N-1} , given that the existence probability of each interaction is t . In other words, the correlation function is decoupled,

$$\langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} = \langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \langle \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}, \tag{30}$$

with probability $1 - t^{N-1}$. Then, we have

$$\begin{aligned}
& -\mathbb{E} \left[\log \left((1 + \tanh \beta \hat{h}_{i,1} \sigma_i) (1 + \tanh \beta \hat{h}_{i,2} \sigma_{i+1}) \right)_{H_{\hat{v}_i}^{\text{chain}}(t)} \right] \\
&= -\mathbb{E} \left[\log \left(1 + \tanh \beta \hat{h}_{i,1} \langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right) \left(1 + \tanh \beta \hat{h}_{i,2} \langle \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right) \right] + \mathcal{O}(t^{N-1}) \\
&= -2\mathbb{E} \left[\log \left(1 + \tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right) \right] + \mathcal{O}(t^{N-1}), \tag{31}
\end{aligned}$$

where we have used Eq. (7) in the last equality. By substituting Eq. (31) into Eq. (29), we obtain

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right) \\
&= \mathbb{E} \left[\log \left(1 + \tanh \beta J_i \langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right) \right] - 2\mathbb{E} \left[\log \left(1 + \tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right) \right] \\
& \quad + \mathbb{E} \left[\log \left(1 + \tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \tanh \beta (H_{i+1} + \hat{h}_{i+1,1}) \right) \right] + \mathcal{O}(t^{N-1}). \tag{32}
\end{aligned}$$

Given that

$$|\tanh \beta J_i \langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}| < 1, \tag{33}$$

$$|\tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}| < 1, \tag{34}$$

$$|\tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \tanh \beta (H_{i+1} + \hat{h}_{i+1,1})| < 1, \tag{35}$$

we can expand the logarithm of the three terms as

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right) \\
&= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \mathbb{E} \left[\left(\tanh \beta J_i \langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right)^n - 2 \left(\tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)} \right)^n \right. \\
&\quad \left. + \left(\tanh \beta J_i \tanh \beta (H_i + \hat{h}_{i,1}) \tanh \beta (H_{i+1} + \hat{h}_{i+1,1}) \right)^n \right] + \mathcal{O}(t^{N-1}), \\
&= - \sum_{n=1}^{\infty} \frac{1}{2n} \mathbb{E} \left[\tanh^{2n} \beta J_i \right] \\
&\quad \left(\mathbb{E} \left[\langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}^{2n} \right] - 2 \mathbb{E} \left[\tanh^{2n} \beta (H_i + \hat{h}_{i,1}) \right] \mathbb{E} \left[\langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}^{2n} \right] + \mathbb{E} \left[\tanh^{2n} \beta (H_i + \hat{h}_{i,1}) \right]^2 \right) \\
&\quad + \mathcal{O}(t^{N-1}), \tag{36}
\end{aligned}$$

where we have used $\mathbb{E}[\tanh^{2n-1} \beta J_i] = 0$ for any positive integer n that follows from $P(J_i) = P(-J_i)$. Again, we use the fact that sites i and $i+1$ are connected with probability t^{N-1} , which implies

$$\mathbb{E} \left[\langle \sigma_i \sigma_{i+1} \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}^{2n} \right] = \mathbb{E} \left[\langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}^{2n} \right]^2 + \mathcal{O}(t^{N-1}). \tag{37}$$

By applying Eq. (37) to Eq. (36), we obtain

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right) \\
&= - \sum_{n=1}^{\infty} \frac{1}{2n} \mathbb{E} \left[\tanh^{2n} \beta J_i \right] \mathbb{E} \left[\langle \sigma_i \rangle_{H_{\hat{v}_i}^{\text{chain}}(t)}^{2n} - \tanh^{2n} \beta (H_i + \hat{h}_{i,1}) \right]^2 + \mathcal{O}(t^{N-1}) \\
&\leq \mathcal{O}(t^{N-1}). \tag{38}
\end{aligned}$$

Thus, we have

$$\begin{aligned}
f_N^{\text{chain}} &= f_N^{\text{chain}}(1) = f_{\text{RS}}^{\text{chain}} + \int_0^1 dt \left(\frac{d}{dt} f_N^{\text{chain}}(t) - \frac{d}{dt} f_N^{\text{chain}}(t) \Big|_{t=0} \right) \\
&\geq f_{\text{RS}}^{\text{chain}} + \int_0^1 dt \mathcal{O}(t^{N-1}) \\
&= f_{\text{RS}}^{\text{chain}} + \mathcal{O}(N^{-1}). \tag{39}
\end{aligned}$$

Finally, taking the thermodynamic limit, we obtain

$$f^{\text{chain}} \geq f_{\text{RS}}^{\text{chain}}, \tag{40}$$

which proves Theorem 1.

B. Proof of Theorem 2

We define the interpolating Hamiltonian as

$$H_N^{\text{ladder}}(t) = - \sum_{i=1}^N L_i \sigma_{i,1} \sigma_{i,2} - \sum_{i=1}^N M_i(t) (K_{i,1} \sigma_{i,1} \sigma_{i+1,1} + K_{i,2} \sigma_{i,2} \sigma_{i+1,2}),$$

$$- \sum_{i=1}^N (1 - M_i(t)) (\hat{l}_{i,1} \sigma_{i,1} \sigma_{i,2} + \hat{l}_{i,2} \sigma_{i+1,1} \sigma_{i+1,2}), \quad (41)$$

$$H_N^{\text{ladder}}(1) = H^{\text{ladder}}, \quad (42)$$

$$H_N^{\text{ladder}}(0) = - \sum_{i=1}^N (L_i + \hat{l}_{i,1} + \hat{l}_{i-1,2}) \sigma_{i,1} \sigma_{i,2}, \quad (43)$$

where $M_i(t)$ are i.i.d. Bernoulli variables with average t , and $\hat{l}_{i,1}$ and $\hat{l}_{i,2}$ are independent copies of the cavity coupling \hat{l} (see Eq. (15)). We define the interpolating quenched free energy as

$$f_N^{\text{ladder}}(t) = - \frac{1}{\beta N} \mathbb{E} \left[\log Z_N^{\text{ladder}}(t) \right]. \quad (44)$$

Note that

$$f_N^{\text{ladder}}(1) = f_N^{\text{ladder}}. \quad (45)$$

Then, we have

$$\begin{aligned} \frac{d}{dt} f_N^{\text{ladder}}(t) &= - \frac{1}{\beta N} \sum_{i=1}^N \mathbb{E} \left[\log \frac{\langle e^{\beta(K_{i,1} \sigma_{i,1} \sigma_{i+1,1} + K_{i,2} \sigma_{i,2} \sigma_{i+1,2})} \rangle_{H_{\setminus i}^{\text{ladder}}(t)}}{\langle e^{\beta(\hat{l}_{i,1} \sigma_{i,1} \sigma_{i,2} + \hat{l}_{i,2} \sigma_{i+1,1} \sigma_{i+1,2})} \rangle_{H_{\setminus i}^{\text{ladder}}(t)}} \right] \\ &= - \frac{1}{\beta} \mathbb{E} \left[\log \frac{\langle e^{\beta(K_{i,1} \sigma_{i,1} \sigma_{i+1,1} + K_{i,2} \sigma_{i,2} \sigma_{i+1,2})} \rangle_{H^{\text{ladder}}(t)}}{\langle e^{\beta(\hat{l}_{i,1} \sigma_{i,1} \sigma_{i,2} + \hat{l}_{i,2} \sigma_{i+1,1} \sigma_{i+1,2})} \rangle_{H^{\text{ladder}}(t)}} \right], \end{aligned} \quad (46)$$

where we have applied the translation invariance property, and $\langle \cdots \rangle_{H_{\setminus i}^{\text{ladder}}(t)}$ denotes the thermal average with respect to $H_{\setminus i}^{\text{ladder}}(t)$. Here, $H_{\setminus i}^{\text{ladder}}(t)$ is defined as

$$\begin{aligned} &H_{\setminus i}^{\text{ladder}}(t) \\ &= H_N^{\text{ladder}}(t) + M_i(t) (K_{i,1} \sigma_{i,1} \sigma_{i+1,1} + K_{i,2} \sigma_{i,2} \sigma_{i+1,2}) + (1 - M_i(t)) (\hat{l}_{i,1} \sigma_{i,1} \sigma_{i,2} + \hat{l}_{i,2} \sigma_{i+1,1} \sigma_{i+1,2}) \end{aligned} \quad (47)$$

Note that $H_{\setminus i}^{\text{ladder}}(t)$ does not contain $K_{i,1}$, $K_{i,2}$, $\hat{l}_{i,1}$, and $\hat{l}_{i,2}$. The lattice structure of $H_{\setminus i}^{\text{ladder}}(t)$ is a two-leg ladder with free boundary conditions.

After some calculations based on Eq. (15), we obtain

$$f_N^{\text{ladder}}(0) + \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} = f_{\text{RS}}^{\text{ladder}}. \quad (48)$$

Then,

$$f_N^{\text{ladder}}(1) = f_{\text{RS}}^{\text{ladder}} + \int_0^1 dt \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right). \quad (49)$$

Thus, to obtain Theorem 2, it is sufficient to prove

$$\lim_{N \rightarrow \infty} \int_0^1 dt \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right) \geq 0. \quad (50)$$

We obtain

$$\begin{aligned} & -\beta \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right) \\ = & \mathbb{E} \left[\log \langle (1 + \sigma_{i,1} \sigma_{i+1,1} \tanh \beta K_{i,1}) (1 + \sigma_{i,2} \sigma_{i+1,2} \tanh \beta K_{i,2}) \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right] \\ & - \mathbb{E} \left[\log \langle (1 + \tanh \beta \hat{l}_{i,1} \sigma_{i,1} \sigma_{i,2}) (1 + \tanh \beta \hat{l}_{i,2} \sigma_{i+1,1} \sigma_{i+1,2}) \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right] \\ & + \mathbb{E} \left[\log \left(1 + \tanh \beta K_{i,1} \tanh \beta K_{i,2} \tanh \beta (L_i + \hat{l}_{i,1}) \tanh \beta (L_{i+1} + \hat{l}_{i+1,1}) \right) \right]. \end{aligned} \quad (51)$$

The proof procedure from this point onwards is identical to that of Theorem 1, albeit with the need for a few additional calculations to expand the logarithm of the three terms in Eq. (51). In the thermal average $\langle \dots \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}$, spins $\sigma_{i,1} \sigma_{i,2}$ and $\sigma_{i+1,1} \sigma_{i+1,2}$ are located at both ends of the two-leg ladder with free boundary conditions, and the probability of each interaction's existence is determined by t . Thus, the correlation functions are decoupled,

$$\langle \sigma_{i,1} \sigma_{i,2} \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} = \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}, \quad (52)$$

$$\langle \sigma_{i,1} \sigma_{i+1,1} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} = \langle \sigma_{i,1} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \langle \sigma_{i+1,1} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} = 0, \quad (53)$$

$$\langle \sigma_{i,2} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} = \langle \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \langle \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} = 0, \quad (54)$$

with probability $1 - (1 - (1 - t)^2)^N$, where we have used $\langle \sigma_{i,1} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} = 0$ owing to spin-flip symmetry.

Then, we have

$$\begin{aligned} & \mathbb{E} \left[\log \langle (1 + \sigma_{i,1} \sigma_{i+1,1} \tanh \beta K_{i,1}) (1 + \sigma_{i,2} \sigma_{i+1,2} \tanh \beta K_{i,2}) \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right] \\ = & \mathbb{E} \left[\log \left(1 + \tanh \beta K_{i,1} \tanh \beta K_{i,2} \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right) \right] + \mathcal{O}((1 - (1 - t)^2)^N), \end{aligned} \quad (55)$$

and

$$\begin{aligned} & -\mathbb{E} \left[\log \langle (1 + \tanh \beta \hat{l}_{i,1} \sigma_{i,1} \sigma_{i,2}) (1 + \tanh \beta \hat{l}_{i,2} \sigma_{i+1,1} \sigma_{i+1,2}) \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right] \\ = & -\mathbb{E} \left[\log \left(1 + \tanh \beta \hat{l}_{i,1} \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right) (1 + \tanh \beta \hat{l}_{i,2} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}) \right] + \mathcal{O}((1 - (1 - t)^2)^N) \\ = & -2 \mathbb{E} \left[\log \left(1 + \tanh \beta K_i^1 \tanh \beta K_i^2 \tanh \beta (L_i + \hat{l}_{i,1}) \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right) \right] + \mathcal{O}((1 - (1 - t)^2)^N), \end{aligned} \quad (56)$$

where we have used Eq. (15) in the last equality. By substituting Eqs. (55) and (56) into Eq. (51), we obtain

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right) \\
&= \mathbb{E} \left[\log \left(1 + \tanh \beta K_{i,1} \tanh \beta K_{i,2} \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right) \right] \\
&\quad - 2 \mathbb{E} \left[\log \left(1 + \tanh \beta K_i^1 \tanh \beta K_i^2 \tanh \beta (L_i + \hat{l}_{i,1}) \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \right) \right] \\
&\quad + \mathbb{E} \left[\log \left(1 + \tanh \beta K_{i,1} \tanh \beta K_{i,2} \tanh \beta (L_i + \hat{l}_{i,1}) \tanh \beta (L_{i+1} + \hat{l}_{i+1,1}) \right) \right] \\
&\quad + \mathcal{O}((1 - (1 - t)^2)^N). \tag{57}
\end{aligned}$$

In addition, given that

$$|\tanh \beta K_{i,1} \tanh \beta K_{i,2} \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}| < 1, \tag{58}$$

$$|\tanh \beta K_i^1 \tanh \beta K_i^2 \tanh \beta (L_i + \hat{l}_{i,1}) \langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}| < 1, \tag{59}$$

$$|\tanh \beta K_{i,1} \tanh \beta K_{i,2} \tanh \beta (L_i + \hat{l}_{i,1}) \tanh \beta (L_{i+1} + \hat{l}_{i+1,1})| < 1, \tag{60}$$

we can expand the logarithm of the three terms as

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right) \\
&= - \sum_{n=1}^{\infty} \frac{1}{2n} \mathbb{E} \left[\tanh^{2n} \beta K_{i,1} \right] \mathbb{E} \left[\tanh^{2n} \beta K_{i,2} \right] \left(\mathbb{E} \left[\langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} \right] \right. \\
&\quad \left. - 2 \mathbb{E} \left[\tanh^{2n} \beta (L_i + \hat{l}_{i,1}) \right] \mathbb{E} \left[\langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} \right] + \mathbb{E} \left[\tanh^{2n} \beta (L_i + \hat{l}_{i,1}) \right]^2 \right) \\
&\quad + \mathcal{O}((1 - (1 - t)^2)^N), \tag{61}
\end{aligned}$$

where we have used $\mathbb{E} \left[\tanh^{2n-1} \beta K_{i,1} \right] = 0$ for any positive integer n that follows from $P(K_{i,1}) = P(-K_{i,1})$. Again, we use the fact that, in the thermal average $\langle \cdots \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}$, both ends of the two-leg ladder are connected with probability $\mathcal{O}((1 - (1 - t)^2)^N)$, which implies

$$\mathbb{E} \left[\langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} \langle \sigma_{i+1,1} \sigma_{i+1,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} \right] = \mathbb{E} \left[\langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} \right]^2 + \mathcal{O}((1 - (1 - t)^2)^N). \tag{62}$$

Then, we obtain

$$\begin{aligned}
& -\beta \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right) \\
&= - \sum_{n=1}^{\infty} \frac{1}{2n} \mathbb{E} \left[\tanh^{2n} \beta K_{i,1} \right] \mathbb{E} \left[\tanh^{2n} \beta K_{i,2} \right] \mathbb{E} \left[\langle \sigma_{i,1} \sigma_{i,2} \rangle_{H_{\hat{v}_i}^{\text{ladder}}(t)}^{2n} - \tanh^{2n} \beta (L_i + \hat{l}_{i,1}) \right]^2 \\
&\quad + \mathcal{O}((1 - (1 - t)^2)^N) \\
&\leq \mathcal{O}((1 - (1 - t)^2)^N). \tag{63}
\end{aligned}$$

Finally, by taking the thermodynamic limit, we obtain

$$\begin{aligned}
\lim_{N \rightarrow \infty} f_N^{\text{ladder}} &= f_{\text{RS}}^{\text{ladder}} + \lim_{N \rightarrow \infty} \int_0^1 dt \left(\frac{d}{dt} f_N^{\text{ladder}}(t) - \frac{d}{dt} f_N^{\text{ladder}}(t) \Big|_{t=0} \right) \\
&\geq f_{\text{RS}}^{\text{ladder}} + \lim_{N \rightarrow \infty} \int_0^1 dt \mathcal{O}((1 - (1 - t)^2)^N) \\
&= f_{\text{RS}}^{\text{ladder}},
\end{aligned} \tag{64}$$

which proves Theorem 2.

IV. DISCUSSIONS

Using the interpolation method, we rigorously proved that the RS cavity method provides lower bounds on the free energies of Ising spin glass models in one dimension, such as a one-dimensional chain and a two-leg ladder. Our proof relies strongly on the property that correlations at both ends are neglected in the thermal average with respect to the one-dimensional interpolating Hamiltonians (25) and (47). Thus, our method does not extend to more than two dimensions.

The interpolation method was effective in one dimension owing to the validity of Eqs. (26) and (48). We emphasize that the left-hand sides of Eqs. (26) and (48) are equivalent to Oguchi's approximate free energies [26] in the case of ferromagnetic interactions. It is known that Oguchi's approximate free energy coincides with the exact solution of the free energy for ferromagnetic systems in one dimension. This excellent property is closely related to the equality of Eqs. (26) and (48).

In one dimension, it is naturally expected that the RS cavity method is rigorous for Ising spin glass models. Indeed, on a one-dimensional chain, the replicated transfer matrix method allows for proving that the RS cavity formula is rigorous in the thermodynamic limit [22]. Thus, an important future direction is to prove the converse bound of the present study, especially for a two-leg ladder.

This study was supported by JSPS KAKENHI Grant Nos. 24K16973 and 23H01432. This study received financial support from the Public\Private R&D Investment Strategic Expansion Program (PRISM) and programs for Bridging the gap between R&D and the IDEal society (society 5.0) and Generating Economic and social value (BRIDGE) from Cabinet Office.

Appendix A: Derivation of RS cavity formula (6) on one-dimensional chain

Let us consider an Ising spin glass model on a one-dimensional chain with free boundary conditions:

$$H_{N,\text{free}}^{\text{chain}} = - \sum_{i=1}^{N-1} J_i \sigma_i \sigma_{i+1} - \sum_{i=1}^N H_i \sigma_i. \quad (\text{A1})$$

First, we assume that, when we take the summation over spin variables from σ_1 to σ_{N-1} , its contribution is represented by a cavity field \hat{g}_{N-1} and a constant factor A_{N-1} as follows (this assumption that the contribution from σ_1 to σ_{N-1} can be expressed in terms of a simple random field corresponds to the RS assumption):

$$\begin{aligned} \mathbb{E} \left[\log Z_{N,\text{free}}^{\text{chain}} \right] &= \mathbb{E} \left[A_{N-1} \log \text{Tr}_{\sigma_N} \left(e^{\beta(\hat{g}_{N-1} + H_N) \sigma_N} \right) \right] \\ &= \mathbb{E} \left[A_{N-1} \log 2 \cosh \beta(\hat{g}_{N-1} + H_N) \right], \end{aligned} \quad (\text{A2})$$

where Tr_{σ_N} is the summation over σ_N . By repeating the same assumption for a system with one more spin number, we obtain

$$\mathbb{E} \left[\log Z_{N+1,\text{free}}^{\text{chain}} \right] = \mathbb{E} \left[A_{N-1} \log \text{Tr}_{\sigma_N, \sigma_{N+1}} \left(e^{\beta(\hat{g}_{N-1} + H_N) \sigma_N + \beta J_N \sigma_N \sigma_{N+1} + \beta H_{N+1} \sigma_{N+1}} \right) \right], \quad (\text{A3})$$

where $\text{Tr}_{\sigma_N, \sigma_{N+1}}$ is the summation over σ_N and σ_{N+1} . Let us introduce the new cavity field \hat{g}_N , which satisfies

$$\tanh \beta \hat{g}_N \stackrel{d}{=} \tanh \beta J_N \tanh \beta(\hat{g}_{N-1} + H_N), \quad (\text{A4})$$

where the equality holds in distribution. Then, we can rewrite $\mathbb{E} \left[\log Z_{N+1,\text{free}}^{\text{chain}} \right]$ as

$$\mathbb{E} \left[\log Z_{N+1,\text{free}}^{\text{chain}} \right] = \mathbb{E} \left[A_{N-1} \log \frac{4 \cosh \beta(\hat{g}_{N-1} + H_N) \cosh \beta J_N \cosh \beta(\hat{g}_N + H_{N+1})}{\cosh \beta \hat{g}_N} \right]. \quad (\text{A5})$$

Thus, we have

$$\mathbb{E} \left[\log Z_{N+1,\text{free}}^{\text{chain}} \right] - \mathbb{E} \left[\log Z_{N,\text{free}}^{\text{chain}} \right] = \mathbb{E} \left[\log \frac{2 \cosh \beta J_N \cosh \beta(\hat{g}_N + H_{N+1})}{\cosh \beta \hat{g}_N} \right]. \quad (\text{A6})$$

By taking the thermodynamic limit $N \rightarrow \infty$, the probability distributions of \hat{g}_{N-1} and \hat{g}_N are expected to converge to the same one, implying

$$\lim_{N \rightarrow \infty} \left(\mathbb{E} \left[\log Z_{N+1,\text{free}}^{\text{chain}} \right] - \mathbb{E} \left[\log Z_{N,\text{free}}^{\text{chain}} \right] \right) = \mathbb{E} \left[\log \frac{2 \cosh \beta J \cosh \beta(\hat{g} + H)}{\cosh \beta \hat{g}} \right], \quad (\text{A7})$$

with

$$\tanh \beta \hat{g} \stackrel{d}{=} \tanh \beta J \tanh \beta (\hat{g}' + H), \quad (\text{A8})$$

where the equality holds in distribution, J is an independent copy of the coupling J_i , H is an independent copy of the magnetic field H_i , and \hat{g}' is an independent copy of \hat{g} .

Finally, because $\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{chain}} \right] - \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{chain}} \right]$ is considered the free energy density per spin, we assume the following relation (see Ref. [27] for a mathematically rigorous treatment):

$$\lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{chain}} \right] = \lim_{N \rightarrow \infty} \left(\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{chain}} \right] - \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{chain}} \right] \right). \quad (\text{A9})$$

In combination with Eqs. (A7) and (A8), we obtain the RS cavity formula (6).

Appendix B: Derivation of RS cavity formula (14) on two-leg ladder

Let us consider an Ising spin glass model on a two-leg ladder with free boundary conditions:

$$H_{N, \text{free}}^{\text{ladder}} = - \sum_{i=1}^N L_i \sigma_{i,1} \sigma_{i,2} - \sum_{i=1}^{N-1} (K_{i,1} \sigma_{i,1} \sigma_{i+1,1} + K_{i,2} \sigma_{i,2} \sigma_{i+1,2}). \quad (\text{B1})$$

First, we assume that, when we take the summation over spin variables from $\sigma_{1,1}$ and $\sigma_{1,2}$ to $\sigma_{N-1,1}$ and $\sigma_{N-1,2}$, its contribution is represented by a cavity coupling \hat{s}_{N-1} and a constant factor B_{N-1} as follows:

$$\begin{aligned} \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{ladder}} \right] &= \mathbb{E} \left[B_{N-1} \log \text{Tr}_{\sigma_N} \left(e^{\beta(\hat{s}_{N-1} + L_N) \sigma_{N,1} \sigma_{N,2}} \right) \right] \\ &= \mathbb{E} \left[B_{N-1} \log 4 \cosh \beta (\hat{s}_{N-1} + L_N) \right], \end{aligned} \quad (\text{B2})$$

where Tr_{σ_N} is the summation over $\sigma_{N,1}$ and $\sigma_{N,2}$. By repeating the same assumption for a system with one more spin number, we obtain

$$\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right] = \mathbb{E} \left[B_{N-1} \log \text{Tr}_{\sigma_N, \sigma_{N+1}} \left(e^{\beta(\hat{s}_{N-1} + L_N) \sigma_{N,1} \sigma_{N,2} + \beta(K_{N,1} \sigma_{N,1} \sigma_{N+1,1} + K_{N,2} \sigma_{N,2} \sigma_{N+1,2}) + \beta L_{N+1} \sigma_{N+1,1} \sigma_{N+1,2}} \right) \right], \quad (\text{B3})$$

where $\text{Tr}_{\sigma_N, \sigma_{N+1}}$ is the summation over $\sigma_{N,1}$, $\sigma_{N,2}$, $\sigma_{N+1,1}$, and $\sigma_{N+1,2}$. Let us introduce the new cavity coupling \hat{s}_N , which satisfies

$$\tanh \beta \hat{s}_N \stackrel{d}{=} \tanh \beta K_{N,1} \tanh \beta K_{N,2} \tanh \beta (\hat{s}_{N-1} + L_N), \quad (\text{B4})$$

where the equality holds in distribution. Then, we can rewrite $\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right]$ as

$$\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right] = \mathbb{E} \left[B_{N-1} \log \frac{16 \cosh \beta (\hat{s}_{N-1} + L_N) \cosh \beta K_{N,1} \cosh \beta K_{N,2} \cosh \beta (\hat{s}_N + L_{N+1})}{\cosh \beta \hat{s}_N} \right] \quad (\text{B5})$$

Thus, we have

$$\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right] - \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{ladder}} \right] = \mathbb{E} \left[\log \frac{4 \cosh \beta K_{N,1} \cosh \beta K_{N,2} \cosh \beta (\hat{s}_N + L_{N+1})}{\cosh \beta \hat{s}_N} \right]. \quad (\text{B6})$$

By taking the thermodynamic limit $N \rightarrow \infty$, the probability distributions of \hat{s}_{N-1} and \hat{s}_N are expected to converge to the same one. This implies

$$\lim_{N \rightarrow \infty} \left(\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right] - \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{ladder}} \right] \right) = \mathbb{E} \left[\log \frac{4 \cosh \beta K_1 \cosh \beta K_2 \cosh \beta (\hat{s} + L)}{\cosh \beta \hat{s}} \right], \quad (\text{B7})$$

with

$$\tanh \beta \hat{s} \stackrel{d}{=} \tanh \beta K_1 \tanh \beta K_2 \tanh \beta (\hat{s}' + L), \quad (\text{B8})$$

where the equality holds in distribution, K_1 is an independent copy of $K_{i,1}$, K_2 is an independent copy of $K_{i,2}$, L is an independent copy of L_i , and \hat{s}' is an independent copy of \hat{s} .

By contrast, given that $\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right] - \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{ladder}} \right]$ is considered the free energy density per spin, we assume the following relation:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{ladder}} \right] = \lim_{N \rightarrow \infty} \left(\mathbb{E} \left[\log Z_{N+1, \text{free}}^{\text{ladder}} \right] - \mathbb{E} \left[\log Z_{N, \text{free}}^{\text{ladder}} \right] \right). \quad (\text{B9})$$

Therefore, we obtain the RS cavity formula (14).

-
- [1] F. Guerra and F. L. Toninelli, The Thermodynamic Limit in Mean Field Spin Glass Models, *Commun. Math. Phys.* **230**, 71 (2002).
 - [2] G. Parisi, A sequence of approximated solutions to the S-K model for spin glasses, *J. Phys. A: Math. Gen.* **13**, L115 (1980).
 - [3] F. Guerra, Broken Replica Symmetry Bounds in the Mean Field Spin Glass Model, *Commun. Math. Phys.* **233**, 1 (2003).
 - [4] M. Talagrand, The Parisi formula, *Ann. Math.* **163**, 221 (2006).
 - [5] D. Panchenko, The Parisi formula for mixed p -spin models, *Ann. Probab.* **42**, 946 (2014).
 - [6] M. Mézard and G. Parisi, The Bethe lattice spin glass revisited, *Euro. Phys. J. B*, **233**, 217 (2001).

- [7] S. Franz and M. Leone, Replica bounds for optimization problems and diluted spin systems, *J. Stat. Phys.* **111**, 535 (2003).
- [8] S. Franz and M. Leone, and F. L. Toninelli, Replica bounds for diluted non-Poissonian spin systems, *J. Phys. A: Math. Gen.* **36**, 10967 (2003).
- [9] D. Panchenko and M. Talagrand, Bounds for diluted mean-fields spin glass models, *Probab. Theory Relat. Fields* **130**, 319 (2004).
- [10] F. Guerra and F. L. Toninelli, The High Temperature Region of the Viana-Bray Diluted Spin Glass Model, *J. Stat. Phys.* **115**, 531 (2004).
- [11] D. Panchenko, Spin glass models from the point of view of spin distributions, *Ann. Probab.* **41**, 1315 (2013).
- [12] D. Panchenko, Structure of 1-RSB asymptotic Gibbs measures in the diluted p-spin models, *J. Stat. Phys.* **155**, 1 (2014).
- [13] D. Panchenko, Hierarchical exchangeability of pure states in mean field spin glass models, *Probab. Theory Relat. Fields* **161**, 619 (2015).
- [14] J. Ding, A. Sly, and N. Sun, Maximum independent sets on random regular graphs, *Acta Math.* 217, 263 (2016).
- [15] M. Lelarge and M. Oulamara, Replica bounds by combinatorial interpolation for diluted spin systems, *J. Stat. Phys.* 173, 917 (2018).
- [16] A. Coja-Oghlan and W. Perkins, Spin systems on Bethe lattices, *Commun. Math. Phys.* **372**, 441 (2019).
- [17] V. Harangi, Improved Replica Bounds for the Independence Ratio of Random Regular Graphs, *J. Stat. Phys.* 190, 60 (2023).
- [18] B. Derrida, J. Vannimenus, and Y. Pomeau, Simple frustrated systems: chains, strips and squares, *J. Phys. C* **11**, 4749 (1978).
- [19] J. M. Luck, M. Funke, and T. M. Nieuwenhuizen, Low-temperature thermodynamics of random-field Ising chains: exact results, *J. Phys. A* **24**, 4155 (1991).
- [20] M. Weigt and R. Monasson, Replica structure of one-dimensional disordered Ising models, *Europhys. Lett.* **36**, 209 (1996).
- [21] A. J. Bray and M. A. Moore, Finite size effects in spin glass overlap functions, *J. Phys. A: Math. Gen.* **18**, L683 (1985).
- [22] C. Lucibello, F. Morone, and T. Rizzo, One-dimensional disordered Ising models by replica and cavity

- methods, Phys. Rev. E **90**, 012140 (2014).
- [23] P. A. Vuilleumat, Thermodynamics of quenched random spin systems, and application to the problem of phase transitions in magnetic (spin) glasses, J. Phys. A **10**, 1319 (1977).
- [24] L. A. Pastur and A. L. Figotin, Theory of disordered spin systems, Teor. Mat. Fiz. **35**, 193 (1978).
- [25] A. Montanari and M. Mézard, *Information, Physics and Computation* (Oxford Univ. Press, 2009).
- [26] A. Oguchi, Approximate Method for the Free Energy, Prog. Theor. Phys. **56**, 1442 (1976).
- [27] M. Aizenman, R. Sims, and S. L. Starr, Extended variational principle for the Sherrington-Kirkpatrick spin-glass model, Phys. Rev. B **68**, 214403 (2003).