

On p -adic Transference Theorem

Chi Zhang

*Key Laboratory of Mathematics Mechanization, NCMIS, Academy of Mathematics
and Systems Science, Chinese Academy of Sciences, Beijing 100190, People's
Republic of China*

and

*School of Mathematical Sciences, University of Chinese Academy of Sciences,
Beijing 100049, People's Republic of China*

zhangchi171@mails.ucas.ac.cn

Abstract

Dual lattice is an important concept of Euclidean lattices. In 2024, Deng gave the definition to the concept of the dual lattice of a p -adic lattice from the duality theory of locally compact abelian groups. He also proved some important properties of the dual lattice of p -adic lattices, which can be viewed as p -adic analogues of the famous Minkowski's first, second theorems and transference theorems for Euclidean lattices. However, he only proved the lower bound of the transference theorems for p -adic lattices. The upper bound is left as an open question. In this paper, we prove the upper bound of the transference theorems for p -adic lattices. We then prove that the dual basis of an orthogonal basis is also an orthogonal basis with respect to the maximum norm.

2010 Mathematics Subject Classification: Primary 11F85.

Key words and phrases: p -adic lattice, Local field, CVP, LVP, Transference Theorem.

1. Transference Theorem

In [2], Deng gave the definition to the concept of the dual lattice of a p -adic lattice and then proved some important properties of the dual lattice of p -adic lattices, including the lower bound of the transference theorems for p -adic lattices. He left the upper bound as an open question.

The lower bound exists even if the p -adic lattice is not of full rank. However, the upper bound does not exist if the p -adic lattice is not of full rank. Consider the following example.

Let $k = \mathbb{Q}_p$ or $k = \mathbb{F}_p((T))$. Let M be the maximum norm on k^m . Let the basis

of Λ be

$$B = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}.$$

Then the dual basis

$$D = \frac{1}{m} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}.$$

Hence

$$\tilde{\lambda}_1(\Lambda) \cdot \tilde{\lambda}_1(\Lambda^*) = \left| \frac{1}{m} \right|_p.$$

We can take $m = p^l$ for any large integer l .

Therefore, in the following subsections, we consider the p -adic lattice which is of full rank.

1.1. Transference Theorem for the Maximum Norm

Let $k = \mathbb{Q}_p$ or $k = \mathbb{F}_p((T))$. Let M be the maximum norm on k^m , i.e., for $v = (v_1, \dots, v_m) \in k^m$,

$$M(v) = \max_{1 \leq i \leq m} |v_i|_p.$$

We consider the upper bound for the transference theorem respect to norm M first. We need the following orthogonalization algorithm.

Algorithm (orthogonalization with orthogonal bases).

Input: an N -orthogonal basis e_1, \dots, e_n of k^n , a lattice $\mathcal{L} = \mathcal{L}(\alpha_1, \dots, \alpha_m)$ in k^n .

Output: an N -orthogonal basis of \mathcal{L} .

1. for $i = 1$ to m do:
2. rearrange $\alpha_i, \dots, \alpha_m$ such that $N(\alpha_i) = \max_{i \leq k \leq m} N(\alpha_k)$,
3. rearrange e_i, \dots, e_n such that $N(a_{ii}e_i) = \max_{i \leq j \leq m} N(a_{ji}e_j)$,
4. for $l = i + 1$ to m do:
5. $\alpha_l \leftarrow \alpha_l - \frac{a_{li}}{a_{ii}}\alpha_i$.

Return $(\alpha_1, \dots, \alpha_m)$.

Theorem 1.1 ([1]). *The algorithm above outputs an N -orthogonal basis of \mathcal{L} .*

Notice that the above theorem is proved only for $k = \mathbb{Q}_p$, but it is easy to see that the theorem also holds for $k = \mathbb{F}_p((T))$.

Now we can prove our main theorem of this subsection.

Theorem 1.2. *Let $k = \mathbb{Q}_p$ or $k = \mathbb{F}_p((T))$. Suppose Λ is any p -adic lattice of full rank in k^n , Λ^* is its dual. Then we have*

$$\tilde{\lambda}_i(\Lambda) \cdot \tilde{\lambda}_{n+1-i}(\Lambda^*) = 1$$

for $1 \leq i \leq n$, where $\tilde{\lambda}_i(\Lambda)$ is the i -th successive maxima of the lattice Λ and $\tilde{\lambda}_{n+1-i}(\Lambda)$ is the $(n+1-i)$ -th successive maxima of the lattice Λ^* with respect to the norm M .

Proof. Suppose $\Lambda = \mathcal{L}(B)$ and $D = (d_1, \dots, d_n)$ is the dual basis of $B = (b_1, \dots, b_n)$. We have $\Lambda^* = \mathcal{L}(D)$. Let the basis of k^n be e_1, \dots, e_n . Then e_1, \dots, e_n is an M -orthogonal basis of k^n because for any vector $v = v_1 e_1 + \dots + v_n e_n \in k^n$, the norm

$$M(v) = \max_{1 \leq i \leq n} |v_i|_p = \max_{1 \leq i \leq n} |v_i e_i|_p.$$

Therefore, we can run the algorithm “orthogonalization with orthogonal bases” to obtain the following M -orthogonal basis of Λ .

$$B = \begin{pmatrix} b_{11} & 0 & \cdots & 0 \\ b_{21} & b_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{pmatrix}$$

where $|b_{ii}|_p = \max \{|b_{jk}|_p : i \leq k \leq j \leq n\}$. Then we have $\tilde{\lambda}_i(\Lambda) = M(b_i) = |b_{ii}|_p$.

Let

$$B^* = \begin{pmatrix} B_{11} & 0 & \cdots & 0 \\ B_{12} & B_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ B_{1n} & B_{2n} & \cdots & B_{nn} \end{pmatrix}$$

be the adjoint matrix of B . We have

$$B_{ii} = \frac{\det B}{b_{ii}}$$

and

$$D = \left(\frac{B^*}{\det B} \right)^\top = \begin{pmatrix} b_{11}^{-1} & * & \cdots & * \\ 0 & b_{22}^{-1} & \cdots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & b_{nn}^{-1} \end{pmatrix}.$$

Write $D = (d_{ij})$. Since

$$|B_{ji}|_p \leq \left| \frac{\det B}{b_{ii}} \right|_p = |B_{ii}|_p,$$

we have $|d_{ij}|_p \leq |d_{jj}|_p$ for all $1 \leq i \leq j \leq n$. Also, we have

$$|d_{nn}|_p = |b_{nn}^{-1}|_p \geq |d_{n-1,n-1}|_p = |b_{n-1,n-1}^{-1}|_p \geq \cdots \geq |d_{11}|_p = |b_{11}^{-1}|_p.$$

Hence d_1, \dots, d_n is an orthogonal basis of Λ^* according to the algorithm. Therefore, we have

$$\tilde{\lambda}_{n+1-i}(\Lambda^*) = M(d_i) = |b_{ii}^{-1}|_p$$

and

$$\tilde{\lambda}_i(\Lambda) \cdot \tilde{\lambda}_{n+1-i}(\Lambda^*) = M(b_i) \cdot M(d_i) = 1$$

for $1 \leq i \leq n$. □

Example 1.3. Let $k = \mathbb{Q}_2$ and $n = 3$. Let the basis of Λ be

$$B = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 2 & 2 \\ 2 & 0 & 4 \end{pmatrix}.$$

We can run the algorithm “orthogonalization with orthogonal bases” to obtain an M -orthogonal basis of Λ ,

$$B' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 2 & -4 & 8 \end{pmatrix}.$$

Hence $\tilde{\lambda}_1(\Lambda) = 1$, $\tilde{\lambda}_2(\Lambda) = \frac{1}{2}$ and $\tilde{\lambda}_3(\Lambda) = \frac{1}{8}$ with respect to the norm M . The dual basis

$$D = \frac{1}{8} \begin{pmatrix} 4 & -4 & 2 \\ 2 & 2 & -1 \\ -2 & 2 & 1 \end{pmatrix}.$$

We can run the algorithm “orthogonalization with orthogonal bases” to obtain an M -orthogonal basis of Λ^* ,

$$D' = \frac{1}{8} \begin{pmatrix} 8 & -8 & 2 \\ 0 & 4 & -1 \\ 0 & 0 & 1 \end{pmatrix}.$$

Hence $\tilde{\lambda}_1(\Lambda^*) = 8$, $\tilde{\lambda}_2(\Lambda^*) = 2$ and $\tilde{\lambda}_3(\Lambda^*) = 1$ with respect to the norm M . We have

$$\tilde{\lambda}_1(\Lambda) \cdot \tilde{\lambda}_3(\Lambda^*) = \tilde{\lambda}_2(\Lambda) \cdot \tilde{\lambda}_2(\Lambda^*) = \tilde{\lambda}_3(\Lambda) \cdot \tilde{\lambda}_1(\Lambda^*) = 1.$$

1.2. Transference Theorem for General Norms

For a general norm N on k^m , there is also an upper bound for the transference theorem. We need the following lemma and proposition.

Lemma 1.4 ([3]). *Let N be a norm on k^m . Then the two norms M and N are equivalent, i.e., there are two positive constants c_1, c_2 such that*

$$c_1 N(v) \leq M(v) \leq c_2 N(v)$$

for any $v \in k^n$.

Proposition 1.5 ([1]). *Let N be a norm on k^m . Let $\mathcal{L} = \mathcal{L}(\alpha_1, \dots, \alpha_n)$ be a lattice of rank n in k^m such that $N(\alpha_1) \geq \dots \geq N(\alpha_n)$. Then $\tilde{\lambda}_1(\mathcal{L}) = N(\alpha_1)$ and $\tilde{\lambda}_i(\mathcal{L}) \leq N(\alpha_i)$ for $2 \leq i \leq n$.*

Notice that the above lemma and proposition is proved only for $k = \mathbb{Q}_p$, but it is easy to see that the proposition also holds for $k = \mathbb{F}_p((T))$.

Now we can prove the transference theorem for a general norm N on k^m .

Theorem 1.6. *Let $k = \mathbb{Q}_p$ or $k = \mathbb{F}_p((T))$. Suppose Λ is any p -adic lattice of full rank in k^n , Λ^* is its dual. Let N be a norm on k . Then we have*

$$\tilde{\lambda}_i(\Lambda) \cdot \tilde{\lambda}_{n+1-i}(\Lambda^*) \leq c'$$

for $1 \leq i \leq n$, where $\tilde{\lambda}_i(\Lambda)$ is the i -th successive maxima of the lattice Λ , $\tilde{\lambda}_{n+1-i}(\Lambda)$ is the $(n+1-i)$ -th successive maxima of the lattice Λ^* with respect to the norm N and c' is a positive constant depending on N .

Proof. Suppose $\Lambda = \mathcal{L}(B)$ and $D = (d_1, \dots, d_n)$ is the dual basis of $B = (b_1, \dots, b_n)$. We have $\Lambda^* = \mathcal{L}(D)$. Without loss of generality, we may assume

$$B = \begin{pmatrix} b_{11} & 0 & \cdots & 0 \\ b_{21} & b_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{pmatrix}$$

and $|b_{ii}|_p = \max \{ |b_{jk}|_p : i \leq k \leq j \leq n \}$. Notice that since e_1, \dots, e_n may not be an N -orthogonal basis of k^n , the above basis is not an N -orthogonal basis of Λ in general. However, we can still perform the transformation in the algorithm “orthogonalization with orthogonal bases” to obtain it.

Suppose that

$$N(b_{s_1}) \geq N(b_{s_2}) \geq \dots \geq N(b_{s_n})$$

where $\{s_1, s_2, \dots, s_n\} = \{1, 2, \dots, n\}$. Then by Proposition 1.5, we have

$$\tilde{\lambda}_i(\Lambda) \leq N(d_{s_i})$$

for $1 \leq i \leq n$.

The dual basis D is the same as in Theorem 1.2. For any $1 \leq i \leq n$, let $1 \leq j \leq i$ be the subscript such that

$$N(d_{s_j}) = \max_{1 \leq k \leq i} N(d_{s_k}).$$

By Proposition 1.5, we have

$$\tilde{\lambda}_{n-i+1}(\Lambda^*) \leq \max_{1 \leq k \leq i} N(d_{s_k}) = N(d_{s_j}).$$

By Lemma 1.4, two norms M and N are equivalent, i.e., there are two positive constants c_1 and c_2 such that

$$c_1 N(v) \leq M(v) \leq c_2 N(v)$$

for any $v \in k^n$. Then we have

$$\begin{aligned} \tilde{\lambda}_i(\Lambda) \cdot \tilde{\lambda}_{n-i+1}(\Lambda^*) &\leq N(b_{s_i}) \cdot N(d_{s_j}) \\ &\leq N(b_{s_j}) \cdot N(d_{s_j}) \\ &\leq c_1^{-2} M(b_{s_j}) \cdot M(d_{s_j}) \\ &= c_1^{-2} \end{aligned}$$

for $1 \leq i \leq n$. We can take $c' = c_1^{-2}$. □

Remark. Since we can choose $c_1 = \max_{1 \leq i \leq m} N(e_i)$, we can choose

$$c' = \left(\max_{1 \leq i \leq m} N(e_i) \right)^{-2}.$$

Example 1.7. Let $K = \mathbb{Q}_2(\pi)$, where π is a root of $f(x) = x^3 - 2$. Let N be the 2-adic absolute value on K . Since f is an Eisenstein polynomial, K is totally ramified of degree 3 and

$$N(\pi) = 2^{-\frac{1}{3}}.$$

Let $e_1 = \pi$, $e_2 = \pi^2$ and $e_3 = 2$. Then we can choose

$$c' = \left(\max_{1 \leq i \leq 3} N(e_i) \right)^{-2} = 2^{\frac{2}{3}}.$$

Let the basis of Λ be

$$B = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 2 & 2 \\ 2 & 0 & 4 \end{pmatrix}.$$

Since $\pi, \pi^2, 2$ is an N -orthogonal basis of K , we can run the algorithm “orthogonalization with orthogonal bases” to obtain an N -orthogonal basis of Λ ,

$$B' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 2 & -4 & 8 \end{pmatrix}.$$

Hence $\tilde{\lambda}_1(\Lambda) = 2^{-\frac{1}{3}}$, $\tilde{\lambda}_2(\Lambda) = 2^{-\frac{5}{3}}$ and $\tilde{\lambda}_3(\Lambda) = \frac{1}{16}$ with respect to the norm N . The dual basis

$$D = \frac{1}{8} \begin{pmatrix} 4 & -4 & 2 \\ 2 & 2 & -1 \\ -2 & 2 & 1 \end{pmatrix}.$$

We can run the algorithm “orthogonalization with orthogonal bases” to obtain an N -orthogonal basis of Λ^* ,

$$D' = \frac{1}{8} \begin{pmatrix} 8 & 0 & 2 \\ 0 & 0 & -1 \\ 0 & 4 & 1 \end{pmatrix}.$$

Hence $\tilde{\lambda}_1(\Lambda^*) = 4$, $\tilde{\lambda}_2(\Lambda^*) = 1$ and $\tilde{\lambda}_3(\Lambda^*) = 2^{-\frac{1}{3}}$ with respect to the norm N . We have

$$\begin{aligned} \tilde{\lambda}_1(\Lambda) \cdot \tilde{\lambda}_3(\Lambda^*) &= 2^{-\frac{2}{3}} \leq 2^{\frac{2}{3}}, \\ \tilde{\lambda}_2(\Lambda) \cdot \tilde{\lambda}_2(\Lambda^*) &= 2^{-\frac{5}{3}} \leq 2^{\frac{2}{3}}, \end{aligned}$$

and

$$\tilde{\lambda}_3(\Lambda) \cdot \tilde{\lambda}_1(\Lambda^*) = \frac{1}{4} \leq 2^{\frac{2}{3}}.$$

2. Dual of Orthogonal Basis

It is interesting to ask whether the dual basis of an orthogonal basis is still an orthogonal basis. It turns out that for the norm M , the answer is affirmative. However, this conjecture does not hold for all norms.

We need the following theorems.

Theorem 2.1 ([1]). *Let N be a norm on k^m . Let $\mathcal{L} = \mathcal{L}(\alpha_1, \dots, \alpha_n)$ be a lattice of rank n in k^m with an N -orthogonal basis $\alpha_1, \dots, \alpha_n$. If β_1, \dots, β_n can be obtained from $\alpha_1, \dots, \alpha_n$ by the following operations:*

1. $\alpha_i \leftarrow k\alpha_i$ for some $k \in \mathbb{Z}_p \setminus p\mathbb{Z}_p$,
2. $\alpha_i \leftrightarrow \alpha_j$,
3. $\alpha_i \leftarrow \alpha_i + k\alpha_j$ for some $k \in \mathbb{Z}_p$ such that $N(k\alpha_j) \leq N(\alpha_i)$,

then β_1, \dots, β_n is also an N -orthogonal basis of \mathcal{L} .

Theorem 2.2 ([1]). *Let N be a norm on k^m . Let \mathcal{L} be a lattice of rank n in k^m . Then any two N -orthogonal bases of \mathcal{L} can be obtained from each other by the following operations:*

1. $\alpha_i \leftarrow k\alpha_i$ for some $k \in \mathbb{Z}_p \setminus p\mathbb{Z}_p$,
2. $\alpha_i \leftrightarrow \alpha_j$,
3. $\alpha_i \leftarrow \alpha_i + k\alpha_j$ for some $k \in \mathbb{Z}_p$ such that $N(k\alpha_j) \leq N(\alpha_i)$.

Notice that the above theorems are proved only for $k = \mathbb{Q}_p$, but it is easy to see that these theorems also hold for $k = \mathbb{F}_p((T))$.

Now we can prove our main theorem of this section.

Theorem 2.3. *Suppose Λ is any p -adic lattice of full rank in k^n , Λ^* is its dual. Suppose $\Lambda = \mathcal{L}(B)$ and D is the dual basis of B . If B is an M -orthogonal basis of Λ , then D is an M -orthogonal basis of Λ^* .*

Proof. According to the proof of Theorem 1.2, if the M -orthogonal basis of Λ is of the form

$$B' = \begin{pmatrix} b'_{11} & 0 & \cdots & 0 \\ b'_{21} & b'_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ b'_{n1} & b'_{n2} & \cdots & b'_{nn} \end{pmatrix}$$

where $|b'_{ii}|_p = \max \left\{ |b'_{jk}|_p : i \leq k \leq j \leq n \right\}$, then the dual basis of B will be of the form

$$D' = \begin{pmatrix} d'_{11} & d'_{12} & \cdots & d'_{1n} \\ 0 & d'_{22} & \cdots & d'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d'_{nn} \end{pmatrix}.$$

where $|d'_{ii}|_p = \max \left\{ |d'_{jk}|_p : 1 \leq j \leq k \leq i \right\}$. Hence it is an M -orthogonal basis of Λ^* .

Now let B be a general M -orthogonal basis of Λ . By Theorem 2.2, B can be obtained from B' by the following operations:

1. $b_i \leftarrow kb_i$ for some $k \in \mathbb{Z}_p \setminus p\mathbb{Z}_p$,
2. $b_i \leftrightarrow b_j$,
3. $b_i \leftarrow b_i + kb_j$ for some $k \in \mathbb{Z}_p$ such that $N(kb_j) \leq N(b_i)$.

Write $B = B'E$ where $E = E_1E_2 \cdots E_s$ and E_i ($1 \leq i \leq s$) is a fundamental column transformation matrix of one of the above three forms. Then

$$D = D'(E^{-1})^\top = D'(E_1^{-1})^\top(E_2^{-1})^\top \cdots (E_s^{-1})^\top.$$

Let

$$B_t = B'E_1E_2 \cdots E_t = (b_1^{(t)}, \dots, b_n^{(t)})$$

and

$$D_t = D'(E_1^{-1})^\top(E_2^{-1})^\top \cdots (E_t^{-1})^\top = (d_1^{(t)}, \dots, d_n^{(t)})$$

be the basis and dual basis after the t -th fundamental column transformation.

Set $B_0 = B'$ and $D_0 = D'$. We claim that $d_1^{(t)}, \dots, d_n^{(t)}$ is an M -orthogonal basis of Λ^* and $M(b_i^{(t)}) \cdot M(d_i^{(t)}) = 1$ for any $1 \leq i \leq n$ and $0 \leq t \leq s$. When $t = 0$ the claim is proved in Theorem 1.2. Suppose that the claim holds for some $0 \leq t \leq s-1$.

If E_{t+1} is the operation 1, then $(E_{t+1}^{-1})^\top$ will be the operation $d_i^{(t)} \leftarrow k^{-1}d_i^{(t)}$, also of the form 1. By Theorem 2.1, $d_1^{(t+1)}, \dots, d_n^{(t+1)}$ is an M -orthogonal basis of Λ^* . On the other hand, we have

$$M(b_i^{(t+1)}) \cdot M(d_i^{(t+1)}) = M(b_i^{(t)}) \cdot M(d_i^{(t)}) = 1.$$

Hence the claim holds for $t+1$.

If E_{t+1} is the operation 2, then $(E_{t+1}^{-1})^\top$ will be the operation $d_i^{(t)} \leftrightarrow d_j^{(t)}$. also of the form 2. By Theorem 2.1, $d_1^{(t+1)}, \dots, d_n^{(t+1)}$ is an M -orthogonal basis of Λ^* . On the other hand, we have

$$M(b_i^{(t+1)}) \cdot M(d_i^{(t+1)}) = M(b_j^{(t)}) \cdot M(d_j^{(t)}) = 1$$

and

$$M(b_j^{(t+1)}) \cdot M(d_j^{(t+1)}) = M(b_i^{(t)}) \cdot M(d_i^{(t)}) = 1.$$

Hence the claim holds for $t+1$.

If E_{t+1} is the operation 3, then

$$N(kb_j^{(t)}) \leq N(b_i^{(t)})$$

and $(E_{t+1}^{-1})^\top$ will be the operation $d_j^{(t)} \leftarrow d_j^{(t)} - kd_i^{(t)}$. Since

$$N(-kd_i^{(t)}) = |k|_p \cdot N(b_i^{(t)})^{-1} \leq N(b_j^{(t)})^{-1} = N(d_j^{(t)}),$$

this operation is also of the form 3. By Theorem 2.1, $d_1^{(t+1)}, \dots, d_n^{(t+1)}$ is an M -orthogonal basis of Λ^* . On the other hand, since $b_1^{(t)}, \dots, b_n^{(t)}$ and $d_1^{(t)}, \dots, d_n^{(t)}$ are

M -orthogonal bases of Λ and Λ^* respectively, we have

$$\begin{aligned} M\left(b_i^{(t+1)}\right) \cdot M\left(d_i^{(t+1)}\right) &= M\left(b_i^{(t)} + kb_j^{(t)}\right) \cdot M\left(d_i^{(t)}\right) \\ &= M\left(b_j^{(t)}\right) \cdot M\left(d_j^{(t)}\right) \\ &= 1. \end{aligned}$$

and

$$\begin{aligned} M\left(b_j^{(t+1)}\right) \cdot M\left(d_j^{(t+1)}\right) &= M\left(b_j^{(t)}\right) \cdot M\left(d_j^{(t)} - kd_i^{(t)}\right) \\ &= M\left(b_j^{(t)}\right) \cdot M\left(d_j^{(t)}\right) \\ &= 1. \end{aligned}$$

Hence the claim holds for $t + 1$.

Therefore, by induction, $D = D_s$ is an M -orthogonal basis of Λ^* . \square

For other norms, here is a simple counterexample:

Example 2.4. Let $K = \mathbb{Q}_2(\pi)$, where π is a root of $f(x) = x^2 - 2$. Let N be the 2-adic absolute value on K . Since f is an Eisenstein polynomial, K is totally ramified of degree 2 and

$$|\pi|_2 = 2^{-\frac{1}{2}}.$$

Let the basis of K be $e_1 = 1$ and $e_2 = \pi$. Let $\Lambda = \mathcal{L}(B)$ where

$$B = (b_1, b_2) = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

Then we have $b_1 = 1 + \pi$ and $b_2 = \pi$. According to the algorithm “orthogonalization with orthogonal bases”, b_1, b_2 is an N -orthogonal basis of Λ .

The dual basis of B is

$$D = (d_1, d_2) = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.$$

Then we have $d_1 = 1$ and $d_2 = \pi - 1$. However, since

$$|d_1 + d_2|_2 = 2^{-\frac{1}{2}} < 1 = \max\{|d_1|_2, |d_2|_2\},$$

d_1, d_2 is not an N -orthogonal basis of Λ^* .

References

- [1] Chi Zhang, Yingpu Deng and Zhaonan Wang, *Norm Orthogonal Bases and Invariants of p -adic Lattices*, arXiv preprint arXiv: 2311.17415, 2023.

- [2] Yingpu Deng, *On p -adic Minkowski's Theorems*, arXiv preprint arXiv: 2401.14023, 2024.
- [3] A.M. Robert, *A Course in p -adic Analysis*, GTM 198, Springer, New York, 2000.
- [4] A. Weil, *Basic Number Theory*, Third edition, Springer, New York, 1974.
- [5] O. Regev, *Lattices in Computer Science*, Lecture notes of a course given in Tel Aviv University, 2004.