

Search for the $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ process at BESIII

M. Ablikim¹, M. N. Achasov^{4,c}, P. Adlarson⁷⁵, O. Afedulidis³, X. C. Ai⁸⁰, R. Aliberti³⁵, A. Amoroso^{74A,74C}, Q. An^{71,58,a}, Y. Bai⁵⁷, O. Bakina³⁶, I. Balossino^{29A}, Y. Ban^{46,h}, H.-R. Bao⁶³, V. Batozskaya^{1,44}, K. Begzsuren³², N. Berger³⁵, M. Berlowski⁴⁴, M. Bertani^{28A}, D. Bettoni^{29A}, F. Bianchi^{74A,74C}, E. Bianco^{74A,74C}, A. Bortone^{74A,74C}, I. Boyko³⁶, R. A. Briere⁵, A. Brueggemann⁶⁸, H. Cai⁷⁶, X. Cai^{1,58}, A. Calcaterra^{28A}, G. F. Cao^{1,63}, N. Cao^{1,63}, S. A. Cetin^{62A}, J. F. Chang^{1,58}, G. R. Che⁴³, G. Chelkov^{36,b}, C. Chen⁴³, C. H. Chen⁹, Chao Chen⁵⁵, G. Chen¹, H. S. Chen^{1,63}, H. Y. Chen²⁰, M. L. Chen^{1,58,63}, S. J. Chen⁴², S. L. Chen⁴⁵, S. M. Chen⁶¹, T. Chen^{1,63}, X. R. Chen^{31,63}, X. T. Chen^{1,63}, Y. B. Chen^{1,58}, Y. Q. Chen³⁴, Z. J. Chen^{25,i}, Z. Y. Chen^{1,63}, S. K. Choi^{10A}, G. Cibinetto^{29A}, F. Cossio^{74C}, J. J. Cui⁵⁰, H. L. Dai^{1,58}, J. P. Dai⁷⁸, A. Dbeyssi¹⁸, R. E. de Boer³, D. Dedovich³⁶, C. Q. Deng⁷², Z. Y. Deng¹, A. Denig³⁵, I. Denysenko³⁶, M. Destefanis^{74A,74C}, F. De Mori^{74A,74C}, B. Ding^{66,1}, X. X. Ding^{46,h}, Y. Ding⁴⁰, Y. Ding³⁴, J. Dong^{1,58}, L. Y. Dong^{1,63}, M. Y. Dong^{1,58,63}, X. Dong⁷⁶, M. C. Du¹, S. X. Du⁸⁰, Y. Y. Duan⁵⁵, Z. H. Duan⁴², P. Egorov^{36,b}, Y. H. Fan⁴⁵, J. Fang^{1,58}, J. Fang⁵⁹, S. S. Fang^{1,63}, W. X. Fang¹, Y. Fang¹, Y. Q. Fang^{1,58}, R. Farinelli^{29A}, L. Fava^{74B,74C}, F. Feldbauer³, G. Felici^{28A}, C. Q. Feng^{71,58}, J. H. Feng⁵⁹, Y. T. Feng^{71,58}, M. Fritsch³, C. D. Fu¹, J. L. Fu⁶³, Y. W. Fu^{1,63}, H. Gao⁶³, X. B. Gao⁴¹, Y. N. Gao^{46,h}, Yang Gao^{71,58}, S. Garbolino^{74C}, I. Garzia^{29A,29B}, L. Ge⁸⁰, P. T. Ge⁷⁶, Z. W. Ge⁴², C. Geng⁵⁹, E. M. Gersabeck⁶⁷, A. Gilman⁶⁹, K. Goetzen¹³, L. Gong⁴⁰, W. X. Gong^{1,58}, W. Gradl³⁵, S. Gramigna^{29A,29B}, M. Greco^{74A,74C}, M. H. Gu^{1,58}, Y. T. Gu¹⁵, C. Y. Guan^{1,63}, A. Q. Guo^{31,63}, L. B. Guo⁴¹, M. J. Guo⁵⁰, R. P. Guo⁴⁹, Y. P. Guo^{12,g}, A. Guskov^{36,b}, J. Gutierrez²⁷, K. L. Han⁶³, T. T. Han¹, F. Hanisch³, X. Q. Hao¹⁹, F. A. Harris⁶⁵, K. K. He⁵⁵, K. L. He^{1,63}, F. H. Heinsius³, C. H. Heinz³⁵, Y. K. Heng^{1,58,63}, C. Herold⁶⁰, T. Holtmann³, P. C. Hong³⁴, G. Y. Hou^{1,63}, X. T. Hou^{1,63}, Y. R. Hou⁶³, Z. L. Hou¹, B. Y. Hu⁵⁹, H. M. Hu^{1,63}, J. F. Hu^{56,j}, S. L. Hu^{12,g}, T. Hu^{1,58,63}, Y. Hu¹, G. S. Huang^{71,58}, K. X. Huang⁵⁹, L. Q. Huang^{31,63}, X. T. Huang⁵⁰, Y. P. Huang¹, Y. S. Huang⁵⁹, T. Hussain⁷³, F. Hölzken³, N. Hüsken³⁵, N. in der Wiesche⁶⁸, J. Jackson²⁷, S. Janchiv³², J. H. Jeong^{10A}, Q. Ji¹, Q. P. Ji¹⁹, W. Ji^{1,63}, X. B. Ji^{1,63}, X. L. Ji^{1,58}, Y. Y. Ji⁵⁰, X. Q. Jia⁵⁰, Z. K. Jia^{71,58}, D. Jiang^{1,63}, H. B. Jiang⁷⁶, P. C. Jiang^{46,h}, S. S. Jiang³⁹, T. J. Jiang¹⁶, X. S. Jiang^{1,58,63}, Y. Jiang⁶³, J. B. Jiao⁵⁰, J. K. Jiao³⁴, Z. Jiao²³, S. Jin⁴², Y. Jin⁶⁶, M. Q. Jing^{1,63}, X. M. Jing⁶³, T. Johansson⁷⁵, S. Kabana³³, N. Kalantar-Nayestanaki⁶⁴, X. L. Kang⁹, X. S. Kang⁴⁰, M. Kavatsyuk⁶⁴, B. C. Ke⁸⁰, V. Khachatryan²⁷, A. Khoukaz⁶⁸, R. Kiuchi¹, O. B. Kolcu^{62A}, B. Kopf³, M. Kuessner³, X. Kui^{1,63}, N. Kumar²⁶, A. Kupsc^{44,75}, W. Kühn³⁷, J. J. Lane⁶⁷, P. Larin¹⁸, L. Lavezzi^{74A,74C}, T. T. Lei^{71,58}, Z. H. Lei^{71,58}, M. Lellmann³⁵, T. Lenz³⁵, C. Li⁴³, C. Li⁴⁷, C. H. Li³⁹, Cheng Li^{71,58}, D. M. Li⁸⁰, F. Li^{1,58}, G. Li¹, H. B. Li^{1,63}, H. J. Li¹⁹, H. N. Li^{56,j}, Hui Li⁴³, J. R. Li⁶¹, J. S. Li⁵⁹, K. Li¹, L. J. Li^{1,63}, L. K. Li¹, Lei Li⁴⁸, M. H. Li⁴³, P. R. Li^{38,k,l}, Q. M. Li^{1,63}, Q. X. Li⁵⁰, R. Li^{17,31}, S. X. Li¹², T. Li⁵⁰, W. D. Li^{1,63}, W. G. Li^{1,a}, X. Li^{1,63}, X. H. Li^{71,58}, X. L. Li⁵⁰, X. Y. Li^{1,63}, X. Z. Li⁵⁹, Y. G. Li^{46,h}, Z. J. Li⁵⁹, Z. Y. Li⁷⁸, C. Liang⁴², H. Liang^{71,58}, H. Liang^{1,63}, Y. F. Liang⁵⁴, Y. T. Liang^{31,63}, G. R. Liao¹⁴, L. Z. Liao⁵⁰, Y. P. Liao^{1,63}, J. Libby²⁶, A. Limphirat⁶⁰, C. C. Lin⁵⁵, D. X. Lin^{31,63}, T. Lin¹, B. J. Liu¹, B. X. Liu⁷⁶, C. Liu³⁴, C. X. Liu¹, F. Liu¹, F. H. Liu⁵³, Feng Liu⁶, G. M. Liu^{56,j}, H. Liu^{38,k,l}, H. B. Liu¹⁵, H. H. Liu¹, H. M. Liu^{1,63}, Huihui Liu²¹, J. B. Liu^{71,58}, J. Y. Liu^{1,63}, K. Liu^{38,k,l}, K. Y. Liu⁴⁰, Ke Liu²², L. Liu^{71,58}, L. C. Liu⁴³, Lu Liu⁴³, M. H. Liu^{12,g}, P. L. Liu¹, Q. Liu⁶³, S. B. Liu^{71,58}, T. Liu^{12,g}, W. K. Liu⁴³, W. M. Liu^{71,58}, X. Liu^{38,k,l}, X. Liu³⁹, Y. Liu^{38,k,l}, Y. Liu⁸⁰, Y. B. Liu⁴³, Z. A. Liu^{1,58,63}, Z. D. Liu⁹, Z. Q. Liu⁵⁰, X. C. Lou^{1,58,63}, F. X. Lu⁵⁹, H. J. Lu²³, J. G. Lu^{1,58}, X. L. Lu¹, Y. Lu⁷, Y. P. Lu^{1,58}, Z. H. Lu^{1,63}, C. L. Luo⁴¹, J. R. Luo⁵⁹, M. X. Luo⁷⁹, T. Luo^{12,g}, X. L. Luo^{1,58}, X. R. Lyu⁶³, Y. F. Lyu⁴³, F. C. Ma⁴⁰, H. Ma⁷⁸, H. L. Ma¹, J. L. Ma^{1,63}, L. L. Ma⁵⁰, M. M. Ma^{1,63}, Q. M. Ma¹, R. Q. Ma^{1,63}, T. Ma^{71,58}, X. T. Ma^{1,63}, X. Y. Ma^{1,58}, Y. Ma^{46,h}, Y. M. Ma³¹, F. E. Maas¹⁸, M. Maggiora^{74A,74C}, S. Malde⁶⁹, Y. J. Mao^{46,h}, Z. P. Mao¹, S. Marcello^{74A,74C}, Z. X. Meng⁶⁶, J. G. Messchendorp^{13,64}, G. Mezzadri^{29A}, H. Miao^{1,63}, T. J. Min⁴², R. E. Mitchell²⁷, X. H. Mo^{1,58,63}, B. Moses²⁷, N. Yu. Muchnoi^{4,c}, J. Muskalla³⁵, Y. Nefedov³⁶, F. Nerling^{18,e}, L. S. Nie²⁰, I. B. Nikolaev^{4,c}, Z. Ning^{1,58}, S. Nisar^{11,m}, Q. L. Niu^{38,k,l}, W. D. Niu⁵⁵, Y. Niu⁵⁰, S. L. Olsen⁶³, Q. Ouyang^{1,58,63}, S. Pacetti^{28B,28C}, X. Pan⁵⁵, Y. Pan⁵⁷, A. Pathak³⁴, P. Patteri^{28A}, Y. P. Pei^{71,58}, M. Pelizaeus³, H. P. Peng^{71,58}, Y. Y. Peng^{38,k,l}, K. Peters^{13,e}, J. L. Ping⁴¹, R. G. Ping^{1,63}, S. Plura³⁵, V. Prasad³³, F. Z. Qi¹, H. Qi^{71,58}, H. R. Qi⁶¹, M. Qi⁴², T. Y. Qi^{12,g}, S. Qian^{1,58}, W. B. Qian⁶³, C. F. Qiao⁶³, X. K. Qiao⁸⁰, J. J. Qin⁷², L. Q. Qin¹⁴, L. Y. Qin^{71,58}, X. P. Qin^{12,g}, X. S. Qin⁵⁰, Z. H. Qin^{1,58}, J. F. Qiu¹, Z. H. Qu⁷², C. F. Redmer³⁵, K. J. Ren³⁹, A. Rivetti^{74C}, M. Rolo^{74C}, G. Rong^{1,63}, Ch. Rosner¹⁸, S. N. Ruan⁴³, N. Salone⁴⁴, A. Sarantsev^{36,d}, Y. Schelhaas³⁵, K. Schoenning⁷⁵, M. Scodeggio^{29A}, K. Y. Shan^{12,g}, W. Shan²⁴, X. Y. Shan^{71,58}, Z. J. Shang^{38,k,l}, J. F. Shangguan¹⁶, L. G. Shao^{1,63}, M. Shao^{71,58}, C. P. Shen^{12,g}, H. F. Shen^{1,8}, W. H. Shen⁶³, X. Y. Shen^{1,63}, B. A. Shi⁶³, H. Shi^{71,58}, H. C. Shi^{71,58}, J. L. Shi^{12,g}, J. Y. Shi¹, Q. Q. Shi⁵⁵, S. Y. Shi⁷², X. Shi^{1,58}, J. J. Song¹⁹, T. Z. Song⁵⁹, W. M. Song^{34,1}, Y. J. Song^{12,g}, Y. X. Song^{46,h,m}, S. Sosio^{74A,74C}, S. Spataro^{74A,74C}, F. Stieler³⁵, Y. J. Su⁶³, G. B. Sun⁷⁶, G. X. Sun¹,

H. Sun⁶³, H. K. Sun¹, J. F. Sun¹⁹, K. Sun⁶¹, L. Sun⁷⁶, S. S. Sun^{1,63}, T. Sun^{51,f}, W. Y. Sun³⁴, Y. Sun⁹,
 Y. J. Sun^{71,58}, Y. Z. Sun¹, Z. Q. Sun^{1,63}, Z. T. Sun⁵⁰, C. J. Tang⁵⁴, G. Y. Tang¹, J. Tang⁵⁹, M. Tang^{71,58},
 Y. A. Tang⁷⁶, L. Y. Tao⁷², Q. T. Tao^{25,i}, M. Tat⁶⁹, J. X. Teng^{71,58}, V. Thoren⁷⁵, W. H. Tian⁵⁹, Y. Tian^{31,63},
 Z. F. Tian⁷⁶, I. Uman^{62B}, Y. Wan⁵⁵, S. J. Wang⁵⁰, B. Wang¹, B. L. Wang⁶³, Bo Wang^{71,58}, D. Y. Wang^{46,h},
 F. Wang⁷², H. J. Wang^{38,k,l}, J. J. Wang⁷⁶, J. P. Wang⁵⁰, K. Wang^{1,58}, L. L. Wang¹, M. Wang⁵⁰, N. Y. Wang⁶³,
 S. Wang^{12,g}, S. Wang^{38,k,l}, T. Wang^{12,g}, T. J. Wang⁴³, W. Wang⁵⁹, W. Wang⁷², W. P. Wang^{35,71,o}, X. Wang^{46,h},
 X. F. Wang^{38,k,l}, X. J. Wang³⁹, X. L. Wang^{12,g}, X. N. Wang¹, Y. Wang⁶¹, Y. D. Wang⁴⁵, Y. F. Wang^{1,58,63},
 Y. L. Wang¹⁹, Y. N. Wang⁴⁵, Y. Q. Wang¹, Yaqian Wang¹⁷, Yi Wang⁶¹, Z. Wang^{1,58}, Z. L. Wang⁷²,
 Z. Y. Wang^{1,63}, Ziyi Wang⁶³, D. H. Wei¹⁴, F. Weidner⁶⁸, S. P. Wen¹, Y. R. Wen³⁹, U. Wiedner³, G. Wilkinson⁶⁹,
 M. Wolke⁷⁵, L. Wollenberg³, C. Wu³⁹, J. F. Wu^{1,8}, L. H. Wu¹, L. J. Wu^{1,63}, X. Wu^{12,g}, X. H. Wu³⁴, Y. Wu^{71,58},
 Y. H. Wu⁵⁵, Y. J. Wu³¹, Z. Wu^{1,58}, L. Xia^{71,58}, X. M. Xian³⁹, B. H. Xiang^{1,63}, T. Xiang^{46,h}, D. Xiao^{38,k,l},
 G. Y. Xiao⁴², S. Y. Xiao¹, Y. L. Xiao^{12,g}, Z. J. Xiao⁴¹, C. Xie⁴², X. H. Xie^{46,h}, Y. Xie⁵⁰, Y. G. Xie^{1,58}, Y. H. Xie⁶,
 Z. P. Xie^{71,58}, T. Y. Xing^{1,63}, C. F. Xu^{1,63}, C. J. Xu⁵⁹, G. F. Xu¹, H. Y. Xu^{66,2,p}, M. Xu^{71,58}, Q. J. Xu¹⁶,
 Q. N. Xu³⁰, W. Xu¹, W. L. Xu⁶⁶, X. P. Xu⁵⁵, Y. C. Xu⁷⁷, Z. P. Xu⁴², Z. S. Xu⁶³, F. Yan^{12,g}, L. Yan^{12,g},
 W. B. Yan^{71,58}, W. C. Yan⁸⁰, X. Q. Yan¹, H. J. Yang^{51,f}, H. L. Yang³⁴, H. X. Yang¹, T. Yang¹, Y. Yang^{12,g},
 Y. F. Yang^{1,63}, Y. F. Yang⁴³, Y. X. Yang^{1,63}, Z. W. Yang^{38,k,l}, Z. P. Yao⁵⁰, M. Ye^{1,58}, M. H. Ye⁸, J. H. Yin¹,
 Z. Y. You⁵⁹, B. X. Yu^{1,58,63}, C. X. Yu⁴³, G. Yu^{1,63}, J. S. Yu^{25,i}, T. Yu⁷², X. D. Yu^{46,h}, Y. C. Yu⁸⁰, C. Z. Yuan^{1,63},
 J. Yuan³⁴, J. Yuan⁴⁵, L. Yuan², S. C. Yuan^{1,63}, Y. Yuan^{1,63}, Z. Y. Yuan⁵⁹, C. X. Yue³⁹, A. A. Zafar⁷³, F. R. Zeng⁵⁰,
 S. H. Zeng⁷², X. Zeng^{12,g}, Y. Zeng^{25,i}, Y. J. Zeng^{1,63}, Y. J. Zeng⁵⁹, X. Y. Zhai³⁴, Y. C. Zhai⁵⁰, Y. H. Zhan⁵⁹,
 A. Q. Zhang^{1,63}, B. L. Zhang^{1,63}, B. X. Zhang¹, D. H. Zhang⁴³, G. Y. Zhang¹⁹, H. Zhang⁸⁰, H. Zhang^{71,58},
 H. C. Zhang^{1,58,63}, H. H. Zhang³⁴, H. H. Zhang⁵⁹, H. Q. Zhang^{1,58,63}, H. R. Zhang^{71,58}, H. Y. Zhang^{1,58}, J. Zhang⁸⁰,
 J. Zhang⁵⁹, J. J. Zhang⁵², J. L. Zhang²⁰, J. Q. Zhang⁴¹, J. S. Zhang^{12,g}, J. W. Zhang^{1,58,63}, J. X. Zhang^{38,k,l},
 J. Y. Zhang¹, J. Z. Zhang^{1,63}, Jianyu Zhang⁶³, L. M. Zhang⁶¹, Lei Zhang⁴², P. Zhang^{1,63}, Q. Y. Zhang³⁴,
 R. Y. Zhang^{38,k,l}, S. H. Zhang^{1,63}, Shulei Zhang^{25,i}, X. D. Zhang⁴⁵, X. M. Zhang¹, X. Y. Zhang⁵⁰, Y. Zhang⁷²,
 Y. Zhang¹, Y. T. Zhang⁸⁰, Y. H. Zhang^{1,58}, Y. M. Zhang³⁹, Yan Zhang^{71,58}, Z. D. Zhang¹, Z. H. Zhang¹,
 Z. L. Zhang³⁴, Z. Y. Zhang⁷⁶, Z. Y. Zhang⁴³, Z. Z. Zhang⁴⁵, G. Zhao¹, J. Y. Zhao^{1,63}, J. Z. Zhao^{1,58}, L. Zhao¹,
 Lei Zhao^{71,58}, M. G. Zhao⁴³, N. Zhao⁷⁸, R. P. Zhao⁶³, S. J. Zhao⁸⁰, Y. B. Zhao^{1,58}, Y. X. Zhao^{31,63}, Z. G. Zhao^{71,58},
 A. Zhemchugov^{36,b}, B. Zheng⁷², B. M. Zheng³⁴, J. P. Zheng^{1,58}, W. J. Zheng^{1,63}, Y. H. Zheng⁶³, B. Zhong⁴¹,
 X. Zhong⁵⁹, H. Zhou⁵⁰, J. Y. Zhou³⁴, L. P. Zhou^{1,63}, S. Zhou⁶, X. Zhou⁷⁶, X. K. Zhou⁶, X. R. Zhou^{71,58},
 X. Y. Zhou³⁹, Y. Z. Zhou^{12,g}, J. Zhu⁴³, K. Zhu¹, K. J. Zhu^{1,58,63}, K. S. Zhu^{12,g}, L. Zhu³⁴, L. X. Zhu⁶³, S. H. Zhu⁷⁰,
 S. Q. Zhu⁴², T. J. Zhu^{12,g}, W. D. Zhu⁴¹, Y. C. Zhu^{71,58}, Z. A. Zhu^{1,63}, J. H. Zou¹, J. Zu^{71,58} (BESIII Collaboration)

¹ *Institute of High Energy Physics, Beijing 100049, People's Republic of China*

² *Beihang University, Beijing 100191, People's Republic of China*

³ *Bochum Ruhr-University, D-44780 Bochum, Germany*

⁴ *Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*

⁵ *Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

⁶ *Central China Normal University, Wuhan 430079, People's Republic of China*

⁷ *Central South University, Changsha 410083, People's Republic of China*

⁸ *China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*

⁹ *China University of Geosciences, Wuhan 430074, People's Republic of China*

¹⁰ *Chung-Ang University, Seoul, 06974, Republic of Korea*

¹¹ *COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*

¹² *Fudan University, Shanghai 200433, People's Republic of China*

¹³ *GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*

¹⁴ *Guangxi Normal University, Guilin 541004, People's Republic of China*

¹⁵ *Guangxi University, Nanning 530004, People's Republic of China*

¹⁶ *Hangzhou Normal University, Hangzhou 310036, People's Republic of China*

¹⁷ *Hebei University, Baoding 071002, People's Republic of China*

¹⁸ *Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*

¹⁹ *Henan Normal University, Xinxiang 453007, People's Republic of China*

²⁰ *Henan University, Kaifeng 475004, People's Republic of China*

²¹ *Henan University of Science and Technology, Luoyang 471003, People's Republic of China*

²² *Henan University of Technology, Zhengzhou 450001, People's Republic of China*

- ²³ *Huangshan College, Huangshan 245000, People's Republic of China*
- ²⁴ *Hunan Normal University, Changsha 410081, People's Republic of China*
- ²⁵ *Hunan University, Changsha 410082, People's Republic of China*
- ²⁶ *Indian Institute of Technology Madras, Chennai 600036, India*
- ²⁷ *Indiana University, Bloomington, Indiana 47405, USA*
- ²⁸ *INFN Laboratori Nazionali di Frascati, (A)INFN Laboratori Nazionali di Frascati, I-00044, Frascati, Italy; (B)INFN Sezione di Perugia, I-06100, Perugia, Italy; (C)University of Perugia, I-06100, Perugia, Italy*
- ²⁹ *INFN Sezione di Ferrara, (A)INFN Sezione di Ferrara, I-44122, Ferrara, Italy; (B)University of Ferrara, I-44122, Ferrara, Italy*
- ³⁰ *Inner Mongolia University, Hohhot 010021, People's Republic of China*
- ³¹ *Institute of Modern Physics, Lanzhou 730000, People's Republic of China*
- ³² *Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia*
- ³³ *Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile*
- ³⁴ *Jilin University, Changchun 130012, People's Republic of China*
- ³⁵ *Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*
- ³⁶ *Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia*
- ³⁷ *Justus-Liebig-Universität Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany*
- ³⁸ *Lanzhou University, Lanzhou 730000, People's Republic of China*
- ³⁹ *Liaoning Normal University, Dalian 116029, People's Republic of China*
- ⁴⁰ *Liaoning University, Shenyang 110036, People's Republic of China*
- ⁴¹ *Nanjing Normal University, Nanjing 210023, People's Republic of China*
- ⁴² *Nanjing University, Nanjing 210093, People's Republic of China*
- ⁴³ *Nankai University, Tianjin 300071, People's Republic of China*
- ⁴⁴ *National Centre for Nuclear Research, Warsaw 02-093, Poland*
- ⁴⁵ *North China Electric Power University, Beijing 102206, People's Republic of China*
- ⁴⁶ *Peking University, Beijing 100871, People's Republic of China*
- ⁴⁷ *Qufu Normal University, Qufu 273165, People's Republic of China*
- ⁴⁸ *Renmin University of China, Beijing 100872, People's Republic of China*
- ⁴⁹ *Shandong Normal University, Jinan 250014, People's Republic of China*
- ⁵⁰ *Shandong University, Jinan 250100, People's Republic of China*
- ⁵¹ *Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China*
- ⁵² *Shanxi Normal University, Linfen 041004, People's Republic of China*
- ⁵³ *Shanxi University, Taiyuan 030006, People's Republic of China*
- ⁵⁴ *Sichuan University, Chengdu 610064, People's Republic of China*
- ⁵⁵ *Soochow University, Suzhou 215006, People's Republic of China*
- ⁵⁶ *South China Normal University, Guangzhou 510006, People's Republic of China*
- ⁵⁷ *Southeast University, Nanjing 211100, People's Republic of China*
- ⁵⁸ *State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China*
- ⁵⁹ *Sun Yat-Sen University, Guangzhou 510275, People's Republic of China*
- ⁶⁰ *Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand*
- ⁶¹ *Tsinghua University, Beijing 100084, People's Republic of China*
- ⁶² *Turkish Accelerator Center Particle Factory Group, (A)Istinye University, 34010, Istanbul, Turkey; (B)Near East University, Nicosia, North Cyprus, 99138, Mersin 10, Turkey*
- ⁶³ *University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China*
- ⁶⁴ *University of Groningen, NL-9747 AA Groningen, The Netherlands*
- ⁶⁵ *University of Hawaii, Honolulu, Hawaii 96822, USA*
- ⁶⁶ *University of Jinan, Jinan 250022, People's Republic of China*
- ⁶⁷ *University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom*
- ⁶⁸ *University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany*
- ⁶⁹ *University of Oxford, Keble Road, Oxford OX13RH, United Kingdom*
- ⁷⁰ *University of Science and Technology Liaoning, Anshan 114051, People's Republic of China*
- ⁷¹ *University of Science and Technology of China, Hefei 230026, People's Republic of China*

⁷² *University of South China, Hengyang 421001, People's Republic of China*

⁷³ *University of the Punjab, Lahore-54590, Pakistan*

⁷⁴ *University of Turin and INFN, (A)University of Turin, I-10125, Turin, Italy; (B)University of Eastern Piedmont, I-15121, Alessandria, Italy; (C)INFN, I-10125, Turin, Italy*

⁷⁵ *Uppsala University, Box 516, SE-75120 Uppsala, Sweden*

⁷⁶ *Wuhan University, Wuhan 430072, People's Republic of China*

⁷⁷ *Yantai University, Yantai 264005, People's Republic of China*

⁷⁸ *Yunnan University, Kunming 650500, People's Republic of China*

⁷⁹ *Zhejiang University, Hangzhou 310027, People's Republic of China*

⁸⁰ *Zhengzhou University, Zhengzhou 450001, People's Republic of China*

^a *Deceased*

^b *Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia*

^c *Also at the Novosibirsk State University, Novosibirsk, 630090, Russia*

^d *Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia*

^e *Also at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany*

^f *Also at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China*

^g *Also at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China*

^h *Also at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China*

ⁱ *Also at School of Physics and Electronics, Hunan University, Changsha 410082, China*

^j *Also at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China*

^k *Also at MOE Frontiers Science Center for Rare Isotopes,*

Lanzhou University, Lanzhou 730000, People's Republic of China

^l *Also at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People's Republic of China*

^m *Also at the Department of Mathematical Sciences, IBA, Karachi 75270, Pakistan*

ⁿ *Also at Ecole Polytechnique Federale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland*

^o *Also at Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*

^p *Also at School of Physics, Beihang University, Beijing 100191, China*

(Dated: June 24, 2024)

Based on 368.5 pb^{-1} of e^+e^- collision data collected at center-of-mass energies 4.914 and 4.946 GeV by the BESIII detector, the $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ process is searched for the first time. No significant signal is observed and the upper limits at the 90% confidence level on the product of the Born cross section $\sigma(e^+e^- \rightarrow \phi\chi_{c1}(3872))$ and the branching fraction $\mathcal{B}[\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi]$ at 4.914 and 4.946 GeV are set to be 0.85 and 0.96 pb, respectively. These measurements provide useful information for the production of the $\chi_{c1}(3872)$ at e^+e^- collider and deepen our understanding about the nature of this particle.

A series of exotic hadron candidates, which can not be accommodated by the potential model [1–3], were observed experimentally in the charmonium energy region during the past decades. They are suggested as good candidates of molecule, quark-gluon hybrid, or tetraquark states [4]. The well-known $\chi_{c1}(3872)$ state was firstly observed by the Belle experiment in the $B^\pm \rightarrow K^\pm\pi^+\pi^-J/\psi$ decay [5], and confirmed subsequently by several other experiments [6–11]. Ten years after its discovery, its spin-parity quantum numbers were finally determined to be $J^{PC} = 1^{++}$ by the LHCb Collaboration [12]. The mass and width are determined to be $M = 3871.65 \pm 0.06 \text{ MeV}/c^2$ and $\Gamma = 1.19 \pm 0.21 \text{ MeV}$

using a Breit-Wigner resonance model [13].

Since the discovery of the $\chi_{c1}(3872)$, there have been tremendous efforts to understand its inner structure. Experimentally, there are intensive studies on the $\chi_{c1}(3872)$ decays currently. The decays of $\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi$ [5, 11, 14], $\gamma J/\psi$ [15–18], $\pi^0\chi_{c1}$ [19], $\omega J/\psi$ [20, 21], $D^{*0}\bar{D}^0$ [15, 22, 23] have been well observed. Theoretically, $\chi_{c1}(3872)$ is interpreted as a good candidate of a meson molecule [24, 25] as its mass is quite near $D^{*0}\bar{D}^0$ mass threshold. On the other hand, its quantum number is 1^{++} . So far, the P -wave excited charmonium state $\chi_{c1}(2P)$ (with $J^{PC} = 1^{++}$) is still missing, and the $\chi_{c1}(3872)$ might be a good candidate for $\chi_{c1}(2P)$ due

to the similar mass as that from potential model prediction [1]. The other interpretations such as a tetraquark candidate [26, 27] is also possible. However, these is still no solid conclusion for the nature of $\chi_{c1}(3872)$.

In complement with decay, the production of $\chi_{c1}(3872)$ offers a new window to understand its nature. In 2014, the BESIII experiment observed the $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$ production [11]. What intriguing is that the $\chi_{c1}(3872)$ might originate from the radiative transition of an excited vector state $Y(4230)$, which for the first time brings together two charmonium-like states and hints commonality for their underlying nature [4]. Recently, the process $e^+e^- \rightarrow \omega\chi_{c1}(3872)$ is observed at BESIII [28], and the production cross section shows potential enhancement near 4.75 GeV. In analogy to γ and ω , the vector meson ϕ has the same J^{PC} and isospin quantum number. Therefore, the process of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ is expected to naturally exist. By investigating the relative production ratio $\sigma_{e^+e^- \rightarrow \phi\chi_{c1}(3872)}/\sigma_{e^+e^- \rightarrow \phi\chi_{c1}}$ [29] and also comparing to $\sigma_{e^+e^- \rightarrow \omega\chi_{c1}(3872)}/\sigma_{e^+e^- \rightarrow \omega\chi_{c1}}$ [28, 30], we gain a deeper understanding of the $\chi_{c1}(3872)$ production and probe the potential $\chi_{c1}(2P)$ core component in the $\chi_{c1}(3872)$ wave function [31]. In addition, the decay $B_s^0 \rightarrow \phi\chi_{c1}(3872)$ has also been observed [32] and has a production rate $\mathcal{B}[B_s^0 \rightarrow \phi\chi_{c1}(3872)] \approx \mathcal{B}[B^0 \rightarrow K^0\chi_{c1}(3872)] \approx \frac{1}{2}\mathcal{B}[B^+ \rightarrow K^+\chi_{c1}(3872)]$ [13]. Together with the $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ process, a more comprehensive understanding of $\chi_{c1}(3872)$ production will be achieved and further to deeply explore its nature [33].

In this article, we search for the $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ process using 368.5 pb^{-1} of data [34] collected with the BESIII detector [35] operated at the BEPCII storage ring [36]. The ϕ meson is reconstructed via K^+K^- decay, while the $\chi_{c1}(3872)$ via $\rho^0 J/\psi$ decay with $\rho^0 \rightarrow \pi^+\pi^-$ and $J/\psi \rightarrow \ell^+\ell^-$ ($\ell = e, \mu$ with close branching fraction [13]). Due to the mass threshold of the $\phi\chi_{c1}(3872)$ system, only the data at the e^+e^- center-of-mass (c.m.) energies $\sqrt{s} = 4.914$ and 4.946 GeV is used.

The BESIII detector [35] records symmetric e^+e^- collisions provided by the BEPCII storage ring [36] in the center-of-mass energy range from 2.0 to 4.95 GeV, with a peak luminosity of $1 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$ achieved at $\sqrt{s} = 3.77$ GeV. BESIII has collected large data samples in this energy region [37, 38]. The cylindrical core of the BESIII detector covers 93% of the full solid angle and consists of a helium-based multilayer drift chamber (MDC), a plastic scintillator time-of-flight system (TOF), and a CsI(Tl) electromagnetic calorimeter (EMC), which are all enclosed in a superconducting solenoidal magnet providing a 1.0 T magnetic field. A muon chamber (MUC) based on resistive plate chambers with 2 cm position resolution provides information for muon identification. The acceptance of charged particles and photons is 93% over 4π solid angle. The charged-particle momentum resolution at 1 GeV/c is 0.5%, and the dE/dx resolution is 6% for electrons from Bhabha scattering. The EMC measures photon energies with a resolution of 2.5% (5%) at

1 GeV in the barrel (end cap) region. The time resolution in the TOF barrel region is 68 ps, while that in the end cap region is 60 ps [39–41].

Monte Carlo (MC) samples, simulated using GEANT4-based software, are used to optimize the selection criteria, determine the detection efficiency and study the potential backgrounds[42]. In the BESIII software framework, KKMC [43] is the generator used to generate charmonium states by including Initial State Radiation (ISR) effects and the spread of the beam energy. We generate 50000 signal MC events of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ at each c.m. energy with a phase space (PHSP) model describing the uniform angular distribution of the final states. The $\chi_{c1}(3872) \rightarrow \rho^0 J/\psi$ decay is described with the PHSP model. Final state radiation of charged particles are simulated with the PHOTOS package [44]. For the possible ISR effect, we model the \sqrt{s} -dependent $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ production cross section with a two-body phase space $(\frac{\sqrt{[s-(M_\phi+M_{\chi_{c1}(3872)})^2] [s-(M_\phi-M_{\chi_{c1}(3872)})^2]}}{2\sqrt{s}})$, M_ϕ , $M_{\chi_{c1}(3872)}$ is the mass of ϕ , $\chi_{c1}(3872)$ [13]. Inclusive MC samples, with a luminosity which are ten times larger than the data sample, are generated at each c.m. energy to study the possible backgrounds. The inclusive MC sample includes the production of open charm processes, the ISR production of vector charmonium(-like) states, and the continuum processes incorporated in KKMC. All particle decays are modelled with EVTGEN [45, 46] using branching fractions taken from the Particle Data Group [13], when available, and the remaining unknown charmonium decays are modelled with LUNDCHARM [47, 48].

Charged tracks detected in the MDC are required to be within a polar angle (θ) range of $|\cos\theta| < 0.93$, where θ is defined with respect to the z -axis, which is the symmetry axis of the MDC. For charged tracks, the distance of closest approach to the interaction point (IP) must be less than 10 cm along the z -axis, $|V_z|$, and less than 1 cm in the transverse plane, $|V_{xy}|$. For each candidate event, the pions from $\chi_{c1}(3872)$ decay (kaons from ϕ decay) and the leptons from J/ψ decay are kinematically well separated. Charged tracks with momenta larger than 1.0 GeV/c in the lab frame are assumed to be leptons, and the others are assumed to be pions or kaons. The energy deposited in the EMC is utilized to separate electrons from muons. For both muon candidates, the deposited energy in the EMC must be less than 0.4 GeV; while for both electrons, it must be larger than 0.8 GeV. To separate pions from kaons, particle identification (PID), which combines measurements of the specific ionization energy loss in the MDC (dE/dx) and the flight time in the TOF to form likelihoods $\mathcal{L}(h)$ ($h = p, K, \pi$) for each hadron h hypothesis is used. Charged kaons (pions) are identified by comparing the likelihoods for the kaon (pion) hypotheses with $\mathcal{L}(K) > \mathcal{L}(\pi)$ ($\mathcal{L}(\pi) > \mathcal{L}(K)$).

For the candidate events with $K^+K^-\pi^+\pi^-\ell^+\ell^-$ detected, referred to as 6-track events, the net charge is

required to be zero. A four-constraint (4C) kinematic fit imposing energy-momentum conservation is applied on the 6-track events to improve resolution and suppress backgrounds. The kinematic fit χ_{4C}^2 is required to be less than 150. The selection criteria are optimized by maximizing the figure-of-merit (FOM) $\epsilon_{\text{sig}}/(\alpha/2 + \sqrt{N_{\text{bkg}}})$, where ϵ_{sig} is the detection efficiency from signal MC events, α is the assumed significance value which is set to 3 and N_{bkg} is the expected number of background events obtained from inclusive MC samples.

To select the J/ψ resonance, a mass window is defined as $[3.070, 3.125]$ GeV/c^2 (mass resolution is 6 MeV/c^2), which roughly covers about $\pm 3\sigma$ of the J/ψ signal. The signal window of ϕ resonance is set as $[0.980, 1.080]$ GeV/c^2 (mass resolution is 7 MeV/c^2) with barely efficiency loss according to signal MC events. To estimate the non- J/ψ background, the J/ψ sideband regions are defined as $[3.010, 3.065]$ and $[3.130, 3.185]$ GeV/c^2 , which are twice as wide as the J/ψ signal region. Due to the constraint of the mass threshold of double kaons, the ϕ sideband is defined as $[1.080, 1.180]$ GeV/c^2 , which is same as wide as the ϕ signal region. After applying all the selection criteria, there is no background in the 6-track events, as indicated by inclusive MC sample. Figure 1 shows the invariant mass distributions for 6 track events from the full data set.




FIG. 1. The distributions of $M(\ell^+\ell^-)$ (a) and $M(K^+K^-)$ (b) for 6-track events. Dots with error bars are 4.914 and 4.946 GeV data, the red histograms are signal MC sample, and the blue filled histograms are inclusive MC sample.

In order to further improve the detection efficiency and thus the signal yield, candidate events with $K^+\pi^+\pi^-\ell^+\ell^-$ or $K^-\pi^+\pi^-\ell^+\ell^-$ detected are also reconstructed, with one of the soft kaons missing due to an inefficiency. A one-constraint (1C) kinematic fit is performed to the 5-track events, by constraining the mass of missing particle to the nominal mass of the kaon [13]. The kinematic fit χ_{1C}^2 is required to be less than 20, which is optimized by maximizing $\text{FOM} = \epsilon_{\text{sig}}/(\alpha/2 + \sqrt{N_{\text{bkg}}})$. Due to the absence of one track in this case, the mass resolution of the event is slightly worse compared to that of the 6-track events. So the J/ψ mass window is defined as $[3.065, 3.130]$ GeV/c^2 (mass resolution is 10 MeV/c^2) and the sideband regions are $[3.000, 3.065]$ and $[3.130, 3.195]$ GeV/c^2 for the 5-track events, which are

twice as wide as the J/ψ signal region. The signal region of ϕ resonance is set to be $[0.980, 1.080]$ GeV/c^2 (mass resolution is 8 MeV/c^2) and the ϕ sideband is defined as $[1.080, 1.180]$ GeV/c^2 , which is as wide as the ϕ signal region. There is a case, about 10% of the total signal events, that $K^\pm\pi^\mp\pi^+\pi^-\ell^+\ell^-$ are detected in the final state, owing to one of the soft kaon decays in the detector. This case is classified as 5-track events and reconstructed by missing a kaon.

By analyzing the inclusive MC samples, we find there are some proton backgrounds remaining in the 5-track events. To reduce the $p \rightarrow \mu$ misidentification background in the $J/\psi \rightarrow \mu^+\mu^-$ channel, the MUC is used to identify muons. At least one of muon candidate should have a hit depth > 30 cm in the MUC. Figure 2 shows the $M(\ell^+\ell^-)$ and $RM(\pi^+\pi^-\ell^+\ell^-)$ distributions for 5-track events from the full data samples, where $RM(\pi^+\pi^-\ell^+\ell^-) = \sqrt{(P_{e^+e^-} - P_{\pi^+\pi^-\ell^+\ell^-})^2}$ is the recoil mass from the $\pi^+\pi^-\ell^+\ell^-$ system, $P_{e^+e^-}$ and $P_{\pi^+\pi^-\ell^+\ell^-}$ are the four-momenta of the initial colliding beams and the $\pi^+\pi^-\ell^+\ell^-$ system.




FIG. 2. The distributions of $M(\ell^+\ell^-)$ (a) and $RM(\pi^+\pi^-\ell^+\ell^-)$ (b) for 5-track events. Dots with error bars are 4.914 and 4.946 GeV data, the red histograms are signal MC sample, and the blue filled histograms are inclusive MC sample.

After applying all the selection criteria, Fig. 3 shows the $M(\pi^+\pi^-J/\psi)$ distribution from 6-track events and 5-track events at each c.m. energy. The invariant mass $M(\pi^+\pi^-J/\psi) = M(\pi^+\pi^-\ell^+\ell^-) - M(\ell^+\ell^-) + M(J/\psi)$ is defined, which partly helps cancel the resolution effect of lepton pairs. Here $M(J/\psi)$ is the nominal mass of J/ψ from the PDG [13]. Through studying the inclusive MC samples at each c.m. energy, no dominant background survives for both 6-track events and 5-tracks events. The study of the J/ψ and ϕ mass sideband events also shows the background level is low and we neglect the background.

The product of the Born cross section of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ and $\mathcal{B}[\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi]$ at c.m. energy \sqrt{s} is calculated with

$$\begin{aligned} & \sigma(e^+e^- \rightarrow \phi\chi_{c1}(3872)) \cdot \mathcal{B}[\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi] \\ &= \frac{N_{\text{sig}}}{\mathcal{L}_{\text{int}}(1+\delta) \frac{1}{|1-\Pi|^2} \epsilon \mathcal{B}_{\text{sub}}}, \end{aligned} \quad (4.1)$$




FIG. 3. The distributions of $M(\pi^+\pi^-J/\psi)$ from (a) 6-track events at $\sqrt{s} = 4.914$ GeV, (b) 5-track events at $\sqrt{s} = 4.914$ GeV, (c) 6-track events at $\sqrt{s} = 4.946$ GeV and (d) 5-track events at $\sqrt{s} = 4.946$ GeV, respectively. Dots with error bars are data, the red histograms are signal MC sample, the blue filled histograms are inclusive MC sample and the green filled histograms are the $\phi - J/\psi$ 2-dimensional sideband.

where N_{sig} is the number of signal events, \mathcal{L}_{int} is the integrated luminosity, ϵ is the selection efficiency, $\frac{1}{1-\Pi^2} = 1.056$ is the vacuum polarization factor taken from QED calculation with an accuracy of 0.05% [49], \mathcal{B}_{sub} is a product of the branching fractions $\mathcal{B}(J/\psi \rightarrow \ell^+\ell^-)$ and $\mathcal{B}(\phi \rightarrow K^+K^-)$, and $(1 + \delta)$ is the radiative correction factor calculated by KKMC [43].

Due to no event observed in the $\chi_{c1}(3872)$ signal region, we report an upper limit on the Born cross section by combining the 5-track events and 6-track events using a counting method. The number of signal events is determined by counting the number of events in the $\chi_{c1}(3872)$ signal region from 5-track events and 6-track events, which is defined as $[3.86, 3.88]\text{GeV}/c^2$. The possible background estimated by the $\chi_{c1}(3872)$ sideband regions $[3.80, 3.85]$ and $[3.90, 3.95]\text{GeV}/c^2$ from 5-track events and 6-track events is subtracted (at $\sqrt{s} = 4.914$ GeV the sideband region is defined as $[3.80, 3.85]\text{GeV}/c^2$ due to the limitation of kinematics). The upper limit on the number of signal yield at the 90% confidence level (C.L.) is obtained using a frequentist method with an unbounded profile likelihood treatment [50]. Assuming the background follows a Poisson distribution and the efficiency (sum of 5-track events and 6-track events) follows a Gaussian distribution with a standard deviation equal to the systematic uncertainty, the upper limit on the product of the Born cross section of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ and the branching fraction of $\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi$ at the 90% C.L. at each c.m. energy are measured and listed

in Table I.

TABLE I. The upper limit on the product of Born cross section of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ and the branching fraction of $\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi$ (denoted as σ_B^{up} (pb)) at 90% C.L. at each c.m. energy. \sqrt{s} (GeV) is the c.m. energy, \mathcal{L}_{int} (pb^{-1}) is the integrated luminosity, N_{obs} is the number of observed events in signal region, N_{sdb} is the number of observed events in sideband region, $N_{\text{signal}}^{\text{up}}$ is the upper limit on the number of observed signal at the 90% C.L., ϵ^5 (%) and ϵ^6 (%) are detection efficiencies for the 5- and 6-track events, respectively, $(1 + \delta)$ is the radiative correction factor.

\sqrt{s}	\mathcal{L}_{int}	N_{obs}	N_{sdb}	$N_{\text{signal}}^{\text{up}}$	$(1 + \delta)$	ϵ^5	ϵ^6	σ_B^{up}
4.914	208.11	0	1	1.70	0.690	19.7	2.8	0.85
4.946	160.37	0	0	2.00	0.755	20.8	7.0	0.96

In the cross section measurement, the systematic uncertainties are mainly from the luminosity measurement, branching fractions, tracking efficiency, PID efficiency, MUC hit depth requirement, kinematic fit, radiative correction factor $(1 + \delta)$, MC decay model and J/ψ mass window.

The uncertainty from luminosity measurement is estimated to be less than 0.66% using large angle Bhabha scattering events [34]. The uncertainties of decay branching fractions are quoted from the PDG [13]. The uncertainty of tracking efficiency for high momentum leptons is assigned to be 1% per track according to the study of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at BESIII [51]. In this measurement, both one kaon events and two kaon events are reconstructed. The total uncertainty of tracking efficiency of the kaon is 1.0% for 5-track events and 2.0% for 6-track events. Considering the uncertainty of PID efficiency is 1% per kaon track at BESIII, the kaon PID uncertainty is 1.0% for 5-track events and 2.0% for 6-track events, too. The systematic uncertainty from pion tracking and PID efficiencies are both 1.0% per pion track [52].

The uncertainty of MUC hit depth is studied using control sample of $e^+e^- \rightarrow \mu^+\mu^-$ [29]. The difference in efficiency between data and MC simulation due to the requirement of μ hit depth in the MUC is taken as the systematic uncertainty. A helix parameters correction method is used to estimate the difference between data and signal MC events caused by kinematic fit. The difference in efficiency with and without correction is taken as the systematic uncertainty. The systematic uncertainty of radiative correction factor is studied by comparing the difference between factors obtained with the two-body phase space model and with a flat cross section line shape. The difference in $(1 + \delta)\epsilon$ is taken as the uncertainty. To estimate the uncertainty due to the MC model, the angular distribution of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ is modelled by a $1 \pm \cos^2(\theta)$ distribution, the efficiency difference with respect to PHSP is taken as the systematic uncertainty. The control sample $e^+e^- \rightarrow \pi^+\pi^-\psi(3686)$ with $\psi(3686) \rightarrow \pi^+\pi^-J/\psi$ [53] is selected to study the system

uncertainty caused by the J/ψ mass window. The difference in efficiency between data and signal MC events due to the mass window is taken as the systematic uncertainty.

In the measurement, two sub-data samples, i.e. the 5-track events and 6-track events are reconstructed. The same source of systematic uncertainty contribute differently to two sub-data samples. To combine the systematic uncertainty, we take the weighted average of the systematic uncertainties in two sub-data samples, which follows:

$$\sigma_{\text{tot}}^2 = \sum_{i=1}^2 \omega_i^2 \sigma_i^2 + 2 \sum_{i < j} \text{cov}(i, j), \quad (5.1)$$

$$\text{cov}(i, j) = \omega_i \omega_j \sigma_i \sigma_j, \quad (5.2)$$

$$\omega_i = \frac{\epsilon_i}{\sum_{i=1}^2 \epsilon_i}, \quad (5.3)$$

where σ_{tot} is the average systematic uncertainty on the cross section, ω_i and σ_i are the weight and systematic uncertainty for the i th sub-data sample, ϵ_i is the efficiency for the i th sub-data sample. For the soft kaon decay events ($K^\pm \pi^\mp \pi^+ \pi^- \ell^+ \ell^-$), its weight contributes to 5-track sub-data sample for J/ψ mass window, MUC hit depth requirement, kinematic fit; otherwise its weight contributes to 6-track sub-data sample.

Assuming all these sources are independent, the total systematic uncertainty in the cross section measurement is obtained by adding them in quadrature. Table II and Table III summarize all the systematic sources and their contributions at 4.914 GeV and 4.946 GeV, respectively.

TABLE II. Systematic uncertainties (%) in the measurement of the Born cross section at 4.914 GeV.

Uncertainty	5 Tracks	6 Tracks	Weighted average
Luminosity	0.7		0.7
Tracking	5.0	6.0	5.2
PID	3.0	4.0	3.2
$\mathcal{B}(J/\psi \rightarrow \ell^+ \ell^-)$	0.6		0.6
$\mathcal{B}(\phi \rightarrow K^+ K^-)$	1.0		1.0
Radiative correction	0.7		0.7
J/ψ mass window	0.1	0.1	0.1
MUC hit depth	2.1	0	1.8
Kinematic fit	0.7	0.4	0.7
MC model	5.2	5.9	5.4
Total	-	-	8.5

In summary, with a data sample corresponding to an integrated luminosity of 368.5 pb^{-1} collected by the

TABLE III. Systematic uncertainties (%) in the measurement of the Born cross section at 4.946 GeV.

Uncertainty	5 Tracks	6 Tracks	Weighted average
Luminosity		0.7	0.7
Tracking	5.0	6.0	5.3
PID	3.0	4.0	3.3
$\mathcal{B}(J/\psi \rightarrow \ell^+ \ell^-)$		0.6	0.6
$\mathcal{B}(\phi \rightarrow K^+ K^-)$		1.0	1.0
Radiative correction		1.9	1.9
J/ψ mass window	0.1	0.1	0.1
MUC hit depth	2.9	0	2.2
Kinematic fit	0.6	0.3	0.5
MC model	2.1	14.0	6.2
Total	-	-	9.4

BESIII detector, the process of $e^+e^- \rightarrow \phi\chi_{c1}(3872)$ is searched for the first time. No significant signal is observed and the upper limits at the 90% C.L. on the product of the Born cross section $\sigma(e^+e^- \rightarrow \phi\chi_{c1}(3872))$ and the branching fraction $\mathcal{B}[\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi]$ at 4.914 and 4.946 GeV are set to be 0.85 and 0.96 pb, respectively. Considering the cross section $\sigma(e^+e^- \rightarrow \phi\chi_{c1}) \sim 2.6 \text{ pb}$ near the production threshold [29], we obtain a rough estimation for the production ratio $\sigma_{\phi\chi_{c1}(3872)}/\sigma_{\phi\chi_{c1}} < 9$. It is in the same order as the relative production ratio $\sigma_{\omega\chi_{c1}(3872)}/\sigma_{\omega\chi_{c1}} \sim 5$ [28, 30]. These measurements provide important inputs to the production of $\chi_{c1}(3872)$ at e^+e^- collider, and help constrain the possible $\chi_{c1}(2P)$ component in the $\chi_{c1}(3872)$ wave function. With the upgrade of the BEPCII [37] project, more data in this energy region is expected and a more comprehensive study of the $\chi_{c1}(3872)$ production will be achieved, which finally help reveal the nature of $\chi_{c1}(3872)$.

ACKNOWLEDGMENTS

The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts Nos. 2020YFA0406300, 2020YFA0406400, 2023YFA1606000; National Natural Science Foundation of China (NSFC) under Contracts Nos. 11635010, 11735014, 11835012, 11935015, 11935016, 11935018, 11961141012, 12025502, 12035009, 12035013, 12061131003, 12192260, 12192261, 12192262, 12192263, 12192264, 12192265, 12221005, 12225509, 12235017; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and

CAS under Contract No. U1832207; CAS Key Research Program of Frontier Sciences under Contracts Nos. QYZDJ-SSW-SLH003, QYZDJ-SSW-SLH040; 100 Talents Program of CAS; Project No. ZR2022JQ02 supported by Shandong Provincial Natural Science Foundation; supported by the China Postdoctoral Science Foundation under Grant No. 2023M742100; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement under Contract No. 894790; German Research Foundation DFG under Contracts Nos. 455635585, Collaborative Research Center CRC 1044, FOR5327, GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia; National Science Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation of Thailand under Contract No. B16F640076; Polish National Science Centre under Contract No. 2019/35/O/ST2/02907; The Swedish Research Council; U. S. Department of Energy under Contract No. DE-FG02-05ER41374.

- [1] E. Eichten *et al.*, “Charmonium: The Model,” *Phys. Rev. D* **17**, 3090 (1978).
- [2] E. Eichten *et al.*, “Charmonium: Comparison with Experiment,” *Phys. Rev. D* **21**, 203 (1980).
- [3] S. Godfrey and N. Isgur, “Mesons in a Relativized Quark Model with Chromodynamics,” *Phys. Rev. D* **32**, 189–231 (1985).
- [4] S. L. Olsen, T. Skwarnicki, and D. Zieminska, “Nonstandard heavy mesons and baryons: Experimental evidence,” *Rev. Mod. Phys.* **90**, 015003 (2018), [arXiv:1708.04012 \[hep-ph\]](#).
- [5] S. K. Choi *et al.* (Belle Collaboration), “Observation of a narrow charmonium-like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays,” *Phys. Rev. Lett.* **91**, 262001 (2003), [arXiv:hep-ex/0309032](#).
- [6] D. Acosta *et al.* (CDF Collaboration), “Observation of the narrow state $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV,” *Phys. Rev. Lett.* **93**, 072001 (2004), [arXiv:hep-ex/0312021](#).
- [7] V. M. Abazov *et al.* (D0 Collaboration), “Observation and properties of the $X(3872)$ decaying to $J/\psi \pi^+ \pi^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV,” *Phys. Rev. Lett.* **93**, 162002 (2004), [arXiv:hep-ex/0405004](#).
- [8] B. Aubert *et al.* (BaBar Collaboration), “Study of the $B \rightarrow J/\psi K^- \pi^+ \pi^-$ decay and measurement of the $B \rightarrow X(3872) K^-$ branching fraction,” *Phys. Rev. D* **71**, 071103 (2005), [arXiv:hep-ex/0406022](#).
- [9] R. Aaij *et al.* (LHCb Collaboration), “Observation of $X(3872)$ production in pp collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **72**, 1972 (2012), [arXiv:1112.5310 \[hep-ex\]](#).
- [10] S. Chatrchyan *et al.* (CMS Collaboration), “Measurement of the $X(3872)$ Production Cross Section Via Decays to $J/\psi \pi^+ \pi^-$ in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **04**, 154 (2013), [arXiv:1302.3968 \[hep-ex\]](#).
- [11] M. Ablikim *et al.* (BESIII Collaboration), “Observation of $e^+e^- \rightarrow \gamma X(3872)$ at BESIII,” *Phys. Rev. Lett.* **112**, 092001 (2014), [arXiv:1310.4101 \[hep-ex\]](#).
- [12] R. Aaij *et al.* (LHCb Collaboration), “Determination of the $X(3872)$ meson quantum numbers,” *Phys. Rev. Lett.* **110**, 222001 (2013), [arXiv:1302.6269 \[hep-ex\]](#).
- [13] R. L. Workman *et al.* (Particle Data Group), “Review of Particle Physics,” *PTEP* **2022**, 083C01 (2022).
- [14] R. Aaij *et al.* (LHCb Collaboration), “Study of the $\psi_2(3823)$ and $\chi_{c1}(3872)$ states in $B^+ \rightarrow (J/\psi \pi^+ \pi^-) K^+$ decays,” *JHEP* **08**, 123 (2020), [arXiv:2005.13422 \[hep-ex\]](#).
- [15] M. Ablikim *et al.* (BESIII Collaboration), “Study of Open-Charm Decays and Radiative Transitions of the $X(3872)$,” *Phys. Rev. Lett.* **124**, 242001 (2020), [arXiv:2001.01156 \[hep-ex\]](#).
- [16] R. Aaij *et al.* (LHCb Collaboration), “Evidence for the decay $X(3872) \rightarrow \psi(2S)\gamma$,” *Nucl. Phys. B* **886**, 665–680 (2014), [arXiv:1404.0275 \[hep-ex\]](#).
- [17] B. Aubert *et al.* (BaBar Collaboration), “Search for $B^+ \rightarrow X(3872)K^+$, $X_{3872} \rightarrow J/\psi\gamma$,” *Phys. Rev. D* **74**, 071101 (2006), [arXiv:hep-ex/0607050](#).
- [18] V. Bhardwaj *et al.* (Belle Collaboration), “Observation of $X(3872) \rightarrow J/\psi\gamma$ and search for $X(3872) \rightarrow \psi'\gamma$ in B decays,” *Phys. Rev. Lett.* **107**, 091803 (2011), [arXiv:1105.0177 \[hep-ex\]](#).
- [19] M. Ablikim *et al.* (BESIII Collaboration), “Observation of the decay $X(3872) \rightarrow \pi^0 \chi_{c1}(1P)$,” *Phys. Rev. Lett.* **122**, 202001 (2019), [arXiv:1901.03992 \[hep-ex\]](#).
- [20] M. Ablikim *et al.* (BESIII Collaboration), “Study of $e^+e^- \rightarrow \gamma\omega J/\psi$ and Observation of $X(3872) \rightarrow \omega J/\psi$,” *Phys. Rev. Lett.* **122**, 232002 (2019), [arXiv:1903.04695 \[hep-ex\]](#).
- [21] P. del Amo Sanchez *et al.* (BaBar Collaboration), “Evidence for the decay $X(3872) \rightarrow J/\psi\omega$,” *Phys. Rev. D* **82**, 011101 (2010), [arXiv:1005.5190 \[hep-ex\]](#).
- [22] B. Aubert *et al.* (BaBar Collaboration), “Study of Resonances in Exclusive B Decays to anti-D(*) D(*) K,” *Phys. Rev. D* **77**, 011102 (2008), [arXiv:0708.1565 \[hep-ex\]](#).
- [23] T. Aushev *et al.* (Belle Collaboration), “Study of the $B \rightarrow X(3872)(\rightarrow D^{*0}\bar{D}^0)K$ decay,” *Phys. Rev. D* **81**, 031103 (2010), [arXiv:0810.0358 \[hep-ex\]](#).
- [24] E. S. Swanson, “The New heavy mesons: A Status report,” *Phys. Rept.* **429**, 243–305 (2006), [arXiv:hep-ph/0601110](#).
- [25] F. K. Guo *et al.*, “Hadronic molecules,” *Rev. Mod. Phys.* **90**, 015004 (2018), [arXiv:1705.00141 \[hep-ph\]](#).
- [26] L. Maiani *et al.*, “Diquark-antidiquarks with hidden or open charm and the nature of $X(3872)$,” *Phys. Rev. D* **71**, 014028 (2005), [arXiv:hep-ph/0412098](#).
- [27] A. Esposito *et al.*, “Four-Quark Hadrons: an Updated Review,” *Int. J. Mod. Phys. A* **30**, 1530002 (2015), [arXiv:1411.5997 \[hep-ph\]](#).
- [28] M. Ablikim *et al.* (BESIII Collaboration), “Observation of a New $X(3872)$ Production Process $e^+e^- \rightarrow \omega X(3872)$,” *Phys. Rev. Lett.* **130**, 151904 (2023), [arXiv:2212.07291 \[hep-ex\]](#).
- [29] M. Ablikim *et al.* (BESIII Collaboration), “Study of $e^+e^- \rightarrow \gamma\phi J/\psi$ from $\sqrt{s} = 4.600$ to 4.951 GeV,” *JHEP* **01**, 132 (2023), [arXiv:2210.13058 \[hep-ex\]](#).
- [30] M. Ablikim *et al.* (BESIII Collaboration), “Study of $e^+e^- \rightarrow \omega X(3872)$ and $\gamma X(3872)$ from 4.66 to 4.95 GeV,” [arXiv:2404.13840 \[hep-ex\]](#).
- [31] M. Takizawa and S. Takeuchi, “ $X(3872)$ as a hybrid state of charmonium and the hadronic molecule,” *PTEP* **2013**, 093D01 (2013), [arXiv:1206.4877 \[hep-ph\]](#).
- [32] A. M. Sirunyan *et al.* (CMS Collaboration), “Observation of the $B_s^0 \rightarrow X(3872)\phi$ decay,” *Phys. Rev. Lett.* **125**, 152001 (2020), [arXiv:2005.04764 \[hep-ex\]](#).
- [33] L. Maiani, Antonio D. Polosa, and V. Riquer, “ $X(3872)$ tetraquarks in B and B_s decays,” *Phys. Rev. D* **102**, 034017 (2020), [arXiv:2005.08764 \[hep-ph\]](#).
- [34] M. Ablikim *et al.* (BESIII Collaboration), “Luminosities and energies of e^+e^- collision data taken between 4.61 GeV and 4.95 GeV

- at BESIII*,” *Chin. Phys. C* **46**, 113003 (2022), [arXiv:2205.04809 \[hep-ex\]](#).
- [35] M. Ablikim *et al.* (BESIII Collaboration), “Design and Construction of the BESIII Detector,” *Nucl. Instrum. Meth. A* **614**, 345–399 (2010), [arXiv:0911.4960 \[physics.ins-det\]](#).
- [36] C. H. Yu *et al.*, “BEPCh Performance and Beam Dynamics Studies on Luminosity,” in *7th International Particle Accelerator Conference* (2016) p. TUYA01.
- [37] M. Ablikim *et al.* (BESIII Collaboration), “Future Physics Programme of BESIII,” *Chin. Phys. C* **44**, 040001 (2020), [arXiv:1912.05983 \[hep-ex\]](#).
- [38] J. D. Lu, Y. J. Xiao, and X. B. Ji, “Online monitoring of the center-of-mass energy from real data at BESIII,” *Radiat. Detect. Technol. Methods* **4**, 337–344 (2020).
- [39] X. Li *et al.*, “Study of MRPC technology for BESIII endcap-TOF upgrade,” *Radiat. Detect. Technol. Methods* **1**, 13 (2017).
- [40] Y. X. Guo *et al.*, “The study of time calibration for upgraded end cap TOF of BESIII,” *Radiat. Detect. Technol. Methods* **1**, 15 (2017).
- [41] P. Cao *et al.*, “Design and construction of the new BESIII endcap Time-of-Flight system with MRPC Technology,” *Nucl. Instrum. Meth. A* **953**, 163053 (2020).
- [42] S. Agostinelli *et al.* (GEANT4), “GEANT4—a simulation toolkit,” *Nucl. Instrum. Meth. A* **506**, 250–303 (2003).
- [43] S. Jadach, B. F. L. Ward, and Z. Was, “Coherent exclusive exponentiation for precision Monte Carlo calculations,” *Phys. Rev. D* **63**, 113009 (2001), [arXiv:hep-ph/0006359](#).
- [44] P. Golonka and Z. Was, “PHOTOS Monte Carlo: A Precision tool for QED corrections in Z and W decays,” *Eur. Phys. J. C* **45**, 97–107 (2006), [arXiv:hep-ph/0506026](#).
- [45] D. J. Lange, “The EvtGen particle decay simulation package,” *Nucl. Instrum. Meth. A* **462**, 152–155 (2001).
- [46] R. G. Ping, “Event generators at BESIII,” *Chin. Phys. C* **32**, 599 (2008).
- [47] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang, and Y. S. Zhu, “Event generator for J/ψ and $\psi(2S)$ decay,” *Phys. Rev. D* **62**, 034003 (2000).
- [48] R. L. Yang, R. G. Ping, and H. Chen, “Tuning and Validation of the Lundcharm Model with J/ψ Decays,” *Chin. Phys. Lett.* **31**, 061301 (2014).
- [49] S. Actis *et al.* (Working Group on Radiative Corrections, Monte Carlo Generators for Low Energies), “Quest for precision in hadronic cross sections at low energy: Monte Carlo tools vs. experimental data,” *Eur. Phys. J. C* **66**, 585–686 (2010), [arXiv:0912.0749 \[hep-ph\]](#).
- [50] W. A. Rolke, A. M. Lopez, and J. Conrad, “Limits and confidence intervals in the presence of nuisance parameters,” *Nucl. Instrum. Meth. A* **551**, 493–503 (2005), [arXiv:physics/0403059](#).
- [51] M. Ablikim *et al.* (BESIII Collaboration), “Precise measurement of the $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ cross section at center-of-mass energies from 3.77 to 4.60 GeV,” *Phys. Rev. Lett.* **118**, 092001 (2017), [arXiv:1611.01317 \[hep-ex\]](#).
- [52] M. Ablikim *et al.* (BESIII Collaboration), “Observation of $\eta' \rightarrow \pi^+\pi^-\pi^+\pi^-$ and $\eta' \rightarrow \pi^+\pi^-\pi^0\pi^0$,” *Phys. Rev. Lett.* **112**, 251801 (2014), [arXiv:1404.0096 \[hep-ex\]](#).
- [53] M. Ablikim *et al.* (BESIII Collaboration), “Cross section measurement of $e^+e^- \rightarrow \pi^+\pi^-$ (3686) from $\sqrt{S} = 4.0076$ to 4.6984 GeV,” *Phys. Rev. D* **104**, 052012 (2021), [arXiv:2107.09210 \[hep-ex\]](#).