

QUANTITATIVE POINTWISE ESTIMATES OF THE COOLING PROCESS FOR INELASTIC BOLTZMANN EQUATION

GAYOUNG AN, JIN WOO JANG, AND DONGHYUN LEE

ABSTRACT. In this paper, we study the homogeneous inelastic Boltzmann equation for hard spheres. We first prove that the solution $f(t, v)$ is bounded pointwise from above by $C_{f_0} \langle t \rangle^3$ and establish that the cooling time is infinite ($T_c = +\infty$) under the condition $f_0 \in L_2^1 \cap L_s^\infty$ for $s > 2$. Away from zero velocity, we further prove that $f(t, v) \leq C_{f_0, |v|} \langle t \rangle$ for $v \neq 0$ at any time $t > 0$. This time-dependent pointwise upper bound is natural in the cooling process, as we expect the density near $v = 0$ to grow rapidly. We also establish an upper bound that depends on the coefficient of normal restitution constant, $\alpha \in (0, 1]$. This upper bound becomes constant when $\alpha = 1$, restoring the known upper bound for elastic collisions [8]. Consequently, through these results, we obtain Maxwellian upper bounds on the solutions at each time.

CONTENTS

1. Introduction	1
1.1. Notations	5
1.2. Known results for the elastic collisions	6
1.3. Known results for the inelastic collisions	7
1.4. Preliminaries	8
1.5. Main results	12
1.6. Outline of the rest of the article	14
2. L^∞ estimates for the inelastic Boltzmann equation	15
3. Maxwellian upper bound for the inelastic Boltzmann equation	24
Acknowledgment	29
References	30

1. INTRODUCTION

The classical Boltzmann equation describes the behavior of a rarefied collisional gas, consisting of a very large number of identical particles undergoing perfectly elastic binary collisions. The velocity distribution function $f(t, x, v)$ is the probable number density of particles at time t , at point x , having speed v . The Boltzmann equation is written as

$$\partial_t f + v \cdot \nabla_x f = Q(f, f), \quad f(0, x, v) = f_0(x, v)$$

for $x \in \Omega \subset \mathbb{R}^3$, $v \in \mathbb{R}^3$ and $t \geq 0$. On the left-hand side of the Boltzmann equation we see a transport operator which describes the translation of a particle distribution without the presence of other forces and collisions. The nonlinear quadratic term, $Q(f, f)$, on the right-hand side is added to account for the collisions




FIGURE 1. Given v, v_* , possible locations of v', v'_* in elastic collisions.

between particles. When two particles, initially traveling at velocities v and v_* , collide with each other, their velocities change due to the interaction. After the collision, the new velocities of the particles are denoted as v' and v'_* , respectively. These post-collisional velocities v' and v'_* are determined by the specific dynamics of the collision process, which can depend on factors such as the masses of the particles and the nature of the interaction between them. We assume that the mass of each particle is identical and normalize it to be 1 throughout the paper. Under elastic collisions, according to the laws of conservation of momentum and energy, the following relationships hold:

$$v + v_* = v' + v'_*, \quad |v|^2 + |v_*|^2 = |v'|^2 + |v'_*|^2. \quad (1.1)$$

From these conservation laws, we derive four constraints on the six variables v' and v'_* , leaving two degrees of freedom. Specifically, the post-collisional velocities v' and v'_* can be expressed as:

$$v' = v - ((v - v_*) \cdot n)n \quad \text{and} \quad v'_* = v_* + ((v - v_*) \cdot n)n, \quad (1.2)$$

using the vector $n \in \mathbb{S}_+^2$. Physically, if gas particles are represented as billiard balls, the vector $n \in \mathbb{S}_+^2$, known as the impact direction, denotes the unit vector connecting the centers of the balls at the moment of collision, as illustrated in Figure 2.

In this paper, we frequently use an alternative representation of the post-collisional velocities v' and v'_* involving another angular variable $\sigma \in \mathbb{S}^2$. This representation is given by:

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2}\sigma \quad \text{and} \quad v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2}\sigma.$$

In Figure 1, we illustrate the possible locations of v' and v'_* for given v and v_* . As shown in the figure, the polar angle of σ along the z -axis, corresponding to $v - v_*$, varies from 0 to π , while the polar angle of n ranges from 0 to $\frac{\pi}{2}$, placing n in \mathbb{S}_+^2 . Interestingly, for fixed v and v_* , v' and v'_* always lie on a single sphere, and the Jacobian determinant between (v, v_*) and (v', v'_*) is always equal to 1. Next, we define the collision operator using the variables v, v_*, v', v'_* , and n as described in

(1.2). It is defined as follows:

$$\begin{aligned} Q(f, f)(t, v) &= \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} B(|v - v_*|, \cos \theta) (f(t, v')f(t, v'_*) - f(t, v)f(t, v_*)) \, dndv_*, \\ &:= Q^+(f, f)(t, v) - f(t, v)Lf(t, v), \end{aligned} \tag{1.3}$$

where

$$\cos \theta = \left(\frac{v - v_*}{|v - v_*|}, n \right), \quad \theta \in \left[0, \frac{\pi}{2} \right].$$

Here, the operators are local in t and x ; we abuse notation by writing $f(t, x, v) = f(t, v)$. The gain term, $Q^+(f, f)(t, v)$, represents an increase in the density of particles with velocity v due to collisions involving velocities v' and v'_* . Because the equations in (1.1) are time-reversible, if particles with velocities v and v_* collide to become v' and v'_* , then particles with velocities v' and v'_* can collide to revert to v and v_* . The loss term, $f(t, v)Lf(t, v)$, indicates a decrease in the density of particles with velocity v due to collisions with particles of velocity v_* , resulting in their disappearance.

Now, we explain the collision kernel. Depending on the assumptions regarding the interaction potentials (e.g., hard-sphere potential or inverse-power-law potential), the collision kernel can be expressed as:


$$B(|v - v_*|, \cos \theta) = b(\cos \theta)|v - v_*|^\gamma$$

for some constant γ . For instance, when particles collide like billiard balls (hard-sphere potential), we write $B(|v - v_*|, \cos \theta) = |(v - v_*) \cdot n|$. If $\int_{\mathbb{S}_+^2} b(\cos \theta) \, dn < \infty$ or if $b(\cos \theta) \in L^\infty(\mathbb{S}_+^2)$, the collision kernel is called a cut-off kernel (or is with angular cutoff), following Grad's angular cutoff criterion in [34]. Otherwise, it is referred to as a non-cutoff kernel (or is without angular cutoff). In the case of a non-cutoff collision kernel, there is a nonintegrable singularity around zero for θ . (Some authors place the singularity near $\pi/2 - \theta \approx 0$.) Recently, the vanishing angular singularity limit from the inverse-power-law potential to the hard-sphere potential has been studied in [45].

By the famous Boltzmann H-theorem, we formally see that the total entropy of the system $-\iint f \log f \, dx \, dv$ will monotonically increase in time. Then we expect that the distribution will converge (in some sense) to an entropy maximizing state. One of the candidates for the states of maximum entropy is so-called the (local) Maxwellian distribution which is given by

$$\mathcal{M}(t, x, v) = \rho(t, x) \frac{e^{-\frac{|v-u(t,x)|^2}{2T(t,x)}}}{(2\pi T(t,x))^{3/2}},$$

where the Boltzmann constant k_B is assumed to be 1. Here, ρ , u , and T correspond to the physical terms such as mass density, macroscopic fluid velocity, and local temperature, respectively. A simple calculation can show that $Q(\mathcal{M}, \mathcal{M})(t, x, v) = 0$. If ρ , u , and T are chosen so that the left-hand side of the Boltzmann equation satisfies $(\partial_t + v \cdot \nabla_x)\mathcal{M} = 0$, (ρ, u, T) solves some macroscopic conservation laws. In this case, the local Maxwellian \mathcal{M} can be written in some specific forms depending on the domain and boundary conditions. We refer [25] for specific forms. In the paper, Desvilletters actually proved that the renormalized solution of the

FIGURE 2. Impact direction, \vec{n} .

Boltzmann equation converges weakly in L^1 to a local time-dependent Maxwellian state in the large-time asymptotics within a bounded domain. It was shown that this local Maxwellian further satisfies $(\partial_t + v \cdot \nabla_x)\mathcal{M} = 0$. Additionally, before this result, Arkeryd proved in [9] the convergence of a renormalized solution to the global Maxwellian. The local stability of the global Maxwellian has now been very well-known. See [1, 2, 35, 55] for the proof of local stability nearby equilibrium for both cutoff and noncutoff cases. The existence of global solutions was firstly proved by Carleman [21] in 1933 for the spatially homogeneous case. Then for the spatially inhomogeneous case, the global existence was proved in the perturbative framework [63] and near vacuum [41]. Then the global existence of renormalized solutions to the Boltzmann equation by DiPerna-Lions [27] in 1989 which does not require size restriction on the initial data. This framework has been extended to the noncutoff case by Alexandre and Villani in [3]. The nonlinear energy method was developed to prove the global existence to the global wellposedness of the Vlasov(-Poisson or -Maxwell)-Boltzmann system in [37, 38, 59]. See also [51] for the energy method for the Boltzmann equation. In [40, 60], the local stability of the Maxwellian for the Landau equation was proved based on the construction of the classical solution via the nonlinear energy method developed in [36]. Regarding the global existence and stability in a bounded domain with general physical boundary conditions (e.g., inflow, bounce-back, specular, and diffuse reflection), see [19, 39, 50].

In contrast to an elastic collision, an inelastic collision is a collision in which kinetic energy is not conserved, and this type of gas is referred to as granular gas. Physically, in each inelastic collision, there is some loss of momentum in the impact direction, though the total momentum remains conserved. Let $0 < \alpha \leq 1$ be the coefficient of normal restitution. For the impact direction $n \in \mathbb{S}_+^2$, the following equations hold:

$$\begin{aligned} (v' - v'_*) \cdot n &= -\alpha(v - v_*) \cdot n, \\ (v' - v'_*) - ((v' - v'_*) \cdot n)n &= (v - v_*) - ((v - v_*) \cdot n)n. \end{aligned} \tag{1.4}$$

If $\alpha = 1$, the collision is elastic, satisfying (1.1). If $\alpha = 0$, the collision is sticky, meaning the particles travel together after the collision. Generally, the coefficient α depends on $|v - v_*|$ and θ , the angle between $v - v_*$ and n . However, in many cases, this dependence is not considered significant, and a constant restitution coefficient

is often used. Further details about inelastic collisions and the Boltzmann equation will be covered in Section 1.4, Preliminaries.

Many physicists and mathematicians are interested in the behavior of solutions to both the elastic and inelastic Boltzmann equations. Numerous numerical, physical, and mathematical studies have been conducted on this topic. In mathematics, the L^∞ norm, weighted L^∞ norm, L^1 norm, and weighted L^1 norm of the Boltzmann equation were among the first to be studied.

The L^∞ norm is particularly important because it allows us to consider the pointwise bounded behavior of the solution, which is much stronger than merely being energy-bounded (i.e., $\int f|v|^2 dv < +\infty$). Based on L^∞ estimates, the upper Maxwellian bounds were established in [32], which are used in the local existence theory of strong solutions as seen in [33, 49]. Additionally, the Hölder continuity modulus and the lower Maxwellian bounds are dependent on the L^∞ norm of the solution, as discussed in [43, 44]. This lower bound is crucial for demonstrating the convergence of solutions to Maxwellian equilibrium, as shown in [26]. One of the most significant properties of inelastic collisions is that they are not time-reversible, making it impossible to determine the direction of entropy with respect to time. In the case of elastic collisions, the maximum entropy principle and Boltzmann's H -theorem ensure that the solution converges to the Maxwellian distribution. However, for inelastic collisions, the absence of the H -theorem makes it more challenging to establish an equilibrium state.

Moreover, the behavior of solutions varies depending on the additional physical forces, such as a heat bath and friction, that are incorporated into the kinetic model. Without external forces, the kinetic energy decreases until all gas particles stop, a state referred to as the *cooling state*. Details about these models are introduced in [65] by Villani. Notably, in [54], Mischler, Mouhot, and Ricard studied the *cooling process* and proved that the solutions converge to a Dirac mass in the weak-* measure sense asymptotically over a long time.

If kinetic energy is lost but the total density is conserved, it is predicted that the number of particles with low velocities will increase over time, while the number of particles with high velocities will decrease. Moreover, as velocity approaches zero, the density function increases rapidly over time (see Figure 3). This motivates our study of the behavior of solutions over time for inelastic collisions without any external forces in this paper.

In this paper, we study the L^∞ norm of the solution of the homogeneous inelastic Boltzmann equation for hard spheres. We will then extend the L^∞ norm to establish upper Maxwellian bounds. During the cooling process, as the solution tends towards $\delta_{v=0}$ and the kinetic energy approaches zero over time, it becomes crucial to derive pointwise upper bounds for the solution.

Before describing the inelastic model, we will briefly review the literature on the behavior of solutions to both the Boltzmann and inelastic Boltzmann equations.

1.1. Notations. Here, we briefly introduce some functions spaces, norms, and our notations. We define the weighted norms

$$\|f\|_{p,s} = \|f(t, \cdot)\|_{p,s} = \left(\int_{\mathbb{R}^3} (f(t, v)(1 + |v|^2)^{\frac{s}{2}})^p dv \right)^{\frac{1}{p}} < \infty$$




FIGURE 3. Velocity distribution for inelastic collisions over time (1 \rightarrow 2).

for $0 \leq s$ and $1 \leq p \leq \infty$. The corresponding weighted spaces are defined as

$$L_s^p = L_s^p(\mathbb{R}^3) = \{f : f \text{ is measurable on } \mathbb{R}^3, \|f\|_{p,s} < \infty\}.$$

If $p = \infty, s = 0$, we have

$$\|f\|_{\infty,0} = \|f\|_{\infty} = \operatorname{ess\,sup}_{v \in \mathbb{R}^3} |f(x)|.$$

The indicator function of a subset A within a set X is a function $\chi_A : X \rightarrow \{0, 1\}$, defined as

$$\chi_A(x) = \begin{cases} 1, & x \in A, \\ 0, & x \notin A \end{cases}$$

for $x \in X$. Throughout the paper, we denote as $A \lesssim B$ to mean that there exists a uniform constant C such that $A \leq CB$. If the constants depend on K , we write $C(K)$.

1.2. Known results for the elastic collisions. In particular, if the velocity distribution function f does not depend on the spatial variable x , the Boltzmann equation is called the (spatially) homogeneous Boltzmann equation. Regarding the homogeneous Boltzmann equation with a cut-off collision kernel, Carleman [22] studied L^∞ estimates for hard spheres. In 1982, Arkeryd extended this result to the general case in [8] and proved that

$$(1 + |v|)^{s'} f(t, v) \in L^\infty(\mathbb{R}_v^3)$$

for $t > 0$ under the assumption that the initial distribution $(1 + |v|)^s f_0 \in L^\infty(\mathbb{R}_v^3)$ for $s > 2$. For $s \leq 5$, this holds for any $s' \leq s$. For $s > 5$, it holds that $s' < s$ under the condition $(1 + |v|^2)^{s_1/2} f \in L^1(\mathbb{R}_v^3)$ for some $s_1 > 2$. In 1997, Bobylev proved that there exists θ_* such that $0 < \theta_* \leq \theta$ and

$$e^{\theta_* |v|^2} f \in L^1(\mathbb{R}_v^3)$$

for $t > 0$ whenever $e^{\theta |v|^2} f_0 \in L^1(\mathbb{R}_v^3)$ for hard spheres in [14]. In [56], Pulvirenti and Wennberg demonstrated that the solution is pointwise bounded from below by a Maxwellian for hard potentials. In 2009, Gamba, Panferov, and Villani proved that if f_0 has a Maxwellian upper bound, then the solutions have a uniform Maxwellian

upper bound for hard potentials in [32]. They applied the comparison principle using a dissipative property of the linear Boltzmann equation. Furthermore, the propagation of upper Maxwellian bounds in the spatially inhomogeneous case, specifically within the unit hypercube with periodic boundary conditions, was discussed.

Regarding the spatially inhomogeneous case with a non-cutoff collision kernel, Silvestre [58, Theorem 1.2] proved that

$$\|f(t, \cdot, \cdot)\|_{L_{x,v}^\infty} \leq C(t) < \infty,$$

where $C(t) < \infty$ for all $t > 0$ and depends only on the bounds of local macroscopic mass, energy, and entropy densities. In [42, Theorem 1.3-(1)], Imbert, Mouhot, and Silvestre obtained

$$\sup_{x,v} f(1 + |v|)^q < \infty$$

for $t > 0$ if $\sup_{x,v} f_0(1 + |v|)^q < \infty$ for $q \geq 0$, applicable to hard and moderately soft potentials. In [43], the authors demonstrated the appearance of a Gaussian lower bound. Briant proved the emergence of an exponential lower bound with the physical Maxwellian diffuse and specular reflection boundary conditions in [17, 18]. In [30], Fournier established the creation and propagation of exponential moments for the homogeneous Boltzmann equation for hard potentials. However, the propagation and creation of a Maxwellian upper bound remain open in the non-cutoff case.

1.3. Known results for the inelastic collisions. When gas particles collide with each other and lose energy, it is termed an inelastic collision, and such a gas is referred to as a granular gas. The behavior of a granular gas differs significantly from that of an ideal gas undergoing elastic collisions. The basic concepts of kinetic theory and exciting phenomena in granular gas dynamics were introduced in [20]. A hydrodynamic description for near elastic particles was explained in [15]. Here, we review various studies on the homogeneous inelastic Boltzmann equation for hard spheres.

In 2004, Bobylev, Gamba, and Panferov [16] obtained the steady velocity distributions of the solutions for large $|v|$ with external forcing. In [31], Gamba, Panferov, and Villani studied the existence, smoothness, uniqueness, and lower bound estimates of the solutions, incorporating an additional diffusion term in the equation. In [54], Mischler, Mouhot, and Ricard developed the Cauchy theory and demonstrated the creation and propagation of exponential moments. Mischler and Mouhot further proved the existence, uniqueness, and time asymptotic convergence of the self-similar solution in [52, 53]. They also proved Haff's law, which explains the precise rate of decay to zero for the granular temperature. Alonso and Lods also investigated a system of viscoelastic particles in [6]. A cooling inelastic Maxwell molecules of a suitable metric in the set of probability measures was studied by Bisi, Carrillo and Toscani in [13]. The Cauchy problem for small data in the space of functions bounded by Maxwellians was studied in [4].

Recently, there have been further developments regarding the homogeneous inelastic Boltzmann equation with a non-cutoff collision kernel. An and Lee [7] proved the non-Maxwellian lower bound of the solutions. Qi [57] studied the well-posedness theory of measure-valued solutions. Additionally, Jang and Qi [46] established the global-in-time existence of measure-valued solutions and demonstrated the creation of polynomial moments.

The collapse phenomenon of inelastic particles has been discussed in [10, 11]. Recently, [28, 29] studied a dynamical system describing this collapse. Additionally, the inelastic collision operator appears in more generalized physical settings, such as the scattering of photons interacting with both the ground-state gas and the excited-state gas in the radiative transfer process (see [24, 48]). Furthermore, [23] examined the Cauchy problem for the inelastic Vlasov–Poisson–Boltzmann system with a soft potential in the near vacuum regime.

1.4. Preliminaries.

1.4.1. *Aftermath of inelastic collisions.* In this section, we explain inelastic collisions. Let v and v_* be the velocities of two particles before a collision, and let v' and v'_* be their velocities after the collision. Recall equation (1.4):

$$\begin{aligned} (v' - v'_*) \cdot n &= -\alpha(v - v_*) \cdot n, \\ (v' - v'_*) - ((v' - v'_*) \cdot n)n &= (v - v_*) - ((v - v_*) \cdot n)n, \end{aligned} \quad (1.5)$$

where α is the coefficient of normal restitution, $0 < \alpha \leq 1$, and the impact direction $n \in \mathbb{S}_+^2$. We define

$$\beta = \frac{1 + \alpha}{2}, \quad \frac{1}{2} < \beta \leq 1. \quad (1.6)$$

Since momentum is conserved, we set

$$v' = v - a(v, v_*, n)n, \quad v'_* = v + a(v, v_*, n)n$$

for some function $a(v, v_*, n)$ and $n \in \mathbb{S}_+^2$. We find $a(v, v_*, n) = \beta((v - v_*) \cdot n)n$, so that v' and v'_* satisfy equation (1.5). The post-collisional velocities v' and v'_* are expressed as:

$$v' = v - \beta((v - v_*) \cdot n)n, \quad v'_* = v_* + \beta((v - v_*) \cdot n)n. \quad (1.7)$$

In Figure 4, we illustrate these coordinates. When v and v_* are fixed, there are two spheres: the possible locations of v' are restricted to the left sphere, while those of v'_* are in the right sphere. Another way to express v' and v'_* is by using $\sigma \in \mathbb{S}^2$, starting with $\frac{v+v_*}{2}$ such that:

$$\begin{aligned} v' &= \frac{v + v_*}{2} + \frac{1 - \beta}{2}(v - v_*) + \frac{\beta}{2}|v - v_*|\sigma, \\ v'_* &= \frac{v + v_*}{2} - \frac{1 - \beta}{2}(v - v_*) - \frac{\beta}{2}|v - v_*|\sigma. \end{aligned} \quad (1.8)$$

In elastic collisions, we only need to define the post-collisional velocities v' and v'_* . However, inelastic collisions are not time-reversible. If v' and v'_* collide, they do not revert to v and v_* . Thus, in inelastic collisions, we define the pre-collisional velocities $'v$ and $'v_*$, which then become v and v_* , respectively, after the collision. In equation (1.5), we substitute v and v_* with $'v$ and $'v'_*$, and $'v$ and $'v'_*$ with v and v_* as follows:

$$\begin{aligned} ('v - 'v'_*) \cdot n &= -\frac{1}{\alpha}(v - v_*) \cdot n, \\ ('v - 'v'_*) - (('v - 'v'_*) \cdot n)n &= (v - v_*) - ((v - v_*) \cdot n)n. \end{aligned} \quad (1.9)$$

In contrast to (1.5), α is replaced by $1/\alpha$ in (1.9). Let γ be

$$\gamma = \frac{1}{2} \left(1 + \frac{1}{\alpha} \right), \quad 1 \leq \gamma < \infty. \quad (1.10)$$




FIGURE 4. Possible locations of v' and v'_* given v and v_* in inelastic collisions.

If we replace α with $1/\alpha$ in (1.7) and (1.8), then $'v, 'v_*$ are expressed as

$$'v = v - \gamma((v - v_*) \cdot n)n, \quad 'v_* = v_* + \gamma((v - v_*) \cdot n)n \quad (1.11)$$

for $n \in \mathbb{S}_+^2$ and

$$\begin{aligned} 'v &= \frac{v + v_*}{2} + \frac{1 - \gamma}{2}(v - v_*) + \frac{\gamma}{2}|v - v_*|\sigma, \\ 'v_* &= \frac{v + v_*}{2} - \frac{1 - \gamma}{2}(v - v_*) - \frac{\gamma}{2}|v - v_*|\sigma \end{aligned} \quad (1.12)$$

for $\sigma \in \mathbb{S}^2$. The Jacobian between (v, v_*) and (v', v'_*) is calculated as

$$\frac{\partial(v', v'_*, -n)}{\partial(v, v_*, n)} = \frac{\partial(v, v_*, -n)}{\partial(v, v_*, n)} = \alpha \quad (1.13)$$

from (1.7) and (1.11). From (1.7) and (1.8), we obtain directly

$$\begin{aligned} &|v'|^2 + |v'_*|^2 - |v|^2 - |v_*|^2 \\ &= -\frac{1 - \alpha^2}{2}((v - v_*) \cdot n)^2 \\ &= -\frac{1 - \alpha^2}{2} \frac{1 - \frac{v - v_*}{|v - v_*|} \cdot \sigma}{2} |v - v_*|^2 \leq 0, \end{aligned} \quad (1.14)$$

and this shows that the loss of kinetic energy depends on the coefficient of normal restitution α , $|v - v_*|$, and the collision angle.

1.4.2. *Inelastic Boltzmann collision operator.* Now we introduce the spatially homogeneous inelastic Boltzmann equation. It goes by

$$\partial_t f = Q(f, f), \quad f(0, v) = f_0(v), \quad (1.15)$$

where $v \in \mathbb{R}^3$, $t \geq 0$ and $f \geq 0$. (The operator is local in t and we abbreviate t for notational convenience.) Taking $\phi(v)$ to be a suitably regular test function, the weak form of the inelastic Boltzmann collision operator $Q(f, g)(t, v)$ is written by

$$\begin{aligned} &\int_{\mathbb{R}^3} Q(f, g)(t, v) \phi(v) dv \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} B(|v - v_*|, \cos \theta) f(t, v_*) g(t, v) (\phi(v') - \phi(v)) dndv_* dv, \end{aligned} \quad (1.16)$$

where

$$\cos \theta = \left(\frac{v - v_*}{|v - v_*|}, n \right), \quad \theta \in \left[0, \frac{\pi}{2} \right]. \quad (1.17)$$

In the case $f = g$ in (1.16), we have

$$\begin{aligned} \int_{\mathbb{R}^3} Q(f, f)(t, v) \phi(v) dv &= \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} B(|v - v_*|, \cos \theta) f(t, v_*) f(t, v) \\ &\quad \times (\phi(v') + \phi(v'_*) - \phi(v) - \phi(v_*)) dndv_* dv. \end{aligned} \quad (1.18)$$

Taking $\phi(v) = 1, |v|^2$, and v_i into above equation, for any $t > 0$, we have

$$\int_{\mathbb{R}^3} f(t, v) dv = \int_{\mathbb{R}^3} f_0(v) dv, \quad \text{and} \quad \int_{\mathbb{R}^3} f(t, v) |v|^2 dv \leq \int_{\mathbb{R}^3} f_0(v) |v|^2 dv \quad (1.19)$$

by (1.14), and

$$\int_{\mathbb{R}^3} f(t, v) (v \cdot e_i) dv = \int_{\mathbb{R}^3} f_0(v) (v \cdot e_i) dv \quad (1.20)$$

for $i = 1, 2, 3$. Here $e_1 = (1, 0, 0)$, $e_2 = (0, 1, 0)$ and $e_3 = (0, 0, 1)$. In this paper, we suppose

$$\int_{\mathbb{R}^3} f_0(v) dv = 1, \quad \int_{\mathbb{R}^3} f_0(v) (v \cdot e_i) dv = 0, \quad \int_{\mathbb{R}^3} f_0(v) |v|^2 dv = 1 \quad (1.21)$$

for $i = 1, 2, 3$, without loss of generality. See also [31, 52–54] in which the authors consider the same initial conditions (1.21). By (1.19)–(1.21), it holds that

$$\begin{aligned} \|f(t, v)\|_{1,0} &= \|f_0(v)\|_{1,0} = 1, \quad \|f(t, v)\|_{1,2} \leq \|f_0(v)\|_{1,2} < +\infty \\ \int_{\mathbb{R}^3} f_0(v) (v \cdot e_i) dv &= \int_{\mathbb{R}^3} f(t, v) (v \cdot e_i) dv = 0 \end{aligned} \quad (1.22)$$

for $t > 0$ and $f_0 \in L_2^1(\mathbb{R}^3)$. Moreover, the initial entropy is denoted as

$$\int_{\mathbb{R}^3} f_0(v) \log f_0(v) dv = H_0. \quad (1.23)$$

Throughout the paper, we will consider a hard sphere model

$$B(|v - v_*|, \cos \theta) = |(v - v_*) \cdot n| = |v - v_*| \cos \theta$$

for $\theta \in [0, \frac{\pi}{2}]$. Since $\delta(v)$ is approximated by C_c^∞ test function, we can take $\phi(v) = \delta(v - v_1)$ in (1.16). By integrating about v , we have

$$\begin{aligned} Q(f, g)(t, v_1) &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v - v_*) \cdot n| f(t, v_*) g(t, v) (\delta(v' - v_1) - \delta(v - v_1)) dndv_* dv \\ &:= Q^+(f, g)(t, v_1) - g(t, v_1) Lf(t, v_1). \end{aligned}$$

By changing of variables $(v, v_*, n) \rightarrow (v', v'_*, -n)$ with Jacobian determinant α , and (1.4),

$$\begin{aligned}
 Q^+(f, g)(t, v_1) &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v - v_*) \cdot n| f(t, v_*) g(t, v) \delta(v' - v_1) \, dndv_* dv \\
 &= \frac{1}{\alpha^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v - v_*) \cdot n| f(t, v'_*) g(t, v) \delta(v - v_1) \, dndv_* dv \\
 &= \frac{1}{\alpha^2} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v_1 - v_*) \cdot n| f(t, v'_*) g(t, v_1) \, dndv_*.
 \end{aligned} \tag{1.24}$$

And we have

$$\begin{aligned}
 Lf(t, v_1) &= \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v_1 - v_*) \cdot n| f(t, v_*) \, dndv_* \\
 &= \pi \int_{\mathbb{R}^3} |v_1 - v_*| f(t, v_*) \, dv_*.
 \end{aligned} \tag{1.25}$$

Before, we parameterize v' and v'_* in two different ways by using impact direction $n \in \mathbb{S}_+^2$ in (1.7), and $\sigma \in \mathbb{S}^2$ in (1.8). In Figure 4, there are n and σ , and we get

$$\frac{v - v_*}{|v - v_*|} \cdot \sigma = \cos(\pi - 2\theta) \quad \text{where} \quad \frac{v - v_*}{|v - v_*|} \cdot n = \cos \theta. \tag{1.26}$$

If we choose and fix the direction of $v - v_*$ as the z -axis, we can consider spherical coordinate of

$$n = (\cos \phi \sin \theta, \sin \phi \sin \theta, \cos \theta), \quad \sigma = (\cos \phi \sin 2\theta, \sin \phi \sin 2\theta, -\cos 2\theta) \tag{1.27}$$

for $\phi \in [0, 2\pi]$, $\theta \in [0, \frac{\pi}{2}]$. By using σ of instead n , we can focus the amplitude, $|v - v_*|$ in $Q^+(f, f)(t, v_1)$ without $\cos \theta$ term in (1.24). Moreover, we define the rescaled σ' as

$$\sigma' = \frac{\beta}{2} |v - v_*| \sigma, \tag{1.28}$$

which is used in the proof of Lemma 2.9. Using

$$d\sigma = 4 \cos \theta \, dn \quad \text{and} \quad d\sigma' = \frac{\beta^2}{4} |v - v_*|^2 d\sigma, \tag{1.29}$$

the weak form of the $Q^+(f, g)(v)$ can be expressed in the variables σ and σ' as follows:

$$\begin{aligned}
 &\int_{\mathbb{R}^3} Q^+(f, g)(t, v) \phi(v) \, dv \\
 &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v - v_*) \cdot n| f(t, v_*) g(t, v) \phi(v') \, dndv_* dv \\
 &= \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*| f(t, v_*) g(t, v) \phi(v') \, d\sigma dv_* dv
 \end{aligned} \tag{1.30}$$

$$= \frac{1}{\beta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{B(\frac{\beta}{2}|v-v_*|)} |v - v_*|^{-1} f(t, v_*) g(t, v) \phi(v') \, d\sigma' dv_* dv, \tag{1.31}$$

where

$$B\left(\frac{\beta}{2}|v - v_*|\right) = \left\{ x \in \mathbb{R}^3 : |x| = \frac{\beta}{2}|v - v_*| \quad \text{for} \quad v, v_* \in \mathbb{R}^3 \right\}.$$

1.4.3. *Carleman type representation for granular gases.* In 1957, Carleman introduced a new representation of the gain term collision operator for the Boltzmann equation with a cutoff kernel. When v' and v are fixed, the set of possible vectors for v'_* forms a hyperplane orthogonal to $v - v'$ and passing through v (see Figure 1). Using this fact, he obtained the gain term integrated over the variables v' and v'_* , where the range is in \mathbb{R}^3 and a hyperplane for each, as described in [22]. Moreover, there is a Carleman-type representation for granular gas presented in [52, Proposition 1.5] by Mischler and Mouhot. To obtain a Carleman-type representation, we define a vector P as:

$$P = \frac{1}{\beta}v' - \left(\frac{1}{\beta} - 1\right)v,$$

which is located on the extension line of v and v' in ratio

$$|P - v| : |P - v'| = 1 : 1 - \beta$$

for $\beta \neq 1$ as in Figure 4. For $\beta = 1$ (elastic collisions), the vector P equals v' . Then P also satisfies $P - v \perp P - v_*$. Next, we introduce a 2-dimensional hyperplane $E_P^{v'-v}$ which is normal to the vector $v' - v$ around the vector P ; it is defined as

$$E_P^{v'-v} = \left\{ x \in \mathbb{R}^3 : (x - P) \perp (v - v') \text{ where } P = \frac{1}{\beta}v' - \left(\frac{1}{\beta} - 1\right)v \right\} \subseteq \mathbb{R}^2. \quad (1.32)$$

We notice that if v and v' are fixed, v'_* is in the hyperplane $E_P^{v'-v}$. Then $Q^+(f, g)(t, v_1)$ is expressed by

$$\begin{aligned} & Q^+(f, g)(t, v_1) \\ &= \int_{\mathbb{R}^3} Q^+(f, g)(t, v) \delta(v - v_1) dv \\ &= \frac{1}{\beta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \delta(v - v_1) g(t, v) |v - v'|^{-1} \int_{v'_* \in E_P^{v'-v}} f(t, v'_*) dE_{v'_*} dv dv' \quad (1.33) \\ &= \frac{1}{\beta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \delta(v' - v_1) g(t, v) |v' - v|^{-1} \int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} dv dv', \end{aligned}$$

where $dE_{v'_*}$ and dE_{v_*} are the Lebesgue measures on the plane $E_P^{v'-v}$ and $E_P^{v'-v}$, respectively. This Carleman-type representation is used crucially for the estimates of the gain term, $Q^+(f, f)$, through the paper.

1.5. Main results. We are now ready to state our main theorems. Given $t > 0$ and $v \in \mathbb{R}^3$, we prove that the solution $f(t, v)$ is bounded pointwise from above by $C_{1, f_0}(1 + t)^3$ in the region including zero velocity, and by $C_{2, f_0}(1 + t)$, away from the zero velocity ($v \neq 0$). We have also derived an upper bound that depends on the coefficient of normal restitution constant $\alpha \in (0, 1]$ and the time $t > 0$. We notice this upper bound becomes smaller as $\alpha \rightarrow 1$, and it becomes a constant when $\alpha = 1$.

Moreover, the first part of the minimum in (1.36) indicates that the density distribution function during the cooling process is very sensitive to the temporal variable when the velocity is near zero. The growth of this bound can be considered to be natural, as shown in Figure 3, during the cooling process. In addition, we

observe that there is a loss of weight in the L^∞ propagation in the first part of the minimum in (1.35). Here, we define the cooling time T_c in the cooling process as

$$\begin{aligned} T_c &:= \inf \{T \geq 0, \mathcal{E}(t) = 0 \ \forall t > T\} \\ &= \sup \{S \geq 0, \mathcal{E}(t) > 0 \ \forall t < S\}, \end{aligned} \quad (1.34)$$

where \mathcal{E} is the total kinetic energy, which is defined as $\mathcal{E}(t) = \int_{\mathbb{R}^3} f(t, v) |v|^2 dv$.

Theorem 1.1. *Let $0 < \alpha \leq 1$ be the coefficient of normal restitution constant, which satisfies (1.4). Assume that (1.21) holds and $f_0 \in L_s^\infty$ for $s > 2$.*

(i) *Then there exist positive constants $C_1(\|f_0\|_{\infty, s})$ and C_2 such that*

$$f(t, v) \leq C_1(\|f_0\|_{\infty, s}) \min \left\{ \frac{(1+t)^3}{1+|v|}, \frac{e^{C_2(\frac{1}{\alpha^2}-1)t}}{(1+|v|)^2} \right\} \quad (1.35)$$

for almost every $v \in \mathbb{R}^3$ and for every $t > 0$.

(ii) *Moreover, there exists a positive constant $C_3(\|f_0\|_{\infty, s})$ such that*

$$f(t, v) \leq C_3(\|f_0\|_{\infty, s}) \min \left\{ \frac{1+t}{|v|^2}, \frac{e^{C_2(\frac{1}{\alpha^2}-1)t}}{(1+|v|)^2} \right\} \quad (1.36)$$

for almost every $v \neq 0 \in \mathbb{R}^3$ and for every $t > 0$.

Corollary 1.2. *Assume that (1.21) holds and $f_0 \in L_s^\infty$ for $s > 2$. Then the cooling time T_c of a solution, defined in (1.34), is infinite.*

Remark 1.3. *Theorem 1.1 holds even for the infinite energy case as long as we have $f(t, \cdot) \in L_1^1$ for any $t \geq 0$; i.e., the finiteness condition on the initial energy in (1.21) can be replaced by $f(t, \cdot) \in L_1^1$ for $t \geq 0$. Either one of these can guarantee that Lf be well-defined in Lemma 2.4.*

Remark 1.4. *In [54], Mischler, Mouhot and Ricard studied the spatially homogeneous inelastic Boltzmann equation for hard spheres, considering a general form of collision rate that includes variable restitution coefficients depending on the kinetic energy and the relative velocity. In Theorem 1.2, they consider the case where the collision rate B is independent of the kinetic energy. In particular, according to Theorem 1.2-(i) and Remark 1.3-(1), for inelastic hard spheres with a constant restitution coefficient, which we studied in this paper, there exists a unique solution under the assumption of (1.21), and the cooling time of the solution is infinite. In Corollary 1.2, we obtain cooling time T_c is infinite, but we also assume $f_0 \in L_s^\infty$.*

Many researchers are interested in the upper Maxwellian bound, lower Maxwellian bound, and the convergence of the system to the Maxwellian state in the elastic Boltzmann equation, where α is 1. Due to the loss of energy in inelastic collisions, we also expect the solution to exhibit thinner velocity tails in the inelastic Boltzmann equation and to approach the Maxwellian upper bound for sufficiently large velocities for all $0 < \alpha \leq 1$. One of the most significant aspects of Theorem 1.1 is that it provides pointwise upper bounds, allowing us to predict the behavior of the solution. Using these upper bound estimates, we will establish the Maxwellian upper bound.

Theorem 1.5. *Let $0 < \alpha \leq 1$ be the coefficient of normal restitution constant, which satisfies (1.5). We assume that (1.21) holds, and that $f_0(v) \leq M_0(v)$ for*

almost every $v \in \mathbb{R}^3$, where $M_0(v) = e^{-a_0|v|^2+c_0}$. Then there exist $a \in (0, a_0)$, $b \in \mathbb{R}$ and $C > 0$, depending on a_0 and c_0 , such that

$$f(t, v) \leq \min \left\{ (1+t)^3, e^{C(\frac{1}{\alpha^2}-1)t} \right\} e^{-a|v|^2+b} \quad (1.37)$$

for almost every $v \in \mathbb{R}^3$ and every $t > 0$.

Remark 1.6. Fix the coefficient of normal restitution constant $0 < \alpha < 1$, and denote $T_\alpha = \alpha^2/(1-\alpha^2)$. Then, by Theorem 1.1-(i), there is a positive constant $\bar{C}(\|f_0\|_{\infty,s})$ such that

$$\operatorname{ess\,sup}_{v \in \mathbb{R}^3} f(t, v)(1+|v|)^2 \leq \bar{C}(\|f_0\|_{\infty,s})$$

for $0 < t < T_\alpha$, under (1.21) and that $f_0 \in L_s^\infty$ for $s > 2$. By Theorem 1.5, there exist $\bar{a} \in (0, a_0)$, $\bar{b} \in \mathbb{R}$, depending on a_0 and c_0 , such that

$$f(t, v) \leq e^{-\bar{a}|v|^2+\bar{b}}$$

for almost every $v \in \mathbb{R}^3$ and $0 < t < T_\alpha$ under $f_0(v) \leq M_0(v)$.

Remark 1.7. In [54], in Proposition 3.2-(ii), Mischler, Mouhot, and Ricard proved that there exist $C, r' > 0$ such that

$$\sup_{t \in (0, T_c)} \int_{\mathbb{R}^N} f(t, v) e^{r'|v|^\eta} dv \leq C$$

if $f_0(v) e^{r|v|^\eta} \in L_0^1(\mathbb{R}^N)$ for $r > 0$ and $\eta \in (0, 2]$, and T_c is defined as (1.34). They also considered a more generalized model that includes inelastic hard spheres with a constant restitution coefficient.

Under elastic collisions, the total entropy remains finite for all $t > 0$ by the H -theorem if it is initially finite. As a consequence, it establishes a uniform lower bound for $Lf(t, v)$ as stated in Lemma 4 of [8], which has been used to provide a pointwise upper-bound of solution via the Grönwall inequality as in [8]. On the other hand, we observe that the inelastic collisions lack an H -theorem and a uniform lower bound for $Lf(t, v)$. So, we derive the weak version for lower bounds of $Lf(t, v)$ in Lemma 2.4. Under elastic collisions, the total entropy remains finite for all $t > 0$ by the H -theorem if it is initially finite. As a consequence, it establishes a uniform lower bound for $Lf(t, v)$ as stated in Lemma 4 of [8], which has been used to provide a pointwise upper-bound of solution via the Grönwall inequality as in [8]. On the other hand, we observe that the inelastic collisions lack an H -theorem and a uniform lower bound for $Lf(t, v)$. Thus, we derive two weaker versions for lower bounds of $Lf(t, v)$ in Lemma 2.4. Since these lower bounds are not uniform with respect to velocity and time, L^∞ norms in Theorem 1.1 depend also on time.

1.6. Outline of the rest of the article. Here we briefly outline the structure of the remainder of the paper. In Section 2, we prove L^∞ estimates. First, we obtain the upper bound of the gain term of the collision operator, $Q^+(f, f)$, in Lemma 2.7 through Lemma 2.12. Next, we apply Grönwall's lemma in Lemma 2.1 to the Boltzmann equation to prove Theorem 1.1. In Section 3, based on a comparison principle in Lemma 3.1, we extend L^∞ estimates to establish the Maxwellian upper bound.

2. L^∞ ESTIMATES FOR THE INELASTIC BOLTZMANN EQUATION

In this Section, we estimate the L^∞ norm of the solution to the inelastic Boltzmann equation. Let $0 < \alpha \leq 1$ be the coefficient of normal restitution constant, which satisfies (1.5). The following Lemma 2.1, which is derived from Grönwall's inequality, will be repeatedly used in the paper.

Lemma 2.1. *Let $h_1(t)$ and $h_2(t)$ be continuous real functions on \mathbb{R}_+ and $h_1(t) > 0$. If*

$$\frac{d}{dt}f + h_1 f \leq h_2$$

for $t > 0$, then

$$f(t) \leq \sup_{0 \leq s \leq t} \frac{h_2(s)}{h_1(s)} + f(0)$$

for $t > 0$.

Proof. From

$$\frac{d}{dt} \left(e^{\int_0^t h_1(s) ds} f \right) \leq e^{\int_0^t h_1(s) ds} h_2(t),$$

we obtain

$$\begin{aligned} f(t) &\leq \int_0^t e^{-\int_\tau^t h_1(s) ds} h_2(\tau) d\tau + f(0) \\ &\leq \left(1 - e^{-\int_0^t h_1(s) ds} \right) \sup_{0 \leq s \leq t} \frac{h_2(s)}{h_1(s)} + f(0) \leq \sup_{0 \leq s \leq t} \frac{h_2(s)}{h_1(s)} + f(0) \end{aligned}$$

for $t > 0$. □

Lemma 2.2. *Assume that (1.21) holds, and that the initial entropy H_0 in (1.23) is finite. Then there exists a positive constant C that depends only on $\|f_0\|_{1,2}$, such that*

$$\int_{\mathbb{R}^3} f(t, v) \log f(t, v) dv \leq Ct \left(\frac{1}{\alpha^2} - 1 \right) + H_0 \quad (2.1)$$

for $t > 0$.

Proof. We denote $f := f(t, v)$, $f_* := f(t, v_*)$, $f' := f(t, v')$, $f'_* := f(t, v'_*)$. From the weak form of the collision operator (1.18), we have

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^3} f \log f dv &= \int_{\mathbb{R}^3} Q(f, f)(t, v) \log f dv \\ &= \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} f f_* \log \frac{f' f'_*}{f f_*} |(v - v_*) \cdot n| dndvdv_* \\ &= \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} f f_* \left(\log \frac{f' f'_*}{f f_*} - \frac{f' f'_*}{f f_*} + 1 \right) |(v - v_*) \cdot n| dndvdv_* \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} (f' f'_* - f f_*) |(v - v_*) \cdot n| dndvdv_*. \end{aligned} \quad (2.2)$$

Taking a change of $(v, v_*, n) \rightarrow (v', v'_*, -n)$ with the Jacobian determinant α and (1.4), we have

$$\begin{aligned} & \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} f' f'_* |(v - v_*) \cdot n| \, dndvdv_* \\ &= \frac{1}{\alpha^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} f f_* |(v - v_*) \cdot n| \, dndvdv_*. \end{aligned} \quad (2.3)$$

By using $\log x - x + 1 \leq 0$ for $x > 0$, and applying (2.3) to (2.2), we have

$$\frac{d}{dt} \int_{\mathbb{R}^3} f \log f \, dv \leq \frac{1}{2} \left(\frac{1}{\alpha^2} - 1 \right) \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} f f_* |(v - v_*) \cdot n| \, dndvdv_*. \quad (2.4)$$

By the assumption (1.21) on the profile f_0 , we obtain

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} f f_* |(v - v_*) \cdot n| \, dndvdv_* \lesssim 1.$$

Lastly, by integrating (2.4) with respect to $t > 0$, we obtain

$$\int_{\mathbb{R}^3} f(t, v) \log f(t, v) \, dv \leq Ct \left(\frac{1}{\alpha^2} - 1 \right) + H_0$$

for some positive constant C that depends only on $\|f_0\|_{1,2}$. \square

Remark 2.3. *The quantitative variants of Boltzmann's H-theorem were proven in [62, 64]. In [26], by using H-theorem, convergence to equilibrium for solutions of the inhomogeneous Boltzmann equation was studied. The H-theorem was also proven for the spatially homogeneous relativistic Boltzmann equation in [47, 61].*

We first estimate $Lf(t, v)$ in (1.25), and later, the lower bound of $Lf(t, v)$ will serve the role of $h_1(t)$ in Lemma 2.1 for the Boltzmann equation.

Lemma 2.4. *Assume that (1.21) holds, and that the initial entropy H_0 in (1.23) is finite. Then there exist positive constants C_1 and C_2 such that*

$$Lf(t, v_1) \geq \max \left\{ \pi |v_1|, C_1 e^{-C_2 \left(\frac{1}{\alpha^2} - 1 \right) t} (1 + |v_1|) \right\} \quad (2.5)$$

for any $v_1 \in \mathbb{R}^3$ and $t > 0$.

Proof. Since $\phi(s) = |s|$ is a convex function, by Jensen's inequality, we have

$$\begin{aligned} Lf(t, v_1) &= \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v_1 - v_*) \cdot n| f(t, v_*) \, dndv_* = \pi \int_{\mathbb{R}^3} |v_1 - v_*| f(t, v_*) \, dv_* \\ &\geq \pi \left| \sum_{i=1,2,3} \left\{ (v_1 \cdot e_i) \int_{\mathbb{R}^3} f(t, v_*) \, dv_* - \int_{\mathbb{R}^3} (v_* \cdot e_i) f(t, v_*) \, dv_* \right\} e_i \right| \\ &= \pi \left| \sum_{i=1,2,3} \{ (v_1 \cdot e_i) \times 1 - 0 \} e_i \right| = \pi |v_1| \end{aligned} \quad (2.6)$$

for $v_1 \in \mathbb{R}^3$.

On the other hand, by Lemma 2.2, we have

$$\begin{aligned}
& \int_{\{|v-v_1|<r\}} f(t, v_1) dv_1 \\
& \leq \int_{\{|v-v_1|<r, f(t, v_1) \leq j\}} f(t, v_1) dv_1 + \int_{\{|v-v_1|<r, f(t, v_1) > j\}} f(t, v_1) dv_1 \\
& \leq \frac{4\pi}{3} jr^3 + \frac{1}{\log j} \left(Ct \left(\frac{1}{\alpha^2} - 1 \right) + H_0 \right)
\end{aligned} \tag{2.7}$$

for $r, j > 0$. Taking

$$\log j = \frac{4}{\|f_0\|_{1,0}} \left(Ct \left(\frac{1}{\alpha^2} - 1 \right) + H_0 \right), \quad r = \left(\frac{3}{8\pi} \|f_0\|_{1,0} \frac{1}{j} \right)^{\frac{1}{3}}$$

in (2.7), we find that

$$\int_{\{|v-v_1|<r\}} f(t, v_1) dv_1 \leq \frac{1}{2} \|f_0\|_{1,0}.$$

Then we have

$$\begin{aligned}
L(f)(t, v) &= \pi \int_{\mathbb{R}^3} f(t, v_1) |v - v_1| dv_1 \geq \pi r \int_{\{|v-v_1| \geq r\}} f(t, v_1) dv_1 \\
&= \pi r \left(\int_{\mathbb{R}^3} f(t, v_1) dv_1 - \int_{\{|v-v_1| < r\}} f(t, v_1) dv_1 \right) \\
&\geq \frac{1}{2} \pi r \|f_0\|_{1,0} = \frac{\pi}{2} \left(\frac{3}{4\pi} \right)^{\frac{1}{3}} \|f_0\|_{1,0}^{\frac{4}{3}} e^{-\frac{4}{3\|f_0\|_{1,0}} (Ct(\frac{1}{\alpha^2}-1)+H_0)} \\
&\geq c_1 e^{-c_2(\frac{1}{\alpha^2}-1)t}
\end{aligned} \tag{2.8}$$

for some positive constants c_1 and c_2 . From (2.6) and (2.8), we conclude (2.5). \square

Remark 2.5. *The finiteness condition on the initial energy in (1.21) can be replaced by $f(t, \cdot) \in L_1^1$. Either of these can guarantee that Lf be well-defined.*

Remark 2.6. *In [8], in Lemma 4, Arkeryd proved that*

$$Lf(t, v) \geq C(C_0, \|f\|_{1,2}, \|f\|_{1,0})(1 + |v|)^\gamma, \quad \gamma \in (0, 1]$$

under the conditions $f \in L_2^1$ and $\int_{\mathbb{R}^3} f \log f dv < C_0$ for the homogeneous Boltzmann equation for hard potentials with angular cutoff. For the hard spheres case, γ is 1. In elastic collisions, it holds that $\int_{\mathbb{R}^3} f \log f dv \leq \int_{\mathbb{R}^3} f_0 \log f_0 dv$ by the H-theorem. However, we cannot control $\int_{\mathbb{R}^3} f \log f dv$ in inelastic collisions, so $Lf(t, v) \gtrsim 1$ does not hold.

Before the estimating $Q^+(f, f)(t, v)$, we will consider Lemma 2.7 and Lemma 2.8.

Lemma 2.7. *Let $f(t, \cdot) \in L_0^1$. For any given fixed $v \in \mathbb{R}^3$ and $t > 0$, we have*

$$\left\| \int_{\mathbb{R}^3} Q^+(f, f)(t, v_1) |v_1 - v|^{-1} dv_1 \right\|_{\infty, 0} \leq 4\pi \|f\|_{1,0}^2.$$

Proof. In (1.30), we replace v, v' with v_1, v'_1 respectively. Next, we choose $\phi(v_1) = |v - v_1|$ for a given fixed $v \in \mathbb{R}^3$. Using (1.29), we have

$$\begin{aligned} & \int_{\mathbb{R}^3} Q^+(f, f)(t, v_1) |v_1 - v|^{-1} dv_1 \\ &= \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} f(t, v_1) f(t, v_*) |v_1 - v_*| \int_{\mathbb{S}^2} |v'_1 - v|^{-1} d\sigma dv_* dv_1 \\ &= \frac{1}{2\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} f(t, v_1) f(t, v_*) \int_{\mathbb{S}^2} |\sigma + A|^{-1} d\sigma dv_* dv_1 \leq 4\pi \|f\|_{1,0}^2, \end{aligned}$$

where A is $(\frac{\beta}{2}|v_1 - v_*|)^{-1}(\frac{v_1 + v_*}{2} + \frac{1-\beta}{2}(v_1 - v_*) - v)$ from (1.8)₁. \square

Lemma 2.8. *For any $v, \bar{v} \in \mathbb{R}^3$, we recall $E_P^{v-\bar{v}}$ from (1.32). We let $d_{v_1} \geq 0$ be the distance from $v_1 \in \mathbb{R}^3$ to the plane $E_P^{v-\bar{v}}$, and*

$$D_j(v_1) = \sqrt{\frac{j}{\pi}} e^{-(\sqrt{j}d_{v_1})^2}. \quad (2.9)$$

Then we have

$$\lim_{j \rightarrow \infty} \int_{\mathbb{R}^3} D_j(v_1) Q^+(f, f)(t, v_1) dv_1 = \int_{v_1 \in E_P^{v-\bar{v}}} Q^+(f, f)(t, v_1) dE_{v_1}.$$

Proof. Note that $D_j(v_1)$ converges to $\delta(v_1)$ in distribution as $j \rightarrow \infty$. Let a be the vector in $E_P^{v-\bar{v}}$ satisfying $dv_1 = da d(d_{v_1})$. We obtain

$$\begin{aligned} & \lim_{j \rightarrow \infty} \int_{\mathbb{R}^3} D_j(v_1) Q^+(f, f)(t, v_1) dv_1 \\ &= \lim_{j \rightarrow \infty} \int_{-\infty}^{+\infty} D_j(v_1) \int_{a \in E_P^{v-\bar{v}}} Q^+(f, f)(t, v_1) da d(d_{v_1}) \\ &= \int_{v_1 \in E_P^{v-\bar{v}}} Q^+(f, f)(t, v_1) dE_{v_1}. \end{aligned}$$

\square

To estimate the gain term collision operator $Q^+(f, f)(t, v)$, we will establish the following Lemma 2.9 and Lemma 2.12. Write a Carleman-type representation of $Q^+(f, f)$ from in 1.33,

$$Q^+(f, f)(t, v_1) = \frac{1}{\beta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \delta(v' - v_1) f(t, v) |v' - v|^{-1} \int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} dv dv',$$

where $E_P^{v'-v}$ is in (1.32). We will estimate

$$\int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} \quad \text{and} \quad \int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1$$

in Lemma 2.9 and Lemma 2.12, respectively.

Lemma 2.9. *Let $f \in L_0^1$. We have*

$$\int_{v_1 \in E_P^{v-\bar{v}}} Q^+(f, f)(t, v_1) dE_{v_1} \leq 2\pi \|f\|_{1,0}^2$$

for any $v, \bar{v} \in \mathbb{R}^3$ and $t > 0$.

Proof. In Figure 4, for given v_1, v_* (with v_1 replacing v in Figure 4 here), the two spheres are given where the sphere on the left represents the possible orbit for v'_1 and the other on the right represents the one for v'_* . Let us denote the sphere on the left as $S(v'_1)$. Then we denote $\chi = 1$ if the plane $E_P^{v-\bar{v}}$ and $S(v'_1)$ intersect and $\chi = 0$ if the plane $E_P^{v-\bar{v}}$ and $S(v'_1)$ do not intersect. For $D_j(v'_1)$ in (2.9), if $\chi = 1$, note that we have

$$\begin{aligned} \int_{S(v'_1)} D_j(v'_1) d\sigma' &= \int_{-\infty}^{+\infty} D_j(v'_1) \int_{a \in S(v'_1) \cap E_P^{v-\bar{v}}} da (dv_1) \\ &\leq \int_{a \in B\left(\frac{\beta|v_1-v_*|}{2}\right)} 1 da \leq \pi\beta|v_1-v_*|, \end{aligned} \quad (2.10)$$

where

$$B\left(\frac{\beta}{2}|v_1-v_*|\right) = \left\{ x \in \mathbb{R}^2 : |x| = \frac{\beta}{2}|v_1-v_*| \text{ for } v_1, v_* \in \mathbb{R}^3 \right\}.$$

In (1.31), we replace v, v' with v_1, v'_1 respectively, and choose $\phi(v_1) = D_j(v_1)$. Then we observe that

$$\begin{aligned} &\int_{\mathbb{R}^3} D_j(v_1) Q^+(f, f)(t, v_1) dv_1 \\ &= \frac{1}{\beta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} f(t, v_1) f(t, v_*) |v_1 - v_*|^{-1} \int_{B\left(\frac{\beta}{2}|v_1-v_*|\right)} D_j(v'_1) d\sigma' dv_1 dv_* \\ &\leq \frac{\pi}{\beta} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} f(t, v_1) f(t, v_*) \chi dv_1 dv_* \leq 2\pi \|f\|_{1,0}^2 \end{aligned}$$

by (2.10). Since the bound is uniform in j , we let $j \rightarrow \infty$ and Lemma 2.8, and end of the proof. \square

We rewrite (1.15) as

$$\partial_t f(t, v_1) + Lf(t, v_1) f(t, v_1) = Q^+(f, f)(t, v_1). \quad (2.11)$$

Lemma 2.10. *Assume that (1.21) holds and $f_0 \in L_s^\infty$ for $s > 2$. Then there exist positive constant $C_1(\|f_0\|_{\infty, s})$ and C_2 such that*

$$\int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} \leq C_1(\|f_0\|_{\infty, s}) \min \left\{ 1 + t, e^{C_2\left(\frac{1}{\alpha^2}-1\right)t} \right\}$$

for any $v, v' \in \mathbb{R}^3$ and $t > 0$.

Proof. From (2.11) and Lemma 2.9, we have

$$\partial_t \int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} \leq \int_{v_* \in E_P^{v'-v}} Q^+(f, f)(t, v_*) dE_{v_*} \leq 2\pi \|f_0\|_{1,0}^2$$

for fixed $v, v' \in \mathbb{R}^3$. Note that we have

$$\begin{aligned} \int_{v_* \in E_P^{v'-v}} f_0(v_*) dE_{v_*} &\leq \|f_0\|_{\infty, s} \int_{\mathbb{R}^2} (1 + |v_*|)^{-s} dv_* \\ &\leq 2\pi \|f_0\|_{\infty, s} \int_0^\infty r(1+r)^{-s} dr \\ &\leq 2\pi \|f_0\|_{\infty, s} \left(\int_0^1 1 dr + \int_1^\infty (1+r)^{-s+1} dr \right) \leq 4\pi \|f_0\|_{\infty, s} \end{aligned} \quad (2.12)$$

for $s > 2$. Thus we have

$$\int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} \leq 2\pi \|f_0\|_{1,0}^2 t + 4\pi \|f_0\|_{\infty, s}. \quad (2.13)$$

Next, we apply (2.8) and Lemma 2.9 to (2.11), and obtain

$$\begin{aligned} \partial_t \int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} + C_1 e^{-C_2(\frac{1}{\alpha^2}-1)t} \int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} \\ \leq 2\pi \|f_0\|_{1,0}^2 \end{aligned} \quad (2.14)$$

for some positive constants C_1 and C_2 . Then we obtain

$$\int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} \leq C_3 (\|f_0\|_{\infty, s}) e^{C_2(\frac{1}{\alpha^2}-1)t} \quad (2.15)$$

for some positive constant $C_3(\|f_0\|_{\infty, s})$, by using Lemma 2.1 and (2.12). Lastly, we combine (2.13) and (2.15). \square

In the proof of Lemma 2.11, we divide the region of v_1 into subsets represented by O_1 , O_2 , and O_3 for $v \neq 0$. Specifically, we define:

$$\begin{aligned} O_2 &= \left\{ v_1 \in \mathbb{R}^3 : |v_1| \geq \frac{|v|}{2} \right\} \cap \left\{ v_1 \in \mathbb{R}^3 : |v - v_1| < \frac{1}{2}|v|^{-p} \right\}, \\ O_3 &= \left\{ v_1 \in \mathbb{R}^3 : |v_1| \geq \frac{|v|}{2} \right\} \cap \left\{ v_1 \in \mathbb{R}^3 : |v - v_1| \geq \frac{1}{2}|v|^{-p} \right\}, \end{aligned} \quad (2.16)$$

for some $p \in \mathbb{R}$. To estimate the region of O_3 in (2.19), we need to restrict the range of p to $p + 1 \leq 2$. For the region of O_2 in (2.18), $2p \geq 1$. This allows us to choose p such that $\frac{1}{2} \leq p \leq 1$. In the proof, we set $p = \frac{1}{2}$.

Lemma 2.11. *Assume that (1.21) holds, and $f \in L_0^\infty$. Then there exists a positive constants $C(\|f\|_{\infty, 0})$ such that*

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C(\|f\|_{\infty, 0})(1 + |v|)^{-1} \quad (2.17)$$

for $v \in \mathbb{R}^3$ and $t > 0$.

Proof. Assume $v \neq 0$ holds. We define

$$O_1 = \left\{ v_1 \in \mathbb{R}^3 : |v_1| < \frac{|v|}{2} \right\},$$

and define O_2 and O_3 as (2.16), where $p = \frac{1}{2}$. Since $|v_1 - v| > |v|/2$ for $v_1 \in O_1$, we have

$$\int_{O_1} f(t, v_1) |v - v_1|^{-1} dv_1 \leq 2|v|^{-1} \|f_0\|_{1,0}.$$

For $v_1 \in O_2$, we have

$$\begin{aligned} \int_{O_2} f(t, v_1) |v - v_1|^{-1} dv_1 \\ \leq \|f\|_{0, \infty} \int_{\{v_1 : |v - v_1| < \frac{1}{2}|v|^{-\frac{1}{2}}\}} |v - v_1|^{-1} dv_1 = \frac{1}{2}\pi \|f\|_{0, \infty} |v|^{-1}. \end{aligned} \quad (2.18)$$

For $v_1 \in O_3$, we have

$$\begin{aligned} \int_{O_3} f(t, v_1) |v - v_1|^{-1} dv_1 &= \int_{O_3} f(t, v_1) (1 + |v_1|^2) \frac{1}{1 + |v_1|^2} |v - v_1|^{-1} dv_1 \\ &\leq \|f_0\|_{1,2} \frac{8\sqrt{|v|}}{4 + |v|^2} \leq 2\|f_0\|_{1,2} |v|^{-1}. \end{aligned} \quad (2.19)$$

This implies that

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \sum_{i=1,2,3} \int_{O_i} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C(\|f\|_{\infty,0}) |v|^{-1} \quad (2.20)$$

for some positive constant $C(\|f\|_{\infty,0})$. On the other hand, we have

$$\begin{aligned} &\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \\ &\leq \int_{\mathbb{R}^3} f(t, v_1) (\chi_{\{|v_1|:|v-v_1|\geq 1\}}(v_1) + \chi_{\{|v_1|:|v-v_1|\leq 1\}}(v_1)) |v - v_1|^{-1} dv_1 \\ &\leq \|f_0\|_{1,0} + 2\pi\|f\|_{\infty,0} \end{aligned} \quad (2.21)$$

for $v \in \mathbb{R}^3$. Therefore we obtain (2.17) from (2.20) and (2.21). \square

In Lemma 2.12, we divide the range of v to account for whether it includes zero or not. We cannot apply Lemma 2.4 if $|v|$ does not have a uniform lower bound. Consequently, the time variable t appears on the left side of (2.23), which includes the case where $v = 0$.

Lemma 2.12. *Assume that (1.21) holds and $f_0 \in L_0^\infty$. Then there exist positive constants $C_1(\|f_0\|_{\infty,0})$, $C_2(\|f_0\|_{\infty,0})$ and C_3 such that*

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C_1(\|f_0\|_{\infty,0}) |v|^{-1} \quad (2.22)$$

for $v \neq 0 \in \mathbb{R}^3$ and $t > 0$, and

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C_2(\|f_0\|_{\infty,0}) (1 + |v|)^{-1} \min \left\{ 1 + t, e^{C_3(\frac{1}{\alpha^2} - 1)t} \right\} \quad (2.23)$$

for $v \in \mathbb{R}^3$ and $t > 0$.

Proof. Assume $v \neq 0$ holds. In the case of $|v_1| < |v|/2$, we have

$$\int_{\mathbb{R}^3} f(t, v_1) \chi_{\{|v_1|:|v_1|<\frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \leq 2|v|^{-1} \|f_0\|_{1,0}. \quad (2.24)$$

For $|v_1| \geq |v|/2 > 0$, we have

$$\begin{aligned} &\partial_t \int_{\mathbb{R}^3} f(t, v_1) \chi_{\{|v_1|:|v_1|\geq\frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \\ &\quad + \int_{\mathbb{R}^3} Lf(t, v_1) f(t, v_1) \chi_{\{|v_1|:|v_1|\geq\frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \\ &= \int_{\mathbb{R}^3} Q^+(f, f)(t, v_1) \chi_{\{|v_1|:|v_1|\geq\frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \end{aligned} \quad (2.25)$$

from (2.11). By applying (2.6) and Lemma 2.7 to (2.25), we obtain

$$\begin{aligned} & \partial_t \int_{\mathbb{R}^3} f(t, v_1) \chi_{\{|v_1| \geq \frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \\ & + \pi \frac{|v|}{2} \int_{\mathbb{R}^3} f(t, v_1) \chi_{\{|v_1| \geq \frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \leq 4\pi \|f_0\|_{1,0}^2. \end{aligned} \quad (2.26)$$

From Lemma 2.11, we have

$$\int_{\mathbb{R}^3} f_0(v_1) |v - v_1|^{-1} dv_1 \leq C_1(\|f_0\|_{\infty,0}) (1 + |v|)^{-1} \quad (2.27)$$

for some positive constant $C_1(\|f_0\|_{\infty,0})$, at $t = 0$. Therefore, by applying Lemma 2.1 and (2.27) to (2.26), we obtain

$$\int_{\mathbb{R}^3} f(t, v_1) \chi_{\{|v_1| \geq \frac{|v|}{2}\}}(v_1) |v - v_1|^{-1} dv_1 \leq |v|^{-1} (8\|f_0\|_{1,0}^2 + C_1(\|f_0\|_{\infty,0})). \quad (2.28)$$

We deduce (2.22) from (2.24) and (2.28).

Now we consider all range of $v \in \mathbb{R}^3$. From (2.11) and Lemma 2.7, we have

$$\partial_t \int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq 4\pi \|f_0\|_{1,0}^2. \quad (2.29)$$

Integrating (2.29), we obtain

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq 4\pi \|f_0\|_{1,0}^2 t + \|f_0\|_{1,0} + 2\pi \|f_0\|_{\infty,0} \quad (2.30)$$

from (2.21). Similarly, if we use (2.8), then we have

$$\begin{aligned} & \partial_t \int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \\ & + C_2 e^{-C_3(\frac{1}{\alpha^2}-1)t} \int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq 4\pi \|f_0\|_{1,0}^2 \end{aligned} \quad (2.31)$$

for some positive constants C_2, C_3 , and obtain

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C_4(\|f_0\|_{\infty,0}) e^{C_3(\frac{1}{\alpha^2}-1)t} \quad (2.32)$$

for some positive constant $C_4(\|f_0\|_{\infty,0})$, by Lemma 2.1 and (2.27). For $0 \leq |v| \leq 1$, since $1 \leq 2(1 + |v|)^{-1}$, there is a positive constant $C_5(\|f_0\|_{\infty,0})$ such that

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C_5(\|f_0\|_{\infty,0}) (1 + |v|)^{-1} \min \left\{ 1 + t, e^{C_3(\frac{1}{\alpha^2}-1)t} \right\}$$

from (2.30) and (2.32). For $|v| > 1$, since $|v|^{-1} < 2(1 + |v|)^{-1}$, there is a positive constant $C_6(\|f_0\|_{\infty,0})$ such that

$$\int_{\mathbb{R}^3} f(t, v_1) |v - v_1|^{-1} dv_1 \leq C_6(\|f_0\|_{\infty,0}) (1 + |v|)^{-1}$$

from (2.22). From the two inequalities above, we derive (2.23). \square

Proof of Theorem 1.1. We recall a Carleman type representation (1.33)

$$\begin{aligned} & Q^+(f, f)(t, v_1) \\ & \leq \frac{1}{\beta^2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \delta(v' - v_1) f(t, v) |v' - v|^{-1} \int_{v_* \in E_P^{v'-v}} f(t, v_*) dE_{v_*} dv dv'. \end{aligned} \quad (2.33)$$

By applying Lemma 2.10 and (2.23) to (2.33), we have

$$Q^+(f, f)(t, v_1) \leq C_1(\|f_0\|_{\infty, s})(1 + |v_1|)^{-1} \min \left\{ (1+t)^2, e^{C_2(\frac{1}{\alpha^2}-1)t} \right\} \quad (2.34)$$

for some positive constants $C_1(\|f_0\|_{\infty, s})$ and C_2 . From (2.11) and (2.34), we have

$$\partial_t f(t, v_1) \leq C_1(\|f_0\|_{\infty, s})(1 + |v_1|)^{-1}(1+t)^2,$$

and by integrating with respect to time, then obtain

$$f(t, v_1) \leq C_3(\|f_0\|_{\infty, s})(1 + |v_1|)^{-1}(1+t)^3 \quad (2.35)$$

for some positive constant $C_3(\|f_0\|_{\infty, s})$.

Now, by applying (2.8) and (2.34) to (2.11), we have

$$\begin{aligned} \partial_t f(t, v_1) + C_4 e^{-C_5(\frac{1}{\alpha^2}-1)t}(1 + |v_1|)f(t, v_1) \\ \leq C_1(\|f_0\|_{\infty, s})(1 + |v_1|)^{-1} e^{C_2(\frac{1}{\alpha^2}-1)t} \end{aligned} \quad (2.36)$$

for some positive constant C_4 and C_5 . Then, by multiplying both sides of (2.36) by $(1 + |v_1|)^2$ and applying Lemma 2.1, then we obtain

$$\operatorname{ess\,sup}_{v \in \mathbb{R}^3} f(t, v)(1 + |v|)^2 \leq C_6(\|f_0\|_{\infty, s})e^{C_7(\frac{1}{\alpha^2}-1)t} \quad (2.37)$$

for some positive constants $C_6(\|f_0\|_{\infty, s})$ and C_7 , under $f_0 \in L_s^\infty$ for $s > 2$.

Let us assume $v_1 \neq 0$. By Lemma 2.10 and (2.22), we have

$$Q^+(f, f)(t, v_1) \leq C_8(\|f_0\|_{\infty, s})|v_1|^{-1}(1+t) \quad (2.38)$$

for some positive constant $C_8(\|f_0\|_{\infty, s})$. By (2.6) and (2.38),

$$\partial_t f(t, v_1) + \pi \frac{|v_1|}{2} f(t, v_1) \leq C_8(\|f_0\|_{\infty, s})|v_1|^{-1}(1+t).$$

Then we obtain

$$\operatorname{ess\,sup}_{v \neq 0} f(t, v)|v|^2 \leq C_9(\|f_0\|_{\infty, s})(1+t) \quad (2.39)$$

for some positive constant $C_9(\|f_0\|_{\infty, s})$, by using Lemma 2.1. We derive (1.35) from (2.35) and (2.37), and derive (1.36) from (2.37) and (2.39). \square

Proof of Corollary 1.2. We assume $T_c < +\infty$, where T_c is in (1.34), and let $r > 0$. Since

$$\mathcal{E}(T_c) = \int_{\mathbb{R}^3} f(T_c, v)|v|^2 dv \geq r^2 \int_{\{|v|>r\}} f(T_c, v) dv$$

and $\mathcal{E}(T_c) = 0$, we obtain

$$\int_{\{|v|>r\}} f(T_c, v) dv = 0. \quad (2.40)$$

Using (1.35) and (2.40), we have

$$\begin{aligned} 1 = \int_{\mathbb{R}^3} f(T_c, v) dv &= \int_{\{|v|<r\}} f(T_c, v) dv \\ &\leq \frac{2\pi^2}{3} r^3 \|f(T_c, \cdot)\|_\infty \leq C(\|f_0\|_{\infty, s}) r^3 (1+T_c)^3 \end{aligned}$$

for some positive constant $C(\|f_0\|_{\infty, s})$. It is a contradiction if we choose $r < (C(\|f_0\|_{\infty, s})(1+T_c)^3)^{-1}$, and we get $T_c = +\infty$. \square

This section on the L^∞ estimates of the solution of the inelastic Boltzmann equation concludes here. Using the estimates, we will prove the formation of the Maxwellian upper-bound in the next section.

3. MAXWELLIAN UPPER BOUND FOR THE INELASTIC BOLTZMANN EQUATION

In this section, we will extend Theorem 1.1 and obtain Maxwellian upper bounds for the inelastic Boltzmann equation for each time. In [32], Gamba, Panferov, and Villani proved Lemma 3.1 for the homogeneous Boltzmann equation. Since (1.16) also holds for the inelastic collision operator, we can prove Lemma 3.1 similarly. Lemma 3.1 is a significant component in the proof of Theorem 1.5.

Lemma 3.1. *Let f, u satisfy*

$$f(t, v) \geq 0 \quad \text{and} \quad u(0, v) \leq 0 \quad (3.1)$$

for $t > 0$ and almost all $v \in \mathbb{R}^3$. Assume that

$$\partial_t u - Q(f, u) \leq 0 \quad \text{on} \quad |v| > R \quad (3.2)$$

and

$$u \leq 0 \quad \text{on} \quad 0 \leq |v| \leq R \quad (3.3)$$

for some $0 < R < \infty$ and $t > 0$. Then, we have $u \leq 0$ for $t > 0$ and almost all $v \in \mathbb{R}^3$.

Proof. We define $u_+(t, v) = \max\{u(t, v), 0\}$. The function $\text{sign } z$ is defined as 1 for $z > 0$, -1 for $z < 0$ and arbitrary fixed value in $[-1, 1]$ for $z = 0$. We have

$$\int_{\mathbb{R}^3} u_+(t, v) - u_+(0, v) \, dv = \int_0^t \int_{\mathbb{R}^3} \partial_t u \times \frac{1}{2} (\text{sign } u(s, v) + 1) \, dv ds.$$

By (1.16), (3.2), (3.3) and $u_+(0, v) = 0$ from (3.1), we have

$$\begin{aligned} \int_{\mathbb{R}^3} u_+(t, v) \, dv &\leq \int_0^t \int_{\mathbb{R}^3} Q(f, u) \frac{1}{2} (\text{sign } u(s, v) + 1) \, dv ds \\ &= \frac{1}{2} \int_0^t \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v - v_*) \cdot n| f(s, v_*) u(s, v) \\ &\quad \times (\text{sign } u(s, v') - \text{sign } u(s, v)) \, dndv_* dv ds \leq 0. \end{aligned}$$

Because $u(s, v)(\text{sign } u(s, v') - \text{sign } u(s, v)) \leq 0$ for every $0 < s < t$, $v \in \mathbb{R}^3$, we obtain $u_+(t, v) \leq 0$ for almost all $v \in \mathbb{R}^3$. \square

To apply Lemma 3.1 to Theorem 1.5, we need to find $R > 0$ in (3.2), where $u(t, v) = f(t, v) - \bar{M}(t, v)$ for some Maxwellian-type time-dependent distribution $\bar{M}(t, v)$. Later, we will determine the specific form of $\bar{M}(t, v)$. In the proof of Lemma 3.2 below, we use the inequality $M(v)M(v_*) \leq M(v')M(v'_*)$ for a global Maxwellian M which holds by (1.14).

Lemma 3.2. *Let $M(v_1) = e^{-a|v_1|^2 + b}$ for $a > 0$. We assume that*

$$\int_{\mathbb{R}^3} f(t, v) e^{2a|v|^2} \, dv \leq C \quad (3.4)$$

for some constant $C > 0$ and $t > 0$. Then, there exists $0 < R < \infty$ such that

$$Q(f, M)(t, v_1) \leq 0$$

for $|v_1| > R$. Here R depends on C and a , but not on b .

Proof. We plug $M(v)$ into $g(t, v)$ of (1.24) to obtain

$$Q^+(f, M)(t, v_1) = \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}_+^2} |(v - v_*) \cdot n| f(t, v_*) M(v) \delta(v' - v_1) \, dndv_* dv.$$

We also recall in (1.17) that

$$\cos \theta = \left(\frac{v - v_*}{|v - v_*|}, n \right), \quad \theta \in \left[0, \frac{\pi}{2} \right],$$

and in (1.6) that

$$\beta = \frac{1 + \alpha}{2}, \quad \frac{1}{2} < \beta \leq 1.$$

From Figure 4, we obtain

$$\cos \theta = \frac{|v_* - v'_*|}{\beta |v - v_*|}, \quad \text{and} \quad \sin \theta = \frac{|v_* - P|}{|v - v_*|} \quad (3.5)$$

for

$$P = \frac{1}{\beta} v' - \left(\frac{1}{\beta} - 1 \right) v, \quad (3.6)$$

and

$$|v' - v_*| \geq (1 - \beta) |v - v_*|. \quad (3.7)$$

By the fact that $M(v)M(v_*) \leq M(v')M(v'_*)$ and (1.27), we have

$$\begin{aligned} & Q^+(f, M)(t, v_1) \\ &= \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*| \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) \, d\sigma dv_* dv. \end{aligned} \quad (3.8)$$

In terms of (v', v'_*) in (1.8), we define

$$A = A(v, v_*, \sigma; \beta) := \left\{ v, v_* \in \mathbb{R}^3, \sigma \in \mathbb{S}^2 : \frac{1}{\beta} - 1 \leq \frac{1}{2} \frac{|v' - v_*|}{|v_* - v'_*|} \right\} \quad (3.9)$$

and

$$A^c = A^c(v, v_*, \sigma; \beta) := \left\{ v, v_* \in \mathbb{R}^3, \sigma \in \mathbb{S}^2 : \frac{1}{\beta} - 1 \geq \frac{1}{2} \frac{|v' - v_*|}{|v_* - v'_*|} \right\}. \quad (3.10)$$

Moreover, we split $Q^+(f, M)(t, v_1)$ into $Q_A^+(f, M)(t, v_1)$ and $Q_{A^c}^+(f, M)(t, v_1)$, where

$$\begin{aligned} & Q_A^+(f, M)(t, v_1) \\ &:= \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*| \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) \mathbf{1}_A \, d\sigma dv_* dv \end{aligned}$$

and

$$\begin{aligned} & Q_{A^c}^+(f, M)(t, v_1) \\ &:= \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*| \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) \mathbf{1}_{A^c} \, d\sigma dv_* dv. \end{aligned}$$

We now estimate $Q_A^+(f, M)(t, v_1)$ for $\frac{1}{2} < \beta < 1$. By (3.5), we have

$$\begin{aligned} & Q_A^+(f, M)(t, v_1) \\ & \leq \frac{1}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*|^{1-\epsilon_1+\epsilon_2} \frac{|v_* - v'_*|^{\epsilon_1} \sin^{\epsilon_2} \theta}{|v_* - P|^{\epsilon_2} \cos^{\epsilon_1} \theta} \mathbf{1}_A \\ & \quad \times \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) d\sigma dv_* dv \end{aligned} \quad (3.11)$$

for $0 < \epsilon_1, \epsilon_2 < 1$. For $v, v_*, \sigma \in A$, we have

$$\begin{aligned} |v_* - P| & \geq |v' - v_*| - |v' - P| \\ & = |v' - v_*| - \left(\frac{1}{\beta} - 1\right) |v_* - v'_*| \geq \frac{1}{2} |v' - v_*|. \end{aligned} \quad (3.12)$$

Applying (3.12) to (3.11), and (3.7) to (3.14), respectively, we have

$$\text{(RHS) of (3.11)} \quad (3.13)$$

$$\begin{aligned} & \leq \frac{2^{\epsilon_2}}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} \frac{|v - v_*|^{1-\epsilon_1+\epsilon_2}}{|v' - v_*|^{\epsilon_2}} |v_* - v'_*|^{\epsilon_1} \frac{\sin^{\epsilon_2} \theta}{\cos^{\epsilon_1} \theta} \\ & \quad \times \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) d\sigma dv_* dv \end{aligned} \quad (3.14)$$

$$\begin{aligned} & \leq \frac{2^{\epsilon_2}}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \left(\frac{1}{1-\beta}\right)^{\epsilon_2} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*|^{1-\epsilon_1} |v_* - v'_*|^{\epsilon_1} \frac{\sin^{\epsilon_2} \theta}{\cos^{\epsilon_1} \theta} \\ & \quad \times \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) d\sigma dv_* dv. \end{aligned} \quad (3.15)$$

By using the fact that $|v - v_*| \leq |v' - v_*| + |v - v'| = |v' - v_*| + |v_* - v'_*|$, $|v'_*| M(v'_*) \lesssim 1$, and $|v_*| e^{-a|v_*|^2} \lesssim 1$, we have

$$\begin{aligned} (3.15) & \lesssim \frac{2^{\epsilon_2}}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \left(\frac{1}{1-\beta}\right)^{\epsilon_2} \int_{\mathbb{R}^3} f(t, v_*) e^{2a|v_*|^2} \\ & \quad \times \int_{\mathbb{S}^2} \frac{\sin^{\epsilon_2} \theta}{\cos^{\epsilon_1} \theta} \int_{\mathbb{R}^3} (1 + |v'|)^{1-\epsilon_1} M(v') \delta(v' - v_1) dv d\sigma dv_*. \end{aligned} \quad (3.16)$$

From (1.8), we can calculate the Jacobian determinant for the map between v and v' for fixed σ as

$$\begin{aligned} \left| \frac{dv'}{dv} \right| & = \left(1 - \frac{\beta}{2}\right)^3 \left(1 + \frac{\beta}{2-\beta} \frac{v - v_*}{|v - v_*|} \cdot \sigma\right) \\ & = \left(1 - \frac{\beta}{2}\right)^3 \left(1 - \frac{\beta}{2-\beta} + \frac{2\beta}{2-\beta} \sin^2 \theta\right) > 0 \end{aligned} \quad (3.17)$$

since (1.26). Applying (3.17) to (3.16), and $\frac{1}{2} < \beta < 1$, we have

$$\text{(RHS) of(3.16)} \tag{3.18}$$

$$\begin{aligned} &\lesssim \frac{2^{\epsilon_2}}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \left(\frac{1}{1-\beta}\right)^{\epsilon_2} \left(1 - \frac{\beta}{2}\right)^{-3} \int_{\mathbb{R}^3} f(t, v_*) e^{2a|v_*|^2} dv_* \\ &\quad \times \int_{\mathbb{R}^3} (1 + |v'|)^{1-\epsilon_1} M(v') \delta(v' - v_1) dv' \\ &\quad \times \int_0^{\frac{\pi}{2}} \sin 2\theta \frac{\sin^{\epsilon_2} \theta}{\cos^{\epsilon_1} \theta} \left(\frac{2\beta}{2-\beta} \sin^2 \theta\right)^{-1} d\theta \\ &\lesssim \left(\frac{1}{1-\beta}\right)^{\epsilon_2} M(v_1) (1 + |v_1|)^{1-\epsilon_1} \int_{\mathbb{R}^3} f(t, v_*) e^{2a|v_*|^2} dv_* \\ &\quad \times \left(\int_0^{\frac{\pi}{4}} \sin^{\epsilon_2-1} \theta d\theta + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cos^{1-\epsilon_1} \theta d\theta \right) \end{aligned} \tag{3.19}$$

for $0 < \epsilon_1, \epsilon_2 < 1$. Taking $\epsilon_2 = 1 - \beta$ and $0 < \epsilon_1 < 1$ in (3.19) and using (3.4), we have

$$Q_A^+(f, M)(t, v_1) \lesssim M(v_1) (1 + |v_1|)^{1-\epsilon_1} \tag{3.20}$$

since $(1 - \beta)^{-(1-\beta)} \lesssim 1$.

Next, we estimate $Q_{A^c}^+(f, M)(t, v_1)$ for $\frac{1}{2} < \beta < 1$. By (3.5), we have

$$\begin{aligned} &Q_{A^c}^+(f, M)(t, v_1) \\ &\leq \frac{1}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} |v - v_*|^{1-\epsilon_1} |v_* - v'_*|^{\epsilon_1} \frac{1}{\cos^{\epsilon_1} \theta} \mathbf{1}_{A^c} \\ &\quad \times \frac{f(t, v_*)}{M(v_*)} M(v'_*) M(v') \delta(v' - v_1) d\sigma dv_* dv \end{aligned} \tag{3.21}$$

for $0 < \epsilon_1 < 1$. By using the fact that $|v - v_*| \leq |v' - v_*| + |v_* - v'_*|$, $|v'_*| M(v'_*) \lesssim 1$, and $|v_*| e^{-a|v_*|^2} \lesssim 1$, we have

$$\begin{aligned} \text{(RHS) of(3.21)} &\lesssim \frac{1}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \int_{\mathbb{R}^3} f(t, v_*) e^{2a|v_*|^2} \\ &\quad \times \int_{\mathbb{S}^2} \frac{1}{\cos^{\epsilon_1} \theta} \int_{\mathbb{R}^3} (1 + |v'|)^{1-\epsilon_1} M(v') \delta(v' - v_1) \mathbf{1}_{A^c} dv d\sigma dv_*. \end{aligned} \tag{3.22}$$

From (3.10), we have

$$\begin{aligned} \frac{1}{\beta} - 1 &\geq \frac{1}{2} \frac{|v' - v_*|}{|v_* - v'_*|} \geq \frac{1}{2} \left(\frac{|v - v_*|}{|v_* - v'_*|} - \frac{|v - v'|}{|v_* - v'_*|} \right) \\ &= \frac{1}{2} \left(\frac{1}{\beta \cos \theta} - 1 \right) > 0 \end{aligned}$$

for $v, v_*, \sigma \notin A$, and it yields that

$$\cos \theta \geq \frac{1}{2 - \beta}. \tag{3.23}$$

From (3.17), and obtain

$$\left| \frac{dv'}{dv} \right| \geq \left(1 - \frac{\beta}{2}\right)^3 \left(1 - \frac{\beta}{2-\beta}\right) = 2 \left(1 - \frac{\beta}{2}\right)^3 \frac{1-\beta}{2-\beta} > 0. \quad (3.24)$$

Applying (3.24) to (3.22), and using (3.23), we have

$$\begin{aligned} (\text{RHS of (3.22)}) &\lesssim \frac{1}{4} \left(\frac{1}{\beta}\right)^{\epsilon_1} \frac{2-\beta}{1-\beta} \left(1 - \frac{\beta}{2}\right)^{-3} \int_{\mathbb{R}^3} f(t, v_*) e^{2a|v_*|^2} dv_* \\ &\quad \times \int_{\mathbb{R}^3} (1 + |v'|)^{1-\epsilon_1} M(v') \delta(v' - v_1) dv' \\ &\quad \times \int_0^{\frac{\pi}{2}} \sin 2\theta \frac{1}{\cos^{\epsilon_1} \theta} \mathbf{1}_{\{\cos \theta \geq \frac{1}{2-\beta}\}} d\theta. \end{aligned} \quad (3.25)$$

Because

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \sin 2\theta \frac{1}{\cos^{\epsilon_1} \theta} \mathbf{1}_{\{\cos \theta \geq \frac{1}{2-\beta}\}} d\theta &\leq 2 \int_0^{\frac{\pi}{2}} \sin \theta \cos^{1-\epsilon_1} \theta \mathbf{1}_{\{\cos \theta \geq \frac{1}{2-\beta}\}} d\theta \\ &\leq 2 \int_{\frac{1}{2-\beta}}^1 t^{1-\epsilon_1} dt \leq 2 \frac{1-\beta}{2-\beta} \end{aligned}$$

for $0 < \epsilon_1 < 1$, from (3.25), we have

$$Q_{A^c}^+(f, M)(t, v_1) \lesssim M(v_1) (1 + |v_1|)^{1-\epsilon_1} \quad (3.26)$$

since $\frac{1}{2} < \beta < 1$. Combining (3.20) with (3.26), there exists some constant $C_1 > 0$ such that

$$\begin{aligned} Q^+(f, M)(f, M) &= Q_A^+(f, M)(t, v_1) + Q_{A^c}^+(f, M)(t, v_1) \\ &\leq C_1 M(v_1) (1 + |v_1|)^{1-\epsilon_1} \end{aligned}$$

for $0 < \epsilon_1 < 1$. From Lemma 2.4, we have

$$Q^-(f, M)(t, v_1) = M(v_1) Lf(t, v_1) \geq \pi M(v_1) |v_1|.$$

Therefore we have

$$Q(f, M)(t, v_1) \leq M(v_1) (C_1 (1 + |v_1|)^{1-\epsilon_1} - \pi |v_1|).$$

Then we choose $R > 0$ sufficiently large such that $Q(f, M) \leq 0$, and it concludes the proof for $0 < \beta < 1$. In Lemma 5 of [32], the authors proved an analogue of this Lemma in the case of elastic collisions, when $\beta = 1$. \square

Now we are ready to prove the Maxwellian upper bound of the solution. We apply Lemma 3.2 to (3.2) and use the inequality

$$f(t, v) \leq C_1 (\|f_0\|_{\infty, s}) \min \left\{ (1+t)^3, e^{C_2 \left(\frac{1}{\alpha^2} - 1\right)t} \right\} \quad (3.27)$$

for some positive constants $C_1(\|f_0\|_{\infty, s})$, C_2 , from Theorem 1.1 to (3.3). Additionally, (3.1) arises from the assumption of the initial bound.

Proof of Theorem 1.5. There exists a positive constant K_1 , which depends on c_0 such that

$$\int_{\mathbb{R}^3} f_0(v) e^{\frac{\alpha_0}{2}|v|^2} dv \leq K_1. \quad (3.28)$$

since $f_0(v) \leq M_0(v)$. By Proposition 3.2-(ii) in [54], and $T_c = +\infty$ in Corollary 1.2 or Remark 1.4, and (3.28), there exist $K_2, a_1 > 0$, which depend on a_0 and K_1 such that

$$\sup_{t \in [0, \infty)} \int_{\mathbb{R}^3} f(t, v) e^{a_1 |v|^2} dv \leq K_2. \quad (3.29)$$

We take $a = \min\{a_0, \frac{a_1}{2}\}$ and $b \geq c_0$ and set

$$u(t, v) = f(t, v_1) - \overline{M}(t, v_1), \quad \text{where } \overline{M}(t, v_1) = (1+t)^3 e^{-a|v_1|^2+b}.$$

Then we have

$$u(0, v) = f(0, v_1) - \overline{M}(0, v_1) \leq f_0(v_1) - M_0(v_1) \leq 0,$$

since $a < a_0$ and $c_0 \leq b$. By (3.29) and Lemma 3.2, there exists a large $R > 0$, which depends on a_1, K_2 such that

$$\partial_t u - Q(f, u)(v_1) = -\partial_t \overline{M} + Q(f, \overline{M})(t, v_1) \leq 0$$

for $|v_1| \geq R$. From (3.27), we have $f(t, v_1) \leq C_1(t+1)^3$, for some constant C_1 depending on c_0 . Taking $b = aR^2 + \log C_1 (\geq c_0)$, it holds that

$$\overline{M}(t, v_1) = C_1(1+t)^3 e^{-a|v_1|^2+aR^2} \geq f(t, v_1)$$

for $|v_1| < R$. Therefore we apply Lemma 3.1 and obtain

$$f(t, v_1) \leq \overline{M}(t, v_1) = (1+t)^3 e^{-a|v_1|^2+b}. \quad (3.30)$$

Moreover, we replace

$$\overline{M}(t, v_1) = (1+t)^3 e^{-a|v_1|^2+b}$$

with

$$\widehat{M}(t, v_1) = e^{C_2(\frac{1}{a^2}-1)t} e^{-a|v_1|^2+b},$$

where the positive constant C_2 is used in (3.27), and depends on c_0 . Then, by following the same argument as above, we obtain

$$f(t, v_1) \leq \widehat{M}(t, v_1) = e^{C_2(\frac{1}{a^2}-1)t} e^{-a|v_1|^2+b} \quad (3.31)$$

instead of (3.30). From (3.30) and (3.31), we conclude

$$f(t, v_1) \leq \min \left\{ (1+t)^3, e^{C_2(\frac{1}{a^2}-1)t} \right\} e^{-a|v_1|^2+b}$$

for almost every $v_1 \in \mathbb{R}^3$ and every $t > 0$. \square

ACKNOWLEDGMENT

The first and the third authors are supported by the National Research Foundation of Korea(NRF) grants RS-2023-00212304 and RS-2023-00219980. The second author is supported by the National Research Foundation of Korea (NRF) grants RS-2023-00210484, RS-2023-00219980, 2022R1G1A1009044 and 2021R1A6A1A10042944, and also by Posco Holdings Inc. and Samsung Electronics Co., Ltd. (IO230412-05903-01).

REFERENCES

- [1] Radjesvarane Alexandre, Yoshinori Morimoto, Seiji Ukai, Chao Jiang Xu, and Tong Yang, *Global existence and full regularity of the Boltzmann equation without angular cutoff*, *Comm. Math. Phys.* **304** (2011), no. 2, 513–581, doi:10.1007/s00220-011-1242-9.
- [2] ———, *The Boltzmann equation without angular cutoff in the whole space: I, Global existence for soft potential*, *J. Funct. Anal.* **262** (2012), no. 3, 915–1010, doi:10.1016/j.jfa.2011.10.007.
- [3] Radjesvarane Alexandre and Cédric Villani, *On the Boltzmann equation for long-range interactions*, *Comm. Pure Appl. Math.* **55** (2002), no. 1, 30–70, doi:10.1002/cpa.10012.
- [4] Ricardo Jose Alonso, *Existence of global solutions to the Cauchy problem for the inelastic Boltzmann equation with near-vacuum data*, *Indiana Univ. Math. J.* **58** (2009), no. 3, 999–1022, doi:10.1512/iumj.2009.58.3506.
- [5] Ricardo Jose Alonso and Bertrand Lods, *Free cooling and high-energy tails of granular gases with variable restitution coefficient*, *SIAM J. Math. Anal.* **42** (2010), no. 6, 2499–2538, doi:10.1137/100793979.
- [6] ———, *Boltzmann model for viscoelastic particles: asymptotic behavior, pointwise lower bounds and regularity*, *Comm. Math. Phys.* **331** (2014), no. 2, 545–591, doi:10.1007/s00220-014-2089-7.
- [7] Gayoung An and Donghyun Lee, *High-velocity tails of the inelastic and the multi-species mixture Boltzmann equations*, *SIAM J. Math. Anal.* **55** (2023), no. 5, 4297–4336, doi:10.1137/22M1529749.
- [8] Leif Arkeryd, *L^∞ estimates for the space-homogeneous Boltzmann equation*, *J. Statist. Phys.* **31** (1983), no. 2, 347–361, doi:10.1007/BF01011586.
- [9] ———, *On the long time behaviour of the Boltzmann equation in a periodic box*, Chalmers Tekniska Högskola/Göteborgs Universitet. Department of Mathematics, 1988.
- [10] Dario Benedetto and Emanuele Caglioti, *The collapse phenomenon in one-dimensional inelastic point particle systems*, *Phys. D* **132** (1999), no. 4, 457–475, doi:10.1016/S0167-2789(99)00056-1.
- [11] Dario Benedetto, Emanuele Caglioti, and Mario Pulvirenti, *A kinetic equation for granular media*, *RAIRO Modél. Math. Anal. Numér.* **31** (1997), no. 5, 615–641, doi:10.1051/m2an/1997310506151.
- [12] Dario Benedetto and Mario Pulvirenti, *On the one-dimensional Boltzmann equation for granular flows*, *M2AN Math. Model. Numer. Anal.* **35** (2001), no. 5, 899–905, doi:10.1051/m2an:2001141.
- [13] Marzia Bisi, José Antonio Carrillo, and Giuseppe Toscani, *Decay rates in probability metrics towards homogeneous cooling states for the inelastic Maxwell model*, *J. Stat. Phys.* **124** (2006), no. 2-4, 625–653, doi:10.1007/s10955-006-9035-9.
- [14] Aleksandr Vasilévich Bobylev, *Moment inequalities for the Boltzmann equation and applications to spatially homogeneous problems*, *J. Statist. Phys.* **88** (1997), no. 5-6, 1183–1214, doi:10.1007/BF02732431.
- [15] Aleksandr Vasilévich Bobylev, José Antonio Carrillo, and Irene Martínez Gamba, *On some properties of kinetic and hydrodynamic equations for inelastic interactions*, *J. Statist. Phys.* **98** (2000), no. 3-4, 743–773, doi:10.1023/A:1018627625800.
- [16] Aleksandr Vasilévich Bobylev, Irene Martínez Gamba, and Vladislav Panferov, *Moment inequalities and high-energy tails for Boltzmann equations with inelastic interactions*, *J. Statist. Phys.* **116** (2004), no. 5-6, 1651–1682, doi:10.1023/B:JOSS.0000041751.11664.ea.
- [17] Marc Briant, *Instantaneous exponential lower bound for solutions to the Boltzmann equation with Maxwellian diffusion boundary conditions*, *Kinet. Relat. Models* **8** (2015), no. 2, 281–308, doi:10.3934/krm.2015.8.281.
- [18] ———, *Instantaneous filling of the vacuum for the full Boltzmann equation in convex domains*, *Arch. Ration. Mech. Anal.* **218** (2015), no. 2, 985–1041, doi:10.1007/s00205-015-0874-x.
- [19] Marc Briant and Yan Guo, *Asymptotic stability of the Boltzmann equation with Maxwell boundary conditions*, *J. Differential Equations* **261** (2016), no. 12, 7000–7079, doi:10.1016/j.jde.2016.09.014.

- [20] Nikolai V. Brilliantov and Thorsten Pöschel, *Kinetic theory of granular gases*, Oxford Graduate Texts, Oxford University Press, Oxford, 2004, doi:10.1093/acprof:oso/9780198530381.001.0001.
- [21] Torsten Carleman, *Sur la théorie de l'équation intégrodifférentielle de Boltzmann*, Acta Math. **60** (1933), no. 1, 91–146, doi:10.1007/BF02398270.
- [22] ———, *Problèmes mathématiques dans la théorie cinétique des gaz*, Publ. Sci. Inst. Mittag-Leffler, vol. 2, Almqvist & Wiksells Boktryckeri AB, Uppsala, 1957.
- [23] Sunho Choi and Seungyeal Ha, *Global existence of classical solutions to the inelastic Vlasov-Poisson-Boltzmann system*, J. Stat. Phys. **156** (2014), no. 5, 948–974, doi:10.1007/s10955-014-1041-8.
- [24] Elena Demattè, *On a kinetic equation describing the behavior of a gas interacting mainly with radiation*, J. Stat. Phys. **190** (2023), no. 7, Paper No. 124, 45, doi:10.1007/s10955-023-03128-0.
- [25] Laurent Desvillettes, *Convergence to equilibrium in large time for Boltzmann and B.G.K. equations*, Arch. Rational Mech. Anal. **110** (1990), no. 1, 73–91, doi:10.1007/BF00375163.
- [26] Laurent Desvillettes and Cédric Villani, *On the trend to global equilibrium for spatially inhomogeneous kinetic systems: the Boltzmann equation*, Invent. Math. **159** (2005), no. 2, 245–316, doi:10.1007/s00222-004-0389-9.
- [27] Ronald J. DiPerna and Pierre-Louis Lions, *On the Cauchy problem for Boltzmann equations: global existence and weak stability*, Ann. of Math. (2) **130** (1989), no. 2, 321–366, doi:10.2307/1971423.
- [28] Théophile Dolmaire and Juan José López Velázquez, *Collapse of inelastic hard spheres in dimension $d \geq 2$* , 2024, arXiv:2402.13803.
- [29] ———, *Properties of some dynamical systems for three collapsing inelastic particles*, 2024, arXiv:2403.16905.
- [30] Nicolas Fournier, *On exponential moments of the homogeneous Boltzmann equation for hard potentials without cutoff*, Comm. Math. Phys. **387** (2021), no. 2, 973–994, doi:10.1007/s00220-021-04205-9.
- [31] Irene Martínez Gamba, Vladislav Panferov, and Cédric Villani, *On the Boltzmann equation for diffusively excited granular media*, Comm. Math. Phys. **246** (2004), no. 3, 503–541, doi:10.1007/s00220-004-1051-5.
- [32] ———, *Upper Maxwellian bounds for the spatially homogeneous Boltzmann equation*, Arch. Ration. Mech. Anal. **194** (2009), no. 1, 253–282, doi:10.1007/s00205-009-0250-9.
- [33] Harold Grad, *Principles of the kinetic theory of gases*, Handbuch der Physik [Encyclopedia of Physics], vol. Bd. 12, Thermodynamik der, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1958, Herausgegeben von S. Flügge, pp. 205–294.
- [34] ———, *Asymptotic theory of the Boltzmann equation. II*, Rarefied Gas Dynamics (Proc. 3rd Internat. Sympos., Palais de l'UNESCO, Paris, 1962), Vol. I, Adv. Appl. Mech., vol. Supplement 2, Academic Press, New York-London, 1963, pp. 26–59.
- [35] Philip Gressman and Robert Strain, *Global classical solutions of the Boltzmann equation without angular cut-off*, Journal of the American Mathematical Society **24** (2011), no. 3, 771–847.
- [36] Yan Guo, *The Landau equation in a periodic box*, Comm. Math. Phys. **231** (2002), no. 3, 391–434, doi:10.1007/s00220-002-0729-9.
- [37] ———, *The Vlasov-Poisson-Boltzmann system near Maxwellians*, Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences **55** (2002), no. 9, 1104–1135.
- [38] ———, *The Vlasov-Maxwell-Boltzmann system near Maxwellians*, Inventiones mathematicae **153** (2003), no. 3, 593–630.
- [39] ———, *Decay and Continuity of the Boltzmann Equation in Bounded Domains*, Archive for rational mechanics and analysis **197** (2010), 713–809.
- [40] ———, *The Vlasov-Poisson-Landau system in a periodic box*, Journal of the American Mathematical Society **25** (2012), no. 3, 759–812.
- [41] Reinhard Illner and Marvin Shinbrot, *The Boltzmann equation: global existence for a rare gas in an infinite vacuum*, Comm. Math. Phys. **95** (1984), no. 2, 217–226.
- [42] Cyril Imbert, Clément Mouhot, and Luis Silvestre, *Decay estimates for large velocities in the Boltzmann equation without cutoff*, J. Éc. polytech. Math. **7** (2020), 143–184, doi:10.5802/jep.113.

- [43] ———, *Gaussian lower bounds for the Boltzmann equation without cutoff*, SIAM J. Math. Anal. **52** (2020), no. 3, 2930–2944, doi:10.1137/19M1252375.
- [44] Cyril Imbert and Luis Silvestre, *The weak Harnack inequality for the Boltzmann equation without cut-off*, 2019, arXiv:1608.07571.
- [45] Jin Woo Jang, Bernhard Kepka, Alessia Nota, and Juan José López Velázquez, *Vanishing angular singularity limit to the hard-sphere Boltzmann equation*, J. Stat. Phys. **190** (2023), no. 4, Paper No. 77, 14, doi:10.1007/s10955-023-03089-4.
- [46] Jin Woo Jang and Kunlun Qi, *Moments creation for the inelastic Boltzmann equation for hard potentials without angular cutoff*, arXiv preprint, 2022, arXiv:2206.09636.
- [47] Jin Woo Jang, Robert M. Strain, and Seok-Bae Yun, *Propagation of uniform upper bounds for the spatially homogeneous relativistic Boltzmann equation*, Arch. Ration. Mech. Anal. **241** (2021), no. 1, 149–186, doi:10.1007/s00205-021-01649-0.
- [48] Jin Woo Jang and Juan José López Velázquez, *LTE and non-LTE solutions in gases interacting with radiation*, J. Stat. Phys. **186** (2022), no. 3, Paper No. 47, 62, doi:10.1007/s10955-022-02888-5.
- [49] Shmuel Kaniell and Marvin Shinbrot, *The Boltzmann equation. I. Uniqueness and local existence*, Comm. Math. Phys. **58** (1978), no. 1, 65–84.
- [50] Chanwoo Kim and Donghyun Lee, *The Boltzmann equation with specular boundary condition in convex domains*, Comm. Pure Appl. Math. **71** (2018), no. 3, 411–504, doi:10.1002/cpa.21705.
- [51] Tai-Ping Liu, Tong Yang, and Shih-Hsien Yu, *Energy method for Boltzmann equation*, Phys. D **188** (2004), no. 3-4, 178–192, doi:10.1016/j.physd.2003.07.011.
- [52] Stéphane Mischler and Clément Mouhot, *Cooling process for inelastic Boltzmann equations for hard spheres. II. Self-similar solutions and tail behavior*, J. Stat. Phys. **124** (2006), no. 2-4, 703–746, doi:10.1007/s10955-006-9097-8.
- [53] ———, *Stability, convergence to self-similarity and elastic limit for the Boltzmann equation for inelastic hard spheres*, Comm. Math. Phys. **288** (2009), no. 2, 431–502, doi:10.1007/s00220-009-0773-9.
- [54] Stéphane Mischler, Clément Mouhot, and Mariano Rodríguez Ricard, *Cooling process for inelastic Boltzmann equations for hard spheres. I. The Cauchy problem*, J. Stat. Phys. **124** (2006), no. 2-4, 655–702, doi:10.1007/s10955-006-9096-9.
- [55] Clément Mouhot and Lukas Neumann, *Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus*, Nonlinearity **19** (2006), no. 4, 969–998, doi:10.1088/0951-7715/19/4/011.
- [56] Ada Pulvirenti and Bernt Wennberg, *A Maxwellian lower bound for solutions to the Boltzmann equation*, Comm. Math. Phys. **183** (1997), no. 1, 145–160, doi:10.1007/BF02509799.
- [57] Kunlun Qi, *Measure valued solution to the spatially homogeneous Boltzmann equation with inelastic long-range interactions*, J. Math. Phys. **63** (2022), no. 2, Paper No. 021503, 22, doi:10.1063/5.0062859.
- [58] Luis Silvestre, *A new regularization mechanism for the Boltzmann equation without cut-off*, Comm. Math. Phys. **348** (2016), no. 1, 69–100, doi:10.1007/s00220-016-2757-x.
- [59] Robert M. Strain, *The Vlasov-Maxwell-Boltzmann system in the whole space*, Comm. Math. Phys. **268** (2006), no. 2, 543–567, doi:10.1007/s00220-006-0109-y.
- [60] Robert M. Strain and Yan Guo, *Almost exponential decay near Maxwellian*, Comm. Partial Differential Equations **31** (2006), no. 1-3, 417–429, doi:10.1080/03605300500361545.
- [61] Robert M. Strain and Seok-Bae Yun, *Spatially homogeneous Boltzmann equation for relativistic particles*, SIAM J. Math. Anal. **46** (2014), no. 1, 917–938, doi:10.1137/130923531.
- [62] Giuseppe Toscani and Cédric Villani, *Sharp entropy dissipation bounds and explicit rate of trend to equilibrium for the spatially homogeneous Boltzmann equation*, Comm. Math. Phys. **203** (1999), no. 3, 667–706, doi:10.1007/s002200050631.
- [63] Seiji Ukai, *On the existence of global solutions of mixed problem for non-linear Boltzmann equation*, Proc. Japan Acad. **50** (1974), 179–184.
- [64] Cédric Villani, *Cercignani’s conjecture is sometimes true and always almost true*, Comm. Math. Phys. **234** (2003), no. 3, 455–490, doi:10.1007/s00220-002-0777-1.
- [65] ———, *Mathematics of granular materials*, J. Stat. Phys. **124** (2006), no. 2-4, 781–822, doi:10.1007/s10955-006-9038-6.

DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH),
SOUTH KOREA

Email address: agy19@postech.ac.kr

DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH),
SOUTH KOREA

Email address: jangjw@postech.ac.kr

DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH),
SOUTH KOREA

Email address: donglee@postech.ac.kr