

Sentences over Random Groups II: Sentences of Minimal Rank

Sobhi Massalha

June 24, 2024

Abstract

Random groups of density $d < \frac{1}{2}$ are infinite hyperbolic, and of density $d > \frac{1}{2}$ are finite. We prove the existence of a uniform quantifier elimination procedure for formulas of minimal rank (probably the superstable part of the theory). Namely, given a minimal rank formula $V(p)$, we prove the existence of a formula $\varphi(p)$ that belongs to the Boolean algebra of two quantifiers, so that the two formulas $V(p)$ and $\varphi(p)$ define the same set over the free group F_k and over a random group of density $d < \frac{1}{2}$. We conclude that any given sentence of minimal rank is a truth sentence over the free group F_k if and only if it is a truth sentence over random groups of density $d < \frac{1}{2}$.

Contents

| | | |
|-----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Limit Groups over Sequences of Random Groups | 4 |
| 3 | Graded Test Sequences over Random Groups | 5 |
| 3.1 | Limit Actions of Sequences over Random Groups | 5 |
| 3.2 | Graded Test Sequences over Random Groups | 6 |
| 4 | Rigid and Solid Limit Groups | 7 |
| 5 | Formal Solutions over Random Groups | 9 |
| 5.1 | The Construction of Graded Formal MR Diagram over Random Groups | 10 |
| 5.2 | The Construction of the Graded Formal MR Diagram over Free Groups | 12 |
| 5.3 | General Properties of the Graded Formal MR Diagram | 14 |
| 6 | The <i>ExtraPS</i> MR Diagram over Random Groups | 18 |
| 6.1 | The Construction of <i>ExtraPS</i> MR Diagram over Random Groups | 19 |
| 6.2 | The Construction of <i>ExtraPS</i> MR Diagram over Free Groups | 21 |
| 6.3 | General Properties of the <i>ExtraPS</i> MR Diagram | 22 |
| 7 | The Construction of the CollapseExtra MR Diagram | 25 |
| 7.1 | General Properties of <i>CollapseExtra</i> MR Diagram | 28 |
| 8 | Approximations of <i>EAE</i>-Sets under Minimal Rank Assumption | 30 |
| 8.1 | First Approximation of <i>EAE</i> (p) | 31 |
| 8.2 | Reducing the Set <i>EAE</i> (p) under Minimal-Rank Assumption | 35 |
| 9 | Proof Systems | 39 |
| 9.1 | The Construction of Proof Systems of Depth 2 under Minimal Rank Assumption | 39 |
| 9.2 | The Construction of Proof Systems of General Depth under Minimal Rank Assumption . . . | 45 |
| 10 | Truth Sentences of Minimal Rank over Random Groups | 49 |

1 Introduction

Around 1945, a well-known question was presented by Alfred Tarski on the first order theory of free groups. He asked if every two non-abelian finitely generated free groups are elementary equivalent. His question was answered affirmatively by Z. Sela through his seminal work, which was published in a series consisting of seven papers titled “Diophantine Geometry” ([Se1, Se2, Se3, Se4, Se5, Se6, Se7]). For a different approach to Tarski’s problem, see also [Kh-My].

Actually, Sela obtained a classification of all the f.g. (finitely generated) groups that are elementary equivalent to a non-abelian f.g. free group. In [Se8], Sela generalized his work to a general torsion-free hyperbolic group.

Hence, all the non-abelian f.g. free groups share the same collection of truth sentences. This collection is called the theory of free groups. In this paper, we will be interested in the groups that can or cannot be distinguished from the free groups by a single sentence.

In the previous paper in this sequence, we have proved that given a first order sentence ψ in the Boolean algebra of universal sentences, the sentence ψ is a truth sentence over non-abelian free groups if and only if ψ is a truth sentence over a random group with overwhelming probability in the (Gromov) density model with density $d < \frac{1}{2}$.

In this paper, we consider general sentences of minimal rank. Our aim in this paper is proving that every given minimal rank sentence is a truth sentence over free groups if and only if it is a truth sentence over a random group of density $d < \frac{1}{2}$.

Theorem. (*Main Theorem*). *Let ψ be a first order sentence of minimal rank. Let Γ be a random group of density $d < \frac{1}{2}$. Then ψ is a truth sentence over the free group F_k if and only if ψ is a truth sentence over Γ (in overwhelming probability).*

In [Se5, Se6], Z. Sela obtained a quantifier elimination procedure for general formulas over free groups. He proved that every definable set belongs to the Boolean algebra of *AE*-sets (All-Exist) over free groups. The tools needed for that quantifier elimination procedure are presented in [Se1, Se2, Se3, Se4].

For our aim in this paper, we prove the existence of a uniform quantifier elimination for formulas of minimal rank. Namely, for a given minimal rank formula $V(p)$, we prove the existence of a formula $\varphi(p)$ that belongs to the Boolean algebra of *AE*-formulas, so that the two formulas $V(p)$ and $\varphi(p)$ are equivalent (i.e., define the same set) over the free group F_k (over which we define the density model; k is the rank of the free group F_k), and are equivalent over almost all the groups of density $d < \frac{1}{2}$.

Theorem. *Let $0 \leq d < \frac{1}{2}$, and let $V(w, p)$ be a minimal rank formula in the Boolean algebra of *AE*-formulas. Let $\pi V(p)$ be the projection of $V(w, p)$. There exists a formula $R(p)$ in the Boolean algebra of *AE*-formulas so that:*

1. *The formulas $\pi V(p)$ and $R(p)$ define the same set over F_k .*
2. *Let Γ be a random group of density d . With overwhelming probability, the formulas $\pi V(p)$ and $R(p)$ define the same set over Γ .*

The uniform quantifier elimination procedure that we present in this paper is essentially the quantifier elimination procedure for formulas over free groups (presented in [Se5]). In other words, the quantifier elimination procedure (for minimal rank formulas) presented in [Se5] is valid over almost all the groups of density $d < \frac{1}{2}$, for every minimal rank formula given in advance.

The paper is organized as follows.

In section §2, we prove that every group obtained as a limit of a convergent sequence over an ascending (w.r.t. the levels of the model) sequence of random groups is a limit group (over free groups). Due to the descending chain condition of limit groups, this observation guarantees that the iterative procedures that we will apply over random groups along the next sections must terminate in a finite time, since these procedures run over ascending sequences of random groups, rather than over a single one.

In section §3, we define graded test sequences over ascending sequences of random groups. The main importance of using test sequences is to preserve the structure of the resolutions that we use, during the constructions explained in the next sections.

Section §4 is devoted to rigid and solid graded limit groups. In this section, we recall the definition of graded formal Makanin-Razborov diagrams, as well as the definitions of rigid and solid graded limit groups (introduced in [Se1]). We extend the definition of exceptional families of solutions for rigid and solid limit groups over an arbitrary group. In addition, we prove the existence of a uniform global bound on the number of distinct exceptional families of a given rigid or solid limit group, over a random group (with overwhelming probability).

The approach to quantifier elimination that we adopt, requires the development of some algebraic objects in advance. Sections 5-7 are devoted to constructing uniform objects, which will be applied in the next sections in order to obtain a uniform quantifier elimination procedure over F_k and over almost all the groups of density $d < \frac{1}{2}$.

In particular, in the classical approach to quantifier elimination for positive formulas over free groups, the notion of formal solutions was used. This notion was generalized in [Se2], in order to prove the existence of formal solutions over Diophantine sets (resolutions) of specializations over free groups. Moreover, for obtaining a quantifier elimination for general formulas over free groups (as presented in [Se5, Se6]), it was necessary to encode the whole collection of formal solutions of every given graded system of equations, in a single Makanin-Razborov diagram, called the graded formal MR diagram of that graded system.

In section §5, we prove that the graded formal Makanin-Razborov diagram of a given graded system of equations, constructed over the fixed free group F_k ([Se2]), encodes also the collections of formal solutions over almost all the groups of density $d < \frac{1}{2}$.

In section §9, we explain the construction of proof systems for a given *EAE*-formula ($\exists \forall \exists$) of minimal rank. The construction of proof systems (tree of stratified sets) was introduced in [Se5] for *EAE*-formulas over free groups.

Given an *EAE*-formula $T(p)$ of minimal rank, we construct finitely many proof systems that uniformly validate the truth of the sentence $T(p_0)$, for each value p_0 in the set defined by the formula $T(p)$ over the free group F_k , and for each value p_0 in the set defined by the formula $T(p)$ over a random group Γ , for almost all the groups Γ of density $d < \frac{1}{2}$.

This validation is interpreted in terms of non-positive Diophantine conditions, together with the requirement of the existence of a full families of exceptional solutions for some (finite) set of solid and rigid limit groups, specified by the (finitely many) proof systems.

Of course, once a quantifier elimination for positive formulas over free groups is given, the substantial difficulty in obtaining a quantifier elimination for general formulas over free groups is due to the existence of inequalities.

The collection of the proof systems that we associate with a given *EAE*-formula (of minimal rank) can be thought of as an attempt to get rid of the inequalities, by collecting them in a formula which belongs to the Boolean algebra of universal formulas. Unfortunately, this cannot be achieved. But in section §8, we will prove that the finitely many proof systems associated with an *EAE*-formula (of minimal rank) can be used in order to construct a formula in the Boolean algebra of *AE*-formulas that is equivalent (define the same set) to the given *EAE*-formula, over F_k and over almost all the groups of density $d < \frac{1}{2}$.

Consequently, we conclude the existence of a uniform quantifier elimination procedure for every given formula of minimal rank, over F_k and over almost all the groups of density $d < \frac{1}{2}$. The method explained in section §8, introduced in [Se5, Se6] for *EAE*-formulas over free groups, is called an approximation of an *EAE*-set.

Thus, the construction of proof systems for a given formula should conceptually precede the approximation of that formula. However, in contrast to [Se5, Se6], in this paper we choose to reverse the order of presentation, because we explain the approximation method from an abstract point of view.

In section §10, we conclude this work by proving that every given sentence of minimal rank, is a truth sentence over the free group F_k if and only if it is a truth sentence over a random group of density $d < \frac{1}{2}$ (with overwhelming probability).

Acknowledgment. I am indebted to my advisor Zlil Sela who introduced me to this problem. He shared his knowledge and ideas with me, and without his input, I couldn't have completed this work.

2 Limit Groups over Sequences of Random Groups

Along all the sections of this paper, we fix an integer $k \geq 2$, a free group F_k , and a basis $a = (a_1, \dots, a_k)$ for F_k . We also fix a real number $0 \leq d < \frac{1}{2}$, and consider the Gromov density model of density d . For the definition of Gromov density model, see section 2 in the previous paper in this series.

According to theorem 10.1 in the previous paper in this series, we know that given a (finite) system of equations $\Sigma(x, a)$, maybe with constants (in F_k), the probability that the random group Γ satisfies the property that every solution of $\Sigma(x, a)$ in Γ can be lifted to a solution of $\Sigma(x, a)$ in the free group F_k equals one. We use this property along all the coming chapters, and for brevity we will refer to it by saying that Γ satisfies the $\Sigma(x, a)$ -l.p. (lifting property). A direct consequence of this fact is that we can lift solutions for all the systems, but gradually and not all at once.

Theorem 2.1. *Let H be the collection of all finite systems of equations. Then, we can order the systems in H in an ascending subcollections $H_1 \subset H_2 \subset H_3 \subset \dots$, so that if we denote by p_l the probability that the random group of level l , Γ_l , satisfies the Σ -l.p. for all $\Sigma \in H_l$, then p_l converges to one as l approaches ∞ .*

Proof. Let $q_l = 1 - p_l$. Of course, q_l equals the probability that there exists some system Σ in H_l , so that the random group of level l , Γ_l , does not satisfy the Σ -l.p.. Given a system $\Sigma \in H$, we denote by q_l^Σ , the probability that Γ_l does not satisfy the Σ -l.p.. Fixing a system Σ , we already know that $q_l^\Sigma \rightarrow 0$ as $l \rightarrow \infty$. Since there exists only a countable amount of finite systems, we can write them in a sequence $\Sigma_1, \Sigma_2, \Sigma_3, \dots$. Now let $1 = l_1 \leq l_2 \leq l_3, \dots$ be an ascending sequence of integers, so that for all $n \in \mathbb{N}$, we have that $q_l^{\Sigma_1} + q_l^{\Sigma_2} \dots + q_l^{\Sigma_n} < \frac{1}{n}$ for all $l \geq l_n$.

For all $l \in \mathbb{N}$, let n be the maximal integer for which $l_n \leq l$, and we define H_l to be $H_l = \{\Sigma_1, \dots, \Sigma_n\}$. Of course, the probability that there exists some system Σ in H_l , so that the random group of level l , Γ_l , does not satisfy the Σ -l.p, q_l , satisfies $q_l \leq q_l^{\Sigma_1} + q_l^{\Sigma_2} \dots + q_l^{\Sigma_n} < \frac{1}{n} \leq \frac{1}{l}$.

Hence $q_l \rightarrow 0$ as $l \rightarrow \infty$. As required. \square

A simple but important conclusion of the last theorem is that every limit group over any sequence of ascending sequence $(\Gamma_l)_l$ of random groups, is actually a limit group over the free group F_k . More precisely:

Theorem 2.2. *For each l , there exists a subcollection \mathcal{N}_l of groups in the level l , \mathcal{M}_l , of the density model, so that the following two conditions are satisfied:*

1. \mathcal{S}_l is negligible, i.e., $\frac{|\mathcal{N}_l|}{|\mathcal{M}_l|} \rightarrow 0$ as l approaches ∞ .
2. For every convergent sequence $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$, where d is some integer, l_n is an increasing sequence of integers, and $\Gamma_{l_n} \in \mathcal{M}_{l_n} \setminus \mathcal{N}_{l_n}$ for all n , the obtained limit group $F_d / \overline{\ker} h_{l_n}$ is a limit group over F_k .

Proof. We start by defining \mathcal{S}_l to be the collection of all groups in level l that does not satisfy the Σ -l.p. for some system in the collection H_l (H_l is defined in the last theorem). Then, by the last theorem, the first condition is satisfied, i.e., $\frac{|\mathcal{N}_l|}{|\mathcal{M}_l|} \rightarrow 0$ as l approaches ∞ . For proving the second condition, let $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$ be a convergent sequence, where d is some integer, l_n is an increasing sequence of integers, and $\Gamma_{l_n} \in \mathcal{M}_{l_n} \setminus \mathcal{N}_{l_n}$. We consider the resulted limit group $F_d / \overline{\ker} h_{l_n}$, and we want to prove that it is a limit group over F_k , i.e., that there exists a convergent sequence of homomorphisms from F_d to F_k , so that this sequence converges to $F_d / \overline{\ker} h_{l_n}$.

Given an integer n_0 , and a collection of elements $A \subset F_d$, we say that the sequence $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$ stabilizes A n_0 -eventually, if for all $w \in A$, we have that $h_{l_n}(w) = 1$ for all $n \geq n_0$, or $h_{l_n}(w) \neq 1$ for all $n \geq n_0$. Note that, by definition, a sequence $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$ is convergent, if for every finite subset $A \subset F_d$, there exists $n_0 = n_0(A)$, so that the sequence stabilizes A n_0 -eventually.

Given an integer m , we denote by B_m the ball of radius m in F_k . For all m , let $n(m)$ be the minimal integer for which the sequence $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$ stabilizes the ball B_m $n(m)$ -eventually, and s.t. each system of equations composed from conjunctions of words in B_m is contained in $H_{n(m)}$.

Now we consider the subsequence $\{h_{l_{n(m)}} : F_d \rightarrow \Gamma_{l_{n(m)}}\}_m$ of the sequence $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$. Obviously, the sequence $\{h_{l_{n(m)}} : F_d \rightarrow \Gamma_{l_{n(m)}}\}_m$ is convergent, and the limit group defined by it is the same limit group defined by the original sequence $\{h_{l_n} : F_d \rightarrow \Gamma_{l_n}\}$.

For all m , let W_m be the collection of words in B_m who are mapped to 1 by $h_{l_{n(m)}}$. Since the system of equations defined by all the words in W_m belongs to $H_{n(m)}$, and since $\Gamma_{l_{n(m)}} \in \mathcal{M}_{l_{n(m)}} \setminus \mathcal{N}_{l_{n(m)}}$, we get

that there exists a lift $\tilde{h}_{l_{n(m)}}$ of $h_{l_{n(m)}}$ to F_k , i.e., $\tilde{h}_{l_{n(m)}}$ is a homomorphism from F_d to F_k , such that $h_{l_{n(m)}} = \pi \circ \tilde{h}_{l_{n(m)}}$, where $\pi = \pi_{l_{n(m)}}$ is the given natural quotient map from F_k onto $\Gamma_{l_{n(m)}}$.

We obtain a sequence of homomorphisms from F_d into F_k , $\{\tilde{h}_{l_{n(m)}} : F_d \rightarrow F_k\}_m$. By the way this sequence was constructed, it is a convergent sequence whose limit is the limit group defined by the sequence $\{h_{l_{n(m)}} : F_d \rightarrow \Gamma_{l_{n(m)}}\}_m$, $F_d/\varinjlim h_{l_n}$. We conclude that $F_d/\varinjlim h_{l_n}$ is indeed a limit group defined over F_k . This implies the second condition in the claim. As desired. \square

The meaning of the last theorem, is that we may assume along all the way that any convergent sequence of homomorphisms from a given f.g. group into a sequence of groups from the density model that are ordered with respect to their levels in the model, converges into a group which is a limit group over F_k and which is defined by lifts of the sequence of the homomorphisms in the original sequence. We may assume this property, without changing the model, i.e., we may drop all the groups in the collection $\cup_l \mathcal{N}_l$ from the model, without changing its (asymptotic) functionality.

Theorem 2.3. (*noetherianity of random groups*). *Let Σ_∞ be an infinite system of equations (with finite number of variables and constants). Let Σ_f be a finite system equivalent to Σ_∞ over F_k (Guba's theorem). Then, the systems Σ_∞ and Σ_f are equivalent over a random group (with overwhelming probability).*

Proof. Let Γ be a random group. Let $x \in \Gamma$ be a solution of Σ_f over Γ . Since every solution of Σ_f over Γ can be lifted to a solution of Σ_f over F_k , there exists a lift \tilde{x} of x to F_k , s.t. \tilde{x} is a solution of Σ_f over F_k . Since Σ_∞ and Σ_f are equivalent over F_k , we have that \tilde{x} is a solution of Σ_∞ over F_k . Hence, its projection x into Γ , is a solution of Σ_∞ over Γ . \square

We end this section by recalling the definition of a sequence of homomorphisms over an ascending sequence of groups in the model, and the fact that the random group of level l is hyperbolic with a hyperbolicity constant linear in l .

Definition 2.4. An *ascending sequence of groups in the model of density d* , is a sequence of groups $\{\Gamma_{l_n}\}_n$, so that:

1. The sequence $\{l_n\}_n$ is an unbounded non-decreasing sequence of integers.
2. For all n , the group Γ_{l_n} belongs to the level l_n of the model (of density d).

Theorem 2.5. (*[Ol1, Ol2]*). *At density $d < \frac{1}{2}$, for every $\epsilon > 0$, with overwhelming probability, the random group Γ of level l is torsion-free $\frac{12l}{(1-2d-\epsilon)^2}$ -hyperbolic.*

In particular, Γ is $\alpha_0 l$ -hyperbolic, where

$$\alpha_0 = \frac{48}{(1-2d)^2}.$$

3 Graded Test Sequences over Random Groups

3.1 Limit Actions of Sequences over Random Groups

Let F_m be a free group with a fixed basis $x = (x_1, \dots, x_m)$.

Theorem 3.1. *Let $\{h_{l_n} : F_m \rightarrow \Gamma_{l_n}\}$ be a sequence of homomorphisms over an ascending sequence of groups in the model. Assume that the stretching factors $\mu_n = \min_{f_n \in \Gamma_{l_n}} \max_{1 \leq u \leq m} d_{\Gamma_{l_n}}(1, h_{l_n}(x_u))$, satisfies that $\mu_n \geq l_n^2$ for all n , where X_n is the Cayley graph of Γ_{l_n} w.r.t. the generating set a . Then, the sequence $\tau_{f_n} \circ h_{l_n}$, where τ_{f_n} is the corresponding inner automorphism of Γ_{l_n} (see section 9 in the previous paper), subconverges to an isometric non-trivial action of F_m on a pointed real tree (Y, y_0) .*

Proof. For all n , let δ_n be the hyperbolicity constant of the group Γ_{l_n} . Since the δ_n is linear in l_n , then,

$$\frac{\delta_n}{\mu_n} \xrightarrow{n \rightarrow \infty} 0.$$

The remaining details follows as in [Se1] Proposition 1.1. \square

Theorem 3.2. *Keep the notation of 3.1, and assume that the sequence h_{l_n} converges into an isometric non-trivial action of F_m on a pointed real tree (Y, y_0) . We denote $K_{h_{l_n}, \infty} = \{g \in F_m : \forall y \in Y \quad gy = y\}$. Let $L_{h_{l_n}, \infty}$ be the quotient group $F_m/K_{h_{l_n}, \infty}$. Then,*

1. $L_{h_{l_n}, \infty}$ is a f.g. group.
2. $L_{h_{l_n}, \infty}$ is torsion-free.
3. If Y is a real line, then $L_{h_{l_n}, \infty}$ is free abelian.
4. If $g \in F_m$ stabilizes (pointwise) a non-trivial tripod in Y , then $h_{l_n}(g) = 1$ eventually.
5. Let $g \in F_m$ be an element that does not belong to $K_{h_{l_n}, \infty}$. Then, $h_{l_n}(g) \neq 1$ eventually.
6. Let $[y_1, y_2] \subset [y_3, y_4]$ be a pair of non-degenerate segments in Y , and assume that $\text{stab}([y_3, y_4])$, the (pointwise) stabilizer of $[y_3, y_4]$ is non-trivial. Then, $\text{stab}([y_3, y_4])$ is abelian, and

$$\text{stab}([y_1, y_2]) = \text{stab}([y_3, y_4]).$$

Proof. The same proof of theorem 9.2 in the previous paper. □

Theorem 3.3. *Keep the notation of 3.2. The group $L_{h_{l_n}, \infty}$ is a limit group. In particular, the homomorphisms h_{l_n} factor through $L_{h_{l_n}, \infty}$ eventually.*

Proof. By theorem 2.2 and theorem 3.2, the group $L_{h_{l_n}, \infty}$ is a limit group. Since limit groups are finitely presented, h_{l_n} must factor through $L_{h_{l_n}, \infty}$ for all large enough n . □

Theorem 3.4. *Keep the notation of 3.2. If the limit group $L_{h_{l_n}, \infty}$ is freely indecomposable, then the homomorphisms h_{l_n} can be shortened eventually, by precomposing with a modular automorphism in the JSJ decomposition of $L_{h_{l_n}, \infty}$.*

Proof. According to theorem 3.3, the homomorphisms h_{l_n} factor through $L_{h_{l_n}, \infty}$ eventually. Hence, the limit action of $L_{h_{l_n}, \infty}$ on the limit real tree Y , is the limit of the actions of $L_{h_{l_n}, \infty}$ on the Cayley graphs of the groups Γ_{l_n} . Hence, the theorem follows by the standard shortening argument presented in [Ri-Se]. □

3.2 Graded Test Sequences over Random Groups

We start by defining a test sequence over an ascending (w.r.t the levels) sequence of groups in the model. The notion of a test sequence of a resolution was introduced in [Se2] over free groups. From now on, we will make an extensive use in the objects defined in [Se2]. In particular, we will assume that the reader is familiar with the definitions of resolutions, graded resolutions, completions, closures, graded completions, graded closures, and covering closures.

Definition 3.5. Let $Glim(y, p, a)$ be a graded limit group (w.r.t. the parameters set p), and let $GRes(y, p, a)$ be a well-structured graded resolution of $Glim(y, p, a)$ that terminates in a group $B(y, p, a)$ which is either p -rigid or p -solid limit group. Let Y be the Cayley graph of $Comp(z, y, p, a)$, the graded completion of the resolution $GRes(y, p, a)$, and denote by d_Y the corresponding metric on Y . Let $\{\Gamma_{l_n}\}$ be an ascending sequence of groups in the model, let X_n be the Cayley graph of Γ_{l_n} w.r.t. to the generating set a , and denote by d_n the corresponding metric on X_n . For all n , let (y_n, p_n, a) be a Γ_{l_n} -exceptional specializations of B (see definitions 4.1 and 4.2), and denote by $GRes(y, p_n, a)$ the induced ungraded resolution (ungraded resolution over Γ_{l_n}). We say that the sequence $\{(y_n, p_n, a)\}$, together with a sequence of automorphisms $\{(\nu_n^i, \tau_n^i)\}$ of the QH subgroups that appear in the decompositions associated with the various levels of the completed resolutions $Comp(GRes)(z, y, p_n, a)$, and a sequence of homomorphisms

$$\{\lambda_n : Comp(GRes)(z, y, p_n, a) \rightarrow \Gamma_{l_n}\},$$

is a *graded test sequence of the resolution $Comp(z, y, p, a)$ over the ascending sequence of groups in the model $\{\Gamma_{l_n}\}$* , if the following conditions hold:

For every index n :

1. The ungraded resolution $GRes(y, p_n, a)$ is non-degenerate.
2. Conditions (i) – (xiv) of Definition 1.20 in [Se2] (w.r.t. the metrics d_n of (Γ_{l_n}, a) , where a is the fixed generating set of Γ_{l_n}).
3. If $g \in Comp(GRes)(z, y, p, a)$, $d_Y(g, 1) \leq l_n$, and g is not elliptic in at least one of the decompositions associated with the various levels of the completed resolution $Comp(GRes)(z, y, p_n, a)$, then

$$l_n^2 \cdot \max d_n(p_n, 1) < d_n(\lambda_n(g), 1).$$

Lemma 3.6. *Keep the assumptions of 3.5. Assume further that the induced ungraded resolution $GRes(y, p_n, a)$ (ungraded resolution over Γ_{l_n}), are non-degenerate. Then, there exist a sequence of automorphisms $\{(\nu_n^i, \tau_n^i)\}$ of the QH subgroups that appear in the decompositions associated with the various levels of the completed resolutions $Comp(GRes)(z, y, p_n, a)$, and a sequence of homomorphisms*

$$\lambda_n : Comp(GRes)(z, y, p_n, a) \rightarrow \Gamma_{l_n},$$

so that these sequences form a graded test sequence of $Comp(z, y, p, a)$ over the ascending sequence of groups in the model $\{\Gamma_{l_n}\}$. Furthermore, for every n , given any two integers $s_1 < s_2$, and given a basis element q_i^t - which is not a peg - of some of the abelian groups that appear in a decomposition associated with some level of $Comp(GRes)(z, y, p_n, a)$, we can choose that $\lambda_n(q_i^t) = h_n^{ms_2+s_1}$ for some integer m , and so that the element $h_n \in \Gamma_{l_n}$ has no non-trivial roots in Γ_{l_n} .

Lemma 3.7. *Let $Glim(y, p, a)$ be a graded limit group, and let $GRes(y, p, a)$ be a well-structured graded resolution of $Glim(y, p, a)$ that terminates in a group $B(y, p, a)$ which is either p -rigid or p -solid limit group. Let $\{\Gamma_{l_n}\}$ be an ascending sequence of groups in the model, and assume that for all n , we are given a graded test sequence $\{\lambda_s^n : Comp(GRes)(z, y, p, a) \rightarrow \Gamma_{l_n}\}_s$ (with corresponding automorphisms $\{(\nu_s^{n,i}, \tau_s^{n,i})\}_s$), of the completed resolution $Comp(GRes)(z, y, p, a)$ over the specific torsion-free hyperbolic group Γ_{l_n} . Then, if for all n , s_n is large enough, then the sequence $\{\lambda_{s_n}^n : Comp(GRes)(z, y, p, a) \rightarrow \Gamma_{l_n}\}_n$ (with the corresponding automorphisms $\{(\nu_{s_n}^{n,i}, \tau_{s_n}^{n,i})\}_n$), is a graded test sequence of $Comp(GRes)(z, y, p, a)$ over the ascending sequence of groups in the model $\{\Gamma_{l_n}\}$.*

The following lemma guarantees that the non-degenerations of a graded resolution over any group in the model, are encoded by the singular locus, defined over F_k , of that resolution.

Lemma 3.8. *Let $Glim(y, p, a)$ be a graded limit group, and let $GRes(y, p, a)$ be a graded resolution of $Glim(y, p, a)$ that terminates in a group $B(y, p, a)$ which is either p -rigid or p -solid limit group. Let $S_1(y, p, a), \dots, S_r(y, p, a)$ be the finitely many limit groups that form the singular locus of the resolution $GRes(y, p, a)$ (constructed over F_k). Let Γ be a group in the model. Then, the Γ -specializations (y_0, p_0, a) of $B(y, p, a)$ for which the induced resolution $GRes(y, p_0, a)$ is degenerated (over Γ), factors through the singular locus of the resolution $GRes(y, p, a)$.*

Proof. By definition, for an ungraded resolution $GRes(y, p_0, a)$ induced from a Γ -specialization (y_0, p_0, a) of B , to be degenerate, the specialization (y_0, p_0, a) has to satisfy one of finitely many systems $\Sigma_1(y, p, a), \dots, \Sigma_s(y, p, a)$ that was constructed over F_k . Given such a Γ -specialization (y_0, p_0, a) , We lift it to a specialization $(\tilde{y}_0, \tilde{p}_0, a)$ of $B(y, p, a)$ that satisfies one of the systems $\Sigma_1(y, p, a), \dots, \Sigma_s(y, p, a)$. \square

4 Rigid and Solid Limit Groups

Let $\Sigma(y, p, a)$ be a p -graded system of equations, that is p is the parameter set, y is the variable set, and the constants appearing in Σ belong to the free group $F_k = F(a)$. According to [Se1], section 10, we construct the graded MR diagram K of the system $\Sigma(y, p, a)$. This is a finite diagram that encodes all the solutions of the graded system $\Sigma(y, p, a)$, so that the JSJ-decompositions associated with the various groups along it, are p -graded (p -JSJ's). That is, the subgroups $\langle a, p \rangle$ of each of the groups along the diagram, is contained in a distinguished vertex of the p -JSJ associated with the corresponding group, and the modular automorphisms of that group, that are induced from the associated p -JSJ, must fix (elementwise) the distinguished vertex (by definition).

Each resolution in the graded MR diagram, terminates in a limit group which is either p -rigid or p -solid. The p -rigid limit groups, denoted usually by $Rgd(y, p, a)$, are by definition, those graded limit groups that admit a trivial p -JSJ. And p -solid limit groups, denoted usually by $Sld(y, p, a)$, are defined as the graded limit groups who admit a non-trivial p -JSJ, but admit an isomorphic shortening quotient.

Every p -rigid or p -solid limit group in the diagram K , admits a finite set (maybe empty) of maximal flexible quotients $Flx(y, p, a)$, which are proper quotients.

Definition 4.1. Let $Rgd(y, p, a)$ be a p -rigid limit group. Let $\Gamma = \langle a : \mathcal{R} \rangle$ be a group (presentation). A homomorphism $h : Rgd(y, p, a) \rightarrow \Gamma$ is called a Γ -rigid homomorphism of $Rgd(y, p, a)$, if it does not factor through a flexible quotient of $Rgd(y, p, a)$. A Γ -rigid homomorphism is called also a Γ -exceptional solution of $Rgd(y, p, a)$.

Definition 4.2. Let $Sld(y, p, a)$ be a p -solid limit group.

If $Sld(y, p, a)$ admits no flexible quotients, then every homomorphism $Sld(y, p, a) \rightarrow \Gamma$ is defined to be a Γ -strictly solid homomorphism of $Sld(y, p, a)$.

In general, let $\eta_j : Sld(y, p, a) \rightarrow Flx_j(y, p, a)$ be the canonical quotient maps between the p -solid group onto each of its (finitely many) maximal flexible quotients. According to [[Se3], Definition 1.5 p. 15, 16], we associate to each one-step resolution $\eta_j : Sld(y, p, a) \rightarrow Flx_j(y, p, a)$ a graded completion $Comp_j(z, y, p, a)$, so that the p -solid group $Sld(y, p, a)$ is mapped canonically into each of these completions $\tau_j : Sld(y, p, a) \rightarrow Comp_j$. Moreover, we consider the graded completion $Comp_T$ of the one-step resolution $Sld(y, p, a) \rightarrow Sld(y, p, a)$ (the identity map). The solid limit group $Sld(y, p, a)$ admits two natural embeddings $\iota_i : Sld(y, p, a) \rightarrow Comp_T$, $i = 1, 2$.

Let $\Gamma = \langle a : \mathcal{R} \rangle$ be a group (presentation). A homomorphism $h : Sld(y, p, a) \rightarrow \Gamma$ is called a Γ -strictly solid homomorphism of $Sld(y, p, a)$, if h is non-degenerate (see lemma 3.8) and does not factor through the map τ_j for all j . Moreover, we define an equivalence relation on the set of homomorphisms $h : Sld(y, p, a) \rightarrow \Gamma$ as follows. Two such homomorphisms h_1, h_2 are called equivalent, if there exists a homomorphism $u : Comp_T \rightarrow \Gamma$ so that the $u \circ \iota_i = h_i$, for $i = 1, 2$. We call the equivalence class of a Γ -strictly solid homomorphism a Γ -strictly solid family. Note that all the non-degenerate homomorphisms in a Γ -strictly solid family, are Γ -strictly solid.

A Γ -strictly solid homomorphism is called also a Γ -exceptional solution of $Sld(y, p, a)$, and a Γ -strictly solid family is called also a Γ -exceptional family of $Sld(y, p, a)$.

When $\Gamma = F_k$, we remove Γ from the notation, and write simply strictly solid or exceptional homomorphisms, and strictly solid or exceptional families.

Theorem 4.3. ([Se3], theorem 2.3). *Let $Rgd(y, p, a)$ be a p -rigid limit group. There exists a global bound $b_R(Rgd)$ for which, for any particular value of the defining parameters p_0 , there are at most b_R distinct rigid homomorphisms $h : Rgd(y, p, a) \rightarrow F_k$ satisfying $h(p) = p_0$.*

Theorem 4.4. *Let $Rgd(y, p, a)$ be a p -rigid limit group. There exists a global bound $b_R(Rgd)$ for which the following property is satisfied. Let Γ_l be a group of level l in the model.*

If l is large enough, then, for any particular value of the defining parameters $p_0 \in \Gamma_l$, there are at most b_R distinct rigid homomorphisms $h : Rgd(y, p, a) \rightarrow \Gamma_l$ satisfying $h(p) = p_0$.

Proof. Let b_R be the one in theorem 4.3, and assume by contradiction that $\Gamma = \Gamma_l$ admits more b_R pairwise distinct Γ -rigid homomorphisms $h_j : Rgd(y, p, a) \rightarrow \Gamma$. For each j , we lift the homomorphism h_j into a homomorphism $\tilde{h}_j : Rgd(y, p, a) \rightarrow F_k$.

Then, since the homomorphisms h_j are pairwise distinct, so are the homomorphisms \tilde{h}_j . Moreover, for all j , the homomorphism \tilde{h}_j is rigid, since h_j is. A contradiction. \square

Theorem 4.5. ([Se3], theorem 2.5). *Let $Sld(y, p, a)$ be a p -solid limit group. There exists a global bound $b_S(Sld)$ for which, for any particular value of the defining parameters p_0 , there are at most b_S distinct strictly solid families of homomorphisms $h : Sld(y, p, a) \rightarrow F_k$ satisfying $h(p) = p_0$.*

Theorem 4.6. *Let $Sld(y, p, a)$ be a p -solid limit group. There exists a global bound $b_S(Sld)$ for which the following property is satisfied. Let Γ_l be a group of level l in the model.*

If l is large enough, then, for any particular value of the defining parameters $p_0 \in \Gamma_l$, there are at most b_S distinct strictly solid families of homomorphisms $h : Sld(y, p, a) \rightarrow \Gamma_l$ satisfying $h(p) = p_0$.

Proof. Let b_S be the one in theorem 4.3, and assume by contradiction that $\Gamma = \Gamma_l$ admits more b_R pairwise distinct Γ -strictly solid families. Let h_j be strictly solid representatives for these families. For each j , we lift the homomorphism h_j into a homomorphism $\tilde{h}_j : Sld(y, p, a) \rightarrow F_k$.

Then, since the homomorphisms h_j represent pairwise distinct Γ -strictly solid families, the homomorphisms \tilde{h}_j must represent pairwise distinct strictly solid families. A contradiction. \square

Lemma 4.7. *Let $B(y, p, a)$ be a p -rigid or a p -solid limit group. Let $Glim(x, y, p, a)$ be a p -graded limit group, and let $GRes(x, y, p, a)$ be a p -graded resolution of $Glim(x, y, p, a)$. Assume that $B(y, p, a)$ is mapped canonically into $Glim(x, y, p, a)$, and denote $\eta_j : B(y, p, a) \rightarrow Glim_j(x, y, p, a)$ the corresponding epimorphisms of $B(y, p, a)$ into the various levels of $GRes(x, y, p, a)$. If $B(y, p, a)$ admits an exceptional specialization (y_0, p_0, a) that factors through the resolution $GRes(x, y, p, a)$, then for all j , the p -JSJ of $B(y, p, a)$ is mapped compatibly into the p -JSJ of $Glim_j(x, y, p, a)$.*

Proof. Assume by contradiction it is not the case. Let $Flx(c, y, p, a) = \{FlxR_1, \dots, FlxR_m\}$ be the collection of the relations of the graded completions of the graded resolutions $B(y, p, a) \rightarrow Flx_1(y, p, a), \dots, B(y, p, a) \rightarrow Flx_m(y, p, a)$, where $Flx_1(y, p, a), \dots, Flx_n(y, p, a)$ are the maximal flexible quotients of $Sld(y, p, a)$. Let $Comp(z, x, y, p, a)$ be the completion of the resolution $GRes(x, y, p, a)$, and let $Comp(z, x, y, p_0, a)$ be an ungraded resolution corresponding to the factorization of the exceptional specialization (y_0, p_0, a) of $B(y, p, a)$ through $Comp(z, x, y, p, a)$.

We Construct the formal MR diagram D of the systems $Flx(c, y, p_0, a)$ w.r.t. the resolution $Comp(z, x, y, p_0, a)$ (see [Se2], section 2, or section §5 below). Since the p -JSJ of $B(y, p, a)$ is not mapped compatibly into the p -JSJ of $Glim_j(x, y, p, a)$, for some j , we conclude that all the test sequences of $Comp(z, x, y, p_0, a)$, must factor through one the terminal formal $FCl(c, x, y, p_0, a)$ closures in the formal MR diagram D . Hence, These formal closures form a covering closure for the resolution $Comp(z, x, y, p_0, a)$. In particular, the specialization (y_0, p_0, a) of $B(y, p, a)$, factors through one of these formal closures. Thus, (y_0, p_0, a) is not an exceptional specialization of $B(y, p, a)$, a contradiction. \square

As stated in the top of this section, all the solutions of the graded system $\Sigma(y, p, a)$, factors through the graded MR diagram K of the graded system $\Sigma(y, p, a)$. By definition, a homomorphism factors through a graded resolution in K , if it can be expressed as a composition of graded modular automorphisms of the different levels of the resolution, the canonical epimorphisms between consecutive levels, and a rigid or a strictly solid homomorphism of the terminal p -rigid or p -solid limit group of the resolution.

Actually, in order to cover all the solutions of the graded system $\Sigma(y, p, a)$ by the graded resolutions of the diagram K , it is not sufficient to consider only strictly solid solutions of the terminal p -solid limit groups. But we need further to consider strictly solid solutions with respect to finitely many sets of graded closures of the graded resolutions of the graded MR diagram associated with each p -solid group that appear in the diagram K . The details of this more general definition are stated in definition 2.12 in [Se3]. This definition can be converted naturally in order to define a Γ -strictly solid solution of a p -solid limit group with respect to a given set of closures for a general group $\Gamma = \langle a : \mathcal{R} \rangle$.

According to theorem 2.13 in [Se3], the number of distinct families of strictly solid homomorphisms of a p -solid group with respect to a given set of closures, is globally bounded. Using this theorem, a similar argument given in the proof of theorem 4.6, proves the existence of a uniform bound on the number of distinct families of Γ -strictly solid homomorphisms of a p -solid group with respect to a given set of closures, for almost all the groups Γ in the model.

Hence, for simplicity along the next sections, we will use the terminology of strictly solid homomorphisms, without stating the corresponding set of closures.

5 Formal Solutions over Random Groups

Along this section, we fix a p -graded resolution $Res(y, p, a)$, and we consider its completion $Comp = Comp(Res)(z, y, p, a)$. We also fix a system of equations $\Sigma(x, y, p, a)$, where x, y are the variables, and p denotes the parameter set (a is the fixed basis of the fixed free group F_k). We denote by B the p -graded base group of $Comp(z, y, p, a)$.

Our aim is to collect all the formal solutions of $\Sigma(x, y, p, a)$, w.r.t. to the resolution $Res(y, p, a)$, over almost all the groups in the model, in a uniform diagram.

For that, we construct an auxiliary graded formal MR diagram that encodes all of these formal solutions. Then, we will use it in order to prove that the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$ over the free group F_k , that encodes all the formal solutions of the system $\Sigma(x, y, p, a)$ over F_k , encodes also all the formal solutions of $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$ over almost all the groups in the model.

5.1 The Construction of Graded Formal MR Diagram over Random Groups

Let \mathfrak{T} be the collection of all the sequences $\{(x_n, z_n, y_n, p_n, a)\}$ for which the following conditions hold:

1. The sequence $\{(x_n, z_n, y_n, p_n, a)\}$ is defined over some ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model.
2. The Γ_{l_n} -specialization (x_n, z_n, y_n, p_n, a) satisfies the system Σ , i.e., $\Sigma(x_n, z_n, y_n, p_n, a) = 1$ (in Γ_{l_n}), for all n .
3. The restricted sequence $\{(z_n, y_n, p_n, a)\}$ is a graded test sequence over the (same) ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model.

The sequences in the collection \mathfrak{T} , factor through finitely many maximal limit groups H (see theorem 2.2). The canonical image of the subgroup $Comp(z, y, p, a)$ in H is denoted briefly by $Comp$. Such an H could be freely decomposable w.r.t. the subgroup $Comp \leq H$. We consider the most refined $Comp$ -free factorization of H . We call the free factors that do not include (a conjugate of) the subgroup $Comp \leq H$, a *standard free factors*.

We keep the standard factors, extend them by their standard MR diagrams, and continue the construction with the free factor that contains $Comp$. We denote this factor by $M = M(x, z, y, p, a)$, which is a limit group. We consider the $Comp$ -JSJ decomposition of the limit group $M(x, z, y, p, a)$, i.e., the JSJ decomposition with respect to the subgroup $Comp$. If this decomposition is trivial, we call the group $M(x, z, y, p, a)$ a *Comp-rigid* group, and terminate the construction for this stage. In this case, the subcollection of \mathfrak{T} consisting of all the sequences $\{(x_n, z_n, y_n, p_n, a)\}$ that factor through M , is denoted by \mathfrak{T}_M .

In the case that the $Comp$ -JSJ of $M(x, z, y, p, a)$ is non-trivial, we denote by \mathfrak{T}_M the subcollection of \mathfrak{T} , consisting of all the sequences $\{(x_n, z_n, y_n, p_n, a)\}$ that satisfy the following conditions:

1. The sequence $\{(x_n, z_n, y_n, p_n, a)\}$, that is defined over an ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model, factors through the group M .
2. For all n , the specialization (x_n, z_n, y_n, p_n, a) cannot be shortened (in the metric of (Γ_{l_n}, a) , where a is the fixed generating set of Γ_{l_n}) by precomposing with a modular automorphism of M that is induced from its $Comp$ -JSJ.

The sequences in the collection \mathfrak{T}_M , factor through finitely many maximal limit quotients of M , that we denote - while abusing the notation - by H . If H is not a strict quotient of M , we call M a *Comp-solid* group, and terminate the construction for this stage.

We continue the construction iteratively. Since all the groups that are obtained along the construction, are limit groups, then, according to the descending chain condition of limit groups, the construction must terminate after finitely many steps. The result is a diagram D that includes finitely many resolutions. Each of these resolutions terminates in a free product $\langle f \rangle * M$, where $\langle f \rangle$ is a free group, and M is either a *Comp-rigid* group or a *Comp-solid* group.

Lemma 5.1. *Assume that $M(x, z, y, p, a)$ is a $Comp$ -rigid group or a $Comp$ -solid group. Then, M admits a structure of a p -graded tower over a p -graded base limit group $Term(M)$, so that:*

1. *up to adding roots to abelian groups, this structure is similar to the one associated with $Comp(z, y, p, a)$,*
2. *this structure is compatible to the one associated with $Comp(z, y, p, a)$, that is, the canonical map $Comp \rightarrow M$ maps the decomposition associated with each level of $Comp$, except maybe the bottom level of $Comp$, compatibly into the decomposition associated with the corresponding level of M ,*
3. *the canonical image of the p -graded base group B of $Comp(z, y, p, a)$ lies entirely in $Term(M)$.*

Proof. Let (x_n, z_n, y_n, p_n, a) be a sequence in \mathfrak{T}_M , that converges into a non-trivial action of $M(x, z, y, p, a)$ on a real tree. Consider the abelian decomposition Π that this action induces for M .

If $Comp$ is contained in a proper subgraph of Π , then the $Comp$ -JSJ of $M(x, z, y, p, a)$ is non-trivial, and, according to the shortening argument over ascending sequence of groups in the model (theorem 3.4), the (generators of some part of the $Comp$ -JSJ of M induced by the) specializations in the sequence (x_n, z_n, y_n, p_n, a) can eventually be shortened by precomposing with modular automorphisms in the $Comp$ -JSJ of $M(x, z, y, p, a)$, a contradiction. Hence, the components of the action of M on the real tree are of the same forms as those of $Comp$. Using the properties of test sequences over ascending sequence of groups in the model, we can apply the analysis in the proofs theorem 3.7, and theorem 3.8 of [Se2], in order to analyze the action of M on the limit real tree.

We deduce that the action of M contains either a single IET, which is the “highest” surface in the decomposition associated with the completion $Comp(z, y, p, a)$, and finitely many point stabilizers, or the action of M is discrete with a unique point stabilizer and one edge, which (iteratively as in the proof of the existence of a formal solution over abelian groups in [Se2], Proposition 1.8) gives a decomposition for M of the form $V *_{Ab_1} Ab$, where Ab is obtained from the “highest” abelian group of the decomposition associated with the completion $Comp(z, y, p, a)$ by adding some roots for non-pegs generators.

Continuing with each of the vertices (point stabilizers) iteratively, this uncovering procedure explains that M admits a tower structure whose upper levels are similar to those of the decomposition associated with the the highest level of the completion $Comp(z, y, p, a)$, up to adding roots for abelian groups along this decomposition.

Finally, this procedure ends up when we get a point stabilizer $Term(M)$ whose intersection with $Comp$ is of the form $\bar{B} * F$, where \bar{B} is the image of B and F is a free group. \square

In addition to lemma 5.1, we want that the p -graded base group B of $Comp(z, y, p, a)$ to be mapped compatibly into the p -graded base group $Term(M)$ of M .

For that, we note that at this point, we can consider the finitely many base groups $Term(M)$ that we have constructed so far, as were given in advance. According to lemma 5.1, each of these groups is a limit group, and in particular a finitely presented group. Hence, according to theorem 2.1, we can assume that if Γ is any group in the model, then every Γ -specialization of $Term(M)$ can be lifted to a specialization of $Term(M)$ in F_k . As a consequence, we obtain the compatibility that we wanted:

Lemma 5.2. *Assume that $M(x, z, y, p, a)$ is a $Comp$ -rigid group or a $Comp$ -solid group. Then, M admits a structure of a graded formal closure of $Comp(z, y, p, a)$ over a p -graded limit group $Term(M)$. That is, M admits a structure of a p -graded tower over a p -graded base group $Term(M)$, so that:*

1. *up to adding roots to abelian groups, this structure is similar to the one associated with $Comp(z, y, p, a)$,*
2. *this structure is compatible to the one associated with $Comp(z, y, p, a)$,*
3. *the canonical image of the p -graded base group B of $Comp(z, y, p, a)$ lies entirely in $Term(M)$, and B is mapped compatibly into $Term(M)$. That is, the minimal subgraph of groups of the p -JSJ of $Term(M)$ containing the canonical image \bar{B} of B , consists only from parts that are taken from the p -JSJ of B .*

Proof. Let (x_n, z_n, y_n, p_n, a) be a sequence in \mathfrak{T}_M . Since the sequence (x_n, z_n, y_n, p_n, a) factors through M , and since the graded limit quotient \bar{B} of B lies entirely in $Term(M)$, we conclude that there exists a Γ -exceptional specialization of B that factors through $Term(M)$, for some group $\Gamma = \Gamma_n$ in the model. We lift this specialization to a specialization of B that factors through $Term(M)$. Note that this lift must be an exceptional specialization of B (over F_k). Hence, according to lemma 4.7, the p -JSJ of $Term(M)$ must be compatible with that of B . We conclude that M is a graded closure of $Comp(z, y, p, a)$ over some p -graded limit group $Term(M)$. \square

We denote by D the diagram obtained by our iterative construction. In light of lemma 5.2, every resolution of the diagram D that we have constructed above, is defined over a graded formal closure of the graded completion $Comp(z, y, p, a)$. And according to the construction of D , every sequence $\{(x_n, z_n, y_n, p_n, a)\}$ of specializations over an ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model, that satisfies the system $\Sigma(x, y, p, a)$, and restricts to a test sequence $\{(z_n, y_n, p_n, a)\}$ over the ascending sequence $\{\Gamma_{l_n}\}$, subfactors through at least one of the resolutions in D . Note that factorization through such a resolution is defined using the canonical

maps, factorization through the standard MR diagrams associated with the standard free factors, together with modular automorphisms induced by the *Comp*-JSJ's of the corresponding (non-standard) groups along the diagram.

The obtained diagram D is called *the auxiliary graded formal MR diagram over random groups of the system $\Sigma(x, y, p, a)$ over the graded resolution $Res(y, p, a)$* .

Our aim is to show that the graded formal MR diagram over F_k of the system $\Sigma(x, y, p, a)$ over the graded resolution $Res(y, p, a)$ (that we explain in the next subsection), encodes all the formal solutions of a random group Γ (in probability 1) for any value of the parameters set $p_0 \in \Gamma$, for which the sentence

$$\forall y \in Res(y, p_0, a) \quad \Sigma(x, y, p_0, a) = 1$$

is a truth sentence over Γ . The precise statement formulating this property, will be given below (theorem 5.10). For that aim, we start with the following lemma:

Lemma 5.3. *Let Γ_l be a group of level l in the model. If l is large enough, then the following property is satisfied in Γ_l .*

Every sequence of specializations $\{(x_n, z_n, y_n, p_n, a)\}$ over Γ_l , for which $\Sigma(x_n, y_n, p_n, a) = 1$ in Γ_l for all n , and the restricted sequence $\{(z_n, y_n, p_n, a)\}$ is a test sequence of $Comp(z, y, p, a)$ over Γ_l , subfactors through the auxiliary graded formal MR diagram over random groups D .

Proof. Assume by contradiction the statement is false. Then, there will be an ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model, so that for all n , there exists a sequence $\{(x_m^n, z_m^n, y_m^n, p_m^n, a)\}_m$ of Γ_{l_n} -specializations, for which:

1. For all m , the Γ_{l_n} -specialization $(x_m^n, z_m^n, y_m^n, p_m^n, a)$ satisfies the system Σ , i.e., $\Sigma(x_m^n, z_m^n, y_m^n, p_m^n, a) = 1$ (in Γ_{l_n}).
2. The restricted sequence $\{(z_m^n, y_m^n, p_m^n, a)\}_m$ is a test sequence of $Comp(z, y, p, a)$ over Γ_{l_n} .
3. For all m , the Γ_{l_n} -specialization $(x_m^n, z_m^n, y_m^n, p_m^n, a)$ does not factor through D .

Hence, using lemma 3.7, there exists a sequence $(m_n)_n$ of integers, so that the sequence

$$\{(x_{m_n}^n, z_{m_n}^n, y_{m_n}^n, p_{m_n}^n, a)\}_n$$

which is defined over the ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model, satisfies the system $\Sigma(x, y, p, a) = 1$, restricts to a test sequence $\{(z_{m_n}^n, y_{m_n}^n, p_{m_n}^n, a)\}_n$ over the ascending sequence $\{\Gamma_{l_n}\}$, but it does not subfactor through D . This contradicts the construction of D . \square

5.2 The Construction of the Graded Formal MR Diagram over Free Groups

In this subsection we will construct the graded formal MR diagram over the free group F_k of the system $\Sigma(x, y, p, a)$ w.r.t. the graded resolution $Res(y, p, a)$. The construction is essentially similar to the construction that we have explained above over random groups.

Let \mathfrak{J} be the collection of all the sequences $\{(x_n, z_n, y_n, p_n, a)\}$ of specializations (over F_k), for which $\Sigma(x_n, z_n, y_n, p_n, a) = 1$ (in F_k) for all n , and for which the restricted sequence $\{(z_n, y_n, p_n, a)\}$ is a test sequence of $Comp(z, y, p, a)$ (over F_k).

The sequences in the collection \mathfrak{J} , factor through finitely many maximal limit groups H . We consider the most refined *Comp*-free factorization of H .

We keep the free factors that do not include (a conjugate of) the subgroup $Comp \leq H$, extend them by their standard MR diagrams, and continue the construction with the free factor that contains $Comp$. We denote this factor by $M = M(x, z, y, p, a)$. We consider the *Comp*-JSJ decomposition of the limit group $M(x, z, y, p, a)$. If $M(x, z, y, p, a)$ is *Comp*-rigid, i.e., if its *Comp*-JSJ is trivial, we terminate the construction for this stage. In this case, the subcollection of \mathfrak{J} consisting of all the sequences $\{(x_n, z_n, y_n, p_n, a)\}$ that factor through M , is denoted by \mathfrak{J}_M .

In the case that the *Comp*-JSJ of $M(x, z, y, p, a)$ is non-trivial, we consider the subcollection \mathfrak{J}_M of \mathfrak{J} , consisting of all sequences $\{(x_n, z_n, y_n, p_n, a)\}$ that factor through M , and so that for all n , the specialization (x_n, z_n, y_n, p_n, a) cannot be shortened (in the metric of (F_k, a) , where a is the fixed generating set of F_k) by

precomposing with a modular automorphism of M that is induced from its $Comp$ -JSJ. The sequences in the collection \mathfrak{J}_M , factor through finitely many maximal limit quotients of M , that we denote again by H . If the obtained H is not a strict quotient of M , i.e., if M is a $Comp$ -solid group, we terminate the construction for this stage.

We continue the construction iteratively. According to the descending chain condition of limit groups, the construction must terminate after finitely many steps. The result is a diagram that we denote by K , that includes finitely many resolutions. Each of these resolutions terminates in a free product $\langle f \rangle * M$, where $\langle f \rangle$ is a free group, and M is either a $Comp$ -rigid group or a $Comp$ -solid group.

Lemma 5.4. *Assume that $M(x, z, y, p, a)$ is a $Comp$ -rigid group or a $Comp$ -solid group. Then, M admits a structure of a graded formal closure of $Comp(z, y, p, a)$ over a p -graded limit group $Term(M)$.*

Proof. The proof is essentially the same as in lemma 5.1 and lemma 5.2. For more details see [Se2] theorems 3.7, and 3.8. \square

Remark 5.5. We note that the above construction of a graded formal MR diagram over F_k of a given graded system $\Sigma(x, y, p, a)$ w.r.t. a given graded resolution $Res(y, p, a)$, can be applied also in the case that the terminal group of $Res(y, p, a)$ is not rigid or solid. This means that we can consider the completion $Comp$ of $Res(y, p, a)$, and then to consider the collection of all sequences $\{(x_n, z_n, y_n, p_n, a)\}$ of specializations, for which $\Sigma(x_n, z_n, y_n, p_n, a) = 1$ for all n , and for which the sequence $\{(z_n, y_n, p_n, a)\}$ is a test sequence for $Comp$ (test sequence over $Comp$ is defined now over any sequence of specializations of $Term(Comp)$).

This way, and using the shortening procedure iterative construction, we can construct a finite diagram K' , so that each of its resolutions terminates in a graded tower (over some p -graded limit group $Glim_t(b, p, a)$ into which the base group of $Comp$ is mapped - maybe non-compatibly) of the initial graded resolution $Res(y, p, a)$, and so that the following property is satisfied. If $\{(x_n, z_n, y_n, p_n, a)\}$ is a sequence of specializations for which $\Sigma(x_n, z_n, y_n, p_n, a) = 1$ for all n , and for which the sequence $\{(z_n, y_n, p_n, a)\}$ is a test sequence for $Comp$, then the sequence $\{(x_n, z_n, y_n, p_n, a)\}$ subfactors through K' . And in particular, each of the terminal groups of the resolutions in K' that contains the subgroup $Comp$ - which are graded towers of the resolution $Res(y, p, a)$ as aforementioned - contains a formal solution $x = x(s, z, y, p, a)$ of the system $\Sigma(x, y, p, a)$, where the s 's stand for the roots that were added for abelian groups along the decomposition of $Comp$.

In order to complete the construction of the graded formal MR diagram over F_k , in contrast of the construction of the auxiliary graded formal MR diagram over random groups, we further extend the resolutions of K . For every resolution in K over a graded formal closure M , we perform the following extension. We consider the subcollection $\mathfrak{B}\mathfrak{J}_M$ of \mathfrak{J}_M consisting of all the sequences $\{(x_n, z_n, y_n, p_n, a)\}$, that satisfies the following:

1. If $Term(M)$ is neither p -rigid nor p -solid, then, for all n , the specialization (x_n, z_n, y_n, p_n, a) restricts to a specialization of $Term(M)$ that cannot be shortened by precomposing with a modular automorphism induced from the p -JSJ of $Term(M)$.
2. If $Term(M)$ is p -rigid or p -solid, then, for all n , the restriction of the specialization (x_n, z_n, y_n, p_n, a) factors through some flexible quotient of $Term(M)$.

The sequences in $\mathfrak{B}\mathfrak{J}_M$ factor through finitely many maximal limit groups that we denote by H . According to [Se1], Lemma10.4, each of the limit quotients H are proper quotients of M . We continue the construction iteratively. According to the descending chain condition of limit groups, this construction must terminate after finitely many steps. The result is a diagram K_M with finitely many resolutions, each of which terminates in a graded formal closure $M'(x, z, y, p, a)$, whose terminal group is either a p -rigid or a p -solid limit group.

Note that factorization through a resolution in the diagram K_M is defined using the p -JSJ's of the base groups of the graded formal closures along that resolution (in addition to the canonical maps, standard MR diagrams, and the $Comp$ -JSJ's of the corresponding free factors along the resolution).

Lemma 5.6. *Let t_0 be a specialization of $Term(M)$, the p -graded base group of M , so that t_0 restricts to an exceptional specialization of B , the p -graded base group of $Comp(z, y, p, a)$. Denote by M_{t_0} the induced ungraded formal closure, and let (x_0, z_0, y_0, p_0, a) be a specialization of M_{t_0} .*

Then, the specialization (x_0, z_0, y_0, p_0, a) factors through one of the resolutions $GRes(x, z, y, p, a)$ of K_M . Moreover, if $Term(M)$ is either a p -rigid or a p -solid group, and t_0 is not an exceptional specialization of

$Term(M)$, then we can assume that the terminal graded formal closure of $GFRes(x, z, y, p, a)$ is a proper quotient of M .

Proof. Since the p -JSJ of $Term(M)$ is compatible with that of B , we may assume without losing the generality that the given solution t_0 is short in the case that $Term(M)$ is neither p -rigid, nor p -solid, and that t_0 factors through a flexible quotient of $Term(M)$, in the case that it is p -rigid or p -solid.

We consider the ungraded formal closure $M_{t_0} = M(x, z, y, p_0, a)$, and the ungraded completion $Comp(z, y, p_0, a)$, that are obtained from M and $Comp(z, y, p, a)$ by specializing their base groups $Term(M)$ and B according to the given solution t_0 , respectively.

Let $\{(x_n, z_n, y_n, p_0, a)\}$ be a test sequence of M_{t_0} . That is, the sequence $\{(z_n, y_n, p_0, a)\}$ is a test sequence of $Comp(z, y, p_0, a)$, and the specialization x_n is the induced specialization of the element x of M_{t_0} , for all n .

Then, the sequence $\{(x_n, z_n, y_n, p_0, a)\}$ belongs to $\mathfrak{B}\mathfrak{J}_M$. Hence, using a simple induction, the sequence $\{(x_n, z_n, y_n, p_0, a)\}$ subfactors through the diagram K_M by the construction of K_M .

This implies that there exists a collection of (boundedly many) ungraded closures $FCl(x, z, y, p_0, a)$, each of which is obtained by specializing the terminal group of some graded closure in K_M (other than M itself), so that this collection forms a covering closure of M_{t_0} , and the Lemma follows. \square

Lemma 5.7. *Let Γ_l be a group of level l in the model. If l is large enough, then the following property is satisfied in Γ_l .*

Let t_0 be a Γ_l -specialization of $Term(M)$, so that t_0 restricts to a Γ_l -exceptional specialization of B . Denote by M_{t_0} the induced ungraded formal closure, and let (x_0, z_0, y_0, p_0, a) be a Γ_l -specialization of M_{t_0} .

Then, the Γ_l -specialization (x_0, z_0, y_0, p_0, a) factors through one of the resolutions $GFRes(x, z, y, p, a)$ of K_M . Moreover, if $Term(M)$ is either a p -rigid or a p -solid group, and t_0 is not a Γ_l -exceptional specialization of $Term(M)$, then we can assume that the terminal graded formal closure of $GFRes(x, z, y, p, a)$ is a proper quotient of M .

Proof. We lift (x_0, z_0, y_0, p_0, a) to a specialization $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ of M in F_k , so that in the case that $Term(M)$ is either p -rigid or p -solid, and the restriction t_0 of the Γ_l -specialization (x_0, z_0, y_0, p_0, a) to $Term(M)$ is not Γ_l -exceptional, then the restriction \tilde{t}_0 of the lift $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ to $Term(M)$ is not F_k -exceptional. Note that the restriction of $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ to B must be an F_k -exceptional solution of B .

Hence, according to lemma 5.6, the lift $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ factors through some resolution $GFRes(x, z, y, p, a)$ of K_M . Thus, the Γ_l -specialization (x_0, z_0, y_0, p_0, a) factors through the resolution $GFRes(x, z, y, p, a)$ too. \square

Finally, We extend each of the resolutions in the diagram K by their corresponding diagram K_M . We call the obtained diagram *the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$* , and we denote it briefly by *GFMRD*.

5.3 General Properties of the Graded Formal MR Diagram

Theorem 5.8. *Let*

$$\forall y \exists x \quad \Sigma(x, y, p, a) = 1 \vee \Psi(x, y, p, a) \neq 1,$$

*be an AE-formula. Let $Res(y, p, a)$ be a graded resolution, and denote its completion by $Comp(z, y, p, a)$. Denote by B the p -rigid or p -solid base group of $Comp(z, y, p, a)$. Let *GFMRD* be the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$. Let $p_0 \in F_k$, and let h_{p_0} be an exceptional solution of B that maps p to p_0 . Denote the ungraded resolution obtained from $Res(y, p, a)$ by specializing its terminal group according to h_{p_0} by $Res(y, p_0, a)$, and the corresponding ungraded completion by $Comp(z, y, p_0, a)$.*

Assume that the sentence:

$$\forall y \in Res(y, p_0, a) \exists x \quad \Sigma(x, y, p_0, a) = 1 \wedge \Psi(x, y, p_0, a) \neq 1,$$

is a truth sentence over F_k .

*Then, there exist some resolutions $GFRes_1(x, z, y, p, a), \dots, GFRes_m(x, z, y, p, a)$ in *GFMRD*, with corresponding terminal graded formal closures $GFCl_1(x, z, y, p, a), \dots, GFCl_m(x, z, y, p, a)$, together with some*

exceptional solutions $g_{p_0}^1, \dots, g_{p_0}^m$ for the p -rigid or p -solid base groups $Term(GFCl_1), \dots, Term(GFCl_m)$, so that:

1. For all $i = 1, \dots, m$, the solution $g_{p_0}^i$ restricts to an exceptional solution of B that belongs to the same exceptional family of h_{p_0} .
2. Every sequence of specializations $\{(x_n, z_n, y_n, p_0, a)\}$, for which the sequence $\{(z_n, y_n, p_0, a)\}$ is a test sequence through $Comp(z, y, p_0, a)$, and $\Sigma(x_n, y_n, p_0, a) = 1 \wedge \Psi(x_n, y_n, p_0, a) \neq 1$ for all n , subfactors through one of the resolutions $GFRes_1(x, z, y, p, a), \dots, GFRes_m(x, z, y, p, a)$.
3. The set of induced ungraded formal closures $FCl_1(x, z, y, p_0, a), \dots, FCl_m(x, z, y, p_0, a)$ forms a covering closure for $Comp(z, y, p_0, a)$.
4. For all $i = 1, \dots, m$, there exists a formal solution $x_i = x_i(s, z, y, p, a) \in GFCl_i$ that factors through the resolution $GFRes_i(x, z, y, p, a)$, so that the words $\Sigma(x_i, y, p, a)$ represent the trivial element in $GFCl_i$, and each of the words $\Psi(x_i, y, p_0, a)$ is non-trivial in $FCl_i(x, z, y, p_0, a)$.

Proof. Let $GFRes_1(x, z, y, p, a), \dots, GFRes_{m'}(x, z, y, p, a)$ be a minimal collection of resolutions in $GFMRD$, through which every sequence of specializations $\{(x_n, z_n, y_n, p_0, a)\}$, for which the sequence $\{(z_n, y_n, p_0, a)\}$ is a test sequence of $Comp(z, y, p_0, a)$, and $\Sigma(x_n, y_n, p_0, a) = 1 \wedge \Psi(x_n, y_n, p_0, a) \neq 1$ for all n , subfactors.

For every $i = 1, \dots, m'$, we consider a (boundedly many) specializations

$$(x_0^{i,1}, z_0^{i,1}, y_0^{i,1}, p_0, a), \dots, (x_0^{i,r(i)}, z_0^{i,r(i)}, y_0^{i,r(i)}, p_0, a)$$

that factor through $GFRes_i(x, z, y, p, a)$, satisfy $\Sigma(x_0^{i,s}, y_0^{i,s}, p_0, a) = 1 \wedge \Psi(x_0^{i,s}, y_0^{i,s}, p_0, a) \neq 1$, and so that the restriction $(z_0^{i,s}, y_0^{i,s}, p_0, a)$ is a specialization of $Comp(z, y, p_0, a)$, for all $s = 1, \dots, r(i)$.

for every $s = 1, \dots, r(i)$, the factorization of $(x_0^{i,s}, z_0^{i,s}, y_0^{i,s}, p_0, a)$ through $GFRes_i(x, z, y, p, a)$, induces a formal solution $x_i^s \in GFCl_i$ (which depends also on the specializations of the other free factors along the resolution), together with an ungraded formal closure $FCl_i^s(x, z, y, p_0, a)$ obtained by specializing the terminal group of the graded formal closure $GFCl_i$, $Term(GFCl_i)$, according to some exceptional solution $g_{p_0}^{i,s}$. Since $\Psi(x_0^{i,s}, y_0^{i,s}, p_0, a) \neq 1$, we deduce point 4.

Finally, if, by contradiction, the set of the obtained formal closures $\{FCl_i^s(x, z, y, p_0, a)\}_{i,s}$ did not form a covering closure for $Comp(z, y, p_0, a)$, then there will exist a test sequence of specializations $\{(z_n, y_n, p_0, a)\}$ of $Comp(z, y, p_0, a)$, so that the sepecialization (z_n, y_n, p_0, a) is not covered by the closures $\{FCl_i^s(x, z, y, p_0, a)\}$, for all n .

For all n , let $x_n \in F_k$ be such that $\Sigma(x_n, y_n, p_0, a) = 1 \wedge \Psi(x_n, y_n, p_0, a) \neq 1$. Then, by construction, the sequence $\{(x_n, z_n, y_n, p_0, a)\}$ subfactors through some $GFRes_i(x, z, y, p, a)$. The ungraded formal closure obtained from this factorization, contains the same ungraded completed resolution $Comp(z, y, p_0, a)$ embedded in some FCl_i^s . Hence (x_n, z_n, y_n, p_0, a) factors through FCl_i^s , a contradiction. \square

The graded formal MR diagram encodes all the formal solutions in the following sense.

Lemma 5.9. *Let $Res(y, p, a)$ be a graded resolution, and denote its completion by $Comp(z, y, p, a)$. Denote by B the p -rigid or p -solid base group of $Comp(z, y, p, a)$. Let $GFMRD$ be the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$.*

Let $GFCl(b, z, y, p, a)$ be a graded closure of $Comp(z, y, p, a)$, and denote the p -graded base limit group of $GFCl(b, z, y, p, a)$ by $Term(GFCl)$. Let x be an element of $GFCl(b, z, y, p, a)$ so that the words $\Sigma(x, z, y, p, a)$ represent the trivial element in $GFCl(b, z, y, p, a)$. Then, for every specialization of $GFCl(b, z, y, p, a)$, that restricts to an exceptional specialization of B , the induced specialization (x_0, z_0, y_0, p_0, a) of the tuple (x, z, y, p, a) , factors through one of the resolutions in $GFMRD$.

Proof. The idea of proving the property of the graded formal MR diagram stated in the lemma, is actually to use the method of its construction.

Let $GFRes_1(x, z, y, p, a), \dots, GFRes_r(x, z, y, p, a)$ be all the graded formal resolutions in $GFMRD$. We consider their completions $Comp(GFRes_1)(w, x, z, y, p, a), \dots, Comp(GFRes_r)(w, x, z, y, p, a)$. Let

$$\Omega_1(w, x, z, y, p, a), \dots, \Omega_r(w, x, z, y, p, a)$$

be the respective (finite) sets of defining relations of $Comp(GFRes_1), \dots, Comp(GFRes_r)$.

Let g_{p_0} be a specialization of $Term(GFCl)$, that maps p to some $p_0 \in F_k$, and restricts to an exceptional specialization of B . The solution g_{p_0} induces an ungraded closure $GFCl(b, z, y, p_0, a)$ that is covered by the given graded closure $GFCl(b, z, y, p, a)$. We emphasize the image of the given element x of $GFCl(b, z, y, p, a)$ in $GFCl(b, z, y, p_0, a)$, by writing $GFCl(x, b, z, y, p_0, a)$.

In a similar way for constructing the graded formal MR diagram, from the whole collection of sequences of specializations $\{(w_n, x_n, b_n, z_n, y_n, p_0, a)\}$, for which the sequences of restricted specializations $\{(x_n, b_n, z_n, y_n, p_0, a)\}$ are test sequences of the ungraded closure $GFCl(x, b, z, y, p_0, a)$ (i.e., $\{(z_n, y_n, p_0, a)\}$ is a test sequence of $Comp(z, y, p, a)$ so that $(x_n, b_n, z_n, y_n, p_0, a)$ is a specialization of $GFCl(x, b, z, y, p_0, a)$ for all n), and that satisfy one of the systems $\Omega_1(w, x, z, y, p_0, a), \dots, \Omega_r(w, x, z, y, p_0, a)$, we construct the formal MR diagram D of the systems $\Omega_1(w, x, z, y, p_0, a), \dots, \Omega_r(w, x, z, y, p_0, a)$ w.r.t. the closure $GFCl(x, b, z, y, p_0, a)$.

Every resolution in D , has to terminate in an ungraded formal closure $GFCl'(w, x, b, z, y, p_0, a)$ of $GFCl(x, b, z, y, p_0, a)$, so that the words in at least one of the systems $\Omega_1(w, x, z, y, p_0, a), \dots, \Omega_r(w, x, z, y, p_0, a)$ represent the trivial element in $GFCl'(w, x, b, z, y, p_0, a)$.

Now we explain the reason that the set of all the terminal closures in the diagram D , forms a covering closure for the resolution $GFCl(x, b, z, y, p_0, a)$. For that, it suffices to show that every test sequence of $GFCl(x, b, z, y, p_0, a)$ subfactors through some terminal formal closure $GFCl'(w, x, b, z, y, p_0, a)$ in D .

Indeed, if $\{(x_n, b_n, z_n, y_n, p_0, a)\}$ is a test sequence of $GFCl(x, b, z, y, p_0, a)$, then $\Sigma(x_n, z_n, y_n, p_0, a) = 1$ for all n , and the restricted sequence $\{(z_n, y_n, p_0, a)\}$ is a test sequence of $Comp(z, y, p, a)$. Hence, by the construction of the graded formal MR diagram $GFMRD$, there exists a sequence of specializations $\{w_{n_s}\}_s$, so that the specialization $(w_{n_s}, x_{n_s}, b_{n_s}, z_{n_s}, y_{n_s}, p_0, a)$ satisfies one of the systems $\Omega_1(w, x, z, y, p, a), \dots, \Omega_r(w, x, z, y, p, a)$, for all s . Hence, by the construction of D , the sequence $(x_n, b_n, z_n, y_n, p_0, a)$ (sub)factors through some terminal formal closure $GFCl'(w, x, b, z, y, p_0, a)$.

Finally, By the construction of D , every terminal formal closure $GFCl'(w, x, b, z, y, p_0, a)$ comes with some formal solution w for one of the systems $\Omega_1(w, x, z, y, p, a), \dots, \Omega_r(w, x, z, y, p, a)$, and the Lemma follows. \square

Theorem 5.10. *Let*

$$\forall y \exists x \quad \Sigma(x, y, p, a) = 1 \vee \Psi(x, y, p, a) \neq 1,$$

be an AE-formula. Let $Res(y, p, a)$ be a graded resolution, and denote its completion by $Comp(z, y, p, a)$. Denote by B the p -rigid or p -solid base group of $Comp(z, y, p, a)$. Let $GFMRD$ be the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$, that was constructed over the free group F_k .

Let Γ_l be a group of level l in the model. If l is large enough, then the following property is satisfied in Γ_l .

Every sequence $\{(x_n, z_n, y_n, p_n, a)\}$ of Γ_l -specializations for which the restricted sequence $\{(z_n, y_n, p_n, a)\}$ is a Γ_l -test sequence of $Comp(z, y, p, a)$, and $\Sigma(x_n, y_n, p_n, a) = 1$ in Γ_l for all n , factors through $GFMRD$.

Moreover, let $p_0 \in \Gamma_l$, and let h_{p_0} be a Γ_l -exceptional solution of B that maps p to p_0 . Denote by $Res(y, p_0, a)$ the ungraded resolution over Γ_l obtained from $Res(y, p, a)$ by specializing its terminal group according to h_{p_0} , and the corresponding Γ_l -ungraded completion by $Comp(z, y, p_0, a)$.

Assume that the sentence:

$$\forall y \in Res(y, p_0, a) \exists x \quad \Sigma(x, y, p_0, a) = 1 \wedge \Psi(x, y, p_0, a) \neq 1,$$

is a truth sentence over Γ_l .

Then, there exist a resolutions $GFRes_1(x, z, y, p, a), \dots, GFRes_m(x, z, y, p, a)$ in $GFMRD$, with corresponding terminal graded formal closures $GFCl_1(x, z, y, p, a), \dots, GFCl_m(x, z, y, p, a)$, together with a Γ_l -exceptional solutions $g_{p_0}^1, \dots, g_{p_0}^m$ for the p -rigid or p -solid base groups $Term(GFCl_1), \dots, Term(GFCl_m)$, so that:

1. *For all $i = 1, \dots, m$, the solution $g_{p_0}^i$ restricts to a Γ_l -exceptional solution of B that belongs to the same Γ_l -exceptional family of h_{p_0} .*
2. *Every sequence of Γ_l -specializations $\{(x_n, z_n, y_n, p_0, a)\}$, for which $\{(z_n, y_n, p_0, a)\}$ is a Γ_l -test sequence through $Comp(z, y, p_0, a)$, and $\Sigma(x_n, y_n, p_0, a) = 1 \wedge \Psi(x_n, y_n, p_0, a) \neq 1$ in Γ_l for all n , factors through one of the resolutions $GFRes_1(x, z, y, p, a), \dots, GFRes_m(x, z, y, p, a)$.*

3. The induced Γ_l -ungraded formal closures $FCl_1(x, z, y, p_0, a), \dots, FCl_m(x, z, y, p_0, a)$ form a Γ_l -covering closure for $Comp(z, y, p_0, a)$.

4. For all $i = 1, \dots, m$, there exists a formal solution $x_i = x_i(s, z, y, p, a) \in GFCl_i$ that factors through the resolution $GFRes_i(x, z, y, p, a)$, so that the words $\Sigma(x_i, y, p, a)$ represent the trivial element in $GFCl_i$, and each of the words $\Psi(x_i, y, p_0, a)$ is non-trivial in $FCl_i(x, z, y, p_0, a)$.

Proof. Denote by D the auxiliary graded formal MR diagram over random groups of the system $\Sigma(x, y, p, a)$ over the graded resolution $Res(y, p, a)$. Let (x_0, z_0, y_0, p_0, a) be a Γ_l -specialization of $Comp(z, y, p, a)$, that factors through a resolution $GFRes(x, z, y, p, a)$ in D . Denote by h_{p_0} the Γ_l -exceptional solution of B induced by (x_0, z_0, y_0, p_0, a) . According to the construction of D , the resolution $GFRes(x, z, y, p, a)$ terminates in a limit group of the form $\langle f \rangle * M$, where $\langle f \rangle$ is a free group, and $M = M(x, z, y, p, a)$ is a graded formal closure for $Comp(z, y, p, a)$. Moreover, there exists a formal solution $x\theta \in \langle f \rangle * M$, that factors through the resolution $GFRes(x, z, y, p, a)$, and a Γ_l -specialization (x'_0, z_0, y_0, p_0, a) of $\langle f \rangle * M$, so that $x\theta$ is mapped to x_0 . Denote by H_{p_0} the restriction of (x'_0, z_0, y_0, p_0, a) to M . Mapping $\langle f \rangle$ into the subgroup F_k of M , we get a formal solution, denoted again $x\theta$, that belongs to M , and that is mapped to x_0 by H_{p_0} . Note that, according to the construction of D , H_{p_0} restricts to h_{p_0} on B .

In particular, $x\theta$ is an element of the graded closure M of $Comp$, so that the words $\Sigma(x\theta, z, y, p, a)$ represent the trivial element in M . Thus, according to lemma 5.9, for every specialization of M in F_k , that restricts to an exceptional solution of B , the induced specialization of $(x\theta, z, y, p, a)$ factors through $GFMRD$.

We lift H_{p_0} to a specialization \tilde{H}_{p_0} of M in F_k . Denote by \tilde{h}_{p_0} the restriction of \tilde{H}_{p_0} to B . In particular, the solution \tilde{h}_{p_0} lifts h_{p_0} , and the specialization $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ in F_k that \tilde{H}_{p_0} induces for the tuple $(x\theta, z, y, p, a)$, lifts the Γ_l -specialization (x_0, z_0, y_0, p_0, a) . Since h_{p_0} is a Γ_l -exceptional solution of B , then \tilde{h}_{p_0} must be an exceptional solution of B over F_k .

Thus, according to lemma 5.9 the induced specialization $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ factors through $GFMRD$. This implies immediately that the original Γ_l -specialization (x_0, z_0, y_0, p_0, a) factors through $GFMRD$. Moreover, if $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ factors through the resolution $GFRes(x, z, y, p, a)$ of $GFMRD$ whose terminal closure is $GFCl(x, z, y, p, a)$, then (x_0, z_0, y_0, p_0, a) factors through the same resolution, and the induced Γ_l -specialization g_{p_0} of the base p -rigid or p -solid group $Term(GFCl)$ of $GFCl(x, z, y, p, a)$ restricts to a Γ_l -exceptional specialization of B that belongs to the same Γ_l -exceptional family of h_{p_0} .

If the Γ_l -specialization g_{p_0} is not itself a Γ_l -exceptional specialization of the base p -rigid or p -solid group $Term(GFCl)$ of $GFCl(x, z, y, p, a)$, then we lift it to a non-exceptional specialization of $Term(GFCl)$ over F_k . This replaces the specialization of $GFCl(x, z, y, p, a)$ induced by the factorization of $(\tilde{x}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ with another one, so that they are the same Γ_l -specializations. Hence, we can assume that g_{p_0} is a Γ_l -exceptional specialization of $Term(GFCl)$.

Now let $\{(x_n, z_n, y_n, p_n, a)\}$ a sequence of Γ_l -specializations, and assume that $\{(z_n, y_n, p_n, a)\}$ is a Γ_l -test sequence through $Comp(z, y, p, a)$, and $\Sigma(x_n, y_n, p_0, a) = 1$ in Γ_l for all n .

Then, according to lemma 5.3, the sequence $\{(x_n, z_n, y_n, p_n, a)\}$ subfactors over Γ_l through some resolution in the auxiliary graded formal MR diagram over random groups, that was denoted by D . Hence, according to the above argument, the sequence $\{(x_n, z_n, y_n, p_n, a)\}$ subfactors through $GFMRD$.

The remaining part of the theorem follows identically as the case over F_k . \square

The following fact is useful when treating a sentence that includes more than one inequality.

Lemma 5.11. *Let $Res(y, p, a)$ be a graded resolution, and denote its completion by $Comp(z, y, p, a)$. Denote by B the p -rigid or p -solid base group of $Comp(z, y, p, a)$. Let $GFMRD$ be the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$ (constructed over F_k).*

Let $GFRes(x, z, y, p, a)$ be a graded formal resolution in $GFMRD$, defined over the terminal closure $GFCl(x, z, y, p, a)$.

Let $\psi_1(x, y, p, a), \dots, \psi_s(x, y, p, a)$ be a finite collection of words, and let $x\theta_1, \dots, x\theta_s$ be formal solutions that factor through the resolution $GFRes(x, z, y, p, a)$. Let (x_0, z_0, y_0, p_0, a) be a specialization of $GFCl(x, z, y, p, a)$, so that the induced specialization $(x\theta_{i,0}, y_0, p_0, a)$, satisfies the inequality

$$\psi_i(x, y, p, a) \neq 1,$$

for all $i = 1, \dots, s$.

Then, there exists a formal solution $x\theta$, that factor through the resolution $GFRes(x, z, y, p, a)$, so that the induced specialization $(x\theta_0, y_0, p_0, a)$, satisfies the inequality $\psi_i(x, y, p, a) \neq 1$, for all $i = 1, \dots, s$.

Proof. Let $GFRes(x, z_0, y_0, p_0, a)$ be the ungraded formal resolution induced from the specialization (x_0, z_0, y_0, p_0, a) and covered by the resolution $GFRes(x, z, y, p, a)$. Let $FComp(t, x, z_0, y_0, p_0, a)$ be the completion of this resolution, and let $(t_n, x_n, z_0, y_0, p_0, a)$ be a test sequence of $GFCComp(t, x, z_0, y_0, p_0, a)$.

By the assumption, for all i , the formal solution $x\theta_i$ factors through $GFRes(x, z, y, p, a)$, so that

$$\psi_i(x\theta_{i,0}, y_0, p_0, a) \neq 1,$$

where $(x\theta_{i,0}, z_0, y_0, p_0, a)$ is the specialization of $(x\theta_i, z, y, p, a)$ induced by (x_0, z_0, y_0, p_0, a) . In particular, the word $\psi_i(x, y, p, a)$, viewed as an element of $FComp(t, x, z_0, y_0, p_0, a)$, must be non-trivial, for all i .

Hence, and since $(t_n, x_n, z_0, y_0, p_0, a)$ is a test sequence, for large enough n , we must have that

$$\psi_i(x_n, y_0, p_0, a) \neq 1.$$

□

Lemma 5.12. *Let $Res(y, p, a)$ be a graded resolution, and denote its completion by $Comp(z, y, p, a)$. Denote by B the p -rigid or p -solid base group of $Comp(z, y, p, a)$. Let $GFMRD$ be the graded formal MR diagram of the system $\Sigma(x, y, p, a)$ w.r.t. the resolution $Res(y, p, a)$ (constructed over F_k), and let $GFRes(x, z, y, p, a)$ be a graded formal resolution in $GFMRD$, defined over the terminal closure $GFCl(x, z, y, p, a)$.*

Let Γ_l be a group of level l in the model. If l is large enough, then the following property is satisfied in Γ_l .

Let $\psi_1(x, y, p, a), \dots, \psi_s(x, y, p, a)$ be a finite collection of words, and let $x\theta_1, \dots, x\theta_s$ be formal solutions that factor through the resolution $GFRes(x, z, y, p, a)$. Let (x_0, z_0, y_0, p_0, a) be a Γ_l -specialization of $GFCl(x, z, y, p, a)$, so that the induced Γ_l -specialization $(x\theta_{i,0}, y_0, p_0, a)$, satisfies the inequality

$$\psi_i(x, y, p, a) \neq 1,$$

for all $i = 1, \dots, s$.

Then, there exists a formal solution $x\theta$, that factor through the resolution $GFRes(x, z, y, p, a)$, so that the induced Γ_l -specialization $(x\theta_0, y_0, p_0, a)$, satisfies the inequality $\psi_i(x, y, p, a) \neq 1$, for all $i = 1, \dots, s$.

Proof. Let $GFRes(x, z_0, y_0, p_0, a)$ be the ungraded formal resolution induced from the Γ_l -specialization (x_0, z_0, y_0, p_0, a) and covered by the resolution $GFRes(x, z, y, p, a)$. Let $FComp(t, x, z_0, y_0, p_0, a)$ be the completion of this resolution, and let $(t_n, x_n, z_0, y_0, p_0, a)$ be a Γ_l -test sequence of $GFCComp(t, x, z_0, y_0, p_0, a)$.

By the assumption, for all i , the formal solution $x\theta_i$ factors through $GFRes(x, z, y, p, a)$, so that

$$\psi_i(x\theta_{i,0}, y_0, p_0, a) \neq 1,$$

where $(x\theta_{i,0}, z_0, y_0, p_0, a)$ is the Γ_l -specialization of $(x\theta_i, z, y, p, a)$ induced by (x_0, z_0, y_0, p_0, a) . In particular, the word $\psi_i(x, y, p, a)$, viewed as an element of $FComp(t, x, z_0, y_0, p_0, a)$, must be non-trivial, for all i .

Hence, and since $(t_n, x_n, z_0, y_0, p_0, a)$ is a Γ_l -test sequence, for large enough n , we must have that $\psi_i(x_n, y_0, p_0, a) \neq 1$. □

6 The *ExtraPS* MR Diagram over Random Groups

Along this section, we fix a collection $WPRS$ consisting of wp -rigid and wp -solid limit groups

$$G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$$

(wp are the parameter set). For all $i = 1, \dots, s$, we fix an integer $e_i \in \{0, \dots, M(G_i)\}$, where $M(G_i)$ is the maximal number of exceptional families of specializations in G_i for a fixed value of the parameters.

We consider a group $PT(y, w, p, a)$ that contains the amalgamated free product of the groups in $WPRS$

$$PT(y, w, p, a) \geq *_{i=1}^s *_{\langle w, p, a \rangle}^{e_i} G_i$$

amalgamated over the subgroup $\langle w, p, a \rangle$, as a subgroup. We consider the collection E of all the specializations (y_0, w_0, p_0, a) of $PT(y, w, p, a)$ for which the restriction of (y_0, w_0, p_0, a) to $*_{j=1}^{e_i} *_{\langle w, p, a \rangle} G_i$ represent a family of representatives of pairwise distinct exceptional families of G_i corresponding to the value $w_0 p_0$ of the parameters, for all $i = 1, \dots, s$.

The collection E factors through finitely many maximal limit groups $PS(y, w, p, a)$. For each $PS(y, w, p, a)$, we consider the subcollection EPS of E consisting of all the specializations in E that factor through $PS(y, w, p, a)$. we associate its standard p -graded MR diagram. We consider a resolution $Res(y, w, p, a)$ in this diagram (rooted at $PS(y, w, p, a)$), through which some specialization from E factors.

We consider the graded completion $Comp(z, y, w, p, a)$ of the resolution $Res(y, w, p, a)$, and we denote the terminal p -rigid or p -solid base group of $Comp(z, y, w, p, a)$ by B .

Lemma 6.1. *For all $i = 1, \dots, s$, all the mappings of G_i into any level of $Comp(z, y, w, p, a)$ are compatible. That is, if \tilde{G}_i is one of the canonical images of G_i in one of the levels of $Comp(z, y, w, p, a)$, whose associated decomposition is Π , for some $i = 1, \dots, s$, then the graph of groups obtained by collapsing the minimal subgraph of Π containing the subgroup $\langle w, p, a \rangle$ into the vertex AP , is compatible with the wp -JSJ of G_i , i.e., the minimal subgraph of it containing the image \tilde{G}_i consists only from parts of the wp -JSJ of G_i .*

Proof. If not, then, using the same argument in the proof of lemma 4.7, we can construct a formal solution(s) in (a covering closure of) $Comp(z, y, w, p, a)$ testifying flexibility of all the specializations of G_i that factor through $Comp(z, y, w, p, a)$, a contradiction. \square

Lemma 6.2. *Let Γ_l be a group in the model. The p -graded MR diagram of the groups $PS(y, w, p, a)$ encodes all the Γ_l -specializations (y_0, w_0, p_0, a) for which the restriction of (y_0, w_0, p_0, a) to $*_{j=1}^{e_i} *_{\langle w, p, a \rangle} G_i$ represent a family of representatives of pairwise distinct Γ_l -exceptional families of G_i corresponding to the value $w_0 p_0$ of the parameters, for all $i = 1, \dots, s$.*

Proof. Any lift of (y_0, w_0, p_0, a) factors through this diagram. \square

6.1 The Construction of *ExtraPS* MR Diagram over Random Groups

Let $Comp(z, y, w, p, a)$ be a graded completion, and assume that the mappings of G_i into $Comp(z, y, w, p, a)$ are compatible, for all $i = 1, \dots, s$ (see lemma 6.1). We are going to construct the auxiliary *ExtraPS* MR diagram over ascending sequences of groups in the model, for the collection $WPRS$ and the resolution $Comp(z, y, w, p, a)$.

We look at the collection \mathfrak{T} of all the sequences of specializations $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ over an ascending sequence of groups $\{\Gamma_{l_n}\}$ in the model, for which:

1. The sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ restricts to a p -graded test sequence of $Comp(z, y, w, p, a)$ over the ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model.
2. For all n , the specialization $(u_n, z_n, y_n, w_n, p_n, a)$ is a specialization of the amalgamated product

$$\left(*_{i=1}^s *_{\langle w, p, a \rangle}^{f_i} G_i \right) *_{\langle w, p, a \rangle} Comp(z, y, w, p, a),$$

where f_i is an integer in $f_i \in \{0, 1\}$.

3. For all n , the specialization $(u_n, z_n, y_n, w_n, p_n, a)$ is *non-collapsed*, i.e., for all $i = 1, \dots, s$, the $f_i + e_i$ specializations of G_i that the specialization $(u_n, z_n, y_n, w_n, p_n, a)$ induces, form a family of representatives of pairwise distinct Γ_{l_n} -exceptional families of G_i corresponding to the value $w_n p_n$ of the parameters (see definitions 4.1 and 4.2).

We apply the method of constructing the auxiliary graded formal MR diagram over random groups, on the collection \mathfrak{T} .

The sequences in the collection \mathfrak{T} , factors through finitely many limit groups $M(u, z, y, w, p, a)$. The canonical image of the subgroup $Comp(z, y, p, a)$ in M is denoted briefly by $Comp$. In light of point 3, the limit group M cannot be freely decomposable w.r.t. the subgroup $Comp \leq M$. We associate to M its relative $Comp$ -JSJ decomposition. Point 3 implies that for all $i = 1, \dots, s$, the mapping of G_i into M is compatible, i.e., the $Comp$ -JSJ of M is compatible with the wp -JSJ of G_i .

If the $Comp$ -JSJ of M is trivial, we call the group $M(u, z, y, w, p, a)$ a $Comp$ -rigid group, and terminate the construction for this stage. In this case, the subcollection of \mathfrak{T} consisting of all the sequences $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ that factors through M , is denoted by \mathfrak{T}_M .

In the case that the $Comp$ -JSJ of $M(u, z, y, w, p, a)$ is non-trivial, we consider the subcollection \mathfrak{T}_M of \mathfrak{T} , consisting of all sequences $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ that factor through M , and so that for all n , the specialization $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ cannot be shortened (in the metric of (Γ_{l_n}, a) , where a is the fixed generating set of Γ_{l_n}) by precomposing with modular automorphism of M that is induced from its $Comp$ -JSJ.

The sequences in the collection \mathfrak{T}_M , factor through finitely many limit quotients of M , that we denote by H . If H is not a proper quotient of M , we call M a $Comp$ -solid group, and terminate the construction for this stage.

We continue iteratively. Since all the groups that are obtained along the construction, are limit groups, then, according to the descending chain condition of limit groups, the construction must terminate after finitely many steps. The result is a diagram D that includes finitely many resolutions. Each of these resolutions terminates in a limit group $M(u, z, y, w, p, a)$ which is either $Comp$ -rigid or $Comp$ -solid.

Lemma 6.3. *Assume that $M(u, z, y, w, p, a)$ is a $Comp$ -rigid group or a $Comp$ -solid group. Then, M admits a structure of a graded closure of $Comp(z, y, w, p, a)$ over a p -graded limit group $Term(M)$. That is, M admits a structure of a p -graded tower over a p -graded base group $Term(M)$, so that:*

1. *up to adding roots to abelian groups, this structure is similar to the one associated with $Comp(z, y, w, p, a)$,*
2. *this structure is compatible to the one associated with $Comp(z, y, w, p, a)$*
3. *the canonical image of the p -graded base group B of $Comp(z, y, w, p, a)$ lies entirely in $Term(M)$, and B is mapped compatibly into $Term(M)$. That is, the minimal subgraph of groups of the p -JSJ of $Term(M)$ containing the canonical image \bar{B} of B , consists only from parts that are taken from the p -JSJ of B .*

Proof. This was proved in lemma 6.3. □

In light of the previous Lemma, every resolution of the diagram D that we have constructed above, is defined over a graded formal closure of the graded completion $Comp(z, y, w, p, a)$. And according to the construction of D , every sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ of non-collapsed specializations over an ascending sequence of groups in the model, that restricts to a test sequence $\{(z_n, y_n, w_n, p_n, a)\}$ over that ascending sequence of groups in the model, subfactors through at least one of the resolutions in D .

We call this diagram *the auxiliary ExtraPS MR diagram over random groups of the collection WPRS over the graded resolution $Comp(z, y, w, p, a)$* . Our aim is to show that the *ExtraPS* MR diagram over F_k of the collection *WPRS* over the graded resolution $Res(y, p, a)$, encodes all the sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ of non-collapsed Γ -specializations, that restricts to a Γ -test sequence $\{(z_n, y_n, w_n, p_n, a)\}$, for a random group Γ (in probability 1). The precise statement formulating this property, will be given below. For that aim, we start with the following lemma:

Lemma 6.4. *Let Γ_l be a group of level l in the model. If l is large enough, then every sequence of non-collapsed Γ_l -specializations $\{(u_n, z_n, y_n, w_n, p_n, a)\}$, that restricts to a Γ_l -test sequence $\{(z_n, y_n, w_n, p_n, a)\}$ of $Comp(z, y, w, p, a)$, subfactors through the auxiliary *ExtraPS* MR diagram over random groups D .*

Proof. Assume by contradiction that the statement is false. Then, there will be an ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model, so that for all n , there exists a sequence $\{(u_m^n, z_m^n, y_m^n, w_m^n, p_m^n, a)\}_m$ of non-collapsed Γ_{l_n} -specializations, so that the restricted sequence $\{(z_m^n, y_m^n, w_m^n, p_m^n, a)\}_m$ is a Γ_{l_n} -test sequence of $Comp(z, y, w, p, a)$, and the specialization $(u_m^n, z_m^n, y_m^n, w_m^n, p_m^n, a)$ does not factor through D for all m .

By the definition of test sequence over an ascending sequence of groups in the model, there exists a sequence $(m_n)_n$ of integers, so that the sequence $\{(u_{m_n}^n, z_{m_n}^n, y_{m_n}^n, w_{m_n}^n, p_{m_n}^n, a)\}_n$, which is defined over the ascending sequence $\{\Gamma_{l_n}\}$ of groups in the model, consists of non-collapsed specializations, restricts to a test sequence $\{(z_{m_n}^n, y_{m_n}^n, w_{m_n}^n, p_{m_n}^n, a)\}_n$ over the ascending sequence $\{\Gamma_{l_n}\}$, but it does not subfactor through D . This contradicts the construction of D . \square

6.2 The Construction of *ExtraPS* MR Diagram over Free Groups

Now we perform the parallel construction over F_k . We consider all the sequences of non-collapsed specializations $\{(u_n, z_n, y_n, w_n p_n, a)\}$ (over F_k), that restrict to test sequences $\{(z_n, y_n, w_n, p_n, a)\}$ of $Comp(z, y, w, p, a)$. We denote the collection of all such sequences $\{(u_n, z_n, y_n, w_n p_n, a)\}$, by \mathfrak{J} . Applying the shortening procedure iterative construction on the sequences in \mathfrak{J} , we construct an MR diagram, in a similar way to the construction that we explained above. The result of this construction over F_k is a finite diagram K , each of its resolutions $Res(u, z, y, w, p, a)$ terminates in a graded closure $M(u, z, y, w, p, a)$, and each of the limit groups along $Res(u, z, y, w, p, a)$ is $Comp$ -freely indecomposable, and admits a $Comp$ -JSJ that is compatible with the w -JSJ of G_i , for all $i = 1, \dots, s$.

For a resolution $Res(u, z, y, w, p, a)$, we consider its terminal graded closure $M(u, z, y, w, p, a)$, and denote its p -graded base group by $Term(M)$. We want to extend the resolution $Res(u, z, y, w, p, a)$ into finitely many resolutions, that terminate in a graded closure over a p -rigid or p -solid limit base group. For that, we consider the subcollection \mathfrak{J}_M of \mathfrak{J} consisting of all the sequences $\{(u_n, z_n, y_n, w_n p_n, a)\}$ for which:

1. If $Term(M)$ is either p -rigid or p -solid, then for all n , the specialization $(u_n, z_n, y_n, w_n p_n, a)$ restricts to a specialization of $Term(M)$ that factors through some flexible quotient of $Term(M)$.
2. Otherwise, for all n , the restriction of the specialization $(u_n, z_n, y_n, w_n p_n, a)$ into $Term(M)$ cannot be shortened by precomposing with a modular automorphism induced by the p -JSJ of $Term(M)$.

The subcollection \mathfrak{J}_M factors through finitely many maximal limit groups H . We continue with H by applying the shortening procedure iterative construction that was presented in the beginning of the construction. According to the descending chain condition of limit groups, this construction must terminate after finitely many steps. The result is a diagram K_M with finitely many resolutions, each of which terminates in a graded closure of $Comp(z, y, w, p, a)$.

Lemma 6.5. *Let $(u_0, z_0, y_0, w_0, p_0, a)$ be a non-collapsed specialization that factors through M . Then, there exists a specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ that factors through M so that:*

1. *The specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ restricts to the same modular family of $Term(M)$ (and thus to the same exceptional family of B).*
2. *The two families of representatives that the specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ induce, represent the same modular families of the group G_i , for all $i = 1, \dots, s$.*
3. *The specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ factors through one of the resolutions of K_M .*

Proof. Let t_0 be a specialization of $Term(M)$, that maps p to some p_0 in F_k . Assume that t_0 is short when $Term(M)$ is neither p -rigid, nor p -solid, and that t_0 factors through an exceptional quotient of $Term(M)$ otherwise. Let M_{t_0} be the induced ungraded closure.

Let Ω be a collection of (positive) Diophantine conditions $\Omega_1(q, u, z, y, w, p_0, a), \dots, \Omega_r(q, u, z, y, w, p_0, a)$ stating that the specialization (u, z, y, w, p_0, a) of M_{t_0} factors through one of the resolutions of K_M . And let Σ be a collection of (positive) Diophantine conditions $\Sigma_1(c, z, y, w, p_0, a), \dots, \Sigma_t(c, z, y, w, p_0, a)$ stating that the specialization (u, z, y, w, p_0, a) of M_{t_0} is collapsed.

We construct two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$ over the ungraded closure M_{t_0} , the first of the collection Ω , and the second for the collection Σ .

Let $(u_n, z_n, y_n, w_n, p_0, a)$ be a test sequence of M_{t_0} . Then, according to the construction of K_M , either $(u_n, z_n, y_n, w_n, p_0, a)$ contains a subsequence that is collapsed, or it subfactors through one of the resolutions of K_M . Hence, the collection of terminal ungraded closures in the two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$, forms a covering closure for M_{t_0} .

Assume that a specialization $(u_0, z_0, y_0, w_0, p_0, a)$ of M_{t_0} is non-collapsed. Then, it cannot factor through $FMRD(\Sigma)$. Hence, it must factor through one of the ungraded closures in $FMRD(\Omega)$. Due to the existence of a formal solution in that ungraded closure, we conclude that $(u_0, z_0, y_0, w_0, p_0, a)$ factors through one of the resolutions of K_M .

Finally, if $(u_0, z_0, y_0, w_0, p_0, a)$ is a specialization of $M_{t'_0}$, for some specialization t'_0 of $Term(M)$ that represents the same modular family of t_0 , then there exists some u''_0 so that the specialization $(u''_0, z_0, y_0, w_0, p_0, a)$ factors through M_{t_0} , and the two families of representatives that the specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u''_0, z_0, y_0, w_0, p_0, a)$ induce, represent the same modular families of the group G_i , for all $i = 1, \dots, s$. \square

We continue by extending each of the terminal graded closures in the diagram K_M , with a new diagram, in a similar way that K_M was constructed. We continue the construction iteratively. According to the descending chain condition of limit groups, this construction must terminate after finitely many steps. The result is an MR diagram, that extends the diagram D , and admits finitely many resolutions, each of which terminates in a graded formal closure $M'(u, z, y, w, p, a)$, whose terminal group is either p -rigid or p -solid limit group.

We call the obtained diagram *the ExtraPS MR diagram of the collection WPRS over the graded resolution $Comp(z, y, w, p, a)$* , and we denote it briefly by *ExtraPSMRD*.

6.3 General Properties of the ExtraPS MR Diagram

Theorem 6.6. *Let WPRS be a given collection of wp-rigid and wp-solid limit groups $G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$. Let $PS(y, w, p, a)$ be a limit quotient of $PT(y, w, p, a) = *_{i=1}^s *_{j=1}^{e_i} G_i$. Let $Res(y, w, p, a)$ be a p -graded resolution of $PS(y, w, p, a)$, and assume that there exists a non-collapsed specialization (y_0, w_0, p_0, a) that factors through $Res(y, w, p, a)$.*

Let $Comp(z, y, w, p, a)$ be the graded completion of $Res(y, w, p, a)$, and denote by B the p -rigid or p -solid base group of $Comp(z, y, w, p, a)$. Let ExtraPSMRD be the ExtraPS MR diagram of the collection WPRS over the resolution $Res(y, w, p, a)$. Let $ExtraPSGCl_1(u, z, y, w, p, a), \dots, ExtraPSGCl_m(u, z, y, w, p, a)$ be the terminal graded closures of the various resolutions of ExtraPSMRD.

For all n , let $(u_n, z_n, y_n, w_n, p_n, a)$ be a non-collapsed specialization of

$$*_{i=1}^s *_{j=1}^{f_i} G_i *_{(w, p, a)} Comp(z, y, w, p, a),$$

where f_i is an integer in $f_i \in \{0, 1\}$, and assume that the restricted sequence $\{(z_n, y_n, w_n, p_n, a)\}$ is a test sequence of $Comp(z, y, w, p, a)$.

Then, for some $k = 1, \dots, m$, and for all n in a subsequence of the integers, there exists a non-collapsed specialization $(u'_n, z_n, y_n, w_n, p_n, a)$, so that:

1. *The specializations $(u_n, z_n, y_n, w_n, p_n, a)$ and $(u'_n, z_n, y_n, w_n, p_n, a)$ restricts to solutions of B that represent the same exceptional family.*
2. *The two families of representatives that $(u_n, z_n, y_n, w_n, p_n, a)$ and $(u'_n, z_n, y_n, w_n, p_n, a)$ induce, represent the same exceptional families of the group G_i , for all $i = 1, \dots, s$.*
3. *The specialization $(u'_n, z_n, y_n, w_n, p_n, a)$ factors through the graded closure $ExtraPSGCl_k(u, z, y, w, p, a)$, so that the restriction of $(u'_n, z_n, y_n, w_n, p_n, a)$ to $Term(ExtraPSGCl_k)$ is an exceptional solution for it.*

Proof. By the construction of *ExtraPSMRD*, the sequence $(u_n, z_n, y_n, w_n, p_n, a)$ subfactors through some resolution of *ExtraPSMRD*, defined over some graded closure $M(u, z, y, w, p, a)$ in the diagram *ExtraPSMRD*. The remaining of the statement follows from lemma 6.1 and lemma 6.3. \square

The *ExtraPS MR diagram* encodes all the non-collapsed specialization of any graded closure of $Comp(z, y, w, p, a)$, in the following sense:

Lemma 6.7. Let $GFCl(b, z, y, w, p, a)$ be a graded closure of $Comp(z, y, w, p, a)$, and denote the p -graded base limit group of $GFCl(b, z, y, w, p, a)$ by $Term(GFCl)$. Assume that the amalgamated free product

$$\ast_{i=1}^s \ast_{j=1}^{f_i} G_i \ast_{(w,p,a)} Comp(z, y, w, p, a),$$

where f_i is an integer in $f_i \in \{0, 1\}$, is mapped into $GFCl(b, z, y, w, p, a)$, so that the subgroup $Comp(z, y, w, p, a)$ is mapped canonically. Let u be the images of the variables x_1, \dots, x_s of the given wp -rigid and wp -solid groups $G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$.

Then, for every specialization of $GFCl(b, z, y, w, p, a)$, that restricts to an exceptional specialization of B , if the induced specialization $(u_0, z_0, y_0, w_0, p_0, a)$ of the tuple (u, z, y, w, p, a) is non-collapsed, then $(u_0, z_0, y_0, w_0, p_0, a)$ factors through one of the resolutions in the *ExtraPS MR diagram* *ExtrPSMRD*.

Proof. Let g_{p_0} be a specialization of $Term(GFCl)$, that maps p to some $p_0 \in F_k$, and restricts to an exceptional specialization of B . The solution g_{p_0} induces an ungraded closure $GFCl_{g_{p_0}}$ that is covered by the given graded closure $GFCl(b, z, y, p, a)$. Note that $GFCl_{g_{p_0}}$ is generated by the (image of the) elements b, z, y, a . In particular the element $u \in GFCl_{g_{p_0}}$ is generated by these elements, $u = u(b, z, y, w, a)$.

Let Ω be a collection of (positive) Diophantine conditions

$$\Omega_1(q, u(b, z, y, w, a), z, y, w, p_0, a), \dots, \Omega_r(q, u(b, z, y, w, a), z, y, w, p_0, a)$$

stating that the specialization $(u(b, z, y, w, a), z, y, w, p_0, a)$ factors through the *ExtraPS MR diagram* *ExtraPSMRD*. And let Σ be a collection of (positive) Diophantine conditions

$$\Sigma_1(c, u(b, z, y, w, a), z, y, w, p_0, a), \dots, \Sigma_t(c, u(b, z, y, w, a), z, y, w, p_0, a)$$

stating that the specialization $(u(b, z, y, w, a), z, y, w, p_0, a)$ is collapsed.

We construct two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$ over the ungraded closure $GFCl_{g_{p_0}}$, the first of the collection Ω , and the second for the collection Σ .

Let $\{(b_n, z_n, y_n, w_n, p_0, a)\}$ be a test sequence of $GFCl_{g_{p_0}}$, and consider the induced specializations u_n of the element $u \in GFCl_{g_{p_0}}$. If for all n , the specialization $(u_n, z_n, y_n, w_n, p_0, a)$ is non-collapsed, then, according to the definition of the *ExtraPS MR diagram* *ExtrPSMRD*, the sequence $\{(u_n, z_n, y_n, w_n, p_0, a)\}$ subfactors through *ExtrPSMRD*. Hence, in this case, the sequence $\{(b_n, z_n, y_n, w_n, p_0, a)\}$ subfactors through $FMRD(\Omega)$.

If for all n , the specialization $(u_n, z_n, y_n, w_n, p_0, a)$ is collapsed, then the sequence $\{(b_n, z_n, y_n, w_n, p_0, a)\}$ subfactors through $FMRD(\Sigma)$.

We deduce that the collection of formal closures of $GFCl_{g_{p_0}}$, that are terminal closures in the diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$, form a covering closure for $GFCl_{g_{p_0}}$.

Finally, let $(b_0, z_0, y_0, w_0, p_0, a)$ be a specialization of $GFCl_{g_{p_0}}$, and assume that the induced specialization $(u_0, z_0, y_0, w_0, p_0, a)$ is non-collapsed. Then, due to the formal solutions in the terminal closures of $FMRD(\Sigma)$, the specialization $(b_0, z_0, y_0, w_0, p_0, a)$ cannot factor through a terminal closure of $FMRD(\Sigma)$. Hence, $(b_0, z_0, y_0, w_0, p_0, a)$ factors through a terminal closure of $FMRD(\Omega)$. The formal solution in that closure, testifies that the specialization $(u_0, z_0, y_0, w_0, p_0, a)$ factors through the *ExtraPS MR diagram* *ExtrPSMRD*. As required. \square

Theorem 6.8. Let *WPRS* be a given collection of wp -rigid and wp -solid limit groups $G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$.

Let $PS(y, w, p, a)$ be a limit quotient of $PT(y, w, p, a) = \ast_{i=1}^s \ast_{j=1}^{e_i} G_i$. Let $Res(y, w, p, a)$ be a p -graded resolution of $PS(y, w, p, a)$, and let $Comp(z, y, w, p, a)$ be the graded completion of $Res(y, w, p, a)$, and denote by B the p -rigid or p -solid base group of $Comp(z, y, w, p, a)$.

Let Γ_l be a group of level l in the model. If l is large enough, then the following properties are satisfied in Γ_l .

Assume that there exists a non-collapsed Γ_l -specialization (y_0, w_0, p_0, a) that factors through $Res(y, w, p, a)$. Then there exists a non-collapsed F_k -specialization $(\tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ that factors through the resolution $Res(y, w, p, a)$.

Moreover, let *ExtraPSMRD* be the *ExtraPS MR diagram* (over F_k) of the collection *WPRS* over the resolution $Res(y, w, p, a)$ (constructed over F_k). Let $ExtraPSGCl_1(u, z, y, w, p, a), \dots, ExtraPSGCl_m(u, z, y, w, p, a)$ be the terminal graded closures of the various resolutions of *ExtraPSMRD*.

For all n , let $(u_n, z_n, y_n, w_n, p_n, a)$ be a non-collapsed Γ_l -specialization of $\ast_{i=1}^s \ast_{j=1}^{f_i} G_i \ast_{(w,p,a)} \text{Comp}(z, y, w, p, a)$,

where f_i is an integer in $f_i \in \{0, 1\}$, so that the restricted sequence (z_n, y_n, w_n, p_n, a) is a Γ_l -test sequence of $\text{Comp}(z, y, w, p, a)$.

Then, for some $k = 1, \dots, m$, and for all n in a subsequence of the integers, there exists a non-collapsed Γ_l -specialization $(u'_n, z_n, y_n, w_n, p_n, a)$, so that:

1. The Γ_l -specializations $(u_n, z_n, y_n, w_n, p_n, a)$ and $(u'_n, z_n, y_n, w_n, p_n, a)$ restrict to solutions of B that represent the same Γ_l -exceptional family.
2. The two families of representatives that $(u_n, z_n, y_n, w_n, p_n, a)$ and $(u'_n, z_n, y_n, w_n, p_n, a)$ induce, represent the same Γ_l -modular families of the group G_i , for all $i = 1, \dots, s$.
3. The Γ_l -specialization $(u'_n, z_n, y_n, w_n, p_n, a)$ factors through the graded closure $\text{ExtraPSGCl}_k(u, z, y, w, p, a)$, so that the restriction of $(u'_n, z_n, y_n, w_n, p_n, a)$ to $\text{Term}(\text{ExtraPSGCl}_k)$ is a Γ_l -exceptional solution for it.

Proof. The first part of the claim follows immediately from that if $(\tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ is any lift of (y_0, w_0, p_0, a) that factors through $\text{Res}(y, w, p, a)$, then, $(\tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ is non-collapsed (over F_k).

Denote by D the auxiliary *ExtraPS* MR diagram of the collection *WPRS* over the resolution $\text{Res}(y, w, p, a)$. Let $(u_0, z_0, y_0, w_0, p_0, a)$ be a Γ_l -specialization of $\text{Comp}(z, y, w, p, a)$, that factors through a resolution $\text{GRes}(u, z, y, w, p, a)$ in D . Denote by h_{p_0} the Γ_l -exceptional solution of B induced by $(u_0, z_0, y_0, w_0, p_0, a)$. According to the construction of D , the resolution $\text{GRes}(u, z, y, w, p, a)$ terminates in a graded closure $M(u, z, y, w, p, a)$. Moreover, there exists an element $u\theta \in M$, that factors through the resolution $\text{GRes}(u, z, y, w, p, a)$ as the image of u , and a Γ_l -specialization $H_{p_0} = (u'_0, z_0, y_0, w_0, p_0, a)$ of M , so that $u\theta$ is mapped to u_0 . Note that, according to the construction of D , H_{p_0} restricts to h_{p_0} on B .

In particular, the amalgamated free product

$$\ast_{i=1}^s \ast_{j=1}^{f_i} G_i \ast_{(w,p,a)} \text{Comp}(z, y, w, p, a),$$

where f_i is an integer in $f_i \in \{0, 1\}$, is mapped into the graded closure M , so that the subgroup $\text{Comp}(z, y, w, p, a)$ is mapped canonically, and the elements $u\theta$ are the images of the variables x_1, \dots, x_s of the given groups $G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$.

Thus, according to lemma 6.7, for every specialization of M in F_k , that restricts to an exceptional solution of B , the induced specialization of $(u\theta, z, y, w, p, a)$ factors through *ExtraPSMRD*.

We lift H_{p_0} to a specialization \tilde{H}_{p_0} of M in F_k . Denote by \tilde{h}_{p_0} the restriction of \tilde{H}_{p_0} to B . In particular, the solution \tilde{h}_{p_0} lifts h_{p_0} , and the specialization $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ in F_k that \tilde{H}_{p_0} induces for the tuple $(u\theta, z, y, w, p, a)$, lifts the Γ_l -specialization $(u_0, z_0, y_0, w_0, p_0, a)$. Since h_{p_0} is a Γ_l -exceptional solution of B , then \tilde{h}_{p_0} must be an exceptional solution of B over F_k . Thus, the induced specialization $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ factors through *ExtraPSMRD*. This implies immediately that the original Γ_l -specialization $(u_0, z_0, y_0, w_0, p_0, a)$ factors through *ExtraPSMRD*. Moreover, if $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ factors through the resolution *ExtraPSGRes* (u, z, y, w, p, a) of *ExtraPSMRD* whose terminal closure is *ExtraPSGCl* (u, z, y, w, p, a) , then $(u_0, z_0, y_0, w_0, p_0, a)$ factors through the same resolution, and the induced Γ_l -specialization g_{p_0} of the base p -rigid or p -solid group $\text{Term}(\text{ExtraPSGCl})$ of *ExtraPSGCl* (u, z, y, w, p, a) restricts to a Γ_l -exceptional specialization of B that belongs to the same Γ_l -exceptional family of h_{p_0} .

If the Γ_l -specialization g_{p_0} is not itself a Γ_l -exceptional specialization of the base p -rigid or p -solid group $\text{Term}(\text{ExtraPSGCl})$ of *ExtraPSGCl* (u, z, y, w, p, a) , then we lift it to a non-exceptional specialization of $\text{Term}(\text{ExtraPSGCl})$ over F_k . This replaces the non-collapsed specialization *ExtraPSGCl* (u, z, y, w, p, a) induced by the factorization of $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ with another non-collapsed one, so that they are the same Γ_l -specializations. Hence, we can assume that g_{p_0} is a Γ_l -exceptional specialization of $\text{Term}(\text{ExtraPSGCl})$.

Now let $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ be a sequence of non-collapsed Γ_l -specializations, and assume that $\{(z_n, y_n, w_n, p_n, a)\}$ is a Γ_l -test sequence through $\text{Comp}(z, y, w, p, a)$. Then, the sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ subfactors through some resolution in the auxiliary *ExtraPS* MR diagram over random groups, that was denoted by D . Hence, according to the above argument, the sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ subfactors through *ExtraPSMRD*.

The remaining part of the theorem follows identically as the case over F_k . \square

7 The Construction of the CollapseExtra MR Diagram

Let $ExtraPSGCl(u, z, y, w, p, a)$ be one of the terminal graded closures in the extra MR diagram $ExtraMRD$ of the collection $WPRS$ over the p -graded resolution $Res(y, w, p, a)$ (constructed in section §6). Let $Comp(z, y, w, p, a)$ be the graded completion of $Res(y, w, p, a)$, and denote its terminal p -rigid or p -solid base group by B . Denote the p -graded base group of $ExtraPSGCl(u, z, y, w, p, a)$, which is either p -rigid or p -solid, by $Term(ExtraPSGCl)$.

Let $\Sigma_i(c, u, z, y, w, p, a)$, $i = 1, \dots, r$, be the finitely many systems that testify that the specialization (u, z, y, w, p, a) is collapsed. We are going to construct a graded formal MR diagram of the systems $\Sigma_i(c, u, z, y, w, p, a)$ over the graded closure $ExtraPSGCl(u, z, y, w, p, a)$.

We consider the collection \mathfrak{J} of all the sequences of specializations $\{(c_n, u_n, z_n, y_n, w_n, p_n, a)\}$, for which the restricted sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ is a sequence of specializations of $ExtraPSGCl(u, z, y, w, p, a)$, the restricted sequence $\{(z_n, y_n, w_n, p_n, a)\}$ is a test sequence of $Comp(z, y, w, p, a)$, and for all n :

1. The restricted specialization (y_n, w_n, p_n, a) is non-collapsed.
2. The specialization $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ restricts to an exceptional solution of the base group $Term(ExtraPSGCl)$ of $ExtraPSGCl$, and to an exceptional solution of B .
3. The specialization c_n explains how the specialization $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ is collapsed, i.e., the elements of one of the systems $\Sigma_i(c_n, u_n, z_n, y_n, w_n, p_n, a)$ represent the trivial word.

The sequences in the collection \mathfrak{J} , factor through finitely many limit groups $H(c, u, z, y, w, p, a)$. The canonical image of the subgroup $Comp(z, y, w, p, a)$ in H is denoted briefly by $Comp$, and the canonical image of the subgroup $ExtraPSGCl(u, z, y, w, p, a)$ in H is denoted by $ExtraPSGCl$. The limit group H could be freely decomposable w.r.t. the subgroup $Comp \leq H$. We consider the most refined $Comp$ -free factorization of H . We keep the free factors that do not include (a conjugate of) the subgroup $Comp \leq H$, extend them by their standard MR diagrams, and continue the construction with the free factor that contains $Comp$. We denote this factor by $M(c, u, z, y, w, p, a)$, which is a limit group. We consider the $Comp$ -JSJ decomposition of the limit group $M(c, u, z, y, w, p, a)$, i.e., the JSJ decomposition with respect to the subgroup $Comp$.

Lemma 7.1. *The canonical image $ExtraPSGCl$ of $ExtraPSGCl(u, z, y, w, p, a)$ is contained in the free factor M of H containing $Comp$. Moreover, the $Comp$ -JSJ of M is compatible with the p -JSJ of $Term(ExtraPSGCl)$.*

Proof. According to the definition of H and the sequences in \mathfrak{T} , there exists a specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ of H that restricts to an exceptional specialization of $Term(ExtraPSGCl)$. Hence, the claim follows using lemma 4.7. \square

If the $Comp$ -JSJ of $M(c, u, z, y, w, p, a)$ is trivial, we call the group $M(c, u, z, y, w, p, a)$ a $Comp$ -rigid group, and terminate the construction for this stage. In this case, the subcollection of \mathfrak{T} consisting of all the sequences $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ that factor through M , is denoted by \mathfrak{J}_M .

In the case that the $Comp$ -JSJ of $M(c, u, z, y, w, p, a)$ is non-trivial, we consider the subcollection \mathfrak{J}_M of \mathfrak{J} , consisting of all the sequences $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ that factor through M , and so that for all n , the specialization $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ cannot be shortened (in the metric of (F_k, a)) by precomposing with modular automorphism of M that is induced from its $Comp$ -JSJ. The sequences in the collection \mathfrak{J}_M , factor through finitely many limit quotients of M , that we denote - while abusing the notation - by H . If H is not a proper quotient of M , we call M a $Comp$ -solid group, and terminate the construction for this stage.

Lemma 7.2. *Let $\{(c_n, u_n, z_n, y_n, w_n, p_n, a)\}$ be a sequence in the collection \mathfrak{J} that factors through M . Then, the shortened sequence $\{((c\theta)_n, (u\theta)_n, z_n, y_n, w_n, p_n, a)\}$ belongs to \mathfrak{J} .*

Proof. For all n , let $((c\theta)_n, (u\theta)_n, z_n, y_n, w_n, p_n, a)$ be a specialization of M that belongs to the same modular family of $(c_n, u_n, z_n, y_n, w_n, p_n, a)$. According to lemma 7.1, and according to the construction of the $ExtraPS$ MR diagram, the two families of representatives, that the specializations $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ and $((c\theta)_n, (u\theta)_n, z_n, y_n, w_n, p_n, a)$ induce, represent the same exceptional families for the group $G_i \in WPRS$, for all $i = 1, \dots, s$. \square

We continue the construction iteratively. Since all the groups that are obtained along the construction, are limit groups, then, according to the descending chain condition of limit groups, the construction must

terminate after finitely many steps. The result is a diagram K that includes finitely many resolutions. Each of these resolutions terminates in a free product $\langle f \rangle * M$, where $\langle f \rangle$ is a free group, and M is either a *Comp*-rigid group or a *Comp*-solid group.

Lemma 7.3. *Assume that $M(c, u, z, y, w, p, a)$ is a *Comp*-rigid group or a *Comp*-solid group. Then, M admits a structure of a graded formal closure of $\text{Comp}(z, y, w, p, a)$ over a p -graded limit group $\text{Term}(M)$. That is, M admits a structure of a p -graded tower over a p -graded base group $\text{Term}(M)$, so that:*

1. *up to adding roots to abelian groups, this structure is similar to the one associated with $\text{Comp}(z, y, w, p, a)$,*
2. *this structure is compatible to the one associated with $\text{Comp}(z, y, w, p, a)$*
3. *the canonical image of the p -graded base group B of $\text{Comp}(z, y, w, p, a)$ lies entirely in $\text{Term}(M)$, and B is mapped compatibly into $\text{Term}(M)$. That is, the minimal subgraph of groups of the p -JSJ of $\text{Term}(M)$ containing the canonical image \bar{B} of B , consists only from parts that are taken from the p -JSJ of B .*

Moreover, the mapping of $\text{Term}(\text{ExtraPSGCl})$ into any level of $\text{Comp}(z, y, w, p, a)$ is compatible.

Proof. The proof of the first part is done in the proof of lemma 5.2.

If the last statement was wrong, then, using the same argument in the proof of lemma 4.7, we can construct a formal solution(s) in (a covering closure of) $\text{Comp}(z, y, w, p, a)$ testifying flexibility of all the specializations of $\text{Term}(\text{ExtraPSGCl})$ that factor through M . \square

Lemma 7.4. *Let t_0 be a specialization of $\text{Term}(M)$, the p -graded base group of M . Consider the induced ungraded closure M_{t_0} , and assume that there exists a specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ that factors through the ungraded closure M_{t_0} and satisfies the conditions 1-3 in the top of the section.*

Then, there exists a test sequence $\{(z_n, y_n, w_n, p_0, a)\}$ of $\text{Comp}(z, y, w, p, a)$, so that for every specialization t'_0 of $\text{Term}(M)$ that belongs to the same modular family of t_0 , the specialization (z_n, y_n, w_n, p_0, a) factors through the ungraded closure $M_{t'_0}$, and the induced specialization $(c'_n, u'_n, z_n, y_n, w_n, p_0, a)$ satisfies the conditions 1-3, for all n .

Proof. Since $\text{Term}(\text{ExtraPSGCl})$ is mapped compatibly into M , and $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ is a specialization of M_{t_0} that restricts to an exceptional solution for $\text{Term}(\text{ExtraPSGCl})$ and B , we conclude that every specialization of $M_{t'_0}$, where t'_0 is a specialization of $\text{Term}(M)$ in the same modular family of t_0 , satisfies the conditions 2-3 in the beginning of the section.

Now a similar argument of the proof of lemma 6.5, there must exist some test sequence $\{(z_n, y_n, w_n, p_0, a)\}$ of $\text{Comp}(z, y, w, p, a)$ consisting of non-collapsed specializations, that factor through M_{t_0} . Hence, if t'_0 is a specialization of $\text{Term}(M)$ that belongs to the same modular family of t_0 , the specialization (z_n, y_n, w_n, p_0, a) factors through $M_{t'_0}$, and the induced specialization $(c'_n, u'_n, z_n, y_n, w_n, p_0, a)$ satisfies the conditions 1-3, for all n . \square

We continue the construction of our graded formal MR diagram. For a resolution $\text{GRes}(c, u, z, y, w, p, a)$ in the diagram K constructed so far, we consider its terminal graded formal closure $M(c, u, z, y, w, p, a)$, and denote its p -graded base group by $\text{Term}(M)$. We want to extend the resolution $\text{GRes}(c, u, z, y, w, p, a)$ into finitely many resolutions, that terminate in a graded closure over a p -rigid or p -solid base limit group. For that, we consider the subcollection \mathfrak{J}_M of \mathfrak{J} consisting of all the sequences of specializations $\{(c_n, u_n, z_n, y_n, w_n, p_n, a)\}$ of M , for which:

1. If $\text{Term}(M)$ is either p -rigid or p -solid, then, for all n , the specialization $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ restricts to a specialization of $\text{Term}(M)$ that factors through some flexible quotient of $\text{Term}(M)$.
2. Otherwise, for all n , the restriction of the specialization $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ into $\text{Term}(M)$ cannot be shortened by precomposing with a modular automorphism induced by the p -JSJ of $\text{Term}(M)$.

The subcollection \mathfrak{J}_M factors through finitely many maximal limit groups H . We continue with H by applying the shortening procedure iterative construction that was presented in the beginning of the construction. According to the descending chain condition of limit groups, this construction must terminate after finitely many steps. The result is a diagram K_M with finitely many resolutions, each of which terminates in a graded formal closure of $\text{Comp}(z, y, w, p, a)$.

Lemma 7.5. *Let $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ be a specialization of M that satisfies the conditions 1-3 in the top of the section. Then, there exists a specialization $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ that factors through M so that:*

1. *The specializations $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ and $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ restricts to the same modular family of $\text{Term}(M)$ (and thus to the same exceptional family of $\text{Term}(\text{ExtraPSGCl})$, and to the same exceptional family of B).*
2. *The two families of representatives that the specializations $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ and $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ induce, represent the same modular families of the group G_i , for all $i = 1, \dots, s$.*
3. *The specialization $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ satisfies the conditions 1-3 in the top of the section, and factors through one of the resolutions $\text{GFRes}(c, u, z, y, w, p, a)$ of K_M .*

Moreover, if $\text{Term}(M)$ is either p -rigid or p -solid, and the restriction of the specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $\text{Term}(M)$ is not exceptional, then we can assume that the terminal graded formal closure of the resolution $\text{GFRes}(c, u, z, y, w, p, a)$ is a proper quotient of M .

Proof. Let t_0 be a specialization of $\text{Term}(M)$, that maps p to some p_0 in F_k . Assume that t_0 is short when $\text{Term}(M)$ is neither p -rigid, nor p -solid, and that t_0 factors through an exceptional quotient of $\text{Term}(M)$ otherwise. Let M_{t_0} be the induced ungraded closure, and assume that $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ is a specialization of M_{t_0} that satisfies the conditions 1-3.

Let Ω be a collection of (positive) Diophantine conditions $\Omega_1(q, c, u, z, y, w, p_0, a), \dots, \Omega_r(q, c, u, z, y, w, p_0, a)$ stating that the specialization (c, u, z, y, w, p_0, a) of M factors through one of the resolutions of K_M (without M itself). And let Σ be a collection of (positive) Diophantine conditions $\Sigma_1(f, c, u, z, y, w, p_0, a), \dots, \Sigma_t(f, c, u, z, y, w, p_0, a)$ stating that the specialization (c, u, z, y, w, p_0, a) of M satisfies that the restriction (z, y, w, p_0, a) is collapsed.

We construct two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$ over the ungraded closure M_{t_0} , the first of the collection Ω , and the second for the collection Σ .

Let $\{(c_n, u_n, z_n, y_n, w_n, p_0, a)\}$ be a test sequence of M_{t_0} . Note that, for all n , if the restriction (z_n, y_n, w_n, p_0, a) is non-collapsed, then the specialization $(c_n, u_n, z_n, y_n, w_n, p_0, a)$ satisfies the conditions 1-3. Thus, according to the construction of K_M , the sequence $\{(c_n, u_n, z_n, y_n, w_n, p_0, a)\}$ either contains a subsequence for which the restricted sequence $\{(z_n, y_n, w_n, p_0, a)\}$ are collapsed, or it subfactors through one of the resolutions of K_M . Hence, the collection of terminal ungraded closures in the two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$, forms a covering closure for M_{t_0} .

Now consider the specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ of M_{t_0} that satisfies conditions 1-3. It cannot factor through $FMRD(\Sigma)$. Hence, it must factor through one of the ungraded closures in $FMRD(\Omega)$. Due to the existence of a formal solution in that ungraded closure, we conclude that the specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ factors through one of the resolutions of K_M .

Finally, if $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ is a specialization of $M_{t'_0}$, for some specialization t'_0 of $\text{Term}(M)$ that represents the same modular family of t_0 , then there exists some $c''_0, u''_0 \in F_k$ so that the specialization $(c''_0, u''_0, z_0, y_0, w_0, p_0, a)$ factors through M_{t_0} , and the two families of representatives that the specializations $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ and $(c''_0, u''_0, z_0, y_0, w_0, p_0, a)$ induce, represent the same modular families of the group G_i , for all $i = 1, \dots, s$. \square

Lemma 7.6. *Let Γ_l be a group of level l in the model. If l is large enough, then the following property is satisfied for Γ_l .*

Let $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ be a Γ_l -specialization of M that satisfies the conditions 1-3 in the top of the section. Then, there exists a Γ_l -specialization $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ that factors through M so that:

1. *The Γ_l -specializations $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ and $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ restricts to the same Γ_l -modular family of $\text{Term}(M)$ (and thus to the same Γ_l -exceptional family of $\text{Term}(\text{ExtraPSGCl})$, and to the same Γ_l -exceptional family of B).*
2. *The two families of representatives that the Γ_l -specializations $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ and $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ induce, represent the same Γ_l -modular families of the group G_i , for all $i = 1, \dots, s$.*
3. *The Γ_l -specialization $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ satisfies the conditions 1-3 in the top of the section, and factors through one of the resolutions $\text{GFRes}(c, u, z, y, w, p, a)$ of K_M .*

Moreover, if $\text{Term}(M)$ is either p -rigid or p -solid, and the restriction of the Γ_l -specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $\text{Term}(M)$ is not Γ_l -exceptional, then we can assume that the terminal graded formal closure of the resolution $\text{GFRes}(c, u, z, y, w, p, a)$ is a proper quotient of M .

Proof. Every lift $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to a specialization of M in F_k , satisfies the conditions 1-3. We lift $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to a specialization $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ of M in F_k , so that in the case that $\text{Term}(M)$ is either p -rigid or p -solid, and the restriction of the Γ_l -specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $\text{Term}(M)$ is not Γ_l -exceptional, the restriction of the lift $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ to $\text{Term}(M)$ is not F_k -exceptional.

Hence, according to lemma 7.5, there exists an F_k -specialization $(\tilde{c}'_0, \tilde{u}'_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ of M that restricts to the modular family of $\text{Term}(M)$ represented by $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$, and factors through one of the resolutions $\text{GFRes}(c, u, z, y, w, p, a)$ of K_M . Let $(c'_0, u'_0, z_0, y_0, w_0, p_0, a)$ be the projection of $(\tilde{c}'_0, \tilde{u}'_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ to Γ_l . Then, $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ is a Γ_l -specialization of M that restricts to the Γ_l -modular family of $\text{Term}(M)$ represented by $(c_0, u_0, z_0, y_0, w_0, p_0, a)$, and factors through the resolution $\text{GFRes}(c, u, z, y, w, p, a)$ of K_M . The other details of the claim follows immediately from this. \square

Finally, we continue by extending each of the terminal graded closures in the diagram K_M , with a new diagram, in a similar way that K_M was constructed. We continue the construction iteratively. According to the descending chain condition of limit groups, this construction must terminate after finitely many steps. The result is an MR diagram, that extends the diagram K , and admits finitely many resolutions, each of which terminates in a graded formal closure $M'(u, z, y, w, p, a)$, whose terminal group is either a p -rigid or a p -solid limit group.

We call the obtained diagram *the CollapseExtra MR diagram of the collection WPRS over the graded closure ExtraPSGCl* (u, z, y, w, p, a) , and we denote it by $\text{CollapseExtraMRD}(\text{ExtraPSGCl})$, or briefly CollapseExtraMRD . We summarize the general properties of the construction in the next theorem.

7.1 General Properties of CollapseExtra MR Diagram

Theorem 7.7. *Let $\text{ExtraPSGCl}(u, z, y, w, p, a)$ be a terminal graded closure in ExtraPSMRD .*

Let CollapseExtraMRD be the CollapseExtra MR diagram of the collection WPRS and the graded closure $\text{ExtraPSGCl}(u, z, y, w, p, a)$.

Let $\{(c_n, u_n, z_n, y_n, w_n, p_n, a)\}$ be a sequence of specializations, for which the restricted sequence $\{(u_n, z_n, y_n, w_n, p_n, a)\}$ is a sequence of specializations of $\text{ExtraPSGCl}(u, z, y, w, p, a)$, the restricted sequence $\{(z_n, y_n, w_n, p_n, a)\}$ is a test sequence of $\text{Comp}(z, y, w, p, a)$, and for all n :

1. *The restriction (y_n, w_n, p_n, a) is non-collapsed.*
2. *The specialization $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ restricts to an exceptional solution of the base group $\text{Term}(\text{ExtraPSGCl})$ of ExtraPSGCl , and to an exceptional solution of B .*
3. *The specialization c_n explains how the specialization $(u_n, z_n, y_n, w_n, p_n, a)$ is collapsed.*

Then, for all n in a subsequence of the integers, there exist $c'_n, u'_n \in F_k$, so that:

1. *The specialization $(c'_n, u'_n, z_n, y_n, w_n, p_n, a)$ factors through one of the terminal graded formal closures $\text{CollapseExtraGFCl}(c, u, z, y, w, p, a)$ of CollapseExtraMRD .*
2. *The specialization $(c'_n, u'_n, z_n, y_n, w_n, p_n, a)$ restricts to an exceptional solution of the p -rigid or p -solid base group $\text{Term}(\text{CollapseExtraGFCl})$ of $\text{CollapseExtraGFCl}(c, u, z, y, w, p, a)$.*
3. *The specialization $(c'_n, u'_n, z_n, y_n, w_n, p_n, a)$ restricts to an exceptional solution of the p -rigid or p -solid base group $\text{Term}(\text{ExtraPSGCl})$ of $\text{ExtraPSGCl}(u, z, y, w, p, a)$, that belongs to the same exceptional family of the restriction of $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ to $\text{Term}(\text{ExtraPSGCl})$.*
4. *The specialization $(c'_n, u'_n, z_n, y_n, w_n, p_n, a)$ restricts to an exceptional solution of B , the p -rigid or p -solid base group of $\text{Comp}(z, y, w, p, a)$, that belongs to the same exceptional family of the restriction of $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ to B .*

5. The specializations $(c_n, u_n, z_n, y_n, w_n, p_n, a)$ and $(c'_n, u'_n, z_n, y_n, w_n, p_n, a)$ are collapsed in the same form.

The *CollapseExtra* MR diagram $\text{CollapseExtra}(\text{ExtraPSGCl})$ encodes all the specializations of $\text{ExtraPSGCl}(u, z, y, w, p, a)$ that are collapsed in a formal way, in the following more precise sense:

Lemma 7.8. *Let $\text{ExtraPSGCl}(u, z, y, w, p, a)$ be a terminal graded closure in ExtraPSMRD .*

*Let CollapseExtraMRD be the *CollapseExtra* MR diagram of the collection $WPRS$ over the graded closure $\text{ExtraPSGCl}(u, z, y, w, p, a)$. Let Λ be a collection of (positive) Diophantine conditions stating that the specialization (u, z, y, w, p_0, a) is collapsed.*

Let $GCl(b, z, y, w, p, a)$ be a graded closure of $\text{Comp}(z, y, w, p, a)$, and denote its p -graded base limit group by $\text{Term}(GCl)$. Assume that the graded closure $\text{ExtraPSGCl}(u, z, y, w, p, a)$ is mapped into $GCl(b, z, y, w, p, a)$, so that the subgroup $\text{Comp}(z, y, w, p, a)$ is mapped canonically. Assume that there exists an element $c \in GCl(b, z, y, w, p, a)$ so that the words of one of the systems $\Lambda_i(c, u, z, y, w, p, a)$ in Λ , $i = 1, \dots, s$, represent the trivial element in $GCl(b, z, y, w, p, a)$.

*Then, every specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ that factors through $GCl(b, z, y, w, p, a)$, and satisfies the conditions 1-3 in the top of the section, factors through the *CollapseExtra* MR diagram CollapseExtraMRD .*

Proof. Assume that there exists a specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ that factors through GCl , and satisfies the conditions 1-3 in the top of the section. Then, the mapping of $\text{Term}(\text{ExtraPSGCl})$ into any level of GCl is compatible.

Let g_{p_0} be the restriction of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $\text{Term}(GCl)$. The solution g_{p_0} induces an ungraded closure $GCl_{g_{p_0}}$ that is covered by the given graded closure $GCl(b, z, y, w, p, a)$. Note that $GCl_{g_{p_0}}$ is generated by the (image of the) elements b, z, y, w, a . In particular the elements $u, c \in GCl_{g_{p_0}}$ are generated by these elements, $u = u(b, z, y, w, a)$, $c = c(b, z, y, w, a)$. Moreover, every specialization $(c'_0, u'_0, z'_0, y'_0, w'_0, p_0, a)$ of $GCl_{g_{p_0}}$ satisfies the conditions 2-3 in the beginning of the section.

Let Ω be a collection of (positive) Diophantine conditions

$$\Omega_1(q, c(b, z, y, w, a), u(b, z, y, w, a), z, y, w, p_0, a), \dots, \Omega_r(q, c(b, z, y, w, a), u(b, z, y, w, a), z, y, w, p_0, a)$$

stating that the specialization $(c(b, z, y, w, a), u(b, z, y, w, a), z, y, w, p_0, a)$ factors through the *CollapseExtra* MR diagram CollapseExtraMRD .

And let Σ be a collection of (positive) Diophantine conditions

$$\Sigma_1(f, c(b, z, y, w, a), u(b, z, y, w, a), z, y, w, p_0, a), \dots, \Sigma_t(f, c(b, z, y, w, a), u(b, z, y, w, a), z, y, w, p_0, a)$$

stating that the specialization (c, u, z, y, w, p_0, a) satisfies that the restriction (z, y, w, p_0, a) is collapsed.

We construct two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$ over the ungraded closure $GCl_{g_{p_0}}$, the first of the collection Ω , and the second for the collection Σ .

Let $\{(b_n, z_n, y_n, w_n, p_0, a)\}$ be a test sequence of $GCl_{g_{p_0}}$, and consider the induced specializations $\{u_n\}$ and $\{c_n\}$ of the elements $u, c \in GCl_{g_{p_0}}$. Then, the sequence $\{(c_n, u_n, c_n, y_n, w_n, p_0, a)\}$ either admits a subsequence for which the restricted specializations $\{(z_n, y_n, w_n, p_0, a)\}$ are collapsed, or it subfactors through one of the resolutions of CollapseExtraMRD . Hence, the collection of terminal ungraded closures in the two formal MR diagrams $FMRD(\Omega)$ and $FMRD(\Sigma)$, forms a covering closure for $GCl_{g_{p_0}}$.

Now consider the specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ that factors through $GCl_{g_{p_0}}$ and satisfies the conditions 1-3. It cannot factor through $FMRD(\Sigma)$, and hence it factors through some terminal ungraded closure of $FMRD(\Omega)$. Due to the formal solution in that ungraded closure, we deduce that $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ factors through CollapseExtraMRD . □

Lemma 7.9. *Keep the notation of 7.8. Let Γ_l be a group of level l in the model. If l is large enough, the the following property is satisfied in Γ_l .*

*Every Γ_l -specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ that factors through $GCl(b, z, y, w, p, a)$, and satisfies the conditions 1-3 in the top of the section, factors through the *CollapseExtra* MR diagram CollapseExtraMRD .*

Proof. We lift $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to a specialization $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ that factors through $GCl(b, z, y, w, p, a)$. Then, the specialization $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ satisfies the conditions 1-3, and thus, according to lemma 7.8, it factors through CollapseExtraMRD . Hence, the Γ_l -specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ factors through CollapseExtraMRD . □

Lemma 7.10. *Let $\text{ExtraPSGCl}(u, z, y, w, p, a)$ be a terminal graded closure in ExtraPSMRD .*

Let CollapseExtraMRD be the CollapseExtra MR diagram of the collection $WPRS$ and the graded closure $\text{ExtraPSGCl}(u, z, y, w, p, a)$.

Let $\text{CollapseExtraGFCl}(c, u, z, y, w, p, a)$ be a terminal graded formal closure in CollapseExtraMRD , and let $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ be a specialization of $\text{CollapseExtraGFCl}(c, u, z, y, w, p, a)$.

Then, every specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ of $\text{ExtraPSGCl}(u, z, y, w, p, a)$, that restricts to a solution of $\text{Term}(\text{ExtraPSGCl})$ that belongs to the same modular family of the restriction of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $\text{Term}(\text{ExtraPSGCl})$, is collapsed.

Proof. Since $(u_0, z_0, y_0, w_0, p_0, a)$ factors through CollapseExtraMRD , we know that it is collapsed. Since $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ are specializations of $\text{ExtraPSGCl}(u, z, y, w, p, a)$ that restrict to the same modular family of $\text{Term}(\text{ExtraPSGCl})$, the two families of representatives that they induce, represent the same modular families of the group G_i , for all $i = 1, \dots, s$. Hence, $(u'_0, z_0, y_0, w_0, p_0, a)$ is collapsed. \square

Lemma 7.11. *Let $\text{ExtraPSGCl}(u, z, y, w, p, a)$ be a terminal graded closure in ExtraPSMRD .*

Let CollapseExtraMRD be the CollapseExtra MR diagram of the collection $WPRS$ and the graded closure $\text{ExtraPSGCl}(u, z, y, w, p, a)$.

Let Γ_l be a group of level l in the model. If l is large enough, the the following property is satisfied in Γ_l .

Let $\text{CollapseExtraGFCl}(c, u, z, y, w, p, a)$ be a terminal graded formal closure in CollapseExtraMRD , and let $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ be a Γ_l -specialization of $\text{CollapseExtraGFCl}(c, u, z, y, w, p, a)$.

Then, every Γ_l -specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ of $\text{ExtraPSGCl}(u, z, y, w, p, a)$, that restricts to a solution of $\text{Term}(\text{ExtraPSGCl})$ that belongs to the same modular family of the restriction of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $\text{Term}(\text{ExtraPSGCl})$, is collapsed.

Proof. The same proof as the case of F_k . \square

8 Approximations of EAE -Sets under Minimal Rank Assumption

Let $WPRS$ be a given collection of wp -rigid and wp -solid limit groups $G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$ (wp is the parameter set). For all $i = 1, \dots, s$, we fix an integer $e_i \in \{0, \dots, M(G_i)\}$, where $M(G_i)$ is the maximal number of exceptional families of specializations in G_i for a fixed value of the parameters. We consider the graded completion $\text{Comp}(z, y, w, p, a)$ of some wp -graded resolution $\text{Res}(y, w, p, a)$, and we denote the terminal p -rigid or p -solid base group of $\text{Comp}(z, y, w, p, a)$ by B . We assume that the amalgamated free product of the groups in the collection $WPRS$

$$*_{i=1}^s \langle w, p, a \rangle * \langle w, p, a \rangle^{e_i} G_i$$

amalgamated over the subgroup $\langle w, p, a \rangle$, is mapped into each of the levels of $\text{Comp}(z, y, w, p, a)$ (in a way that respects the canonical maps between those levels). We assume further that $\text{Comp}(z, y, w, p, a)$ admits some specialization (z_0, y_0, w_0, p_0, a) so that the restriction of (y_0, w_0, p_0, a) to $* \langle w, p, a \rangle^{e_i} G_i$ represent a family of representatives of pairwise distinct exceptional families of G_i corresponding to the value $w_0 p_0$ of the parameters, for all $i = 1, \dots, s$.

Let $\text{AbsSys}(c, z, y, w, p, a)$ be a finite collection of systems of equations. We add to the systems in $\text{AbsSys}(c, z, y, w, p, a)$, the finitely many systems stating that the specialization (z, y, w, p, a) of $\text{Comp}(z, y, w, p, a)$ is collapsed, i.e., that its restriction to one of the groups $G_1(x_1, w, p, a), \dots, G_s(x_s, w, p, a)$ that are mapped into $\text{Comp}(z, y, w, p, a)$ is not exceptional.

We construct the p -graded formal MR diagram of the systems $\text{AbsSys}(c, z, y, w, p, a)$ w.r.t. the resolution $\text{Comp}(z, y, w, p, a)$. We denote this diagram by PosGFMRD . We also construct the ExtraPS MR diagram ExtraPSMRD of the collection $WPRS$ over the resolution $\text{Com}(z, y, w, p, a)$.

Definition 8.1. Let (z_0, y_0, w_0, p_0, a) be a specialization of $\text{Comp}(z, y, w, p, a)$ (constructed with respect to some proof system PS). We say that (z_0, y_0, w_0, p_0, a) is an *abstract validPS statement*, if the following conditions hold:

1. The specialization (z_0, y_0, w_0, p_0, a) is non-collapsed.
2. For all $i = 1, \dots, s$, the family of representatives of the exceptional families of the wp -rigid or wp -solid group G_i , that the specialization (z_0, y_0, w_0, p_0, a) induces, forms a full family w.r.t. the value $w_0 p_0$ of the parameters.
3. There exists no $x_0 \in F_k$, so that the specialization $(x_0, z_0, y_0, w_0, p_0, a)$ solves each of the systems in $AbsSys(x, z, y, w, p, a)$.

The set of all the values $p_0 \in F_k$, for which there exists an abstract validPS statement (z_0, y_0, w_0, p_0, a) , is denoted by $EAE(p)$.

Definition 8.2. Let Γ be a group in the model. Let (z_0, y_0, w_0, p_0, a) be a specialization of $Comp(z, y, w, p, a)$ (constructed with respect to some proof system PS). We say that (z_0, y_0, w_0, p_0, a) is an *abstract Γ -validPS statement*, if the following conditions hold:

1. The Γ -specialization (z_0, y_0, w_0, p_0, a) is non-collapsed.
2. For all $i = 1, \dots, s$, the family of representatives of the Γ -exceptional families of the wp -rigid or wp -solid group G_i , that the Γ -specialization (z_0, y_0, w_0, p_0, a) induces, forms a full family w.r.t. the value $w_0 p_0$ of the parameters.
3. There exists no $x_0 \in \Gamma$, so that the specialization $(x_0, z_0, y_0, w_0, p_0, a)$ solves each of the systems in $AbsSys(x, z, y, w, p, a)$.

The set of all the values $p_0 \in \Gamma$, for which there exists an abstract Γ -validPS statement (z_0, y_0, w_0, p_0, a) , is denoted by $EAE_\Gamma(p)$.

Our aim in this section, is to reduce the sets $EAE(p)$ and $EAE_\Gamma(p)$ uniformly, for almost all the groups Γ , into the Boolean algebra of AE -sets. That is, to construct a formula $\varphi(p)$ in the Boolean algebra of AE -formulas, that defines each of these sets over the corresponding groups. We achieve this aim using a process that terminates in finitely many steps. In each step, we define a formula in the Boolean algebra of AE -formulas that approximates our sets $EAE(p)$ and $EAE_\Gamma(p)$ uniformly, on one hand. And on the other hand, every specialization $p = p_0$ in $EAE(p)$ or in $EAE_\Gamma(p)$, must be collected (over the corresponding group) by a formula defined in some step along the process that we are going to explain below.

8.1 First Approximation of $EAE(p)$

Theorem 8.3. *Let $p_0 \in F_k$. Assume that there exists a specialization (z_0, y_0, w_0, p_0, a) that factors through $Comp(z, y, w, p, a)$, and satisfies the following conditions:*

1. (z_0, y_0, w_0, p_0, a) restricts to an exceptional specialization (b_0, p_0, a) of B .
2. (z_0, y_0, w_0, p_0, a) does not factor through $PosGFMRD$.
3. For every terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in $ExtraPSMRD$, for every $u_0 \in F_k$, if $(u_0, z_0, y_0, w_0, p_0, a)$ factors through $ExtraPSGCl(u, z, y, w, p, a)$, then there exists some $u'_0 \in F_k$, so that:
 - (a) The specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ restrict to the same exceptional family of $Term(ExtraPSGCl)$.
 - (b) The specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ factors through the collapse-extra MR diagram associated with the graded closure $ExtraPSGCl(u, z, y, w, p, a)$.

Then $p_0 \in EAE(p)$. The set of all the specializations $p = p_0$ of the parameters, for which there exists some specialization (z_0, y_0, w_0, p_0, a) who satisfies the conditions in the theorem, is denoted by $TSPS(p)$.

Proof. We consider the ungraded completion $Comp(z, y, w, p_0, a)$ induced by (z_0, y_0, w_0, p_0, a) .

For every terminal graded formal closure $PosGFCl(c, z, y, w, p, a)$, in $PosGFMRD$, we consider its p -rigid or p -solid base group $Term(PosGFCl)$. If $Term(PosGFCl)$ admits an exceptional solution h_{p_0} , so that the restriction of h_{p_0} to B is an exceptional solution of B that belongs to the same exceptional family of (b_0, p_0, a) , we list the ungraded closure $PosGFCl_{h_{p_0}}$ induced by (some exceptional solution in the exceptional family of) h_{p_0} . The collection of all such (boundedly many) ungraded closures, is denoted by Pos_{p_0} .

For every terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$, in $ExtraPSMRD$, we consider its p -rigid or p -solid base group $Term(ExtraPSGCl)$. If $Term(ExtraPSGCl)$ admits an exceptional solution h_{p_0} , so that the restriction of h_{p_0} to B is an exceptional solution of B that belongs to the same exceptional family of (b_0, p_0, a) , we list the ungraded closure $ExtraPSCl_{h_{p_0}}$ induced by (some exceptional solution in the exceptional family of) h_{p_0} . The collection of all such (boundedly many) ungraded closures, is denoted by $ExtraPS_{p_0}$. We denote the subcollection of $ExtraPS_{p_0}$ consisting of the ungraded closures through which the specialization (z_0, y_0, w_0, p_0, a) factors, for some u_0 , by $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$.

By the assumption, and according to the construction of the collapse-extra MR diagram, for every ungraded closure $ExtraPSCl_{h_{p_0}}$ in $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$, who is covered by the graded closure $ExtraPSGCl(u, z, y, w, p, a)$, there exists a specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ of some terminal graded formal closure $CollapseExtraGFCl(c, u, z, y, w, p, a)$ in the collapse-extra MR diagram $CollapseExtraMRD$ associated with $ExtraPSGCl(u, z, y, w, p, a)$, so that the restriction of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to B belongs to the exceptional family represented by (b_0, p_0, a) , the restriction of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $Term(ExtraPSGCl)$ belongs to the exceptional family represented by h_{p_0} , and the restriction of $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ to $Term(CollapseExtraGFCl)$ is some exceptional solution which we denote by g_{p_0} . We choose such a terminal graded formal closure $CollapseExtraGFCl(c, u, z, y, w, p, a)$ and a specialization $(c_0, u_0, z_0, y_0, w_0, p_0, a)$ of it, and denote the induced ungraded closure by $CollapseExtraFCl_{g_{p_0}}$. For every $ExtraPSCl_{h_{p_0}}$ in $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$, we have associated an ungraded closure $CollapseExtraFCl_{g_{p_0}}$. The collection of all the (boundedly many) ungraded closures $CollapseExtraFCl_{g_{p_0}}$ that we obtain, is denoted briefly by $CollapseExtra_{p_0}$.

Let $ExtraPSCl_{h_{p_0}}$ be an ungraded closure in $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$ who is covered by the graded closure $ExtraPSGCl(u, z, y, w, p, a)$, and let $CollapseExtraFCl_{g_{p_0}}$ be the ungraded closure in $CollapseExtra_{p_0}$ associated with $ExtraPSCl_{h_{p_0}}$. Then, according to the construction of the collapse extra MR diagram, every specialization $(u'_0, z'_0, y'_0, w'_0, p_0, a)$ that factors through $CollapseExtraFCl_{g_{p_0}}$, restricts to an exceptional specialization of $Term(ExtraPSGCl)$ that belongs to the same exceptional family of h_{p_0} . Hence, according to lemma 7.10, if $(z'_0, y'_0, w'_0, p_0, a)$ is a specialization that factors through $CollapseExtraFCl_{g_{p_0}}$, then, for every $u''_0 \in F_k$, if $(u''_0, z'_0, y'_0, w'_0, p_0, a)$ is a specialization of $ExtraPSGCl(u, z, y, w, p, a)$ that restricts to an exceptional family of $Term(ExtraPSGCl)$ that belongs to the same exceptional family of h_{p_0} , then $(u''_0, z'_0, y'_0, w'_0, p_0, a)$ is collapsed.

We consider now the collection of ungraded closures $CollapseExtra_{p_0}$, in one hand, and on the other hand, we consider the collection of ungraded closures Pos_{p_0} and $ExtraPS_{p_0} \setminus ExtraPS_{(z_0, y_0, w_0, p_0, a)}$. Each of these closures, is in particular an ungraded closure of the ungraded completion $Comp(z, y, w, p_0, a)$. However, the specialization (z_0, y_0, w_0, p_0, a) of $Comp(z, y, w, p_0, a)$, factors through each of the closures in $CollapseExtra_{p_0}$, but it does not factor through any of the closures in Pos_{p_0} and $ExtraPS_{p_0} \setminus ExtraPS_{(z_0, y_0, w_0, p_0, a)}$. Thus, there exists a test sequence of specializations (z_n, y_n, w_n, p_0, a) of $Comp(z, y, w, p_0, a)$, so that for all n , the specialization (z_n, y_n, w_n, p_0, a) factors through each of the closures in $CollapseExtra_{p_0}$, and does not factor through any of the closures in Pos_{p_0} and $ExtraPS_{p_0} \setminus ExtraPS_{(z_0, y_0, w_0, p_0, a)}$. We are going to prove that (z_n, y_n, w_n, p_0, a) is eventually a validPS statement.

By the construction of the sequence (z_n, y_n, w_n, p_0, a) , and since, for all n , the specialization (z_n, y_n, w_n, p_0, a) restricts to an exceptional solution of B that belongs to the same exceptional family of (b_0, p_0, a) , we have that for all n , the specialization (z_n, y_n, w_n, p_0, a) does not factor through the diagram $PosGFMRD$, nor through any of the terminal graded closures $ExtraPSGCl(u, z, y, w, p, a)$ in $ExtraPSMRD$ that does not cover some closure in $ExtraPS_{p_0}$, and if it factors through an ungraded closure $ExtraPSCl_{h_{p_0}}$ in $ExtraPS_{p_0}$, then $ExtraPSCl_{h_{p_0}}$ belongs to $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$.

By the constructions of $PosGFMRD$ diagram, this implies that the specialization (z_n, y_n, w_n, p_0, a) includes (eventually) only exceptional solutions of the corresponding wp -rigid and wp -solid limit groups that are mapped into $Comp(z, y, w, p, a)$, i.e., the specialization (z_n, y_n, w_n, p_0, a) is non-collapsed.

Assume by contradiction that for a subsequence n of the integers, the sequence (z_n, y_n, w_n, p_0, a) con-

sists of non-validPS statements. Let $u_n \in F_k$ be a sequence of specializations testifying this property, i.e., $(u_n, z_n, y_n, w_n, p_0, a)$ is non-collapsed, for all n . In particular, according to theorem 6.6, there exists some terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in the diagram $ExtraPSMRD$, so that for all n in a subsequence of the integers, there exists $u'_n \in F_k$, so that $(u'_n, z_n, y_n, w_n, p_0, a)$ is a specialization of $ExtraPSGCl(u, z, y, w, p, a)$, and:

1. The specialization $(u'_n, z_n, y_n, w_n, p_0, a)$ restricts to an exceptional specialization of B that belongs to the same exceptional family of (b_0, p_0, a) .
2. The specialization $(u'_n, z_n, y_n, w_n, p_0, a)$ restricts to an exceptional specialization $h_{p_0}^n$ of $Term(ExtraPSGCl)$.
3. The specialization $(u'_n, z_n, y_n, w_n, p_0, a)$ is a non-collapsed.

Denote by $ExtraPSCl_{h_{p_0}^n}$ the ungraded closure obtained by specializing the base group $Term(ExtraPSGCl)$ of $ExtraPSGCl(u, z, y, w, p, a)$ according to the restriction $h_{p_0}^n$ of $(u'_n, z_n, y_n, w_n, p_0, a)$ to it. By the construction of the test sequence (z_n, y_n, w_n, p_0, a) , the specialization (z_n, y_n, w_n, p_0, a) does not factor through any of the closures $ExtraPS_{p_0} \setminus ExtraPS_{(z_0, y_0, w_0, p_0, a)}$. Thus, the ungraded closure $ExtraPSCl_{h_{p_0}^n}$ must be represented by some ungraded closure $ExtraPSCl_{h_{p_0}}$, that is covered by the graded closure $ExtraPSGCl(u, z, y, w, p, a)$, and belongs to $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$. That is, $h_{p_0}^n$ and h_{p_0} belong to the same exceptional family of $Term(ExtraPSGCl)$, for all n .

However, by construction, the specialization (z_n, y_n, w_n, p_0, a) factors through the ungraded closure $CollapseExtraFCl_{g_{p_0}}$ associated with $ExtraPSCl_{h_{p_0}}$, so $(u'_n, z_n, y_n, w_n, p_0, a)$ must be collapsed, for all n , a contradiction. \square

Theorem 8.4. *Let Γ be a group of level l in the model. If l is large enough, then the following property is satisfied for Γ .*

Let $p_0 \in \Gamma$. Assume that there exists a Γ -specialization (z_0, y_0, w_0, p_0, a) that factors through $Comp(z, y, w, p, a)$, and satisfies the following conditions:

1. (z_0, y_0, w_0, p_0, a) restricts to a Γ -exceptional specialization (b_0, p_0, a) of B .
2. (z_0, y_0, w_0, p_0, a) does not factor through $PosGFMRD$.
3. *For every terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in $ExtraPSMRD$, for every $u_0 \in \Gamma$, if $(u_0, z_0, y_0, w_0, p_0, a)$ factors through $ExtraPSGCl(u, z, y, w, p, a)$, then there exists some $u'_0 \in \Gamma$, so that:*
 - (a) *The Γ -specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ restrict to the same Γ -exceptional family of $Term(ExtraPSGCl)$.*
 - (b) *The Γ -specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ factors through the collapse-extra MR diagram associated with the graded closure $ExtraPSGCl(u, z, y, w, p, a)$.*

Then $p_0 \in EAE_\Gamma(p)$. The set of all the Γ -specializations $p = p_0$ of the parameters, for which there exists some specialization (z_0, y_0, w_0, p_0, a) who satisfies the conditions in the theorem, is denoted by $TSPS_\Gamma(p)$.

Proof. In a similar way of the proof of theorem 8.3, we consider the ungraded completion $Comp(z, y, w, p_0, a)$ induced by (z_0, y_0, w_0, p_0, a) .

We list, in a collection denoted Pos_{p_0} , the (boundedly many) ungraded closures $PosGFCl_{h_{p_0}}$ who are covered by terminal graded formal closures $PosGFCl(c, z, y, w, p, a)$ in $PosGFMRD$, so that h_{p_0} is a Γ -exceptional solution of $Term(PosGFCl)$ whose restriction to B belongs to the same Γ -exceptional family of (b_0, p_0, a) .

We list, in a collection denoted $ExtraPS_{p_0}$, the (boundedly many) ungraded closures $ExtraPSCl_{h_{p_0}}$ who are covered by terminal graded closures $ExtraPSGCl(u, z, y, w, p, a)$ in $ExtraPSMRD$, so that h_{p_0} is a Γ -exceptional solution of $Term(ExtraPSGCl)$ whose restriction to B belongs to the same Γ -exceptional family of (b_0, p_0, a) . The subcollection of $ExtraPS_{p_0}$ consisting of the ungraded closures through which the specialization (z_0, y_0, w_0, p_0, a) factors, is denoted by $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$.

By the assumption, and according to the construction of the collapse-extra MR diagram, lemma 7.6 and lemma 7.11, for every ungraded closure $ExtraPSCl_{h_{p_0}}$ in $ExtraPS_{(z_0, y_0, w_0, p_0, a)}$, who is covered by the

graded closure $ExtraPSGCl(u, z, y, w, p, a)$, we can associate an ungraded closure $CollapseExtraFCl_{g_{p_0}}$, that is induced from some Γ -exceptional solution g_{p_0} of the base p -rigid or p -solid group of some graded formal closure $CollapseExtraGFCl(c, u, z, y, w, p, a)$ in the collapse-extra MR diagram $CollapseExtraMRD$ associated with $ExtraPSGCl(u, z, y, w, p, a)$, so that the restriction of g_{p_0} to B is a Γ -exceptional solution of B that belongs to the Γ -exceptional family of (b_0, p_0, a) , and that for every Γ -specialization $(z'_0, y'_0, w'_0, p_0, a)$ that factors through $CollapseExtraFCl_{g_{p_0}}$, and for every $u''_0 \in \Gamma$, if $(u''_0, z'_0, y'_0, w'_0, p_0, a)$ is a specialization of $ExtraPSGCl(u, z, y, w, p, a)$ that restricts to an exceptional family of $Term(ExtraPSGCl)$ that belongs to the exceptional family of h_{p_0} , then $(u''_0, z'_0, y'_0, w'_0, p_0, a)$ is collapsed. The collection of all such (boundedly many) associated ungraded closures $CollapseExtraFCl_{g_{p_0}}$, is denoted briefly by $CollapseExtra_{p_0}$.

Now since the Γ -specialization (z_0, y_0, w_0, p_0, a) of $Comp(z, y, w, p_0, a)$, factors through each of the ungraded closures in ungraded closures $CollapseExtra_{p_0}$, but does not factor through any of the ungraded closures in Pos_{p_0} and $ExtraPS_{p_0} \setminus ExtraPS_{(z_0, y_0, w_0, p_0, a)}$, then, there must exist a Γ -test sequence (z_n, y_n, w_n, p_0, a) of $Comp(z, y, w, p_0, a)$, so that for all n , the Γ -specialization (z_n, y_n, w_n, p_0, a) factors through each of the closures in $CollapseExtra_{p_0}$, and does not factor through any of the closures in Pos_{p_0} and $ExtraPS_{p_0} \setminus ExtraPS_{(z_0, y_0, w_0, p_0, a)}$. We are going to prove that (z_n, y_n, w_n, p_0, a) is eventually a Γ -validPS statement.

According to theorems 5.10 and 6.8, the remaining part of the proof is identical to the case over F_k presented in theorem 8.3. \square

Theorem 8.5. *Let $p_0 \in F_k$. There exists a formula $\varphi(p)$ that belongs to the Boolean algebra of AE-formulas, so that:*

1. *The formula $\varphi(p)$ defines the set $TSPS(p)$.*
2. *Let Γ be a group of level l in the model. If l is large enough, then the formula $\varphi(p)$ defines the set $TSPS_\Gamma(p)$.*

Proof. We show how to construct $\varphi(p)$, and explain the reason that it defines the sets $TSPS(p)$ and $TSPS_\Gamma(p)$.

Let (z_0, y_0, w_0, p_0, a) be a $[\Gamma]$ -specialization that factors through $Comp(z, y, w, p, a)$.

The second condition of theorem 8.3 [theorem 8.4], is equivalent to that the $[\Gamma]$ -specialization (z_0, y_0, w_0, p_0, a) does not factor through any of the terminal graded formal closures of the diagrams $NonRgdMRD$ and $NonSldMRD$.

The third condition of theorem 8.3[theorem 8.4], is equivalent to the following: for every graded terminal closure

$ExtraPSCl(u, z, y, w, p, a)$ of $ExtraMRD$, there exists a $[\Gamma]$ -full family of representatives h_1, \dots, h_r , for the $[\Gamma]$ -exceptional families of its base group $Term(ExtraPSCl)$ corresponding to $p = p_0$, so that, for all $i = 1, \dots, r$, if the $[\Gamma]$ -specialization (z_0, y_0, w_0, p_0, a) factors through the ungraded closure covered by $ExtraPSCl(u, z, y, w, p, a)$ and induced from h_i , and u_0^i is the corresponding induced $[\Gamma]$ -specialization of the variables u , then the $[\Gamma]$ -specialization $(u_0^i, z_0, y_0, w_0, p_0, a)$ factors through some terminal graded formal closure in $CollapseExtraMRD$ associated with $ExtraPSCl(u, z, y, w, p, a)$.

For simplicity, we denote by E_1, \dots, E_r all the terminal graded closures in $ExtraPSMRD$, and by C_1, \dots, C_s all the graded formal terminal closures in $CollapseExtraMRD$. We denote also the terminal graded formal closures of $PosGFMRD$ by S_1, \dots, S_q .

Hence, given any $p = p_0 \in F_k$, if for every terminal graded closure E_i in $ExtraMRD$, the exact number n_i of exceptional families of solutions of its p -rigid or p -solid base group corresponding to the value $p = p_0$ of the parameters is known, then there will exist a specialization (z_0, y_0, w_0, p_0, a) of $Comp(z, y, w, p, a)$ that satisfies the conditions in theorem 8.3, if and only if the following sentence is true:

$$\begin{aligned}
& \exists z_0, y_0, w_0, \bigvee_{i=1}^r h_1^i, \dots, h_{n_i}^i \bigvee_{i=1}^r c_1^i, \dots, c_{n_i}^i, \quad \text{s.t.} \\
& [(z_0, y_0, w_0, p_0, a) \text{ restricts to an exceptional solution for } B], \text{ and} \\
& \left[\bigwedge_{i=1}^q \forall c_0 S_i(c_0, z_0, y_0, w_0, p_0, a) \neq 1 \right], \text{ and} \\
& \left[\bigwedge_{i=1}^r \bigwedge_{j=1}^{n_i} h_j^i \text{ is an exceptional solution for } \text{Term}(E_i) \right], \text{ and} \\
& \left[\bigwedge_{i=1}^r \bigwedge_{j \neq j'} h_j^i \text{ and } h_{j'}^i \text{ represent distinct exceptional families of } \text{Term}(E_i) \right], \text{ and} \\
& \left[\bigwedge_{i=1}^r \bigwedge_{j=1}^{n_i} \forall u_0 \left[E_{h_j^i}(u_0, z_0, y_0, w_0, p_0, a) \implies \bigvee_{k=1}^s C_k(c_j^i, u_0, z_0, y_0, w_0, p_0, a) \right] \right].
\end{aligned}$$

Since the numbers n_i admits a uniform upper bound for all the values $p = p_0 \in F_k$ [$p = p_0 \in \Gamma$], and since the values $p = p_0$ for which a given p -rigid or p -solid limit group admits exactly a given number of $[\Gamma]$ -exceptional families corresponding to $p = p_0$ form a set that belongs to the Boolean algebra of AE -sets, a finite disjunction that runs on all the possible values of the numbers n_i , and for each possibility it lists the corresponding formula similar to the aforementioned one, defines the desired formula $\varphi(p)$. \square

8.2 Reducing the Set $EAE(p)$ under Minimal-Rank Assumption

Now we explain how theorem 8.5 can be applied iteratively, in order to collect all the values $p_0 \in F_k$ [resp. $p_0 \in \Gamma$], for which there exist some validPS statement [resp. Γ -validPS statement] that factors through the completed resolution $Comp(z, y, w, p, a)$. For that aim, we use the method of constructing quotient resolutions presented in [Se4].

Theorem 8.6. *Let $Rlim(y, p, a)$ be a minimal rank p -graded limit group, and let $Res(y, p, a)$ be a graded resolution of $Rlim(y, p, a)$. Let $Comp(z, y, p, a)$ be the graded completion of $Res(y, p, a)$, and denote its base p -rigid or p -solid group by B . Let $GCl(u, z, y, p, a)$ be a graded closure for $Comp(z, y, p, a)$. Let $\Sigma = \{\Sigma_i(c, u, z, y, p, a)\}_{i=1, \dots, r}$ be a finite collection of systems. Let \mathcal{A} be the collection of all the specializations (u_0, z_0, y_0, p_0, a) for which:*

1. *The specialization (z_0, y_0, p_0, a) factors through $GCl(u, z, y, p, a)$.*
2. *The specialization (u_0, z_0, y_0, p_0, a) restricts to an exceptional specialization of $\text{Term}(GCl)$ and to an exceptional specialization of B .*
3. *There exists some $c_0 \in F_k$, so that $\Sigma_i(c_0, u_0, z_0, y_0, p_0, a)$ for some $i = 1, \dots, r$.*

Then, there exists finitely many p -graded completed resolutions $Comp_1^2(t, c, u, z, y, p, a), \dots, Comp_s^2(t, c, u, z, y, p, a)$, so that for all $j = 1, \dots, s$:

1. *The graded closure $GCl(u, z, y, p, a)$ is mapped canonically into $Comp_j^2(t, c, u, z, y, p, a)$.*
2. *The words of the system $\Sigma_i(c, u, z, y, p, a)$, for some $i = 1, \dots, r$, represent the trivial element in $Comp_j^2(t, c, u, z, y, p, a)$.*
3. *The image of $\text{Term}(GCl)$ is contained in the base group $\text{Term}(Comp_j^2)$, and is mapped compatibly, i.e., the minimal subgraph of $\text{Term}(Comp_j^2)$ containing the image of $\text{Term}(GCl)$, in the p -JSJ of $\text{Term}(Comp_j^2)$, consists only from parts of the p -JSJ of $\text{Term}(GCl)$.*
4. *The complexity of $Comp_j^2(t, c, u, z, y, p, a)$ is not larger than the complexity of $Comp(z, y, p, a)$ (definition 1.16 in [Se5]).*
5. *If the complexity of $Comp_j^2(t, c, u, z, y, p, a)$ equals the complexity of $Comp(z, y, p, a)$, then $Comp_j^2(t, c, u, z, y, p, a)$ is a graded formal closure $GFCl(c, u, z, y, p, a)$ of $Comp(z, y, p, a)$.*

Moreover, for every specialization (u_0, z_0, y_0, p_0, a) in the collection \mathcal{A} , there exists a specialization (u'_0, z_0, y_0, p_0, a) that belongs to \mathcal{A} , so that (u'_0, z_0, y_0, p_0, a) factors through one of the resolutions

$$\text{Comp}_1^2(t, c, u, z, y, p, a), \dots, \text{Comp}_s^2(t, c, u, z, y, p, a),$$

and the specializations (u_0, z_0, y_0, p_0, a) and (u'_0, z_0, y_0, p_0, a) restrict to the same exceptional family of $\text{Term}(GCl)$.

We call the resolutions $\text{Comp}_1^2, \dots, \text{Comp}_s^2$, the quotient resolutions of the systems Σ w.r.t. the graded closure $GCl(u, z, y, p, a)$.

Proof. The collection \mathcal{A} factors through finitely many maximal limit groups $QRlim(c, u, z, y, p, a)$. Since $Rlim(y, p, a)$ is assumed to be a minimal rank graded limit group, if $QRlim(c, u, z, y, p, a)$ is p -freely decomposable, then the factor containing the subgroup $\langle a, p \rangle$ must contain the whole image of the graded completion $\text{Comp}(z, y, p, a)$. Since the specializations in \mathcal{A} restrict to exceptional specializations for $\text{Term}(GCl)$, we conclude that the whole image of the graded closure $GCl(u, z, y, p, a)$ is contained in the free factor of $QRlim(c, u, z, y, p, a)$ containing the subgroup $\langle a, p \rangle$, in the case that $QRlim(c, u, z, y, p, a)$ is p -freely decomposable. Hence, By mapping the factors that does not intersect a conjugate of the subgroup $\langle a, p \rangle$ to the trivial element, we may assume that $QRlim(c, u, z, y, p, a)$ is p -freely indecomposable.

Now we denote by zup_b^1 , a set of generators for the second level of the graded closure $GCl(u, z, y, p, a)$. We recognize the limit group $QRlim(c, u, z, y, p, a)$ as a zup_b^1 -graded limit group $QRlim(c, u, z, y, p, a) = QRlim(c, u, z, y, zup_b^1, a)$. The subcollection of specializations in \mathcal{A} that factor through $QRlim$, induces a zup_b^1 -graded MR diagram of the zup_b^1 -graded limit group $QRlim$. Let $QRes(c, u, z, y, zup_b^1, a)$ be a resolution in this MR diagram.

Let Q_1, \dots, Q_t be some surviving surfaces (definitions 1.7 and 1.10 in [Se4]) that appear in the p -JSJ of the first level of the given resolution $Res(y, p, a)$. That is, Q_i is mapped monomorphically onto a subgroup of finite index of a QH subgroup of the j -th level zup_b^1 -graded limit group $Glim_j(c, u, z, y, zup_b^1, a)$, for some j . If Q is a surviving surface, and j is the minimal integer for which Q is mapped monomorphically onto a subgroup of finite index of a QH subgroup of the j -th level zup_b^1 -graded limit group $Glim_j(c, u, z, y, zup_b^1, a)$, we denote by Q' its isomorphic image in $Glim_j$.

Let j be an integer and assume that all the images Q'_1, \dots, Q'_t appear in the j -th level of the zup_b^1 -graded limit group $Glim_j(c, u, z, y, zup_b^1, a)$. Assume further that for all $i = 1, \dots, t$, there is no surviving surface Q , other than Q_i , that is mapped non-trivially into a conjugate of Q'_i , and that j is the minimal integer for which the group $Glim_j$ contains the subgroup Q' for some surviving surface Q .

According to the proof of theorem 1.7 in [Se4], we can isolate the images of the surviving surfaces Q_1, \dots, Q_t along the subresolution $QRes_j$, consisting of the first j levels of $QRes$. We map the isolated QH subgroups Q'_1, \dots, Q'_t into the distinguished vertex containing zup_b^1 , and continue iteratively.

We finish up with finitely many zup_b^1 -graded resolutions, denoted again by $QRes(c, u, z, y, zup_b^1, a)$, in which all the surviving surfaces are isolated. Each of the terminal groups of the new resolutions is either zup_b^1 -rigid or zup_b^1 -solid, and these resolutions cover all the specializations in the collection \mathcal{A} .

We continue by considering the subcollection \mathcal{A}_1 of \mathcal{A} that consists of the specializations (u_0, z_0, y_0, p_0, a) for which there exists some $c_0 \in F_k$, so that $(c_0, u_0, z_0, y_0, p_0, a)$ is a zup_b^1 -exceptional specialization of $T_1(c, u, z, y, zup_b^1, a)$, the terminal group of $QRes(c, u, z, y, zup_b^1, a)$.

The specializations in the collection \mathcal{A}_1 factors through finitely many maximal limit groups $QRlim^2$. We denote by zup_b^2 , a set of generators for the second level of the graded closure $GCl(u, z, y, p, a)$. We recognize each of the maximal limit groups $QRlim^2$ as a zup_b^2 -graded limit group. For each resolution in the zup_b^2 -graded MR diagram of $QRlim^2$, we use a similar construction to the one described above, in order to isolate all the surviving surfaces that appear in the second level of the graded closure $GCl(u, z, y, p, a)$. And we continue iteratively.

This process terminates in finitely many resolutions denoted again by $QRes(c, u, z, y, p, a)$. Each of the resolutions terminate in p -rigid or p -solid group, and these resolutions cover all the specializations in the collection \mathcal{A} . Moreover, according to the construction, and using Propositions 1.11 and 1.12 of [Se4], all the surviving surfaces w.r.t. some obtained resolution $QRes(c, u, z, y, p, a)$, in all the levels of the given resolution $Res(y, p, a)$, are isolated along the resolution $QRes(c, u, z, y, p, a)$.

We consider the completion Comp^2 of some obtained resolution $QRes(c, u, z, y, zup_b^1, a)$. We are going now to modify the completed resolution Comp^2 . For each of the abelian vertex groups in the decomposition Λ_1 associated with the top level of Comp^2 , we fix a basis. We map the QH subgroups that appear in Λ_1

and does not intersect a conjugate of the image of GCl non-trivially, into the next level of $Comp^2$. For each generator of an abelian vertex in Λ_1 , if the cyclic subgroup generated by this generator, does not intersect a conjugate of the image of GCl non-trivially, we map it to the next level of $Comp^2$.

We continue modifying the various levels of $Comp^2$ successively, until we reach the terminal group of $Comp^2$, that we do not change. According to the analysis of quotient resolutions in the minimal rank case, that appear in Lemmas 1.3, 1.5, 1.11, and 1.12 in [Se4], the quadratic complexity of the (sub)resolution (of) $Comp^2$, is no larger than that of the given resolution $Comp$. In the case that of equality, we can modify the tower structure of $Comp^2$ by “dropping” some levels that their decompositions do not contain any isolated surviving surface, and by “merging” remaining levels, so that the quadratic complexity do not change, and the abelian rank of the obtained resolution is no larger than the abelian rank of the given resolution $Comp$. In the case that of equality in the abelian ranks too, the obtained resolution $Comp^2$ will be a graded formal closure of $Comp$. \square

Lemma 8.7. *Let $Rlim(y, p, a)$ be a minimal rank limit p -graded limit group, and let $Res(y, p, a)$ be a graded resolution of $Rlim(y, p, a)$. Let $Comp(z, y, p, a)$ be the graded completion of $Res(y, p, a)$, and denote its base p -rigid or p -solid group by B . Let $GCl(u, z, y, p, a)$ be a graded closure for $Comp(z, y, p, a)$. Let $\Sigma = \{\Sigma_i(c, u, z, y, p, a)\}_{i=1, \dots, r}$ be a finite collection of systems. Let \mathcal{A} be the collection of all the specializations (u_0, z_0, y_0, p_0, a) for which:*

1. *The specialization (u_0, z_0, y_0, p_0, a) factors through $GCl(u, z, y, p, a)$.*
2. *The specialization (u_0, z_0, y_0, p_0, a) restricts to an exceptional specialization of $Term(GCl)$ and to an exceptional specialization of B .*
3. *There exists some $c_0 \in F_k$, so that $\Sigma_i(c_0, u_0, z_0, y_0, p_0, a)$ for some $i = 1, \dots, r$.*

Let $Comp_1^2(t, c, u, z, y, p, a), \dots, Comp_s^2(t, c, u, z, y, p, a)$ be the p -graded completed resolutions from 8.6.

Let Γ be a group in the model. Then, the following property is satisfied for Γ . Let $\Gamma\mathcal{A}$ be the collection of all the Γ -specializations (u_0, z_0, y_0, p_0, a) for which:

1. *The Γ -specialization (z_0, y_0, p_0, a) factors through $GCl(u, z, y, p, a)$.*
2. *The Γ -specialization (u_0, z_0, y_0, p_0, a) restricts to an exceptional specialization of $Term(GCl)$ and to an exceptional specialization of B .*
3. *There exists some $c_0 \in \Gamma$, so that $\Sigma_i(c_0, u_0, z_0, y_0, p_0, a)$ for some $i = 1, \dots, r$.*

Then, the resolutions $Comp_1^2(t, c, u, z, y, p, a), \dots, Comp_s^2(t, c, u, z, y, p, a)$ cover the collection $\Gamma\mathcal{A}$.

For every Γ -specialization (u_0, z_0, y_0, p_0, a) in the collection $\Gamma\mathcal{A}$, there exists a Γ -specialization (u'_0, z_0, y_0, p_0, a) that belongs to $\Gamma\mathcal{A}$, so that (u'_0, z_0, y_0, p_0, a) factors through one of the resolutions

$$Comp_1^2(t, c, u, z, y, p, a), \dots, Comp_s^2(t, c, u, z, y, p, a),$$

and the Γ -specializations (u_0, z_0, y_0, p_0, a) and (u'_0, z_0, y_0, p_0, a) restrict to the same Γ -exceptional family of $Term(GCl)$.

Proof. Let (u_0, z_0, y_0, p_0, a) be a Γ -specialization in $\Gamma\mathcal{A}$, and let $c_0 \in \Gamma$ be so that $\Sigma_i(c_0, u_0, z_0, y_0, p_0, a)$ for some $i = 1, \dots, r$. We lift $(c_0, u_0, z_0, y_0, p_0, a)$ to a specialization $(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ in F_k , so that $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ is a specialization of GCl , and $\Sigma_i(\tilde{c}_0, \tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a) = 1$.

Hence, the specialization $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ belongs to the collection \mathcal{A} , and according to theorem 8.6, there exists a specialization $(\tilde{u}'_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ that belongs to \mathcal{A} , so that $(\tilde{u}'_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ factors through one of the resolutions $Comp_1^2(t, c, u, z, y, p, a), \dots, Comp_s^2(t, c, u, z, y, p, a)$, and the specializations $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ and $(\tilde{u}'_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ restrict to the same exceptional family of $Term(GCl)$. We denote by (u'_0, z_0, y_0, p_0, a) , the projection of $(\tilde{u}'_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ to Γ . Then, (u'_0, z_0, y_0, p_0, a) belongs to the collection $\Gamma\mathcal{A}$ and factors through the resolution $Comp_j^2(t, c, u, z, y, p, a)$.

If the induced Γ -specialization $(c_0, u''_0, z'_0, y'_0, p_0, a)$ of the terminal p -rigid or p -solid group $Term(Comp_j^2)$ of $Comp_j^2$ is not Γ -exceptional, we lift it to a non-exceptional specialization $(\tilde{c}_0, \tilde{u}''_0, \tilde{z}'_0, \tilde{y}'_0, \tilde{p}_0, a)$ of $Term(Comp_j^2)$ in F_k . The specialization $(u''_0, z'_0, y'_0, p_0, a)$ belongs to $\Gamma\mathcal{A}$ and hence $(\tilde{u}''_0, \tilde{z}'_0, \tilde{y}'_0, \tilde{p}_0, a)$ belongs to \mathcal{A} . Thus,

and according to the construction of the resolutions $Comp_1^2, \dots, Comp_s^2$, there exists some resolution $Comp_{j'}^2$, for $j' \neq j$, whose upper levels are identical to that of $Comp_j^2$ but its terminal group is a p -rigid or p -solid limit quotient of the $Term(Comp_j^2)$, so that there exists a specialization $(\tilde{u}_0^{(3)}, \tilde{z}_0'', \tilde{y}_0'', \tilde{p}_0, a)$ of $Term(Comp_j^2)$ that restrict to the same exceptional family of $Term(GCl)$ represented by $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$. In particular, the specialization $(\tilde{z}_0'', \tilde{y}_0'', \tilde{p}_0, a)$ restrict to the same exceptional family of B represented by $(\tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$. Hence, there exists some $\tilde{u}_0^{(4)} \in F_k$, so that the specialization $(\tilde{u}_0^{(4)}, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$ factors through $Comp_{j'}$, and restricts to the same exceptional family of $Term(GCl)$ represented by $(\tilde{u}_0, \tilde{z}_0, \tilde{y}_0, \tilde{p}_0, a)$. We denote by $u_0^{(4)}$ the projection of $\tilde{u}_0^{(4)}$ to Γ . Then, the Γ -specialization $(u_0^{(4)}, z_0, y_0, p_0, a)$ factors through $Comp_{j'}$, and restricts to the same Γ -exceptional family represented by (u_0, z_0, y_0, p_0, a) . A finite induction argument, implies that we can assume that the induced Γ -specialization of $Term(Comp_{j'}^2)$ is Γ -exceptional. \square

For each terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in the $ExtraPS$ MR diagram $ExtraPSMRD$ of the collection $WPRS$ over the resolution $Comp(z, y, w, p, a)$, we consider the finitely many systems $\Sigma = \{\Sigma_i(c, u, z, y, w, p, a)\}_i$ stating that the specializations of the variables u in the specialization (u, z, y, w, p, a) are collapsed. Let $Comp_1^2(t, c, u, z, y, w, p, a), \dots, Comp_s^2(t, c, u, z, y, w, p, a)$ be the quotient resolutions of the systems Σ w.r.t. the graded closure $ExtraPSGCl(u, z, y, w, p, a)$.

Theorem 8.8. 1. Let (z_0, y_0, w_0, p_0, a) be a validPS statement that factors through the completed resolution $Comp(z, y, w, p, a)$. If $p_0 \notin TSPS(p)$, then (z_0, y_0, w_0, p_0, a) factors through some quotient resolution $Comp^2(t, c, u, z, y, w, p, a)$, associated with some terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in the $ExtraPS$ MR diagram $ExtraPSMRD$, so that the complexity of $Comp^2$ is strictly smaller than that of $Comp(z, y, w, p, a)$.

2. Let (z_0, y_0, w_0, p_0, a) be a Γ -validPS statement that factors through the completed resolution $Comp(z, y, w, p, a)$. If $p_0 \notin TSPS_\Gamma(p)$, then (z_0, y_0, w_0, p_0, a) factors through some quotient resolution $Comp^2(t, c, u, z, y, w, p, a)$, associated with some terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in the $ExtraPS$ MR diagram $ExtraPSMRD$, so that the complexity of $Comp^2$ is strictly smaller than that of $Comp(z, y, w, p, a)$.

Proof. We start with the first part of the claim. Since the specialization (z_0, y_0, w_0, p_0, a) is a validPS statement, it does not factor through the diagram $PosGFMRD$. Let $ExtraPSGCl(u, z, y, w, p, a)$ be a terminal graded closure in the $ExtraPS$ MR diagram $ExtraPSMRD$, and assume that there exists some $u_0 \in F_k$, so that $(u_0, z_0, y_0, w_0, p_0, a)$ factors through $ExtraPSGCl(u, z, y, w, p, a)$.

Since (z_0, y_0, w_0, p_0, a) is a validPS statement, the specialization u_0 must be collapsed in the specialization $(u_0, z_0, y_0, w_0, p_0, a)$. Hence, and according to theorem 8.6, there exists some $u'_0 \in F_k$, so that $(u'_0, z_0, y_0, w_0, p_0, a)$ factors through some quotient resolution $Comp^2(t, c, u, z, y, w, p, a)$ associated with $ExtraPSGCl(u, z, y, w, p, a)$, and the specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ restrict to the same exceptional family of $Term(ExtraPSGCl)$.

If the complexity of $Comp^2$ equals that of $Comp(z, y, w, p, a)$, then $Comp^2$ is a graded formal closure of $Comp(z, y, w, p, a)$. Hence, according to lemma 7.8, in this case, the specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ factors through the $CollapseExtra$ MR diagram $CollapseExtraMRD$ associated with the graded closure $ExtraPSGCl(u, z, y, w, p, a)$.

Hence, and since $p_0 \notin TSPS(p)$, there must exist some terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in the $ExtraPS$ MR diagram $ExtraPSMRD$, and some $u_0 \in F_k$, so that $(u_0, z_0, y_0, w_0, p_0, a)$ factors through $ExtraPSGCl(u, z, y, w, p, a)$ and for all $u'_0 \in F_k$, for which the specializations $(u_0, z_0, y_0, w_0, p_0, a)$ and $(u'_0, z_0, y_0, w_0, p_0, a)$ restrict to the same exceptional family of $Term(ExtraPSGCl)$, the specialization $(u'_0, z_0, y_0, w_0, p_0, a)$ does not factor through any of the quotient resolutions $Comp^2$ associated with $ExtraPSGCl(u, z, y, w, p, a)$ and have the same complexity as that of $Comp(z, y, w, p, a)$. That is, (z_0, y_0, w_0, p_0, a) must factor through some quotient resolution $Comp^2$ associated with $ExtraPSGCl(u, z, y, w, p, a)$, with complexity strictly smaller than that of $Comp(z, y, w, p, a)$.

The second statement follows by a similar argument that uses 8.7 and 7.9 in place of theorems 8.6 and 7.8. \square

In light of theorem 8.8, for every terminal graded closure $ExtraPSGCl(u, z, y, w, p, a)$ in the $ExtraPS$ MR diagram $ExtraPSMRD$ of the collection $WPRS$ over the resolution $Comp(z, y, w, p, a)$, we collect those quotient resolutions $Comp^2$ associated with $ExtraPSGCl(u, z, y, w, p, a)$, who admit a strictly smaller

complexity than that of $Comp(z, y, w, p, a)$, and for which there exists some non-collapsed specialization (z_0, y_0, w_0, p_0, a) of $Comp(z, y, w, p, a)$ that factors through $Comp^2$. According to theorem 8.5, the sets $TSPS^2(p)$ and $TSPS^2_\Gamma(p)$, that correspond to the new resolution $Comp^2$, are defined by one formula $\varphi^2(p)$ in the Boolean algebra of AE -formulas, for every group Γ that belongs to sufficiently high level of the model.

We Continue iteratively. By the definition of the complexity of a graded resolution, this process must terminate after finitely many steps. In each step, we have a formula $\varphi^j(p)$ in the Boolean algebra of AE -formulas, that defines a subset of $EAE(p)$ and a subset of $EAE_\Gamma(p)$, for every group Γ that belongs to sufficiently high level of the model. Moreover, all the quotient resolutions in the final step of this procedure, must be of maximal complexity. Hence, every $p_0 \in EAE(p)$ for which there exists some validPS statement that factors through the initial resolution $Comp(z, y, w, p, a)$, must be defined by some formula $\varphi^j(p)$ along the process. Similarly, if Γ belongs to sufficiently high level in the model, then, every $p_0 \in EAE_\Gamma(p)$ for which there exists some Γ -validPS statement that factors through the initial resolution $Comp(z, y, w, p, a)$, must be defined by some formula $\varphi^j(p)$ along the process.

9 Proof Systems

Assume that we have a $EAE(p)$ set over F_k defined by the formula:

$$\exists w \forall y \exists x \quad \Sigma(x, y, w, p, a) = 1 \wedge \Psi(x, y, w, p, a) \neq 1.$$

Given a group Γ , we denote by $EAE_\Gamma(p)$, the set of Γ -specializations $p_0 \in \Gamma$ of the parameters p , that are defined by the above formula when applied on Γ . We also denote by $AE(w, p)$ and by $AE_\Gamma(w, p)$ the sets defined by the formula

$$\forall y \exists x \quad \Sigma(x, y, w, p, a) = 1 \wedge \Psi(x, y, w, p, a) \neq 1,$$

over F_k and over Γ , correspondingly. If $(w_0, p_0) \in AE(w, p)$, we say that w_0 is a *witness* for $p_0 \in EAE(p)$. Note that for every $p_0 \in EAE(p)$ there must exist some witness $w_0 \in F_k$ so that $(w_0, p_0) \in AE(w, p)$. Similarly, if $(w_0, p_0) \in AE_\Gamma(w, p)$, we say that w_0 is a *witness* for $p_0 \in EAE_\Gamma(p)$, and we note that for every $p_0 \in EAE_\Gamma(p)$ there must exist some witness $w_0 \in \Gamma$ so that $(w_0, p_0) \in AE_\Gamma(w, p)$.

Our aim in this section, is proving quantifier elimination. That is, we will show that the sets $EAE(p)$ and $EAE_\Gamma(p)$ belong to the Boolean algebra of AE -sets. Moreover, we show that there exists a formula $\varphi(p)$ in the Boolean algebra of AE -formulas, that defines each of these sets uniformly (over the corresponding groups), for almost all the groups Γ . For that aim, we bring the situation to the setting of section §8. For that, we construct finitely many (graded) *proof systems*. A proof system, roughly, is a finite tree whose branches consist of alternating graded MR diagrams and graded formal MR diagrams. Using the graded formal MR diagrams along a proof system, we can apply the formal solutions of our fixed system $\Sigma(x, y, w, p, a)$ on the specializations that factor through a graded resolution in the graded MR diagram that was placed in the previous stage in the proof system. Those specializations for which all of the formal solutions applied on them do not satisfy our fixed set of inequalities $\Psi(x, y, w, p, a) \neq 1$ (the remaining y 's), are passed to the next graded MR diagram in the proof system, and so on.

9.1 The Construction of Proof Systems of Depth 2 under Minimal Rank Assumption

Let $GFMRD_1$ be the graded formal MR diagram of the system $\Sigma(x, y, w, p, a)$ over the wp -graded resolution $F_y * F(w, p, a)$. According to the construction of $GFMRD_1$, every graded formal resolution $GFRes^1(x, y, w, p, a)$ in it, is defined over a graded formal closure of $F_y * F(w, p, a)$, which is a limit group of the form $WPH(h_1, w, p, a) * F_y$, where $WPH(h_1, w, p, a)$ is a wp -rigid or a wp -solid limit group.

We apply all the modular automorphisms of the decompositions associated with the various groups along the resolution $GFRes^1(x, y, w, p, a)$, and we specialize the free factors that do not intersect a conjugate of the subgroup $\langle y, w, p, a \rangle$ in all the possible ways. That way, each word $\psi_i(x, y, w, p, a)$ in the system $\Psi(x, y, w, p, a)$, induces a set of elements $\lambda_i^1(h_1, y, w, p, a)$ in the group $WPH(h_1, w, p, a) * F_y$. We think on the induced set $\lambda_i^1(h_1, y, w, p, a)$ as a system of equations imposed on the variables (h_1, y, w, p, a) . By Guba's

theorem, for all i , the system $\lambda_i^1(h_1, y, w, p, a)$ is equivalent to a finite system $\lambda_i^1(h_1, y, w, p, a)$ over the free group F_k .

Denote by \mathfrak{J}^2 the collection of all the specializations $(h_{1,0}, y_0, w_0, p_0, a)$ of $WPH(h_1, w, p, a) * F_y$, so that $(h_{1,0}, y_0, w_0, p_0, a)$ restricts to an exceptional specialization of $WPH(h_1, w, p, a)$, and solves the system $\lambda_i^1(h_1, y, w, p, a)$, for some i . From the collection \mathfrak{J}^2 , we construct a wp -graded MR diagram, denoted $GRemMRD_2$. Note that the wp -graded limit group $WPH(h_1, w, p, a)$ is mapped compatibly into each of the wp -graded limit groups in this MR diagram $GRemMRD_2$. Every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in the diagram $GRemMRD_2$, terminates in a group $WPHG(g_1, h'_1, w, p, a)$ which is either wp -rigid or wp -solid. The variables g_1, h'_1 stand for the canonical images of the variables y, h_1 correspondingly in the terminal group $WPHG(g_1, h'_1, w, p, a)$ of the resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$. We denote the collection of all the resolutions in the diagram $GRemMRD_2$ briefly by $GRem_2$.

We denote the first level groups in $GRemMRD_2$, i.e., the finitely many maximal limit groups associated with the collection \mathfrak{J}^2 , by $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$.

Lemma 9.1. *Let Γ be a group in the model. Then, $\lambda_i^1(h_1, y, w, p, a)$ is equivalent to $\lambda_i^1(h_1, y, w, p, a)$ over Γ . Moreover, if $(h_{1,0}, y_0, w_0, p_0, a)$ is a Γ -specialization of $WPH(h_1, w, p, a) * F_y$, so that $(h_{1,0}, y_0, w_0, p_0, a)$ restricts to a Γ -exceptional specialization of $WPH(h_1, w, p, a)$, and solves the system $\lambda_i^1(h_1, y, w, p, a)$, for some i , then $(h_{1,0}, y_0, w_0, p_0, a)$ factors through one of the resolutions in $GRem_2$.*

Proof. Let $(h_{1,0}, y_0, w_0, p_0, a)$ be a Γ -solution for the system $\lambda_i^1(h_1, y, w, p, a)$. We lift $(h_{1,0}, y_0, w_0, p_0, a)$ to a solution $(\tilde{h}_{1,0}, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ of the system $\lambda_i^1(h_1, y, w, p, a)$ in F_k . Since $\lambda_i^1(h_1, y, w, p, a)$ and $\lambda_i^1(h_1, y, w, p, a)$ are equivalent over F_k , then $(\tilde{h}_{1,0}, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ is a solution for $\lambda_i^1(h_1, y, w, p, a)$ in F_k . Hence, the projection of $(\tilde{h}_{1,0}, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ to Γ , which is $(h_{1,0}, y_0, w_0, p_0, a)$, is a solution for $\lambda_i^1(h_1, y, w, p, a)$ in Γ .

Assume further that $(h_{1,0}, w_0, p_0, a)$ is a Γ -exceptional specialization of $WPH(h_1, w, p, a)$. Then $(\tilde{h}_{1,0}, \tilde{w}_0, \tilde{p}_0, a)$ is an exceptional specialization of $WPH(h_1, w, p, a)$ in F_k , and the specialization $(\tilde{h}_{1,0}, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ belongs to the collection \mathfrak{J}^2 . Hence, by the definition of $GRem_2$, the specialization $(\tilde{h}_{1,0}, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ factors through some resolution in $GRem_2$, and thus, the Γ -specialization $(h_{1,0}, y_0, w_0, p_0, a)$ factors through that resolution. If the Γ -specialization of the terminal group $WPHG(g_1, h'_1, w, p, a)$ of that resolution is not Γ -exceptional, we lift it to a non-exceptional specialization of $WPHG(g_1, h'_1, w, p, a)$. This lift must belong to the collection \mathfrak{J}^2 , and we can continue the factorization through the flexible quotients of $WPHG(g_1, h'_1, w, p, a)$. Hence, a finite induction implies the desired. \square

Lemma 9.2.

1. *Let $(h_{1,0}, w_0, p_0, a)$ be an exceptional specialization of $WPH(h_1, w, p, a)$, and assume that it factors through one of the groups $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$. Then, every specialization of $WPH(h_1, w, p, a)$ that belongs to the exceptional family represented by $(h_{1,0}, w_0, p_0, a)$, factors through one of the groups $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$.*
2. *Let Γ be a group in the model. Let $(h_{1,0}, w_0, p_0, a)$ be a Γ -exceptional specialization of $WPH(h_1, w, p, a)$, and assume that it factors through one of the groups $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$. Then, every Γ -specialization of $WPH(h_1, w, p, a)$ that belongs to the Γ -exceptional family represented by $(h_{1,0}, w_0, p_0, a)$, factors through one of the groups $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$.*

Proof. Assume that the specialization $(h_{1,0}, y_0, w_0, p_0, a)$ factors through one of the groups

$$\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a),$$

for some $y_0 \in F_k$. Then, the specialization $(h_{1,0}, y_0, w_0, p_0, a)$ is a solution for the system $\lambda_i^1(h_1, y, w, p, a)$, for some i . By the definition of the set of elements $\lambda_i^1(h_1, y, w, p, a)$ of the group $WPH(h_1, w, p, a) * F_y$, if θ is a modular automorphism of the wp -graded limit group $WPH(h_1, w, p, a)$, then, applying θ on the set of elements $\lambda_i^1(h_1, y, w, p, a)$, does not change this set. Hence, if $(h_{1,0}\theta, y_0, w_0, p_0, a)$ is the specialization of $WPH(h_1, w, p, a) * F_y$ obtained by precomposing θ with the specialization $(h_{1,0}, y_0, w_0, p_0, a)$, then $(h_{1,0}\theta, y_0, w_0, p_0, a)$ is a solution for the system $\lambda_i^1(h_1, y, w, p, a)$.

For the second part of the claim, let $(h_{1,0}, y_0, w_0, p_0, a)$ be a Γ -specialization of one of the groups $\lambda^1 WPGL_j(h_1, y, w, p, a)$, for some $j = 1, \dots, d$ and some $y_0 \in \Gamma$. We lift $(h_{1,0}, y_0, w_0, p_0, a)$ to a specialization $(\tilde{h}_{1,0}, \tilde{y}_0, \tilde{w}_0, \tilde{p}_0, a)$ of $\lambda^1 WPGL_j(h_1, y, w, p, a)$ in F_k . Then, by the above argument, every specialization of

$WPH(h_1, w, p, a)$ that belongs to the same exceptional family represented by $(\tilde{h}_{1,0}, \tilde{w}_0, \tilde{p}_0, a)$, factors through one of the groups $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$. Hence, every Γ -specialization of $WPH(h_1, w, p, a)$ that belongs to the same Γ -exceptional family represented by $(h_{1,0}, w_0, p_0, a)$, factors through one of the groups $\lambda^1 WPGL_1(h_1, y, w, p, a), \dots, \lambda^1 WPGL_d(h_1, y, w, p, a)$. \square

Before we continue the construction of the proof systems, we stop for an approximation of the sets $EAE(p)$ and $EAE_\Gamma(p)$ that is uniform for all the groups Γ in the model.

Lemma 9.3. *We denote:*

1. Let $U_1(p)$ be the set of all $p_0 \in F_k$, for which there exists $w_0 \in F_k$, so that for one of the groups $WPH(h_1, w, p, a)$, there exists $h_{1,0} \in F_k$, so that $(h_{1,0}, w_0, p_0, a)$ is an exceptional specialization of $WPH(h_1, w, p, a)$, and for every limit group $\lambda^1 WPGL_j(h_1, y, w, p, a)$ associated with $WPH(h_1, w, p, a)$, there exists no $y_0 \in F_k$, so that $(h_{1,0}, y_0, w_0, p_0, a)$ is a specialization of $\lambda^1 WPGL_j(h_1, y, w, p, a)$.
2. Given a group Γ in the model, we let $U_1^\Gamma(p)$ be the set of all $p_0 \in \Gamma$, for which there exists $w_0 \in \Gamma$, so that for one of the groups $WPH(h_1, w, p, a)$, there exists $h_{1,0} \in \Gamma$, so that $(h_{1,0}, w_0, p_0, a)$ is a Γ -exceptional specialization of $WPH(h_1, w, p, a)$, and for every limit group $\lambda^1 WPGL_j(h_1, y, w, p, a)$ associated with $WPH(h_1, w, p, a)$, there exists no $y_0 \in \Gamma$, so that $(h_{1,0}, y_0, w_0, p_0, a)$ is a Γ -specialization of $\lambda^1 WPGL_j(h_1, y, w, p, a)$.

Then,

1. $U_1(p) \subset EAE(p)$, and there exists a formula $\varphi_1(p)$ in the Boolean algebra of AE-formulas, that defines the set $U_1(p)$.
2. Let Γ be a group in the model. Then, $U_1^\Gamma(p) \subset EAE_\Gamma(p)$, and the same formula $\varphi_1(p)$ above, defines the set $U_1^\Gamma(p)$ in Γ .

Proof. The existence of a formula $\varphi_1(p)$ that satisfies the claim is obvious. The fact that $U_1(p) \subset EAE(p)$ follows from the definition of the limit groups $\lambda^1 WPGL_j(h_1, y, w, p, a)$ as the maximal limit groups through which the specializations in the collection \mathfrak{J}^2 factor, and from lemma 5.11.

Now let Γ be a group in the model. We prove that $U_1^\Gamma(p) \subset EAE_\Gamma(p)$. Let $(h_{1,0}, w_0, p_0, a)$ be a Γ -exceptional specialization of $WPH(h_1, w, p, a)$, and assume it does not factor through any of the limit groups $\lambda^1 WPGL_j(h_1, y, w, p, a)$ associated with $WPH(h_1, w, p, a)$. Hence, according to lemma 9.1, every Γ -specialization $(h_{1,0}, y_0, w_0, p_0, a)$ of $WPH(h_1, w, p, a) * F_y$, does not solve any of the systems $\lambda_i^1(h_1, y, w, p, a)$. Hence, and using lemma 5.12, we deduce that $p_0 \in EAE_\Gamma(p)$. \square

Now we consider a wp -graded resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRem_2$, who terminates in a group $WPHG(g_1, h'_1, w, p, a)$ which is either wp -rigid or wp -solid. Let $Comp^2 = Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$ be the completion of the resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$, and denote its base group by $B = WPHG(g_1, h'_1, w, p, a)$.

We construct the wp -graded formal MR diagram $GFMRD_2$ of the system $\Sigma(x, y, w, p, a)$ over the resolution $Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$. Every graded formal resolution $GFRes^2(x, z, h_1, y, w, p, a)$ in $GFMRD_2$, terminates in a graded formal closure $GFCl(x, z, h_1, y, w, p, a)$ of $Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$.

We fix a generating set $(h_2, g_1, h'_1, w, p, a)$ for the terminal group $Term(GFCl)$ of $GFCl(x, z, h_1, y, w, p, a)$, and denote $Term(GFCl) = WPHGH(h_2, g_1, h'_1, w, p, a)$. The elements (g_1, h'_1, w, p, a) of $WPHGH(h_2, g_1, h'_1, w, p, a)$ are the canonical images of the elements (g_1, h'_1, w, p, a) of B in $Term(GFCl)$. Note that, since $GFCl(x, z, h_1, y, w, p, a)$ is a graded formal closure for the resolution $Comp(\lambda^1 WPGRes)$, it is generated, as a group, by $Term(GFCl)$ and (the canonical image of) $Comp(\lambda^1 WPGRes)$, together with a finite collection of elements s , that are roots for some elements in the abelian vertex groups that appear in the decompositions associated with the various levels of $Comp(\lambda^1 WPGRes)$.

We apply all the modular automorphisms of the decompositions associated with the various groups along the resolution $GFRes^2(x, z, h_1, y, w, p, a)$, and we specialize the free factors that do not intersect a conjugate of the subgroup $\langle z, h_1, y, w, p, a \rangle$ in all the possible ways. That way, each word $\psi_i(x, y, w, p, a)$ in the system $\Psi(x, y, w, p, a)$, induces a set of elements $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ in the group $GFCl(x, z, h_1, y, w, p, a)$. We think on the induced set $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ as a system of equations imposed on the variables

$(s, z, h_2, g_1, h'_1, y, w, p, a)$. By Guba's theorem, for all i , the system $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ is equivalent to a finite system $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ over the free group F_k .

Let \mathfrak{J}^3 be the collection of all the specializations $(s_0, z_0, h_{2,0}, g_{1,0}, h'_{1,0}, y_0, w_0, p_0, a)$ of $GFCl$ that restrict to exceptional specializations of $Term(GFCl)$, and satisfy one of the systems $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$, for some i . The specializations in the collection \mathfrak{J}^3 , factor through finitely many maximal limit groups $\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a)$.

Lemma 9.4. *Let Γ be a group in the model. Then, $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ is equivalent to $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ over Γ . Moreover, if $(s_0, z_0, h_{2,0}, g_{1,0}, h'_{1,0}, y_0, w_0, p_0, a)$ is a Γ -specialization of $GFCl$, so that $(s_0, z_0, h_{2,0}, g_{1,0}, h'_{1,0}, y_0, w_0, p_0, a)$ restricts to a Γ -exceptional specialization of $Term(GFCl)$, and solves the system $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$, for some i , then $(s_0, z_0, h_{2,0}, g_{1,0}, h'_{1,0}, y_0, w_0, p_0, a)$ factors through one of the groups $\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a)$.*

Proof. Identical to lemma 9.1. □

Lemma 9.5.

1. Let $(h_{2,0}, g_{1,0}, h'_{1,0}, w_0, p_0, a)$ be an exceptional specialization of $WPHGH(h_2, g_1, h'_1, w, p, a)$, and assume that it factors through one of the groups

$$\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a).$$

Then, every specialization of $WPHGH(h_2, g_1, h'_1, w, p, a)$ that belongs to the exceptional family represented by $(h_{2,0}, g_{1,0}, h'_{1,0}, w_0, p_0, a)$, factors through one of the groups

$$\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a).$$

2. Let Γ be a group in the model. Let $(h_{2,0}, g_{1,0}, h'_{1,0}, w_0, p_0, a)$ be a Γ -exceptional specialization of $WPHGH(h_2, g_1, h'_1, w, p, a)$, and assume that it factors through one of the groups

$$\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a).$$

Then, every Γ -specialization of $WPHGH(h_2, g_1, h'_1, w, p, a)$ that belongs to the Γ -exceptional family represented by $(h_{2,0}, g_{1,0}, h'_{1,0}, w_0, p_0, a)$, factors through one of the groups

$$\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a).$$

Proof. Similar to lemma 9.2. □

Lemma 9.6. *Let $GFCl_1, \dots, GFCl_r$ be a collection of terminal graded formal closures in the graded formal MR diagram $GFMRD_2$ associated with the resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$.*

1. Let $(g_{1,0}, h'_{1,0}, w_0, p_0, a)$ be an exceptional specialization of the wp -graded base group B of $Comp(\lambda^1 WPGRes)$, and for all $l = 1, \dots, r$, let $(h_{2,0}^l, g_{1,0}^l, h'_{1,0}^l, w_0, p_0, a)$ be an exceptional specialization of the wp -graded base group $Term(GFCl)$ of $GFCl_j$, so that the restriction $(g_{1,0}^l, h'_{1,0}^l, w_0, p_0, a)$ belongs to the exceptional family of B represented by $(g_{1,0}, h'_{1,0}, w_0, p_0, a)$.

Assume that the collection of the induced ungraded closures

$$GFCl_{l,0} = GFCl_{l,(h_{2,0}^l, g_{1,0}^l, h'_{1,0}^l, w_0, p_0, a)},$$

$l = 1, \dots, r$, form a covering closure for the induced ungraded resolution

$$Comp_0^2 = Comp(\lambda WPGRes)_{(g_{1,0}, h'_{1,0}, w_0, p_0, a)}.$$

Assume further that for all $l = 1, \dots, r$, the specialization $(h_{2,0}^l, g_{1,0}^l, h'_{1,0}^l, w_0, p_0, a)$ cannot be extended to a specialization of any of the groups $\lambda^2 WPGL_j(s, z, h_2, g_1, h'_1, y, w, p, a)$ associated with $GFCl_l$.

Then, for every $y_0 \in F_k$ that factors through the ungraded resolution $Comp_0^2$, there exists some $x_0 \in F_k$, so that

$$\Sigma(x_0, y_0, w_0, p_0, a) = 1 \wedge \Psi(x_0, y_0, w_0, p_0, a) \neq 1.$$

2. Let Γ be a group in the model.

Let $(g_{1,0}, h'_{1,0}, w_0, p_0, a)$ be a Γ -exceptional specialization of the wp -graded base group B of $\text{Comp}(\lambda\text{WPGRes})$, and for all $l = 1, \dots, r$, let $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ be a Γ -exceptional specialization of the wp -graded base group $\text{Term}(GFCl)$ of $GFCl_j$, so that the restriction $(g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ belongs to the Γ -exceptional family of B represented by $(g_{1,0}, h'_{1,0}, w_0, p_0, a)$.

Assume that the collection of the induced ungraded closures

$$GFCl_{l,0} = GFCl_{l,(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)},$$

$l = 1, \dots, r$, form a covering closure for the induced ungraded resolution

$$\text{Comp}_0^2 = \text{Comp}(\lambda\text{WPGRes})_{(g_{1,0}, h'_{1,0}, w_0, p_0, a)}.$$

Assume further that for all $l = 1, \dots, r$, the Γ -specialization $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ cannot be extended to a Γ -specialization of any of the groups $\lambda^2\text{WPGL}_j(s, z, h_2, g_1, h'_1, y, w, p, a)$ associated with $GFCl_l$.

Then, for every $y_0 \in \Gamma$ that factors through the ungraded resolution Comp_0^2 , there exists some $x_0 \in \Gamma$, so that

$$\Sigma(x_0, y_0, w_0, p_0, a) = 1 \wedge \Psi(x_0, y_0, w_0, p_0, a) \neq 1.$$

Proof. For the first statement, let $y_0 \in F_k$ so that it factors through the ungraded resolution Comp_0^2 . Since the collection $GFCl_{1,0}, \dots, GFCl_{r,0}$ is a covering closure for Comp_0^2 , y_0 must factor through $GFCl_{l,0}$, for some $l = 1, \dots, r$. Since $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ does not factor through any of the groups $\lambda^2\text{WPGL}_j(s, z, h_2, g_1, h'_1, y, w, p, a)$ associated with $GFCl_l$, and using lemma 5.11, we deduce the desired.

For the second statement, let $y_0 \in \Gamma$ so that it factors through the ungraded resolution Comp_0^2 . Since the collection $GFCl_{1,0}, \dots, GFCl_{r,0}$ is a covering closure for Comp_0^2 , y_0 must factor through $GFCl_{l,0}$, for some $l = 1, \dots, r$. Since $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ does not factor through any of the groups $\lambda^2\text{WPGL}_j(s, z, h_2, g_1, h'_1, y, w, p, a)$ associated with $GFCl_l$, then according to lemma 9.4, and using lemma 5.12, we deduce the desired. \square

We assume that the first level limit groups $\lambda^1\text{WPGL}_1(h_1, y, w, p, a), \dots, \lambda^1\text{WPGL}_d(h_1, y, w, p, a)$ of the wp -graded MR diagram $G\text{RemMRD}_2$ are of minimal rank. That is, we assume that for all $j = 1, \dots, d$, if the group $\lambda^1\text{WPGL}_j(h_1, y, w, p, a)$ admits a restricted epimorphism onto $F * F_k$, for some free group F , so that the subgroup $\langle w, p, a \rangle$ is mapped entirely into F_k , then F is the trivial group.

From now until the end of this section, we continue under this assumption.

This assumption, allows us to apply theorem 8.6 in order to construct from the collection \mathfrak{J}^3 of all the specializations $(s_0, z_0, h_{2,0}, g_{1,0}, h'_{1,0}, y_0, w_0, p_0, a)$ of $GFCl$ that restrict to exceptional specializations of $\text{Term}(GFCl) = \text{WPHGH}(h_2, g_1, h'_1, w, p, a)$, and satisfy one of the systems $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$, for some i , to construct the quotient resolutions $\text{Comp}_1^3(t, s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \text{Comp}_s^3(t, s, z, h_2, g_1, h'_1, y, w, p, a)$ of the systems $\lambda_i^2(s, z, h_2, g_1, h_1, y, w, p, a)$ w.r.t. the wp -graded closure $GFCl$. According to the construction of the quotient resolutions in this case, $\text{Comp}_1^3, \dots, \text{Comp}_s^3$ are quotients of the graded formal closure $GFCl$. And in the case that a quotient resolution Comp_j^3 is of maximal complexity, i.e., of complexity that equals the complexity of $\text{Comp}(\lambda^1\text{WPGRes})$, then, the completed resolution Comp_j^3 can differ from $GFCl$ only in its base group, i.e., Comp_j^3 is obtained from $GFCl$ by adding some relations to $\text{Term}(GFCl)$.

The collection of quotient resolutions associated with the graded formal closure $GFCl$ that are not of maximal complexity, is denoted by $G\text{Rem}_3$.

Lemma 9.7.

1. Let $(g_{1,0}, h'_{1,0}, w_0, p_0, a)$ be an exceptional specialization of the wp -rigid or wp -solid base group B of $\text{Comp}(\lambda^1\text{WPGRes})$, and assume that $(w_0, p_0) \in \text{AE}(w, p)$. Then, there exists a collection of terminal graded formal closures $GFCl_1, \dots, GFCl_r$ in the graded formal MR diagram $G\text{FMRD}_2$ associated with the resolution $\lambda^1\text{WPGRes}(h_1, y, w, p, a)$, together with exceptional specializations $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$, $l = 1, \dots, r$, of their corresponding wp -rigid or wp -solid base groups $\text{Term}(GFCl_l)$, so that:

- (a) The restriction $(g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ belongs to the exceptional family of B represented by $(g_{1,0}, h'_{1,0}, w_0, p_0, a)$, for all $l = 1, \dots, r$.

- (b) The induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{r,0}$ form a covering closure for the induced ungraded completion $Comp_0^2$.
- (c) For every maximal complexity quotient resolution $Comp_j^3$ that is associated with $GFCl_l$, there is no specialization of the wp -rigid or wp -solid base group $Term(Comp_j^3)$ of $Comp_j^3$, that restricts to the exceptional family of $Term(GFCl_l)$ represented by $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$.

2. Let Γ be a group in the model.

Let $(g_{1,0}, h_{1,0}^l, w_0, p_0, a)$ be a Γ -exceptional specialization of the wp -rigid or wp -solid base group B of $Comp(\lambda WPGRes)$, and assume that $(w_0, p_0) \in AE_\Gamma(w, p)$. Then, there exists a collection of terminal graded formal closures $GFCl_1, \dots, GFCl_r$ in the graded formal MR diagram $GFMRD_2$ associated with the resolution $\lambda WPGRes(h_1, y, w, p, a)$, together with Γ -exceptional specializations $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$, $l = 1, \dots, r$, of their corresponding wp -rigid or wp -solid base groups $Term(GFCl_l)$, so that:

- (a) The restriction $(g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$ belongs to the Γ -exceptional family of B represented by $(g_{1,0}, h_{1,0}^l, w_0, p_0, a)$, for all $l = 1, \dots, r$.
- (b) The induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{r,0}$ form a covering closure for the induced ungraded completion $Comp_0^2$.
- (c) For every maximal complexity quotient resolution $Comp_j^3$ that is associated with $GFCl_l$, there is no Γ -specialization of the wp -rigid or wp -solid base group $Term(Comp_j^3)$ of $Comp_j^3$, that restricts to the Γ -exceptional family of $Term(GFCl_l)$ represented by $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$.

Proof. For the first statement, since $(w_0, p_0) \in AE(w, p)$, then, according to theorem 5.8, there exists a collection of terminal graded formal closures $GFCl_1, \dots, GFCl_r$ in $GFMRD_2$, together with exceptional specializations $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$, $l = 1, \dots, r$, of their corresponding wp -graded base groups $Term(GFCl_l)$, so that the conditions (a) and (b) are satisfied, and the elements of the system $\lambda_i^2(s, z, h_2, g_1, h_1', y, w, p, a)$ are not all trivial in the induced ungraded formal closure $GFCl_{l,0}$, for all i and all $l = 1, \dots, r$. Assume by contradiction that for some maximal complexity quotient resolution $Comp_j^3$, the base group $Term(Comp_j^3)$ admits a specialization that belongs to the modular family of $Term(GFCl_l)$ represented by $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$, and let $Comp_{j,0}^3$ be the induced ungraded resolution. Hence, since $Comp_j^3$ is of maximal complexity, and according to lemma 9.5, we deduce that the elements of the system $\lambda_i^2(s, z, h_2, g_1, h_1', y, w, p, a)$ are not all trivial in $Comp_{j,0}^3$, for all i . This contradicts the construction of $Comp_j^3$ being a quotient resolution for the systems $\lambda_i^2(s, z, h_2, g_1, h_1', y, w, p, a)$ w.r.t. the wp -graded closure $GFCl_l$.

For the second statement, since $(w_0, p_0) \in AE_\Gamma(w, p)$, then, according to theorem 5.10, there exists a collection of terminal graded formal closures $GFCl_1, \dots, GFCl_r$ in $GFMRD_2$, together with Γ -exceptional specializations $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$, $l = 1, \dots, r$, of their corresponding wp -graded base groups $Term(GFCl_l)$, so that the conditions (a) and (b) are satisfied, and the elements of the system $\lambda_i^2(s, z, h_2, g_1, h_1', y, w, p, a)$ are not all trivial in the induced ungraded formal closure $GFCl_{l,0}$, for all i and all $l = 1, \dots, r$. Assume by contradiction that for some maximal complexity quotient resolution $Comp_j^3$, the base group $Term(Comp_j^3)$ admits a Γ -specialization that belongs to the modular family of $Term(GFCl_l)$ represented by $(h_{2,0}^l, g_{1,0}^l, h_{1,0}^l, w_0, p_0, a)$, and let $Comp_{j,0}^3$ be the induced ungraded resolution. Hence, since $Comp_j^3$ is of maximal complexity, and according to lemma 9.5, we deduce that the elements of the system $\lambda_i^2(s, z, h_2, g_1, h_1', y, w, p, a)$ are not all trivial in $Comp_{j,0}^3$, for all i . This contradicts the construction of $Comp_j^3$ being a quotient resolution for the systems $\lambda_i^2(s, z, h_2, g_1, h_1', y, w, p, a)$ w.r.t. the wp -graded closure $GFCl_l$. \square

Definition 9.8. A *proof system of depth 2* is to make the following ordered choice:

1. To choose one of the terminal graded formal closures $WPH(h_1, w, p, a) * F_y$ in $GFMRD_1$.
2. For every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRemMRD_2$, to choose an integer $0 \leq n \leq M(WPHG)$, where $WPHG(g_1, h_1', w, p, a)$ is the wp -rigid or wp -solid base group of the completed resolution $Comp^2 = Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$, and $M(WPHG)$ is the maximal possible number of exceptional families of specializations of $WPHG$ for a fixed value of the parameters $w_0, p_0 \in F_k$.
3. For every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRemMRD_2$ for which the integer n was chosen, and for every $1 \leq i \leq n$, to choose finitely many terminal graded formal closures $GFCl_1^i, \dots, GFCl_{m_i}^i$ in the

graded formal MR diagram $GFMRD_2$ associated with the resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$, where $1 \leq m_i \leq \Sigma \{M(Term(GFCl)) : GFCl \text{ is a terminal closure in } GFMRD_2\}$, and $M(Term(GFCl))$ is the maximal possible number of exceptional families of specializations of the wp -rigid or wp -solid base group $Term(GFCl)$ for a fixed value of the parameters $w_0, p_0 \in F_k$.

We say that a proof system PS as above, verifies $(w_0, p_0) \in AE(w, p)$, if for every graded formal closure $GFCl_j^i$ listed in the proof system PS , there exists an exceptional specialization $(h_{2,0}^{i,j}, g_{1,0}^{i,j}, h_{1,0}^{i,j}, w_0, p_0, a)$ of $Term(GFCl_j^i)$, so that

1. For all i, j , the specializations $(h_{1,0}^{i,j}, w_0, p_0, a)$ represent the same exceptional family of the group $WPH(h_1, w, p, a)$ that was specified by the proof system PS .
2. For all i , all the specializations $(g_{1,0}^{i,j}, h_{1,0}^{i,j}, w_0, p_0, a)$ represent the same exceptional family of the group $WPHG(g_1, h'_1, w, p, a)$ for which the graded formal closures $GFCl_j^i$ were associated by the proof system PS .
3. For all i , the induced ungraded closures $GFCl_{1,0}^i, \dots, GFCl_{m_i,0}^i$ form a covering closure for the induced ungraded resolution $Comp_0^2$, where $Comp^2 = Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$ is the completion of the resolution of $GRemMRD_2$ for which the graded formal closures $GFCl_j^i$ were associated by the proof system PS .
4. For every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRemMRD_2$ for which the integer n was chosen by the proof system PS , and for every $1 \leq i_1 \neq i_2 \leq n$, the exceptional specializations $(g_{1,0}^{i_1,j}, h_{1,0}^{i_1,j}, w_0, p_0, a)$ and $(g_{1,0}^{i_2,j}, h_{1,0}^{i_2,j}, w_0, p_0, a)$ represent distinct exceptional families of the terminal wp -graded group $WPHG(g_1, h'_1, w, p, a)$ of $\lambda^1 WPGRes$.
5. For every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRemMRD_2$ for which the integer n was chosen by the proof system PS , the terminal group $WPHG(g_1, h'_1, w, p, a)$ of $\lambda^1 WPGRes$, admits exactly n distinct exceptional families corresponding to the value $w_0 p_0$ of the parameters.
6. For all i, j , the specialization $(h_{2,0}^{i,j}, g_{1,0}^{i,j}, h_{1,0}^{i,j}, w_0, p_0, a)$ does not factor through any of the maximal limit groups $\lambda^2 WPGL_1(s, z, h_2, g_1, h'_1, y, w, p, a), \dots, \lambda^2 WPGL_{d_2}(s, z, h_2, g_1, h'_1, y, w, p, a)$ associated with the graded formal closure $GFCl_j^i$.

Such a tuple

$$\left(\left(h_{2,0}^{i,j}, g_{1,0}^{i,j}, h_{1,0}^{i,j} \right)_{i,j}, w_0, p_0, a \right)$$

that satisfies the conditions listed above, is called a *validPS statement* (of depth 2).

9.2 The Construction of Proof Systems of General Depth under Minimal Rank Assumption

The construction that we explained so far, suffices for encoding “proofs” for that $(w_0, p_0) \in AE(w, p)$ for a subset of the parameters in $AE(w, p)$ (those that admit a validPS statement of depth 2 or of depth 1 - see definition 9.8 and lemma 9.3 correspondingly). In order to encode “proofs” for all the parameters in $AE(w, p)$, we continue the construction with each of the quotient resolutions

$$Comp_1^3(t, s, z, h_2, g_1, h'_1, y, w, p, a), \dots, Comp_s^3(t, s, z, h_2, g_1, h'_1, y, w, p, a).$$

According to lemma 9.7, we can continue only with those quotient resolutions that are not of maximal complexity. We summarize the construction so far, and explain how to continue it iteratively.

Let $Comp^2 = Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$ be the completion of the wp -graded resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in the diagram $GRemMRD_2$, who terminates in the wp -rigid or wp -solid group $WPHG(g_1, h'_1, w, p, a)$. Let $GFMRD_2$ be the wp -graded formal MR diagram of the system $\Sigma(x, y, w, p, a)$ over the resolution $Comp^2$. Let $GFRes^2(x, z, h_1, y, w, p, a)$ be a graded formal resolution in $GFMRD_2$, who terminates in a graded formal closure $GFCl(x, z, h_1, y, w, p, a)$ of $Comp^2$ whose wp -rigid or wp -solid base

group is base $Term(GFCl) = WPHGH(h_2, g_1, h'_1, w, p, a)$. Let $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ be the finitely many systems associated with $GFCl$ using the resolution $GFRes^2$ and the elements ψ_i of the inequalities Ψ . Let \mathfrak{J}^3 be the collection of all the specializations $(s_0, z_0, h_{2,0}, g_{1,0}, h'_{1,0}, y_0, w_0, p_0, a)$ of $GFCl$ that restrict to exceptional specializations of $Term(GFCl)$, and satisfy one of the systems $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$, for some i . The specializations in the collection \mathfrak{J}^3 , factors through finitely many maximal limit groups $\lambda^2 WPGL_j(s, z, h_2, g_1, h'_1, y, w, p, a)$.

Let $Comp_i^3(t, s, z, h_2, g_1, h'_1, y, w, p, a)$ be the finitely many quotient resolutions of the systems $\lambda_i^2(s, z, h_2, g_1, h'_1, y, w, p, a)$ w.r.t. the wp -graded closure $GFCl$, that are not of maximal complexity, i.e., the complexity of each $Comp_i^3$ is strictly smaller than the complexity of $Comp^2$. The collection of the quotient resolutions $Comp_i^3$ is denoted by $GRem_3$.

And we continue the construction iteratively with each quotient resolution $Comp^3$ in $GRem_3$.

In light of the strict reduction in the complexities, the procedure must terminate after finitely many steps.

Definition 9.9. Let $d \geq 2$ be an integer. A (general) *proof system* is to make the following ordered choice:

1. To choose one of the terminal graded formal closures $WPH(h_1, w, p, a) * F_y$ in $GFMRD_1$.
2. For every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRemMRD_2$, to choose an integer $0 \leq n \leq M(WPHG)$, where $WPHG(g_1, h'_1, w, p, a)$ is the wp -rigid or wp -solid base group of the completed resolution $Comp^2 = Comp(\lambda^1 WPGRes)(z, h_1, y, w, p, a)$, and $M(WPHG)$ is the maximal possible number of exceptional families of specializations of $WPHG$ for a fixed value of the parameters $w_0, p_0 \in F_k$.
3. For every resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$ in $GRemMRD_2$ for which the integer n was chosen, and for every $1 \leq i \leq n$, to choose finitely many terminal graded formal closures $GFCl_1^i, \dots, GFCl_{m_i}^i$ in the graded formal MR diagram $GFMRD_2$ associated with the resolution $\lambda^1 WPGRes(h_1, y, w, p, a)$, where $1 \leq m_i \leq \Sigma \{M(Term(GFCl)) : GFCl \text{ is a terminal closure in } GFMRD_2\}$, and $M(Term(GFCl))$ is the maximal possible number of exceptional families of specializations of the wp -rigid or wp -solid base group $Term(GFCl)$ for a fixed value of the parameters $w_0, p_0 \in F_k$.
4. And by induction on $l \geq 2$, if for the resolution $Comp^l$ in $GRem_l$ who terminates in $WP(HG)^{l-1}$ the integer n was chosen, and if for some $1 \leq i \leq n$ the graded formal closure $GFCl$ in the diagram $GFMRD_l$ associated with $Comp^l$ was chosen, then, for every resolution $Comp^{l+1}$ in $GRem_{l+1}$ associated with $GFCl$ who terminates in $WP(HG)^l$, to choose an integer $0 \leq m \leq M(WP(HG)^l)$, and for every $1 \leq j \leq m$, to choose finitely many terminal graded formal closures $GFCl_1, \dots, GFCl_s$ in the diagram $GFMRD_{l+1}$ associated with $Comp^{l+1}$, where

$$1 \leq s \leq \Sigma \{M(Term(GFCl')) : GFCl' \text{ is a terminal closure in } GFMRD_{l+1}\}.$$

We say that a proof system PS as above, is of *depth* d , if d is the maximal integer with the property that some graded formal closure $GFCl$ listed in the proof system PS , is associated with a resolution $Comp^d$ in $GRem_d$.

We say that a proof system PS of depth d , *verifies* $(w_0, p_0) \in AE(w, p)$, if for every graded formal closure $GFCl$ listed in the proof system PS and is deepest w.r.t. the proof system PS , there exists an exceptional specialization f_0 of $Term(GFCl)$, so that:

1. The restrictions of all the specializations f_0 to the group $WPH(h_1, w, p, a)$ that was specified by the proof system PS , represent the same exceptional family of it.
2. For every resolution $Comp^l$ that was listed in PS and for all $1 \leq i \leq n$, where n is the chosen integer for $Comp^l$, all the induced specializations of the terminal group $WP(HG)^{l-1}$ of $Comp^l$ represent the same exceptional family of $WP(HG)^{l-1}$.
3. For every resolution $Comp^l$ that was listed in PS and for all $1 \leq i \leq n$, where n is the chosen integer for $Comp^l$, if the chosen collection of terminal graded formal closures in the diagram $GFMRD_l$ associated with $Comp^l$, is $GFCl_1, \dots, GFCl_s$, then the induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{s,0}$ form a covering closure for the induced ungraded resolution $Comp_0^l$.

4. For every resolution $Comp^l$ that was listed in PS and for all $1 \leq i_1 \neq i_2 \leq n$, where n is the chosen integer for $Comp^l$, the induced two exceptional families of the terminal group $WP(HG)^{l-1}$ of $Comp^l$ from the directions of i_1 and i_2 in the proof system PS , are distinct exceptional families of $WP(HG)^{l-1}$.
5. For every resolution $Comp^l$ that was listed in PS and for which the integer n was chosen, the terminal group $WP(HG)^{l-1}$ of $Comp^l$ admits exactly n distinct exceptional families corresponding to the value $w_0 p_0$ of the parameters.
6. Each of the specializations f_0 does not factor through any of the maximal limit groups $\lambda^t WPGL_j$ associated with the graded formal closure $GFCl$.

Such a tuple $((f_0), w_0, p_0, a)$, denoted for brevity (f_0, w_0, p_0, a) , that satisfies the conditions listed above, is called a *validPS statement* (of depth d).

By the construction of proof systems that we have explained in this section, there exist only finitely many proof systems. According to the construction that we have explained in this section, for every $p_0 \in EAE(p)$, there exists a witness $w_0 \in F_k$ and a proof system PS , so that the proof system PS verifies that $(w_0, p_0) \in AE(w, p)$. Similarly, for every group Γ in the model, and for every $p_0 \in EAE_\Gamma(p)$, there exist a witness $w_0 \in \Gamma$ and a proof system PS , so that PS verifies that $(w_0, p_0) \in AE(w, p)$.

We fix a proof system PS . We extend each validPS statement $((f_0), w_0, p_0, a)$ in the following two ways:

1. Let $Comp$ be one of the quotient resolutions listed in the proof system PS . Denote by $WPHG$ the wp -rigid or wp -solid base group of $Comp$. Assume that n is the integer chosen for the resolution $Comp$, and let $1 \leq i \leq n$. Let $GFCl_1, \dots, GFCl_s$ be the collection of terminal graded formal closures in the diagram $GFMRD_i$ (associated with $Comp$) that was chosen for $Comp$ and i . For every non-cyclic maximal (pegged) abelian subgroup A that appears in one of the decompositions along the various levels of $Comp$, and for each $j = 1, \dots, s$, one can find or associate a free abelian group \mathbb{Z}^{m_A} , a subgroup of finite index $C_A^j \leq \mathbb{Z}^{m_A}$ and a parameterized coset of C_A^j (with integer parameter). We denote by $lcm(A)$ the least common multiple of the indices of the subgroups C_A^1, \dots, C_A^s in \mathbb{Z}^{m_A} .

The value of a representative for the associated coset depends only on the (modular family of a) specialization of the base group $Term(GFCl_j)$. In other words, once we have fixed a specialization f_0^j for $Term(GFCl_j)$, the coset $C_{A,0}^j$ is determined. Actually, if the abelian group A does not appear in the terminal level of $Comp$ (i.e., A does not appear in the wp -graded JSJ of $WPHG$), then the coset C_A^j depends only on the graded closure $GFCl_j$ itself, and does not depend on the specific specialization f_0^j (i.e., $C_A^j = C_{A,0}^j$). Moreover, the entries of the elements in the coset $C_A^j = C_{A,0}^j$ correspond to the indices of the cyclic subgroups generated by the images of a fixed sub-basis of the abelian group A , in the maximal cyclic subgroup generated by the peg peg_A of A in the next level of the graded closure $GFCl_j$, where peg_A is a fixed generator for the cyclic edge group connecting A to the next level of $Comp$. However, in the case that the abelian group A appears in the terminal level of $Comp$, the entries of the elements in the coset $C_{A,0}^j$ correspond to the indices of the cyclic subgroups generated by the images of a fixed sub-basis of the abelian group A , in the maximal cyclic subgroup generated by the maximal root of the image of peg_A in the next level of the induced ungraded closure $GFCl_{j,0}$, under the specializations of $Comp$ that factor through the induced ungraded closure $GFCl_{j,0}$.

If further f_0^1, \dots, f_0^s are specializations of the base groups $Term(GFCl_1), \dots, Term(GFCl_s)$, and in addition the restrictions of f_0^1, \dots, f_0^s to the base group $WPHG$ of $Comp$ represent the same exceptional family, then in order for the collection of the induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{s,0}$ to form a covering closure for the corresponding ungraded resolution $Comp_0$, the union of the induced cosets $C_{A,0}^1, \dots, C_{A,0}^s$ must cover the entire group \mathbb{Z}^{m_A} for every maximal abelian group A that appears along the decompositions in the resolution $Comp$.

Hence, for each maximal abelian group A that appears along the decompositions in the resolution $Comp$, we add a new variable r_A that will be mapped to the image of peg_A when A does not appear in the terminal level of $Comp$, or to $Rpeg_{A,0}^q$ otherwise, where $Rpeg_{A,0}$ is the maximal root of the image of peg_A and q is co-prime to each of the indices of the subgroups C_A^j in \mathbb{Z}^{m_A} . In order for the collection of the induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{s,0}$ to form a covering closure for the corresponding ungraded resolution $Comp_0$, we must be able to extend each of the specializations $(r_{A,0}^{l_1}, \dots, r_{A,0}^{l_{m_A}})$ of the

fixed sub-basis of the abelian group A , for all $l_1, \dots, l_{m_A} = 0, \dots, lcm(A)$, to a specialization that factor through one of the ungraded closures $GFCl_{1,0}, \dots, GFCl_{s,0}$.

By the definition of a validPS statement, for every resolution $Comp^l$ that was listed in PS and for all $1 \leq i \leq n$, where n is the chosen integer for $Comp^l$, if the chosen collection of terminal graded formal closures in the diagram $GFMRD_l$ associated with $Comp^l$ is $GFCl_1, \dots, GFCl_s$, then the induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{s,0}$ form a covering closure for the induced ungraded resolution $Comp_0^l$. For each such $Comp^l$, and each $1 \leq i \leq n$, we add the corresponding specializations $r_{A,0}$ for the new variables r_A . Moreover, we add specializations for new variables $AbDem$. These specializations demonstrate that each of the (finitely many) corresponding specializations $(r_{A,0}^{l_1}, \dots, r_{A,0}^{l_{m_A}})$ of the fixed sub-bases of the maximal abelian groups A that appear in $Comp^l$, factors through one of the closures $GFCl_{1,0}, \dots, GFCl_{s,0}$.

2. We add specializations for new variables $SameFamily$ that demonstrates that the restrictions of all the specializations f_0 in a given validPS statement, to the group $WPH(h_1, w, p, a)$ that was specified by the proof system PS , represent the same modular family of it, and demonstrates that for every resolution $Comp^l$ that was listed in PS and for all $1 \leq i \leq n$, where n is the chosen integer for $Comp^l$, all the induced specializations of the terminal group $WP(HG)^{l-1}$ of $Comp^l$ represent the same modular family of $WP(HG)^{l-1}$.

We denote the extended validPS statements by the extended tuples $((SameFamily_0), (AbDem_0), (r_0), (f_0), w_0, p_0, a)$, or briefly by $(SameFamily_0, AbDem_0, r_0, f_0, w_0, p_0, a)$. We want to prove that the given $EAE(p)$ set, can be defined over F_k and over the random groups Γ by a uniform formula which belongs to the Boolean algebra of AE -formulas. For that, we start by collecting all the validPS statements and Γ -validPS statements in a finite uniform collection of limit groups.

We consider the collection \mathfrak{J} of all the specializations $((SameFamily_0), (AbDem_0), (r_0), (f_0), w_0, p_0, a)$ (over F_k) that satisfy all the conditions in the definition of validPS statement, except maybe that for some resolution $Comp^l$ that was listed in PS and for which the integer n was chosen, the terminal group $WP(HG)^{l-1}$ of $Comp^l$ admits more than n distinct exceptional families corresponding to the value $w_0 p_0$ of the parameters. The collection \mathfrak{J} factors through finitely many maximal limit groups $PS(SameFamily, AbDem, r, f, w, p, a)$.

Lemma 9.10.

1. Every validPS statement factors through the groups $PS(SameFamily, AbDem, r, f, w, p, a)$.
2. Let Γ be a group in the model. Every Γ -validPS statement factors through the groups $PS(SameFamily, AbDem, r, f, w, p, a)$.

Proof. The first claim follows directly from the definition of the groups $PS(SameFamily, AbDem, r, f, w, p, a)$. For the second one, let $((SameFamily_0), (AbDem_0), (r_0), (f_0), w_0, p_0, a)$ be a Γ -validPS statement. Let $((\widetilde{SameFamily_0}), (\widetilde{AbDem_0}), (\tilde{r}_0), (\tilde{f}_0), \tilde{w}_0, \tilde{p}_0, a)$ be a lift of $(SameFamily_0, (AbDem_0), r_0, f_0, w_0, p_0, a)$ to a specialization in F_k so that:

1. each \tilde{f}_0 is a specialization for the corresponding graded formal closure $GFCl$ listed in the proof system PS ,
2. for every resolution $Comp^l$ that was listed in PS and for all $1 \leq i \leq n$, where n is the chosen integer for $Comp^l$, the specializations $\widetilde{AbDem_0}$ demonstrates that each of the corresponding specializations \tilde{r}_0 of the fixed sub-bases of the maximal abelian groups that appear in $Comp^l$, factor through one of the induced ungraded closures $GFCl_{1,0}, \dots, GFCl_{s,0}$ (induced from the lifted specializations (\tilde{f}_0)).
3. The specializations $(\widetilde{SameFamily_0})$ demonstrates that the restrictions of all the specializations \tilde{f}_0 , to the group $WPH(h_1, w, p, a)$ that was specified by the proof system PS , represent the same modular family of it, and demonstrates that for every resolution $Comp^l$ that was listed in PS and for all $1 \leq i \leq n$, where n is the chosen integer for $Comp^l$, all the induced specializations of the terminal group $WP(HG)^{l-1}$ of $Comp^l$ represent the same modular family of $WP(HG)^{l-1}$.

Note that if some specialization \tilde{r}_0 equals to $\tilde{r}_0 = x_0^q$ for some non-trivial $x_0 \in F_k$, then its projection r_0 to Γ is also a q -th power (Γ is torsion-free and the corresponding ungraded resolutions are non-degenerate). Hence, and since all the lifts of a Γ -exceptional specialization of a given rigid or solid group to F_k is also exceptional, we deduce that the lifted specialization $\left((\widetilde{SameFamily}_0), (\widetilde{AbDem}_0), (\tilde{r}_0), (\tilde{f}_0), \tilde{w}_0, \tilde{p}_0, a \right)$ factors through the groups $PS(SameFamily, AbDem, r, f, w, p, a)$. \square

Finally, for every group $PS(SameFamily, AbDem, r, f, w, p, a)$, we associate its standard p -graded MR diagram, and consider all the resolutions $GRes(SameFamily, AbDem, r, f, w, p, a)$ in these diagrams, through which some specialization from the collection \mathfrak{J} factors. For each such resolution, we apply the method presented in section §8, and this implies the desired.

Theorem 9.11. *Let $\zeta(p)$ be an EAE formula*

$$\zeta(p) : \quad \exists w \forall y \exists x \quad \Sigma(x, y, w, p, a) = 1 \wedge \Psi(x, y, w, p, a) \neq 1.$$

Let $EAE(p)$ be the set of values $p = p_0 \in F_k$ defined by the formula $\zeta(p)$.

Let Γ be a group of level l in the model, and let $EAE_\Gamma(p)$ be the set of values $p = p_0 \in \Gamma$ defined by the formula $\zeta(p)$.

Then, there exists a formula $\varphi(p)$ in the Boolean algebra of AE-formulas, so that:

1. *The set $EAE(p)$ is defined by the formula $\varphi(p)$.*
2. *If l is large enough, then the set $EAE_\Gamma(p)$ is defined by the formula $\varphi(p)$.*

Theorem 9.12. *Let $\zeta V(p)$ be an EAE formula*

$\zeta V(p) :$

$$\exists w \forall y \exists x \quad (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1) .$$

Let $EAEV(p)$ be the set of values $p = p_0 \in F_k$ defined by the formula $\zeta V(p)$.

Let Γ be a group of level l in the model, and let $EAEV_\Gamma(p)$ be the set of values $p = p_0 \in \Gamma$ defined by the formula $\zeta V(p)$.

Then, there exists a formula $\varphi(p)$ in the Boolean algebra of AE-formulas, so that:

1. *The set $EAEV(p)$ is defined by the formula $\varphi(p)$.*
2. *If l is large enough, then the set $EAEV_\Gamma(p)$ is defined by the formula $\varphi(p)$.*

Proof. For proving theorem 9.11, we have constructed finitely many proof systems PS for the formula $\zeta(p)$ that encode validPS statements and Γ -validPS statements for all the values $p = p_0$ for which the sentence $\zeta(p_0)$ is as truth sentence over the corresponding group.

For constructing a sentence $\varphi(p)$ that defines the sets $EAEV(p)$ and $EAEV_\Gamma(p)$, we construct finitely many proof systems for the formula $\zeta V(p)$, in a similar way to the one explained in this section.

The only change that we should perform in the construction for this case, is that instead of constructing (during the corresponding steps) a graded formal MR diagram of the system $\Sigma(x, y, w, p, a)$ for each of the resolutions of the graded MR diagram constructed in the previous step, we construct, for each $j = 1, \dots, r$, the graded formal MR diagram of the system $\Sigma_j(x, y, w, p, a)$ for each of the resolutions of the graded MR diagram constructed in the previous step. Then, the graded MR diagrams in the next step, corresponds to the specializations for which there is no formal solution that verifies the corresponding set of inequalities $\Psi_j(x, y, w, p, a) \neq 1$. \square

10 Truth Sentences of Minimal Rank over Random Groups

Lemma 10.1. *Let $V(p)$ be a formula in the Boolean algebra of AE-formulas.*

Then, there exists a formula $V'(p)$ which is a finite union of EA-formulas, AE-formulas, and intersections of an EA-formula and an AE-formula, so that $V(p)$ and $V'(p)$ are tautologically equivalent, i.e., for every group Γ , the formulas $V(p)$ and $V'(p)$ define the same set over Γ .

Proof. A finite union or a finite intersection, of AE -formulas is tautologically an AE -formula. Similarly, A finite union or a finite intersection, of EA -formulas is tautologically an EA -formula. \square

Lemma 10.2. *The projection of an EA -formula is tautologically (over any group) an EA -formula. The projection of the intersection of an EA -formula and an AE -formula, is tautologically a projection of an AE -formula.*

Proof. For the first part of the statement, let $EA(w, p)$ be a given EA -formula:

$EA(w, p) :$

$$\exists y \forall x (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1) .$$

The projection of $EA(w, p)$ is by definition the following formula

$\pi EA(p) :$

$$\exists w \exists y \forall x (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1) ,$$

which is an EA -formula.

Now let $V(w, p)$ be the intersection of an an EA -formula and an AE -formula. Then, $V(w, p)$ has the form:

$V(w, p) :$

$$(\forall y \exists x (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1)) \wedge (\exists t \forall u (\Sigma'_1(u, t, w, p, a) = 1 \wedge \Psi'_1(u, t, w, p, a) \neq 1) \vee \dots \vee (\Sigma'_{r'}(u, t, w, p, a) = 1 \wedge \Psi'_{r'}(u, t, w, p, a) \neq 1)) .$$

Hence, the projection of $V(w, p)$ is given by:

$\pi V(p) :$

$$\exists w (\forall y \exists x (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1)) \wedge (\exists t \forall u (\Sigma'_1(u, t, w, p, a) = 1 \wedge \Psi'_1(u, t, w, p, a) \neq 1) \vee \dots \vee (\Sigma'_{r'}(u, t, w, p, a) = 1 \wedge \Psi'_{r'}(u, t, w, p, a) \neq 1)) ,$$

which is tautologically equivalent to:

$$\exists w \exists t \forall y \forall u \exists x ((\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1)) \wedge ((\Sigma'_1(u, t, w, p, a) = 1 \wedge \Psi'_1(u, t, w, p, a) \neq 1) \vee \dots \vee (\Sigma'_{r'}(u, t, w, p, a) = 1 \wedge \Psi'_{r'}(u, t, w, p, a) \neq 1)) .$$

Hence, iterative replacements of intersections of the form $A \wedge (B \vee C)$, by unions of the form $(A \wedge B) \vee (A \wedge C)$, in the formula

$$((\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1)) \wedge ((\Sigma'_1(u, t, w, p, a) = 1 \wedge \Psi'_1(u, t, w, p, a) \neq 1) \vee \dots \vee (\Sigma'_{r'}(u, t, w, p, a) = 1 \wedge \Psi'_{r'}(u, t, w, p, a) \neq 1)) ,$$

implies the required. \square

Lemma 10.3. *Let $V(w, p)$ be a formula in the Boolean algebra of AE -formulas.*

Then, there exists finitely many AE -formulas $V_1(w, p), \dots, V_s(w, p)$, and an EA -formula $H(p)$, so that the projection of $V(w, p)$ is tautologically equivalent to the union of $H(p)$ with the projections of the AE -formulas $V_1(w, p), \dots, V_s(w, p)$:

$$\pi V(p) : \quad H(p) \vee \pi V_1(p) \vee \dots \vee \pi V_s(p) .$$

Proof. According to lemma 10.1, the formula $V(w, p)$ is a finite union of EA -formulas, AE -formulas, and intersections of an EA -formula and an AE -formula.

Hence, the projection of the formula $V(w, p)$ is tautologically the union of the projections of these EA -formulas, AE -formulas, and the intersections of an EA -formula and an AE -formula.

According to lemma 10.2, and since the union of EA -formulas is tautologically an EA -formula, we obtain the required. \square

Definition 10.4. Let $V(w, p)$ be a formula in the Boolean algebra of AE -formulas. Let $V_1(w, p), \dots, V_s(w, p)$, and $H(p)$ be as in 10.3. We call the formula $V(w, p)$ a *minimal rank formula*, if all the graded limit groups along the construction of the proof systems for each of the EAE -formulas $\pi V_1(p), \dots, \pi V_s(p)$, are of minimal rank (see section §9). Note that the construction of proof systems depends only on the fixed free group F_k and the given AE -formula.

Theorem 10.5. *Let $V(w, p)$ be a minimal rank formula in the Boolean algebra of AE -formulas. Let $\pi V(p)$ be the projection of $V(w, p)$. Then, there exists a formula $R(p)$ in the Boolean algebra of AE -formulas, so that:*

1. The formulas $\pi V(p)$ and $R(p)$ define the same set over F_k .
2. Let Γ be a random group. With overwhelming probability, the formulas $\pi V(p)$ and $R(p)$ define the same set over Γ .

Proof. Let $V_1(w, p), \dots, V_s(w, p)$, and $H(p)$ be as in lemma 10.3, and write

$$\pi V(p) : \quad H(p) \vee \pi V_1(p) \vee \dots \vee \pi V_s(p).$$

According to theorem 9.12, for every $i = 1, \dots, s$, there exists a formula $\varphi_i(p)$ in the Boolean algebra of AE -formulas, so that:

1. The formulas $\pi V_i(p)$ and $\varphi_i(p)$ define the same set over F_k .
2. The formulas $\pi V_i(p)$ and $\varphi_i(p)$ define the same set over Γ (with overwhelming probability).

Hence, the formula

$$R(p) : \quad H(p) \vee \varphi_1(p) \vee \dots \vee \varphi_s(p),$$

is as required. \square

Definition 10.6. Let ψ be a first order sentence.

1. If ψ belongs to the Boolean algebra of AE -sentences, then, we say that ψ is a *minimal rank sentence*, if ψ , as a formula with empty set of parameters, is a minimal rank formula (definition 10.4).
2. If ψ is an EAE -sentence, then ψ is called a minimal rank sentence if it is the projection of a minimal rank formula.
3. If ψ is an AEA -sentence, then ψ is called a minimal rank sentence if $\neg\psi$ is a minimal rank sentence.
4. Assume that the sentence ψ has the (prenex) normal form

$$\forall y_1 \exists x_1 \dots \exists x_{n-1} \forall y_n \exists x_n \\ (\Sigma_1(x_i, y_i, w, a) = 1 \wedge \Psi_1(x_i, y_i, w, a) \neq 1) \vee \dots \vee (\Sigma_r(x_i, y_i, w, a) = 1 \wedge \Psi_r(x_i, y_i, w, a) \neq 1).$$

We denote $y = y_n, x = x_n, w = x_{n-1}$, and we denote by p the union of the variables sets y_1, x_1, \dots, y_{n-1} . We consider the AE -formula

$$V(w, p) : \\ \forall y \exists x (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1),$$

and we assume that $V(w, p)$ is a minimal rank formula (definition 10.4). Note that

$$\psi = \forall y_1 \exists x_1 \dots \forall y_{n-1} \pi V(p).$$

Let $R(p)$ be the formula obtained from theorem 10.5 corresponding to $V(w, p)$. In light of theorem 10.5, the sentence ψ is equivalent over F_k and over random groups (with overwhelming probability) to the sentence

$$\psi' : \quad \forall y_1 \exists x_1 \dots \forall y_{n-1} R(p).$$

Since $R(p)$ belongs to the Boolean algebra of AE -formulas, ψ' is tautologically equivalent to a sentence of the form

$$\psi' = \forall y_1 \exists x_1 \dots \forall y_{n-1} \forall u \exists t \forall z R'(p, u, t, z),$$

(note the consecutive universal quantifiers $\forall y_{n-1} \forall u$). The sentence ψ is called a minimal rank sentence if ψ' is a minimal rank sentence (defined by induction).

5. Assume that the sentence ψ has the normal form

$$\begin{aligned} & \exists y_1 \forall x_1 \dots \forall x_{n-1} \exists y_n \forall x_n \\ & (\Sigma_1(x_i, y_i, w, a) = 1 \wedge \Psi_1(x_i, y_i, w, a) \neq 1) \vee \dots \vee (\Sigma_r(x_i, y_i, w, a) = 1 \wedge \Psi_r(x_i, y_i, w, a) \neq 1). \end{aligned}$$

We denote $y = y_n$, $x = x_n$, $w = x_{n-1}$, and we denote by p the union of the variables sets y_1, x_1, \dots, y_{n-1} . We consider the AE -formula

$V(w, p)$:

$$\neg (\exists y \forall x (\Sigma_1(x, y, w, p, a) = 1 \wedge \Psi_1(x, y, w, p, a) \neq 1) \vee \dots \vee (\Sigma_r(x, y, w, p, a) = 1 \wedge \Psi_r(x, y, w, p, a) \neq 1)),$$

and we assume that $V(w, p)$ is a minimal rank formula (definition 10.4). Note that

$$\psi = \exists y_1 \forall x_1 \dots \exists y_{n-1} \neg \pi V(p).$$

In a similar way to the previous point of the definition, using $V(w, p)$ and theorem 10.5, we construct a sentence ψ' that is equivalent to ψ over F_k and over random groups, so that ψ' is of the form

$$\psi' = \exists y_1 \forall x_1 \dots \exists y_{n-1} \neg (\forall u \exists t \forall z R'(p, u, t, z)).$$

And we call the sentence ψ a minimal rank sentence if ψ' is a minimal rank sentence (defined by induction).

Theorem 10.7. *Let ψ be a first order sentence of minimal rank. Then, there exists a minimal rank sentence $R(\psi)$ that belongs to the Boolean algebra of AE -sentences, so that:*

1. *The sentences ψ and $R(\psi)$ are equivalent over F_k .*
2. *Let Γ be a random group. With overwhelming probability, the sentences ψ and $R(\psi)$ are equivalent over Γ .*

Theorem 10.8. *Let ψ be a minimal rank sentence in the Boolean algebra of AE -sentences. Let Γ be a random group. Then, ψ is a truth sentence over F_k if and only if ψ is a truth sentence over Γ .*

Proof. If ψ is an EA -sentence, and ψ is true over F_k , then, according to the first paper, ψ is a truth sentence over Γ . Hence, it suffices to prove that if ψ is an AE -sentence that is true over F_k , then ψ is true over Γ . Indeed, we construct the proof system for the sentence ψ (as a EAE -formula with empty set of parameters) over F_k . According to the construction of proof systems in section §9, the proof system of ψ constructed over F_k , validates the truthiness of ψ over Γ (with overwhelming probability). \square

Theorem 10.9. *Let ψ be a first order sentence of minimal rank. Let Γ be a random group. Then, ψ is a truth sentence over F_k if and only if ψ is a truth sentence over Γ .*

Proof. According to theorem 10.7, there exists a sentence ψ' in the Boolean algebra of AE -sentences that is equivalent to ψ over F_k and over Γ . According to theorem 10.8, ψ' is true over F_k if and only if it is true over Γ . Hence, ψ is true over F_k if and only if ψ is true over Γ . \square

References

- [Kh-My] Kharlampovich, Olga; Myasnikov, Alexei (2006). “Elementary theory of free non-abelian groups”. *Journal of Algebra*. 302 (2): 451–552. <https://doi.org/10.1016/j.jalgebra.2006.03.033>
- [O11] Ollivier, Y. “Sharp phase transition theorems for hyperbolicity of random groups”. *Geom. funct. anal.* 14, 595–679 (2004). <https://doi.org/10.1007/s00039-004-0470-y>
- [O12] Yann Ollivier. “Some small cancellation properties of random groups”. *International Journal of Algebra and Computation*, 2007, 17, pp.37-51.
- [Ri-Se] Rips, E., Sela, Z. “Structure and rigidity in hyperbolic groups I”. *Geometric and Functional Analysis* 4, 337–371 (1994). <https://doi.org/10.1007/BF01896245>
- [Se1] Z. Sela, “Diophantine geometry over groups I: Makanin-Razborov diagrams”, *Publications Mathématiques de l’IHES* 93(2001), 31-105.
- [Se2] Z. Sela, “Diophantine geometry over groups II: Completions, closures and formal solutions”, *Israel Jour. of Math.* 134(2003), 173-254.
- [Se3] Z. Sela, “Diophantine geometry over groups III: Rigid and solid solutions”, *Israel Jour. of Math.* 147(2005), 1-73.
- [Se4] Z. Sela, “Diophantine geometry over groups IV: An iterative procedure for validation of a sentence”, *Israel Jour. of Math.* 143(2004), 1-130.
- [Se5] Z. Sela, “Diophantine geometry over groups V: Quantifier elimination”, divided into two parts - *Israel Jour. of Math.* 150(2005), 1-197, and *GAF* 16(2006), 537-706.
- [Se6] Z. Sela, “Diophantine geometry over groups V2: quantifier elimination II.” *GAF*, *Geom. funct. anal.* 16, 537–706 (2006).
- [Se7] Z. Sela, “Diophantine geometry over groups VI: the elementary theory of a free group.” *GAF*, *Geom. funct. anal.* 16, 707–730 (2006).
- [Se8] Z. Sela, “Diophantine geometry over groups. VII. The elementary theory of a hyperbolic group”, *Proc. Lond. Math. Soc.* (3) 99(1), 217–273 (2009).