

APPROXIMATION OF SHAPE OPTIMIZATION PROBLEMS WITH NON-SMOOTH PDE CONSTRAINTS: AN OPTIMALITY CONDITIONS POINT OF VIEW

L. BETZ*

Abstract. This paper is concerned with a shape optimization problem governed by a non-smooth PDE, i.e., the nonlinearity in the state equation is not necessarily differentiable. We follow the functional variational approach of [40] where the set of admissible shapes is parametrized by a large class of continuous mappings. This methodology allows for both boundary and topological variations. It has the advantage that one can rewrite the shape optimization problem as a control problem in a function space. To overcome the lack of convexity of the set of admissible controls, we provide an essential density property. This permits us to show that each parametrization associated to the optimal shape is the limit of global optima of non-smooth distributed optimal control problems. The admissible set of the approximating minimization problems is a convex subset of a Hilbert space of functions. Moreover, its structure is such that one can derive strong stationary optimality conditions [6]. The present manuscript provides the basis for the investigations from [5], where necessary conditions in form of an optimality system have been recently established.

Key words. Optimal control of non-smooth PDEs, shape optimization, topological variations, functional variational approach, approximation scheme of fixed domain type

AMS subject classifications. 49Q10, 35Q93, 49N99.

1. Introduction. Shape optimization problems arise in many engineering applications [7] and are often formulated as minimization problems governed by one or more PDEs or VIs [35]. Their main particularity is the fact that these equations are solved on unknown domains [31, 35]. The goal is to find the optimal shape, i.e., that domain for which the distance to a certain desired state is minimized. Thus, shape optimization problems exhibit similarities with optimal control problems [42], the essential difference and difficulty being that the admissible control set consists of variable geometries. Such problems are highly nonconvex, which makes their investigation challenging both from the theoretical and the computational point of view.

There is an enormous amount of literature devoted to the study of optimal design problems, see e.g. the classical contributions [31, 35] and the references therein. To keep the depiction concise, we just focus on the works that deal with these problems at a theoretical level, with an emphasis on optimality systems in qualified form. These resemble the classical Karush-Kuhn-Tucker conditions, and in the case of optimal control problems, they involve an adjoint equation. In [34], topology and boundary variations are combined for the first time in order to derive necessary optimality conditions. In [2], the considered shape and topology optimization problem is governed by a cone constraint and the existence of Lagrange multipliers is shown, the respective optimality conditions being expressed in the form of a complementarity system. Methods for computing the shape derivative of the cost functional without resorting to the differentiability of the control-to-state map have been developed in [15, 17, 36]. The techniques therein are applied for shape optimization problems governed by smooth PDEs.

Shape optimization problems with non-smooth constraints have been investigated at a theoretical level mostly with respect to existence of optimal shapes [8, 9] and

*Faculty of Mathematics, University of Wuerzburg, Germany

sensitivity analysis [12,25,33,35]. While there is an increasing number of contributions concerning optimal shape design problems governed by VIs, see [8,9,12,19,20,25] and the references therein, there are no papers known to the author that address the case where the governing equation is a *non-smooth PDE*. In [19,20], the authors resort to smoothing techniques and optimality systems in qualified form are obtained just for the smoothed problem [19], or, for the original problem [20], but only if certain assumptions are imposed on the converging sequences. If smoothing is not involved, optimality conditions for the non-smooth shape optimization problem do not involve an adjoint equation, unless linearity w.r.t. direction is assumed [25]. Otherwise, these are stated just in a primal form [12], i.e., the respective optimality condition only asserts the non-negativity of the shape derivative of the reduced objective functional in feasible directions.

As in most of the literature, the approaches in all the aforementioned papers (see also the references therein) are based on variations of the geometry. One of the most common notions in this context are the shape derivative [35] and the topological derivative [29].

A more novel technique to derive optimality conditions in qualified form, where general functional variations instead of geometrical ones are involved, can be found in [41]. Therein, an optimal design problem governed by a linear PDE with Neumann boundary conditions is investigated. By means of the implicit parametrization theorem [28,38] combined with Hamiltonian systems [39], the equivalence of the shape and topology optimization problem with an optimal control problem in function space is established, provided that the set of admissible geometries is generated by a certain class of continuous functions. The respective control problem in function space is amenable to the derivation of optimality conditions by a classical Lagrange multipliers approach. The same functional variational method will be adopted (to some extent) in the present work.

The aim of this paper is to provide a first essential step towards the derivation of optimality systems for the optimal shape associated to the following *non-smooth* shape optimization problem

$$\left. \begin{array}{l} \min_{\Omega \in \mathcal{O}, E \subset \Omega} \int_E (y_\Omega(x) - y_d(x))^2 dx + \alpha \int_\Omega dx, \\ \text{s.t.} \quad -\Delta y_\Omega + \beta(y_\Omega) = f \quad \text{a.e. in } \Omega, \\ \quad \quad y_\Omega = 0 \quad \text{on } \partial\Omega. \end{array} \right\} \quad (P_\Omega)$$

The main particularities of (P_Ω) are:

- the non-smooth character of β , which is supposed to be locally Lipschitz continuous and directionally differentiable, but *not* necessarily *differentiable*, e.g., $\beta = \max\{\cdot, 0\}$ (we underline that we do not intend to replace β by a differentiable function);
- the fact that the governing PDE is solved on the unknown (variable) domain Ω (which plays the role of the control).

The admissible control set \mathcal{O} consists of (unknown) subdomains of a given, fixed domain D . These are the so-called *admissible shapes*. In this manuscript, they are generated by a class of continuous functions, see Definition 2.5. The holdall domain $D \subset \mathbb{R}^2$ is bounded, of class $C^{1,1}$, and $f \in L^2(D)$. In the current work, the non-smoothness β can be thought of as a piecewise linear mapping. In [6], which

specifically deals with the lack of differentiability (see below), we will consider much more general situations. The symbol $-\Delta : H_0^1(D) \rightarrow H^{-1}(D)$ denotes the Laplace operator in the distributional sense; note that $H_0^1(D)$ is the closure of the set $C_c^\infty(D)$ w.r.t. the $H^1(D)$ -norm. The desired state y_d is an L^2 -function, which is defined on the observation set E (Assumption 2.3). The parameter α appearing in the objective is supposed to satisfy $\alpha \geq 0$.

The main purpose of this manuscript is to develop an approximation scheme for (P_Ω) by means of more approachable optimal control problems where the underlying control space consists of functions, see (P_ε) below. We show that each parametrization associated to the optimal shape of (P_Ω) is the limit of global minimizers of (P_ε) (Corollary 5.4). The approximating control problem is governed by a non-smooth PDE in the holdall domain D . Its structure is such that, under certain requirements on the given data, it permits the derivation of strong stationary optimality systems [6], i.e., systems that are equivalent to the first-order necessary optimality condition. In the recent work [5] we carry out the passage to the limit $\varepsilon \rightarrow 0$ in these optimality conditions.

The starting point of our analysis is the functional variational method introduced by [26], where the admissible set \mathcal{O} is generated by a large class of functions [40]. We point out that this is different from the level set method of [30]. It allows one to switch from the shape optimization problem to a control problem in a function space, the latter being more amenable to a rigorous mathematical investigation. This technique involves the description of the boundary of the unknown domain as the solution to a Hamiltonian system and it is based on the implicit function theorem [28, 38] and Poincaré-Bendixson theory [14, Ch.10]. The idea that the admissible shapes are parametrized by so called shape functions turned out to be successful in various papers [11, 21–24, 26, 27, 40, 41]. The variations used in all these works have no prescribed geometric form (as usual in the literature) and the methodology therein provides a unified analytic framework allowing for both boundary and topological variations. Penalization methods were developed in [40] and extended in [21, 22]. We mention in particular [22] where the shape and topology optimization problem governed by a PDE with Neumann boundary conditions is equivalently rewritten as a control problem in a function space where the state equation is solved on the fixed domain while the boundary condition appears penalized in the objective. The functional variational approach was also the fundament for the derivation of optimality conditions for optimal shape design problems governed by linear PDEs with Neumann boundary conditions in [41]. We also refer to the more recent contributions [24] (clamped plates) and [23] (Navier-Stokes equations).

Given a continuous function g on the holdall domain D , we define

$$\Omega_g := \text{int}\{x \in D : g(x) \leq 0\}. \quad (1.1)$$

To ensure that $E \subset \Omega_g$, one requires

$$g \leq 0 \quad \text{in } E.$$

Two other key properties which need to be imposed on the parametrization (of the unknown domain) are

$$|\nabla g| + |g| > 0 \quad \text{in } D, \quad g > 0 \quad \text{on } \partial D.$$

If $g \in C^2(\bar{D})$, these imply that $\partial\Omega_g$ is a finite union of closed disjoint C^2 curves, without self intersections, that do not intersect ∂D [40, Prop. 2]. Together, the above conditions give rise to the definition of the set of admissible shape functions, called \mathcal{F}_{sh} , see (2.3) and Remark 2.7 below. We note that Ω_g may have many connected components. The admissible shape (domain) that we use in the definition of \mathcal{O} below, see (2.4), is the component that contains the subdomain E . Since this may not be simply connected, the approach we discuss in this paper is related to topological optimization too.

By proceeding like this, we arrive at a reformulation of (P_Ω) in terms of a control problem in a function space. This reads as follows

$$\left. \begin{array}{l} \min_{g \in \mathcal{F}_{\text{sh}}} \int_E (y_g(x) - y_d(x))^2 dx + \alpha \int_D 1 - H(g) dx, \\ \text{s.t.} \quad -\Delta y_g + \beta(y_g) = f \quad \text{in } \Omega_g, \\ \quad \quad \quad y_g = 0 \quad \text{on } \partial\Omega_g, \end{array} \right\} \quad (P)$$

where $H : \mathbb{R} \rightarrow \{0, 1\}$ is the Heaviside function, see (2.5) below.

We underline the difficulty of the problem (P) . The application of traditional optimization methods is excluded, as we deal with a control problem governed by a non-smooth PDE with the additional challenges that the admissible set \mathcal{F}_{sh} is non-convex, while the control does not appear on the right-hand side of the non-smooth PDE, but in the definition of the variable domain on which this is solved. Even in the case of classical control problems (where the domain is fixed), the non-smooth character is particularly challenging, as the standard KKT theory cannot be directly employed if the differentiability of the control-to-state map is not available. When we exclude the prominent smoothening techniques from [3], the derivation of (strong stationary) optimality conditions for non-smooth control problems in the presence of control constraints [4, 43] is restricted to certain situations (so-called 'constraint qualifications'). As far as we know, there isn't any literature on the topic of optimality conditions for non-smooth optimal control problems with non-convex admissible set that does not involve smoothening. We emphasize that we do not intend to resort to such techniques. In the subsequent papers [5, 6] that continue the findings from the present manuscript, all the problems examined ((P_Ω) , (P) and (P_ε)) feature the same non-differentiable mapping β .

In order to deal with the control problem (P) , and thus with the shape optimization problem (P_Ω) , we introduce an approximation scheme of fixed domain type [26, 27]. That is, we extend the state equation on the whole reference domain D . We only smoothen the Heaviside function, since this has jumps, unlike the non-smoothness β which is supposed to be continuous. By proceeding like this, we preserve the non-smooth character, and arrive, for $\varepsilon > 0$ small, fixed, at the following approximating optimal control problem:

$$\left. \begin{array}{l} \min_{g \in \mathcal{F}} \int_E (y(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(g))(x) dx + \frac{1}{2} \|g - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2 \\ \text{s.t.} \quad -\Delta y + \beta(y) + \frac{1}{\varepsilon} H_\varepsilon(g)y = f + \varepsilon g \quad \text{in } D, \\ \quad \quad \quad y = 0 \quad \text{on } \partial D. \end{array} \right\} \quad (P_\varepsilon)$$

In the objective of (P_ε) , \bar{g}_{sh} is a local optimum of (P) which we intend to approximate by local optima of (P_ε) (adapted penalization [3]), in the sense that \bar{g}_{sh} will arise as

the limit of a sequence of local minima of (P_ε) (Theorem 5.2). The mapping H_ε is the regularization of the Heaviside function, cf. (4.2) below. We underline the fact that we replaced the non-convex set \mathcal{F}_{sth} with a convex subset of the Hilbert-space $\mathcal{W} := L^2(D) \cap H^s(D \setminus \bar{E})$, $s \in (1, 2]$, namely

$$\mathcal{F} := \{g \in \mathcal{W} : g \leq 0 \text{ a.e. in } E\}. \quad (1.2)$$

The formulation of (P_ε) is slightly reminiscent of the minimization problem recently studied in [4] with the additional feature that the control enters the state equation in a non-linear fashion (via the mapping H_ε). However, the structure of (P_ε) allows us to state conditions on the given data under which the derivation of a complete (so-called strong stationary) optimality system is possible [6]. For more comments regarding the particularities of (P_ε) , see section 4.

Our main goal is to show that local optima of (P) can be approximated by local optima of (P_ε) (Theorem 5.2). One of the fundamental challenges will arise from the fact that in (P_ε) we replaced the non-convex set \mathcal{F}_{sth} from the original problem (P) by \mathcal{F} . To bridge this gap, we prove that \mathcal{F}_{sth} is dense in \mathcal{F} w.r.t. the $L^2(D)$ -norm (section 3.1). Since the control-to-state operator of (P) is defined on \mathcal{F}_{sth} only, not on \mathcal{F} , the meanwhile standard arguments of the adapted penalization method [3] must be accordingly changed. This requires establishing certain convergences of the control-to-state operator $g \mapsto y_g$ of (P) (subsection 4.2), which are more difficult to obtain compared to the classical case where the domain is fixed and the control is distributed.

The involved techniques, especially the density $\mathcal{F}_{\text{sth}} \xrightarrow{d} \mathcal{F}$, may be employed in the study of other similar problems (Remark 3.2).

The manuscript is organized as follows. After introducing the precise assumptions at the beginning of section 2, subsection 2.1 deals with the definition of the admissible set \mathcal{O} . Then, in subsection 2.2, we reformulate the non-smooth shape optimization problem (P_Ω) as a non-smooth control problem in a function space, i.e., as (P) . Here, we also introduce the notion of local optimum for the latter. Section 3 is dedicated to the density property $\mathcal{F}_{\text{sth}} \xrightarrow{d} \mathcal{F}$, which is essential for making the connection between (P) and the approximating non-smooth control problem with fixed domain (P_ε) . This is introduced in section 4. In subsection 4.1 we show that the unique solution of the state equation in (P) is the limit of solutions to state equations in (P_ε) . Subsection 4.2 is dedicated to the convergence properties of the control-to-state maps of (P) and (P_ε) . In subsection 4.3 we establish that (P_ε) admits optimal solutions. Finally, section 5 contains the main results of this paper, namely Theorem 5.2 and Corollary 5.4. Theorem 5.2 states that the approximation of (P) by (P_ε) is meaningful in the sense that local minima of (P) arise as limits of local optima of (P_ε) . As a consequence, each parametrization of the optimal shape of (P_Ω) is approximable by global minimizers of (P_ε) (Corollary 5.4). The findings in this last section provide the basis for the investigations with respect to the derivation of qualified optimality systems for (P_Ω) , see [5].

2. The shape optimization problem. We begin this section by stating the precise assumptions on the non-smoothness β appearing in the state equation in (P_Ω) . We also mention the requirements needed for the observation set E . Then, in the upcoming subsections we will introduce the set of admissible shapes \mathcal{O} for (P_Ω) . This will allow us to 'rewrite' (P_Ω) as an optimal control problem in a function space (Proposition 2.11).

ASSUMPTION 2.1 (The non-smoothness). *The function $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is monotone increasing and locally Lipschitz continuous in the following sense: For all $M > 0$, there exists a constant $L_M > 0$ such that*

$$|\beta(z_1) - \beta(z_2)| \leq L_M |z_1 - z_2| \quad \forall z_1, z_2 \in [-M, M].$$

REMARK 2.2. *At this stage, we do not make any assumptions concerning the limited differentiability properties of β that were advertised in the introduction. These do not play any role in the present paper, but in [6], which is a continuation of the present work. We just mention that, therein, β is supposed to be semi-differentiable only, i.e., the right and left side derivative of β may not be equal at certain points, e.g. $\beta = \max\{\cdot, 0\}$ or $\beta = |\cdot|$. This is precisely one of the main challenges in [6].*

By Assumption 2.1, it is straight forward to see that the Nemytskii operator $\beta : L^\infty(D) \rightarrow L^\infty(D)$ is well-defined. Moreover, this is Lipschitz continuous on bounded sets in the following sense: for every $M > 0$, there exists $L_M > 0$ so that

$$\|\beta(y_1) - \beta(y_2)\|_{L^q(D)} \leq L_M \|y_1 - y_2\|_{L^q(D)} \quad \forall y_1, y_2 \in \overline{B_{L^\infty(D)}(0, M)}, \quad \forall 1 \leq q \leq \infty. \quad (2.1)$$

ASSUMPTION 2.3 (The observation set). *The set $E \subset D$ is a subdomain with boundary of measure zero. Moreover, $\text{dist}(\overline{E}, \partial D) > 0$, where*

$$\text{dist}(\overline{E}, \partial D) := \inf_{(x_1, x_2) \in \overline{E} \times \partial D} \text{dist}(x_1, x_2).$$

We point out that the last condition in Assumption 2.3 ensures that the set of admissible shape functions \mathcal{F}_{sh} is not empty, cf. (2.3) below.

In the rest of the paper, one tacitly supposes that Assumptions 2.1 and 2.3 are always fulfilled without mentioning them every time.

2.1. The admissible set. As pointed out in the introduction, we will work in the framework of [21, 22, 26, 27, 40, 41] (see also the references therein), where the admissible set \mathcal{O} consists of a family of subdomains of D that are generated by a certain class of continuous functions, called \mathcal{F}_{sh} , see (2.3) below. For a general mapping $g \in C(\overline{D})$, we will use the notation Ω_g to describe the following open subset of the holdall domain D :

$$\Omega_g := \text{int}\{x \in D : g(x) \leq 0\}. \quad (2.2)$$

Note that Ω_g is not necessarily connected and the choice of g is not unique, in fact there are infinitely many functions generating the same Ω_g , see for instance the end of the proof of Lemma 2.10. While Ω_g is an open set and may have many connected components, the admissible shape (domain) that we use in the definition of \mathcal{O} below, see (2.4), is the component that contains the subdomain E . Its existence is guaranteed by imposing that the parametrization g satisfies $g \leq 0$ in E .

DEFINITION 2.4 (The set of admissible shape functions). *We define*

$$\mathcal{F}_{\text{sh}} := \{g \in C^2(\overline{D}) : g(x) \leq 0 \quad \forall x \in E, \quad |\nabla g(x)| + |g(x)| > 0 \quad \forall x \in D, \quad (2.3) \\ g(x) > 0 \quad \forall x \in \partial D, \quad g(x) > 0 \quad \forall x \notin \overline{\Omega_g}\}.$$

\mathcal{F}_{sh} will be later the control set for the optimization problem (P) below, as we will switch from the topology of subsets in the Euclidean space to the one in a function

space; note that (P_Ω) and (P) are 'equivalent' in the sense of Proposition 2.11. Instead of C^2 functions, we may take C^k mappings in (2.3), $k \in \mathbb{N}_+$ (this is possible due to Proposition 3.1). Switching to C^k shape functions impacts only the regularity of the boundary of Ω_g , which is as smooth as g is (Lemma 2.9). Since we want to keep the admissible set \mathcal{F}_{sh} as large as possible and at the same time ensure the H^2 regularity of the state y_g in (P) , we choose to stick to the C^2 -setting.

DEFINITION 2.5 (The set of admissible domains). *The admissible set for the shape optimization problem (P_Ω) is given by*

$$\mathcal{O} := \{\text{the component of } \Omega_g \text{ that contains the set } E : g \in \mathcal{F}_{\text{sh}}\}. \quad (2.4)$$

We note that the family \mathcal{O} is rich, it consists of domains that may not be simply connected, that is, topological and boundary variations are both included in our investigations.

REMARK 2.6 (The relevant component). *We point out that, for each $g \in \mathcal{F}_{\text{sh}}$, the existence of a component of Ω_g that contains E is guaranteed by the inequality $g \leq 0$ in E , cf. (2.3). In fact, this is the only component of Ω_g that is relevant when solving the state equation in (P_Ω) , since, in the objective, the integral term involving the state is taken over E . In other words, it doesn't make sense to solve the state equation on components of Ω_g that do not contain E , since the associated states do not appear in the objective.*

We also remark that, given a relevant component $\Omega \in \mathcal{O}$, there is an infinite number of mappings $g \in \mathcal{F}_{\text{sh}}$ so that $\Omega_g = \Omega$, see Lemma 2.10 below.

REMARK 2.7 (Comments regarding the definition of \mathcal{F}_{sh}). *Let us underline that all the conditions appearing in the definition of \mathcal{F}_{sh} are meaningful, as they guarantee that the admissible shapes have the properties stated in Lemma 2.9 below. These are mostly used for (2.6), in the proof of Proposition 4.6 and in subsection 4.2. Lemma 2.9 will also play a central role when establishing the limit adjoint equation in [5]. The inequality $g \leq 0$ in E implies that all admissible shapes contain the set E , as already mentioned.*

Since $g \in C^2(\bar{D})$, the requirements $|\nabla g| + |g| > 0$ in D and $g > 0$ on ∂D ensure that the set $\{x \in D : g(x) = 0\}$ is a finite union of disjoint closed C^2 curves, without self intersections, not intersecting ∂D [40, Prop. 2].

REMARK 2.8. *The condition $g(x) > 0 \forall x \notin \overline{\Omega_g}$ from (2.3) may be dropped in the present paper. This is however essential when one wants to describe the boundary of Ω_g as the set $\{x \in D : g(x) = 0\}$. Note that, when g is smooth and the first three inequalities in (2.3) are true, $\{x \in D : g(x) = 0\}$ may contain, in addition to $\partial\Omega_g$, other closed curves [40, Prop. 2]. We point out that in this paper, we only need that $g(x) > 0$ for a.a. $x \notin \overline{\Omega_g}$, which is in this case automatically fulfilled, cf. the proof of Lemma 2.9 below. In the recent work [5] concerned with optimality conditions (which is based on the findings from the present manuscript), the characterization $\partial\Omega_g = \{g = 0\}$ is however fundamental. In order to avoid redefining \mathcal{F}_{sh} there, we include now the requirement $g(x) > 0 \forall x \notin \overline{\Omega_g}$ in (2.3).*

The fact that we work in two dimensions only is due to the viability of the result in next lemma. If the three-dimensional case is considered, we need to include more conditions in \mathcal{F}_{sh} [28, 38].

LEMMA 2.9 (Properties of admissible shapes and Ω_g). *Let $g \in \mathcal{F}_{\text{sh}}$ and denote by $\Omega \in \mathcal{O}$ the relevant component of Ω_g , that is, the component that contains E . Then,*

1. Ω is a domain of class C^2 ;
2. $\partial\Omega_g = \{x \in D : g(x) = 0\}$ and $\Omega_g = \{x \in D : g(x) < 0\}$;
3. $\mu\{x \in D : g(x) = 0\} = 0$.

Proof. We observe that $\partial\Omega_g \subset \{x \in D : g(x) = 0\}$. As $g \in C^2(\bar{D})$ and since it satisfies the gradient condition and the condition on the boundary of D in (2.3), one deduces from [40, Prop. 2] that $\{x \in D : g(x) = 0\}$ is a finite union of disjoint closed C^2 curves, without self intersections, disjoint from ∂D . This implies the first claim.

The identity $\partial\Omega_g = \{x \in D : g(x) = 0\}$ is a consequence of the fact that $g > 0$ outside $\bar{\Omega}_g$ and it follows by a contradiction argument. Then, because of (2.2), it is straightforward to conclude $\Omega_g = \{x \in D : g(x) < 0\}$.

The last assertion is a direct consequence of the first two. It can also be deduced directly from [18, Lemma A.4] and the gradient condition in the definition of \mathcal{F}_{sh} . \square

2.2. Reformulation of (P_Ω) as an optimal control problem in a function space. In this subsection we rewrite (P_Ω) as an optimal control problem where the admissible set no longer consists of subsets of D , but of functions. In other words, we change the topology, so that (P_Ω) becomes amenable to optimal control methods. This is possible in light of the definition of the set of admissible shapes, see (2.4). The reformulation of (P_Ω) reads

$$\left. \begin{aligned} \min_{g \in \mathcal{F}_{\text{sh}}} \quad & \int_E (y_g(x) - y_d(x))^2 dx + \alpha \int_D 1 - H(g) dx, \\ \text{s.t.} \quad & -\Delta y_g + \beta(y_g) = f \quad \text{in } \Omega_g, \\ & y_g = 0 \quad \text{on } \partial\Omega_g, \end{aligned} \right\} \quad (P)$$

where $H : \mathbb{R} \rightarrow \{0, 1\}$ is the Heaviside function

$$H(v) := \begin{cases} 0, & \text{if } v \leq 0, \\ 1 & \text{if } v > 0. \end{cases} \quad (2.5)$$

Regarding the second term in the objective, we note that

$$\int_D 1 - H(g) dx = \int_{\Omega_g} dx \quad \forall g \in \mathcal{F}_{\text{sh}}, \quad (2.6)$$

in view of Lemma 2.9 and since $g > 0$ outside $\bar{\Omega}_g$, cf. (2.3).

LEMMA 2.10. *For each $\Omega \in \mathcal{O}$ there exists an infinite number of mappings $g \in \mathcal{F}_{\text{sh}}$ so that $\Omega_g = \Omega$.*

Proof. Since $\Omega \in \mathcal{O}$, there exists $\hat{g} \in \mathcal{F}_{\text{sh}}$ so that Ω is the component of $\Omega_{\hat{g}}$ that contains E , cf. (2.4). From [40, Prop. 2], see also the proof of Lemma 2.9, we know that $\Omega_{\hat{g}}$ has a finite number of components, and we define

$$\delta := \min_{\omega \neq \Omega, \omega \text{ is a component of } \Omega_{\hat{g}}} \text{dist}(\bar{\Omega}, \bar{\omega}) > 0.$$

Note that $\text{dist}(\bar{\Omega}, \bar{\omega}) > 0$ for each component ω of $\Omega_{\hat{g}}$ which is different from Ω (the boundary of $\Omega_{\hat{g}}$ consists of disjoint closed curves [40, Prop. 2], see also the proof of Lemma 2.9). Now, define the C^2 function

$$g := \hat{g} + c \text{dist}(\bar{\Omega}, \cdot)^3,$$

with $c > \frac{-\min_{x \in D} \widehat{g}}{\delta^3}$. Note that $c > 0$, since $\min_{x \in D} \widehat{g} \leq 0$, otherwise $\widehat{g} > 0$ in D and the proof is complete. We observe that

$$g = \widehat{g} \text{ in } \overline{\Omega}, \quad g > 0 \text{ in } D \setminus \overline{\Omega},$$

since $\widehat{g} \geq 0$ outside $\overline{\Omega_{\widehat{g}}}$. As $\widehat{g} \in \mathcal{F}_{\text{sh}}$, we can conclude from the above that $g \in \mathcal{F}_{\text{sh}}$ with $\Omega_g = \Omega$. Finally, we see that one can define an infinite number of mappings with this property, say $g_n := g + n \text{ dist}(\overline{\Omega_g}, \cdot)^3$, $n \in \mathbb{N}$. \square

The shape optimization problem (P_{Ω}) with the admissible set \mathcal{O} from Definition 2.5 is equivalent to (P) in the following sense.

PROPOSITION 2.11. *Let $\Omega^* \in \mathcal{O}$ be an optimal shape of (P_{Ω}) . Then, each of the functions $g^* \in \mathcal{F}_{\text{sh}}$ that satisfy $\Omega_{g^*} = \Omega^*$ is a global minimizer of (P) . Conversely, if $g^* \in \mathcal{F}_{\text{sh}}$ minimizes (P) , then the component of Ω_{g^*} that contains E is an optimal shape for (P_{Ω}) .*

Proof. Let $\Omega^* \in \mathcal{O}$ be an optimal shape for (P_{Ω}) with \mathcal{O} as in (2.4), i.e.,

$$\int_E (y_{\Omega^*}(x) - y_d(x))^2 dx + \alpha \int_{\Omega^*} dx \leq \int_E (y_{\Omega}(x) - y_d(x))^2 dx + \alpha \int_{\Omega} dx \quad \forall \Omega \in \mathcal{O}.$$

Now, let $g^* \in \mathcal{F}_{\text{sh}}$ with $\Omega_{g^*} = \Omega^*$ be fixed (note that, according to Lemma 2.10, there are infinitely many mappings with this property). Then, $y_{\Omega^*} = y_{g^*}$ and

$$\int_E (y_{g^*}(x) - y_d(x))^2 dx + \alpha \int_{\Omega_{g^*}} dx \leq \int_E (y_{\Omega}(x) - y_d(x))^2 dx + \alpha \int_{\Omega} dx \quad \forall \Omega \in \mathcal{O}.$$

Let $g \in \mathcal{F}_{\text{sh}}$ be arbitrary and fixed and denote by Ω the component of Ω_g that contains E . Testing with this particular Ω in the above inequality yields

$$\begin{aligned} \int_E (y_{g^*}(x) - y_d(x))^2 dx + \alpha \int_{\Omega_{g^*}} dx &\leq \int_E (y_g(x) - y_d(x))^2 dx + \alpha \int_{\Omega} dx \\ &\leq \int_E (y_g(x) - y_d(x))^2 dx + \alpha \int_{\Omega_g} dx, \end{aligned}$$

where we used that $y_g = y_{\Omega}$ in E . Since $g^* \in \mathcal{F}_{\text{sh}}$ and $g \in \mathcal{F}_{\text{sh}}$ was arbitrary, this proves the first statement, see (2.6).

To show the converse assertion, assume that $g^* \in \mathcal{F}_{\text{sh}}$ satisfies

$$\int_E (y_{g^*}(x) - y_d(x))^2 dx + \alpha \int_{\Omega_{g^*}} dx \leq \int_E (y_g(x) - y_d(x))^2 dx + \alpha \int_{\Omega_g} dx \quad \forall g \in \mathcal{F}_{\text{sh}}. \quad (2.7)$$

We denote by ω_{g^*} the component of Ω_{g^*} that contains E . This implies

$$y_{\omega_{g^*}} = y_{g^*} \text{ in } E.$$

Let $\Omega \in \mathcal{O}$ be arbitrary but fixed. Again, by Lemma 2.10, we can define $\tilde{g} \in \mathcal{F}_{\text{sh}}$ so that $\Omega_{\tilde{g}} = \Omega$. Then, $y_{\Omega} = y_{\tilde{g}}$. In view of (2.7), where we test with \tilde{g} , we have

$$\begin{aligned} \int_E (y_{\omega_{g^*}}(x) - y_d(x))^2 dx + \alpha \int_{\omega_{g^*}} dx &\leq \int_E (y_{g^*}(x) - y_d(x))^2 dx + \alpha \int_{\Omega_{g^*}} dx \\ &\leq \int_E (y_{\Omega}(x) - y_d(x))^2 dx + \alpha \int_{\Omega} dx. \end{aligned}$$

Since $\omega_{g^*} \in \mathcal{O}$ and $\Omega \in \mathcal{O}$ was arbitrary, this proves the second assertion. \square

From now on, all our findings concern the optimal control problem formulated as (P). We focus not only on global minimizers, but on the much larger class of local optima in the $L^2(D)$ -sense, which are defined as follows.

DEFINITION 2.12. *We say that $\bar{g}_{\text{sh}} \in \mathcal{F}_{\text{sh}}$ is locally optimal for (P) in the $L^2(D)$ sense, if there exists $r > 0$ such that*

$$\mathcal{J}(\bar{g}_{\text{sh}}) \leq \mathcal{J}(g) \quad \forall g \in \mathcal{F}_{\text{sh}} \text{ with } \|g - \bar{g}_{\text{sh}}\|_{L^2(D)} \leq r, \quad (2.8)$$

where

$$\mathcal{J}(g) := \int_E (\mathcal{S}(g)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H(g))(x) dx$$

is the reduced cost functional associated to the control problem (P) and

$$\mathcal{S} : g \in \mathcal{F}_{\text{sh}} \mapsto y_g \in H_0^1(\Omega_g) \cap H^2(\Omega_g)$$

denotes the control-to-state map. This will be introduced below, see Definition 4.8.

REMARK 2.13. *Clearly, in the study of local optima in the $L^2(D)$ sense of (P) we cover global optima, and thus, the associated (global) optimal shapes of (P_Ω) , cf. Proposition 2.11.*

Note that the set of local optima in the $H^1(D)$ -sense is larger than the set of local optima in the $L^2(D)$ -sense. However, we choose to work with the concept of local optima in the $L^2(D)$ -sense, in view of the essential density property from Proposition 3.1 below. This does not hold w.r.t. the $H^1(D)$ norm because of the boundary condition on ∂D in (2.3). Proposition 3.1 is however essential in the proof of Theorem 5.2, as a closer inspection shows; it ensures that $\{\hat{g}_\varepsilon\} \subset \mathcal{F}_{\text{sh}}$ is contained in the ball of local optimality of \bar{g}_{sh} . If this ball of local optimality is taken w.r.t. the $H^1(D)$ -norm, then it is necessary that Proposition 3.1 is also true w.r.t. the $H^1(D)$ -norm, which is not the case, as explained above.

REMARK 2.14 (Relation between the notion of local optimality in Definition 2.12 and 'local' optimal shape). *If two admissible shapes are 'close' to each other, this does not mean that the same is true for their parametrizations, since, given $g \in \mathcal{F}_{\text{sh}}$, there is an infinity of functions in \mathcal{F}_{sh} generating $\Omega_g \in \mathcal{O}$ (see Lemma 2.10). In fact, for each $g \in \mathcal{F}_{\text{sh}}$, there is a sequence $\{g_n\} \subset \mathcal{F}_{\text{sh}}$ so that $\Omega_g = \Omega_{g_n}$, $n \in \mathbb{N}$, and at the same time $\|g_n - g\|_{L^2(D)} \rightarrow \infty$ as $n \rightarrow \infty$ (cf. the end of the proof of Lemma 2.10). On the other hand, if $\|g_n - g\|_{L^2(D)} \rightarrow 0$, $g_n, g \in \mathcal{F}_{\text{sh}}$, we have*

$$\mu\{x \in D : g > 0 \text{ and } g_n \leq 0\} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

$$\mu\{x \in D : g < 0 \text{ and } g_n \geq 0\} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

see Lemma A.1. In light of Lemma 2.9, this accounts to

$$\mu(\Omega_{g_n} \setminus \Omega_g) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

$$\mu(\Omega_g \setminus \Omega_{g_n}) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

It is an open question how to define the notion of 'local optimal shape' so that it makes sense in connection with the L^2 local optimality of the associated parametrizations; this may involve the Hausdorff-Pompeiu distance or other notions of distances between two sets. The above convergences may be a good starting point for further investigations on this particular topic. At this stage, it is however clear that our method covers all global optimal shapes (Proposition 2.11 and Corollary 5.4).

3. Density of the set of admissible shape functions in $L^p(D)$, $p \in [1, \infty)$.

As already mentioned in the introduction, we cannot tackle (P) in a direct manner, so that we will consider an approximation scheme, cf. sections 4 and 5 below. One of the main challenges that arise in the investigation of (P) is the structure of the set \mathcal{F}_{sh} . It is a non-convex cone, while the governing PDE is non-smooth and the control appears as the parametrization of the unknown domain. The admissible set \mathcal{F}_{sh} has however a density property, which allows us to examine the approximating optimal control problem (P_ε) below on a convex subset of a Hilbert space, see (4.1).

The focus of this subsection is to prove the aforementioned density property, which is stated in the following (in a slightly more general form):

PROPOSITION 3.1. *Let*

$$\tilde{\mathcal{F}} := \{g \in L^2(D) : g \leq 0 \text{ a.e. in } E\}.$$

For each $g \in \tilde{\mathcal{F}}$, there exists a sequence $\{g_m\} \subset \mathcal{F}_{\text{sh}} \cap C^\infty(\bar{D})$ so that

$$g_m \rightarrow g \quad \text{in } L^2(D) \quad \text{as } m \rightarrow \infty.$$

REMARK 3.2. *The assertion in Proposition 3.1 remains true if $L^2(D)$ is replaced by $L^p(D)$, $p \in [1, \infty)$, see (3.4) and (3.8) below.*

It has different applications for all sort of problems where the solution operator is defined on \mathcal{F}_{sh} only, i.e., where conditions as those in Lemma 2.9 need to be fulfilled by the admissible shapes, see e.g. [22] and the references therein.

The proof of Proposition 3.1 will be conducted in three steps, as follows. We introduce the sets

$$\mathcal{F}_c := \{g \in C^\infty(\bar{D}) : g(x) \leq 0 \forall x \in E\},$$

$$\mathcal{F}_c^+ := \{g \in C^\infty(\bar{D}) : g(x) \leq 0 \forall x \in E, g(x) > 0 \forall x \in \partial D\} \quad (3.1)$$

and prove Proposition 3.1 by showing the set of densities

$$\begin{aligned} \mathcal{F}_c &\stackrel{d}{\hookrightarrow} \tilde{\mathcal{F}} \quad (\text{Lemma 3.3 below}), \\ \mathcal{F}_c^+ &\stackrel{d}{\hookrightarrow} \mathcal{F}_c \quad (\text{Lemma 3.4 below}), \\ \mathcal{F}_{\text{sh}} \cap C^\infty(\bar{D}) &\stackrel{d}{\hookrightarrow} \mathcal{F}_c^+ \quad (\text{Lemma 3.5 below}) \end{aligned} \quad (3.2)$$

w.r.t. the $L^2(D)$ norm.

LEMMA 3.3. *For each $g \in \tilde{\mathcal{F}}$, there exists a sequence $\{g_m\} \subset \mathcal{F}_c$ so that*

$$g_m \rightarrow g \quad \text{in } L^2(D) \quad \text{as } m \rightarrow \infty.$$

Proof. Let $g \in L^2(D)$ with $g \leq 0$ a.e. in E be arbitrary, but fixed and denote by $\chi_D g : \mathbb{R}^2 \rightarrow \mathbb{R}$ the extension of g by zero outside D . For every $m \in \mathbb{N}$ we define the $L^2(D)$ function $\hat{g}_m : \mathbb{R}^2 \rightarrow \mathbb{R}$ as

$$\hat{g}_m := \begin{cases} \min\{g, 0\} & \text{on } E_m, \\ \chi_D g & \text{otherwise,} \end{cases} \quad (3.3)$$

where

$$E_m := \{v \in \mathbb{R}^2 : d(v, E) \leq 1/m\},$$

and d is the distance induced by the l_∞ norm; note that $E_m \subset D$ for m large enough, by Assumption 2.3. Since $g \leq 0$ a.e. in E , we have

$$\|\widehat{g}_m - g\|_{L^2(D)}^2 = \|\min\{g, 0\} - g\|_{L^2(E_m \setminus E)}^2 = \int_{E_m \setminus E} (-\max\{g, 0\})^2 \rightarrow 0 \quad \text{as } m \rightarrow \infty, \quad (3.4)$$

since $(\max\{g, 0\})^2 \in L^1(D)$ and $\mu(E_m \setminus E) \rightarrow 0$ as $m \rightarrow \infty$, cf. e.g. [1, Lemma A.1.17]. Now we define the C^∞ function $g_m : \mathbb{R}^2 \rightarrow \mathbb{R}$ as

$$g_m(v) := \int_{\mathbb{R}^2} \widehat{g}_m(v - \frac{1}{m}s) \psi(s) ds,$$

where $\psi \in C_c^\infty(\mathbb{R}^2)$, $\psi \geq 0$, $\text{supp } \psi \subset [-1, 1]^2$ and $\int_{\mathbb{R}^2} \psi(s) ds = 1$. Then, by the definition of E_m and \widehat{g}_m , it holds

$$g_m(v) = \int_{[-1, 1]^2} \widehat{g}_m(v - \frac{1}{m}s) \psi(s) ds \leq 0 \quad \forall v \in E,$$

i.e., $g_m \in \mathcal{F}_c$. In view of (3.4), the proof is now complete. \square

LEMMA 3.4. *For each $g \in \mathcal{F}_c$, there exists a sequence $\{g_m\} \subset \mathcal{F}_c^+$, so that*

$$g_m \rightarrow g \quad \text{in } L^2(D) \quad \text{as } m \rightarrow \infty.$$

Proof. We recall that

$$\mathcal{F}_c := \{g \in C^\infty(\bar{D}) : g(x) \leq 0 \forall x \in E\}$$

and

$$\mathcal{F}_c^+ := \{g \in C^\infty(\bar{D}) : g(x) \leq 0 \forall x \in E, g(x) > 0 \forall x \in \partial D\}. \quad (3.5)$$

Let $g \in \mathcal{F}_c$ be arbitrary but fixed and $m \in \mathbb{N}$. We define $h_m \in C^\infty(\bar{D})$ as

$$h_m(x) := \begin{cases} -2 \min\{\min_{x \in \partial D} g, 0\} + 1/m & \text{for } x \in \partial D, \\ \in (0, -2 \min\{\min_{x \in \partial D} g, 0\} + 1/m) & \text{for } x \in D \text{ where } d(\partial D, x) \in (0, 1/m), \\ 0 & \text{for } x \in D \text{ where } d(\partial D, x) \geq 1/m. \end{cases} \quad (3.6)$$

Then, the mapping

$$g_m := g + h_m \in C^\infty(\bar{D}) \quad (3.7)$$

satisfies

$$\|g_m - g\|_{L^2(D)} = \|h_m\|_{L^2(D_m)} \rightarrow 0 \quad \text{as } m \rightarrow \infty, \quad (3.8)$$

where we abbreviate

$$D_m := \{x \in D : d(\partial D, x) \in (0, 1/m)\}.$$

Let us now check if g_m fulfils the inequalities appearing in the definition of \mathcal{F}_c^+ .

(i) $g_m \leq 0$ in E : By Assumption 2.3, we can choose m large enough so that $1/m < d(\bar{E}, \partial D)$. This implies that $h_m = 0$ in \bar{E} , which gives in turn $g_m = g \leq 0$ in \bar{E} .

(ii) $g_m > 0$ on ∂D : By (3.7), we have for $x \in \partial D$:

$$g_m(x) \geq \min\{\min_{x \in \partial D} g, 0\} - 2 \min\{\min_{x \in \partial D} g, 0\} + 1/m = -\min\{\min_{x \in \partial D} g, 0\} + 1/m \geq 1/m,$$

where for the first estimate we used the definition of h_m , i.e., (3.6). This allows us to conclude the desired assertion. \square

In the proof of the next lemma we are going to use the fact that, for an open set $A \subset \mathbb{R}^2$, it holds

$$\bar{A} = \overline{\text{int } A}. \quad (3.9)$$

This assertion can be easily checked, and it is true for general topological spaces.

LEMMA 3.5. *For each $g \in \mathcal{F}_c^+$, there exists a sequence $\{g_m\} \subset \mathcal{F}_{\text{sh}} \cap C^\infty(\bar{D})$, so that*

$$g_m \rightarrow g \quad \text{in } C(\bar{D}) \quad \text{as } m \rightarrow \infty.$$

Proof. We recall the definitions of \mathcal{F}_c^+ and \mathcal{F}_{sh} :

$$\mathcal{F}_c^+ := \{g \in C^\infty(\bar{D}) : g(x) \leq 0 \forall x \in E, g(x) > 0 \forall x \in \partial D\}, \quad (3.10)$$

$$\begin{aligned} \mathcal{F}_{\text{sh}} := \{g \in C^2(\bar{D}) : g(x) \leq 0 \forall x \in E, |\nabla g(x)| + |g(x)| > 0 \forall x \in D, g(x) > 0 \forall x \in \partial D, \\ g(x) > 0 \forall x \notin \bar{\Omega}_g\}. \end{aligned} \quad (3.11)$$

For the definition of Ω_g , see (2.2). Let $g \in \mathcal{F}_c^+$ be arbitrary but fixed. According to Sard's theorem [32, Thm. 7.2], the set

$$\{g(x) : \nabla g(x) = 0\} \subset \mathbb{R} \quad \text{has measure zero.} \quad (3.12)$$

Since g is continuous and $g > 0$ on ∂D , it holds $\min_{x \in \partial D} g(x) > 0$. In light of (3.12), there exists $\delta_1 \in (0, \min_{x \in \partial D} g(x))$ so that

$$\nabla g(x) \neq 0 \quad \forall x \in g^{-1}(\delta_1).$$

Note that $g^{-1}(\delta_1)$ may be empty, in which case the above assertion is still valid. Sard's theorem, see (3.12), further ensures the existence of a sequence $\{\delta_m\}$ satisfying

$$\delta_m \in (0, \min\{\delta_{m-1}, \frac{1}{m}\}), \quad m \in \mathbb{N}$$

and

$$\nabla g(x) \neq 0 \quad \forall x \in g^{-1}(\delta_m) \quad \forall m \in \mathbb{N}. \quad (3.13)$$

We notice that the above constructed sequence satisfies

$$\min_{x \in \partial D} g(x) > \delta_1 > \delta_2 > \dots > \delta_m > 0, \quad \delta_m \rightarrow 0 \quad \text{as } m \rightarrow \infty. \quad (3.14)$$

Further, we define $f_m \in C^\infty(\bar{D})$ as

$$f_m := \begin{cases} \in (0, 1) & \text{for } x \in \bar{D} \setminus \overline{\Omega_{g-\delta_m}}, \\ 0 & \text{for } x \in \overline{\Omega_{g-\delta_m}}, \end{cases} \quad (3.15)$$

where we recall that, according to (2.2), the set $\Omega_{g-\delta_m}$ is defined as

$$\Omega_{g-\delta_m} = \text{int}\{x \in D : g(x) - \delta_m \leq 0\}. \quad (3.16)$$

Note that functions f_m as in (3.15) exist, since we can choose a compact subset of \mathbb{R}^2 , disjoint with \bar{D} , so that $f_m = 1$ there. Then, the mapping

$$g_m := g - \delta_m + \frac{1}{m} f_m \in C^\infty(\bar{D}) \quad (3.17)$$

satisfies

$$\|g_m - g\|_{C(\bar{D})} \rightarrow 0 \quad \text{as } m \rightarrow \infty, \quad (3.18)$$

see (3.14). Let us now check if g_m fulfils the inequalities appearing in the definition of \mathcal{F}_{sh} .

(i) $g_m \leq 0$ in E : Since $g \in \mathcal{F}_c^+$, it holds $g \leq 0$ in \bar{E} , which implies that

$$\bar{E} \subset \{x \in D : g(x) - \delta_m \leq 0\}.$$

In view of (3.9) and (3.16), we infer

$$\bar{E} \subset \overline{\Omega_{g-\delta_m}},$$

which yields $f_m = 0$ in \bar{E} , see (3.15). Thus, by (3.17), $g_m = g - \delta_m \leq 0$ in \bar{E} .

(ii) $g_m > 0$ on ∂D : From (3.14) one deduces that for $x \in \partial D$ it holds

$$g(x) - \delta_m > g(x) - \min_{x \in \partial D} g \geq 0.$$

From (3.17) and (3.15) we now conclude that $g_m > 0$ on ∂D .

(iii) $g_m(x) > 0 \forall x \in \bar{D} \setminus \overline{\Omega_{g_m}}$: By (3.17) and (3.15) one knows that

$$g_m(x) > 0 \forall x \in \bar{D} \setminus \overline{\Omega_{g-\delta_m}}. \quad (3.19)$$

Next we prove the identity

$$\overline{\Omega_{g-\delta_m}} = \overline{\Omega_{g_m}}. \quad (3.20)$$

Let $x \in \overline{\Omega_{g-\delta_m}}$. Then, by (3.16), (3.15) and (3.17), we have that $x \in D$, $g(x) - \delta_m = g_m(x) \leq 0$; note that $x \notin \partial D$, due to (3.14), which implies $g > \delta_m$ on ∂D . Therefore,

$$\overline{\Omega_{g-\delta_m}} \subset \{x \in D : g_m(x) \leq 0\} \subset \{x \in D : g(x) - \delta_m \leq 0\}. \quad (3.21)$$

Hence,

$$\overline{\text{int } \overline{\Omega_{g-\delta_m}}} \subset \overline{\Omega_{g_m}} \subset \overline{\Omega_{g-\delta_m}}, \quad (3.22)$$

see (2.2). From (3.9) we infer

$$\overline{\text{int } \overline{\Omega_{g-\delta_m}}} = \overline{\Omega_{g-\delta_m}},$$

and, in light of (3.22), we have shown (3.20). Thanks to (3.19), this proves the desired assertion.

(iv) $|\nabla g_m(x)| + |g_m(x)| > 0 \forall x \in D$: In view of (iii), we only need to show that

$$|\nabla g_m(x)| + |g_m(x)| > 0 \forall x \in \overline{\Omega_{g_m}}.$$

By (3.17), (3.15), and (3.20), it holds

$$g_m = g - \delta_m \text{ in } \overline{\Omega_{g_m}}.$$

Note that, due to the regularity and the definition of f_m , see (3.15), its gradient vanishes on $\overline{\Omega_{g-\delta_m}}$, which coincides with the set $\overline{\Omega_{g_m}}$, cf. (3.20). This means that

$$\nabla g_m = \nabla g \quad \text{in } \overline{\Omega_{g_m}},$$

see (3.17). Now, thanks to (3.13), $g_m(x) = 0$ implies $\nabla g_m(x) \neq 0$, if $x \in \overline{\Omega_{g_m}}$. The proof is now complete. \square

4. The approximating optimal control problem. In this section, $\bar{g}_{\text{sh}} \in \mathcal{F}_{\text{sh}}$ is a local optimum of (P) in the $L^2(D)$ -sense. Inspired by a classical adapted penalization scheme [3] in combination with the fixed domain methodology from [26], our approximating minimization problem (P_ε) preserves the non-smoothness. We point out that the non-convex set \mathcal{F}_{sh} from (P) is now replaced by a convex subset of a Hilbert space. As shown by one of the main results in the next section (Theorem 5.2), this replacement is reasonable thanks to the findings from the previous section (Proposition 3.1).

Let $\varepsilon > 0$ be fixed. We consider the following approximating control problem

$$\left. \begin{aligned} \min_{g \in \mathcal{F}} \quad & \int_E (y(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(g))(x) dx + \frac{1}{2} \|g - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2 \\ \text{s.t.} \quad & -\Delta y + \beta(y) + \frac{1}{\varepsilon} H_\varepsilon(g)y = f + \varepsilon g \quad \text{in } D, \\ & y = 0 \quad \text{on } \partial D, \end{aligned} \right\} (P_\varepsilon)$$

where \mathcal{W} is the Hilbert space $L^2(D) \cap H^s(D \setminus \bar{E})$, $s \in (1, 2]$, endowed with the norm

$$\|\cdot\|_{\mathcal{W}}^2 := \|\cdot\|_{L^2(D)}^2 + \|\cdot\|_{H^s(D \setminus \bar{E})}^2,$$

and

$$\mathcal{F} := \{g \in \mathcal{W} : g \leq 0 \text{ a.e. in } E\}. \quad (4.1)$$

DEFINITION 4.1. *The non-linearity $H_\varepsilon : \mathbb{R} \rightarrow [0, 1]$ is defined as follows*

$$H_\varepsilon(v) := \begin{cases} 0, & \text{if } v \leq 0, \\ \frac{v^2(3\varepsilon - 2v)}{\varepsilon^3}, & \text{if } v \in (0, \varepsilon), \\ 1, & \text{if } v \geq \varepsilon. \end{cases} \quad (4.2)$$

REMARK 4.2. *The mapping H_ε introduced in (4.2) is Lipschitz continuous and continuously differentiable. It is obtained as a regularization of the Heaviside function*

from (2.5) and it is one possible choice that has the aforementioned properties. Note that Heaviside functions and their regularizations play an essential role in the context of shape optimization via fixed domain approaches, see for instance [26, 27] and the references therein.

REMARK 4.3 (Additional term in the state equation). *We notice that the state equation in (P_ε) is an approximating extension of the state equation in (P) from Ω_g to D [26]. We point out the presence of the additional term $+\varepsilon g$ on the right hand side, which is essential for the proof of our main strong stationarity result in [6], see [6, Rem. 3.16] for details. In all the other upcoming investigations, this can be dropped.*

REMARK 4.4 (Objective in (P_ε)). *The presence of the H^s term in the objective of (P_ε) ensures the existence of optimal solutions. We underline that it is necessary to consider the $H^s(D \setminus \bar{E})$ norm (not the full $H^s(D)$ norm), in order to conclude complete (i.e., strong stationary) optimality conditions for local optima of (P_ε) , see [6]. If $\mathcal{W} = H^s(D)$, certain sign conditions for the adjoint state are not available, and the optimality system for (P_ε) reduces to a weaker one, namely the limit optimality system obtained via smoothening methods.*

Moreover, it is essential that $s > 1$ in order to obtain the convergence of the states in Theorem 5.2, see also Lemma 4.10. This condition ensures that the embedding $H^s(D \setminus \bar{E}) \hookrightarrow L^\infty(D \setminus \bar{E})$ is true [16, p.88]. It is also necessary when deriving the final optimality system for the optimal shape of (P_Ω) in [5]. On the other side, the upper bound $s \leq 2$ is only needed to guarantee the inclusion $\mathcal{F}_{\text{sh}} \subset \mathcal{W}$, so that the last term in the objective of (P_ε) is well-defined, since $\bar{g}_{\text{sh}} \in \mathcal{F}_{\text{sh}}$.

4.1. Solvability of the state equations in (P_ε) and (P) . We start this subsection with a result on the unique solvability of the state equation in the approximating control problem. For convenience, we recall here the governing PDE in (P_ε) :

$$\begin{aligned} -\Delta y + \beta(y) + \frac{1}{\varepsilon} H_\varepsilon(g)y &= f + \varepsilon g \quad \text{in } D, \\ y &= 0 \quad \text{on } \partial D. \end{aligned} \tag{4.3}$$

LEMMA 4.5. *For any right-hand side $g \in L^2(D)$, the equation (4.3) admits a unique solution $y \in H_0^1(D) \cap H^2(D)$. This satisfies*

$$\|y\|_{H_0^1(D) \cap C(\bar{D})} \leq c_1 + c_2 \|g\|_{L^2(D)},$$

where $c_1, c_2 > 0$ are independent of ε , β , and g . The control-to-state mapping $S_\varepsilon : L^2(D) \rightarrow H_0^1(D) \cap H^2(D)$ is Lipschitz continuous on bounded sets, i.e., for every $M > 0$, there exists $L_{M,\varepsilon} > 0$ so that

$$\|S_\varepsilon(g_1) - S_\varepsilon(g_2)\|_{H_0^1(D) \cap H^2(D)} \leq L_{M,\varepsilon} \|g_1 - g_2\|_{L^2(D)} \quad \forall g_1, g_2 \in \overline{B_{L^2(D)}(0, M)}. \tag{4.4}$$

Proof. According to [42, Thm. 4.8], the equation (4.3) admits a unique solution $y \in H_0^1(D) \cap C(\bar{D})$ with

$$\|y\|_{H^1(D) \cap L^\infty(D)} \leq c \|f + \varepsilon g - \beta(0)\|_{L^2(D)},$$

where $c > 0$ is independent of ε , β , and of g . Then, by [10, Lem. 9.17], the $H^2(D)$ regularity follows.

To show the desired Lipschitz continuity of the control-to-state map, let $g_1, g_2 \in L^2(D)$ be arbitrary but fixed and abbreviate $y_i := S_\varepsilon(g_i)$, $i = 1, 2$. As a consequence of the monotonicity of β and the Lipschitz continuity and non-negativity of H_ε , cf. (4.2), one obtains

$$\|y_1 - y_2\|_{H_0^1(D)}^2 \leq \varepsilon \|g_1 - g_2\|_{L^2(D)} \|y_1 - y_2\|_{L^2(D)} \quad (4.5)$$

$$+ \frac{1}{\varepsilon} \|H_\varepsilon(g_1) - H_\varepsilon(g_2)\|_{L^2(D)} \|y_2\|_{C(\bar{D})} \|y_1 - y_2\|_{L^2(D)} \quad (4.6)$$

$$\leq \|g_1 - g_2\|_{L^2(D)} \|y_1 - y_2\|_{L^2(D)} \left(\varepsilon + \frac{L_{H_\varepsilon} c}{\varepsilon} \|f + \varepsilon g_2 - \beta(0)\|_{L^2(D)} \right), \quad (4.7)$$

where $c > 0$ is independent of ε , β , and of g_2 . This means that $S_\varepsilon : L^2(D) \rightarrow H_0^1(D)$ is Lipschitz continuous on bounded sets. In view of the $C^{1,1}$ regularity of D , the same can be concluded for $S_\varepsilon : L^2(D) \rightarrow H_0^1(D) \cap H^2(D)$. \square

Next we aim at highlighting the connection between (4.3) and the state equation in (P).

PROPOSITION 4.6 (Solvability of the state equation in (P)). *Let $g \in C(\bar{D})$ be fixed so that Ω_g is a domain of class C. Moreover, assume that $g > 0$ a.e. in $D \setminus \Omega_g$. Then,*

$$S_\varepsilon(g)|_{\Omega_g} \rightharpoonup y_g \quad \text{in } H^1(\Omega_g) \quad \text{as } \varepsilon \searrow 0,$$

where S_ε is the control-to-state map associated to (4.3) and $y_g \in H_0^1(\Omega_g)$ is the unique solution to

$$\begin{aligned} -\Delta y + \beta(y) &= f && \text{in } \Omega_g, \\ y &= 0 && \text{on } \partial\Omega_g. \end{aligned} \quad (4.8)$$

REMARK 4.7. *All the admissible domains $\Omega_g \in \mathcal{O}$ (Definition 2.4) satisfy (together with their associated parametrization) the hypotheses of Proposition 4.6. See Lemma 2.9 and (2.3).*

Proof. We follow the arguments from the proof of [26, Thm. 1]. For the beginning, let $\varepsilon > 0$ be arbitrary but fixed. We abbreviate for simplicity $y_\varepsilon := S_\varepsilon(g)$ and, by multiplying (4.3) with y_ε , we infer

$$\|y_\varepsilon\|_{H_0^1(D)}^2 + \int_D \beta(y_\varepsilon) y_\varepsilon \, dx + \frac{1}{\varepsilon} \int_D H_\varepsilon(g) y_\varepsilon^2 \, dx \leq \int_D (f + \varepsilon g) y_\varepsilon \, dx. \quad (4.9)$$

Since β is monotonically increasing and $H_\varepsilon(g) \geq 0$, by assumption, we deduce from (4.9) the estimate

$$\|y_\varepsilon\|_{H_0^1(D)}^2 + \underbrace{\int_D [\beta(y_\varepsilon) - \beta(0)] y_\varepsilon \, dx}_{\geq 0} \leq \int_D (f + \varepsilon g) y_\varepsilon \, dx - \int_D \beta(0) y_\varepsilon \, dx.$$

By applying Young's inequality on the right-hand side, we derive uniform bounds w.r.t. ε so that we can extract a weakly convergent subsequence

$$y_\varepsilon \rightharpoonup \hat{y} \quad \text{in } H_0^1(D). \quad (4.10)$$

In view of (2.1) and the compact embedding $H^1(D) \hookrightarrow L^2(D)$, we then have

$$\beta(y_\varepsilon) \rightarrow \beta(\widehat{y}) \quad \text{in } L^2(D); \quad (4.11)$$

note that $\|y_\varepsilon\|_{L^\infty(D)} \leq c$, where $c > 0$ is independent of ε , by Lemma 4.5. Further, multiplying (4.9) with ε implies

$$\int_D H_\varepsilon(g)y_\varepsilon^2 dx \rightarrow 0 \text{ as } \varepsilon \searrow 0.$$

In light of (4.2), it holds $H_\varepsilon(g) \rightarrow H(g)$ in $L^q(D)$, $q \in [1, \infty)$, where $H : \mathbb{R} \rightarrow \{0, 1\}$ stands for the Heaviside function (2.5). Thus, by (4.10) and the compact embedding $H^1(D) \hookrightarrow L^q(D)$, one obtains

$$\int_{D \setminus \Omega_g} \widehat{y}^2 dx = \int_D H(g)\widehat{y}^2 dx = \lim_{\varepsilon \searrow 0} \int_D H_\varepsilon(g)y_\varepsilon^2 dx = 0,$$

where the first identity is due to (2.2), (2.5), and $g > 0$ a.e. in $D \setminus \Omega_g$, by assumption. Hence, $\widehat{y} = 0$ a.e. in $D \setminus \Omega_g$. As $\widehat{y} \in H_0^1(D)$, cf. (4.10), it can be extended by zero on \mathbb{R}^2 (while preserving the H^1 regularity). Since Ω_g is of class C , by assumption, the result in [37, Thm. 2.3] implies that the weak limit from (4.10) satisfies

$$\widehat{y} \in H_0^1(\Omega_g). \quad (4.12)$$

Testing (4.3) with $\phi \in C_c^\infty(\Omega_g)$ further implies

$$\int_{\Omega_g} \nabla y_\varepsilon \nabla \phi dx + \int_{\Omega_g} \beta(y_\varepsilon) \phi dx = \int_{\Omega_g} (f + \varepsilon g) \phi dx,$$

since $H_\varepsilon(g) = 0$ on Ω_g (cf. definition (2.2) and (4.2)). Passing to the limit $\varepsilon \searrow 0$, where one relies on (4.10), (4.11), then results in

$$\int_{\Omega_g} \nabla \widehat{y} \nabla \phi dx + \int_{\Omega_g} \beta(\widehat{y}) \phi dx = \int_{\Omega_g} f \phi dx.$$

Since $\widehat{y} \in H_0^1(\Omega_g)$, cf. (4.12), we have $\widehat{y}|_{\Omega_g} = y_g$ and the proof is now complete. \square

The result in Proposition 4.6 allows us to introduce

DEFINITION 4.8 (The control-to-state map associated to the state equation in (P)).
We define

$$\mathcal{S} : g \in \mathcal{F}_{\text{sh}} \mapsto y_g \in H_0^1(\Omega_g) \cap H^2(\Omega_g), \quad (4.13)$$

where y_g solves the equation (4.8) on the component of Ω_g containing E , i.e., $\Omega_g \in \mathcal{O}$ (with a little abuse of notation, we do not make a difference between the notation for Ω_g and the notation for its relevant component; we just write $\Omega_g \in \mathcal{O}$ when we mean the latter, cf. Remark 2.6).

Note that the additional $H^2(\Omega_g)$ regularity is due to the fact that each admissible shape Ω_g is of class C^2 (Lemma 2.9.1) and [10, Lem. 9.17].

REMARK 4.9. According to Definition 4.8, $\mathcal{S}(g)$ exists only as an element of $H_0^1(\Omega_g)$. Whenever we write $\mathcal{S}(g)$ as an element of $H^1(D)$ in what follows, we think of its extension by zero outside $\Omega_g \in \mathcal{O}$.

4.2. Convergence properties. This subsection is dedicated to the study of other limit behaviours of solutions to (4.3) when $\varepsilon \rightarrow 0$. In addition to the convergence from Proposition 4.6, we will need in the proof of our main result (Theorem 5.2) two convergence results that are contained in Lemmas 4.10 and 4.12 below. Note that, from now on, we simply write $\Omega_g \in \mathcal{O}$ when we talk about the relevant component of Ω_g (Remark 2.6).

LEMMA 4.10. *Let $\{g_\varepsilon\} \subset \mathcal{F}$ and $g \in \mathcal{F}_{\text{sh}}$ so that*

$$g_\varepsilon \rightarrow g \quad \text{in } L^2(D) \cap L^\infty(D \setminus \bar{E}) \quad \text{as } \varepsilon \searrow 0.$$

Then, for each compact subset K of Ω_g , there exists $\varepsilon_0 > 0$, independent of x , so that

$$g_\varepsilon \leq 0 \quad \text{a.e. in } K, \quad \forall \varepsilon \in (0, \varepsilon_0]. \quad (4.14)$$

Moreover,

$$S_\varepsilon(g_\varepsilon) \rightarrow S(g) \quad \text{in } H_0^1(D) \quad \text{as } \varepsilon \searrow 0.$$

Proof. Since g is continuous, we have

$$g(x) \leq -\delta \quad \forall x \in K \subset \subset \Omega_g$$

for some $\delta > 0$. As $g_\varepsilon \rightarrow g$ in $L^\infty(D \setminus \bar{E})$, by assumption, there exists $\varepsilon > 0$, small, independent of x , so that

$$g_\varepsilon(x) \leq g(x) + \delta/2 \leq -\delta/2 < 0 \quad \text{a.e. in } K \cap (D \setminus \bar{E}).$$

Since $g_\varepsilon \in \mathcal{F}$, we can thus conclude (4.14).

Let us now show the desired convergence. We abbreviate for simplicity $y_\varepsilon := S_\varepsilon(g_\varepsilon)$ and, by arguing exactly as in the proof of Proposition 4.6, we show that there exists a weakly convergent subsequence (denoted by the same symbol) with

$$y_\varepsilon \rightharpoonup \hat{y} \quad \text{in } H_0^1(D) \quad \text{as } \varepsilon \searrow 0, \quad (4.15)$$

$$\beta(y_\varepsilon) \rightarrow \beta(\hat{y}) \quad \text{in } L^2(D) \quad \text{as } \varepsilon \searrow 0. \quad (4.16)$$

Moreover,

$$\int_D H_\varepsilon(g_\varepsilon) y_\varepsilon^2 dx \rightarrow 0 \quad \text{as } \varepsilon \searrow 0. \quad (4.17)$$

We also notice that, by Lemma 4.5, there exists a constant $C > 0$, independent of ε , so that

$$\|y_\varepsilon\|_{L^\infty(D)} \leq C \quad \forall \varepsilon > 0. \quad (4.18)$$

As a result of $g_\varepsilon \rightarrow g$ in $L^2(D)$ combined with Lemma A.1, we have

$$\mu\{x \in D : g < 0 \text{ and } g_\varepsilon \geq 0\} \rightarrow 0 \quad \text{as } \varepsilon \searrow 0,$$

$$\mu\{x \in D : g > 0 \text{ and } g_\varepsilon - \varepsilon \leq 0\} \rightarrow 0 \quad \text{as } \varepsilon \searrow 0$$

and with (4.18), it follows

$$\lim_{\varepsilon \rightarrow 0} \int_{\{g < 0, g_\varepsilon \geq 0\}} H_\varepsilon(g_\varepsilon) y_\varepsilon^2 dx + \lim_{\varepsilon \rightarrow 0} \int_{\{g > 0, g_\varepsilon \leq \varepsilon\}} H_\varepsilon(g_\varepsilon) y_\varepsilon^2 dx = 0.$$

Thus, by (4.17), Lemma 2.9.3, (4.2) and (4.15) combined with the compact embedding $H^1(D) \hookrightarrow L^2(D)$, one deduces

$$\begin{aligned} 0 &= \lim_{\varepsilon \rightarrow 0} \int_D H_\varepsilon(g_\varepsilon) y_\varepsilon^2 dx = \lim_{\varepsilon \rightarrow 0} \int_{\{g < 0, g_\varepsilon < 0\}} H_\varepsilon(g_\varepsilon) y_\varepsilon^2 dx + \lim_{\varepsilon \rightarrow 0} \int_{\{g > 0, g_\varepsilon > \varepsilon\}} H_\varepsilon(g_\varepsilon) y_\varepsilon^2 dx \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\{g > 0, g_\varepsilon > \varepsilon\}} y_\varepsilon^2 dx = \lim_{\varepsilon \rightarrow 0} \int_{\{g > 0\}} y_\varepsilon^2 dx = \lim_{\varepsilon \rightarrow 0} \int_{D \setminus \Omega_g} y_\varepsilon^2 dx \\ &= \int_{D \setminus \Omega_g} \widehat{y}^2 dx. \end{aligned} \tag{4.19}$$

Hence, $\widehat{y} = 0$ a.e. in $D \setminus \Omega_g$, and in view of [37, Thm. 2.3] (applied for the relevant component of Ω_g), this implies that

$$\widehat{y} \in H_0^1(\Omega_g).$$

Note that here we use that $\Omega_g \in \mathcal{O}$ is a domain of class C (even C^2 , see Lemma 2.9.1). Testing (4.3) w.r.h.s. g_ε with $\phi \in C_c^\infty(\Omega_g)$, $\Omega_g \in \mathcal{O}$, further yields

$$\int_{\Omega_g} \nabla y_\varepsilon \nabla \phi dx + \int_{\Omega_g} \beta(y_\varepsilon) \phi dx + \int_{\Omega_g} \frac{1}{\varepsilon} H_\varepsilon(g_\varepsilon) y_\varepsilon \phi dx = \int_{\Omega_g} (f + \varepsilon g_\varepsilon) \phi dx.$$

For a fixed $\phi \in C_c^\infty(\Omega_g)$ there exists a compact subset \widetilde{K} of Ω_g so that $\phi \in C_c^\infty(\widetilde{K})$. Hence, by (4.14) and (4.2), the third term in the above variational identity vanishes for $\varepsilon > 0$ small enough, independent of x (dependent on \widetilde{K} , and thus on ϕ). Passing to the limit $\varepsilon \searrow 0$, where one relies on (4.15), (4.16), and the uniform boundedness of $\{g_\varepsilon\}$ in $L^2(D)$, then results in

$$\int_{\Omega_g} \nabla \widehat{y} \nabla \phi dx + \int_{\Omega_g} \beta(\widehat{y}) \phi dx = \int_{\Omega_g} f \phi dx.$$

As $\widehat{y} \in H_0^1(\Omega_g)$, and since (4.8) is uniquely solvable, we conclude $\widehat{y} = \mathcal{S}(g)$ and, thanks to (4.15), the proof is now complete. \square

ASSUMPTION 4.11. *For the desired state we require $y_d \in H_0^1(E)$ with $\Delta y_d \in L^2(E)$. Moreover,*

$$f \geq \chi_E(-\Delta y_d + \beta(y_d)) + \chi_{D \setminus E} \beta(0) \quad \text{a.e. in } D, \tag{4.20}$$

or, alternatively,

$$f \leq \chi_E(-\Delta y_d + \beta(y_d)) + \chi_{D \setminus E} \beta(0) \quad \text{a.e. in } D. \tag{4.21}$$

LEMMA 4.12. *If Assumption 4.11 is satisfied, then*

$$\liminf_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(g_\varepsilon)(x) - y_d(x))^2 dx \geq \liminf_{\varepsilon \rightarrow 0} \int_E (\mathcal{S}(g_\varepsilon)(x) - y_d(x))^2 dx$$

for each sequence $\{g_\varepsilon\} \subset \mathcal{F}_{\text{sb}}$ that is uniformly bounded in $L^2(D)$.

Proof. Let $\varepsilon > 0$ be arbitrary but fixed. Arguing as in Lemma 4.5 we see that the equation

$$\begin{aligned} -\Delta y_\varepsilon + \beta(y_\varepsilon) + \frac{1}{\varepsilon} H_\varepsilon(g_\varepsilon) y_\varepsilon &= f \quad \text{a.e. in } D, \\ y_\varepsilon &= 0 \quad \text{on } \partial D \end{aligned} \quad (4.22)$$

admits a unique solution $y_\varepsilon := \widehat{S}_\varepsilon(g_\varepsilon) \in H_0^1(D) \cap H^2(D)$. Since $\{g_\varepsilon\}$ is uniformly bounded in $L^2(D)$, we have

$$\|y_\varepsilon - S_\varepsilon(g_\varepsilon)\|_{H_0^1(D)} \leq \varepsilon \|g_\varepsilon\|_{L^2(D)} \rightarrow 0 \quad \text{as } \varepsilon \searrow 0.$$

Thus, it suffices to show

$$\liminf_{\varepsilon \rightarrow 0} \int_E (\widehat{S}_\varepsilon(g_\varepsilon)(x) - y_d(x))^2 dx \geq \liminf_{\varepsilon \rightarrow 0} \int_E (S(g_\varepsilon)(x) - y_d(x))^2 dx. \quad (4.23)$$

We begin by noticing that the extension of y_d on D by zero, say $y_{d,0} \in H_0^1(D)$, solves

$$\begin{aligned} -\Delta \phi + \beta(\phi) &= \chi_E(-\Delta y_d + \beta(y_d)) + \chi_{D \setminus E} \beta(0) \quad \text{a.e. in } D, \\ \phi &= 0 \quad \text{on } \partial D, \end{aligned} \quad (4.24)$$

where we used that $D \setminus \bar{E}$ is open and that ∂E has measure zero, by Assumption 2.3. Further, we recall that, Ω_{g_ε} may have many components and, by Lemma 2.10, there exists $\tilde{g}_\varepsilon \in \mathcal{F}_{\text{sb}}$ so that the entire $\Omega_{\tilde{g}_\varepsilon}$ is just the relevant component of Ω_{g_ε} (the one that contains the set E , cf. Remark 2.6). Then we define $\tilde{y}_\varepsilon \in H_0^1(D) \cap H^2(D)$ as the unique solution to

$$\begin{aligned} -\Delta \tilde{y}_\varepsilon + \beta(\tilde{y}_\varepsilon) + \frac{1}{\varepsilon} H(\tilde{g}_\varepsilon) \tilde{y}_\varepsilon &= f \quad \text{a.e. in } D, \\ \tilde{y}_\varepsilon &= 0 \quad \text{on } \partial D, \end{aligned} \quad (4.25)$$

cf. the proof of Lemma 4.5. Note that the only difference to (4.22) is that we now consider the Heaviside function (2.5) acting on \tilde{g}_ε instead of $H_\varepsilon(g_\varepsilon)$. We underline that, in view of the end of the present proof, it is important to work here with $H(\tilde{g}_\varepsilon)$, not $H(g_\varepsilon)$.

Now we want to compare $y_\varepsilon, \tilde{y}_\varepsilon$ and y_d in order to obtain

$$\liminf_{\varepsilon \rightarrow 0} \|y_\varepsilon - y_d\|_{L^2(E)}^2 \geq \liminf_{\varepsilon \rightarrow 0} \|\tilde{y}_\varepsilon - y_d\|_{L^2(E)}^2. \quad (4.26)$$

(I) If (4.20) in Assumption 4.11 is true, then it holds

$$\tilde{y}_\varepsilon \geq y_{d,0} \quad \text{a.e. in } D, \quad (4.27)$$

where we keep in mind that

$$H(\tilde{g}_\varepsilon) y_{d,0} = \chi_{D \setminus E} H(\tilde{g}_\varepsilon) y_{d,0} = 0 \quad \text{a.e. in } D,$$

in light of $\tilde{g}_\varepsilon \leq 0$ in E and (2.5). Moreover, by comparing (4.22) and (4.24) we see that

$$y_\varepsilon \geq y_{d,0} \quad \text{a.e. in } D, \quad (4.28)$$

where this time we use

$$H_\varepsilon(g_\varepsilon)y_{d,0} = \chi_{D \setminus E} H_\varepsilon(g_\varepsilon)y_{d,0} = 0 \quad \text{a.e. in } D,$$

cf. (4.2). Further, from (4.22) we know that

$$\begin{aligned} -\Delta y_\varepsilon + \beta(y_\varepsilon) + \frac{1}{\varepsilon} H(\tilde{g}_\varepsilon)y_\varepsilon &= f + \frac{1}{\varepsilon} (H(\tilde{g}_\varepsilon) - H_\varepsilon(g_\varepsilon))y_\varepsilon \quad \text{a.e. in } D, \\ y_\varepsilon &= 0 \quad \text{on } \partial D. \end{aligned} \quad (4.29)$$

Due to (4.28), it holds

$$y_\varepsilon \geq 0 \quad \text{a.e. in } D \setminus E.$$

Owing to $\tilde{g}_\varepsilon \geq g_\varepsilon$ (by construction, see the proof of Lemma 2.10) combined with the fact that H is monotone increasing and (4.2), we have

$$H(\tilde{g}_\varepsilon) \geq H(g_\varepsilon) \geq H_\varepsilon(g_\varepsilon) \quad \text{in } D.$$

The two above inequalities together with (2.5), (4.2), and $\tilde{g}_\varepsilon, g_\varepsilon \leq 0$ in E , imply

$$(H(\tilde{g}_\varepsilon) - H_\varepsilon(g_\varepsilon))y_\varepsilon = \chi_{D \setminus E} \underbrace{(H(\tilde{g}_\varepsilon) - H_\varepsilon(g_\varepsilon))}_{\geq 0} y_\varepsilon \geq 0 \quad \text{a.e. in } D.$$

Hence, comparing (4.29) with (4.25) yields

$$y_\varepsilon \geq \tilde{y}_\varepsilon \quad \text{a.e. in } D.$$

From (4.28) and (4.27) we then arrive at (4.26).

(II) The case when (4.21) in Assumption 4.11 is true is treated completely analogously. Instead of (4.27) we have

$$\tilde{y}_\varepsilon \leq y_{d,0} \quad \text{and} \quad y_\varepsilon \leq y_{d,0} \quad \text{a.e. in } D,$$

which means that

$$y_\varepsilon \leq 0 \quad \text{a.e. in } D \setminus E.$$

Therefore,

$$(H(\tilde{g}_\varepsilon) - H_\varepsilon(g_\varepsilon))y_\varepsilon \leq 0 \quad \text{a.e. in } D.$$

Arguing as in (I) then gives in turn

$$y_\varepsilon \leq \tilde{y}_\varepsilon \leq y_{d,0} \quad \text{a.e. in } D,$$

whence (4.26) follows.

Thus, based on Assumption 4.11, we have just finished showing (4.26). The last part of the proof is concerned with the convergence

$$\tilde{y}_\varepsilon - \mathcal{S}(g_\varepsilon) \rightarrow 0 \quad \text{in } H_0^1(D) \quad \text{as } \varepsilon \searrow 0.$$

We abbreviate $z_\varepsilon := \mathcal{S}(g_\varepsilon)$. In view of Definition 4.8 and (4.8), z_ε is the unique solution to

$$\begin{aligned} -\Delta z_\varepsilon + \beta(z_\varepsilon) &= f \quad \text{in } \Omega_{g_\varepsilon}, \\ z_\varepsilon &= 0 \quad \text{on } \partial\Omega_{g_\varepsilon}, \end{aligned} \quad (4.30)$$

where $\Omega_{g_\varepsilon} \in \mathcal{O}$. Recall that $\tilde{g}_\varepsilon \in \mathcal{F}_{\text{sth}}$ was defined so that the entire $\Omega_{\tilde{g}_\varepsilon}$ is just the relevant component of Ω_{g_ε} (the one on which the above PDE is solved). Note that $\mathcal{S}(g_\varepsilon) = \mathcal{S}(\tilde{g}_\varepsilon)$. Thus, $z_\varepsilon \in H_0^1(D)$ (see Remark 4.9) satisfies

$$\begin{aligned} -\Delta z_\varepsilon + \beta(z_\varepsilon) + \frac{1}{\varepsilon} H(\tilde{g}_\varepsilon) z_\varepsilon &= \chi_{\Omega_{\tilde{g}_\varepsilon}} f + \chi_{D \setminus \Omega_{\tilde{g}_\varepsilon}} \beta(0) \quad \text{a.e. in } D, \\ z_\varepsilon &= 0 \quad \text{on } \partial D. \end{aligned} \quad (4.31)$$

We subtract (4.31) from (4.25) and test the resulting equation with $\tilde{y}_\varepsilon - z_\varepsilon \in H_0^1(D)$. This results in

$$\|\tilde{y}_\varepsilon - z_\varepsilon\|_{H_0^1(D)}^2 + \int_D \frac{1}{\varepsilon} H(\tilde{g}_\varepsilon) (\tilde{y}_\varepsilon - z_\varepsilon)^2 \leq \int_{D \setminus \Omega_{\tilde{g}_\varepsilon}} [f - \beta(0)] (\tilde{y}_\varepsilon - z_\varepsilon) dx, \quad (4.32)$$

by the monotonicity of β (Assumption 2.1). Further, the fact that $H(\tilde{g}_\varepsilon) = 0$ in $\Omega_{\tilde{g}_\varepsilon}$ and $z_\varepsilon = 0$ in $D \setminus \Omega_{\tilde{g}_\varepsilon}$ imply

$$\|\tilde{y}_\varepsilon - z_\varepsilon\|_{H_0^1(D)}^2 + \int_{D \setminus \Omega_{\tilde{g}_\varepsilon}} \frac{1}{\varepsilon} H(\tilde{g}_\varepsilon) \tilde{y}_\varepsilon^2 dx \leq \int_{D \setminus \Omega_{\tilde{g}_\varepsilon}} [f - \beta(0)] \tilde{y}_\varepsilon dx.$$

In light of (2.3) and Lemma 2.9.2-3 it holds $D \setminus \Omega_{\tilde{g}_\varepsilon} = \{\tilde{g}_\varepsilon > 0\}$ up to a set of measure zero (that is, $\partial\Omega_{\tilde{g}_\varepsilon}$). Thus, by (2.5) and Young's inequality, we get

$$\|\tilde{y}_\varepsilon - z_\varepsilon\|_{H_0^1(D)}^2 + \int_{D \setminus \Omega_{\tilde{g}_\varepsilon}} \frac{1}{\varepsilon} \tilde{y}_\varepsilon^2 dx \leq \int_{D \setminus \Omega_{\tilde{g}_\varepsilon}} \frac{\varepsilon}{2} [f - \beta(0)]^2 dx + \int_{D \setminus \Omega_{\tilde{g}_\varepsilon}} \frac{1}{2\varepsilon} \tilde{y}_\varepsilon^2 dx. \quad (4.33)$$

From (4.33) we infer

$$\tilde{y}_\varepsilon - z_\varepsilon \rightarrow 0 \quad \text{in } H_0^1(D) \quad \text{as } \varepsilon \searrow 0.$$

Thanks to (4.26), this leads to

$$\liminf_{\varepsilon \rightarrow 0} \|y_\varepsilon - y_d\|_{L^2(E)}^2 \geq \liminf_{\varepsilon \rightarrow 0} \|\tilde{y}_\varepsilon - y_d\|_{L^2(E)}^2 = \liminf_{\varepsilon \rightarrow 0} \|z_\varepsilon - y_d\|_{L^2(E)}^2,$$

whence (4.23) follows. The proof is now complete. \square

4.3. Existence of optimal solutions for (P_ε) . In the proof of our main result (Theorem 5.2), we will need that the approximating control problem (P_ε) admits optimal solutions (when the admissible set \mathcal{F} is restricted to a certain ball). For this, we gather some useful convergences in the next lemma. Throughout this subsection, $\varepsilon > 0$ is arbitrary but fixed, and the solution operator to (4.3) is denoted by S_ε , cf. Lemma 4.5.

LEMMA 4.13. *Let $\{g_n\} \subset \mathcal{F}$ with*

$$g_n \rightharpoonup g \quad \text{in } \mathcal{W}, \quad \text{as } n \rightarrow \infty.$$

Then,

$$H_\varepsilon(g_n) \rightarrow H_\varepsilon(g) \quad \text{in } L^q(D), \quad q \in [1, \infty) \quad \text{as } n \rightarrow \infty, \quad (4.34)$$

$$S_\varepsilon(g_n) \rightarrow S_\varepsilon(g) \quad \text{in } H^2(D) \cap H_0^1(D), \quad \text{as } n \rightarrow \infty. \quad (4.35)$$

Proof. We begin by observing that, due to $\{g_n\} \subset \mathcal{F}$, we have $g \in \mathcal{F}$, and thus, it holds

$$H_\varepsilon(g_n) = 0 = H_\varepsilon(g) \quad \text{a.e. in } E, \quad \forall n. \quad (4.36)$$

In light of the compact embedding $H^s(D \setminus \bar{E}) \hookrightarrow L^2(D \setminus E)$, and the continuity of H_ε , we also have

$$H_\varepsilon(g_n) \rightarrow H_\varepsilon(g) \quad \text{in } L^2(D \setminus E), \quad \text{as } n \rightarrow \infty.$$

This implies a.e. convergence on a subsequence and since $H_\varepsilon(g_n) \in [0, 1]$, we may now deduce (4.34), by Lebesgue's dominated convergence theorem. To show (4.35), we abbreviate for simplicity $y_n := S_\varepsilon(g_n)$. As a result of Lemma 4.5, the sequence $\{y_n\}$ is uniformly bounded in $H^2(D) \cap H_0^1(D)$, and we can thus extract a weakly convergent subsequence denoted by the same symbol so that

$$y_n \rightharpoonup \tilde{y} \quad \text{in } H^2(D) \cap H_0^1(D) \quad \text{as } n \rightarrow \infty.$$

Passing to the limit $n \rightarrow \infty$ in (4.3), where one uses (4.34), then yields

$$\begin{aligned} -\Delta \tilde{y} + \beta(\tilde{y}) + \frac{1}{\varepsilon} H_\varepsilon(g) \tilde{y} &= f + \varepsilon g \quad \text{a.e. in } D, \\ \tilde{y} &= 0 \quad \text{on } \partial D. \end{aligned} \quad (4.37)$$

From Lemma 4.5 we finally deduce $\tilde{y} = S_\varepsilon(g) \in H^2(D) \cap H_0^1(D)$ and the proof of (4.35) is complete. \square

By means of the direct method of calculus of variations, we can now show the following result.

PROPOSITION 4.14. *The approximating optimal control problem (P_ε) admits at least one global minimizer in \mathcal{F} .*

5. Correlation between (P_ε) and shape optimization. This section contains the main results of the present paper. One of them states that local optima in the $L^2(D)$ -sense of (P) (Definition 2.12) can be approximated by local minimizers of the approximating control problems (P_ε) introduced in the last section. As a consequence, we deduce that each parametrization of the optimal shape of (P_Ω) is the limit of global optima of (P_ε) (Corollary 5.4). These two findings allowed us to continue the research on the topic of optimality systems for (P_Ω) in the recent work [5].

Similarly to Definition 2.12, we introduce the notion of local optimality for the approximating control problem.

DEFINITION 5.1. *Let $\varepsilon > 0$ be fixed and $\bar{g}_{\text{sh}} \in \mathcal{F}_{\text{sh}}$. We say that $\bar{g}_\varepsilon \in \mathcal{F}$ is locally optimal for (P_ε) in the $L^2(D)$ sense, if there exists $r > 0$ such that*

$$j_\varepsilon(\bar{g}_\varepsilon) \leq j_\varepsilon(g) \quad \forall g \in \mathcal{F} \quad \text{with } \|g - \bar{g}_\varepsilon\|_{L^2(D)} \leq r, \quad (5.1)$$

where

$$j_\varepsilon(g) := \int_E (S_\varepsilon(g)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(g))(x) dx + \frac{1}{2} \|g - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2$$

is the reduced cost functional associated to the control problem (P_ε) .

THEOREM 5.2. *Let $\bar{g}_{\text{sh}} \in \mathcal{F}_{\text{sh}}$ be a local minimizer of (P) in the sense of Definition 2.12. Then, under Assumption 4.11, there exists a sequence of local minimizers $\{\bar{g}_\varepsilon\}$ of (P_ε) such that*

$$\bar{g}_\varepsilon \rightarrow \bar{g}_{\text{sh}} \quad \text{in } \mathcal{W}, \quad \text{as } \varepsilon \searrow 0.$$

Moreover,

$$S_\varepsilon(\bar{g}_\varepsilon) \rightarrow \mathcal{S}(\bar{g}_{\text{sh}}) \quad \text{in } H_0^1(D), \quad \text{as } \varepsilon \searrow 0.$$

Proof. The proof is a variation of well-known ideas [3]. However, let us point out that we need to take into account that the control-to-state map \mathcal{S} is defined on \mathcal{F}_{sh} , not on \mathcal{F} . To bridge the gap between these two sets, we will make use of the essential Proposition 3.1 and the convergence results from subsection 4.2.

Denote by $\overline{B_{L^2(D)}(\bar{g}_{\text{sh}}, r)}$, $r > 0$, the (closed) ball of local optimality of $\bar{g}_{\text{sh}} \in \mathcal{F}_{\text{sh}}$. Let $\varepsilon > 0$ be fixed. Thanks to Lemma 4.13, we can show by the direct method of calculus of variations that the optimization problem

$$\min_{g \in \mathcal{F} \cap \overline{B_{L^2(D)}(\bar{g}_{\text{sh}}, r/2)}} j_\varepsilon(g) \tag{5.2}$$

admits a global minimizer

$$\bar{g}_\varepsilon \in \mathcal{F} \cap \overline{B_{L^2(D)}(\bar{g}_{\text{sh}}, r/2)}.$$

We recall here that \mathcal{W} is the Hilbert space $L^2(D) \cap H^s(D \setminus \bar{E})$, $s \in (1, 2]$, endowed with the norm

$$\|\cdot\|_{\mathcal{W}}^2 := \|\cdot\|_{L^2(D)}^2 + \|\cdot\|_{H^s(D \setminus \bar{E})}^2,$$

and

$$\mathcal{F} := \{g \in \mathcal{W} : g \leq 0 \text{ a.e. in } E\}, \tag{5.3}$$

cf. (4.1). Note that the set $\mathcal{F} \cap \overline{B_{L^2(D)}(\bar{g}_{\text{sh}}, r/2)}$ is weakly closed, i.e., for $\{z_n\} \subset \mathcal{F} \cap \overline{B_{L^2(D)}(\bar{g}_{\text{sh}}, r/2)}$ we have

$$z_n \rightharpoonup z \quad \text{in } \mathcal{W} \Rightarrow z \in \mathcal{F} \cap \overline{B_{L^2(D)}(\bar{g}_{\text{sh}}, r/2)}.$$

Next, we recall that the reduced cost functional associated to the control problem (P) is given by

$$\mathcal{J}(g) = \int_E (\mathcal{S}(g)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H(g))(x) dx, \quad g \in \mathcal{F}_{\text{sh}}.$$

Due to Proposition 4.6 (see also Remark 4.7), it holds

$$\begin{aligned}
\mathcal{J}(\bar{g}_{\text{sh}}) &= \lim_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_{\text{sh}})(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_{\text{sh}}))(x) dx \\
&\geq \limsup_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_\varepsilon))(x) dx \\
&\quad + \frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2,
\end{aligned} \tag{5.4}$$

where for the last inequality we relied on the fact that \bar{g}_ε is a global minimizer of (5.2) and that \bar{g}_{sh} is admissible for (5.2). Thanks to Proposition 3.1, for each $\varepsilon > 0$, there exists $\hat{g}_\varepsilon \in \mathcal{F}_{\text{sh}}$ so that

$$\|\hat{g}_\varepsilon - \bar{g}_\varepsilon\|_{L^2(D)} \leq \frac{r\varepsilon}{2} \min\left\{\frac{1}{L_{H_\varepsilon}}, \frac{1}{L_{\|\bar{g}_{\text{sh}}\|_{L^2(D)} + r, \varepsilon}}\right\} \leq r\varepsilon/2, \tag{5.5}$$

where $L_{H_\varepsilon} > 0$ and $L_{\|\bar{g}_{\text{sh}}\|_{L^2(D)} + r, \varepsilon} > 0$ are the Lipschitz constant of H_ε and S_ε (see (4.4)), respectively; note that the second estimate in (5.5) is true for ε small enough. Then,

$$\|H_\varepsilon(\hat{g}_\varepsilon) - H_\varepsilon(\bar{g}_\varepsilon)\|_{L^2(D)} \leq r\varepsilon/2, \tag{5.6}$$

and according to (4.4), we also have

$$\|S_\varepsilon(\hat{g}_\varepsilon) - S_\varepsilon(\bar{g}_\varepsilon)\|_{L^2(D)} \leq L_{\|\bar{g}_{\text{sh}}\|_{L^2(D)} + r, \varepsilon} \|\hat{g}_\varepsilon - \bar{g}_\varepsilon\|_{L^2(D)} \leq r\varepsilon/2. \tag{5.7}$$

Here we note that

$$\|\hat{g}_\varepsilon\|_{L^2(D)}, \|\bar{g}_\varepsilon\|_{L^2(D)} \leq \|\bar{g}_{\text{sh}}\|_{L^2(D)} + r, \tag{5.8}$$

since \bar{g}_ε is admissible for (5.2) and in light of (5.5).

Going back to (5.4), we can thus write

$$\begin{aligned}
\mathcal{J}(\bar{g}_{\text{sh}}) &= \lim_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_{\text{sh}})(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_{\text{sh}}))(x) dx \\
&\geq \limsup_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_\varepsilon))(x) dx \\
&\quad + \frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2 \\
&\geq \liminf_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_\varepsilon))(x) dx \\
&\quad + \frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2 \\
&= \liminf_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\hat{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\hat{g}_\varepsilon))(x) dx \\
&\quad + \frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2,
\end{aligned} \tag{5.9}$$

where we relied on (5.6) and (5.7). Thanks to Lemma 4.12 combined with (5.8) and

$H_\varepsilon \leq H$ (see (4.2) and (2.5)), (5.9) can be continued as

$$\begin{aligned}
\mathcal{J}(\bar{g}_{\text{sh}}) &\geq \liminf_{\varepsilon \rightarrow 0} \int_E (\mathcal{S}(\widehat{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H(\widehat{g}_\varepsilon))(x) dx \\
&\quad + \frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2 \\
&\geq \int_E (\mathcal{S}(\bar{g}_{\text{sh}})(x) - y_d(x))^2 dx + \alpha \int_D (1 - H(\bar{g}_{\text{sh}}))(x) dx \\
&= \mathcal{J}(\bar{g}_{\text{sh}}).
\end{aligned} \tag{5.10}$$

The last estimate in (5.10) is due to the fact that $\widehat{g}_\varepsilon \in \mathcal{F}_{\text{sh}}$ is in the ball of local optimality of \bar{g}_{sh} (in light of (5.5) and since \bar{g}_ε is admissible for (5.2)). From (5.9) and (5.10) we can now conclude

$$\begin{aligned}
&\lim_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_\varepsilon))(x) dx + \frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2 \\
&= \mathcal{J}(\bar{g}_{\text{sh}}) = \int_E (\mathcal{S}(\bar{g}_{\text{sh}})(x) - y_d(x))^2 dx + \alpha \int_D (1 - H(\bar{g}_{\text{sh}}))(x) dx \\
&= \lim_{\varepsilon \rightarrow 0} \int_E (S_\varepsilon(\bar{g}_\varepsilon)(x) - y_d(x))^2 dx + \alpha \int_D (1 - H_\varepsilon(\bar{g}_\varepsilon))(x) dx,
\end{aligned} \tag{5.11}$$

where the last identity follows by estimating as in (5.9)-(5.10), without taking the term $\frac{1}{2} \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{\mathcal{W}}^2$ into account. By (5.11), we finally deduce

$$\bar{g}_\varepsilon \rightarrow \bar{g}_{\text{sh}} \quad \text{in } \mathcal{W}. \tag{5.12}$$

To show that \bar{g}_ε is a local minimizer of (P_ε) , let $v \in \mathcal{F}$ with $\|v - \bar{g}_\varepsilon\|_{L^2(D)} \leq r/4$ be arbitrary, but fixed. Then, by (5.12), we have for $\varepsilon > 0$ small enough

$$\|v - \bar{g}_{\text{sh}}\|_{L^2(D)} \leq \|v - \bar{g}_\varepsilon\|_{L^2(D)} + \|\bar{g}_\varepsilon - \bar{g}_{\text{sh}}\|_{L^2(D)} \leq r/2.$$

As \bar{g}_ε is global optimal for (5.2), this means that $j_\varepsilon(\bar{g}_\varepsilon) \leq j_\varepsilon(v)$, whence the desired local optimality of \bar{g}_ε follows (Definition 5.1).

Since $s > 1$, we have $\mathcal{W} \hookrightarrow L^2(D) \cap L^\infty(D \setminus \bar{E})$, and Lemma 4.10 with (5.12) implies

$$S_\varepsilon(\bar{g}_\varepsilon) \rightharpoonup \mathcal{S}(\bar{g}_{\text{sh}}) \quad \text{in } H_0^1(D), \text{ as } \varepsilon \searrow 0.$$

The proof is now complete. \square

REMARK 5.3 (Approximation of global optima of (P)). *A short inspection of the proof of Theorem 5.2 shows that, if \bar{g}_{sh} is a global minimizer, then the associated sequence $\{\bar{g}_\varepsilon\}$ inherits this property, i.e., it consists of global optima of (P_ε) . Here we pay attention to the fact that (5.8) is replaced by*

$$\|\widehat{g}_\varepsilon\|_{L^2(D)}, \|\bar{g}_\varepsilon\|_{L^2(D)} \leq c, \tag{5.13}$$

where $c > 0$ is a constant independent of ε (this can be proven by arguing as in the proof of (5.4)). Thus, we can still make use of Lemma 4.12 to deduce (5.10).

As a consequence of Theorem 5.2 and Proposition 2.11 we then arrive at the following **COROLLARY 5.4** (Approximation of the optimal shape). *Suppose that Assumption 4.11 is satisfied. Let $\Omega^* \in \mathcal{O}$ be an optimal shape for (P_Ω) . Then, for each $g^* \in \mathcal{F}_{\text{sh}}$ with $\Omega_{g^*} = \Omega^*$, there exists a sequence of global minimizers $\{\bar{g}_\varepsilon\}$ of (P_ε) such that*

$$\bar{g}_\varepsilon \rightarrow g^* \quad \text{in } \mathcal{W}, \text{ as } \varepsilon \searrow 0.$$

Moreover,

$$S_\varepsilon(\bar{g}_\varepsilon) \rightharpoonup \mathcal{S}(g^*) \quad \text{in } H_0^1(D), \text{ as } \varepsilon \searrow 0.$$

Acknowledgment. This work was financially supported by the DFG grant BE 7178/3-1 for the project "Optimal Control of Viscous Fatigue Damage Models for Brittle Materials: Optimality Systems".

The author thanks Prof. Dan Tiba (Institute of Mathematics of the Romanian Academy) for useful discussions during the writing of this paper.

Appendix A. A useful convergence.

LEMMA A.1. *Let \mathcal{M} be a measurable set and assume that $h_n \rightarrow h$ in $L^1(\mathcal{M})$ as $n \rightarrow \infty$. Then,*

$$\mu\{x \in \mathcal{M} : h(x) > 0 \text{ and } h_n(x) \leq 0\} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Proof. The proof is inspired by the proof of [13, Lem. A.2] and uses the result established in [13, Lem. A.1]. We abbreviate $\mathcal{M}^+ := \{x \in \mathcal{M} : h(x) > 0\}$ and $\mathcal{M}_n := \{x \in \mathcal{M} : h(x) > 0 \text{ and } h_n(x) \leq 0\}$. If the desired assertion is not true, then there exists $\gamma > 0$ and a subsequence $\{n_k\}$ so that

$$\mu(\mathcal{M}_{n_k}) \geq \gamma \quad \forall k \in \mathbb{N}.$$

From [13, Lem. A.1] we know that

$$\int_{\mathcal{M}_{n_k}} h \, dx \geq \delta \quad \forall k \in \mathbb{N}$$

for some $\delta > 0$. Further, it holds

$$\begin{aligned} \|\max\{h_{n_k}, 0\} - \max\{h, 0\}\|_{L^1(\mathcal{M}^+)} &\geq \int_{\mathcal{M}_{n_k}} |\max\{h_{n_k}, 0\} - \max\{h, 0\}| \, dx \\ &= \int_{\mathcal{M}_{n_k}} h \, dx \geq \delta \quad \forall k \in \mathbb{N}, \end{aligned}$$

where, for the identity, we made use of the definition of \mathcal{M}_n . Since $h_n \rightarrow h$ in $L^1(\mathcal{M})$, by assumption, we get a contradiction and the proof is now complete. \square

REFERENCES

- [1] H.W. Alt. *Lineare Funktionalanalysis*. Springer Verlag, Berlin, 2006.
- [2] S. Amstutz and M. Ciligot-Travain. Optimality conditions for shape and topology optimization subject to a cone constraint. *SIAM Journal on Control and Optimization*, 48(6):4056–4077, 2010.
- [3] V. Barbu. *Optimal control of variational inequalities*. Pitman (Advanced Publishing Program), Boston, 1984.
- [4] L. Betz. Strong stationarity for a highly nonsmooth optimization problem with control constraints. *Mathematical Control and Related Fields*, 13(4):1500–1528, 2023.
- [5] L. Betz. Necessary conditions for the optimal control of a shape optimization problem with non-smooth PDE constraints. *arXiv:2409.15039*, 2024.
- [6] L. Betz. Optimal control of a non-smooth elliptic PDE with non-linear term acting on the control. *arXiv:2407.06726*, 2024.
- [7] M. Delfour and J.-P. Zolesio. *Shapes and Geometries*. SIAM, Philadelphia, 2001.
- [8] Z. Denkowski and S. Migórski. Optimal shape design for elliptic hemivariational inequalities in nonlinear elasticity. In *Variational calculus, optimal control and applications*, volume 124 of *Internat. Ser. Numer. Math.* Birkhäuser, Basel, 1998.
- [9] Z. Denkowski and S. Migórski. Optimal shape design problems for a class of systems described by hemivariational inequalities. *J. Global Optim.*, 12(1):37–59, 1998.

- [10] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Springer Verlag, Berlin, 2001.
- [11] A. Halanay, C. Murea, and D. Tiba. Some properties of the period for certain ordinary differential systems and applications to topology optimization of variational inequalities. *Banach Center Publications*, 127:129–146, 2024.
- [12] C. Heinemann and K. Sturm. Shape optimization for a class of semilinear variational inequalities with applications to damage models. *SIAM J. Math. Anal.*, 48(5):3579–3617, 2016.
- [13] R. Herzog, C. Meyer, and G. Wachsmuth. B- and strong stationarity for optimal control of static plasticity with hardening. *SIAM J. Optim.*, 23(1):321–352, 2013.
- [14] M. W. Hirsch, S. Smale, and R. L. Devaney. *Differential Equations, Dynamical Systems, and an Introduction to Chaos*. Academic Press, 2012.
- [15] K. Ito, K. Kunisch, and G.H. Peichl. Variational approach to shape derivatives. *ESAIM: Control, Optimisation and Calculus of Variations*, 14(3):517–539, 2008.
- [16] W. Kabbalo. *Aufbaukurs Funktionalanalysis und Operatortheorie*. Springer Verlag, Berlin Heidelberg, 2014.
- [17] H. Kasumba and K. Kunisch. On computation of the shape Hessian of the cost functional without shape sensitivity of the state variable. *J. Optim. Theory Appl.*, 162:779–804, 2014.
- [18] D. Kinderlehrer and G. Stampacchia. *An Introduction to Variational Inequalities and Their Applications*. SIAM, Philadelphia, 2000.
- [19] A. Laurain, M. Winckler, and I. Yousept. Shape optimization for superconductors governed by H(curl)- elliptic variational inequalities. *SIAM J. Control and Optimiz.*, 59(3):2247–2272, 2021.
- [20] D. Luft, V. H. Schulz, and K. Welker. Efficient techniques for shape optimization with variational inequalities using adjoints. *SIAM J. Optimiz.*, 30(3), 2020.
- [21] C. Murea and D. Tiba. Topological optimization via cost penalization. *Topological Methods in Nonlinear Analysis*, 54(2B):1023–1050, 2019.
- [22] C. Murea and D. Tiba. Periodic Hamiltonian systems in shape optimization problems with Neumann boundary conditions. *Journal of Differential Equations*, 321:1–39, 2022.
- [23] C. Murea and D. Tiba. Penalization of stationary Navier-Stokes equations and applications in topology optimization. *arXiv:2212.04796*, 2023.
- [24] C. Murea and D. Tiba. Topology optimization and boundary observation for clamped plates. *arXiv:2310.11905*, 2023.
- [25] P. Neittaanmaki, J. Sokolowski, and J.P. Zolesio. Optimization of the domain in elliptic variational inequalities. *Appl. Math. and Optimiz.*, 18(1):85–98, 1998.
- [26] P. Neittaanmäki, A. Pennanen, and D. Tiba. Fixed domain approaches in shape optimization problems with Dirichlet boundary conditions. *Inverse Problems*, 25(5):055003, 2009.
- [27] P. Neittaanmäki and D. Tiba. Fixed domain approaches in shape optimization problems. *Inverse Problems*, 28(9):093001, 2012.
- [28] M.R. Nicolai and D. Tiba. Implicit functions and parametrizations in dimension three: generalized solutions. *Discrete Contin. Dyn. Syst.*, 35(6):2701–2710, 2015.
- [29] A. A. Novotny and J. Sokolowski. *Topological Derivatives in Shape Optimization*. Springer, Berlin, 2013.
- [30] S. Osher and J.A. Sethian. Fronts propagating with curvature-dependent speed: algorithms based on Hamilton-Jacobi formulations. *J. Comput. Phys.*, 79(1):12–49, 1988.
- [31] O. Pironneau. *Optimal Shape Design for Elliptic Systems*. Springer Verlag, Berlin, 1984.
- [32] A. Sard. The measure of the critical values of differentiable maps. *Bulletin of the American Mathematical Society*, 48:883–890, 1942.
- [33] J. Sokolowski. Shape sensitivity analysis of nonsmooth variational problems. *Boundary Control and Boundary Variations. Proceedings of the IFIP WG 7.2 Conference, Lecture Notes in Control and Information Sciences, Springer Verlag, Berlin*, 100:265–285, 1988.
- [34] J. Sokolowski and A. Zochowski. Optimality conditions for simultaneous topology and shape optimization. *SICON*, 42(4):1198–1221, 2003.
- [35] J. Sokolowski and J.-P. Zolesio. *Introduction to Shape Optimizations. Shape Sensitivity Analysis*. Springer, Berlin, 1992.
- [36] K. Sturm. Minimax Lagrangian approach to the differentiability of nonlinear PDE constrained shape functions without saddle point assumption. *SICON*, 53(4):2017–2039, 2015.
- [37] D. Tiba. Domains of class C: properties and applications. *Annals of the University of Bucharest*, 4(LXII):89–102, 2013.
- [38] D. Tiba. The implicit function theorem and implicit parametrizations. *Ann. Acad. Rom. Sci. Ser. Math. Appl.*, 5(1-2):193–208, 2013.
- [39] D. Tiba. Iterated Hamiltonian type systems and applications. *J. Differential Equations*, 264(8):5465–5479, 2018.

- [40] D. Tiba. A penalization approach in shape optimization. *Atti Accad. Pelorit. Pericol. Cl. Sci. Fis. Mat. Nat.*, 96(1), 2018.
- [41] D. Tiba. Optimality conditions and Lagrange multipliers for shape and topology optimization problems. *Atti Accad. Pelorit. Pericol. Cl. Sci. Fis. Mat. Nat.*, 101(1), 2023.
- [42] F. Tröltzsch. *Optimal Control of Partial Differential Equations*, volume 112 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, 2010. Theory, methods and applications, Translated from the 2005 German original by Jürgen Sprekels.
- [43] G. Wachsmuth. Strong stationarity for optimal control of the obstacle problem with control constraints. *SIAM Journal on Optimization*, 24(4):1914–1932, 2014.