

ON THE CLASSIFICATION OF INOUE SURFACES

ZAHRAA KHALED AND ANDREI TELEMAN

ABSTRACT. We prove that any Inoue surface admits a unique holomorphic connection. Using this result we show that two Inoue surfaces $S = H \times \mathbb{C}/G$, $S' = H \times \mathbb{C}/G'$ are biholomorphic if and only if G, G' are conjugate in the group of affine transformations of $H \times \mathbb{C}$. This result allows us to prove explicit classification theorems for Inoue surfaces: Let \mathcal{M} be the set of $\mathrm{SL}(3, \mathbb{Z})$ -matrices M with a real eigenvalue $\alpha > 1$ and two non-real eigenvalues, and let \mathcal{N}^\pm the set of $\mathrm{GL}(2, \mathbb{Z})$ -matrices N with a real eigenvalue $\alpha > 1$ and $\det(N) = \pm 1$.

We prove that:

- (1) For any $\mathrm{GL}(3, \mathbb{Z})$ -similarity class $\mathfrak{M} \in \mathcal{M}/\sim$, there exists exactly two biholomorphism classes of type I Inoue surfaces.
- (2) (+) For any $\mathrm{GL}(2, \mathbb{Z})$ similarity class $\mathfrak{N} = [N] \in \mathcal{N}^+/\sim$ and positive integer $r \in \mathbb{N}^*$, we have a finite set of deformation classes of type II Inoue surfaces. This set is parameterised by the quotient of $\mathbb{Z}^2/(I_2 - N)\mathbb{Z}^2 + r\mathbb{Z}^2$ by an action of the "positive centraliser" $Z_{\mathrm{GL}(2, \mathbb{Z})}^+(N)$ of N in $\mathrm{GL}(2, \mathbb{Z})$. The set of biholomorphism types corresponding to a deformation class, endowed with its natural topology, can be identified with either \mathbb{C}^* or \mathbb{C} .
- (-) For any $\mathrm{GL}(2, \mathbb{Z})$ -similarity class $\mathfrak{N} = [N] \in \mathcal{N}^-/\sim$ and positive integer $r \in \mathbb{N}^*$, we have a finite set of biholomorphism classes of type III Inoue surfaces. This set is parameterised by the quotient of $\mathbb{Z}^2/(I_2 + N)\mathbb{Z}^2 + r\mathbb{Z}^2$ by an action of $Z_{\mathrm{GL}(2, \mathbb{Z})}^+(N)$.

In both cases $Z_{\mathrm{GL}(2, \mathbb{Z})}^+(N)$ -can be identified with the stabiliser of N in $\mathrm{PGL}(2, \mathbb{Z})$ and is an infinite cyclic group (see section 5).

Taking into account the Latimer-MacDuffee theorem, it follows that:

- (1) For any polynomial $\chi \in \chi(\mathcal{M})$ we have only finitely many biholomorphism classes of type I surfaces.
- (2) For any pair $(\chi, r) \in \chi(\mathcal{N}^+) \times \mathbb{N}^*$ we have only finitely many deformation classes of type II Inoue surfaces.
- (3) For any pair $(\chi, r) \in \chi(\mathcal{N}^-) \times \mathbb{N}^*$ we have only finitely many biholomorphism classes of type III Inoue surfaces.

CONTENTS

1. Introduction	2
2. The construction of Inoue surfaces	5
2.1. Type I Inoue surfaces	5
2.2. Type II Inoue surfaces	7
2.3. Type III Inoue surfaces	10
3. Fundamental properties of Inoue surfaces	11
3.1. Line bundles on Inoue surfaces	11
3.2. The tangent bundle of an Inoue surface	12
3.3. Vanishing theorems	13
4. The classification of Inoue surfaces	17

2000 *Mathematics Subject Classification.* 32J15, 32Q57.

We are grateful to Matei Toma for his useful suggestions concerning the formalism used in the construction of type I Inoue surfaces.

4.1. The classification of type I Inoue surfaces	18
4.2. The classification of type II Inoue surfaces	21
4.3. The classification of type III Inoue surfaces	39
5. Appendix: The positive centraliser $Z_{\mathrm{GL}(2,\mathbb{Z})}^+(N)$	40
References	43

1. INTRODUCTION

In his renowned article [In], Inoue introduced three types of non-Kählerian surfaces obtained as quotients of $U := H \times \mathbb{C}$ (where $H := \{w \in \mathbb{C} \mid \Im(w) > 0\}$) by certain groups of affine transformations acting properly discontinuously on U . Moreover Inoue proves that any surface S belonging to one of his three families is a class VII surface with $b_2 = 0$ and has the following geometric properties:

- (A) S admits a holomorphic foliation, or, equivalently, the tangent sheaf admits an invertible subsheaf.
- (B) S contains no curve, or, equivalently, for any nontrivial invertible sheaf \mathcal{N} on S , one has $H^0(S, \mathcal{N}) = 0$.

The fundamental theorem [In, Theorem p. 280] states that, conversely, any class VII surface with $b_2 = 0$ having these two properties is biholomorphic to an Inoue surface, i.e. to a surface belonging to one of Inoue's three families.

Inoue surfaces have many interesting properties (see for instance [Wall], [Te]) and play a fundamental role in the theory of surfaces (see for instance [BHPV]) so, naturally, they are abundantly mentioned in the literature. However, to our surprise, we did not find in the literature an explicit classification theorem for Inoue surfaces (as we have for instance for primary Hopf surfaces [We]).

Also to our surprise we realised that the problem is more difficult and more interesting than we expected. The first difficulty we encountered comes from the traditional notations used in the literature for these surfaces. For instance, the "traditional" notation used for Inoue surfaces of the first type is S_M , where $M \in \mathrm{SL}(3, \mathbb{Z})$ is a matrix with $\mathrm{Spec}_{\mathbb{C}}(M) \supsetneq \mathrm{Spec}_{\mathbb{R}}(M) \subset]1, +\infty[$. This notation suggests that for any such matrix M one has a well defined complex surface S_M . This is not true. In fact, in order to construct an Inoue surface of the first type, one also has to fix a non-real eigenvalue $\beta \in \mathrm{Spec}_{\mathbb{C}}(M) \setminus \mathbb{R}$ and a pair $(a, b) \in \mathbb{R}^3 \times \mathbb{C}^3$ consisting of a real eigenvector for the real eigenvalue $\alpha = |\beta|^{-2} \in \mathrm{Spec}_{\mathbb{R}}(M)$ and an eigenvector for β . Having the triple (β, a, b) one can recover M , but not vice-versa. Therefore the notation which is appropriate for classification purposes is $S_{a,b}^{\beta}$, not S_M .

For the first type of surfaces our final result (see Theorem 4.7, Remark 4.8) shows that biholomorphism classes of first type Inoue surfaces correspond bijectively to pairs (\mathfrak{M}, β) consisting of a $\mathrm{GL}(3, \mathbb{Z})$ -similarity class of $\mathrm{SL}(3, \mathbb{Z})$ -matrices of the considered type and a non-real eigenvalue $\beta \in \mathrm{Spec}(\mathfrak{M}) \setminus \mathbb{R}$. Therefore to any similarity class \mathfrak{M} of such matrices correspond *two* biholomorphism types of type 1 Inoue surfaces. This shows that *giving M does not even specify a biholomorphism type of Inoue surfaces*.

Note that, by a renowned Theorem of Latimer and MacDuffee (see [LMD], [Ta]), for any degree n polynomial $\chi \in \mathbb{Z}[X]$ which is irreducible in $\mathbb{Q}[X]$, the set of $\mathrm{GL}(n, \mathbb{Z})$ -similarity classes of matrices with integer coefficients and characteristic polynomial χ correspond bijectively to the set of ideal classes of the order $\mathbb{Z}[\alpha]$,

where α is a root of χ . Therefore, by a fundamental theorem in Number Theory [BoSh, Theorem 3, p. 128], this set is finite¹.

Therefore, for any polynomial $X^3 - \theta_2 X^2 + \theta_1 X - 1 \in \mathbb{Z}[X]$ with a root $\alpha \in]1, +\infty[$ and two non-real roots, we have finitely many biholomorphism classes of type I Inoue surfaces, namely two biholomorphism classes for any equivalence class of ideals of the order $\mathbb{Z}[\alpha]$.

For surfaces of the second type, the situation is even more complicated: the "traditional" notation for a type II Inoue surface is $S_{N,p,q,r,t}^+$, where $N \in \text{SL}(2, \mathbb{Z})$ has $\text{Spec}(N) \subset (0, +\infty) \setminus \{1\}$ and $(p, q, r, t) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}^* \times \mathbb{C}$. This notation suggests that any 5-tuple (N, p, q, r, t) as above yields a well defined surface denoted $S_{N,p,q,r,t}^+$. This again is not true; the construction also needs a pair $(a, b) \in \mathbb{R}^2 \times \mathbb{R}^2$, where a is an eigenvector for the eigenvalue $\alpha \in \text{Spec}(N) \cap]1, +\infty[$ and b is an eigenvector for α^{-1} .

For this class of surfaces we will adopt the notation $S_{a,b,c,t}^{\alpha,r}$, where

- $t \in \mathbb{C}$, $r \in \mathbb{N}^*$ and $\alpha \in]1, +\infty[$ is S^+ -admissible, i.e. $\alpha + \alpha^{-1} \in \mathbb{N}$.
- $(a, b) \in \mathbb{R}^2 \times \mathbb{R}^2$ is a linearly independent pair which is α -compatible, i.e.

$$N(\alpha, a, b) := \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}^{-1} \in M_2(\mathbb{Z}).$$

- $c \in \mathbb{R}^2$ is (a, b, r) -compatible in the sense of the compatibility condition (2).

The set of (a, b, r) -compatible vectors is identified with \mathbb{Z}^2 via a bijection which depends on (a, b, r) . Therefore we will replace Inoue's pair $(p, q) \in \mathbb{Z}^2$ by the pair of coefficients $c = (c_1, c_2)$ which intervenes effectively in the construction of the surface as a quotient [In, p. 276].

For any S^+ -admissible $\alpha \in]1, +\infty[$, let $\mathcal{N}_\alpha \subset \text{SL}(2, \mathbb{Z})$ be the set of $\text{SL}(2, \mathbb{Z})$ matrices with $\alpha \in \text{Spec}(N)$. By the Latimer and MacDuffee Theorem, the set of similarity classes $\mathcal{N}_\alpha / \sim$ is finite.

The pair (α, r) is a biholomorphism invariant. We will identify the set of biholomorphism classes of type II Inoue surfaces $S_{a,b,c,t}^{\alpha,r}$ with $N(\alpha, a, b) \in \mathcal{N}_\alpha$ with a quotient set $\mathcal{Q}_{\alpha,r}$ of the set of parameters (a, b, c, t) , where (a, b, c) satisfies the above compatibility conditions (see Theorem 4.26). We have a natural map

$$\Psi_{\alpha,r} : \mathcal{Q}_{\alpha,r} \rightarrow \mathcal{N}_\alpha / \sim, [S_{a,b,c,t}^{\alpha,r}] \mapsto [N(\alpha, a, b)].$$

Let $\mathfrak{N} \in \mathcal{N}_\alpha / \sim$, choose $N = N(\alpha, a, b) \in \mathcal{N}_\alpha$, and let $Z_{\text{GL}(2, \mathbb{Z})}^+(N)$ be "the positive centraliser of N " in $\text{GL}(2, \mathbb{Z})$, which is an infinite cyclic group, see the definition formula (74) and section 5. We will prove (see Theorem 4.34) that the fibre $\Psi_{\alpha,r}^{-1}(\mathfrak{N})$ over \mathfrak{N} comes with a natural topology and that the set of connected components of this fibre can be naturally identified with the quotient of

$$\mathbb{Z}_{N,r} := \mathbb{Z}^2 / (I_2 - N)\mathbb{Z}^2 + r\mathbb{Z}^2$$

by the group $Z_{\text{GL}(2, \mathbb{Z})}^+(N)$ acting on $\mathbb{Z}_{N,r}$ by $K * [p] := [\varepsilon_K K p]$.

Moreover, we will prove that any connected components of the fibre $\Psi_{\alpha,r}^{-1}(\mathfrak{N})$ can be identified to either \mathbb{C}^* or \mathbb{C} , the latter case occurring when condition (C) stated in Proposition 4.37 is satisfied. Using Inoue's [In, Proposition 4 p. 278] one can prove easily that two type II Inoue surfaces $S_{a,b,c,t}^{\alpha,r}$, $S_{a',b',c',t'}^{\alpha',r'}$ are deformation equivalent if and only if $(\alpha, r) = (\alpha', r')$ and the images of the the 4-tuples (a, b, c, t) , (a', b', c', t') in the quotient space $\mathcal{Q}_{\alpha,r}$ belong to the same connected component.

Therefore, for any pair (α, r) as above with α admissible, we have only finitely many deformation classes of type II Inoue surfaces, namely a finite number of

¹This finiteness theorem is usually stated for a maximal order, i.e. for the ring of integers O_K of an algebraic number field K . In general the order $\mathbb{Z}[\alpha]$ is not maximal (see Example 4.1).

similarity classes of $\mathrm{SL}(2, \mathbb{Z})$ -matrices with eigenvalue α , and a finite number of deformation classes for each such similarity class.

At the end of section 4.2 we treat in detail the cases $\theta = 3$ ($\alpha = \frac{3+\sqrt{5}}{2}$) and $\theta = 4$ ($\alpha = 2 + \sqrt{3}$) specifying the number of deformation classes corresponding to any $r \in \mathbb{N}^*$ and the isomorphism type of the connected components of the space of isomorphism classes. The main tools are: the algorithmic version of the Latimer-MacDuffee theorem of [BVdM] and our Theorem 5.4 which specifies a generator of the positive centraliser $Z_{\mathrm{GL}(2, \mathbb{Z})}^+(N)$ for a large class of $\mathrm{SL}(2, \mathbb{Z})$ matrices.

We will use similar methods for the classification of type III surfaces, i.e. of the surfaces denoted traditionally by $S_{N,p,q,r}^-$. For this surfaces we will adopt the notation $S_{a,b,c}^{\alpha,r}$, where

- $r \in \mathbb{N}^*$ and $\alpha \in]1, +\infty[$ is S^- -admissible, i.e. $\alpha - \alpha^{-1} \in \mathbb{N}$.
- $(a, b) \in \mathbb{R}^2 \times \mathbb{R}^2$ is a linearly independent pair which is α -compatible, i.e.

$$N(\alpha, a, b) := \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha^{-1} \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}^{-1} \in M_2(\mathbb{Z}).$$

- $c \in \mathbb{R}^2$ is (a, b, r) -compatible in the sense of (11).

We define $\mathcal{N}_\alpha^- := \{N \in \mathrm{GL}(2, \mathbb{Z}) \mid \det(N) = -1, \alpha \in \mathrm{Spec}(N)\}$ and we let $\mathcal{Q}_{\alpha,r}$ denote the set of biholomorphism classes of type III Inoue surfaces $S_{a,b,c}^{\alpha,r}$ with $N(\alpha, a, b) \in \mathcal{N}_\alpha^-$. Our main result for this class of surfaces is Theorem 4.41, which states that the fibre of the natural map $\mathcal{Q}_{\alpha,r} \rightarrow \mathcal{N}_\alpha^-$ over a similarity class $\mathfrak{N} = [N(\alpha, a, b)] \in \mathcal{N}_\alpha^- / \sim$ can be identified with the quotient of

$$\mathbb{Z}_{N,r} := \mathbb{Z}^2 / (I_2 + N)\mathbb{Z}^2 + r\mathbb{Z}^2$$

by the group $Z_{\mathrm{GL}(2, \mathbb{Z})}^+(N)$ acting on $\mathbb{Z}_{N,r}$ by $K * [p] := [\varepsilon_K K p]$. The set $\mathcal{N}_\alpha^- / \sim$ of similarity classes associated with α is again finite, so, for any S^- -admissible α one obtains a finite number of biholomorphism classes of type III Inoue surfaces.

Note that ‘‘positive centraliser’’ $Z_{\mathrm{GL}(2, \mathbb{Z})}^+(N)$ associated with a matrix $N \in \mathcal{N}_\alpha$ ($N \in \mathcal{N}_\alpha^-$), where α is S^+ - (respectively S^- -)admissible, is always infinite cyclic (see section 5, where it is shown that specifying a generator of this group is an interesting number theoretical problem).

We will begin our article with a new presentation of Inoue’s constructions; for each one of the three types of Inoue surfaces we will precise the entire data system needed for the construction of a surface and we will use the notation $S_{\mathfrak{p}}$ for the complex surface associated with the data system (parameter) \mathfrak{p} varying in a set \mathfrak{P} which we will describe explicitly. Noting that surfaces of different types cannot be biholomorphic (see Corollary 3.6), it follows that the classification problem for Inoue surfaces reduces to giving – for every fixed type – an explicit description of the quotient set \mathfrak{P}/R , where R is the equivalence relation

$$R := \{(\mathfrak{p}', \mathfrak{p}'') \in \mathfrak{P} \times \mathfrak{P} \mid S_{\mathfrak{p}'} \simeq S_{\mathfrak{p}''}\}.$$

The main difficulty is to understand in detail the equivalence relation R .

Let $\mathrm{Aff}(\mathbb{C}^2)$, $\mathrm{T}(\mathbb{C}^2)$ be the groups of affine transformations, respectively translations, of \mathbb{C}^2 , $U := H \times \mathbb{C}$ and let $\mathrm{Aff}(U)$, $\mathrm{T}(U)$ be the subgroups

$$\mathrm{Aff}(U) := \{f \in \mathrm{Aff}(\mathbb{C}^2) \mid f(U) = U\}, \quad \mathrm{T}(U) := \{f \in \mathrm{T}(\mathbb{C}^2) \mid f(U) = U\}$$

consisting of those affine transformations (respectively translations) which fix U .

An elementary computation shows that

Remark 1.1. *We have*

$$\begin{aligned} \Gamma(U) &= \left\{ \begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{f} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix} \mid u \in \mathbb{R}, \zeta \in \mathbb{C} \right\}. \\ \text{Aff}(U) &= \left\{ \begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{f} \begin{pmatrix} \mu & 0 \\ \lambda & \nu \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix} \mid \mu \in \mathbb{R}_{>0}, \nu \in \mathbb{C}^*, u \in \mathbb{R}, (\lambda, \zeta) \in \mathbb{C}^2 \right\}. \end{aligned}$$

Any Inoue surface is the quotient $S = U/\Gamma$, where Γ is a subgroup of the group $\text{Aff}(U)$ acting properly discontinuously on U . Our classification results are based on the following theorem:

Theorem 4.1. *Let $S' = U/\Gamma'$, $S'' = U/\Gamma''$ be Inoue surfaces, $f : S' \rightarrow S''$ be a biholomorphism and $\tilde{f} : U \rightarrow U$ be a lift of f . Then $\tilde{f} \in \text{Aff}(U)$.*

The proof of this theorem is based on a unicity theorem: the tangent bundle of any Inoue surface admits a unique holomorphic connection (see Theorem 3.9 in this article and also [Du, Proposition 3.1] which concerns torsion free holomorphic connections).

Theorem 4.1 gives a purely algebraic interpretation of the set of isomorphism classes of Inoue surfaces:

Corollary 4.2. *Two Inoue surfaces $S' = U/\Gamma'$, $S'' = U/\Gamma''$ are biholomorphic if and only if the subgroups Γ' , Γ'' of the group $\text{Aff}(U)$ belong to the same conjugacy class.*

Using this result we will obtain, for each type of Inoue surfaces, a purely algebraic interpretation of the equivalence relation R on the parameter set \mathfrak{P} , which will allow us to describe the quotient set \mathfrak{P}/R .

2. THE CONSTRUCTION OF INOUE SURFACES

2.1. Type I Inoue surfaces. We begin with the following definition:

Definition 2.1. *A complex number β will be called admissible if $\beta \in \mathbb{C}^* \setminus \mathbb{R}$, $|\beta| < 1$, and $|\beta|^{-2}$, β and $\bar{\beta}$ are the roots of a polynomial $P(X) \in \mathbb{Z}[X]$ of the form*

$$P(X) = X^3 - \theta_2 X^2 + \theta_1 X - 1 \text{ with } \theta_1, \theta_2 \in \mathbb{Z}.$$

Equivalently, $\beta \in \mathbb{C} \setminus \mathbb{R}$ with $|\beta| < 1$ is admissible if

$$\begin{cases} |\beta|^{-2} + 2\Re(\beta) \in \mathbb{Z} \\ 2|\beta|^{-2}\Re(\beta) + |\beta|^2 \in \mathbb{Z}. \end{cases}$$

Remark 2.2. *The polynomial $P(X) = X^3 - \theta_2 X^2 + \theta_1 X - 1 \in \mathbb{Z}[X]$ associated with an admissible complex number β is irreducible in $\mathbb{Q}[X]$.*

Proof. It suffices to prove that $\alpha := |\beta|^{-2}$ is not rational. Suppose that $\alpha \in \mathbb{Q}$. $P(X)$ is a monic polynomial with integer coefficients, so its root α is integer over \mathbb{Z} . Since \mathbb{Z} is integrally closed in \mathbb{Q} , it follows that $\alpha \in \mathbb{Z}$, so $P(X)$ decomposes as $P(X) = (X - \alpha)(X^2 + (\alpha - \theta_2)X + \alpha(\alpha - \theta_2) + \theta_1)$. This implies $\alpha(\alpha(\alpha - \theta_2) + \theta_1) = 1$, so $\alpha^{-1} = |\beta|^2 \in \mathbb{Z}$, which contradicts the assumption $|\beta| < 1$. ■

Definition 2.3. *Let β be an admissible complex number. A pair $(a, b) \in \mathbb{R}^3 \times \mathbb{C}^3$ will be called β -compatible if a , b and \bar{b} are linearly independent over \mathbb{C} and*

$$M(\beta, a, b) := \begin{pmatrix} a_1 & b_1 & \bar{b}_1 \\ a_2 & b_2 & \bar{b}_2 \\ a_3 & b_3 & \bar{b}_3 \end{pmatrix} \begin{pmatrix} |\beta|^{-2} & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \bar{\beta} \end{pmatrix} \begin{pmatrix} a_1 & b_1 & \bar{b}_1 \\ a_2 & b_2 & \bar{b}_2 \\ a_3 & b_3 & \bar{b}_3 \end{pmatrix}^{-1} \in M_3(\mathbb{Z})$$

We put:

$$\mathcal{P}_\beta = \{(a, b) \in \mathbb{R}^3 \times \mathbb{C}^3 \mid (a, b) \text{ is } \beta\text{-compatible}\}.$$

Note that

Remark 2.4. *Let (a, b) be a β -compatible pair. Then*

- (1) $M(\beta, a, b) \in \mathrm{SL}(3, \mathbb{Z})$.
- (2) $\mathrm{Spec}(M(\beta, a, b)) = \{|\beta|^{-2}, \beta, \bar{\beta}\}$.
- (3) *The eigenspaces of $M(\beta, a, b)$ in \mathbb{C}^3 are:*

$$E_{|\beta|^{-2}} = \mathbb{C}a, \quad E_{\beta} = \mathbb{C}b, \quad E_{\bar{\beta}} = \mathbb{C}\bar{b}.$$

For a pair $(a, b) \in \mathbb{R}^3 \times \mathbb{C}^3$ we define the affine transformations $g_i(a, b) \in \mathrm{Aff}(U)$, $1 \leq i \leq 3$, by

$$g_i(a, b) : (w, z) \rightarrow (w + a_i, z + b_i).$$

Definition 2.5. *Let β be an admissible complex number and $(a, b) \in \mathcal{P}_{\beta}$. We define $g_0(\beta) \in \mathrm{Aff}(U)$ by*

$$g_0(\beta) : (w, z) \rightarrow (|\beta|^{-2}w, \beta z)$$

and we define $G(\beta, a, b)$ to be the group generated by $g_0(\beta)$ and $(g_i(a, b))_{1 \leq i \leq 3}$.

Note that $G(\beta, a, b)$ acts properly discontinuously on U [In]. The argument uses essentially the compatibility property of the pair (a, b) .

Definition 2.6. *Let β be an admissible complex number and $(a, b) \in \mathcal{P}_{\beta}$. We define the (first type) Inoue surface $S_{a,b}^{\beta}$ by*

$$S_{a,b}^{\beta} = U/G(\beta, a, b)$$

where $G(\beta, a, b)$ is the subgroup of $\mathrm{Aff}(U)$ generated by the affine automorphisms $g_0(\beta)$, $(g_i(a, b))_{1 \leq i \leq 3}$.

Therefore, for type I Inoue surfaces, the space of parameters \mathfrak{P} considered in the introduction is

$$\mathfrak{P} = \{(\beta, a, b) \in \mathbb{C} \times \mathbb{R}^3 \times \mathbb{C}^3 \mid \beta \text{ is admissible, } (a, b) \text{ is } \beta\text{-compatible}\}.$$

Note that

Remark 2.7. [In, p. 274] *The generators $g_0 := g_0(\beta)$, $g_i := g_i(a, b)$ of $G(\beta, a, b)$ satisfy the commutation relations*

$$\begin{aligned} g_i g_j &= g_j g_i \quad i, j = 1, 2, 3, \\ g_0 g_i g_0^{-1} &= g_1^{m_{i1}} g_2^{m_{i2}} g_3^{m_{i3}} \quad i = 1, 2, 3, \end{aligned}$$

where m_{ij} are the entries of the matrix $M := M(\beta, a, b)$.

Using Remark 2.7 we obtain the following simple

Remark 2.8. *With the notations introduced above we have:*

- (1) *Any element $g \in G(\beta, a, b)$ can be written in a unique way as*

$$g = g_3(a, b)^{k_3} g_2(a, b)^{k_2} g_1(a, b)^{k_1} g_0(\beta)^{k_0} \quad \text{with } k_i \in \mathbb{Z}.$$

- (2) *We have*

$$G(\beta, a, b) \cap \mathrm{T}(U) = \langle g_1(a, b), g_2(a, b), g_3(a, b) \rangle \simeq \mathbb{Z}^3.$$

Proof. The first statement follows using the fact that the subgroup

$$\langle g_1(a, b), g_2(a, b), g_3(a, b) \rangle \subset G(\beta, a, b)$$

is abelian and normal in $G(\beta, a, b)$. The second statement follows from the first taking into account that $|\beta| < 1$. ■

2.2. Type II Inoue surfaces. In this section we will need the following subgroups of the group $\text{Aff}(U)$ of affine transformations of $U := H \times \mathbb{C}$:

$$\begin{aligned} \text{Aff}_1(U) &:= \left\{ \begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{g} \begin{pmatrix} \mu & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix} \middle| \mu \in \mathbb{R}_{>0}, u \in \mathbb{R}, \lambda \in \mathbb{C}, \zeta \in \mathbb{C} \right\}, \\ \text{Aff}_1^1(U) &:= \left\{ \begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{g} \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix} \middle| u \in \mathbb{R}, \lambda \in \mathbb{C}, \zeta \in \mathbb{C} \right\}, \\ \text{T}^0(U) &:= \left\{ \begin{pmatrix} w \\ z \end{pmatrix} \longrightarrow \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} 0 \\ \zeta \end{pmatrix} \middle| \zeta \in \mathbb{C} \right\}. \end{aligned}$$

Note that

Remark 2.9. (1) $\text{T}^0(U)$, $\text{T}(U)$, $\text{Aff}_1(U)$, $\text{Aff}_1^1(U)$ are normal subgroups of $\text{Aff}(U)$.
(2) $\text{T}^0(U)$ is central in $\text{Aff}_1(U)$, so also in $\text{Aff}_1^1(U)$.

We shall see that the subgroups which intervene in the construction of type II Inoue surfaces are all contained in $\text{Aff}_1(U)$.

Definition 2.10. A real number α will be called S^+ -admissible if $\alpha = \frac{\theta + \sqrt{\theta^2 - 4}}{2}$, where $\theta \in \mathbb{N}_{\geq 3}$.

Therefore α is S^+ -admissible if and only if $\alpha \in]1, +\infty[$ and $\alpha + \alpha^{-1} \in \mathbb{N}$, and if and only if it is the larger root of a quadratic equation $t^2 - \theta t + 1$ with $\theta \in \mathbb{N}_{\geq 3}$.

Definition 2.11. Let $\alpha \in \mathbb{R}$ be S^+ -admissible. A linearly independent pair $(a, b) \in \mathbb{R}^2 \times \mathbb{R}^2$ will be called α -compatible if

$$N(\alpha, a, b) := \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}^{-1} \in M_2(\mathbb{Z}). \quad (1)$$

Remark 2.12. Let α be as above and (a, b) be an α -compatible pair. Then

- (1) $N(\alpha, a, b) \in \text{SL}(2, \mathbb{Z})$.
- (2) $\text{Spec}(N(\alpha, a, b)) = \{\alpha, \alpha^{-1}\}$.
- (3) The eigenspaces of $N(\alpha, a, b)$ in \mathbb{R}^2 are $E_\alpha = \mathbb{R}a$, $E_{\alpha^{-1}} = \mathbb{R}b$.

We put:

$$\begin{aligned} \mathcal{P}_\alpha &:= \{(a, b) \in \mathbb{R}^2 \times \mathbb{R}^2 \mid (a, b) \text{ is } \alpha\text{-compatible}\}, \\ \mathcal{N}_\alpha &:= \{N \in \text{SL}(2, \mathbb{Z}) \mid \alpha \in \text{Spec}(N)\}, \end{aligned}$$

and we denote by $\eta_\alpha : \mathcal{P}_\alpha \rightarrow \mathcal{N}_\alpha$ the map given by $\eta_\alpha(a, b) := N(\alpha, a, b)$.

Definition 2.13. Let (a, b) be an α -compatible pair and $r \in \mathbb{N}^*$.

- (1) A vector $c \in \mathbb{R}^2$ will be called (a, b, r) -compatible if, putting $N := N(\alpha, a, b)$, we have

$$(I_2 - N) \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) - \frac{b \wedge a}{2} \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} \in \frac{b \wedge a}{r} \mathbb{Z}^2. \quad (2)$$

If this is the case, we define $p(a, b, c, r) \in \mathbb{Z}^2$ by

$$(I_2 - N) \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) - \frac{b \wedge a}{2} \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} = \frac{b \wedge a}{r} p(a, b, c, r). \quad (3)$$

The set of (a, b, r) -compatible vectors will be denoted by $\mathcal{C}_{a, b, r}$.

- (2) If (a, b) is α -compatible and c is (a, b, r) -compatible we will also say that the triple (a, b, c) is (α, r) -compatible. Therefore, (a, b, c) is (α, r) -compatible if and only if

$$(a) \quad N(\alpha, a, b) := \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}^{-1} \in M_2(\mathbb{Z}),$$

(b) Putting $N = (n_{ij})_{i,j} = N(\alpha, a, b)$, we have

$$(I_2 - N) \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) - \frac{b \wedge a}{2} \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} \in \frac{b \wedge a}{r} \mathbb{Z}^2.$$

Remark 2.14. Our compatibility condition (2) is equivalent to Inoue's compatibility condition [In, formula (17) p. 276], which reads

$$(I_2 - N)c - e \in \frac{b \wedge a}{r} \mathbb{Z}^2,$$

where

$$\begin{pmatrix} e_1 \\ e_2 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} n_{11}(n_{11} - 1)a_1 b_1 + n_{12}(n_{12} - 1)a_2 b_2 \\ n_{21}(n_{21} - 1)a_1 b_1 + n_{22}(n_{22} - 1)a_2 b_2 \end{pmatrix} + \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} b_1 a_2.$$

Using the identities $Na = \alpha a$, $Nb = \alpha^{-1}b$, one obtains easily:

$$\begin{pmatrix} e_1 \\ e_2 \end{pmatrix} = \frac{1}{2} \left[\begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} - N \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} + (b \wedge a) \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} \right],$$

which proves the claim.

The Inoue surfaces of type II are quotients of U by a group of affine transformations constructed using an (α, r) -compatible triple and a complex parameter $t \in \mathbb{C}$:

Remark 2.15. Let $(a, b) \in \mathcal{P}_\alpha$ and $r \in \mathbb{N}^*$.

- (1) The set $\mathcal{C}_{a,b,r}$ is a $\frac{b \wedge a}{r}(I_2 - N)^{-1}(\mathbb{Z}^2)$ -torsor.
- (2) The map $\pi_{a,b,r} : \mathcal{C}_{a,b,r} \rightarrow \mathbb{Z}^2$, $\pi_{a,b,r}(c) := p(a, b, c, r) \in \mathbb{Z}^2$ is bijective and satisfies the identity

$$\pi_{a,b,r} \left(c + \frac{b \wedge a}{r} (I_2 - N)^{-1} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} \right) = \pi_{a,b,r}(c) + \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}.$$

Note that the two components of $p(a, b, c, r)$ correspond to the integers p, q in Inoue's notation.

Let $(a, b, c) \in \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$ and $r \in \mathbb{N}^*$. We define the affine transformations $g_i = g_i(a, b, c)$ ($1 \leq i \leq 2$), $g_3 = g_3(a, b, r) \in \text{Aff}(U)$, by

$$g_i \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} w + a_i \\ z + b_i w + c_i \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ b_i & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} a_i \\ c_i \end{pmatrix}, \quad g_3 \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} w \\ z + \frac{b \wedge a}{r} \end{pmatrix}. \quad (4)$$

Definition 2.16. Let α, r be as above and (a, b, c) be an (α, r) -compatible triple. We define $G(a, b, c, r)$ to be the subgroup of $\text{Aff}(U)$ generated by the affine transformations $g_1 = g_1(a, b, c)$, $g_2 = g_2(a, b, c)$, $g_3 = g_3(a, b, r)$.

For $t \in \mathbb{C}$ we also define $g_0 = g_0(\alpha, t) \in \text{Aff}(U)$ by

$$g_0(\alpha, t) \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} \alpha w \\ z + t \end{pmatrix}$$

and the subgroup $G(\alpha, a, b, c, r, t) \subset \text{Aff}(U)$ by

$$G(\alpha, a, b, c, r, t) := \langle g_0, g_1, g_2, g_3 \rangle.$$

Recall [In, p. 276] that $G(\alpha, a, b, c, r, t)$ acts properly discontinuously on U . The argument uses essentially the (α, r) -compatibility condition of the triple (a, b, c) . We define

Definition 2.17. Let α, r be as above and (a, b, c) be an (α, r) -compatible triple. The type II Inoue surface associated with the parameters (α, a, b, c, r, t) is

$$S_{a,b,c,t}^{\alpha,r} := U / G(\alpha, a, b, c, r, t).$$

Therefore, for type II Inoue surfaces, the space of parameters \mathfrak{P} considered in the introduction is

$$\mathfrak{P} = \{(\alpha, a, b, c, r, t) \in]1, +\infty[\times \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{N}^* \times \mathbb{C} \mid \alpha \text{ is } S^+ \text{-admissible, and } (a, b, c) \text{ is } (\alpha, r) \text{-compatible}\}.$$

Remark 2.18. [In, p. 276] *Let (a, b, c) be an (α, r) -compatible triple. Putting $p := p(a, b, c, r) \in \mathbb{Z}^2$, we have the commutation relations*

$$g_1^{-1} g_2^{-1} g_1 g_2 = g_3^r, \quad g_i g_3 = g_3 g_i \text{ for } 0 \leq i \leq 2, \quad g_0 g_i g_0^{-1} = g_1^{n_{i1}} g_2^{n_{i2}} g_3^{p_i},$$

where $N = (n_{ij})_{i,j} = N(\alpha, a, b)$.

Remark 2.19. *The subgroup $Z(a, b, r) := \langle g_3 \rangle \subset G(\alpha, a, b, c, r, t)$ is central in $G(\alpha, a, b, c, r, t)$ and $G(a, b, c, r)$ is a normal subgroup of $G(\alpha, a, b, c, r, t)$.*

(1) *We have*

$$Z(a, b, r) = G(a, b, c, r) \cap \mathbb{T}(U) = G(\alpha, a, b, c, r, t) \cap \mathbb{T}(U) \quad (5)$$

The proof uses the fact that b_1, b_2 are linearly independent over \mathbb{Q} (which follows easily taking into account that b is an eigenvector of a rational matrix associated with an irrational eigenvalue).

(2) *We have*

$$G(a, b, c, r) = G(\alpha, a, b, c, r, t) \cap \text{Aff}_1^1(U). \quad (6)$$

(3) *We have inclusions $G(\alpha, a, b, c, r, t) \subset \text{Aff}_1(U)$, $G(a, b, c, r) \subset \text{Aff}_1^1(U)$, and the induced morphism*

$$G(\alpha, a, b, c, r, t)/G(a, b, c, r) \rightarrow \text{Aff}_1(U)/\text{Aff}_1^1(U) \quad (7)$$

between the corresponding quotient groups is a monomorphism. Identifying the quotient $\text{Aff}_1(U)/\text{Aff}_1^1(U)$ with $\mathbb{R}_{>0}$ via the obvious isomorphism

$$\text{Aff}_1(U)/\text{Aff}_1^1(U) \xrightarrow{\cong} \mathbb{R}_{>0}$$

induced by $\begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{g} \begin{pmatrix} \mu & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix} \mapsto \mu$, we obtain a group isomorphism

$$G(r, \alpha, a, b, c, t)/G(a, b, c, r) = \langle [g_0] \rangle \xrightarrow{\cong} \langle \alpha \rangle.$$

onto the cyclic group $\langle \alpha \rangle \subset \mathbb{R}_{>0}$ which maps the class $[g_0]_{G(a,b,c,r)}$ to α .

Using Remark 2.18, we obtain:

Remark 2.20. *One has $[G(a, b, c, r), G(a, b, c, r)] = \langle g_3^r \rangle \subset \langle g_3 \rangle$, in particular one has an isomorphism*

$$\langle g_3 \rangle / [G(a, b, c, r), G(a, b, c, r)] \simeq \mathbb{Z}_r.$$

Remark 2.21. *Any element $g \in G(a, b, c, r)$ can be written in a unique way in the form*

$$g = g_1^{n_1} g_2^{n_2} g_3^k \quad (8)$$

with $n_1, n_2, k \in \mathbb{Z}$; any element $g \in G(\alpha, a, b, c, r, t)$ can be written in a unique way in the form

$$g = g_0^l g_1^{n_1} g_2^{n_2} g_3^k \quad (9)$$

with $n_1, n_2, k, l \in \mathbb{Z}$.

2.3. Type III Inoue surfaces. We start with the analogue of Definition 2.10 for type III surfaces:

Definition 2.22. A real number $\alpha \in]1, +\infty[$ will be called S^- -admissible if $\alpha = \frac{\theta + \sqrt{\theta^2 + 4}}{2}$, where $\theta \in \mathbb{N}^*$.

In other words α is S^- -admissible if it coincides with the positive root of a quadratic equation of the form $t^2 - \theta t - 1 = 0$ with $\theta \in \mathbb{N}^*$. Let α be S^- -admissible.

Definition 2.23. A linearly independent pair $(a, b) \in \mathbb{R}^2 \times \mathbb{R}^2$ will be called α -compatible if, putting $P_{ab} = \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}$, we have

$$N(\alpha, a, b) := P_{ab} \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha^{-1} \end{pmatrix} P_{ab}^{-1} \in M_2(\mathbb{Z}). \quad (10)$$

The space of α -compatible pairs in the sense of Definition 2.23 will be denoted by \mathcal{P}_α^- . For $(a, b) \in \mathcal{P}_\alpha^-$ we have

$$N(\alpha, a, b) \in \mathrm{GL}_-(2, \mathbb{Z}) := \{N \in \mathrm{GL}(2, \mathbb{Z}) \mid \det(N) = -1\}, \quad \mathrm{Tr}(N(\alpha, a, b)) = \theta,$$

and the eigenspaces of $N(\alpha, a, b)$ are $E_\alpha = \mathbb{R}a$, $E_{-\alpha^{-1}} = \mathbb{R}b$. For an S^- -admissible $\alpha \in]1, +\infty[$ we will use the notation:

$$\mathcal{N}_\alpha^- := \{N \in \mathrm{GL}_-(2, \mathbb{Z}) \mid \alpha \in \mathrm{Spec}(N)\}.$$

Definition 2.24. Let (a, b) be an α -compatible pair and $r \in \mathbb{N}^*$. An element $c \in \mathbb{R}^2$ is called (a, b, r) -compatible if, putting $N := N(\alpha, a, b)$, we have

$$(I_2 + N) \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) - \frac{a \wedge b}{2} \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} \in \frac{a \wedge b}{r} \mathbb{Z}^2. \quad (11)$$

If this is the case, we define $p(a, b, c, r) \in \mathbb{Z}^2$ by

$$(I_2 + N) \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) - \frac{a \wedge b}{2} \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} = \frac{a \wedge b}{r} \begin{pmatrix} p_1 \\ p_2 \end{pmatrix},$$

and we'll also say that (a, b, c) is an (α, r) -compatible triple. We will denote by $\mathcal{C}_{a,b,r}$ the space of (a, b, r) -compatible vectors and by $\mathcal{T}_{\alpha,r}$ the space of (α, r) -compatible triples.

Note that, as mentioned for type II surfaces, our compatibility condition is equivalent to Inoue's formula [In, (20), p. 279].

Definition 2.25. Let α, r be as above and (a, b, c) be an (α, r) -compatible triple. We define $G(a, b, c, r)$ to be the subgroup of $\mathrm{Aff}(U)$ generated by the affine transformations $g_1 = g_1(a, b, c)$, $g_2 = g_2(a, b, c)$, $g_3 = g_3(a, b, r)$ defined in (4). We also define $g_0 = g_0(\alpha) \in \mathrm{Aff}(U)$ by

$$g_0(\alpha) \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} \alpha w \\ -z \end{pmatrix}$$

and the subgroup $G(\alpha, a, b, c, r) \subset \mathrm{Aff}(U)$ by

$$G(\alpha, a, b, c, r) := \langle g_0, g_1, g_2, g_3 \rangle.$$

Recall [In, p. 279] that $G(\alpha, a, b, c, r)$ acts properly discontinuously on U . We define

Definition 2.26. Let α, r be as above and (a, b, c) be an (α, r) -compatible triple. The type III Inoue surface associated with the parameters (α, a, b, c, r) is

$$S_{a,b,c}^{\alpha,r} := U / G(\alpha, a, b, c, r).$$

Remark 2.27. [In, p. 279] Let (a, b, c) be an (α, r) -compatible triple. Putting $p := p(a, b, c, r) \in \mathbb{Z}^2$, we have the commutation relations

$$g_1^{-1} g_2^{-1} g_1 g_2 = g_3^r, \quad g_i g_3 = g_3 g_i \text{ for } 1 \leq i \leq 2, \quad g_0 g_3 g_0^{-1} = g_3^{-1}, \quad g_0 g_i g_0^{-1} = g_1^{n_{i1}} g_2^{n_{i2}} g_3^{p_i}$$

where $N = (n_{ij})_{i,j} = N(\alpha, a, b)$.

The statements of Remark 2.19 extend to type III surfaces with one notable exception: the cyclic subgroup $\langle g_3 \rangle$ is still central in the group $G(a, b, c, r)$, but not in $G(\alpha, a, b, c, r)$, because g_3 does not commute with g_0 . The obvious analogues of Remarks 2.20, 2.21 hold for type III surfaces.

3. FUNDAMENTAL PROPERTIES OF INOUE SURFACES

3.1. Line bundles on Inoue surfaces. Let X be a class VII surface, $x_0 \in X$ and $\pi_1(X, x_0)$ be the fundamental group of the pair (X, x_0) . Since $b_1(X) = 1$, the torsion free quotient $H_1(X, \mathbb{Z})/\text{Tors}(H_1(X, \mathbb{Z}))$ is always isomorphic to \mathbb{Z} . If $b_2(X) = 0$, the natural map

$$\text{Hom}(H_1(X, \mathbb{Z}), \mathbb{C}^*) = \text{Hom}(\pi_1(X, x_0), \mathbb{C}^*) \rightarrow \text{Pic}(X)$$

is an isomorphism, and the image of the natural monomorphism

$$\text{Hom}\left(\frac{H_1(X, \mathbb{Z})}{\text{Tors}(H_1(X, \mathbb{Z}))}, \mathbb{C}^*\right) \rightarrow \text{Hom}(H_1(X, \mathbb{Z}), \mathbb{C}^*) = \text{Pic}(X)$$

is precisely the identity component $\text{Pic}^0(X)$ of $\text{Pic}(X)$. Fixing an isomorphism $H_1(X, \mathbb{Z})/\text{Tors}(H_1(X, \mathbb{Z})) \xrightarrow{\cong} \mathbb{Z}$ gives an isomorphism

$$\text{Hom}(\mathbb{Z}, \mathbb{C}^*) = \mathbb{C}^* \xrightarrow{\cong} \text{Hom}\left(\frac{H_1(X, \mathbb{Z})}{\text{Tors}(H_1(X, \mathbb{Z}))}, \mathbb{C}^*\right) = \text{Pic}^0(X).$$

Any Inoue surface S is the quotient U/Γ where Γ is a group of affine transformations acting properly discontinuously on $U := H \times \mathbb{C} \subset \mathbb{C}^2$. We choose $u_0 \in U$, we put $x_0 = [u_0] \in S$ and we identify $\pi_1(S, x_0)$ with Γ in the obvious way. The canonical isomorphism $\text{Hom}(\pi_1(S, x_0), \mathbb{C}^*) = \text{Hom}(\Gamma, \mathbb{C}^*) \rightarrow \text{Pic}(S)$ is given explicitly by

$$\text{Hom}(\Gamma, \mathbb{C}^*) \ni \rho \mapsto [L_\rho] \in \text{Pic}(S),$$

where

$$L_\rho := U \times_{\mathbb{C}/\rho} \Gamma.$$

Here $U \times_{\mathbb{C}/\rho} \Gamma$ stands for the quotient of the trivial line bundle $U \times \mathbb{C}$ over U by the group Γ acting by

$$(g, ((w, z), \zeta)) \mapsto (g(w, z), \rho(g)\zeta).$$

For an Inoue surface S we also have a natural choice of an isomorphism

$$H_1(S, \mathbb{Z})/\text{Tors}(H_1(S, \mathbb{Z})) \rightarrow \mathbb{Z}$$

obtained using Inoue's description of the group $H_1(S, \mathbb{Z})$ (see [In, p. 274, 276, 279]):

Identifying $H_1(S, \mathbb{Z})$ with the abelianization $\Gamma/[\Gamma, \Gamma]$ of Γ , we see that, for all three types of Inoue surfaces, the classes $[g_i]$, $1 \leq i \leq 3$, generate $\text{Tors}(H_1(S, \mathbb{Z}))$, whereas the map

$$H_1(S, \mathbb{Z}) = \Gamma/[\Gamma, \Gamma] \ni [g_0^{k_0}][g_1^{k_1}][g_2^{k_2}][g_3^{k_3}] \mapsto k_0 \in \mathbb{Z}$$

is an epimorphism with kernel $\text{Tors}(H_1(S, \mathbb{Z}))$, so it induces an isomorphism

$$\kappa_0 : \frac{H_1(S, \mathbb{Z})}{\text{Tors}(H_1(S, \mathbb{Z}))} \rightarrow \mathbb{Z}.$$

For an Inoue surface S we will always use the isomorphism κ_0 defined above to identify the torsion free quotient $H_1(S, \mathbb{Z})/\text{Tors}(H_1(S, \mathbb{Z}))$ with \mathbb{Z} . The corresponding isomorphism

$$\mathbb{C}^* = \text{Hom}(\mathbb{Z}, \mathbb{C}^*) \xrightarrow{\cong} \text{Hom}\left(\frac{H_1(S, \mathbb{Z})}{\text{Tors}(H_1(S, \mathbb{Z}))}, \mathbb{C}^*\right) = \text{Pic}^0(S)$$

is given explicitly by

$$\zeta \mapsto [L_{\rho_\zeta}],$$

where $\rho_\zeta : \Gamma \rightarrow \mathbb{C}^*$ is defined by

$$\rho_\zeta(g_i) = 1 \text{ for } 1 \leq i \leq 3, \quad \rho_\zeta(g_0) = \zeta.$$

We will put $L_\zeta := L_{\rho_\zeta}$ to save on notations.

The main result of [In] is the following fundamental theorem:

Theorem 3.1. (Inoue) *Let S be an Inoue surface. Then S has no complex curve.*

In other words an Inoue surface has no positive effective divisor. This implies:

Corollary 3.2. *Let S be an Inoue surface and \mathcal{L} be a non-trivial invertible sheaf on S . Then $H^0(S, \mathcal{L}) = 0$.*

Indeed, if $H^0(S, \mathcal{L}) \neq 0$, the zero locus of a section $s \in H^0(S, \mathcal{L}) \setminus \{0\}$ must be empty, so s is nowhere vanishing, so it defines a global isomorphism $\mathcal{O}_X \rightarrow \mathcal{L}$.

3.2. The tangent bundle of an Inoue surface. Denoting by (e_w, e_z) the canonical basis of \mathbb{C}^2 , note that in all cases the line subbundle $U \times \mathbb{C}e_z \subset T_U = U \times \mathbb{C}^2$ is invariant under the tangent map g_* for any $g \in \Gamma$. Therefore we obtain a line subbundle M of T_S defined as the quotient of $U \times \mathbb{C}e_z$ by the induced Γ -action on the tangent bundle T_U . Putting $L := T_S/M$, we obtain a short exact sequence

$$0 \rightarrow M \hookrightarrow T_S \rightarrow L \rightarrow 0, \quad (12)$$

which will be called *the canonical exact sequence* for the tangent bundle of S .

Note that:

Remark 3.3. *For Inoue surfaces of type I, the line subbundle $U \times \mathbb{C}e_w \subset T_U$ is also Γ -invariant and its Γ -quotient is a complement of M in T_S , so it can be identified with L . Therefore, in this case, the canonical exact sequence (12) splits giving a direct sum decomposition $T_S = M \oplus L$.*

Identifying the line subbundles $U \times \mathbb{C}e_z \subset T_U$, $U \times \mathbb{C}e_w \subset T_U$ with the trivial line bundle $U \times \mathbb{C}$, and taking into account the explicit formulae for the generators g_i of Γ in each case, we see that M, L can be identified respectively with the quotients

$$(U \times \mathbb{C})/\rho_M \Gamma, \quad (U \times \mathbb{C})/\rho_L \Gamma$$

where the group morphisms $\rho_M, \rho_L \in \text{Hom}(\Gamma, \mathbb{C}^*)$ are given by the table below

type	ρ	$\rho(g_0)$	$\rho(g_1)$	$\rho(g_2)$	$\rho(g_3)$
I	ρ_M	β	1	1	1
	ρ_L	α	1	1	1
II	ρ_M	1	1	1	1
	ρ_L	α	1	1	1
III	ρ_M	-1	1	1	1
	ρ_L	α	1	1	1

(13)

Using the notation introduced above, this shows that

Remark 3.4. For any Inoue surface S we have a natural identification $L = L_\alpha$. For an Inoue surface of type I (respectively II, III), we have a natural identification $M = L_\beta$ (respectively $M = L_1 = S \times \mathbb{C}$, $M = L_{-1}$).

Note also that (12) gives natural identifications

$$K_S^* = \det(T_S) = M \otimes L = \begin{cases} L_{\alpha\beta} & \text{if } S \text{ is of type I} \\ L_\alpha & \text{if } S \text{ is of type II} \\ L_{-\alpha} & \text{if } S \text{ is of type III} \end{cases}. \quad (14)$$

Remark 3.5. For an Inoue surface of type II, the canonical exact sequence (12) becomes

$$0 \rightarrow \mathcal{O}_S \rightarrow \mathcal{T}_S \rightarrow \mathcal{K}_S^* \rightarrow 0, \quad (15)$$

and for an Inoue surface of type III, the canonical exact sequence (12) becomes

$$0 \rightarrow \mathcal{L}_{-1} \rightarrow \mathcal{T}_S \rightarrow \mathcal{K}_S^* \otimes \mathcal{L}_{-1} \rightarrow 0. \quad (16)$$

In both cases the extension class $h \in \text{Ext}^1(\mathcal{L}, \mathcal{M}) = H^1(S, \mathcal{K}_S)$ is non-trivial.

Proof. Suppose first that S is of type II. In this case, taking duals in the short exact sequence (15) we obtain

$$0 \rightarrow \mathcal{K}_S \rightarrow \Omega_S \rightarrow \mathcal{O}_S \rightarrow 0, \quad (17)$$

and the extension classes of (15), (17) coincide. But (17) does not split because, S being a class VII surface, we have $h_S^{10} := \dim(H^0(S, \Omega_S)) = 0$.

Suppose now that S is of type III. As noted in [In, p. 279], S admits a double cover \tilde{S} which is an Inoue surface of type II. Moreover, the canonical short exact sequence of \tilde{S} is the pull-back of the canonical short exact sequence of S . Since the latter is not split, the canonical short exact sequence of S cannot be split. ■

Corollary 3.6. 1. A Inoue surface

- (1) is of type I if and only if its tangent bundle has two (different) line subbundles.
- (2) is of type II if and only if it has its tangent bundle has a unique line subbundle and this line subbundle is trivial.
- (3) is of type III if and only if it has its tangent bundle has a unique line subbundle and this line subbundle is non-trivial.

2. If two Inoue surfaces are biholomorphic, then they have the same type.

Proof. The first statements follow by Remarks 3.3, 3.5. The second statement follows from the first. ■

3.3. Vanishing theorems. For a vector bundle E (a locally free sheaf \mathcal{E}) we will denote by $\text{End}_0(E)$ (respectively $\mathcal{E}nd_0(\mathcal{E})$) the bundle (sheaf) of trace-free endomorphisms of E (respectively \mathcal{E}). The goal of this section is the following vanishing theorem:

Theorem 3.7. Let S be an Inoue surface. Then

- (1) $H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) = 0$.
- (2) $H^0(S, \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S)) = 0$.

Proof. If S is of type I, we have

$$\begin{aligned} \mathcal{T}_S &\simeq \mathcal{M} \oplus \mathcal{L}, \quad \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S) \simeq \mathcal{M}^{*\otimes 2} \oplus \mathcal{L}^{*\otimes 2} \oplus \mathcal{K}_S^{\oplus 2}, \\ \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S) &= (\mathcal{L}^{*\otimes 2} \otimes \mathcal{M}) \oplus (\mathcal{M}^{*\otimes 2} \otimes \mathcal{L}) \oplus \mathcal{M}^{*\oplus 3} \oplus \mathcal{L}^{*\oplus 3}. \end{aligned}$$

We apply Corollary 3.2 to the invertible sheaves \mathcal{M}^* , \mathcal{L}^* , $\mathcal{M}^{*\otimes 2}$, $\mathcal{L}^{*\otimes 2}$, $\mathcal{L}^{*\otimes 2} \otimes \mathcal{M}$, $\mathcal{M}^{*\otimes 2} \otimes \mathcal{L}$ and \mathcal{K}_S . These sheaves are non-trivial: this follows using (13) taking

into account that $1 \notin \{\alpha^{-1}, \beta^{-1}, \alpha^{-2}, \beta^{-2}, \alpha^{-2}\beta, \beta^{-2}\alpha, \alpha^{-1}\beta^{-1}\}$.

Suppose that S is of type II.

(1) Since

$$\mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S) = \mathcal{K}_S \otimes (\mathcal{O}_S \text{id}_{\mathcal{T}_S} \oplus \mathcal{E}nd_0(\mathcal{T}_S)) \simeq \mathcal{K}_S \oplus (\mathcal{K}_S \otimes \mathcal{E}nd_0(\mathcal{T}_S)),$$

and \mathcal{K}_S is non-trivial, it suffices to prove that $H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd_0(\mathcal{T}_S)) = 0$. We will make use of Lemma 3.8 proved below applied to the exact sequence (15). Tensorizing by \mathcal{K}_S the last two lines of the first diagram, we obtain the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{K}_S \otimes \mathcal{F}_0 & \xrightarrow{i_{\mathcal{F}_0}} & \mathcal{K}_S \otimes \mathcal{E}nd_0(\mathcal{T}_S) & \xrightarrow{(qj)_0} & \mathcal{O}_S \longrightarrow 0 \\ & & \downarrow (u,v)_0 & & \downarrow \circ j & & \downarrow \text{id} \\ 0 & \longrightarrow & \mathcal{K}_S & \xrightarrow{\circ j} & \mathcal{K}_S \otimes \mathcal{T}_S & \xrightarrow{q^\circ} & \mathcal{O}_S \longrightarrow 0 \end{array} \quad (18)$$

with exact rows. We obtain the following commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \rightarrow & H^0(S, \mathcal{K}_S \otimes \mathcal{F}_0) & \hookrightarrow & H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd_0(\mathcal{T}_S)) & \rightarrow & \mathbb{C} \xrightarrow{\partial'} H^1(S, \mathcal{K}_S \otimes \mathcal{F}_0) \\ & & \downarrow & & \downarrow & & \downarrow \text{id} \\ 0 & \longrightarrow & H^0(S, \mathcal{K}_S) & \longrightarrow & H^0(S, \mathcal{K}_S \otimes \mathcal{T}_S) & \longrightarrow & \mathbb{C} \xrightarrow{\partial''} H^1(S, \mathcal{K}_S) \end{array}, \quad (19)$$

where ∂' , ∂'' are the connecting morphisms associated with the two exact sequences in (18). Since the lower exact sequence in (18) is non-split (it is obtained by tensorizing (15) with \mathcal{K}_S), it follows that ∂'' is injective, so ∂' is also injective. Therefore the monomorphism

$$H^0(S, \mathcal{K}_S \otimes \mathcal{F}_0) \rightarrow H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd_0(\mathcal{T}_S))$$

is an isomorphism, so it suffices to prove that

$$H^0(S, \mathcal{K}_S \otimes \mathcal{F}_0) = 0. \quad (20)$$

Tensorizing by \mathcal{K}_S the left hand column in the first diagram of Lemma 3.8, we obtain the short exact sequence

$$0 \rightarrow \mathcal{K}_S^{\otimes 2} \rightarrow \mathcal{K}_S \otimes \mathcal{F}_0 \rightarrow \mathcal{K}_S \rightarrow 0.$$

Since $H^0(S, \mathcal{K}_S^{\otimes 2}) = H^0(S, \mathcal{K}_S) = 0$, the associated cohomology exact sequence gives $H^0(S, \mathcal{K}_S \otimes \mathcal{F}_0) = 0$, as claimed.

(2) Tensorizing the exact sequence (17) by $\mathcal{E}nd(\mathcal{T}_S)$, we obtain the exact sequence

$$0 \rightarrow \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S) \rightarrow \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S) \rightarrow \mathcal{E}nd(\mathcal{T}_S) \rightarrow 0, \quad (21)$$

which gives the long exact cohomology sequence

$$\begin{array}{ccccccc} 0 & \rightarrow & H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) & \rightarrow & H^0(S, \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S)) & \rightarrow & H^0(S, \mathcal{E}nd(\mathcal{T}_S)) \xrightarrow{\Delta} \\ & & & & & & \xrightarrow{\Delta} H^1(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) \rightarrow \dots, \end{array} \quad (22)$$

where Δ is the connecting morphism associated with (21).

Since $H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) = 0$ by (1), the vanishing of $H^0(S, \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S))$ follows from the following claims which will be proved below:

Claim: $\ker(\Delta : H^0(S, \mathcal{E}nd(\mathcal{T}_S)) \rightarrow H^1(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S))) = 0$.

For this Claim we use the following commutative diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S) & \longrightarrow & \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S) & \longrightarrow & \mathcal{E}nd(\mathcal{T}_S) \longrightarrow 0 \\
& & \downarrow \text{id}_{\mathcal{K}_S} \otimes \text{Tr} & & \downarrow \text{id}_{\Omega_S} \otimes \text{Tr} & & \downarrow \text{Tr} \\
0 & \longrightarrow & \mathcal{K}_S & \longrightarrow & \Omega_S & \longrightarrow & \mathcal{O}_S \longrightarrow 0
\end{array}, \quad (23)$$

where the vertical morphisms on the left are give by:

$$(\text{id}_{\Omega_S} \otimes \text{Tr})(\omega \otimes f) = \omega \otimes \text{Tr}(f) = \text{Tr}(f)\omega$$

By the functoriality of the connecting morphism, we obtain the commutative diagram

$$\begin{array}{ccc}
0 & \longrightarrow & H^0(S, \mathcal{E}nd(\mathcal{T}_S)) \xrightarrow{\Delta} H^1(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) \\
& & \downarrow H^0(\text{Tr}) \qquad \qquad \qquad \downarrow H^1(\text{id}_{\mathcal{K}_S} \otimes \text{Tr}) \\
0 & \longrightarrow & H^0(S, \mathcal{O}_S) = \mathbb{C} \xrightarrow{\partial} H^1(S, \mathcal{K}_S)
\end{array}$$

where ∂ is the connecting morphism associated with the lower short exact sequence in (23). We know that this exact sequence is non-split, so $\partial(1) \neq 0$, which shows that ∂ is injective.

Therefore the claim will be proved if we show that $H^0(\text{Tr})$ is injective. Using the cohomology long exact sequence associated with the short exact sequence

$$0 \rightarrow \mathcal{E}nd_0(\mathcal{T}_S) \hookrightarrow \mathcal{E}nd(\mathcal{T}_S) \xrightarrow{\text{Tr}} \mathcal{O}_S \rightarrow 0$$

we see that

$$\ker(H^0(\text{Tr}) : H^0(S, \mathcal{E}nd(\mathcal{T}_S)) \rightarrow H^0(S, \mathcal{O}_S)) = H^0(S, \mathcal{E}nd_0(\mathcal{T}_S)).$$

Therefore our claim will be proved if we show that $H^0(S, \mathcal{E}nd_0(\mathcal{T}_S)) = 0$. Using the central exact row in the first diagram in Lemma 3.8 and the vanishing of $H^0(S, \mathcal{K}_S^*)$, it suffices to prove that $H^0(S, \mathcal{F}_0) = 0$. This follows as in the proof of (1) using the vanishing of $H^0(S, \mathcal{K}_S)$, the commutative diagram

$$\begin{array}{ccc}
0 & & 0 \\
\downarrow & & \downarrow \\
H^0(S, \mathcal{K}_S) & \xrightarrow{\text{id}} & H^0(S, \mathcal{K}_S) \\
\downarrow & & \downarrow \\
H^0(S, \Omega_S) & \longrightarrow & H^0(S, \mathcal{F}_0) \\
\downarrow & & \downarrow \\
\mathbb{C} = H^0(S, \mathcal{O}_S) & \xrightarrow{\cong} & H^0(S, \mathcal{O}_S) = \mathbb{C} \\
\downarrow \partial_1 & & \downarrow \partial_2 \\
H^1(S, \mathcal{K}_S) & \xrightarrow{\text{id}} & H^1(S, \mathcal{K}_S)
\end{array} \quad (24)$$

associated with the second diagram in Lemma 3.8 and the injectivity of ∂_1 (which follows by Remark 3.5).

Finally suppose that S is of type III. In this case we can complete the proof in two ways: First method: we use again the canonical exact sequence and the associated commutative diagrams given by Lemma 3.8. Second method: we use

the double cover $\sigma : \tilde{S} \rightarrow S$ (with \tilde{S} of type II) as in the proof of Remark 3.4 and we note that

$$\begin{aligned}\sigma^*(\mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) &= \mathcal{K}_{\tilde{S}} \otimes \mathcal{E}nd(\mathcal{T}_{\tilde{S}}), \\ \sigma^*(\Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S)) &= \Omega_{\tilde{S}} \otimes \mathcal{E}nd(\mathcal{T}_{\tilde{S}}).\end{aligned}$$

Since σ is surjective, the natural pull-back morphisms

$$H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S)) \rightarrow H^0(\tilde{S}, \sigma^*(\mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S))) = H^0(\tilde{S}, \mathcal{K}_{\tilde{S}} \otimes \mathcal{E}nd(\mathcal{T}_{\tilde{S}}))$$

$$H^0(S, \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S)) \rightarrow H^0(\tilde{S}, \sigma^*(\Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S))) = H^0(\tilde{S}, \Omega_{\tilde{S}} \otimes \mathcal{E}nd(\mathcal{T}_{\tilde{S}}))$$

are injective, so the vanishing of $H^0(S, \mathcal{K}_S \otimes \mathcal{E}nd(\mathcal{T}_S))$, $H^0(S, \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S))$ follow from the vanishing of $H^0(\tilde{S}, \mathcal{K}_{\tilde{S}} \otimes \mathcal{E}nd(\mathcal{T}_{\tilde{S}}))$, $H^0(\tilde{S}, \Omega_{\tilde{S}} \otimes \mathcal{E}nd(\mathcal{T}_{\tilde{S}}))$. \blacksquare

Lemma 3.8. *Let X be a complex manifold and let*

$$0 \rightarrow \mathcal{M} \xrightarrow{j} \mathcal{E} \xrightarrow{q} \mathcal{Q} \rightarrow 0 \quad (25)$$

be an exact sequence of locally free sheaves, where \mathcal{M} , \mathcal{Q} are of rank 1. Let

$$qj : \mathcal{E}nd(\mathcal{E}) \rightarrow \mathcal{H}om(\mathcal{M}, \mathcal{Q}) = \mathcal{M}^* \otimes \mathcal{Q}, \quad (qj)_0 : \mathcal{E}nd_0(\mathcal{E}) \rightarrow \mathcal{H}om(\mathcal{M}, \mathcal{Q}) = \mathcal{M}^* \otimes \mathcal{Q}$$

be the morphisms defined $\varphi \mapsto q \circ \varphi \circ j$. Put

$$\mathcal{F} := \ker(qj), \quad \mathcal{F}_0 = \mathcal{F} \cap \mathcal{E}nd_0(\mathcal{E}) = \ker(qj)_0.$$

- (1) The sheaf morphism $jq : \mathcal{H}om(\mathcal{Q}, \mathcal{M}) \rightarrow \mathcal{E}nd(\mathcal{E})$ defined by $\varphi \mapsto j \circ \varphi \circ q$ takes values in \mathcal{F}_0 , so it defines a morphism $(jq)_0 : \mathcal{H}om(\mathcal{Q}, \mathcal{M}) \rightarrow \mathcal{F}_0$.
- (2) \mathcal{F} comes with natural morphisms $u : \mathcal{F} \rightarrow \mathcal{E}nd(\mathcal{M})$, $v : \mathcal{F} \rightarrow \mathcal{E}nd(\mathcal{Q})$ defined as follows: for any $x \in X$ and $\varphi \in \mathcal{F}_x$, we have $\varphi(\mathcal{M}_x) \subset \mathcal{M}_x$, so φ induces morphisms $u(\varphi) \in \mathcal{E}nd(\mathcal{M}_x)$, $v(\varphi) \in \mathcal{E}nd(\mathcal{E}_x/\mathcal{M}_x) = \mathcal{E}nd(\mathcal{Q}_x)$.
- (3) The restriction $(u, v)_0$ of the obtained morphism

$$(u, v) : \mathcal{F} \rightarrow \mathcal{E}nd(\mathcal{M}) \oplus \mathcal{E}nd(\mathcal{Q}) = \mathcal{O}_X \text{id}_{\mathcal{F}} \oplus \mathcal{O}_X \text{id}_{\mathcal{Q}}$$

to \mathcal{F}_0 takes values in the trivial invertible sheaf $\mathcal{O}_X(\text{id}_{\mathcal{M}}, -\text{id}_{\mathcal{Q}})$.

- (4) The composition of the morphism $(j \circ) : \mathcal{H}om(\mathcal{E}, \mathcal{M}) \rightarrow \mathcal{E}nd(\mathcal{E})$ with the trace-free morphism $\mathcal{E}nd(\mathcal{E}) \rightarrow \mathcal{E}nd_0(\mathcal{E})$ takes values in \mathcal{F}_0 , so it induces a morphism

$$(j \circ)_0 : \mathcal{H}om(\mathcal{E}, \mathcal{M}) \rightarrow \mathcal{F}_0.$$

- (5) We have commutative diagrams with exact lines and columns

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{H}om(\mathcal{Q}, \mathcal{M}) & \xrightarrow{\text{id}} & \mathcal{H}om(\mathcal{Q}, \mathcal{M}) & \longrightarrow & 0 \\ & & \downarrow (jq)_0 & & \downarrow i_{\mathcal{F}_0} \circ (jq)_0 & & \downarrow \\ 0 & \longrightarrow & \mathcal{F}_0 & \xleftarrow{i_{\mathcal{F}_0}} & \mathcal{E}nd_0(\mathcal{E}) & \xrightarrow{(qj)_0} & \mathcal{H}om(\mathcal{M}, \mathcal{Q}) \longrightarrow 0 \quad (26) \\ & & \downarrow (u, v)_0 & & \downarrow \circ j & & \downarrow \text{id} \\ 0 & \longrightarrow & \mathcal{O}_X(\text{id}_{\mathcal{M}}, -\text{id}_{\mathcal{Q}}) \simeq \mathcal{O}_X \text{id}_{\mathcal{M}} & \xrightarrow{\circ j} & \mathcal{H}om(\mathcal{M}, \mathcal{E}) & \xrightarrow{q \circ} & \mathcal{H}om(\mathcal{M}, \mathcal{Q}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

$$\begin{array}{ccc}
0 & & 0 \\
\downarrow & & \downarrow \\
\mathcal{H}om(\mathcal{Q}, \mathcal{M}) & \xrightarrow{\text{id}} & \mathcal{H}om(\mathcal{Q}, \mathcal{M}) \\
\downarrow \circ q & & \downarrow (jq)_0 \\
\mathcal{H}om(\mathcal{E}, \mathcal{M}) & \xrightarrow{(j \circ)_0} & \mathcal{F}_0 \\
\downarrow \frac{1}{2} \circ j & & \downarrow (u, v)_0 \\
\mathcal{H}om(\mathcal{M}, \mathcal{M}) = \mathcal{O}_X \text{id}_{\mathcal{M}} & \xrightarrow{\cong} & \mathcal{O}_X(\text{id}_{\mathcal{M}}, -\text{id}_{\mathcal{Q}}) \\
\downarrow & & \downarrow \\
0 & & 0
\end{array} \tag{27}$$

Proof. The statement follows from the analogue statement for an exact sequence of holomorphic bundles, which is a consequence of the analogue statement for an exact sequence $0 \rightarrow M \xrightarrow{j} E \xrightarrow{q} Q \rightarrow 0$ of vector spaces. The latter follows easily using a basis (u, v) of E with $u \in M$. ■

Using the second statement in Theorem 3.7, we obtain

Theorem 3.9. *Let $S = U/\Gamma$ be an Inoue surface. The tangent bundle T_S admits a unique holomorphic connection. The pull-back of this connection to U coincides with the trivial connection on $T_U = U \times \mathbb{C}^2$, in particular this connection is flat and torsion free.*

Proof. For the existence: the trivial connection ∇_0 on $T_U = U \times \mathbb{C}^2$ is holomorphic, flat, torsion free and Γ -invariant, so it descends to a flat, torsion free holomorphic connection on the quotient $S = U/\Gamma$.

For the unicity: if non-empty, the space of holomorphic connections on T_S is an affine space with model vector space $H^0(S, \Omega_S \otimes \mathcal{E}nd(\mathcal{T}_S))$. But this vector space vanishes by Theorem 3.7. ■

4. THE CLASSIFICATION OF INOUE SURFACES

Recall that any Inoue surface is the quotient $S = U/\Gamma$, where Γ is a subgroup of the group $\text{Aff}(U)$ acting properly discontinuously on U (see section 1).

Let $S' = U/\Gamma'$, $S'' = U/\Gamma''$ be Inoue surfaces and $p' : U \rightarrow S'$, $p'' : U \rightarrow S''$ be the corresponding covering projections. Let $f : S' \rightarrow S''$ be a biholomorphism. Since p'' and $f \circ p'$ are both universal covers of S'' , it follows that there exists a biholomorphism $\tilde{f} : U \rightarrow U$ lifting f , i.e. such that $p'' \circ \tilde{f} = f \circ p'$. This gives the commutative diagram

$$\begin{array}{ccc}
U & \xrightarrow{\tilde{f}} & U \\
\downarrow p' & & \downarrow p'' \\
S' = U/\Gamma' & \xrightarrow{f} & U/\Gamma'' = S''
\end{array} \tag{28}$$

and shows that, for such a lift \tilde{f} , we have:

$$\tilde{f} \circ \Gamma' \circ \tilde{f}^{-1} = \Gamma''. \tag{29}$$

An important role in our following results will be played by the following theorem, which reduces the classification of Inoue surfaces to an algebraic problem:

Theorem 4.1. *Let $S' = U/\Gamma'$, $S'' = U/\Gamma''$ be Inoue surfaces, $f : S' \rightarrow S''$ be a biholomorphism and $\tilde{f} : U \rightarrow U$ be a lift of f . Then $\tilde{f} \in \text{Aff}(U)$.*

Proof. Let ∇' , ∇'' be the (unique) holomorphic connections on the tangent bundles $T_{S'}$, $T_{S''}$ respectively. By Theorem 3.9 we know that $p'^*(\nabla') = p''^*(\nabla'') = \nabla_0$.

The pull-back $f^*(\nabla'')$ is a holomorphic connection on S' , so, by our unicity result, it coincides with ∇' . Therefore

$$\begin{aligned} \nabla_0 &= p'^*(\nabla') = p'^*(f^*(\nabla'')) = (f \circ p')^*(\nabla'') = \\ &= (p'' \circ \tilde{f})^*(\nabla'') = \tilde{f}^*(p''^*(\nabla'')) = \tilde{f}^*(\nabla_0). \end{aligned} \quad (30)$$

Therefore \tilde{f} is ∇_0 -affine. This implies that the tangent map \tilde{f}_* preserves the space $\langle \frac{\partial}{\partial w}, \frac{\partial}{\partial z} \rangle_{\mathbb{C}}$ of ∇_0 -parallel holomorphic vector fields on U . Therefore the Jacobian matrix of \tilde{f} is constant on U , so, since U is connected, \tilde{f} is the restriction to U of an affine transformation (which fixes U). ■

Taking into account Theorem 4.1 and formula (29), we obtain the following result, which reduces the classification of Inoue surfaces to a purely algebraic problem:

Corollary 4.2. *Two Inoue surfaces $S' = U/\Gamma'$, $S'' = U/\Gamma''$ are biholomorphic if and only if the subgroups Γ' , Γ'' of the group $\text{Aff}(U)$ belong to the same conjugacy class.*

Our next goal: based on this corollary, we will obtain explicit descriptions of isomorphism classes of Inoue surfaces of a fixed type.

4.1. The classification of type I Inoue surfaces. The goal of this section is Theorem 4.7 stated at the end of the section, which reduces the classification of type I Inoue surfaces to a purely algebraic problem.

Recall from section 2.1 that the space of parameters for type I Inoue surfaces is

$$\mathfrak{P} = \{(\beta, a, b) \in \mathbb{C} \times \mathbb{R}^3 \times \mathbb{C}^3 \mid \beta \text{ is admissible, } (a, b) \text{ is } \beta\text{-compatible}\}.$$

Taking into account Corollary 4.2, our problem becomes:

Problem. Let β, β' be admissible complex numbers. Let $(a, b) \in \mathcal{P}_\beta$ be a β -compatible pair, and $(a', b') \in \mathcal{P}_{\beta'}$ a β' -compatible pair (see section 2.1). Describe explicitly the quotient of \mathfrak{P} by the equivalence relation

$$(\beta, a, b) \sim (\beta', a', b') \text{ if } G(\beta, a, b), G(\beta', a', b') \text{ are conj. in } \text{Aff}(U).$$

Remark 4.3. *Let β be an admissible complex number. The formula*

$$(K, (\mu, \nu)) \cdot (a, b) := K \cdot ((\mu, \nu) \cdot (a, b)) := (K\mu a, K\nu b)$$

defines an action of the product $\text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^ \times \mathbb{C}^*)$ on the space \mathcal{P}_β .*

Proposition 4.4. *Let $(\beta, a, b), (\beta', a', b') \in \mathfrak{P}$. The following conditions are equivalent:*

- (1) $(\beta, a, b) \sim (\beta', a', b')$.
- (2) $\beta = \beta'$ and $(a', b') \in \text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*)(a, b)$.

Therefore the quotient set \mathfrak{P}/\sim is identified with the orbit space

$$\mathfrak{P}/\text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*) = \coprod_{\substack{\beta \in \mathbb{C} \\ \text{admissible}}} \mathcal{P}_\beta / \text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*).$$

Proof. Put $G := G(\beta, a, b)$, $G' := G(\beta', a', b')$,

$$g_i := g_i(a, b), g'_i := g_i(a', b') \text{ for } 1 \leq i \leq 3, g_0 := g_0(\beta), g'_0 := g_0(\beta').$$

First, suppose G, G' are conjugate in $\text{Aff}(U)$, and let $\theta \in \text{Aff}(U)$,

$$\theta \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} \mu & 0 \\ \lambda & \nu \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix} \text{ with } \mu \in \mathbb{R}_+^*, \nu \in \mathbb{C}^*, u \in \mathbb{R}, (\lambda, \zeta) \in \mathbb{C}^2$$

be such that

$$G(\beta', a', b') = \theta G(\beta, a, b) \theta^{-1}. \quad (31)$$

Formula (31) shows in particular that

$$g' := \theta \circ g_0 \circ \theta^{-1} \in G'. \quad (32)$$

We can write

$$g' = g_3^{l_{k_3}} g_2^{l_{k_2}} g_1^{l_{k_1}} g_0^{l_{k_0}}$$

with $k_i \in \mathbb{Z}$. Comparing the linear parts of g' and $\theta \circ g_0 \circ \theta^{-1}$ we obtain

$$\mu^{-1} \lambda (|\beta|^{-2} - \beta) = 0, |\beta|^{-2} = |\beta'|^{-2k_0}, \beta = \beta'^{k_0}, \quad (33)$$

in particular $\lambda = 0$ (i.e. the linear part of θ is diagonal) and (since $|\beta| < 1, |\beta'| < 1$) $k_0 \in \mathbb{N}^*$. Changing the roles of G and G' we obtain a formula of the form $\beta' = \beta^{l_0}$ with $l_0 \in \mathbb{N}^*$. It follows $\beta = \beta^{k_0 l_0}$, so $k_0 = l_0 = 1$ and $\beta = \beta'$.

Note first that (31) implies

$$\theta(G \cap \text{T}(U))\theta^{-1} = (\theta G \theta^{-1}) \cap (\theta \text{T}(U) \theta^{-1}) = (\theta G \theta^{-1}) \cap \text{T}(U) = G' \cap \text{T}(U). \quad (34)$$

Taking into account Remark 2.8(2) it follows that $\gamma'_i := \theta g_i \theta^{-1}$ belongs to

$$G' \cap \text{T}(U) = \langle g'_1, g'_2, g'_3 \rangle$$

for $1 \leq i \leq 3$, and $(\gamma'_1, \gamma'_2, \gamma'_3)$ is a system of generators of this free abelian group. In other words there exists $K \in \text{GL}(3, \mathbb{Z})$ such that

$$\gamma'_i = g_1^{l_{k_{i1}}} g_2^{l_{k_{i2}}} g_3^{l_{k_{i3}}} \text{ for } 1 \leq i \leq 3. \quad (35)$$

On the other hand, since $\lambda = 0$, we have

$$\gamma'_i \begin{pmatrix} w \\ z \end{pmatrix} = (\theta g_i \theta^{-1}) \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} w + \mu a_i \\ z + \nu b_i \end{pmatrix}. \quad (36)$$

Comparing (35) with (36) we obtain $a' = K^{-1} \mu a$, $b' = K^{-1} \nu b$, with $K^{-1} \in \text{GL}(3, \mathbb{Z})$, which proves the claim.

2. Conversely, suppose that $\beta = \beta'$ and $(a', b') \in \text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*)(a, b)$. Therefore there exists $(K, (\mu, \nu)) \in \text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*)$ such that

$$(a', b') := K \cdot ((\mu, \nu) \cdot (a, b)). \quad (37)$$

The linear transformation ϕ defined by the diagonal matrix $\begin{pmatrix} \mu & 0 \\ 0 & \nu \end{pmatrix}$ belongs to $\text{Aff}(U)$ and (37) shows that

$$\phi g_i \phi^{-1} = g_1^{l_{i1}} g_2^{l_{i2}} g_3^{l_{i3}} \text{ for } 1 \leq i \leq 3,$$

where $L = (l_{ij}) := K^{-1}$. Noting that $\phi g_0 \phi^{-1} = g_0 = g'_0$, it follows that

$$(\phi g_0 \phi^{-1}, \phi g_1 \phi^{-1}, \phi g_2 \phi^{-1}, \phi g_3 \phi^{-1})$$

is a system of generators of G' , so $\phi G \phi^{-1} = G'$. ■

Proposition 4.5. *Let β be an admissible complex number. Let $(a, b), (a', b') \in P_\beta$ be β -compatible pairs. The following conditions are equivalent:*

- (1) $(a', b') \in (\text{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*))(a, b)$

(2) $M(\beta, a, b) \sim M(\beta, a', b')$, where \sim denotes the similarity relation on $\mathrm{SL}(3, \mathbb{Z})$, i.e. the equivalence relation defined by conjugation with matrices in $\mathrm{GL}(3, \mathbb{Z})$.

Proof. (1) \Rightarrow (2): Suppose that $(a', b') \in \mathrm{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*)(a, b)$. Therefore there exists $(K, (\mu, \nu)) \in \mathrm{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*)$ such that

$$(a', b') := K \cdot ((\mu, \nu) \cdot (a, b)) := (K\mu a, K\nu b), \quad (38)$$

Then,

$$\begin{pmatrix} a'_1 & b'_1 & \bar{b}'_1 \\ a'_2 & b'_2 & \bar{b}'_2 \\ a'_3 & b'_3 & \bar{b}'_3 \end{pmatrix} = K \begin{pmatrix} a_1 & b_1 & \bar{b}_1 \\ a_2 & b_2 & \bar{b}_2 \\ a_3 & b_3 & \bar{b}_3 \end{pmatrix} \begin{pmatrix} \mu & 0 & 0 \\ 0 & \nu & 0 \\ 0 & 0 & \bar{\nu} \end{pmatrix},$$

so

$$M(\beta, a', b') = KM(\beta, a, b)K^{-1},$$

which proves the claim.

(2) \Rightarrow (1): Suppose that $M(\beta, a, b) \sim M'(\beta, a', b')$. Therefore there exists $K \in \mathrm{GL}(3, \mathbb{Z})$ such that $M(\beta, a, b) = KM'(\beta, a', b')K^{-1}$, so comparing the eigenspaces of the two matrices, it follows that $Ka' \in \mathbb{R}^*a$, $Kb' \in \mathbb{C}^*b$, so there exists $(\mathfrak{l}, \mathfrak{m}) \in \mathbb{R}^* \times \mathbb{C}^*$ such that

$$Ka' = \mathfrak{l}a, \quad Kb' = \mathfrak{m}b. \quad (39)$$

If $\mathfrak{l} > 0$, it follows that $(a', b') := K^{-1} \cdot ((\mathfrak{l}, \mathfrak{m}) \cdot (a, b))$, so the claim is verified. If $\mathfrak{l} < 0$, then $(-K)a' = (-\mathfrak{l})a$, $(-K)b' = (-\mathfrak{m})b$. Therefore

$$(a', b') := -K^{-1} \cdot ((-\mathfrak{l}, -\mathfrak{m}) \cdot (a, b)),$$

with $(-K^{-1}, ((-\mathfrak{l}, -\mathfrak{m})) \in \mathrm{GL}(3, \mathbb{Z}) \times (\mathbb{R}_+^* \times \mathbb{C}^*)$. \blacksquare

Combining Propositions 4.4, 4.5 and taking into account Corollary 4.2, we obtain:

Corollary 4.6. *Let β, β' be admissible complex numbers. Let $(a, b) \in P_\beta$ be a β -compatible pair, and let $(a', b') \in P_{\beta'}$ be a β' -compatible pair.*

Then

$$S_{a,b}^\beta \simeq S_{a',b'}^{\beta'} \iff \begin{cases} \beta = \beta' \\ \text{and} \\ M(\beta, a, b) \sim M(\beta', a', b') \end{cases},$$

where \sim denotes the similarity relation on $\mathrm{SL}(3, \mathbb{Z})$, i.e. the equivalence relation defined by conjugation with matrices in $\mathrm{GL}(3, \mathbb{Z})$.

Now put

$$\mathcal{M} := \{M \in \mathrm{SL}(3, \mathbb{Z}) \mid \mathrm{Spec}_{\mathbb{C}}(M) \supseteq \mathrm{Spec}_{\mathbb{R}}(M) \subset]1, +\infty[\}.$$

For a similarity class $\mathfrak{M} = [M] \in \mathcal{M}/\sim$, put $\mathrm{Spec}_{\mathbb{R}}(\mathfrak{M}) = \mathrm{Spec}_{\mathbb{R}}(M)$, $\mathrm{Spec}_{\mathbb{C}}(\mathfrak{M}) = \mathrm{Spec}_{\mathbb{C}}(M)$.

Note that knowing the similarity class of $M(\beta, a, b)$, β is only determined up to conjugation. With this remark, Corollary 4.6 becomes:

Theorem 4.7. *Biholomorphism classes of Inoue surfaces of type 1 correspond bijectively to elements of the set*

$$\left\{ (\mathfrak{M}, \beta) \mid \mathfrak{M} \in \mathcal{M}/\sim, \beta \in \mathrm{Spec}_{\mathbb{C}}(\mathfrak{M}) \setminus \mathrm{Spec}_{\mathbb{R}}(\mathfrak{M}) \right\} = \coprod_{\mathfrak{M} \in \mathcal{M}/\sim} (\mathrm{Spec}_{\mathbb{C}}(\mathfrak{M}) \setminus \mathrm{Spec}_{\mathbb{R}}(\mathfrak{M})).$$

The pair corresponding to the surface $S_{a,b}^\beta$ is $([M(\beta, a, b)], \beta)$.

Therefore,

Remark 4.8. *The set of biholomorphism classes of Inoue surfaces of type I comes with a natural surjective map onto the set \mathcal{M}/\sim of similarity class of matrices $M \in \mathcal{M}$. For any similarity class \mathfrak{M} we have two biholomorphism classes of type I Inoue surfaces.*

As explained in the introduction, for any polynomial $P(X) = X^3 - \theta_2 X^2 + \theta_1 X - 1 \in \mathbb{Z}[X]$ with a real root $\alpha \in]1, +\infty[$ and two non-real roots, the set of $\mathrm{GL}(3, \mathbb{Z})$ -similarity classes of matrices $M \in \mathrm{SL}(3, \mathbb{Z})$ with $\chi_M(X) = P(X)$ correspond bijectively to equivalence classes of ideals in the order $\mathbb{Z}[\alpha]$. This follows from the Latimer-MacDuffee Theorem taking into account Remark 2.2. Since the set of equivalence classes of ideals in an order is finite [BoSh, Theorem 3, p. 128], we obtain the following finiteness theorem:

Theorem 4.9. *To any polynomial $P(X) = X^3 - \theta_2 X^2 + \theta_1 X - 1 \in \mathbb{Z}[X]$ with a real root $\alpha \in]1, +\infty[$ and two non-real roots, correspond exactly $2h_{\mathbb{Z}[\alpha]}$ isomorphism classes of type I Inoue surfaces, where $h_{\mathbb{Z}[\alpha]}$ is the number of equivalence classes of ideals in the order $\mathbb{Z}[\alpha]$.*

Note that in general the order $\mathbb{Z}[\alpha] \subset K := \mathbb{Q}(\alpha)$ is not maximal, i.e. it does not necessarily coincide with the ring \mathcal{O}_K of integers of the algebraic number field K . We are grateful to Stéphane Louboutin for explaining us the following example:

Example 4.1. The polynomial $P_1(X) = X^3 - 2X^2 - 2X - 1$ has a single real root $\varepsilon \approx 2.83117$ and its discriminant is $\Delta_1 = -83$. Put $K = \mathbb{Q}(\varepsilon)$. Using [Mil, formula (8) p. 38] and the equality $\mathrm{disc}(1, \varepsilon, \varepsilon^2) = \Delta_1$ (see [Mil, p. 37]), we obtain

$$\Delta_1 = [\mathcal{O}_K : \mathbb{Z}[\varepsilon]]^2 \mathrm{disc}(\mathcal{O}_K).$$

Since Δ_1 is a prime number, it follows $\mathrm{disc}(\mathcal{O}_K) = \Delta_1 = -83$ and $[\mathcal{O}_K : \mathbb{Z}[\varepsilon]] = 1$, i.e. $\mathbb{Z}[\varepsilon] = \mathcal{O}_K$. Now note that $\alpha := \varepsilon^2 \approx 8.01556$ is the only real root of the polynomial $P_2(X) = X^3 - 8X^2 - 1$ whose discriminant is $\Delta_2 = 25\Delta_1$. Noting that $\alpha\varepsilon - 8\alpha - 1 = 0$, it follows $\mathbb{Q}(\alpha) = \mathbb{Q}(\varepsilon) = K$, and formula [Mil, formula (8) p. 38] applied to α gives $\Delta_2 = [\mathcal{O}_K : \mathbb{Z}[\alpha]]^2 \mathrm{disc}(\mathcal{O}_K)$, so $[\mathcal{O}_K : \mathbb{Z}[\alpha]] = 5$.

4.2. The classification of type II Inoue surfaces. Recall from section 2.2 that the space of parameters for type II Inoue surfaces is

$$\mathfrak{P} = \{(\alpha, a, b, c, r, t) \in]1, +\infty[\times \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{N}^* \times \mathbb{C} \mid \alpha \text{ is } S^+ \text{-admissible, and } (a, b, c) \text{ is } (\alpha, r) \text{-compatible}\}.$$

We begin with the simple

Remark 4.10. *Let $\mathfrak{p} = (\alpha, a, b, c, r, t)$, $\mathfrak{p}' = (\alpha', a', b', c', r', t') \in \mathfrak{P}$ such that $S_{\mathfrak{p}} \simeq S_{\mathfrak{p}'}$. Then $\alpha = \alpha'$, $r = r'$.*

Proof. By Corollary 4.2, there exists $\theta \in \mathrm{Aff}(U)$ such that

$$\theta G(\alpha, a, b, c, r, t) \theta^{-1} = G(\alpha', a', b', c', r', t').$$

In particular $\theta g_0(\alpha, t) \theta^{-1} = g'$ for an element $g' \in G(\alpha', a', b', c', r', t')$. Using Remark 2.21 and the explicit computation of the right hand term of formula (9), we obtain $\alpha = \alpha^l$ with $l \in \mathbb{Z}$. Since $\alpha, \alpha' > 1$, we have $l \in \mathbb{N}^*$. Changing the roles, we obtain $\alpha' = \alpha^{l'}$ with $l' \in \mathbb{N}^*$. It follows $ll' = 1$, so $l = l' = 1$, so $\alpha = \alpha'$.

We have

$$G(a, b, c, r) = G(\alpha, a, b, c, r, t) \cap \mathrm{Aff}_1^1, \quad G(a', b', c', r') = G(\alpha', a', b', c', r', t') \cap \mathrm{Aff}_1^1.$$

Since Aff_1^1 is normal in $\mathrm{Aff}(U)$ and $\theta G(\alpha, a, b, c, r, t) \theta^{-1} = G(\alpha', a', b', c', r', t')$, it follows

$$\theta G(a, b, c, r) \theta^{-1} = G(a', b', c', r'). \quad (40)$$

This implies

$$\theta(G(a, b, c, r) \cap T(U))\theta^{-1} = G(a', b', c', r') \cap (\theta T(U)\theta^{-1}) = G(a', b', c', r') \cap T(U),$$

so

$$\theta\langle g_3(a, b, r) \rangle\theta^{-1} = \langle g_3(a', b', r') \rangle.$$

On the other hand formula (40) also implies

$$\theta[G(a, b, c, r), G(a, b, c, r)]\theta^{-1} = [G(a', b', c', r'), G(a', b', c', r')].$$

Therefore the isomorphism $\langle g_3(a, b, r) \rangle \rightarrow \langle g_3(a', b', r') \rangle$ induced by the interior automorphism ι_θ maps the subgroup $[G(a, b, c, r), G(a, b, c, r)]$ of $\langle g_3(a, b, r) \rangle$ onto the subgroup $[G(a', b', c', r'), G(a', b', c', r')]$ of $\langle g_3(a', b', r') \rangle$. It follows that ι_θ induces an isomorphism

$$\begin{array}{ccc} \langle g_3(a, b, r) \rangle / [G(a, b, c, r), G(a, b, c, r)] & \xrightarrow{\cong} & \mathbb{Z}_r \\ \downarrow \cong & & \\ \langle g_3(a', b', r') \rangle / [G(a', b', c', r'), G(a', b', c', r')] & \xrightarrow{\cong} & \mathbb{Z}_{r'} \end{array} .$$

On the right we have used the isomorphisms given by Remark 2.20. So $r = r'$. ■

Remark 4.10 shows that, for any pair $(\alpha, r) \in]1, +\infty[\times \mathbb{N}^*$ with α S^+ -admissible, the subset $\mathfrak{P}_{\alpha, r}$ obtained from \mathfrak{P} by fixing α and r is saturated with respect to the equivalence relation defined by the condition $S_{\mathfrak{p}} \simeq S_{\mathfrak{p}'}$.

Therefore we will fix such a pair (α, r) and we will study the induced equivalence relation on the subset $\mathfrak{P}_{\alpha, r}$, which is obviously identified with the product

$$\mathcal{A}_{\alpha, r} := \mathcal{T}_{\alpha, r} \times \mathbb{C},$$

where $\mathcal{T}_{\alpha, r}$ is the space of (α, r) -compatible triples in the sense of Definition 2.13.

We will endow $\mathcal{T}_{\alpha, r}$ with its natural topology, namely with the coarsest topology which makes the maps

$$(a, b, c) \mapsto N(\alpha, a, b) \in \mathrm{SL}(2, \mathbb{Z}), (a, b, c) \mapsto \pi_{a, b, r}(c) \in \mathbb{Z}^2, (a, b, c) \mapsto (a, b) \in \mathbb{R}^2$$

continuous. With respect to this topology,

$$\{ \{ (\mu, \rho) \mapsto (\mu a, \rho b, \mu \rho c) \mid \mu, \rho \in]1 - \varepsilon, 1 + \varepsilon[\} \mid \varepsilon \in]0, 1[\}$$

is a fundamental system of neighbourhoods of $(a, b, c) \in \mathcal{T}_{\alpha, r}$, and we endow $\mathcal{A}_{\alpha, r}$ with the product topology. The quotient spaces we define will be endowed with the quotient topologies.

Remark 4.11. *Taking into account Corollary 4.2, our problem becomes:*

Problem. Let $(\alpha, r) \in]1, +\infty[\times \mathbb{N}^*$ with α S^+ -admissible. Describe explicitly the quotient of $\mathcal{A}_{\alpha, r} := \mathcal{T}_{\alpha, r} \times \mathbb{C}$ by the equivalence relation

$$(a, b, c, t) \sim (a', b', c', t') \text{ if } G(\alpha, a, b, c, r, t), G(\alpha, a', b', c', r, t') \text{ are conj. in } \mathrm{Aff}(U).$$

4.2.1. *Parameterizing the groups defining type II Inoue surfaces.* We will first study the quotient of $\mathcal{A}_{\alpha, r}$ by the finer equivalence relation

$$(a, b, c, t) \approx (a', b', c', t') \text{ if } G(\alpha, a, b, c, r, t) = G(\alpha, a', b', c', r, t').$$

The quotient set $\mathcal{A}_{\alpha, r} / \approx$ is obviously identified with the set of all subgroups $G \subset \mathrm{Aff}(U)$ defining type II Inoue surfaces.

Remark 4.12. Let $(a, b) \in \mathcal{P}_\alpha$, $c, c' \in \mathcal{C}_{a,b,r}$, $t, t' \in \mathbb{C}$. We have the equivalences

$$g_i(a, b, c') = g_i(a, b, c)g_3(r, a, b)^{s_i} \quad (1 \leq i \leq 2) \Leftrightarrow c' = c + \frac{b \wedge a}{r} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}. \quad (41)$$

$$g_0(\alpha, t') = g_0(\alpha, t)g_3(r, a, b)^{s_0} \Leftrightarrow t' = t + \frac{b \wedge a}{r} s_0. \quad (42)$$

This shows that the action

$$\left(\begin{pmatrix} s_0 \\ s_1 \\ s_2 \end{pmatrix}, (a, b, c, t) \right) \mapsto \left(a, b, c + \frac{b \wedge a}{r} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}, t + \frac{b \wedge a}{r} s_0 \right) \quad (43)$$

of the group \mathbb{Z}^3 on the parameter space $\mathcal{A}_{\alpha,r}$ is compatible with the equivalence relation \approx , i.e. we have

$$\forall \begin{pmatrix} s_0 \\ s_1 \\ s_2 \end{pmatrix} \in \mathbb{Z}^3 \quad \forall (a, b, c, t) \in \mathcal{A}_{\alpha,r}, \quad \left(a, b, c + \frac{b \wedge a}{r} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}, t + \frac{b \wedge a}{r} s_0 \right) \approx (a, b, c, t).$$

We will also need the action of \mathbb{Z}^2 on $\mathcal{T}_{\alpha,r}$ obtained by omitting the fourth component in (43).

For a triple $(a, b, c) \in \mathcal{T}_{\alpha,r}$ and $t \in \mathbb{C}$, put $[c]_{ab} := c + \frac{b \wedge a}{r} \mathbb{Z}^2$, $[t]_{ab} := t + \frac{b \wedge a}{r} \mathbb{Z}$ and define

$$\mathcal{C}_{a,b,r} := \mathcal{C}_{a,b,r} / \frac{b \wedge a}{r} \mathbb{Z}^2,$$

$$\mathcal{T}_{\alpha,r} := \{(a, b, [c]_{ab}) \mid (a, b) \in \mathcal{P}_\alpha, [c]_{ab} \in \mathcal{C}_{a,b,r}\} \simeq \mathcal{T}_{\alpha,r} / \mathbb{Z}^2,$$

$$\mathcal{A}_{\alpha,r} := \left\{ (a, b, [c]_{ab}, [t]_{ab}) \mid (a, b) \in \mathcal{P}_\alpha, [c]_{ab} \in \mathcal{C}_{a,b,r}, [t]_{ab} \in \mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right\} \simeq \mathcal{A}_{\alpha,r} / \mathbb{Z}^3.$$

Using Remark 2.15, we see that the bijection $\pi_{a,b,r} : \mathcal{C}_{a,b,r} \rightarrow \mathbb{Z}^2$ induces a bijection

$$\mathfrak{p}_{a,b,r} : \mathcal{C}_{a,b,r} \rightarrow \mathbb{Z}^2 / (I_2 - N) \mathbb{Z}^2 \simeq \mathbb{Z}_{e_1} \oplus \mathbb{Z}_{e_2},$$

where e_1, e_2 are the elementary divisors of the matrix $I_2 - N$.

Remark 4.13. The quotient $\mathcal{C}_{a,b,r} = \mathcal{C}_{a,b,r} / \frac{b \wedge a}{r} \mathbb{Z}^2$ is always finite. It reduces to a singleton if and only if $\theta = 3$.

Proof. The second statement follows from $\det(I_2 - N) = (1 - \alpha)(1 - \alpha^{-1}) = 2 - \theta$ recalling that $\theta \geq 3$. \blacksquare

We will regard $\mathcal{T}_{\alpha,r}, \mathcal{A}_{\alpha,r}$ as spaces over the set \mathcal{P}_α of α -compatible pairs. The fibre of $\mathcal{T}_{\alpha,r}$ over a pair (a, b) is identified with the finite set $\mathcal{C}_{a,b,r} \simeq \mathbb{Z}_{e_1} \oplus \mathbb{Z}_{e_2}$, whereas the fibre of $\mathcal{A}_{\alpha,r}$ over pair (a, b) is $\mathcal{C}_{a,b,r} \times (\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z}) \simeq (\mathbb{Z}_{e_1} \oplus \mathbb{Z}_{e_2}) \times \mathbb{C}^*$

The next step is to introduce a natural $\mathrm{GL}(2, \mathbb{Z})$ -action on the new parameter space $\mathcal{A}_{\alpha,r} \simeq \mathcal{A}_{\alpha,r} / \mathbb{Z}^3$. We start with the following elementary lemma:

Lemma 4.14. Let $(a, b, c) \in \mathcal{T}_{\alpha,r}$ and $K \in \mathrm{GL}(2, \mathbb{Z})$. Put $g_i := g_i(a, b, c)$ for $1 \leq i \leq 2$, $g_3 = g_3(a, b, r)$ and

$$G_1 := g_1^{k_{11}} g_2^{k_{12}}, \quad G_2 := g_1^{k_{21}} g_2^{k_{22}}, \quad G_3 := g_3^{\det(K)}. \quad (44)$$

Then $G_i = g_i(A, B, C)$ and $G_3 = g_3(A, B, r)$ (in the sense of formulae (4)), where

$$A = Ka, \quad B = Kb, \quad C = \frac{1}{2} \begin{pmatrix} A_1 B_1 \\ A_2 B_2 \end{pmatrix} + K \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) + \frac{b \wedge a}{2} \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix}. \quad (45)$$

Proposition 4.15. Let $(a, b, c) \in \mathcal{T}_{\alpha,r}$ and $K \in \mathrm{GL}(2, \mathbb{Z})$. Then

- (1) The formulae (45) define an element $(A, B, C) \in \mathcal{T}_{\alpha, r}$ which will be denoted $K \cdot (a, b, c)$.
- (2) The triple $(A, B, [C]_{AB}) \in \mathcal{T}_{\alpha, r}$ depends only on the matrix K and the triple $(a, b, [c]_{ab}) \in \mathcal{T}_{\alpha, r}$. Moreover, the formula

$$K \cdot (a, b, [c]_{ab}) := (A, B, [C]_{AB})$$

defines an action of $\mathrm{GL}(2, \mathbb{Z})$ on the space $\mathcal{T}_{\alpha, r}$.

The complicated formula for C is justified by the equivalence explained in Lemma 4.14 stated above. Note that formula $(K, (a, b, c)) \mapsto K \cdot (a, b, c)$ does *not* define a $\mathrm{GL}(2, \mathbb{Z})$ -action on $\mathcal{T}_{\alpha, r}$. Therefore factorising the space of triples $\mathcal{T}_{\alpha, r}$ by \mathbb{Z}^2 is an important step in our constructions.

Proof. (1) It's easy to see that (A, B) remains an α -compatible pair. The corresponding matrix is

$$N(\alpha, A, B) = KN(\alpha, a, b)K^{-1}.$$

It remains to check that C is (A, B, r) compatible, i.e. that, putting

$$\mathfrak{N} = (\mathfrak{n}_{ij}) := N(\alpha, A, B),$$

we have

$$(I_2 - \mathfrak{N}) \left(C - \frac{1}{2} \begin{pmatrix} A_1 B_1 \\ A_2 B_2 \end{pmatrix} \right) - \frac{B \wedge A}{2} \begin{pmatrix} \mathfrak{n}_{11} \mathfrak{n}_{12} \\ \mathfrak{n}_{21} \mathfrak{n}_{22} \end{pmatrix} \in \frac{B \wedge A}{r} \mathbb{Z}^2. \quad (46)$$

Putting $p := p(a, b, c, r)$, the first term on the left in (46) is

$$\begin{aligned} (I_2 - \mathfrak{N}) \left(C - \frac{1}{2} \begin{pmatrix} A_1 B_1 \\ A_2 B_2 \end{pmatrix} \right) &= K(I_2 - N) \left(c - \frac{1}{2} \begin{pmatrix} a_1 b_1 \\ a_2 b_2 \end{pmatrix} \right) + \frac{b \wedge a}{2} (I_2 - \mathfrak{N}) \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} = \\ &= b \wedge a \left[K \left(\frac{1}{2} \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} + \frac{1}{r} p \right) + \frac{1}{2} (I_2 - \mathfrak{N}) \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} \right]. \end{aligned}$$

Put $\varepsilon_K := \det(K) \in \{\pm 1\}$. The element P (a priori in \mathbb{R}^2) associated with (A, B, C, r) via formula (3) is

$$P = \varepsilon_K \left[Kp + \frac{r}{2} \left(K \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} + (I_2 - \mathfrak{N}) \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} - \varepsilon_K \begin{pmatrix} \mathfrak{n}_{11} \mathfrak{n}_{12} \\ \mathfrak{n}_{21} \mathfrak{n}_{22} \end{pmatrix} \right) \right] \quad (47)$$

It suffices to prove that

$$K \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} + \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} - \mathfrak{N} \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} - \varepsilon_K \begin{pmatrix} \mathfrak{n}_{11} \mathfrak{n}_{12} \\ \mathfrak{n}_{21} \mathfrak{n}_{22} \end{pmatrix} \in 2\mathbb{Z}^2.$$

Applying the elementary Lemma 4.16 stated below to the pairs (K, N) , (\mathfrak{N}, K) , we obtain

$$\begin{aligned} K \begin{pmatrix} n_{11} n_{12} \\ n_{21} n_{22} \end{pmatrix} + \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} - \begin{pmatrix} (KN)_{11} (KN)_{12} \\ (KN)_{21} (KN)_{22} \end{pmatrix} &\in 2\mathbb{Z}^2, \\ \mathfrak{N} \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} + \varepsilon_K \begin{pmatrix} \mathfrak{n}_{11} \mathfrak{n}_{12} \\ \mathfrak{n}_{21} \mathfrak{n}_{22} \end{pmatrix} - \begin{pmatrix} (\mathfrak{N}K)_{11} (\mathfrak{N}K)_{12} \\ (\mathfrak{N}K)_{21} (\mathfrak{N}K)_{22} \end{pmatrix} &\in 2\mathbb{Z}^2. \end{aligned}$$

It suffices to note that $\mathfrak{N}K = K N K^{-1} K = KN$.

- (2) For the first claim of (2) note that replacing $(a, b, c) \in \mathcal{T}_{\alpha, r}$ by (a, b, c') where $c' = \frac{b \wedge a}{r} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}$ in the formula for $K \cdot (a, b, c)$, gives the triple (A, B, C') , where

$$C' = C + \frac{b \wedge a}{r} K \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = C + \frac{B \wedge A}{r} \det(K)^{-1} K \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} \in C + \frac{B \wedge A}{r} \mathbb{Z}^2.$$

For the second claim of (2) we have to prove that for any $(a, b, c) \in \mathcal{T}_{\alpha, r}$ and $K, L \in \mathrm{GL}(2, \mathbb{Z})$, the triples $L \cdot (K \cdot (a, b, [c]_{ab}))$, $(LK) \cdot (a, b, [c]_{ab})$ coincide in $\mathcal{T}_{\alpha, r}$.

Put

$$\begin{aligned}\mathbb{G}_1 &:= G_1^{l_{11}} G_2^{l_{12}} = (g_1^{k_{11}} g_2^{k_{12}})^{l_{11}} (g_1^{k_{21}} g_2^{k_{22}})^{l_{12}}, \\ \mathbb{G}_2 &:= G_1^{l_{21}} G_2^{l_{22}} = (g_1^{k_{11}} g_2^{k_{12}})^{l_{21}} (g_1^{k_{21}} g_2^{k_{22}})^{l_{22}}, \\ \mathbb{G}_3 &:= G_3^{\det(L)} = g_3^{\det(LK)}.\end{aligned}$$

By Lemma 4.14 applied to $(A, B, C) \in \mathcal{T}_{\alpha, r}$ and $L \in \mathrm{GL}(2, \mathbb{Z})$, we have $\mathbb{G}_i = g_i(\mathfrak{A}, \mathfrak{B}, \mathfrak{C})$, where

$$(\mathfrak{A}, \mathfrak{B}, \mathfrak{C}) = L \cdot (A, B, C) = L \cdot (K \cdot (a, b, c)).$$

Put $R = LK$. Using the commutation formulae $[g_1, g_2] = g_3^r$ we obtain

$$\mathbb{G}_1 = g_1^{r_{11}} g_2^{r_{12}} g_3^{s_1(L, K)}, \quad \mathbb{G}_2 = g_1^{r_{21}} g_2^{r_{22}} g_3^{s_2(L, K)}$$

where $s_i(L, K) \in \mathbb{Z}$. Put $(\mathfrak{A}', \mathfrak{B}', \mathfrak{C}') := R(a, b, c)$, $\mathbb{G}'_i := g_i(\mathfrak{A}', \mathfrak{B}', \mathfrak{C}')$. We have

$$\mathfrak{A}' = (LK)a = LA = \mathfrak{A}, \quad \mathfrak{B}' = (LK)b = LB = \mathfrak{B}.$$

By Lemma 4.14, we have

$$\mathbb{G}'_1 = g_1^{r_{11}} g_2^{r_{12}}, \quad \mathbb{G}'_2 = g_1^{r_{21}} g_2^{r_{22}},$$

in other words for $1 \leq i \leq 2$ we have

$$g_i(\mathfrak{A}, \mathfrak{B}, \mathfrak{C}) = g_i(\mathfrak{A}', \mathfrak{B}', \mathfrak{C}') g_3(a, b, r)^{s_i(L, K)} = g_i(\mathfrak{A}', \mathfrak{B}', \mathfrak{C}') g_3(\mathfrak{A}, \mathfrak{B}, r)^{\det(R) s_i(L, K)}.$$

Taking into account the equivalence (41), this proves that

$$\mathfrak{C} = \mathfrak{C}' + \frac{\mathfrak{B} \wedge \mathfrak{A}}{r} \begin{pmatrix} \det(R) s_1(L, K) \\ \det(R) s_2(L, K) \end{pmatrix},$$

which proves the claim. ■

Lemma 4.16. *Let $K, L \in \mathrm{GL}(2, \mathbb{Z})$. Then*

$$\begin{aligned}L \begin{pmatrix} k_{11} k_{12} \\ k_{21} k_{22} \end{pmatrix} + \det(K) \begin{pmatrix} l_{11} l_{12} \\ l_{21} l_{22} \end{pmatrix} - \begin{pmatrix} (LK)_{11} (LK)_{12} \\ (LK)_{21} (LK)_{22} \end{pmatrix} = \\ = 2 \left(\frac{l_{11} - l_{11}^2}{2} k_{11} k_{12} + \frac{l_{12} - l_{12}^2}{2} k_{21} k_{22} \right) - 2k_{12} k_{21} \begin{pmatrix} l_{11} l_{12} \\ l_{21} l_{22} \end{pmatrix},\end{aligned}\tag{48}$$

which obviously belongs to $2\mathbb{Z}^2$ if $K, L \in \mathrm{GL}(2, \mathbb{Z})$.

Proof. Direct computation. ■

For a 4-tuple $((a, b, c), t) \in \mathcal{A}_{\alpha, r}$ we put

$$K \cdot (a, b, c, t) = (A, B, C, t)$$

We need: the class of $K \cdot (a, b, c, t) = (A, B, C, t) \bmod \frac{B \wedge A}{r} \mathbb{Z}^3$ depends only on K and the class $[a, b, c, t]$. This follows from the formula:

$$\begin{aligned}K \cdot \left(a, b, c + \frac{b \wedge a}{r} \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}, t + \frac{b \wedge a}{r} s_0 \right) \\ = (A, B, C + \frac{B \wedge A}{r} \det(K) K \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}, t + \frac{B \wedge A}{r} \det(K) s_0).\end{aligned}$$

Note also that the formula

$$K \cdot (a, b, [c]_{ab}, [t]_{ab}) := (A, B, [C]_{AB}, [t]_{AB})$$

defines a $\mathrm{GL}(2, \mathbb{Z})$ -action on $\mathcal{A}_{\alpha, r}$.

Theorem 4.17. *Let $(a, b, c, t), (a', b', c', t') \in \mathcal{A}_{\alpha, r}$. The following conditions are equivalent:*

- (1) $(a, b, c, t) \approx (a', b', c', t')$.
(2) $(a', b', [c']_{a'b'}, [t']_{a'b'}) \in \text{GL}(2, \mathbb{Z})(a, b, [c]_{ab}, [t]_{ab})$.

Therefore the quotient set $\mathcal{A}_{\alpha, r}/\approx$ is identified with the orbit space $\mathcal{A}_{\alpha, r}/\text{GL}(2, \mathbb{Z})$.

Proof. (1) \Rightarrow (2): Suppose that $(a, b, c, t) \approx (a', b', c', t')$, in other words we have $G(\alpha, a, b, c, r, t) = G(\alpha, a', b', c', r, t')$, in particular

$$g_0(\alpha, t'), g_3(a', b', r), g_i(a', b', c') \quad (1 \leq i \leq 2)$$

is a system of generators of $G(\alpha, a, b, c, r, t)$.

Put $G = G(\alpha, a, b, c, r, t) = G(\alpha, a', b', c', r, t')$. By Remark 2.19 we know that

$$(g_1(a, b, c), g_2(a, b, c), g_3(a, b, r)), (g_1(a', b', c'), g_2(a', b', c'), g_3(a', b', r))$$

are both system of generators of the group

$$G \cap \text{Aff}_1^1(U) = G(a, b, c, r) = G(a', b', c', r)$$

and $g_3(a, b, r), g_3(a', b', r)$ generate the same normal subgroup

$$Z := Z(a, b, r) = Z(a', b', r) = \langle g_3(a, b, r) \rangle = \langle g_3(a', b', r) \rangle.$$

The two pairs

$$([(g_1(a, b, c)]_Z, [(g_2(a, b, c)]_Z), [(g_1(a', b', c')]_Z, [(g_2(a', b', c')]_Z)$$

of congruence classes with respect to Z are both systems of generators of the quotient group $(G \cap \text{Aff}_1^1(U))/Z \simeq \mathbb{Z}^2$. It follows that there exists $K = (k_{ij}) \in \text{GL}(2, \mathbb{Z})$ such that

$$\begin{aligned} [g_1(a', b', c')]_Z &= [g_1(a, b, c)]_Z^{k_{11}} [g_2(a, b, c)]_Z^{k_{12}}, \\ [g_2(a', b', c')]_Z &= [g_1(a, b, c)]_Z^{k_{21}} [g_2(a, b, c)]_Z^{k_{22}}. \end{aligned}$$

Therefore

$$\begin{aligned} g_1(a', b', c') &= g_1(a, b, c)^{k_{11}} g_2(a, b, c)^{k_{12}} g_3(a, b, r)^{s_1}, \\ g_2(a', b', c') &= g_1(a, b, c)^{k_{21}} g_2(a, b, c)^{k_{22}} g_3(a, b, r)^{s_2} \end{aligned} \quad (49)$$

with $s_1, s_2 \in \mathbb{Z}$. Put

$$G_1 := g_1(a, b, c)^{k_{11}} g_2(a, b, c)^{k_{12}}, \quad G_2 := g_1(a, b, c)^{k_{21}} g_2(a, b, c)^{k_{22}}. \quad (50)$$

By (49) it follows that

$$[G_1, G_2] = [g_1(a', b', c'), g_2(a', b', c')] = g_3(a', b', r)^r.$$

On the other hand, taking into account that $[g_1(a, b, c), g_2(a, b, c)] = g_3(a, b, r)^r$ and the definitions of G_1, G_2 , direct computations give $[G_1, G_2] = g_3(a, b, r)^{\det(K)r}$. Therefore

$$G_3 := g_3(a', b', r) = g_3(a, b, r)^{\det(K)}. \quad (51)$$

For comparing $g_0(\alpha, t)$ with $g_0(\alpha, t')$ recall that, by Remark 2.19, their classes with respect to the subgroup $G \cap \text{Aff}_1^1(U) = G(a, b, c, r) = G(a', b', c', r)$ coincide, so

$$g_0(\alpha, t') = g_0(\alpha, t) g_1(a, b, c)^{k_1} g_2(a, b, c)^{k_2} g_3(a, b, r)^{s_0}$$

with $k_1, k_2, s_0 \in \mathbb{Z}$. Putting $k = (k_1, k_2)$, $ka = k_1 a_1 + k_2 a_2$, $kb = k_1 b_1 + k_2 b_2$, direct computations give

$$\begin{aligned} g_0(\alpha, t) g_1(a, b, c)^{k_1} g_2(a, b, c)^{k_2} g_3(a, b, r)^{s_0} \begin{pmatrix} w \\ z \end{pmatrix} &= \\ \left(\begin{array}{c} \alpha(w + ka) \\ z + (kb)w + \frac{k_1(k_1-1)}{2} a_1 b_1 + k_1 k_2 b_1 a_2 + \frac{k_2(k_2-1)}{2} a_2 b_2 + kc + t + \frac{s_0}{r} b \wedge a \end{array} \right). \end{aligned}$$

Taking into account the explicit formula for $g_0(\alpha, t')$, we obtain $ka = kb = 0$, so $k = 0$. Therefore we have

$$g_0(\alpha, t') = g_0(\alpha, t)g_3(a, b, r)^{s_0} \quad (52)$$

By Lemma 4.14 we see that formulae (50), (51) give

$$G_i = g_i(A, B, C) \text{ for } 1 \leq i \leq 2, \quad G_3 = g_3(A, B, r)$$

where $(A, B, C) = K \cdot (a, b, c)$. On the other hand (49) can be written as

$$\begin{aligned} g_1(a', b', c') &= g_1(A, B, C)g_3(A, B, r)^{\det(K)s_1}, \\ g_2(a', b', c') &= g_2(A, B, C)g_3(A, B, r)^{\det(K)s_2}, \end{aligned} \quad (53)$$

which, by Remark 4.12, gives $c' = C + \frac{B \wedge A}{r} \det(K)s \in [C]_{AB}$. Finally, formula (52) shows that $t' = t + \frac{B \wedge A}{r} \det(K)s_0 \in [t]_{AB}$.

Therefore, $a' = A = Ka$, $b' = B = Kb$ and $[c']_{AB} = [C]_{AB}$, $[t']_{AB} = [t]_{AB}$ which, taking into account that $K \cdot (a, b, c, t) = (A, B, C, t)$, proves the claim.

(2) \Rightarrow (1):

It suffices to note that, if $(a', b', [c']_{a'b'}, [t']_{a'b'}) = K \cdot (a, b, [c]_{ab}, [t]_{ab})$ with $K \in \text{GL}(2, \mathbb{Z})$, then $G(\alpha, a', b', c', r, t') \subset G(\alpha, a, b, c, r, t)$. \blacksquare

Therefore, by Theorem 4.17, the quotient set $\mathcal{A}_{\alpha, r} / \approx$ (which parameterises the subgroups $G \subset \text{Aff}(U)$ defining type II Inoue surfaces) is identified with the orbit space $\mathcal{A}_{\alpha, r} / \text{GL}(2, \mathbb{Z})$. This orbit space comes with a natural surjective map

$$\mathcal{A}_{\alpha, r} / \text{GL}(2, \mathbb{Z}) \rightarrow \mathcal{P}_{\alpha} / \text{GL}(2, \mathbb{Z}) \rightarrow \mathcal{N}_{\alpha} / \text{GL}(2, \mathbb{Z})$$

onto the set of $\text{GL}(2, \mathbb{Z})$ -similarity classes of matrices $N \in \mathcal{N}_{\alpha}$ (see section 2.2).

4.2.2. A classification theorem for type II Inoue surfaces. Let $(\alpha, r) \in]1, +\infty[\times \mathbb{N}^*$ with α S^+ -admissible. We now come back to our initial problem, which is: describe explicitly the quotient of $\mathcal{A}_{\alpha, r} := \mathcal{T}_{\alpha, r} \times \mathbb{C}$ by the equivalence relation

$$(a, b, c, t) \sim (a', b', c', t') \text{ if } G(\alpha, a, b, c, r, t), G(\alpha, a', b', c', r, t') \text{ are conj. in } \text{Aff}(U).$$

Elementary computations give the following:

Lemma 4.18. *Let $(a, b, c, t), (a', b', c', t') \in \mathcal{A}_{\alpha, r}$ and put*

$$\begin{cases} g_0 = g_0(\alpha, t), & g_i = g_i(a, b, c), & g_3 = g_3(a, b, r), \\ g'_0 = g_0(\alpha, t'), & g'_i = g_i(a', b', c'), & g'_3 = g_3(a', b', r). \end{cases}$$

Let $k = (k_1, k_2) \in \mathbb{Z}^2$, $\begin{pmatrix} s_0 \\ s_1 \\ s_2 \end{pmatrix} \in \mathbb{Z}^3$ and $\tau \in \text{Aff}_1^1(U)$, $\begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{\tau} \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix}$.

The following conditions are equivalent:

$$(i) \quad \tau g'_0 \tau^{-1} = g_0 g_1^{k_1} g_2^{k_2} g_3^{s_0}, \quad \tau \circ g'_i \circ \tau^{-1} = g_i g_3^{s_i} \text{ for } 1 \leq i \leq 2, \quad \tau \circ g'_3 \circ \tau^{-1} = g_3.$$

(ii) We have:

$$\begin{cases} \lambda &= \frac{kb}{\alpha-1}. \\ u &= \frac{\alpha(ka)}{1-\alpha}. \\ a' &= a. \\ b' &= b. \\ c' &= c - \frac{\alpha(ka)}{\alpha-1}b - \frac{kb}{\alpha-1}a + \frac{b \wedge a}{r}s, \text{ where } s := \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}. \\ t' &= t + \frac{\alpha(ka)(kb)}{1-\alpha} + kc + \frac{k_1(k_1-1)a_1b_1 + k_2(k_2-1)a_2b_2}{2} + k_1k_2b_1a_2 + \frac{s_0}{r}b \wedge a. \end{cases}$$

Lemma 4.19. (1) *Putting*

$$k \cdot c := c - \frac{\alpha(ka)}{\alpha-1}b - \frac{kb}{\alpha-1}a,$$

$$k \cdot_c t := t + \frac{\alpha(ka)(kb)}{1-\alpha} + kc + \frac{k_1(k_1-1)a_1b_1 + k_2(k_2-1)a_2b_2}{2} + k_1k_2b_1a_2,$$

we have the identities

$$k \cdot c = c + b \wedge a (I_2 - N)^{-1} \begin{pmatrix} -k_2 \\ k_1 \end{pmatrix}, \quad (54)$$

$$k \cdot_c t = t + b \wedge a \left(k (I_2 - N)^{-1} \left(\frac{1}{2} \begin{pmatrix} n_{11}n_{12} - k_2 \\ n_{21}n_{22} + k_1 \end{pmatrix} + \frac{1}{r} \pi_{a,b,r}(c) \right) + \frac{k_1k_2}{2} \right). \quad (55)$$

(2) Put $\delta := \det(I_2 - N) = 2 - \theta$. If $k \in \delta\mathbb{Z}^2$, then

$$k (I_2 - N)^{-1} \begin{pmatrix} n_{11}n_{12} - k_2 \\ n_{21}n_{22} + k_1 \end{pmatrix} + k_1k_2 \in 2\mathbb{Z}, \quad (56)$$

$$k (I_2 - N)^{-1} \left(\frac{1}{2} \begin{pmatrix} n_{11}n_{12} - k_2 \\ n_{21}n_{22} + k_1 \end{pmatrix} + \frac{1}{r} \mathbb{Z}^2 \right) + \frac{k_1k_2}{2} \subset \frac{1}{r} \mathbb{Z}. \quad (57)$$

Proof. (1) For the first formula it suffices to apply $I_2 - N$ to $k \cdot c$ taking into account that $(I_2 - N)a = (1 - \alpha)a$, $(I_2 - N)b = (1 - \alpha^{-1})b$.

For the second formula in (1), we have

$$\begin{aligned} & \frac{\alpha(ka)(kb)}{1-\alpha} + kc + \frac{k_1(k_1-1)a_1b_1 + k_2(k_2-1)a_2b_2}{2} + k_1k_2b_1a_2 = \\ &= \frac{k_1(k_1-1)}{2}a_1b_1 + k_1k_2b_1a_2 + \frac{k_2(k_2-1)}{2}a_2b_2 + kc - \frac{\alpha}{\alpha-1}(ka)(kb) = \\ &= \frac{1}{2}(k_1^2a_1b_1 + k_2^2a_2b_2 + k_1k_2(a_1b_2 + a_2b_1) - k_1k_2(a_1b_2 + a_2b_1) - k_1a_1b_1 - k_2a_2b_2) + \\ & \quad + \frac{1}{2}2k_1k_2b_1a_2 + kc - \frac{\alpha}{\alpha-1}(ka)(kb) = \\ &= \frac{1}{2}((ka)(kb) + k_1k_2b \wedge a) + kc - \frac{\alpha}{\alpha-1}(ka)(kb) - k_1a_1b_1 - k_2a_2b_2 = \\ &= kc + \frac{1}{2}(k_1k_2b \wedge a - k_1a_1b_1 - k_2a_2b_2) - \frac{\alpha+1}{2(\alpha-1)}(ka)(kb) = \\ & \quad k \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \frac{1}{2}k_1k_2b \wedge a - \frac{\alpha+1}{2(\alpha-1)}(ka)(kb) \end{aligned}$$

On the other hand, applying Lemma 4.20 to $(I_2 - N)^{-1}$ and noting that

$$(I_2 - N)^{-1}(a) = \frac{1}{1-\alpha}a, \quad (I_2 - N)^{-1}(b) = \frac{\alpha}{\alpha-1}b,$$

we obtain:

$$-\frac{\alpha+1}{\alpha-1}(ka)(kb) = (b \wedge a) \left(k (I_2 - N)^{-1} \begin{pmatrix} -k_2 \\ k_1 \end{pmatrix} \right).$$

Therefore

$$\begin{aligned} & \frac{\alpha(ka)(kb)}{1-\alpha} + kc + \frac{k_1(k_1-1)a_1b_1 + k_2(k_2-1)a_2b_2}{2} + k_1k_2b_1a_2 = \\ &= k \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \frac{k_1k_2}{2}b \wedge a + \frac{b \wedge a}{2} \left(k (I_2 - N)^{-1} \begin{pmatrix} -k_2 \\ k_1 \end{pmatrix} \right). \end{aligned}$$

It suffices to recall that:

$$c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} = (I_2 - N)^{-1} \left(\frac{b \wedge a}{2} \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \frac{b \wedge a}{r} \pi_{a,b,r}(c) \right)$$

by the definition of $p(a, b, c, r) = \pi_{a,b,r}(c)$.

(2) Note first that (57) follows easily from (56) by noting that $\delta(I_2 - N)^{-1}$ is a matrix with integer entries. Therefore it's enough to prove (56). Let $l \in \mathbb{Z}^2$ such that $k = \delta l$. The claim follows by elementary computations using the formulae:

$$\begin{aligned} k(I_2 - N)^{-1} &= \delta l(I_2 - N)^{-1} = l {}^t \text{cof}(I_2 - N) = l \begin{pmatrix} 1 - n_{22} & n_{12} \\ n_{21} & 1 - n_{11} \end{pmatrix}, \\ l_1 l_2 (n_{22} - n_{11} + \delta) &= l_1 l_2 (n_{22} - n_{11} + 2 - (n_{11} + n_{22})) = 2l_1 l_2 (1 - n_{11}) \in 2\mathbb{Z}, \\ l_i^2 &\equiv l_i \pmod{2}, \quad \delta \equiv \theta \equiv n_{11} + n_{22} \pmod{2} \\ n_{12} n_{22} (1 + n_{21} - n_{11}), & \quad n_{21} n_{11} (-1 + n_{12} - n_{22}) \in 2\mathbb{Z}. \end{aligned}$$

The latter formula follows using $\det(N) = 1$. ■

Lemma 4.20. *Let $M \in M_2(\mathbb{C})$ and $a = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}$, $b = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$ be linearly independent in \mathbb{C}^2 such that $Ma = \alpha a$, $Mb = \beta b$ and $x = (x_1, x_2) \in \mathbb{C}^2$. Then*

$$(\beta - \alpha)(xa)(xb) = (a \wedge b)xM \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}.$$

Proof. We apply the Cramer rule to the system

$$\begin{cases} a_1 x_1 + a_2 x_2 &= xa \\ b_1 x_1 + b_2 x_2 &= xb \end{cases}.$$

in order to express x_1, x_2 in terms of xa and xb . ■

Proposition 4.21. *Let $(a, b) \in \mathcal{P}_\alpha$.*

(1) *The formula $(k, c) \mapsto k \cdot c$ defines an action of \mathbb{Z}^2 on $\mathcal{C}_{a,b,r}$ satisfying the identity*

$$\pi_{a,b,r}(k \cdot c) = \pi_{a,b,r}(c) + r \begin{pmatrix} -k_2 \\ k_1 \end{pmatrix}. \quad (58)$$

(2) *The formula $\left(k, \begin{pmatrix} [c]_{ab} \\ [t]_{ab} \end{pmatrix}\right) \mapsto \begin{pmatrix} [k \cdot c]_{ab} \\ [k \cdot c \ t]_{ab} \end{pmatrix}$ defines an action of \mathbb{Z}^2 on the product $\mathcal{C}_{a,b,r} \times (\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z})$. This action descends to an action of the finite group $\mathbb{Z}^2/d\mathbb{Z}^2 = (\mathbb{Z}_d)^2$, where $d := |\det(I_2 - N)| = \theta - 2$.*

Proof. (1) This follows from formula (54).

(2) Note first that the pair $([k \cdot c]_{ab}, [k \cdot c \ t]_{ab})$ depends only on the classes $[c]_{ab}$, $[t]_{ab}$ and k , so the right hand term is well defined. To prove that the given formula defines an action, put:

$$\mathfrak{c} := k \cdot c, \quad \mathfrak{t} := k \cdot c \ t, \quad \mathfrak{C} := l \cdot \mathfrak{c}, \quad \mathfrak{T} := l \cdot \mathfrak{t}, \quad \mathfrak{C}' := (k + l) \cdot c, \quad \mathfrak{T}' := (k + l) \cdot c \ t.$$

By (1) we have $\mathfrak{C}' = \mathfrak{C}$, and, using the definition of $k \cdot c \ t$, elementary computations give

$$\mathfrak{T}' = \mathfrak{T} + k_2 l_1 b \wedge a \in \mathfrak{T} + \frac{b \wedge a}{r}\mathbb{Z}, \quad (59)$$

which proves the claim. For the second claim in (2), note that, by formulae (54), (55), (57) stated in Lemma 4.19, we have

$$k \cdot c - c \in (b \wedge a)\mathbb{Z}^2 \subset \frac{b \wedge a}{r}\mathbb{Z}^2, \quad k \cdot c \ t - t \in \frac{b \wedge a}{r}\mathbb{Z}$$

for any $k \in d\mathbb{Z}^2$, so $d\mathbb{Z}^2$ acts trivially on $\mathcal{C}_{a,b,r} \times (\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z})$. ■

For a triple $(a, b, c) \in \mathcal{T}_{\alpha, r}$ and $t \in \mathbb{C}$ we will also use the notation

$$k \cdot (a, b, c, t) = (a, b, \mathbf{c}, \mathbf{t}) \text{ where } (\mathbf{c}, \mathbf{t}) := k \cdot (c, t).$$

Formula (59) shows that the formulae

$$(k, (c, t)) \mapsto k \cdot (c, t), \quad (k, (a, b, c, t)) = k \cdot (a, b, c, t)$$

do not define \mathbb{Z}^2 -actions on the respective spaces. However, Proposition 4.21 shows that the formulae

$$(k, ([c]_{ab}, [t]_{ab})) \mapsto k \cdot ([c]_{ab}, [t]_{ab}) := ([\mathbf{c}]_{ab}, [\mathbf{t}]_{ab}),$$

$$(k, (a, b, [c]_{ab}, [t]_{ab})) \mapsto k \cdot (a, b, [c]_{ab}, [t]_{ab}) := (a, b, [\mathbf{c}]_{ab}, [\mathbf{t}]_{ab})$$

define \mathbb{Z}^2 actions which descend to \mathbb{Z}_d^2 -actions on the spaces $\mathcal{C}_{a, b, r} \times (\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z})$, $\mathcal{A}_{\alpha, r}$ respectively.

On the latter space $\mathcal{A}_{\alpha, r}$ we have now defined a $\mathrm{GL}(2, \mathbb{Z})$ -action and a \mathbb{Z}^2 -action (which induces a \mathbb{Z}_d^2 -action). These actions *do not commute*. We have

Remark 4.22. Let $(a, b, c) \in \mathcal{T}_{\alpha, r}$, $t \in \mathbb{C}$, $k = (k_1, k_2) \in \mathbb{Z}^2$, $K = (k_{ij}) \in \mathrm{GL}(2, \mathbb{Z})$. Then the 4-tuples $K \cdot (k \cdot (a, b, c, t))$, $(kK^{-1}) \cdot (K \cdot (a, b, c, t))$ define the same elements in $\mathcal{A}_{\alpha, r}$. More precisely

(1) We have the identity

$$K \cdot (k \cdot c) = (kK^{-1}) \cdot (K \cdot c).$$

(2) Putting

$$(\mathbf{c}, \mathbf{t}) := k \cdot (c, t), \quad (A, B, C) = K \cdot (a, b, c), \quad (A, B, \mathfrak{C}) = K \cdot (a, b, c)$$

we have

$$([\mathfrak{C}]_{AB}, [\mathbf{t}]_{AB}) = (kK^{-1}) \cdot ([C]_{AB}, [t]_{AB}).$$

Proof. (1) Follows by elementary computations.

(2) Put $k' := kK^{-1}$. Note that, by (1), the first component of $k' \cdot (C, t)$ is \mathfrak{C} . Define $\mathfrak{t}' \in \mathbb{C}$ by $k' \cdot (C, t) = (\mathfrak{C}, \mathfrak{t}')$. We have to prove that $\mathfrak{t}, \mathfrak{t}'$ are congruent mod $\frac{B \wedge A}{r}$. Elementary computations give:

$$\begin{aligned} \mathfrak{t}' - \mathfrak{t} &= \frac{b \wedge a}{2} \left(k' \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} + k'_1 k'_2 \det K - k_1 k_2 \right) \\ &= \frac{b \wedge a}{2} ((k'_1 - k_1'^2)k_{11}k_{12} + (k'_2 - k_2'^2)k_{21}k_{22} - 2k'_2 k'_1 k_{12}k_{21}) \\ &\in \frac{B \wedge A}{r} r\mathbb{Z} \subset \frac{B \wedge A}{r} \mathbb{Z}. \end{aligned} \quad (60)$$

■

Proposition 4.21 and Remark 4.22 show that:

Proposition 4.23. Let $\varphi : \mathrm{GL}(2, \mathbb{Z}) \rightarrow \mathrm{Aut}(\mathbb{Z}^2)$, $\psi : \mathrm{GL}(2, \mathbb{Z}) \rightarrow \mathrm{Aut}(\mathbb{Z}_d^2)$ be the group morphisms defined by $\varphi(K)(k) = kK^{-1}$, $\psi(K)([k]) = [k]K^{-1}$. The formula

$$(K, k) \cdot (a, b, [c]_{ab}, [t]_{ab}) := K \cdot (k \cdot (a, b, [c]_{ab}, [t]_{ab})),$$

defines an action of the semi-direct product² $\mathrm{GL}(2, \mathbb{Z}) \rtimes_{\varphi} \mathbb{Z}^2$ on the space $\mathcal{A}_{\alpha, r}$ which descends to an action of $\mathrm{GL}(2, \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}_d^2$.

²In general, for groups N, H and a group morphism $\varphi : H \rightarrow \mathrm{Aut}(N)$, the semi-direct product $H \rtimes_{\varphi} N$ is $H \times N$ endowed with the composition law $(h, n) * (h', n') := (hh', \varphi(h'^{-1})(n)n')$.

Now note that the action

$$(\mu, \rho) \cdot (a, b, c, t) := (\mu a, \rho b, \mu \rho c, \mu \rho t)$$

of the product $\mathbb{R}_+^* \times \mathbb{R}^*$ on $\mathcal{A}_{\alpha, r}$ induces an $\mathbb{R}_+^* \times \mathbb{R}^*$ -action on $\mathcal{A}_{\alpha, r}$ which commutes with the $(\mathrm{GL}(2, \mathbb{Z}) \times_{\varphi} \mathbb{Z}^2)$ -action defined above. Therefore we obtain an action of the product group

$$(\mathbb{R}_+^* \times \mathbb{R}^*) \times (\mathrm{GL}(2, \mathbb{Z}) \times_{\varphi} \mathbb{Z}^2)$$

on $\mathcal{A}_{\alpha, r}$ given explicitly by

$$((\mu, \rho), (K, k)) \cdot (a, b, [c]_{ab}, [t]_{ab}) := (K, k) \cdot (\mu a, \rho b, [\mu \rho c]_{ab}, [\mu \rho t]_{ab}),$$

which descends to an action of $(\mathbb{R}_+^* \times \mathbb{R}^*) \times (\mathrm{GL}(2, \mathbb{Z}) \times_{\psi} \mathbb{Z}_d^2)$

Our main theorem is

Theorem 4.24. *Let $(a, b, c, t), (a', b', c', t') \in \mathcal{A}_{\alpha, r}$. The groups $G(\alpha, a, b, c, r, t), G(\alpha, a', b', c', r, t')$ are conjugate in $\mathrm{Aff}(U)$ if and only if*

$$\begin{aligned} (a', b', [c']_{a'b'}, [t']_{a'b'}) \in & ((\mathbb{R}_+^* \times \mathbb{R}^*) \times (\mathrm{GL}(2, \mathbb{Z}) \times_{\psi} \mathbb{Z}_d^2)) \cdot (a, b, [c]_{ab}, [t]_{ab}). \\ & = ((\mathbb{R}_+^* \times \mathbb{R}^*) \times (\mathrm{GL}(2, \mathbb{Z}) \times_{\varphi} \mathbb{Z}^2)) \cdot (a, b, [c]_{ab}, [t]_{ab}). \end{aligned} \quad (61)$$

Proof. Put $G := G(\alpha, a, b, c, r, t), G' := G(\alpha, a', b', c', r, t')$,

$$\begin{cases} g_0 = g_0(\alpha, t), & g_i = g_i(a, b, c), & g_3 = g_3(a, b, r), \\ g'_0 = g_0(\alpha, t'), & g'_i = g_i(a', b', c'), & g'_3 = g_3(a', b', r). \end{cases}$$

First, suppose G, G' are conjugate in $\mathrm{Aff}(U)$, and let $\theta \in \mathrm{Aff}(U)$,

$$\theta \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} \mu & 0 \\ \lambda & \nu \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix}$$

(with $\mu \in \mathbb{R}_+^*, \nu \in \mathbb{C}^*, u \in \mathbb{R}, \lambda, \zeta \in \mathbb{C}$) be such that $\theta G' \theta^{-1} = G$.

It follows that, putting, $\gamma_i := \theta g'_i \theta^{-1}$ for $0 \leq i \leq 3$, $(\gamma_0, \gamma_1, \gamma_2, \gamma_3)$ is a system of generators of G . Note that this system has the properties:

- (C1) γ_3 is a generator of the cyclic group $G \cap \mathrm{T}(U)$, in other words $\gamma_3 \in \{g_3, g_3^{-1}\}$.
- (C2) $(\gamma_1, \gamma_2, \gamma_3)$ is a system of generators of $G(a, b, c, r) = G \cap \mathrm{Aff}_1^r(U)$ with the property $[\gamma_1, \gamma_2] = \gamma_3^r$.
- (C3) The first diagonal element of the linear part of γ_0 is α .

By Lemma 4.25 proved below, it follows that there exists $K \in \mathrm{GL}(2, \mathbb{Z}), k =$

$$(k_1, k_2) \in \mathbb{Z}^2 \text{ and } \begin{pmatrix} s_0 \\ s_1 \\ s_2 \end{pmatrix} \in \mathbb{Z}^3 \text{ such that}$$

$$\gamma_0 = g_0 g_1^{k_1} g_2^{k_2} g_3^{s_0}, \gamma_1 = g_1^{k_{11}} g_2^{k_{12}} g_3^{s_1}, \gamma_2 = g_1^{k_{21}} g_2^{k_{22}} g_3^{s_2}, \gamma_3 = g_3^{\det(K)}. \quad (62)$$

Identifying the (2,1)-entries of the matrices associated with the linear parts of the two sides in the equalities $\theta g'_i \theta^{-1} = g_1^{k_{i1}} g_2^{k_{i2}} g_3^{s_i}$ for $1 \leq i \leq 2$, we obtain $\nu \mu^{-1} b'_i \in \mathbb{R}$, so $\nu \mu^{-1} \in \mathbb{R}$, i.e. $\nu \in \mathbb{C}^* \cap \mathbb{R} = \mathbb{R}^*$.

Put

$$G_0 := g_0, G_1 := g_1^{k_{11}} g_2^{k_{12}}, G_2 := g_1^{k_{21}} g_2^{k_{22}}, G_3 := g_3^{\det(K)}.$$

By Lemma 4.25 (G_0, G_1, G_2, G_3) is a system of generators of G and, by Lemma 4.14, we have

$$G_0 = g_0(\alpha, t), G_i = g_i(A, B, C), G_3 = g_3(A, B, r),$$

where $(A, B, C) = K \cdot (a, b, c)$.

On the other hand we can write $\theta = \tau \circ \delta$ where

$$\tau \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \lambda \mu^{-1} & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix}, \delta \begin{pmatrix} w \\ z \end{pmatrix} = \begin{pmatrix} \mu w \\ \nu z \end{pmatrix}.$$

Note that $\tau \in \text{Aff}_1^1(U)$. We have

$$\gamma_i = \tau g_i'' \tau^{-1} \text{ for } 0 \leq i \leq 3 \quad (63)$$

where $g_i'' := \delta g_i' \delta^{-1}$. Direct computations show that

$$g_0'' = g_0(\alpha, t''), \quad g_i'' = g_i(a'', b'', c'') \text{ for } 1 \leq i \leq 2, \quad g_3'' = g_3(a'', b'', r),$$

where $a'' = \mu a'$, $b'' = \nu \mu^{-1} b'$, $c'' = \nu c'$, $t'' = \nu t'$. Since $\nu \mu^{-1} \in \mathbb{R}^*$, we can write

$$(a'', b'', c'', t'') = (\mu, \nu \mu^{-1}) \cdot (a', b', c', t'). \quad (64)$$

Taking into account (62) we obtain

$$\begin{aligned} \tau g_0'' \tau^{-1} &= G_0 g_1^{k_1} g_2^{k_2} g_3^{s_0}, \quad \tau g_1'' \tau^{-1} = G_1 G_3^{s_1 \det(K)}, \\ \tau g_2'' \tau^{-1} &= G_2 G_3^{s_2 \det(K)}, \quad \tau g_3'' \tau^{-1} = G_3. \end{aligned} \quad (65)$$

Since g_1 and g_2 commute modulo $\langle g_3^r \rangle$ (see Remark 2.18) it follows that, putting $L := K^{-1}$, $l = (l_1, l_2) := kL$, we have

$$g_1 = G_1^{l_{11}} G_2^{l_{12}} g_3^{r m_1}, \quad g_2 = G_1^{l_{21}} G_2^{l_{22}} g_3^{r m_2}, \quad g_1^{k_1} g_2^{k_2} = G_1^{l_1} G_2^{l_2} g_3^{r m}$$

and the formula $\tau g_0'' \tau^{-1} = G_0 g_1^{k_1} g_2^{k_2} g_3^{s_0}$ can be written as

$$\tau g_0'' \tau^{-1} = G_0 G_1^{l_1} G_2^{l_2} g_3^{s_0 + r m}.$$

Therefore (65) becomes

$$\begin{aligned} \tau g_0'' \tau^{-1} &= G_0 G_1^{l_1} G_2^{l_2} G_3^{(s_0 + r m) \det(K)}, \quad \tau g_1'' \tau^{-1} = G_1 G_3^{s_1 \det(K)}, \\ \tau g_2'' \tau^{-1} &= G_2 G_3^{s_2 \det(K)}, \quad \tau g_3'' \tau^{-1} = G_3. \end{aligned} \quad (66)$$

By Lemma 4.18, it follows that

$$a'' = A, \quad b'' = B, \quad ([c'']_{a''b''}, [t'']_{a''b''}) = l \cdot ([C]_{AB}, t_{AB})$$

so, by (64),

$$(\mu, \nu \mu^{-1}) \cdot (a', b', [c']_{a'b'}, [t']_{a'b'}) = l \cdot (A, B, [C]_{AB}, [t]_{AB}),$$

i.e.

$$(\mu, \nu \mu^{-1}) \cdot (a', b', [c']_{a'b'}, [t']_{a'b'}) = l \cdot (K \cdot (a, b, [c]_{ab}, [t]_{ab})),$$

which completes the proof of the first implication.

For the converse, note first that, if

$$(a', b', [c']_{a'b'}, [t']_{a'b'}) \in ((\mathbb{R}_+^* \times \mathbb{R}^*) \times (\text{GL}(2, \mathbb{Z}) \times_{\varphi} \mathbb{Z}^2)) \cdot (a, b, [c]_{ab}, [t]_{ab}),$$

then there exists $(\mu, \rho) \in \mathbb{R}_+^* \times \mathbb{R}^*$, and $(K, k) \in (\text{GL}(2, \mathbb{Z}) \times_{\varphi} \mathbb{Z}^2)$ such that

$$(\mu, \rho) \cdot (a', b', [c']_{a'b'}, [t']_{a'b'}) = (K, k) \cdot (a, b, [c]_{ab}, [t]_{ab}). \quad (67)$$

We have to show that there exists $\theta \in \text{Aff}(U)$ such that

$$\theta G(\alpha, a', b', c', r, t') \theta^{-1} \subset G(\alpha, a, b, c, r, t).$$

Put $g_0'' := g_0(\alpha, t'')$, $g_i'' := g_i(a'', b'', c'')$ for $1 \leq i \leq 2$, $g_3'' := g_3(a'', b'', r)$, where

$$(a'', b'', c'', t'') := (\mu, \rho) \cdot (a', b', c', t').$$

An easy computation shows that

$$g_i'' = \delta g_i' \delta^{-1} \text{ for } 0 \leq i \leq 3, \quad (68)$$

where $\delta \in \text{Aff}(U)$ is given by $\delta(w, z) = (\mu w, \mu \rho z)$.

On the other hand, by (67) we have

$$\begin{aligned} (a'', b'', [c'']_{a''b''}, [t'']_{a''b''}) &= k K^{-1} \cdot (K \cdot (a, b, [c]_{ab}, [t]_{ab})) = \\ &= (K, k) \cdot (a, b, [c]_{ab}, [t]_{ab}) = l \cdot (A, B, [C]_{AB}, [t]_{AB}), \end{aligned} \quad (69)$$

where $l := kK^{-1} = (l_1, l_2)$. Formula (69) shows that there exists $s_0 \in \mathbb{Z}$, $s = \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} \in \mathbb{Z}^2$ such that

$$\begin{cases} a'' = A. \\ b'' = B. \\ c'' = C - \frac{\alpha(lA)}{\alpha-1}B - \frac{lB}{\alpha-1}A + \frac{A \wedge B}{r}s. \\ t'' = t + \frac{\alpha(lA)(lB)}{1-\alpha} + lC + \frac{l_1(l_1-1)A_1B_1 + l_2(l_2-1)A_2B_2}{2} + l_1l_2B_1A_2 + \frac{s_0}{r}B \wedge A. \end{cases}$$

Let $\tau \in \text{Aff}_1^1(U)$ be given by

$$\begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{\tau} \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ 0 \end{pmatrix}, \quad \lambda := \frac{lB}{\alpha-1}, \quad u := \frac{\alpha(lA)}{1-\alpha}.$$

Using the implication $ii) \Rightarrow i)$ of Lemma 4.18, it follows that:

$$\begin{aligned} \tau g_0'' \tau^{-1} &= g_0(\alpha, t) g_1(A, B, C)^{l_1} g_2(A, B, C)^{l_2} g_3(A, B, r)^{s_0}, \\ \tau g_i'' \tau^{-1} &= g_i(A, B, C) g_3(A, B, r)^{s_i} \text{ for } 1 \leq i \leq 2, \\ \tau g_3'' \tau^{-1} &= g_3(A, B, r). \end{aligned} \quad (70)$$

Combining (68) and (70) and putting $\theta = \tau \circ \delta$, we obtain

$$\begin{aligned} \theta g_0' \theta^{-1} &= g_0(\alpha, t) g_1(A, B, C)^{l_1} g_2(A, B, C)^{l_2} g_3(A, B, r)^{s_0}, \\ \theta g_i' \theta^{-1} &= g_i(A, B, C) g_3(A, B, r)^{s_i}, \quad 1 \leq i \leq 2, \\ \theta g_3' \theta^{-1} &= g_3(A, B, r). \end{aligned}$$

This shows that $\theta G(\alpha, a', b', c', r, t) \theta^{-1} = G(\alpha, A, B, C, r, t)$. On the other hand, since $(A, B, [C]_{AB}, [t]_{AB}) = K \cdot (a, b, [c]_{ab}, [t]_{ab})$, it follows by Theorem 4.17 that $G(\alpha, A, B, C, r, t) = G(\alpha, a, b, c, r, t)$.

Therefore $\theta G(\alpha, a', b', c', r, t) \theta^{-1} = G(\alpha, a, b, c, r, t)$, which completes the proof. \blacksquare

Lemma 4.25. *The set of system of generators $(\gamma_0, \gamma_1, \gamma_2, \gamma_3)$ of G satisfying the properties (C1), (C2), (C3) stated in the proof of Theorem 4.24 is*

$$\left\{ (g_0 g_1^{k_1} g_2^{k_2} g_3^{s_0}, g_1^{k_{11}} g_2^{k_{12}} g_3^{s_1}, g_1^{k_{21}} g_2^{k_{22}} g_3^{s_2}, g_3^{\det(K)}) \mid K \in \text{GL}(2, \mathbb{Z}), k \in \mathbb{Z}^2, \begin{pmatrix} s_0 \\ s_1 \\ s_2 \end{pmatrix} \in \mathbb{Z}^3 \right\}.$$

Proof. This follows using Remark 2.21 by elementary computations. \blacksquare

Taking into account Remark 4.11, Theorem 4.24 gives the following geometric interpretation of the set of biholomorphism classes of type II Inoue surfaces associated with a pair x :

Theorem 4.26. *Let $(\alpha, r) \in]1, +\infty[\times \mathbb{N}^*$ with α S^+ -admissible. The set of biholomorphism classes of type II Inoue surfaces associated with (α, r) is naturally identified with the quotient space*

$$\mathcal{Q}_{\alpha, r} := \mathcal{A}_{\alpha, r} / (\mathbb{R}_+^* \times \mathbb{R}^*) \times (\text{GL}(2, \mathbb{Z}) \rtimes_{\psi} \mathbb{Z}_d^2).$$

4.2.3. *The fibre over a similarity class of $\text{SL}(2, \mathbb{Z})$ -matrices.* In section 2.2 we have put $\mathcal{N}_{\alpha} := \{N \in \text{SL}(2, \mathbb{Z}) \mid \alpha \in \text{Spec}(N)\}$. Consider the *surjective* maps

$$\begin{array}{ccc} \mathcal{A}_{\alpha, r} & \xrightarrow{\psi_{\alpha, r}} & \mathcal{N}_{\alpha} \\ & \searrow \phi_{\alpha, r} & \nearrow \eta_{\alpha} \\ & \mathcal{P}_{\alpha} & \end{array} \quad \left| \begin{array}{l} \phi_{\alpha, r}(a, b, [c]_{ab}, [t]_{ab}) \quad := \quad (a, b), \\ \eta_{\alpha}(a, b) \quad := \quad N(\alpha, a, b), \\ \psi_{\alpha, r} \quad := \quad \eta_{\alpha} \circ \phi_{\alpha, r}, \end{array} \right.$$

and let the group

$$\mathcal{G} := (\mathbb{R}_+^* \times \mathbb{R}^*) \times (\mathrm{GL}(2, \mathbb{Z}) \times_{\psi} \mathbb{Z}_d^2)$$

act on the spaces $\mathcal{P}_\alpha, \mathcal{N}_\alpha$ by

$$((\mu, \rho), (K, [k])) \cdot (a, b) := (\mu K a, \rho K b), \quad ((\mu, \rho), (K, [k])) \cdot N := K N K^{-1}.$$

The maps $\phi_{\alpha, r}, \eta_\alpha, \psi_{\alpha, r}$ are obviously \mathcal{G} -equivariant, so they induce maps

$$\begin{array}{ccc} \mathcal{Q}_{\alpha, r} := \mathcal{A}_{\alpha, r} / \mathcal{G} & \xrightarrow{\Psi_{\alpha, r}} & \mathcal{N}_\alpha / \mathcal{G} = \mathcal{N}_\alpha / \mathrm{GL}(2, \mathbb{Z}) = \mathcal{N}_\alpha / \sim \\ & \searrow \Phi_{\alpha, r} & \nearrow H_\alpha \\ & \mathcal{P}_\alpha / \mathcal{G} & \xlongequal{\quad} \mathcal{P}_\alpha / (\mathbb{R}_+^* \times \mathbb{R}^*) \times \mathrm{GL}(2, \mathbb{Z}) \end{array}$$

between the respective \mathcal{G} -quotients. We are interested in the fibres of $\Psi_{\alpha, r}$.

Using the same method as is the proof of the implication (2) \Rightarrow (1) of Proposition 4.5, we obtain easily:

Remark 4.27. *The map $H_\alpha : \mathcal{P}_\alpha / (\mathbb{R}_+^* \times \mathbb{R}^*) \times \mathrm{GL}(2, \mathbb{Z}) \rightarrow \mathcal{N}_\alpha / \sim$ induced by η_α is bijective, in particular for any $(a, b) \in \mathcal{P}_\alpha$ we have*

$$\Psi_{\alpha, r}^{-1}([N(\alpha, a, b)]) = \Phi_{\alpha, r}^{-1}([(a, b)]).$$

We will describe the fibres $\Phi_{\alpha, r}^{-1}([(a, b)])$ of $\Phi_{\alpha, r}$ using the general

Remark 4.28. *Let G be a group, X, B be topological spaces endowed with G -actions by homeomorphisms, $\phi : X \rightarrow B$ a G -equivariant map, and $\Phi : X/G \rightarrow B/G$ the induced map between the quotients. Let $b \in B$. The map*

$$\iota_b : \phi^{-1}(b)/G_b \rightarrow \Phi^{-1}([b]_G), \quad \iota_b([x]_{G_b}) := [x]_G$$

is a continuous bijection. If the map $\phi^{-1}(b) \rightarrow \phi^{-1}(Gb)/G$ induced by the inclusion $\phi^{-1}(b) \hookrightarrow \phi^{-1}(Gb)$ is open, then ι_b is a homeomorphism.

It is easy to see that the openness condition in Remark 4.28 is satisfied, so:

Proposition 4.29. *Let $\mathcal{G}_{(a, b)}$ be the stabiliser of (a, b) in \mathcal{G} . The obvious map*

$$\left(\{(a, b)\} \times \mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right) \right) / \mathcal{G}_{(a, b)} \rightarrow \Phi_{\alpha, r}^{-1}([(a, b)]) = \Psi_{\alpha, r}^{-1}([N(\alpha, a, b)])$$

is a homeomorphism.

Now note that $\{(a, b)\} \times \mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right)$ can be obviously identified with the product $\mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right)$; we let $\mathcal{G}_{(a, b)}$ act on the product $\mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right)$ via this identification and we endow its first factor $\mathcal{C}_{a, b, r}$ with the $\mathcal{G}_{(a, b)}$ -action which makes the first projection

$$p_1 : \mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right) \rightarrow \mathcal{C}_{a, b, r}.$$

$\mathcal{G}_{(a, b)}$ -equivariant. More precisely, for $((\mu, \rho), (K, [k])) \in \mathcal{G}_{a, b}$ and $([c]_{ab}, [t]_{ab}) \in \mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right)$ we have

$$((\mu, \rho), (K, [k])) \cdot ([c]_{ab}, [t]_{ab}) = ([\mu\rho\mathfrak{C}]_{ab}, [\mu\rho\mathfrak{t}]_{ab}), \quad (71)$$

$$((\mu, \rho), (K, [k])) \cdot ([c]_{ab}) = [\mu\rho\mathfrak{C}]_{ab}, \quad (72)$$

where the pair $(\mathfrak{C}, \mathfrak{t})$ is defined by the equality $(K a, K b, \mathfrak{C}, \mathfrak{t}) = K \cdot (k \cdot (a, b, c, t))$.

Applying Remark 4.28 this time to the $\mathcal{G}_{(a, b)}$ -equivariant map p_1 , and taking into account Proposition 4.29, we obtain:

Proposition 4.30. *Let (a, b) be an α -compatible pair. The fibre $\Phi_{\alpha, r}^{-1}([(a, b)])$ comes with a canonical surjective map*

$$\Pi : \Phi_{\alpha, r}^{-1}([(a, b)]) = \left(\mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right) \right) / \mathcal{G}_{(a, b)} \rightarrow \mathcal{C}_{a, b, r} / \mathcal{G}_{(a, b)}$$

induced by p_1 . Its fibre over an orbit $\mathcal{G}_{(a, b)}[c]_{ab}$ is canonically identified with the quotient

$$\left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right) / \mathcal{G}_{(a, b), [c]_{ab}}$$

of $\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z}$ by the stabiliser $\mathcal{G}_{(a, b), [c]_{ab}}$ of $[c]_{ab}$ in $\mathcal{G}_{(a, b)}$.

Therefore, since the quotient topology on the finite set $\mathcal{C}_{a, b, r} / \mathcal{G}_{(a, b)}$ is discrete and any quotient of $\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \simeq \mathbb{C}^*$ is obviously connected, it follows that

Corollary 4.31. *Let (a, b) be an α -compatible pair. The connected components of the fibre $\Phi_{\alpha, r}^{-1}([(a, b)])$ are parameterised by the finite set $\mathcal{C}_{a, b, r} / \mathcal{G}_{(a, b)}$. The connected component corresponding to an orbit $\mathcal{G}_{(a, b)}[c]_{ab}$ is canonically identified with $\left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right) / \mathcal{G}_{(a, b), [c]_{ab}}$.*

Our next goal is to describe explicitly the stabiliser $\mathcal{G}_{(a, b)}$ and the quotient $\mathcal{C}_{a, b, r} / \mathcal{G}_{(a, b)}$ which parameterises the set of connected components of the fibre $\Phi_{\alpha, r}^{-1}([(a, b)])$.

We obviously have:

$$\mathcal{G}_{(a, b)} = \{((\mu, \rho), (K, [k])) \in \mathcal{G} \mid Ka = \mu^{-1}a, Kb = \rho^{-1}b\}. \quad (73)$$

Put $N := N(\alpha, a, b)$. The formulae $Ka = \mu^{-1}a, Kb = \rho^{-1}b$ with $(\mu, \rho) \in \mathbb{R}_+^* \times \mathbb{R}^*$ show that K belongs to the subgroup

$$Z_{\text{GL}(2, \mathbb{Z})}^+(N) := \{K \in \text{GL}(2, \mathbb{Z}) \mid KN = NK, Ka \in \mathbb{R}_+^* a\} \quad (74)$$

of the centraliser

$$\begin{aligned} Z_{\text{GL}(2, \mathbb{Z})}(N) &:= \{K \in \text{GL}(2, \mathbb{Z}) \mid KN = NK\} \\ &= \{K \in \text{GL}(2, \mathbb{Z}) \mid a, b \text{ are eigenvectors for } K\} \end{aligned}$$

of N in the group $\text{GL}(2, \mathbb{Z})$. Note that $Z_{\text{GL}(2, \mathbb{Z})}^+(N)$ is infinite cyclic, see section 5.

$Z_{\text{GL}(2, \mathbb{Z})}^+(N)$ comes with an obvious group morphism $\vartheta : Z_{\text{GL}(2, \mathbb{Z})}^+(N) \rightarrow \mathbb{R}_+^*$ defined by the condition

$$Ka = \vartheta(K)a$$

(i.e. $\vartheta(K)$ is the eigenvalue of K corresponding to the eigenvector a). Note that the eigenvalue of K corresponding to the eigenvector b will be $\varepsilon_K \vartheta(K)^{-1}$, where $\varepsilon_K := \det(K) \in \{\pm 1\}$. Therefore formula (73) becomes

$$\mathcal{G}_{(a, b)} = \{(\vartheta(K)^{-1}, \varepsilon_K \vartheta(K)), (K, [k]) \mid K \in Z_{\text{GL}(2, \mathbb{Z})}^+(N), [k] \in \mathbb{Z}_d^2\}.$$

Note that, by (71) we have

Remark 4.32. *The map*

$$Z_{\text{GL}(2, \mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2 \rightarrow \mathcal{G}_{(a, b)}, (K, [k]) \mapsto ((\vartheta(K)^{-1}, \varepsilon_K \vartheta(K)), (K, [k]))$$

is a group isomorphism, so $\mathcal{G}_{(a, b)}$ is naturally isomorphic to the semidirect product $Z_{\text{GL}(2, \mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2$. Via this isomorphism, $Z_{\text{GL}(2, \mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2$ acts on the spaces $\mathcal{C}_{a, b, r}, \mathcal{C}_{a, b, r} \times \left(\mathbb{C} / \frac{b \wedge a}{r} \mathbb{Z} \right)$ by the formulae

$$(K, [k]) * ([c]_{ab}) = [\varepsilon_K \mathfrak{C}]_{ab}, \quad (75)$$

$$(K, [k]) * ([c]_{ab}, [t]_{ab}) = ([\varepsilon_K \mathfrak{C}]_{ab}, [\varepsilon_K \mathfrak{t}]_{ab}), \quad (76)$$

where the pair $(\mathfrak{C}, \mathfrak{t})$ is defined by the equality $(Ka, Kb, \mathfrak{C}, \mathfrak{t}) = K \cdot (k \cdot (a, b, c, t))$.

Proposition 4.33. *Via the bijection $\mathcal{C}_{a,b,r} \xrightarrow{\cong} \mathbb{Z}^2/(I_2 - N)\mathbb{Z}^2$ induced by $\pi_{a,b,r}$ the subgroup $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ of the semi-direct product $Z_{\text{GL}(2,\mathbb{Z})}^+(N) \ltimes_{\psi} \mathbb{Z}_d^2$ acts on $\mathbb{Z}^2/(I_2 - N)\mathbb{Z}^2$ by*

$$K * [p] = \frac{r}{2} \left((\varepsilon_K K - I_2) \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \varepsilon_K (I_2 - N) \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \right) + \varepsilon_K K[p]. \quad (77)$$

The first term on the right belongs to $r\mathbb{Z}^2$.

Proof. By Remark 4.32 we have $K \cdot [c]_{ab} = [\varepsilon_K C^K]_{ab}$, where

$$\begin{aligned} C^K &= \frac{1}{2} \begin{pmatrix} (Ka)_1(Kb)_1 \\ (Ka)_2(Kb)_2 \end{pmatrix} + K \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \frac{b \wedge a}{2} \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \\ &= \frac{\varepsilon_K}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} + K \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \frac{b \wedge a}{2} \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix}. \end{aligned} \quad (78)$$

Put $P := \pi_{a,b,r}(\varepsilon_K C^K)$. Using Remark 2.15, we obtain:

$$\begin{aligned} \frac{b \wedge a}{r}(P - p) &= (I_2 - N) \left(\varepsilon_K C^K - c \right) = \\ &= (I_2 - N) \left(\frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} + \varepsilon_K K \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \varepsilon_K \frac{b \wedge a}{2} \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} - c \right) \\ &= (I_2 - N) \left((\varepsilon_K K - I_2) \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \varepsilon_K \frac{b \wedge a}{2} \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \right) \\ &= (I_2 - N)(\varepsilon_K K - I_2) \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \varepsilon_K \frac{b \wedge a}{2} (I_2 - N) \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \\ &= (\varepsilon_K K - I_2)(I_2 - N) \left(c - \frac{1}{2} \begin{pmatrix} a_1b_1 \\ a_2b_2 \end{pmatrix} \right) + \varepsilon_K \frac{b \wedge a}{2} (I_2 - N) \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \\ &= b \wedge a \left((\varepsilon_K K - I_2) \left(\frac{1}{2} \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \frac{1}{r}p \right) + \frac{\varepsilon_K}{2} (I_2 - N) \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \right), \end{aligned}$$

which proves the first claim.

For the second claim, we have to prove that

$$\begin{aligned} &(\varepsilon_K K - I_2) \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \varepsilon_K (I_2 - N) \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} = \\ &= \varepsilon_K \left(K \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \varepsilon_N \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} \right) - \varepsilon_K \left(N \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} + \varepsilon_K \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} \right) \in 2\mathbb{Z}^2. \end{aligned}$$

By Lemma 4.16 we know that

$$\begin{aligned} K \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \varepsilon_N \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} &\equiv \begin{pmatrix} (KN)_{11}(KN)_{12} \\ (KN)_{21}(KN)_{22} \end{pmatrix} \pmod{2\mathbb{Z}^2}, \\ N \begin{pmatrix} k_{11}k_{12} \\ k_{21}k_{22} \end{pmatrix} + \varepsilon_K \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} &\equiv \begin{pmatrix} (NK)_{11}(NK)_{12} \\ (NK)_{21}(NK)_{22} \end{pmatrix} \pmod{2\mathbb{Z}^2}. \end{aligned}$$

Since $KN = NK$, this proves the claim. \blacksquare

Theorem 4.34. *Let α be S^+ -admissible and (a, b) be an α -compatible pair. The quotient $\mathcal{C}_{a,b,r}/\mathcal{G}_{(a,b)}$ which parameterises the connected components of the fibre $\Phi_{\alpha,r}^{-1}([(a, b)]) = \Psi_{\alpha,r}^{-1}([N(\alpha, a, b)])$ can be naturally identified with the quotient of*

$$\mathbb{Z}_{N,r} := \mathbb{Z}^2/(I_2 - N)\mathbb{Z}^2 + r\mathbb{Z}^2$$

by the group $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ acting on $\mathbb{Z}_{N,r}$ by $K * [p] := [\varepsilon_K Kp]$.

Proof. We have identified $\mathcal{G}_{(a,b)}$ with the semi-direct product $Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2$, which obviously fits in the short exact sequence

$$0 \rightarrow \mathbb{Z}_d^2 \rightarrow Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2 \rightarrow Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rightarrow \{1\}.$$

It follows that the quotient $\mathcal{C}_{a,b,r}/\mathcal{G}_{(a,b)}$ can be identified with the quotient of $\mathcal{C}_{a,b,r}/\mathbb{Z}_d^2$ by $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$. Taking into account Proposition 4.21 (2), we see that, via the bijection $\mathcal{C}_{a,b,r} \xrightarrow{\cong} \mathbb{Z}^2/(I_2 - N)\mathbb{Z}^2$ induced by $\pi_{a,b,r}$, the quotient $\mathcal{C}_{a,b,r}/\mathbb{Z}_d^2$ is identified with $\mathbb{Z}^2/(I_2 - N)\mathbb{Z}^2 + r\mathbb{Z}^2$. On the other hand, since the first term in (77) belongs to $r\mathbb{Z}^2$, Proposition 4.33 shows that the induced action of $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ on $\mathcal{C}_{a,b,r}/\mathbb{Z}_d^2$ is given by $K * [p] = [\varepsilon_K K p]$, as claimed. \blacksquare

Remark 4.35. *The cyclic subgroup $\langle N \rangle \subset Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ generated by N acts trivially on $\mathbb{Z}_{N,r}$ via $*$.*

For the description of the connected components of the fibre $\Phi_{\alpha,r}^{-1}([(a,b)])$: By Proposition 4.30, the connected component corresponding to the $\mathcal{G}_{(a,b)}$ -orbit of $[c]_{ab} \in \mathcal{C}_{a,b,r}$ is naturally identified with the quotient $(\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z})/\mathcal{G}_{(a,b),[c]_{ab}}$. Although it is difficult to compute the stabiliser $\mathcal{G}_{(a,b),[c]_{ab}}$ explicitly in the general case, we have a good control of its image in the automorphism group of the complex manifold $\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z}$. The subgroup $T \subset \text{Aut}_h(\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z})$ generated by translations associated with torsion elements and the inversion automorphism ι can be identified with the semi-direct product $\mu_2 \rtimes \text{Tors}$, where

$$\text{Tors} = \frac{b \wedge a}{r}\mathbb{Q} / \frac{b \wedge a}{r}\mathbb{Z} \simeq \mathbb{Q}/\mathbb{Z} \simeq \{e^{2\pi i q} \mid q \in \mathbb{Q}\} \subset S^1$$

is the torsion subgroup of $\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z}$, and $\mu_2 := \{\pm 1\}$ acts on Tors by $\varepsilon \cdot \tau := \varepsilon \tau$. Therefore T fits in the short exact sequence

$$0 \rightarrow \text{Tors} \rightarrow T \xrightarrow{\sigma} \mu_2 \rightarrow \{1\},$$

where $\sigma(\tau) = 1$ (-1) if and only if τ preserves (respectively interchanges) the two ends of $\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z} \simeq \mathbb{C}^*$.

Remark 4.36. *Let $H \subset T$ be a finite subgroup.*

- (1) *If $\sigma(H) = \{1\}$, H is a finite (hence cyclic) subgroup of Tors .*
- (2) *If $\sigma(H) = \mu_2$, H is a semi-direct product $\mu_2 \times C$, where $C \subset \text{Tors}$ is a finite cyclic subgroup.*

Proposition 4.37. *Let (a,b) be an α -compatible pair, $[c]_{ab} \in \mathcal{C}_{a,b,r}$. and H the image of the stabiliser $\mathcal{G}_{(a,b),[c]_{ab}}$ in $\text{Aut}_h(\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z})$. Then*

- (1) *H is a finite subgroup of T .*
- (2) *Put $p := \pi_{a,b,r}(c)$. We have $\sigma(H) = \mu_2$ if and only if there exists $L \in Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ with $\varepsilon_L = -1$ such that*

$$\exists L \in Z_{\text{GL}(2,\mathbb{Z})}^+(N) \text{ such that } \varepsilon_L = -1 \text{ and } (I_2 - \varepsilon_L L)p \in r\mathbb{Z}^2 + (I_2 - N)\mathbb{Z}^2. \quad (C)$$

Proof. (1) Formula (76) shows that any element of $\mathcal{G}_{(a,b),[c]_{ab}}$ acts on $\mathbb{C}/\frac{b \wedge a}{r}\mathbb{Z}$ by a formula of the form $[t]_{ab} \mapsto \pm[k] \cdot [t]_{ab}$, where $[k]$ belongs to the finite set \mathbb{Z}_d^2 .

(2) Taking into account formula (76), we see that $\sigma(H) = \mu_2$ if and only if there exists $([k], K) \in Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2$ leaving $[c]_{ab}$ invariant such that $\varepsilon_K = -1$.

An element $([k], K) \in Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rtimes_{\psi} \mathbb{Z}_d^2$ leaves $[c]_{ab}$ invariant, if and only if, putting $L := K^{-1}$, we have

$$[k] \cdot [c]_{ab} = L * [c]_{ab},$$

i.e, using the bijection induced by $\pi_{a,b,r}$, Remark 2.15, and formulae (54),(77),

$$p + r \begin{pmatrix} -k_2 \\ k_1 \end{pmatrix} \equiv \frac{r}{2} \left((\varepsilon_L L - I_2) \begin{pmatrix} n_{11}n_{12} \\ n_{21}n_{22} \end{pmatrix} + \varepsilon_L(I_2 - N) \begin{pmatrix} l_{11}l_{12} \\ l_{21}l_{22} \end{pmatrix} \right) + \varepsilon_L L p$$

mod $(I_2 - N)\mathbb{Z}^2$. It suffices to recall that the first term o the right belongs to $r\mathbb{Z}^2$ by Proposition 4.33. \blacksquare

Now note that the quotient of \mathbb{C}^* by the group of translations associated with a cyclic subgroup $C \subset \mathbb{C}^*$ is biholomorphic with \mathbb{C}^* , whereas the quotient of \mathbb{C}^* by $\langle \iota \rangle$ is biholomorphic with \mathbb{C} .

Therefore

Corollary 4.38. *Let (a, b) be an α -compatible pair. The connected components of the fibre $\Phi_{\alpha,r}^{-1}([(a, b)])$ are biholomorphic to either \mathbb{C}^* or \mathbb{C} . The connected component associated with an orbit $\mathcal{G}_{(a,b)}[c]_{ab}$, where $[c]_{ab} \in \mathcal{C}_{a,b,r}$, is biholomorphic with \mathbb{C} if and only if, putting $p := \pi_{a,b,r}(c)$, condition (C) stated in Proposition 4.37 is satisfied.*

Example 4.2. The case $\theta = 3$, i.e. $\alpha = \frac{3+\sqrt{5}}{2}$.

As explained in the proof of Theorem 5.4 (3), using [BVdM, Section 4] it follows that any $N \in \text{SL}(2, \mathbb{Z})$ with $\text{Tr}(N) = 3$ is similar to the matrix $N_0 := \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$. Therefore the set $\mathcal{N}_\alpha / \sim$ of similarity classes of such matrices reduces to the singleton $\{\mathfrak{N}_0\} = \{[N_0]\}$.

In this case we have $\det(I_2 - N) = -1$, so the quotient $\mathbb{Z}_{N,r} = \mathbb{Z}^2 / (I_2 - N)\mathbb{Z}^2 + r\mathbb{Z}^2$ is a singleton for any $r \in \mathbb{N}^*$. By Theorem 4.34 it follows that, for any fixed $r \in \mathbb{N}^*$, there exists a unique deformation class of type II Inoue surfaces $S_{a,b,c,t}^{\alpha,r}$ with $\text{Tr}(N(\alpha, a, b)) = 3$.

By Theorem 5.4 we know that $Z_{\text{GL}(2,\mathbb{Z})}^+(N_0)$ is generated by the matrix $N'_0 := \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ which satisfies $N_0'^2 = N_0$ with $\det(N'_0) = -1$.

By Corollary 4.38, it follows that for any fixed $r \in \mathbb{N}^*$, the unique connected component of the space of isomorphism classes of type II Inoue surfaces of the form $S_{a,b,c,t}^{\alpha,r}$ is naturally biholomorphic to \mathbb{C} .

Example 4.3. The case $\theta = 4$, i.e. $\alpha = 2 + \sqrt{3}$.

There exists exactly two matrices with determinant 1 and trace 4 which are reduced in the sense of [BVdM, Definition 4.2], namely

$$N' = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}, \quad N'' = \begin{pmatrix} 1 & 1 \\ 2 & 3 \end{pmatrix}.$$

On the other hand the reduction operator P defined on [BVdM, p. 10] maps N' to N'' . Therefore there exists a unique cycle of reduced matrices with determinant 1 and trace 4, so, by [BVdM, Theorem 4.3], a unique similarity class of such matrices.

Therefore the set $\mathcal{N}_\alpha / \sim$ of similarity classes of such matrices reduces to the singleton $\{\mathfrak{N}'\} = \{[N']\}$. The elementary divisors of the matrix

$$(I_2 - N' | rI_2) = \begin{pmatrix} 0 & -2 & r & 0 \\ -1 & -2 & 0 & r \end{pmatrix} \in M_{2,4}(\mathbb{Z})$$

are $\varepsilon_1 = 1$, $\varepsilon_2 = \gcd(2, r)$, so the quotient $\mathbb{Z}_{N',r} = \mathbb{Z}^2 / (I_2 - N')\mathbb{Z}^2 + r\mathbb{Z}^2$ intervening in Theorem 4.34 is isomorphic as a \mathbb{Z} -module to $\mathbb{Z}_{\gcd(2,r)}$.

On the other hand by Theorem 5.4 the positive centraliser $Z_{\text{GL}(2,\mathbb{Z})}^+(N')$ is generated by N' , so it acts trivially on $\mathbb{Z}_{N',r}$ by Remark 4.35. Therefore, by Theorem 4.34, for any $r \in \mathbb{N}^*$ there are exactly $\gcd(2, r)$ deformation classes of type II Inoue

surfaces $S_{a,b,c,t}^{\alpha,r}$, and, by Corollary 4.38, both connected components of the space of isomorphism classes of such type II Inoue surfaces is naturally biholomorphic to \mathbb{C}^* .

4.3. The classification of type III Inoue surfaces. The classification of type III Inoue surfaces can be obtained using the same method as for type II surfaces: For a triple $(a, b, c) \in \mathcal{T}_{\alpha,r}$ put $[c]_{ab} := c + \frac{b \wedge a}{r} \mathbb{Z}^2$ and we define

$$\begin{aligned} \mathcal{C}_{a,b,r} &:= \mathcal{C}_{a,b,r} / \frac{b \wedge a}{r} \mathbb{Z}^2, \\ \mathcal{T}_{\alpha,r} &:= \{(a, b, [c]_{ab}) \mid (a, b) \in \mathcal{P}_\alpha, [c]_{ab} \in \mathcal{C}_{a,b,r}\} \simeq \mathcal{T}_{\alpha,r} / \mathbb{Z}^2. \end{aligned}$$

We let $\mathrm{GL}(2, \mathbb{Z})$ act on $\mathcal{T}_{\alpha,r}$ via the same formula (45) as for type II surfaces, and with this definition the analogue of Theorem 4.17 for type III surfaces remains true. Therefore the set of subgroups $G \subset \mathrm{Aff}(U)$ defining type III Inoue surfaces is identified with the quotient $\mathcal{T}_{\alpha,r} / \mathrm{GL}(2, \mathbb{Z})$.

On the other hand we have to use a different formula for defining a \mathbb{Z}^2 -action on $\mathcal{T}_{\alpha,r}$; this new formula is provided by the following analogue of Lemma 4.18:

Lemma 4.39. *Let $(a, b, c), (a', b', c') \in \mathcal{T}_{\alpha,r}$ and put*

$$\begin{cases} g_0 = g_0(\alpha), & g_i = g_i(a, b, c), & g_3 = g_3(a, b, r), \\ g'_0 = g_0(\alpha), & g'_i = g_i(a', b', c'), & g'_3 = g_3(a', b', r) \end{cases}.$$

Let $k = (k_1, k_2) \in \mathbb{Z}^2$, $\begin{pmatrix} s_0 \\ s_1 \\ s_2 \end{pmatrix} \in \mathbb{Z}^3$ and $\tau \in \mathrm{Aff}_1^1(U)$, $\begin{pmatrix} w \\ z \end{pmatrix} \xrightarrow{\tau} \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} w \\ z \end{pmatrix} + \begin{pmatrix} u \\ \zeta \end{pmatrix}$.

The following conditions are equivalent:

- (i) $\tau g'_0 \tau^{-1} = g_0 g_1^{k_1} g_2^{k_2} g_3^{s_0}$, $\tau \circ g'_i \circ \tau^{-1} = g_i g_3^{s_i}$ for $1 \leq i \leq 2$, $\tau \circ g'_3 \circ \tau^{-1} = g_3$.
- (ii) We have:

$$\begin{cases} \lambda &= \frac{kb}{\alpha-1}. \\ u &= \frac{\alpha(ka)}{1-\alpha}. \\ 2\zeta &= \lambda u(1+\alpha) - \frac{k_1(k_1-1)}{2} a_1 b_1 - \frac{k_2(k_2-1)}{2} a_2 b_2 \\ &\quad - k_1 k_2 b_1 a_2 - kc - \frac{s_0}{r} b \wedge a. \\ a' &= a. \\ b' &= b. \\ c' &= c + \frac{kb}{1+\alpha} a + \frac{\alpha(ka)}{1-\alpha} b + \frac{b \wedge a}{r} s, \text{ where } s := \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}. \end{cases}$$

Note that, whereas for type II surfaces, both conditions in Lemma 4.18 were independent of the coefficient ζ of τ , this is no longer true for type III surfaces. In Lemma 4.39 both conditions do depend on ζ .

We obtain an action

$$(k, c) \mapsto k \cdot c := c + \frac{kb}{1+\alpha} a + \frac{\alpha(ka)}{1-\alpha} b = c + b \wedge a (I_2 + N)^{-1} \begin{pmatrix} -k_2 \\ k_1 \end{pmatrix} \quad (79)$$

on $\mathcal{C}_{a,b,r}$ and an induced action $(k, [c]_{ab}) \mapsto k \cdot [c]_{ab} := [k \cdot c]_{ab}$ on $\mathcal{C}_{a,b,r}$.

Noting that for type III surfaces we have $\det(I_2 + N) = \theta \in \mathbb{N}^*$, we see that the obtained \mathbb{Z}^2 action on $\mathcal{T}_{\alpha,r}$ descends to a \mathbb{Z}_θ^2 -action. In the same way as for type II surfaces we obtain actions of the semi-direct product $\mathrm{GL}(2, \mathbb{Z}) \rtimes_\psi \mathbb{Z}_\theta^2$ and of the product

$$\mathcal{G} := (\mathbb{R}_+^* \times \mathbb{R}^*) \times (\mathrm{GL}(2, \mathbb{Z}) \rtimes_\psi \mathbb{Z}_\theta^2)$$

on $\mathcal{T}_{\alpha,r}$, and the following classification theorem

Theorem 4.40. *Let α be S^- -admissible and $(a, b, c), (a', b', c') \in \mathcal{T}_{\alpha, r}$. The groups $G(\alpha, a, b, c, r), G(\alpha, a', b', c', r)$ are conjugate in $\text{Aff}(U)$ if and only if*

$$\begin{aligned} (a', b', [c']_{a'b'}) &\in ((\mathbb{R}_+^* \times \mathbb{R}^*) \times (\text{GL}(2, \mathbb{Z}) \times_{\psi} \mathbb{Z}_0^2)) \cdot (a, b, [c]_{ab}). \\ &= ((\mathbb{R}_+^* \times \mathbb{R}^*) \times (\text{GL}(2, \mathbb{Z}) \times_{\varphi} \mathbb{Z}^2)) \cdot (a, b, [c]_{ab}). \end{aligned}$$

Let now

$$\Psi_{\alpha, r} : \mathcal{Q}_{\alpha, r}^- := \mathcal{T}_{\alpha, r}/\mathcal{G} \rightarrow \mathcal{N}_{\alpha}/\mathcal{G} = \mathcal{N}_{\alpha}/\text{GL}(2, \mathbb{Z}) = \mathcal{N}_{\alpha}/\sim,$$

$$\Phi_{\alpha, r} : \mathcal{Q}_{\alpha, r}^- \rightarrow \mathcal{P}_{\alpha, r}^-/\mathcal{G} = \mathcal{P}_{\alpha, r}^-/(\mathbb{R}_+^* \times \mathbb{R}^*) \times \text{GL}(2, \mathbb{Z})$$

be the obvious maps. Using now the methods of section 4.2.3, we obtain

Theorem 4.41. *Let α be S^- -admissible and (a, b) be an α -compatible pair. The quotient $\mathcal{C}_{\alpha, b, r}/\mathcal{G}_{(a, b)}$ which parameterises the connected components of the fibre*

$$\Phi_{\alpha, r}^{-1}([(a, b)]) = \Psi_{\alpha, r}^{-1}([N(\alpha, a, b)])$$

can be naturally identified with the quotient of

$$\mathbb{Z}_{N, r} := \mathbb{Z}^2/(I_2 + N)\mathbb{Z}^2 + r\mathbb{Z}^2$$

by the group $Z_{\text{GL}(2, \mathbb{Z})}^+(N)$ acting on $\mathbb{Z}_{N, r}$ by $K * [p] := [\varepsilon_K K p]$.

5. APPENDIX: THE POSITIVE CENTRALISER $Z_{\text{GL}(2, \mathbb{Z})}^+(N)$

The ‘‘positive centraliser’’ $Z_{\text{GL}(2, \mathbb{Z})}^+(N)$ associated with a matrix $N \in \mathcal{N}_{\alpha}$ ($N \in \mathcal{N}_{\alpha}^-$), where α is S^+ - (respectively S^- -) admissible plays a crucial role in our results. This group is always infinite cyclic. This follows from:

Proposition 5.1. *Let $N \in \text{GL}(2, \mathbb{Z})$ with $\text{Spec}_{\mathbb{R}}(N) \cap]1, +\infty[\neq \emptyset$. Fix $\alpha \in \text{Spec}(N) \cap]1, +\infty[$ and an eigenvector $a \in \mathbb{R}^2 \setminus \{0\}$ for the eigenvalue α .*

- (1) *The group $Z_{\text{SL}(2, \mathbb{Z})}^+(N) := \{K \in \text{SL}(2, \mathbb{Z}) \mid KN = NK, Ka \in \mathbb{R}_+^* a\}$ is infinite cyclic.*
- (2) *Suppose $\alpha \notin \mathbb{Q}$. Then $Z_{\text{GL}(2, \mathbb{Z})}^+(N) := \{K \in \text{GL}(2, \mathbb{Z}) \mid KN = NK, Ka \in \mathbb{R}_+^* a\}$ is also infinite cyclic.*

Proof. (1) Note first that the group morphism

$$\vartheta|_{Z_{\text{SL}(2, \mathbb{Z})}^+(N)} : Z_{\text{SL}(2, \mathbb{Z})}^+(N) \rightarrow \mathbb{R}_+^*$$

is injective, so it induces an isomorphism $Z_{\text{SL}(2, \mathbb{Z})}^+(N) \xrightarrow{\cong} \vartheta(Z_{\text{SL}(2, \mathbb{Z})}^+(N)) \subset \mathbb{R}_+^*$. Recall that any closed proper subgroup of \mathbb{R} is either trivial or infinite cyclic. Since the Lie groups $(\mathbb{R}_+^*, \cdot), (\mathbb{R}, +)$ are isomorphic, the same will hold for the subgroups of (\mathbb{R}_+^*, \cdot) .

Since the subgroup $\vartheta(Z_{\text{SL}(2, \mathbb{Z})}^+(N))$ is non-trivial (because it contains $\alpha > 1$) and is proper (because is countable), it suffices to prove that $\vartheta(Z_{\text{SL}(2, \mathbb{Z})}^+(N))$ is closed in \mathbb{R}_+^* .

Let b be an eigenvector for the second eigenvalue $\beta = \det(N)\alpha^{-1}$ of N . For any $u \in \mathbb{R}_+^*$ let

$$K_{(a, b)}^u := \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} \begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix}^{-1} \in M_2(\mathbb{R})$$

be the (unique) real $(2, 2)$ -matrix which admits a, b as eigenvectors with eigenvalues u, u^{-1} respectively.

We obviously have

$$\vartheta(Z_{\text{SL}(2, \mathbb{Z})}^+(N)) = \{u \in \mathbb{R}_+^* \mid K_{(a, b)}^u \in M_2(\mathbb{Z})\}$$

which shows that $\vartheta(Z_{\text{SL}(2,\mathbb{Z})}^+(N))$ is closed in \mathbb{R}_+^* , because the map $u \mapsto K_{(a,b)}^u$ is continuous and $M_2(\mathbb{Z})$ is closed in $M_2(\mathbb{R})$.

(2) We claim that, under our assumptions, the group morphism

$$\vartheta : Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rightarrow \mathbb{R}_+^*$$

is still injective. Indeed, we have

$$\ker(\vartheta) \subset \{I_2, L\},$$

where $L \in M_2(\mathbb{R})$ is the matrix of the endomorphism of \mathbb{R}^2 which admits a, b as eigenvectors with eigenvalues $1, -1$ respectively. We will show that $L \notin M_2(\mathbb{Q})$ so this matrix cannot be an element of $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$. Indeed, if L belonged to $M_2(\mathbb{Q})$, its eigenspace $\mathbb{R}a$ associated with the (rational) eigenvalue 1 would admit a rational generator $q = \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} \in \mathbb{Q}^2 \setminus \{0\}$. Since $\mathbb{R}a$ is also the eigenspace of N associated with the eigenvalue α , it would follow $Nq = \alpha q$ so, since $N \in M_2(\mathbb{Q})$, it would follow $\alpha \in \mathbb{Q}$ which contradicts our assumption. This proves that ϑ is injective as claimed.

Since ϑ is injective, $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ is isomorphic to a subgroup of $(\mathbb{R}_+^*, \cdot) \simeq (\mathbb{R}, +)$, so it is torsion free. On the other hand, this abelian group fits in the short exact sequence

$$\{1\} \rightarrow Z_{\text{SL}(2,\mathbb{Z})}^+(N) \rightarrow Z_{\text{GL}(2,\mathbb{Z})}^+(N) \xrightarrow{\det} \{\pm 1\} \rightarrow 1,$$

which, taking into account (1), shows that it is a finitely generated abelian group of rank 1. Since it is torsion free, it is infinite cyclic. \blacksquare

Note that

Remark 5.2. *The natural group morphisms*

$$Z_{\text{SL}(2,\mathbb{Z})}^+(N) \rightarrow \text{PSL}(2,\mathbb{Z}), \quad Z_{\text{GL}(2,\mathbb{Z})}^+(N) \rightarrow \text{PGL}(2,\mathbb{Z})$$

are injective and their images coincide with the centralisers

$$Z_{\text{PSL}(2,\mathbb{Z})}([N]), \quad Z_{\text{PGL}(2,\mathbb{Z})}([N])$$

of $[N]$ in $\text{PSL}(2,\mathbb{Z})$, respectively $\text{PGL}(2,\mathbb{Z})$.

Therefore Lemma 5.1 gives:

Corollary 5.3. *In the conditions of Lemma 5.1, the centraliser $Z_{\text{PSL}(2,\mathbb{Z})}(N)$ of $[N]$ in $\text{PSL}(2,\mathbb{Z})$ is infinite cyclic. If $\alpha \notin \mathbb{Q}$, the centraliser $Z_{\text{PGL}(2,\mathbb{Z})}([N])$ of N in $\text{PGL}(2,\mathbb{Z})$ is also infinite cyclic.*

This result answers a question discussed by experts on the Mathoverflow forum, see [Centers of Elements in \$\text{SL}\(2,\mathbb{Z}\)\$ on Mathoverflow](#).

A natural problem: given a matrix N satisfying the assumptions of Proposition 5.1, specify a generator of the cyclic group $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$. The following result answer this question in a particular case:

Theorem 5.4. *Let $N \in \text{SL}(2,\mathbb{Z})$ with $\theta := \text{Tr}(N) \geq 3$. Suppose that*

$$\gcd(n_{12}, n_{21}, n_{22} - n_{11}) = 1.$$

Then:

- (1) $Z_{\text{SL}(2,\mathbb{Z})}^+(N) = \{N^k \mid k \in \mathbb{Z}\}$, so $Z_{\text{SL}(2,\mathbb{Z})}^+(N)$ is the cyclic subgroup of $\text{SL}(2,\mathbb{Z})$ generated by N .
- (2) If $\theta > 3$, then $Z_{\text{GL}(2,\mathbb{Z})}^+(N) = Z_{\text{SL}(2,\mathbb{Z})}^+(N)$.
- (3) If $\theta = 3$, then $Z_{\text{GL}(2,\mathbb{Z})}^+(N)$ is generated by a matrix $N' \in \text{GL}(2,\mathbb{Z})$ with $N'^2 = N$, $\det(N') = -1$.

Proof. (1) Since the characteristic polynomial of N has no multiple roots in \mathbb{C} , it follows that by a classical theorem in matrix theory [We, Theorem 4, p. 27] that

$$Z_{M_{2,2}(\mathbb{C})}(N) = \{P(N) \mid P(X) \in \mathbb{C}[X]\} = \langle I_2, N \rangle_{\mathbb{C}}.$$

Therefore

$$Z_{M_{2,2}(\mathbb{Q})}(N) = Z_{M_{2,2}(\mathbb{C})}(N) \cap M_{2,2}(\mathbb{Q}) = \langle I_2, N \rangle_{\mathbb{C}} \cap M_{2,2}(\mathbb{Q}) = \langle I_2, N \rangle_{\mathbb{Q}},$$

so

$$\begin{aligned} Z_{M_{2,2}(\mathbb{Z})}(N) &= Z_{M_{2,2}(\mathbb{Q})}(N) \cap M_{2,2}(\mathbb{Z}) = \\ &= \{xN + yI_2 \mid (x, y) \in \mathbb{Q}^2, xN + yI_2 \in M_{2,2}(\mathbb{Z})\}. \end{aligned} \quad (80)$$

Let $x = \frac{m}{n}$, $y = \frac{p}{q}$ with $\gcd(m, n) = \gcd(p, q) = 1$ be such that $L := xN + yI_2$ is an integer matrix. We have

$$\frac{m}{n}n_{12} = l_{12} \in \mathbb{Z}, \quad \frac{m}{n}n_{21} = l_{21} \in \mathbb{Z}, \quad \frac{m}{n}(n_{22} - n_{11}) = l_{22} - l_{11},$$

so $n \mid mn_{12}$, $n \mid mn_{21}$, $n \mid m(n_{22} - n_{11})$. But $\gcd(n, m) = 1$, so $n \mid n_{12}$, $n \mid n_{21}$ and $n \mid (n_{22} - n_{11})$, so $n \mid \gcd(n_{12}, n_{21}, (n_{22} - n_{11}))$. We assumed $\gcd(n_{12}, n_{21}, (n_{22} - n_{11})) = 1$, so $n = \pm 1$, so $x \in \mathbb{Z}$, so $yI_2 = L - xN \in M_{2,2}(\mathbb{Z})$, so $y \in \mathbb{Z}$. Therefore, by (80), we obtain

$$Z_{M_{2,2}(\mathbb{Z})}(N) = \{xN + yI_2 \mid (x, y) \in \mathbb{Z}^2\} = \langle I_2, N \rangle_{\mathbb{Z}}. \quad (81)$$

Putting $\alpha := \frac{\theta + \sqrt{\theta^3 - 4}}{2}$, we obtain $\det(xN + yI_2) = (x\alpha + y)(x\alpha^{-1} + y) = x^2 + \theta xy + y^2$, so, by (81),

$$\begin{aligned} Z_{\text{SL}(2, \mathbb{Z})}(N) &= \{xN + yI_2 \mid (x, y) \in \mathbb{Z}^2, x^2 + \theta xy + y^2 = 1\}, \\ Z_{\text{GL}(2, \mathbb{Z})}(N) &= \{xN + yI_2 \mid (x, y) \in \mathbb{Z}^2, x^2 + \theta xy + y^2 = \pm 1\}. \end{aligned} \quad (82)$$

For $k \in \mathbb{Z}$ let $(a_k, b_k) \in \mathbb{Z}^2$ be defined by the condition $N^k = a_k N + b_k I_2$, and let $(u_n)_{n \in \mathbb{N}}$ be the integer sequence defined by the order 2 recurrence

$$u_0 = 0, \quad u_1 = 1, \quad u_{k+2} = \theta u_{k+1} - u_k.$$

Using the formulae $N^2 = \theta N - I_2$, $(N^{-1})^2 = \theta N^{-1} - I_2$, it is easy to prove that

$$(a_k, b_k) = \begin{cases} (u_k, -u_{k-1}) & \text{for } k \geq 1 \\ (-u_{-k}, u_{-k+1}) & \text{for } k \leq 0. \end{cases}$$

By [AA, Theorem 5.6.1, p. 130] and [AA, section 6.3], it follows that, for $\theta \geq 3$, the set of solutions of the diophantine equation $x^2 + \theta xy + y^2 = 1$ is precisely

$$\{\pm(-u_n, u_{n+1}) \mid n \in \mathbb{N}\} \cup \{\pm(u_{n+1}, -u_n) \mid n \in \mathbb{N}\} = \{\pm(a_k, b_k) \mid k \in \mathbb{Z}\}.$$

Therefore, the first formula in (82) combined with $N^k = a_k N + b_k I_2$ gives

$$Z_{\text{SL}(2, \mathbb{Z})}(N) = \{\pm N^k \mid k \in \mathbb{Z}\},$$

which implies that $Z_{\text{SL}(2, \mathbb{Z})}^+(N) = \{N^k \mid k \in \mathbb{Z}\}$. This proves (1).

(2) follows by second formula in (82) taking into account that, by [AA, Theorem 6.3.1, p. 150], for $\theta > 3$, the diophantine equation $x^2 + \theta xy + y^2 = -1$ has no solution.

(3) Using [BVdM, Theorem 4.3] it follows that any matrix $N \in \text{SL}(2, \mathbb{Z})$ with $\text{Tr}(N) = 3$ is similar to $N_0 := \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$. Indeed, there exists a unique integer matrix with trace 3 and determinant 1 which is reduced in the sense of [BVdM, Definition 4.2]. Note that the matrix $N'_0 := \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ is a square root of N_0 , belongs to $Z_{\text{GL}(2, \mathbb{Z})}^+(N_0)$ and has $\det(N'_0) = -1$. Let K be a generator of the cyclic group

$Z_{\text{GL}(2,\mathbb{Z})}^+(N_0)$, and let $s \in \mathbb{Z}$ such that $N'_0 = K^s$. Since $K^2 \in Z_{\text{SL}(2,\mathbb{Z})}^+(N_0)$ (which is generated by N_0), there exists $t \in \mathbb{Z}$ such that $K^2 = N_0^t$. It follows

$$N_0 = N_0'^2 = (K^s)^2 = K^{2s} = (K^2)^s = N_0^{ts},$$

so $ts = 1$, so $s = \pm 1$, which proves that $N'_0 = K^s$ is also a generator of $Z_{\text{GL}(2,\mathbb{Z})}^+(N_0)$. ■

REFERENCES

- [AA] T. Andreescu, D. Andrica, Quadratic Diophantine Equations, Developments in Mathematics vol. 40, Springer, New York Heidelberg Dordrecht London, 2015.
- [BHPV] W. Barth, K. Hulek, Ch. Peters, A. Van de Ven, Compact Complex Surfaces, Springer-Verlag, Berlin-Heidelberg (2004).
- [BVdM] A. Behn, A.B. Van der Merwe, An algorithmic version of the theorem by Latimer and MacDuffee for 2×2 integral matrices, *Invar Algebra and its Applications* 346 (2002) 1–14.
- [BoSh] Z. I. Borevich, I. R. Shafarevich, Number Theory, Academic Press, New York San Francisco London, 1966.
- [Du] S. Dumitrescu, Connexions affines et projectives sur les surfaces complexes compactes, *Math. Z.* 264, 301–316 (2010).
- [In] M. Inoue, On Surfaces of Class VII₀, *Inventiones math.* 24, 269 - 310 (1974).
- [LMD] C. G. Latimer and C. C. MacDuffee, A Correspondence Between Classes of Ideals and Classes of Matrices, *Ann. of Math.*, vol. 34 (1933), 313-316.
- [Mil] J. Milne, Algebraic Number Theory, Lecture Notes, [Algebraic Number Theory](#).
- [Ta] O. Taussky, On a Theorem of Latimer and Macduffee, *Canadian Journal of Mathematics*, Volume 1, Issue 3 (1949), 300-302.
- [We] J. Wehler, Versal deformations of Hopf surfaces, *Journal für die reine und angewandte Mathematik* 328 (1981), 22-32.
- [Wed] J. H. M. Wedderburn, Lectures on Matrices, Amer. Math. Soc. Colloquium Publications, vol. 17 (1934).

AIX MARSEILLE UNIVERSITÉ, CNRS, I2M, UMR 7373, 13453 MARSEILLE, FRANCE, EMAIL: ZAHRAA.KHALED@UNIV-AMU.FR, ANDREI.TELEMEN@UNIV-AMU.FR