

Universality of scaled particle spectra in ultrarelativistic heavy-ion collisions

Cicero D. Muncinelli,^{1,*} Fernando G. Gardim,^{2,3,†} David D. Chinellato,⁴ Gabriel S. Denicol,⁵ Andre V. Giannini,^{6,7} Matthew Luzum,⁸ Jorge Noronha,³ Tiago Nunes da Silva,⁹ Jun Takahashi,¹ and Giorgio Torrieri¹
(The ExTrEMe Collaboration)

¹*Universidade Estadual de Campinas (Unicamp),*

R. Sérgio Buarque de Holanda, 777, Campinas, Brazil, 13083-859

²*Instituto de Ciência e Tecnologia, Universidade Federal de Alfenas, 37715-400 Poços de Caldas, MG, Brazil*

³*Illinois Center for Advanced Studies of the Universe & Department of Physics,*

University of Illinois at Urbana-Champaign, Urbana, IL 61801-3003, USA

⁴*Stefan Meyer Institute for Subatomic Physics of the Austrian Academy of Sciences, Wiesingerstraße 4 1010 Vienna, Austria*

⁵*Instituto de Física, Universidade Federal Fluminense,*

Av. Milton Tavares de Souza, Niterói, Brazil, 24210-346,

⁶*Faculdade de Ciências Exatas e Tecnologia, Universidade Federal da Grande Dourados, Dourados, MS, Brazil, 79804-970*

⁷*Departamento de Física, Universidade do Estado de Santa Catarina, 89219-710 Joinville, SC, Brazil*

⁸*Instituto de Física, Universidade de São Paulo,*

R. do Matão, 1371, São Paulo, Brazil, 05508-090

⁹*Departamento de Física, Centro de Ciências Físicas e Matemáticas, Universidade Federal de Santa Catarina, Campus Universitário Reitor João David Ferreira Lima, Florianópolis, Brazil, 88040-900*

(Dated: June 24, 2024)

We propose a new observable derived from a centrality-dependent scaling of transverse particle spectra. By removing the global scales of total particle number and mean transverse momentum, we isolate the shape of the spectrum. In hydrodynamic simulations, while the multiplicity and mean transverse momentum fluctuate significantly, the scaled spectrum is found to be almost constant even at an event-by-event level and after resonance decays. This universality survives when averaging over events in each centrality bin before scaling. We then investigate the presence of this scaling in experimental data from the ALICE collaboration in Pb-Pb, Xe-Xe, and p-Pb collisions. We find a remarkable universality in the experimentally observed scaled spectra at low transverse momentum, compatible with hydrodynamic predictions. The data show a minor breaking of universality at large transverse momentum and hints of evolution with the system size that are not seen in simulations. Our results motivate further theoretical and experimental investigations of this new observable to bring to light the collective and non-collective behavior encoded in the transverse particle spectrum of different collision systems.

I. INTRODUCTION

High-energy nuclear collisions investigate the fundamental properties of bulk nuclear matter under extreme conditions, shedding light on the nature of strong interactions. These collisions yield a large number of particles; in the case of lead-lead collisions at the Large Hadron Collider (LHC), up to 35,000 particles can be produced. By studying the properties and correlations between these particles, one can infer significant information about the dynamics of the generated state [1]. The experimental results obtained from relativistic heavy-ion collisions performed at the Relativistic Heavy Ion Collider (RHIC) and the LHC indicate that the resulting state exhibits a phase with characteristics resembling those of a relativistic viscous fluid composed of quarks and gluons [2], with a temperature of a few billion Kelvin [3], in a volume less than 10^{-45} m³, known as the quark-gluon plasma (QGP). In proton-lead collisions, where the reaction takes place in a volume one thousand

times smaller than in lead-lead collisions, the matter produced still exhibits the characteristics of a QGP fluid when a large number of particles are produced [1, 4–7].

The transverse momentum distribution of particles measured in heavy ion collisions reflects the collective behavior of the relativistic fluid produced at the early stages of the collision. In this regard, the azimuthal transverse momentum distribution of charged hadrons, $dN/d\phi$, encoded in the flow coefficients v_n [8–11], has been the most insightful observable as it quantifies the final state momentum anisotropy of nuclear collisions. In fluid-dynamical models, this anisotropy emerges due to the hydrodynamic response to the initial-state geometry of the system, and its magnitude reflects how viscous the QGP is [12].

Naturally, the azimuthally integrated transverse momentum distribution, dN/dp_T , must also carry relevant information about the fluid produced in these collisions. So far, this has been mostly explored using only two moments of this distribution: the number of charged particles, N_{ch} , and the mean transverse momentum¹, $\langle p_T \rangle$.

* c250891@dac.unicamp.br

† fernando.gardim@unifal-mg.edu.br

¹To obtain $\langle p_T \rangle$, additional modeling [13–15] is used to extrapolate

These integrated observables are sensitive to the transport properties of the fluid, such as bulk viscosity [16], and are usually included in global fits of hydrodynamic models through Bayesian analyses [17–19]. Nevertheless, it remains unclear what new information about the QGP’s hydrodynamic evolution is concealed in the full differential p_T -spectra.

In this paper, we propose a new observable derived from a centrality-dependent scaling of transverse particle spectra to unravel fundamental features of fluid-dynamical models that have not been pointed out before. For this purpose, we isolate the shape of the spectra by removing the global scales of total particle number, N , and $\langle p_T \rangle$. In practice, this is done by considering the following normalized transverse momentum distribution:

$$U(x_T) \equiv \frac{\langle p_T \rangle}{N} \frac{dN}{dp_T} = \frac{1}{N} \frac{dN}{dx_T}, \quad (1)$$

where $x_T = p_T / \langle p_T \rangle$. We demonstrate that in hydrodynamic simulations, while the multiplicity and mean transverse momentum fluctuate significantly event by event, the scaled spectrum is almost identical even at an event-by-event level and after resonance decays, displaying a nearly universal shape. We further show that this feature of hydrodynamic models survives even when averaging over events before scaling, though it can depend somewhat on the collision system – thus, this prediction of hydrodynamic models can, in principle, be checked experimentally. This is done here by performing the scaling on experimental data from the ALICE collaboration in Pb-Pb, Xe-Xe, and p-Pb collisions. Indeed, we observe that the experimentally measured scaled spectra at low transverse momentum display a remarkable universal behavior, which is in qualitative agreement with the hydrodynamic prediction. However, the data show a minor breaking of this universality at large transverse momentum (marking the onset of semi-hard physics) and hints of evolution with the system size that are not seen in simulations.

II. EVENT-BY-EVENT RESULTS IN HYBRID SIMULATIONS

To understand the properties of the scaled transverse momentum spectra proposed in (1), we first present the results of hybrid simulations for Pb-Pb collisions at 2.76 TeV, which involve: (i) initial conditions from the TRENTo model [20], (ii) a pre-equilibrium phase modeled using $\text{K}\phi\text{MP}\phi\text{ST}$ [21], (iii) 2+1 viscous hydrodynamics simulations performed using MUSIC [22, 23] with the equation of state [24], (iv) Cooper-Frye procedure [25] and particlization via the hypersurface sampler iSS [26],

and (v) hadronic transport modeling using UrQMD [27]. For these simulations, TRENTo and viscous parameters were obtained from the Bayesian analysis in [17]. Detailed information about these simulations can be found in [28]. In the present work, 2334 minimum bias events were used, with oversampling.

Ultrarelativistic nucleus-nucleus collisions produce a variety of particles, including pions, kaons, and protons, which are the most abundant species. The relative proportions of these particles vary based on the collision energy. Pions are the most abundant particles, comprising roughly 80% of the total yield [29] due to their low mass threshold. Kaons and protons are produced in smaller quantities, with kaons being lighter and more abundant than protons, accounting for approximately 10% and 5%, respectively [30]. Therefore, for simplicity, we present in this work only the results for charged pions, though we have checked that kaon and proton spectra display similar universal behavior. In Figure 1 (left), event-by-event p_T spectra for Pb-Pb collisions at 2.76 TeV across different centralities are shown, distinguished by different colors. It is evident that dN/dp_T changes with centrality and that there is a large variation within each centrality class. However, once the scaling proposed in Eq. (1) is applied to the p_T spectra, all events approximately collapse onto the same curve, making them indistinguishable from one another. We employed the scaled spectra for each event, denoted as U_{ev} , as defined by equation (1), incorporating the multiplicity of each event, N_{ev} , and its average transverse momentum $\langle p_T \rangle_{\text{ev}}$. The scaled spectra shown in Fig. 1 (right) reveal universal behavior for any event, not just between centralities, which could be interpreted as a signature of the hydrodynamic phase. Additionally, we computed the standard deviation of U_{ev} for each x_T and observed that the deviations are less than 5% up to $x_T = 4$.




FIG. 1. (left) Presents the pion, $\pi^+ + \pi^-$, p_T -spectra for Pb-Pb at 2.76 TeV minimum bias, in the hybrid simulations. (right) Scaled spectra for each event, $U_{\text{ev}}(x_T) = (\langle p_T \rangle_{\text{ev}} / N_{\text{ev}}) dN_{\text{ev}} / dp_T$, computed using Eq. 1, for the events in the left plot. Centrality-independent universal behavior in $U_{\text{ev}}(x_T)$ is observed.

olate the spectrum down to $p_T = 0$ GeV/c.

III. VARYING SYSTEMS AND ENERGY

It is natural to wonder if the universality observed above holds for different energies and systems. Furthermore, it is crucial to expand our analysis to include small systems, as there is an ongoing debate about whether the same features observed in large systems apply to small systems [7, 31]. Given the consistent trend observed across events, we will focus on analyzing $U(x_T)$ per centrality, as this is the most accessible way to obtain experimental data. We verified this with two new sets of simulations of 1000 events each: p-Pb and Pb-Pb at 5.02 TeV, using parameters from [32, 33], (see Figure 2). After fixing the nuclei and collision energy, we still observe agreement in U across centralities, even for peripheral collisions of 70-80%, as shown in the top panel of Figure 2. Nevertheless, the shape of U changes when the model and collision system change. We use throughout this paper open symbols to represent centralities up to 40%, and filled symbols for the other centralities, the peripheral ones. Results up to 90-100% centrality are not shown, but they lead to the same conclusion.

To accurately compare the values of U in different centralities, we computed the ratio between the U of a specific centrality and the one obtained in the most central collision category. We avoided double interpolation by interpolating the x_T -spectra for the 0-5% bin and using it to calculate the ratio in the x_T from the numerator. In the three bottom plots of Figure 2, one can see that all U curves are almost identical, with small deviations at low and high x_T . These deviations are expected since the hybrid simulation incorporates decay and hadronic scattering, which can slightly modify some hydrodynamic signatures. At low x_T , particularly for pions, the spectra undergo some modifications due to decay. At high x_T , there is an additional contribution from hadron scattering [13]. The boundary covering 99% of pions produced is indicated by vertical lines, providing a reliable region where universality is observed in hydrodynamic models, regardless of energy and system size.

If systems of different sizes with equal collision energy exhibit the same multiplicity, it can be suggested that their x_T -spectra should be identical. This corresponds to the scenario involving Pb-Pb 70-80% and p-Pb 0-5% at 5.02 TeV. These collisions are represented by red diamonds and green circles in Figure 2 (a). It is evident that the corresponding $U(x_T)$ values for these collisions do not exhibit agreement. Having the same multiplicity does not necessarily mean having the same energy density and gradients, which means it will not lead to the same hydrodynamic expansion.

Finally, we have examined the spectra of kaons and protons and reached similar conclusions. Further investigation of these particles will be presented in future work.




FIG. 2. (a) The scaled spectra, Eq. (1), for $\pi^+ + \pi^-$ in Pb-Pb at 2.76 TeV and 5.02 TeV, and p-Pb 5.02 TeV, computed using hybrid simulations. Once the system is fixed, there is universality across centralities. (b) The ratio between U coming from four centralities and U for the most central collisions in Pb-Pb 2.76 TeV. (c) Ratio between U coming from four centralities and U for the most central collisions in Pb-Pb 5.02 TeV. (d) Ratio between U coming from four centralities and U for the most central collisions in p-Pb 5.02 TeV. We note that 99% of the pions are produced to the left of the vertical gray lines. The ExTReME simulations (b) used the parameters from [28], and the hybrid simulation presented in (c) and (d) used Duke parameters [32, 33].

IV. EXPERIMENTAL RESULTS ON THE SCALED SPECTRA

Since our simulations predict quasi-universal behavior for the scaled spectrum, it is crucial to investigate whether similar behavior for $U(x_T)$ is present in the available experimental data across various energies and systems. In the case of heavy-ion collisions, we used ALICE Pb-Pb data at 2.76 TeV [30] and 5.02 TeV [29] as well as Xe-Xe data at 5.44 TeV [34]. For p-Pb collisions at 5.02 TeV, we used ALICE data from [29]. To obtain $U(x_T)$, it is essential to have information about the p_T -spectra down to 0 GeV/c, as the computation of the global quantities, $\langle p_T \rangle$ and N_{ch} , relies on this information. Since there are no measurements for $p_T < 0.1$ GeV/c for pions, further modeling is used to access these scales through extrapolations performed by the ALICE collaboration [30, 35].

Before comparing the scaled spectra within the same system, it is important to emphasize that the experimen-

tal data contains physical effects that our simulations did not consider. This includes the impact of jets on the medium, which affects not only high-momentum particles but also those with low momentum. Furthermore, the medium response to jets can differ due to variables such as size, centralities, and atomic mass number. Additionally, the p_T -spectrum for large momentum fails to correspond to thermal behavior and requires a description using a power-law function [36, 37], indicating that it does not solely arise from hydrodynamics.

The results for different centralities, ranging from central to more peripheral regions, are presented in Figure 3, with error bars derived from the p_T -spectra. For a fixed collision system and energy, represented by different colors in the plot, it is possible to observe the emergence of a universal x_T -spectrum, Figure 3 (a). In the case of p-Pb, although universality is seen across centralities, $U(x_T)$ is not the same as in heavy-ion collisions. The ALICE results, combined with our findings from hybrid simulations, suggest that universality may depend on energy and system size.

The comparison between $U(x_T)$ across centralities for the Pb-Pb and Xe-Xe measurements, as shown in Figure 3 (b), (c), and (d), reveals universality within the 0-40% centrality range. However, agreement quickly diminishes beyond this centrality range. This change in the behavior of U may indicate that the matter created in these different regimes exhibits distinct physics. As expected, there is a deviation for large $p_T > 3 \langle p_T \rangle$, which is roughly independent of the system. However, the universality remains up to 99% of the pions created, as indicated by the vertical gray lines in Figure 3. This region may serve as an indication of where (semi-)hard physics begins to govern the production of particles.

The $U(x_T)$ spectra for p-Pb are shown in the bottom plot (e) of Figure 3. For large x_T , all centralities are compatible; however, for x_T close to 1 there is a significant deviation starting from 40% centrality, with a stronger deviation for more peripheral events. This breaking in scaling does not occur in our simulations, indicating that for events with low multiplicity, i.e. centralities greater than 40%, there is no compatibility between the universality of the scaled spectra of a fluid and the experimental data. Therefore, the new observable proposed in this paper gives further evidence for the hydrodynamic behavior of p-Pb collisions at high multiplicity, providing information about collectivity in small systems independently from anisotropic flow measurements.

The behavior of $U(x_T)$ for kaons and protons in ALICE data was also examined, although not shown here, and the conclusions presented in this paper are unaffected.

V. CONCLUSION

The present work introduces a new observable, the scaled transverse momentum spectrum $U(x_T)$, see (1), which serves as a new signature of the collective hy-




FIG. 3. (a) The scaled spectra $U(x_T)$, as defined in Eq. (1), for $\pi^+ + \pi^-$ in Pb-Pb at 2.76 TeV [30] and 5.02 TeV [29], Xe-Xe 5.44 TeV [34], and p-Pb 5.02 TeV from experimental data [29]. Once the system is fixed, there is universality across centralities. (b) The ratio between U coming from four centralities and U for the most central collisions in Pb-Pb 2.76 TeV. (c) Ratio between U coming from four centralities and U for the most central collisions in Pb-Pb 5.02 TeV. (d) Ratio between U coming from four centralities and U for the most central collisions in Xe-Xe 5.44 TeV. (e) Ratio between U coming from four centralities and U for the most central collisions in p-Pb 5.02 TeV. We note that 99% of the pions are produced to the left of the vertical gray lines.

drodynamic behavior of the quark-gluon plasma, independently from anisotropic flow. When hydrodynamics dominate the description of matter, the resulting scaled spectrum reveals a unified pattern in all centralities for fixed energy and system type. Decay and hadronic scatterings minimally affect this universal behavior. Our conclusions were achieved using the standard modeling of ultrarelativistic collisions, with parameters obtained from Bayesian analyses for Pb-Pb and p-Pb collisions at energies of 2.76 TeV and 5.02 TeV across different centralities.

We focused this work on pion spectra because they constitute the predominant fraction of observed particles in ultrarelativistic nuclear collisions, accounting for ap-

proximately 80% of the total yield. The scaled spectra in our simulations were computed for three systems: Pb-Pb at 2.76 and 5.02 TeV and p-Pb at 5.02 TeV. For these systems, $U(x_T)$ displayed universal behavior for all centralities up to $p_T < 3.5 \langle p_T \rangle$ on an event-by-event basis. However, we found in our simulations that the shape of $U(x_T)$ can change with center-of-mass energy and the type of collision system. Further systematic studies of this dependence can be straightforwardly done using modern emulation techniques [19].

The x_T spectrum in (1) can be easily obtained from the experimental data of the p_T spectra. Using ALICE data, we computed the scaled spectra for charged pions in Pb-Pb collisions at 2.76 and 5.02 TeV, p-Pb collisions at 5.02 TeV, and also in Xe-Xe collisions at 5.44 TeV. Universality in U up to $x_T < 3$ is observed for centralities below 40% in Pb-Pb and Xe-Xe collisions. Beyond this point, universal behavior ceases, suggesting that additional physics beyond hydrodynamics may be needed as one moves toward more peripheral events. This finding is consistent with the overall success of hydrodynamic-based hybrid model calculations in describing various observables, including anisotropic flow, average transverse momentum, and multiplicity. The breakdown of universality in experimental data at large x_T is expected due to the onset of non-hydrodynamic physics phenomena [38, 39].

The observable proposed in this work brings new light to investigating collectivity in ultrarelativistic collisions. In p-Pb collisions, where there is an ongoing debate about the formation of quark-gluon plasma and the validity of hydrodynamics in its description, we find using ALICE data that universal behavior holds for high multiplicity events, up to 40% centrality, with pronounced deviations when $x_T \approx 1$. This provides new evidence that a hydrodynamically-expanding quark-gluon plasma is formed in high multiplicity p-Pb events. In this regard, we note that our results are compatible with a previous study of the inferred thermodynamic properties of p-Pb collisions in the same regime [6]. Clearly, it would be interesting to extend the analysis done here also to p-p collisions.

This scaling is not trivial to describe using simple models amenable to a closed form analytical formula. In particular, we find that both the blast-wave [13] and the Tsallis [40] distribution do not reproduce the scaling when the model parameters (temperature, flow velocity, chemical potential and Tsallis q) are varied without considerable fine-tuning. Also, an intermediate p_T scale

where partons are thermalized but hadronize via recombination rather than Cooper-Frye does not seem to be borne out by the scaling, since $\langle p_T \rangle$ is firmly below this scale and the scaling is borne out across species.

Understanding the origin of the universality of scaled particle spectra in experimental data is crucial to determining whether this feature is indeed a robust signature of the hydrodynamic behavior of QCD matter or just a general consequence of collectivity without hydrodynamics. The scaled spectrum can be investigated in low center-of-mass-energy collisions to determine the importance of the hydrodynamic phase in such systems. Furthermore, since near a critical point hydrodynamics is modified by the presence of additional slow degrees of freedom [41], it may be interesting to investigate the scaled spectrum also in the context of the ongoing search for the QCD critical point [42]. Finally, our new observable can be instrumental to attest to the existence (or not) of hydrodynamic droplets in other systems, such as in $e^+ - e^-$ collisions [43–45], or the presence of collective effects inside individual jets [46, 47]. Such investigations may lead to new fundamental questions about the formation and evolution of hot and dense QCD matter in extreme conditions.

ACKNOWLEDGMENTS

G.S.D. thanks J.-F. Paquet for fruitful discussions. F.G.G. is supported by CNPq (Conselho Nacional de Desenvolvimento Científico) through 306762/2021-8 and by the Fulbright Program. F.G.G. and T. N.d.S. are supported by CNPq through the INCT-FNA grant 312932/2018-9 and the Universal Grant 409029/2021-1. M.L. was supported by FAPESP projects 2017/05685-2 and 2018/24720-6, by project INCT-FNA Proc. No. 464898/2014-5, and by CAPES - Finance Code 001. G.S.D. acknowledges support from CNPq and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), grant No. E-26/202.747/2018. J.T. was supported by FAPESP projects 2017/05685-2 and CNPq through 309174/2020-1. J.N. is partially supported by the U.S. Department of Energy, Office of Science, Office for Nuclear Physics under Award No. DE- SC0023861. C.D.M. was supported by FAPESP project 2017/05685-2 through the 2022/16166-4 grant, and by CNPq through the PIBIC grant 122740/2023-8. G.T. acknowledges support from Bolsa de produtividade CNPQ 305731/2023-8, Bolsa de pesquisa FAPESP 2023/06278-2.

[1] W. Busza, K. Rajagopal, and W. van der Schee, *Ann. Rev. Nucl. Part. Sci.* **68**, 339 (2018), arXiv:1802.04801 [hep-ph].
 [2] U. Heinz and R. Snellings, *Ann. Rev. Nucl. Part. Sci.* **63**, 123 (2013), arXiv:1301.2826 [nucl-th].

[3] F. G. Gardim, G. Giacalone, M. Luzum, and J.-Y. Ollitrault, *Nature Phys.* **16**, 615 (2020), arXiv:1908.09728 [nucl-th].
 [4] B. Abelev *et al.* (ALICE), *Phys. Lett. B* **719**, 29 (2013), arXiv:1212.2001 [nucl-ex].

- [5] V. Khachatryan *et al.* (CMS), JHEP **09**, 091 (2010), arXiv:1009.4122 [hep-ex].
- [6] F. G. Gardim, R. Krupczak, and T. N. da Silva, Phys. Rev. C **109**, 014904 (2024), arXiv:2212.11710 [nucl-th].
- [7] J. Noronha, B. Schenke, C. Shen, and W. Zhao (2024) arXiv:2401.09208 [nucl-th].
- [8] J.-Y. Ollitrault, Phys. Rev. D **46**, 229 (1992).
- [9] F. G. Gardim, F. Grassi, M. Luzum, and J.-Y. Ollitrault, Phys. Rev. Lett. **109**, 202302 (2012), arXiv:1203.2882 [nucl-th].
- [10] M. Luzum and H. Petersen, J. Phys. G **41**, 063102 (2014), arXiv:1312.5503 [nucl-th].
- [11] J. Adam *et al.* (ALICE), Phys. Rev. Lett. **117**, 182301 (2016), arXiv:1604.07663 [nucl-ex].
- [12] M. Luzum and P. Romatschke, Phys. Rev. C **78**, 034915 (2008), [Erratum: Phys.Rev.C 79, 039903 (2009)], arXiv:0804.4015 [nucl-th].
- [13] E. Schnedermann, J. Sollfrank, and U. W. Heinz, Phys. Rev. C **48**, 2462 (1993), arXiv:nucl-th/9307020.
- [14] B. I. Abelev *et al.* (STAR), Phys. Rev. C **79**, 034909 (2009), arXiv:0808.2041 [nucl-ex].
- [15] A. Guillen and J.-Y. Ollitrault, Phys. Rev. C **103**, 064911 (2021), arXiv:2012.07898 [nucl-th].
- [16] S. Ryu, J. F. Paquet, C. Shen, G. S. Denicol, B. Schenke, S. Jeon, and C. Gale, Phys. Rev. Lett. **115**, 132301 (2015), arXiv:1502.01675 [nucl-th].
- [17] J. E. Bernhard, J. S. Moreland, S. A. Bass, J. Liu, and U. Heinz, Phys. Rev. C **94**, 024907 (2016), arXiv:1605.03954 [nucl-th].
- [18] G. Nijs, W. van der Schee, U. Gürsoy, and R. Snellings, Phys. Rev. C **103**, 054909 (2021), arXiv:2010.15134 [nucl-th].
- [19] D. Everett *et al.* (JETSCAPE), Phys. Rev. C **103**, 054904 (2021), arXiv:2011.01430 [hep-ph].
- [20] J. S. Moreland, J. E. Bernhard, and S. A. Bass, Phys. Rev. C **92**, 011901 (2015), arXiv:1412.4708 [nucl-th].
- [21] A. Kurkela, A. Mazeliauskas, J.-F. Paquet, S. Schlichting, and D. Teaney, Phys. Rev. Lett. **122**, 122302 (2019), arXiv:1805.01604 [hep-ph].
- [22] B. Schenke, S. Jeon, and C. Gale, Phys. Rev. C **82**, 014903 (2010), arXiv:1004.1408 [hep-ph].
- [23] J.-F. Paquet, C. Shen, G. S. Denicol, M. Luzum, B. Schenke, S. Jeon, and C. Gale, Phys. Rev. C **93**, 044906 (2016), arXiv:1509.06738 [hep-ph].
- [24] P. Huovinen and P. Petreczky, Nucl. Phys. A **837**, 26 (2010), arXiv:0912.2541 [hep-ph].
- [25] F. Cooper and G. Frye, Phys. Rev. D **10**, 186 (1974).
- [26] C. Shen, Z. Qiu, H. Song, J. Bernhard, S. Bass, and U. Heinz, Comput. Phys. Commun. **199**, 61 (2016), arXiv:1409.8164 [nucl-th].
- [27] S. A. Bass *et al.*, Prog. Part. Nucl. Phys. **41**, 255 (1998), arXiv:nucl-th/9803035.
- [28] T. Nunes da Silva, D. Chinellato, M. Hippert, W. Serenone, J. Takahashi, G. S. Denicol, M. Luzum, and J. Noronha, Phys. Rev. C **103**, 054906 (2021), arXiv:2006.02324 [nucl-th].
- [29] S. Acharya *et al.* (ALICE), Phys. Rev. C **101**, 044907 (2020), arXiv:1910.07678 [nucl-ex].
- [30] B. Abelev *et al.* (ALICE), Phys. Rev. C **88**, 044910 (2013), arXiv:1303.0737 [hep-ex].
- [31] J. L. Nagle and W. A. Zajc, Ann. Rev. Nucl. Part. Sci. **68**, 211 (2018), arXiv:1801.03477 [nucl-ex].
- [32] J. S. Moreland, J. E. Bernhard, and S. A. Bass, Phys. Rev. C **101**, 024911 (2020), arXiv:1808.02106 [nucl-th].
- [33] T. N. da Silva, D. D. Chinellato, A. V. Giannini, M. N. Ferreira, G. S. Denicol, M. Hippert, M. Luzum, J. Noronha, and J. Takahashi (ExTrEMe), Phys. Rev. C **107**, 044901 (2023), arXiv:2211.10561 [nucl-th].
- [34] S. Acharya *et al.* (ALICE), Eur. Phys. J. C **81**, 584 (2021), arXiv:2101.03100 [nucl-ex].
- [35] B. B. Abelev *et al.* (ALICE), Phys. Lett. B **728**, 25 (2014), arXiv:1307.6796 [nucl-ex].
- [36] R. Hagedorn, Riv. Nuovo Cim. **6N10**, 1 (1983).
- [37] B. B. Abelev *et al.* (ALICE), Eur. Phys. J. C **73**, 2662 (2013), arXiv:1307.1093 [nucl-ex].
- [38] S. Acharya *et al.* (ALICE), Phys. Lett. B **776**, 249 (2018), arXiv:1702.00804 [nucl-ex].
- [39] A. M. Sirunyan *et al.* (CMS), JHEP **05**, 284 (2021), arXiv:2102.13080 [hep-ex].
- [40] Z. Tang, Y. Xu, L. Ruan, G. van Buren, F. Wang, and Z. Xu, Phys. Rev. C **79**, 051901 (2009), arXiv:0812.1609 [nucl-ex].
- [41] M. Stephanov and Y. Yin, Phys. Rev. D **98**, 036006 (2018), arXiv:1712.10305 [nucl-th].
- [42] X. An *et al.*, Nucl. Phys. A **1017**, 122343 (2022), arXiv:2108.13867 [nucl-th].
- [43] A. Badaea, A. Baty, P. Chang, G. M. Innocenti, M. Maggi, C. McGinn, M. Peters, T.-A. Sheng, J. Thaler, and Y.-J. Lee, Phys. Rev. Lett. **123**, 212002 (2019), arXiv:1906.00489 [hep-ex].
- [44] Y. C. Chen *et al.* (Belle), Phys. Rev. Lett. **128**, 142005 (2022), arXiv:2201.01694 [hep-ex].
- [45] Y.-C. Chen *et al.*, (2023), arXiv:2312.05084 [hep-ex].
- [46] A. Baty, P. Gardner, and W. Li, Phys. Rev. C **107**, 064908 (2023), arXiv:2104.11735 [hep-ph].
- [47] A. Hayrapetyan *et al.* (CMS), (2023), arXiv:2312.17103 [hep-ex].