

Computing True Parameter Values Using Monte Carlo Integration in Simple and Complex Simulation Designs

Ashley I. Naimi, PhD^{1,*}

David Benkeser, PhD²

Jacqueline E. Rudolph, PhD³

¹ Department of Epidemiology, Emory University.

² Department of Biostatistics, Emory University.

³ Department of Epidemiology, Johns Hopkins University.

*Correspondence: Department of Epidemiology
Rollins School of Public Health
Emory University
1518 Clifton Road
Atlanta, GA 30322
ashley.naimi@emory.edu

Conflicts of Interest: None

Acknowledgements: We thank Dr Tim Morris for helpful comments on a previous version of this manuscript.

Sources of Funding: AIN was supported by NIH grant number R01 HD102313.

Target Journal:	<i>Epidemiol</i>
Text word count:	2,063
Abstract word count:	178
Number of Figures:	2
Number of Tables:	0
Number of References:	15
Running head:	Monte Carlo Integration for Simulation Estimands

Abstract

Simulation studies are used to evaluate and compare the properties of statistical methods in controlled experimental settings. In most cases, performing a simulation study requires knowledge of the true value of the parameter, or estimand, of interest. However, in many simulation designs, the true value of the estimand is difficult to compute analytically. Here, we illustrate the use of Monte Carlo integration to compute true estimand values in simple and complex simulation designs. We provide general pseudocode that can be replicated in any software program of choice to demonstrate key principles in using Monte Carlo integration in two scenarios: a simple three variable simulation where interest lies in the marginally adjusted odds ratio; and a more complex causal mediation analysis where interest lies in the controlled direct effect in the presence of mediator-outcome confounders affected by the exposure. We discuss general strategies that can be used to minimize Monte Carlo error, and to serve as checks on the simulation program to avoid coding errors. R programming code is provided illustrating the application of our pseudocode in these settings.

KEY WORDS: Epidemiologic methods; Monte Carlo Simulation; Monte Carlo integration; Numeric integration; Causal Inference; Statistics.

Introduction

Simulation studies are often employed to evaluate and compare the properties of statistical methods in controlled experimental settings. Simulation studies can be used towards a wide range of aims, from basic conceptual or didactic purposes,¹ to formal research questions about the performance of different analytic methods.² Monte Carlo simulation studies are so named because they belong to a general category of techniques known as Monte Carlo methods,³ a class of methods that rely on *in silico* (pseudo) random number generation to find solutions to problems that may otherwise be intractable.⁴

The design of Monte Carlo simulation studies often involves several steps. These include clearly articulating the aims of the simulation study, the data generating mechanisms that will be used, the estimand of interest, the methods that will be used to estimate the estimand, and the performance measures used to evaluate the properties of the methods under study.² Simulation studies proceed by repeatedly simulating data sets from the selected data generating mechanism. Each simulated data set is analyzed using the selected methods. The performance of the selected methods is compared by aggregating results across multiple simulated data sets. For example, to approximate the bias of an estimator, we can compute the empirical average (across all simulated data sets) difference between the point estimates and the truth. To approximate the coverage of a method for building a confidence interval, we can compute the empirical proportion (out of all simulated data sets) of intervals that contain the true parameter value. This makes clear the importance for the researcher to know the precise true value of the estimand that is implied by the selected data generating process.

In ideal settings, the true value of the estimand will be directly ascertainable from the parameter values used in the data generating mechanism. For example, if the data generating mechanism involves simulating a binary outcome Y using a logistic regression model that includes a single binary exposure A , and a single continuous confounder C , $P(Y = 1 | A, C) = \text{expit}\{\beta_0 + \beta_1 A + \beta_2 C\}$. To simulate data from this data generating distribution, we would generally need to select values for β_0 , β_1 , and β_2 . If our simulation study is interested in the conditionally adjusted odds ratio for the exposure-outcome association, then the true estimand value is easily computed as $\exp(\beta_1)$, where

β_1 is the researcher-selected value.

In such simple examples, it is possible to read off the true estimand value directly from the data generating process. However, in many settings, simulation studies are used to evaluate estimators of estimands whose values cannot be so easily ascertained. Importantly, in such settings, the true estimand value may not be easily computed as a single parameter, or simple combination of parameters, used to generate the data. Here, we outline the use of Monte Carlo integration for solving the true estimand value. We illustrate the approach using two example scenarios: a data generating mechanism where the estimand of interest is the marginally adjusted odds ratio; and a causal mediation setting where interest lies in the controlled direct effect as the estimand. We provide pseudocode for each example that can be used to construct a program implementing Monte Carlo integration in a programming language of choice. We also provide R code for each example in an associated [GitHub](#) repository.

Example 1: Marginally Adjusted Odds Ratio as the Estimand

The first example involves estimating the marginally adjusted odds ratio using methods such as inverse probability weighting or marginal standardization^{5,6}. The causal diagram representing our data generating mechanism⁷ is displayed in Figure 1:




Figure 1: Causal diagram depicting the relationships between an exposure A , outcome Y , and confounder C .

A parametric approach to simulating data compatible with the causal diagram in Figure 1 is to first simulate C from, e.g., a Normal distribution with mean μ and variance σ^2 . Then based on the

simulated values of C , simulate the exposure A from a logistic regression model with $P(A = 1 | C) = \text{expit}(\alpha_0 + \alpha_1 C)$, for some researcher-selected values α_0, α_1 . Finally, the outcome is simulated from a logistic regression model

$$P(Y = 1 | X, C) = \text{expit}\{\beta_0 + \beta_1 A + \beta_2 C\} . \quad (1)$$

As noted above, if the true parameter of interest was the conditionally adjusted odds ratio, its value could easily be obtained as $\exp(\beta_1)$. However, if our interest lies in evaluating the performance of methods that estimate the marginally adjusted odds ratio, we would have to compute the true value of this estimand entailed by the model.

There are well known and important differences between marginally versus conditionally adjusted odds ratios,⁸⁻¹⁰ due to the fact that the odds ratio is a noncollapsible measure of effect.¹¹ Noncollapsibility is distinct from confounding,¹² which means that for a given logistic regression model with the same set of correctly chosen confounders, the marginally versus conditionally adjusted odds ratio may be numerically different. This means that, in the context of equation 1, while exponentiating the value of the β_1 coefficient will provide us with a numerical value for the conditionally adjusted odds ratio, the marginally adjusted odds ratio may be a different value.

Under ideal conditions, one would rely on algebraic derivations to obtain an analytic solution to the marginally adjusted odds ratio. Computing a marginally adjusted odds ratio analytically from a model such as equation 1 could be accomplished via integration, which would require solving for:

$$\mu(a) = \int_c \text{expit}\{\beta_0 + \beta_1 a + \beta_2 c\} \frac{1}{(2\pi)^{1/2}\sigma} \exp\left\{-\frac{(c-\mu)^2}{2\sigma^2}\right\} dc ,$$

for $a = 0, 1$. The solution to this integral yields a marginal probability of the outcome that would be observed if $A = a$. One can then construct and odds from this probability as: $\frac{\mu(1)}{1-\mu(1)}$. The marginally adjusted odds ratio can then be computed as:

$$\psi = \frac{\mu(1)}{1-\mu(1)} \bigg/ \frac{\mu(0)}{1-\mu(0)}$$

However, even in this simple setting with a single variable C , analytically solving for this integral is challenging. One could employ the use of numeric integration available in many statistical software packages (See [GitHub](#) repository). However, this approach would not scale well to settings with more than one C variable.

Instead, we can use Monte Carlo integration to solve for ψ , as follows:

Algorithm 1 Pseudocode for Implementing Monte Carlo Integration to Solve for the True Marginally Adjusted Odds Ratio in a Simple Three Node Causal Diagram

- 1: set random number generator seed value
- 2: set large sample size N
- 3: simulate $i \in 1 \dots N$ observations C_i from a Normal distribution with mean μ and variance σ^2
- 4: compute $\hat{\mu}_i(x) = \text{expit}(\beta_0 + \beta_1 x + \beta_2 C_i)$ for both $x = 0$ and $x = 1$. Thus, $\hat{\mu}_i(0) = \text{expit}(\beta_0 + \beta_2 C_i)$ and $\hat{\mu}_i(1) = \text{expit}(\beta_0 + \beta_1 + \beta_2 C_i)$.
- 5: the approximated value of $\mu(x)$ is given by the mean of $\hat{\mu}_i(x)$ over all N simulated observations, $\hat{\mu}(x) = \frac{1}{N} \sum_{i=1}^N \hat{\mu}_i(x)$. The approximated value of ψ is given by

$$\hat{\psi} = \frac{[\hat{\mu}(1)/(1 - \hat{\mu}(1))]}{[\hat{\mu}(0)/(1 - \hat{\mu}(0))]}$$

The approximated value of ψ obtained from the pseudocode above can be used as the true estimand value for a simulation study evaluating the properties of an estimator seeking to quantify the marginally adjusted odds ratio. However, it is important to note that this "true" estimand value depends on: (i) the specifications for the distribution of C ; and (ii) the parameter values for the logistic regression model generating Y in this case, $\beta_0, \beta_1, \beta_2$. If the distribution of C changes, or if the regression model coefficients change, this pseudoalgorithm should be run again under these new settings. Additionally, the value of ψ will be subject to error that depends on the sample size used in the Monte Carlo integration. For this reason, N should be chosen to be as large as possible under the allocated computing constraints. Several iterations of the pseudo-code could be run to confirm that the sample is large enough to lead to only minimal changes in the approximated value.

Of note, in step 5 of the above pseudocode, one should compute $\hat{\mu}_i(a = 0)$ and $\hat{\mu}_i(a = 1)$ directly from the logistic model for the outcome, and not to convert these probabilities to binary outcomes with a draw from the Bernoulli distribution. Doing so prevents the introduction of additional Monte Carlo error, and reduces the variability of the Monte Carlo integration technique.


Example 2: Controlled Direct Effect as the Estimand

The second example consists of a more complex causal mediation analysis where interest may lie in conducting a simulation study to evaluate the properties of methods used to estimate the controlled direct effect (CDE). Here, we define the CDE of interest on the difference scale, but other scales could be entertained:

$$\psi = E(Y^{a,m} - Y^{a^*,m})$$

In the above, $Y^{a,m}$ is the potential outcome that would be observed if the exposure A were set to some value a and the mediator M were set to some value m , while $Y^{a^*,m}$ is the corresponding value that would be observed under some referent a^* . The causal diagram representing our data generating mechanism is displayed in Figure 2.

Figure 2: Causal diagram depicting the relationships between an exposure A , a mediator M , a confounder C , a mediator-outcome confounder affected by the exposure L , and an outcome Y .



In this example, the true value of the controlled direct effect is difficult to compute analytically. It consists of a function of the magnitudes of the following paths in Figure 2:

$$A \rightarrow Y$$

$$A \rightarrow M \rightarrow Y$$

$$A \rightarrow L \rightarrow Y$$

In particular, the specific function that should be used to combine these magnitudes will depend on how M , L , and Y are distributed, and whether the effect magnitudes for each arrow are on the multiplicative or additive scales. In fact, in many settings, it simply may not be possible to analytically solve for the true controlled direct effect. However, we can use Monte Carlo integration as follows:

Algorithm 2 Pseudocode for Implementing Monte Carlo Integration to Solve for the True Controlled Direct Effect in a Causal Mediation Analysis DAG with a Mediator-Outcome Confounder Affected by the Exposure.

- 1: set random number generator seed value.
- 2: set large sample size N .
- 3: simulate $i \in 1 \dots N$ observations C_i and U_i from a distribution of choice.
- 4: construct a_1 and a_0 for all N observations corresponding to exposed (e.g., $a_1 = 1$) and referent (e.g., $a_0 = 0$) states.
- 5: simulate two L_i variables, one corresponding to a_1 ($L_i^{a_1}$) and one corresponding to a_0 ($L_i^{a_0}$), for all N observations. This would be from a regression model that includes the simulated U_i and the specified a_x value, along with relevant parameters.
- 6: construct m for all N observations corresponding to the specific mediator value of interest in the controlled direct effect contrast (e.g., $m = 0$ for all N).
- 7: simulate $Q_i^{a_x, m}$ for all $i \in 1 \dots N$ observations from the model for the outcome:

$$Q_i^{a_x, m} = f(\beta_0 + \beta_1 a_x + \beta_2 C_i + \beta_3 m + \beta_4 L_i + \beta_5 U_i),$$

where $f(\bullet)$ represents a specified link function of interest.

- 8: take the mean of $Q_i^{a_x, m}$ over all N to obtain $\mu_{a_x, m} = \frac{1}{N} \sum_{i=1}^N Q_i^{a_x, m}$, and construct a difference measure as:

$$\tilde{\psi} = \mu_{a_1, m} - \mu_{a_0, m}$$

Again, the value of $\tilde{\psi}$ obtained from the pseudocode above can be used as the true estimand value for the controlled direct effect under the specified simulation settings. The magnitude of this true value will once again depend on the specific parameter values, distributions, and functional forms selected to generate all variables used in the outcome model. If any of these are changed, a separate $\tilde{\psi}$ should be quantified. Additionally, as a check of the algorithm's validity, one could use Monte Carlo integration to estimate the true conditional association between the exposure-outcome, mediator-outcome, or confounder-outcome relationships, or any other known quantity in the data generating mechanism under study.

Discussion

Monte Carlo simulation studies are commonly used to evaluate the properties of estimators in a range of different scientific settings. However, estimator properties such as bias, mean squared error, and confidence interval coverage require a numerical value for the true estimand of interest. In many settings, this true value cannot be obtained analytically, and Monte Carlo integration can be used instead.

There are several important issues to consider when designing a program to implement Monte Carlo integration to compute estimand values. First, general recommendations suggest that the selected sample size used to compute the true value should be as large as is computationally feasible to minimize Monte Carlo error to the smallest value possible. However, in practice a degree of Monte Carlo error could be tolerated. For example, if increasing the Monte Carlo sample size past a certain boundary κ only changes the estimand value at the 5th or greater decimal place, it might be advisable to set the Monte Carlo sample size to κ to avoid expending resources for little meaningful gain. Second, it is important to note that the source of Monte Carlo error in any simulation study is derived from programming calls to the system's random number generator. For example, in the R programming language, functions used to generate observations from a distribution of interest [e.g., `rnorm()`, `runif()`, or `rexp()`], or functions used to sample from a set of observations, will contribute to Monte Carlo error. For this reason, one strategy that should be used to minimize Monte Carlo error in the estimand value is to minimize these calls where possible (e.g., Step 5 in Algorithm 1, or Step 7 in Algorithm 2).

Several classes of simulation study design exist. In fully parametric simulation studies, each component of the data generating mechanism is specified parametrically using researcher defined functions. For example, confounders, exposures, and outcomes will all be drawn from known distributions with pre-specific relationships between each. Alternatively, plasmode simulations can be conducted in which a portion of the data generating mechanism (e.g., confounders) can be constructed using actual data sampled from a population of interest.¹³ Doing so preserves the empirical associations between variables drawn from the sample, thus providing a closer approximation between simulated and real data. To compute the true value using Monte Carlo integration with

a plasmode design using data from a fixed sample size, one could resample the original data with replacement to obtain a sufficiently large sample that will minimize Monte Carlo error. Finally, synthetic simulations have recently been developed that rely on machine learning to simulate data whose variables and joint distributions between them closely approximate an index dataset sampled from a real population of interest, but where the effect of the exposure of interest is specified *a priori*.^{14,15} In principle, Monte Carlo integration can be used in all three settings to derive the true estimand of interest, or to serve as a verification step in the process of simulating complex data. Indeed, the Monte Carlo method is a general methodological strategy that has been used across the sciences to solve simple and complex problems. With increasingly complex analyses being carried out in epidemiology and the health sciences, the Monte Carlo method, and Monte Carlo intergation more specifically, will be a useful tool to gain insights into complex questions.

References

1. Rudolph JE, Fox MP, and Naimi AI. Simulation as a tool for teaching and learning epidemiologic methods. *American Journal of Epidemiology*. 2021; **190**:900–907.
2. Morris TP, White IR, and Crowther MJ. Using simulation studies to evaluate statistical methods. *Statistics in Medicine*. 2019; **38**:2074–2102.
3. Metropolis N and Ulam S. The Monte Carlo method. *J Am Stat Assoc*. 1949; **44**:335–341.
4. McCracken DD. The monte carlo method. *Scientific American*. 1955; **192**:90–97.
5. Naimi, AI, Cole SR, Kennedy EH. Introduction to G Methods. *International Journal of Epidemiology*. 2017; **46**:756-762.
6. Naimi, AI, Whitcomb BW. Estimating Risk Ratios and Risk Differences Using Regression. *American Journal of Epidemiology*. 2020; **189**:508-510.
7. Fox MP, Nianogo R, Rudolph JE, and Howe CJ. Illustrating how to simulate data from directed acyclic graphs to understand epidemiologic concepts. *American Journal of Epidemiology*. 2022; **191**:1300–1306.
8. Neuhaus JM and Jewell NP. A geometric approach to assess bias due to omitted covariates in generalized linear models. *Biometrika*. 1993; **80**:807–815.
9. Daniel R, Zhang J, and Farewell D. Making apples from oranges: Comparing noncollapsible effect estimators and their standard errors after adjustment for different covariate sets. *Biom J*. 2021; **63**:528–557.
10. Pang M, Kaufman JS, and Platt RW. Studying noncollapsibility of the odds ratio with marginal structural and logistic regression models. *Stat Methods Med Res*. 2013; .
11. Greenland S. Collapsibility. In: Gail MH and Bénichou J (Eds.) *Encyclopedia of Epidemiologic Methods*. John Wiley & Sons, Ltd. 2005; .
12. Pang M, Kaufman JS, and Platt RW. Mixing of confounding and non-collapsibility: a notable deficiency of the odds ratio. *Am J Cardiol*. 2013; **111**:302–303.
13. Franklin JM, Schneeweiss S, Polinski JM, and Rassen JA. Plasmode simulation for the evaluation of pharmacoepidemiologic methods in complex healthcare databases. *Comput Stat Data Anal*. 2014; **72**:219–226.
14. Athey S, Imbens GW, Metzger J, and Munro E. Using wasserstein generative adversarial networks for the design of monte carlo simulations. *Journal of Econometrics*. 2021; :105076.

15. Parikh H, Varjao C, Xu L, and Tchetgen ET. Validating causal inference methods. *International Conference on Machine Learning*. 2022; **PMLR**.