

ON A BERS THEOREM FOR  $SL(3, \mathbb{C})$ 

CHRISTIAN EL EMAM AND NATHANIEL SAGMAN

ABSTRACT. For  $S$  a closed surface of genus at least 2, let  $\text{Hit}_3(S)$  be the Hitchin component of representations to  $SL(3, \mathbb{R})$ , equipped with the Labourie-Loftin complex structure. We construct a mapping class group equivariant holomorphic map from a large open subset of  $\text{Hit}_3(S) \times \overline{\text{Hit}_3(S)}$  to the  $SL(3, \mathbb{C})$ -character variety that restricts to the identity on the diagonal and to Bers' simultaneous uniformization on  $\mathcal{T}(S) \times \overline{\mathcal{T}(S)}$ . The open subset contains  $\text{Hit}_3(S) \times \overline{\mathcal{T}(S)}$  and  $\mathcal{T}(S) \times \overline{\text{Hit}_3(S)}$ , and the image includes the holonomies of  $SL(3, \mathbb{C})$ -opers.

The map is realized by associating pairs of Hitchin representations to immersions into  $\mathbb{C}^3$  that we call complex affine spheres, which are equivalent to certain conformal harmonic maps into  $SL(3, \mathbb{C})/SO(3, \mathbb{C})$  and to new objects called bi-Higgs bundles. Complex affine spheres are obtained by solving a second-order complex elliptic PDE that resembles both the Beltrami and Tzitzéica equations. To study this equation we establish analytic results that should be of independent interest.

## CONTENTS

1. Introduction	2
1.1. The main result	3
1.2. The space of complex affine spheres	4
1.3. Bi-Higgs bundles, opers, and future work	6
1.4. Analytic results	7
1.5. Outline of the paper	8
1.6. Related work	9
1.7. Acknowledgements	9
2. Preliminaries	9
2.1. Teichmüller space	9
2.2. Holomorphic Riemannian manifolds	10
2.3. Complex metrics	11
2.4. Bers' theorem and Bers metrics	13
3. Complex affine spheres	14
3.1. Complex affine immersions	15
3.2. Blaschke affine immersions	16
3.3. The fundamental theorem for complex affine immersions	18
3.4. Complex affine spheres	20
3.5. Motivating examples	24
4. Complex Lie derivatives and transport	25
4.1. Complex Lie derivatives	26
4.2. Transport through complex vector fields	28
4.3. Transport of real analytic tensors	31
5. Complex elliptic operators	34
5.1. The Laplacian of a complex metric	34

5.2. Interior elliptic estimates	35
5.3. Proof of Theorem F	38
5.4. Proof of Theorem G	42
6. The holomorphic extension of the Labourie-Loftin parametrization	45
6.1. The Labourie-Loftin parametrization	45
6.2. Global analysis and the Gauss equation	45
6.3. The space $\text{CAS}(S)$	48
6.4. The action by $\text{MCG}(S)$ and the $\mathbb{C}^*$ twistor action	51
6.5. Proof of Theorem A	52
6.6. Proof of Theorem C	56
7. Bi-Higgs bundles and opers	57
7.1. Complex harmonic maps	57
7.2. Complex bi-gradings	58
7.3. bi-Higgs bundles	60
7.4. Bi-Hitchin bi-Higgs bundles	62
7.5. From bi-Higgs bundles to complex affine spheres	64
7.6. Opers	66
Appendix A. On the Beltrami operator	68
A.1. Proof of Proposition 5.7	68
A.2. Holomorphic dependence	70
A.3. Proof of Proposition 2.9	73
References	74

## 1. INTRODUCTION

Given a closed oriented surface  $S$  of genus  $g \geq 2$ , let  $\mathcal{T}(S)$  be the Teichmüller space of marked complex structures on  $S$ . By the uniformization theorem, every point in  $\mathcal{T}(S)$  determines a unique conjugacy class of Fuchsian representations of  $\pi_1(S)$  into  $\text{PSL}(2, \mathbb{R})$ .

Embedding  $\text{PSL}(2, \mathbb{R})$  inside its complexification  $\text{PSL}(2, \mathbb{C})$ , any Fuchsian representation produces a representation to  $\text{PSL}(2, \mathbb{C})$  that preserves a round circle in  $\mathbb{CP}^1$ . If we deform such a representation inside the character variety  $\chi(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  in a controlled manner, the resulting representation is quasi-Fuchsian, which means it preserves a Jordan curve in  $\mathbb{CP}^1$ . Quasi-Fuchsian representations have a number of nice properties; for example, quasi-Fuchsian representations are discrete and faithful, and hence the quotient of the action of a quasi-Fuchsian representation on  $\mathbb{H}^3$  is a quasi-Fuchsian 3-manifold. Given a quasi-Fuchsian representation  $\rho$  preserving a Jordan curve  $\gamma$ ,  $\rho$  also preserves and acts properly discontinuously on the two connected domains  $\Omega_\rho^+, \Omega_\rho^- \subset \mathbb{CP}^1$  that are complementary to  $\gamma$ . The resulting quotients  $\Omega_\rho^+/\rho$  and  $\Omega_\rho^-/\rho$  represent points in  $\mathcal{T}(S)$  and  $\mathcal{T}(\bar{S})$  respectively, where  $\mathcal{T}(\bar{S})$  is the Teichmüller space of the oppositely oriented surface  $\bar{S}$ . On the other hand, Bers proved that one can associate a quasi-Fuchsian representation to any pair of oppositely oriented complex structures, in a way that reverses the process above. Moreover, if we denote by  $\mathcal{QF}(S) \subset \chi(\pi_1(S), \text{PSL}(2, \mathbb{C}))$  the open subset of quasi-Fuchsian representations, and give  $\mathcal{T}(S)$  its natural complex structure, then Bers' simultaneous uniformization theorem (Theorem 2.7 below) says that this construction descends to a biholomorphism

$$\mathcal{T}(S) \times \mathcal{T}(\bar{S}) \rightarrow \mathcal{QF}(S).$$

In another direction, for every  $n \geq 2$ , given a Fuchsian representation into  $PSL(2, \mathbb{R})$ , we can turn it into a representation to  $PSL(n, \mathbb{R})$  by post-composing with Kostant's principle embedding from  $PSL(2, \mathbb{R}) \rightarrow PSL(n, \mathbb{R})$  (see [Kos59] for details on this embedding). Any deformation of that representation inside  $\chi(\pi_1(S), PSL(n, \mathbb{R}))$  is called a Hitchin representation, and by definition such representations fill up a connected component  $\text{Hit}_n(S)$  of  $\chi(\pi_1(S), PSL(n, \mathbb{R}))$  called a Hitchin component. Introduced in Hitchin's seminal paper [Hit92], Hitchin representations are Anosov (hence discrete and faithful) [Lab06], and are central objects of study in higher Teichmüller theory.

For  $n$  odd,  $SL(n, \mathbb{R}) = PSL(n, \mathbb{R})$ , so for the rest of the paper we use  $SL(3, \cdot)$  over  $PSL(3, \cdot)$ . Let  $\mathcal{M}_3(S)$  be the holomorphic vector bundle over  $\mathcal{T}(S)$  whose fiber over a class of a Riemann surface is identified with its space of holomorphic cubic differentials, i.e., it consists of pairs  $[c, Q]$ , where  $c$  is a complex structure on  $S$  and  $Q$  is a holomorphic cubic differential on  $(S, c)$ . Labourie in [Lab07] and Loftin in [Lof01] independently constructed a mapping class group equivariant and real analytic diffeomorphism

$$\mathcal{L} : \mathcal{M}_3(S) \xrightarrow{\sim} \text{Hit}_3(S) .$$

Both authors build this map using hyperbolic affine spheres. Labourie also gave a way to see  $\mathcal{L}$  using minimal surfaces in symmetric spaces [Lab17], which generalizes to other higher Teichmüller spaces for Lie groups of rank 2 (see [CTT19]).

Similar to the case  $n = 2$ , we can deform the Hitchin locus inside  $\chi(\pi_1(S), SL(3, \mathbb{C}))$ , and moreover we can inquire to what extent a Bers' phenomenon might hold for  $SL(3, \mathbb{C})$ , when we give  $\text{Hit}_3(S)$  the complex structure induced by  $\mathcal{L}$ . That is, one could—perhaps optimistically—try to holomorphically identify points in  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$  with representations to  $SL(3, \mathbb{C})$ . In fact, embedding  $\mathcal{M}_3(S)$  as the “diagonal” inside  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$ , by analytic continuing power series, one can find neighbourhoods  $U$  of this diagonal and  $V$  of the Hitchin representations inside  $\chi(\pi_1(S), SL(3, \mathbb{C}))$  on which  $\mathcal{L}$  extends uniquely to a biholomorphism

$$\mathcal{L}_{\mathbb{C}} : U \rightarrow V .$$

In this paper, we introduce maps from surfaces to  $\mathbb{C}^3$ , called complex affine spheres, that provide a concrete and geometrically meaningful realization of  $\mathcal{L}_{\mathbb{C}}$ . We use our affine spheres to show that  $\mathcal{L}_{\mathbb{C}}$  extends surprisingly far into  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$ , well past the diagonal as shown in Theorem A below, providing a more substantial and interesting Bers' phenomenon for  $SL(3, \mathbb{C})$ . These results should have incarnations for more general higher Teichmüller spaces.

**1.1. The main result.** The Labourie-Loftin parametrization via hyperbolic affine spheres comes from the classical theory of affine differential geometry. See the book [NS94] for references. It follows, from work of Cheng-Yau in [CY77] and [CY86] on affine differential geometry, and Goldman in [Gol90] and Choi-Goldman in [CG93] on geometric structures, that every Hitchin representation to  $SL(3, \mathbb{R})$  is uniquely associated with an equivariant hyperbolic affine sphere. Labourie and Loftin proved that every equivariant hyperbolic affine sphere is determined by the data of a complex structure on  $S$  and an accompanying holomorphic cubic differential (see also Wang [Wan91]), determining the map  $\mathcal{L}$  above.

We define (positive hyperbolic) complex affine spheres to be immersions of surfaces inside  $\mathbb{C}^3$  satisfying a geometric condition and structural equation analogous to that of the real affine spheres in  $\mathbb{R}^3$  (see Definition 3.14 and the subsection below). We show in Corollary 3.22 that a complex affine sphere is equivalent to a pair of complex structures on  $S$  with opposite orientations (not necessarily conjugate), holomorphic cubic differentials for the two complex structures, and a solution to an equation that depends on the complex structures

and differentials. Through this formalism, we explicitly construct and study  $\mathcal{L}_{\mathbb{C}}$ . Denote the mapping class group by  $\text{MCG}(S)$ . The following is proved in Section 6.

**Theorem A.** *The Labourie-Loftin parametrization extends uniquely to a  $\text{MCG}(S)$ -equivariant holomorphic map*

$$\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S}) \supset \Omega \xrightarrow{\mathcal{L}_{\mathbb{C}}} \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$$

where  $\Omega$  is a  $\text{MCG}(S)$ -invariant open subset containing the diagonal,  $\mathcal{M}_3(S) \times \mathcal{T}(\overline{S})$ , and  $\mathcal{T}(S) \times \mathcal{M}_3(\overline{S})$ . The image consists of holonomies of complex affine spheres. Moreover,  $\mathcal{L}_{\mathbb{C}}$  agrees with Bers' simultaneous uniformization map on  $\mathcal{T}(S) \times \mathcal{T}(\overline{S})$ .

Above, we are using  $\mathcal{T}(S)$  to denote the embedding of  $\mathcal{T}(S)$  into  $\mathcal{M}_3(S)$  as the zero section. The fact that  $\Omega$  contains  $\mathcal{M}_3(S) \times \mathcal{T}(\overline{S})$  and  $\mathcal{T}(\overline{S}) \times \mathcal{M}_3(\overline{S})$  is very interesting to us because, a priori,  $\mathcal{L}_{\mathbb{C}}$  is not expected to extend this far from the diagonal. In fact,  $\Omega$  can be constructed to be invariant under what we call the  $\mathbb{C}^*$  twistor action, defined by  $\zeta \cdot ([c_1, Q_1], [\overline{c_2}, \overline{Q_2}]) = ([c_1, \zeta Q_1], [\overline{c_2}, \zeta^{-1} \overline{Q_2}])$ , which also shows that  $\Omega$  is large. Note we need to be precise when talking about holomorphicity around the character variety (see Remarks 6.4 and 7.16).

We partially address what  $\mathcal{L}_{\mathbb{C}}$  does on  $\mathcal{M}_3(S) \times \mathcal{T}(\overline{S})$  and its partner in Theorem E below. See Section 1.3 for further comments.

We do not know whether the representations in the image of  $\mathcal{L}_{\mathbb{C}}$  are Anosov. Note that real affine spheres descend to maps from  $\tilde{S} \rightarrow \mathbb{RP}^2$ , which then pull back convex projective structures to  $S$ . In light of this point, it would be interesting to understand if one can use complex affine spheres to see geometric structures.

**1.2. The space of complex affine spheres.** We prove Theorem A by constructing a finite dimensional moduli space of complex affine spheres through which  $\mathcal{L}_{\mathbb{C}}$  factors.

The first ingredient in the construction is the notion of complex metrics on  $S$  (discussed in detail in Section 2). A complex metric on  $S$  is a non-degenerate bilinear symmetric form on the complexified tangent bundle  $\mathbb{C}TS$ , and it is called positive if it satisfies a further non-degeneracy condition (see Definition 2.4). Given a positive complex metric  $g$ , one can associate two complex structures  $c_1$  and  $\overline{c_2}$  to  $S$  of opposite orientations, and one can always locally find coordinates  $z$  for  $c_1$  and  $\overline{w}$  for  $\overline{c_2}$ , and a function  $\lambda$  to  $\mathbb{C}^*$  such that

$$g = \lambda dz d\overline{w}.$$

In [BE22], the authors formulate a covariant differential calculus for positive complex metrics, analogous to that of Riemannian metrics, in which we have Levi-Civita connections, curvature, Laplacians, and other usual objects. Finally, for every pair of oppositely oriented complex structures  $(c_1, \overline{c_2})$ , one can find a special positive complex metric  $\mathbf{h}_{(c_1, \overline{c_2})}$  of constant curvature  $-1$  in that conformal class, which is called the Bers metric, and every positive complex metric is conformally equivalent to a Bers metric.

Toward building  $\mathcal{L}_{\mathbb{C}}$ , in Section 3 we study (positive hyperbolic) complex affine spheres that can be deformed to real affine spheres. Complex affine spheres carry a number of tensors that generalize those from the real theory, such as (complex) Blaschke metrics and Pick tensors, satisfying Gauss and Codazzi-type equations. If the Blaschke metric has conformal class  $(c_1, \overline{c_2})$ , then the Codazzi equation is equivalent to the fact that the Pick tensor is of the form  $C = Q_1 + \overline{Q_2}$ , with  $Q_1$  and  $\overline{Q_2}$  holomorphic cubic differentials for  $c_1$  and  $\overline{c_2}$  respectively (see Theorem 3.20). For a distinctive subset of complex affine spheres, including those deformable to the real locus, solving the Gauss equation corresponds to

finding complex solutions  $u$  to the equation

$$G(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) := \Delta_h u - e^{2u} + \frac{1}{4}h(Q_1, \bar{Q}_2) \cdot e^{-4u} + 1. \quad (1)$$

where  $h = \mathbf{h}_{(c_1, \bar{c}_2)}$ . Note that for a “diagonal” point  $(c, \bar{c}, Q, \bar{Q})$ , Equation (1) reduces to the ordinary Tzitzéica equation for real hyperbolic affine spheres, as studied by Labourie and Loftin.

Let  $\mathcal{C}(S)$  be the space of complex structures on  $S$ , let  $\mathcal{CD}(S)$  be the bundle of cubic differentials over  $\mathcal{C}(S)$ , and let  $\mathcal{C}(\bar{S})$  and  $\mathcal{CD}(\bar{S})$  be the analogous spaces for  $\bar{S}$ . The product  $\mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$  is a complex Banach manifold, and when we take the quotient by  $\text{Diff}_0(S) \times \text{Diff}_0(\bar{S})$ , where  $\text{Diff}_0(S)$  is the space of diffeomorphisms of  $S$  isotopic to the identity, we get  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$ . Our distinctive subset of affine spheres is parametrized by

$$\text{CAS}(S) = \{(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) \in \mathcal{CD}(S) \times \mathcal{CD}(\bar{S}) \times C^\infty(S, \mathbb{C}) : G(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) = 0\}.$$

We say that a complex affine sphere  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u)$  is infinitesimally rigid if linearization of  $G$  in the direction of  $u$  is an isomorphism. We denote the space of infinitesimally rigid complex affine spheres by  $\text{CAS}^*(S)$ . The space  $\text{CAS}^*(S)$  is a complex Fréchet manifold for which the natural projection  $\hat{\pi}: \text{CAS}^*(S) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$  is a local biholomorphism and the holonomy map  $hol: \text{CAS}^*(S) \rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$  is holomorphic (see Section 6.2).

An important issue concerning the spaces  $\text{CAS}(S)$  and  $\text{CAS}^*(S)$ , which stems from a general issue around complex-metric related objects, is that their quotients under the diagonal action of  $\text{Diff}_0(S)$  are infinite-dimensional. To contend with this matter, in Section 4 we introduce notions of complex Lie derivative and transport through paths of complex vector fields, which allow to construct local deformations through “double isotopies” (also see the introduction to Section 4). Defining the transport amounts to solving a Cauchy problem (see Equation (18)), which is analytically not so well-behaved. Precisely, the Cauchy problem fails Hörmander’s criterion for local solvability (see Section 4.2).

Using complex Lie derivatives and transport, we manage to take a finite-dimensional quotient  $\mathbb{C}\text{AS}(S)$  of the space  $\text{CAS}^*(S)$  for which the fibers roughly correspond to double isotopies of the cubic differentials. The space  $\mathbb{C}\text{AS}(S)$  keeps track of the geometric information of the complex affine spheres, as summarized in the following theorem (corresponding to the statements in Section 6.3 of the text).

**Theorem B.** *The space  $\mathbb{C}\text{AS}(S)$  admits the structure of a finite dimensional complex manifold for which the quotient  $\text{CAS}^*(S) \rightarrow \mathbb{C}\text{AS}(S)$  is holomorphic. Moreover,  $hol$  descends holomorphically to a map  $hol: \mathbb{C}\text{AS}(S) \rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$  and  $\hat{\pi}$  descends to a local biholomorphism  $\pi: \mathbb{C}\text{AS}(S) \rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$ .*

As we show in Section 6, the map  $\mathcal{L}_\mathbb{C}$  in Theorem 1.1 is the composition of  $hol$  with a local inverse of  $\pi$ .

A natural question is whether  $\mathcal{L}_\mathbb{C}$  admits a global extension. The following statement provides evidence towards a negative answer.

**Theorem C.** *There is no holomorphic map*

$$\mathcal{CD}(S) \times \mathcal{CD}(\bar{S}) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\bar{S}) \times C^\infty(S, \mathbb{C})$$

*with image in  $\text{CAS}(S)$  that inverts  $\hat{\pi}$  and that coincides with the real affine spheres parametrization on the real locus.*

In particular,  $\hat{\pi}: \text{CAS}^*(S) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$  has no globally defined right inverse extending the real affine spheres parametrization.

For input data  $(c, \bar{c}, Q, -\bar{Q})$ , Equation (1) is exactly the structural equation for minimal Lagrangians in  $\mathbb{C}\mathbb{H}^2$ . Theorem C is a consequence of the fact that for  $(c, \bar{c}, Q, -\bar{Q})$ , as  $Q$  grows larger the linearization of Equation (1) eventually develops a kernel, which was proved by Huang-Loftin-Lucia in their paper about minimal Lagrangians [HLL13]. More discussion in Section 6.6.

**1.3. Bi-Higgs bundles, opers, and future work.** Labourie and Baraglia have shown that real hyperbolic affine spheres are equivalent to certain minimal surfaces in the  $SL(3, \mathbb{R})$  Riemannian symmetric space (see [Lab07, Section 9] and [Bar10, Section 3.4]). Moreover, if an affine sphere  $\sigma$  is equivalent to a minimal surface  $f$ , then the Higgs bundle of  $f$  encodes the Pick tensor of  $\sigma$ . Naturally, one might expect complex affine spheres to be equivalent to minimal surfaces in some  $SL(3, \mathbb{C})$ -homogeneous space.

While working on this paper, by first studying complex affine spheres, we came to realize a generalization of the theory of harmonic maps, minimal surfaces, and Higgs bundles. Specifically, harmonic maps into holomorphic Riemannian symmetric spaces such as  $SL(n, \mathbb{C})/SO(n, \mathbb{C}) =: \mathbb{Y}_n$  give rise to new objects that we call bi-Higgs bundles. In Section 7.3, we define a class of harmonic maps into  $\mathbb{Y}_3$ , which we call bi-Hitchin conformal harmonic maps, which are equivalent to certain bi-Higgs bundles, and which come along with a pair of cubic differentials. We prove the following in Sections 7.4 and 7.5. See Section 7 for more definitions and precise details.

**Theorem D.** *There is a one-to-one correspondence, up to conjugation, between equivariant positive hyperbolic complex affine spheres and bi-Hitchin conformal harmonic maps to  $\mathbb{Y}_3$ , which preserves the cubic differentials. When the affine sphere is deformable to a real affine sphere within the infinitesimally rigid locus, the holonomies are the same.*

Regarding the holonomies, one can certainly remove the additional assumption, which we keep to simplify the exposition.

A complete development of the theory of bi-Higgs bundles would take us too far afield from the aim of this paper, so we have chosen to defer a more extended treatment to an upcoming paper. As well, writing this paper from the perspective of complex affine spheres has its advantages: firstly, complex affine spheres might shed more light on the geometry of representations to  $SL(3, \mathbb{C})$ , and also, Equation (1), our main tool in proving Theorem A, appears to arise more naturally using complex affine spheres (see the end of Section 7.3).

Using bi-Higgs bundles, we also observe that a certain subset of  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$  corresponds to meaningful objects: opers.  $SL(n, \mathbb{C})$ -opers are gauge theoretic objects that arise from globalizing differential equations on Riemann surfaces (the full definition is in Section 7.6), and they are equivalent to  $\mathbb{C}\mathbb{P}^1$ -structures when  $n = 2$ . The modern theory of opers stems from Drinfeld-Sokolov [DS84] and the work of Beilinson-Drinfeld on the Geometric Langlands Conjecture [BD96]. In the latter work, Beilinson-Drinfeld showed that, over a Riemann surface  $(S, c)$  with canonical bundle  $\mathcal{K}$ , every connected component of the moduli space of  $SL(n, \mathbb{C})$ -opers is parametrized by the Hitchin base  $\oplus_{i=2}^n H^0(S, \mathcal{K}^i)$ .

**Theorem E.** *Given  $[c, S] \in \mathcal{M}_3(S)$ ,  $\mathcal{L}_{\mathbb{C}}([c, Q], [\bar{c}, 0])$  is represented by the holonomy of an  $SL(3, \mathbb{C})$ -oper on  $(S, c)$  with Hitchin basepoint  $(0, Q)$ , and  $\mathcal{L}_{\mathbb{C}}([c, 0], [\bar{c}, \bar{Q}])$  is represented by the holonomy of an oper on  $(S, \bar{c})$  with Hitchin basepoint  $(0, \bar{Q})$ .*

Since Bers' simultaneous uniformization procedure goes through constructing  $\mathbb{C}\mathbb{P}^1$ -structures with quasi-Fuchsian holonomies, it feels quite natural that opers appear in the image of  $\mathcal{L}_{\mathbb{C}}$ . As well, from our understanding, it's been an open question whether opers have a geometric interpretation; Theorems A and E seem to answer that question positively in a special case.

The relations between complex harmonic maps and opers seem worthy of further study. As we mentioned above, we have not addressed the general case of what  $\mathcal{L}_{\mathbb{C}}$  does on  $\mathcal{M}_3(S) \times \mathcal{T}(\bar{S})$  and  $\mathcal{T}(S) \times \mathcal{M}_3(\bar{S})$ . These loci suggest a notion of “bi-opers,” which we plan to develop in future work. As well, it’s worth pointing out that the  $\mathbb{C}^*$  twistor action is really an artifact of a well-known  $\mathbb{C}^*$ -action from non-abelian Hodge theory. In [Gai14], Gaiotto conjectured, motivated by physics, that the Hitchin component and the space of opers are holomorphically identified by taking a “conformal limit” through this  $\mathbb{C}^*$ -action. This conjecture was proved for closed surfaces in [Dum+21a] (see also [CW19]). Thus, bi-Higgs bundles seem to provide another perspective on Gaiotto’s conjecture and conformal limits, which we’re eager to explore. We discuss this topic in more detail in Section 7.6.

**1.4. Analytic results.** As indicated above, proving Theorem A requires studying Equation (1). In Section 5, we compute in local coordinates that if  $g = \lambda dz d\bar{w}$ , and  $\mu$  is the Beltrami form of  $w$  as a function of  $z$ , i.e.,  $\partial_{\bar{z}} w = \mu \partial_z w$ , then the Laplace operator of  $g$  is

$$\Delta_g = \frac{1}{\lambda \partial_{\bar{z}} w} \partial_{\bar{z}} (\partial_z - \bar{\mu} \partial_{\bar{z}}). \quad (2)$$

Note that the factor  $\partial_z - \bar{\mu} \partial_{\bar{z}}$  is the complex conjugation of the Beltrami operator. Since  $|\mu| < 1$ ,  $\Delta_g$  is a second order complex elliptic operator, i.e., it’s principle symbol never vanishes. The general class of such PDE’s is not so well studied (see [AIM09, §7.8] and [Beg94, §5] for references), and hence we have to develop some general results ourselves. We record our two most substantial and important results as theorems below.

Drawing on the Ahlfors-Bers analysis for the Beltrami operator, we prove interior elliptic estimates for complex operators such as  $\Delta_g$  (Proposition 5.5), from which we deduce Fredholm properties (Proposition 5.9). Our first main analytic theorem shows that complex affine spheres with real analytic input data are real analytic. Using this theorem, we’re able to use our transport method, defined in Section 4, on complex affine spheres in the proofs of Theorems A and B (we also use it in the proof of Theorem G below).

**Theorem F.** *Assume that  $g$  is a real analytic positive complex metric on  $U \subset \mathbb{C}$ , let  $V \subset \mathbb{C} \times \mathbb{C}^2$  be an open domain, let  $F : U \times V \rightarrow \mathbb{C}$  be a real analytic function, and let  $L$  be a semi-linear second order complex elliptic operator of the form*

$$Lu(z) = \Delta_g u + F(z, u, Du).$$

*If a function  $u \in C^2(U, \mathbb{C})$  with  $(u, Du)(U) \subset V$  satisfies  $Lu = 0$ , then  $u$  is real analytic.*

We prove Theorem F by adapting a proof of the analogous result for real elliptic PDE’s, which is attributed to Kato [Kat96], Hashimoto [Has06], and Blatt [Bla20]. There are some genuine differences in our complex setting (see Remark 5.11), but still the idea pushes through.

Theorem G below is the key result in extending  $\Omega$  to  $\mathcal{M}_3(S) \times \mathcal{T}(\bar{S})$  and  $\mathcal{M}_3(\bar{S}) \times \mathcal{T}(S)$ : it is used to show that we have infinitesimally rigid complex affine spheres lying over this locus. For the theorem below, see the more precise statement in Theorem 5.23.

**Theorem G.** *The operator  $L_{\mathbf{h}_{(c_1, \bar{c}_2)}} : C^\infty(S, \mathbb{C}) \rightarrow C^\infty(S, \mathbb{C})$  defined by*

$$L_{\mathbf{h}_{(c_1, \bar{c}_2)}} u = \Delta_{\mathbf{h}_{(c_1, \bar{c}_2)}} u - 2u$$

*is invertible for an open and dense subset of elements  $(c_1, \bar{c}_2) \in \mathcal{C}(S) \times \mathcal{C}(\bar{S})$ , including all the pairs  $(c_1, \bar{c}_2)$  such that  $c_1$  and  $\bar{c}_2$  induce the same real analytic structure on  $S$ .*

Theorem G can be recast as saying that a dense class of Bers metrics are infinitesimally rigid in their conformal class. In [BE22], it is proved that Bers metrics form a connected

component of the space of positive complex metrics of constant curvature  $-1$ ; this result is consistent with that one.

The proof of Theorem G falls outside the standard analytic toolkit: we use a continuity argument, relying on the transport technique defined in Section 4 and on one of the main theorems proved in the first author's work on variations of positive complex metrics [Ema23, Theorem A], to connect pairs  $(c_1, \overline{c_2})$  inducing the same real analytic structure to pairs  $(c, \overline{c})$ , for which the result of Theorem G is elementary.

In fact, we believe that  $L_{h_{(c_1, \overline{c_2})}}$  in Theorem G is an isomorphism for all Bers metrics, but we didn't need to prove the result in this generality. While Theorem G shows that 2 is not in the spectrum, we do not know whether  $u \mapsto \Delta_h u - ku$  is an isomorphism for every  $k > 0$ . Studying the spectrum of  $\Delta_h$  further seems interesting.

**1.5. Outline of the paper.** Toward understanding Theorem A, on a first reading one might want to focus on Sections 2, 3, and 6, and take the more analytic results from Sections 4 and 5 (such as Theorems F and G) for granted. Theorems A, B, and C are proved in Section 6, while in Section 7, a lot of which can be read independently of Sections 3-6, we prove Theorem D, which we then use to prove Theorem E.

In Section 2 we give the preliminaries on Teichmüller theory and complex metrics, among other things. Section 3 contains the geometry that underpins the rest of the paper: we develop a general theory of complex affine immersions, define complex affine spheres, and investigate their basic properties. We also give a number of explicit examples of complex affine immersions and complex affine spheres. Key results include Theorem 3.20, which shows that every positive hyperbolic complex affine sphere determines a pair of complex structures and cubic differentials, and Corollary 3.22, which shows that if one can solve Equation (1) built out of that data, then one can recover a complex affine sphere.

In Section 4, a lot of which can be read independently of the rest of the paper, we develop the notion of complex Lie derivatives and transport along complex vector fields. The main takeaways toward our main theorems are Theorem 4.11 and Proposition 4.13, which are used in the proof of Theorem F, and then eventually used to show in the proofs of Theorems A and C that local double isotopy deformations preserve the holonomy classes of infinitesimally rigid complex affine spheres.

In Section 5, which is independent of the theory of affine spheres, we carry out our general analysis of complex elliptic operators. We first derive our expression for  $\Delta_g$ , prove interior elliptic estimates, and Fredholm properties. We then prove Theorem F and Theorem G.

With the essential tools developed, in Section 6 we return to complex affine spheres. Here we formalize the strategy outlined in Section 1.2, construct the moduli space of infinitesimally rigid complex affine spheres (thus proving Theorem B), and then give the proof of Theorem A. We then prove Theorem C in Section 6.6.

As indicated above, in Section 7, we define harmonic maps to holomorphic Riemannian manifolds, show how bi-Higgs bundles arise out of them, and prove Theorem D. After giving a brief introduction toopers in Section 7.6, we use Theorem D to prove Theorem E.

To prove the interior elliptic interior estimates for  $\Delta_g$  in Section 5, and to show that the map  $\mathcal{L}_C$  is holomorphic in Section 6, we need a number of results on the Beltrami equation that should be known to experts but which we couldn't find in the literature. So we prove these results in an appendix. Theorem A.5 should be independently valuable: extending a classical result of Ahlfors and Bers [AB60, Theorem 11], Theorem A.5 is the statement that the Riemann Mapping Theorem solutions to the Beltrami equation vary holomorphically in higher Sobolev spaces.



**1.6. Related work.** There have been a number of works related to representations of surface groups into complex Lie groups close to the Hitchin locus, although the perspectives of all of the works so far are distinct from ours. See for example [DS20], [DS21], [ADL21], [Ale+23], [Dav23], and in particular [DS21], where Dumas-Sanders find a local continuation of Bers' theorem in a setting that falls more in line with the theory of Anosov representations.

We could frame the first half of Section 3 as the formulation of a basic theory of affine real submanifolds of dimension  $n$  inside  $\mathbb{C}^{n+1}$ , which then leads to the definition of complex affine spheres. We'd like to point out that there is an established theory of complex affine differential geometry, but centered around complex hypersurfaces in  $\mathbb{C}^{n+1}$ , developed in [Abe91] and [DVV88] among other sources.

There might be some interesting relations between bi-Higgs bundles and Hodge's thesis [Hod06], in which Hodge places some results about quasi-Fuchsian representations in a Higgs bundle context, and also with the latter work [LS20].

After writing this paper, we were notified about the paper [Kim07], in which Kim studies a complex extension of the Riemannian Laplacian, which seems to agree with our Bers Laplacians, and studies their determinants. Our analytic results, such as elliptic estimates and Fredholmness for our Laplacians, Theorem F, and especially Theorem G, can be seen as contributing to the study initiated in [Kim07]. We also mention that Kim proves that the complex Laplacians vary holomorphically in a  $C^2$  sense, but for our aims, we still need to prove (via Theorem A.5) that our operators vary holomorphically in our specific Sobolev spaces.

Around the end of writing this paper, we learned of independent work of Rungi and Tamburelli [RT24], who introduce complex minimal Lagrangian surfaces in the bi-complex hyperbolic space, which generalize minimal Lagrangian surfaces in  $\mathbb{C}\mathbb{H}^2$ , hyperbolic affine spheres in  $\mathbb{R}^3$ , and Bers embeddings in the holomorphic space form  $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \setminus \Delta$ . In [RT24], the authors define what they call  $SL(3, \mathbb{C})$ -bi-complex Higgs bundles, and they give a parameterization of a subset of  $SL(3, \mathbb{C})$ -Anosov representations by an open set in  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$ , from which they deduce that this subset of representations is endowed with a bi-complex structure. The approach of their work seems to give a different point of view and it would be interesting to analyze the interplay further. Nevertheless, we underline that the two works are completely independent.

**1.7. Acknowledgements.** We thank Kari Astala, Simon Blatt, and Peter Smillie for helpful discussions and correspondence. N.S. thanks the organizers and participants of the conference *Advances in Higgs Bundles*, which took place at the Brin Mathematics Research Center at the University of Maryland in April 2024, for conversations and sharing references from which this work benefited. We also thank the Geometry and Topology group in Luxembourg for the support and for creating such a nice working environment.

Both authors are funded by the FNR grant O20/14766753, *Convex Surfaces in Hyperbolic Geometry*.

## 2. PRELIMINARIES

For the rest of the paper, let  $S$  be a closed oriented surface of genus  $g \geq 2$ , and let  $\bar{S}$  be that same surface with the opposite orientation. Throughout the paper, we use  $W^{k,p}(\cdot)$ ,  $k \geq 1$ ,  $1 \leq p \leq \infty$ , for Sobolev spaces. As we will see, the analysis of the equations that we will eventually encounter require precise choices of Sobolev spaces.

**2.1. Teichmüller space.** Fix a basepoint complex structure on  $S$  that's compatible with the orientation, say  $c_0$ , and let  $\mathcal{K}$  be the canonical bundle. A **Beltrami form**  $\mu$  on  $S$  is an

$L^\infty$ -measurable section of the bundle  $\mathcal{K}^* \otimes \mathcal{K}^{-1}$  with  $L^\infty$  norm strictly less than 1. Recall that marked complex structures are parametrized by Beltrami forms. Explicitly, given a Beltrami form  $\mu$  on  $(S, c_0)$ , the Measurable Riemann Mapping Theorem produces a unique pair consisting of a compatible complex structure  $c_\mu$  together with a quasiconformal map  $f_\mu : (S, c_0) \rightarrow (S, c_\mu)$  (see Section A.3 for further comments). We say that a complex structure is in  $W^{k,p}$  if the Beltrami form is  $W^{k,p}$ . For a positive integer  $l > 1$ , denote by  $\mathcal{C}^l(S)$  the space of  $W^{l,\infty}$  complex structures on  $S$  that are compatible with the orientation. We write  $\mathcal{C}(S)$  for the  $C^\infty$  complex structures. Let  $\text{Diff}_+(S)$  be the space of orientation preserving diffeomorphisms of  $S$  and let  $\text{Diff}_0(S)$  be the normal subgroup of diffeomorphisms isotopic to the identity. The quotient  $\text{Diff}_+(S)/\text{Diff}_0(S)$  is the mapping class group  $\text{MCG}(S)$ .

**Definition 2.1.** The **Teichmüller space** of marked complex structures on  $S$  is the quotient  $\mathcal{T}(S) = \mathcal{C}(S)/\text{Diff}_0(S)$ .

For  $l < \infty$ ,  $\mathcal{C}^l(S)$  is a complex Banach manifold modelled on the space of  $W^{l,\infty}$  Beltrami forms, and for  $l = \infty$ ,  $\mathcal{C}(S)$  is a complex Fréchet manifold, with semi-norms coming from the  $W^{l,\infty}$  norms (by Sobolev embedding theorems, this Fréchet structure is biholomorphic to all of the more common ones). The following is due to Earle-Eells. [EE69].

**Theorem 2.2** (Earle-Eells). *Under the quotient topology,  $\mathcal{T}(S)$  inherits a complex structure with respect to which the map from  $\mathcal{C}(S) \rightarrow \mathcal{T}(S)$  is holomorphic and a principle  $\text{Diff}_0(S)$ -fiber bundle.*

Throughout the paper, we insist on viewing  $\mathcal{C}(S)$  as a Fréchet manifold, so that we can directly apply the result of Earle-Eells.

Note that the complex structure on the Teichmüller space  $\mathcal{T}(\bar{S})$  of the oppositely oriented surface  $\bar{S}$ , identifies with the complex structure conjugate to the usual one on  $\mathcal{T}(S)$ .

**2.2. Holomorphic Riemannian manifolds.** Let  $\mathbb{X}$  be a complex manifold with almost complex structure  $\mathbb{J}$ . A **holomorphic Riemannian metric**  $\langle \cdot, \cdot \rangle$  on  $\mathbb{X}$  is a holomorphic assignment of a symmetric non-degenerate  $\mathbb{C}$ -bilinear form on each fiber of the holomorphic tangent bundle of  $\mathbb{X}$ , or more concisely a holomorphic nowhere degenerate section of  $\text{Sym}_2(T^{1,0}(\mathbb{X}))$ . These objects have been widely studied in literature with different approaches, see for instance [DZ09].

A holomorphic affine connection  $D$  on  $T\mathbb{X}$  is an affine connection such that  $D\mathbb{J} = 0$  and for local holomorphic vector fields  $Z_1, Z_2$  one has  $D_{\mathbb{J}Z_1}Z_2 = \mathbb{J}(D_{Z_1}Z_2)$ . Holomorphic Riemannian manifolds  $(\mathbb{X}, \langle \cdot, \cdot \rangle)$  have a natural notion of **Levi-Civita connection**  $D$ , which is a holomorphic affine connection on  $T\mathbb{X}$  that makes the metric parallel. Such a connection induces a natural notion of  $\mathbb{C}$ -multilinear curvature tensor and, for all complex vector subspace  $V < T_p\mathbb{X}$  such that  $\dim_{\mathbb{C}}V = 2$  and that  $\langle \cdot, \cdot \rangle|_V$  is non-degenerate, one can define the sectional curvature.

Given a complex semisimple Lie group, its complex Killing form extends to the whole tangent bundle by left or right translation and hence determines a holomorphic Riemannian metric  $\langle \cdot, \cdot \rangle$ . For  $G = \text{SL}(n, \mathbb{C})$ , at  $A \in \text{SL}(n, \mathbb{C})$ , the metric is given by, for  $V, W \in T_A\text{SL}(n, \mathbb{C})$ ,  $\langle V, W \rangle|_A = 2n\text{tr}(A^{-1}VA^{-1}W)$ . We highlight two examples that arise from taking quotients of complex Lie groups.

- (1)  $\mathbb{C}^n$  with the canonical symmetric  $\mathbb{C}$ -bilinear form given by  $\langle \underline{z}, \underline{w} \rangle = \sum_{k=1}^n z_k w_k$ , for  $\underline{z} = (z_1, \dots, z_n), \underline{w} = (w_1, \dots, w_n) \in \mathbb{C}^n$ .  $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$  is the unique simply connected complete holomorphic Riemannian metric with constant sectional curvature zero [BE22, Theorem 2.6].

(2) The hyperboloid

$$\mathbb{X}_n = \{\underline{z} \in \mathbb{C}^{n+1} \mid \langle \underline{z}, \underline{z} \rangle = -1\}$$

is a complex submanifold of  $\mathbb{C}^n$  and it inherits a holomorphic Riemannian metric, under which it identifies with  $SO(n+1, \mathbb{C})/SO(n, \mathbb{C})$ . For  $n \geq 2$ ,  $\mathbb{X}_n$  is the unique complete simply connected holomorphic Riemannian manifold with constant sectional curvature -1 [BE22, Theorem 2.6]. Moreover, for all  $p, q$  with  $p+q = n$ ,  $\mathbb{H}^{p,q} = SO(p, q)/SO(p) \times SO(q)$  embeds isometrically in  $\mathbb{X}_n$  in a unique way up to composition with ambient isometries.

Observe that, since all non-degenerate bilinear forms on  $\mathbb{C}^n$  are isomorphic, one can define several isometric models for  $\mathbb{C}^{n+1}$  and  $\mathbb{X}_n$  by changing the ambient bilinear form. For instance, in Example 3.24, we will refer to the isometric model for  $(\mathbb{C}^{n+1}, \langle \cdot, \cdot \rangle)$  given by  $\mathbb{C}^{n,1}$ , namely  $\mathbb{C}^{n+1}$  endowed with the bilinear form  $\langle \underline{z}, \underline{w} \rangle_{n,1} = -z_{n+1}w_{n+1} + \sum_{k=1}^n z_k w_k$ . Via the isometry between these two spaces one gets that  $\mathbb{X}_n$  is equivalent to  $(\hat{\mathbb{X}}_n, \langle \cdot, \cdot \rangle_{n,1})$  with

$$\hat{\mathbb{X}}_n = \{\underline{z} \in \mathbb{C}^{n+1} \mid \langle \underline{z}, \underline{z} \rangle_{n,1} = -1\}. \quad (3)$$

Finally,  $\mathbb{X}_2$  models the space of oriented geodesics of hyperbolic space  $\mathbb{H}^3$ , which will be relevant below. Indeed, identifying  $SO(3, \mathbb{C})$  with  $PSL(2, \mathbb{C})$  and viewing  $\mathbb{X}_2$  as  $PSL(2, \mathbb{C})/SO(2, \mathbb{C})$ ,  $SL(2, \mathbb{C})$  acts transitively on the space of oriented geodesics, and the stabilizer of any oriented geodesic is conjugate to  $SO(2, \mathbb{C})$ . By identifying a geodesic in  $\mathbb{H}^3$  with its endpoints in  $\mathbb{C}\mathbb{P}^1$ , we obtain a biholomorphism from

$$\mathbb{X}_2 \rightarrow \mathbb{G} := \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \setminus \Delta.$$

Under this biholomorphism, the pushed-forward holomorphic Riemannian metric has the following description: given any complex affine chart  $(U, z)$  on  $\mathbb{C}\mathbb{P}^1$ , the metric on  $(U \times U \setminus \Delta, (z_1, z_2))$  has the form

$$\langle \cdot, \cdot \rangle_{\mathbb{G}} = -\frac{4}{(z_1 - z_2)^2} dz_1 \cdot dz_2.$$

See [BE22, Section 2.4] for details. Several relations between immersions of surfaces into  $\mathbb{H}^3$  and  $\mathbb{G}$  are treated in [ES22].

**2.3. Complex metrics.** Complex metrics first appeared in a paper by the first author on immersions of (real) smooth manifolds into holomorphic Riemannian manifolds [BE22]. Throughout the paper, given a smooth manifold  $M$ , we denote the complexified tangent bundle  $TM \otimes_{\mathbb{R}} \mathbb{C}$  by  $\mathbb{C}TM$ , and likewise for the complexified cotangent bundle  $\mathbb{C}T^*M$ .

**Definition 2.3.** A **complex metric** on a smooth (real) manifold  $M$  is a smooth section of  $\text{Sym}^2(\mathbb{C}TM)$  that determines a non-degenerate symmetric bilinear form in each fiber.

Of course, the bilinear extension of a Riemannian metric to the complexified tangent bundle is a complex metric. Throughout the paper, we will view a complex metric on a manifold  $M$  as the same thing as an invariant complex metric on the universal cover  $\widetilde{M}$ , and we won't distinguish notation.

A lot of the usual constructions in Riemannian geometry extend to the setting of complex metrics and the proofs apply verbatim. Here are a few objects and facts that we will use in this paper.

- (1) Every complex metric  $g$  has a natural **Levi-Civita connection**, namely an affine connection  $\nabla^g: \Gamma(\mathbb{C}TM) \rightarrow \Gamma(\text{End}(\mathbb{C}TM))$  uniquely determined by the fact that it is torsion-free and compatible with the metric, i.e.  $\nabla^g g = 0$ .

- (2) Through  $g$  and its Levi-Civita connection  $\nabla^g$ , one can define the (0, 4)-type and the (1, 3)-type **curvature tensors**, which we both denote by  $R^g$  when there is no ambiguity, as

$$R^g(X, Y, Z, W) = g(R^g(X, Y)Z, W) := g(\nabla_X^g \nabla_Y^g Z - \nabla_Y^g \nabla_X^g Z - \nabla_{[X, Y]}^g Z, W)$$

for all  $X, Y, Z, W \in \Gamma(\mathbb{C}TM)$ , which is  $\mathbb{C}$ -multilinear and has the usual symmetries as in the Riemannian case. If  $X$  and  $Y$  span a  $g$ -non-degenerate 2-plane in  $\mathbb{C}TM$ , we have the sectional curvature

$$K(X, Y) = \frac{R^g(X, Y, Y, X)}{g(X, X)g(Y, Y) - (g(X, Y))^2}.$$

When  $M$  is a surface, this sectional curvature is called the **Gauss curvature** and defines a function  $K_g : M \rightarrow \mathbb{C}$ .

- (3) For any  $C^1$  function  $f : M \rightarrow \mathbb{C}$ , one can define the **gradient** of  $f$  as the unique vector field  $\nabla^g f$  defined by  $df = g(\nabla^g f, \cdot)$ . If  $f$  is also  $C^2$ , we can define the **Hessian** and the **Laplacian** of  $f$  as

$$\begin{aligned} \text{Hess}_g(f)(X, Y) &= g(\nabla_X^g \nabla^g f, Y), \\ \Delta_g(f) &= \text{tr}_g(\text{Hess}_g(f)). \end{aligned}$$

- (4) Assume  $M$  is a surface, and let  $\varrho : M \rightarrow \mathbb{C}^*$ . Then, the conformal metric  $\hat{g} = \varrho \cdot g$  satisfies  $\Delta_{\hat{g}} = \frac{1}{\varrho} \Delta_g$  and

$$K_{\hat{g}} = \frac{1}{\varrho} (K_g - \frac{1}{2} \Delta_g \log(\varrho)). \quad (4)$$

While the function  $\log(\varrho)$  is only locally well-defined, up to choices,  $\Delta_g \log(\varrho)$  is globally well-defined.

In the following, we restrict to the case of surfaces  $M = S$ . The fact that a complex metric  $g$  on  $S$  is non-degenerate implies that  $g$  has two isotropic directions at each point. In other words, for each point  $p \in S$ , the set  $\{v \in \mathbb{C}T_p S \mid g_p(v, v) = 0\}$  consists of two complex lines of  $\mathbb{C}T_p S$ , corresponding to two points in  $\mathbb{P}\mathbb{C}T_p S$ . Observe that  $\mathbb{P}T_p S$  embeds into the sphere  $\mathbb{P}\mathbb{C}T_p S$  as an equatorial circle, which cuts  $\mathbb{P}\mathbb{C}T_p S$  into two connected components.

**Definition 2.4.** A complex metric  $g$  is **positive** if for each point  $p \in S$ , no isotropic direction of  $g_p$  lies in  $\mathbb{P}T_p S$ , and the two isotropic directions are contained in distinct components of  $\mathbb{P}\mathbb{C}T_p S \setminus \mathbb{P}T_p S$ .

Riemannian metrics complexify to positive complex metrics: the isotropic directions are the eigen-lines of the almost complex structure and are hence antipodal.

We denote the space of  $C^\infty$  – resp.  $W^{l,2}$  – positive complex metrics on  $S$  by  $\mathcal{CM}(S)$  – resp.  $\mathcal{CM}^l(S)$ . Being open subsets of the space of  $C^\infty$  – resp.  $W^{l,2}$  – sections of  $\text{Sym}^2(\mathbb{C}TS)$ , one can see that  $\mathcal{CM}(S)$  is a complex Fréchet manifold and  $\mathcal{CM}^l(S)$  is a complex Banach manifold.

We further define  $\mathcal{CM}_{-1}(S)$  and  $\mathcal{CM}_{-1}^l(S)$  to be the subsets of positive complex metrics of constant curvature  $-1$  in  $\mathcal{CM}(S)$  and  $\mathcal{CM}^l(S)$  respectively.

The isotropic directions of a positive complex metric determine a pair of oppositely oriented complex structures  $c_1$  and  $\bar{c}_2$  on  $S$ . In fact, if  $z$  and  $\bar{w}$  are local holomorphic coordinates for  $c_1$  and  $\bar{c}_2$  respectively, then  $g(\partial_{\bar{z}}, \partial_{\bar{z}}) = g(\partial_w, \partial_w) = 0$ , and  $g$  can be locally written as  $g = \lambda dz d\bar{w}$ , for some non-vanishing complex valued function  $\lambda$  (see [BE22, Section 6.1] for details).

This defines a projection map

$$(c_+, c_-): \mathcal{CM}(S) \rightarrow \mathcal{C}(S) \times \mathcal{C}(\overline{S})$$

whose fibers correspond to conformal classes of positive complex metrics.

The locally defined basis for  $\mathbb{C}TS$  given by  $\{\partial_w, \partial_{\overline{z}}\}$  will be particularly useful for computations, so we record a few elementary facts here. Let  $\nabla^g$  be the Levi-Civita connection for  $g$ , and let  $\Pi_1$  and  $\Pi_2$  be the projections of  $\mathbb{C}TS$  onto the isotropic directions of  $g$  for  $c_1$  and  $\overline{c_2}$  respectively, which are locally spanned by  $\partial_{\overline{z}}$  and  $\partial_w$  respectively.

**Lemma 2.5.** *For any locally defined complex vector field  $X$ ,  $\nabla_X^g \partial_{\overline{z}}$  is parallel to  $\partial_{\overline{z}}$ , and  $\nabla_X^g \partial_w$  is parallel to  $\partial_w$ .*

*Proof.* By metric compatibility,  $g(\nabla_X^g \partial_{\overline{z}}, \partial_{\overline{z}}) = \frac{1}{2} \nabla_X^g g(\partial_{\overline{z}}, \partial_{\overline{z}}) = 0$ , and likewise for  $\partial_w$ .  $\square$

**Lemma 2.6.**  $\nabla_{\overline{z}}^g \partial_w = \Pi_2([\partial_{\overline{z}}, \partial_w])$  and  $\nabla_w^g \partial_{\overline{z}} = \Pi_1([\partial_w, \partial_{\overline{z}}])$ .

*Proof.* Since  $\nabla^g$  is torsion free,  $\nabla_{\overline{z}}^g \partial_w - \nabla_w^g \partial_{\overline{z}} = [\partial_{\overline{z}}, \partial_w]$ . We apply Lemma 2.5.  $\square$

**2.4. Bers' theorem and Bers metrics.** In the introduction we defined quasi-Fuchsian representations. Bers' Simultaneous Uniformization Theorem is more formally stated below. If  $\rho$  is quasi-Fuchsian, preserving a Jordan curve  $\gamma \subset \mathbb{CP}^1$ , the domains complementary to  $\gamma$  are called the domains of discontinuity. Recall that, identifying  $\pi_1(S)$  with the group of covering transformations of  $\tilde{S} \rightarrow S$ , given an action  $\rho$  of  $\pi_1(S)$  on a space  $X$ , we say that  $f: \tilde{S} \rightarrow X$  is  $\rho$ -**equivariant** if for all  $\gamma \in \pi_1(S)$ ,  $f \circ \gamma = \rho(\gamma) \circ f$ .

**Theorem 2.7** (Bers' Simultaneous Uniformization Theorem [Ber60]). *For all  $(c_1, \overline{c_2}) \in \mathcal{C}^l(S) \times \mathcal{C}^l(\overline{S})$ , there exists a quasi-Fuchsian representation  $\rho: \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ , unique up to conjugation, with domains of discontinuity  $\Omega_\rho^+$  and  $\Omega_\rho^-$ , together with unique  $\rho$ -equivariant holomorphic diffeomorphisms*

$$\mathbf{f}_+(c_1, \overline{c_2}): (\tilde{S}, c_1) \rightarrow \Omega_\rho^+, \quad \overline{\mathbf{f}}_-(c_1, \overline{c_2}): (\tilde{S}, \overline{c_2}) \rightarrow \Omega_\rho^-.$$

*This correspondence determines a biholomorphism*

$$\mathcal{T}(S) \times \mathcal{T}(\overline{S}) \xrightarrow{\sim} \mathrm{QF}(S).$$

Fix a pair  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ . The two maps  $f_1 = \mathbf{f}_+(c_1, \overline{c_2})$  and  $\overline{f_2} = \overline{\mathbf{f}}_-(c_1, \overline{c_2})$  define an immersion  $(f_1, \overline{f_2}): \tilde{S} \rightarrow \mathbb{G} \supset \Omega_\rho^+ \times \Omega_\rho^-$ , and the pull-back of  $\langle \cdot, \cdot \rangle_{\mathbb{G}}$  is the positive complex metric

$$\mathbf{h}_{(c_1, \overline{c_2})} := -\frac{4}{(f_1 - \overline{f_2})^2} df_1 d\overline{f_2} \quad (5)$$

that, as a result of Theorem 6.7 in [BE22], has constant curvature  $-1$ . We refer to the metrics obtained in this fashion as **Bers metrics**. In [BE22], the first author with Bonsante proved that one can uniformize positive complex metrics via Bers metrics.

**Proposition 2.8** (Theorem 6.11 in [BE22]). *A complex metric  $g$  is positive if and only if there exists a (unique) Bers metric  $\mathbf{h}_{(c_1, \overline{c_2})}$  and a function  $\varrho: S \rightarrow \mathbb{C}^*$  such that*

$$g = \varrho \cdot \mathbf{h}_{(c_1, \overline{c_2})}.$$

As proved in [BE22], in the smooth category, Bers metrics form a connected component of the subset of smooth complex metrics of constant curvature  $-1$ , and  $(c_1, \overline{c_2}) \mapsto \mathbf{h}_{(c_1, \overline{c_2})}$  defines a bijective correspondence between pairs of complex structures and Bers metrics. If the complex structures are  $W^{l, \infty}$ , then the Bers metric, viewed as a metric on  $S$ , is a  $W^{l, 2}$

bilinear form on  $\mathbb{C}TS$ . Indeed, briefly, by Proposition 5.7 stated later on, the maps  $\mathbf{f}_+$  and  $\overline{\mathbf{f}}_-$  are  $W_{loc}^{l+1,p}$  for some small  $p > 2$ , hence  $W_{loc}^{l+1,2}$ , and routinely estimating the formula (5) using that  $W^{l,2}(U)$  is a Banach algebra for any  $U \subset \mathbb{C}$  (see Proposition 5.17), one sees that the Bers metric on  $\tilde{S}$  is  $W_{loc}^{l,2}$ . Descending to  $S$  gives the result. A few times in the paper, especially in Section 6, we'll need to know the following.

**Proposition 2.9.** *The map  $\mathcal{C}^l(S) \times \mathcal{C}^l(\overline{S}) \rightarrow \mathcal{CM}^l(S)$ , defined by  $(c_1, \overline{c_2}) \mapsto \mathbf{h}_{(c_1, \overline{c_2})}$ , with image in  $\mathcal{CM}_{-1}^l(S)$ , is holomorphic.*

From the definition of the Fréchet structures on  $\mathcal{C}(S) \times \mathcal{C}(\overline{S})$  and  $\mathcal{CM}(S)$ , it follows that the map  $(c_1, \overline{c_2}) \mapsto \mathbf{h}_{(c_1, \overline{c_2})}$  is holomorphic in these spaces as well. Proposition 2.9 follows from the analytic dependence of solutions to the Beltrami equation in higher Sobolev spaces; since we couldn't find a source for the results we need, we prove these results, as well as Proposition 2.9, in the Appendix A.

Finally, we record some results from [Ema23] on holomorphically perturbing positive complex metrics, which we will use in the proof of Theorem G in Section 5.3. For all  $c \in \mathcal{C}(S)$  (resp.  $\overline{c} \in \mathcal{C}(\overline{S})$ ), denote by  $\text{QD}(c)$  (resp.  $\text{QD}(\overline{c})$ ) the space of holomorphic quadratic differentials for  $c$  (resp.  $\overline{c}$ ). If  $g$  is a positive complex metric, then by openness of the positive property, for all  $q_1 \in \text{QD}(c)$  and  $\overline{q_2} \in \text{QD}(\overline{c})$  small enough,  $g + q_1$  and  $g + \overline{q_2}$  are positive metrics as well.

**Proposition 2.10.** *Let  $g$  be a positive complex metric, with  $(c_1, \overline{c_2}) = (\mathbf{c}_+, \mathbf{c}_-)(g)$ , and  $q_1 \in \text{QD}(c)$  and  $\overline{q_2} \in \text{QD}(\overline{c})$  so that  $g + q_1$  and  $g + \overline{q_2}$  are both positive. Then*

$$K_g = K_{g+q_1}, \quad \text{and} \quad \mathbf{c}_+(g + q_1) = c_1,$$

and

$$K_g = K_{g+\overline{q_2}}, \quad \text{and} \quad \mathbf{c}_-(g + \overline{q_2}) = \overline{c_2}.$$

This Proposition is stated as Proposition 3.8 in [Ema23] for Bers metrics, but the proof works verbatim for any positive complex metric. Moreover, in [Ema23] it is proved that deformations from  $h$  to  $h + q_1$  and  $h + \overline{q_2}$  descend to give complex charts on Teichmüller space.

**Theorem 2.11** (Theorems A and B in [Ema23]). *Let  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ ,  $h = \mathbf{h}_{(c_1, \overline{c_2})}$ , and let  $U_{(c_1, \overline{c_2})} \subset \text{QD}(c_1)$  be an open neighborhood of zero such that  $\mathbf{h}_{(c_1, \overline{c_2})} + q_1$  is a Bers metric. The map*

$$\begin{aligned} U &\rightarrow \mathcal{T}(\overline{S}) \\ q_1 &\mapsto [\mathbf{c}_-(h + q_1)] \end{aligned}$$

is a biholomorphism onto its image. The analogous statement holds for deformations of the form  $\mathbf{h}_{(c_1, \overline{c_2})} \mapsto \mathbf{h}_{(c_1, \overline{c_2})} + \overline{q_2}$ .

Although we won't use this fact in the paper, it is worthwhile to note that the charts from the theorem above are affine reparametrizations of the Schwarzian parametrization.

### 3. COMPLEX AFFINE SPHERES

In this section we develop the theory of complex affine immersions and complex affine spheres as an extension of the theory of (real) affine immersions and affine spheres. We don't recall the real theory here, but we highlight it as a special case of our construction (see Example 3.23, and also Section 6.1 later on). Sources on the real theory include [NS94], [Lof01], and [Lab07].

In the section below, if regularity is unspecified, we will assume that all the objects are regular enough for the operations to be well-defined.

**3.1. Complex affine immersions.** Let  $\mathbb{X}$  be a complex manifold of complex dimension  $n = m + 1$  with almost complex structure  $\mathbb{J}$  and torsion free holomorphic affine connection  $D$ . Starting in Section 3.4,  $\mathbb{X} = \mathbb{C}^3$ , equipped with the Levi-Civita connection of its holomorphic Riemannian metric, so it might be helpful to have  $\mathbb{X} = \mathbb{C}^n$  in mind as the typical target space. We denote by  $\text{Aff}(\mathbb{X}, D)$  the space of biholomorphisms of  $\mathbb{X}$  that preserve  $D$ .

Now, let  $M$  be a (real) smooth manifold of dimension  $m$ , let  $\widetilde{M}$  be its universal cover, and identify  $\pi_1(M)$  with the deck group of the covering map. From now on until Section 3.3, fix a representation  $\rho: \pi_1(M) \rightarrow \text{Aff}(\mathbb{X}, D)$  and let  $\sigma: \widetilde{M} \rightarrow \mathbb{X}$  be a equivariant map. We refer to  $\rho$  as the **holonomy** of  $\sigma$ . Note that one can see immersions  $M \rightarrow \mathbb{X}$  as equivariant immersions  $\widetilde{M} \rightarrow \mathbb{X}$  with the trivial holonomy  $\rho \equiv id_{\mathbb{X}}$ .

The tangent map  $\sigma_*: T\widetilde{M} \rightarrow \sigma^*(T\mathbb{X})$  extends to a  $\mathbb{C}$ -linear map  $\sigma_*: \mathbb{C}T\widetilde{M} \rightarrow \sigma^*(T\mathbb{X})$  by defining  $\sigma_*(v + iw) := \sigma_*(v) + \mathbb{J}(\sigma_*(w))$ . In the following, with  $\sigma_*$  we will always denote the extension to  $\mathbb{C}TM$ .

**Definition 3.1.** We say that an immersion  $\sigma: \widetilde{M} \rightarrow \mathbb{X}$  is **admissible** if the induced map  $\sigma_*: \mathbb{C}T\widetilde{M} \rightarrow \sigma^*T\mathbb{X}$  is injective.

Admissibility means that  $\sigma_*$  maps  $\mathbb{R}$ -linearly independent vectors in  $T_p\widetilde{M}$  to  $\mathbb{C}$ -linearly independent vectors in  $\sigma^*(T\mathbb{X})$ . The admissibility condition allows the construction of several geometric tensors associated with the immersion, imitating the theory of hypersurfaces in smooth manifolds.

Since  $\sigma$  is  $\rho$ -equivariant,  $\pi_1(M)$  acts on the bundle  $\sigma^*(T\mathbb{X})$  through  $\rho$  preserving both  $\sigma^*D$  and  $\sigma^*\mathbb{J}$ : the quotient defines a bundle  $E_\rho \rightarrow M$  that inherits both a connection, which we still denote with  $\sigma^*D$ , and a complex structure that we denote with  $i$  to relax the notation. Assuming  $\sigma$  is admissible,  $\mathbb{C}TM$  can be seen as a complex subbundle of  $E_\rho$  with complex co-rank 1. We abuse notation slightly and write this subbundle as  $\mathbb{C}TM \subset E_\rho$  when the context is clear.

**Definition 3.2.** A  $\rho$ -equivariant **complex affine immersion** of  $\widetilde{M}$  in  $\mathbb{X}$  is a pair  $(\sigma, \xi)$  where  $\sigma: \widetilde{M} \rightarrow \mathbb{X}$  is a  $\rho$ -equivariant admissible immersion, and  $\xi$  is a smooth section of  $E_\rho$  called the **transversal vector field**, such that  $\mathbb{C}T\widetilde{M} \oplus \text{Span}_{\mathbb{C}}(\xi) = E_\rho$ .

Compare with [NS94, Definition II.1.1].

A **complex affine connection**  $\nabla$  on  $M$  is a  $\mathbb{C}$ -bilinear affine connection on the bundle  $\mathbb{C}TM$ . Observe that all affine connections on  $TM$  can be extended by linearity to complex affine connections on  $\mathbb{C}TM$ . Notice that this notion, defined on smooth manifolds, is very different from the notion of holomorphic connection on a complex manifold introduced above.

According to the splitting  $\mathbb{C}TM \oplus \text{Span}_{\mathbb{C}}(\xi)$ , we get that for all  $X \in \Gamma(TM)$ ,  $Y \in \Gamma(\mathbb{C}TM)$

$$\begin{cases} (\sigma^*D)_X Y & =: \nabla_X Y + g(X, Y) \cdot \xi \\ (\sigma^*D)_X \xi & =: -S(X) + \tau(X) \cdot \xi. \end{cases} \quad (6)$$

**Remark 3.3.** One can see that:

- for all  $X, Y \in \Gamma(\mathbb{C}TM)$ , with  $X = X_1 + iX_2$ ,  $\nabla_X Y := \nabla_{X_1} Y + i\nabla_{X_2} Y$  is a torsion-free complex affine connection on  $\mathbb{C}TM$ ;
- $g$  extends to a symmetric  $\mathbb{C}$ -bilinear form on  $\mathbb{C}TM$ , which, by analogy with the theory of real affine spheres, we call the **affine fundamental form**;

- $\mathbb{S}$  is linear, and it extends to a section of  $\text{End}(\mathbb{C}TM)$ , called the **shape operator**;
- $\tau$  is linear as well, and it extends to an element in  $\Omega^1(\mathbb{C}TM)$ , called **transversal connection form**.

**3.2. Blaschke affine immersions.** In this subsection we construct preferred transversal vector fields called affine normals. The theory here is roughly parallel to [NS94, §II.3]. However, unlike the real case, we will see that there are topological obstructions to the existence of such an affine normal. First of all, let us see how the affine tensors defined in the previous subsection change according to a change in the transversal vector field.

**Lemma 3.4.** *Let  $(\sigma, \xi)$  be a complex affine immersion, with  $\nabla, g, \mathbb{S}, \tau$  as in Equation (6). Let  $\hat{\xi} = \alpha\xi + \eta$  be a different transversal vector field for  $\sigma$ , with  $\alpha \in C^\infty(M, \mathbb{C}^*)$  and  $\eta \in \Gamma(\mathbb{C}TM)$ . Then, the complex affine immersion  $(\sigma, \hat{\xi})$  induces  $\hat{\nabla}, \hat{g}, \hat{\mathbb{S}}, \hat{\tau}$  according to:*

- (1)  $\hat{g} = \frac{1}{\alpha}g$ ,
- (2)  $\hat{\nabla}_X Y = \nabla_X Y - \frac{1}{\alpha}g(X, Y)Z$ ,
- (3)  $\hat{\mathbb{S}}(X) = \alpha\mathbb{S}(X) - \nabla_X \eta + \tau(X)\eta + \frac{\partial_X \alpha + g(X, \eta)}{\alpha}\eta$ ,
- (4)  $\hat{\tau}(X) = \tau(X) + \frac{\partial_X \alpha + g(X, \eta)}{\alpha}$ .

*Proof.* For  $X, Y, \eta \in \Gamma(TM)$ , the proof follows by direct computation, as shown for the real case in [NS94, Proposition II.2.5]. The equations for  $X, Y \in \Gamma(\mathbb{C}TM)$  follow by  $\mathbb{C}$ -linearity, while the result for  $\eta \in \Gamma(\mathbb{C}TM)$  follows by applying transformations repeatedly, namely:

$$\begin{aligned} \xi &\rightsquigarrow \xi + \text{Re}\left(\frac{\eta}{\alpha}\right) \rightsquigarrow -i\left(\xi + \text{Re}\left(\frac{\eta}{\alpha}\right)\right) + \text{Im}\left(\frac{\eta}{\alpha}\right) = -i\xi - i\frac{\eta}{\alpha} \rightsquigarrow \\ &\rightsquigarrow i\alpha\left(-i\xi - i\frac{\eta}{\alpha}\right) = \alpha\xi + \eta. \end{aligned}$$

□

From now on, assume that  $\mathbb{X}$  carries a **holomorphic  $D$ -flat volume form**  $\omega$ , i.e. a nowhere vanishing holomorphic  $n$ -form  $\omega$  such that  $D\omega = 0$ , and that our actions  $\rho$  preserve this volume form. We denote by  $\text{Aff}(\mathbb{X}, D, \omega)$  the space of biholomorphisms of  $\mathbb{X}$  that preserve both  $D$  and  $\omega$ .

Given  $(\sigma, \xi)$  a complex affine immersion of  $\widetilde{M}$  into  $\mathbb{X}$ , we define a nowhere vanishing section  $\theta$  of  $\bigwedge^k \mathbb{C}T^*\widetilde{M}$  on  $\widetilde{M}$  by

$$\theta(X_1, \dots, X_m) = (\sigma^*\omega)(X_1, \dots, X_m, \xi). \quad (7)$$

We call  $\theta$  the **induced complex volume form**. By equivariance, the induced complex volume form  $\theta$  descends to a volume form on  $M$ .

**Proposition 3.5.** *For any  $X \in \mathbb{C}TM$ ,  $\nabla_X \theta = \tau(X)\theta$ .*

*Proof.* The proof for  $X \in \Gamma(TS)$  is identical to the proof of the analogous result for real affine immersions [NS94, Proposition II.1.4], then the statement follows by  $\mathbb{C}$ -linearity of  $\nabla\theta$  and  $\tau$ . □

Proposition 3.5 and Equation (6) tell us that  $\theta$  is  $\nabla$ -parallel if and only if  $(\sigma^*D)\xi$  preserves the sub-bundle  $\mathbb{C}T\widetilde{M}$ .

We say that  $\sigma$  is **non-degenerate** if, given a transversal vector field  $\xi$ , the affine fundamental form  $g$  is nowhere degenerate, i.e., it is a complex metric. By Lemma 3.4, all choices of affine transversal sections determine conformally equivalent fundamental forms, so this notion of non-degeneracy is independent of our choice of the transversal section.



**Remark 3.6.** Given a complex metric  $g$  on a smooth manifold  $\widetilde{M}$ , on each simply connected chart with local coordinates  $(x_i)_{i=1}^m$  one can define a local **metric volume form**  $dV_g = \sqrt{\det(g(\partial_{x_i}, \partial_{x_j}))_{i,j}} \cdot dx_1 \wedge \cdots \wedge dx_n$ , which is well-defined up to a sign. Since  $\widetilde{M}$  is simply connected, there always exists a global metric volume form on  $\widetilde{M}$ .

**Definition 3.7.** Let  $\rho: \pi_1(M) \rightarrow \text{Aff}(\mathbb{X}, D, \omega)$  be a homomorphism. A  $\rho$ -equivariant **Blaschke affine immersion** of  $\widetilde{M}$  in  $\mathbb{X}$  is an (equivariant) affine immersion  $(\sigma, \xi)$  such that:

- (1) the affine fundamental form  $g$  is nowhere degenerate, and there exists a global metric volume form  $dV_g$  on  $M$ ,
- (2)  $\nabla\theta = 0$ , hence  $\tau = 0$  by Proposition 3.5,
- (3)  $\theta = dV_g$ .

The transversal vector field  $\xi$  is called **affine normal** and  $g$  is called **Blaschke metric**.

**Proposition 3.8.** *Let  $M = \widetilde{M}$  be simply connected. Given a non-degenerate affine immersion  $(\sigma, \xi)$  of  $\widetilde{M}$  into  $\mathbb{X}$ , there exists a transversal vector field  $\hat{\xi}$ , unique up to a sign, such that  $(\sigma, \hat{\xi})$  is a Blaschke affine immersion.  $(\sigma, \hat{\xi})$  is called the **Blaschke normalization** of  $(\sigma, \xi)$ .*

*Proof.* This proof is totally analogous to that of the corresponding result in the real case (see [NS94, Theorem II.3.1]), but we include the full proof to show where the topological assumption is used.

Since  $\widetilde{M}$  is simply connected, the nondegenerate affine fundamental form  $g$  of  $(\sigma, \xi)$  admits a global metric volume form  $dV_g$ . Given  $\alpha \in C^\infty(M, \mathbb{C}^*)$  and  $\eta \in \Gamma(\mathbb{C}TM)$ , if we transform  $\xi$  to  $\hat{\xi} = \alpha\xi + \eta$ , the metric volume form  $dV_g$  and the induced complex volume form  $\theta$  of  $(\sigma, \xi)$  change according to  $dV_{\hat{g}} = \frac{1}{\alpha}dV_g$  and

$$\hat{\theta}(X, Y) = \omega(X, Y, \alpha\xi + \eta) = \omega(X, Y, \alpha\xi) = \alpha\theta(X, Y).$$

Since  $M$  is simply connected, there exists a function  $\alpha \in C^\infty(\widetilde{M}, \mathbb{C}^*)$ , unique up to multiplication by  $-1$ , such that

$$\alpha^2 = \frac{dV_g}{\theta}.$$

Choosing such  $\alpha$  guarantees that the condition (3) holds, so it remains to pick  $\eta$  so that (2) holds. Since  $g$  is non-degenerate, there exists a unique complex vector field  $\eta$  such that, as

$$g(\eta, \cdot) = -(\alpha\tau + d\alpha).$$

Using item (4) from Lemma 3.4, we see that  $\hat{\xi} = \alpha\xi + \eta$  satisfies the requirements of the proposition. Observe that if we exchange  $\alpha$  with  $-\alpha$ , then  $\eta$  becomes  $-\eta$ , and hence  $\xi$  is unique up to multiplication by  $\pm 1$ .  $\square$

**Remark 3.9.** Looking into the proof of Proposition 3.8, one can see that, unlike the real case, for general  $M$  and for a general representation  $\rho: \pi_1(M) \rightarrow \text{Aff}(\mathbb{X}, D, \omega)$ , a  $\rho$ -equivariant non-degenerate affine immersion  $(\sigma, \xi)$  might not admit a  $\rho$ -equivariant Blaschke normalization. In fact, one needs to assume that  $(M, g)$  admits a global volume form (equivalently, a  $\pi_1(M)$ -invariant metric volume form for  $\tilde{g}$  on  $\widetilde{M}$ ), and - even assuming this - the function  $\frac{dV_g}{\theta}$  might not admit a  $\pi_1(M)$ -invariant square root.

Nevertheless, the obstructions only depend on the topological properties of  $g$ ,  $dV_g$ , and  $\theta$ , so, if  $\sigma$  is a deformation of a real affine immersion, then it admits a Blaschke normalization.

**Example 3.10** (Graph immersions). Let  $U$  a domain in  $\mathbb{R}^n$  and  $F : \mathbb{R}^n \rightarrow \mathbb{C}$  a  $C^1$  function. A graph immersion is an admissible immersion  $\sigma : U \rightarrow \mathbb{C}^{n+1}$  of the form

$$\sigma(x_1, \dots, x_n) = (x_1, \dots, x_n, F(x_1, \dots, x_n)).$$

The choice of transversal vector field  $\xi = (0, \dots, 0, 1)$  yields  $\mathbf{S} = 0$  and  $\tau = 0$ .

**Example 3.11** (Complex paraboloids). As a special example of a graph immersion, we have what we call complex paraboloids. Consider, for instance,  $F(x, y) = (x, y, \frac{1}{2}(x^2 + e^{i\theta}y^2))$ . Taking  $e^{i\theta} = 1$  gives the well-known elliptic paraboloid, and taking  $e^{i\theta} = -1$  gives a hyperbolic paraboloid. It is not hard to compute that  $\xi = (0, 0, e^{-i\theta})$  produces a transversal vector field with  $\tau = 0$  and  $\theta = dV_g$ .

**3.3. The fundamental theorem for complex affine immersions.** From now on, we will assume that  $\mathbb{X} = \mathbb{C}^{n+1}$  endowed with the Levi-Civita connection of the holomorphic Riemannian metric, say  $D$ . The standard equations from affine differential geometry admit complex extensions.

**Proposition 3.12.** *Consider an equivariant complex affine immersion  $(\sigma, \xi)$  of  $\widetilde{M}$  into  $(\mathbb{C}^{m+1}, D)$  inducing the data  $\nabla, g, \mathbf{S}, \tau$  on  $M$  as in Equation (6).*

*The following equations hold for all  $X, Y, Z \in \Gamma(\mathbb{C}TM)$ .*

$$\text{Gauss equation:} \quad R^\nabla(X, Y)Z = g(Y, Z)\mathbf{S}(X) - g(X, Z)\mathbf{S}(Y), \quad (8)$$

$$\text{Codazzi equation 1:} \quad (\nabla_X g)(Y, Z) + \tau(X)g(Y, Z) = (\nabla_Y g)(X, Z) + \tau(Y)g(X, Z), \quad (9)$$

$$\text{Codazzi equation 2:} \quad (\nabla_X \mathbf{S})(Y) - \tau(X)\mathbf{S}(Y) = (\nabla_Y \mathbf{S})(X) - \tau(Y)\mathbf{S}(X), \quad (10)$$

$$\text{Ricci equation:} \quad g(X, \mathbf{S}(Y)) - g(\mathbf{S}(X), Y) = d\tau(X, Y). \quad (11)$$

*Proof.* The corresponding results hold for (real) affine immersions between real manifolds [NS94, Theorem II.2.1], as a result of the flatness of  $D$ . Indeed, one can see explicitly from Equation (6) that for all  $X, Y, Z \in \Gamma(TM)$

$$\begin{aligned} R^D(X, Y)Z &= R^\nabla(X, Y)Z - g(Y, Z)\mathbf{S}(X) + g(X, Z)\mathbf{S}(Y) \\ &\quad + (\nabla_X g)(Y, Z)\xi - (\nabla_Y g)(X, Z)\xi + g(Y, Z)\tau(X) - g(X, Z)\tau(Y), \\ R^D(X, Y)\xi &= -(\nabla_X \mathbf{S})(Y) + (\nabla_Y \mathbf{S})(X) - \tau(X)\mathbf{S}(Y) + \tau(Y)\mathbf{S}(X) \\ &\quad - g(X, \mathbf{S}(Y))\xi + g(Y, \mathbf{S}(X))\xi + d\tau(X, Y)\xi, \end{aligned} \quad (12)$$

hence Equations (8)-(9)-(10)-(11) follow by  $R^D(X, Y)Z = 0 = R^D(X, Y)\xi$ . Finally, the statement follows for all  $X, Y, Z \in \Gamma(\mathbb{C}TS)$  by  $\mathbb{C}$ -linearity of  $\nabla$  and of the tensors.  $\square$

Conversely, the immersion data  $\nabla, g, \mathbf{S}$ , and  $\tau$  determine uniquely an affine immersion, as shown in the following statement.

**Theorem 3.13.** *Let  $M$  be a smooth real  $m$ -manifold. Let  $\nabla : \Gamma(\mathbb{C}TM) \rightarrow \Gamma(\text{End}_{\mathbb{C}}(\mathbb{C}TM))$  be a torsion-free affine connection, let  $g$  be a complex metric,  $\mathbf{S}$  an endomorphism of  $\mathbb{C}TM$ , and  $\tau \in \Omega^1(\mathbb{C}TM)$  such that Equations (8)-(9)-(10)-(11) hold.*

*Then, there exists a  $\pi_1(M)$ -equivariant complex affine immersion  $(\sigma, \xi) : \widetilde{M} \rightarrow \mathbb{C}^{m+1}$  with affine connection  $\nabla$ , affine fundamental form  $g$ , shape operator  $\mathbf{S}$ , and transversal connection form  $\tau$ . Such affine immersions are unique up to composition with elements in  $\text{Aff}(\mathbb{C}^{m+1}, D) = \text{GL}(m+1, \mathbb{C}) \times \mathbb{C}^{m+1}$ .*

*Moreover, if  $\tau = 0$  and  $\nabla dV_g = 0$ ,  $(\sigma, \xi)$  can be chosen to be an equivariant Blaschke affine immersion for  $\mathbb{C}^{m+1}$  endowed with the canonical volume form  $\omega = dz_1 \wedge \dots \wedge dz_{n+1}$ , and its holonomy takes values in  $\text{SL}(m+1, \mathbb{C}) \times \mathbb{C}^{m+1}$ .*

For this statement, too, the computation follows as in the real case [NS94, Theorem II.8.1], up to a few adjustments. We recall the essential ideas and point out the main differences.

*Proof. Step 1: Existence for  $M = \widetilde{M}$ .* Let  $N: \mathbb{C} \times \widetilde{M} \rightarrow \widetilde{M}$  be the trivial bundle, and fix  $\hat{\xi} \in \Gamma(N)$  a nowhere zero section. Consider the bundle  $V = \mathbb{C}T\widetilde{M} \oplus N$  and the connection  $\hat{D}$  defined for all  $X \in \Gamma(T\widetilde{M})$  and  $Y \in \Gamma(\mathbb{C}T\widetilde{M})$  as

$$\begin{cases} \hat{D}_X Y &= \nabla_X Y + g(X, Y)\hat{\xi} \\ \hat{D}_X \hat{\xi} &= -S(X) + \tau(X)\hat{\xi} \\ \hat{D}_X(i\hat{\xi}) &= -iS(X) + i\tau(X)\hat{\xi}. \end{cases}$$

Then, one gets the same equation as in (12) after replacing  $D$  with  $\hat{D}$ , and  $R^{\hat{D}}(X, Y)(i\hat{\xi}) = iR^{\hat{D}}(X, Y)(\hat{\xi})$ , hence the hypotheses of the theorem imply that  $R^{\hat{D}} = 0$ . Moreover,  $\hat{D}iY = i\hat{D}Y$ , so there exists a bundle isomorphism

$$\Phi: (\mathbb{C}T\widetilde{M} \oplus N, \hat{D}) \rightarrow (\mathbb{C}^{m+1} \times \widetilde{M}, D)$$

with  $D$  denoting the standard connection on the trivial bundle, and  $\Phi$  being a  $\mathbb{C}$ -linear isomorphism fiberwise.

For all  $X \in \Gamma(\mathbb{C}T\widetilde{M})$ , we can see  $\Phi(X_p) =: (p, \alpha(X_p))$  with  $\alpha$  a  $\mathbb{C}^{m+1}$ -valued 1-form on  $\widetilde{M}$ . With a little abuse of notation, we will say that  $\alpha = \Phi|_{\mathbb{C}T\widetilde{M}}$ . Then, for all  $X, Y \in \Gamma(T\widetilde{M})$ ,  $D_X(\Phi(Y)) = \alpha(Y)$ , and

$$d\alpha(X, Y) = \partial_X(\alpha(Y)) - \partial_Y(\alpha(X)) - \alpha([X, Y]) = \Phi(\nabla_X Y - \nabla_Y X - [X, Y]) = 0$$

as a result of  $\nabla$  being torsion-free. Hence, there exists  $\sigma: \widetilde{M} \rightarrow \mathbb{C}^{m+1}$  such that  $d\sigma = \alpha$ . As a consequence,  $\sigma_*(X) = d\sigma(X) = \Phi(X)$  so  $\sigma_*(X) = 0$  iff  $X = 0$ , implying that  $\sigma$  is an admissible immersion.

Moreover, if we consider the pull-back bundle  $\sigma^*(\mathbb{C}^{m+1})$  with the pull-back connection  $\sigma^*D$  and transversal vector field  $\xi$  defined by  $\sigma_*(\xi) = \Phi(\hat{\xi})$ , we get that, for all  $X \in \Gamma(T\widetilde{M})$  and  $Y \in \Gamma(\mathbb{C}T\widetilde{M})$ , we have at each point that

$$\begin{cases} \sigma_*((\sigma^*D)_X Y) = D_X(\sigma_*(Y)) = D_X(\Phi(Y)) = \Phi(\hat{D}_X Y) = \Phi(\nabla_X Y + g(X, Y)\hat{\xi}) = \sigma_*(\nabla_X Y) + g(X, Y)\sigma_*(\hat{\xi}) \\ \sigma_*((\sigma^*D)_X \xi) = D_X(\sigma_*\xi) = D_X(\Phi(\hat{\xi})) = \Phi(\hat{D}_X \hat{\xi}) = -\Phi(S(X)) + \tau(X)\Phi(\hat{\xi}) = \sigma_*(-S(X) + \tau(X)\hat{\xi}), \end{cases}$$

which implies that  $(\sigma, \xi)$  has immersion data given by  $\nabla, g, S, \tau$ .

*Step 2: Uniqueness and the equivariant case.* Let  $(\sigma, \xi)$  and  $(\sigma', \xi')$  be two affine immersions with the same immersion data  $\nabla, g, S, \tau$ . We prove that there exists  $A \in GL(m+1, \mathbb{C}) \times \mathbb{C}^{m+1}$  such that  $\sigma' = A \circ \sigma$  and  $\sigma'_*\xi' = A(\sigma_*\xi)$ .

Define the function  $L: \widetilde{M} \rightarrow GL(m+1, \mathbb{C})$  as the unique matrix such that, for all  $X \in \Gamma(\mathbb{C}T\widetilde{M})$ ,  $\sigma'_*(X) = L \circ \sigma_*(X)$  and  $\sigma'_*(\xi') = L(\sigma_*\xi)$ .

Then (recalling that  $\sigma_*((\sigma^*D)_X Y) = \partial_X(d\sigma(Y)) = \partial_X(\sigma_*(Y))$ ), we have that for all  $X \in \Gamma(T\widetilde{M})$ ,  $Y \in \Gamma(\mathbb{C}T\widetilde{M})$ ,

$$\begin{aligned} (\partial_X L)(\sigma_* Y) &= \partial_X((L \circ \sigma_*)(Y)) - L(\partial_X(\sigma_* Y)) = \partial_X(\sigma'_*(Y)) - L(\partial_X(\sigma_* Y)) \\ &= \sigma'_*(\nabla_X Y) + g(X, Y)\xi' - L(\sigma_*(\nabla_X Y)) - g(X, Y)L(\xi) = 0, \\ (\partial_X L)(\sigma_* \xi) &= \partial_X(L(\sigma_* \xi)) - L(\partial_X(\sigma_* \xi)) = \partial_X(L(\sigma_* \xi)) - L(\partial_X(\sigma_* \xi)) - L(\partial_X(\sigma_* \xi)) \\ &= -\sigma'_*(S(X)) + \tau(X)\sigma'_*(\xi) + L(\sigma_*(S(X))) + \tau(X)L(\sigma_*\xi) = 0, \end{aligned}$$

hence  $dL = 0$ , implying that  $L$  is a constant. Moreover,  $d(\sigma' - L\sigma) = 0$ , hence there exists  $b \in \mathbb{C}^{m+1}$  such that  $\sigma' = L \cdot \sigma + b$ , proving uniqueness. The equivariant case follows from uniqueness in the usual way.

*Step 3: Blaschke immersions.* Let  $\omega = dz_1 \wedge \cdots \wedge dz_{m+1}$  be the standard  $D$ -flat volume form on  $\mathbb{C}^{m+1}$ . Assume  $\tau = 0$ , and that  $\nabla dV_g = 0$  and construct a corresponding equivariant affine immersion  $(\sigma, \xi): \widetilde{M} \rightarrow \mathbb{C}^{m+1}$ , and let  $\theta = (\sigma^*\omega)(\cdot, \xi)$  be the pull-back form. Then, by Proposition 3.5,  $\nabla\theta = 0$ . Assuming  $\nabla(dV_g) = 0$  as well, we get that  $dV_g = C\theta$  for some constant  $C$  as a consequence of the fact that  $\bigwedge^n \mathbb{C}T^*\widetilde{M}$  is rank-1. Up to replacing now  $(\sigma, \xi)$  with the affinely equivalent affine immersion  $(\frac{1}{C}\sigma, \frac{1}{C}\xi)$ , we can assume that  $C = 1$ , therefore  $(\sigma, \xi)$  is a Blaschke affine immersion in  $(\mathbb{C}^{m+1}, D, \omega)$ . If  $(\sigma, \xi)$  is equivariant, its holonomy must have image in  $\mathrm{SL}(m+1, \mathbb{C}) \times \mathbb{C}^{m+1}$ . □

**3.4. Complex affine spheres.** We now restrict to the case where  $M$  is a closed oriented surface, and accordingly now write  $M = S$ , and  $\mathbb{X} = \mathbb{C}^3$  with the standard affine connection  $D$  and volume form  $\omega = dz_1 \wedge dz_2 \wedge dz_3$ . We have that  $\mathrm{Aff}(\mathbb{C}^3, D, \omega) = \mathrm{SL}(3, \mathbb{C}) \times \mathbb{C}^3$ .

**Definition 3.14.** Let  $\rho: \pi_1(M) \rightarrow \mathrm{Aff}(\mathbb{C}^3, D, \omega)$ , and let  $(\sigma, \xi)$  be a  $\rho$ -equivariant Blaschke affine immersion. We say that  $(\sigma, \xi)$  is a (proper) **complex affine sphere** if the shape operator has the form  $S = \lambda \cdot id$ , where  $\lambda \in \mathbb{C}^*$ . A complex affine sphere  $(\sigma, \xi)$  is **positive and hyperbolic** if  $\lambda = -1$  and the Blaschke metric  $g$  is a positive complex metric.

**Remark 3.15.** As shown in [BE22, Theorem 6.5], every positive complex metric on an oriented surface  $S$  admits a metric volume form compatible with the orientation. We will assume that the affine normal is compatible with the induced volume form.

**Remark 3.16.** If an affine sphere  $(\sigma, \xi)$  has shape operator in the form  $S = \lambda \cdot id$  with  $\lambda$  a function, by Codazzi Equation in Proposition 3.12 one gets that  $d\lambda = 0$ , hence  $\lambda$  is a constant.

**Remark 3.17.** If  $(\sigma, \xi)$  is a proper complex affine sphere, then all of the affine normals meet at a single point. Indeed, one can see that the map  $p \mapsto \sigma(p) + \frac{1}{\lambda}\xi(p)$  is a constant.

**Convention.** By Remark 3.17, up to composing  $\sigma$  with a translation, from now on we will assume that the meeting point of the affine normals is  $\underline{0} \in \mathbb{C}^3$ , hence

$$\sigma = -\frac{1}{\lambda}\xi \tag{13}$$

and the data of  $\sigma$  and  $\lambda$  include the data of  $\xi$ . In this setting, if  $\sigma$  is an equivariant affine sphere, then its holonomy fixes  $\underline{0}$ , and hence takes values in  $\mathrm{SL}(3, \mathbb{C})$ .

Pick tensors play a prominent role in the real theory, and we turn to them now. Given  $g$ , we have both the induced connection  $\nabla$  and the Levi-Civita connection of  $g$ , say  $\nabla^g$ . The difference tensor on  $\mathbb{C}TS$  is

$$K(X, Y) = K_X(Y) = \nabla_X Y - \nabla_X^g Y.$$

Since both connections have no torsion,  $K$  is symmetric. The **Pick tensor** is the complex 3-form defined by lowering indices,

$$C(X, Y, Z) = -2g(K_X Y, Z).$$

The proposition below can be proved as in the real case for real vector fields (see [NS94, Section II.4]), and by  $\mathbb{C}$ -linearity we get the general case.

**Proposition 3.18.** *Let  $(\sigma, \xi)$  be an equivariant proper complex affine sphere. Then, for all  $X, Y, Z, W \in \Gamma(\mathbb{C}TS)$ , we have that*

- (1)  $C = \nabla g$ ,
- (2)  $\text{tr}_g K_X = 0$ ,
- (3)  $C, \nabla C, \nabla K, \nabla^g C$ , and  $\nabla^g K$  are symmetric in all of their entries,
- (4)  $g((\nabla_X^g K)(Y, Z), W) = g((\nabla_X^g K)(Y, W), Z)$ ,
- (5)  $R^\nabla(X, Y)Z = R^g(X, Y)Z + [K_X, K_Y](Z)$ .

We use only items (2) and (3), but the other results above help motivate some aspects of the proof of Theorem 3.20 below. Using that  $\nabla^g C$  is symmetric, we show that positive hyperbolic affine spheres come with holomorphic cubic differentials, and thus we see the link to Theorem A.

**Proposition 3.19.** *Let  $(\sigma, \xi)$  be an equivariant positive hyperbolic complex affine sphere in  $\mathbb{C}^3$ . Let  $g$  be the Blaschke metric, with conformal classes given by  $(c_1, \bar{c}_2) \in \mathcal{C}(S) \times \mathcal{C}(\bar{S})$ . Then,*

$$C = Q_1 + \bar{Q}_2,$$

where  $Q_1$  is a  $c_1$ -holomorphic cubic differential and  $\bar{Q}_2$  is a  $\bar{c}_2$ -holomorphic cubic differential.

*Proof.* Let  $z$  and  $\bar{w}$  denote local coordinates for  $c_1$  and  $\bar{c}_2$  respectively. By Proposition 3.18,  $C$  is symmetric, so it decomposes as

$$C = pdz^3 + qdz^2 \cdot d\bar{w} + rdz \cdot d\bar{w}^2 + sd\bar{w}^3,$$

where  $p, q, r, s$  are local  $\mathbb{C}$ -valued functions. By (2) in Proposition 3.18,  $\text{tr}_g(K_{\partial_{\bar{z}}}) = 0$ , hence

$$0 = \text{tr}_g(C(\partial_{\bar{z}}, \cdot, \cdot)) = 2 \frac{1}{g(\partial_{\bar{z}}, \partial_w)} C(\partial_{\bar{z}}, \partial_{\bar{z}}, \partial_w) = \frac{\partial_{\bar{z}} \bar{w} \cdot \partial_w z}{h(\partial_{\bar{z}}, \partial_w)} r,$$

hence  $r = 0$ . Similarly, from  $\text{tr}_g(K_{\partial_w}) = 0$  we deduce that  $q = 0$ .

We are left to prove that  $p$  is  $c_1$ -holomorphic and  $s$  is  $\bar{c}_2$ -holomorphic. First, by Lemma 2.5, for all  $X \in \Gamma(\mathbb{C}TS)$ ,  $\nabla_X^g \partial_{\bar{z}} \in \text{Span}(\partial_{\bar{z}})$  and  $\nabla_X^g \partial_w \in \Gamma(\text{Span}(\partial_w))$ . As a result,

$$(\nabla^g dz)(\partial_{\bar{z}}) = d(dz(\partial_{\bar{z}})) - dz(\nabla^g \partial_{\bar{z}}) = 0 - 0 = 0.$$

Moreover, by Cartan's formula, for all  $X, Y \in \Gamma(\mathbb{C}TS)$ ,

$$\begin{aligned} (\nabla_X^g dz)(Y) - (\nabla_Y^g dz)(X) &= \partial_X(dz(Y)) - dz(\nabla_X^g Y) - \partial_Y(dz(X)) + dz(\nabla_Y^g X) = \\ &= \partial_X(dz(Y)) - \partial_Y(dz(X)) - dz([X, Y]) = (ddz)(X, Y) = 0, \end{aligned}$$

hence  $\nabla^g dz$  is commutative and  $\nabla^g dz \in \Gamma(\text{Span}(dz^2))$ . In the same fashion,  $\nabla^g d\bar{w} \in \Gamma(\text{Span}(d\bar{w}^2))$ . Iterating the derivative, one can see that for all  $m \in \mathbb{Z}_+$

$$\nabla^g(dz^m) \in \Gamma(\text{Span}(dz^{m+1})) \quad \nabla^g(d\bar{w}^m) \in \Gamma(\text{Span}(d\bar{w}^{m+1})).$$

Now, by (3) in Proposition 3.18, we have that  $\nabla^g C$  is symmetric. Hence, from

$$\nabla^g C = dp \otimes dz^3 + p \nabla^g(dz^3) + ds \otimes d\bar{w}^3 + s \nabla^g(d\bar{w}^3)$$

we finally observe that

$$\begin{aligned} \partial_{\bar{z}} p \cdot (\partial_w z)^3 &= (\nabla_{\partial_{\bar{z}}}^g C)(\partial_w, \partial_w, \partial_w) = (\nabla_{\partial_w}^g C)(\partial_w, \partial_w, \partial_{\bar{z}}) = 0, \\ \partial_w s \cdot (\partial_{\bar{z}} \bar{w})^3 &= (\nabla_{\partial_w}^g C)(\partial_{\bar{z}}, \partial_{\bar{z}}, \partial_{\bar{z}}) = (\nabla_{\partial_{\bar{z}}}^g C)(\partial_{\bar{z}}, \partial_{\bar{z}}, \partial_w) = 0, \end{aligned}$$

implying that  $\partial_{\bar{z}} p = \partial_w s \equiv 0$ . □

Conversely, the data of  $g$  and  $C$  completely characterize the complex affine immersion, as described by the following theorem.

**Theorem 3.20.** *Let  $g$  be a positive complex metric on a closed oriented surface  $S$ , with conformal class  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ , and let  $Q_1$  and  $\overline{Q_2}$  be holomorphic cubic differentials for  $c_1$  and  $\overline{c_2}$  respectively.*

*There exists an equivariant positive hyperbolic complex affine sphere  $\sigma: \tilde{S} \rightarrow (\mathbb{C}^3, D, \omega)$  with Blaschke metric  $g$ , Pick tensor  $C = Q_1 + \overline{Q_2}$ , and with holonomy in  $\mathrm{SL}(3, \mathbb{C})$  if and only if  $g$  and  $C$  satisfy*

$$K_g = -1 + \frac{1}{4}g(Q_1, \overline{Q_2}). \quad (14)$$

*Moreover, such a complex affine sphere is unique up to composition with elements in  $\mathrm{SL}(3, \mathbb{C})$ .*

In local holomorphic coordinates  $z$  and  $\overline{w}$  for  $c_1$  and  $\overline{c_2}$  respectively, if  $g = \lambda dz d\overline{w}$  and  $Q_1 = \varphi dz^3$ ,  $\overline{Q_2} = \overline{\psi} d\overline{w}^3$ , then

$$\frac{1}{4}g(Q_1, \overline{Q_2}) = 2\frac{\varphi \overline{\psi}}{\lambda^3}.$$

As well, take note that, in terms of the Pick tensor,  $g(Q_1, \overline{Q_2}) = \frac{1}{2}g(C, C)$ , since  $Q_1$  and  $\overline{Q_2}$  are  $g$ -isotropic.

*Proof.* As above, let  $z$  and  $\overline{w}$  be coordinates for  $c_1$  and  $\overline{c_2}$  respectively, and  $g = \lambda dz d\overline{w}$ ,  $Q_1 = \varphi dz^3$ , and  $\overline{Q_2} = \overline{\psi} d\overline{w}^3$ .

Let  $\nabla$  be the affine connection on  $\mathbb{C}TS$  defined by  $C = -2g(\nabla - \nabla^g, \cdot)$ . It suffices to show that  $g, \nabla, S = id$ , and  $\tau = 0$  satisfy the hypothesis of Theorem 3.13, as well as  $\nabla dV_g = 0$ , to produce an equivariant complex affine sphere. As in Remark 3.17, by translating we can ensure that the holonomy lies in  $\mathrm{SL}(3, \mathbb{C})$ .

Observe that Equations (10) and (11) hold trivially. We prove that Equation (9) holds. Since  $\nabla^g g = 0$ , for all  $X, Y, Z \in \Gamma(\mathbb{C}TS)$ ,

$$\begin{aligned} (\nabla_X g)(Y, Z) - (\nabla_Y g)(X, Z) &= (\nabla_X g)(Y, Z) - (\nabla_Y g)(X, Z) - (\nabla_X^g g)(Y, Z) - (\nabla_Y^g g)(X, Z) \\ &= -g(\nabla_X Y, Z) - g(Y, \nabla_X Z) + g(\nabla_Y X, Z) + g(X, \nabla_Y Z) \\ &= \frac{1}{2}(C(X, Y, Z) + C(X, Z, Y) - C(Y, X, Z) - C(Y, Z, X)) = 0, \end{aligned}$$

where in the last line we used the symmetry of  $C$ .

We next prove that  $\nabla dV_g = 0$ . Suggestively defining the symmetric tensor  $K$  by  $K_X Y = \nabla_X Y - \nabla_X^g Y$ , we observe that for all  $X$ ,

$$0 = C(\cdot, \partial_{\overline{z}}, \partial_w) = -2g(K \cdot \partial_{\overline{z}}, \partial_w),$$

and hence  $K_X \partial_{\overline{z}} = K_{\partial_{\overline{z}}} X \in \Gamma(\mathrm{Span}(\partial_w))$ . Similarly,  $K_X \partial_w = K_{\partial_w} X \in \Gamma(\mathrm{Span}(\partial_{\overline{z}}))$ . One can compute the tensor  $K$  explicitly:

$$\begin{aligned} K_{\partial_w} \partial_w &= -\frac{1}{2} \frac{(\partial_w z)^3}{g(\partial_{\overline{z}}, \partial_w)} \varphi \partial_{\overline{z}}, \\ K_{\partial_{\overline{z}}} \partial_{\overline{z}} &= -\frac{1}{2} \frac{(\partial_{\overline{z}} \overline{w})^3}{g(\partial_{\overline{z}}, \partial_w)} \overline{\psi} \partial_w, \\ K_{\partial_{\overline{z}}} \partial_w &= K_{\partial_w} \partial_{\overline{z}} = 0. \end{aligned} \quad (15)$$

Observe that, for all 1-forms  $\theta$ , one has  $K_X(\theta) := \nabla_X \theta - \nabla_X^g \theta = -\theta(K_X)$ . Hence,  $K_{\partial_w} dz = -dz(K_{\partial_w}) = 0$ , implying that  $K dz \in \mathrm{Span}(d\overline{w}^2)$  and  $K d\overline{w} \in \mathrm{Span}(dz^2)$ .

As a result, writing  $dV_g$  in the form  $dV_g = \delta dz \wedge d\overline{w}$ , we have that for all  $X \in \Gamma(\mathbb{C}TS)$

$$\nabla_X dV_g = \nabla_X dV_g - \nabla_X^g dV_g = \delta K_X(dz) \wedge d\overline{w} + \delta dz \wedge K_X(d\overline{w}) = 0,$$

as desired.

Now we prove that Equation (8) holds. First of all, observe that  $\nabla^g C$  is symmetric in all of its entries. In fact,

$$\nabla^g C = \partial_z \varphi dz^4 + \varphi \nabla^g(dz^3) + \partial_{\bar{w}} \bar{\psi} d\bar{w}^4 + \bar{\psi} \nabla^g(d\bar{w}^3)$$

and, as seen in the proof of Proposition 3.19,  $\nabla^g(dz^3) \in \Gamma(\text{Span}(dz^4))$  and  $\nabla^g(d\bar{w}^3) \in \Gamma(\text{Span}(d\bar{w}^4))$ . As a consequence,  $\nabla^g K$  is symmetric in all of its entries because one can easily check that

$$(\nabla_X^g C)(Y, Z, W) - (\nabla_Y^g C)(X, Z, W) = -2g((\nabla_X^g K)(Y, Z), W) + 2g((\nabla_Y^g K)(X, Z), W)$$

for all  $X, Y, Z \in \Gamma(\mathbb{C}TS)$ . Now, from  $\nabla = \nabla^g + K$ , we have that

$$\begin{aligned} R^\nabla(X, Y)Z &= R^g(X, Y)Z + (\nabla_X^g K)(Y, Z) - (\nabla_Y^g K)(X, Z) + [K_X, K_Y]Z \\ &= R^g(X, Y)Z + [K_X, K_Y]Z, \end{aligned}$$

so the Gauss Equation (8) for  $S = -id$  is equivalent to

$$R^g(X, Y)Z = -[K_X, K_Y]Z - g(Y, Z)X + g(X, Z)Y,$$

hence to

$$g(R^g(X, Y)Z, W) = -g([K_X, K_Y]Z, W) - g(Y, Z)g(X, W) + g(X, Z)g(Y, W) \quad (16)$$

for all  $X, Y, Z, W \in \Gamma(\mathbb{C}TS)$ . It is now easy to check, using that  $K$  is  $g$ -self-adjoint, that in Equation (16) the right-hand side has the same symmetries as the left-hand side, so it is sufficient to check that the equality holds for  $X = W = \partial_{\bar{z}}$  and  $Y = Z = \partial_w$ . Finally writing out the Gauss curvature as

$$K_g = \frac{g(R^g(\partial_{\bar{z}}, \partial_w)\partial_w, \partial_{\bar{z}})}{-(g(\partial_{\bar{z}}, \partial_w))^2} = \frac{g(K_{\partial_{\bar{z}}} K_{\partial_w} \partial_w, \partial_{\bar{z}}) - g(K_{\partial_w} K_{\partial_{\bar{z}}} \partial_w, \partial_{\bar{z}}) - (g(\partial_{\bar{z}}, \partial_w))^2}{(g(\partial_{\bar{z}}, \partial_w))^2},$$

Equation (15) returns that the Gauss equation (8) is equivalent to

$$K_g = \frac{g(K_{\partial_w} \partial_w, K_{\partial_{\bar{z}}} \partial_{\bar{z}})}{(g(\partial_{\bar{z}}, \partial_w))^2} - 1 = \frac{1}{4} \varphi \bar{\psi} \cdot \frac{(\partial_w z)^3 (\partial_{\bar{z}} \bar{w})^3}{(g(\partial_{\bar{z}}, \partial_w))^3} - 1 = 2 \frac{\varphi \bar{\psi}}{\lambda^3} - 1.$$

□

From now on, we will be interested only in Blaschke metrics  $g$  of the form  $g = e^{2u}h$ , with  $h$  being a Bers metric and  $u : S \rightarrow \mathbb{C}$  a function. Any Blaschke metric that's connected to a Riemannian metric can be written in this way.

**Example 3.21.** We are not providing a full construction, but there are in fact positive complex metrics  $g$  that have constant curvature  $-1$  and that are conformal multiples of Bers metrics by a conformal multiple that does not admit a logarithm; by 3.20,  $g$  and  $C = 0$  define a positive hyperbolic complex affine sphere. To construct such a metric, one can imagine starting from a Fuchsian type immersion  $(f, \bar{f}) : \tilde{S} \rightarrow \mathbb{G}$  and applying to  $f$  a  $2\pi$ -grafting around a closed non-separating curve  $\tilde{\gamma}$ : precomposing with a suitable isotopy before grafting, we get a new developing map  $\hat{f} : \tilde{S} \rightarrow \mathbb{C}\mathbb{P}^1$  with the same holonomy and such that  $\hat{f}(p) \neq \bar{f}(p)$  for all  $p \in \tilde{S}$ , hence  $(\hat{f}, \bar{f}) : \tilde{S} \rightarrow \mathbb{G}$  is an equivariant immersion. The induced complex metric  $g$  now has curvature  $-1$  by Theorem 6.7 in [BE22], and it is such that, assuming  $\dot{\gamma} \neq 0$ ,  $g(\dot{\gamma}, \dot{\gamma})$  defines a homotopically nontrivial loop on  $\mathbb{C}^*$ , defining an example as above.

As a consequence of Theorem 3.20 and of the formula for the curvature of conformal metrics in Section 2.3, we have the following corollary.

**Corollary 3.22.** *Let  $h = \mathbf{h}_{(c_1, \overline{c_2})}$  be a Bers metric, with  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ . Let  $Q_1$  and  $\overline{Q_2}$  be holomorphic cubic differentials for  $c_1$  and  $\overline{c_2}$  respectively. Then  $g = e^{2u} \cdot h$ , with  $u: S \rightarrow \mathbb{C}$ , and  $C = Q_1 + \overline{Q_2}$  are the Blaschke metric and the Pick tensor of an equivariant positive hyperbolic complex affine sphere in  $\mathbb{C}^3$  if and only if*

$$G(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) := \Delta_h u - e^{2u} + \frac{1}{4}h(Q_1, \overline{Q_2}) \cdot e^{-4u} + 1 = 0. \quad (17)$$

**3.5. Motivating examples.** Our first two examples show that complex affine spheres generalize both real affine spheres and the quasiconformal maps from the Bers theorem.

**Example 3.23** (Real affine spheres). Affine differential geometry is a special case of the theory outlined here.

By embedding  $\mathbb{R}^3$  inside  $\mathbb{C}^3$ , any immersion  $\sigma: \tilde{S} \rightarrow \mathbb{R}^3$  can be viewed as an admissible immersion  $\sigma: \tilde{S} \rightarrow \mathbb{C}^3$ . Moreover, if we can pick a section  $\xi$  of  $\sigma^*T\mathbb{R}^3$  that is transverse to  $TS$ , in which case  $\sigma$  is an affine immersion in the traditional sense [NS94, Definition II.1.1], then  $\sigma: \tilde{S} \rightarrow \mathbb{C}^3$  is a complex affine immersion. If we demand that the image of  $\sigma$  is locally strictly convex, then one can find a global real transverse normal. Choosing the transverse normal to be the complexification of a real transverse normal, all of the tensors that come into play are complexifications of real tensors. Moreover, the Blaschke metric is Riemannian, the Pick tensor is real, and the immersion data from Theorem 3.20 is always of the form  $(c, \overline{c}, Q, \overline{Q})$ . Equation (17) is a real semilinear elliptic PDE,

$$\Delta_h u = e^{2u} - \frac{1}{4}h(Q, \overline{Q})e^{-4u} - 1,$$

which is the ordinary Tzitzéica equation and is known to admit a unique real solution [Lof01, Proposition 4.0.2], [Lab07, Lemma 4.1.2]. If one takes that real solution and integrates it out to an affine sphere using Theorem 3.20, then by uniqueness that affine sphere lives in an isometrically and totally geodesically embedded copy of  $\mathbb{R}^3$ .

**Example 3.24** (Bers maps). Consider  $\mathbb{C}^{2,1}$ , namely  $\mathbb{C}^3$  with the complex bilinear form  $\langle \cdot, \cdot \rangle$  defined by  $\langle \underline{z}, \underline{w} \rangle = z_1 w_1 + z_2 w_2 - z_3 w_3$ , and define  $\hat{\mathbb{X}}_2 = \{\underline{z} \in \mathbb{C}^3 \mid \langle \underline{z}, \underline{z} \rangle = -1\}$  with the inherited holomorphic Riemannian metric. With reference to Section 2.2, one has  $\hat{\mathbb{X}}_2 \cong \mathbb{X}_2 \cong \mathbb{G}$ .

We show that positive hyperbolic complex affine spheres with Pick tensor  $C = 0$  correspond to equivariant immersions into  $\hat{\mathbb{X}}_2$  with pull-back metric being the Blaschke metric (which, by Theorem 3.20, has curvature -1). Moreover, in the identification of  $\text{PSL}(2, \mathbb{C})$  with  $\text{Isom}_0(\hat{\mathbb{X}}_2) = \text{Isom}_0(\mathbb{C}^{2,1}) = \text{SO}(2, 1, \mathbb{C})$ , if the Blaschke metric is a Bers metric, then the holonomy is quasi-Fuchsian, which shows that the construction is compatible with Bers' theorem.

As a consequence of Theorem 6.7 in [BE22], for all positive complex metric  $g$  of curvature -1 on  $S$ , there exists a  $(\pi_1(S), \text{SO}(3, \mathbb{C}))$ -equivariant admissible immersion  $\sigma: \tilde{S} \rightarrow \hat{\mathbb{X}}_2$  with pull-back metric  $g$ . Moreover, such an immersion is totally geodesic, namely  $(\sigma^*D^{\hat{\mathbb{X}}_2})_X Y = \nabla_X^g Y$  where  $D^{\hat{\mathbb{X}}_2}$  denotes the Levi-Civita connection of  $\hat{\mathbb{X}}_2$ . On the other hand,  $N_{\underline{z}} = i\underline{z}$  is a normal vector field for  $\hat{\mathbb{X}}_2$  inside  $\mathbb{C}^{2,1}$  (namely, at each point it is the unique vector - up to a sign - with norm 1 orthogonal to  $T\hat{\mathbb{X}}_2$ ). As a result, one can easily see that  $(\sigma^*D)_X N = iX$  and the second fundamental form of the immersion inside  $\mathbb{C}^3$  is  $-ig \otimes N = g \otimes \underline{\sigma(p)}$ , where  $\underline{\sigma(p)}$  denotes  $\sigma(p)$  seen as a vector of  $\mathbb{C}^3$ . Hence, denoting  $\xi_p = \sigma^*(\underline{\sigma(p)})$ , we have that

$$\begin{cases} (\sigma^*D)_X Y &= \nabla_X^g Y + g(X, Y)\xi \\ (\sigma^*D)_X \xi &= X. \end{cases}$$



Fix the canonical volume form  $\omega$  on  $\mathbb{C}^{2,1}$  given by the determinant. One can easily see that  $\omega(\cdot, \cdot, \underline{z})$  on  $\hat{\mathbb{X}}_2$  is a volume, i.e. it is compatible with the holomorphic Riemannian metric. Up to composing  $\sigma$  with the map  $-id$  on  $\hat{\mathbb{X}}_2$  (which does not lie in  $SO(2, 1, \mathbb{C})$ ), we have that  $dV_g = \sigma^*(\omega(\cdot, \cdot, z))$ . We therefore conclude that  $\sigma$  is a positive hyperbolic complex affine sphere. Since the affine connection coincides with  $\nabla^g$ , the Pick tensor is zero, and by uniqueness, we conclude that this is the unique positive hyperbolic complex affine sphere with Blaschke metric  $g$  and Pick tensor  $C = 0$ . Finally, in the identification of  $PSL(2, \mathbb{C}) \cong Isom_0(\mathbb{G}) \cong Isom_0(\hat{\mathbb{X}}_2) \cong SO(2, 1, \mathbb{C})$ , one has trivially that the holonomy  $\pi_1(S) \rightarrow SO(2, 1, \mathbb{C})$  of the positive hyperbolic complex affine sphere with Blaschke metric being the Bers metric  $\mathbf{h}_{(c_1, \bar{c}_2)}$  and Pick tensor  $C = 0$  is the quasi-Fuchsian representation  $\rho(c_1, \bar{c}_2)$ .

**Example 3.25** (Complex Tzitzéica surfaces). The Tzitzéica surface plays an important role in studying high energy harmonic maps to symmetric spaces [DW15]. We can explicitly construct a complex analogue. We won't refer to this example again, but we hope it conveys that one can make a lot of examples of positive hyperbolic complex affine spheres by hand.

Let  $z = z(x, y), w = w(x, y) : \mathbb{R}^2 \rightarrow \mathbb{C}$  be diffeomorphisms, and  $\zeta = e^{2\pi\frac{i}{3}}$  the third root of unity. For each  $c \in \mathbb{C}^*$ , we consider the surface given by,

$$f(x, y) = c(e^{z+\bar{w}}, e^{\zeta^2 z + \zeta \bar{w}}, e^{\zeta z + \zeta^2 \bar{w}}).$$

Note that the image lies inside the complex surface

$$\{z = (z_1, z_2, z_3) \in \mathbb{C}^3 : z_1 z_2 z_3 = c\}.$$

When  $z = w$ , the parametrization becomes

$$f(x, y) = c(e^{\operatorname{Re}z}, e^{\operatorname{Re}z\zeta^{-1}}, e^{\operatorname{Re}z\zeta^{-2}}),$$

and if we assume as well that  $c \in \mathbb{R}$ , then this surface lives in  $\mathbb{R}^3$  and is called the Tzitzéica surface. In this real case, the map  $f$  describes a connected component of the surface in  $\mathbb{R}^3$  defined implicitly by  $xyz = c$ . For this reason, when  $z \neq w$ , we call our surfaces complex Tzitzéica surfaces. We leave it to the reader to compute that  $f$  defines a positive hyperbolic complex affine sphere, and that the Blaschke metric is  $2dzd\bar{w}$  and the Pick tensor is  $C = dz^3 + d\bar{w}^3$ .

#### 4. COMPLEX LIE DERIVATIVES AND TRANSPORT

The space of complex structures on a surface is an infinite-dimensional Banach manifold. Still, its quotient up to isotopy, namely Teichmüller space, is finite-dimensional and the holonomy map factors through it. Generally speaking, taking the quotient up to isotopy is a powerful tool for constructing finite-dimensional configuration spaces while preserving their geometric significance. This is also the case for the moduli space of real hyperbolic affine spheres (discussed in Section 6.1).

Since  $\mathcal{C}(S) \times \mathcal{C}(\bar{S})$  is the complexification of  $\mathcal{C}(S)$ , the natural group acting on  $\mathcal{C}(S) \times \mathcal{C}(\bar{S})$  is the product  $\operatorname{Diff}_0(S) \times \operatorname{Diff}_0(S)$ , which returns a finite dimensional quotient, namely  $\mathcal{T}(S) \times \mathcal{T}(\bar{S})$  (note that if we naively act by  $\operatorname{Diff}_0(S)$  on the left or right or diagonally, we get an infinite dimensional quotient). The map  $(c_1, \bar{c}_2) \mapsto \mathbf{h}_{(c_1, \bar{c}_2)}$  pushes forward the action of  $\operatorname{Diff}_0(S) \times \operatorname{Diff}_0(S)$  onto the space of Bers metrics, but the meaning of the induced quotient is not clear. For instance, if  $\phi \in \operatorname{Diff}_0(S)$ , it is not straightforward to see in what sense or why  $-\frac{4}{(z-\bar{w})^2} dzd\bar{w}$  and  $-\frac{4}{(z-\phi^*(\bar{w}))^2} dz\phi^*(d\bar{w})$  should be identified.

In this section, we define the notions of complex Lie derivative and of transport of tensors through paths of complex vector fields. As we will see, these tools allow us to study the

$\text{Diff}_0(S) \times \text{Diff}_0(S)$  action and will motivate the construction of the finite-dimensional space  $\text{CAS}(S)$  in Section 6.3. Moreover, this approach will be a key tool in proving Theorem G.

**4.1. Complex Lie derivatives.** Let  $M$  be a smooth manifold. Denote by

$$\text{Tens}^{p,q}(M) = \Gamma((TM)^{p\otimes} \otimes (T^*M)^{q\otimes})$$

the space of  $(p, q)$ -type tensors on  $M$ . The Lie derivative along a vector field can be seen as a bilinear map

$$\begin{aligned} \Gamma(TM) \times \text{Tens}^{p,q}(M) &\rightarrow \text{Tens}^{p,q}(M), \\ (X, \alpha) &\mapsto \mathcal{L}_X \alpha. \end{aligned}$$

We extend it  $\mathbb{C}$ -bilinearly to a map

$$\Gamma(\mathbb{C}TM) \times (\text{Tens}^{p,q}(M) \otimes \mathbb{C}) \rightarrow \text{Tens}^{p,q}(M) \otimes \mathbb{C}.$$

That is, we extend by the rule

$$\mathcal{L}_Z \alpha = \mathcal{L}_{\text{Re}Z}(\text{Re}(\alpha)) - \mathcal{L}_{\text{Im}Z}(\text{Im}(\alpha)) + i\mathcal{L}_{\text{Re}Z}(\text{Im}(\alpha)) + i\mathcal{L}_{\text{Im}Z}(\text{Re}(\alpha)).$$

We call  $\mathcal{L}_Z \alpha$  the **complex Lie derivative of  $\alpha$  along the complex vector field  $Z$** .  $\mathcal{L}_Z$  is characterized by the following algebraic properties:

- (1) for all  $C^1$  function  $f: S \rightarrow \mathbb{C}$ ,  $\mathcal{L}_Z(f) = \partial_Z f$ ,
- (2) for all complex vector fields  $W \in \Gamma(\mathbb{C}TM)$ ,  $\mathcal{L}_Z W = [Z, W]$ ,
- (3) for all complex differential forms  $\omega \in \Omega^r(M) \otimes \mathbb{C}$ ,  $\mathcal{L}_Z \omega = d\omega(Z, \cdot) + d(\omega(Z, \cdot))$ ,
- (4) for all  $\alpha, \beta$  smooth complex tensors,  $\mathcal{L}_Z(\alpha \otimes \beta) = (\mathcal{L}_Z \alpha) \otimes \beta + \alpha \otimes (\mathcal{L}_Z \beta)$ ,
- (5) one has the recursive formula for every  $\alpha \in \text{Tens}^{p,q}(M) \otimes \mathbb{C}$  given by

$$\begin{aligned} (\mathcal{L}_Z \alpha)(\omega_1, \dots, \omega_p, W_1, \dots, W_q) &= \partial_Z(\alpha(\omega_1, \dots, \omega_p, W_1, \dots, W_q)) - \alpha(\mathcal{L}_Z \omega_1, \dots, \omega_p, W_1, \dots, W_q) - \dots \\ &\quad - \alpha(\omega_1, \dots, \mathcal{L}_Z \omega_p, W_1, \dots, W_q) - \alpha(\omega_1, \dots, \omega_p, \mathcal{L}_Z W_1, \dots, W_q) - \dots \\ &\quad - \alpha(\omega_1, \dots, \omega_p, W_1, \dots, \mathcal{L}_Z W_q), \end{aligned}$$

where  $\omega_1, \dots, \omega_p \in \Omega^1(\mathbb{C}TM)$ , and  $W_1, \dots, W_q \in \Gamma(\mathbb{C}TM)$ .

**Remark 4.1.** We show a way to define the **complex Lie derivative of a complex structure**  $c_1 \in \mathcal{C}(S)$  for which  $\mathcal{L}_Z c_1 \in T_{c_1} \mathcal{C}(S)$  lies in the kernel of the projection  $\mathcal{C}(S) \rightarrow \mathcal{T}(S)$ .

We define it as follows. Let  $Z \in \Gamma(\mathbb{C}TS)$  and let  $z$  be any local holomorphic coordinate for  $c_1$ , and decompose  $Z$  with respect to the basis  $\{\partial_z, \partial_{\bar{z}}\}$ , hence  $Z = pr_{\partial_z}(Z) + pr_{\partial_{\bar{z}}}(Z)$ . We define

$$\mathcal{L}_Z c_1 := \mathcal{L}_{pr_{\partial_z}(Z) + \overline{pr_{\partial_z}(Z)}} c_1$$

where the RHS is a standard real Lie derivative, corresponding to the infinitesimal deformation through the isotopy  $\phi_t$  with  $\dot{\phi}_0 = pr_{\partial_z}(Z) + \overline{pr_{\partial_z}(Z)}$ .

A motivation for this choice is that

$$\mathcal{L}_Z z = \mathcal{L}_{pr_{\partial_z}(Z)} z = \mathcal{L}_{pr_{\partial_z}(Z) + \overline{pr_{\partial_z}(Z)}} z = \left. \frac{d}{dt} \right|_{t=0} (z \circ \phi_t),$$

and also  $\mathcal{L}_Z dz = \mathcal{L}_{\dot{\phi}_0} dz$ . Moreover, by the properties of the complex Lie derivative, one has that for all  $c_1$ -holomorphic  $n$ -th differential  $\alpha$ ,  $\mathcal{L}_Z \alpha = \mathcal{L}_{\dot{\phi}_0} \alpha$ .

Using the same approach on  $\overline{S}$ , one defines the complex Lie derivative of any element in  $\mathcal{C}(\overline{S})$ .

**Example 4.2.** Here is a key example to motivate the complex Lie derivative. Let  $S$  be a closed surface, let  $\phi_t : [0, 1] \times S \rightarrow S$  be an isotopy, and let  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$  with Bers metric  $\mathbf{h}_{(c_1, \overline{c_2})}$ . Let  $z$  and  $\overline{w}$  be local coordinates for  $c_1$  and  $\overline{c_2}$  respectively, and consider the path of Bers metrics given locally by

$$h_t = \mathbf{h}_{(c_1, \phi_t^*(\overline{c_2}))} = -\frac{4}{(z - \phi_t^*(\overline{w}))^2} dz \cdot (\phi_t^*(d\overline{w})).$$

The variation  $\dot{h}_t = \frac{d}{dt} \Big|_0 h_t$  defines a bilinear form on  $\mathbb{C}TS$ . We claim that this variation can be seen as a complex Lie derivative of  $h_t$ . Indeed, let  $\overline{w}_t := \phi_t^*(\overline{w})$  be a local coordinate for  $\phi_t^*(\overline{c_2})$ , denote  $\dot{\phi}_t =: \gamma_t \partial_z + \overline{\gamma}_t \partial_{\overline{z}}$ , and consider the complex vector field

$$Z_t := \left( \gamma_t \frac{\partial_z \overline{w}_t}{\partial_{\overline{z}} \overline{w}_t} + \overline{\gamma}_t \right) \partial_{\overline{z}}.$$

One can easily check that this description of  $Z_t$  does not depend on the choice of the local holomorphic coordinates  $z$  and  $\overline{w}_t$ , so  $Z_t \in \Gamma(\mathbb{C}TS)$ .

By construction, one has that:

$$\begin{aligned} \mathcal{L}_{Z_t} c_1 &= 0, \\ \mathcal{L}_{Z_t} \phi_t^*(\overline{c_2}) &= \gamma_t \partial_z \overline{w}_t + \overline{\gamma}_t \partial_{\overline{z}} \overline{c_2} = \mathcal{L}_{\dot{\phi}_t} \phi_t^*(\overline{c_2}). \end{aligned}$$

In particular,  $\mathcal{L}_{Z_t} z = 0$ ,  $\mathcal{L}_{Z_t} dz = 0$ ,  $\mathcal{L}_{Z_t} \overline{w}_t = \mathcal{L}_{\dot{\phi}_t} \overline{w}_t$  and  $\mathcal{L}_{Z_t} d\overline{w}_t = \mathcal{L}_{\dot{\phi}_t} d\overline{w}_t$ . In conclusion, by applying the properties of the complex Lie derivative, one gets that

$$\dot{h}_t = \mathcal{L}_{Z_t} h_t.$$

Recall the quotient map  $\mathcal{C}(S) \rightarrow \mathcal{T}(\overline{S})$ , which we label as  $\Psi$ . As Example 4.2 suggests, Lie derivatives through complex vector fields allow to deform simultaneously pairs of complex structures. This is made clearer by the following Proposition which can be seen as a complex extension of the natural description for  $T\mathcal{C}(S)$  through the bundle  $\Psi$ .

**Proposition 4.3.** *Let  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$  and consider the map*

$$d_{c_1} \Psi \oplus d_{\overline{c_2}} \overline{\Psi} : T_{c_1} \mathcal{C}(S) \times T_{\overline{c_2}} \mathcal{C}(\overline{S}) \rightarrow T_{[c_1]} \mathcal{T}(S) \times T_{[\overline{c_2}]} \mathcal{T}(\overline{S}).$$

The map

$$\begin{aligned} \Gamma(\mathbb{C}TS) &\rightarrow T_{c_1} \mathcal{C}(S) \times T_{\overline{c_2}} \mathcal{C}(\overline{S}) \\ Z &\mapsto (\mathcal{L}_Z c_1, \mathcal{L}_Z \overline{c_2}) \end{aligned}$$

defines a linear isomorphism onto the kernel of  $d\Psi_{c_1} \oplus d\overline{\Psi}_{\overline{c_2}}$ .

As a result, one can see

$$T_{(c_1, \overline{c_2})} \mathcal{C}(S) \times \mathcal{C}(\overline{S}) \cong T_{[c_1]} \mathcal{T}(S) \times T_{[\overline{c_2}]} \mathcal{C}(\overline{S}) \times \Gamma(\mathbb{C}TS).$$

*Proof.* Use the notation of Remark 4.1 and Example 4.2. The map is injective: since  $\text{Span}(\partial_{\overline{z}}) \cap \text{Span}(\partial_w) = \{0\}$ ,  $pr_{\partial_z} Z = 0 = pr_{\partial_{\overline{w}}} Z$  implies  $Z = 0$ .

By Remark 4.1, we see that the image of the map is in fact contained in the kernel.

Let  $\dot{\phi}_1, \dot{\phi}_2 \in T_{id} \text{Diff}(S)$ , hence  $\mathcal{L}_{\dot{\phi}_1} c_1 \in T_{c_1} \mathcal{C}(S)$  and  $\mathcal{L}_{\dot{\phi}_2} \overline{c_2} \in T_{\overline{c_2}} \mathcal{C}(\overline{S})$ . As in Example 4.2, one can find  $Z_1, Z_2 \in \Gamma(\mathbb{C}TS)$  such that

$$\begin{aligned} \mathcal{L}_{Z_1} c_1 &= \mathcal{L}_{\dot{\phi}_1} c_1 & \text{and} & & \mathcal{L}_{Z_1} \overline{c_2} &= 0, \\ \mathcal{L}_{Z_1} c_1 &= 0 & \text{and} & & \mathcal{L}_{Z_1} \overline{c_2} &= \mathcal{L}_{\dot{\phi}_2} \overline{c_2}, \end{aligned}$$

hence

$$(\mathcal{L}_{Z_1+Z_2} c_1, \mathcal{L}_{Z_1+Z_2} \overline{c_2}) = (\mathcal{L}_{\dot{\phi}_1} c_1, \mathcal{L}_{\dot{\phi}_2} \overline{c_2}),$$

proving surjectivity.  $\square$

**Corollary 4.4.** *Let  $h_t = \mathbf{h}_{(c_1^t, \overline{c_2^t})}$  be a  $C^1$  path of Bers metrics with constant holonomy, then  $\dot{h}_t = \mathcal{L}_{Z_t} h_t$ , where  $Z_t$  is such that  $(\dot{c}_1^t, \dot{\overline{c_2^t}}) = (\mathcal{L}_{Z_t} c_1^t, \mathcal{L}_{Z_t} \overline{c_2^t})$ .*

*Proof.* Applying the same method as in example 4.2, denoting by  $z_t$  and  $\overline{w_t}$  complex coordinates of  $c_1^t$  and  $\overline{c_2^t}$ , we get

$$\frac{d}{dt} h_t = \frac{8(\partial_{Z_t} z_t - \partial_{Z_t} \overline{w_t})}{(z_t - \overline{w_t})^3} dz_t d\overline{w_t} - \frac{4}{(z_t - \overline{w_t})^2} ((\mathcal{L}_{Z_t} dz) \cdot d\overline{w_t} + dz \cdot (\mathcal{L}_{Z_t} d\overline{w_t})) = \mathcal{L}_{Z_t} h_t.$$

$\square$

**4.2. Transport through complex vector fields.** Unlike real Lie derivatives, realizing complex Lie derivatives as infinitesimal deformations is difficult. More precisely, given a path of complex vector fields  $Z_t \in \Gamma(\mathbb{C}TM)$ ,  $t \in [0, T]$  and a tensor  $\alpha$ , we are interested in solutions  $\alpha_t$  to the system

$$\begin{cases} \mathcal{L}_{Z_t} \alpha_t = \dot{\alpha}_t \\ \alpha_0 = \alpha, \end{cases} \quad (18)$$

and understanding the existence and uniqueness for this Cauchy problem (18) is not in general straightforward. A simple case is as follows: on the unit disk, for  $g(z, t) \partial_{\overline{z}}$  a time-dependent vector field and  $f$  a given function, find  $f_t$  such that

$$\begin{cases} g(z, t) \partial_{\overline{z}} f_t(z) = \frac{d}{dt} f_t(z) \\ f_0(z) = f(z). \end{cases} \quad (19)$$

As we will see below, by the Cauchy-Kovalevskaya and Holmgren theorems, when  $g$  and  $f$  are real analytic, we can find a unique short time solution. But in fact, as can be verified through Hörmander's classical criterion, the problem (19) is not locally solvable, which means one can find  $C_0^\infty$  input data for which no short time solution even exists. For more on local solvability, see [Fol95] and [Tre70].

**Definition 4.5.** We will call a solution to (18) a **transport of  $\alpha$  through  $Z_\bullet$**  and we denote it by  $\alpha_t =: \mathcal{T}_{Z_t} \alpha$ .

We will also use the notation  $\mathcal{T}_{Z_t} c$  to denote paths of complex structures such that  $\frac{d}{dt} \mathcal{T}_{Z_t} c = \mathcal{L}_{Z_t} (\mathcal{T}_{Z_t} c)$ , in the sense of Remark 4.1.

For the rest of this subsection, we assume that one can solve the problem (18), and in the next subsection we discuss existence in the real analytic case.

**Example 4.6.** In fact, the theory of Bers metrics suggests that we shouldn't hope for long time existence of the transport either. Using the unit disk as a toy model, consider a map  $(f_1, \overline{f_2}) : \mathbb{D} \rightarrow \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \setminus \Delta$  for which there exists points  $p, q \in \mathbb{D}$  such that  $f_1(p) = \overline{f_2}(q)$ . Then let  $\phi_t$  be any isotopy of the disk taking  $p$  to  $q$  at a time  $t_0$ . For  $t < t_0$ , we have a family of positive complex metrics given by

$$-\frac{4}{(f_1 - \phi_t^* \overline{f_2})^2} df_1 \cdot (\phi_t^* d(\overline{f_2})),$$

which is tangent to a complex Lie derivative at time zero (see Example 4.2), but which also blows up at  $p$  as  $t$  increases to  $t_0$ .

**Remark 4.7.** By the definition of complex Lie derivative, we can deduce some natural properties of the transport.

(1) If  $\omega \in \Omega^p(S) \otimes \mathbb{C}$  is transportable through  $Z_\bullet$ , then

$$\mathcal{T}_{Z_t}(d\omega) = d(\mathcal{T}_{Z_t}\omega).$$

(2) For all complex tensors  $\alpha, \beta$  which can be transported through  $Z_\bullet$  one has

$$\mathcal{T}_{Z_t}(\alpha \otimes \beta) = (\mathcal{T}_{Z_t}\alpha) \otimes (\mathcal{T}_{Z_t}\beta).$$

(3) If  $\alpha$  in  $Tens^{p,q}(S) \otimes \mathbb{C}$ ,  $W_1, \dots, W_q \in \Gamma(\mathbb{C}TS)$ , and  $\omega_1, \dots, \omega_p \in \Omega^1(\mathbb{C}TS)$  can all be transported through  $Z_\bullet$ , then

$$\mathcal{T}_{Z_t}(\alpha(\omega_1, \dots, \omega_p, W_1, \dots, W_q)) = (\mathcal{T}_{Z_t}\alpha)(\mathcal{T}_{Z_t}(\omega_1), \dots, \mathcal{T}_{Z_t}(\omega_p), \mathcal{T}_{Z_t}(W_1), \dots, \mathcal{T}_{Z_t}(W_q)).$$

**Remark 4.8.** Let  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ , let  $\phi_t^1, \phi_t^2$  be isotopies. Let  $Z_t \in \Gamma(\mathbb{C}TS)$  be such that  $\frac{d}{dt}((\phi_t^1)^*c_1, (\phi_t^2)^*\overline{c_2}) = (\mathcal{L}_{Z_t}c_1^t, \mathcal{L}_{Z_t}\overline{c_2}^t)$  as in Proposition 4.3. Then, for all  $c_1$ -holomorphic (resp.  $\overline{c_2}$ -holomorphic) tensor  $\alpha$ , one has  $(\phi_t^1)^*\alpha = \mathcal{T}_{Z_t}\alpha$  (resp.  $(\phi_t^2)^*\alpha = \mathcal{T}_{Z_t}\alpha$ ). Moreover, by Corollary 4.4,  $\mathbf{h}_{((\phi_t^1)^*c_1, (\phi_t^2)^*\overline{c_2})} = \mathcal{T}_{Z_t}\mathbf{h}_{(c_1, \overline{c_2})}$ .

Here are a few more remarks about the transport through complex vector fields. Observe that the hypothesis of the following proposition holds for  $Z_t \in \Gamma(\mathbb{C}TS)$  corresponding to  $\frac{d}{dt}((\phi_t^1)^*c_1, (\phi_t^2)^*\overline{c_2})$  and for  $\mathcal{T}_t(\mathbf{h}_{(c_1, \overline{c_2})}) = \mathbf{h}_{((\phi_t^1)^*c_1, (\phi_t^2)^*\overline{c_2})}$ .

**Proposition 4.9.** Denote by  $\mathcal{T}_t = \mathcal{T}_{Z_t}$  the transport of tensors through the path of vector fields  $t \mapsto Z_t \in \Gamma(\mathbb{C}TS)$ ,  $t \in [0, T]$ . Assume that  $g$  is a positive complex metric on  $S$  for which  $\mathcal{T}_t g$  exists and is a positive complex metric, and that  $(\mathbf{c}_+, \mathbf{c}_-)(\mathcal{T}_t g) = (\mathcal{T}_t(\mathbf{c}_+(g)), \mathcal{T}_t(\mathbf{c}_-(g)))$ .

- (1) Up to reducing  $T$ , there exists a local orthonormal frame  $(e_1, e_2)$  for  $g$  which admits a transport through  $Z_\bullet$  and such that  $(\mathcal{T}_t e_1, \mathcal{T}_t e_2)$  is orthonormal for  $\mathcal{T}_t g$ .  
(2) If  $\alpha \in Tens^{p,q}(S) \otimes \mathbb{C}$  can be transported through  $Z_\bullet$ ,

$$\mathcal{T}_t(g(\alpha, \alpha)) = (\mathcal{T}_t g)(\mathcal{T}_t \alpha, \mathcal{T}_t \alpha).$$

(3) If  $X, Y \in \Gamma(\mathbb{C}TS)$  can be transported through  $Z_\bullet$ ,

$$\mathcal{T}_t(\nabla_X^g Y) = \nabla_{\mathcal{T}_t X}^{\mathcal{T}_t g}(\mathcal{T}_t Y),$$

where  $\nabla^{\mathcal{T}_t(g)}$  denotes the Levi-Civita connection of  $\mathcal{T}_t(g)$ .

- (4) The curvature changes according to  $\mathcal{T}_t(\mathbf{R}^g) = \mathbf{R}^{\mathcal{T}_t g}$  and  $\mathcal{T}_t(\mathbf{K}_g) = \mathbf{K}_{\mathcal{T}_t g}$ .  
(5) If  $\alpha$  can be transported through  $Z_\bullet$ ,

$$\mathcal{T}_t(\nabla^g \alpha) = \nabla^{\mathcal{T}_t g}(\mathcal{T}_t \alpha).$$

(6) If  $f \in C^r(S, \mathbb{C})$  can be transported through  $Z_\bullet$ , higher-order derivatives change according to

$$\mathcal{T}_t((\nabla^g)^r f) = (\nabla^{\mathcal{T}_t g})^r(\mathcal{T}_t f).$$

In particular, the Hessian and the Laplacian change according to

$$\mathcal{T}_t(\text{Hess}_g(f)) = \text{Hess}_{\mathcal{T}_t g}(\mathcal{T}_t f),$$

$$\mathcal{T}_t(\Delta_g(f)) = \Delta_{\mathcal{T}_t g}(\mathcal{T}_t f).$$

*Proof.* In the proof, we denote  $g_t = \mathcal{T}_t g$  and  $(c_1, \overline{c_2}) = (\mathbf{c}_+, \mathbf{c}_-)(g)$ , with  $z$  and  $\overline{w}$  local holomorphic coordinates for  $c_1$  and  $\overline{c_2}$  respectively. Observe that since  $\mathcal{T}_t c_1$  and  $\mathcal{T}_t \overline{c_2}$  correspond, by Proposition 4.3, to deformation by isotopies, up to shrinking the charts and reducing  $T$ , we can take  $z_t = \mathcal{T}_t z$  and  $\overline{w}_t = \mathcal{T}_t \overline{w}$  local holomorphic coordinates for  $\mathcal{T}_t c_1$  and  $\mathcal{T}_t \overline{c_2}$ .

- (1) We first determine a transportable orthonormal coframe. Use  $g_t$  to denote also the dual bilinear forms on  $\mathbb{C}T^*S$ . By hypothesis,  $g_t(dz_t, dz_t) = g_t(d\bar{w}_t, d\bar{w}_t) = 0$ . It is therefore easy to check that

$$\theta_t^1 = \frac{dz_t + d\bar{w}_t}{\sqrt{2g_t(dz_t, d\bar{w}_t)}} \quad \theta_t^2 = i \frac{dz_t - d\bar{w}_t}{\sqrt{2g_t(dz_t, d\bar{w}_t)}}$$

determine an orthonormal coframe for  $g_t$ , and satisfy  $\frac{d}{dt}\theta_t^j = \mathcal{L}_{Z_t}\theta_t^j$ , hence  $\theta_t^j = \mathcal{T}_t\theta_0^j$ . (Here the square roots exist up to shrinking to a contractible neighborhood.)

Denote with  $(e_1^t, e_2^t)$  the dual frame to  $(\theta_t^1, \theta_t^2)$ , which is therefore  $g_t$ -orthonormal. Then, applying  $\frac{d}{dt} - \mathcal{L}_{Z_t}$  to the equations  $\theta_t^j(e_k^t) = \delta_k^j$ , we get that  $\frac{d}{dt}e_k^t - \mathcal{L}_{Z_t}e_k^t$  lies in the kernel of both  $\theta_t^1$  and  $\theta_t^2$ , therefore it is zero and  $e_k^t = \mathcal{T}_te_k^0$ .

- (2) Let  $(e_1, e_2)$  be a  $g$ -orthonormal frame as in item 1. Denote the dual frame  $(\theta^1, \theta^2)$ , which is therefore such that there exist transports for which  $(\mathcal{T}_t\theta_t^1, \mathcal{T}_t\theta_t^2)$  is an orthonormal frame for  $g_t$ . Then,

$$\begin{aligned} & \mathcal{T}_t \left( g \left( \theta^{m_1} \otimes \theta^{m_p} \otimes e_{n_1} \otimes e_{n_q}, \theta^{m'_1} \otimes \theta^{m'_p} \otimes e_{n'_1} \otimes e_{n'_q} \right) \right) \\ &= \delta_{m_1, m'_1} \cdots \delta_{m_p, m'_p} \cdot \delta_{n_1, n'_1} \cdots \delta_{n_q, n'_q} \\ &= (\mathcal{T}_t g) \left( \mathcal{T}_t(\theta^{m_1} \otimes \cdots \otimes \theta^{m_p} \otimes e_{n_1} \otimes \cdots \otimes e_{n_q}), \mathcal{T}_t(\theta^{m'_1} \otimes \cdots \otimes \theta^{m'_p} \otimes e_{n'_1} \otimes \cdots \otimes e_{n'_q}) \right). \end{aligned}$$

To conclude the proof, observe that applying  $\frac{d}{dt} - \mathcal{L}_{Z_t}$  to the basis decomposition  $\mathcal{T}_t\alpha = \sum_{I,J} a_{I,J}^t \mathcal{T}_t(\theta^I) \otimes \mathcal{T}_t(e_J)$ , with  $I, J$  multi-indices, one gets that  $a_{I,J}^t = \mathcal{T}_ta_{I,J}^0$ , hence the statement follows by linearity.

- (3) Use the notation as in the previous steps. As shown in [BE22], for complex metrics one has a natural notion of Levi-Civita connection forms that extend the Riemannian notion. So the metric  $g_t$  and the orthonormal frame  $(\mathcal{T}_te_1, \mathcal{T}_te_2)$  coframe  $(\mathcal{T}_t\theta^1, \mathcal{T}_t\theta^2)$  induce the forms  $(\omega_i^j)_t$  defined by

$$\nabla^{g_t} \mathcal{T}_t(e_i) = \sum_j (\omega_i^j)_t \otimes \mathcal{T}_t(e_j),$$

and which are characterized as the unique 1-forms such that

$$\begin{cases} \mathcal{T}_t(d\theta^i) = -\sum_j (\omega_j^i)_t \wedge \mathcal{T}_t(\theta^j) \\ (\omega_j^i)_t = -(\omega_i^j)_t. \end{cases}$$

Applying  $\frac{d}{dt} - \mathcal{L}_{Z_t}$  to the equation on top, one gets by Remark 4.7 that

$$\left( \frac{d}{dt} - \mathcal{L}_{Z_t} \right) (\omega_j^i)_t \wedge (\mathcal{T}_t\theta^j) = 0 = - \left( \frac{d}{dt} - \mathcal{L}_{Z_t} \right) (\omega_j^i)_t \wedge (\mathcal{T}_t\theta^i),$$

hence  $(\frac{d}{dt} - \mathcal{L}_{Z_t})(\omega_j^i)_t = 0$ , thus  $\mathcal{T}_t(\nabla^g e_j) = \nabla^{\mathcal{T}_t g}(\mathcal{T}_t e_j)$  for all  $j$ . Finally if  $X \in \Gamma(CTS)$  admits a transport through  $Z_\bullet$ , and  $X = \sum_j a_j e_j$ , we obtain once again that  $\mathcal{T}_t X = \sum_j \mathcal{T}_t a_j \mathcal{T}_t e_j$  and

$$\begin{aligned} \mathcal{T}_t(\nabla^g X) &= \mathcal{T}_t \left( \sum_j da_j \otimes e_j + \sum_j a_j \otimes \nabla^g e_j \right) \\ &= \sum_j d(\mathcal{T}_t a_j) \otimes (\mathcal{T}_t e_j) + \sum_j \mathcal{T}_t(a_j) \otimes \nabla^{\mathcal{T}_t(g)}(\mathcal{T}_t e_j) = \nabla^{\mathcal{T}_t(g)} \mathcal{T}_t(X). \end{aligned}$$

which concludes the proof.

(4) The proof follows easily by applying iteratively the previous step to the tensor

$$R^g(X, Y)Z = \nabla_X^g \nabla_Y^g Z - \nabla_Y^g \nabla_X^g Z - \nabla_{\nabla_X^g Y - \nabla_Y^g X}^g Z.$$

With the notation as in the previous steps, we also see that

$$\mathcal{T}_t(K_g) = \mathcal{T}_t(g(R_g(e_1, e_2)e_2, e_1)) = (\mathcal{T}_t g)(R_{\mathcal{T}_t g}(\mathcal{T}_t e_1, \mathcal{T}_t e_2)\mathcal{T}_t e_2, \mathcal{T}_t e_1) = K_{(\mathcal{T}_t g)}.$$

(5) By the first item and Remark 4.7, the statement is true if  $\alpha$  is a 1-form since

$$(\mathcal{T}_t(\nabla^g \alpha))(\mathcal{T}_t e_j) = \mathcal{T}_t((\nabla^g \alpha)e_j) = \mathcal{T}_t(d(\alpha(e_j))) - \mathcal{T}_t(\alpha(\nabla^g e_j)) = (\nabla^{\mathcal{T}_t(g)} \mathcal{T}_t(\alpha))(\mathcal{T}_t e_j).$$

The proof then follows for all  $(p, q)$ -type real analytic tensors  $\alpha$  using the formula

$$\begin{aligned} (\nabla^g \alpha)(\omega_1, \dots, \omega_p, Y_1, \dots, Y_q) &= d(\alpha(\omega_1, \dots, \omega_p, W_1, \dots, W_q)) - \alpha(\nabla^g \omega_1, \dots, \omega_p, W_1, \dots, W_q) - \dots - \\ &\quad - \alpha(\omega_1, \dots, \nabla^g \omega_p, W_1, \dots, W_q) - \alpha(\omega_1, \dots, \omega_p, \nabla^g W_1, \dots, W_q) - \dots - \\ &\quad - \alpha(\omega_1, \dots, \omega_p, W_1, \dots, \nabla^g W_q) \end{aligned}$$

for  $\omega_k \in \{\theta^1, \theta^2\}$  and  $W_k \in \{e_1, e_2\}$ .

(6) The statement is true for  $r = 1$ :

$$g_t(\mathcal{T}_t(\nabla^g f), \mathcal{T}_t e_k) = \mathcal{T}_t(g(\nabla^g f, \mathcal{T}_t e_k)) = \mathcal{T}_t(df(Y)) = df_t(\mathcal{T}_t e_k) = g_t(\nabla^{g_t} f_t, \mathcal{T}_t e_k).$$

where  $g_t = \mathcal{T}_t(g)$ ,  $f_t = \mathcal{T}_t(f)$ ,  $Y_t = \mathcal{T}_t(Y)$ . The proof then follows directly by induction on  $r$ .

To see the equality for the Laplace-Beltrami operator, one can just compute that

$$\mathcal{T}_t(\Delta_g f) = \sum_j \mathcal{T}_t(\text{Hess}_g(f)(e_j, e_j)) = \sum_j \text{Hess}_{\mathcal{T}_t g}(\mathcal{T}_t f)(\mathcal{T}_t e_j, \mathcal{T}_t e_j) = \Delta_{(\mathcal{T}_t g)}(\mathcal{T}_t f).$$

□

**Remark 4.10.** The condition on the conformal type of the complex metric is used only in the proof of the first item of the Proposition. In fact, one can prove the other items for complex metrics on manifolds provided that an analog of step 1 is assumed as a hypothesis, namely that there exist local orthonormal frames that admit transports that are orthonormal for the transported metric.

**4.3. Transport of real analytic tensors.** A useful class of cases for which the problem (18) admits a unique solution is for the transport of real analytic complex tensors through real analytic complex vector fields. Real analytic objects on  $M$  have some nice density properties, which extend some results also more generally in the smooth setting. Here we recall a few results:

- The space of real analytic sections of a real analytic bundle over  $M$  is dense in the space of sections (see [KM97, §30.12]). In particular, real analytic complex vector fields are dense in the space of complex vector fields.
- The space  $C^\omega(M, M)$  of real analytic mappings from  $M$  to  $M$  is locally modeled on the space of real analytic vector fields by locally applying the inverse of the exponential map for any real analytic Riemannian metric on  $M$  (see [KM97, §37]). In particular, it is locally path-connected through paths of the form  $t \mapsto \phi_t$  such that  $t \mapsto \phi_t(p)$  is piecewise analytic for all  $p \in M$ .
- The space  $\text{Diff}^\omega(M)$  of analytic diffeomorphisms of  $M$  is an open subset of  $C^\omega(M, M)$ , and it is therefore locally path-connected through paths of the same type.
- $\text{Diff}^\omega(M)$  is dense in  $\text{Diff}(M)$  and the inclusion is a homotopy equivalence (see [Tsu09]). In particular,  $\text{Diff}_0^\omega(M)$  is dense in  $\text{Diff}_0(M)$ .

As we will see, the following theorem will be very useful also in the non-analytic setting because of the nice density properties above.

**Theorem 4.11.** *Let  $M$  be a real analytic manifold. Let  $t \mapsto Z_t$  be a path of (local) real analytic complex vector fields on  $M$  that is piecewise real analytic with respect to  $t \in [0, T]$ . Let  $\alpha$  be a real analytic (local) complex tensor on  $M$ .*

*Then, for all  $p \in M$  there exists a neighborhood  $U_p \subset M$ ,  $T_0 \in (0, T]$ , and a unique path of tensors  $t \mapsto \alpha_t$ ,  $t \in [0, T_0]$ , such that on  $U_p$*

$$\begin{cases} \dot{\alpha}_t = \mathcal{L}_{Z_t} \alpha_t \\ \alpha_0 = \alpha. \end{cases}$$

*A posteriori the path  $t \mapsto \alpha_t$  is piecewise real analytic with respect to  $t$  and each  $\alpha_t$  is a real analytic tensor.*

*Moreover, if  $M$  is compact, one can take  $M = U_p$ .*

*Proof.* This theorem is mainly an application of the Cauchy-Kovalevskaya Theorem which allows to find the unique real analytic solution. Uniqueness then follows by Holmgren's theorem, which says that any solution to the problem must be real analytic.

*Step 1.* We first prove the statement for a real analytic function  $\alpha$ . In a real analytic chart  $(x_1, \dots, x_n)$  centered in  $p$ , solving the equation  $\dot{\alpha}_t = \mathcal{L}_{Z_t} \alpha_t$  with prescribed  $\alpha_0$  is equivalent to solving the Cauchy problem (with real coefficients)

$$\begin{cases} \dot{u}_1^t = \sum_j a_j^t \partial_{x_j} u_1^t - b_j^t \partial_{x_j} u_2^t + a_j^t \partial_{x_j} \operatorname{Re}(\alpha_0) - b_j^t \partial_{x_j} \operatorname{Im}(\alpha_0) \\ \dot{u}_2^t = \sum_j b_j^t \partial_{x_j} u_1^t + a_j^t \partial_{x_j} u_2^t + b_j^t \partial_{x_j} \operatorname{Re}(\alpha_0) + a_j^t \partial_{x_j} \operatorname{Im}(\alpha_0) \\ u_1^0 = u_2^0 = 0, \end{cases}$$

where  $u_1^t + iu_2^t := \alpha_t - \alpha_0$ , and  $Z_t = \sum_j (a_j^t + ib_j^t) \partial_{x_j}$ .

By the Cauchy-Kovalevskaya Theorem, there exists a unique real analytic solution  $(u_1^t, u_2^t)$  to this problem in a neighborhood  $U_0$  of the point and for small enough time.

*Step 2.* We prove existence for tensors. Consider again a local analytic chart  $(x_1, \dots, x_n)$  around  $p$ . By the previous step, there exists a neighborhood  $U_p$  contained in the chart and  $T_0 > 0$  such that for all  $t \in [0, T_0]$ :

- there exists a unique  $x_j^t: U_p \rightarrow \mathbb{C}$  such that  $\frac{d}{dt} x_j^t = \partial_{Z_t} x_j^t$  and  $x_j^0 = x_j$ ,
- $dx_1^t, \dots, dx_n^t$  are  $\mathbb{C}$ -linearly independent 1-forms.

By the properties of the complex Lie derivative,  $\mathcal{L}_{Z_t} dx_j^t = \frac{d}{dt} (dx_j^t)$ . Now, denote by  $e_1^t, \dots, e_n^t \in \Gamma(\mathbb{C}TU_p)$  the dual frame corresponding to  $dx_1^t, \dots, dx_n^t$ . Then,  $e_j^0 = \partial_{x_j}$ , and

$$\begin{aligned} 0 &= \mathcal{L}_{Z_t} (dx_j^t(e_k^t)) = (\mathcal{L}_{Z_t} dx_j^t)(e_k^t) + dx_j^t(\mathcal{L}_{Z_t} e_k^t), \\ 0 &= \frac{d}{dt} (dx_j^t(e_k^t)) = d\dot{x}_j^t(e_k^t) + dx_j^t(\dot{e}_k^t) = (\mathcal{L}_{Z_t} dx_j^t)(e_k^t) + dx_j^t(\dot{e}_k^t), \end{aligned}$$

so  $\frac{d}{dt} (e_k^t) = \mathcal{L}_{Z_t} e_k^t$ .

Finally, let  $I = (i_1, \dots, i_p)$ ,  $J = (j_1, \dots, j_q)$  be multindices. Denote  $e_I^t = e_{i_1}^t \otimes \dots \otimes e_{i_p}^t$  and  $dx_J^t = dx_{j_1}^t \otimes \dots \otimes dx_{j_q}^t$ . One can easily check that  $\frac{d}{dt} e_I^t = \mathcal{L}_{Z_t} e_I^t$  and  $\frac{d}{dt} dx_J^t = \mathcal{L}_{Z_t} dx_J^t$ .

Finally, for any analytic tensor  $\alpha_0 = \sum_{I, J} \alpha_{I, J} \otimes dx_J$  tensor, let  $\alpha_{I, J}^t$  be the unique function defined for small times in a neighborhood of  $p_0$  such that  $\dot{\alpha}_{I, J}^t = \partial_{Z_t} \alpha_{I, J}^t$ ; then it is straightforward to see, using the properties of the complex Lie derivative, that

$$\frac{d}{dt} \alpha_t = \mathcal{L}_{Z_t} \alpha_t,$$



with  $\alpha_t = \sum_{I,J} \alpha_{I,J}^t e_I^t \otimes dx_J^t$ .

*Step 3.* We prove uniqueness. Assume that  $\alpha_t$  and  $\beta_t$  satisfy  $\dot{\alpha}_t = \mathcal{L}_{Z_t} \alpha_t$ ,  $\dot{\beta}_t = \mathcal{L}_{Z_t} \beta_t$  and  $\alpha_0 = \beta_0$ . With respect to the bases above we can write  $\alpha_t = \sum_{I,J} \alpha_{I,J}^t e_I^t \otimes dx_J^t$  and  $\beta_t = \sum_{I,J} \beta_{I,J}^t e_I^t \otimes dx_J^t$  for some functions  $\alpha_{I,J}^t$  and  $\beta_{I,J}^t$ . Using the properties of the complex Lie derivative, we deduce that  $\frac{d}{dt} \alpha_{I,J}^t = \partial_{Z_t} \alpha_{I,J}^t$  and  $\frac{d}{dt} \beta_{I,J}^t = \partial_{Z_t} \beta_{I,J}^t$  with  $\alpha_{I,J}^0 = \beta_{I,J}^0$ , hence  $\alpha_{I,J}^t = \beta_{I,J}^t$  by Step 1 and by the Cauchy-Kovalevskaya Theorem.

*Step 4.* If  $M$  is compact, one obtains a solution for small time in a neighborhood of any given point, and the solutions coincide on the intersections of such neighborhoods by uniqueness. Extracting a finite subcover from such neighborhoods, one can find analytic solutions on the whole  $M$  for all  $t \in [0, T']$  for some  $T' > 0$  small enough.  $\square$

**Corollary 4.12.** *Let  $M$  be a real analytic manifold. Let  $t \mapsto Z_t$  be a path of (local) real analytic complex vector fields on  $M$  that is piecewise real analytic with respect to  $t \in [0, T]$ .*

*Then, for all  $C^1$  tensors  $\alpha$ , the problem (18) has at most one solution.*

*Proof.* It follows by the linearity of the equation and by the fact that by Theorem 4.11,  $\alpha_t \equiv 0$  is the unique solution for  $\alpha = 0$ .  $\square$

We can therefore transport real analytic complex affine spheres.

**Proposition 4.13.** *Let  $\sigma: \tilde{S} \rightarrow \mathbb{C}^3$  be an equivariant positive hyperbolic complex affine sphere with analytic Blaschke metric  $g$  and Pick tensor  $C = Q_1 + \overline{Q}_2$ . Let  $(\mathbf{c}_+, \mathbf{c}_-)(g) =: (c_1, \overline{c}_2) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ ,  $Q_1 \in \text{CD}(c_1)$ , and  $\overline{Q}_2 \in \text{CD}(\overline{c}_2)$ .*

*Let  $\phi_t \in \text{Diff}_0^\omega(S)$  be an analytic isotopy. Then, there exists  $T_0$ , and a real analytic vector field  $Z_t \in \Gamma(\mathbb{C}TS)$ ,  $t \in [0, T_0]$  such that: for all  $t \in [0, T_0]$ ,  $g_t = \mathcal{I}_{Z_t} g$  is a positive complex metric with  $(\mathbf{c}_+, \mathbf{c}_-)(g_t) = (c_1, \phi_t^*(\overline{c}_2))$ , and there exists an equivariant positive hyperbolic complex affine sphere with Blaschke metric  $g_t$  and Pick tensor  $C_t = Q_1 + \phi_t^*(\overline{Q}_2)$ .*

*Moreover, the holonomy is constant.*

*Proof.* Since  $g$  is real analytic, its isotropic directions vary in a real analytic way on the surface, hence  $c_1$  and  $\overline{c}_2$  are real analytic.

Let  $z$  and  $\overline{w}$  be local holomorphic coordinates for  $c_1$  and  $\overline{c}_2$ . Choose  $T_0$  such that all the following transports are well-defined.

As in Example 4.2, denoting  $\phi_t = \gamma_t \partial_z + \overline{\gamma}_t \partial_{\overline{z}}$ , one has that  $Z_t = \left( \gamma_t \frac{\partial_z \overline{w}_t}{\partial_{\overline{z}} \overline{w}_t} + \overline{\gamma}_t \right) \partial_{\overline{z}}$  is such that  $\mathcal{I}_{Z_t} dz = dz$  and  $\mathcal{I}_{Z_t} d\overline{w} = \phi_t^*(d\overline{w})$ . As a result, if  $g = \lambda dz d\overline{w}$ , then  $g_t = \mathcal{I}_{Z_t} g = (\mathcal{I}_{Z_t} \lambda) \cdot dz \cdot (\phi_t^* d\overline{w})$ , so  $(\mathbf{c}_+, \mathbf{c}_-)(g_t) = (c_1, \phi_t^*(\overline{c}_2))$ .

By Proposition 4.9,  $g_t$  and  $C_t$  satisfy Equation (14), hence by Theorem 3.20 we conclude that there exists an equivariant positive hyperbolic complex affine sphere with Blaschke metric  $g_t$  and Pick tensor  $C_t$ .

To prove that the holonomy stays constant, we need to look at the construction of the corresponding flat connection as in the proof of Theorem 3.13. Let  $N: \mathbb{C} \times \tilde{S} \rightarrow \tilde{S}$  be the trivial bundle with generator  $\hat{\xi} \equiv 1$  and let  $E = \mathbb{C}T\tilde{S} \oplus N$ . Then the data  $g_t$  and  $C_t$  as above define a flat connection

$$\begin{cases} \hat{D}_X^t Y &= \nabla_X^t Y + g_t(X, Y) \hat{\xi} \\ \hat{D}_X^t \hat{\xi} &= X \\ \hat{D}_X^t (i\hat{\xi}) &= iX, \end{cases}$$

where  $\nabla^t$  is the affine connection on  $\mathbb{C}TS$  defined by  $C_t =: g_t(\nabla^t - \nabla^{g_t}, \cdot)$ . Since  $\dot{C}_t = \mathcal{L}_{Z_t} C_t$  and  $\dot{g}_t = \mathcal{L}_{Z_t} g_t$ , one can immediately see that  $\mathcal{L}_{Z_t} \nabla^t = \dot{\nabla}^t$ , where we denote  $(\mathcal{L}_{Z_t} \nabla^t)_X Y = \mathcal{L}_{Z_t}(\nabla_X^t Y) - \nabla_{\mathcal{L}_{Z_t} X}^t Y - \nabla_X^t(\mathcal{L}_{Z_t} Y)$ .

For a general section  $s = X + b\hat{\xi}$  of the bundle  $E$ , denote  $\mathcal{L}_{Z_t} s = \mathcal{L}_{Z_t}(X) + (\mathcal{L}_{Z_t} b)\hat{\xi}$ . Fix a section  $s_0 = X_0 + b_0\hat{\xi}$  such that  $D^0 s_0 = 0$ . Then, since the  $D^t$ 's are analytic, both  $X_0$  and  $b_0$  are analytic, and denoting

$$s_t := \mathcal{I}_{Z_t} X_0 + (\mathcal{I}_{Z_t} b_0)\hat{\xi},$$

a straightforward computation shows that

$$\frac{d}{dt}(D^t s_t) = \dot{D}^t s_t + D^t \dot{s}_t = \mathcal{L}_{Z_t}(D^t s_t).$$

Hence, since  $D^0 s_0 = 0$ , applying Theorem 4.11 we conclude by uniqueness that  $D^t s_t = 0$  for all  $t$ , hence  $s_t$  is  $D^t$ -parallel.

Finally, fixing a basis of  $D^0$ -parallel sections  $(s_0^k)$ , we have that  $(s_t^k)$  with  $s_t^k = \mathcal{I}_{Z_t} s_0^k$  is a basis of  $D^t$ -parallel sections. The holonomy of  $D^t$ , and of the corresponding complex affine sphere is given by the matrices  $(m_t^{jk}(\gamma))_{jk}$  where  $\gamma \in \pi_1(S)$  and

$$\gamma^* s_t^j = \sum_k m_t^{jk}(\gamma) s_t^k.$$

Since  $Z_t$  is  $\pi_1(S)$ -invariant we have

$$\frac{d}{dt}(\gamma^* s_t^j) = \mathcal{L}_{Z_t}(\gamma^* s_t^j) = \sum_k m_t^{jk}(\gamma) \mathcal{L}_{Z_t}(s_t^k) = \sum_k m_t^{jk}(\gamma) \frac{d}{dt} s_t^k,$$

from which we conclude that  $\frac{d}{dt}(m_t^{jk}(\gamma)) = 0$ , hence the holonomy is constant.  $\square$

## 5. COMPLEX ELLIPTIC OPERATORS

Here we derive the local expression for the Laplacian of a complex metric, investigate the analytic properties of our complex elliptic operators, and prove Theorems F and G.

We keep the following notation. For a constant  $C$  depending on quantities  $a_1, \dots, a_n$ , we write  $C(a_1, \dots, a_n)$ . In this notation, the value of such a constant may change in the course of a proof. Analogous to Section 2, for  $U$  a domain inside  $\mathbb{C}$ , or the surface  $S$ , we denote  $L^p$  spaces and Sobolev spaces by  $L^p(U)$  and  $W^{k,p}(U)$  respectively, and local spaces by  $L_{loc}^p(U)$  and  $W_{loc}^{k,p}(U)$ . We use  $\|\cdot\|_{L^p(U)}$ ,  $\|\cdot\|_{W^{k,p}(U)}$  to denote the norm. When  $U = \mathbb{C}$ , we'll just write  $\|\cdot\|_p$  or  $\|\cdot\|_{k,p}$ . We view Beltrami forms defined on domains in  $\mathbb{C}$  as functions with respect to the standard coordinate  $z$  and use the terms function and form interchangeably. We'll often work locally in disks inside  $\mathbb{C}$  so for this we set the notation, for  $r > 0$ ,  $\mathbb{D}_r = \{z \in \mathbb{C} : |z| < r\}$ , and  $\mathbb{D}_1 = \mathbb{D}$ .

**5.1. The Laplacian of a complex metric.** Throughout this subsection, let  $z$  and  $w$  be coordinates on the unit disk  $\mathbb{D}$ . We view  $z$  as the standard coordinate and  $w$  as a  $C^2$  reparametrization of  $\mathbb{D}$ ,  $w = w(z)$ . Let  $\mu$  denote the Beltrami form, defined by  $\mu = \frac{\partial_{\bar{z}} w}{\partial_z w}$ , which satisfies  $|\mu| < 1$ . Let  $g$  be a  $C^2$  positive complex metric on  $\mathbb{D}$  that is locally of the form  $g(z) = \lambda(z) dz d\bar{w}$ , where  $\lambda \in C^2(\mathbb{D}, \mathbb{C}^*)$ . We compute the Laplacian as a function of  $z$ . First we record a lemma. Let  $\nabla^g$  be the Levi-Civita connection of  $g$ .

**Lemma 5.1.**  $\nabla_{\partial_w}^g \partial_{\bar{z}} = \partial_w z \cdot \partial_{\bar{z}} \bar{\mu} \cdot \partial_{\bar{z}}$ .

*Proof.* Recall from Lemma 2.6 that if  $\Pi_1$  is the projections of  $\mathbb{C}T\mathbb{D}$  onto the the line spanned by  $\partial_{\bar{z}}$ , with kernel spanned by  $\partial_w$ , then  $\nabla_{\partial_w}^g \partial_{\bar{z}} = \Pi_1([\partial_w, \partial_{\bar{z}}])$ . As well, note that  $\frac{\partial_w \bar{z}}{\partial_w z} = -\bar{\mu}$ , as can be seen from the expression

$$0 = \partial_{\bar{w}} w = \partial_{\bar{w}} \bar{z} \partial_{\bar{z}} w + \partial_{\bar{w}} z \partial_z w.$$

Thus, writing out

$$\partial_w = \partial_w z \cdot \partial_z + \partial_w \bar{z} \cdot \partial_{\bar{z}} = \partial_w z (\partial_z - \bar{\mu} \partial_{\bar{z}}), \quad (20)$$

we get that

$$\begin{aligned} [\partial_w, \partial_{\bar{z}}] &= \partial_{\bar{z}}(\partial_w z \bar{\mu}) \cdot \partial_{\bar{z}} - \partial_{\bar{z}} \partial_w z \cdot \partial_z \\ &= \partial_{\bar{z}}(\partial_w z \bar{\mu}) \cdot \partial_{\bar{z}} - \frac{\partial_{\bar{z}} \partial_w z}{\partial_w z} \partial_w - \partial_{\bar{z}} \partial_w z \bar{\mu} \partial_{\bar{z}} \\ &= \partial_w z \partial_{\bar{z}} \bar{\mu} \partial_{\bar{z}} - \frac{\partial_{\bar{z}} \partial_w z}{\partial_w z} \partial_w, \end{aligned}$$

so  $\nabla_{\partial_w}^g \partial_{\bar{z}} = \partial_w z \partial_{\bar{z}} \bar{\mu} \partial_{\bar{z}}$ .  $\square$

**Proposition 5.2.** *In the holomorphic coordinate  $z$ , the Laplacian  $\Delta_g$  is expressed as*

$$\Delta_g = \frac{4}{\lambda \partial_z w} \partial_{\bar{z}} (\partial_z - \bar{\mu} \partial_{\bar{z}}).$$

*Proof.* In the  $g$ -isotropic basis  $\{\partial_{\bar{z}}, \partial_w\}$  for the complexified tangent bundle  $\mathbb{C}T\mathbb{D}$ , for all  $C^2$  functions  $f: \mathbb{D} \rightarrow \mathbb{C}$ ,

$$\Delta_g f = \frac{2}{g(\partial_w, \partial_{\bar{z}})} \text{Hess}_g(f)(\partial_w, \partial_{\bar{z}}) = \frac{2}{g(\partial_w, \partial_{\bar{z}})} (\partial_w \partial_{\bar{z}} f - (\nabla_{\partial_w}^g \partial_{\bar{z}}) f).$$

Hence, by Lemma 5.1 and Equation (20),

$$\begin{aligned} \Delta_g &= \frac{2}{g(\partial_w, \partial_{\bar{z}})} (\partial_w \partial_{\bar{z}} - (\partial_w z)(\partial_{\bar{z}} \bar{\mu}) \partial_{\bar{z}}) \\ &= \frac{2 \partial_w z}{g(\partial_w, \partial_{\bar{z}})} (\partial_z \partial_{\bar{z}} - \bar{\mu} \partial_{\bar{z}} \partial_{\bar{z}} - \partial_{\bar{z}} \bar{\mu} \partial_{\bar{z}}) \\ &= \frac{4}{\lambda \partial_z w} (\partial_{\bar{z}} (\partial_z - \bar{\mu} \partial_{\bar{z}})). \end{aligned}$$

$\square$

**Remark 5.3.** Observe that, from Equation (20), we equivalently have

$$\Delta_g = \frac{4}{\lambda \partial_z w} \partial_{\bar{z}} \left( \frac{1}{\partial_w z} \partial_w \right).$$

We'll use this particular expression in Section 7.

**5.2. Interior elliptic estimates.** Here we deduce interior elliptic estimates for complex Laplacians via estimates for the Beltrami operator. We make reference to the **Beurling transform**  $T$ , defined on  $C_0^\infty(\mathbb{C})$  by the principal value integral

$$T(u)(z) = \lim_{\varepsilon \rightarrow 0} -\frac{1}{\pi} \int_{|\zeta - z| > \varepsilon} \frac{u(\zeta)}{(\zeta - z)^2} dx dy.$$

Via the Calderon-Zygmund theory, the continuity properties of  $T$  are well understood.

**Proposition 5.4.** *For all  $1 < p < \infty$ ,  $T$  extends to a continuous linear operator from  $L^p(\mathbb{C}) \rightarrow L^p(\mathbb{C})$ , which is an isometry when  $p = 2$ . The operator norm of  $T: L^p(\mathbb{C}) \rightarrow L^p(\mathbb{C})$  varies continuously with  $p$ .*

For a source, see [IT92, Chapter 4].

Henceforth, we set  $C_p$  to be the operator norm of  $T : L^p(\mathbb{C}) \rightarrow L^p(\mathbb{C})$ . As in the previous subsection, let  $g$  be a positive complex metric on  $\mathbb{D}$  of the form  $g = \lambda dz d\bar{w}$ , and let  $\mu$  be the Beltrami form of  $w$ . We now choose  $k > 2$  and  $l > k + 2$ . We assume that  $\mu$  is  $W^{l,\infty}$ . We use  $W^{k+2,p}$  for the Sobolev spaces on which our linear maps act. For a linear differential operator  $L$ , we define  $\|L\|_{W^{k,p}(U)}$  to be the sum of the  $W^{k,p}(U)$  norms of the coefficients, and we say that  $L$  is  $W^{k,p}$  (resp.  $W_{loc}^{k,p}$ ) if all of the coefficients are  $W^{k,p}$  (resp.  $W_{loc}^{k,p}$ ). The main estimate is as follows.

**Proposition 5.5.** *Let  $L$  be a  $W_{loc}^{k+1,\infty}$  linear second order operator with the same principle symbol as  $\Delta_g$ . Let  $u \in L_{loc}^p(\mathbb{C})$  and  $v \in W_{loc}^{k,p}(\mathbb{C})$ , with  $u$  satisfying  $Lu = v$  in the weak sense. Then  $u \in W_{loc}^{k+2,p}(\mathbb{C})$  and for all  $r < R < \infty$  and  $p$  such that  $\|\mu\|_\infty C_p < 1$ , there exists  $C = C(\|L\|_{W^{k+1,\infty}(\mathbb{D}_R)}, \|\mu\|_{W^{k+1,\infty}(\mathbb{D}_R)}, r, R, k, p)$  such that*

$$\|u\|_{W^{k+2,p}(\mathbb{D}_r)} \leq C(\|u\|_{W^{k,p}(\mathbb{D}_R)} + \|v\|_{W^{k,p}(\mathbb{D}_R)}).$$

**Remark 5.6.** If  $(c_1, \bar{c}_2)$  are  $W^{l,\infty}$  complex structures, then the associated Bers metric  $h$  is  $W^{l,2}$ , as discussed around Proposition 2.9. For  $l > k + 2$ , by Morrey's inequality,  $h$  is  $k + 1$  times differentiable with Hölder continuous  $(k + 1)^{th}$ -derivatives, hence  $\Delta_h$  is  $W^{k+1,\infty}$ , and the hypothesis of Proposition 5.5 applies to  $L = \Delta_h$ .

Since the Laplacian factors as a product of first order complex elliptic operators, Proposition 5.5 essentially follows from the interior elliptic estimates for the Beltrami operator, stated below.

**Proposition 5.7.** *Let  $\mu \in W_{loc}^{k,\infty}(\mathbb{C})$ ,  $u \in L_{loc}^p(\mathbb{C})$ , and  $v \in W_{loc}^{k,p}(\mathbb{C})$ , with  $u$  satisfying*

$$\left(\partial_{\bar{z}} - \mu\partial_z\right)u = v \tag{21}$$

*in the weak sense. Then  $u \in W_{loc}^{k+1,p}$ , and for all  $r < R < \infty$  and  $p$  such that  $\|\mu\|_\infty C_p < 1$ , there exists  $C = C(\|\mu\|_{W^{k,\infty}(\mathbb{D}_R)}, r, R, k, p)$  such that*

$$\|u\|_{W^{k+1,p}(\mathbb{D}_r)} \leq C(\|u\|_{W^{k,p}(\mathbb{D}_R)} + \|v\|_{W^{k,p}(\mathbb{D}_R)}).$$

Proposition 5.7 is certainly well-known to experts, although we couldn't locate a precise reference. We provide the proof in Appendix A (although at some point we refer to [Nic07] instead of repeating a standard argument). For reference, in Lemmas 5.21 and 5.22 of [AIM09], it is explained how to obtain higher regularity for  $u \in W^{1,p}$ , assuming  $\mu, v \in C_0^\infty(\mathbb{C})$ . As well, Schauder estimates for the Beltrami equation are proved in Chapter 15 of [AIM09]. We need elliptic estimates in particular over Schauder estimates for the proof of Theorem F and the proof of Proposition 2.9.

*Proof of Proposition 5.5.* We first assume that  $L = \Delta_g$ . By a routine density and reflexivity argument (see [Nic07, Corollary 10.3.10]), it suffices to prove the estimate of Proposition 5.5 under the assumption  $u$  is smooth. Note as well that, by complex conjugation, we obtain the conclusion of Proposition 5.7 for the conjugate Beltrami operator  $\left(\partial_z - \bar{\mu}\partial_{\bar{z}}\right)$  as well (we stated the estimate for the ordinary Beltrami operator since that is more familiar and we'll use that result, too, later on). The hypothesis  $\Delta_g u = v$  means that  $u' := \left(\partial_z - \bar{\mu}\partial_{\bar{z}}\right)u$  satisfies  $\partial_{\bar{z}}u' = \lambda\overline{\partial_z wv}$ . Let  $0 < r < s < R$ . By elliptic regularity for  $\partial_{\bar{z}}$ , we obtain

$$\begin{aligned} \|u'\|_{W^{k+1,p}(\mathbb{D}_s)} &\leq C(s, R, k)(\|u'\|_{W^{k,p}(\mathbb{D}_R)} + \|\lambda\overline{\partial_z wv}\|_{W^{k,p}(\mathbb{D}_R)}) \\ &\leq C(\|L\|_{W^{k,\infty}(\mathbb{D}_R)}, s, R, k)(\|u'\|_{W^{k,p}(\mathbb{D}_R)} + \|v\|_{W^{k,p}(\mathbb{D}_R)}). \end{aligned}$$

Note that  $\|u'\|_{W^{k,p}(\mathbb{D}_R)} \leq C(\|\mu\|_{W^{k,\infty}(\mathbb{D}_R)})\|u\|_{W^{k+1,p}(\mathbb{D}_R)}$ , so we can turn the inequality above into

$$\|u'\|_{W^{k+1,p}(\mathbb{D}_s)} \leq C(\|L\|_{W^{k,\infty}(\mathbb{D}_R)}, s, R, k)(\|u\|_{W^{k+1,p}(\mathbb{D}_R)} + \|v\|_{W^{k,p}(\mathbb{D}_R)}). \quad (22)$$

Now, applying Proposition 5.7 to the equation  $(\partial_z - \bar{\mu}\partial_{\bar{z}})u = u'$ , we have

$$\|u\|_{W^{k+2,p}(\mathbb{D}_r)} \leq C(\|\mu\|_{W^{k+1,\infty}(\mathbb{D}_R)}, s, r, k, p)(\|u\|_{W^{k+1,p}(\mathbb{D}_s)} + \|u'\|_{W^{k+1,p}(\mathbb{D}_s)}). \quad (23)$$

Composing the estimates (22) and (23), we arrive at

$$\|u\|_{W^{k+2,p}(\mathbb{D}_r)} \leq C(\|L\|_{W^{k+1,\infty}(\mathbb{D}_R)}, r, R, k, p)(\|u\|_{W^{k+1,p}(\mathbb{D}_R)} + \|v\|_{W^{k,p}(\mathbb{D}_R)}). \quad (24)$$

We want to adjust the constants in order to replace the  $\|u\|_{W^{k+1,p}(\mathbb{D}_R)}$  on the right with  $\|u\|_{W^{k,p}(\mathbb{D}_R)}$ . By the interpolation inequality [GT01, Theorem 7.28], there exists  $C(R, k) > 0$  such that for every  $\varepsilon > 0$ ,

$$\|u\|_{W^{k+1,p}(\mathbb{D}_R)} \leq C(R, k)(\varepsilon\|u\|_{W^{k+2,p}(\mathbb{D}_R)} + \varepsilon^{-(k-1)}\|u\|_{W^{k,p}(\mathbb{D}_R)}). \quad (25)$$

Inserting (25) into (24), for  $\varepsilon > 0$  small enough we can rearrange to obtain

$$\|u\|_{W^{k+2,p}(\mathbb{D}_r)} \leq C(\|L\|_{W^{k+1,\infty}(\mathbb{D}_R)}, r, R, k, p)(\|u\|_{W^{k,p}(\mathbb{D}_R)} + \|v\|_{W^{k,p}(\mathbb{D}_R)}), \quad (26)$$

which is the statement of the proposition for  $L = \Delta_g$ .

Finally, we establish the result for arbitrary  $L$ . Toward this, we just need to control  $\|\Delta_g u\|_{W^{k,p}(\mathbb{D}_R)}$  in terms of  $\|Lu\|_{W^{k,p}(\mathbb{D}_R)}$  and  $\|u\|_{W^{k,p}(\mathbb{D}_R)}$ . The only needed observation is that our hypothesis guarantees that  $(L - \Delta_g)$  is an operator of order at most 1, and therefore

$$\begin{aligned} \|\Delta_g u\|_{W^{k,p}(\mathbb{D}_R)} &\leq \|Lu\|_{W^{k,p}(\mathbb{D}_R)} + \|(L - \Delta_g)u\|_{W^{k,p}(\mathbb{D}_R)} \\ &\leq \|Lu\|_{W^{k,p}(\mathbb{D}_R)} + C(\|L\|_{W^{k,\infty}(\mathbb{D}_R)}, \|\mu\|_{W^{k+1,\infty}(\mathbb{D}_R)})\|u\|_{W^{k+1,p}(\mathbb{D}_R)}. \end{aligned}$$

We then insert into the last inequality into the  $v$  slot of (26) and use the interpolation inequality as above to turn the  $\|u\|_{W^{k+1,p}(\mathbb{D}_R)}$  term into a  $\|u\|_{W^{k,p}(\mathbb{D}_R)}$  term.  $\square$

Having finished Proposition 5.5, we deduce the expected consequences that should come along with elliptic estimates. From bootstrapping the estimate of Proposition 5.5, we have the following.

**Proposition 5.8** (Regularity). *Let  $U \subset \mathbb{C}$  and  $V \subset \mathbb{C} \times \mathbb{C}^2$  be open domains, let  $F : U \times V \rightarrow \mathbb{C}$  be a  $W^{k+1,\infty}_{loc}$  function, and let  $L$  be a  $W^{k+1,\infty}_{loc}$  semi-linear second order complex elliptic operator of the form*

$$Lu(z) = \Delta_g u + F(z, u, Du).$$

*If  $u \in W^{k,p}_{loc}(U)$  satisfies  $Lu = 0$ , then for  $p$  such that  $\|\mu\|_{\infty} C_p < 1$ ,  $u$  is in  $W^{k+2,p}_{loc}(U)$ .*

It of course follows that, if  $L$  is  $C^\infty$ , then  $u$  is  $C^\infty$ .

*Proof.* Since  $u \in W^{k,p}_{loc}(U)$  the function  $G(z) = F(z, u(z), Du(z))$  is  $W^{k-1,p}_{loc}(U)$ . Therefore,  $\Delta_g u \in W^{k-1,p}_{loc}(U)$ . Applying the interior elliptic estimate from Proposition 5.5 with  $\Delta_g$ , we obtain that  $u$  is  $W^{k+1,p}_{loc}$ . This implies that  $G$  and hence  $\Delta_g u$  are  $W^{k,p}_{loc}$ . Elliptic estimates then give that  $u$  is  $W^{k+2,p}_{loc}$ .  $\square$

On closed surfaces, we have the expected Fredholm property. Given a positive complex metric  $g$ , locally of the form  $g = \lambda dz d\bar{w}$ , with  $\partial_{\bar{z}} w = \mu \partial_z w$ , we say that  $g$  has Beltrami form  $\mu$ .

**Proposition 5.9** (Fredholmness). *Let  $S$  be a closed surface,  $g$  a positive complex metric on  $S$  with induced  $W^{l,\infty}$  complex structures  $(c_1, \bar{c}_2)$ , and with Beltrami form  $\mu \in W^{l,\infty}$ . For  $l > k + 2$ , let*

$$L : W^{k+2,p}(S) \rightarrow W^{k,p}(S)$$

*be a linear operator whose principle symbol agrees with that of  $\Delta_h$ . For  $p$  such that  $\|\mu\|_\infty C_p < 1$ ,  $L$  is a Fredholm operator. If  $g$  is deformable to a Bers metric, the Fredholm index is zero.*

*Proof.* Once one has the estimate of Proposition 5.5, one can carry out the usual “real” proof of Fredholmness verbatim (for example, see [Nic07, Theorem 10.4.7]). If  $g$  is deformable to a Bers metric, then  $L$  is homotopic through Fredholm operators to the Euclidean Laplacian, so its index is zero.  $\square$

**Remark 5.10.** We believe it should be possible to prove analogous results for more general second order complex linear elliptic operators, for example those of the form

$$a\partial_z^2 + b\partial_z\partial_{\bar{z}} + c\partial_{\bar{z}}^2,$$

where  $a, b, c$  are complex-valued functions satisfying

$$a\zeta^2 + b\zeta\bar{\zeta} + c\bar{\zeta}^2 \neq 0$$

for all  $\zeta \in \mathbb{C}^*$ , using more general factorizations (see [AIM09, §7.8]), or higher order singular integral operators, as developed by Begehr (see [Beg94, §5]). We worked with the Laplacian rather than a more general operator because doing so allows for more expeditious proofs that use fewer tools and also highlights special features of the Laplacian.

The same remark applies to Theorem F.

**5.3. Proof of Theorem F.** We prove Theorem F. Let  $g$  be a real analytic positive complex metric on an open domain  $U \subset \mathbb{C}$ , locally of the form  $g = \lambda dz d\bar{w}$ . Recalling the setup, let  $V \subset \mathbb{C} \times \mathbb{C}^2$  be an open domain, let  $F : U \times V \rightarrow \mathbb{C}$  be a real analytic function, and let  $L$  be a semi-linear second order complex elliptic operator of the form

$$Lu(z) = \Delta_g u + F(z, u, Du).$$

Throughout the subsection, suppose we are given  $u \in C^2(U, \mathbb{C})$  with  $(u, Du)(U) \subset V$  that satisfies  $Lu = 0$ . The statement of Theorem F is that  $u$  is real analytic.

To prove the theorem, we will follow Kato’s strategy for establishing real analyticity of smooth solutions to real analytic real elliptic second order equations [Kat96], which was carried out in the general case by Hashimoto [Has06] and Blatt [Bla20].

**Remark 5.11.** We have to control a few extra terms compared to the real case. The reason is as follows. For any real semi-linear elliptic second order PDE, around any point one can find a small neighbourhood in which we have a change of coordinates that transforms the principle symbol to the Euclidean Laplacian. For complex elliptic PDE, such local transformations do not exist. For example, the reader can verify that no such transformation exists for

$$\partial_z\partial_{\bar{z}} + c\partial_{\bar{z}}^2,$$

where  $c$  is a constant of norm strictly less than 1.

**Remark 5.12.** In [Bla20], Blatt proves the result analogous to Theorem F for fully non-linear second-order elliptic operators. Thus, in comparison to [Bla20], in other ways our proof also appears simpler. The proof here could surely be pushed through to fully non-linear settings.

We now begin the preparations toward the proof. A multi-index is a tuple of positive integers  $\alpha = (m_1, \dots, m_N)$ , and we write  $|\alpha| = \sum_{i=1}^N m_i$ . Given (real or complex) coordinates  $x_1, \dots, x_n$  and a multi-index  $\alpha = (m_1, \dots, m_N)$ , we set  $\partial^\alpha = \partial_{x_{m_1}} \dots \partial_{x_{m_N}}$ . By Proposition 5.8, we know that any solution  $u$  to  $Lu = 0$  is  $C^\infty$ , so we can apply Cauchy's method of majorants to  $u$ .

**Theorem 5.13** (Cauchy's method of majorants). *Given a domain  $U \subset \mathbb{R}^n$ , a  $C^\infty$  function  $f : U \rightarrow \mathbb{C}$  is real analytic if and only if for any relatively compact sub-domain  $V$  and all multi-indices  $\alpha$ , there exist constants  $A, C > 0$  such that*

$$\|\partial^\alpha f\|_{L^\infty(V)} \leq CA^{|\alpha|} |\alpha|!.$$

Since real analyticity is a local property, we're welcome to work locally and assume  $U = \mathbb{D}$ , and furthermore we can assume that our function  $u$  admits uniform  $C^\infty$  bounds in  $\mathbb{D}$ . By repeated applications of Morrey's inequality, it suffices to prove an  $L^2$  bound rather than an  $L^\infty$  bound: namely, when working on the unit disk  $\mathbb{D}$ , one need only prove that for any  $0 < r < 1$ ,

$$\|\partial^\alpha u\|_{L^2(\mathbb{D}_r)} \leq CA^{|\alpha|} |\alpha|!, \quad (27)$$

where again  $A, C > 0$  are fixed constants, which now depend on  $r$ . We also need the higher order chain rule.

**Proposition 5.14** (Faa di Bruno). *Let  $K = \mathbb{R}$  or  $\mathbb{C}$  and let  $f : K^{m_1} \rightarrow K$  and  $g : K^{m_2} \rightarrow K^{m_1}$  be two functions. Then for any multi-index  $\alpha$ ,*

$$\partial^\alpha (f \circ g) = P_{m_1, m_2}^\alpha((\partial^\gamma f(g(x)))_{|\gamma| \leq |\alpha|}, (\partial^\gamma g)_{|\gamma| \leq |\alpha|}),$$

where  $P_{m_1, m_2}^\alpha$  is a linear combination with positive coefficients of terms of the form

$$\partial_{x_{i_1}} \dots \partial_{x_{i_1}} f(g(x)) \partial^{\gamma_1} g_{i_1} \dots \partial^{\gamma_1} g_{i_1}$$

with  $1 \leq m_2 \leq |\alpha|$  and  $|\gamma_1| + \dots + |\gamma_{m_2}| = |\alpha|$ .

The expressions are given by Faa di Bruno's formula, but we do not need the explicit form. When  $m_1 = m_2 = 1$ , we write  $P_{m_1, m_2}^\alpha = P^\alpha$ . With this notation established, we can present a criterion, extracted from the main proof in [Bla20], which will we use to show that our function satisfies the Cauchy estimates.

**Proposition 5.15.** *Let  $A, C, E > 0$  and let  $L_N$  be a sequence of positive numbers satisfying the recursive inequality*

$$L_{N+1} \leq CP^{|\alpha|}((E^k A^k k!)_{k \leq |\alpha|}, (L_k)_{k \leq |\alpha|}) + CN^2 L_{N-1} + CN L_N. \quad (28)$$

Then there exists  $B, D > 0$  such that

$$L_N \leq BD^N N!.$$

*Proof.* This is shown implicitly in [Bla20]: in [Bla20, Section 3.3], Blatt constructs an analytic function whose  $(N+1)^{th}$  Taylor coefficient at 0 can be seen to dominate the right hand side of (28). Cauchy estimates apply to the analytic function constructed in [Bla20, Section 3.3].  $\square$

Blatt provides a way to reduce the multivariate to the univariate case, which we'll use in order to apply Proposition 5.15.

**Lemma 5.16** (Lemma 2.9 in [Bla20]). *Suppose we are given real numbers  $(a_\gamma)_{|\gamma| \leq |\alpha|}, (b_\gamma)_{|\gamma| \leq |\alpha|}$  with the condition that  $a_\gamma$  and  $b_\gamma$  depend only on the length  $|\gamma|$  of the multi-index  $\gamma$ , and not the specific  $\gamma$ . Then,*

$$P_{m_1, m_2}^\alpha((a_\gamma)_{|\gamma| \leq |\alpha|}, (b_\gamma)_{|\gamma| \leq |\alpha|}) = P^\alpha((a_\gamma)_{|\gamma| \leq |\alpha|}, (b_\gamma)_{|\gamma| \leq |\alpha|}).$$

In the proofs of Lemma 5.18 and Theorem F below, we will use that  $W^{k,2}$  spaces are Banach algebras under multiplication.

**Proposition 5.17.** *There exists  $C = C(r, k) > 0$  such that for any two compactly supported  $W^{k,2}$  functions  $f_1, f_2$*

$$\|f_1 f_2\|_{W^{k,2}(\mathbb{D}_r)} \leq C \|f_1\|_{W^{k,2}(\mathbb{D}_r)} \|f_2\|_{W^{k,2}(\mathbb{D}_r)}.$$

As another adjustment for convenience, we replace  $\|\cdot\|_{W^{k,2}(\mathbb{D}_r)}$  with  $C(r, k)\|\cdot\|_{W^{k,2}(\mathbb{D}_r)}$  so that multiplication in  $W^{k,2}(\mathbb{D}_r)$  has norm at most 1.

Preliminary results aside, fix  $r > 0$ . As in [Kat96], [Has06], and [Bla20], we define a cutoff function  $\eta$  in  $C_0^\infty(\mathbb{D}, [0, 1])$  that is equal to 1 in  $\mathbb{D}_r$ . We prove the estimates (27) for  $\partial^\alpha u$  by induction on the length  $N$  of the multi-index  $\alpha$ . We will actually estimate things like  $\eta^N \partial^\alpha u$  in a subdisk  $\mathbb{D}_R$ ,  $R > r$ , containing the support of  $\eta$ , and then simply restrict to  $\mathbb{D}_r$ . Since the support does not go past  $\mathbb{D}_R$ , we will abuse notation slightly and write  $\|\cdot\|_k$  for  $\|\cdot\|_{W^{k,2}(\mathbb{D}_R)}$ . The point of multiplying by  $\eta^N$  is that we don't have to deal with passing to subdisks: if  $v$  is any regular enough function, with an  $s < r$  such that  $\mathbb{D}_s$  also contains the support of  $\eta$ , then,

$$\|\eta v\|_{k+2} \leq C(\|L_0(\eta v)\|_k + \|\eta v\|_k).$$

The following notation will be helpful. For  $N = 0$  and 1, set  $\tilde{L}_N = \max_{|\alpha|=N} \|\partial^\alpha u\|_k$  and for  $N \geq 2$  put

$$\tilde{L}_N = \max_{|\alpha|=N} \|\eta^{N-2} \partial^\alpha u\|_k.$$

Further set

$$L_N = \tilde{L}_{N+1} + \tilde{L}_N + 1.$$

We record one estimate before stepping into the proof of Theorem F.

**Lemma 5.18.** *Let  $K = \mathbb{R}$  or  $\mathbb{C}$ . For  $\alpha$  a multi-index of length  $N$  and  $f$  a uniformly bounded  $C^\infty$  function on a domain in  $K^n$ ,*

$$\|\eta^{|\alpha|} \partial^\alpha f(x, u, Du)\|_k \leq P^{|\alpha|}((E^{|\gamma|} \|D^{|\gamma|} f\|_k)_{|\gamma| \leq |\alpha|}, (L_j)_{j \leq |\alpha|}),$$

where  $E = (1 + \|\eta\|_k)^2$ .

This lemma is analogous to Lemma 2.10 from [Bla20], which concerns functions of the form  $f(x, u, Du, D^2u)$ . We give the proof, but refer to [Bla20] for one detail.

*Proof.* We assume  $N \geq 2$ , since the cases  $N = 0$  and 1 are simpler. We apply Proposition 5.14 to  $f \circ g$ , where  $g(x) = (x, u, Du)$ , and obtain

$$\eta^{|\alpha|} \partial^\alpha f(x, u, Du) = \eta^{|\alpha|} P_{n, 2n+1}^\alpha((\partial^\gamma f(g(x)))_{|\gamma| \leq |\alpha|}, (\partial^\gamma g)_{|\gamma| \leq |\alpha|}). \quad (29)$$

We estimate the  $W^{k,2}$  norms for all expressions of the form

$$\eta^{|\alpha|} \partial_{x_{i_1}} \dots \partial_{x_{i_l}} f(g(x)) \partial^{\gamma_1} g_{i_1} \dots \partial^{\gamma_l} g_{i_l},$$

where  $1 \leq l \leq |\alpha|$ ,  $|\gamma_1| + \dots + |\gamma_l| = |\alpha|$ . The main task is to estimate

$$\eta^{|\alpha|} \partial^{\gamma_1} g_{i_1} \dots \partial^{\gamma_l} g_{i_l} = \eta^{|\gamma_1|} \partial^{\gamma_1} g_{i_1} \dots \eta^{|\gamma_l|} \partial^{\gamma_l} g_{i_l}. \quad (30)$$

To understand the derivatives, take note that for  $i = 1, \dots, n$   $\partial^\gamma g_i = 1$  for  $|\gamma| = 1$  and 0 for  $|\gamma| > 1$ . Via the Banach algebra property and the triangle inequality, we thus obtain

$$\|\eta^{|\gamma_j|} \partial^{\gamma_j} g_{i_j}\|_k \leq \|\eta^{|\gamma_j|}\|_k + \|\eta^2\|_k \sup_{|\beta|=|\gamma_j|} \|\eta^{|\gamma_j|-2} \partial^\beta u\|_m + \|\eta\|_k \sup_{|\beta|=|\gamma_j|+1} \|\eta^{|\gamma_j|-1} \partial^\beta u\|_k.$$



By the Banach algebra property, all three terms  $\|\eta^{|\gamma_j|}\|_k$ ,  $\|\eta\|_k$ , and  $\|\eta^2\|_k$ , are crudely controlled by  $E^{|\gamma_j|}$ , and hence

$$\|\eta^{|\gamma_j|} \partial^{\gamma_j} g_{i_j}\|_k \leq E^{|\gamma_j|} L_{|\gamma_j|}.$$

Returning to (30), by the Banach algebra property again,

$$\eta^{|\alpha|} \partial_{x_{i_1}} \dots \partial_{x_{i_l}} f(g(x)) \partial^{\gamma_1} g_{i_1} \dots \partial^{\gamma_l} g_{i_l} \leq E^l \|\partial_{x_{i_1} \dots x_{i_l}}^l f(g(x))\|_k L_{|\gamma_1|} \dots L_{|\gamma_l|}.$$

And going back to (29), we have

$$\|\eta^{|\alpha|} \partial^\alpha f(x, u, Du)\|_k \leq \eta^{|\alpha|} P_{n, 2n+1}^{|\alpha|} ((E^{|\gamma|} \|D^{|\gamma|} f\|_k)_{|\gamma| \leq |\alpha|}, (L_j)_{j \leq |\alpha|}).$$

Applying Lemma 5.16 then yields the result.  $\square$

*Proof of Theorem F.* Let  $u$  be a solution to  $Lu = 0$ , which we know is  $C^\infty$ . Let's write the equation  $\Delta_g u + F(x, u, Du) = 0$  as  $L_0 + \lambda \overline{\partial}_z w \hat{F}(x, u, Du) = 0$ , where now  $L_0 = \partial_z \partial_{\bar{z}} - \bar{\mu} \partial_{\bar{z}}^2$ , and  $\hat{F}$  contains all of the first order information of  $L$ . To simplify things, let's just abuse notation and redefine  $F$  to be  $\lambda \overline{\partial}_z w \hat{F}$ . Getting rid of the conformal factor and the first order terms from  $\Delta_g$  makes for a slightly cleaner computation below.

Since  $F$  is real analytic, there exists  $A > 0$  such that for all multi-indices of length  $k$ ,  $\|\partial^\alpha F\|_m \leq A^k k!$  (to simplify further we enlarge  $A$  to absorb the usual constant  $C$ ).

As we discussed above, we go by induction on the length of the multi-index, say  $N$ . The base case is trivial. For the induction step, let  $\alpha$  be a multi-index of length  $N$ , and  $\beta$  a multi-index of length 2. Write

$$\eta^N \partial^{\alpha+\beta} u = \partial^\beta (\eta^N \partial^\alpha u) + [\eta^N, \partial^\beta] \partial^\alpha u.$$

The elliptic estimate of Proposition 5.5 then returns

$$\begin{aligned} \|\eta^N \partial^{\alpha+\beta} u\|_k &\leq \|\partial^\beta \eta^N \partial^\alpha u\|_k + \|[\eta^N, \partial^\beta] \partial^\alpha u\|_k \\ &\leq \|\eta^N \partial^\alpha u\|_{k+2} + \|[\eta^N, \partial^\beta] \partial^\alpha u\|_k \\ &\leq C(\|L_0(\eta^N \partial^\alpha u)\|_k + \|\eta^N \partial^\alpha u\|_k) + \|[\eta^N, \partial^\beta] \partial^\alpha u\|_k. \end{aligned}$$

Similar to above, we express  $L_0(\eta^N \partial^\alpha u)$

$$L_0(\eta^N \partial^\alpha u) = \eta^N L_0(\partial^\alpha u) + [L_0, \eta^N] \partial^\alpha u.$$

We further unfold, using that  $L_0(u) + F(x, u, Du) = 0$ ,

$$L_0(\partial^\alpha u) = \partial^\alpha L_0 u + [L_0, \partial^\alpha] u = \partial^\alpha F + \partial^\alpha(\bar{\mu}) \partial_{\bar{z}}^2 u.$$

Moreover,

$$\|L_0(\eta^N \partial^\alpha u)\|_k \leq \|\eta^N \partial^\alpha F\|_k + \|\partial^\alpha(\bar{\mu})\|_k \|\partial_{\bar{z}}^2 u\|_k + \|[L_0, \eta^N] \partial^\alpha u\|_k.$$

Returning to the main estimate, we have

$$\begin{aligned} \|\eta^N \partial^{\alpha+\beta} u\|_k &\leq C(\|\eta^N \partial^\alpha F\|_k + \|[L_0, \eta^N] \partial^\alpha u\|_k + \|\eta^N \partial^\alpha u\|_k + \|\eta^N \partial_{\bar{z}}^2 u\|_k) + \|[\eta^N, \partial^\beta] \partial^\alpha u\|_k \\ &=: J_1 + J_2 + J_3 + J_4 + J_5. \end{aligned}$$

We treat each  $J_i$  individually. Firstly, for  $J_1$  we apply Lemma 5.18 to see that

$$J_1 \leq P^{|\alpha|} ((E^k A^k k!)_{k \leq |\alpha|}, (L_k)_{k \leq |\alpha|}).$$

$J_3$  and  $J_4$  are both straightforwardly bounded by  $L_{N-1}$ . To handle  $J_2$  and  $J_5$ , first take note of the formula, for any sufficiently regular function  $v$ , for coordinates  $x_1, \dots, x_n$ , writing  $\partial_{ij} = \partial_{x_i} \partial_{x_j}$ ,

$$\partial_{ij}(\eta^N v) = (N(N-1)\eta^{N-2} \partial_i \eta \partial_j \eta + N\eta^{N-1} \partial_{ij} \eta) v + N\eta^{N-1} (\partial_i \eta \partial_j v + \partial_j \eta \partial_i v) + \eta^N \partial_{ij} v,$$

from which we derive the identity

$$[\eta^N, \partial_{ij}] \partial^\alpha u = (N(N-1)\eta^{N-2} \partial_i \eta \partial_j \eta + N\eta^{N-1} \partial_{ij} \eta) \partial^\alpha u + N\eta^{N-1} (\partial_i \eta \partial_j v + \partial_j \eta \partial_i \partial^\alpha u). \quad (31)$$

Splitting  $L_0$  as  $L_0 = \partial_z \partial_{\bar{z}} - \bar{\mu} \partial_{\bar{z}}^2$ ,

$$J_2 + J_5 \leq C \left( \| [\partial_z \partial_{\bar{z}}, \eta^N] \partial^\alpha u \|_k + \| [\partial_{\bar{z}}^2, \eta^N] \partial^\alpha u \|_k + \| [\partial^\beta, \eta^N] \partial^\alpha u \|_k \right).$$

Via the formula (31), and the Banach algebra property, all three of the terms above can be controlled by  $CN(N-1)L_{N-1} + CNL_N$ , and therefore

$$J_2 + J_5 \leq CN(N-1)L_{N-1} + CNL_N.$$

Putting everything together we have

$$\begin{aligned} L_{N+1} &\leq P^{|\alpha|} ((E^k A^k k!)_{k \leq |\alpha|}, (L_k)_{k \leq |\alpha|}) + CNL_{N-1} + CN(N-1)L_{N-1} + CNL_N \\ &= P^{|\alpha|} ((E^k A^k k!)_{k \leq |\alpha|}, (L_k)_{k \leq |\alpha|}) + CN^2 L_{N-1} + CNL_N. \end{aligned}$$

We then apply Proposition 5.15 to see that  $L_N \leq BD^N N!$ , for some constants  $B, D > 0$ , and Proposition 5.13 then implies that  $u$  is real analytic.  $\square$

**5.4. Proof of Theorem G.** The aim of this section is to prove Theorem 5.23, which directly implies Theorem G. Toward this goal, we first prove some general results. We define more generally, for any positive complex metric  $g$ ,  $L_{g,\lambda} : W^{k+2,p}(S) \rightarrow W^{k,p}(S)$  by  $L_{g,\lambda} u = \Delta_g u - \lambda u$ . By Proposition 5.9, when  $g$  is deformable to a Bers metric,  $L_{g,\lambda}$  has index zero and is invertible if and only if  $\lambda$  is not an eigenvalue for  $\Delta_g$ .

**Lemma 5.19.** *Let  $g$  be a positive complex metric such that  $\Delta_g$  has Fredholm index zero,  $\lambda \in \mathbb{C}$ , and suppose that  $L_{g,\lambda}$  is an isomorphism. Then there exists a neighbourhood  $U$  of  $g$  inside  $\mathcal{CM}(S)$  such that for all  $g' \in U$ ,  $L_{g',\lambda}$  is an isomorphism.*

*Proof.* Suppose that the conclusion of the lemma fails, so that we can find positive metrics  $g_n$  converging to  $g$  such that  $L_{g_n,\lambda}$  is not an isomorphism. Since for  $n$  large, each  $L_{g_n,\lambda}$  is Fredholm of index zero, this means that for every  $n$  we can find a non-zero  $u_n \in W^{k+2,p}(S)$  such that

$$L_{g_n,\lambda} u_n = 0.$$

Since each  $L_{g_n}^\lambda$  is linear, we can rescale each  $u_n$  so that  $\|u_n\|_{W^{k+2,p}} = 1$ . By the Rellich-Kondrachov theorem,  $u_n$  subconverges in  $W^{k,p}$  to some non-zero  $u_\infty$  that satisfies  $L_{g,\lambda} u = 0$  weakly. By Proposition 5.5,  $u \in W^{k+2,p}(S)$ , and we've contradicted the fact that  $L_{g,\lambda}$  is an isomorphism.  $\square$

We will say that  $c_1 \in \mathcal{C}(S)$  and  $\bar{c}_2 \in \mathcal{C}(\bar{S})$  are **co-real analytic** (or that  $(c_1, \bar{c}_2)$  is a co-real analytic pair) if they induce the same real analytic structure on  $S$ . Equivalently,  $c_1$  and  $\bar{c}_2$  are co-real analytic if the Beltrami form of one with respect to the other is real analytic (by [AB60, Theorem 3]). Observe that the diagonal action of  $\text{Diff}_0(S)$  preserves this condition.

By the density of  $\text{Diff}_0^\omega(S)$  in  $\text{Diff}_0(S)$ , the projection  $\mathcal{C}(S) \times \mathcal{C}(\bar{S}) \rightarrow \mathcal{T}(S) \times \mathcal{T}(\bar{S})$  is such that co-real analytic pairs in each fiber are dense in the fiber.

We will use the following result about the spectrum of the Laplacian of Bers metrics from co-real analytic complex structures, which is independently interesting.

**Proposition 5.20.** *Let  $(c_1, \bar{c}_2), (c'_1, \bar{c}'_2) \in \mathcal{C}(S) \times \mathcal{C}(\bar{S})$  be two pairs of co-real analytic complex structures on  $S$  that project to the same point in  $\mathcal{T}(S) \times \mathcal{T}(\bar{S})$ .*

*Then  $L_{\mathbf{h}_{(c_1, \bar{c}_2)}, \lambda}$  is an isomorphism if and only if  $L_{\mathbf{h}_{(c'_1, \bar{c}'_2)}, \lambda}$  is an isomorphism.*

*Proof.* We will actually prove the contrapositive statements, namely  $\lambda$  lies in the spectrum of  $\Delta_{\mathbf{h}_{(c_1, \overline{c_2})}}$  if and only if  $\lambda$  lies in the spectrum of  $\Delta_{\mathbf{h}_{(c'_1, \overline{c'_2})}, \lambda}$ .

Fix on  $S$  the real analytic structure induced by  $c_1$  and  $\overline{c_2}$  and denote by  $\text{Diff}^\omega(S)$  the corresponding subgroup of real analytic diffeomorphisms.

Since, for all  $\phi \in \text{Diff}(S)$ ,  $L_{\mathbf{h}_{(c_1, \overline{c_2})}, \lambda}$  is an isomorphism if and only if  $L_{\mathbf{h}_{(\phi^*c_1, \phi^*\overline{c_2})}, \lambda}$  is an isomorphism, we can assume that  $c_1 = c'_1$  and that  $\overline{c'_2}$  and  $\overline{c_2}$  lie in the same  $\text{Diff}_0^\omega(S)$ -orbit.

As we mentioned in the introduction to Section 4.3,  $\text{Diff}_0^\omega(S)$  is locally analytically path connected, so we are left to show the following: if  $\lambda$  is an eigenvalue for  $\Delta_{\mathbf{h}_{(c_1, \overline{c_2})}}$ , for all real analytic path  $\phi_t \in \text{Diff}_0^\omega(S)$ ,  $t \in [0, 1]$ ,  $\lambda$  is an eigenvalue for  $\Delta_{\mathbf{h}_{(c_1, \phi_t^*\overline{c_2})}}$  for all  $t \in [0, 1]$ .

Denote  $h_t = \mathbf{h}_{(c_1, \phi_t^*\overline{c_2})}$  and

$$A^\lambda = \{t \in [0, 1] : \lambda \text{ lies in the spectrum of } \Delta_{h_t}\}.$$

$A_\lambda$  contains 0 and, by Lemma 5.19,  $A^\lambda$  is closed. Let  $t_0$  be the maximum of the connected component of  $A^\lambda$  containing 0, and assume for the sake of contradiction that  $t_0 < 1$ . Denoting by  $z$  and  $\overline{w}$  local coordinates of  $c_1$  and  $\overline{c_2}$  respectively, as well as  $\dot{\phi}_t = \xi_t \partial_z + \overline{\xi}_t \partial_{\overline{z}}$ , define the vector field  $Z_t := \left( \xi_{t+t_0} \frac{\partial_z \overline{w}_{t+t_0}}{\partial_z \overline{w}_t} + \overline{\xi}_{t+t_0} \right) \partial_{\overline{z}}$  as in Example 4.2, so that, in the notation of Definition 4.5,  $\mathcal{I}_{Z_t}(\mathbf{h}_{(c_1, \phi_{t_0}^*(\overline{c_2}))}) = \mathbf{h}_{(c_1, \phi_{t_0+t}^*(\overline{c_2}))}$ . Now, let  $u$  be in the kernel of  $L_{h_t, \lambda}$ . By Theorem F,  $u$  is a real analytic function. By Theorem 4.11, there exists  $t_1$  with  $t_0 < t_1 \leq 1$  such that the analytic transport  $u_t = \mathcal{I}_{Z_t} u$  is defined for all  $t \in [0, t_1 - t_0]$  (with  $u_t$  being non-zero), and, by Proposition 4.9, for all  $t \in [0, t_0 - t_1]$

$$\Delta_{\mathbf{h}_{(c_1, \phi_{t_0+t}^*(\overline{c_2}))}} u_t = \Delta_{\mathbf{h}_{(c_1, \phi_{t_0}^*(\overline{c_2}))}} (\mathcal{I}_{Z_t} u) = \mathcal{I}_{Z_t} \left( \Delta_{\mathbf{h}_{(c_1, \phi_{t_0}^*(\overline{c_2}))}} u \right) = \lambda \mathcal{I}_{Z_t} u = \lambda u_t,$$

hence  $[t_0, t_1] \subset A^\lambda$ , contradicting the maximality of  $t_0$ .  $\square$

For the case  $\lambda = 2$ , we can say more. Recall we're setting  $L_h = L_{h,2}$ .

**Proposition 5.21.** *Let  $h = \mathbf{h}_{(c_1, \overline{c_2})}$  be a smooth Bers metric and  $q$  a small-enough  $c_1$ -holomorphic quadratic differential such that  $h + q$  is a Bers metric. Then,  $L_h$  is an isomorphism if and only if  $L_{h+q}$  is an isomorphism.*

*Proof.* Notice that it is sufficient to prove that if  $L_h$  is an isomorphism, then  $L_{h+q}$  is an isomorphism. Indeed, the converse implication follows by seeing  $h = (h + q) - q$ .

Observe that there exists  $\varepsilon = \varepsilon(h, q)$  such that, for all smooth  $\varrho: S \rightarrow \mathbb{C}^*$  for which  $\|1 - \varrho\|_\infty < \varepsilon$ ,  $\varrho h + q$  is a positive complex metric. In fact, one can see explicitly from the computation of Lemma 6.1 in [Ema23] that the condition for  $\varrho h + q$  to be positive is

$$\left| \overline{\mu} + \frac{1}{\varrho} \frac{q(\partial_z, \partial_z)}{h(\partial_z, \partial_z)} \right| < 1,$$

where  $\mu = \frac{\partial_z \overline{w}}{\partial_z \overline{w}}$ , and  $z$  and  $\overline{w}$  are local holomorphic coordinates for  $c_1$  and  $\overline{c_2}$  respectively.

We prove that the image of  $L_{h+q}$  contains  $C^\infty(S, \overline{\mathbb{C}})$ . It will follow from Fredholmness that  $L_{h+q}$  is surjective, and hence an isomorphism because it has index zero.

Since  $L_h$  is an isomorphism, for all  $f \in C^\infty(S, \mathbb{C})$  there exists  $\dot{u}$  such that  $L_h(\dot{u}) = f$ . By Proposition 5.8,  $\dot{u} \in C^\infty(S, \mathbb{C})$ . Now, assume  $t$  is small enough so that  $(1 + t\dot{u})h + q$  is a positive complex metric, and set  $h'_t$  to be the Bers metric in the conformal class of  $(1 + t\dot{u})h + q$ , so there exists a path of functions  $\sigma_t$ , with  $\sigma_0 = 1$ , such that

$$(1 + t\dot{u})h + q = \sigma_t h'_t.$$

One can see explicitly, with computations as in [Ema23, Remark 3.9], that  $\mathbf{c}_+(h'_t) = c_1$  and that the Beltrami form of  $\mathbf{c}_-(h'_t)$  with respect to  $c_1$  is  $\mu + \frac{q(\partial_z, \partial_z)}{(1+t\dot{u})h(\partial_z, \partial_z)}$ , so  $\mathbf{c}_-(h'_t)$  depends smoothly on  $t$ , hence, by Proposition 2.9,  $h'_t$  depends smoothly on  $t$  and so does  $\sigma_t$ .

By Proposition 2.10, we have that  $K_{(1+t\dot{u})h} = K_{\sigma_t h'_t}$ , hence, using Equation (4),

$$\frac{1}{(1+t\dot{u})} \left( -1 - \frac{1}{2} \Delta_h(\log(1+t\dot{u})) \right) = \frac{1}{\sigma_t} \left( -1 - \frac{1}{2} \Delta_{h'_t}(\log \sigma_t) \right),$$

and, taking the time derivative at 0,

$$\Delta_h \dot{u} - 2\dot{u} = \Delta_{h+q} \dot{\sigma} - 2\dot{\sigma}, \quad (32)$$

where we used that the coefficients of the operators  $\Delta_{h'_t}$  change smoothly with respect to  $t$ , so that  $\left( \frac{d}{dt} \Big|_0 \Delta_{h'_t} \right) (0) = 0$ . As a result, we get that  $f = \Delta_{h+q} \dot{\sigma} - 2\dot{\sigma}$ , hence  $\dot{\sigma}$  is smooth by Proposition 5.8, proving the surjectivity of  $L_{h+q}$ .  $\square$

**Lemma 5.22.** *Let  $h = \mathbf{h}_{(c_1, \overline{c_2})}$  be a Bers metric and  $q$  a small-enough  $c_1$ -holomorphic quadratic differential such that  $h + q = \mathbf{h}_{(c_1, \overline{c_2}' )}$  is a Bers metric. If  $c_1, \overline{c_2}$  are co-real analytic, then  $c_1, \overline{c_2}'$  are co-real analytic.*

*Proof.* As usual, denote  $h = \lambda dz d\overline{w}$ ,  $h + q = \lambda' dz d\overline{\eta}$  and  $q = \varphi dz^2$  in local coordinates  $z, \overline{w}, \overline{\eta}$  for  $c_1, \overline{c_2}, \overline{c_2}'$  respectively.

From  $\lambda dz d\overline{w} + q = \lambda' dz d\overline{\eta}$  we get that

$$\frac{\partial_z \overline{\eta} dz}{\partial_z \overline{\eta} d\overline{z}} = \frac{\lambda \partial_z \overline{w} + \varphi dz}{\lambda \partial_z \overline{w} d\overline{z}},$$

which is real analytic.  $\square$

Theorem G now follows from the following statement.

**Theorem 5.23.** *The set*

$$\mathcal{B}^{reg} = \{(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S}) \mid L_{\mathbf{h}_{(c_1, \overline{c_2})}} : W^{k+2,p}(S) \rightarrow W^{k,p}(S) \text{ is invertible}\}$$

*is an open dense subset of  $\mathcal{C}(S) \times \mathcal{C}(\overline{S})$  which contains all the pairs of co-real analytic complex structures. The intersection of  $\mathcal{B}^{reg}$  with any fiber of the projection  $\mathcal{C}(S) \times \mathcal{C}(\overline{S}) \rightarrow \mathcal{T}(S) \times \mathcal{T}(\overline{S})$  is connected.*

*Proof.* We first prove that  $\mathcal{B}^{reg}$  contains all the co-real analytic pairs.

Fix  $[c_1] \in \mathcal{T}(S)$ . Define  $Y \subseteq \{[c_1]\} \times \mathcal{T}(\overline{S})$  to be the subset of elements admitting a co-real analytic pair of representatives  $(c_1, \overline{c_2})$  such that  $L_{\mathbf{h}_{(c_1, \overline{c_2})}}$  is invertible. Conversely, define  $N \subseteq \{[c_1]\} \times \mathcal{T}(\overline{S})$  to be the subset of elements admitting a co-real analytic pair of representatives  $(c_1, \overline{c_2})$  such that  $L_{\mathbf{h}_{(c_1, \overline{c_2})}}$  is *not* invertible.

By Proposition 5.20,  $Y$  and  $N$  are complementary.  $Y$  contains  $([c_1], [\overline{c_1}])$ : in fact, for the Riemannian hyperbolic metric  $h = \mathbf{h}_{(c_1, \overline{c_1})}$ ,  $L_h$  is the complexification of a real operator  $W^{k+2,p}(S, \mathbb{R}) \rightarrow W^{k,p}(S, \mathbb{R})$ , which is invertible by a simple maximum-principle argument.

Moreover, both  $Y$  and  $N$  are open: by Theorem 2.11, Proposition 5.21, and Lemma 5.22, if  $c_1$  and  $\overline{c_2}$  are co-real analytic and  $([c_1], [\overline{c_2}])$  lies in  $Y$  (resp  $N$ ), then one can parametrize a local neighborhood of it in  $Y$  (resp  $N$ ) whose elements are of the form  $([c_1], [\overline{c_2}^q])$  where  $\mathbf{h}_{(c_1, \overline{c_2})} + q =: \mathbf{h}_{(c_1, \overline{c_2}^q)}$  and  $q$  is a small-enough  $c_1$ -holomorphic quadratic differential. We conclude that  $Y = \{[c_1]\} \times \mathcal{T}(\overline{S})$  so, by the generality of  $c_1$ , we get that all the co-real analytic pairs lie in  $\mathcal{B}^{reg}$ .

By Lemma 5.19,  $\mathcal{B}^{reg}$  is open.

Finally, let  $F$  be a fiber of  $\mathcal{C}(S) \times \mathcal{C}(\bar{S}) \rightarrow \mathcal{T}(S) \times \mathcal{T}(\bar{S})$ . By looking at any smooth trivialization of  $\mathcal{C}(S) \times \mathcal{C}(\bar{S}) \cong \text{Diff}_0(S) \times \text{Diff}_0(S) \times \mathcal{T}(S) \times \mathcal{T}(\bar{S})$ , the density of  $\text{Diff}_0^\omega(S)$  in  $\text{Diff}_0(S)$  implies that any element of  $F \cap \mathcal{B}^{reg}$  admits a connected neighborhood inside it containing a co-real analytic pair; the statement follows from the fact that co-real analytic pairs in  $F \cap \mathcal{B}^{reg}$  determine a connected subset.  $\square$

## 6. THE HOLOMORPHIC EXTENSION OF THE LABOURIE-LOFTIN PARAMETRIZATION

**6.1. The Labourie-Loftin parametrization.** We recall the definition of Hitchin representations from the introduction. Firstly, irreducibly embedding  $\iota : \text{PSL}(2, \mathbb{R}) \rightarrow \text{PSL}(3, \mathbb{R}) = \text{SL}(3, \mathbb{R})$ , we can turn a Fuchsian representation  $\rho_0 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{R})$  into a representation to  $\text{SL}(3, \mathbb{R})$ . A representation  $\rho : \pi_1(S) \rightarrow \text{SL}(3, \mathbb{R})$  is **Hitchin** if it can be deformed to a representation of the form  $\iota \circ \rho_0$  through representations to  $\text{SL}(3, \mathbb{R})$ . The space of Hitchin representations inside  $\chi(\pi_1(S), \text{SL}(3, \mathbb{R}))$  determines a connected component, called a Hitchin component and labelled  $\text{Hit}_3(S)$ .

Recall that a domain  $D \subset \mathbb{RP}^2$  is convex if it can be seen as a convex domain inside an affine chart. By work of Goldman [Gol90] and Choi-Goldman [CG93], every Hitchin representation  $\rho : \pi_1(S) \rightarrow \text{SL}(3, \mathbb{R})$  acts properly discontinuously on a bounded convex domain  $D \subset \mathbb{RP}^2$ , identifying  $S$  with  $D / \rho(\pi_1(S))$ . Labourie and Loftin both observed that, by a result conjectured by Calabi [Cal72] and attributed to Cheng-Yau [CY77] [CY86] and Calabi-Nirenberg (unpublished), there exists a unique  $\rho$ -equivariant hyperbolic affine sphere centered at zero and asymptotic to the positive cone of  $D$  in  $\mathbb{R}^3$ .

Labourie and Loftin both prove that the Gauss equation for real affine spheres

$$\Delta_h u = e^{2u} - \frac{1}{4}h(Q, \bar{Q})e^{-4u} - 1$$

has a unique (real) solution, and thus one can parametrize a real analytic moduli space of real affine spheres by the bundle  $\mathcal{M}_3(S)$  from the introduction (which we define precisely below). By taking the holonomies of affine spheres, one gets the Labourie-Loftin parametrization

$$\mathcal{L} : \mathcal{M}_3(S) \xrightarrow{\sim} \text{Hit}_3(S).$$

In the same spirit, we now return to complex affine spheres and our Equation (17) from Section 3 in order to build an open subset  $\Omega \subset \mathcal{M}_3(S) \times \mathcal{M}_3(S)$  containing the diagonal and the map  $\mathcal{L}_{\mathbb{C}} : \Omega \rightarrow \chi(\pi(S), \text{SL}(3, \mathbb{C}))$  that extends  $\mathcal{L}$ .

**6.2. Global analysis and the Gauss equation.** In this subsection we parametrize a class of  $C^\infty$  positive hyperbolic affine spheres as a complex Fréchet manifold. In order to make the global analysis more convenient, we first work over a family of Banach manifolds, and then we take a projective limit.

Recall  $\mathcal{C}^l(S), \mathcal{C}(S)$ , etc., defined in Section 2. Let  $\mathcal{CD}^l(S)$  be the bundle over  $\mathcal{C}^l(S)$  whose fiber over  $c \in \mathcal{C}^l(S)$  is the space  $H^0(S, \mathcal{K}_c^3)$  of  $c$ -holomorphic cubic differentials. Similarly, we have the analogous bundle  $\mathcal{CD}(S) \rightarrow \mathcal{C}(S)$ . It follows from work of Bers [Ber61, Theorem 2] that  $\mathcal{CD}^l(S)$  (resp.  $\mathcal{CD}(S)$ ) is a holomorphic vector bundle over  $\mathcal{C}^l(S)$  (resp.  $\mathcal{C}(S)$ ).  $\text{Diff}_0(S)$  acts on  $\mathcal{CD}(S)$  via the ordinary action on the basis  $\mathcal{C}(S)$  and by pull-back on the fibers. Again from [Ber61], the quotient of  $\mathcal{CD}(S)$  by this action is a holomorphic vector bundle over  $\mathcal{T}(S)$ , which is our bundle  $\mathcal{M}_3(S)$  from the introduction. We will denote by  $\mathcal{CD}^l(\bar{S}), \mathcal{CD}(\bar{S})$ , and  $\mathcal{M}_3(\bar{S})$  the analogous spaces for the oppositely oriented surface  $\bar{S}$ .

For  $k > 2$  and  $l > k + 2$ , we consider the function

$$\begin{aligned} G: \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times W^{k+2,2}(S) &\rightarrow W^{k,2}(S) \\ G(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) &:= \Delta_{\mathbf{h}_{(c_1, \overline{c_2})}} u - e^{2u} + \frac{1}{4} \mathbf{h}_{(c_1, \overline{c_2})}(Q_1, \overline{Q_2}) \cdot e^{-4u} + 1 \end{aligned} \quad (33)$$

whose zeroes, by Corollary 3.22, correspond to positive hyperbolic complex affine spheres with Blaschke metric of the form  $g = e^{2u} \mathbf{h}_{(c_1, \overline{c_2})}$ . Moreover, we define the space

$$\text{CAS}(S) := \{(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) \in \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) \times C^\infty(S, \mathbb{C}) : G(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) = 0\}, \quad (34)$$

with the induced subspace topology.

Note that by Proposition 5.8, if  $c_1, \overline{c_2}, Q_1, \overline{Q_2}$  are  $C^\infty$ , then any  $W^{k+2,2}$  solution  $u$  to  $G(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) = 0$  is actually  $C^\infty$ , so in the definition of  $\text{CAS}(S)$  one can replace the  $C^\infty$  above by  $W^{k+2,2}$ .

By Example 3.23, the space of real hyperbolic affine spheres corresponds to the subspace

$$\text{RAS}(S) := \{(c, \overline{c}, Q, \overline{Q}, u) \in \text{CAS}(S) \mid u \text{ is a real function}\}.$$

The map  $G: \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times W^{k+2,2}(S) \rightarrow W^{k,2}(S)$  is holomorphic: by Hartogs' theorem on separate holomorphicity, we're permitted to check on each coordinate separately. Since holomorphicity in  $u$  is clear, we are left to prove it in the direction of  $\mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S})$ . By Proposition 2.9, the association  $(c_1, \overline{c_2}) \rightarrow \mathbf{h}_{(c_1, \overline{c_2})}$  is holomorphic. It follows from the Koszul formula for the Levi-Civita connection that for fixed  $u$ ,  $(c_1, \overline{c_2}) \mapsto \Delta_{\mathbf{h}_{(c_1, \overline{c_2})}} u$  is holomorphic as well. Similarly,  $(c_1, \overline{c_2}, Q_1, \overline{Q_2}) \mapsto \mathbf{h}_{(c_1, \overline{c_2})}(Q_1, \overline{Q_2})$ , being a composition of holomorphic functions, is holomorphic.

Fix a point  $\sigma = (c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) \in \text{CAS}(S)$  and consider the linearization  $L_\sigma$  of  $G$  at  $\sigma$  in the direction of  $u$ , which is a linear map from  $T_u W^{k+2,2}(S, \mathbb{C}) = W^{k+2,2}(S, \mathbb{C})$  to  $T_0 W^{k,2}(S, \mathbb{C}) = W^{k,2}(S, \mathbb{C})$ , where the equal sign is an implicit identification that we will always make. The evaluation of the map  $L_\sigma$  at a tangent vector  $v \in W^{k+2,2}(S, \mathbb{C})$  is obtained by computing

$$L_\sigma(v) = \left. \frac{d}{dt} \right|_{t=0} G(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u + tv),$$

which returns

$$L_\sigma(v) = \Delta_{\mathbf{h}_{(c_1, \overline{c_2})}} v - 2ve^{2u} - ve^{-4u} \mathbf{h}_{(c_1, \overline{c_2})}(Q_1, \overline{Q_2}). \quad (35)$$

By Propositions 5.8 and 5.9,  $L_\sigma$  defines a Fredholm operator of index zero on each  $W^{k+2,2}$ , and it is an isomorphism on  $C^\infty$  if and only if it defines an isomorphism for every  $W^{k+2,2}$ .

**Definition 6.1.** We say that  $\sigma \in \text{CAS}(S)$  is **infinitesimally rigid** if  $L_\sigma$  is an isomorphism. We denote the subspace of infinitesimally rigid elements by  $\text{CAS}^*(S)$ .

First of all, we observe that  $\text{CAS}^*(S)$  is non-empty and contains the real affine spheres.

**Proposition 6.2.**  $\text{RAS}(S)$  is contained in  $\text{CAS}^*(S)$ .

*Proof.* For all  $\sigma \in \text{RAS}(S)$ , the linear map  $L_\sigma$  restricts to a map  $W^{k+2,2}(S, \mathbb{R}) \rightarrow W^{k,2}(S, \mathbb{R})$ , and Labourie showed in his study of real affine spheres that such a restriction is bijective [Lab07, Lemma 4.1.2].  $L_\sigma$  is the complex bilinear extension of the map from  $W^{k+2,2}(S, \mathbb{R}) \rightarrow W^{k,2}(S, \mathbb{R})$ , and hence for any  $v \in W^{k+2,2}(S)$ ,  $L_\sigma(v) = L_\sigma(\text{Re}(v)) + iL_\sigma(\text{Im}(v))$ , and it follows that  $L_\sigma$  is injective on  $W^{k,\infty}(S, \mathbb{C})$ . Since  $L_\sigma$  is Fredholm of degree zero, it is bijective.  $\square$

Denote by

$$\begin{aligned} \hat{\pi} : \text{CAS}^*(S) &\rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) \\ \sigma = (c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) &\mapsto (c_1, \overline{c_2}, Q_1, \overline{Q_2}) \end{aligned}$$

the natural projection. By transversality theory, we have the following.

**Proposition 6.3.** *The space  $\text{CAS}^*(S)$  is a complex Fréchet manifold and  $\hat{\pi} : \text{CAS}^*(S) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\overline{S})$  is a local biholomorphism.*

*Proof.* For  $k > 2$  and  $l > k + 2$ , define  $\text{CAS}_{l,k}^*(S)$  to be the subset of elements  $\sigma \in \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times W^{k+2,2}(S, \mathbb{C})$  such that  $G(\sigma) = 0$  and its linearization  $L_\sigma : W^{k+2,2}(S, \mathbb{C}) \rightarrow W^{k,2}(S, \mathbb{C})$  is an isomorphism. Also denote by  $\hat{\pi} : \text{CAS}_{l,k}^*(S) \rightarrow \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S})$  the projection that extends the projection from the smooth setting. The map

$$\begin{aligned} \Phi : \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times W^{k+2,2}(S, \mathbb{C}) &\rightarrow \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times W^{k,2}(S, \mathbb{C}) \\ (c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) &\mapsto (c_1, \overline{c_2}, Q_1, \overline{Q_2}, G(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u)) \end{aligned}$$

is holomorphic and maps onto the submanifold of the target described by  $\mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times \{0\}$ . By assumption, for all  $\sigma \in \text{CAS}_{l,k}^*(S)$ ,  $L_\sigma$  is an isomorphism, and hence so is  $d_\sigma \Phi$ . By the Transversality Theorem for Complex Banach manifolds,  $\text{CAS}_{l,k}^*(S)$  is a complex Banach submanifold of  $\mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S}) \times W^{k+2,2}(S, \mathbb{C})$ .

Since  $G|_{\text{CAS}_{k,l}^*(S)} \equiv 0$ ,  $d_\sigma \Phi = (d_\sigma \hat{\pi}, 0)$ , and hence  $d_\sigma \hat{\pi} : \text{CAS}_{l,k}^*(S) \rightarrow \mathcal{CD}^l(S) \times \mathcal{CD}^l(\overline{S})$  is bijective, which implies that  $\hat{\pi}$  is a local biholomorphism.

From now on, set  $l = k + 3$ , and  $\text{CAS}_k^*(S) = \text{CAS}_{k,k+3}^*(S)$ . Then,  $\text{CAS}_k^*(S)$  parametrizes complex affine spheres with lower regularity, and the intersection is  $\bigcap_{k' \geq k} \text{CAS}_{k'}^*(S)$  is  $\text{CAS}^*(S)$ . To put a Fréchet structure on  $\text{CAS}^*(S)$ , fix an open cover of  $\text{CAS}_k^*(S)$  by open subsets  $U_k$  over which  $\hat{\pi}$  projects biholomorphically to open subsets  $V_k \subset \mathcal{CD}^k(S) \times \mathcal{CD}^k(\overline{S})$ , thus giving a compatible atlas for the complex Banach structure on  $\text{CAS}_k^*(S)$ . For all  $k' > k$ , the intersection with  $\text{CAS}_{k'}^*(S)$  gives complex charts  $U_{k'} = U_k \cap \text{CAS}_{k'}^*(S) \rightarrow V_{k'} = V_k \cap (\mathcal{CD}^{k'+3}(S) \times \mathcal{CD}^{k'+3}(\overline{S}))$ . Finally, the restriction to

$$U_k \cap \text{CAS}^*(S) = \bigcap_{k' \geq k} U_{k'} \rightarrow \bigcap_{k' \geq k} V_{k'} = V_k \cap (\mathcal{CD}(S) \times \mathcal{CD}(\overline{S}))$$

defines Fréchet charts for  $\text{CAS}^*(S)$  for which  $\hat{\pi}$  is holomorphic.

The construction above can be formalized through a projective limit construction for complex Banach manifolds. As explained in [DGV15], in the smooth category, one can take a projective limit of Banach manifolds and the limit is a Fréchet manifold, and moreover a projective family of locally diffeomorphic mappings between projective systems gives rise to a local diffeomorphism between the two limiting Fréchet manifolds (see Propositions 3.13, 3.18, and Corollary 3.29 in [DGV15]). The proofs go through verbatim in the holomorphic category, although we don't need to refer to [DGV15] for the whole thing since the construction above is explicit.  $\square$

**Remark 6.4.** We need to make a brief digression on the character variety and holomorphicity, which concerns Theorem A, Proposition 6.5 and Theorem 6.10. The space  $\text{Hom}(\pi_1(S), \text{SL}(3, \mathbb{C}))$  embeds inside  $\text{SL}(3, \mathbb{C})^{2g}$  as a complex manifold, and the  $\text{SL}(3, \mathbb{C})$ -action by conjugation is by biholomorphisms. We set  $\chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$  to be the Hausdorffification of the quotient with respect to this action. In this paper, we take a naive standpoint and say that a map  $F : M \rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$ , with  $M$  a complex (Fréchet) manifold, is holomorphic if, around every point of  $M$ , it locally lifts to holomorphic maps

$M \supset U \rightarrow \text{Hom}(\pi_1(S), \text{SL}(3, \mathbb{C}))$ . As a result, if  $N \rightarrow M$  is a holomorphic map between complex (Fréchet) manifolds, then the composition  $N \rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$  is holomorphic.

The quotient of the subset of irreducible representations in  $\text{Hom}(\pi_1(S), \text{SL}(3, \mathbb{C}))$  carries the structure of a (singular) complex analytic space (see [Mar]), and if  $F$  has image in this subset, then it is holomorphic in the sense of complex analytic spaces.

We don't know whether the holonomy of a positive hyperbolic complex affine sphere is irreducible, but a lot of the families of points in the set  $\Omega$  from Theorem A have irreducible holonomy. Indeed, by openness of irreducibility, we have the condition for free around the diagonal and the quasi-Fuchsian space  $\mathcal{T}(S) \times \mathcal{T}(\bar{S})$ . By Theorem E, points of the form  $(c, \bar{c}, Q_1, 0)$  and  $(c, \bar{c}, 0, Q_2)$ , correspond to opers, which are known to have irreducible holonomy (see [BD96] and [Wen15, Proposition 4.8]). As we alluded to in the introduction, points coming from  $\mathcal{M}_3(S) \times \mathcal{T}(\bar{S})$  should correspond to some generalization of opers, so we strongly believe that these points should be irreducible (see Remark 7.16 for more on this).

**Proposition 6.5.** *The holonomy map  $hol: \text{CAS}^*(S) \rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$  is holomorphic.*

*Proof.* The complex affine sphere  $\sigma$  parametrized by  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) \in \text{CAS}^*(S)$  gives rise to structural data  $g = e^{2u} \mathbf{h}_{(c_1, \bar{c}_2)}$ ,  $\nabla = -\frac{1}{2}g^{-1}(Q_1 + \bar{Q}_2) + \nabla^g$ ,  $S = -id$ , and  $\tau = 0$ . Going through Theorem 3.13, out of this data we obtain a rank 3 complex vector bundle over  $S$  that carries a flat  $\text{SL}(3, \mathbb{C})$ -connection  $D^\sigma$  that satisfies the structural equation given at the beginning of the proof of Theorem 3.20.

Using Proposition 2.9, the correspondence  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) \mapsto e^{2u} \mathbf{h}_{(c_1, \bar{c}_2)}$  is holomorphic, and it is routinely checked that  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) \mapsto -\frac{1}{2}g^{-1}(Q_1 + \bar{Q}_2) + \nabla^g$  is holomorphic as well. Using the formula from Theorem 3.20, it is clear that  $\sigma = (c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) \mapsto D_\sigma$  defines a holomorphic map to the complex affine space that models flat  $\text{SL}(3, \mathbb{C})$  connections.

It is then well known that the holonomy map from the space of flat  $\text{SL}(3, \mathbb{C})$ -connections to the character variety is holomorphic [Gun67, §9]. Intuitively, if  $D$  is a holomorphically varying flat connection, the  $D$ -parallel section passing by a fixed element in the bundle varies holomorphically as well: by fixing a point  $p$  on  $S$  and a basis on one fiber, one gets that the holonomy homomorphism  $\pi_1(S) \rightarrow \text{SL}(3, \mathbb{C})$  induced by the corresponding basis of  $D$ -parallel sections varies holomorphically inside  $\text{Hom}(\pi_1(S), \text{SL}(3, \mathbb{C}))$ , hence  $hol$  is holomorphic in the sense of Remark 6.4.  $\square$

**6.3. The space  $\text{CAS}(S)$ .** We are now ready to define a finite-dimensional quotient of  $\text{CAS}^*(S)$  that keeps track of the geometric information of the corresponding complex affine spheres.

For the rest of the section, we will take local trivialisations of the bundle  $\mathcal{C}(S) \rightarrow \mathcal{T}(S)$  obtained from local holomorphic sections  $s: \mathcal{T}(S) \supset U \rightarrow \mathcal{C}(S)$  such that each element in the image is a real analytic complex structure compatible with a fixed real analytic structure on  $S$ . We show a way to construct such a section in Remark 6.6 below. In this way, we can trivialize open subsets of  $\mathcal{C}(S)$  as  $U \times \text{Diff}_0(S)$ , with  $U$  projecting injectively to  $\mathcal{T}(S)$ , in such a way that the projection  $U \times \text{Diff}_0(S) \rightarrow U$  is holomorphic (recall Theorem 2.2), and the trivialization restricts to  $\mathcal{C}^\omega(S) \cap (U \times \text{Diff}_0(S)) = U \times \text{Diff}_0^\omega(S)$ . By trivializing  $\mathcal{CD}(S) \rightarrow \mathcal{C}(S)$  over such an open subset, we also obtain local trivialisations for the bundle  $\mathcal{CD}(S) \rightarrow \mathcal{M}_3(S)$ .

**Remark 6.6.** We can construct such sections using Theorem 2.11: starting from a real analytic representative  $c$ , let  $h = \mathbf{h}_{(c,c)}$  be the (real analytic) hyperbolic metric in its conformal class, then define a local section by taking complex structures of the form  $c_+(h + \bar{q})$



with  $q \in \mathcal{K}_c^2(S)$  and  $\|q\|_h < \frac{1}{2}$  (the corresponding Beltrami form for  $c_+(h + \bar{q})$  is in fact  $\frac{\bar{q}}{h}$ , see [Ema23, Remark 2.9]).

Now, define the map

$$\begin{aligned} [\hat{\pi}]: \text{CAS}^*(S) &\rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S}) \\ (c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) &\mapsto ([c_1, Q_1], [\bar{c}_2, \bar{Q}_2]). \end{aligned}$$

Continuous deformations within the fibers of  $[\hat{\pi}]$  can actually be better understood using the transport formalism.

**Proposition 6.7.** *Let  $\sigma_t = (c_1^t, \bar{c}_2^t, Q_1^t, \bar{Q}_2^t, u_t) \in \text{CAS}^*(S)$  be a  $C^1$  path contained in a fiber of  $[\hat{\pi}]$ . Then, for all  $t$ , there exists  $Z_t \in \Gamma(\mathbb{C}TS)$  such that*

$$\dot{\sigma}_t = (\mathcal{L}_{Z_t} c_1^t, \mathcal{L}_{Z_t} \bar{c}_2^t, \mathcal{L}_{Z_t} Q_1^t, \mathcal{L}_{Z_t} \bar{Q}_2^t, \mathcal{L}_{Z_t} u_t).$$

*Proof.* To simplify the notation, we do the computation in  $t = 0$ , then the statement follows for the whole path by reparametrizing. Denote  $\sigma_0 = (c_1, \bar{c}_2, Q_1, \bar{Q}_2, u)$ . Since  $\sigma_t$  is contained in a fiber of  $[\hat{\pi}]$ , there exist  $\phi_1^t, \phi_2^t \in \text{Diff}_0(S)$  such that

$$(c_1^t, Q_1^t) = ((\phi_1^t)^* c_1, (\phi_1^t)^* Q_1), \quad (\bar{c}_2^t, \bar{Q}_2^t) = ((\phi_2^t)^* \bar{c}_2, (\phi_2^t)^* \bar{Q}_2).$$

By Proposition 4.3 and Remark 4.1, there exist  $Z \in \Gamma(\mathbb{C}TS)$  such that

$$\frac{d}{dt} \Big|_0 ((\phi_1^t)^* c_1, (\phi_2^t)^* \bar{c}_2, (\phi_1^t)^* Q_1, (\phi_2^t)^* \bar{Q}_2) = (\mathcal{L}_Z c_1, \mathcal{L}_Z \bar{c}_2, \mathcal{L}_Z Q_1, \mathcal{L}_Z \bar{Q}_2).$$

Applying both  $\frac{d}{dt}$  and  $\mathcal{L}_Z$  to Gauss equation  $G(\sigma_t) = 0$  in  $t = 0$ , by Proposition 4.9 one gets that

$$\begin{aligned} 0 &= \Delta_{\mathbf{h}_{(c_1, \bar{c}_2)}} \left( \frac{d}{dt} \Big|_0 u - \mathcal{L}_Z u \right) - 2e^{2u} \left( \frac{d}{dt} \Big|_0 u - \mathcal{L}_Z u \right) - e^{-4u} \mathbf{h}_{(c_1, \bar{c}_2)}(Q_1, \bar{Q}_2) \left( \frac{d}{dt} \Big|_0 u - \mathcal{L}_Z u \right) \\ &= L_\sigma \left( \frac{d}{dt} \Big|_0 u - \mathcal{L}_Z u \right). \end{aligned}$$

Since  $\sigma \in \text{CAS}^*(S)$ ,  $L_\sigma$  is injective, hence  $\frac{d}{dt} \Big|_0 u = \mathcal{L}_Z u$ .  $\square$

The last proposition motivates the definition of the following quotient for  $\text{CAS}^*(S)$ .

**Definition 6.8.** Define the equivalence relation  $\sim$  on  $\text{CAS}^*(S)$  by  $\sigma \sim \sigma'$  if and only if  $\sigma$  and  $\sigma'$  lie in the same connected component of a fiber of  $[\hat{\pi}]$ .

Define

$$\mathbb{C}\text{AS}(S) := \text{CAS}^*(S) / \sim$$

as the quotient by  $\sim$ . We denote the quotient map by  $p_\sim: \text{CAS}^*(S) \rightarrow \mathbb{C}\text{AS}(S)$ .

By construction,  $\hat{\pi}$  descends to a map

$$\begin{aligned} \pi: \mathbb{C}\text{AS}(S) &\rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S}) \\ [\sigma] &\mapsto [\hat{\pi}(\sigma)]. \end{aligned}$$

**Proposition 6.9.**  *$\mathbb{C}\text{AS}(S)$  has a structure of finite-dimensional complex manifold such that the quotient map  $p_\sim: \text{CAS}^*(S) \rightarrow \mathbb{C}\text{AS}(S)$  is holomorphic, and the map  $\pi: \mathbb{C}\text{AS}(S) \rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$  is a local biholomorphism.*

*Proof.* Consider local trivializations of  $\mathcal{CD}(S) \rightarrow \mathcal{M}_3(S)$  as in Remark 6.6.

By Proposition 6.3, for all  $\sigma \in \text{CAS}^*(S)$ , we can restrict  $\hat{\pi}$  to a biholomorphism in a neighborhood of  $\sigma$ , namely

$$\hat{\pi}: \hat{U}_\sigma \xrightarrow{\sim} A_1 \times B_1 \times A_2 \times B_2 \subset \mathcal{CD}(S) \times \mathcal{CD}(\bar{S}),$$

where  $U_\sigma$  is a connected neighborhood of  $\sigma$ , while  $A_1 \subset \mathcal{M}_3(S)$ ,  $A_2 \subset \mathcal{M}_3(\bar{S})$ , and  $B_1, B_2 \subset \text{Diff}_0(S)$  are open subsets.

Denoting with  $p_\sim: \text{CAS}^*(S) \rightarrow \mathbb{CAS}(S)$  the quotient map, we therefore have the following commutative diagram:

$$\begin{array}{ccc} \hat{U}_\sigma & \xrightarrow{\hat{\pi} \sim} & A_1 \times B_1 \times A_2 \times B_2 \\ \downarrow p_\sim & \searrow [\hat{\pi}] & \downarrow \text{projection} \\ U_\sigma := p_\sim(\hat{U}_\sigma) & \xrightarrow{\pi \sim} & A_1 \times A_2 \end{array} \quad (36)$$

It is easy to check that the collection of subsets  $\{U_\sigma\}_{\sigma \in \text{CAS}^*(S)}$  of  $\mathbb{CAS}(S)$  defines a base for a topology on  $\mathbb{CAS}(S)$ . Moreover, using the one-to-one restrictions of  $\pi$  to each  $U_\sigma$  and the fact that the projection is holomorphic, we get a holomorphic atlas on  $\mathbb{CAS}(S)$  by pulling back the complex structure on  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$ : by the diagram (36),  $\pi$  is a local biholomorphism and  $p_\sim$  is holomorphic.  $\square$

The following theorem shows that  $\mathbb{CAS}(S)$  preserves the information about the holonomy.

**Theorem 6.10.** *The holonomy map*

$$\begin{aligned} \text{hol}: \mathbb{CAS}(S) &\rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C})) \\ [\sigma] &\mapsto \text{hol}(\sigma) \end{aligned}$$

*is well-defined and holomorphic.*

*Proof.* If we prove that the holonomy map descends in a well-defined way to  $\mathbb{CAS}(S)$ , then it is holomorphic for free: by taking local charts for  $p_\sim$  as in the diagram (36), one has local holomorphic inverses of  $p_\sim$  in the form  $([c_1, Q_1], [\bar{c}_2, Q_2]) \mapsto ([c_1, Q_1], \phi_0, [\bar{c}_2, Q_2], \phi'_0)$  around each point in  $\mathbb{CAS}(S)$ . The composition with the holonomy map  $\text{CAS}^*(S) \rightarrow \chi(\pi_1(S), \text{SL}(3, \mathbb{C}))$ , which is holomorphic by Proposition 6.5, gives precisely the map  $\text{hol}$  in the statement of the theorem.

Using the diagram (36), we see that the connected components of the fibers of  $p_\sim$  in  $\text{CAS}^*(S)$  are open (and connected by construction). To prove that  $\text{hol}$  is well-defined on  $\mathbb{CAS}(S)$ , it is therefore sufficient to prove that  $\text{hol}$  is locally constant on the fibers. Moreover, the holonomy is clearly constant if we deform the complex affine sphere by pre-composition with an isotopy (inducing a diagonal isotopy in the projection to  $\mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$ ), and, in the notation of diagram (36), the density of analytic diffeomorphisms implies that it is sufficient to prove that the holonomy is constant on subsets of the form

$$F_0 := \hat{\pi}^{-1} \left( \{(c'_1, Q'_1)\} \times \{\phi'\} \times \{(\bar{c}'_2, \bar{Q}'_2)\} \times (\text{Diff}_0^\omega(S) \cap B_2) \right),$$

where  $c'_1$ ,  $\phi'$ , and  $\bar{c}'_2$  are real analytic. Observe that, by Theorem F, the complex affine spheres in  $F_0$  are of the form  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u)$  where  $u$  is a real analytic function.

We can therefore adapt an argument that we have used in the proof of Proposition 5.20. Fix  $\sigma_0 = (c_1, Q_1, \bar{c}_2, \bar{Q}_2, u) \in F_0$ , and let  $\rho$  be its holonomy (we will have  $c_1 = \phi'(c'_1)$  according to the notation above). Let  $\phi_t$ ,  $t \in [0, 1]$ , be a piecewise analytic path in  $\text{Diff}_0^\omega(S)$  such that  $(c_1, Q_1, \phi_t^*(\bar{c}_2), \phi_t^*(\bar{Q}_2)) \in \hat{\pi}(F_0)$ , hence there exists  $u_t$  such that  $\sigma_t = (c_1, Q_1, \phi_t^*(\bar{c}_2), \phi_t^*(\bar{Q}_2), u_t) \in F_0$ . Denote  $A_\rho = \{t \in [0, 1] \mid \text{hol}(\sigma_t) = \rho\}$ , which is a closed

subset of  $[0, 1]$ . Let  $T$  be the maximum of the connected component of  $A_\rho$  which contains 0, and assume by contradiction that  $T < 1$ ; then, as shown in Example 4.2, there exists for all  $t \in [0, 1]$  a (real analytic and piecewise real analytic with respect to  $t$ ) complex vector field  $Z_t$  such that  $\mathcal{T}_{Z_t}c_1 = c_1$  and  $\mathcal{T}_{Z_t}\overline{c_2} = \phi_t^*\overline{c_2}$ . By Theorem 4.11 and in the notation of Definition 4.5, there exists  $T' > T$  such that for all  $t \in [T, T']$  there exists  $u_t = \mathcal{T}_{Z_{t-T}}u_T$  which is the transport of  $u_T$  through  $Z_{\bullet-T}$ . Moreover, by Proposition 4.13,  $\sigma_t := ((c_1, Q_1, \phi_t^*(\overline{c_2}), \phi_t^*(\overline{Q_2}), u_t) \in \text{CAS}(S)$  for all  $t \in [T, T']$  and it has the same holonomy as  $\sigma_T$ , namely  $\rho$ : up to shrinking  $T'$ , we can assume that  $\sigma_t \in \text{CAS}^*(S)$  (which is an open subset of  $\text{CAS}(S)$ ) for all  $t \in [T, T']$ , contradicting the maximality of  $T$ , and proving that  $A_\rho = [0, 1]$ . By the generality of the path, we conclude the proof.  $\square$

In conclusion, we have the following commutative diagram of holomorphic maps:

$$\begin{array}{ccc} \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) & \xleftarrow{[\hat{\pi}]} & \text{CAS}^*(S) \\ \text{Diff}_0(S) \times \text{Diff}_0(\overline{S}) \downarrow \text{action} & & \downarrow p \sim \\ \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S}) & \xleftarrow{\pi} & \mathbb{CAS}(S) \xrightarrow{\text{hol}} \chi(\pi_1(S), \text{SL}(3, \mathbb{C})) \end{array}$$

**6.4. The action by  $\text{MCG}(S)$  and the  $\mathbb{C}^*$  twistor action.** We digress to discuss two actions on the spaces  $\text{CAS}(S)$ ,  $\text{CAS}^*(S)$ ,  $\mathbb{CAS}(S)$ , and  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$  related to Theorem A. Recall the groups  $\text{Diff}_+(S)$ ,  $\text{Diff}_0(S)$ , and  $\text{MCG}(S)$  from Section 2.

Given a positive hyperbolic complex affine sphere  $\sigma: \tilde{S} \rightarrow \mathbb{C}^3$  with Blaschke metric  $g$  and Pick tensor  $C$ , for all  $\phi \in \text{Diff}_+(S)$  one gets that  $\sigma \circ \phi$  is a positive hyperbolic complex affine sphere with Blaschke metric  $\phi^*g$  and Pick tensor  $\phi^*C$ . Moreover, if  $\sigma$  has holonomy  $\rho$ , then  $\sigma \circ \phi$  has holonomy  $\rho \circ \pi_1(\phi)$  where  $\pi_1(\phi): \pi_1(\tilde{S}) \rightarrow \pi_1(S)$  is the induced map in homotopy.

In the notation of the space  $\text{CAS}(S)$ ,  $\text{Diff}_+(S)$  acts (on the right) on  $\text{CAS}(S)$  through

$$\phi \cdot (c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) = (\phi^*(c_1), \phi^*(\overline{c_2}), \phi^*(Q_1), \phi^*(\overline{Q_2}), \phi^*u).$$

In fact,  $\phi$  defines an isometry between  $\phi^*(\mathbf{h}_{(c_1, \overline{c_2})}) = \mathbf{h}_{(\phi^*(c_1), \phi^*(\overline{c_2}))}$  and  $\mathbf{h}_{(c_1, \overline{c_2})}$  conjugating the respective Levi-Civita connections, hence for all Bers metric  $h$

$$\phi^*(h(Q_1, \overline{Q_2})) = (\phi^*h)(\phi^*Q_1, \phi^*\overline{Q_2}) \quad \text{and} \quad \phi^*(\Delta_h u) = \Delta_{\phi^*h}\phi^*u,$$

so  $G(\phi \cdot \sigma) = \phi^*(G(\sigma)) = 0$ .

Observe that, for the same reason,  $\phi^*(L_\sigma v) = L_{\phi \cdot \sigma}(\phi^*v)$ , so the action of  $\text{Diff}_+(S)$  restricts to  $\text{CAS}^*(S)$ . Moreover, the action is holomorphic since it is a restriction to a complex Banach submanifold of a holomorphic action on  $\mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) \times W^{k,p}(S)$  (it is actually  $\mathbb{C}$ -linear on the last factor). Restricting to  $\mathcal{CD}(S) \times \mathcal{CD}(\overline{S})$ , it is well known that this action descends to a holomorphic action of  $\text{MCG}(S)$  on  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$ .

**Proposition 6.11.** *The action of  $\text{Diff}_+(S)$  on  $\text{CAS}(S)$  descends to an action of  $\text{MCG}(S)$  on  $\mathbb{CAS}(S)$  and the maps*

$$\begin{array}{ccc} & \mathbb{CAS}(S) & \\ \pi \swarrow & & \searrow \text{hol} \\ \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S}) & & \chi(\pi_1(S), \text{SL}(3, \mathbb{C})) \end{array}$$

are  $\text{MCG}(S)$ -equivariant.

*Proof.* Since the  $\text{Diff}_0(S)$ -orbit of an element in  $\text{CAS}^*(S)$  is path connected, it is contained in a fiber of the projection  $[\hat{\pi}]: \text{CAS}^*(S) \rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$ , so it descends to an action of  $\text{MCG}(S)$  on  $\mathbb{CAS}(S)$ . The rest of the statement follows easily from the above remarks.  $\square$

The other action on  $\text{CAS}(S)$  that we want to highlight is the  $\mathbb{C}^*$ -action defined by

$$\zeta \cdot (c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) = (c_1, \overline{c_2}, \zeta Q_1, \frac{1}{\zeta} \overline{Q_2}, u),$$

which is easily well-defined, as a result of the fact that  $h(\zeta Q_1, \frac{1}{\zeta} \overline{Q_2}) = h(Q_1, \overline{Q_2})$ , and holomorphic. Similarly to above, this action restricts to an action on  $\mathcal{CD}(S) \times \mathcal{CD}(\overline{S})$  that descends to  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$ . These actions are strongly related to a  $\mathbb{C}^*$  action on the moduli space of flat connections (the connection will be explained in Section 7), and in accordance with that action we call it the  $\mathbb{C}^*$  **twistor action**. Observe that the  $\mathbb{C}^*$  twistor action restricts to a well-known  $\mathbb{S}^1$  action on the space of real affine spheres. It is also clear that  $L_\sigma = L_{\zeta \cdot \sigma}$ , hence the action restricts to  $\text{CAS}^*(S)$ .

Finally, if  $\sigma_0$  and  $\sigma_1$  lie in a path-connected component of a fiber of  $[\hat{\pi}]$ , with  $\sigma_t$  denoting a connecting path, then  $\zeta \cdot \sigma_0$  and  $\zeta \cdot \sigma_1$  are connected by  $\zeta \cdot \sigma_t$  and the whole path is contained in a fiber of  $[\hat{\pi}]$ . In conclusion, we get the following.

**Proposition 6.12.** *The  $\mathbb{C}^*$  twistor action on  $\text{CAS}(S)$  descends to a holomorphic action on  $\text{CAS}(S)$  and the map*

$$\pi: \text{CAS}(S) \rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$$

*is equivariant with respect to the  $\mathbb{C}^*$  actions on  $\text{CAS}(S)$  and  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$ .*

**Remark 6.13.** It is important to observe that the  $\text{MCG}(S)$  action and the  $\mathbb{C}^*$  twistor action commute both on  $\text{CAS}(S)$  and  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$ .

**6.5. Proof of Theorem A.** We are now able to prove Theorem A. The main strategy of the proof consists in taking an inverse of the projection  $\pi: \text{CAS}(S) \rightarrow \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$  defined on a big open subset  $\Omega$  of  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$  and then composing it with *hol*.

Recalling the space  $\text{RAS}(S)$  from Section 6.2, define

$$\mathbb{R}\text{AS}(S) := \text{RAS}(S) / \text{Diff}_0(S).$$

As a result of Theorems 6.9 and 6.10, we have two holomorphic maps from  $\text{CAS}(S)$ , namely

$$\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S}) \xleftarrow{\pi} \text{CAS}(S) \xrightarrow{\text{hol}} \chi(\pi_1(S), \text{SL}(3, \mathbb{C})). \quad (37)$$

On the real locus, the maps above restrict to

$$\mathcal{M}_3(S) \cong \Delta_{\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})} \xleftarrow{\sim} \mathbb{R}\text{AS}(S) \xrightarrow{\sim} \text{Hit}_3(S), \quad (38)$$

and the composition of the inverse of the LHS with the RHS coincides with the Labourie-Loftin parametrization  $\mathcal{L}$ .

For all  $(c_1, \overline{c_2}, Q_1, 0) \in \mathcal{CD}(S) \times \mathcal{C}(\overline{S})$ , one can easily see that  $g = \mathbf{h}_{(c_1, \overline{c_2})}$  and  $C = Q_1$  satisfy the hypothesis of Theorem 3.20, so  $\sigma = (c_1, \overline{c_2}, Q_1, 0, 0) \in \text{CAS}(S)$ . Moreover, the linearization  $L_\sigma$  of the Gauss equation (33) is  $L_\sigma = \Delta - 2id$ , which, in the notation of Theorem 5.23, is an isomorphism if and only if  $(c_1, \overline{c_2}) \in \mathcal{B}^{reg}$ : as a result we have a holomorphic map

$$\begin{aligned} \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) \supset W_+ := \{(c_1, \overline{c_2}, Q_1, 0) \mid (c_1, \overline{c_2}) \in \mathcal{B}^{reg}\} &\rightarrow \text{CAS}^*(S) \\ (c_1, \overline{c_2}, Q_1, 0) &\mapsto (c_1, \overline{c_2}, Q_1, 0, 0), \end{aligned} \quad (39)$$

which has  $\hat{\pi}$  as left inverse. Notice that  $W_+$  is a finite-rank vector bundle of  $\mathcal{B}^{reg}$ , and, as a result of Theorem 5.23,  $W_+$  is an open dense subset of  $\mathcal{CD}(S) \times \mathcal{C}(\overline{S})$ . By using the fact that  $\hat{\pi}$  is a local biholomorphism and a left inverse of (39), one can easily see that (39) is a holomorphic embedding with a closed image (as a subset of  $\text{CAS}^*(S)$ ) because it

is a continuous, injective, and closed map. Moreover, by theorem 5.23, the fibers of the projection  $W_+ \rightarrow \mathcal{M}_3(S) \times \mathcal{T}(\bar{S})$  are connected, so (39) descends to a well-defined map

$$s_1: \mathcal{M}_3(S) \times \mathcal{T}(\bar{S}) \xrightarrow{\sim} \mathcal{R} \subset \mathbb{C}\mathbb{A}\mathbb{S}(S), \quad (40)$$

which is injective and continuous (as one can easily see by restricting to local charts as in (36)) and whose image we denote by  $\mathcal{R}$ . Moreover,  $s_1$  is a holomorphic embedding since  $\pi$  is a left inverse and a local biholomorphism. By construction, one can also immediately see that  $s_1$  is equivariant for both the  $\text{MCG}(S)$  action and the  $\mathbb{C}^*$  twistor action.

In the same fashion, the embedding

$$\begin{aligned} W_- := \{((c_1, 0), (\bar{c}_2, \bar{Q}_2)) \mid (c_1, \bar{c}_2) \in \mathcal{B}^{reg}\} &\rightarrow \mathbb{C}\mathbb{A}\mathbb{S}^*(S) \\ (c_1, \bar{c}_2, 0, Q_2) &\mapsto (c_1, \bar{c}_2, 0, \bar{Q}_2, 0) \end{aligned} \quad (41)$$

descends to the holomorphic embedding

$$s_2: \mathcal{T}(S) \times \mathcal{M}_3(\bar{S}) \xrightarrow{\sim} \bar{\mathcal{R}} \subset \mathbb{C}\mathbb{A}\mathbb{S}(S). \quad (42)$$

which is equivariant for both  $\text{MCG}(S)$  and  $\mathbb{C}^*$ , and whose image we denote by  $\bar{\mathcal{R}}$ .

It is easy to check that, if  $\sigma: \tilde{S} \rightarrow \mathbb{C}^3$  is the complex affine sphere corresponding to the data  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u)$ , then  $\bar{\sigma}$  has conjugate holonomy and corresponds to  $(c_2, \bar{c}_1, Q_2, \bar{Q}_1, \bar{u})$ , motivating the notation for  $\bar{\mathcal{R}}$ .

Finally, observe that  $s_1$  and  $s_2$  coincide on  $\mathcal{T}(S) \times \mathcal{T}(\bar{S})$ , defining a diffeomorphism between  $\mathcal{T}(S) \times \mathcal{T}(\bar{S})$  and  $\mathcal{R} \cap \bar{\mathcal{R}}$ , which, by Example 3.24, corresponds to Bers embeddings in  $\hat{\mathbb{X}}_2$ . As a consequence,

$$hol \circ (s_1 \cap s_2): \mathcal{T}(S) \times \mathcal{T}(\bar{S}) \rightarrow \chi(S, SL(3, \mathbb{C}))$$

coincides with Bers parametrization of the quasi-Fuchsian space, after identifying  $\text{PSL}(2, \mathbb{C})$  with  $\text{SO}(2, 1, \mathbb{C}) \cong \text{Isom}_0(\hat{\mathbb{X}}_2) \subset \text{SL}(3, \mathbb{C})$ .

Theorem A will follow as a corollary of the following theorem.

**Theorem 6.14.** *There exists an open subset  $\Omega \subset \mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$  containing the diagonal,  $\mathcal{M}_3(S) \times \mathcal{T}(\bar{S})$ , and  $\mathcal{T}(S) \times \mathcal{M}_3(\bar{S})$  over which  $\pi$  admits a global biholomorphic inverse  $\varsigma: \Omega \rightarrow \mathbb{C}\mathbb{A}\mathbb{S}(S)$  which extends  $s_1$  and  $s_2$  above, and the real parametrization  $\Delta \rightarrow \text{RAS}(S)$ .*

*Moreover,  $\Omega$  can be chosen to be invariant for the actions of both  $\text{MCG}(S)$  and  $\mathbb{C}^*$ .*

Before proving Theorem 6.14, we show how it implies Theorem A.

*Proof of Theorem A.* Consider the inverse  $\varsigma: \Omega \rightarrow \mathbb{C}\mathbb{A}\mathbb{S}(S)$  for  $\pi$  given by Theorem 6.14 and compose it with the holonomy map to get  $hol \circ \varsigma: \mathbb{C}\mathbb{A}\mathbb{S}(S) \rightarrow \chi(S, SL(3, \mathbb{C}))$ . By construction, this map is holomorphic and  $\text{MCG}(S)$ -equivariant. Moreover,  $\mathcal{L} = hol \circ \varsigma|_{\Delta}$ , and  $hol \circ \varsigma|_{\mathcal{T}(S) \times \mathcal{T}(\bar{S})} = hol \circ (s_1 \cap s_2)$  which, as shown above, coincides with Bers' Theorem after identifying  $\text{PSL}(2, \mathbb{C})$  with  $\text{SO}(2, 1, \mathbb{C})$ .  $\square$

The main tool in the proof of Theorem 6.14 is the generalized inverse function theorem, which says that if a local diffeomorphism restricts to a diffeomorphism along a submanifold whose image is a submanifold, then it restricts to a diffeomorphism between connected neighborhoods of them (see Exercise 1.8.14 in [GP10] and [Wan]). Theorem 6.14 is much easier to prove if we don't demand that  $\Omega$  is  $\mathbb{C}^*$  and  $\text{MCG}(S)$ -invariant. However, the proof for the action-invariant version will require some attention on some topological issues and on the choice of the neighborhoods. In particular, to construct the open subsets, it's useful to recall the  $\text{MCG}(S)$ -invariant Riemannian metric on  $\mathcal{M}_3(S)$  defined in [KZ17] and [Lab17], which we call  $\underline{g}$  in the proof below. As a result,  $\underline{g} \oplus \underline{g}$  defines a Riemannian metric on  $\mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$  for which the factors  $\mathcal{M}_3(S) \times \{0\}$  and  $\{0\} \times \mathcal{M}_3(\bar{S})$  are totally geodesic.

Recall a subset  $U$  of a vector space is star-shaped if there is a point  $v \in U$ , called the center, such that for every point  $v' \in U$ , the segment between  $v$  and  $v'$  is contained in  $U$ . We call a subset of  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$  geodesically star-shaped if it's star-shaped with respect to geodesics for  $g \oplus g$ .

*Proof of Theorem 6.14.* By the generalized inverse function theorem mentioned above, there exist  $V_1, V_2 \subset \mathbb{C}\mathcal{A}\mathcal{S}(S)$ , and  $U_1, U_2 \subset \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$  open subsets such that  $\pi$  restricts to the biholomorphisms

$$\begin{aligned} \mathcal{R} \cap \overline{\mathcal{R}} &\subset V_1 \xrightarrow{\sim} U_1 \supset \mathcal{T}(S) \times \mathcal{T}(\overline{S}) \\ \mathbb{R}\mathcal{A}\mathcal{S}(S) &\subset V_2 \xrightarrow{\sim} U_2 \supset \Delta. \end{aligned}$$

We denote the inverse maps by  $\varsigma_1$  and  $\varsigma_2$  respectively.

Up to restricting these open subsets, we can make the following assumptions on  $U_1$  and  $U_2$ , which we denote by *assumptions* (\*).

*Assumption* (\*) on  $U_1$ .

The intersections between  $U_1$  and submanifolds of the form  $\mathcal{C}\mathcal{D}([c_1]) \times \mathcal{C}\mathcal{D}([\overline{c}_2]) \subset \mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S})$ , which can be seen as open subsets in a vector space, are star-shaped open subsets with center in  $([c_1, 0], [\overline{c}_2, 0])$ . Such open subsets can be obtained by taking local trivializations of the bundle  $\mathcal{M}_3(S) \times \mathcal{M}_3(\overline{S}) \rightarrow \mathcal{T}(S) \times \mathcal{T}(\overline{S})$ .

*Assumption* (\*) on  $U_2$ .

If both  $m = ([c_1, \mathbb{Q}_1], [\overline{c}_2, \overline{\mathbb{Q}}_2])$  and  $\zeta \cdot m = ([c_1, \zeta \mathbb{Q}_1], [\overline{c}_2, \frac{1}{\zeta} \overline{\mathbb{Q}}_2])$  lie in  $U_2$ , then  $\varsigma_2(\zeta \cdot m) = \zeta \cdot \varsigma_2(m)$ : to realize this, observe that on  $\mathbb{C}^* \times U_2$  the functions  $(\zeta, m) \mapsto \varsigma_2(\zeta \cdot m)$  and  $(\zeta, m) \mapsto \zeta \cdot \varsigma_2(m)$  are holomorphic maps that agree on the totally real submanifold  $\mathbb{S}^1 \times \Delta$  as a result of  $\varsigma_2|_{\Delta}$  being  $\mathbb{S}^1$ -invariant, so, up to shrinking  $U_2$ , they coincide on some open subset  $W_0 \times U_2$  (hence on its intersection with the preimage of  $U_2$  itself).

Moreover, we assume that the intersections between  $U_2$  and the submanifolds of the form  $\{[c_1, \mathbb{Q}_1]\} \times \mathcal{M}_3(\overline{S})$  are geodesically star-shaped for the metric  $g \oplus g$  with center in  $([c_1, \mathbb{Q}_1], [\overline{c}_1, \overline{\mathbb{Q}}_1])$ : this can be constructed (within the previous refinements) through the exponential map and the fact that the factors  $\mathcal{M}_3(S) \times \{0\}$  and  $\{0\} \times \mathcal{M}_3(\overline{S})$  are totally geodesic makes the construction more simple.

*Step 1.* We prove that if  $\hat{V}_i \rightarrow \hat{U}_i$  is another biholomorphic restriction with  $U_i$  satisfying assumption (\*), then its inverse coincides with the inverse of  $V_i \rightarrow U_i$  on the intersection  $U_i \cap \hat{U}_i$ . We prove it for  $i = 1$ , but the proof works in the very same fashion for  $i = 2$ . If  $([c_1, \mathbb{Q}_1], [c_2, \overline{\mathbb{Q}}_2]) \in U_1 \cap \hat{U}_1$ , then by assumption (\*) the path  $([c_1, t\mathbb{Q}_1], [c_2, t\overline{\mathbb{Q}}_2])$  lies in the intersection for all  $t \in [0, 1]$  (for  $i = 2$ , take the geodesic contained in the leaf  $\{[c_1, \mathbb{Q}_1]\} \times \mathcal{M}_3(\overline{S})$  connecting it to the intersection point with  $\Delta$ ). Let  $\hat{\varsigma}_1$  be the inverse of  $\hat{V}_1 \rightarrow \hat{U}_1$ . Now, the set

$$\{t \in [0, 1] \mid \hat{\varsigma}_1([c_1, t\mathbb{Q}_1], [c_2, t\overline{\mathbb{Q}}_2]) = \varsigma_1([c_1, t\mathbb{Q}_1], [c_2, t\overline{\mathbb{Q}}_2])\}$$

contains  $t = 0$  and is closed; if by contradiction its maximum  $T$  were less than 1, inverting  $\pi$  in a neighborhood of  $\varsigma_1([c_1, T\mathbb{Q}_1], [c_2, T\overline{\mathbb{Q}}_2])$  we get that such a  $T$  can be extended a bit further, reaching a contradiction. We conclude that the inverses coincide in the intersection.

*Step 2.* We extend the  $U_i$ 's a bit further.

In fact, observe that, for all  $\phi \in \text{MCG}(S)$ ,  $\phi \cdot U_i$  satisfies assumption (\*) where the inverses are given by  $\phi \circ \varsigma \circ \phi^{-1}: \phi \cdot U_i \rightarrow \phi \cdot V_i$ .

Moreover, for all  $\zeta \in \mathbb{C}^*$ ,  $\zeta \cdot U_1$  satisfies assumption (\*) and an inverse of  $\pi$  is given by  $\zeta \circ \varsigma \circ \frac{1}{\zeta}: \zeta \cdot U_1 \rightarrow \zeta \cdot V_1$ . As a result of the previous step, we get that the inverses match

on the intersections. Therefore, recalling that the  $MCG(S)$  and the  $\mathbb{C}^*$  twistor actions commute,  $\pi$  admits a biholomorphic inverse on the restrictions

$$\begin{aligned} V'_1 &:= \bigcup_{\phi \in MCG(S)} \bigcup_{\zeta \in \mathbb{C}^*} \phi \cdot \zeta \cdot V_1 \rightarrow \bigcup_{\phi \in MCG(S)} \bigcup_{\zeta \in \mathbb{C}^*} \phi \cdot \zeta \cdot U_1 := U'_1 \\ V'_2 &:= \bigcup_{\phi \in MCG(S)} \phi \cdot V_2 \rightarrow \bigcup_{\phi \in MCG(S)} \phi \cdot U_1 := U'_1, \end{aligned}$$

whose inverses we still denote by  $\varsigma_1$  and  $\varsigma_2$  respectively.

Observe that  $V'_1 \rightarrow U'_1$  restricts to  $\mathcal{R} \rightarrow \mathcal{M}_3(S) \times \mathcal{T}(\overline{S})$  and  $\overline{\mathcal{R}} \rightarrow \mathcal{T}(S) \times \mathcal{M}_3(\overline{S})$  because for all  $[(c_1, \overline{c_2}, Q_1, 0, 0)] \in \mathcal{R}$  – resp.  $[(c_1, \overline{c_2}, 0, \overline{Q_2}, 0)] \in \overline{\mathcal{R}}$  – there exists  $\zeta$  small enough such that  $[(c_1, \overline{c_2}, \zeta Q_1, 0, 0)]$  – resp.  $[(c_1, \overline{c_2}, 0, \zeta \overline{Q_2}, 0)]$  – lies in  $V_1$ .

Notice that a priori it does not seem clear whether  $\varsigma_1$  and  $\varsigma_2$  agree on  $U'_1 \cap U'_2$ .

*Step 3.* We glue  $U'_1$  and a neighborhood of  $\Delta$ .

The restriction  $\pi|_{V'_1}$  coincides with the real analytic map  $\mathbb{R}AS(S) \rightarrow \Delta$  on the intersection  $V'_1 \cap \mathbb{R}AS(S)$ : by uniqueness,  $\varsigma_1$  must coincide with  $\varsigma_2$  in a neighborhood  $W \subset U'_1$  of  $\Delta \cap U'_1$ . Extend now  $W$  out of  $U'_1$  to an open neighborhood of  $\Delta$  (to construct this extension, one could take a thin enough neighborhood of  $\Delta$  disjoint from the closure of  $U'_1 \setminus W$  and then unionize it with  $W$ ), which we still denote by  $W$ . Up to shrinking it, we can also assume  $W$  to be contained inside  $U'_2$ , and to satisfy assumption (\*). By construction, the restriction of  $\varsigma_2$  on  $W$  coincides with  $\varsigma_1$  on  $W \cap U'_1$ .

Finally, by the fact that  $U'_2$  is  $MCG(S)$ -invariant and that  $\varsigma_2$  is  $MCG(S)$ -equivariant,  $\phi \cdot W \subset U'_2$  for all  $\phi \in MCG(S)$  and  $\varsigma_2$  is well-defined on

$$U''_2 := \bigcup_{\phi \in MCG(S)} \phi \cdot W,$$

whose image we denote by  $V''_2 := \varsigma(U''_2) \subset V'_2$ .

Moreover,  $\varsigma_1$  and  $\varsigma_2$  coincide on  $U''_2 \cap U'_1$ : in fact, if by contradiction there existed a point in which they were different, then, since both sections are  $MCG(S)$ -equivariant, one would find a point in  $W \cap U'_1$  in which they are different, giving a contradiction.

We call  $\varsigma$  the inverse constructed by patching  $\varsigma_1$  and  $\varsigma_2$  on  $U'_1 \cup U''_2$ .

*Step 4.* We prove that  $\varsigma$  can be extended to

$$\Omega := U'_1 \cup \bigcup_{\zeta \in \mathbb{C}^*} \zeta \cdot U''_2,$$

concluding the proof.

On each  $\zeta \cdot U''_2$ , consider the inverse  $\varsigma_2^\zeta := \zeta \circ \varsigma_2 \circ \frac{1}{\zeta}: \zeta \cdot U''_2 \rightarrow \zeta \cdot V''_2$  for  $\pi$ .

For all  $\zeta \in \mathbb{C}^*$ , on  $\zeta \cdot U''_2 \cap U'_1$ ,  $\varsigma_2^\zeta$  and  $\varsigma_1$  coincide, since if they didn't then one would find a point of  $U''_2 \cap U'_1$  in which they don't coincide, because  $\varsigma_1$  is  $\mathbb{C}^*$ -equivariant.

We prove that different  $\varsigma_2^\zeta$ 's coincide on the intersection. Recall that on  $U_2$  the inverse  $\varsigma_2$  is such that  $\varsigma_2(\zeta \cdot m) = \zeta \cdot \varsigma_2(m)$  when both  $m$  and  $\zeta \cdot m$  lie in  $U_2$ ; since the  $MCG(S)$  and the  $\mathbb{C}^*$  actions commute, this property holds also on  $U'_2$ , hence on  $U''_2$ . Finally, if  $\varsigma_2^\zeta$  and  $\varsigma_2^{\hat{\zeta}}$  did not coincide on  $(\zeta \cdot U''_2) \cap (\hat{\zeta} \cdot U''_2)$ , then (by applying  $\frac{1}{\zeta}$ ) we would have that  $\varsigma_2$  and  $\varsigma_2^{\hat{\zeta}}$  do not coincide on  $U'' \cap (\hat{\zeta} \cdot U''_2)$ , contradicting the assumption (\*) for  $U$ . □

**6.6. Proof of Theorem C.** Here we prove Theorem C. As stated in Section 1.2, to do so we analyze one of the simpler families of points in  $\mathcal{CD}(S) \times \mathcal{CD}(\overline{S})$ , namely rays

$$[0, \infty) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}), t \mapsto (c, \overline{c}, tQ, -t\overline{Q}).$$

The Gauss equation (17) is

$$\Delta_h u = e^{2u} + \frac{t^2}{4} e^{-4u} h(Q, \overline{Q}) - 1, \quad (43)$$

which, when restricted to real functions  $u$ , coincides with the structural equation for minimal Lagrangian immersions into the 2-dimensional complex hyperbolic space  $\mathbb{C}\mathbb{H}^2$ . Minimal Lagrangians in  $\mathbb{C}\mathbb{H}^2$  have been studied in a number of works, such as [LM13] and [HLL13], and we can use known results directly.

We need only part (3) of the following theorem about real solutions, which is proved by a calculation using Gauss-Bonnet, but we include parts (1) and (2) because they shed light on the theory of complex affine spheres. Note that in previous works on Equation (43), authors have considered only real solutions. Here we are allowed to have complex solutions. So when stating previous results, we will be sure to specify that they apply only for real solutions.

**Theorem 6.15** (Theorem 1.1 in Huang-Loftin-Lucia [HLL13]). *Let  $c$  be a complex structure on  $S$  and  $Q$  a  $c$ -holomorphic cubic differential. There are constants  $T_0 = T_0(c, Q) > 0$  and  $T = T(c, Q) > T_0$  such that the following hold.*

- (1) *For any  $t \in [0, T_0)$ , there are at least two real solutions to (43).*
- (2) *For  $t = T_0$ , there is a real solution to (43).*
- (3) *For  $t > T$ , there are no real solutions to (43).*

One hint toward Theorem 6.15 is that Equation (43) strongly resembles the better-known Gauss equation for minimal surfaces in  $\mathbb{H}^3$ , which is not particularly well behaved (see [Uhl83]). Heuristically, issues should arise because the linearization at a solution  $\sigma$  is expected to have a kernel for  $t$  large. Indeed, at a solution  $\sigma = (c, \overline{c}, tQ, t\overline{Q}, u)$ , the linearization is given by

$$L_{\sigma_t}(v) = \Delta_h v - v(e^u - t^2 e^{-4u} h(Q, \overline{Q})). \quad (44)$$

For  $Q$  very large, the expression in the brackets in (44) is negative. If we pretend that expression is a negative constant, then Equation (44) becomes the equation for eigenfunctions of the Laplacian, which could have a non-zero solution. The authors in [HLL13] realize this expectation by showing that the linearization does have non-empty kernel at the solution at time  $T_0$ .

As in [HLL13], one can construct a nice path of real solutions around  $(c, \overline{c}, 0, 0, 0)$ . Define  $G_{\mathbb{R}} : \mathcal{CD}(S) \times C^\infty(S, \mathbb{R}) \rightarrow C^\infty(S, \mathbb{R})$  by  $G_{\mathbb{R}}(c, Q, u) = G(c, \overline{c}, Q, -\overline{Q}, u)$ . Taking  $Q = 0$  and  $u = 0$  and extending  $G_{\mathbb{R}}$  to all of the Banach manifolds  $\mathcal{CD}^l(S) \times W^{k+2,2}(S, \mathbb{R})$ , the linearization of  $G_{\mathbb{R}}$  is an isomorphism in the direction of  $u$ , and it follows that there is a unique continuous path into  $C^\infty(S, \mathbb{R})$  of real solutions  $t \mapsto u_t$  of  $G(c, \overline{c}, tQ, -t\overline{Q}, u_t) = 0$ , for  $t$  in some small interval  $(-\varepsilon, \varepsilon)$ .

Theorem C follows as a consequence of the following Proposition.

**Proposition 6.16.** *There exists no global holomorphic map  $\varsigma : \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\overline{S}) \times C^\infty(S, \mathbb{C})$  whose image is contained in  $\text{CAS}(S)$ , and which inverts  $\pi$  and maps any  $(c, \overline{c}, 0, 0)$  to  $(c, \overline{c}, 0, 0, 0)$ .*

*Proof.* Denote  $\sigma_t = (c, \overline{c}, tQ, -t\overline{Q}, u_t) := \varsigma(c, \overline{c}, tQ, -t\overline{Q})$ . By the fact that  $L_{\sigma_0}$  is invertible,  $\sigma_0 \in \text{CAS}^*(S)$ , and since  $\text{CAS}^*(S) \rightarrow \mathcal{CD}(S) \times \mathcal{CD}(\overline{S})$  is a local homeomorphism, we get



that, for  $t > 0$  small enough,  $u_t$  must be the real solution to  $G = 0$  mentioned above, just because they can be continuously deformed to  $t = 0$ . Moreover, since  $\zeta$  is holomorphic, the path  $\zeta(c, \bar{c}, tQ, -t\bar{Q}, u_t)$  is real analytic with respect to  $t$ , hence at each point  $p \in S$ ,  $u_t(p)$  being real for small  $t$  implies it being real for all  $t > 0$ . This would produce a solution  $u_t$  for  $t > T$ , contradicting Theorem 6.15.  $\square$

**Remark 6.17.** We expect that, as one can possibly see by writing the connection forms explicitly or by means of the perspective of [RT24], complex affine spheres projecting to  $(c, \bar{c}, Q, -\bar{Q}) \in \mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$  with Riemannian Blaschke metric have the same holonomy as the corresponding minimal Lagrangian immersion in  $\mathbb{CH}^2$ .

## 7. BI-HIGGS BUNDLES AND OPERS

With the proof of Theorem A complete, we move on to study further connections. In this section we define complex harmonic maps into  $SL(n, \mathbb{C})/SO(n, \mathbb{C})$  and bi-Higgs bundles, and relate them to complex affine spheres via Theorem D. We use Theorem D to prove Theorem E. We won't give the basic definitions on harmonic maps and Higgs bundles, and they won't be needed, although having some prior knowledge would certainly be helpful. For sources that are in line with our presentation, see [Gui18] and [Li19].

**7.1. Complex harmonic maps.** Let  $(S, c)$  be a closed Riemann surface of genus  $g \geq 2$  and let  $G$  be a real semisimple Lie group with finite center. The non-abelian Hodge correspondence for  $G$  associates reductive representations from  $\pi_1(S) \rightarrow G$  to  $G$ -Higgs bundles over  $S$ . In a nutshell, from the representation one builds an equivariant harmonic map  $f$  to a Riemannian symmetric space of  $G$ , and then the holomorphic derivative of  $f$  determines the data of a Higgs bundle.

For  $G = SL(n, \mathbb{R})$ , the symmetric space  $Y_n$  is modelled by the space of inner products on  $\mathbb{R}^n$  that induce the unit inner product on  $\wedge^n \mathbb{R}^n = \mathbb{R}$ , and, for  $SL(n, \mathbb{C})$ , the symmetric space models Hermitian forms on  $\mathbb{C}^n$  that induce the unit form on  $\wedge^n \mathbb{C}^n = \mathbb{C}$ . The embedding of symmetric spaces is realized by taking the sesquilinear extensions of inner products. Upon doing this embedding, a  $SL(n, \mathbb{R})$ -Higgs bundle can be seen as a  $SL(n, \mathbb{C})$ -Higgs bundle that's compatible with an  $SO(n, \mathbb{C})$ -structure.

We go a different route, and embed  $Y_n$  into the homogeneous space  $SL(n, \mathbb{C})/SO(n, \mathbb{C})$ , which we denote by  $\mathbb{Y}_n$ , and which we'll see produces a different Higgs bundle-like object. Concretely, the space  $\mathbb{Y}_n$  models non-degenerate bilinear forms on  $\mathbb{C}^n$  that induce 1 on  $\wedge^n \mathbb{C}^n = \mathbb{C}$ , and in our models the embedding from  $Y_n \rightarrow \mathbb{Y}_n$  is realized by taking the bilinear extension of an inner product. Since  $SO(n, \mathbb{C})$  is a complex Lie group, the complex Killing form on  $SL(n, \mathbb{C})$  descends to a holomorphic Riemannian metric  $\langle \cdot, \cdot \rangle$  on  $\mathbb{Y}_n$ . Furthermore, the involution of the Lie algebra given by  $A \mapsto -A^T$  determines a holomorphic involution of  $\mathbb{Y}_n$ , and hence we refer to  $(\mathbb{Y}_n, \langle \cdot, \cdot \rangle)$  as a holomorphic Riemannian symmetric space.

More abstractly, let  $K$  be a maximal compact subgroup of  $G$ , let  $K^{\mathbb{C}}$  and  $G^{\mathbb{C}}$  be the complexifications, and let  $K'$  be a maximal compact of  $G^{\mathbb{C}}$ . Instead of embedding  $G/K$  into  $G^{\mathbb{C}}/K'$ , we've taken  $G/K$  to  $G^{\mathbb{C}}/K^{\mathbb{C}}$ .

We're almost ready to define our notion of harmonic maps to  $\mathbb{Y}_n$ , but first we'll add one more layer of complication (or, rather, complexification). Given  $(c_1, \bar{c}_2)$ , let  $g$  be a positive complex metric in the conformal class of the Bers metric  $h_{(c_1, \bar{c}_2)}$ . The derivative  $df$  defines a section of the endomorphism bundle  $\mathbb{C}T^*\tilde{S} \otimes f^*T\mathbb{Y}_n$ , the pullback complex metric  $f^*\langle \cdot, \cdot \rangle$  – which we will often just denote by  $\langle \cdot, \cdot \rangle$  – descends to  $S$ , and, as would work in a Riemannian setting, the Levi-Civita connections  $\nabla^g$  for  $g$  on  $\mathbb{C}T^*S$  and  $\nabla^f$  for  $f^*\langle \cdot, \cdot \rangle$  on

$f^*T\mathbb{Y}_n$  determine a torsion-free connection  $\hat{\nabla}^f := \nabla^{g \otimes f^* \langle \cdot, \cdot \rangle}$  on the endomorphism bundle  $\mathbb{C}T^*S \otimes f^*T\mathbb{Y}_n$  that makes  $g^* \otimes f^* \langle \cdot, \cdot \rangle$  parallel.

**Definition 7.1.** An equivariant map  $f : (\tilde{S}, c_1, \bar{c}_2) \rightarrow (\mathbb{Y}_n, \langle \cdot, \cdot \rangle)$  is **complex harmonic** if, for any positive complex metric  $g$  in the conformal class of  $\mathbf{h}_{(c_1, \bar{c}_2)}$ ,

$$\mathrm{tr}_g \nabla^{g \otimes f^* \langle \cdot, \cdot \rangle} df = 0. \quad (45)$$

Note that Equation (45) is conformally invariant—if it holds for one  $g$ , then it holds for every  $g$  in the conformal class. When  $f$  lands in the Riemannian symmetric space and  $g$  is the complexification of a Riemannian metric, Definition 7.1 agrees with the usual definition of harmonic maps. As we've discussed in the introduction, we'll develop the theory of complex harmonic maps in future work, and for now we just cover what we need.

**7.2. Complex bi-gradings.** To prepare for discussing bi-Higgs bundles, we introduce complex bi-gradings. Recall that if  $g$  is a positive complex metric with complex structures  $c_1$  and  $\bar{c}_2$  with local holomorphic coordinates  $z$  and  $\bar{w}$  respectively, we have the projections  $\Pi_1, \Pi_2$  of  $\mathbb{C}TS$  with respect to the decomposition  $\mathrm{Span}(dz) \oplus \mathrm{Span}(\bar{d}w)$ , corresponding to the isotropic directions of the dual metric  $g$  in  $\mathbb{C}T^*S$ . These projections induce a splitting on  $\Omega^1(S)$  that we write as

$$\Omega^1(S) = \Omega_{c_1}^{1,0}(S) \oplus \Omega_{c_2}^{0,1}(S),$$

where  $\Omega_{c_1}^{1,0} = \Pi_1 \Omega^1$  and  $\Omega_{c_2}^{0,1} = \Pi_2 \Omega^1$ . To understand these operators, we record that for a 1-form  $\omega$ ,

$$\Pi_1 \omega = \frac{\omega(\partial_w)}{\partial_w z} dz, \quad \Pi_2 \omega = \frac{\omega(\partial_{\bar{z}})}{\partial_{\bar{z}} \bar{w}} d\bar{w}, \quad (46)$$

which can be verified directly. Note that, when  $c_1 = c_2$ , the two splittings agree and are the ordinary splitting of  $\Omega^1$  into  $(1, 0)$  and  $(0, 1)$  forms. We further define  $\Omega_{c_1, c_2}^{1,1} = \Omega_{c_1}^{1,0} \otimes \Omega_{c_2}^{0,1}$ . We will write  $\Omega_{c_1, c_2}^{p,q}(S)$ , where  $(p, q)$  can be  $(1, 0)$ ,  $(0, 1)$ , or  $(1, 1)$ , and for vector bundles  $E$ ,  $\Omega_{c_1, c_2}^{p,q}(E)$  will be the associated space of  $E$ -valued  $(p, q)$ -forms.

Writing  $d$  for the ordinary exterior derivative on functions, we split it as

$$d = \Pi_1 \circ d + \Pi_2 \circ d =: \partial_{c_1} + \bar{\partial}_{c_2}.$$

Both  $\partial_{c_1}$  and  $\bar{\partial}_{c_2}$  extend to  $\Omega^1(S)$  by the rules  $\partial_{c_1} = d \circ \Pi_2$  and  $\bar{\partial}_{c_2} = d \circ \Pi_1$ . Equivalently, we have

$$\partial_{c_1}(fdz) = 0, \quad \partial_{c_1}(f\bar{d}w) = \partial_{c_1} f \wedge \bar{d}w = d(f\bar{d}w),$$

and

$$\partial_{c_1}(f\bar{d}w) = 0, \quad \bar{\partial}_{c_2}(fdz) = \bar{\partial}_{c_2} f \wedge dz = d(fdz).$$

In the Riemannian setting, a map  $f$  is harmonic if and only if its holomorphic derivative lies in the kernel of  $(\hat{\nabla}^f)^{0,1}$ . Using the complex bi-grading, we obtain the analogous result for complex harmonic maps. Denote  $\nabla^g, \nabla^f, \hat{\nabla}^f$  as above.

**Proposition 7.2.**  $f : (\tilde{S}, c_1, \bar{c}_2) \rightarrow (\mathbb{Y}_n, \langle \cdot, \cdot \rangle)$  is harmonic if and only if  $(\Pi_2 \circ \hat{\nabla}^f) \partial_{c_1} f = 0$ .

By reversing the roles of our complex metrics, we see as well that harmonicity is equivalent to  $(\Pi_1 \circ \hat{\nabla}^f) \bar{\partial}_{c_2} f = 0$ . Proposition 7.2 follows from the lemma below, which we'll see to be quite useful.

**Lemma 7.3.** Denoting  $g = \lambda dz d\bar{w}$ ,

$$(\Pi_2 \circ \hat{\nabla}^f) \partial_{c_1} f = (\mathrm{tr}_g \hat{\nabla}^f df) \lambda dz \otimes \bar{d}w.$$

*Proof.* First, we expand  $\text{tr}_g \hat{\nabla} df$  in the isotropic basis. Recalling from the proof of Lemma 5.1 that  $\nabla_{\partial_{\bar{z}}}^g \partial_w = \frac{\partial_{\bar{z}} \partial_w z}{\partial_w z} \partial_w$ ,

$$\begin{aligned} \text{tr}_g \hat{\nabla}^f df &= \frac{2}{g(\partial_{\bar{z}}, \partial_w)} (\nabla_{\partial_{\bar{z}}}^f (df(\partial_w)) - df(\nabla_{\partial_{\bar{z}}}^g \partial_w)) \\ &= \frac{2\partial_w z}{g(\partial_{\bar{z}}, \partial_w)} (\nabla_{\partial_{\bar{z}}}^f (df(\partial_w)) - \frac{\partial_{\bar{z}} \partial_w z}{\partial_w z} df(\partial_w)) \\ &= \frac{2\partial_w z}{g(\partial_{\bar{z}}, \partial_w)} \nabla_{\partial_{\bar{z}}}^f \left( \frac{df(\partial_w)}{\partial_w z} \right). \end{aligned}$$

Before expanding  $(\Pi_2 \circ \hat{\nabla}^f) \partial_{c_1} f$ , observe that  $\nabla_{\partial_{\bar{z}}}^g dz = 0$ . Indeed,

$$(\nabla_{\partial_{\bar{z}}}^g dz)(\partial_z) = \partial_{\bar{z}}(dz(\partial_z)) - dz(\nabla_{\partial_{\bar{z}}}^g \partial_z) = 0,$$

and

$$(\nabla_{\partial_{\bar{z}}}^g dz)(\partial_{\bar{z}}) = \partial_{\bar{z}}(dz(\partial_{\bar{z}})) - dz(\nabla_{\partial_{\bar{z}}}^g \partial_{\bar{z}}) = 0,$$

where we used the parallelity from Lemma 2.5 to say  $dz(\nabla_{\partial_{\bar{z}}}^g \partial_z) = 0$  and  $dz(\nabla_{\partial_{\bar{z}}}^g \partial_{\bar{z}}) = 0$ . Thus, using Equation (46),

$$\begin{aligned} (\Pi_2 \circ \hat{\nabla}^f) \partial_{c_1} f &= \hat{\nabla}_{\partial_{\bar{z}}}^f \left( \frac{df(\partial_w)}{\partial_w z} dz \right) \otimes \frac{d\bar{w}}{\partial_{\bar{z}} \bar{w}} \\ &= \left( \nabla_{\partial_{\bar{z}}}^f \left( \frac{df(\partial_w)}{\partial_w z} \right) \cdot dz + \frac{df(\partial_w)}{\partial_w z} \nabla_{\partial_{\bar{z}}}^g dz \right) \otimes \frac{d\bar{w}}{\partial_{\bar{z}} \bar{w}} \\ &= \nabla_{\partial_{\bar{z}}}^f \left( \frac{df(\partial_w)}{\partial_w z} \right) \otimes dz \otimes \frac{d\bar{w}}{\partial_{\bar{z}} \bar{w}}. \end{aligned}$$

Rearranging yields

$$(\Pi_2 \circ \hat{\nabla}^f) \partial_{c_1} f = (\text{tr}_g \hat{\nabla}^f df) \cdot \frac{2\partial_{\bar{z}} \bar{w} \partial_w z}{g(\partial_{\bar{z}}, \partial_w)} dz \otimes d\bar{w} = (\text{tr}_g \hat{\nabla}^f df) \lambda dz \otimes d\bar{w},$$

as desired.  $\square$

We also have a notion of Hopf differentials for complex harmonic maps. Decomposing  $\langle \cdot, \cdot \rangle$  according to the splitting  $\text{Span}(dz d\bar{w}) \oplus \text{Span}(dz^2) \oplus \text{Span}(d\bar{w}^2)$ , we get

$$f^* \langle \cdot, \cdot \rangle = \|df\|_f^2 g + q_1 + \bar{q}_2 \quad (47)$$

with  $q_1 = \langle \partial_{c_1} f, \partial_{c_1} f \rangle$  and  $\bar{q}_2 = \langle \bar{\partial}_{c_2} f, \bar{\partial}_{c_2} f \rangle$ . By analogy with the Riemannian case, we say that  $f$  is **weakly conformal** if  $f^* \langle \cdot, \cdot \rangle$  is a non-zero multiple of  $g$ , hence of  $\mathbf{h}_{(c_1, \bar{c}_2)}$ , on the complement of its degenerate set. Note that  $f$  is weakly conformal if and only if  $q_1 = 0$  and  $\bar{q}_2 = 0$ .

**Proposition 7.4.** *If  $f$  is harmonic,  $q_1$  and  $\bar{q}_2$  are holomorphic quadratic differentials for  $c_1$  and  $\bar{c}_2$  respectively.*

*Proof.* By the expansion of  $(\Pi_2 \circ \hat{\nabla}^f) \partial_{c_1} f$  in Lemma 7.3, we see that  $f$  harmonic implies  $\hat{\nabla}_{\partial_{\bar{z}}}^f \left( \frac{df(\partial_w)}{\partial_w z} dz \right) = 0$ . Also, recalling that  $g(dz, dz) = 0$  and  $\nabla_{\partial_{\bar{z}}}^g dz = 0$ , we get that  $q_1 = \varphi dz^2$  is such that

$$\begin{aligned} \partial_{\bar{z}} \varphi &= \partial_{\bar{z}} \langle (\partial_{c_1} f)(\partial_z), (\partial_{c_1} f)(\partial_z) \rangle = 2 \langle D_{\partial_{\bar{z}}}((\partial_{c_1} f)(\partial_z)), (\partial_{c_1} f)(\partial_z) \rangle \\ &= 2 \langle (\hat{\nabla}_{\partial_{\bar{z}}}^f (\partial_{c_1} f))(\partial_z), ((\partial_{c_1} f)(\partial_z)) \rangle = 0. \end{aligned}$$

The proof is analogous for  $\bar{q}_2$ .  $\square$

**7.3. bi-Higgs bundles.** Now we turn complex harmonic maps into Higgs bundle data. A lot of the content here mirrors the setup for ordinary Higgs bundles, which is very well explained in [Li19, Section 2], so we're not going to linger on every detail.

Given a representation  $\rho : \pi_1(S) \rightarrow \mathrm{SL}(n, \mathbb{C})$ , we form the bundle  $E_\rho = \tilde{S} \times \mathbb{C}^n / \rho$  over  $S$  associated with the action of  $\rho$  on  $\mathbb{C}^n$ , which carries a flat  $\mathrm{SL}(n, \mathbb{C})$ -connection  $D$ . The bundle  $\tilde{S} \times \mathbb{Y}_n / \rho$  associated with the action of  $\rho$  on  $\mathbb{Y}_n$  identifies with a bundle of non-degenerate  $\mathbb{C}$ -bilinear forms on  $E_\rho$ . A  $\rho$ -equivariant map  $f : \tilde{S} \rightarrow \mathbb{Y}_n$  is equivalent to a section of this latter bundle, and hence produces exactly a non-degenerate  $\mathbb{C}$ -bilinear form  $H$  on  $(E_\rho, D)$ .

Given such a bilinear form  $H$ , we can uniquely decompose  $D$  as

$$D = \nabla^H + \Psi^H, \quad (48)$$

where  $\nabla^H$  is a connection on  $E_\rho$  satisfying  $\nabla^H H = 0$  and  $\Psi^H \in \Omega^1(\mathrm{End}_0(E_\rho))$ , is  $H$ -self-adjoint, namely, for all sections  $s_1, s_2$ ,  $H(\Psi^H s_1, s_2) = H(s_1, \Psi^H s_2)$  as forms. Note that, using the Maurer-Cartan form,  $\Psi^H$  identifies with the derivative  $df$  of the equivariant map  $f$  engendered by  $H$ , and the action of  $\nabla^H$  on endomorphisms of  $E_\rho$  identifies with that of the Levi-Civita connection on the pullback bundle of the tangent space of  $\mathbb{Y}_n$  (the computation is identical to that of [Li19, Sections 2.23 and 2.24] and also those found in [Gui18, Section 7]).

Next, given complex structures  $(c_1, \overline{c_2})$ , we decompose the expression (48) according to  $\Omega^1 = \Omega_{c_1}^{1,0} \oplus \Omega_{c_2}^{0,1}$ : we split  $\nabla^H$  as

$$\nabla^H = \partial_1^H + \overline{\partial}_2^H,$$

and  $\Psi^H$  as

$$\Psi^H = \Psi_1^H + \overline{\Psi}_2^H.$$

By Proposition 7.2, the harmonic map equation (45) translates to two equations,

$$\overline{\partial}_2^H \Psi_1^H = 0 \quad \text{and} \quad \partial_1^H \overline{\Psi}_2^H = 0, \quad (49)$$

and the flatness of  $D$  becomes the expression

$$F(\partial_1^H + \overline{\partial}_2^H) + [\Psi_1^H, \overline{\Psi}_2^H] = 0. \quad (50)$$

It's important to underline that  $\overline{\partial}_2^H$  is not a del-bar operator on  $\Omega^1(S)$  (unless  $c_1 = c_2$ ), since it does not satisfy the required Leibniz rule. Indeed, for  $s$  a section  $E$  and  $u$  a function,

$$\overline{\partial}_2^H(us) = (\overline{\partial}_{c_2} u) \otimes s + u \overline{\partial}_2^H s.$$

and  $\overline{\partial}_{c_2} u$ , defined as above, does not locally return  $\partial_{\overline{z}} u d\overline{z}$ . This all leads to the definitions below

**Definition 7.5.** Let  $E$  be a complex vector bundle over  $S$  and let  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$ .

A  $(c_1, \overline{c_2})$ -**del operator**  $\partial_1$  on  $E$  is an operator  $\partial_1 : \Omega^0(E) \rightarrow \Omega^1(E)$ , such that, for all  $s$  in  $\Omega^0(E)$  and  $u$  a function,

$$\partial_1(us) = (\partial_{c_1} u) \otimes s + u \partial_1 s. \quad (51)$$

Similarly, a  $(c_1, \overline{c_2})$ -**del-bar operator**  $\overline{\partial}_2$  on  $E$  is an operator  $\overline{\partial}_2 : \Omega^0(E) \rightarrow \Omega^1(E)$  such that, for all  $s$  a section and  $u$  a function,

$$\overline{\partial}_2(us) = (\overline{\partial}_{c_2} u) \otimes s + u \overline{\partial}_2 s. \quad (52)$$

Observe that  $\nabla = \partial_1 + \bar{\partial}_2$  is a connection on  $E$ .  $\partial_1$  and  $\bar{\partial}_2$  extend naturally to operators  $\Omega^1(E) \rightarrow \Omega^2(E)$ , which we still denote by  $\partial_1$  and  $\bar{\partial}_2$  respectively, by

$$\begin{aligned}\partial_1(\omega \otimes s) &= \partial_{c_1}\omega \otimes s - \omega \wedge \partial_1 s, \\ \bar{\partial}_2(\omega \otimes s) &= \bar{\partial}_{c_2}\omega \otimes s - \omega \wedge \bar{\partial}_2 s,\end{aligned}$$

which satisfy the analog of (51) and (52) and which are such that  $\partial_1 \circ \partial_1 = 0$  and  $\bar{\partial}_2 \circ \bar{\partial}_2 = 0$ . Moreover, the curvature form of  $\partial_1 + \bar{\partial}_2$  is precisely given by

$$F(\partial_1 + \bar{\partial}_2) = (\partial_1 + \bar{\partial}_2) \circ (\partial_1 + \bar{\partial}_2) = \partial_1 \circ \bar{\partial}_2 + \bar{\partial}_2 \circ \partial_1 .$$

**Definition 7.6.** A **bi-Higgs bundle**  $(E, \bar{\partial}_1, \partial_2, \Psi_1, \bar{\Psi}_2)$  over  $(S, c_1, \bar{c}_2)$  is a complex vector bundle  $E$  over  $S$  equipped with a  $(c_1, \bar{c}_2)$ -del operator  $\partial_1$  and a  $(c_1, \bar{c}_2)$ -del bar operator  $\bar{\partial}_2$ , together with sections  $\Psi_1$  of  $\Omega_{c_1}^{1,0}(\text{End}(E))$  and  $\bar{\Psi}_2$  of  $\Omega_{c_2}^{0,1}(\text{End}(E))$  called Higgs fields that satisfy

$$\bar{\partial}_2 \Psi_1 = 0, \quad \partial_1 \bar{\Psi}_2 = 0,$$

and

$$F(\partial_1 + \bar{\partial}_2) + [\Psi_1, \bar{\Psi}_2] = 0.$$

Now, given  $(c_1, \bar{c}_2)$ , we write  $\mathcal{K}_{c_1}$  and  $\overline{\mathcal{K}_{c_2}}$  for the canonical bundles of  $c_1$  and  $\bar{c}_2$  respectively. We denote by  $\mathcal{K}_{c_1}^i$  the  $i^{\text{th}}$  tensor power of  $\mathcal{K}_{c_1}$ , and likewise for  $\overline{\mathcal{K}_{c_2}}$ . We define the **bi-Hitchin base**

$$bH(c_1, \bar{c}_2) = \bigoplus_{i=2}^n H^0(S, \mathcal{K}_{c_1}^i) \oplus \bigoplus_{i=2}^n H^0(\bar{S}, \overline{\mathcal{K}_{c_2}}^i)$$

and the **bi-Hitchin map** from the set of bi-Higgs bundles to the bi-Hitchin base to be the map

$$(E, \bar{\partial}_1, \partial_2, \Psi_1, \bar{\Psi}_2) \mapsto \left( \left( \frac{3}{\sqrt{2}} \text{tr}(\Psi_1^2), \dots, \frac{3}{\sqrt{2}} \text{tr}(\Psi_1^n) \right), \left( \frac{3}{\sqrt{2}} \text{tr}(\bar{\Psi}_2^2), \dots, \frac{3}{\sqrt{2}} \text{tr}(\bar{\Psi}_2^n) \right) \right).$$

The normalization is unimportant and we pick this particular one to state our theorem more cleanly. By the formulas (46), the differentials are indeed holomorphic (we omit the computation, since we won't use the result). We comment that when  $c_1 = c_2$ , and  $H$  is the complexification of a Riemannian metric,  $\overline{\text{tr}(\Psi_1^i)} = \text{tr}(\bar{\Psi}_2^i)$  for all  $i$ , so the image of the bi-Hitchin map lies in the "diagonal." As well, depending on the normalization of the Killing form, the Hopf differentials of  $f$ ,  $q_1$  and  $\bar{q}_2$ , are scalar multiples of  $\text{tr}(\Psi_1^2)$  and  $\text{tr}(\bar{\Psi}_2^2)$  respectively. So a harmonic map to  $\mathbb{Y}_n$  is weakly conformal if and only if the  $H^0(S, \mathcal{K}_{c_1}^2)$  and  $H^0(\bar{S}, \overline{\mathcal{K}_{c_2}}^2)$  components of the bi-Hitchin map vanish. Specializing to the case  $n = 3$ , we deduce:

**Proposition 7.7.** *An equivariant weakly conformal harmonic map from a closed surface  $f : (\tilde{S}, c_1, \bar{c}_2) \rightarrow (\mathbb{Y}_3, \langle \cdot, \cdot \rangle)$  determines a point in  $\mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$  in the fiber over  $(c_1, \bar{c}_2)$ .*

There's a striking peculiarity in the construction above: unlike the Hitchin equation, the expression (50) is not an equation for a harmonic metric. That is, once one prescribes all of the holomorphic data, there is nothing to solve for. As a result, it's not clear how to take a point in the bi-Hitchin base and produce a bi-Higgs bundle with that same data (and then there's the further question of whether such a bi-Higgs bundle arises from a harmonic map). In our situation, the Gauss equation that we've already derived, together with Hitchin's seminal construction from [Hit92], suggest an ansatz.

**7.4. Bi-Hitchin bi-Higgs bundles.** Here we prove the first statement of Theorem D by constructing a bi-Higgs bundle arising from a conformal harmonic map to  $\mathbb{Y}_3$  that is equivalent to the information of a positive hyperbolic complex affine sphere. More precisely, let  $(c_1, \overline{c_2}) \in \mathcal{C}(S) \times \mathcal{C}(\overline{S})$  with induced Bers metric  $h = \mathbf{h}_{(c_1, \overline{c_2})}$  and  $Q_1$  and  $\overline{Q_2}$  cubic differentials on  $(S, c_1)$  and  $(\overline{S}, \overline{c_2})$  respectively. We show in this subsection that if one can solve the Gauss equation

$$\Delta_h u = e^{2u} - \frac{1}{4}h(Q_1, \overline{Q_2})e^{-4u} - 1, \quad (53)$$

then one can build a bi-Higgs bundle that arises from a conformal harmonic map, and then in the next subsection we'll show that this bi-Higgs bundle can be used to produce the complex affine sphere associated with (53).

Let  $E$  be the complex vector bundle  $E = \mathcal{K}_{c_1} \oplus \mathcal{O} \oplus \mathcal{K}_{c_1}^{-1}$ . As in Section 7.2, we split  $d$  as  $d = \partial_{c_1} + \overline{\partial}_{c_2}$ . The operator  $\overline{\partial}_{c_2}$  determines a  $(c_1, \overline{c_2})$ -del-bar operator  $\overline{\partial}_2$  on  $E$  uniquely characterized by the Leibniz rule (52) and by having  $(dz, 0, 0)$ ,  $(0, 1, 0)$ , and  $(0, 0, dz^{-1})$  in its kernel. By analogy with the ordinary Higgs bundles theory, we take

$$\Psi_1 = \begin{pmatrix} 0 & 0 & \frac{1}{\sqrt{2}}Q_1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

To cook up the rest of a bi-Higgs bundle, note that if we were in the case  $c_1 = c_2$  and  $Q_1 = Q_2$ , then taking the adjoints of  $\overline{\partial}_2$  and  $\Psi_1$  with respect to a Hermitian metric on  $E$  would return for us the data of a  $(c_1, \overline{c_2})$ -del operator and an  $\text{End}(E)$ -valued form. Recalling that a Hermitian metric on a bundle  $E_0$  can be seen as an isomorphism from  $E_0^* \rightarrow \overline{E_0}^*$ , our idea below is to build an isomorphism from  $E \rightarrow \overline{\mathcal{K}_{c_2}^{-1}} \oplus \mathcal{O} \oplus \overline{\mathcal{K}_{c_2}}$ , and then to mimic the adjoint construction in order to take objects on the latter bundle and put them on  $E$ .

On the bundle  $\overline{\mathcal{K}_{c_2}^{-1}} \oplus \mathcal{O} \oplus \overline{\mathcal{K}_{c_2}}$ , we have a  $(c_1, \overline{c_2})$ -del operator  $\hat{\partial}_1$ , characterized by the rule (51) and sending  $(d\overline{w}^{-1}, 0, 0)$ ,  $(0, 1, 0)$ , and  $(0, 0, d\overline{w})$  to 0. We set  $U_0$  to be a non-vanishing section of  $\mathcal{K}_{c_1} \otimes \overline{\mathcal{K}_{c_2}}$  locally of the form  $U_0 = \lambda dz \otimes d\overline{w}$ , where  $\lambda$  is the density of the Bers metric  $h = \mathbf{h}_{(c_1, \overline{c_2})}$ . We then consider a function  $u : S \rightarrow \mathbb{C}^*$  and set  $U = \frac{1}{2}e^{2u}U_0$ . We define an isomorphism

$$I_u : \mathcal{K}_{c_1} \oplus \mathcal{O} \oplus \mathcal{K}_{c_1}^{-1} \rightarrow \overline{\mathcal{K}_{c_2}^{-1}} \oplus \mathcal{O} \oplus \overline{\mathcal{K}_{c_2}}, \quad I_u = \begin{pmatrix} U^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & U \end{pmatrix}.$$

Now, we define our  $(c_1, \overline{c_2})$ -del operator on  $E$  by

$$\partial_1^u = I_u^{-1} \circ \hat{\partial}_1 \circ I_u,$$

and our second Higgs field by

$$\overline{\Psi}_2^u = I_u^{-1} \circ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{2}}\overline{Q_2} & 0 & 0 \end{pmatrix} \circ I_u = \begin{pmatrix} 0 & U & 0 \\ 0 & 0 & U \\ \frac{1}{\sqrt{2}}U^{-2}\overline{Q_2} & 0 & 0 \end{pmatrix}.$$

Above, the first matrix-valued form is valued in  $\overline{\mathcal{K}_{c_2}^{-1}} \oplus \mathcal{O} \oplus \overline{\mathcal{K}_{c_2}}$ .

**Proposition 7.8.** *The data  $(E, \partial_1^u, \overline{\partial}_2, \Psi_1, \overline{\Psi}_2^u)$  defines a bi-Higgs bundle.*

In order to prove this, we will need the following lemma.

**Lemma 7.9.**  $\Delta_h \lambda = 2$

*Proof.* A simple way to prove this statement is to remark that it holds for Riemannian metrics and then to use a holomorphic continuation argument. Explicitly, we can argue as follows. For the local complex metric  $dz \cdot d\bar{w}$ , one has that  $\nabla \partial_{\bar{z}} = d \log(\partial_{\bar{z}} \bar{w}) \otimes \partial_{\bar{z}}$ , as one can see in the local frame  $\{\partial_z, \partial_{\bar{z}}\}$ . Using this, one gets straightforwardly that the curvature of  $dz \cdot d\bar{w}$  is zero from  $R(\partial_z, \partial_{\bar{z}}) \partial_{\bar{z}} = 0$ . The statement then follows by applying Equation (4) to the metric  $\lambda dz d\bar{w}$ , which has curvature -1.  $\square$

*Proof of Proposition 7.8.* It follows from holomorphicity of  $Q_1$  and  $\overline{Q_2}$ , that  $\overline{\partial_2} \Psi_1 = 0$  and  $\partial_1^u \overline{\Psi_2^u} = 0$ . Indeed, for  $\Psi_1$ , holomorphicity just follows from the fact that  $\overline{\partial_{c_2}}$  is zero on  $c_1$ -holomorphic functions and  $\overline{\partial_2} dz = 0$ . For  $\overline{\Psi_2^u}$ ,

$$\partial_1^u \overline{\Psi_2^u} = (I_u^{-1} \circ \hat{\partial}_1 \circ I_u) \circ (I_u^{-1} \circ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{2}} \overline{Q_2} & 0 & 0 \end{pmatrix} \circ I_u) = I_u^{-1} \circ \hat{\partial}_1 \left( \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{2}} \overline{Q_2} & 0 & 0 \end{pmatrix} \right) \circ I_u = 0,$$

where the middle term vanishes from reasoning similar to above.

We now compute the curvature equation (50) in a local basis  $(dz, 0, 0), (0, 1, 0), (0, 0, \frac{1}{dz})$ . Recall that

$$F(\partial_1^u + \overline{\partial_2}) = (\partial_1^u + \overline{\partial_2}) \circ (\partial_1^u + \overline{\partial_2}) = \partial_1^u \circ \overline{\partial_2} + \overline{\partial_2} \circ \partial_1^u$$

hence, since  $\overline{\partial_2}$  on this local basis, we are left to compute  $\overline{\partial_2} \circ I_u^{-1} \circ \hat{\partial}_1 \circ I_u$  with respect to it. Recalling Remark 5.3 and Lemma 7.9, we get

$$\begin{aligned} (\overline{\partial_2} \circ I_u^{-1} \circ \hat{\partial}_1 \circ I_u)(dz) &= \overline{\partial_2} \left( \lambda e^{2u} \frac{\partial_w(\frac{1}{\lambda e^{2u}})}{\partial_{\bar{z}} \bar{w}} dz \otimes dz \right) \\ &= \frac{\lambda}{4} \Delta_h(\log(\lambda) + 2u)(dz \wedge d\bar{w}) \otimes dz \\ &= \left( \frac{1}{2} \Delta_g u + \frac{1}{2} \right) (dz \wedge d\bar{w}) \otimes dz. \end{aligned}$$

The analogous computation for  $\frac{1}{dz}$  shows that in the basis  $(dz, 0, 0), (0, 1, 0), (0, 0, \frac{1}{dz})$  the curvature form is

$$F(\partial_1^u + \overline{\partial_2}) = \begin{pmatrix} \frac{1}{2} \Delta_h u + \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} \Delta_h u - \frac{1}{2} \end{pmatrix} \lambda dz \wedge d\bar{w}.$$

As well, writing locally  $Q_1 = \varphi_1 dz^2$  and  $\overline{Q_2} = \overline{\varphi_2} d\bar{w}^2$ , we see directly,

$$[\Psi_1, \overline{\Psi_2^u}] = \begin{pmatrix} -\frac{1}{2} \lambda e^{2u} + \frac{\varphi_1 \overline{\varphi_2}}{\lambda^2} e^{-4u} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} \lambda e^{2u} - \frac{\varphi_1 \overline{\varphi_2}}{\lambda^2} e^{-4u} \end{pmatrix} dz \wedge d\bar{w}.$$

Dividing by the 2-form  $\lambda dz \wedge d\bar{w}$ , we get that  $F(\partial_1^u + \overline{\partial_2}) + [\Psi_1, \overline{\Psi_2^u}] = 0$  if and only if  $u$  solves Equation (53). Putting everything together, we've established the result.  $\square$

**Definition 7.10.** When Equation (53) has a solution  $u$ , we call  $(E, \partial_1^u, \overline{\partial_2}, \Psi_1, \overline{\Psi_2^u})$  a **bi-Hitchin bi-Higgs bundle**.

**Remark 7.11.** Since Equation (53) can have multiple solutions, this construction gives multiple bi-Higgs bundles lying over the same point in the bi-Hitchin base.

We now show that such bi-Hitchin bi-Higgs bundles arise from conformal harmonic maps in  $\mathbb{Y}_n$ . Set  $\hat{D} = \partial_1^u + \overline{\partial}_2 + \Psi_1 + \overline{\Psi}_2^u$ . The equations characterizing a bi-Higgs bundle imply that  $\hat{D}$  is flat.

**Proposition 7.12.** *Keeping in the setting above, if  $u$  solves (53), the  $\mathbb{C}$ -bilinear form on  $E$  given by*

$$H = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

*gives rise to a conformal harmonic map  $f: \tilde{S} \rightarrow \mathbb{Y}_n$  whose bi-Higgs bundle is  $(E, \partial_1^u, \overline{\partial}_2, \Psi_1, \overline{\Psi}_2^u)$ .*

*Proof.* Setting  $\nabla^H = \partial_1^u + \overline{\partial}_2$ , it is easily computed working in a frame that  $\nabla^H H = 0$ . One can check by hand that  $\Psi^H$  is  $H$ -self adjoint.

Picking a point on  $S$ ,  $H$  determines a map  $f$  to the symmetric space  $\mathbb{Y}_n$ , equivariant with respect to the holonomy representation of  $\hat{D}$  at that point. By uniqueness of the decomposition (48), and our comments near the top of the subsection 7.3,  $\Psi^H$  has an identification with the derivative of  $f$ . Using Proposition 7.2,  $f$  is harmonic if and only if  $\overline{\partial}_2 \Psi_1 = 0$ , which holds by construction, together with the flatness condition. Reversing the construction of the previous subsection, we see the bi-Higgs bundle is  $(E, \partial_1^u, \overline{\partial}_2, \Psi_1, \overline{\Psi}_2^u)$ . Since  $\Psi_1$  never vanishes,  $df$  never vanishes, and hence  $f$  is genuinely conformal.  $\square$

We refer to  $f$  as the **bi-Hitchin conformal harmonic map**. Since positive hyperbolic complex affine spheres and bi-Hitchin conformal harmonic maps are both determined by input data  $(c_1, \overline{c}_2, Q_1, Q_2)$  and a solution to Equation (53), we have proved that they are in one-to-one correspondence. To complete the proof of Theorem D, we show in the next subsection that the holonomies agree on the special locus of Theorem D.

**7.5. From bi-Higgs bundles to complex affine spheres.** There is a known correspondence between equivariant real affine spheres in  $\mathbb{R}^3$  and conformal harmonic maps into the Riemannian symmetric space  $Y_3$  that are equivariant for Hitchin representations. Since the holonomies of infinitesimally rigid complex affine spheres vary holomorphically, and the same should hold for infinitesimally rigid conformal harmonic maps to  $\mathbb{Y}_3$ , the result should be true. Using this reasoning, we prove the second statement of Theorem D.

We follow Section 3.4 from [Bar10]; the constructions go through with very minor change. Let  $\sigma$  be a complex affine sphere corresponding to a point  $(c_1, \overline{c}_2, Q_1, Q_2, u) \in \text{CAS}(S)$ , with Blaschke connection  $\nabla$ , Blaschke metric  $g$ , and associated flat connection  $D$ . Using  $\xi$  for the transverse normal, we split  $\sigma^* T \mathbb{C}^3 = \mathbb{C} T S \oplus \text{Span}_{\mathbb{C}}(\xi)$ . Recall from Section 3.4 that  $\sigma$  is equivalent to the section  $\xi$  of  $\sigma^* T \mathbb{C}^3$ ,  $z \mapsto \xi(z)$ .

As well, let  $(E, \partial_1^u, \overline{\partial}_2, \Psi_1, \overline{\Psi}_2^u)$  be the bi-Hitchin bi-Higgs bundle associated with  $\sigma$ , with flat connection  $\hat{D}$ . Let  $s$  be the tautological section of  $\mathcal{O}$ , sitting inside  $E$ ,  $s = [0, 1, 0]^T$ . Evaluating  $\hat{D}$  on  $s$ , we get  $\hat{D}s = [U, 0, 1]^T$ . Since  $U$  is non-vanishing,  $s$  thus determines an isomorphism of complex vector bundles

$$B_0 : \mathbb{C} T S \rightarrow E/\mathcal{O}, \quad Z \mapsto [\hat{D}_Z s]_{\mathcal{O}},$$

where the notation  $[\cdot]_{\mathcal{O}}$  means the equivalence class mod  $\mathcal{O}$ . The splitting  $E = (\mathcal{K}_{c_1} \oplus \mathcal{K}_{c_1}^{-1}) \oplus \mathcal{O}$  induces an isomorphism from  $E/\mathcal{O}$  to  $(\mathcal{K}_{c_1} \oplus \mathcal{K}_{c_1}^{-1})$ , and hence  $B_0$  determines an isomorphism from  $\mathbb{C} T S$  to  $(\mathcal{K}_{c_1} \oplus \mathcal{K}_{c_1}^{-1})$ . We further extend to an isomorphism

$$B : \sigma^* T \mathbb{C}^3 = \mathbb{C} T S \oplus \text{Span}_{\mathbb{C}}(\xi) \rightarrow E = (\mathcal{K}_{c_1} \oplus \mathcal{K}_{c_1}^{-1}) \oplus \mathcal{O}$$



by, in each fiber, sending  $\xi$  to  $s$  and extending linearly. The last part of Theorem D is immediate from the following.

**Proposition 7.13.** *For  $\sigma$  deformable to a real affine sphere through the infinitesimally rigid locus, the pullback connection  $B^*\hat{D}$  is equal to  $D$ , up to an  $\mathrm{SL}(3, \mathbb{C})$ -automorphism.*

*Proof.* Firstly, we point out that the space of  $C^\infty$  connections on a rank  $n$  complex vector bundle over a surface is modelled on a complex Fréchet manifold consisting of  $C^\infty$  sections of the rank  $n$  complex affine bundle of smooth endomorphism-valued 1-forms. We pull back all bundles to the universal cover  $\tilde{S}$ .

Recall from the proof of Theorem 3.20 that we constructed  $\sigma$  by taking the trivial bundle  $N : \mathbb{C} \times \tilde{S} \rightarrow \tilde{S}$ , defining structural data on  $\mathbb{C}T\tilde{S} \oplus N$ , and then choosing an isomorphism to the trivial  $\mathbb{C}^3$ -bundle over  $S$ . Undoing this isomorphism, we view every  $D$  as a connection on the bundle  $V = \mathbb{C}T\tilde{S} \oplus N$ , and  $B$  as a map from  $V$  to  $E$  that preserves the relevant splittings.  $B^*\hat{D}$  is now a connection on  $V$ .

Note that, by the results of Section 6, specifically the argument of Proposition 6.3, solutions to Equation (53) for infinitesimally rigid affine spheres depend holomorphically in  $C^\infty(S, \mathbb{C})$  on the input data (namely points  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2)$  in  $\mathcal{CD}(S) \times \mathcal{CD}(\bar{S})$ ). As explained in the proof of Proposition 6.5, it follows that the structural data on  $V$  from the proof of Theorem 3.20, and hence the flat connections, vary holomorphically.

We show that also the  $B^*\hat{D}$ 's vary holomorphically. Toward this, first recall our basepoint complex structure  $c_0$  from Section 2, and set  $E_0 := \mathcal{K}_{c_0} \oplus \mathcal{O} \oplus \mathcal{K}_{c_0}^{-1}$ . Given any other complex structure  $c_1$  coming from a Beltrami form  $\mu$ , the diffeomorphism  $f_\mu : (S, c_0) \rightarrow (S, c_1)$  (recall the notation from Section 2.1) induces an isomorphism of complex vector bundles from  $E_0$  to  $\mathcal{K}_{c_1} \oplus \mathcal{O} \oplus \mathcal{K}_{c_1}^{-1}$ . Moreover, it follows from Theorem A.5 in the appendix that, working in a local frame for  $E_0$ , as we vary the input data, the components of the  $(f_\mu)_*\hat{D}$ 's vary holomorphically. Now, composing the isomorphisms from  $V \rightarrow E$  with our identifications from  $E \rightarrow E_0$ , the composite isomorphisms depend only on  $(f_\mu)_*\hat{D}$ , and hence these isomorphisms vary holomorphically as  $C^\infty$  sections of  $\mathrm{Hom}(V, E_0)$ . We conclude that once the  $B^*\hat{D}$ 's are pulled back to  $V$  they form a holomorphically varying family of flat connections.

For real data  $(c, \bar{c}, Q, \bar{Q}, u) \in \mathrm{RAS}(S) \subset \mathrm{CAS}(S)$ ,  $D$  is written explicitly in [DW15] among other sources,  $B^*\hat{D}$  is computed in [Bar10, Section 3.4], and the isomorphisms from  $V \rightarrow \sigma^*T\mathbb{C}^3$  can be chosen so that the two connections agree (note that slightly different normalizations are chosen, but if we choose the same ones, then things match up). That is, the two families of affine connections on  $V$  agree on  $\mathrm{RAS}(S)$ . We deduce by holomorphicity that the two connections agree when they come from input data in the infinitesimally rigid locus.  $\square$

As one can see, we only used the extra hypothesis on  $\sigma$  to circumvent computing the connections explicitly. The connection should always agree, but we prefer to carry out this computation in a more general setting related to bi-Higgs bundles in a future work.

Note that if we pull back  $E$  to  $\tilde{S} \rightarrow S$  and then trivialize this pullback bundle, upon picking a point  $p \in S$ ,  $s$  determines an equivariant map  $\sigma_s : \tilde{S} \rightarrow \mathbb{C}^3$ . Going back into the proof of Theorem 3.13, Theorem 3.20 shows that, because Equation (53) is satisfied by our input data, the equivariant map  $\sigma_s$  determined by  $s$  is a positive hyperbolic complex affine sphere. Moreover, the structural data of  $\sigma_s$  agrees with that of  $\sigma$ , and hence the uniqueness of Theorem 3.13 shows that  $\sigma_s$  is identified with  $\sigma$ .

**7.6. Opers.** As previously discussed, when given the data  $(c_1, \bar{c}_2, Q_1, 0)$ , we can always solve Equation (53) with the function  $u = 0$ . Here we'll show using bi-Higgs bundles that positive hyperbolic complex affine spheres of the form  $(c_1, \bar{c}_2, Q_1, \bar{Q}_2, u) = (c, \bar{c}, Q, 0, 0)$ , are equivalent to opers. As above, let  $(S, c)$  be a Riemann surface with canonical bundle  $\mathcal{K}$ .

**Definition 7.14.** An  $\mathrm{SL}(n, \mathbb{C})$ -**oper** on  $(S, c)$  is the data of a holomorphic vector bundle  $V \rightarrow S$  carrying a holomorphic connection  $\nabla$  inducing the trivial connection on  $\det(V)$ , together with a holomorphic full filtration

$$0 = V_0 \subset V_1 \subset \cdots \subset V_n = V$$

such that

- (1)  $\nabla V_i \subset V_{i+1} \otimes \mathcal{K}$ , and
- (2) for  $i = 1, \dots, n-1$ , the induced  $\mathcal{O}$ -linear map  $V_i/V_{i-1} \rightarrow V_{i+1}/V_i \otimes \mathcal{K}$  is an isomorphism.

The holonomy of the oper is the holonomy of the holomorphic connection  $\nabla$ .

For a general introduction to opers, from a differential geometric standpoint, see [Wen15]. Opers generalize  $n^{\mathrm{th}}$  order linear differential operators: if one tries to globalize an  $n^{\mathrm{th}}$  order operator on  $S$  in a way that respects the coordinate changes, one ends up with an oper. See [Dum+21b, Section 2.11] for details and explanation. As well, as mentioned in the introduction, for  $n = 2$  opers are equivalent to complex projective structures.

Beilinson and Drinfeld defined an explicit family of opers, parametrized by the Hitchin base  $\oplus_{i=2}^n H^0(S, \mathcal{K}^i)$ . Let  $E$  be the holomorphic vector bundle

$$\mathcal{K}^{-\frac{(n-1)}{2}} \oplus \mathcal{K}^{-\frac{(n-3)}{2}} \oplus \cdots \oplus \mathcal{K}^{\frac{n-3}{2}} \oplus \mathcal{K}^{\frac{n-1}{2}},$$

with standard holomorphic structure  $\bar{\partial}_E$  induced by  $c$  and let  $\Psi_0$  be the principal nilpotent Higgs field

$$\Psi_0 = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{pmatrix}.$$

Let  $H_0$  be the Hermitian metric that solves Hitchin's equations for the stable Higgs bundle  $(E, \bar{\partial}_E, \Psi_0)$ . That is,  $H_0$  is the unique Hermitian metric such that

$$\bar{\partial}_E + \bar{\partial}_E^{*H_0} + \Psi_0 + \Psi_0^{*H_0}$$

is a flat connection, where  $\bar{\partial}_E^{*H_0}$  and  $\Psi_0^{*H_0}$  are the Hermitian adjoints. Writing

$$\mathrm{End}_j(E) = \oplus_{|i-k|=j} \mathrm{Hom}(\mathcal{K}^{\frac{n+1}{2}-i}, \mathcal{K}^{\frac{n+1}{2}-k}),$$

the  $\mathrm{End}_j(E)$ 's yield a  $\mathbb{Z}$ -grading of  $\mathrm{End}(E)$ . We interpret sections of  $H^0(S, \mathcal{K}^i)$  as sections of  $\mathrm{End}_j(E) \otimes \mathcal{K}$ .

**Definition 7.15. Beilinson-Drinfeld's oper** associated with  $q = (q_2, \dots, q_n) \in \oplus_{i=2}^n H^0(S, \mathcal{K}^i)$  is the holomorphic vector bundle  $(E, \bar{\partial}_E)$  equipped with the holomorphic connection

$$\bar{\partial}_E + \bar{\partial}_E^{*H_0} + \Psi_0 + \frac{1}{\sqrt{2}}q_2 + \frac{1}{\sqrt{2}}q_3 + \cdots + \frac{1}{\sqrt{2}}q_n + \Psi_0^{*H_0} \quad (54)$$

and filtration

$$0 \subset \mathcal{K}^{-\frac{(n-1)}{2}} \subset \mathcal{K}^{-\frac{(n-1)}{2}} \oplus \mathcal{K}^{-\frac{(n-3)}{2}} \subset \dots \subset \mathcal{K}^{-\frac{(n-1)}{2}} \oplus \dots \oplus \mathcal{K}^{\frac{n-1}{2}} = E.$$

Note that there are different ways of defining Beilinson-Drinfeld's oper in the literature, but they are all gauge equivalent (see [Dum+21a, Theorem 3.2]). We choose the  $2^{1/2}$  normalization so that it matches up our construction. When  $n = 2$ , the  $q_2$  is a multiple of the Schwarzian of the associated complex projective structure. Observe that the holomorphic connection (54) is exactly the flat connection associated with the bi-Higgs bundle

$$(E, \bar{\partial}_E, \bar{\partial}_E^{*H_0}, \Psi_0 + q_2 + q_3 + \dots + q_n, \Psi_0^{*H_0}).$$

With the definitions given, Theorem E is immediate from Theorem G.

*Proof of Theorem E.* Fix the data  $(E, \bar{\partial}_E, \Psi_0, H_0)$  as above for  $n = 3$ . Let  $\sigma$  be the positive hyperbolic complex affine sphere with immersion data  $(c, \bar{c}, Q, 0, 0)$ . If we apply our construction from Section 7.4 to  $\sigma$ , we get the bi-Hitchin bi-Higgs bundle

$$(E, \bar{\partial}_E, \bar{\partial}_E^{*H_0}, \Psi_0 + \frac{1}{\sqrt{2}} Q, \Psi_0^{*H_0}),$$

and that  $H_0$  corresponds to a conformal harmonic map to the symmetric space  $\mathbb{Y}_n$ . From Theorem D we conclude that the holonomy of the complex affine sphere  $\sigma$  agrees with that of the flat connection

$$\bar{\partial}_E + \bar{\partial}_E^{*H_0} + \Psi_0 + \frac{1}{\sqrt{2}} Q + \Psi_0^{*H_0},$$

which is exactly the connection (54). This establishes the  $(c, \bar{c}, Q, 0, 0)$  case. Reversing  $c$  with  $\bar{c}$  we get the analogous result, and moreover we obtain the statement of Theorem E.  $\square$

Finally, in Section 1.3 we promised a discussion related to the  $\mathbb{C}^*$  twistor action. We assume some background on Higgs bundles here. Let  $(E, \bar{\partial}_E, \Psi)$  be a stable Higgs bundle over a Riemann surface  $(S, c)$  with Hermitian metric  $H$  solving Hitchin's equations, i.e., which makes the connection

$$D = \nabla^H + \Psi + \Psi^{*H}$$

flat. From this data, we can build a  $\mathbb{C}^*$ -family of flat connections via

$$D_\zeta = \nabla^H + \zeta \Psi + \zeta^{-1} \Psi^{*H}, \quad \zeta \in \mathbb{C}^*. \quad (55)$$

The family of flat connections (55) is called a real twistor line. For every  $R \in \mathbb{C}^*$ , we can find a unique Hermitian metric  $H_R$  so that  $D_R = \nabla_{H_R} + R\Psi + R\Psi^{*H_R}$  is flat, and then we can further apply the  $\mathbb{C}^*$ -action. Gaiotto proposed to leave the ratio  $\hbar = \zeta R^{-1}$  constant, and then to consider the family of flat connections

$$D_{\hbar, R} = \nabla_{H_R} + \hbar^{-1} \Psi + \hbar R^2 \Psi^{*H_R}.$$

The gauge equivalence class of the limit as  $R \rightarrow 0$ , if it exists, is called the  $\hbar$ -conformal limit of  $(E, \bar{\partial}_E, \Psi)$ . Gaiotto conjectured that for a Higgs bundle in the Hitchin component (on a punctured surface), the  $\hbar$ -conformal limit always exists and the limit is an oper. This conjecture was proved on closed surfaces in [Dum+21a] (see also [CW19]).

Now, by Theorem E, after doing a standard gauge transformation (as found in [Dum+21a, Section 3]), one sees that the  $\mathbb{C}^*$ -family described by (55) corresponds exactly to the family of complex affine spheres with immersion data given by

$$(c, \bar{c}, \zeta^3 Q, \zeta^{-3} \bar{Q}, u), \quad \zeta \in \mathbb{C}^*.$$

Since  $(c, \bar{c}, Q, 0, 0)$  and  $(c, \bar{c}, 0, \bar{Q}, 0)$  correspond toopers, paths in  $\Omega \subset \mathcal{M}_3(S) \times \mathcal{M}_3(\bar{S})$  from the real locus to (the classes of) these points should be related to conformal limits.

The  $\mathbb{C}^*$ -action on  $\text{CAS}^*(S)$ ,

$$\zeta \cdot (c_1, \overline{c_2}, Q_1, \overline{Q_2}, u) \mapsto (c_1, \overline{c_2}, \zeta Q_1, \zeta^{-1} \overline{Q_2}, u),$$

seems to give rise to a new type of  $\mathbb{C}^*$  family in the space of flat connections: if  $(c_1, \overline{c_2}, Q_1, \overline{Q_2}, u)$  corresponds to a bi-Higgs bundle  $(E, \partial_1^u, \overline{\partial_2}, \Psi_1, \overline{\Psi_2^u})$ , then we have

$$D_\zeta = (\partial_1^u + \overline{\partial_2}) + \zeta^3 \Psi_1 + \zeta^{-3} \overline{\Psi_2^u}, \quad \zeta \in \mathbb{C}^* .$$

It is then natural to expect that by taking a conformal limit, analogous to Gaiotto's, one can arrive at the holonomy of a point in  $\text{CAS}^*(S)$  that projects to the loci  $\mathcal{M}_3(S) \times \mathcal{T}(\overline{S})$  or  $\mathcal{T}(S) \times \mathcal{M}_3(\overline{S})$ , and that this holonomy should be some complex analog of an oper. We hope to develop this picture in a future work.

**Remark 7.16.** Relating to Remark 6.4, a proof that opers have irreducible holonomy can be found in [Wen15, Proposition 4.8]. For a point  $(c_1, \overline{c_2}, Q_1, 0)$ , we have written out the associated flat connection, and the proof from [Wen15, Proposition 4.8] appears to be replicable for our connections. We prefer to work out such results in a more general context in future work.

#### APPENDIX A. ON THE BELTRAMI OPERATOR

Here we prove Propositions 5.7 and 2.9. The main input toward Proposition 2.9 is the holomorphic dependence of solutions of the Beltrami equation, Theorem A.5, which we prove in Section A.2. Throughout, we set  $L_1^\infty(\mathbb{C})$  to be the open unit ball of  $L^\infty(\mathbb{C})$ , which we think of as the space of Beltrami forms on  $\mathbb{C}$ . Recall from Section 5 that if we have a constant  $C$  depending on quantities  $a_1, \dots, a_n$ , we write  $C(a_1, \dots, a_n)$ , and we're using  $\partial^\alpha$  for multi-indices and  $|\alpha|$  for the length of a multi-index. As well, we use  $\|\cdot\|_{L^p(U)}$  and  $\|\cdot\|_{W^{k,p}(U)}$  for  $L^p$  and Sobolev norms over a domain  $U \subset \mathbb{C}$ . When  $U = \mathbb{C}$ , we just write  $\|\cdot\|_p$  and  $\|\cdot\|_{k,p}$ .

**A.1. Proof of Proposition 5.7.** Here we provide details on Proposition 5.7, the interior elliptic estimates for the Beltrami operator. First we recall the Beurling transform  $T$ , which we already defined in Section 5 and on  $C_0^\infty(\mathbb{C})$  by the principal value integral

$$T(u)(z) = \lim_{\varepsilon \rightarrow 0} -\frac{1}{\pi} \int_{|\zeta - z| > \varepsilon} \frac{u(\zeta)}{(\zeta - z)^2} dx dy,$$

and the **Cauchy transform**  $P$ , defined on  $L^p(\mathbb{C})$ ,  $2 < p < \infty$ , by

$$P(u)(z) = -\frac{1}{\pi} \int_{\mathbb{C}} u(\zeta) \left( \frac{1}{\zeta - z} - \frac{1}{\zeta} \right) dx dy.$$

We restate Proposition 5.4 about the Beurling transform here.

**Proposition A.1.** *For all  $1 < p < \infty$ ,  $T$  extends to a continuous linear operator from  $L^p(\mathbb{C}) \rightarrow L^p(\mathbb{C})$ , which is an isometry when  $p = 2$ . The operator norm of  $T : L^p(\mathbb{C}) \rightarrow L^p(\mathbb{C})$  varies continuously with  $p$ .*

As in Section 5, we use  $C_p$  to denote the norm of  $T : L^p(\mathbb{C}) \rightarrow L^p(\mathbb{C})$ . For the Cauchy transform, we take note of the following results, which can be verified immediately.

**Proposition A.2.** *For  $2 < p < \infty$  and  $u \in L^p(\mathbb{C})$ , in the distributional sense,  $\partial_{\overline{z}}(P(u)) = u$ , and  $\partial_z(P(u)) = Tu$ .*

The following lemma represents the main step in the proof of Proposition 5.7.

**Lemma A.3.** *Let  $1 < p < \infty$ ,  $k \geq 0$ , and let  $u, v$  be  $C_0^\infty(\mathbb{C})$  functions supported in an open set  $U \subset \mathbb{C}$  and satisfying*

$$\left(\partial_{\bar{z}} - \mu\partial_z\right)u = v. \quad (56)$$

For  $p$  such that  $\|\mu\|_\infty C_p < 1$ , there exists  $C = C(\|\mu\|_{W^{k,\infty}(U)}, k, p)$  such that

$$\|u\|_{k+1,p} \leq C(\|u\|_{k,p} + \|v\|_{k,p}).$$

*Proof.* Let  $u \in C_0^\infty(\mathbb{C})$  and set  $h = \partial_{\bar{z}}u$ , so that by Proposition A.2,  $u = Ph$  and  $\partial_z u = Th$ . Using both of these identities, Equation (56) becomes the integral equation

$$v = \partial_{\bar{z}}(Ph) - k\partial_z(Ph) = h - \mu Th. \quad (57)$$

We claim that for all  $k \geq 0$

$$\|h\|_{k,p} \leq C(\|\mu\|_{W^{k,\infty}(U)}, k, p)\|v\|_{k,p},$$

for  $\|\mu\|_\infty C_p < 1$ , and we prove the result by induction on  $k$ . For  $k = 0$ , taking  $L^p$  norms on (57) gives

$$\|h\|_p \leq \|v\|_p + \|\mu\|_\infty C_p \|h\|_p,$$

and hence for  $\|\mu\|_\infty C_p < 1$ , we can rearrange to find that

$$\|h\|_p \leq (1 - \|\mu\|_\infty C_p)^{-1} \|v\|_p.$$

For the induction step, let  $\alpha$  be any multi-index with  $|\alpha| = k$ . Applying  $\partial^\alpha$  to (57), which commutes with  $T$ , we have

$$\partial^\alpha v = \partial^\alpha h - \mu T(\partial^\alpha h) + Q(\mu, h), \quad (58)$$

where  $Q$  is a polynomial in  $\partial^\beta \mu$ ,  $\partial^\gamma(Th) = T(\partial^\gamma h)$ , for  $0 \leq |\gamma| < |\alpha|$ ,  $1 \leq |\beta| \leq |\alpha|$ ,  $\beta + \gamma = \alpha$ , computed straightforwardly using the product rule.  $\|Q(\mu, h)\|_p$  is rather crudely bounded:

$$\|Q(\mu, h)\|_p \leq \sum_{\beta, \gamma} \|\partial^\beta \mu\|_\infty C_p \|\partial^\gamma h\|_p \leq C(k) \|\mu\|_{W^{k,\infty}(U)} C_p \|h\|_{k-1,p}.$$

Then using the induction hypothesis  $\|h\|_{k-1,p} \leq \|v\|_{k-1,p}$ , together with the trivial bound  $\|\cdot\|_{k-1,p} \leq \|\cdot\|_{k,p}$ , we extrapolate the final bound

$$\|Q(\mu, h)\|_p \leq C(k) \|\mu\|_{W^{k,\infty}(U)} C_p \|v\|_{k,p}.$$

Returning to (58), we deduce

$$\|\partial^\alpha h\|_p \leq \|\mu\|_\infty C_p \|\partial^\alpha h\|_p + (C(k) \|\mu\|_{W^{k,\infty}(U)} C_p + 1) \|v\|_{k,p}.$$

Under the assumption  $\|\mu\|_\infty C_p < 1$ , doing the same rearrangement as in the  $k = 0$  step returns the desired bound for  $\|\partial^\alpha h\|_p$ . Summing up over all multi-indices of length  $k$  completes the induction.

To finish the proof of the lemma, we turn our estimate on  $h$  into an estimate on  $u$  using elliptic regularity for  $\partial_{\bar{z}}$  relative to disks containing the support of  $u$ :

$$\|u\|_{k+1,p} \leq C(\|u\|_{k,p} + \|h\|_{k,p}) \leq C(\|u\|_{k,p} + \|v\|_{k,p}).$$

□

*Proof of Proposition 5.7.* The result follows swiftly from Lemma A.3 and standard approximation arguments, using the density of  $C_0^\infty(\mathbb{C})$  in all of the Sobolev spaces in question. One can follow the proof of Theorem 10.3.6 and Corollary 10.3.10 in [Nic07], as long as one remembers to restrict the range of  $p$  to those such that  $\|\mu\|_\infty C_p < 1$ . □

**A.2. Holomorphic dependence.** The main input toward Proposition 2.9 is a holomorphic dependence result for solutions to the Beltrami equation, which we prove in this subsection. To state the result properly, we recall the Measurable Riemann Mapping Theorem. We holomorphically embed  $\mathbb{C}$  inside  $\mathbb{CP}^1$  in the standard way.

**Theorem A.4** (Measurable Riemann Mapping Theorem). *Let  $\mu \in L_1^\infty(\mathbb{C})$ . There exists a quasiconformal homeomorphism from  $\mathbb{CP}^1 \rightarrow \mathbb{CP}^1$  with Beltrami form  $\mu$ , unique up to postcomposing by Möbius transformations.*

By the uniqueness condition, given  $\mu \in L_1^\infty(\mathbb{C})$ , there exists a unique quasiconformal homeomorphism  $f^\mu : \mathbb{C} \rightarrow \mathbb{C}$  fixing 0 and 1 (this is the restriction of the map from  $\mathbb{CP}^1 \rightarrow \mathbb{CP}^1$  fixing  $0, 1, \infty$ ). By Proposition 5.7, if  $\mu \in W_{loc}^{l,\infty}(\mathbb{C})$ , then  $f^\mu \in W_{loc}^{l+1,p}(\mathbb{C})$  for  $p$  such that  $\|\mu\|_\infty C_p < 1$ . Our result on holomorphic dependence is as follows.

**Theorem A.5.** *Let  $\Lambda$  be a complex manifold and*

$$\begin{aligned} \Lambda &\rightarrow L_1^\infty(\mathbb{C}) \cap W^{l,\infty}(\mathbb{C}) \\ t &\mapsto \mu(t), \end{aligned}$$

*a family of Beltrami forms with  $\|\mu(t)\|_\infty \leq \delta$  for some  $\delta < 1$ , which varies holomorphically with respect to the  $W^{l,\infty}$ -norm. For  $p$  such that  $\delta C_p < 1$ , the map from  $\Lambda \rightarrow W_{loc}^{l+1,p}(\mathbb{C})$  given by  $t \mapsto f^{\mu(t)}$  is holomorphic.*

Above, we interpret  $W_{loc}^{l+1,p}(\mathbb{C})$  as a complex Fréchet space. For  $l = 0$ , this is a classical result of Ahlfors and Bers [AB60, Theorem 11] (a proof is also found in [AIM09, §5.7]). To prove this higher regularity generalization we essentially go through the proof from [AB60], but use Proposition 5.7 to bootstrap certain estimates along the way.

Throughout this subsection, we fix the number  $\delta$  of Theorem A.5, and assume that all Beltrami forms considered here have  $L^\infty(\mathbb{C})$ -norm at most  $\delta$ , and we work with  $p$  such that  $\delta C_p < 1$ . Moving toward the proof, we first explain the proof of the  $l = 0$  case from [AB60].

**Lemma A.6** (Lemma 21 in [AB60]). *Let  $\mu \in L_1^\infty(\mathbb{C})$ ,  $a \in L^\infty(\mathbb{C})$ , and for a real or complex parameter  $s$ , let  $\alpha(s)$  be a continuously varying family of  $L^\infty(\mathbb{C})$  functions such that  $\|\alpha(s)\|_\infty$  is uniformly bounded in  $s$ , and  $\lim_{s \rightarrow 0} \|\alpha(s)\|_\infty = 0$ . We assume that  $s$  is chosen small enough so that  $\mu(s) = \mu + sa + s\alpha(s)$  satisfies  $\|\mu(s)\|_\infty \leq \delta$ . Then,*

$$\lim_{s \rightarrow 0} \frac{f^{\mu+sa+s\alpha(s)} - f^\mu}{s}$$

*exists in  $W_{loc}^{1,p}(\mathbb{C})$ , does not depend on  $\alpha$  and defines an element of  $W_{loc}^{1,p}(\mathbb{C})$ .*

As in Ahlfors-Bers, we denote the  $W_{loc}^{1,p}$ -limit by

$$\theta^{\mu,a} = \lim_{s \rightarrow 0} \frac{f^{\mu+sa+s\alpha(s)} - f^\mu}{s}.$$

After their Lemma 21, the authors show the following. In the lemma below,  $\mu, a$ , and  $\alpha(s)$  are as above.

**Lemma A.7** (Part (iii) of Lemma 22 in [AB60]). *Assume we are also given uniformly bounded sequences  $(\mu_n)_{n=1}^\infty \subset L_1^\infty(\mathbb{C})$  and  $(a_n)_{n=1}^\infty \subset L^\infty(\mathbb{C})$  such that  $\lim_{n \rightarrow \infty} \mu_n = \mu$  and  $\lim_{n \rightarrow \infty} a_n = a$  in  $L^\infty(\mathbb{C})$ . Then,*

$$\lim_{n \rightarrow \infty} \theta^{\mu_n, a_n} = \theta^{\mu, a}$$

*in  $W_{loc}^{1,p}(\mathbb{C})$ .*

With these two results, they prove that  $f^\mu$  depends differentiably on  $\mu$ , in their chosen Sobolev spaces.

**Theorem A.8** (Theorem 10 in [AB60]). *Let  $t = (t_1, \dots, t_n)$  and  $s = (s_1, \dots, s_n)$  be real or complex parameters, and suppose that we are given a family of Beltrami forms  $\mu(t)$  such that for all  $t$  in some open set,*

$$\mu(t+s) = \mu(t) + \sum_{i=1}^n a_i(t) s_i + |s| \alpha(t, s),$$

where  $a_1, \dots, a_n$  are uniformly bounded functions in  $L^\infty(\mathbb{C})$  satisfying  $a_i(t+s) - a_i(t) \rightarrow 0$  in  $L^\infty(\mathbb{C})$  as  $s \rightarrow 0$ , and  $\alpha(t, s)$  is a uniformly bounded family of functions in  $L^\infty(\mathbb{C})$  satisfying  $\lim_{s \rightarrow 0} \|\alpha(t, s)\|_\infty = 0$  for all  $t$ . Then  $f^{\mu(t+s)}$  admits the development

$$f^{\mu(t+s)} = f^{\mu(t)} + \sum_{i=1}^n \theta^{\mu(t), a_i(t)} s_i + |s| \gamma(t, s), \quad (59)$$

where  $\gamma(t, s)$  is a continuously varying family of functions in  $W_{loc}^{1,p}(\mathbb{C})$ , uniformly bounded in  $W_{loc}^{1,p}(\mathbb{C})$ , and such that, for every  $t$ ,  $\gamma(t, s)$  tends to 0 in  $W_{loc}^{1,p}(\mathbb{C})$  as  $s \rightarrow 0$ .

This result is more or less immediate from the two lemmas above. Indeed, Lemma A.6 shows that one can differentiate  $f^{\mu(t)}$  in  $t$ , which returns  $\sum_{i=1}^n \theta^{\mu(t), a_i(t)}$ , and Lemma A.7 shows that the remainder term in the first order Taylor expansion in  $W_{loc}^{1,p}(\mathbb{C})$  has the right decay. See the proof of [AB60, Theorem 10] for the quick proof.

With their Theorem 10, Ahlfors and Bers deduce holomorphic dependence with ease: assuming that  $\mu(t)$  depends holomorphically on complex parameters  $(t_1, \dots, t_n)$ , then the formula (59) shows that, as an element of  $W_{loc}^{1,p}(\mathbb{C})$ ,  $f^{\mu(t)}$  satisfies the Cauchy-Riemann equations in every  $t_1, \dots, t_n$ , and hence  $t \mapsto f^{\mu(t)}$  is holomorphic in  $W_{loc}^{1,p}(\mathbb{C})$ .

Now, for Theorem A.5, we need to establish the analogue of Theorem 10 from [AB60]. We package the needed ingredients into the lemma below. Given  $\mu \in L_1^\infty(\mathbb{C})$  and  $a \in L^\infty(\mathbb{C})$ , we keep the same definition of  $\theta^{\mu, a}$ , which we already know defines an element of  $W_{loc}^{1,p}(\mathbb{C})$ .

**Lemma A.9.** *In the setting of Lemma 21 from [AB60], now assume that  $\mu, a, \alpha(s) \in W^{l,\infty}(\mathbb{C})$ , that  $s \mapsto \alpha(s)$  is continuous as a map to  $W^{l,\infty}(\mathbb{C})$ , and that  $\|\alpha(s)\|_{l,\infty}$  is uniformly bounded in  $s$ , and tends to 0 as  $s \rightarrow 0$ . Then,*

- (1)  $\theta^{\mu, a} \in W_{loc}^{l+1,p}(\mathbb{C})$ ,
- (2) and

$$\theta^{\mu, a} = \lim_{s \rightarrow 0} \frac{f^{\mu+sa+s\alpha(s)} - f^\mu}{s}$$

in  $W_{loc}^{l+1,p}(\mathbb{C})$ .

Moreover, given uniformly bounded sequences  $(\mu_n)_{n=1}^\infty \subset L_1^\infty(\mathbb{C}) \cap W^{l,\infty}(\mathbb{C})$  and  $(a_n)_{n=1}^\infty \subset W^{l,\infty}(\mathbb{C})$  such that  $\lim_{n \rightarrow \infty} \mu_n = \mu$  and  $\lim_{n \rightarrow \infty} a_n = a$  in  $W^{l,\infty}(\mathbb{C})$ , we have that

$$\lim_{n \rightarrow \infty} \theta^{\mu_n, a_n} = \theta^{\mu, a}$$

in  $W_{loc}^{l+1,p}(\mathbb{C})$ .

Theorem A.5 follows from Lemma A.9 by the exact same argument as in [AB60]: if  $\mu(t)$  depends on a complex parameter  $t = (t_1, \dots, t_n)$ , then for small  $s = (s_1, \dots, s_n)$  we can now write

$$f^{\mu(t+s)} = f^{\mu(t)} + \sum_{i=1}^n \theta^{\mu(t), a_i(t)} s_i + |s| \gamma(t, s), \quad (60)$$

with every term the same function as in Theorem A.8, but also lying in  $W_{loc}^{l+1,p}(\mathbb{C})$ , and with  $\gamma(t, s)$  tending to 0 in  $W_{loc}^{l+1,p}(\mathbb{C})$  as  $s \rightarrow 0$  for every  $t$ . That is,  $f^{\mu(t)}$  depends differentiably in  $t$  as an element of  $W_{loc}^{l+1,p}(\mathbb{C})$ . And, as in [AB60, Theorem 11], if the  $\mu(t)$ 's vary holomorphically in  $W^{l,\infty}(\mathbb{C})$ , then by the formula (60),  $f^{\mu(t)}$  satisfies the Cauchy-Riemann equations in  $W_{loc}^{l+1,p}(\mathbb{C})$ , which yields the statement of Theorem A.5. So, it remains to prove Lemma A.9.

*Proof of Lemma A.9.* We bootstrap on the estimates implied by Lemmas A.6 and A.7, using Proposition 5.7. To show our bounds and convergence in  $W_{loc}^{1,p}(\mathbb{C})$ , it suffices to prove that for any fixed ball  $U$ , that we have bounds and convergence in  $W^{l+1,p}(U)$ . Recall that at the beginning of this subsection we declared  $p$  to be close enough to 2 so that  $\delta C_p < 1$ , so in particular we can always apply Proposition 5.7. As well, choosing  $s$  small, we can always assume that  $f^{\mu+sa+s\alpha(s)}$  is uniformly bounded in  $W_{loc}^{1+1,p}(\mathbb{C})$ , independently of  $s$ .

Note that (as is observed in [AB60]),

$$(\partial_{\bar{z}} - \mu\partial_z)\theta^{\mu,a} = a\partial_z f^\mu. \quad (61)$$

Thus, since  $a \in W^{l,\infty}(\mathbb{C})$  and  $f^\mu \in W_{loc}^{l+1,p}(\mathbb{C})$ ,

$$(\partial_{\bar{z}} - \mu\partial_z)\theta^{\mu,a} \in W_{loc}^{l,p}(\mathbb{C}).$$

The first assertion of Lemma A.9 then follows routinely: by Proposition 5.7, for any ball  $V$  containing  $U$ ,

$$\|\theta^{\mu,a}\|_{W^{2,p}(U)} \leq C(\|\mu\|_{l,\infty}, U, V, 1, p)(\|\theta^{\mu,a}\|_{W^{1,p}(V)} + \|(\partial_{\bar{z}} - \mu\partial_z)\theta^{\mu,a}\|_{W^{1,p}(V)}),$$

so  $\theta^{\mu,a} \in W_{loc}^{2,p}(\mathbb{C})$ . Bootstrapping that estimate as much as we can, we obtain  $\theta^{\mu,a} \in W_{loc}^{l+1,p}(\mathbb{C})$ .

For the second assertion, setting

$$\hat{\chi}(s) = \frac{f^{\mu+sa+s\alpha(s)} - f^\mu}{s},$$

observe that

$$(\partial_{\bar{z}} - \mu\partial_z)\hat{\chi}(s) = (a + s\alpha(s))\partial_z f^{\mu+sa+s\alpha(s)},$$

which is uniformly bounded in  $W_{loc}^{l,p}(\mathbb{C})$ , since for small  $s$  the  $W^{l,p}(U)$  norm is dominated by  $(\|a\|_{l,\infty} + 1)\|f\|_{W^{l+1,p}(U)}$ . Therefore, by Proposition 5.7, since we know by Lemma A.6 that  $\hat{\chi}(s)$  is uniformly bounded in  $W_{loc}^{1,p}(\mathbb{C})$ , we get by Proposition 5.7 that it is uniformly bounded in  $W_{loc}^{2,p}(\mathbb{C})$ . Iterating the bootstrap as above, we obtain that  $\hat{\chi}(s)$  is uniformly bounded in  $W_{loc}^{l+1,p}(\mathbb{C})$ . Moreover the function

$$\chi(s) = \frac{f^{\mu+sa+s\alpha(s)} - f^\mu}{s} - \theta^{\mu,a},$$

which we know tends to 0 in  $W_{loc}^{1,p}(\mathbb{C})$  as  $s \rightarrow 0$ , is uniformly bounded in  $W_{loc}^{l+1,p}(\mathbb{C})$ . We further compute that

$$(\partial_{\bar{z}} - \mu\partial_z)\chi(s) = (a + s\alpha(s))\partial_z f^{\mu+sa+s\alpha(s)} - a\partial_z f^\mu,$$

from which we deduce that  $(\partial_{\bar{z}} - \mu\partial_z)\chi(s)$  tends to 0 in  $W_{loc}^{l,p}(\mathbb{C})$  as  $s \rightarrow 0$ , since

$$\begin{aligned} \|(\partial_{\bar{z}} - \mu\partial_z)\chi\|_{W^{l,p}(U)} &\leq \|\alpha(s)\|_{l,\infty} \|f^{\mu+sa+s\alpha(s)}\|_{W^{l+1,p}(U)} + \|a\|_{l,\infty} (\|f^{\mu+sa+s\alpha(s)} - f^\mu\|_{W^{l+1,p}(U)}) \\ &= \|\alpha(s)\|_{l,\infty} \|f^{\mu+sa+s\alpha(s)}\|_{W^{l+1,p}(U)} + \|a\|_{l,\infty} |s| \cdot \|\hat{\chi}\|_{W^{l+1,p}(U)}, \end{aligned}$$

and both terms above tend to 0 as  $s \rightarrow 0$ . Bootstrapping via Proposition 5.7 again, we deduce that  $\chi(s)$  tends to 0 in  $W_{loc}^{1+1,p}(\mathbb{C})$  as  $s \rightarrow 0$ , which proves the second assertion.



Finally, for the “moreover” statement, we compute, using (61), that

$$\begin{aligned} (\partial_{\bar{z}} - \mu\partial_z)(\theta^{\mu,a} - \theta^{\mu_n,a_n}) &= (\partial_{\bar{z}} - \mu\partial_z)\theta^{\mu,a} - (\partial_{\bar{z}} - \mu_n\partial_z)\theta^{\mu_n,a_n} + (\mu_n - \mu)\partial_z\theta^{\mu_n,a_n} \\ &= a\partial_z f^\mu - a_n\partial_z f^{\mu_n} + (\mu_n - \mu)\partial_z\theta^{\mu_n,a_n}, \end{aligned}$$

and hence for  $n$  large,

$$\|(\partial_{\bar{z}} - \mu\partial_z)(\theta^{\mu,a} - \theta^{\mu_n,a_n})\|_{W^{l,p}(U)} \leq (\|a\|_{l,\infty} + 1)\|f^\mu - f^{\mu_n}\|_{W^{l,p}(U)} + \|\mu_n - \mu\|_{l,\infty}\|\theta^{\mu_n,a_n}\|_{W^{l+1,p}(U)}.$$

Since the bounds on  $\|\theta^{\mu_n,a_n}\|_{W^{l+1,p}(U)}$  depend only on the  $W^{l,\infty}(\mathbb{C})$  norms of  $\mu_n$  and  $a_n$ , they are uniformly controlled. It follows that

$$\lim_{n \rightarrow \infty} \|(\partial_{\bar{z}} - \mu\partial_z)(\theta^{\mu,a} - \theta^{\mu_n,a_n})\|_{W^{l,p}(U)} = 0.$$

By yet another bootstrap using Proposition 5.7, we conclude that  $\theta^{\mu_n,a_n}$  tends to  $\theta^{\mu,a}$  in  $W_{loc}^{l+1,p}(\mathbb{C})$ , as desired.  $\square$

As discussed above, verifying Lemma A.9 completes the proof of Theorem A.5.

**A.3. Proof of Proposition 2.9.** Before explaining how to deduce Proposition 2.9 from Theorem A.5, we first need to recall how to prove Theorem 2.7, our version of Bers’ Simultaneous Uniformization Theorem, from the Measurable Riemann Mapping Theorem. Given  $(c_1, \bar{c}_2) \in \mathcal{C}^l(S) \times \mathcal{C}^l(\bar{S})$ , let us first assume that  $c_2$  is the basepoint with respect to which we defined  $\mathcal{C}(S)$ , and let’s holomorphically identify the universal cover of  $(S, c_2)$  with  $\mathbb{H}$ . Then  $c_1$  is represented by a Beltrami form  $\mu$  on  $(S, c_2)$ , which we lift to an invariant Beltrami form on the universal cover, and still denote by  $\mu$ . One defines the Beltrami form  $\hat{\mu}$  by  $\hat{\mu} = \mu$  on  $\mathbb{H}$ , and  $\hat{\mu} = 0$  on  $\bar{\mathbb{H}}$ . Applying the Measurable Riemann Mapping Theorem, we obtain the quasiconformal map  $f^{\hat{\mu}}$ , which we view as a homeomorphism of  $\mathbb{C}\mathbb{P}^1$ . The maps  $f_+$  and  $f_-$  from the statement of Theorem 2.7 are defined, for  $z \in \mathbb{H}$ , by  $f_+(z) = f^{\hat{\mu}}(z)$  and  $f_-(z) = f^{\hat{\mu}}(\bar{z})$ .

For the general case, we change the basepoint to  $c_2$ , and then apply the procedure above. Let’s explain how this works. To make the basepoint explicit, let us now denote by  $\mathcal{C}_{c_0}^l(S)$  the space of  $W^{l,\infty}$  complex structures based at a complex structure  $c_0$ . Let  $c$  be some other complex structure, and  $f_c : (S, c_0) \rightarrow (S, c)$  the diffeomorphism relating the two structures. Given a Beltrami form  $\mu$  giving rise via the Measurable Riemann Mapping Theorem to a complex structure  $c_\mu \in \mathcal{C}_{c_0}^l(S)$  with quasiconformal map  $f_\mu : (S, c_0) \rightarrow (S, c_\mu)$ , one can think of  $c_\mu$  as a complex structure based on  $(S, c)$  by taking the Beltrami form  $\mu_c \in \mathcal{C}_c^l(S)$  of  $(f_\mu \circ f_c^{-1})$ . We use notation like  $f_c$  and  $f_\mu$  (maps between surfaces) to distinguish from  $f^\mu$  (a self-mapping of  $\mathbb{C}\mathbb{P}^1$ ). Going back to Bers’ Theorem, one starts with  $(c_1, \bar{c}_2) \in \mathcal{C}^l(S) \times \mathcal{C}^l(\bar{S})$ , changes the basepoint to  $c_2$  in order to turn the data of  $(c_1, \bar{c}_2)$  into that of a Beltrami form on  $(S, c_2)$ , and then follows the procedure above.

In all of this discussion, one needs to keep track of the regularity. Given  $c \in \mathcal{C}_{c_0}^l(S)$ ,  $f_c$  is in  $W^{l+1,p}(\mathbb{C})$ , for  $p$  slightly larger than 2. If  $\mu$  is any Beltrami form as above with map  $f^\mu$ , by the formula for the Beltrami form of a composition (see [IT92, Proposition 4.13]), the Beltrami form of the map  $f_\mu \circ f_c^{-1}$  is  $W^{l,\infty}$ . Moreover, the maps produced by Bers’ Theorem are  $W^{l+1,p}$ , for  $p > 2$  depending on  $c$  and  $\mu$ .

**Lemma A.10.** *The change of basepoint map  $\mathcal{C}_{c_0}^l(S) \rightarrow \mathcal{C}_c^l(S)$ , described on the level of Beltrami forms by  $\mu \mapsto \mu_c$ , is holomorphic.*

Recall from classical Teichmüller theory that, in the notation above,  $f_c$  is constructed by taking the Beltrami form  $\mu_0$  on  $(S, c_0)$  representing  $c$ , identifying the universal cover with  $\mathbb{H}$  and lifting  $\mu_0$  there, and extending  $\mu_0$  by reflection across  $\bar{\mathbb{H}}$  to obtain a new Beltrami form

$\mu_0^*$ . One applies the Measurable Riemann Mapping Theorem to obtain the solution to the Beltrami equation  $f^{\mu_0^*}$ , which then conjugates the Fuchsian group uniformizing  $(S, c_0)$  to the Fuchsian group uniformizing  $(S, c)$ . Taking the quotients, we obtain the quasiconformal map  $f_c : (S, c_0) \rightarrow (S, c)$ .

*Proof.* Let  $U$  be a fundamental domain in  $\mathbb{H}$  for the action of the Fuchsian group uniformizing  $(S, c_0)$ . The  $W^{l, \infty}$  norm of an element in  $\mathcal{C}_{c_0}^l(S)$  can be computed by lifting to the universal cover and computing the  $W^{l, \infty}$  norm in  $U$ .

For any holomorphic variation  $\lambda \mapsto \mu_\lambda$ , let  $f_\lambda$  be the lift of  $f_{\mu_\lambda}$  to  $\mathbb{H}$ . By Theorem A.5,  $\lambda \mapsto f_\lambda$  defines a holomorphic map into  $W_{loc}^{l+1, p}(\mathbb{C})$ , for  $p > 2$  sufficiently close to 2. It is readily checked that  $\lambda \mapsto f_\lambda \circ f_c^{-1}$  is holomorphic as well, and moreover that the map to  $\mathcal{C}_c^l(S)$  is holomorphic, since taking the Beltrami form of a  $W_{loc}^{l+1, p}(\mathbb{C})$  quasiconformal map is a holomorphic operation.  $\square$

*Proof of Proposition 2.9.* By Hartogs' theorem on separate analyticity, it suffices to verify holomorphicity on variations of the form  $\lambda \mapsto (c_1^\lambda, \overline{c_2})$  and  $\lambda \mapsto (c_1, \overline{c_2}^\lambda)$ . Since the argument will be symmetric, we'll just carry out the first verification. Given a variation  $\lambda \mapsto (c_1^\lambda, \overline{c_2})$ , we change the basepoint of our space of complex structures to be  $c_2$ , so that our variation is equivalent to a holomorphic variation of Beltrami forms  $\lambda \mapsto \mu_\lambda$  for  $c_2$ . By Lemma A.10, for  $p$  close enough to 2, this change of basepoint operation is holomorphic.

Next, as in the proof of Bers' Theorem, we lift to  $\mathbb{H}$ , construct  $\widetilde{\mu}_\lambda$ , and take the quasiconformal maps  $f^{\widetilde{\mu}_\lambda}$ . As in the proof of Lemma A.10, if  $U$  is a fundamental domain in  $\mathbb{H}$  for the action of the Fuchsian group uniformizing  $(S, c_2)$ , the claim reduces to showing that  $\lambda \mapsto (f^{\widetilde{\mu}_\lambda})^* \langle \cdot, \cdot \rangle_{\mathbb{G}}$  is holomorphic as a map to  $W^{l, 2}(U)$ , where we recall that  $\langle \cdot, \cdot \rangle_{\mathbb{G}}$  is the holomorphic Riemannian metric on  $\mathbb{G}$ .

Toward this, by Theorem A.5, for  $p > 2$  such that  $\|\mu_\lambda\|_\infty C_p < 1$ , the variation  $\lambda \mapsto f^{\widetilde{\mu}_\lambda} \in W^{l+1, p}(U)$  is holomorphic. Since  $U$  is relatively compact,  $W^{l+1, p}(U)$  embeds into  $W^{l+1, 2}(U)$  linearly and continuously, thus holomorphically. Finally, by the formula (5), the pullback map from  $W^{l+1, 2}(U)$  to  $W^{l, 2}(U)$  taking  $f \mapsto f^* \langle \cdot, \cdot \rangle_{\mathbb{G}}$  is holomorphic, and the result follows.  $\square$

#### REFERENCES

- [AB60] Lars Ahlfors and Lipman Bers. "Riemann's mapping theorem for variable metrics". In: *Ann. of Math. (2)* 72 (1960), pp. 385–404. ISSN: 0003-486X. DOI: 10.2307/1970141. URL: <https://doi.org/10.2307/1970141>.
- [Abe91] Kinetsu Abe. "Affine differential geometry of complex hypersurfaces". In: *Geometry and topology of submanifolds, III (Leeds, 1990)*. World Sci. Publ., River Edge, NJ, 1991, pp. 1–31. ISBN: 981-02-0310-1.
- [ADL21] Daniele Alessandrini, Colin Davalo, and Qionglin Li. *Projective Structures with (Quasi-)Hitchin Holonomy*. 2021. arXiv: 2110.15407 [id='math.GT'].
- [AIM09] Kari Astala, Tadeusz Iwaniec, and Gaven Martin. *Elliptic partial differential equations and quasiconformal mappings in the plane*. Vol. 48. Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 2009, pp. xviii+677. ISBN: 978-0-691-13777-3.
- [Ale+23] Daniele Alessandrini et al. *Fiber bundles associated with Anosov representations*. 2023. arXiv: 2303.10786 [math.GT].
- [Bar10] David Baraglia. *G2 geometry and integrable systems*. 2010. arXiv: 1002.1767 [math.DG].

- [BD96] A. A. Beilinson and V. G. Drinfeld. “Quantization of Hitchin’s fibration and Langlands’ program”. In: *Algebraic and geometric methods in mathematical physics (Kaciveli, 1993)*. Vol. 19. Math. Phys. Stud. Kluwer Acad. Publ., Dordrecht, 1996, pp. 3–7. ISBN: 0-7923-3909-6.
- [BE22] Francesco Bonsante and Christian El Emam. “On immersions of surfaces into  $SL(2, \mathbb{C})$  and geometric consequences”. In: *Int. Math. Res. Not. IMRN* 12 (2022), pp. 8803–8864. ISSN: 1073-7928,1687-0247. DOI: 10.1093/imrn/rnab189. URL: <https://doi.org/10.1093/imrn/rnab189>.
- [Beg94] Heinrich G. W. Begehr. *Complex analytic methods for partial differential equations*. An introductory text. World Scientific Publishing Co., Inc., River Edge, NJ, 1994, pp. x+273. ISBN: 981-02-1550-9. DOI: 10.1142/2162. URL: <https://doi.org/10.1142/2162>.
- [Ber60] Lipman Bers. “Simultaneous uniformization”. In: *Bull. Amer. Math. Soc.* 66 (1960), pp. 94–97. ISSN: 0002-9904. DOI: 10.1090/S0002-9904-1960-10413-2. URL: <https://doi.org/10.1090/S0002-9904-1960-10413-2>.
- [Ber61] Lipman Bers. “Holomorphic differentials as functions of moduli”. In: *Bull. Amer. Math. Soc.* 67 (1961), pp. 206–210. ISSN: 0002-9904. DOI: 10.1090/S0002-9904-1961-10569-7. URL: <https://doi.org/10.1090/S0002-9904-1961-10569-7>.
- [Bla20] Simon Blatt. *On the analyticity of solutions to non-linear elliptic partial differential systems*. 2020. arXiv: 2009.08762 [math.AP].
- [Cal72] Eugenio Calabi. “Complete affine hyperspheres. I”. In: *Symposia Mathematica, Vol. X (Convegno di Geometria Differenziale, INDAM, Rome, 1971 & Convegno di Analisi Numerica, INDAM, Rome, 1972)*. Academic Press, London-New York, 1972, pp. 19–38.
- [CG93] Suhyoung Choi and William M. Goldman. “Convex real projective structures on closed surfaces are closed”. In: *Proc. Amer. Math. Soc.* 118.2 (1993), pp. 657–661. ISSN: 0002-9939. DOI: 10.2307/2160352. URL: <https://doi.org/10.2307/2160352>.
- [CTT19] Brian Collier, Nicolas Tholozan, and Jérémy Toullisse. “The geometry of maximal representations of surface groups into  $SO_0(2, n)$ ”. In: *Duke Math. J.* 168.15 (2019), pp. 2873–2949. ISSN: 0012-7094,1547-7398. DOI: 10.1215/00127094-2019-0052. URL: <https://doi.org/10.1215/00127094-2019-0052>.
- [CW19] Brian Collier and Richard Wentworth. “Conformal limits and the Białynicki-Birula stratification of the space of  $\lambda$ -connections”. In: *Adv. Math.* 350 (2019), pp. 1193–1225. ISSN: 0001-8708,1090-2082. DOI: 10.1016/j.aim.2019.04.034. URL: <https://doi.org/10.1016/j.aim.2019.04.034>.
- [CY77] Shiu Yuen Cheng and Shing Tung Yau. “On the regularity of the Monge-Ampère equation  $\det(\partial^2 u / \partial x_i \partial x_j) = F(x, u)$ ”. In: *Comm. Pure Appl. Math.* 30.1 (1977), pp. 41–68. ISSN: 0010-3640,1097-0312. DOI: 10.1002/cpa.3160300104. URL: <https://doi.org/10.1002/cpa.3160300104>.
- [CY86] Shiu Yuen Cheng and Shing-Tung Yau. “Complete affine hypersurfaces. I. The completeness of affine metrics”. In: *Comm. Pure Appl. Math.* 39.6 (1986),

- pp. 839–866. ISSN: 0010-3640,1097-0312. DOI: 10.1002/cpa.3160390606. URL: <https://doi.org/10.1002/cpa.3160390606>.
- [Dav23] Colin Davalo. *Nearly geodesic immersions and domains of discontinuity*. 2023. arXiv: 2303.11260 [math.DG].
- [DGV15] C. T. J. Dodson, George Galanis, and Efstathios Vassiliou. *Geometry in a Fréchet Context: A Projective Limit Approach*. London Mathematical Society Lecture Note Series. Cambridge University Press, 2015.
- [DS20] David Dumas and Andrew Sanders. “Geometry of compact complex manifolds associated to generalized quasi-Fuchsian representations”. In: *Geom. Topol.* 24.4 (2020), pp. 1615–1693. ISSN: 1465-3060,1364-0380. DOI: 10.2140/gt.2020.24.1615. URL: <https://doi.org/10.2140/gt.2020.24.1615>.
- [DS21] David Dumas and Andrew Sanders. “Uniformization of compact complex manifolds by Anosov homomorphisms”. In: *Geom. Funct. Anal.* 31.4 (2021), pp. 815–854. ISSN: 1016-443X,1420-8970. DOI: 10.1007/s00039-021-00572-6. URL: <https://doi.org/10.1007/s00039-021-00572-6>.
- [DS84] V. G. Drinfeld and V. V. Sokolov. “Lie algebras and equations of Korteweg-de Vries type”. In: *Current problems in mathematics, Vol. 24. Itogi Nauki i Tekhniki. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow*, 1984, pp. 81–180.
- [Dum+21a] Olivia Dumitrescu et al. “From the Hitchin section to opers through nonabelian Hodge”. In: *J. Differential Geom.* 117.2 (2021), pp. 223–253. ISSN: 0022-040X,1945-743X. DOI: 10.4310/jdg/1612975016. URL: <https://doi.org/10.4310/jdg/1612975016>.
- [Dum+21b] Olivia Dumitrescu et al. “From the Hitchin section to opers through nonabelian Hodge”. In: *J. Differential Geom.* 117.2 (2021), pp. 223–253. ISSN: 0022-040X,1945-743X. DOI: 10.4310/jdg/1612975016. URL: <https://doi.org/10.4310/jdg/1612975016>.
- [DVV88] F. Dillen, L. Vrancken, and L. Verstraelen. “Complex affine differential geometry”. In: *Atti Accad. Peloritana Pericolanti. Cl. Sci. Fis. Mat. Natur.* 66 (1988), pp. 231–260. ISSN: 0365-0359,1825-1242.
- [DW15] David Dumas and Michael Wolf. “Polynomial cubic differentials and convex polygons in the projective plane”. In: *Geom. Funct. Anal.* 25.6 (2015), pp. 1734–1798. ISSN: 1016-443X,1420-8970. DOI: 10.1007/s00039-015-0344-5. URL: <https://doi.org/10.1007/s00039-015-0344-5>.
- [DZ09] Sorin Dumitrescu and Abdelghani Zeghib. “Global rigidity of holomorphic Riemannian metrics on compact complex 3-manifolds”. In: *Math. Ann.* 345.1 (2009), pp. 53–81. ISSN: 0025-5831,1432-1807. DOI: 10.1007/s00208-009-0342-8. URL: <https://doi.org/10.1007/s00208-009-0342-8>.
- [EE69] Clifford J. Earle and James Eells. “A fibre bundle description of Teichmüller theory”. In: *Journal of Differential Geometry* 3.1-2 (1969), pp. 19–43. DOI: 10.4310/jdg/1214428816. URL: <https://doi.org/10.4310/jdg/1214428816>.
- [Ema23] Christian El Emam. *A metric uniformization model for the Quasi-Fuchsian space*. 2023. arXiv: 2307.07388 [math.DG].

- [ES22] Christian El Emam and Andrea Seppi. “On the Gauss map of equivariant immersions in hyperbolic space”. In: *J. Topol.* 15.1 (2022), pp. 238–301. ISSN: 1753-8416,1753-8424.
- [Fol95] Gerald B. Folland. *Introduction to partial differential equations*. Second. Princeton University Press, Princeton, NJ, 1995, pp. xii+324. ISBN: 0-691-04361-2.
- [Gai14] Davide Gaiotto. “Opers and TBA”. In: (Mar. 2014). arXiv: 1403.6137 [hep-th].
- [Gol90] William M. Goldman. “Convex real projective structures on compact surfaces”. In: *J. Differential Geom.* 31.3 (1990), pp. 791–845. ISSN: 0022-040X,1945-743X. URL: <http://projecteuclid.org/euclid.jdg/1214444635>.
- [GP10] Victor Guillemin and Alan Pollack. *Differential topology*. Reprint of the 1974 original. AMS Chelsea Publishing, Providence, RI, 2010, pp. xviii+224. ISBN: 978-0-8218-5193-7. DOI: 10.1090/chel/370. URL: <https://doi.org/10.1090/chel/370>.
- [GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Reprint of the 1998 edition. Springer-Verlag, Berlin, 2001, pp. xiv+517. ISBN: 3-540-41160-7.
- [Gui18] Olivier Guichard. “An introduction to the differential geometry of flat bundles and of Higgs bundles”. In: *The geometry, topology and physics of moduli spaces of Higgs bundles*. Vol. 36. Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap. World Sci. Publ., Hackensack, NJ, 2018, pp. 1–63. ISBN: 978-981-3229-08-2.
- [Gun67] R. C. Gunning. *Lectures on vector bundles over Riemann surfaces*. University of Tokyo Press, Tokyo; Princeton University Press, Princeton, NJ, 1967, pp. v+243.
- [Has06] Yoshiaki Hashimoto. “A remark on the analyticity of the solutions for non-linear elliptic partial differential equations”. In: *Tokyo J. Math.* 29.2 (2006), pp. 271–281. ISSN: 0387-3870. DOI: 10.3836/tjm/1170348166. URL: <https://doi.org/10.3836/tjm/1170348166>.
- [Hit92] N. J. Hitchin. “Lie groups and Teichmüller space”. In: *Topology* 31.3 (1992), pp. 449–473. ISSN: 0040-9383. DOI: 10.1016/0040-9383(92)90044-I. URL: [https://doi.org/10.1016/0040-9383\(92\)90044-I](https://doi.org/10.1016/0040-9383(92)90044-I).
- [HLL13] Zheng Huang, John Loftin, and Marcello Lucia. “Holomorphic cubic differentials and minimal Lagrangian surfaces in  $\mathbb{C}\mathbb{H}^2$ ”. In: *Math. Res. Lett.* 20.3 (2013), pp. 501–520. ISSN: 1073-2780,1945-001X. DOI: 10.4310/MRL.2013.v20.n3.a8. URL: <https://doi.org/10.4310/MRL.2013.v20.n3.a8>.
- [Hod06] Thomas Hodge. “Hyperkahler Geometry and Teichmüller Space”. In: (Jan. 2006).
- [IT92] Y. Imayoshi and M. Taniguchi. *An introduction to Teichmüller spaces*. Translated and revised from the Japanese by the authors. Springer-Verlag, Tokyo, 1992, pp. xiv+279. ISBN: 4-431-70088-9. DOI: 10.1007/978-4-431-68174-8. URL: <https://doi.org/10.1007/978-4-431-68174-8>.
- [Kat96] Keiichi Kato. “New idea for proof of analyticity of solutions to analytic non-linear elliptic equations”. In: *SUT J. Math.* 32.2 (1996), pp. 157–161. ISSN: 0916-5746.

- [Kim07] Young-Heon Kim. “Holomorphic extensions of Laplacians and their determinants”. In: *Adv. Math.* 211.2 (2007), pp. 517–545. ISSN: 0001-8708,1090-2082. DOI: 10.1016/j.aim.2006.09.009. URL: <https://doi.org/10.1016/j.aim.2006.09.009>.
- [KM97] Andreas Kriegl and Peter W. Michor. *The convenient setting of global analysis*. Vol. 53. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1997, pp. x+618. ISBN: 0-8218-0780-3. DOI: 10.1090/surv/053. URL: <https://doi.org/10.1090/surv/053>.
- [Kos59] Bertram Kostant. “The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group”. In: *Amer. J. Math.* 81 (1959), pp. 973–1032. ISSN: 0002-9327,1080-6377. DOI: 10.2307/2372999. URL: <https://doi.org/10.2307/2372999>.
- [KZ17] In Kang Kim and Genkai Zhang. “Kähler metric on the space of convex real projective structures on surface”. In: *Journal of Differential Geometry* 106.1 (2017), pp. 127–137.
- [Lab06] François Labourie. “Anosov flows, surface groups and curves in projective space”. In: *Invent. Math.* 165.1 (2006), pp. 51–114. ISSN: 0020-9910,1432-1297. DOI: 10.1007/s00222-005-0487-3. URL: <https://doi.org/10.1007/s00222-005-0487-3>.
- [Lab07] François Labourie. “Flat projective structures on surfaces and cubic holomorphic differentials”. In: *Pure Appl. Math. Q.* 3.4 (2007), pp. 1057–1099. ISSN: 1558-8599,1558-8602. DOI: 10.4310/PAMQ.2007.v3.n4.a10. URL: <https://doi.org/10.4310/PAMQ.2007.v3.n4.a10>.
- [Lab17] François Labourie. “Cyclic surfaces and Hitchin components in rank 2”. In: *Ann. of Math. (2)* 185.1 (2017), pp. 1–58. ISSN: 0003-486X,1939-8980. DOI: 10.4007/annals.2017.185.1.1. URL: <https://doi.org/10.4007/annals.2017.185.1.1>.
- [Li19] Qionglng Li. “An introduction to Higgs bundles via harmonic maps”. In: *SIGMA Symmetry Integrability Geom. Methods Appl.* 15 (2019), Paper No. 035, 30. ISSN: 1815-0659. DOI: 10.3842/SIGMA.2019.035. URL: <https://doi.org/10.3842/SIGMA.2019.035>.
- [LM13] John Loftin and Ian McIntosh. “Minimal Lagrangian surfaces in  $\mathbb{C}\mathbb{H}^2$  and representations of surface groups into  $SU(2, 1)$ ”. In: *Geom. Dedicata* 162 (2013), pp. 67–93. ISSN: 0046-5755,1572-9168. DOI: 10.1007/s10711-012-9717-1. URL: <https://doi.org/10.1007/s10711-012-9717-1>.
- [Lof01] John C. Loftin. “Affine spheres and convex  $\mathbb{R}\mathbb{P}^n$ -manifolds”. In: *Amer. J. Math.* 123.2 (2001), pp. 255–274. ISSN: 0002-9327,1080-6377. URL: [http://muse.jhu.edu/journals/american\\_journal\\_of\\_mathematics/v123/123.2loftin.pdf](http://muse.jhu.edu/journals/american_journal_of_mathematics/v123/123.2loftin.pdf).
- [LS20] Brice Loustau and Andrew Sanders. *Bi-Lagrangian structures and Teichmüller theory*. 2020. arXiv: 1708.09145 [id='math.DG'].
- [Mar] Arnaud Maret. *A note on character varieties*. <https://arnaudmaret.com/files/character-varieties.pdf>.
- [Nic07] Liviu I. Nicolaescu. *Lectures on the geometry of manifolds*. Second. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007, pp. xviii+589. ISBN: 978-981-277-862-8; 981-277-862-4. DOI: 10.1142/9789812770295. URL: <https://doi-org.clsproxy.library.caltech.edu/10.1142/9789812770295>.

- [NS94] K. Nomizu and T. Sasaki. *Affine Differential Geometry: Geometry of Affine Immersions*. Cambridge Tracts in Mathematics. Cambridge University Press, 1994. ISBN: 9780521441773. URL: <https://books.google.lu/books?id=1EUVHyjQANcC>.
- [RT24] Nicholas Rungi and Andrea Tamburelli. *Complex Lagrangian minimal surfaces, bi-complex Higgs bundles and  $SL(3, \mathbb{C})$ -quasi-Fuchsian representations*. 2024. arXiv: 2406.14945 [id='math.DG'].
- [Tre70] François Treves. “On local solvability of linear partial differential equations”. In: *Bulletin of the American Mathematical Society* 76.3 (1970), pp. 552–571.
- [Tsu09] Takashi Tsuboi. “On the group of real analytic diffeomorphisms”. In: *Ann. Sci. Éc. Norm. Supér. (4)* 42.4 (2009), pp. 601–651. ISSN: 0012-9593,1873-2151. DOI: 10.24033/asens.2104. URL: <https://doi.org/10.24033/asens.2104>.
- [Uhl83] Karen K. Uhlenbeck. “Closed minimal surfaces in hyperbolic 3-manifolds”. In: *Seminar on minimal submanifolds*. Vol. 103. Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ, 1983, pp. 147–168. ISBN: 0-691-08324-X; 0-691-08319-3.
- [Wan] Zuoqin Wang. *Lecture notes*. <http://staff.ustc.edu.cn/~wangzuoq/Courses/21F-Manifolds/Notes/Lec10.pdf>.
- [Wan91] Chang Ping Wang. “Some examples of complete hyperbolic affine 2-spheres in  $\mathbf{R}^3$ ”. In: *Global differential geometry and global analysis (Berlin, 1990)*. Vol. 1481. Lecture Notes in Math. Springer, Berlin, 1991, pp. 271–280. ISBN: 3-540-54728-2. DOI: 10.1007/BFb0083648. URL: <https://doi.org/10.1007/BFb0083648>.
- [Wen15] Richard A. Wentworth. *Higgs bundles and local systems on Riemann surfaces*. 2015. arXiv: 1402.4203 [math.DG].

CHRISTIAN EL EMAM: UNIVERSITY OF LUXEMBOURG, MAISON DU NOMBRE, 6 AVENUE DE LA FONTE, L-4364 ESCH-SUR-ALZETTE, LUXEMBOURG.

*Email address:* christian.elemam@uni.lu

NATHANIEL SAGMAN: UNIVERSITY OF LUXEMBOURG, MAISON DU NOMBRE, 6 AVENUE DE LA FONTE, L-4364 ESCH-SUR-ALZETTE, LUXEMBOURG.

*Email address:* nathaniel.sagman@uni.lu