BACKWARD BEHAVIOR AND DETERMINING FUNCTIONALS FOR CHEVRON PATTERN EQUATIONS

V. K. KALANTAROV¹, H. V. KALANTAROVA², AND O. VANTZOS³

ABSTRACT. The paper is devoted to the study of the backward behavior of solutions of the initial boundary value problem for the chevron pattern equations under homogeneous Dirichlet's boundary conditions. We prove that, as $t \to \infty$, the asymptotic behavior of solutions of the considered problem is completely determined by the dynamics of a finite set of functionals. Furthermore, we provide numerical evidence for the blow-up of certain solutions of the backward problem in finite time in 1D.

1. INTRODUCTION

We consider the system of equations

(1.1)
$$\tau \partial_t A = A + \Delta A - \phi^2 A - |A|^2 A - 2ic_1 \phi \partial_y A + i\beta A \partial_y \phi,$$

(1.2)
$$\partial_t \phi = -\mathbb{L}\phi - h\phi + \phi |A|^2 - c_2 \operatorname{Im} [A^* \partial_y A], \quad (x, y) \in \Omega, \quad t > 0,$$

where the linear operator \mathbb{L} is given by

$$\mathbb{L} := -D_1 \partial_x^2 - D_2 \partial_y^2,$$

and $\Omega \subset \mathbb{R}^2$ is a bounded domain with a sufficiently smooth boundary $\partial \Omega$. The system is complemented by the following boundary conditions

(1.3)
$$A\Big|_{\partial\Omega} = 0, \quad \phi\Big|_{\partial\Omega} = 0, \quad t > 0,$$

and the initial conditions

(1.4)
$$A\Big|_{t=0} = A_0, \quad \phi\Big|_{t=0} = \phi_0, \quad (x,y) \in \Omega.$$

In the context of this study,

(1.5)
$$D_1 > 0, \ D_2 > 0, \ c_1 \ge 0, \ c_2 \ge 0, \ h \ge 0, \ \beta \in \mathbb{R}$$

are given parameters, the complex valued function A, where A^* denotes its complex conjugate, and the real valued function ϕ are unknown functions. This coupled system of equations was proposed by Rossberg et al. [26], [27] to model the formation and evolution of patterns in nematic liquid crystals subject to convection driven by an electric field, sometimes accompanied with magnetic field. These works were motivated by a number of experimental studies [18], [25], [24]. A typical experimental setup used in these

Date: June 24, 2024.



FIGURE 1. Cross-section of a nematic liquid crystal contained between two parallel transparent plates. Nematic liquid crystals are composed of rod-like molecules, that are free to flow and orient themselves in 3D space, when external electric \vec{E} and magnetic fields \vec{H} are applied.

investigations consists of a nematic liquid crystal confined between two planar transparent glass plates, placed parallel at a distance of approximately $50\mu m$ from each other. Then AC-voltage of varying frequency is applied across the plates, sometimes in conjunction with a magnetic field parallel to the plates. The results of the described paradigmatic experiment, revealed that through use of either microscopes (with or without polarizers) or via laser diffraction various pattern formations are observable: Williams domain, Grid-Pattern, chevron patterns, etc.. The formation of these patterns is governed by experimentally easily controllable parameters, the amplitude and the frequency of the applied voltage and the magnitude of magnetic field. The advantage of liquid crystals over other fluids in these experiments besides its transparency and its phase transition temperatures being close to the room temperature, which makes detection of patterns easier, [3], is the fact that their molecules are locally oriented along desired direction. In [28], it is reported that the described system has a tendency to form chevron structures after a transition to defect chaos and that chevron patterns are generally observed when the isotropy of the system is spontaneously broken. this effect is achieved either by increasing the voltage or by applying a weak magnetic field.

The model (1.1)-(1.2) is obtained by coupling the real Ginzburg-Landau equation, where A represents the phase, direction and amplitude of the periodic pattern and equation (1.2), where ϕ represents the angle between director and the x-axis and also by using symmetries, smallness and regularity observations from experiments as assumptions. In this model the parameter τ is a function of the various physical timescales of the problem,

 β measures the interaction between the gradient of the director field and the phase of the rolls, the parameters D_1 and D_2 are the coefficients of the anisotropic diffusion of the director field for the liquid crystal particles, the dampening parameter h measures the tendency of the director field to align with the magnetic field H, corresponding to $\phi = 0$ and the parameters c_1 and c_2 control the torque that the director field and the wave vector of the rolls exert on each other.

In [27], Rossberg et al. generated chevron patterns in simulations of (1.1)-(1.2). For more numerical studies of the model refer to [29], [21].

The mathematical model (1.1)-(1.2) has also been studied qualitatively. In [19], the authors prove the following well-posedness result:

Theorem 1.1. ([19]) If $c_1 \in [0,1)$ or $c_1 \geq 2c_2$, h > 0, then the initial boundary value problem (1.1)-(1.4), where $A_0, \phi_0 \in L^2(\Omega)$, has a unique weak solution

(1.6)
$$A, \phi \in C([0,T]; L^2(\Omega)); H^1_0(\Omega)), \quad \forall T > 0,$$

such that

(1.7)
$$\|\phi(t)\| \le M_0, \quad \|A(t)\| \le M_0, \quad \forall t > 0,$$

and

(1.8)
$$\int_0^T \|\nabla \phi(t)\|^2 dt \le M_T, \quad \int_0^T \|\nabla A(t)\|^2 dt \le M_T, \ \forall T > 0,$$

where M_0 denotes a generic constant depending only on $||A_0||$, $||\phi_0||$ and $|\Omega|$, and where M_T is a generic constant, which depends also on T, besides $||A_0||$, $||\phi_0||$ and $|\Omega|$. In other words this problem generates a continuous semigroup S(t), $t \ge 0$, in the phase space $V^0 := L^2(\Omega) \times L^2(\Omega)$. Moreover, this semigroup is bounded dissipative in the phase space V^0 .

In [19], the authors also proved that the problem (1.1)-(1.4) possesses a global attractor, i.e., there exists a compact, invariant set that attracts uniformly each bounded set of the phase space V^0 . Furthermore, it is proved that this semigroup has a global attractor that belongs to $V^1 := H_0^1(\Omega) \times H_0^1(\Omega)$. Later, this result is improved in [20], in which it is proved that under the same assumptions of the Theorem 1.1 the semigroup

 $S(t): V^0 \to V^0, t \ge 0$, generated by the problem (1.1)-(1.4) possesses an exponential attractor.

In [20], the authors also prove the global stabilization of the zero steady state of the system (1.1)-(1.2) by finitely many Fourier modes:

Theorem 1.2. The solution $[A(t), \phi(t)]$ of the problem

(1.9)
$$\tau \partial_t A = A + \Delta A - \phi^2 A - |A|^2 A - 2ic_1 \phi \partial_y A + i\beta A \partial_y \phi,$$
$$-\mu \sum_{k=1}^{N} (A, w_k) w_k$$

(1.10)
$$\partial_t \phi = -\mathbb{L}\phi - h\phi + \phi |A|^2 - c_2 Im [A^* \partial_y A],$$

(1.11)
$$A|_{\partial\Omega} = \phi|_{\partial\Omega} = 0, \quad A|_{t=0} = A_0, \quad \phi|_{t=0} = \phi_0,$$

tends to zero stationary state with an exponential rate in V^1 , if

$$\mu \ge 1 \quad and \quad \lambda_{N+1}^{-1} < \delta_0,$$

k=1

where $0 < \lambda_1 \leq \lambda_2 \leq \ldots \leq \lambda_{N+1} \leq \ldots$ are the eigenvalues of the Laplace operator under the homogeneous Dirichlet boundary condition,

$$\delta_0 := 2(1-c_1)/(2+c_2)$$
 and $c_1 < 1$.

In this work, our primary objective is to demonstrate that the long time behavior of solutions to the initial boundary value problem for the system of equations (1.1)-(1.2) can be uniquely determined by finite number of determining functionals. Furthermore, we study the backward in time behavior of solutions to the system (1.1)-(1.2).

After covering some preliminaries in section 2, we devote section 3 of the paper to the backward in time behavior of solutions to the system (1.1)-(1.2). There are numerous papers dedicated to the backward behavior of solutions to nonlinear evolution equations (see, e.g. Bardos and Tartar [2], Dascaliuc [12], Kukavica and Malcok [22], Guo and Titi [15], Vukadinovic [31] and references therein). In this section, we establish the backward uniqueness of the problem (1.1)-(1.4) and show that some solutions of the backward problem blow up in a finite time. We follow this with further numerical evidence of finite-time blow up in section 4, using a convex-concave splitting scheme with explicit tracking of blow up times during each discrete time step.

Section 5 of the paper is dedicated to studying the existence of determining functionals for the system of chevron pattern equations (1.1)-(1.2). The intensive study over the last decades of finite dimensional asymptotic behavior of dissipative dynamical systems generated by nonlinear PDEs has been inspired by the pioneering works of Foias and Prodi [13] and Ladyzhenskaya [23]. They proved that asymptotic behavior of solutions to the initial boundary value problem for 2D Navier-Stokes equations is completely determined by asymptotic behavior of first N Fourier modes (for N large enough). For subsequent investigations of the finite-dimensional behavior of dynamical systems generated by the initial boundary value problem for dissipative PDEs (existence of finite-dimensional global attractors, inertial manifolds, exponential attractors, finite number of modes, nodes and local volume elements determining asymptotic behavior of solutions) we refer to [1], [14], [16], [30] and the references therein. The concept of determining functionals (determining interpolant operators) is introduced in [10], [11]. This concept provided a unified approach for investigating parameters that uniquely determine the asymptotic behavior of solutions to dissipative PDEs. A further extension of this unified approach for studying the long-time behavior of various nonlinear dissipative PDEs was developed in subsequent investigations of the long-time behaviour of solutions to initial boundary value problems for Navier-Stokes equations, systems of nonlinear parabolic equations, weakly damped nonlinear wave equations, nonlinear wave equation with nonlinear damping, von Karman equations and some other dissipative systems (see, e.g., Chueshov [6], [5], [7], Chueshov and Kalantarov [9], Chueshov and Lasiecka [8], Holst and Titi [17] and references therein.)

The main result of this section is Theorem 5.1, which establishes the existence of a finite number of functionals that uniquely determine the asymptotic behavior of solutions to the problem (1.1)-(1.4).

2. Preliminaries

In this section, we introduce the notation that will be used in what follows and recapitulate some definitions and known results.

- C will denote a generic positive constant,
- $(\cdot, \cdot) := (\cdot, \cdot)_{L^2(\Omega)}, \|\cdot\| := \|\cdot\|_{L^2(\Omega)},$
- $V^0 := L^2(\Omega) \times L^2(\Omega)$ is the Hilbert space with the inner product

 $\langle [u_1, v_1], [u_2, v_2] \rangle_{V^0} := (u_1, u_2) + (v_1, v_2),$

• $V^1 := H^1_0 \times H^1_0$ denotes the Hilbert space with the inner product

$$\langle [u_1, v_1], [u_2, v_2] \rangle_{V^1} := (\nabla u_1, \nabla u_2) + (\nabla v_1, \nabla v_2).$$

Definition 2.1. ([4]) Let V and H be reflexive Banach spaces and V be continuously and densely embedded into H. The *completeness defect* of a set $\mathcal{L} = \{\ell_j : j = 1, ..., N\}$ of linear functionals on V with respect to H is the following number

$$\epsilon_{\mathcal{L}}(V,H) := \sup\{ \|w\|_H : w \in V, \, \ell_j(w) = 0, \, \ell_j \in \mathcal{L}, \, \|w\|_V \le 1 \}.$$

 $\epsilon_{\mathcal{L}}(V, H) = 0$ if and only if the class of functionals \mathcal{L} is complete in V, i.e., the condition $\ell_j(w) = 0$ for all $\ell_j \in \mathcal{L}$ implies w = 0.

Theorem 2.2. ([4]) Let $\epsilon_{\mathcal{L}} = \epsilon_{\mathcal{L}}(V, H)$ be the completeness defect of a set $\mathcal{L} = \{\ell_j : j = 1, ..., N\}$ of linear functionals on V with respect to H. Then there exists a constant $C_{\mathcal{L}}$ such that

$$\|w\|_{H} \le C_{\mathcal{L}} \eta_{\mathcal{L}}(w) + \epsilon_{\mathcal{L}} \|w\|_{V} \quad for \ any \quad w \in V,$$

where $\eta_{\mathcal{L}}(w) = \max\{|\ell_j(w)| : j = 1, ..., N\}.$

Lemma 2.3. (see, e.g., [30]) Let g, h and y be three positive locally integrable functions on the interval (t_0, ∞) , such that y' (or equivalently $\frac{dy}{dt}$) is locally

integrable on the same interval. Suppose that the following inequalities are satisfied

$$y'(t) \leq g(t)y(t) + h(t), \quad t \in [t_0, \infty),$$

$$\int_t^{t+r} g(s)ds \leq a_1, \quad \int_t^{t+r} h(s)ds \leq a_2, \quad \int_t^{t+r} y(s)ds \leq a_3, \quad \forall t \in [t_0, \infty),$$
here r as an are moniton constants

where r, a_1, a_2, a_3 are positive constants. Then

$$y(t+r) \le (\frac{a_3}{r} + a_2)e^{a_1}, \ t \in [t_0, \infty).$$

Theorem 2.4. (see, e.g. [30]) Suppose that \mathbf{A} is a positive definite, unbounded, self adjoint linear operator in H with domain $V^2 := D(\mathbf{A})$ and let the nonlinear operator $\mathbf{B}(\cdot) : (0,T) \times V^1 \to H$, with $V^1 := D(\mathbf{A}^{\frac{1}{2}})$ satisfy the condition

(2.1)
$$\|\mathbf{B}(\mathbf{u}(t))\|_{H} \le k(t) \|\mathbf{u}(t)\|_{V^{1}}, \quad k(t) \in L^{2}(0,T).$$

If

(2.2)
$$\mathbf{u} \in L^{\infty}(0,T;V^1) \cap L^2(0,T;V^2)$$

is a solution of the equation

(2.3)
$$\mathbf{u}'(t) + \mathbf{A}\mathbf{u}(t) = \mathbf{B}(\mathbf{u}(t)), \ t \in (0,T),$$

that satisfies the condition $\mathbf{u}(T) = 0$, then $\mathbf{u}(t) = 0$, $\forall t \in [0, T]$.

Definition 2.5. A set of continuous linear functionals $\mathcal{L} = \{l_j : j = 1, \ldots, N\}$ on V is said to be a set of asymptotically (V, H, W)-determining functionals for the problem (2.5) if for any two solutions $u, v \in W$ the condition

$$\lim_{t \to \infty} \int_t^{t+1} |l_j(u(\tau)) - l_j(v(\tau))|^2 d\tau = 0 \text{ for } j = 1, \dots, N,$$

implies that

$$\lim_{t \to \infty} \|u(t) - v(t)\|_{H} = 0.$$

Theorem 2.6. ([6], Theorem 4.1.) Suppose that

• A is a positive definite operator with a discrete spectrum in a Hilbert space H, the basis $\{w_k\}$ of which consists of eigenvectors of the operator A:

$$\mathbf{A}w_k = \lambda_k w_k, \quad (w_k, w_j) = \delta_{kj}, \ 0 < \lambda_1 \le \lambda_2 \le \cdots, \quad \lim_{k \to \infty} \lambda_k = \infty,$$

 the nonlinear operator B(·) : D(A^{1/2}) → D(A^{-γ}), 0 ≤ γ < 1/2 satisfies the condition

(2.4)
$$\|\mathbf{A}^{-\gamma}(\mathbf{B}(\mathbf{u}_1) - \mathbf{B}(\mathbf{u}_2))\| \le M(\rho) \|\mathbf{A}^{\frac{1}{2}}(\mathbf{u}_1 - \mathbf{u}_2)\|,$$

for all $\mathbf{u}_j \in \mathbf{D}(\mathbf{A}^{\frac{1}{2}})$ such that $\|\mathbf{A}^{\frac{1}{2}}\mathbf{u}_j\| \leq \rho, \ \rho > 0$, and $M(\rho) > 0$ is a continuous function on \mathbb{R}^+ ,

• the problem

(2.5)
$$\begin{cases} \mathbf{u}'(t) + \mathbf{A}\mathbf{u}(t) = \mathbf{B}(\mathbf{u}(t)), & t > 0, \\ \mathbf{u}(0) = \mathbf{u}_0, \end{cases}$$

has a unique solution in the class of functions $\mathcal{W} = C([0,\infty);H) \cap C([0,\infty);D(A^{\frac{1}{2}}))$ and it is point dissipative, i.e.,

$$\|\mathbf{A}^{\frac{1}{2}}\mathbf{u}(t)\| \le R, \quad \forall t \ge t_0(u), \quad \forall \mathbf{u}(t) \in \mathcal{W}.$$

Suppose also that $\mathcal{L} = \{\ell_j, j = 1, ..., N\}$ is a set of linear continuous functionals defined on $V^1 = D(\mathbf{A}^{\frac{1}{2}})$. Then \mathcal{L} is a set of asymptotically (V^1, H, \mathcal{W}) - determining functionals of the problem (2.5) whenever

$$\varepsilon_{\mathcal{L}}(V^1, H) < \varepsilon_0(\gamma, R) \equiv \sqrt{\frac{1+2\gamma}{1-2\gamma}} \left[(1+2\gamma)M(R) \right]^{\frac{1}{2\gamma-1}}.$$

Definition 2.7. We call a pair of functions $[A, \phi]$ a weak solution of the initial boundary value problem for the system of equations (1.1)-(1.2) under the homogeneous Dirichlet's boundary condition, if $[A, \phi] \in \mathcal{W} = C([0, \infty); V^0) \cap C((0, \infty); V^1)$ and the system of equations (1.1)-(1.2) is satisfied in the sense of distributions.

3. BACKWARD BEHAVIOR

In this section, we will prove the backward uniqueness of a solution to the problem (1.1)-(1.4) and demonstrate that some solutions of the backward problem blow up in finite time. For 1D chevron pattern equations, we will illustrate that solutions not belonging to the global attractor of the semigroup generated by the initial boundary value problem blow up in a finite time T > 0.

The backward uniqueness of solution to the problem follows from the corresponding result for an abstract differential equation in a Hilbert space H of the form (2.5).

Suppose that $[A, \phi]$ and $[\tilde{A}, \tilde{\phi}]$ are solutions of the system (1.1)-(1.2) corresponding to initial data $[A_0, \phi_0]$ and $[\tilde{A}_0, \tilde{\phi}_0]$ respectively, then $[a, \Phi] := [A - \tilde{A}, \phi - \tilde{\phi}]$ is a solution of the following system

(3.1)
$$\tau \partial_t a = a + \Delta a - \phi^2 a - \tilde{A}(\phi + \tilde{\phi})\Phi - |A|^2 a - (Aa^* + \tilde{A}^*a)\tilde{A} - 2ic_1[\Phi \partial_y A + \tilde{\phi} \partial_y a] + i\beta[a\partial_y \phi + \tilde{A}\partial_y \Phi],$$

(3.2)
$$\partial_t \Phi = -\mathbb{L}\Phi - h\Phi + \Phi |A|^2 + \tilde{\phi}(Aa^* + \tilde{A}^*a) - c_2 \mathrm{Im} \left[a^* \partial_y A + \tilde{A}^* \partial_y a\right].$$

We can write this system in the form (2.5) in a Hilbert space $V^0 = L^2(\Omega) \times L^2(\Omega)$ with $\mathbf{u} := \begin{bmatrix} a \\ \Phi \end{bmatrix}$,

(3.3)
$$\mathbf{A} := \begin{bmatrix} -\frac{1}{\tau}\Delta & 0\\ 0 & \mathbb{L} \end{bmatrix}, \quad \mathbf{B}(a,\Phi) := \begin{bmatrix} \frac{1}{\tau}f_1(a,\Phi)\\ f_2(a,\Phi) \end{bmatrix},$$

where

(3.4)
$$f_1(a,\Phi) := a - \phi^2 a - (\phi^2 - \tilde{\phi}^2)\tilde{A} - |A|^2 A + |\tilde{A}|^2 \tilde{A} - 2ic_1[\Phi \partial_y A + \tilde{\phi} \partial_y a] + i\beta [a \partial_y \phi + \tilde{A} \partial_y \Phi],$$

(3.5)
$$f_2(a, \Phi) := -h\Phi + \Phi |A|^2 + \tilde{\phi}(|A|^2 - |\tilde{A}|^2) - c_2 \mathrm{Im} \left[a^* \partial_y A + \tilde{A}^* \partial_y a \right]$$

$$V^{1} := D(\mathbf{A}^{\frac{1}{2}}) = H^{1}_{0}(\Omega) \times H^{1}_{0}(\Omega)$$

and

(3.6)
$$V^2 := D(\mathbf{A}) = H^2(\Omega) \cap H^1_0(\Omega) \times H^2(\Omega) \cap H^1_0(\Omega)$$

It is clear that the operator \mathbf{A} is a positive definite self-adjoint operator in $V^0 := L^2(\Omega) \times L^2(\Omega)$, with the domain $D(\mathbf{A})$ and the nonlinear term $\mathbf{B}(a, \Phi)$ defined in (3.3) satisfies condition (2.1). Thus the statement of the Theorem 2.4 holds true also for the system (1.1)-(1.2). In other words, solution of the initial boundary value problem for the backward chevron pattern equations, i.e., solution of the problem

(3.7)
$$-\frac{\partial A}{\partial t} = A + \Delta A - \phi^2 A - |A|^2 A - 2ic_1 \phi \frac{\partial A}{\partial y} + i\beta A \frac{\partial \phi}{\partial y},$$

(3.8)
$$-\frac{\partial\phi}{\partial t} = D_1 \frac{\partial^2\phi}{\partial x^2} + D_2 \frac{\partial^2\phi}{\partial y^2} - h\phi + \phi |A|^2 - c_2 \mathrm{Im} \left[A^* \frac{\partial A}{\partial y}\right],$$

(3.9)
$$A|_{\partial\Omega} = \phi|_{\partial\Omega} = 0, \ A|_{t=0} = A_0, \ \phi|_{t=0} = \phi_0$$

is unique.

Theorem 2.4 also guarantees backward uniqueness of 1D version of the system (1.1)-(1.2), i.e., uniqueness of a weak solution to the problem

(3.10) $\partial_t A + A + \partial_x^2 A - \phi^2 A = |A|^2 A, \quad x \in (0, L), \quad t > 0,$

(3.11)
$$\phi_t + D_1 \partial_x^2 \phi - h\phi + |A|^2 \phi = 0, \quad x \in (0, L), \quad t > 0,$$

(3.12)
$$A|_{x=0} = A|_{x=L} = \phi|_{x=0} = \phi|_{x=L} = 0,$$

(3.13) $A|_{t=0} = A_0, \quad \phi|_{t=0} = \phi_0.$

It is not difficult to see that some solutions of the initial boundary value problem (3.10)-(3.13) blow up in finite time.

In fact suppose that $[A, \phi]$ is a local solution of the problem corresponding to the initial data $[A_0, \phi_0]$. Multiplying the equation (3.10) by A in L^2 sense we get

$$\frac{1}{2}\frac{d}{dt}\|A(t)\|^2 = -\|A(t)\|^2 + \|\partial_x A(t)\|^2 + (\phi^2(t), |A(t)|^2) + \int_0^L |A(x,t)|^4 dx.$$

Thanks to the Cauchy-Schwarz inequality we have

$$\int_0^L |A(x,t)|^2 dx \le \sqrt{L} \left(\int_0^L |A(x,t)|^4 dx \right)^{\frac{1}{2}}.$$

Thus employing the last inequality and the Poincare inequality we obtain that the function $\Psi(t) = ||A(t)||^2$ satisfies the differential inequality

$$\Psi'(t) \ge -2\Psi(t) + 2\lambda_1\Psi(t) + 2L^{-1}\Psi^2(t).$$

It is clear that if $\Psi_0 := \Psi(0) = ||A_0||^2 > (1 - \lambda_1)L$, then the solution of this inequality tends to infinity in a finite time

$$T_0 \le \int_{\Psi_0}^{\infty} \frac{ds}{2(L^{-1}s^2 + (\lambda_1 - 1)s)},$$

i.e., the solution of the problem (3.10) blows up in a finite time.

4. NUMERICAL BLOW-UP OF THE BACKWARD SYSTEM

To illustrate numerically the blow-up of ||A|| in the 1D backward system (3.10), we consider the case where $\phi = 0$ and note immediately that the resulting equation $\partial_t A = -\partial_x^2 A + (|A|^2 - 1)A$ can not be integrated numerically in a straightforward manner because of the 'wrong' sign of the Laplacian $\partial_x^2 A$. It is a basic result in numerical analysis that a direct numerical integration of this equation would be dominated by spurious high frequency modes.

To progress, we notice that the equation, in its weak form, can be thought of as the maximizing L^2 -gradient flow

$$(\partial_t A, u) = \mathcal{F}'(A)(u), \quad u \in H^1_0([0, L]),$$

 $\mathcal{F}(A) = \frac{1}{2} \|\partial_x A\|^2 + \frac{1}{4} \||A|^2 - 1\|^2.$

Given that the main challenge with integrating this gradient flow numerically is the uncontrollable divergence of the gradient $\partial_x A$, we consider instead the maximizing gradient flow of the functional \mathcal{F} in the H_{ϵ}^1 -norm

$$||A||_{H^{1}_{\epsilon}}^{2} := ||A||^{2} + \epsilon ||\partial_{x}A||^{2}, \quad \epsilon > 0,$$

which can be written as:

$$(\partial_t A, u) + \epsilon(\partial_x \partial_t A, \partial_x u) = \mathcal{F}'(A)(u) \,.$$

This regularization controls the 'rate of ascent' of A in H^1 instead of L^2 , and so penalizes the rapid divergence of the gradient that accompanies the introduction of spurious oscillatory modes. In strong form, it yields the regularized PDE

(4.1)
$$\partial_t A - \epsilon \partial_x^2 \partial_t A = -\partial_x^2 A + (|A|^2 - 1)A.$$

Proceeding with the numerical integration of (4.1), we assume that $A(t^k) = A^k$ and apply *operator splitting* to do the integration in two steps: first integrate the non-linear part

(4.2)
$$\partial_t \hat{A} - (|\hat{A}|^2 - 1)\hat{A} = 0, \quad \hat{A}(t^k) = A^k,$$

to calculate $\hat{A}^{k+1} = \hat{A}(t^{k+1}), t^{k+1} = t^k + \tau$, and then the linear part

(4.3)
$$(id - \epsilon \partial_x^2) \partial_t A = -\partial_x^2 A, \quad A(t^k) = \hat{A}^{k+1},$$

to update A to the next time step.

The first step (4.2) can be integrated exactly:

$$\hat{A}(x,t) = \frac{A^k(x)}{\sqrt{|A^k(x)|^2 + (1 - |A^k(x)|^2)e^{2(t-t^k)}}}.$$

This blows up for $|A^k(x)| > 1$ at the time

$$t = t^k + \frac{1}{2} \log \left(\frac{|A^k(x)|^2}{|A^k(x)|^2 - 1} \right) \,.$$

It follows that

(4.4)
$$\hat{A}^{k+1} = \frac{A^k}{\sqrt{|A^k|^2 + (1 - |A^k|^2)e^{2\tau}}},$$

as long as the (global) time step τ is chosen so that

$$\tau < \tau_{blow} = \inf_{x} \frac{1}{2} \log \left(\frac{\max(1, |A^{k}(x)|^{2})}{\max(1, |A^{k}(x)|^{2}) - 1} \right) \,.$$

Having selected a suitable τ , in particular ensuring that $\tau < \epsilon$, the second step (4.3) can be discretized implicitly

$$(id - \epsilon \partial_x^2) \frac{A - \hat{A}^{k+1}}{\tau} = -\partial_x^2 A$$

$$\Rightarrow A^{k+1} = \left(id - (\epsilon - \tau)\partial_{xx}^2\right)^{-1} \left(id - \epsilon \partial_{xx}^2\right) \hat{A}^{k+1} + \frac{1}{2} \left(id - \epsilon \partial_{xx}^2\right) \hat{A}^{$$

The operator on the right-hand side is positive definite (for $\tau < \epsilon$) and is straightforward to calculate numerically with spectral methods (FFT). Moreover, all the eigenvalues of the operator have modulus smaller than 1, which reaffirms the control of high frequency modes by the regularization.

In figure 2, we show the results of applying the scheme to the initial condition $A_0(x) = 5 \sin^3(20\pi x/L) + 2 \sin^3(12\pi x/L) - \sin^3(4\pi x/L)$ with L = 10. We discretize the interval [0, L] with 1000 uniformly distributed points, and we track the evolution of A up to the time t_{blow} , when the adaptive time step $\tau \approx 0$ up to machine precision ($\sim 10^{-16}$). We repeat the numerical experiment with different values for the regularizing parameter





FIGURE 2. Evolution of an oscillating initial condition under the regularized PDE (4.1) with $\epsilon = 0.1$, via the operator splitting scheme (4.2)-(4.3). The evolution is tracked up to the time t_{blow} , when the adaptive time step $\tau \approx 0$ up to machine precision (~ 10⁻¹⁶).



FIGURE 3. Effect of varying the regularizing parameter ϵ . In the top plot, we see the norm ||A|| blowing up at progressively earlier times for smaller values of ϵ . In the bottom plot, we look at the log-log plot of blow-up time vs the parameter ϵ , which appears to be $t_{blow} \sim \epsilon^{0.21}$ as $\epsilon \to 0$.

5. Determining functionals

In this section, we will show that finitely many linear functionals defined on the phase space uniquely determine the asymptotic behavior of solutions. To this end, first we derive a uniform a priory estimates of solutions to the problem in the phase space $V^1 := H_0^1(\Omega) \times H_0^1(\Omega)$. Multiplying the equation (1.1) with $-\Delta A^*$ in $L^2(\Omega)$ and employing the

Multiplying the equation (1.1) with $-\Delta A^*$ in $L^2(\Omega)$ and employing the Cauchy-Schwarz inequality we get

(5.1)
$$\frac{\tau}{2} \frac{d}{dt} \|\nabla A\|^{2} + \|\Delta A\|^{2} + (\phi^{2}, |\nabla A|^{2}) + (|A|^{2}, |\nabla A|^{2}) \\ \leq \|\nabla A\|^{2} + 2|(A\partial_{x}A^{*}, \phi\partial_{x}\phi)| + 2|(A\partial_{y}A^{*}, \phi\partial_{y}\phi)| \\ + 2c_{1}\|\phi\partial_{y}A\|\|\Delta A\| + |\beta|\|A\partial_{y}\phi\|\|\Delta A\|$$

Utilizing the inequality

.

(5.2)
$$\|\nabla v\|^2 \le \|v\| \|\Delta v\|, \quad \forall v \in H^2(\Omega) \cap H^1_0(\Omega),$$

the Ladyzhenskaya inequality

(5.3)
$$\|v\|_{L^4(\Omega)}^2 \le \sqrt{2} \|v\| \|\nabla v\|, \quad \forall v \in H_0^1(\Omega),$$

the interpolation inequality,

(5.4)
$$\|\nabla v\|_{L^4(\Omega)} \le d_0 \|\Delta v\|^{\frac{1}{2}} \|\nabla v\|^{\frac{1}{2}}, \quad \forall v \in H^2(\Omega) \cap H^1_0(\Omega),$$

the Young's inequality

(5.5)
$$ab \leq \frac{\varepsilon}{p}a^p + \frac{1}{q\varepsilon^{\frac{1}{p-1}}}b^q$$
, for all $a, b, \varepsilon > 0$, with $q = p/(p-1), 1 ,$

we derive the following inequality from (5.1)

(5.6)
$$\begin{aligned} \frac{\tau}{2} \frac{d}{dt} \|\nabla A\|^2 + \|\Delta A\|^2 + (\phi^2, |\nabla A|^2) + (|A|^2, |\nabla A|^2) \\ &\leq 3 \|\Delta A\|^2 + \frac{1}{4\varepsilon} \|A\|^2 + 2|(A\partial_x A^*, \phi \partial_x \phi)| \\ &+ 2|(A\partial_y A^*, \phi \partial_y \phi)| + \frac{c_1^2}{\varepsilon} \|\phi \partial_y A\|^2 + \frac{\beta^2}{4\varepsilon} \|A\partial_y \phi\|^2. \end{aligned}$$

We obtain the following estimates for the terms on the right hand side of (5.6) by using the following inequality

(5.7)
$$\nu_0 \|\Delta\phi\|^2 \le \|\mathbb{L}\phi\|^2 \le \nu_1 \|\Delta\phi\|^2, \quad \nu_0 > 0, \quad \nu_1 > 0,$$

in addition to (5.2), (5.3), (5.4) and (5.5):

in addition to (5.2), (5.3), (5.4) and (5.5):

$$2|(A\partial_{x}A^{*},\phi\partial_{x}\phi)| + 2|(A\partial_{y}A^{*},\phi\partial_{y}\phi)| \leq 4||A||_{L^{4}(\Omega)}||\phi||_{L^{4}(\Omega)}||\nabla\phi||_{L^{4}(\Omega)}||\nabla A||_{L^{4}(\Omega)} \leq 8||A||^{\frac{1}{2}}||\nabla A||^{\frac{1}{2}}||\phi||^{\frac{1}{2}}||\nabla \phi||^{\frac{1}{2}}||\nabla \phi||^{\frac{1}{2}}||\Delta \phi||^{\frac{1}{2}}||\nabla A||^{\frac{1}{2}}||\Delta A||^{\frac{1}{2}} \leq \frac{\varepsilon}{4}||\Delta \phi||^{2} + \frac{\varepsilon}{4}||\Delta A||^{2} + C(\varepsilon)||A||^{2}||\phi||^{2}||\nabla A||^{4} + \frac{||\nabla \phi||^{4}}{4},$$

$$\begin{split} \frac{c_1^2}{\varepsilon} \|\phi \partial_y A\|^2 &\leq \frac{c_1^2}{\varepsilon} \|\phi\|_{L^4(\Omega)}^2 \|\nabla A\|_{L^4(\Omega)}^2 \\ &\leq \frac{c_1^2 d_0^2}{\varepsilon} \|\phi\|_{L^4(\Omega)}^2 \|\Delta A\| \|\nabla A\| \leq \frac{c_1^2 d_0^2}{\varepsilon} \|\phi\|_{L^4(\Omega)}^2 \|A\|^{1/2} \|\Delta A\|^{3/2} \\ &\leq \frac{3\varepsilon}{4} \|\Delta A\|^2 + \frac{c_1^8 d_0^8}{4\varepsilon^7} \|\phi\|_{L^4(\Omega)}^8 \|A\|^2 \\ &\leq \frac{3\varepsilon}{4} \|\Delta A\|^2 + C(\varepsilon, c_1, d_0) \|\phi\|^4 \|\nabla \phi\|^4 \|A\|^2 \end{split}$$

By following the same sequence of steps, we can similarly get

$$\begin{aligned} \frac{\beta^2}{4\varepsilon} \|A\partial_y\phi\|^2 &\leq \frac{\beta^2}{4\varepsilon} \|A\|_{L^4(\Omega)}^2 \|\nabla\phi\|_{L^4(\Omega)}^2 \\ &\leq \frac{3\varepsilon}{4} \|\Delta\phi\|^2 + C(\varepsilon,\beta,d_0) \|A\|^4 \|\nabla A\|^4 \|\phi\|^2. \end{aligned}$$

Thus we infer from (5.6) that

14

$$(5.8) \quad \frac{\tau}{2} \frac{d}{dt} \|\nabla A\|^{2} + (1 - 4\varepsilon) \|\Delta A\|^{2} + (\phi^{2}, |\nabla A|^{2}) + (|A|^{2}, |\nabla A|^{2}) \\ \leq \frac{\varepsilon}{4\nu_{0}} \|\mathbb{L}\phi\|^{2} + \frac{1}{4\varepsilon} \|A\|^{2} + C(\varepsilon) \|A\|^{2} \|\phi\|^{2} \|\nabla A\|^{4} + \frac{\|\nabla \phi\|^{4}}{4} \\ + C(\varepsilon, c_{1}, d_{0}) \|\phi\|^{4} \|\nabla \phi\|^{4} \|A\|^{2} + C(\varepsilon, \beta, d_{0}) \|A\|^{4} \|\nabla A\|^{4} \|\phi\|^{2}.$$

Next, we multiply the equation (1.2) in $L^2(\Omega)$ with $-\mathbb{L}\phi$:

(5.9)
$$\frac{1}{2} \frac{d}{dt} (\mathbb{L}\phi, \phi) + \|\mathbb{L}\phi\|^2 + h(\mathbb{L}\phi, \phi) \\ \leq (|\phi||A|^2, |\mathbb{L}\phi|) + |c_2|(|A||\partial_y A|, |\mathbb{L}\phi|)$$

Arguing similarly to the calculation of the estimate (5.8) and employing the interpolation inequality

$$||v||_{L^6(\Omega)} \le d_1 ||\nabla v||^{\frac{2}{3}} ||v||^{\frac{1}{3}},$$

we obtain:

$$\begin{aligned} (|\phi||A|^{2}, |\mathbb{L}\phi|) &\leq \varepsilon \|\mathbb{L}\phi\|^{2} + \frac{1}{4\varepsilon} \|\phi\|_{L^{6}}^{2} \|A\|_{L^{6}}^{4} \\ &\leq \varepsilon \|\mathbb{L}\phi\|^{2} + \frac{d_{1}^{6}}{4\varepsilon} \|\nabla\phi\|^{4/3} \|\phi\|^{2/3} \|\nabla A\|^{8/3} \|A\|^{4/3} \\ &\leq \varepsilon \|\mathbb{L}\phi\|^{2} + C(\varepsilon, d_{1})(\|\phi\|^{2} \|\nabla\phi\|^{4} + \|A\|^{2} \|\nabla A\|^{4}), \end{aligned}$$

$$\begin{aligned} |c_2|(|A||\partial_y A|, |\mathbb{L}\phi|) &\leq \varepsilon \|\mathbb{L}\phi\|^2 + \frac{c_2^2}{4\varepsilon}(|A|^2, |\partial_y A|^2) \\ &\leq \varepsilon \|\mathbb{L}\phi\|^2 + \frac{c_2^2}{4\varepsilon}\|A\|_{L^4(\Omega)}\|\partial_y A\|_{L^4(\Omega)}^2 \\ &\leq \varepsilon \|\mathbb{L}\phi\|^2 + \frac{\sqrt{2}c_2^2d_0^2}{4\varepsilon}\|A\|\|\nabla A\|^2\|\Delta A\| \\ &\leq \varepsilon (\|\mathbb{L}\phi\|^2 + \|\Delta A\|^2) + C(\varepsilon, c_2, d_0)\|A\|^2\|\nabla A\|^2. \end{aligned}$$

By using the above estimates, we infer from (5.9)

(5.10)
$$\frac{1}{2} \frac{d}{dt} (\mathbb{L}\phi, \phi) + (1 - 2\varepsilon) \|\mathbb{L}\phi\|^2 - \varepsilon \|\Delta A\|^2 + h(\mathbb{L}\phi, \phi) \\ \leq C(\varepsilon, d_1) (\|\phi\|^2 \|\nabla \phi\|^4 + \|A\|^2 \|\nabla A\|^4) + C(\varepsilon, c_2, d_0) \|A\|^2 \|\nabla A\|^4.$$

Finally adding (5.8) and (5.10) we get

$$(5.11) \quad \frac{d}{dt} \left[\frac{\tau}{2} \| \nabla A \|^{2} + \frac{1}{2} (\mathbb{L}\phi, \phi) \right] + (1 - 5\varepsilon) \| \Delta A \|^{2} \\ + (1 - 2\varepsilon - \frac{\varepsilon}{4\nu_{0}}) \| \mathbb{L}\phi \|^{2} + (\phi^{2}, |\nabla A|^{2}) + (|A|^{2}, |\nabla A|^{2}) \\ \leq \frac{1}{4\varepsilon} \| A \|^{2} + C(\varepsilon) \| A \|^{2} \| \phi \|^{2} \| \nabla A \|^{4} + \frac{\| \nabla \phi \|^{4}}{4} + C(\varepsilon, c_{1}, d_{0}) \| \phi \|^{4} \| \nabla \phi \|^{4} \| A \|^{2} \\ + C(\varepsilon, \beta, d_{0}) \| A \|^{4} \| \nabla A \|^{4} \| \phi \|^{2} + C(\varepsilon, d_{1}) (\| \phi \|^{2} \| \nabla \phi \|^{4} + \| A \|^{2} \| \nabla A \|^{4}) \\ + C(\varepsilon, c_{2}, d_{0}) \| A \|^{2} \| \nabla A \|^{4}$$

Since the semigroup generated by the problem is dissipative in V^0 there exist $T_0 > 0$ and $M_0 > 0$ such that

$$||A(t)||^2 + ||\phi(t)||^2 \le M_0, \quad \forall t \ge T_0.$$

Thus, by choosing $\varepsilon = \min\{\frac{1}{10}, \frac{2\nu_0}{8\nu_0+1}\}$ we obtain from (5.11) that

$$\frac{d}{dt}[\tau \|\nabla A\|^2 + (\mathbb{L}\phi,\phi)] + \|\Delta A\|^2 \|\mathbb{L}\phi\|^2
\leq \frac{1}{4\varepsilon} \|A\|^2 + C(\varepsilon, M_0, \beta, d_0, d_1)(\|\nabla A\|^4 + \|\nabla \phi\|^4).$$

Finally, by using the uniform Gronwall Lemma 2.3, we infer from the last inequality that there exists $R_0 > 0$, depending only on the parameters of the system, such that

(5.12)
$$\|\nabla A\|^2 + \|\nabla \phi\|^2 \le R_0, \quad \forall t \ge T_1$$

i.e., the semigroup generated by the problem is bounded dissipative in $V^1 = H_0^1(\Omega) \times H_0^1(\Omega)$.

It remains to show that the condition (2.4) is satisfied, i.e.,

(5.13)
$$||f_j(a,\Phi)||_{H^{-\gamma}} \le M(R_0)(||\nabla a|| + ||\nabla \Phi||), \quad j = 1,2,$$

with some $\gamma \in (0, 1)$.

In fact, employing the estimate (5.12), the following inequality (see, e.g., [1], Chp. 1)

$$\|v\|_{H^{-\delta}(\Omega)} \le d_0 \|v\|_{L^{2/(1+\delta)}(\Omega)}$$

which is valid for $\delta \in (0,1)$, and the fact that $H^1(\Omega)$ is continuously embedded into $L^{\frac{2}{\delta}}(\Omega)$ we can bound the last term on the right hand side of (3.4)

$$\begin{aligned} \|i\beta[\tilde{A}\partial_y\Phi]\|_{H^{-\delta}(\Omega)} &\leq d_0|\beta|\|\tilde{A}\partial_y\Phi\|_{L^{2/(1+\delta)}} \\ &\leq d_0|\beta|\|\tilde{A}\|_{L^{2/\delta}}\|\partial_y\Phi\| \leq Cd_0\|\nabla\tilde{A}\|\|\partial_y\Phi\| \leq C(R_0)\|\nabla\Phi\|. \end{aligned}$$

Similarly we can show that

$$\|i\beta[a\partial_y\phi]\|_{H^{-\delta}(\Omega)} \le C(R_0)\|\nabla a\|$$

Utilizing the same arguments we can get necessary bounds for all terms on the right-hand side of (3.4) and (3.5) and obtain the desired estimates (5.13). Thus thanks to the Theorem 2.6 we have

Theorem 5.1. The set \mathcal{L} of functionals on V^1 is asymptotically (V^1, V^0, \mathcal{W}) determining functionals for the problem (1.1)-(1.4) provided the defect $\varepsilon(V^1, V^0)$ is small enough.

Remark 5.2. Similar to the proof of the Theorem 5.1 we can prove that the set \mathcal{L} of functionals on V^1 is asymptotically (V^1, V^0, \mathcal{W}) – determining functionals for the problem 1D problem

(5.14)
$$\begin{cases} \tau \partial_t A = A + \partial_x^2 A - \phi^2 A - |A|^2 A, \ x \in (0, L), t > 0, \\ \partial_t \phi = \partial_x^2 \phi - h \phi + \phi |A|^2, \ x \in (0, L), t > 0, \\ A|_{x=0} = A|_{x=L} = \phi|_{x=0} = \phi|_{x=L} = 0, \\ A|_{t=0} = A_0, \ \phi|_{t=0} = \phi_0 \end{cases}$$

provided the defect $\varepsilon(V^1, \mathbf{H})$ is small enough. Let us note that dissipativity of the semigroup $S(t), t \ge 0$ generated by the problem (5.14), as well as existence of and exponential attractor for the semigroup $S(t), t \ge 0$ is established in [20].

References

- A. V. Babin and M. I. Vishik, Attractors of evolution equations, Studies in Mathematics and its Applications, vol. 25, North-Holland Publishing Co., Amsterdam, 1992, Translated and revised from the 1989 Russian original by Babin. MR 1156492
- [2] C. Bardos and L. Tartar, Sur l'unicité rétrograde des équations paraboliques et quelques questions voisines, Arch. Rational Mech. Anal. 50 (1973), 10–25. MR 338517
- [3] Á. Buka, T. Börzsönyi, N. Éber, and T. Tóth-Katona, *Patterns in the bulk and at the interface of liquid crystals*, Coherent Structures in Complex Systems (D. Reguera, J. M. Rubí, and L. L. Bonilla, eds.), Lecture Notes in Physics, vol. 567, Springer, Berlin, Heidelberg, 2001, p. 125.
- [4] A. Yu. Chebotarev, Finite-dimensional controllability of systems of Navier-Stokes type, Differ. Uravn. 46 (2010), no. 10, 1495–1503. MR 2798716
- [5] I. D. Chueshov, A remark on sets of defining elements for reaction-diffusion systems, Mat. Zametki 63 (1998), no. 5, 774–784. MR 1683651
- [6] _____, The theory of functionals that uniquely determine the asymptotic dynamics of infinite-dimensional dissipative systems, Uspekhi Mat. Nauk 53 (1998), no. 4(322), 77–124. MR 1668050
- [7] Igor Chueshov, Dynamics of quasi-stable dissipative systems, Universitext, Springer, Cham, 2015. MR 3408002
- [8] Igor Chueshov and Irena Lasiecka, Long-time behavior of second order evolution equations with nonlinear damping, Mem. Amer. Math. Soc. 195 (2008), no. 912, viii+183. MR 2438025
- [9] Igor D. Chueshov and Varga K. Kalantarov, Determining functionals for nonlinear damped wave equations, Mat. Fiz. Anal. Geom. 8 (2001), no. 2, 215–227. MR 1859881
- [10] B. Cockburn, D. A. Jones, and E. S. Titi, *Determining degrees of freedom for non-linear dissipative equations*, C. R. Acad. Sci. Paris Sér. I Math. **321** (1995), no. 5, 563–568. MR 1356554

- [11] _____, Estimating the number of asymptotic degrees of freedom for nonlinear dissipative systems, Math. Comp. 66 (1997), no. 219, 1073–1087. MR 1415799
- [12] R. Dascaliuc, On backward-time behavior of the solutions to the 2-d space periodic navier-stokes equations, Annales de l'Institut Henri Poincaré C, Analyse non linéaire, vol. 22, 2005, pp. 385–401.
- [13] C. Foiaş and G. Prodi, Sur le comportement global des solutions non-stationnaires des équations de Navier-Stokes en dimension 2, Rend. Sem. Mat. Univ. Padova 39 (1967), 1–34. MR 223716
- [14] C. Foias, O. Manley, R. Rosa, and R. Temam, Navier-Stokes equations and turbulence, Encyclopedia of Mathematics and its Applications, vol. 83, Cambridge University Press, Cambridge, 2001. MR 1855030
- [15] Y. Guo and E. S. Titi, On the backward behavior of some dissipative evolution equations, Phys. D 306 (2015), 34–47. MR 3367572
- [16] J. K. Hale and G. Raugel, Regularity, determining modes and Galerkin methods, J. Math. Pures Appl. (9) 82 (2003), no. 9, 1075–1136. MR 2012806
- [17] M. J. Holst and E. S. Titi, *Determining projections and functionals for weak solu*tions of the Navier-Stokes equations, Recent developments in optimization theory and nonlinear analysis (Jerusalem, 1995), Contemp. Math., vol. 204, Amer. Math. Soc., Providence, RI, 1997, p. 125.
- [18] S. Kai, K. Hayashi, and Y. Hidaka, Pattern forming instability in homeotropically aligned liquid crystals, The Journal of Physical Chemistry 100 (1996), no. 49, 19007– 19016.
- [19] H. Kalantarova, V. Kalantarov, and O. Vantzos, Global behavior of solutions to chevron pattern equations, J. Math. Phys. 61 (2020), no. 6, 061511, 13. MR 4112645
- [20] _____, Chevron pattern equations: exponential attractor and global stabilization, Vietnam J. Math. 49 (2021), no. 3, 901–918. MR 4298702
- [21] S. Komineas, H. Zhao, and L. Kramer, Modulated structures in electroconvection in nematic liquid crystals, Physical Review E 67 (2003), no. 3, 031701.
- [22] Igor Kukavica and Mehmet Malcok, Backward behavior of solutions of the Kuramoto-Sivashinsky equation, J. Math. Anal. Appl. 307 (2005), no. 2, 455–464. MR 2142437
- [23] O. A. Ladyzhenskaya, Finding minimal global attractors for the Navier-Stokes equations and other partial differential equations, Uspekhi Mat. Nauk 42 (1987), no. 6(258), 25–60, 246. MR 933994
- [24] S. Nasuno, O. Sasaki, S. Kai, and W. Zimmermann, Secondary instabilities in electroconvection in nematic liquid crystals, Physical Review A 46 (1992), no. 8, 4954.
- [25] H. Richter, A. Buka, and I. Rehberg, *Electrohydrodynamic convection in a homeotrop*ically aligned nematic sample, Physical Review E 51 (1995), no. 6, 5886.
- [26] A. G. Rossberg, The amplitude formalism for pattern forming systems with spontaneously broken isotropy and some applications, Ph.D. thesis, Universität Bayreuth, Fakultät für Mathematik, Physik und Informatik, 1998.
- [27] A. G. Rossberg, A. Hertrich, L. Kramer, and W. Pesch, Weakly nonlinear theory of pattern-forming systems with spontaneously broken isotropy, Physical review letters 76 (1996), no. 25, 4729.
- [28] A. G. Rossberg and L. Kramer, Pattern formation from defect chaos—a theory of chevrons, Physica D: Nonlinear Phenomena 115 (1998), no. 1-2, 19–28.
- [29] H. Sakaguchi and A. Matsuda, Chevron patterns and defect lattices in an anisotropic model for electroconvection, Phys. D 238 (2009), no. 1, 1–8. MR 2571961
- [30] R. Temam, Infinite-dimensional dynamical systems in mechanics and physics, 2 ed., Applied Mathematical Sciences, vol. 68, Springer-Verlag, New York, 1997.
- [31] J. Vukadinović, On the backwards behavior of the solutions of the 2D periodic viscous Camassa-Holm equations, J. Dynam. Differential Equations 14 (2002), no. 1, 37–62. MR 1878244

 $^1\mathrm{Department}$ of Mathematics, Koç University, Rumelifeneri Yolu, Sariyer, Istanbul, Turkey

²Springer-Verlag, Heidelberg, Germany

³VANTZOS RESEARCH SMPC, ATHENS, GREECE