

ON SMALL SYLOW NUMBERS OF FINITE GROUPS

XIAOFANG GAO, IGOR LIMA, AND RULIN SHEN

ABSTRACT. Let G be a finite group and $n_p(G)$ the number of Sylow p -subgroups of G . In this paper, we prove if $n_p(G) < p^2$ then almost all numbers $n_p(G)$ are a power of a prime.

1. INTRODUCTION

Let G be a finite group and p a prime. We denote by $n_p(G)$ the number of Sylow p -subgroups of G , which is called Sylow p -number of G (hereinafter referred to as Sylow number). The influence of the number of Sylow subgroups in finite groups on group structure is a very meaningful research topic. We called $n_p(G)$ a solvable Sylow number if its ℓ -part was congruent to 1 modulo p for any prime ℓ . In 1967, Hall [1] studied the number of Sylow subgroups in finite groups, and proved that solvable groups have solvable Sylow numbers, and 22 is never a Sylow 3-number and 21 a Sylow 5-number. In 1995, Zhang [2] proved that a finite group G is p -nilpotent if and only if p is prime to every Sylow number of G . It is easy to see that $n_p(H) \leq n_p(G)$ for $H \leq G$, however $n_p(H)$ does not divide $n_p(G)$ in general. In 2003, Navarro [3] proved that if G is p -solvable, then $n_p(H)$ divides $n_p(G)$ for every $H \leq G$. In 2016 [4], Li and Liu classified finite non-abelian simple group with only solvable Sylow numbers. We say that a group G satisfies $DivSyl(p)$ if $n_p(H)$ divides $n_p(G)$ for every $H \leq G$. In 2018, Guo and Vdovin [5] generalized the results of Navarro, and proved that G satisfies $DivSyl(p)$ provided every non-abelian composition factor of G satisfies $DivSyl(p)$. Recently, Wu [6] proved that almost all finite simple groups do not satisfy $DivSyl(p)$.

In this paper, we consider the numbers of Sylow subgroups $n_p(G) < p^2$, and prove that almost all numbers $n_p(G)$ are a power of a prime. Our main theorem is the following.

Theorem 1.1. *Let G be a finite group and p a prime. Then the number $n_p(G)$ of Sylow p -subgroups of G is less than p^2 if and only if $n_p(G)$ is a power of a prime being less than p^2 or one of the following cases:*

- (1) $n_{13}(G) = 12^2$,
- (2) $n_p(G) = 1 + p$, where p is odd and not Mersenne,
- (3) $n_p(G) = 1 + (p - 3)p/2$, where $p > 3$ is Fermat,
- (4) $n_p(G) = 1 + (p + 3)p/2$, where $p > 3$ is Mersenne.

Key words and phrases. Sylow Subgroup, Non-abelian simple subgroup, Maximal subgroup. Project supported by the NSF of China (Grant No. 12161035).

Theorem 1.2. *Let G be a finite group and p not a Mersenne prime that is greater than 3. Assume that the number $n_p(G)$ of Sylow p -subgroups of G is less than p^2 . Then G is p -solvable if and only if $n_p(G)$ is a power of a prime.*

2. PRELIMINARY RESULTS

In this section we introduce and prove some useful results.

Lemma 2.1 (Zsigmondy's Theorem [7]). *Let a and n be integers greater 1. Then there exists a prime divisor q of $a^n - 1$ such that q does not divide $a^j - 1$ for all $0 < j < n$, except exactly the following cases:*

- (1) $n = 2$, $a = 2^s - 1$, where $s \geq 2$.
- (2) $n = 6$, $a = 2$.

Note that the prime divisor q of $a^n - 1$ as described above called a primitive prime or Zsigmondy prime.

Lemma 2.2. *Let $p = 2^n + 1$ be a Fermat prime and q prime. If $p^2 - 3p + 2 = 2q^a$ where a is a positive integer, then $(n, p, q, a) = (1, 3, 1, 1)$.*

Proof. Since $p = 2^n + 1$, $p^2 - 3p + 2 = 2^{2n} - 2^n = 2q^a$, this implies that $2^{n-1}(2^n - 1) = q^a$. Note that $(2^{n-1}, 2^n - 1) = 1$ and q^a has only one prime divisor, so either $2^{n-1} = 1$ or $2^n - 1 = 1$. Thus $n = q = a = 1$ and $p = 3$. \square

Lemma 2.3. *Let p be an odd prime and $e = \min\{k \geq 1 \mid q^{2k} \equiv 1 \pmod{p}\}$. Then p is a primitive prime divisor of $q^e - 1$ or $q^{2e} - 1$.*

Proof. Since $p \mid (q^{2e} - 1)$, p divides $q^e + 1$ or $q^e - 1$. If $p \mid (q^e - 1)$, then e is odd by the minimality of e , this implies that p is a primitive prime divisor of $q^e - 1$. Now assume that p divides $q^e + 1$, then $q^e - 1$ is not divisible by p since $(q^e - 1, q^e + 1) = 2$. We claim that p is a primitive prime divisor of $q^{2e} - 1$. If this was not the case, then there would exist a minimal integer k with $k < 2e$ such that $p \mid (q^k - 1)$, this implies that $k \mid 2e$. By the minimality of e , k would be odd, implying $k \mid e$. Write $e = ks$ where $s > 1$ (since p does not divide $q^e - 1$). Then

$$q^e - 1 = q^{ks} - 1 = (q^k - 1)((q^k)^{s-1} + (q^k)^{s-2} + \cdots + q^k + 1).$$

Note that $p \mid (q^k - 1)$, so p would divide $q^e - 1$, a contradiction. \square

Lemma 2.4 (Theorem 2.1 of [1]). *Let G have a normal subgroup K and let P be a Sylow p -subgroup of G . Then*

$$n_p(G) = n_p(G/K)n_p(K)n_p(C)$$

where $C = N_{PK}(P \cap K)/P \cap K$.

Lemma 2.5 (Lemma 2.7 of [4]). *Let G be a finite group, H a normal subgroup of G , and p a prime. If p does not divide $|G/H|$ then $n_p(H/Z(H)) = n_p(G)$.*

Lemma 2.6. *Let S be a non-abelian simple group, let P be a Sylow p -subgroup of S . If $n_p(S) < p^2$, then $P \cong C_p$.*

Proof. Write $n := n_p(S)$. As $n \equiv 1 \pmod{p}$ we can say $n = 1 + rp$ with $r < p$. Note that $n = |S : N_S(P)|$, we consider the permutation representation of S on cosets of $N_S(P)$. Since S is simple, S is isomorphic to a subgroup of S_n . From $n < p^2$ we deduce that S_n has no element of order p^2 , and hence the exponent of Sylow p -subgroup of S_n is p . Observe that the p -part of $|S_n|$ is

$$|S_n|_p = p^{\lfloor \frac{n}{p} \rfloor + \lfloor \frac{n}{p^2} \rfloor + \lfloor \frac{n}{p^3} \rfloor + \dots} = p^{\lfloor \frac{1+rp}{p} \rfloor} = p^r,$$

so we can get that the Sylow p -subgroup of S_n is an elementary abelian p -group, and then $P \cong C_p^a$ for some integer a . Let P_1 be a Sylow p -subgroup of S distinct from P , then $N_P(P_1) = P \cap P_1$ because otherwise $N_P(P_1) \not\leq P_1$, P_1 would be a proper subgroup of $N_P(P_1)P_1$, which would be a p -group since $P_1 \trianglelefteq N_P(P_1)$, contradicting P_1 is a Sylow p -subgroup of S . So the number of conjugates of P_1 under P is $|P : P_1 \cap P| < n < p^2$, thus $|P : P_1 \cap P| = p$. By using [8, Brodkey Theorem 5.28] we have $P_1 \cap P = 1$ because S is simple. Thus $P \cong C_p$. \square

Lemma 2.7. *Let S be a simple group, and let $M \cong C_n.N$ be an extension of C_n by N , which is a maximal subgroup of S . If the Sylow p -subgroup P of S is contained in C_n , then $n_p(S) = |S : M|$.*

Proof. Since $P \leq C_n$, P is a characteristic subgroup of C_n . Note that $C_n \trianglelefteq M$, so $P \trianglelefteq M$. Implying that $N_M(P) = M$. On the other hand, $P \leq N_M(P) \leq N_S(P) < S$. By the maximality of M , we have $N_S(P) = M$. Thus $n_p(S) = |S : M|$. \square

3. THE NUMBER OF SYLOW SUBGROUPS OF SIMPLE GROUPS

In this section, we will discuss the number of Sylow p -subgroups P of finite simple groups and use the classification theorem for finite simple groups to prove Theorem 3.1. As usual, $\pi(S)$ denotes the set of prime divisors of $|S|$, by $\Phi_n(q)$ we denote the n th cyclotomic polynomial, by function ϕ denotes Euler's totient function.

Theorem 3.1. *Let S be a finite simple group and $n_p(S) = 1 + rp$ for some integer r . Then $n_p(S) < p^2$ if and only if one of the following holds:*

- (1) $r = 1$, $S \cong \text{PSL}_2(p)$.
- (2) $r = 11$, $S \cong \text{PSL}_3(3)$ and $p = 13$.
- (3) $r = \frac{p-3}{2}$, $S \cong \text{PSL}_2(p-1)$, where $p > 3$ is a Fermat prime.
- (4) $r = \frac{p+3}{2}$, $S \cong \text{PSL}_2(p+1)$ with $p > 3$ is a Mersenne prime.

We will analyse the different possibilities for S by using the classification of finite simple groups. Note that $n_p(S) < p^2$ implies that a Sylow p -subgroup of S is isomorphic to C_p by Lemma 2.6, thus the p -part of $|S|$ is p . We assume $p \in \pi(S)$ such that $n_p(S) < p^2$ and let P be a Sylow p -subgroup of S . We will divide the proof of Theorem 3.1 into several propositions.

Proposition 3.2. *If S is a sporadic simple group or $S \cong {}^2F_4(2)'$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Since S is a simple group, $N_S(P) < S$, it follows that there is a maximal subgroup M of S such that $N_S(P) \leq M < S$. Thus $|S : N_S(P)| \geq |S : M|$, it is not less than the degree of minimal permutation representation of simple group S . With the help of [10], $n_p(S) \geq p^2$, a contradiction. \square

Proposition 3.3. *Let $S \cong A_n$ with $n \geq 5$. Then $n_p(S) < p^2$ if and only if $S \cong A_5$ and $p = 5$.*

Proof. Suppose now that $S \cong A_n$ with $n \geq 5$. Obviously, $p \leq n < 2p$ since $P \cong C_p$, thus $p \neq 2$. Applying [11] we have

$$\begin{aligned} n_p(S) &= \frac{n!}{(n-p)!p(p-1)} = \frac{n(n-1)\cdots(n-p+2)(n-p+1)}{p(p-1)} \\ &\geq (n-2)(n-3)\cdots(n-(p-2))(n-(p-1)) \geq (p-2)!. \end{aligned}$$

Assume $p \geq 7$, then $(p-2)! > p^2$, this implies that $n_p(S) > p^2$, a contradiction. Thus $p = 5$. If $n = 5$ then $n_5(A_5) = 6 < 25$ and if $n \geq 6$ then

$$n_p(S) = \frac{n!}{20(n-5)!} = \frac{n(n-1)(n-2)(n-3)(n-4)}{20} \geq \frac{6!}{20}.$$

In the second case, $n_p(S) > 25 = p^2$. It remains to deal with the case $p = 3$, that is $n = 5$ because $3 \leq n < 6$, it is easy to check that $n_3(A_5) = 10 > 3^2$, a contradiction. \square

Proposition 3.4. *Let $S \cong \text{PSL}_n(q)$ with $q = r^f$. Then $n_p(S) < p^2$ if and only if the following hold:*

- (1) $S \cong \text{PSL}_2(p)$ and $n_p(S) = 1 + p$.
- (2) $S \cong \text{PSL}_3(3)$, $p = 13$ and $n_{13}(S) = 144$.
- (3) $S \cong \text{PSL}_2(p-1)$ and $n_p(S) = 1 + \frac{(p-3)p}{2}$ where $p > 3$ is a Fermat prime.
- (4) $S \cong \text{PSL}_2(p+1)$ and $n_p(S) = 1 + \frac{(p+3)p}{2}$ where $p > 3$ is a Mersenne prime.

Proof. Suppose that $S \cong \text{PSL}_n(q)$ with $q = r^f$, we have

$$|S| = \frac{q^{\frac{n(n-1)}{2}}(q-1)(q^2-1)(q^3-1)\cdots(q^n-1)}{(q-1)(n, q-1)}.$$

Assume $(p, 2r) \neq 1$. If $p = 2$ then $r = 2$ because if this was not the case, then $S \cong \text{PSL}_2(q)$ with q odd and P would not be isomorphic to C_2 . Therefore, $S \cong \text{PSL}_n(2^f)$. Observe that $P \cong C_2$, so $S \cong \text{PSL}_2(2)$, a contradiction. If $p = r$ is odd then $S \cong \text{PSL}_2(p)$, it is not difficult to get that $|N_S(P)| = p(p-1)/2$, thus $n_p(S) = 1 + p$.

Next, we assume that $(p, 2r) = 1$. Let p be a primitive prime divisor of $q^e - 1$, that is,

$$e = \min\{k \geq 1 \mid q^k \equiv 1 \pmod{p}\}.$$

If $e \geq 2$, then p does not divide $q-1$ and $n/2 < e \leq n$ because $P \cong C_p$. By Lemma 2.5, $n_p(\text{PSL}_n(q)) = n_p(\text{GL}_n(q))$ since $|\text{GL}_n(q) : \text{SL}_n(q)|$ is not divisible by p . Note that $\text{GL}_e(q) \leq \text{GL}_n(q)$, thus $n_p(\text{GL}_e(q)) \leq n_p(\text{GL}_n(q)) < p^2$. Now, we are only concerned with the number of Sylow p -subgroups of $\text{GL}_e(q)$. Let $G := \text{GL}_e(q)$ and $Q \in \text{Syl}_p(G)$. In the light of [12, Lemma 2.5], $|N_G(Q)| = e(q^e - 1)$, implying that

$$(1) \quad n_p(G) = \frac{q^{\frac{e(e-1)}{2}}(q-1)(q^2-1)\cdots(q^{e-1}-1)}{e}.$$

Since p is a primitive prime divisor of $q^e - 1$, $p \mid \Phi_e(q)$, it follows that

$$p^2 \leq (\Phi_e(q))^2 < (4q^{\phi(e)})^2$$

by [13, Lemma 2.1(f)].

If $e \geq 6$ then $q^{e-1} - 1 \geq e$ and $q^{\frac{e(e-1)}{2}} > 16q^{2(e-1)}$. Therefore

$$n_p(G) \geq q^{\frac{e(e-1)}{2}} > 16q^{2(e-1)} \geq (4q^{\phi(e)})^2 > p^2,$$

which is a contradiction. Thus $e \in \{5, 4, 3, 2\}$.

If $e = 5$, then by [13, Lemma 2.1(f)] and equality (1), we get

$$\begin{aligned} n_p(G) &= \frac{q^{10}(q-1)(q^2-1)(q^3-1)(q^4-1)}{5} \geq q^{10}(q-1)^4(q+1)^2(q^2+1) \\ &\geq q^{14} > 16q^8 = (4q^{\Phi(5)})^2 > (\Phi_5(q))^2 \geq p^2, \end{aligned}$$

a contradiction.

Assume $e = 4$. If $q \geq 3$, then by [13, Lemma 2.1(f)] and equality (1),

$$n_p(G) = \frac{q^6(q-1)(q^2-1)(q^3-1)}{4} \geq q^8 \geq (4q^{\phi(4)})^2 > p^2$$

and if $q = 2$ then $p = 5$, $n_p(G) = 336 > 5^2$. These cases are impossible.

We next assume that $e = 3$, the equality (1) implies that

$$n_p(G) = \frac{q^3(q-1)(q^2-1)}{3} \geq \frac{q^4(q-1)^2}{3}.$$

Note that $p^2 \leq (\Phi_3(q))^2 = (q^2 + q + 1)^2 \leq 9q^4$. Obviously, if $q \geq 7$, then $n_p(G) \geq q^4(q-1)^2/3 \geq 9q^4 \geq p^2$. Since p is a primitive prime divisor of $q^3 - 1$, we can get that if $q = 5, 4, 3$ or 2 then $p = 31, 7, 13$ or 7 and $n_p(G) = 4000, 960, 144$ or 8 , respectively. Therefore $(q, p) = (3, 13)$ or $(q, p) = (2, 7)$. Recall that $n/2 < 3 \leq n$, thus $n = 3, 4$ or 5 . Therefore we only need to check the number of Sylow 13-subgroups of $\text{PSL}_3(3)$, $\text{PSL}_4(3)$ and $\text{PSL}_5(3)$ and the number of Sylow 7-subgroups of $\text{PSL}_3(2)$, $\text{PSL}_4(2)$ or $\text{PSL}_5(2)$. By using [14] $n_p(S) > p^2$ except $n_7(\text{PSL}_3(2)) = 8$ and $n_{13}(\text{PSL}_3(3)) = 144 = 1 + 11 \cdot 13$.

Suppose $e = 2$, then $n_p(G) = \frac{q(q-1)}{2}$ and $n = 2$ or 3 . Since p is a primitive prime divisor of $q^2 - 1$, thus $p \mid q + 1$. We write $kp = q + 1$ for some integer k . Assume $k \geq 2$. If $q \geq 5$ then

$$n_p(G) = \frac{q(q-1)}{2} \geq \frac{(q+1)^2}{4} \geq p^2.$$

We are left to examine $q = 2, 3$ or 4 . In these cases, $(q, p) = (2, 3)$ or $(4, 5)$ because p divides $q + 1$ and $(p, 2r) = 1$. Therefore we only need to calculate $n_3(\text{PSL}_3(2))$, $n_5(\text{PSL}_2(4))$ and $n_5(\text{PSL}_3(4))$. Simple calculations by using GAP [14] show that $S \cong \text{PSL}_2(4)$ with $n_5(S) = 6$ because $n_5(\text{PSL}_3(4)) = 2016$, $n_3(\text{PSL}_3(2)) = 28$, and both of them are greater than p^2 . Now assume that $k = 1$, we have $p = q + 1$, thus $n_p(G) = \frac{(p-1)(p-2)}{2} < p^2$, so we cannot determine the relationship between p^2 and the number $n_p(S)$ of Sylow p -subgroups of S by considering $n_p(G)$. Therefore, we need to consider the simple group S . Note that p is an odd prime integer, so $q = 2^f$ and $p = 2^f + 1$ is Fermat. Moreover, $S \cong \text{PSL}_2(2^f)$ or $\text{PSL}_3(2^f)$. If $S \cong \text{PSL}_2(2^f)$, then S has a maximal subgroup M of type $D_{2(q+1)}$. Thus there is a Sylow p -subgroup P of S such that $P \leq M$. Obviously, $P \in \text{Syl}_p(M)$ and $P \trianglelefteq M$.

By the maximality of M , $M = N_M(P) = N_S(P)$. Therefore,

$$n_p(S) = \frac{|S|}{|M|} = \frac{q(q-1)}{2} = \frac{(p-1)(p-2)}{2} = 1 + rp,$$

where $r = \frac{p-3}{2}$. Let $S \cong \text{PSL}_3(2^f)$. In the light of [19, Lemma 1.2(1)], the order of maximal torus of S is $\frac{(q-1)^2}{(3, q-1)}$, $\frac{q^2-1}{(3, q-1)}$ or $\frac{q^2+q+1}{(3, q-1)}$, it implies that the Sylow p -subgroup P of S is contained in the maximal torus of order $\frac{q^2-1}{(3, q-1)}$. Thus, $\frac{q^2-1}{(3, q-1)}$ divides $|C_S(P)|$. By [19, Table 4], S does not have element of order $2p$, so that the order $|C_S(P)|$ of the centralizer of P in S is $\frac{q^2-1}{(3, q-1)}$. Note that $N_S(P)/C_S(P)$ is isomorphic to a subgroup of Weyl group S_3 . It is also isomorphic to a subgroup of $\text{Aut}(P)$, which is a cyclic group of order 2^f . Thus $N_P(S)/C_P(S) \cong C_2$ (Since S is simple, it is not a p -nilpotent group, so that $|N_S(P)| \neq |C_S(P)|$). Thus $|N_S(P)| = \frac{2(q^2-1)}{(3, q-1)}$. Therefore,

$$n_p(S) = \frac{|S|}{|N_S(P)|} = \frac{q^3(q^3-1)}{2} > q^2 + 2q + 1 = p^2,$$

a contradiction.

Now we consider $e = 1$, i.e., $p \mid q-1$. Since $P \cong C_p$, $n \leq 3$. Let $S \cong \text{PSL}_2(q)$. If $q \in \{5, 7, 9, 11\}$, then, by $(p, 2r) = 1$ and p divides $q-1$, we only need to consider $n_3(\text{PSL}_2(7))$ and $n_5(\text{PSL}_2(11))$. By applying [14], we have $n_3(\text{PSL}_2(7))$ and $n_5(\text{PSL}_2(11))$ are greater than p^2 . If $q \geq 13$ is odd, then S has a maximal subgroup M isomorphic to D_{q-1} . Clearly, S has a Sylow p -subgroup P that is contained in $C_{\frac{q-1}{2}}$. Moreover, $C_{\frac{q-1}{2}} \triangleleft M$. By Lemma 2.7, $n_p(S) = |S|/|M| = \frac{q(q+1)}{2}$. Recalled that $p \mid q-1$, we can say $q-1 = kp$ for some integer k . If $k \geq 2$ then $n_p(S) \geq \frac{(q-1)^2}{k^2} = p^2$. Hence, $p = q-1$. Note that p is prime and q is odd, so $p = 2$, contradicting the fact that $(p, 2r) = 1$. Suppose $q = 2^f \geq 4$, then $M := D_{2(q-1)}$ is a maximal subgroup of S . Similarly, $N_S(P) = M$ and $n_p(S) = \frac{q(q+1)}{2}$. Since $n_p(S) < p^2$ and $p \mid (q-1)$, $p = q-1 = 2^f - 1$ is a Mersenne prime number. Therefore,

$$n_p(S) = \frac{(p+1)(p+2)}{2} = 1 + rp,$$

where $r = \frac{p+3}{2}$. Let $S \cong \text{PSL}_3(q)$. Observe that $\text{PSL}_2(q) \leq \text{PSL}_3(q)$, thus we only need to consider $\text{PSL}_3(2^f)$ where $2^f = p+1 \geq 4$ because otherwise $n_p(\text{PSL}_3(q)) \geq n_p(\text{PSL}_2(q)) \geq p^2$ by above discussion. We claim that $\text{PSL}_3(2^f) = \text{SL}_3(2^f)$. If this was not the case, then $Z(\text{SL}_3(2^f))$ would be isomorphic to $C_{(3, 2^f-1)} = C_3$, thus $p = 3$ and $S \cong \text{PSL}_3(4)$, by using GAP [14] again, $n_3(\text{PSL}_3(4)) = 280 > 9$, a contradiction. By [15, Table 8.3] S has a maximal subgroup M of type $(q-1)^2 : S_3 = p^2 : S_3$. Contradicting the fact that $P \cong C_p$. \square

Proposition 3.5. *Let $S \cong \text{PSU}_n(q)$ where $q = r^f \geq 2$ and $n \geq 3$. Then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Suppose now that $S \cong \text{PSU}_n(q)$. Then

$$|S| = \frac{q^{\frac{n(n-1)}{2}}(q+1)(q^2-1)(q^3+1) \cdots (q^n - (-1)^n)}{(n, q+1)(q+1)}.$$

Obviously, p is coprime to $2r$ since otherwise $n = 2$, contradicting $n \geq 3$.

If $n \geq 4$, then by Lemma 2.6, p does not divide $q + 1$. Let

$$e = \min\{k \geq 1 \mid (-q)^k \equiv 1 \pmod{p}\}.$$

Since $P \cong C_p$, $n/2 < e \leq n$. Lemma 2.5 implies that $n_p(\text{PSU}_n(q)) = n_p(\text{GU}_n(q)) < p^2$ because $\text{SU}_n(q) \trianglelefteq \text{GU}_n(q)$ and $|\text{GU}_n(q) : \text{SU}_n(q)|$ is not divisible by p . In the light of [16, Theorem 3.9], $\text{GU}_e(q) \leq \text{GU}_n(q)$, so $n_p(\text{GU}_e(q)) \leq n_p(\text{GU}_n(q)) < p^2$. In order to prove our result, it is enough to prove that $n_p(\text{GU}_e(q)) \geq p^2$. For the convenience, let $G := \text{GU}_e(q)$ and let Q denote the Sylow p -subgroup of G . Using [18, Theorem 3] we can obtain

$$|N_G(Q)| = |C_{(q^e - (-1)^e)_p} \cdot |C_{(q^e - (-1)^e)_{p'}} \cdot |C_e| = e \cdot (q^e - (-1)^e).$$

Therefore,

$$(2) \quad n_p(G) = \frac{q^{\frac{e(e-1)}{2}}(q+1)(q^2-1) \cdots (q^{e-1} - (-1)^{e-1})}{e}.$$

Clearly, $q^{e-1} + 1 \geq q^{e-1} - 1 \geq e$ since $e \geq 3$. By the above formula (2), we have

$$(3) \quad n_p(G) \geq q^{\frac{e(e-1)}{2}}(q+1)(q^2-1) \cdots (q^{e-2} - (-1)^{e-2}).$$

If e is even then p divides $q^e - 1$ and if e is odd then p divides $q^e + 1$. Thus, the order of q in the multiplicative group $(\mathbb{Z}/p\mathbb{Z})^*$ is either e or $e/2$ when e is even, and it is at most $2e$ when e is odd by Lemma 2.3. Implying that the order of q in this group is at most $2e$. By the properties of Euler's totient function and [13, Lemma 2.1(f)], we have

$$p^2 \leq (\Phi_{2e}(q))^2 < (4q^{\phi(2e)})^2 \leq 16q^{4e-2}.$$

We will examine $n_p(G)$ by discussing e . Assume $e \geq 9$, we have

$$n_p(G) \geq q^{\frac{e(e-1)}{2}} \geq 16q^{4(e-1)} \geq p^2,$$

a contradiction. If $e = 8$ or 7 , then by the equality (3), it is easy to check that $n_p(G) \geq 16q^{4e-2}$, these cases are impossible. Let $e = 6$, so that p is a primitive prime divisor of $q^6 - 1$ or $q^3 - 1$, this follows that $p^2 \leq (4q^{\phi(3)})^2 = (4q^{\phi(6)})^2 = 16q^4$ with the help of [13, Lemma 2.1(f)]. On the other hand,

$$n_p(G) \geq q^{15}(q+1)(q^2-1)(q^3+1)(q^4-1) > q^{15} > 16q^4 > p^2,$$

by the equality (2), contradicting $n_p(G) < p^2$. Assume $e = 5$, the formula (2) implies that

$$n_p(G) = \frac{q^{10}(q+1)(q^2-1)(q^3+1)(q^4-1)}{5} \geq q^{16}(q-1)^2(q^2-q+1).$$

Recall that $p^2 \leq (\Phi_{10}(q))^2 < (4q^{\phi(10)})^2 = 16q^8$ by [13, Lemma 2.1(f)]. Thus, $n_p(G) \geq q^{16}(q-1)^2(q^2-q+1) > 16q^8 > p^2$. Assume now that $e = 4$, in other words, p is a primitive prime divisor of $q^4 - 1$ or $q^2 - 1$, so that $p^2 \leq (4q^{\phi(2)})^2 \leq (4q^{\phi(4)})^2 = 16q^4$ by using [13, Lemma 2.1(f)] again. In the light of the formula (2),

$$n_p(G) = \frac{q^6(q+1)(q^2-1)(q^3+1)}{4} \geq q^6(q^3+1)(q-1) > 16q^4 > p^2,$$

which yields a contradiction. Finally, assume $e = 3$, then $n_p(G) = \frac{q^3(q+1)(q^2-1)}{3}$ by the formula (2). Since $p^2 \leq (\Phi_6(q))^2 = (q^2 - q + 1)^2 \leq 3q^4$, we have

$$n_p(G) = \frac{q^3(q+1)(q^2-1)}{3} \geq 3q^4 \geq p^2$$

if $q \geq 4$. Finally, if $q = 3$ then $p = 7$ by the definition of e and $(p, 2r) = 1$. Moreover, $n_7(G) = 288$. If $q = 2$ then $p = 3$ and $n_p(G) = 24$. Both of them are greater than p^2 , we deduce a contradiction.

Next, we suppose that $n = 3$, that is, $S \cong \text{PSU}_3(q)$ where $q > 2$. In the light of [6, Lemma 2.2], $n_p(S) = n_p(\text{SU}_3(q)) < p^2$. Write $H := \text{SU}_3(q)$ and Q the Sylow p -subgroup of H , we have

$$|H| = q^3(q-1)(q+1)^2(q^2-q+1).$$

We claim that $p \nmid q+1$. If this was not the case, then $p = 3$ because the Sylow p -subgroup of S is isomorphic to C_p , since H has a maximal subgroup of type $(q+1)^2 : S_3$ (see [15, Table 8.5]), this implies that 27 would be a divisor of $|H|$, and hence $3^2 \mid |S|$, contradicting $P \cong C_p$. Assume $p \mid q^2 - q + 1$, by [15, Table 8.5], H has a maximal subgroup M of type $(q^2 - q + 1) : 3$, thus H has a Sylow p -subgroup Q such that $Q \leq M$ because $(|H : M|, q^2 - q + 1) = 1$. Since $Q \trianglelefteq_c C_{q^2-q+1} \leq M$, Q is normal in M . It is obvious that Q is not normal in H since p does not divide $q+1$ and H has an unique normal subgroup $Z(H)$ with $Z(H) = (q+1, 3)$. Thus $M = N_H(Q) \geq N_M(Q) = M$ by the maximality of M , therefore

$$n_p(H) = |H : ((q^2 - q + 1) : 3)| = \frac{q^3(q-1)(q+1)^2}{3} \geq q^4(q-1).$$

Clearly, if $q \geq 4$ then $q^4(q-1) \geq 3q^4 \geq (q^2+1)^2 \geq (q^2-q+1)^2 \geq p^2$ and if $q = 3$ then $p = 7$, $n_7(H) = 288 > 49$, a contradiction. Finally, if $p \mid q-1$, then $p^2 \leq (q-1)^2 \leq 2q^2$. By applying [15, Table 8.5] again, H has a maximal subgroup M isomorphic to $[q^3] : (q^2-1)$. Note that Q is a Sylow p -subgroup of M because $|H : M|$ is not divisible by p . We claim that $N_M(Q) = C_{q^2-1}$. In fact, we assume that $N := N_M(Q) \cap [q^3]$ is a nontrivial r -group, thus $N \trianglelefteq N_M(Q)$ since $[q^3] \trianglelefteq M$. This implies that $Q \leq N_M(Q) \leq N_M(N)$. Observe that $(|N|, |Q|) = 1$, so $mn = nm$ for any $m \in Q, n \in N$, that is, H has an element with order rp , where p is a primitive prime divisor of $q-1$. Since p is an odd prime and $p \mid q-1, q-1 \neq 2^k$ for some integer k . By applying [19, Table 4], H does not have an element of order rp , a contradiction. Therefore, $N = 1$, and $N_M(Q) = C_{q^2-1}$. Consequently,

$$n_p(M) = |M : N_M(Q)| = q^3 \geq 2q^2 \geq p^2.$$

This implies that $n_p(H) \geq n_p(M) \geq p^2$, a contradiction. \square

Proposition 3.6. *If $S \cong \text{PSp}_{2n}(q)$ where $q = r^f \geq 2$ and $n \geq 2$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Assume $S \cong \text{PSp}_{2n}(q)$, where $n \geq 2$ and $q = r^f \geq 2$. Then

$$|S| = \frac{q^{n^2}(q^2-1)(q^4-1)\cdots(q^{2n}-1)}{(2, q-1)}.$$

Since $P \cong C_p, (p, 2r) = 1$. Set

$$e = \min\{k \geq 1 \mid q^{2k} \equiv 1 \pmod{p}\},$$

then p is a primitive prime divisor of $q^e - 1$ or $q^{2e} - 1$ by Lemma 2.3. Clearly, $n/2 < e \leq n$ (because $P \cong C_p$). By [16, Theorems 3.7 and 3.8], S has a subgroup isomorphic to $\text{Sp}_{2e}(q)$. Note that $n_p(\text{Sp}_{2e}(q)) \leq n_p(S) < p^2$, so we can prove our

proposition by considering $n_p(\mathrm{Sp}_{2e}(q))$. For convenience, let $G := \mathrm{Sp}_{2e}(q)$ and $Q \in \mathrm{Syl}_p(G)$, then

$$|G| = q^{e^2}(q^2 - 1)(q^4 - 1) \cdots (q^{2e} - 1).$$

We first suppose that p is a primitive prime divisor of $q^{2e} - 1$, Clearly, $e = 2e/(2e, 2)$. By [12, Theorem 2.6],

$$|N_G(Q)| = 2e(q^e + (-1)^{2e}) = 2e(q^e + 1),$$

so that

$$(4) \quad n_p(G) = \frac{q^{e^2}(q^2 - 1)(q^4 - 1) \cdots (q^{2e-2} - 1)(q^e - 1)}{2e}.$$

If $e \geq 4$, then $q^e - 1 \geq 2e$ and $q^{e^2}(q^2 - 1)(q^4 - 1)(q^6 - 1) \geq q^{e^2+3}(q^6 - 1) \geq 16q^{4e-2}$, thus

$$n_p(G) \geq q^{e^2}(q^2 - 1)(q^4 - 1)(q^6 - 1) \geq 16q^{4e-2}.$$

Recall that p is a primitive prime divisor of $q^{2e} - 1$,

$$p^2 \leq (\Phi_{2e}(q))^2 < (4q^{\phi(2e)})^2 \leq 16q^{4e-2}$$

by [13, Lemma 2.1(f)]. This follows that $n_p(G) \geq 16q^{4e-2} \geq p^2$, a contradiction. Assume $e = 3$ then

$$n_p(G) = \frac{q^9(q^2 - 1)(q^4 - 1)(q^3 - 1)}{6} \geq q^{12}.$$

[13, Lemma 2.1(f)] shows that $p^2 \leq (\Phi_6(q))^2 \leq (4q^2)^2 = 16q^4 \leq q^{12}$, thus $p^2 \leq n_p(G)$. We now assume $e = 2$ then

$$n_p(G) = \frac{q^4(q^2 - 1)^2}{4}$$

by the formula (4) and $p^2 \leq (\Phi_4(q))^2 \leq (4q^{\phi(4)})^2 = 16q^4$. It is not difficult to get that $n_p(G) \geq p^2$ if $q \geq 3$. We are left to examine the case $q = 2$, meaning that p is a primitive prime divisor of $2^4 - 1$, and thus $p = 5$ and $n_p(G) = 36$. Therefore, $n_p(G) > 25 = p^2$. Finally, if $e = 1$, then $n = 1$ since $n < 2e \leq 2n$, thus $S \cong \mathrm{PSp}_2(q) \cong \mathrm{PSL}_2(q)$, which has been considered in the case of linear groups.

We next suppose that p is a primitive prime divisor of $q^e - 1$, thus e is odd by the minimality of e . Observe that $e = e/(e, 2)$. Using [12, Theorem 2.6] again, we obtain

$$\begin{aligned} n_p(G) &= \frac{q^{e^2}(q^2 - 1)(q^4 - 1) \cdots (q^{2e-2} - 1)(q^{2e} - 1)}{2e(q^e - 1)} \\ &= \frac{q^{e^2}(q^2 - 1)(q^4 - 1) \cdots (q^{2e-2} - 1)(q^e + 1)}{2e}. \end{aligned}$$

If $e \geq 3$ then $q^e + 1 \geq q^e - 1 \geq 2e$, so that

$$n_p(G) \geq q^{e^2}(q^2 - 1)(q^4 - 1) \geq (q^2 + 1)(q + 1)^2 q^{e^2} \geq 16q^{2e-2}.$$

Obviously, $p^2 \leq (\Phi_e(q))^2 < (4q^{\phi(e)})^2 \leq 16q^{2e-2}$. Thus $n_p(G) \geq p^2$. Since e is odd, thus $e = 1$ and then $n = 1$. In this case $S \cong \mathrm{PSp}_2(q) \cong \mathrm{PSL}_2(q)$, which has been considered before. \square

Proposition 3.7. *If $S \cong \text{P}\Omega_{2n+1}(q)$ where $q = r^f$ is odd and $n \geq 2$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Suppose $S \cong \text{P}\Omega_{2n+1}(q)$ where $q = r^f$ is odd, then $|\text{P}\Omega_{2n+1}(q)| = |\text{PSp}_{2n}(q)|$. If $p = 2$, then $n = 1$ since $P \cong C_2$, a contradiction. If $p \neq 2$ then, by [20, Corollary 2.2], $|N_{\text{P}\Omega_{2n+1}(q)}(P_1)| = |N_{\text{PSp}_{2n}(q)}(P_2)|$ where P_1, P_2 are Sylow p -subgroups of $\text{P}\Omega_{2n+1}(q)$ and $\text{PSp}_{2n}(q)$, respectively. Thus $n_p(\text{P}\Omega_{2n+1}(q)) = n_p(\text{PSp}_{2n}(q)) \geq p^2$ by Proposition 3.6, a contradiction. \square

Proposition 3.8. *Assume $S \cong \text{P}\Omega_{2n}^+(q)$, where $q = r^f$ and $n \geq 3$. Then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong \text{P}\Omega_{2n}^+(q)$, then

$$|S| = \frac{q^{n(n-1)}(q^n - 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2n-2} - 1)}{(4, q^n - 1)}.$$

It is not difficult to check that $(p, 2r) = 1$ by Lemma 2.6. Since $\Omega_{2n}^+(q) \leq \text{GO}_{2n}^+(q)$ and p does not divide $|\text{GO}_{2n}^+(q) : \Omega_{2n}^+(q)|$, $n_p(\text{GO}_{2n}^+(q)) = n_p(S) < p^2$ with the help of Lemma 2.5. Let $G := \text{GO}_{2n}^+(q)$. Then

$$|G| = 2q^{n(n-1)}(q^n - 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2n-2} - 1).$$

Set

$$e = \min\{k \geq 1 \mid q^{2k} \equiv 1 \pmod{p}\}.$$

Since $|G : S|$ is not divisible by p , the Sylow p -subgroup Q of G is also a cyclic subgroup of order p by Lemma 2.6. Therefore, $n - 1 < 2e \leq 2n - 2$ and p is a primitive prime divisor of $q^e - 1$ or $q^{2e} - 1$ by Lemma 2.3.

Assume that p is a primitive prime divisor of $q^{2e} - 1$. In the light of [16, Theorem 3.12], G has a maximal subgroup of type $\text{GO}_{2k}^+(q) \times \text{GO}_{2m}^+(q)$, with $k + m = n$ and $0 < k < n$. If $e < n - 1$, then G has a subgroup H isomorphic to $\text{GO}_{2(e+1)}^+(q)$. Obviously, the Sylow p -subgroup of H is not trivial because

$$|H| = 2q^{(e+1)e}(q^{e+1} - 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2e} - 1)$$

and $p \mid q^{2e} - 1$. Note that p is a primitive prime divisor of $q^{2e} - 1$ and $n_p(G) \geq n_p(H) = n_p(\text{P}\Omega_{2(e+1)}^+(q))$. Let S_1 denote $\text{P}\Omega_{2(e+1)}^+(q)$ and Q_1 denote its Sylow p -subgroup. In the light of [20, Lemma 2.4] we have

$$|N_{S_1}(Q_1)| = 2e(q^e + 1)(q + 1)/(q^{e+1} - 1, 4),$$

so that

$$(5) \quad n_p(S_1) = \frac{q^{(e+1)e}(q^{e+1} - 1)(q - 1)(q^4 - 1)(q^6 - 1) \cdots (q^{2e-2} - 1)(q^e - 1)}{2e}.$$

Recall that $p^2 \leq (\Phi_{2e}(q))^2 < (4q^{\phi(2e)})^2 \leq 16q^{4e-2}$. Let $e \geq 4$, we have $q^{(e+1)e}(q^{e+1} - 1)(q^4 - 1) \geq (q^{e+1} - 1)q^{e^2+e+3} \geq 16q^{4e-2}$ and $q^e - 1 \geq 2e$. Thus

$$n_p(S_1) \geq q^{(e+1)e}(q^{e+1} - 1)(q^4 - 1) \geq 16q^{4e-2} \geq p^2,$$

therefore, $n_p(S) = n_p(G) \geq n_p(H) = n_p(S_1) > p^2$, we get a contradiction. Next if $e = 3$ then

$$n_p(S_1) = q^{12}(q^4 - 1)(q - 1)(q^4 - 1)(q^3 - 1)/6$$

and if $e = 2$ then

$$n_p(S_1) = q^6(q^3 - 1)(q - 1)(q^2 - 1)/4$$

by the formula (5). By $p^2 \leq (\Phi_{2e}(q))^2 < (4q^{\phi(2e)})^2$, it is easy to check that $n_p(S_1) \geq p^2$ unless $e = q = 2$, in which case $p = 5$ and $n_p(S_1) = 336 > 5^2$ (since p is a primitive prime divisor of $q^4 - 1$). Hence $n_p(G) \geq n_p(H) = n_p(S_1) \geq p^2$, a contradiction. Finally let $e = 1$, then from $n - 1 < 2e \leq 2n - 2$ deduce that $n = 2$, this contradicts the assumption. Next if $e = n - 1$, that is p is a primitive prime divisor of $q^{2(e-1)} - 1$. We can use [20, Lemma 2.4] again to calculate $n_p(S)$ directly. We have

$$|N_S(P)| = 2(q^{n-1} + 1)(q + 1)(n - 1)/(4, q^n - 1),$$

and hence

$$(6) \quad n_p(S) = \frac{q^{n(n-1)}(q^n - 1)(q - 1)(q^4 - 1) \cdots (q^{2n-4} - 1)(q^{n-1} - 1)}{2(n - 1)}.$$

If $n \geq 5$, then $q^{n-1} - 1 \geq 2(n - 1)$, so that

$$n_p(S) \geq q^{n(n-1)}(q^n - 1) \geq 16q^{4n-4} \geq (4q^{\phi(2n-2)})^2 > (\Phi_{2n-2}(q))^2 \geq p^2,$$

a contradiction. Similarly to the previous case, if $n = 4$ then

$$n_p(S) = q^{12}(q^4 - 1)(q - 1)(q^3 - 1)(q^4 - 1)/6$$

and if $n = 3$ then

$$n_p(S) = q^6(q^3 - 1)(q - 1)(q^2 - 1)/4$$

by the quality (6). Using [13, Lemma 2.1(f)], we have $p^2 < 16q^4$, this implies that $n_p(S) > p^2$ except for $(n, q) = (3, 2)$. Suppose $(n, q) = (3, 2)$, then p is a primitive prime divisor of $2^4 - 1$, implying that $p = 5$ and $n_p(S) = 336 > p^2$, a contradiction.

Next, we suppose that p is a primitive prime divisor of $q^e - 1$. Then, by the minimality of e , e is odd. Moreover, if p divides $q^n - 1$, then n must be odd because the Sylow p -subgroup of G is a cyclic group of order p and $n < 2n - 2$. Thus, in this case, p is a primitive prime divisor of $q^n - 1$. Therefore, $e \leq n$. [16, Theorem 3.12] implies that G admits a subgroup H of type $\text{GL}_n(q)$. Note that

$$|H| = q^{n(n-1)/2}(q - 1)(q^2 - 1)(q^3 - 1) \cdots (q^n - 1)$$

and $p \mid q^e - 1$, where $e \leq n$, thus the Sylow p -subgroup of G is contained in H . In particular,

$$n_p(G) \geq n_p(H) \geq n_p(\text{SL}_n(q)) = n_p(\text{PSL}_n(q)).$$

By Proposition 3.4, we can get $n_p(G) \geq p^2$ unless $(n, p, q) = (3, 13, 3)$ and $n = 2$, namely $G = \text{GO}_6^+(3)$ and $S = \text{P}\Omega_6^+(3) \cong \text{PSL}_4(3)$ (Since $n \geq 3$). In the light of Proposition 3.4, $n_{13}(S) \geq 13^2$, a contradiction. \square

Proposition 3.9. *Assume $S \cong \text{P}\Omega_{2n}^-(q)$, where $q = r^f$ and $n \geq 2$. Then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong \text{P}\Omega_{2n}^-(q)$. Then

$$|S| = \frac{q^{n(n-1)}(q^n + 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2n-2} - 1)}{(4, q^n + 1)}.$$

By Lemma 2.6, we can get that p is coprime to $2r$, that is p is an odd integer. Similarly as Proposition 3.8,

$$n_p(\mathrm{P}\Omega_{2n}^-(q)) = n_p(\Omega_{2n}^-(q)) = n_p(\mathrm{GO}_{2n}^-(q)) < p^2$$

because $|\mathrm{GO}_{2n}^-(q) : \Omega_{2n}^-(q)|$ is not divisible by p . Now we are concerned with studying $n_p(\mathrm{GO}_{2n}^-(q))$. Let $G := \mathrm{GO}_{2n}^-(q)$, then

$$|G| = 2q^{n(n-1)}(q^n + 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2n-2} - 1).$$

In particular, the Sylow p -subgroup Q of G is also isomorphic to C_p because p does not divide $|\mathrm{GO}_{2n}^-(q) : S|$. Set

$$e = \min\{k \geq 1 \mid q^{2k} \equiv 1 \pmod{p}\},$$

Thus $n - 1 < 2e \leq 2n - 2$ by Lemma 2.6, and p is a primitive prime divisor of $q^e - 1$ or $q^{2e} - 1$ with the help of Lemma 2.3. Let $e = 1$, then, $n = 2$ since $n - 1 < 2e \leq 2n - 2$. Thus, $G \cong \mathrm{GO}_4^-(q)$ and $S \cong \mathrm{P}\Omega_4^-(q) \cong \mathrm{PSL}_2(q^2)$. Moreover, $p \mid (q^2 - 1)$. By Proposition 3.4 we have $S \cong \mathrm{PSL}_2(p - 1)$ or $S \cong \mathrm{PSL}_2(p + 1)$. Thus $q^2 = p - 1$ or $q^2 = p + 1$. Since $q^2 - 1$ is divisible by p and $(p, 2r) = 1$, $p = q^2 - 1$, thus $q = 2$ and $p = 3$, and hence $S \cong \mathrm{P}\Omega_4^-(2) \cong \mathrm{PSL}_2(4)$. By Proposition 3.4 we have $n_3(S) > p^2$, a contradiction. In the rest of the proof we will assume that $e \geq 2$.

Assume p is a primitive prime divisor of $q^e - 1$, then $e \leq n - 1$ and e is odd by the minimality of e , so that $e \geq 3$. [16, Theorem 3.11] shows that G has a maximal subgroup of type $\mathrm{GO}_{2k}^+(q) \times \mathrm{GO}_{2m}^-(q)$, with $k + m = n$ and $0 < k < n$. Thus, we can consider the subgroup H of type $\mathrm{GO}_{2e}^+(q)$. Note that $p \mid (q^e - 1)$ and

$$|H| = 2q^{e(e-1)}(q^e - 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2e-2} - 1),$$

so the Sylow p -subgroup of G is contained in H . Moreover, $n_p(G) \geq n_p(H)$. By Proposition 3.8, $n_p(G) \geq n_p(H) = n_p(\mathrm{P}\Omega_{2e}^+(q)) \geq p^2$ since $e \geq 3$, a contradiction.

Now suppose that p is a primitive prime divisor of $q^{2e} - 1$. Then $e \leq n - 1$. Assume $e < n - 1$, we consider the subgroup H of type $\mathrm{GO}_{2(e+1)}^+(q)$. Note that $p \mid (q^{2e} - 1)$ and

$$|H| = 2q^{(e+1)e}(q^{e+1} - 1)(q^2 - 1)(q^4 - 1) \cdots (q^{2e} - 1).$$

Thus H contains the Sylow p -subgroup of G . In particular, $e + 1 \geq 3$. By using Proposition 3.8,

$$n_p(G) \geq n_p(H) = n_p(\mathrm{P}\Omega_{2(e+1)}^+(q)) \geq p^2$$

(Since p does not divide $|H : \Omega_{2(e+1)}^+(q)|$, we have $n_p(H) = n_p(\mathrm{P}\Omega_{2(e+1)}^+(q))$), a contradiction. Suppose that $e = n - 1$, that is, p is a primitive prime divisor of $q^{2n-2} - 1$, so that $p \mid (q^{n-1} + 1)$. Assume q is odd. By using [16, Theorem 3.11] G has a maximal subgroup M of type $[q^{2(n-1)}] \cdot (C_{q-1} \times \mathrm{GO}_{2(n-1)}^-(q))$. Since $Q \cong C_p$ and $p \nmid |M|$ (Recall that $Q \in \mathrm{Syl}_p(G)$), any Sylow p -subgroup of M is a Sylow p -subgroup of G . For convenience, we write

$$H := [q^{2(n-1)}], \quad K := C_{q-1} \times \mathrm{GO}_{2(n-1)}^-(q)$$

and $T \in \mathrm{Syl}_p(M)$. We claim that $H \cap N_M(T) = 1$. If this was not the case, then $N := H \cap N_M(T)$ would not be a trivial r -group, and $N \trianglelefteq N_M(T)$ since $H \trianglelefteq M$. We deduce that $T \leq N_M(T) \leq N_M(N)$. Thus the commutator of T and N would be trivial because $(|N|, |T|) = 1$ (Since $N^T = N$, $T^N = T$ and $(|N|, |T|) = 1$).

Therefore M would have an element of order rp , contradicting [19, Table 4], which shows that M does not have an element of order rp . Hence $H \cap N_M(T) = 1$ and $N_M(T) \leq K$. It follows that

$$n_p(M) = |M : N_M(T)| \geq |M : K| = q^{2(n-1)}.$$

Recall that $p \mid q^{n-1} + 1$, we write $kp = q^{n-1} + 1$ for some positive integer k . Since both p and q are odd numbers, $k \geq 2$. Hence

$$p^2 = \frac{q^{2n-2} + 2q^{n-1} + 1}{k^2} \leq \frac{3q^{2n-2}}{k^2} \leq q^{2(n-1)} \leq n_p(M).$$

Therefore, $n_p(G) \geq n_p(M) \geq p^2$, a contradiction. Assume q is even, i.e., $q = 2^f$. By [16, Theorem 3.11], G has a maximal subgroup of type $\mathrm{Sp}_{2(n-1)}(q)$. Obviously, $Q \leq M$ and

$$n_p(G) \geq n_p(\mathrm{Sp}_{2(n-1)}(q)) = n_p(\mathrm{PSp}_{2(n-1)}(q)) > p^2$$

unless $n = 2$ by Proposition 3.6. Finally, let $n = 2$, then $G \cong \mathrm{GO}_4^-(q)$ and $S \cong \mathrm{P}\Omega_4^-(q) \cong \mathrm{PSL}_2(q^2)$. By Proposition 3.4, $q^2 = p - 1$ or $q^2 = p + 1$, this implies that $p = q^2 + 1$ or $p = q^2 - 1$, contradicting the fact that $p \mid (q + 1)$.

Finally, we suppose that $p \mid q^n + 1$. If n is odd, then G has a subgroup $H := \mathrm{GU}_n(q)$ by [16, Theorem 3.11]. Obviously, H contains the Sylow p -subgroup of G and

$$n_p(G) \geq n_p(H) \geq n_p(\mathrm{PSU}_n(q)) \geq p^2$$

by Proposition 3.5, a contradiction. Assume n is even. Write $n = 2^a \cdot d$ with d odd. [16, Theorem 3.11] implies that G has a subgroup H isomorphic to $\mathrm{GU}_d(q^{2^a})$. It is easy to check that the Sylow p -subgroup of H is not trivial and $n_p(G) \geq n_p(H)$. If $d \geq 3$, then by Proposition 3.5,

$$n_p(G) \geq n_p(H) \geq n_p(\mathrm{PSU}_d(q^{2^a})) \geq p^2,$$

a contradiction. Let $d = 1$, that is $G \cong \mathrm{GO}_{2,2}^-(q)$ and $p \mid (q^{2^a} + 1)$. Similarly as above, G has a subgroup H of type $\mathrm{GO}_{2,2}^-(q^{2^{a-1}})$. Note that

$$n_p(S) = n_p(G) \geq n_p(H) = n_p(\Omega_4^-(q^{2^{a-1}})) = n_p(\mathrm{PSL}_2(q^{2^a})).$$

Since $n_p(S) < p^2$, by Proposition 3.4, $q^{2^a} + 1 = p$ is a Fermat prime number, so that q is even. On the other hand, G also has a subgroup M of type $\mathrm{GO}_8^-(u)$ where $u = q^{2^{a-2}}$ and it is easy to see that P is contained in the simple group $A := \mathrm{P}\Omega_8^-(u)$. Moreover, $p = u^4 + 1$. Since $n_p(S) = n_p(G) \geq n_p(M) = n_p(A)$ and every maximal torus of A has order

$$\frac{1}{(2, q^4 + 1)} (q^{n_1} - 1)(q^{n_2} - 1) \cdots (q^{n_k} - 1)(q^{l_1} + 1)(q^{l_2} + 1) \cdots (q^{l_t} + 1)$$

for appropriate partition $4 = n_1 + n_2 + \cdots + n_k + l_1 + l_2 + \cdots + l_t$ of 4, where t is odd. This implies that P is a maximal torus of A and $C_A(P) = P$. Note that $N_A(P)/C_A(P)$ is isomorphic to a subgroup of $\mathrm{Aut}(P) \cong C_{u^4}$, thus $|N_A(P)| \leq u^4(u^4 + 1)$. It follows that

$$\begin{aligned} n_p(A) &= \frac{|A|}{|N_A(P)|} \geq \frac{u^{12}(u^4 + 1)(u^2 - 1)(u^4 - 1)(u^6 - 1)}{u^4(u^4 + 1)} \\ &= u^8(u^2 - 1)(u^4 - 1)(u^6 - 1) > u^{17}. \end{aligned}$$

Since $p = u^4 + 1$, $p^2 = (u^4 + 1)^2 \leq 3u^8 < u^{17}$, it implies that $n_p(A) > p^2$. Therefore, $n_p(S) \geq n_p(A) > p^2$, a contradiction. \square

Next we will consider the exceptional group of Lie type.

Proposition 3.10. *Let $S \cong Sz(q)$ where $q = 2^{2m+1} \geq 8$. Then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong Sz(q)$. Then

$$|S| = q^2(q-1)(q - \sqrt{2q} + 1)(q + \sqrt{2q} + 1).$$

Since $P \cong C_p$, $p \neq 2$. Assume $p \mid (q-1)$. By [15, Table 8.16] S has a maximal subgroup M of type $D_{2(q-1)}$. Obviously, $P \leq C_{q-1} \leq D_{2(q-1)}$. Thus

$$n_p(S) = \frac{|S|}{|M|} = \frac{q^2(q^2+1)}{2}$$

with the help of Lemma 2.7. Note that $p^2 \leq (q-1)^2$ and $q \geq 8$. Therefore $n_p(S) = \frac{q^2(q^2+1)}{2} \geq p^2$, a contradiction. Suppose $p \mid (q - \sqrt{2q} + 1)$, then

$$n_p(S) = \frac{q^2(q-1)(q^2 + \sqrt{2q} + 1)}{4} \geq q^2(q + \sqrt{2q} + 1)$$

by [15, Table 8.16] and Lemma 2.7. Since $p \mid (q - \sqrt{2q} + 1)$, $p^2 \leq (q - \sqrt{2q} + 1)^2 \leq (q+1)^2 \leq 4q^2$. Implying that $n_p(S) \geq q^2(q + \sqrt{2q} + 1) \geq 4q^2 \geq p^2$ since $q \geq 8$, it is impossible. Finally, we assume that $p \mid (q + \sqrt{2q} + 1)$. Then $p^2 \leq (q + \sqrt{2q} + 1)^2 \leq 5q^2$. Similarly as above,

$$n_p(S) = \frac{q^2(q-1)(q - \sqrt{2q} + 1)}{4} = \frac{q^2((q - \sqrt{2q})q + (\sqrt{2q} - 1))}{4} \geq q^3$$

because $q - \sqrt{2q} \geq 4$, it follows that $n_p(S) \geq q^3 \geq 5q^2 \geq p^2$, a contradiction. \square

Proposition 3.11. *Assume $S \cong G_2(q)$ where $q \geq 3$. Then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong G_2(q)$. Then

$$|S| = q^6(q-1)^2(q+1)^2(q^2+q+1)(q^2-q+1).$$

Since $P \cong C_p$, p is coprime to $q(q^2-1)$. Thus, $p \mid (q^2+q+1)$ or $p \mid (q^2-q+1)$. By [15, Tables 8.30, 8.41 and 8.42] S has subgroups $SU_3(q)$ and $SL_3(q)$. Assume p divides q^2+q+1 , then the Sylow p -subgroup P of S is contained in $SL_3(q)$, so that $n_p(S) \geq n_p(SL_3(q)) = n_p(PSL_3(q)) \geq p^2$ unless $S \cong G_2(3)$ by [6, Lemma 2.2] and Proposition 3.4. Suppose $S \cong G_2(3)$, then $p = 13$. Note that S is simple, so P is not normal in G , implying that there exists a maximal subgroup M of S such that $N_S(P) \leq M$, thus $n_p(S) = |S : N_S(P)| \geq |S : M|$. By applying [10], $n_p(S) \geq |S : M| > 13^2$, it is impossible. Suppose p divides q^2-q+1 . Then $P \leq SU_3(q)$ and $n_p(S) \geq n_p(SU_3(q)) = n_p(PSU_3(q)) \geq p^2$ with the help of [6, Lemma 2.2] and Proposition 3.5, a contradiction. \square

Proposition 3.12. *If $S \cong {}^2G_2(q)$ where $q = 3^{2m+1} \geq 27$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Suppose $S \cong {}^2G_2(q)$. Then

$$|S| = q^3(q+1)(q-1)(q-\sqrt{3q}+1)(q+\sqrt{3q}+1).$$

We first suppose that p divides $q+1$. By [15, Table 8.43], S has a maximal subgroup H of type $(2^2 \times D_{\frac{q+1}{2}}) : 3$. Since $|S : H|$ is not divisible by p , the Sylow p -subgroup P of S is contained in H . Moreover, $P \leq D_{\frac{q+1}{2}}$. Clearly, P is normal in $D_{\frac{q+1}{2}}$, so P is also normal in $K := 2^2 \times D_{\frac{q+1}{2}}$. This implies that $K = N_K(P) \leq N_S(P) < S$. Thus $N_S(P) = K$ or H , and

$$n_p(S) = |S : N_S(P)| \geq |S : H| = \frac{q^3(q^2 - q + 1)(q - 1)}{6}.$$

Since $q \geq 27$, $n_p(S) > q^3 \geq 3q^2 \geq (q+1)^2 \geq p^2$, a contradiction.

If $p \mid q-1$. Applying [15, Table 8.43] again, S has a maximal subgroup M isomorphic to $[q^3] : C_{q-1}$. Let Q be a Sylow p -subgroup of M . Then $N_M(Q) = C_{q-1}$ by using [19, Table 5], thus $n_p(M) = |M : N_M(Q)| = |M : C_{q-1}| = q^3$. Since $n_p(S) \geq n_p(M)$ and $q^3 > (q-1)^2 \geq p^2$, $n_p(S) \geq p^2$, a contradiction.

We now suppose that $p \mid q - \sqrt{3q} + 1$. By using [15, Table 8.43] again, S has a maximal subgroup M of type $C_{q-\sqrt{3q}+1} : C_6$. In the light of Lemma 2.7, $N_S(P) = M$. Therefore,

$$n_p(S) = |S : M| = \frac{q^3(q^2 - 1)(q + \sqrt{3q} + 1)}{6} \geq q^4.$$

Note that $p^2 \leq (q - \sqrt{3q} + 1)^2 \leq 4q^2 < q^4$, so $n_p(S) \geq p^2$. Finally, we assume that $p \mid q + \sqrt{3q} + 1$. Similarly as above, we have

$$n_p(S) = \frac{q^3(q^2 - 1)(q - \sqrt{3q} + 1)}{6} \geq q^4.$$

Observe that $p^2 \leq (q + \sqrt{3q} + 1)^2 \leq 9q^2$ since $q \geq 27$. Thus $n_p(S) \geq q^4 \geq 9q^2 \geq p^2$. These cases are impossible. \square

Proposition 3.13. *If $S \cong {}^3D_4(q)$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Suppose $S \cong {}^3D_4(q)$, then

$$|S| = q^{12}(q^2 + q + 1)^2(q^2 - q + 1)^2(q^4 - q^2 + 1)(q + 1)^2(q - 1)^2.$$

Since the Sylow p -subgroup P of S is isomorphic to C_p , p divides $q^4 - q^2 + 1$. By [15, Table 8.51], S has a maximal subgroup M of type $C_{q^4 - q^2 + 1}.C_4$. In the light of Lemma 2.7 we have

$$n_p(S) = |S : M| = \frac{q^{12}(q^2 + q + 1)^2(q^2 - q + 1)^2(q + 1)^2(q - 1)^2}{4} \geq 8q^8.$$

Clearly, $p^2 \leq (q^4 - q^2 + 1)^2 \leq 8q^8$, so $n_p(S) \geq p^2$, a contradiction. \square

Proposition 3.14. *If $S \cong F_4(q)$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Assume $S \cong F_4(q)$, then

$$|S| = q^{24}(q^6 - 1)^2(q^2 - 1)^2(q^2 + 1)^2(q^4 + 1)(q^4 - q^2 + 1).$$

Since the Sylow p -subgroup P of S is isomorphic to C_p , either $p \mid q^4 + 1$ or $p \mid q^4 - q^2 + 1$ by Lemma 2.6, so that $p^2 \leq (q^4 + 1)^2 \leq 3q^8$. Since $N_S(P) < S$, there exists a maximal subgroup M of S such that $N_S(P) \leq M$. Therefore, $n_p(S) = |S : N_S(P)| \geq |S : M|$, which is not less than the degree of minimal permutation representation of $F_4(q)$. In the light of [22, Theorem 2],

$$n_p(S) \geq \frac{(q^{12} - 1)(q^4 + 1)}{(q - 1)} \geq q^{15} > 3q^8.$$

Thus $n_p(S) \geq p^2$, a contradiction. \square

Proposition 3.15. *If $S \cong {}^2F_4(q)$ where $q = 2^{2m+1} \geq 8$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong {}^2F_4(q)$. Then

$$|S| = q^{12}(q^4 - 1)^2(q^2 - q + 1)(q^2 + q + 1 + \sqrt{2q}(q + 1))(q^2 + q + 1 - \sqrt{2q}(q + 1)).$$

Since $P \cong C_p$, p is a divisor of one of the following numbers: $q^2 + q + 1 + \sqrt{2q}(q + 1)$, $q^2 + q + 1 - \sqrt{2q}(q + 1)$ or $q^2 - q - 1$. This implies that $p^2 \leq (q^2 + q + 1 + \sqrt{2q}(q + 1))^2 \leq 25q^4$. On the other hand, since S is a simple group, $N_S(P)$ must be contained in one of maximal subgroups of S . Therefore, $|S : N_S(P)|$ is not less than the degree of minimal permutation representation of ${}^2F_4(q)$. By using [23, Theorem 5],

$$n_p(S) = |S : N_S(P)| \geq (q^6 + 1)(q^3 + 1)(q + 1) > q^{10} > 25q^4 \geq p^2,$$

a contradiction. \square

Proposition 3.16. *If $S \cong E_6(q)$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong E_6(q)$. Then

$$|S| = \frac{q^{36}(q^{12} - 1)(q^9 - 1)(q^8 - 1)(q^6 - 1)(q^5 - 1)(q^2 - 1)}{(3, q - 1)}.$$

Obviously, p does not divide $q - 1$ since the Sylow p -subgroup P of S is isomorphic to C_p , that is $p \nmid (3, q - 1)$. We first assume that $p \mid (q^9 - 1)$. By using [24, Table 9] S has a subgroup H of type $\text{PSL}_3(q^3)$, thus $n_p(S) \geq n_p(H) \geq p^2$ by Proposition 3.4, it is impossible. Suppose that p is a prime divisor of $(q^{12} - 1)(q^8 - 1)(q^6 - 1)(q^5 - 1)(q^2 - 1)$, then $p^2 \leq (q^6 + 1)^2 < 3q^{12}$. On the other hand, $n_p(S)$ is not less than the degree of the minimal permutation representation of $E_6(q)$ since $N_S(P) < S$. With the help of [25, Theorem 1] we have

$$n_p(S) \geq \frac{(q^9 - 1)(q^8 + q^4 + 1)}{q - 1} > q^{16} > 3q^{12} > p^2,$$

which is a contradiction. \square

Proposition 3.17. *If $S \cong {}^2E_6(q)$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong {}^2E_6(q)$. Then

$$|S| = \frac{q^{36}(q^{12}-1)(q^9+1)(q^8-1)(q^6-1)(q^5+1)(q^2-1)}{(3, q+1)}.$$

Observe that p does not divide $(3, q+1)$. We first suppose that p divides q^9-1 . By [24, Table 10] S has a subgroup H of type $\text{PSU}_3(q^3)$, so that $n_p(S) \geq n_p(H) \geq p^2$ with the help of Proposition 3.5, a contradiction. We next suppose that p divides $(q^{12}-1)(q^8-1)(q^6-1)(q^5+1)(q^2-1)$, then $p^2 \leq (q^6+1)^2 \leq 3q^{12}$. Similarly as above, $n_p(S)$ is not less than the smallest index of a maximal subgroup of S . In the light of [23, Theorem 4]

$$n_p(S) \geq \frac{(q^{12}-1)(q^6-q^3+1)(q^4+1)}{q-1} > q^{15} > 3q^{12} \geq p^2,$$

a contradiction. \square

Proposition 3.18. *If $S \cong E_7(q)$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. Let $S \cong E_7(q)$. Then

$$|S| = q^{63}(q^{18}-1)(q^{14}-1)(q^{12}-1)(q^{10}-1)(q^8-1)(q^6-1)(q^2-1).$$

Since p is a prime divisor of $|S|$, $p^2 \leq (q^9+1)^2 < 3q^{18}$. S is a simple group, so $N_S(P)$ must be contained in one maximal subgroups of S . Therefore, $n_p(S) = |S : N_P(S)|$ is not less than the smallest index of a maximal subgroup of S . By [25, Theorem 2],

$$n_p(S) \geq \frac{(q^{14}-1)(q^9+1)(q^5+1)}{(q-1)} \geq q^{27} > 3q^{18} > p^2,$$

which is a contradiction. \square

Proposition 3.19. *If $S \cong E_8(q)$, then there does not exist $p \in \pi(S)$ such that $n_p(S) < p^2$.*

Proof. If $S \cong E_8(q)$, then

$$|S| = q^{120}(q^{30}-1)(q^{24}-1)(q^{20}-1)(q^{18}-1)(q^{14}-1)(q^{12}-1)(q^8-1)(q^2-1).$$

Clearly, $p^2 \leq (q^{15}+1)^2 < 3q^{30}$ and $n_p(S)$ is not less than the smallest index of a maximal subgroup of S . By using [25, Theorem 3]

$$n_p(S) \geq \frac{(q^{20}-1)(q^{12}+1)(q^{10}+1)(q^6+1)}{(q-1)} > q^{47} > 3q^{30} > p^2,$$

a contradiction. \square

4. PROOF OF THEOREM 1.1

In this section, we will prove Theorem 1.1 by the following theorem due to M. Hall.

Theorem 4.1 (Theorem 2.2 of [1]). *The number $n_p(G)$ of Sylow p -subgroups in a finite group G is the product of factors of the following two kinds:*

- (1) the number s_p of Sylow p -subgroups in a simple group X ;

(2) a prime power q^t where $q^t \equiv 1 \pmod{p}$.

Let G be a finite group with $n_p(G) < p^2$. Assume that $n_p(G)$ is not a power of prime. In order to prove Theorem 1.1, it is enough to prove that $n_p(G) = n_p(S)$ for some simple group S with $n_p(S) < p^2$. By Theorem 4.1 we have

$$n_p(G) = q_1^{t_1} q_2^{t_2} \cdots q_m^{t_m} \cdot n_p(S_1) n_p(S_2) \cdots n_p(S_n)$$

where S_i is a simple group for $1 \leq i \leq n$ and $q_j^{t_j} \equiv 1 \pmod{p}$ for $1 \leq j \leq m$. Obviously, if $q_j^{t_j}$ and $n_p(S_i)$ are not equal to 1 then they are greater than p , for any i, j . Since $n_p(G) < p^2$ is not a power of prime, $q_1^{t_1} \cdots q_m^{t_m} = 1$ and there exists at most one index i such that $n_p(S_i) \neq 1$. Therefore $n_p(G) = n_p(S_i)$ where S_i is a simple group with $n_p(S_i) < p^2$.

Conversely, a power of a prime r which is congruent to 1 mod p can become a Sylow p -number. In fact it is one of a Frobenius group with the complement of the cyclic group of order p and the kernel of elementary abelian r -group. \square

5. THE PROOF OF THEOREM 1.2

In this section, we will prove Theorem 1.2. Recall that, if G is a p -solvable group, meaning that it admits a series of normal subgroups $1 = V_0 < \cdots < V_n = G$ such that each factor V_{i+1}/V_i is either a p -group or a p' -group. It is easy to see that normal subgroups and the quotient groups of p -solvable group are also p -solvable.

Lemma 5.1. *Let $a = 1 + r_1p$, $b = 1 + r_2p < p^2$ where r_1, r_2 are integers. If $a|b$, then $a = 1$ or $a = b$.*

Proof. We assume that $a \neq 1$, that is $r_1 \neq 0$. Since a divides b , we can write $1 + r_2p = d(1 + r_1p)$ where d is an integer. Thus $r_2 = \frac{d-1}{p} + dr_1$. Note that r_2, dr_1 are integers, so $\frac{d-1}{p}$ is also an integer, this implies that $d = 1$ or $d > p$. If $d > p$, then $1 + r_2p = d(1 + r_1p) > p^2$, a contradiction. Therefore $d = 1$. \square

First, we prove that if $n_p(G)$ is a power of a prime then G is p -solvable. To prove our result, we can prove that if G is not a p -solvable group, then $n_p(G)$ is not a power of a prime. Since G is not a p -solvable group, G is nonsolvable. Therefore, G has a nonsolvable chief factor S^m such that p divides $|S|^m$ where S is a nonabelian simple group and $m \geq 1$. Obviously, $n_p(S) \neq 1$. In the light of Lemma 2.4 we have $n_p(S)$ divides $n_p(G)$. Moreover, we can get $n_p(S) = n_p(G)$ by Lemma 5.1. On the other hand, $n_p(G) < p^2$, thus $n_p(S)$ satisfies one of items (1), (2) and (3) of Theorem 3.1. If $n_p(S) = 1 + p = q^a$ for some prime q and integer a , then $q = 2$, a is a prime number, that is $p = 2^a - 1$ is a Mersenne prime number, a contradiction. If $n_p(G) = 1 + \frac{(p-3)p}{2} = q^a$ where p is a Fermat prime number, q is prime and a is an integer, then $p^2 - 3p + 2 = 2q^a$. By Lemma 2.2, we have $p = 3$, contradicting $p > 3$. Therefore, $n_p(G)$ is not a power of a prime. This proves that if G is not a p -solvable group then $n_p(G)$ is not a power of a prime.

Conversely, let G be a p -solvable group. We assume that G is a minimal order such that $n_p(G)$ is not a power of a prime. Write $n_p(G) = q_1^{t_1} q_2^{t_2} \cdots q_k^{t_k}$ is the prime factorization of $n_p(G)$. We claim that G is nonsolvable. If this were not the case,

then $q_i^{t_i} \equiv 1 \pmod{p}$ for any $i = 1, \dots, k$ (see [26]). By our assumption that $n_p(G)$ is not a power of a prime, $n_p(G)$ has at least two prime divisors. This implies that $n_p(G) > p^2$, a contradiction. It is not difficult to see that G is not simple because otherwise $p \notin \pi(G)$, so that $n_p(G) = 1$, a contradiction.

Let M be a maximal normal subgroup of G , then either G/M is a cyclic group of order p or $G/M \cong S$ where p does not divide $|S|$. Thus $n_p(G/M) = 1$. By Lemma 2.4 we have

$$n_p(G) = n_p(G/M)n_p(M)n_p(N_{PM}(P \cap M)/P \cap M),$$

where P is a Sylow p -subgroup of G . Observe that $n_p(G)$ is divisible by $n_p(M)$, so $n_p(M) = 1$ or $n_p(M) = n_p(G)$ with the help of Lemma 5.1. Since M is a p -solvable group and $|M| < |G|$, by the minimality of G , $n_p(M) = 1$. Thus

$$n_p(G) = n_p(N_{PM}(P \cap M)/P \cap M).$$

Assume p does divide $|G/M|$, then $P \leq M$. Moreover, $P \leq M$. Thus

$$N_{PM}(P \cap M)/P \cap M = N_M(P)/P = M/P,$$

this implies that $p \notin \pi(M/P)$, thus $n_p(G) = n_p(M/P) = 1$, contradicting the fact that $n_p(G)$ is not a power of a prime.

Assume $G/M \cong C_p$, then $P \cap M$ is a Sylow p -subgroup of M . Since $n_p(M) = 1$, $P \cap M \leq M$. Obviously, $P \cap M \leq P$ because M is normal in G , thus $N_{PM}(P \cap M) = PM$. Therefore

$$N_{PM}(P \cap M)/P \cap M = PM/P \cap M.$$

Now we have a normal series

$$P \cap M \leq M \leq PM,$$

thus $M/P \cap M \leq PM/P \cap M$. Since p does not divide $|M/M \cap P|$, the Sylow p -subgroup of $PM/M \cap P$ is $P/M \cap P$, which is isomorphic to C_p because $P/M \cap P \cong MP/M = G/M$. Note that

$$PM/P \cap M \cong (M/M \cap P)(P/M \cap P) \cong (M/M \cap P)C_p$$

and p does not divide $M/(M \cap P)$, thus $PM/M \cap P$ is a p -solvable group. Recall that $n_p(G) = n_p(PM/M \cap P)$, the minimality of G implies that $|PM/M \cap P| = |G|$. Therefore $G = MP$ and $P \cap M = 1$, that is

$$G \cong M \rtimes P \cong M \rtimes C_p.$$

Observe that $(|M|, |P|) = 1$, so the action of P on M is coprime. For any $r \in \pi(M)$, there exists a P -invariant Sylow r -subgroup R of M by [21, 8.2.3(a)], this implies that $RP \leq G$. In the light of [3, Theorem A], $n_p(RP)$ divides $n_p(G)$, thus $n_p(RP) = 1$ or $n_p(RP) = n_p(G)$ with the help of Lemma 5.1. Assume $n_p(G) = n_p(RP)$. Since RP is a p -solvable group, by the minimality of G we can get $RP = G$, contradicting the nonsolvability of G . Hence $n_p(RP) = 1$, implying that $RP = R \times P$. Since

$$M = \prod_{r \in \pi(M)} R,$$

$G = M \times P$, and hence $n_p(G) = 1$, a contradiction. \square

REFERENCES

- [1] M. Hall. *On the number of Sylow subgroups in a finite group*. Journal of Algebra, 7(3):363-371, 1967.
- [2] J. P. Zhang. *Sylow numbers of finite groups*. Journal of Algebra, 176(1):111-123, 1995.
- [3] G. Navarro. *Number of Sylow subgroups in p -solvable groups*. Proceeding of the American Mathematical Society, 131(10):3019-3020, 2003.
- [4] T. Li and Y. Liu. *Mersenne primes and solvable Sylow numbers*. Journal of Algebra and its Application, 15(9):1650163, 2016.
- [5] W. Guo and E. P. Vdovin. *Number of Sylow subgroups in finite groups*. Journal of Group Theory, 21(4):695-712, 2018.
- [6] Z. Wu. *On the number of Sylow subgroups in finite simple groups*. Journal of Algebra and Its Applications, 20(07):2150115, 2021.
- [7] K. Zsigmondy. *Zur theorie der potenzreste*. Monatshefte für Mathematik, 3:265-284, 1892.
- [8] I. M. Isaacs. *Algebra: a graduate course, volume 100*. American Mathematical Society, 2009.
- [9] R. Brauer and W. F. Reynolds. *On a problem of E. Artin*. Annals of Mathematics, 68(3):713-720, 1958.
- [10] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson. *Atlas of finite groups*. Oxford University Press, Eynsham, 1985.
- [11] L. Weisner. *On the Sylow Subgroups of the Symmetric and Alternating Groups*. American Journal of Mathematics, 47(2):121-124, 1925.
- [12] F. Gross. *Hall subgroups of order not divisible by 3*. The Rocky Mountain Journal of Mathematics, 23(2):569-591, 1993.
- [13] S. P. Glasby, F. Lübeck, A. C. Niemeyer, and C. E. Praeger. *Primitive prime divisors and the n -th cyclotomic polynomial*. Journal of the Australian Mathematical Society, 102:122-135, 2017.
- [14] The GAP Group. *GAP-Groups, Algorithms, and Programming*. Version 4.7.8.
- [15] J. N. Bray, D. F. Holt, and C. M. Roney-Dougal. *The maximal subgroups of the low-dimensional finite classical groups, volume 407*. Cambridge university press, 2013.
- [16] R. Wilson. *The finite simple groups, volume 147*. Springer, 2009.
- [17] R. W. Carter. *Simple Groups of Lie Type*. John Wiley and Sons, New York, 1972.
- [18] A. V. Vasil'ev. *Normalizers of Sylow subgroups in finite linear and unitary groups*. Algebra and Logic, 59(1):1-18, 2020.
- [19] A. V. Vasiliev and E. P. Vdovin. *An adjacency criterion for the prime graph of a finite simple group*. Algebra and Logic, 44(6):381-406, 2005.
- [20] N. Ahanjideh and A. Iranmanesh. *On the Sylow normalizers of some simple classical groups*. Bulletin of the Malaysian Mathematical Sciences Society, 35(2):459-467, 2012.
- [21] H. Kurzweil and B. Stellmacher. *The Theory of Finite Groups: An Introduction*. Springer, 2004.
- [22] A. V. Vasilyev. *Minimal permutation representations of finite simple exceptional groups of types G_2 and F_4* . Algebra and Logic, 35(6):371-383, 1996.
- [23] A. V. Vasilyev. *Minimal permutation representations of finite simple exceptional twisted groups*. Algebra and Logic, 37(1):9-20, 1998.
- [24] D. A. Craven. *The maximal subgroups of the exceptional groups $F_4(q)$, $E_6(q)$ and ${}^2E_6(q)$ and related almost simple groups*. Inventiones Mathematicae, 234:637-719, 2023.
- [25] A. V. Vasilyev. *Minimal permutation representations of finite simple exceptional groups of types E_6 , E_7 , and E_8* . Algebra and Logic, 36(5):302-310, 1997.
- [26] P. Hall. *A note on soluble groups*. Journal of the London Mathematical Society, s1-3(2):98-105, 1928.

XIAOFANG GAO. DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE BRASÍLIA, CAMPUS UNIVERSITÁRIO, DARCY RIBEIRO, BRASÍLIA-DF, 70910-900, BRAZIL.

Email address: `gaoxiaofang2020@hotmail.com`

IGOR LIMA. DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE BRASÍLIA, CAMPUS UNIVERSITÁRIO, DARCY RIBEIRO, BRASÍLIA-DF, 70910-900, BRAZIL.

Email address: `igor.matematico@gmail.com`

RULIN SHEN. DEPARTMENT OF MATHEMATICS, HUBEI MINZU UNIVERSITY, ENSHI, HUBEI PROVINCE, 445000, P. R. CHINA.

Email address: `shenrulin@hotmail.com`