

TORSION OF RATIONAL ELLIPTIC CURVES OVER THE CYCLOTOMIC EXTENSIONS OF \mathbb{Q}

OMER AVCI

ABSTRACT. Let E be an elliptic curve defined over \mathbb{Q} . Let $p > 3$ be a prime such that $p - 1$ is not divisible by 3, 4, 5, 7, 11. In this article we classify the groups that can arise as $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ up to isomorphism. The method illustrates techniques for eliminating possible structures that can appear as a subgroup of $E(\mathbb{Q}^{ab})_{\text{tors}}$.

1. INTRODUCTION AND NOTATION

Let K denote a number field, and let E be an elliptic curve over K . The Mordell-Weil theorem states that the set of K -rational points on E forms a finitely generated abelian group. Specifically, denoting $E(K)$ as the set of K -rational points on E , we have

$$E(K) \cong \mathbb{Z}^r \oplus E(K)_{\text{tors}}.$$

Here, r is a nonnegative integer called the rank, and $E(K)_{\text{tors}}$ is a finite group known as the torsion subgroup of E over K .

In fact, due to a theorem of Merel [15], there is a bound on the size of the torsion subgroup that depends only on the degree of K over \mathbb{Q} . Thus, there is a finite list of torsion subgroups that can appear as $E(K)_{\text{tors}}$ as K varies over number fields of a fixed degree d and E/K varies. This means that there are only finitely many torsion subgroups (up to isomorphism) that can appear as $E(K)_{\text{tors}}$ when K has a fixed degree and E is defined over K .

Determining the possible groups for a fixed degree d has been solved by Mazur for $d = 1$ in [2], by Kenku [8], Kamienny, Momose [9] for $d = 2$, and by Derickx, and independently by Etropolski, Morrow, and Zureick-Brown for $d = 3$ in [10].

Another variant of this problem is to determine the torsion subgroups (up to isomorphism) that can appear as $E(K)_{\text{tors}}$ when K has a fixed degree and E is defined over \mathbb{Q} . Clearly, the set of torsion subgroups in this problem is a subset of the set of torsion subgroups in the previous problem. This version of the problem is solved by Najman when $d = 2, 3$. A subcase of the problem is solved by Chou when $d = 4$ and K/\mathbb{Q} is abelian in [4]. For all quartic fields, it is solved by González-Jiménez and Najman in [11].

In our work we will classify $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ for the primes $p > 3$ such that $p - 1$ is not divisible by 3, 4, 5, 7, 11. By doing so we will be also classifying $E(\mathbb{Q}(\mu_{p^\infty}))_{\text{tors}}$ for those primes. Our result is the following theorem. We will give the proof in Section 3.

Theorem 1.1. *Let E/\mathbb{Q} be an elliptic curve. Let $p > 3$ be a prime such that $p - 1$ is not divisible by 3, 4, 5, 7, 11. Then $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ is either one of the groups from Mazur's Theorem, or one of the following groups $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ or $\mathbb{Z}/16\mathbb{Z}$.*

It is also possible to consider the possibilities of torsion subgroups when K is an infinite extension. In this case, the Mordell-Weil theorem does not apply, so we do not know whether $E(K)_{\text{tors}}$ is a finite group. For some special cases of K , it is possible to prove that $E(K)_{\text{tors}}$ is finite and can take only finitely many possibilities where E/\mathbb{Q} varies. For example, Fujita classified all the possibilities of $E(\mathbb{Q}(2^\infty))_{\text{tors}}$ in [12], where $\mathbb{Q}(2^\infty)$ is the compositum of all quadratic fields. Similarly, Daniels, Lozano-Robledo, Najman, and Sutherland classified all the possibilities of $E(\mathbb{Q}(3^\infty))_{\text{tors}}$ in [13], where $\mathbb{Q}(3^\infty)$ is the compositum of all cubic fields.

One recent development is due to Chou, who classified all the possibilities of $E(\mathbb{Q}^{ab})_{\text{tors}}$ in [1], where \mathbb{Q}^{ab} is the maximal abelian extension of \mathbb{Q} . Before this, it was known that $E(\mathbb{Q}^{ab})_{\text{tors}}$ is finite for any E/\mathbb{Q} due to a theorem of Ribet [14]. However, this did not imply that $E(\mathbb{Q}^{ab})_{\text{tors}}$ could take only finitely many structures because Merel's bound in [15] cannot be used to bound the order of this torsion subgroup in an infinite extension. Chou's result enabled us to classify $E(K)_{\text{tors}}$ for any abelian Galois number field K as E/\mathbb{Q} varies. All possible groups are subgroups of those appearing in Theorem 1.2 of Chou. The remaining task is to eliminate groups from this list that are not realized for any E/\mathbb{Q} .

Theorem 1.2 (Chou, [1]). *Let E/\mathbb{Q} be an elliptic curve. Then $E(\mathbb{Q}^{ab})_{\text{tors}}$ is isomorphic to one of the following groups*

$$E(\mathbb{Q}^{ab})_{\text{tors}} \simeq \begin{cases} \mathbb{Z}/N\mathbb{Z} & \text{with } 1 \leq N \leq 19, \text{ or } N = 21, 25, 27, 37, 43, 67, 163, \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z} & \text{with } 1 \leq N \leq 9, \\ \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3N\mathbb{Z} & \text{with } N = 1, 2, 3, \\ \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4N\mathbb{Z} & \text{with } N = 1, 2, 3, 4, \\ \mathbb{Z}/5\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z} \\ \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \\ \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z} \end{cases}$$

In González-Jiménez and Najman's result in [11], it is shown that if K is any number field with extension degree not divisible by 2, 3, 5, 7, then $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$ for any elliptic curve E/\mathbb{Q} . We enhance this approach by relaxing the condition from not divisible by 2 to not divisible by 4. Similar to this result, we will show that many of the torsion points are rational points, though not necessarily on the original elliptic curve E itself, but possibly on a quadratic twist of E instead. This modification allows us to restrict the possible group structures using Mazur's theorem.

Let μ_n , for positive integer n , be the set of n -th roots of unity. Let ζ_p be the p -th primitive root of unity. For a prime number p , we define a set μ_{p^∞} as the set of all complex numbers ω for which there exists non-negative integer k such that $\omega^{p^k} = 1$. Note that $\mathbb{Q}(\mu_{p^\infty})$ is the set \mathbb{Q} extended with all $p^{n^{\text{th}}}$ primitive roots of unity.

Gužvić and Krijan [5] showed that $E(\mathbb{Q}(\mu_{p^\infty}))_{\text{tors}} = E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ for all primes $p > 3$. This work enables us to classify $E(\mathbb{Q}(\mu_{p^\infty}))_{\text{tors}}$ for the primes p such that $p - 1$ is not divisible by 3, 4, 5, 7, 11. Also, Gužvić and Vukorepa [6] classified all the possible torsion subgroups that can appear as $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ for $p = 5, 7, 11$. In our result, we extend the classification of $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ for an infinite family of primes.

2. TORSION OF RATIONAL ELLIPTIC CURVES

In this section, we will present some theorems about the classification of $E(K)_{\text{tors}}$.

Theorem 2.1 (Mazur, [2]). *Let E/\mathbb{Q} be an elliptic curve. Then*

$$E(\mathbb{Q})_{\text{tors}} \simeq \begin{cases} \mathbb{Z}/N\mathbb{Z} & \text{with } 1 \leq N \leq 10 \text{ or } N = 12 \text{ or,} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z} & \text{with } 1 \leq N \leq 4. \end{cases}$$

All of the 15 torsion subgroups appear infinitely often.

Theorem 2.2. (Najman, [3]) *Let E/\mathbb{Q} be an elliptic curve defined over \mathbb{Q} , and let K be a quadratic number field. Then*

$$E(K)_{\text{tors}} \simeq \begin{cases} \mathbb{Z}/N\mathbb{Z} & \text{with } 1 \leq N \leq 10 \text{ or } N = 12, 15, 16, \text{ or} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z} & \text{with } 1 \leq N \leq 6 \text{ or} \\ \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3N\mathbb{Z} & \text{with } N = 1, 2, \text{ only if } K = \mathbb{Q}(\sqrt{-3}) \text{ or} \\ \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} & \text{only if } K = \mathbb{Q}(\sqrt{-1}). \end{cases}$$

Each of these groups, except for $\mathbb{Z}/15\mathbb{Z}$, appears as the torsion structure over a quadratic field for infinitely many rational elliptic curves E . The elliptic curves 50B1 and 50A3 have 15-torsion over $\mathbb{Q}(\sqrt{5})$, 50B2 and 450B4 have 15-torsion over $\mathbb{Q}(\sqrt{-15})$. These are the only rational curves having non-trivial 15-torsion over any quadratic field.

Theorem 2.3. (Chou, [1], Corollary to Theorem 1.2) *Let E/\mathbb{Q} be an elliptic curve. Let K be an abelian Galois number field. Then $E(K)_{\text{tors}}$ is isomorphic to one of the following groups:*

$$E(K)_{\text{tors}} \simeq \begin{cases} \mathbb{Z}/N\mathbb{Z} & \text{with } 1 \leq N \leq 19, \text{ or } N = 21, 25, 27, 37, 43, 67, 163, \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z} & \text{with } 1 \leq N \leq 9, \\ \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3N\mathbb{Z} & \text{with } N = 1, 2, 3, \\ \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4N\mathbb{Z} & \text{with } N = 1, 2, 3, 4, \\ \mathbb{Z}/5\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z} \\ \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \\ \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z} \end{cases}$$

Lemma 2.4 (Chou, [4]). *Let E be an elliptic curve defined over K , α square-free in K . If n is odd, then there exists an isomorphism*

$$E(K(\sqrt{\alpha}))[n] \cong E(K)[n] \oplus E_{\alpha}(K)[n]$$

where E_{α} denotes the α -quadratic twist of E .

3. ELIMINATING POSSIBLE TORSION

In this section, we will present some methods used to eliminate possible torsion subgroups that appear in Corollary 2.3.

Lemma 3.1. *Let K be a Galois number field of degree n . Let E/\mathbb{Q} be an elliptic curve. Let p be an odd prime. Let $E(K)[p^\infty] \cong \mathbb{Z}/p^m\mathbb{Z}$ for some positive integer m . If $\gcd(n, p^m) = 1$ then $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$. If $\gcd(n, p^m) = 2$ then $E(K)[p^\infty] = E(\mathbb{Q}(\sqrt{d}))[p^\infty]$ for some square-free integer d satisfying $\sqrt{d} \in K$.*

Proof. E is isomorphic over rational to some elliptic curve in short Weierstrass form, so we can assume $E : y^2 = x^3 + Ax + B$ for some $A, B \in \mathbb{Q}$. Let $P \in E(K)[p^\infty]$ be a point of order p^m . Let $\sigma \in \text{Gal}(K/\mathbb{Q})$ be a Galois action. Then $P^\sigma = aP$ for some $a \in (\mathbb{Z}/p^m\mathbb{Z})^\times$ because $P^\sigma \in E(K)[p^\infty]$. If we apply σ action n times, we obtain $P^{\sigma^n} = P$ since $\sigma^n = \text{id}$. If we apply σ action $\phi(p^m)$ times, we obtain $P^{\sigma^{\phi(p^m)}} = P$ since $a^{\phi(p^m)} \equiv 1 \pmod{p^m}$.

If $\gcd(n, \phi(p^m)) = 1$ then by Euclidean algorithm we obtain $P^\sigma = P$ for all $\sigma \in \text{Gal}(K/\mathbb{Q})$. Therefore P is a rational point. Since P generates $E(K)[p^\infty]$, we obtain $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$.

Similarly if $\gcd(n, \phi(p^m)) = 2$ then by Euclidean algorithm we obtain $P^{\sigma^2} = P$ for all $\sigma \in \text{Gal}(K/\mathbb{Q})$. Therefore $a \equiv \pm 1 \pmod{\phi(p^m)}$ and $P^\sigma = \pm P$. Then

$$(x^\sigma, y^\sigma) = P^\sigma = \pm P = (x, \pm y)$$

This implies $x^\sigma = x$ for all $\sigma \in \text{Gal}(K/\mathbb{Q})$ hence it is rational. This implies y^2 is also rational therefore $y \in \mathbb{Q}(\sqrt{d})$ for some square-free integer d satisfying $\sqrt{d} \in K$. This means that $P \in E(\mathbb{Q}(\sqrt{d}))[p^\infty]$ and P generates $E(\mathbb{Q}(\sqrt{d}))[p^\infty]$ so we have $E(K)[p^\infty] = E(\mathbb{Q}(\sqrt{d}))[p^\infty]$. \square

Corollary 3.2. *Let K be a Galois number field of degree n . Let E/\mathbb{Q} be an elliptic curve. Let p be an odd prime and $E(K)[p^\infty] \cong \mathbb{Z}/p^m\mathbb{Z}$ for some positive integer m . If $\gcd(n, p^m) \leq 2$ then $E(K)[p^\infty] \cong E_d(\mathbb{Q})[p^\infty]$ for some square-free integer d (possibly $d = 1$) satisfying $\sqrt{d} \in K$.*

Proof. We have shown that $E(K)[p^\infty] = E(\mathbb{Q}(\sqrt{d}))[p^\infty]$ for some square-free integer d satisfying $\sqrt{d} \in K$. We also have $E(\mathbb{Q}(\sqrt{d}))[p^\infty] = E(\mathbb{Q}(\sqrt{d}))[p^m]$. We can use Lemma 2.4 to obtain

$$E(\mathbb{Q}(\sqrt{d}))[p^m] = E(\mathbb{Q})[p^m] \oplus E_d(\mathbb{Q})[p^m].$$

This means that either $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ or $E(K)[p^\infty] \cong E_d(\mathbb{Q})[p^\infty]$ holds. \square

Lemma 3.3. *Let K be a finite Galois extension of \mathbb{Q} with $[K : \mathbb{Q}] = n$ for some positive integer n . Let E/\mathbb{Q} be an elliptic curve. Then,*

- (i) *if the extension degree n is not divisible by 3, $E(K)_{\text{tors}} \not\cong \mathbb{Z}/N\mathbb{Z}$ for $N = 19, 21, 27, 163$, and $E(K)_{\text{tors}} \not\cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z}$ for $N = 7, 9$,*
- (ii) *if the extension degree n is not divisible by 3 and 4, then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/N\mathbb{Z}$ for $N = 13, 37$,*
- (iii) *if the extension degree n is not divisible by 3 and 7, then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/43\mathbb{Z}$,*
- (iv) *if the extension degree n is not divisible by 3 and 11, then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/67\mathbb{Z}$.*
- (v) *if the extension degree n is not divisible by 4, then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/17\mathbb{Z}$.*
- (vi) *if the extension degree n is not divisible by 4, and the fields $\mathbb{Q}(\sqrt{5})$ and $\mathbb{Q}(\sqrt{-15})$ are not subfields of K , then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/15\mathbb{Z}$.*
- (vii) *if the extension degree n is not divisible by 5, then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/11\mathbb{Z}$.*
- (viii) *if the extension degree n is not divisible by 4 and 5 then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/25\mathbb{Z}$.*

Proof. Let us show that $E(K)_{\text{tors}} \cong \mathbb{Z}/N\mathbb{Z}$ is not possible for the N values except 15 and 21, given in the lemma with the suitable assumption about divisors of n . All such N are prime or prime powers so we can use Corollary 3.2. It can be seen that with the assumption about n , we have $\gcd(n, \phi(N)) \leq 2$ therefore $E(K)_{\text{tors}} \cong E_d(\mathbb{Q})_{\text{tors}}$ for some square-free integer d satisfying $\sqrt{d} \in K$. Since E_d is also an elliptic curve over \mathbb{Q} , from Mazur's theorem we can see that $E_d(\mathbb{Q})_{\text{tors}} \cong \mathbb{Z}/N\mathbb{Z}$ is not possible.

The case $E(K)_{\text{tors}} \cong \mathbb{Z}/21\mathbb{Z}$ needs more careful consideration. By using Corollary 3.2, we obtain $E(K)[3] \cong E_{d_1}(\mathbb{Q})[3]$ and $E(K)[7] \cong E_{d_2}(\mathbb{Q})[7]$ for some square-free integers d_1, d_2 satisfying $\sqrt{d_1}, \sqrt{d_2} \in K$. Clearly $E_{d_1}(\mathbb{Q})[3]$ is a subgroup of $E_{d_1}(\mathbb{Q})[21]$ and $E_{d_2}(\mathbb{Q})[7]$ is a subgroup of $E_{d_2}(\mathbb{Q})[21]$. If $d_1 = d_2$ then we get $E_{d_1} \cong \mathbb{Z}/21\mathbb{Z}$ which is not possible due to Mazur's theorem since E_{d_1} is defined over \mathbb{Q} . Otherwise, by using Lemma 2.4, we obtain

$$E_{d_1}(\sqrt{d_2})[21] \cong E_{d_1}(\mathbb{Q})[21] \oplus E_{d_2}(\mathbb{Q})[21].$$

Note that $E_{d_1}(\mathbb{Q})[3] \cong \mathbb{Z}/3\mathbb{Z}$ and $E_{d_2}(\mathbb{Q})[7] \cong \mathbb{Z}/7\mathbb{Z}$ since $E(K)_{\text{tors}} \cong \mathbb{Z}/21\mathbb{Z}$. This means that $E_{d_1}(\sqrt{d_2})[21]$ has a subgroup isomorphic to $\mathbb{Z}/21\mathbb{Z}$ and it is also a subgroup of $E_{d_1}(\sqrt{d_2})_{\text{tors}}$. Notice that E_{d_1} is defined over \mathbb{Q} and we are looking its torsion over a quadratic field. By Theorem 2.2, we can see that it can not have a subgroup isomorphic to $\mathbb{Z}/21\mathbb{Z}$. Hence $E(K)_{\text{tors}} \cong \mathbb{Z}/21\mathbb{Z}$ is not possible.

Similarly, the case $E(K)_{\text{tors}} \cong \mathbb{Z}/15\mathbb{Z}$ also requires special attention. By using same arguments, we obtain $E_{d_1}(\mathbb{Q}(\sqrt{d_2}))_{\text{tors}}$ has a subgroup of 15 elements for some square-free integers d_1, d_2 satisfying $\sqrt{d_1}, \sqrt{d_2} \in K$. Since E_{d_1} is defined over \mathbb{Q} , by Theorem 2.2, we get $d_2 = 5$ or $d_2 = -15$ but this is not possible since we assumed both $\sqrt{5}$ and $\sqrt{-15}$ does not belong to K .

Finally, we need to show that $E(K)_{\text{tors}} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z}$ is not possible for $N = 7, 9$. Assume otherwise, then from Corollary 3.2, we get $E(K)[N] \cong E_d(\mathbb{Q})[N]$ for some square-free d satisfying $\sqrt{d} \in K$. Since $E(K)$ has full 2-torsion, $E_d(K)$ also has full 2-torsion. If the cubic $x^3 + Ax + B$ is irreducible over \mathbb{Q} , then the extension degree n if K must be divisible by 3 which contradicts with our assumption. This means that the cubic must have a rational root so there is a rational 2-torsion point of E and E_d . Recall that $E(K)[N] \cong E_d(\mathbb{Q})[N] \cong \mathbb{Z}/N\mathbb{Z}$. Then, there must be a subgroup of $E_d(\mathbb{Q})$ of order $2N$, but this contradicts with Mazur's theorem. \square

Lemma 3.4. *Let K be a finite Galois extension of \mathbb{Q} with $[K : \mathbb{Q}] = n$ for some positive integer n . Let E/\mathbb{Q} be an elliptic curve. If the extension degree n is not divisible by 3 and 4, then $E(K)_{\text{tors}} \not\cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/16\mathbb{Z}$.*

Proof. Assume otherwise to obtain contradiction. Let $P = (x_0, y_0) \in E(K)_{\text{tors}}$ be a point of order 16. For any Galois action $\sigma \in \text{Gal}(K/\mathbb{Q})$, the point $P^\sigma \in E(K)_{\text{tors}}$ also has order 16. There are 16 points in $E(K)_{\text{tors}}$ of order 16, so their x -coordinates form an 8-element set under the assumption that E is in short Weierstrass form. This means that $\{x_0^\sigma : \sigma \in \text{Gal}(K/\mathbb{Q})\}$ has at most 8 elements. Now we will analyze the extension $\mathbb{Q}(x_0)$, which is obtained by extending the rational numbers by the x -coordinate of the point P . The order of the set we defined as the orbit of x -coordinate of P under Galois actions is equal to the extension degree of $\mathbb{Q}(x_0)$, which is a divisor of n since $\mathbb{Q}(x_0) \subset K$. This means that set can not have order 3 or 4. If it has order at most 2, this means that $\mathbb{Q}(x_0) \subset \mathbb{Q}(\sqrt{d})$. Then, the extension degree of $\mathbb{Q}(x_0)$ cannot be 3, 4, 6, 8 because we assumed n is not divisible by 3 and 4. This leaves only the possibilities 1, 2, 5, 7 for the extension degree. If it is 1 or 2, we will proceed with the point P in rest of the proof. Otherwise, there must be another point $Q = (x_1, y_1) \in E(K)_{\text{tors}}$ of order 16 such that $x_1 \neq x_0^\sigma$ for any $\sigma \in \text{Gal}(K/\mathbb{Q})$. Then,

the sets $\{x_0^\sigma : \sigma \in \text{Gal}(K/\mathbb{Q})\}$ and $\{x_1^\sigma : \sigma \in \text{Gal}(K/\mathbb{Q})\}$ are disjoint and are subsets of the set of x -coordinates of the 16-torsion points. This implies that number of elements in those two orbits, does not exceed 8 in total. If the first orbit has 5 or 7 elements, then the second orbit has at most 3 elements. It also cannot have exactly 3 elements since n is not divisible by 3. Ultimately, the second orbit has at most two elements, so the extension degree of $\mathbb{Q}(x_1)$ is at most 2. In this case we can work with Q instead of P .

Without loss of generality, assume that $\mathbb{Q}(x_0)$ has extension degree at most 2. We know that $[\mathbb{Q}(x_0, y_0) : \mathbb{Q}(x_0)]$ is at most 2. This means that the extension degree of $\mathbb{Q}(x_0, y_0) = \mathbb{Q}(P)$ is a divisor of 4. It is also a divisor of n . This means that the extension degree of $\mathbb{Q}(P)$ is at most 2.

Now, consider the extension $\mathbb{Q}(E[2])$ which is the extension obtained by the coordinates of the 2-torsion points. Assume that E is in short Weierstrass form, so $\mathbb{Q}(E[2])$ is the splitting field of the cubic $x^3 + Ax + B$ where $A, B \in \mathbb{Q}$. Note that all 2-torsion points belong to K so $\mathbb{Q}(E[2])$ is a subfield of K , hence the extension degree of $\mathbb{Q}(E[2])$ is a divisor of n . If the cubic is irreducible, then the extension degree of $\mathbb{Q}(E[2])$ is either 3 or 6, which implies n is divisible by 3, but this contradicts our assumptions. Therefore, the cubic must have a rational root and the extension degree of $\mathbb{Q}(E[2])$ is at most 2.

We have shown that there must be a point $P \in E(K)_{\text{tors}}$ of order 16 such that $\mathbb{Q}(P)$ is an extension of degree at most 2. We also showed that the extension $\mathbb{Q}(E[2])$ has degree at most 2. Then the extension degree of $\mathbb{Q}(E[2], P)$ must be a divisor of 4. It also divides n , so the extension degree is at most 2. This proves that all 2-torsion points and P belongs to a subfield $\mathbb{Q}(\sqrt{d})$ of K where d is a square-free integer (possibly $d = 1$). Finally, we can observe that, all points in $E(K)_{\text{tors}}$ can be generated by 2-torsion point and a point of order 16. Therefore, $E(K)_{\text{tors}} = E(\mathbb{Q}(\sqrt{d}))_{\text{tors}}$. But this contradicts with the Theorem 2.2. □

Now, we will proceed to the proof of our main theorem.

Proof of Theorem 1.1. Since $\mathbb{Q}(\zeta_p)$ is an abelian Galois number field of degree $p - 1$, we know that $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ is isomorphic to one of the groups given in the Corollary 2.3. Recall that, the groups there are all possible subgroups of the groups that can be realized as $E(\mathbb{Q}^{ab})_{\text{tors}}$. In our proof, we will show that most of those groups are not realizable.

Firstly, let us prove that $\mathbb{Z}/N\mathbb{Z} \times \mathbb{Z}/N\mathbb{Z}$ cannot be a subgroup of $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$, when $N = 3, 4, 5, 6, 8$. Due to Weil pairing, if $\mathbb{Z}/N\mathbb{Z} \times \mathbb{Z}/N\mathbb{Z}$ is a subgroup of $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$, then $\zeta_N \in \mathbb{Q}(\zeta_p)$. This is not the case for $N = 3, 4, 5, 6, 8$, since $p \geq 11$ is a prime.

As a combined result of Lemma 3.2, by assuming $p - 1$ is not divisible by 3, 4, 5, 7, 11, we can use all items of the lemma except the result about $\mathbb{Z}/15\mathbb{Z}$. This result requires more conditions. We know that when p is a prime, $\mathbb{Q}(\zeta_p)$ has a unique quadratic subfield. If $p \equiv 1 \pmod{p}$, that unique quadratic subfield is $\mathbb{Q}(\sqrt{p})$. If $p \equiv 3 \pmod{p}$, as in our theorem, the unique quadratic subfield is $\mathbb{Q}(\sqrt{-p})$. This means that, $\mathbb{Q}(\sqrt{5})$ and $\mathbb{Q}(\sqrt{-15})$ are not subfields of $\mathbb{Q}(\zeta_p)$, so we can use that result of the lemma too. Then, we have shown that $E(\mathbb{Q}(\zeta_p))_{\text{tors}} \not\cong \mathbb{Z}/N\mathbb{Z}$ for $N = 11, 13, 15, 17, 19, 21, 25, 27, 37, 43, 67, 163$.

By combining Lemma 3.2 and Lemma 3.4, we can show that $E(\mathbb{Q}(\zeta_p))_{\text{tors}} \not\cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z}$ for $N = 7, 8, 9$. Only remaining possibilities for $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ are the groups in the Mazur's theorem and the three remaining groups given in the statement of Theorem 1.1. □

4. REALIZATION OF TORSION SUBGROUPS

In this section, we will discuss the realization of the three torsion subgroups appearing in Theorem 1.1, which are different from those in Mazur's theorem. Our first result shows that we must look for these torsion subgroups in the quadratic subfield. The next two theorems show how to find the desired torsion subgroups in quadratic fields.

Theorem 4.1. *Let E/\mathbb{Q} be an elliptic curve. Let $p > 3$ be a prime such that $p - 1$ is not divisible by 3, 4, 5, 7, 11. If $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ is one of the groups $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ or $\mathbb{Z}/16\mathbb{Z}$, then $E(\mathbb{Q}(\zeta_p))_{\text{tors}} = E(\mathbb{Q}(\sqrt{-p}))_{\text{tors}}$.*

Proof. Firstly, let $E(\mathbb{Q}(\zeta_p))_{\text{tors}} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2N\mathbb{Z}$ for $N = 7, 9$. Only quadratic subfield of $\mathbb{Q}(\zeta_p)$ is $\mathbb{Q}(\sqrt{-p})$. Combining this fact with Lemma 3.2, we obtain $E(\mathbb{Q}(\sqrt{-p}))[N] = E(\mathbb{Q}(\zeta_p))[N]$. Assume that E is in short Weierstrass form, then $\mathbb{Q}(E[2])$ is the splitting field of the cubic $x^3 + Ax + B$. Since $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ has full 2-torsion, the field $\mathbb{Q}(E[2])$ must be a subfield of $\mathbb{Q}(\zeta_p)$. Then the cubic cannot be irreducible since $p - 1$ is not divisible by 3. This means that either cubic has three rational roots or it has one rational root and two other roots from $\mathbb{Q}(\sqrt{-p})$. This means that $E(\mathbb{Q}(\sqrt{-p}))_{\text{tors}}$ has also full 2-torsion. Ultimately, $E(\mathbb{Q}(\zeta_p))_{\text{tors}} = E(\mathbb{Q}(\sqrt{-p}))_{\text{tors}}$.

Secondly, let $E(\mathbb{Q}(\zeta_p))_{\text{tors}} \cong \mathbb{Z}/16\mathbb{Z}$. Let $P = (x_0, y_0) \in E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ be a point of order 16. Then, extension degree of $\mathbb{Q}(x_0)$ is equal to the number of elements in the set $\{x_0^\sigma : \sigma \in \text{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{Q})\}$ and also it divides n . In total there is exactly 8 point of order 16 in the $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ and their x -coordinates form a set of 4 elements. This means that $\{x_0^\sigma : \sigma \in \text{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{Q})\}$ has at most 4 elements. It cannot have 3 or 4 elements because we assumed n is not divisible by them. This means that $\mathbb{Q}(x_0)$ is at most degree 2 extension so $x_0 \in \mathbb{Q}(\sqrt{-p})$. Also, $[\mathbb{Q}(x_0, y_0) : \mathbb{Q}(x_0)] \leq 2$ so extension degree of $\mathbb{Q}(x_0, y)$ is a divisor of 4, but it also divides n so it is at most 2 degree extension. This implies $P \in E(\mathbb{Q}(\sqrt{-p}))_{\text{tors}}$. Since P generates $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$, we obtain $E(\mathbb{Q}(\zeta_p))_{\text{tors}} = E(\mathbb{Q}(\sqrt{-p}))_{\text{tors}}$. \square

This theorem shows that if $E(\mathbb{Q}(\zeta_p))_{\text{tors}}$ is not one of the groups in Mazur's theorem, then we should only consider the quadratic extension and look for torsion points here. Those three groups appears in the Theorem 2.2, so we know that they are realizable over infinitely many elliptic curves. But we are interested about whether they are realizable over the quadratic fields $\mathbb{Q}(\sqrt{-p})$ for the primes p such that $p - 1$ is not divisible by 3, 4, 5, 7, 11. By using the next two theorems we can show that they are realizable over that quadratic fields for some prime p .

Theorem 4.2. *(Jeon, Kim, Lee, [7], Theorem 3.2) Put $d = d(t) = 8t^3 - 8t^2 + 1$ with $t \in \mathbb{Q}$. Let E be an elliptic curve defined by the equation*

$$E : y^2 + (1 - c)xy - by = x^3 - bx^2$$

where $b = \frac{t^3(2t^2 - 3t + 1)}{(t^2 - 3t + 1)^2}$ and $c = -\frac{t(2t^2 - 3t + 1)}{t^2 - 3t + 1}$ with $t \neq 0, \frac{1}{2}, 1$. Then the torsion subgroup of E over $\mathbb{Q}(\sqrt{d})$ is $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$.

In this theorem, if we choose $t = -75/242$, then we get $d = -59 \times 14^2/11^6$ which means that E has torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$ over $\mathbb{Q}(\sqrt{-59})$. This torsion subgroup is a subgroup of $E(\mathbb{Q}(\zeta_{59}))_{\text{tors}}$. By Theorem 1.1, we can observe that $E(\mathbb{Q}(\zeta_{59}))_{\text{tors}} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$ since it cannot be a larger group.

Theorem 4.3. (Jeon, Kim, Lee, [7], Theorem 3.3) Put $d = d(t) = \frac{t^2-1}{t^2+3}$ with $t \in \mathbb{Q}$. Let E be an elliptic curve defined by the equation

$$E : y^2 + (1 - c)xy - (c + c^2)y = x^3 - (c + c^2)x^2$$

where $c = \frac{1-t^2}{t^4+3t^2}$ with $t \neq -1, 0, 1$. Then the torsion subgroup of E over $\mathbb{Q}(\sqrt{d})$ is equal to $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$.

In this theorem, if we choose $t = \frac{248354}{307104}$, then we get $d = -47(\frac{5^2 \times 17 \times 31}{37 \times 7933})^2$. This means that our curve described above has torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ over $\mathbb{Q}(\sqrt{-47})$. This torsion subgroup is a subgroup of $E(\mathbb{Q}(\zeta_{47}))_{\text{tors}}$. By Theorem 1.1, we can observe that $E(\mathbb{Q}(\zeta_{47}))_{\text{tors}} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ since it cannot be a larger group.

These two elliptic curves show that the group structures $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$ and $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ are realizable. However, note that we have shown they are realizable for specific values of p . We do not know if they are realizable for all such primes p .

Realization of $\mathbb{Z}/16\mathbb{Z}$ as torsion subgroup structure problem is remains open. We know that, in order to make it realized, we need to find a prime p with $E(\mathbb{Q}(\sqrt{-p}))_{\text{tors}} \cong \mathbb{Z}/16\mathbb{Z}$. Due to Theorem 1.3 in Banwait and Derickx [16], we know that it is not realizable for any primes $p < 800$ satisfying $p \equiv 3 \pmod{4}$, which includes the primes in the statement of our Theorem 1.1. For the primes $p \equiv 1 \pmod{4}$, we are looking to see if there is any prime p with $E(\mathbb{Q}(\sqrt{p}))_{\text{tors}} \cong \mathbb{Z}/16\mathbb{Z}$. It is realizable for $p = 41$ and that is the only prime $p < 800$ with the desired torsion subgroup. As an example, the elliptic curve 419430bm5 satisfies the desired condition. This combined with Corollary 2.3, implies that for some rational elliptic curve E , the torsion subgroup $E(\mathbb{Q}(\zeta_{41}))_{\text{tors}}$ is isomorphic to $\mathbb{Z}/16\mathbb{Z}$.

REFERENCES

- [1] M. Chou, *Torsion of rational elliptic curves over the maximal abelian extension of \mathbb{Q}* , Pacific J. Math. 302 (2019), No. 2, 481–509.
- [2] B. Mazur, *Rational isogenies of prime degree*, Inventiones Math. 44 (1978), 129 - 162.
- [3] F. Najman, *Torsion of rational elliptic curves over cubic fields and sporadic points on $X_1(n)$* , Math. Res. Letters, 23 (2016) 245-272.
- [4] M. Chou, *Torsion of rational elliptic curves over quartic Galois number fields*, J. Number Theory 160 (2016), 603-628.
- [5] T. Gužvić, I. Krijan, *Torsion groups of elliptic curves over some infinite abelian extensions of \mathbb{Q}* , submitted.
- [6] T. Gužvić, B. Vukorepa, *Torsion groups of elliptic curves over $\mathbb{Q}(\mu_{p^\infty})$* , submitted.
- [7] D. Jeon, C. H. Kim and Y. Lee, *Families of elliptic curves over cubic number fields with prescribed torsion subgroups*, Math. Comp. 80 (2011), 579-591.
- [8] S. Kamienny, *Torsion points on elliptic curves and q -coefficients of modular forms*, Invent. Math. 109 (1992), 221-229.
- [9] M.A. Kenku, F. Momose, *Torsion points on elliptic curves defined over quadratic fields*, Nagoya Math. J. 109 (1988), 125-149.
- [10] M. Derickx, A. Etropolski, J. Morrow, M. van Hoeij, D. Zureick-Brown, *Sporadic torsion on elliptic curves*, preprint.
- [11] E. González-Jiménez, F. Najman, *Growth of torsion groups of elliptic curves upon base change*, preprint.
- [12] Y. Fujita, *Torsion subgroups of elliptic curves in elementary abelian 2-extensions of \mathbb{Q}* , J. Number Theory 114 (2005), 124-134.
- [13] H. Daniels, Á. Lozano-Robledo, F. Najman, A. Sutherland, *Torsion subgroups of rational elliptic curves over the compositum of all cubic fields*, submitted, to appear in Math. Comp.
- [14] K. Ribet, *Torsion points of abelian varieties in cyclotomic extensions*, Enseign. Math. 27, 315-319 (1981).
- [15] L. Merel, *Bornes pour la torsion des courbes elliptiques sur les corps de nombres*, Invent. Math. 124 (1996), no. 1-3, 437-449.

- [16] B.S. Banwait, M. Derickx, *Torsion subgroups of elliptic curves over quadratic fields and a conjecture of Granville*, preprint.

DEPT. OF MATHEMATICS, BOGAZICI UNIVERSITY, ISTANBUL, TURKEY
Email address: omeravci372742@gmail.com