

ON THE TOPOLOGY AND GEOMETRY OF CERTAIN 13-MANIFOLDS

WEN SHEN

ABSTRACT. This paper gives the classifications of certain manifolds \mathcal{M} of dimension 13 up to diffeomorphism, homeomorphism, and homotopy equivalence, whose cohomology rings are isomorphic to $H^*(\mathbb{C}P^3 \times S^7; \mathbb{Z})$. Moreover, we prove that either \mathcal{M} or $\mathcal{M} \# \Sigma^{13}$ admits a metric of non-negative sectional curvature where Σ^{13} is a certain exotic sphere of dimension 13.

1. INTRODUCTION

Classification of the manifolds with a given cohomology ring (up to diffeomorphism, homeomorphism, or homotopy) is one of the central problems in geometric topology. The Poincaré theorem is a special case. For a more complicated cohomology ring, there also exist many investigations. In the general case, Wall [26, 28] studied $(s - 1)$ -connected $2s$ -manifolds and $(s - 1)$ -connected $(2s + 1)$ -manifolds. For concrete cases with specified dimensions, we also list some results. Barden [3] classified simply connected 5-manifolds, and Kreck and Su [19] classified certain nonsimply connected 5-manifolds. Wall [27], Jupp [12] classified simply connected 6-manifolds. For the dimension 7, Kreck and Stolz [17, 18] gave a classification for the 7-manifolds modeled on Aloff-Wallach spaces [2]. Subsequently, a homotopy classification of Aloff-Wallach spaces was given by Kruggel [20]. Besides, there are other classifications for the 7-manifolds with certain cohomology rings, such as [6] [16]. More recently, Fang and Shen [7] considered the 13-manifolds, named Bazaikin spaces [4].

In this paper, we consider manifolds that look like $\mathbb{C}P^3 \times S^7$. More precisely, the manifold, denoted by \mathcal{M} , is a 1-connected closed smooth manifold with the same cohomology ring as $\mathbb{C}P^3 \times S^7$, where $\mathbb{C}P^3$ is the complex projective space. Many examples of these 13-manifolds arise from the following construction [30]: From an S^1 action on $S^7 \times S^7$ given by $\theta \circ (x, y) = (e^{il\theta}x, e^{-ik\theta}y)$, $(k, l) = 1$, we get the S^1 -bundle

2020 *Mathematics Subject Classification.* Primary 57R19, 57R50, 53C20.

$S^1 \rightarrow S^7 \times S^7 \rightarrow \mathcal{M}$. In addition, Kerin [13] gave an example with almost positive curvature, which is an S^7 -bundle over $\mathbb{C}P^3$.

Then, we make some conventions. Throughout the rest of the paper, let $H^i(-)$ (or $H_i(-)$) denote the integral cohomology (or homology) group $H^i(-; \mathbb{Z})$ (or $H_i(-; \mathbb{Z})$) unless other statements are given. We always choose the cup product x^2 as the generator of $H^4(\mathcal{M})$, where x is a generator of $H^2(\mathcal{M})$.

Definition 1.1. Let $x : \mathcal{M} \rightarrow \mathbb{C}P^\infty$ denote the generator x of $H^2(\mathcal{M})$. We call that \mathcal{M} admits a *restriction lift* if there exists a lift $f : \mathcal{M} \rightarrow \mathbb{C}P^4$ such that the following diagram is homotopy commutative

$$\begin{array}{ccc} & & \mathbb{C}P^4 \\ & \nearrow f & \downarrow \\ \mathcal{M} & \xrightarrow{x} & \mathbb{C}P^\infty \end{array}$$

Now, we are ready to present the main theorems of this paper.

Theorem 1.2. Assume that $\mathcal{M}, \mathcal{M}'$ are nonspin and admit *restriction lifts*. Then

- 1) \mathcal{M} is diffeomorphic to $\mathcal{M}' \# \Sigma^{13}$ if and only if $p_1(\mathcal{M}) = p_1(\mathcal{M}')$. If $p_1(\mathcal{M}) = p_1(\mathcal{M}') \neq 0 \pmod{3}$, \mathcal{M} is diffeomorphic to \mathcal{M}' .
- 2) \mathcal{M} is homeomorphic to \mathcal{M}' if and only if $p_1(\mathcal{M}) = p_1(\mathcal{M}')$.
- 3) \mathcal{M} is homotopy equivalent to \mathcal{M}' if and only if $p_1(\mathcal{M}) = p_1(\mathcal{M}') \pmod{24}$.

Applying the 1) and 2) of Theorem 1.2, we have

Corollary 1.3. If \mathcal{M} is nonspin and admits a *restriction lift*, then the underlying topological manifold of \mathcal{M} is the total space of a topological S^7 -bundle over $\mathbb{C}P^3$; if \mathcal{M} additionally satisfies $p_1(\mathcal{M}) \neq 0 \pmod{3}$, then it is the total space of a S^7 -bundle over $\mathbb{C}P^3$.

On the one hand, many geometers use curvatures to detect the topology of manifolds. On the other hand, one of the most interesting problems in Riemannian geometry is whether a smooth manifold admits a metric of non-negative sectional curvature or positive sectional curvature. Here, we use the topology of manifolds to investigate the metrics. From Theorem 1.2, Corollary 1.3 and Lemma 7.1, we have

Theorem 1.4. Let \mathcal{M} be nonspin and admit a *restriction lift*. If $p_1(\mathcal{M}) \neq 0 \pmod{3}$, \mathcal{M} admits a metric of non-negative sectional curvature; otherwise, $\mathcal{M} \# \Sigma^{13}$ admits a metric of non-negative sectional curvature for some homotopy sphere Σ^{13} .

For the general spin case, the same theorem as Theorem 1.2 is unclear. However, for some special spin manifolds, we have

Theorem 1.5. For any homotopy 7-spheres Σ^7 and Σ'^7 , $\mathbb{C}P^3 \times \Sigma^7$ is diffeomorphic to $\mathbb{C}P^3 \times \Sigma'^7$.

Obviously, $\mathbb{C}P^3 \times \Sigma^7 \stackrel{\text{diff}}{\cong} \mathbb{C}P^3 \times S^7$. Thus we have

Corollary 1.6. For any homotopy sphere Σ^7 , $\mathbb{C}P^3 \times \Sigma^7$ admits a metric of non-negative sectional curvature.

We mainly use the surgery theory [15] to prove the classification theorems, i.e. Theorem 1.2 and 1.5. In Section 2, we establish a suitable bordism theory for the considered spaces in the smooth category. In Section 3, we consider the surgery obstruction. Then, we compute a certain bordism group in Section 4. To classify the considered manifolds up to homotopy equivalence, we give a model in Section 5. Finally, the proofs of the classification theorems are presented in Section 6. At last, we discuss the metrics of the sphere bundles of some 8-dimensional vector bundles in Section 7.

2. THE NORMAL 5-SMOOTHING

Let $\pi : B \rightarrow \text{BO}$ be a fibration over the classifying space BO . As described in [15], given a smooth manifold M with the Gauss map $\mathcal{N}_M : M \rightarrow \text{BO}$, a normal B -structure of M is a lift $\tilde{\mathcal{N}}_M : M \rightarrow B$ of \mathcal{N}_M . Moreover, $\tilde{\mathcal{N}}_M$ is called a normal ℓ -smoothing when it is a $(\ell + 1)$ homotopy equivalence.

Let \mathcal{M} admit a *restriction lift* $f : \mathcal{M} \rightarrow \mathbb{C}P^4$. Denote the first Pontryagin class of \mathcal{M} by $p_1(\mathcal{M}) = sx^2$ $s \in \mathbb{Z}$, the second Stiefel-Whitney class by $w_2(\mathcal{M}) = sx \pmod{2}$. Let \mathcal{H} denote the Hopf bundle over $\mathbb{C}P^4$. If $s \geq 0$, let ξ be the complementary bundle of the Whitney sum $s\mathcal{H}$; if $s < 0$, let ξ be the Whitney sum $-s\mathcal{H}$. Formally $\xi = -s\mathcal{H}$. Note that the Chern classes

$$c_1(\xi) = -sx \quad c_2(\xi) = \frac{s(s+1)}{2}x^2,$$

the first Pontryagin class $p_1(\xi) = -sx^2$. Let $\mathcal{N}_{\mathcal{M}}$ denote both the Gauss map $\mathcal{M} \rightarrow \text{BO}$ and the stable normal bundle of \mathcal{M} . Obviously the difference bundle $\mathcal{N}_{\mathcal{M}} - f^*\xi$ has a $\text{BO}\langle 8 \rangle$ -structure. Considering the classifying map

$$\pi : B = \text{BO}\langle 8 \rangle \times \mathbb{C}P^4 \rightarrow \text{BO}$$

of the bundle $\gamma_8 \times \xi$ where γ_8 is the universal bundle over $\text{BO}\langle 8 \rangle$, we claim that \mathcal{M} has a normal B -structure, i.e., the Gauss map $\mathcal{N}_{\mathcal{M}}$ lifts to $\tilde{\mathcal{N}}_{\mathcal{M}} : \mathcal{M} \rightarrow B$ which is furthermore a 7-equivalence.

All n -dimensional manifolds with the normal B -structure form a bordism group $\Omega_n^{O(8)}(\xi)$ which is isomorphic to the homotopy group $\pi_n(\text{MB})$ by the Pontryagin-Thom isomorphism. Here, $\text{MB} = \text{MO}\langle 8 \rangle \wedge \text{M}\xi$; $\text{M}\xi$ and $\text{MO}\langle 8 \rangle$ are the Thom spectra of the bundles ξ and γ_8 .

3. SURGERY OBSTRUCTION

Assume that the W^{14} has the $B = \text{BO}\langle 8 \rangle \times \text{CP}^4$ -structure with the boundary $\mathcal{M} \sqcup -\mathcal{M}'$ which is the disjoint union. From surgery theory [29], we can take appropriate W such that $\phi : W \rightarrow B$ is a 7-equivalence, and the embedding $i : \mathcal{M} \rightarrow W$ induces the isomorphisms $H_n(\mathcal{M}) \cong H_n(W)$ for $0 \leq n \leq 6$. From the long exact sequences for the pair (W, \mathcal{M}) and the triple $(W, \partial W, \mathcal{M})$, we have that $H_7(W) \cong H_7(W, \mathcal{M}) \oplus \mathbb{Z}$ is torsion free.

Considering the surgery obstruction for replacing W with an h -cobordism, we first transform the elements in $H_7(W, \mathcal{M})$ into the kernel of the homomorphism $\phi_* : \pi_7(W) \rightarrow \pi_7(B)$.

$$\begin{array}{ccccccc} \pi_8(B) = \mathbb{Z} & \longrightarrow & \pi_8(B, W) & \longrightarrow & \pi_7(W) & \xrightarrow{\phi_*} & \pi_7(B) = 0 \\ & & \downarrow & \cong & \downarrow h_* & & \downarrow \\ H_8(W) = 0 & \longrightarrow & H_8(B) = \mathbb{Z}^2 & \longrightarrow & H_8(B, W) & \longrightarrow & H_7(W) \longrightarrow H_7(B) = 0 \end{array}$$

By chasing the above commutative diagram, we have

Lemma 3.1. $\ker \phi_* = \pi_7(W) \cong H_7(W) \oplus \mathbb{Z} \cong H_7(W, \mathcal{M}) \oplus \mathbb{Z}^2$.

By smooth embedding theorem [22], every element $u \in H_7(W, \mathcal{M})$ can be represented by the embedding $S^7 \rightarrow W$. From [14], any 7-dimensional vector bundle over S^7 is trivial. Hence, the normal bundle of this embedding is trivial. Thus, u can be represented by the embedding $\bar{u} : S^7 \times D^7 \rightarrow W$. Let F be the homotopy fiber of $\pi : B \rightarrow \text{BO}$. Due to $\pi_7(F) = 0$, the embedding \bar{u} is a compatible embedding. So, we can replace the W with an h -cobordism.


Lemma 3.2. If \mathcal{M} is B -cobordant to \mathcal{M}' , then there is a smooth h -cobordism $(W, \mathcal{M}, \mathcal{M}')$.

4. THE COMPUTATIONS OF CERTAIN BORDISM GROUPS

Recall that $\Omega_*^{O(8)}(\xi) \cong \pi_*(\text{MB})$ where $\text{MB} = \text{MO}\langle 8 \rangle \wedge \text{M}\xi$, $\text{M}\xi$ and $\text{MO}\langle 8 \rangle$ are the Thom spectra of the bundles ξ and γ_8 .

There is an Atiyah-Hirzebruch spectral sequence (AHSS) for $\pi_*(\text{MB})$

$$E_2^{p,q} \cong H_p(\text{M}\xi; \pi_q(\text{MO}\langle 8 \rangle)) \implies \text{MO}\langle 8 \rangle_*(\text{M}\xi) \cong \pi_*(\text{MO}\langle 8 \rangle \wedge \text{M}\xi).$$


 FIG 1. $\text{Ext}_{\mathcal{A}_2}^{s,t}(\mathbb{Z}_2, \mathbb{Z}_2)$

Recall the E_2 -terms of the Adams spectral sequence (ASS) for $\text{MO}\langle 8 \rangle$ as in Fig.1. In dimension $i \leq 14$ [8, 11], $\pi_i(\text{MO}\langle 8 \rangle)$ is as follows.

i	0	1	2	3	4	5	6	7	8	9
$\pi_i(\text{MO}\langle 8 \rangle)$	\mathbb{Z}	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_{24}	0	0	\mathbb{Z}_2	0	$\mathbb{Z} \oplus \mathbb{Z}_2$	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$

i	10	11	12	13	14
$\pi_i(\text{MO}\langle 8 \rangle)$	\mathbb{Z}_6	0	\mathbb{Z}	\mathbb{Z}_3	\mathbb{Z}_2

In the AHSS for $\Omega_{13}^{\text{O}(8)}(\xi)$, all nontrivial E_2 -terms are $E_2^{0,13} = \mathbb{Z}_3$ and $E_2^{4,9} = \mathbb{Z}_2^2$. Let $G = E_\infty^{4,9}$. Obviously the 2-primary part ${}_2\pi_{13}(\text{MB})$ is isomorphic to $G = \mathbb{Z}_2^n$ where $n = 0, 1$ or 2 . Next, we will observe the G in the ASS

$$E_2^{s,t}(\text{ASS}) = \text{Ext}_{\mathcal{A}_2}^{s,t}(H^*(\text{MB}; \mathbb{Z}_2), \mathbb{Z}_2) \implies {}_2\pi_{t-s}(\text{MB})$$

$$\text{Ext}^{s,t} = \text{Ext}_{\mathcal{A}_2}^{s,t}(H^*(\text{MB}; \mathbb{Z}_2), \mathbb{Z}_2) \cong \text{Ext}_{\mathcal{A}_2}^{s,t}(H^*(\text{M}\xi; \mathbb{Z}_2), \mathbb{Z}_2)$$

where $H^*(\text{M}\xi; \mathbb{Z}_2) \cong H^*(\text{CP}^4; \mathbb{Z}_2) \cup U$ as \mathbb{Z}_2 -vector space, U is the stable Thom class of the bundle ξ . In particular, for any $a \in H^*(\text{CP}^4; \mathbb{Z}_2)$

$$\text{Sq}^n(aU) = \sum_{i=0}^n \text{Sq}^i a \cup \text{Sq}^{n-i} U$$

$$\text{Sq}^i U = w_i(\xi)U$$

where $w_i(\xi)$ is the i th Stiefel-Whitney of bundle ξ .

By the Algebraic Atiyah-Hirzebruch spectral sequence (AAHSS) (also see [7])

$$E_1^{s,m,n} = \text{Ext}_{\mathcal{A}_2}^{s,s+m}(\mathbb{Z}_2, \mathbb{Z}_2) \otimes H^n(\text{M}\xi; \mathbb{Z}_2) \implies \text{Ext}^{s,s+m+n}$$

we get quickly the results shown in Fig.2 and the following proposition




FIG 2. Partial E_1 -terms of $Ext_{\mathcal{A}_2}^{s,t}(H^*(M\xi; \mathbb{Z}_2), \mathbb{Z}_2)$

Proposition 4.1. More details for the elements of Fig.2 are

- Case 1: $w_2(\xi) = w_4(\xi) = 0$. Only $h_1 c_0 x^2 U, h_0^s h_2 \omega x U$ ($s = 0, 1$) survive to $E_\infty^{*,13-n,n}$.
- Case 2: $w_2(\xi) \neq 0, w_4(\xi) = 0$. Only $h_1 c_0 x^2 U$ and $h_1 \omega x^2 U$ do not survive to $E_\infty^{*,13-n,n}$.
- Case 3: $w_2(\xi) = 0, w_4(\xi) \neq 0$. Only $h_1 c_0 x^2 U$ survive to $E_\infty^{*,13-n,n}$.
- Case 4: $w_2(\xi) \neq 0, w_4(\xi) \neq 0$. Only $h_1 c_0 x^2 U$ and $h_1 \omega x^2 U$ do not survive to $E_\infty^{*,13-n,n}$.

For all cases, $\tau x U \in E_\infty^{*,14-n,n}$ in the AAHSS.

Proof. Let d_r be the differential of the AAHSS

$$d_r : E_r^{s,m,n} \rightarrow E_r^{s+1,m+r-1,n-r}$$

All $E_1^{s,m,n}$ -terms are shown in Fig.2 for $n + m = 12, 13$.

Case 1: By $\text{Sq}^2(xU) = x^2U$, we have $d_2(h_1^s \omega x^2 U) = h_1^{s+1} \omega x U$ for $s = 0, 1$. Since $h_0^2 h_2 \omega x U = h_1^3 \omega x U$, $d_2(h_1^2 \omega x^2 U) = h_1^3 \omega x U$. By the degrees of differentials d_r , we have $E_\infty^{*,13-n,n} = E_3^{*,13-n,n}$.

Case 2: By $\text{Sq}^2(x^2U) = x^3U$, we have $d_2(c_0 x^3 U) = h_1 c_0 x^2 U$ and $d_2(\omega x^3 U) = h_1 \omega x^2 U$.

Similarly, Cases 3 and 4 follows from the Sq^2, Sq^4 actions. By the degrees of differentials d_r , $h_0^s \tau x U \in E_\infty^{3+s,14-n,n}$. \square

The Fig.2 and Proposition 4.1 show $Ext_{\mathcal{A}_2}^{s,13+s}(H^*(M\xi; \mathbb{Z}_2), \mathbb{Z}_2)$ and $Ext_{\mathcal{A}_2}^{s,12+s}(H^*(M\xi; \mathbb{Z}_2), \mathbb{Z}_2)$ for $s \geq 0$ completely. Some nontrivial Adams differentials are given in the following lemma.

Lemma 4.2. For cases 1, 2, 4, $d_2(\tau xU) = h_2\omega xU$ in the ASS.

Proof. Let $\xi|_2$ be the restriction bundle over $\mathbb{C}P^1$ of ξ , $M\xi|_2$ be its Thom spectrum. On the one hand, by the AHSS

$$E_2^{p,q}(|_2) = H_p(M\xi|_2; \pi_q(\text{MO}\langle 8 \rangle)) \implies \pi_{p+q}(\text{MO}\langle 8 \rangle \wedge M\xi|_2)$$

we have ${}_2\pi_{13}(\text{MO}\langle 8 \rangle \wedge M\xi|_2) = 0$. On the other hand, for all cases as described in Proposition 4.1, $Ext_{\mathcal{A}_2}^{5+s,18+s}(H^*(M\xi|_2; \mathbb{Z}_2), \mathbb{Z}_2)$ is generated by $h_0^s h_2 \omega xU$ ($s \geq 0$); $Ext_{\mathcal{A}_2}^{3,17}(H^*(M\xi|_2; \mathbb{Z}_2), \mathbb{Z}_2)$ is generated by τxU , and $Ext_{\mathcal{A}_2}^{s,14+s}(H^*(M\xi|_2; \mathbb{Z}_2), \mathbb{Z}_2) = 0$ ($0 \leq s \leq 2$). Thus $d_2(\tau xU) = h_2\omega xU$ in the ASS for the spectrum $\text{MO}\langle 8 \rangle \wedge M\xi|_2$. By the naturality and Proposition 4.1, the desired result follows. \square

Summarizing Fig.2, Proposition 4.1, and Lemma 4.2, we have

Lemma 4.3. For cases 1, 3, ${}_2\pi_{13}(\text{MB}) = \mathbb{Z}_2\langle h_1 c_0 x^2 U \rangle$.

For cases 2, 4, ${}_2\pi_{13}(\text{MB}) = 0$.

Finally, we consider the 3-primary part ${}_3\pi_{13}(\text{MB})$.

Lemma 4.4. (1) ${}_3\pi_{13}(\text{MB}) \cong \mathbb{Z}_3$ or 0.

(2) A nontrivial element with order 3 in $\Omega_{13}^{O(8)}(\xi)$ is represented by an exotic sphere Σ^{13} .

Proof. By the AHSS, it is easy to get (1). Let $\iota : S^0 \rightarrow M\xi$ be the unit. Then, we have the morphism of the spectral sequences

$$\begin{array}{ccc} \bar{E}_2^{p,q} = H_p(S^0; \pi_q(\text{MO}\langle 8 \rangle)) & \implies & \pi_{p+q}(\text{MO}\langle 8 \rangle) \\ \downarrow & & \downarrow \iota_* \\ E_2^{p,q} = H_p(M\xi; \pi_q(\text{MO}\langle 8 \rangle)) & \implies & \pi_{p+q}(\text{MO}\langle 8 \rangle \wedge M\xi) \end{array}$$

If $E_\infty^{0,13} = E_2^{0,13}$, then a nontrivial element with order 3 in $\Omega_{13}^{O(8)}(\xi)$ lies in the image of ι_* . By [14], an exotic sphere Σ^{13} is a generator of $\pi_{13}(\text{MO}\langle 8 \rangle)$. So, the (2) follows. \square

Now, we calculate ${}_3\pi_{13}(\text{MB})$ clearly by the Adams spectral sequence.

Lemma 4.5. If $p_1(\xi) \neq 0 \pmod{3}$, ${}_3\pi_{13}(\text{MB}) = 0$.

Proof. From [1] and [11], $\pi_{13}(\text{MO}\langle 8 \rangle) = \mathbb{Z}_3$ is generated by

$$b_0 h_{1,0} \in Ext_{\mathcal{A}}^{3,16}(H^*(\text{MO}\langle 8 \rangle; \mathbb{Z}_3), \mathbb{Z}_3)$$

in the ASS, where \mathcal{A} is the mod 3 Steenrod algebra. By the proof of the (2) of Lemma 4.4, $b_0 h_{1,0} U$ is nontrivial in $Ext_{\mathcal{A}}^{3,16}(H^*(\text{MB}; \mathbb{Z}_3), \mathbb{Z}_3)$ if ${}_{3\pi_{13}}(\text{MB}) = \mathbb{Z}_3$, where U is the stable Thom class of $M\xi$. Note that

- (1) $\mathcal{P}^1 U = p_1(\xi)U \pmod{3}$ where \mathcal{P}^1 is the reduced Steenrod power in the mod 3 Steenrod algebra \mathcal{A} [21].
- (2) $h_{1,0}$ is the dual of \mathcal{P}^1 .

If $p_1(\xi) \neq 0 \pmod{3}$, then $\delta(b_0 p_1(\xi)U) = b_0 h_{1,0} U$ in the cobar complex of $Ext_{\mathcal{A}}^{*,*}(H^*(\text{MB}; \mathbb{Z}_3), \mathbb{Z}_3)$. In other words, $b_0 h_{1,0} U$ is trivial. Hence, the desired result follows from the (1) of Lemma 4.4. \square

5. THE S^7 -FIBRATIONS OVER CP^3

For a fibration $E \rightarrow B$ with fiber F , we say “a fibration $F \rightarrow E \rightarrow B$ ” instead of “an F -fibration”. For the S^{n-1} -fibrations, there exists a classifying space BG_n . Furthermore, by the natural sequence

$$\cdots \rightarrow \text{BG}_{n-1} \rightarrow \text{BG}_n \xrightarrow{r_n^G} \text{BG}_{n+1} \rightarrow \cdots$$

we have the telescope BG , where the homotopy fiber of $r_n^G : \text{BG}_n \rightarrow \text{BG}_{n+1}$ is $(n-2)$ -connected [23]. By taking the sphere bundle, we have a natural map $\alpha_n : \text{BO}_n \rightarrow \text{BG}_n$, which induces the map $\alpha : \text{BO} \rightarrow \text{BG}$. The classifying spaces BO and BG for the stable bundle theories are infinite loop spaces and the natural map between them is an infinite loop map by theorems of Boardman and Vogt [5]. In fact, they are homotopy equivalent to the 0th spaces in the Ω -spectra of the multiplicative cohomology theories KO and KG respectively. Thus, there is a natural map between the Ω -spectra KO and KG , which is induced by $\alpha : \text{BO} \rightarrow \text{BG}$. In addition, the homotopy fiber G/O of $\alpha : \text{BO} \rightarrow \text{BG}$ is also an infinite loop space.

Lemma 5.1. The following items hold:

- (1) $\widetilde{KO}^0(\text{CP}^3) \cong [\text{CP}^3, \text{BO}_8]$, $\widetilde{KG}^0(\text{CP}^3) \cong [\text{CP}^3, \text{BG}_8]$.
- (2) $\widetilde{KO}^0(\text{CP}^3) \rightarrow \widetilde{KG}^0(\text{CP}^3)$ is an epimorphism.

Proof. Let $V = O, G$. The homotopy fiber of $r_{8*}^V : \text{BV}_8 \rightarrow \text{BV}$ is 6-connected at least. Then, it is easy to get (1).

It is well known that $\pi_1(G/O) = \pi_3(G/O) = \pi_5(G/O) = 0$ [25]. By the obstruction theory, any map from CP^3 to BG can be lifted to BO . Thus, the (2) follows. \square

By Sanderson [24], we have $\widetilde{KO}^0(\text{CP}^2) = \mathbb{Z}$ and $\widetilde{KO}^0(\text{CP}^3) \cong \mathbb{Z}$. Furthermore, the stable class of a vector bundle over CP^3 is determined by its first Pontrjagin class.

Lemma 5.2. Let η, η' be any 8-dimensional vector bundles over $\mathbb{C}P^3$. The S^7 -bundle of η is fiber homotopy equivalent to the S^7 -bundle of η' if $p_1(\eta) = p_1(\eta') \pmod{24}$.

Proof. For the cofibration sequence $S^2 \rightarrow \mathbb{C}P^2 \rightarrow S^4 \rightarrow \Sigma S^2$, we have the following exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & [S^4, \text{BO}] & \longrightarrow & [\mathbb{C}P^2, \text{BO}] & \longrightarrow & [S^2, \text{BO}] & \longrightarrow & 0 \\ \downarrow & & \downarrow \alpha_4 & & \downarrow & & \downarrow \alpha_2 & & \downarrow \\ [S^3, \text{BG}] & \xrightarrow{\rho_3} & [S^4, \text{BG}] & \longrightarrow & [\mathbb{C}P^2, \text{BG}] & \longrightarrow & [S^2, \text{BG}] & \xrightarrow{\rho_2} & \dots \end{array}$$

Note that α_2 is an isomorphism, $\alpha_4 : \mathbb{Z} \rightarrow \mathbb{Z}_{24}$ is an epimorphism, and $[\mathbb{C}P^2, \text{BO}] = \mathbb{Z}$. By [31], $\rho_3 : \mathbb{Z}_2 \rightarrow \mathbb{Z}_{24}$ is a monomorphism. Therefore, $[\mathbb{C}P^2, \text{BO}] = \mathbb{Z} \rightarrow [\mathbb{C}P^2, \text{BG}] = \mathbb{Z}_{24}$ is an epimorphism. By the exact sequences for the cofibration sequence $\mathbb{C}P^2 \rightarrow \mathbb{C}P^3 \rightarrow S^6 \rightarrow \Sigma \mathbb{C}P^2$, we have that $[\mathbb{C}P^3, \text{BO}] = \mathbb{Z} \rightarrow [\mathbb{C}P^3, \text{BG}] = \mathbb{Z}_{24}$ is an epimorphism. So, the desired result follows. \square

6. PROOFS OF THEOREMS 1.2 AND 1.5

Proof of Theorem 1.2: For any nonspin manifold \mathcal{M} with a *restriction lift*, the corresponding bordism group $\pi_{13}(\text{MB})$ as in Section 2 satisfies the case 2 or 4 as in Lemma 4.3. Let \mathcal{M}' be another considered manifold so that $p_1(\mathcal{M}) = p_1(\mathcal{M}')$. Then, they lie in the same bordism group.

By Lemma 4.4 and 3.2, \mathcal{M} is diffeomorphic to $\mathcal{M}' \# \Sigma^{13}$. Especially, by Lemma 4.5, \mathcal{M} is diffeomorphic to \mathcal{M}' if $p_1(\mathcal{M}) \neq 0 \pmod{3}$.

In the topological category, we also have the same lemma as Lemma 3.2. So \mathcal{M} is homeomorphic to \mathcal{M}' by Lemma 4.4.

Note that the first Pontrjagin class of the stable class of the standard Hopf bundle over $\mathbb{C}P^3$ is $x^2 \in H^4(\mathbb{C}P^3)$. Hence, for any $k \in \mathbb{Z}$, there exists an 8-dimensional vector bundle η over $\mathbb{C}P^3$ so that $p_1(\eta) = kx^2$. Besides, the total space $\overline{\mathcal{M}}$ of the S^7 -bundle of η has the first Pontrjagin class $(4+k)x^2$. By the 1) and 2) of Theorem 1.2, Corollary 1.3 follows. By Lemma 5.2 and Corollary 1.3, \mathcal{M} is homotopy equivalent to \mathcal{M}' if $p_1(\mathcal{M}) = p_1(\mathcal{M}') \pmod{24}$. Furthermore, $p_1 \pmod{24}$ is a homotopy invariant [10]. \square

Proof of Theorem 1.5: Since $\Omega_7^{\text{O}(8)} \cong \pi_7(\text{MO}\langle 8 \rangle)$, there exists an 8-manifold W with $\text{BO}\langle 8 \rangle$ -structure so that $\partial W = \Sigma^7$ for any homotopy 7-sphere Σ^7 . $\mathbb{C}P^3 \times \Sigma^7$ has a normal B -structure as in Section 2. It is obvious that $\mathbb{C}P^3 \times W$ also has a normal B -structure. Thus, $\mathbb{C}P^3 \times \Sigma^7$ equals to $\mathbb{C}P^3 \times \Sigma^7$ as the bordism class in $\Omega_{13}^{\text{O}(8)}(\xi)$. Hence, Theorem 1.5 follows by Lemma 3.2. \square

7. A METRIC OF NON-NEGATIVE SECTIONAL CURVATURE

The real K -theory of projective spaces are well known, see [24]. The ring $\widetilde{KO}(\mathbb{C}P^n)$ is generated by $r(\mathcal{H}) - 2$, where \mathcal{H} is the complex canonical line bundle over $\mathbb{C}P^n$, $r : K \rightarrow KO$. Let $n = 3$. Note that

$$\mathbb{C}P^3 = U(4)/(U(1) \times U(3)) = G/H$$

and $r(\mathcal{H})$ is homogeneous, i.e. it is a bundle of the form $(G \times \mathbb{R}^2)/H$, where H acts on \mathbb{R}^2 by means of a real linear representation $\rho \in \text{Rep}_{\mathbb{R}}(H)$. In particular, $\rho(H)$ lies in $O(2)$.

Since $[\mathbb{C}P^3, \text{BO}(8)] \cong \widetilde{KO}(\mathbb{C}P^3)$, any 8-dimensional vector bundle over $\mathbb{C}P^3$ is homogeneous. Then we take its sphere bundle, i.e. it is a bundle of the form $(G \times S^7)/H$, where H acts on S^7 by means of a real linear representation ρ' such that $\rho'(H)$ lies in $O(8)$.

Lemma 7.1. Let η be any 8-dimensional vector bundle over $\mathbb{C}P^3$. The total space of the S^7 -bundle of η admits a metric of non-negative sectional curvature.

Proof. As in [9], we can endow $G \times S^7$ with the product metric of \langle, \rangle_G and the standard sphere metric. By O'Neill's Theorem on Riemannian submersions, the lemma follows. \square

REFERENCES

- [1] Aikawa, T. 3-dimensional cohomology of the mod p Steenrod algebra, Math. Scand. 47 (1980), 91-115.
- [2] Aloff, S., Wallach, N. An infinite family of 7-manifolds admitting positively curved Riemannian structures. Bull. Am. Math. Soc. 81 (1975), 93-97.
- [3] Barden, D. Simply connected five-manifolds. Ann. of Math. (2) 82 (1965), 365-385.
- [4] Bazaikin, Y. V. On a certain family of closed 13-dimensional Riemannian manifolds of positive curvature. Sib. Math. J. 37, No. 6 (1996), 1219-1237.
- [5] Boardman, J. M.; Vogt, R. Homotopy invariant algebraic structures, Lecture Notes in Mathematics 347 Springer-Verlag (1973).
- [6] Crowley, D.; Nordström, J. The classification of 2-connected 7-manifolds. Proc. Lond. Math. Soc. (3) 119 (2019), no. 1, 1-54.
- [7] Fang, F.; Shen, W. Topological classification of Bazaikin spaces. appearing. <https://doi.org/10.48550/arXiv.2307.13303>.
- [8] Giambalvo, V. On $\langle 8 \rangle$ -cobordism. Ill. J. Math. 15 (1971), 533-541.
- [9] González-Álvarez, D. Nonnegative curvature on stable bundles over compact rank one symmetric spaces. Adv. Math. 307 (2017), 53-71.
- [10] Hirzebruch, F. A Riemann-Roch theorem for differentiable manifolds. Séminaire Bourbaki, expose 177, 1959.
- [11] Hovey, M. A.; Ravenel, D. C. The 7-connected cobordism ring at $p=3$. Trans. Amer. Math. Soc. 347 (1995), no. 9, 3473-3502.

- [12] Jupp, P. E. Classification of certain 6-manifolds. Proc. Cambridge Philos. Soc. 73 (1973), 293-300.
- [13] Kerin, M. Some new examples with almost positive curvature. Geom. Topol. 15 (2011), no. 1, 217-260.
- [14] Kervaire, M. A.; Milnor, J. W. Groups of homotopy spheres. I. Ann. of Math. (2) 77 (1963), 504-537.
- [15] Kreck, M. Surgery and duality. Ann. of Math. (2) 149 (1999), no. 3, 707-754.
- [16] Kreck, M. On the classification of 1-connected 7-manifolds with torsion free second homology. J. Topol. 11 (2018), no. 3, 720-738.
- [17] Kreck, M.; Stolz, S. A diffeomorphism classification of 7-dimensional homogeneous Einstein manifolds with $SU(3) \times SU(2) \times U(1)$ -symmetry. Ann. of Math. (2) 127 (1988), no. 2, 373-388.
- [18] Kreck, M.; Stolz, S. Some nondiffeomorphic homeomorphic homogeneous 7-manifolds with positive sectional curvature. J. Differential Geom. 33 (1991), no. 2, 465-486.
- [19] Kreck, M.; Su, Y. On 5-manifolds with free fundamental group and simple boundary links in S^5 . Geom. Topol. 21 (2017), no. 5, 2989-3008.
- [20] Kruggel, B. A homotopy classification of certain 7-manifolds. Trans. Amer. Math. Soc. 349 (1997), no. 7, 2827-2843.
- [21] Milnor, J. W. Characteristic classes, Mimeographed Notes, Princeton Univ, Princeton, N. J., 1957.
- [22] Milnor, J. W. Lectures on the cobordism theorem. Notes by L. Siebenmann and J. Sondow Princeton University Press, Princeton, N.J. 1965.
- [23] Rudyak, Y. B. On Thom Spectra, Orientability, and Cobordism, Springer Monographs in Mathematics, Corrected 2nd printing, Springer, 2008.
- [24] Sanderson, B. J. Immersions and embeddings of projective spaces. Proc. London Math. Soc. (3) 14 (1964), 137-153.
- [25] Sullivan, D. On the Hauptvermutung for manifolds. Bull. Amer. Math. Soc. 73 (1967), 598-600.
- [26] Wall, C. T. C. Classification of $(n-1)$ -connected $2n$ -manifolds. Ann. of Math. (2) 75 (1962), 163-189.
- [27] Wall, C. T. C. Classification problems in differential topology. V. On certain 6-manifolds. Invent. Math. 1 (1966), 355-374.
- [28] Wall, C. T. C. Classification problems in differential topology, VI: Classification of $(s-1)$ -connected $(2s+1)$ -manifolds, Topology 6 (1967) 273-296.
- [29] Wall, C. T. C. Surgery on Compact Manifolds, London Math. Soc. Monographs 1, Academic Press, New York, 1970.
- [30] Wang, M. Y.; Ziller, W. Einstein metrics on principal torus bundles. J. Differential Geom. 31 (1990), no. 1, 215-248.
- [31] Wang, X. On the classification of certain 1-connected 7-manifolds and related problems. Q. J. Math. 73 (2022), no. 2, 711-727.

DEPARTMENT OF MATHEMATICS, CAPITAL NORMAL UNIVERSITY, BEIJING, P.R.CHINA

Email address: shenwen121212@163.com