

# WEIGHTED HARDY-RELLICH TYPE INEQUALITIES: IMPROVED BEST CONSTANTS AND SYMMETRY BREAKING

CRISTIAN CAZACU AND IRINA FIDEL

ABSTRACT. When studying the weighted Hardy-Rellich inequality in  $L^2$  with the full gradient replaced by the radial derivative the best constant becomes trivially larger or equal than in the first situation. Our contribution is to determine the new sharp constant and to show that for some part of the weights is strictly larger than before. In some cases we emphasize that the extremals functions of the sharp constant are not radially symmetric.

*2020 Mathematics Subject Classification:* 35A23, 35R45, 35Q40, 35B09, 34A40, 34K38.

*Key words:* Hardy-Rellich inequalities, optimal constants, spherical harmonics.

## 1. INTRODUCTION

The celebrated  $L^2$  *Hardy inequality* (e.g. [10, 13]) states that for  $N \geq 3$  and  $u \in C_c^\infty(\mathbb{R}^N)$  it holds

$$(1) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq C_H \int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx, \quad C_H(N) := \frac{(N-2)^2}{4},$$

where the constant  $C_H(N)$  is sharp.

*Rellich inequality* (e.g. [14]) asserts that for  $N \geq 5$  and  $u \in C_c^\infty(\mathbb{R}^N)$  we have

$$(2) \quad \int_{\mathbb{R}^N} |\Delta u|^2 dx \geq C_R(N) \int_{\mathbb{R}^N} \frac{|u|^2}{|x|^4} dx, \quad C_R(N) := \left( \frac{N(N-4)}{4} \right)^2,$$

with the best constant  $C_R(N)$ . The Hardy and Rellich inequalities are important tools widely used in the analysis of partial differential operators and equations of harmonic and biharmonic-type.

The *Hardy-Rellich inequality* has been studied more recently (see, e.g. [15, 7, 4]). This is in fact an improved Hardy inequality (with a larger optimal constant) applied to classes of vector fields originated from potential gradients, which arises in fluid mechanics. For  $N \geq 3$  and  $u \in C_0^\infty(\mathbb{R}^N)$  this leads to

$$(3) \quad \int_{\mathbb{R}^N} |\Delta u|^2 dx \geq C_{HR}(N) \int_{\mathbb{R}^N} \frac{|\nabla u|^2}{|x|^2} dx,$$

with the best constant

$$C_{HR}(N) := \begin{cases} \frac{N^2}{4}, & N \geq 5 \\ 3, & N = 4 \\ \frac{25}{36}, & N = 3. \end{cases}$$

The Hardy-Rellich inequality was firstly extended in [15] to more general singular weights of the form

$$(4) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx \geq C(N, m) \int_{\mathbb{R}^N} |\nabla u|^2 |x|^{m-2} dx, \quad \forall u \in C_c^\infty(\mathbb{R}^N),$$

where the authors proved that for any  $N \geq 5$  and any  $4 - N < m \leq 0$  the best constant is

$$(5) \quad C(N, m) = \min_{k=0,1,2,\dots} \frac{\left( \frac{(N-4+m)(N-m)}{4} + k(N+k-2) \right)^2}{\left( \frac{N-4+m}{2} \right)^2 + k(N+k-2)}.$$

In particular, according to the computations in [15, pag. 453], if  $\frac{N+4-2\sqrt{N^2-N+1}}{3} \leq m \leq 0$  then

$$C(N, m) = \left( \frac{N-m}{2} \right)^2$$

otherwise, if  $4 - N < m < \frac{N+4-2\sqrt{N^2-N+1}}{3}$  then

$$C(N, m) < \left( \frac{N-m}{2} \right)^2.$$

To ensure the integrability of the singular term in inequality (4) we need to impose that  $|x|^{m-2} \in L_{loc}^1(\mathbb{R}^N)$  which gives us the constraint

$$(6) \quad m > 2 - N \quad (\text{or } m + N - 2 > 0).$$

The weighted inequality (4) was later extended in [7] to all the cases  $N \geq 1$  and  $m > 2 - N$ . Optimal constants of the cases which were not covered in [15] were solved in [7, Theorem 6.1]. Next, we emphasize a brief presentation of these additional cases:

- If  $N = 1$  and  $m \in (1, \frac{7}{3}] \cup [3, \infty)$  then  $C(1, m) = \left( \frac{1-m}{2} \right)^2$ .
- If  $N = 1$  and  $m \in (\frac{7}{3}; 3)$  then  $C(1, m) \leq \left( \frac{1-m}{2} \right)^2$  (there are values  $m$  for which the inequality is strict).
- If  $N \geq 1$  and  $m = 4 - N$  then  $C(N, m) = \min\{(N-2)^2, N-1\}$ .
- If  $N \geq 2$  and  $\frac{N+4-2\sqrt{N^2-N+1}}{3} \leq m$  then  $C(N, m) = \left( \frac{N-m}{2} \right)^2$ .
- If  $2 \leq N \leq 3$  and  $2 - N < m < \frac{N+4-2\sqrt{N^2-N+1}}{3}$ , or  $N \geq 4$  and  $2 - N < m \leq 4 - N$ , then  $C(N, m) = \frac{\left( \frac{(N-4+m)(N+m)}{4} + N-1 \right)^2}{\left( \frac{N-4+m}{2} \right)^2 + N-1}$

- If  $N = 3$  and  $m \geq \frac{N+4-2\sqrt{N^2-N+1}}{3}$  or  $N \geq 4$  and  $m > 4 - N$  the best constant requires a further subdivision, based on very technical expressions, which could be consulted in [7, Th. 6.1, pag. 52].

Subsequent extensions of the weighted Hardy-Rellich type inequalities with remainder terms have been done recently in [16] and [17] by applying factorization methods. Also recent improvements when adding magnetic fields have been established in [3, 11]. See also very recent results on the Hardy-Rellich inequalities in [1, 12] and in [9] (in the context of solenoidal vector fields) and the references there in.

Overall for any  $N \geq 1$  and  $m > 2 - N$  always happens that the best constant in (4) does not pass the threshold

$$(7) \quad C(N, m) \leq \left( \frac{N - m}{2} \right)^2.$$

In this paper we study a new weighted Hardy-Rellich type inequality which, to our knowledge, has not been treated yet in the literature, by replacing the full gradient in (4) with the radial derivative  $\partial_r u := \frac{x}{|x|} \cdot \nabla u$  on the right hand side, namely

$$(8) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx \geq \tilde{C}(N, m) \int_{\mathbb{R}^N} |x \cdot \nabla u|^2 |x|^{m-4} dx, \quad \forall u \in C_c^\infty(\mathbb{R}^N).$$

where  $\tilde{C}(N, m)$  denotes the best constant in (8). Since  $|\partial_r u| \leq |\nabla u|$  notice that inequality (8) is a relaxation of (4) and in balance with that the best constant might be larger or equal, i.e.  $C(N, m) \leq \tilde{C}(N, m)$ .

Our purposes are to supply explicitly the best constant  $\tilde{C}(N, m)$  for the full range of parameters  $N \geq 1$  and  $m > 2 - N$ , to emphasize situations in which we get an improvement in (8) with respect to (4) i.e.  $\tilde{C}(N, m) > C(N, m)$  and to put in evidence the radially symmetry breaking for the optimal approximations of the sharp constants.

## 2. MAIN RESULT

On the purpose to state the main result we need to introduce a cut-off function  $g \in C^\infty([0, \infty))$ , with  $0 \leq g \leq 1$ , such that

$$(9) \quad g(r) = \begin{cases} 1, & 0 \leq r \leq 1, \\ 0, & r \geq 2. \end{cases}$$

The main result of this paper is the following

**Theorem 2.1.** Let  $N \geq 1$  and  $m > 2 - N$ . Then

$$(10) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx \geq \tilde{C}(N, m) \int_{\mathbb{R}^N} |x \cdot \nabla u|^2 |x|^{m-4} dx. \quad \forall u \in C_c^\infty(\mathbb{R}^N),$$

where the optimal constant  $\tilde{C}(N, m)$  is given as follows.

If  $N = 1$  and  $m > 1$  or if  $N \geq 2$  and  $m \in [2 - \sqrt{(N-1)^2 + 1}, 2 + \sqrt{(N-1)^2 + 1}]$ , then

$$\tilde{C}(N, m) = \left( \frac{N-m}{2} \right)^2,$$

which is approximated by the sequence  $\{u_\epsilon\}_{\epsilon>0}$  given by

$$(11) \quad u_\epsilon(x) = |x|^{-\frac{N+m-4}{2}+\epsilon} g(|x|).$$

If  $N \geq 2$  and  $m \in (2 - N, 2 - \sqrt{(N-1)^2 + 1})$ , then

$$\tilde{C}(N, m) = \frac{((m-2)^2 - N^2)^2}{4(N+m-4)^2}$$

which is approximated by the sequence  $\{u_\epsilon\}_{\epsilon>0}$  given by

$$(12) \quad u_\epsilon(x) = |x|^{-\frac{N+m-4}{2}+\epsilon} g(|x|) \phi_1(x),$$

where  $\phi_1$  is a spherical harmonic function of degree 1 with  $\|\phi_1\|_{L^2(S^{N-1})} = 1$ .

If  $N \geq 2$  and  $m \in (2 + \sqrt{(N-1)^2 + 1}, \infty)$ , then

$$\tilde{C}(N, m) = \min_{l \leq k(m)} \tilde{C}(N, m, l), \quad \tilde{C}(N, m, l) := \frac{(-N+m-2l)^2 (2l+m+N-4)^2}{4(m+N-4)^2}$$

where  $k(m)$  is defined later in (31)-(44). The constant  $\tilde{C}(N, m)$  is approximated by the sequence given by

$$(13) \quad u_\epsilon(x) = |x|^{-\frac{N+m-4}{2}+\epsilon} g(|x|) \phi_{l_{\min}}(x),$$

where  $\phi_{l_{\min}}$  is a spherical harmonic function of degree  $l_{\min}$  such that  $\|\phi_{l_{\min}}\|_{L^2(S^{N-1})} = 1$  and  $l_{\min} := \arg \min_{l \leq k(m)} \tilde{C}(N, m, l)$ .

**Remark 2.2.** Notice also that in all the situations above

$$\tilde{C}(N, m) \leq \left( \frac{N-m}{2} \right)^2,$$

but there are cases (see for instance  $N = 1$  and  $m \in (\frac{7}{3}, 3)$ ) when our best constant  $\tilde{C}(N, m)$  in (8) improves with respect to the best constant  $C(N, m)$  in (4).

**Remark 2.3.** The approximating sequences  $u_\epsilon$  in (11), (13), (12) do not belong to the space  $C_c^\infty(\mathbb{R}^N)$  but they are in the energy space of the inequality (10), i.e. both terms in (10) are finite for  $u_\epsilon$ . So, by regularizing  $u_\epsilon$  near the origin one can show that the constants remain sharp for functions  $u \in C_c^\infty(\mathbb{R}^N)$ , see for instance similar arguments in [5, 6].

## 3. PROOF OF THE MAIN RESULT

We will use spherical coordinates instead of cartesian coordinates. This coordinates transformation is given by

$$x \in \mathbb{R}^N \setminus \{0\} \mapsto (r, \sigma) \in (0, \infty) \times S^{N-1}, \quad r = |x|, \quad \sigma = \frac{x}{|x|},$$

where  $S^{N-1}$  is the  $N - 1$ -dimensional sphere with respect to the Hausdorff measure in  $\mathbb{R}^N$ . We will use the formula of the Laplacian in spherical coordinates

$$(14) \quad \Delta = \partial_{rr}^2 + \frac{N-1}{r} \partial_r + \frac{1}{r^2} \Delta_{S^{N-1}},$$

where  $\partial_r$  and  $\partial_{rr}^2$  are first and second order partial derivatives with respect to the radial component  $r$  whereas (for fixed  $r$ ) the Laplace-Beltrami operator with respect to the metric tensor on  $S^{N-1}$  and respectively the spherical gradient are given by

$$\Delta_{S^{N-1}} u(r\sigma) = \Delta \left[ u \left( \frac{x}{|x|} \right) \right]_{|x=\sigma}, \quad \nabla_{S^{N-1}} u(r\sigma) = \nabla \left[ u \left( \frac{x}{|x|} \right) \right]_{|x=\sigma}.$$

Applying the spherical harmonics decomposition we can expand  $u \in C_c^\infty(\mathbb{R}^N)$  as

$$u(x) = u(r\sigma) = \sum_{k=0}^{\infty} u_k(r) \phi_k(\sigma).$$

The set of functions  $\{\phi_k\}_{k \geq 0}$  are spherical harmonics of degree  $k$  which consists in an orthogonal basis in  $L^2(S^{N-1})$ . These functions satisfy the properties

$$(15) \quad \begin{cases} -\Delta_{S^{N-1}} \phi_k = c_k \phi_k \text{ on } S^{N-1}, \\ -\int_{S^{N-1}} \Delta_{S^{N-1}} \phi_k \phi_l d\sigma = \int_{S^{N-1}} \nabla_{S^{N-1}} \phi_k \cdot \nabla_{S^{N-1}} \phi_l d\sigma \\ = c_k \int_{S^{N-1}} \phi_k \phi_l d\sigma = c_k \delta_{lk}, \quad k, l \in \mathbb{N}, \end{cases}$$

where  $c_k = k(k + N - 2)$ ,  $k \geq 0$  are the eigenvalues of the Laplace-Beltrami operator  $\Delta_{S^{N-1}}$ , where  $\delta_{lk}$  represents the Kronecker symbol, see e.g. [8] for more detailed properties of spherical harmonics.

Since in view of (14)

$$\Delta \left[ u_k(|x|) \phi_k \left( \frac{x}{|x|} \right) \right] = \left( \Delta_r u_k(r) - \frac{c_k}{r^2} u_k(r) \right) \phi_k(\sigma),$$

due to (15) similar computations as in [4] lead to

$$(16) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx = \sum_{k=0}^{\infty} \int_0^{\infty} \left( |\Delta_r u_k|^2 + \frac{c_k^2}{r^4} u_k^2 - \frac{2c_k}{r^2} u_k \Delta_r u_k \right) r^{N+m-1} dr.$$

Next, we will write  $u'_k$  and  $u''_k$  to express both first and second derivatives of the Fourier coefficients  $\{u_k\}_k$ .

We need to compute  $\int_0^{\infty} |\Delta_r u_k| r^{N+m-1}$  and  $\int_0^{\infty} \frac{\Delta_r u_k}{r^2} u_k(r) r^{N+m-1} dr$ .

Integration by parts leads to

$$(17) \quad \int_0^\infty |\Delta_r u_k|^2 r^{N+m-1} dr = (N-1)(1-m) \int_0^\infty |u'_k(r)| r^{N+m-3} dr + \\ + \int_0^\infty |u''_k(r)| r^{N+m-1} dr,$$

and

$$(18) \quad \int_0^\infty \frac{\Delta_r u_k}{r^2} u_k(r) r^{N+m-1} dr = \frac{1}{2} \left( -2 \int_0^\infty |u'_k(r)| r^{N+m-3} dr \right) + \\ + (N+m-4)(m-2) \int_0^\infty |u_k(r)| r^{N+m-5} dr.$$

Then (16) becomes

$$(19) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx = \sum_{k=0}^\infty \left( \int_0^\infty |u''_k|^2 r^{N+m-1} dr + \right. \\ \left. + (2c_k + (N-1)(1-m)) \int_0^\infty |u'_k|^2 r^{N+m-3} dr \right. \\ \left. + (c_k^2 - c_k(m-2)(N+m-4)) \int_0^\infty |u_k|^2 r^{N+m-5} dr \right).$$

and more easily, since  $\partial_r u = \frac{x}{|x|} \cdot \nabla u$ , we get

$$(20) \quad \int_{\mathbb{R}^N} |x \cdot \nabla u|^2 |x|^{m-4} dx = \sum_{k=0}^\infty \int_0^\infty |u'_k|^2 r^{N+m-3} dr.$$

**3.1. The case  $N = 1$  and  $m > 1$ .** The inequality (10) reduces to:

$$(21) \quad \int_{\mathbb{R}} |u''|^2 r^m dr \geq \tilde{C}(1, m) \int_{\mathbb{R}} |u'|^2 r^{m-2} dr, \quad \forall u \in C_c^\infty(\mathbb{R}).$$

The relation (21) comes from

$$\begin{aligned} \int_{\mathbb{R}} (u')^2 r^{m-2} dr &= \int_{\mathbb{R}} (u')^2 \left( \frac{r^{m-1}}{m-1} \right)' dr \\ &= -\frac{2}{m-1} \int_{\mathbb{R}} u' u'' r^{m-1} dr \\ &= -\frac{2}{m-1} \int_{\mathbb{R}} u' r^{\frac{m}{2}} u'' r^{\frac{m-2}{2}} dr \\ &\leq -\frac{2}{m-1} \left( \int_{\mathbb{R}} (u'')^2 r^m dr \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}} (u')^2 r^{m-2} dr \right)^{\frac{1}{2}}. \end{aligned}$$

Integration by parts and Cauchy-Schwarz inequality lead that (21) hold with the constant

$$(22) \quad \tilde{C}(1, m) \geq \left( \frac{m-1}{2} \right)^2.$$

**3.2. The case  $N \geq 2$ .** In this case it remains to compare the right hand sides in (19)-(20). We will apply the well-known 1-d weighted Hardy inequalities (see, e.g. [15, page 454], [2, Prop. 2.4])

$$(23) \quad \int_0^\infty |u_k''|^2 r^{N+m-1} dr \geq \left( \frac{N+m-2}{2} \right)^2 \int_0^\infty |u_k'|^2 r^{N+m-3} dr,$$

$$(24) \quad \int_0^\infty |u_k'|^2 r^{N+m-3} dr \geq \left( \frac{N+m-4}{2} \right)^2 \int_0^\infty |u_k|^2 r^{N+m-5} dr.$$

**3.2.1. The case A:**  $(2-m)(N+m-4) \geq 0$ . We distinguish two sub-cases as follows.

**A1)**  $2-m \geq 0$  and  $N+m-4 \geq 0$ . These imply  $4-N \leq m \leq 2$ .

**A2)**  $2-m \leq 0$  and  $N+m-4 \leq 0$ . Since  $N \geq 2$  it follows that  $m = N = 2$ .

Gluing both situations we can summarize that case **A** is equivalent to

$$(25) \quad 4-N \leq m \leq 2.$$

Then we obtain from (19)-(20) and (23) that

$$(26) \quad \begin{aligned} \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx &\geq \sum_{k=0}^\infty \left( \left( \frac{N+m-2}{2} \right)^2 \int_0^\infty |u_k'|^2 r^{N+m-3} dr \right. \\ &\quad \left. + (2c_k + (N-1)(1-m)) \int_0^\infty |u_k'|^2 r^{N+m-3} dr \right) \\ &\geq \left( \left( \frac{N+m-2}{2} \right)^2 + (N-1)(1-m) \right) \times \\ &\quad \times \sum_{k=0}^\infty \int_0^\infty |u_k'|^2 r^{N+m-3} dr \\ &= \left( \frac{N-m}{2} \right)^2 \sum_{k=0}^\infty \int_0^\infty |u_k'|^2 r^{N+m-3} dr \\ &= \left( \frac{N-m}{2} \right)^2 \int_{\mathbb{R}^N} |x \cdot \nabla u|^2 |x|^{m-4} dx. \end{aligned}$$

Hence

$$(27) \quad \tilde{C}(N, m) \geq \left( \frac{N-m}{2} \right)^2.$$

*The case B:*  $(2-m)(N+m-4) < 0$ . **B1)**  $2-m > 0$  **and**  $N+m-4 < 0$ . These combined with (6) are equivalent to

$$(28) \quad 2 - N < m < 4 - N.$$

**B2)**  $m-2 < 0$  **and**  $N+m-4 > 0$ . These are equivalent to

$$(29) \quad m > 2.$$

We first look at the spherical part (which contains the terms multiplied with  $c_k$ ) in the relation (19). Taking into account (24) and the fact that  $c_k \geq 0$  for any  $k \geq 0$  we successively have

$$(30) \quad \begin{aligned} & 2c_k \int_0^\infty |u'_k|^2 r^{N+m-3} dr + (c_k^2 - c_k(m-2)(N+m-4)) \int_0^\infty |u_k|^2 r^{N+m-5} dr \\ & \geq c_k \left( 2 \left( \frac{N+m-4}{2} \right)^2 + c_k - (m-2)(N+m-4) \right) \int_0^\infty |u_k|^2 r^{N+m-5} dr \end{aligned}$$

In view of this let us denote

$$(31) \quad I_k(m, N) := 2 \left( \frac{N+m-4}{2} \right)^2 + c_k - (m-2)(N+m-4).$$

In view of identity (19), Hardy inequalities (23)-(24) and (30)-(31) we get

$$(32) \quad \begin{aligned} \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx & \geq \left( \frac{N-m}{2} \right)^2 \sum_{k=0}^\infty \int_0^\infty |u'_k|^2 r^{N+m-3} dr + \\ & + \sum_{k=0}^\infty c_k I_k(m, N) \int_0^\infty |u_k|^2 r^{N+m-5} dr. \end{aligned}$$

Next we want to investigate when  $I_k(m, N) \geq 0$ . We start with estimating

$$\begin{aligned} I_1(m, N) & = 2 \left( \frac{N+m-4}{2} \right)^2 + N - 1 - (m-2)(N+m-4) \\ & = -\frac{1}{2}m^2 + 2m + \frac{1}{2}N^2 - N - 1. \end{aligned}$$

The equation  $I_1(m, N) = 0$  in the unknown  $m$  has the discriminant  $\Delta = (N-1)^2 + 1 \geq 0$  and the roots

$$m_{1,2} = 2 \pm \sqrt{(N-1)^2 + 1}$$



which imply

$$I_1(m, N) \geq 0, \quad \text{iff } m \in [2 - \sqrt{(N-1)^2 + 1}, 2 + \sqrt{(N-1)^2 + 1}].$$

Therefore, we conclude that in the case **B1**) we have

$$I_1(m, N) \geq 0, \quad \text{iff } m \in [2 - \sqrt{(N-1)^2 + 1}, 4 - N)$$

whereas in the case **B2**) we have

$$I_1(m, N) \geq 0, \quad \text{iff } m \in (2, 2 + \sqrt{(N-1)^2 + 1}].$$

Since  $\{I_k\}$  is an increasing sequence with respect to  $k$  we get that

$$(33) \quad I_k(m, N) \geq 0, \quad \forall k \geq 1,$$

for any  $m$  satisfying

$$(34) \quad m \in [2 - \sqrt{(N-1)^2 + 1}, 4 - N) \cup (2, 2 + \sqrt{(N-1)^2 + 1}].$$

This together with (19) and (32) yield to

$$(35) \quad \tilde{C}(N, m) \geq \left(\frac{N-m}{2}\right)^2, \quad \forall m \text{ as in (34)}.$$

It remains to analyze the complementary "bad cases" of (34) for which  $I_1(m, N) < 0$ :

$$(36) \quad m \in \underbrace{(2 - N, 2 - \sqrt{(N-1)^2 + 1})}_{\text{remaining cases of B1)} \cup \underbrace{(2 + \sqrt{(N-1)^2 + 1}, \infty)}_{\text{remaining cases of B2)}.$$

If  $k = 2$  we obtain

$$\begin{aligned} I_2(m, N) &= 2 \left(\frac{N+m-4}{2}\right)^2 + 2N - (m-2)(N+m-4) \\ &= -\frac{m^2}{2} + 2m + \frac{N^2}{2}. \end{aligned}$$

Therefore

$$(37) \quad I_2(m, N) \geq 0, \quad \text{iff } m \in [2 - \sqrt{N^2 + 4}, 2 + \sqrt{N^2 + 4}].$$

This leads to

$$(38) \quad I_k(m, N) \geq 0, \quad \forall k \geq 2, \quad m \in (2 - N, 2 - \sqrt{(N-1)^2 + 1}),$$

which cover the remaining cases of **B1**).

Inequality (10) in the remaining cases of B1):  $m \in (2 - N, 2 - \sqrt{(N - 1)^2 + 1})$ . The right hand side in (19) can be bounded from below in terms of a parameter  $\varepsilon > 0$  (which will be well precised later) as follows, in view of (32) and (24):

$$\begin{aligned}
\int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx &\geq \sum_{k=0, k \neq 1}^{\infty} \left( \left( \frac{N-m}{2} \right)^2 \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr + \right. \\
&\quad \left. + c_k I_k(m, N) \int_0^{\infty} |u_k|^2 r^{N+m-5} dr \right) + \\
&\quad + \left( \frac{N-m}{2} \right)^2 \int_0^{\infty} |u'_1|^2 r^{N+m-3} + c_1 I_1(m, N) \int_0^{\infty} |u_1|^2 r^{N+m-5} dr \\
&\geq \sum_{k=0, k \neq 1}^{\infty} \left( \frac{N-m}{2} \right)^2 \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr + \\
&\quad + \sum_{k=0, k \neq 1}^{\infty} c_k I_k(m, N) \int_0^{\infty} |u_k|^2 r^{N+m-5} dr + \\
&\quad + \left( \left( \frac{N-m}{2} \right)^2 - \varepsilon \right) \int_0^{\infty} |u'_1|^2 r^{N+m-3} + \\
&\quad + \left( (2c_1 + \varepsilon) \left( \frac{N+m-4}{2} \right)^2 \right. \\
(39) \quad &\quad \left. + c_1^2 - c_1(m-2)(m+N-4) \right) \int_0^{\infty} |u_1|^2 r^{N+m-5} dr.
\end{aligned}$$

The coefficient of the integral term  $\int_0^{\infty} |u_1|^2 r^{N+m-5} dr$  in (39) becomes

$$\begin{aligned}
(40) \quad a_1(N, m, \varepsilon) &:= \left( (2(N-1) + \varepsilon) \left( \frac{N+m-4}{2} \right)^2 \right. \\
&\quad \left. + (N-1)^2 - (N-1)(m-2)(m+N-4) \right).
\end{aligned}$$

We choose  $\varepsilon_1 > 0$  such that  $a_1(N, m, \varepsilon_1) = 0$  and we get

$$(41) \quad \varepsilon_1 = \frac{2(N-1)(m^2 - N^2 - 4m + 2N + 2)}{(m+N-4)^2}.$$

Indeed  $\varepsilon_1 > 0$  because the inequality  $m^2 - N^2 - 4m + 2N + 2 > 0$  holds iff

$$m \in (-\infty, 2 - \sqrt{(N-1)^2 + 1}) \cup (2 + \sqrt{(N-1)^2 + 1}, \infty),$$

set which contains the remaining cases  $m \in (2 - N, 2 - \sqrt{(N - 1)^2 + 1})$  of the case **B1**). From (41) we obtain that

$$\left(\frac{N - m}{2}\right)^2 - \varepsilon_1 = \frac{((m - 2)^2 - N^2)^2}{4(N + m - 4)^2} > 0.$$

Coming back to (39) we have

$$(42) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx \geq \frac{((m - 2)^2 - N^2)^2}{4(N + m - 4)^2} \sum_{k=0}^{\infty} \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr,$$

and therefore

$$(43) \quad \tilde{C}(N, m) \geq \frac{((m - 2)^2 - N^2)^2}{4(N + m - 4)^2}.$$

Inequality (10) in the remaining cases of B2):  $m \in (2 + \sqrt{(N - 1)^2 + 1}, \infty)$ . We want to apply the same idea as in the previous case. For any  $m \in (2 + \sqrt{(N + 1)^2 + 1}, \infty)$  we define the number

$$(44) \quad k(m) := \min \{k \in \mathbb{N} \mid I_k(m, N) \geq 0\}.$$

Then

$$(45) \quad \begin{aligned} I_1(m, N) &< I_2(m, N) < \dots < I_{k(m)-1}(m, N) < 0 \\ I_k(m, N) &\geq I_{k(m)}(m, N) \geq 0, \text{ for all } k \geq k(m). \end{aligned}$$

The inequality (32) will be rewritten as:

$$\begin{aligned} \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx &\geq \left(\frac{N - m}{2}\right)^2 \left( \sum_{k \geq k(m)} \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr + \int_0^{\infty} |u'_0|^2 r^{N+m-3} dr \right) \\ &+ \sum_{k=1}^{k(m)-1} c_k I_k(m, N) \int_0^{\infty} |u_k|^2 r^{N+m-5} dr + \\ &= \left(\frac{N - m}{2}\right)^2 \left( \sum_{k \geq k(m)} \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr + \int_0^{\infty} |u'_0|^2 r^{N+m-3} dr \right) + \\ &+ \sum_{k=1}^{k(m)-1} \left( \left(\left(\frac{N - m}{2}\right)^2 - \varepsilon_k\right) \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr \right. \\ &\left. + c_k I_k(m, N) \int_0^{\infty} |u_k|^2 r^{N+m-5} dr + \varepsilon_k \int_0^{\infty} |u'_k|^2 r^{N+m-3} dr \right). \end{aligned}$$

Then

$$\begin{aligned}
\int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx &\geq \left( \frac{N-m}{2} \right)^2 \left( \sum_{k \geq k(m)} \int_0^\infty |u'_k|^2 r^{N+m-3} dr + \int_0^\infty |u'_0|^2 r^{N+m-3} dr \right) + \\
&+ \sum_{k=1}^{k(m)-1} \left( \left( \left( \frac{N-m}{2} \right)^2 - \varepsilon_k \right) \int_0^\infty |u'_k|^2 r^{N+m-3} dr + \right. \\
(46) \quad &\left. + \left( c_k I_k(m, N) + \varepsilon_k \left( \frac{N+m-4}{2} \right)^2 \right) \int_0^\infty |u_k|^2 r^{N+m-5} dr \right).
\end{aligned}$$

We will denote the coefficient of the zero order term above by

$$(47) \quad a_k(N, m, \varepsilon_k) := c_k I_k(m, N) + \varepsilon_k \left( \frac{N+m-4}{2} \right)^2.$$

We choose  $\varepsilon_k$  such that  $a_k(N, m, \varepsilon_k) = 0$  and we obtain

$$(48) \quad \varepsilon_k = \frac{-4c_k I_k(m, N)}{(N+m-4)^2} > 0.$$

For consistency, we want to make sure that  $\left( \frac{N-m}{2} \right)^2 - \varepsilon_k$  is positive for every  $k, m$  and  $N$ . Indeed,

$$\begin{aligned}
(49) \quad \left( \frac{N-m}{2} \right)^2 - \varepsilon_k &= \left( \frac{N-m}{2} \right)^2 + \frac{4c_k I_k(m, N)}{(N+m-4)^2} \\
&= \frac{(-N+m-2k)^2 (2k+m+N-4)^2}{4(m+N-4)^2} > 0.
\end{aligned}$$

As a consequence of (46) we get

$$(50) \quad \int_{\mathbb{R}^N} |\Delta u|^2 |x|^m dx \geq \min_{l \leq k(m)} \left( \left( \frac{N-m}{2} \right)^2 - \varepsilon_l \right) \sum_{k=0}^\infty \int_0^\infty |u'_k|^2 r^{N+m-3} dr,$$

where  $\varepsilon_l$  is as in (48). So, in view of (49) we have

$$\tilde{C}(N, m) \geq \min_{l \leq k(m)} \tilde{C}(N, m, l), \quad \tilde{C}(N, m, l) := \frac{(-N+m-2l)^2 (2l+m+N-4)^2}{4(m+N-4)^2}.$$

**3.2.2. Optimality.** In this section we aim to prove that all the lower bound constants obtained in (22), (27), (35), (43) and (49) are the sharp constants. For that it suffices to prove the existence of approximating sequences in (10) for the quoted constants.

### Step I. The cases with radially symmetric approximations:

- $N = 1$  and  $m > 1$
- $N \geq 2$  and  $m$  as in the case **A**) (condition (25))

- $N \geq 2$  and  $m$  as in the "good" cases **B**) (condition (34))

The above cases can be treated similarly because the same sequence with radial symmetry can be built to approach the constants (22), (27) and (35).

To prove that let us consider the radially symmetric sequence

$$u_\epsilon(x) = |x|^{-\frac{N+m-4}{2}+\epsilon} g(|x|) = r^{-\frac{N+m-4}{2}+\epsilon} g(r) =: U_\epsilon(r)$$

with  $g$  given in (9). Replacing  $u$  with  $u_\epsilon$  in (19) and arguing as in [4], since the spherical part is missing we obtain

$$\begin{aligned} \int_{\mathbb{R}^N} |\Delta u_\epsilon|^2 |x|^m dx &= \left( \int_0^\infty r^{N-1+m} |U_\epsilon''(r)|^2 dr + \right. \\ &\quad \left. + (N-1)(1-m) \int_0^\infty r^{N+m-3} |U_\epsilon'(r)|^2 dr \right) \end{aligned}$$

and

$$\int_{\mathbb{R}^N} |x \cdot \nabla u_\epsilon|^2 |x|^{m-4} dx = \int_0^\infty |U_\epsilon'(r)|^2 r^{N+m-3} dr.$$

From the definition of  $U_\epsilon$  we have

$$\begin{aligned} &\int_0^\infty r^{N+m-3} |U_\epsilon'(r)|^2 dr = \\ &= \int_0^\infty r^{N+m-3} \left( \left( -\frac{N+m-4}{2} + \epsilon \right)^2 r^{-(N+m-2)+2\epsilon} g^2(r) + r^{-(N+m-4)+2\epsilon} g'(r)^2 \right) dr \\ &+ \int_0^\infty r^{N+m-3} \left( 2 \left( -\frac{N+m-4}{2} \right) + 2\epsilon \right) r^{-(N+m-3)+2\epsilon} g'(r) g(r) dr \\ (51) \quad &= \frac{1}{2\epsilon} \left( -\frac{N-4+m}{2} + \epsilon \right)^2 + \mathcal{O}(1). \end{aligned}$$

Also since

$$U_\epsilon''(r) = \left( -\frac{N-4+m}{2} + \epsilon \right) \left( -\frac{N-2+m}{2} + \epsilon \right) r^{-\frac{N+m}{2}+\epsilon} g(r) + \mathcal{O}(1),$$

we obtain

$$\begin{aligned} &\int_0^\infty r^{N+m-1} |U_\epsilon''(r)|^2 dr = \\ (52) \quad &= \frac{1}{2\epsilon} \left( -\frac{N+m-4}{2} + \epsilon \right)^2 \left( -\frac{N+m-2}{2} + \epsilon \right)^2 + \mathcal{O}(1). \end{aligned}$$

Due to (51) and (52) we obtain

$$\begin{aligned}
& \frac{\int_{\mathbb{R}^N} |\Delta u_\epsilon|^2 |x|^m dx}{\int_{\mathbb{R}^N} |x \cdot \nabla u_\epsilon|^2 |x|^{m-4} dx} = \\
&= \frac{\frac{1}{2\epsilon} \left( \left( -\frac{N+m-4}{2} + \epsilon \right)^2 \left( -\frac{N+m-2}{2} + \epsilon \right)^2 + (N-1)(1-m) \left( -\frac{N+m-4}{2} + \epsilon \right)^2 \right) + \mathcal{O}(1)}{\frac{1}{2\epsilon} \left( -\frac{N+m-4}{2} + \epsilon \right)^2 + \mathcal{O}(1)} \\
&= \frac{\left( -\frac{N+m-4}{2} + \epsilon \right)^2 \left( \left( -\frac{N+m-2}{2} + \epsilon \right)^2 + (N-1)(1-m) \right) + \mathcal{O}(\epsilon)}{\left( -\frac{N+m-4}{2} + \epsilon \right)^2 + \mathcal{O}(\epsilon)} \\
&\searrow \left( \frac{N+m-2}{2} \right)^2 + (N-1)(1-m) = \frac{N^2 - 2Nm + m^2}{4} = \left( \frac{N-m}{2} \right)^2,
\end{aligned}$$

as  $\epsilon \searrow 0$ .

**Step II. The "bad" cases of B) and non radially symmetric optimal approximations:**

- $N \geq 2$  and  $m$  as in the "bad" cases of **B1)**  
(i.e.  $m \in (2-N, 2 - \sqrt{(N-1)^2 + 1})$ )
- $N \geq 2$  and  $m$  as in the "bad" cases of **B2)**  
(i.e.  $m \in (2 + \sqrt{(N-1)^2 + 1}, \infty)$ )

In the first "bad" case  $N \geq 2$  and  $m \in (2-N, 2 - \sqrt{(N-1)^2 + 1})$  we consider the sequence

$$u_\epsilon(x) = |x|^{-\frac{N+m-4}{2} + \epsilon} g(|x|) \phi_1 \left( \frac{x}{|x|} \right) = r^{-\frac{N+m-4}{2} + \epsilon} g(r) \phi_1(\sigma) =: U_\epsilon(r) \phi_1(\sigma)$$

with  $g$  as in (9).

Replace once more  $u$  with the above  $u_\epsilon$  in (19) we obtain

$$\begin{aligned}
\int_{\mathbb{R}^N} |\Delta u_\epsilon|^2 |x|^m dx &= \left( \int_0^\infty r^{N-1+m} |U_\epsilon''(r)|^2 dr + \right. \\
&\quad \left. + (2c_1 + (N-1)(1-m)) \int_0^\infty r^{N+m-3} |U_\epsilon'(r)|^2 dr + \right. \\
&\quad \left. + (c_1^2 - c_1(m-2)(N+m-4)) \int_0^\infty r^{N+m-5} |U_\epsilon(r)|^2 dr \right)
\end{aligned}$$

and

$$\int_{\mathbb{R}^N} |x \cdot \nabla u_\epsilon|^2 |x|^{m-4} dx = \int_0^\infty |U_\epsilon'(r)|^2 r^{N+m-3} dr.$$

Due to (51), (52) and the fact that

$$(53) \quad \int_0^\infty r^{N+m-5} |U_\epsilon(r)|^2 dr = \frac{1}{2\epsilon} + \mathcal{O}(1),$$

we have

$$\begin{aligned}
\int_{\mathbb{R}^N} |\Delta u_\epsilon|^2 |x|^m dx &= \frac{1}{2\epsilon} \left( \left( -\frac{N+m-4}{2} + \epsilon \right)^2 \left( -\frac{N+m-2}{2} + \epsilon \right)^2 + \right. \\
&\quad \left. + (2c_1 + (N-1)(1-m)) \left( -\frac{N+m-4}{2} + \epsilon \right)^2 + \right. \\
(54) \quad &\quad \left. + c_1^2 - c_1(m-2)(N+m-4) \right) + \mathcal{O}(1)
\end{aligned}$$

and

$$(55) \quad \int_{\mathbb{R}^N} |x \cdot \nabla u_\epsilon|^2 |x|^{m-4} dx = \frac{1}{2\epsilon} \left( -\frac{N+m-4}{2} + \epsilon \right)^2 + \mathcal{O}(1).$$

Due to (54) and (55) we successively obtain

$$\begin{aligned}
&\frac{\int_{\mathbb{R}^N} |\Delta u_\epsilon|^2 |x|^m dx}{\int_{\mathbb{R}^N} |x \cdot \nabla u_\epsilon|^2 |x|^{m-4} dx} = \\
&\searrow \left( -\frac{N+m-2}{2} \right)^2 + 2c_1 + (N-1)(1-m) + \frac{4(c_1^2 - c_1(m-2)(N+m-4))}{(N+m-4)^2} \\
&= \frac{((m-2)^2 - N^2)^2}{4(N+m-4)^2}
\end{aligned}$$

as  $\epsilon \searrow 0$ .

For the second "bad" case  $N \geq 2$  and  $m \in (2 + \sqrt{(N-1)^2 + 1}, \infty)$  we consider the sequence

$$(56) \quad u_\epsilon(x) = |x|^{-\frac{N+m-4}{2} + \epsilon} g(|x|) \phi_{l_{\min}}(x),$$

where  $\phi_{l_{\min}}$  is a spherical harmonic function of degree  $l_{\min}$  such that

$\|\phi_{l_{\min}}\|_{L^2(S^{N-1})} = 1$  and  $l_{\min} := \arg \min_{l \leq k(m)} \tilde{C}(N, m, l)$ , where  $\tilde{C}(N, m, l) := \frac{(-N+m-2l)^2(2l+m+N-4)^2}{4(m+N-4)^2}$  and  $k(m)$  was defined in (31)-(44).

Similarly (but more technically) as above one can show that

$$\frac{\int_{\mathbb{R}^N} |\Delta u_\epsilon|^2 |x|^m dx}{\int_{\mathbb{R}^N} |x \cdot \nabla u_\epsilon|^2 |x|^{m-4} dx} \searrow \tilde{C}(N, m, l)$$

as  $\epsilon \searrow 0$ . The details are let to the reader.

The optimality is showed and the proof of the main result is complete now.

**Acknowledgments.** First author was partially supported by a grant of the Ministry of Research, Innovation and Digitization, CNCS-UEFISCDI Romania, project number PN-III-P1-1.1-TE-2021-1539, within PNCDI III. This work started during the elaboration of the dissertation thesis of the second author, within the master program "Advanced Studies in Mathematics" at the University of Bucharest.

## REFERENCES

- [1] N. Bez, S. Machihara and T. Ozawa, *Revisiting the Rellich inequality*, Math. Z. 303 (2023), no. 2, Paper No. 49, 11 pp.
- [2] B. Cassano and F. Pizzichillo, *Self-adjoint extensions for the Dirac operator with Coulomb-type spherically symmetric potentials*. Lett. Math. Phys. 108 (2018), 2635–2667.
- [3] B. Cassano, L. Cossetti and L. Fanelli, *Improved Hardy-Rellich inequalities*. Commun. Pure Appl. Anal. 21 (2022), no. 3, 867–889.
- [4] C. Cazacu, *A new proof of the Hardy-Rellich inequality in any dimension*, Proc. Roy. Soc. Edinburgh Sect. A, 150 (2020), no. 6, 2894-2904.
- [5] C. Cazacu, J. Flynn and N. Lam, *Short proofs of refined sharp Caffarelli-Kohn-Nirenberg inequalities*, J. Differential Equations 302 (2021), 533–549.
- [6] C. Cazacu, L. Ignat and D. Manea, *The Hardy inequality and large time behaviour of the heat equation on  $\mathbb{R}^{N-k} \times (0, \infty)^k$* , J. Differential Equations 375 (2023), 682–705.
- [7] N. Ghoussoub, N. and A. Moradifam, *Bessel pairs and optimal Hardy and Hardy-Rellich inequalities*, Math. Ann. 349 (2011)
- [8] H. Groemer, *Geometric Applications of Fourier Series and Spherical Harmonics*, Cambridge University Press, 1996
- [9] N. Hamamoto, *Solenoidal improvement of Rellich-Hardy inequalities with power weights*. Calc. Var. Partial Differential Equations 63 (2024), no. 4, Paper No. 84, 25 pp.
- [10] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*. Reprint of the 1952 edition. Cambridge University Press, Cambridge, 1988.
- [11] N. Lam and G. Lu, *Improved  $L^p$ -Hardy and  $L^p$ -Rellich inequalities with magnetic fields*. Vietnam J. Math. 51 (2023), no. 4, 971–984.
- [12] A. Michelat and T. Rivière, *Weighted Eigenvalue Problems for Fourth-Order Operators in Degenerating Annuli*. Preprint arXiv:2306.04609.
- [13] B. Opic, A. Kufner, *Hardy-type inequalities*. Pitman Res. Notes Math. Ser., 219. Longman Scientific & Technical, Harlow, 1990.
- [14] F. Rellich, *Halbbeschränkte Differentialoperatoren höherer Ordnung*. (German) Proceedings of the International Congress of Mathematicians, 1954, Amsterdam, vol. III, pp. 243–250
- [15] A. Tertikas A. and Zographopoulos, *Best constant in the Hardy-Rellich inequalities and related improvements*, Adv. Math. 206 (2007)
- [16] Barbatis and A. Tertikas, *Sobolev improvements on sharp Rellich inequalities*, to appear in J. Spectral Theory, arXiv:2312.00433.
- [17] B. Rosenzweig and J. Stanfill, *private communication*.



CRISTIAN CAZACU: FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF BUCHAREST, 14 ACADEMIEI STREET, 010014 BUCHAREST, ROMANIA & GHEORGHE MIHOC-CAIUS IACOB INSTITUTE OF MATHEMATICAL STATISTICS AND APPLIED MATHEMATICS OF THE ROMANIAN ACADEMY, No.13 CALEA 13 SEPTEMBRIE, SECTOR 5, 050711 BUCHAREST, ROMANIA

*Email address:* cristian.cazacu@fmi.unibuc.ro

IRINA FIDEL: SFANTUL VASILE SECONDARY SCHOOL, No. 145 REPUBLICII, 100389 PLOIESTI, ROMANIA, & FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF BUCHAREST, 14 ACADEMIEI STREET, 010014 BUCHAREST, ROMANIA

*Email address:* irina.fidel@s.unibuc.ro