

On the Elliptic Sombor and Euler Sombor indices of Corona product of certain graphs

B. KIRANA¹, M.C. SHANMUKHA^{2,*}, A. USHA³

¹ Department of Mathematics, KVG College of Engineering, Sullia, 574327 and affiliated to Visvesvaraya Technological University, Belagavi-590018, India

² Department of Mathematics, PES Institute of Technology and Management, Shivamogga, 577204 and affiliated to Visvesvaraya Technological University, Belagavi, 590018, India

³ Department of Mathematics, Alliance School of Applied Mathematics, Alliance University, Bangalore, 562106, India

e-mail*: mcsanmukha@gmail.com

Abstract

Elliptic Sombor and Euler Sombor indices are recently defined topological indices using Sombor index. Elliptic sombor index is defined as $ESO(G) = \sum_{uv \in E(G)} (d_u + d_v) \sqrt{d_u^2 + d_v^2}$ and Euler Sombor index is defined as $EU(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2 + d_u d_v}$, where d_u and d_v are degrees of vertices u and v in graph G . In this article, we compute the elliptic Sombor and Euler Sombor indices of some resultant graphs. Using the operations join and Corona product on standard graphs like path, cycle and complete graphs.

Keywords: Elliptic Sombor index, Euler Sombor index, Join, Corona product

1 Introduction

Degree-based topological indices are fundamental tools in the fields of mathematical chemistry and graph theory, serving as quantitative measures of a molecular graph structure. A molecular graph represents a chemical compound where vertices correspond to atoms and edges correspond to chemical bonds. The degree of a vertex in this context refers to the number of edges connected to it, representing the number of bonds of an atom. The primary utility of these indices lie in their ability to predict various physical, chemical, and biological properties of molecules [1,2].

The Sombor index, introduced by Gutman in 2021, is a topological index derived from graph theory and Euclidean geometry. This index has gained significant traction in the fields such as chemistry and pharmacology for its utility in characterizing

molecular structures and predicting biological activity. Let us delve into the formal definition, properties, and an example calculation of the Sombor index [3, 4].

The general form of a vertex degree-based index is a function which is chosen such that it satisfies symmetry property. The edge uv representation in 2-dimensional coordinate system is called the degree-point of the edge uv . The Euclidean distance between the degree-point (d_u, d_v) and the origin O is $\sqrt{d_u^2 + d_v^2}$ which is the definition of Sombor index [5].

In 2023, Gutman et al., [6] introduced another version of Sombor index called elliptic Sombor index referring to the orbits of planets in the solar system which takes elliptic orbits with the sun as focus point. The concept of an elliptic Sombor index can be an extension of the traditional Sombor index by incorporating elliptic distance rather than the Euclidean distance between degree-points.

In 2024, Gutman et al., [7] showed that the lengths of semi major and the semi minor axes are equal in an ellipse. Leonard Euler found the approximate perimeter of the ellipse as $\pi\sqrt{2(d_u^2 + d_v^2)(d_u + d_v)^2}$. Using these relations, Euler Sombor index was proposed as $\sqrt{d_u^2 + d_v^2 + d_u \cdot d_v}$. Algebraically, there is a geometric analogy of Sombor and Euler Sombor indices.

Bibhas et al., [8] studied structural properties of corona graphs including the statistics of signed links and types of signed triangles and degree distribution. They analyzed algebraic conflict of signed corona graphs generated by seed graphs. Sheeba et al., [9] established the explicit expressions of Y -index of different types of corona product of graphs. Khalid et al., [10] studied certain degree-based topological indices such as Randić index, Zagreb indices, multiplicative Zagreb indices, Narumi–Katayama index, atom-bond connectivity index, augmented Zagreb index, geometric-arithmetic index, harmonic index, and sum-connectivity index for the bistar graphs and the corona product. Sheeja et al., [11] established the explicit expressions for the SK index over different types of corona products on graphs are presented. Iswadi et al., [12] determined the metric dimension of corona product, and lower bound of metric dimension of join of graphs. Dhananjaya et al., [13] derived ABC index and the GA index of several corona products of graphs composed of path, cycle, and complete graphs. Arif et al., [14] computed the Sombor index of some graph operations namely, join and corona product of two graphs, for the standard graphs path, cycle, and complete graphs.

Motivated by the above studies on graph operations, this study focusses on computing the elliptic Sombor and Euler Sombor indices of some resultant graphs using the operations join and Corona product on standard graphs like path, cycle and complete graphs.

Graph operations, such as join, union, Cartesian product, and others, are funda-

mental tools in graph theory that significantly influence topological indices [15–18]. Given two disjoint graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$, the join $G_1 + G_2$ (See Figure 1) is defined as:

$$G_1 + G_2 = (V_1 \cup V_2, E_1 \cup E_2 \cup \{(v_1, v_2) \mid v_1 \in V_1, v_2 \in V_2\})$$

Let G be a connected graph with n vertices and H be a graph with at least two vertices, the corona product of G and H , is defined as a graph which is formed by taking n copies of H and connecting i^{th} vertex of G to the vertices of H in each copy, and it is denoted by $G \odot H$ (See Figure 2).




Figure 1: Join of two paths P_2 and P_3 .




Figure 2: Corona product of two graphs K_3 and P_2 .

Elliptic Sombor and Euler Sombor indices are defined as follows [6, 7]

$$ESO(G) = \sum_{uv \in E(G)} (d_u + d_v)(\sqrt{d_u^2 + d_v^2}) \quad (1)$$

$$EU(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2 + d_u d_v} \quad (2)$$

2 Main Results

In this section, Elliptic Sombor and Euler Sombor indices of $P_r + P_s$, $C_r + C_s$, $K_r + K_s$, $P_r \odot P_s$, $C_r \odot C_s$ are obtained.

If P_r and P_s are two paths of order $|V(G)| = r + s$ and size $|E(G)| = (r - 1) + (s - 1) + rs = r + s + rs - 2$. In the following theorem, the Elliptic Sombor and Euler Sombor indices of two path graphs P_r and P_s are computed.

Theorem 2.1. *Elliptic Sombor index of $P_r + P_s$ is given by*

$$ESO(P_r + P_s) = \begin{cases} 108\sqrt{2}, r = s = 2 \\ 70 + 2\sqrt{2}(s + 1)^2 + 32\sqrt{2}(s - 3) \\ + 4(s + 4)\sqrt{9 + (s + 1)^2} \\ + 2(s - 2)(s + 5)\sqrt{16 + (s + 1)^2}; s > 2, r = 2 \\ 70 + 2\sqrt{2}(r + 1)^2 + 32\sqrt{2}(r - 3) \\ + 4(r + 4)\sqrt{9 + (r + 1)^2} \\ + 2(r - 2)(s + 5)\sqrt{16 + (r + 1)^2}; r > 2, s = 2 \\ 2(2s + 3)\sqrt{(s + 1)^2 + (s + 2)^2} \\ + (r - 3)(2s + 4(s + 2)\sqrt{2} + 2(2r + 3)\sqrt{(r + 1)^2 + (r + 2)^2}) \\ + (s - 3)(2r + 4)(r + 2)\sqrt{2} + 4(r + s + 2)\sqrt{(r + 1)^2 + (s + 1)^2}; r, s > 2 \end{cases}$$

Proof. Consider the join of two graphs P_r and P_s . If $r, s > 1$, based on the degrees of vertices of $P_r + P_s$, there are 4 types of vertices having degrees $s + 1, s + 2, r + 1$ and $r + 2$. Then by considering the degrees of the vertices, there are 10 types of edge partitions of $P_r + P_s$ as shown in the Table 1.

Table 1: Number of edges for different cases in $P_r + P_s$.

Case	Number of edges	Condition
$(s + 1, s + 1)$	1	$r = 2$
	0	$r > 2$
$(s + 1, s + 2)$	0	$r = 2$
	2	$r > 2$
$(s + 2, s + 2)$	0	$r = 2$
	$r - 3$	$r > 2$
$(r + 1, r + 1)$	1	$s = 2$
	0	$s > 2$
$(r + 1, r + 2)$	0	$s = 2$
	2	$s > 2$
$(s + 2, s + 2)$	0	$s = 2$
	$s - 3$	$s > 2$
$(s + 1, r + 1)$	4	Always
$(s + 1, r + 2)$	0	$s = 2$
	$2(s - 2)$	$s > 2$
$(s + 2, s + 1)$	0	$r = 2$
	$2(r - 2)$	$r > 2$
$(s + 2, r + 2)$	0	$r = 2$ or $s = 2$
	$(r - 2)(s - 2)$	$r, s > 2$

Case 1 $r=s=2$

$$ESO(P_r + P_s) = 2(s + 1)\sqrt{(s + 1)^2 + (s + 1)^2} + 2(r + 1)\sqrt{(r + 1)^2 + (r + 1)^2},$$

$$+ 4(s + 1 + r + 1)\sqrt{(s + 1)^2 + (r + 1)^2}$$

$$ESO(P_r + P_s) = 2(s + 1)\sqrt{2(s + 1)^2} + 2(r + 1)\sqrt{2(r + 1)^2}$$

$$+ 4(r + s + 2)\sqrt{(s + 1)^2 + (r + 1)^2},$$

Substitute $r = 2, s = 2$ we get

$$ESO(P_r + P_s) = 108\sqrt{2}.$$

Case 2 $r=2, s > 2$

$$\begin{aligned}
ESO(P_r + P_s) &= 2(s+1)\sqrt{2(s+1)^2} + 2(r+1+r+2)\sqrt{(r+1)^2 + (r+2)^2} \\
&\quad + (s-3)2(r+2)\sqrt{2(r+2)^2} + 4(r+1+s+1)\sqrt{(r+1)^2 + (s+1)^2} \\
&\quad + 2(s-2)(r+2+s+1)\sqrt{(r+2)^2 + (s+1)^2},
\end{aligned}$$

$$\begin{aligned}
ESO(P_r + P_s) &= 2(s+1)^2\sqrt{2} + 2(2r+3)\sqrt{(r+1)^2 + (r+2)^2} \\
&\quad + 2(s-3)(r+2)(r+2)\sqrt{2} + 4(r+s+2)\sqrt{(r+1)^2 + (s+1)^2} \\
&\quad + 2(s-2)(r+s+3)\sqrt{(r+2)^2 + (s+1)^2},
\end{aligned}$$

take $r = 2$ we obtain

$$\begin{aligned}
ESO(P_r + P_s) &= 2\sqrt{2}(s+1)^2 + 2 \times 7\sqrt{3^2 + 4^2} + 32(s-3)\sqrt{2} \\
&\quad + 4(s+4)\sqrt{9 + (s+1)^2} + 2(s-2)(s+5)\sqrt{16 + (s+1)^2}, \\
ESO(P_r + P_s) &= 70 + 2\sqrt{2}(s+1)^2 + 32(s-3)\sqrt{2} + 4(s+4)\sqrt{9 + (s+1)^2} \\
&\quad + 2(s-2)(s+5)\sqrt{16 + (s+1)^2}.
\end{aligned}$$

Case 3 $s=2, r > 2$

$$\begin{aligned}
ESO(P_r + P_s) &= 2(r+1)\sqrt{2(r+1)^2} + 2(s+1+s+2)\sqrt{(s+1)^2 + (s+2)^2} \\
&\quad + (r-3)2(s+2)\sqrt{2(s+2)^2} + 4(r+1+s+1)\sqrt{(r+1)^2 + (s+1)^2} \\
&\quad + 2(r-2)(r+1+s+2)\sqrt{(r+1)^2 + (s+2)^2},
\end{aligned}$$

$$\begin{aligned}
ESO(P_r + P_s) &= 2(r+1)^2\sqrt{2} + 2(2s+3)\sqrt{(s+1)^2 + (s+2)^2} \\
&\quad + 2(r-3)(s+2)^2\sqrt{2} + 4(r+s+2)\sqrt{(r+1)^2 + (s+1)^2} \\
&\quad + 2(r-2)(r+s+3)\sqrt{(s+2)^2 + (r+1)^2},
\end{aligned}$$

Substitute $s = 2$

$$\begin{aligned}
ESO(P_r + P_s) &= 2\sqrt{2}(r+1)^2 + 2 \times 7\sqrt{3^2 + 4^2} + 32(r-3)\sqrt{2} \\
&\quad + 4(r+4)\sqrt{9 + (r+1)^2} + 2(r-2)(s+5)\sqrt{16 + (r+1)^2}, \\
ESO(P_r + P_s) &= 70 + 2\sqrt{2}(r+1)^2 + 32\sqrt{2}(r-3) + 4(r+4)\sqrt{9 + (r+1)^2} \\
&\quad + 2(r-2)(s+5)\sqrt{16 + (r+1)^2}.
\end{aligned}$$

Case 4 $r, s > 2$

$$\begin{aligned}
 ESO(P_r + P_s) = & 2(2s + 3)\sqrt{(s + 1)^2 + (s + 2)^2} + 2(r - 3)(s + 2)\sqrt{2(s + 2)^2} \\
 & + 2(2r + 3)\sqrt{(r + 1)^2 + (r + 2)^2} + 2(s - 3)(r + 2)\sqrt{2(r + 2)^2} \\
 & + 4(r + s + 2)\sqrt{(r + 1)^2 + (s + 1)^2} + 2(s - 2)(r + s + 3) \\
 & \sqrt{(r + 2)^2 + (s + 1)^2} + (r - 2)(s - 2)(r + s + 4)\sqrt{(r + 2)^2 + (s + 2)^2} \\
 & + 2(r - 2)(r + s + 3)\sqrt{(r + 1)^2 + (s + 2)^2}.
 \end{aligned}$$

□

Theorem 2.2. *The Euler Sombor index of $P_r + P_s$ is given by*

$$EU(P_r + P_s) = \begin{cases} 18\sqrt{3}, r = s = 2 \\ 2\sqrt{37} + (5s - 11)\sqrt{3} + 4\sqrt{(s + 1)^2 + 3s + 12} \\ + 2(s - 2)\sqrt{(s + 1)^2 + 4s + 20}; s > 2, r = 2 \\ 2\sqrt{37} + (5s - 11)\sqrt{3} + 4\sqrt{(r + 1)^2 + 3r + 12} \\ + 2(r - 2)\sqrt{(r + 1)^2 + 4r + 20}; r > 2, s = 2 \\ 2\sqrt{(s + 1)^2 + (s + 2)^2 + (s + 1)(s + 2)} + (r - 3)\sqrt{3(s + 2)^2} \\ + 2\sqrt{(r + 1)^2 + (r + 2)^2 + (r + 1)(r + 2)} + (s - 3)\sqrt{3(r + 2)^2} \\ + 4\sqrt{(r + 1)^2 + (s + 1)^2 + (r + 1)(s + 1)} \\ + 2(s - 2)\sqrt{(r + 2)^2 + (s + 1)^2 + (r + 2)(s + 1)} \\ + (r - 2)(s - 2)\sqrt{(r + 2)^2 + (s + 2)^2 + (r + 2)(s + 2)} \\ + 2(r - 2)\sqrt{(r + 1)^2 + (s + 2)^2 + (r + 1)(s + 2)}; r, s > 2 \end{cases}$$

Proof. Using edge partition of join of two path graphs P_r and P_s from Table 1, Euler Sombor index is calculated using the equation 2. The following cases are considered.

Case 1 $r=s=2$

$$\begin{aligned}
 EU(P_r + P_s) = & \sqrt{(s + 1)^2 + (s + 1)^2 + (s + 1)(s + 1)} \\
 & + \sqrt{(r + 1)^2 + (r + 1)^2 + (r + 1)(r + 1)} \\
 & + 4\sqrt{(s + 1)^2 + (r + 1)^2 + (s + 1)(r + 1)},
 \end{aligned}$$

$EU(P_r + P_s) = \sqrt{3(s+1)^2} + \sqrt{3(r+1)^2} + 4\sqrt{(s+1)^2 + (r+1)^2 + (s+1)(r+1)}$,
 take $r = 2, s = 2$ we get

$$EU(P_r + P_s) = 18\sqrt{3}.$$

Case 2 $r=2, s > 2$

$$\begin{aligned} EU(P_r + P_s) &= \sqrt{2(s+1)^2 + (s+1)(s+1)} + 2\sqrt{(r+1)^2 + (r+2)^2 + (r+1)(r+2)} \\ &\quad + (s-3)\sqrt{2(r+2)^2 + (r+2)(r+2)} \\ &\quad + 4\sqrt{(r+1)^2 + (s+1)^2 + (r+1)(s+1)} \\ &\quad + 2(s-2)\sqrt{(r+2)^2 + (s+1)^2 + (r+2)(s+1)}, \end{aligned}$$

$$\begin{aligned} EU(P_r + P_s) &= \sqrt{3}[(s+1) + (r+2)(s-3)] + 2\sqrt{(r+1)^2 + (r+2)^2 + (r+1)(r+2)} \\ &\quad + 4\sqrt{(r+1)^2 + (s+1)^2 + (r+1)(s+1)} \\ &\quad + 2(s-2)\sqrt{(r+2)^2 + (s+1)^2 + (r+2)(s+1)}, \end{aligned}$$

Substitute $r = 2$

$$EU(P_r + P_s) = 2\sqrt{37} + (5s - 11)\sqrt{3} + 4\sqrt{(s+1)^2 + 3s + 12} + 2(s-2)\sqrt{(s+1)^2 + 4s + 20}.$$

Case 3 $s = 2, r > 2$

$$\begin{aligned} EU(P_r + P_s) &= \sqrt{2(r+1)^2 + (r+1)(r+1)} + 2\sqrt{(s+1)^2 + (s+2)^2 + (s+1)(s+2)} \\ &\quad + (r-3)\sqrt{3(r+2)^2} + 4\sqrt{(r+1)^2 + (s+1)^2 + (r+1)(s+1)} \\ &\quad + 2(r-2)\sqrt{(s+2)^2 + (r+1)^2 + (s+2)(r+1)}, \end{aligned}$$

$$\begin{aligned} EU(P_r + P_s) &= \sqrt{3}[(r+1) + (s+2)(r-3)] + 2\sqrt{(s+1)^2 + (s+2)^2 + (s+1)(s+2)} \\ &\quad + 4\sqrt{(r+1)^2 + (s+1)^2 + (r+1)(s+1)} \\ &\quad + 2(r-2)\sqrt{(s+2)^2 + (r+1)^2 + (r+2)(s+1)}, \end{aligned}$$

Substitute $s = 2$

$$\begin{aligned} EU(P_r + P_s) &= 2\sqrt{37} + (5s - 11)\sqrt{3} + 4\sqrt{(r+1)^2 + 3r + 12} \\ &\quad + 2(r-2)\sqrt{(r+1)^2 + 4r + 20}. \end{aligned}$$

Case 4 $r, s > 2$

$$\begin{aligned}
EU(P_r + P_s) = & 2\sqrt{(s+1)^2 + (s+2)^2 + (s+1)(s+2)} + (r-3)\sqrt{3(s+2)^2} \\
& + 2\sqrt{(r+1)^2 + (r+2)^2 + (r+1)(r+2)} + (s-3)\sqrt{3(r+2)^2} \\
& + 4\sqrt{(r+1)^2 + (s+1)^2 + (r+1)(s+1)} \\
& + 2(s-2)\sqrt{(r+2)^2 + (s+1)^2 + (r+2)(s+1)} \\
& + (r-2)(s-2)\sqrt{(r+2)^2 + (s+2)^2 + (r+2)(s+2)} \\
& + 2(r-2)\sqrt{(r+1)^2 + (s+2)^2 + (r+1)(s+2)}.
\end{aligned}$$

□

Theorem 2.3. *Let r, s be two positive integers such that $r, s \geq 3$. Then Elliptic Sombor index and Euler Sombor index of $C_r + C_s$ are given by*

$$ESO(C_r + C_s) = 2\sqrt{2}(s+2)^2 + s(r+2)^2] + rs(r+s+4)\sqrt{(r+2)^2 + (s+2)^2}.$$

$$EU(C_r + C_s) = 2\sqrt{3}(r+s+rs) + rs\sqrt{(r+2)^2 + (s+2)^2 + (r+2)(s+2)}.$$

Proof. Consider join of two cycles C_r and C_s which is graph with order $|V(C_r + C_s)| = r + s$ and size $|E(C_r + C_s)| = r + s + rs$. There are two types of degree vertices such as $s+2$ and other type has $r+2$ in the resultant graph. Thus, there are three edge partitions

Table 2: Edge partitions of $C_r + C_s$.

Case	Number of edges
$(s+2, s+2)$	r
$(s+2, r+2)$	rs
$(r+2, r+2)$	s

Using above edge partitions and the formula

$$\begin{aligned}
ESO(C_r + C_s) = & r(s+2+s+2)\sqrt{2(s+2)^2} + rs(r+2+s+2)\sqrt{(r+2)^2 + (s+2)^2} \\
& + s(r+2+r+2)\sqrt{2(r+2)^2},
\end{aligned}$$

$$ESO(C_r + C_s) = 2\sqrt{2}r(s+2)^2 + rs(r+s+4)\sqrt{(r+2)^2 + (s+2)^2} + 2\sqrt{2}s(r+2)^2,$$

$$ESO(C_r + C_s) = 2\sqrt{2}((r(s+2)^2 + s(r+2)^2) + rs(r+s+4)\sqrt{(r+2)^2 + (s+2)^2}).$$

$$\begin{aligned}
EU(C_r + C_s) &= r\sqrt{3(s+2)^2} + rs\sqrt{(r+2)^2 + (s+2)^2 + (r+2)(s+2)} + s\sqrt{3(r+2)^2}, \\
EU(C_r + C_s) &= \sqrt{3}[r(s+2) + s(r+2)] + rs\sqrt{(r+2)^2 + (s+2)^2 + (r+2)(s+2)}, \\
EU(C_r + C_s) &= 2\sqrt{3}[r + s + rs] + rs\sqrt{(r+2)^2 + (s+2)^2 + (r+2)(s+2)}.
\end{aligned}$$

□

Theorem 2.4. *If two complete graphs K_r and K_s are joined, the resultant is a complete graph K_{r+s} with order $|V(K_{r+s})| = r + s$ and size $|E(K_{r+s})| = \frac{(r+s)(r+s-1)}{2}$. Then*

$$\begin{aligned}
ESO(K_r + K_s) &= \sqrt{2}(r+s)(r+s-1)^3. \\
EU(K_r + K_s) &= \frac{\sqrt{3}}{2}(r+s)(r+s-1)^2.
\end{aligned}$$

Proof. Each vertex of join of two graphs K_r and K_s has degree $r+s-1$. So there is only one type of edge that is $(r+s-1, r+s-1)$. Using equation (1), Elliptic Sombor index is

$$\begin{aligned}
ESO(K_r + K_s) &= ESO(K_{r+s}) \\
&= \frac{(r+s)(r+s-1)}{2}(r+s-1+r+s-1)\sqrt{2(r+s-1)^2} \\
&= \sqrt{2}(r+s)(r+s-1)^3.
\end{aligned}$$

Using equation (2), Euler Sombor index is

$$\begin{aligned}
EU(K_r + K_s) &= EU(K_{r+s}) \\
&= \frac{(r+s)(r+s-1)}{2}\sqrt{3(r+s-1)^2} \\
&= \sqrt{3}\frac{(r+s)(r+s-1)^2}{2}.
\end{aligned}$$

□

Theorem 2.5. *Let C_r and K_S be the cycle and complete graph respectively. Then*

Elliptic Sombor index and Euler sombor index of $C_r + K_s$ are

$$ESO(C_r + K_s) = 2\sqrt{2}r(s+2)^2 + rs(r+2s+1)\sqrt{(s+2)^2 + (r+s-1)^2} \\ + C(s,2)(r+s-1)\sqrt{2}.$$

$$EU(C_r + K_s) = r(s+2)\sqrt{3} + rs\sqrt{(s+2)^2 + (r+s-1)^2} + (s+2)(r+s-1) \\ + C(s,2)(r+s-1)\sqrt{3}.$$

Proof. The graph $C_r + K_s$ contains $r + s$ number of vertices and $r + rs + C(s, 2)$ number of edges. Here C_r has r number of edges and K_s has $C(s, 2)$ number of edges. There are three types of edge partitions that are tabulated in Table 3.

Table 3: Number of edges for different cases in $C_r + K_s$.

Case	Number of edges
$(s+2, s+2)$	r
$(s+2, s+r-1)$	rs
$(r+s-1, r+s-1)$	$C(s, 2)$

$$ESO(C_r + K_s) = r(s+2+s+2)\sqrt{2(s+2)^2} + rs(s+2+r+s-1)\sqrt{(s+2)^2 + (r+s-1)^2} \\ + C(s,2)(r+s-1+r+s-1)\sqrt{2(r+s-1)^2},$$

$$ESO(C_r + K_s) = 2\sqrt{2}r(s+2)^2 + rs(r+2s+1)\sqrt{(s+2)^2 + (r+s-1)^2} \\ + 2\sqrt{2}C(s,2)(r+s-1)^2.$$

$$EU(C_r + K_s) = r\sqrt{3(s+2)^2} + rs\sqrt{(s+2)^2 + (r+s-1)^2} + (s+2)(r+s-1) \\ + C(s,2)\sqrt{3(r+s-1)^2},$$

$$EU(C_r + K_s) = \sqrt{3}r(s+2) + rs\sqrt{(s+2)^2 + (r+s-1)^2} + (s+2)(r+s-1) \\ + \sqrt{3}C(s,2)(r+s-1).$$

□

Theorem 2.6. *Elliptic Sombor index of corona product of two paths are given by*

$$ESO(P_r \odot P_s) = \begin{cases} 34\sqrt{2} + 20\sqrt{13}, r = s = 2 \\ 20\sqrt{13} + 2\sqrt{2}(s+1)^2 + 4(s+3)\sqrt{4+(s+1)^2} \\ + 2(s-2)(s+4)\sqrt{9+(s+1)^2}; s > 2, r = 2 \\ 70 + 20\sqrt{13} + 8\sqrt{2}r + 32\sqrt{2}(r-3) + 24(r-2)\sqrt{5}; r > 2, s = 2 \\ 18\sqrt{2}r(s-3) + 2\sqrt{2}(r-3)(s+2)^2 + 10\sqrt{13}r + 4(s+3)\sqrt{4+(s+1)^2} \\ + 2(r-1)(s+4)\sqrt{4+(s+2)^2} + 2(s-2)(s+4)\sqrt{9+(s+1)^2} \\ + (r-2)(s-2)(s+5)\sqrt{9+(s+2)^2} + 2(2s+3)\sqrt{(s+1)^2+(s+2)^2} \\ ; r, s > 2 \end{cases}$$

Proof. Consider the graph $P_r \odot P_s$ having $|V(P_r \odot P_s)| = rs + r$ and $|E(P_r \odot P_s)| = 2rs - 1$. There are four types of degrees of vertices of graph $P_r \odot P_s$ namely 2, 3, $s+1$, $s+2$. Using edge partition listed in Table 4 and formula(1),

Case 1 $r=s=2$

$$ESO(P_r \odot P_s) = r(2+2)\sqrt{2^2+2^2} + (s+1+s+1)\sqrt{2(s+1)^2} \\ + 4(s+1+2)\sqrt{(s+1)^2+2^2},$$

Substitute $r = s = 2$

$$ESO(P_r \odot P_s) = 34\sqrt{2} + 20\sqrt{13}.$$

Case 2 $r = 2, s > 2$

$$ESO(P_r \odot P_s) = (s+1+s+1)\sqrt{2(s+1)^2} + 2r(2+3)\sqrt{2^2+3^2} \\ + 4(2+s+1)\sqrt{2^2+(s+1)^2} + 2(s-2)(3+s+1)\sqrt{3^2+(s+1)^2}$$

Substitute $r = 2$

$$ESO(P_r \odot P_s) = 20\sqrt{13} + 2\sqrt{2}(s+1)^2 + 4(s+3)\sqrt{4+(s+1)^2} \\ + 2(s-2)(s+4)\sqrt{9+(s+1)^2}.$$

Case 3 $s = 2, r > 2$

Table 4: Number of edges for different cases in $P_r \odot P_s$.

Edge partitions	Number of edges	Condition
(2, 2)	r 0	$s = 2$ $s > 2$
(3, 3)	0 $r(s - 3)$	$s = 2$ $s > 2$
($s + 1, s + 1$)	1 0	$r = 2$ $s > 2$
($s + 2, s + 2$)	0 $n - 3$	$r = 2$ $r > 2$
(2, 3)	0 $2r$	$s = 2$ $s > 2$
(2, $s + 1$)	4	always
(2, $s + 2$)	0 $2(r - 2)$	$r = 2$ $r > 2$
(3, $s + 1$)	0 $2(s - 2)$	$s = 2$ $s > 2$
(3, $s + 2$)	0 $(r - 2)(s - 2)$	$r = 2$ $r > 2$
($s + 1, s + 2$)	0 2	$r = 2$ $s > 2$

$$\begin{aligned}
ESO(P_r \odot P_s) = & r(2 + 2)\sqrt{2^2 + 2^2} + (r - 3)(s + 2 + s + 2)\sqrt{2(s + 2)^2} \\
& + 4(2 + s + 1)\sqrt{2^2 + (s + 1)^2} + 2(r - 2)(s + 2 + 2)\sqrt{2^2 + (s + 2)^2} \\
& + (r - 2)(s - 2)(3 + s + 2)\sqrt{3^2 + (s + 2)^2} \\
& + 2(s + 1 + s + 2)\sqrt{(s + 1)^2 + (s + 2)^2}
\end{aligned}$$

Substitute $s = 2$, we get

$$ESO(P_r \odot P_s) = 70 + 20\sqrt{138}\sqrt{2}r + 32\sqrt{2}(r - 3) + 24\sqrt{5}(r - 2).$$

Case 4 $r, s > 2$

$$\begin{aligned}
ESO(P_r \odot P_s) = & r(s-3)(3+3)\sqrt{3^2+3^2} + (r-3)(s+2+s+2)\sqrt{2(s+2)^2} \\
& + 2r(2+3)\sqrt{2^2+3^2} + 4(s+1+2)\sqrt{(s+1)^2+2^2} \\
& + 2(r-1)(s+2+2)\sqrt{(s+2)^2+2^2} + 2(s-2)(s+1+3)\sqrt{(s+1)^2+3^2} \\
& + (r-2)(s-2)(s+2+3)\sqrt{(s+2)^2+3^2} \\
& + 2(s+1+s+2)\sqrt{(s+1)^2+(s+2)^2}.
\end{aligned}$$

□

Theorem 2.7. Euler Sombor index of corona product of two paths are given by

$$EU(P_r \odot P_s) = \begin{cases} 7\sqrt{3} + 4\sqrt{19}, r = s = 2 \\ 4\sqrt{19} + (s+1)\sqrt{3} + 4\sqrt{4 + (s+1)^2 + 2(s+1)} \\ + 2(s-2)\sqrt{9 + (s+1)^2 + 3(s+1)}; s > 2, r = 2 \\ 4\sqrt{19} + 2\sqrt{3}r + 8\sqrt{2}r + 32\sqrt{2}(r-3) + 24(r-2)\sqrt{5}; r > 2, s = 2 \\ 3\sqrt{3}r(s-3) + \sqrt{3}(r-3)(s+2) + 2\sqrt{19}r + 4\sqrt{4 + (s+1)^2 + 2(s+1)} \\ + 2(r-1)\sqrt{4 + (s+2)^2 + 2(s+2)} + 2(s-2)\sqrt{9 + (s+1)^2 + 3(s+1)} \\ + (r-2)(s-2)\sqrt{9 + (s+2)^2 + 3(s+2)} \\ + 2\sqrt{(s+1)^2 + (s+2)^2 + (s+1)(s+2)} \end{cases}$$

Proof. Using the edge partitions in Table 4 and formula (2) we have

Case 1 $r=s=2$

$$EU(P_r \odot P_s) = r\sqrt{2^2+2^2+2^2} + \sqrt{3(s+1)^2} + 4\sqrt{(s+1)^2+2^2+2(s+1)},$$

Substitute $r = s = 2$

$$EU(P_r \odot P_s) = 7\sqrt{3} + 4\sqrt{19}.$$

Case 2 $r = 2, s > 2,$

$$\begin{aligned}
EU(P_r \odot P_s) = & \sqrt{3(s+1)^2} + 2r\sqrt{2^2+3^2+2 \times 3} \\
& + 4\sqrt{2^2+(s+1)^2+2(s+1)} + 2(s-2)\sqrt{3^2+(s+1)^2+3(s+1)},
\end{aligned}$$

Substitute $r = 2$

$$\begin{aligned}
EU(P_r \odot P_s) &= 4\sqrt{19} + \sqrt{3}(s+1) + 4\sqrt{4 + (s+1)^2 + 2(s+1)} \\
&\quad + 2(s-2)\sqrt{9 + (s+1)^2 + 3(s+1)}. \text{Case 3 } s = 2, r > 2 \\
EU(P_r \odot P_s) &= r\sqrt{2^2 + 2^2 + 2^2} + (r-3)\sqrt{3(s+2)^2} \\
&\quad + 4\sqrt{2^2 + (s+1)^2 + 2(s+1)} + 2(r-2)\sqrt{2^2 + (s+2)^2 + 2(s+2)} \\
&\quad + (r-2)(s-2)\sqrt{3^2 + (s+2)^2 + 3(s+2)} \\
&\quad + 2\sqrt{(s+1)^2 + (s+2)^2 + (s+1)(s+2)},
\end{aligned}$$

Substitute $s = 2$, we get

$$EU(P_r \odot P_s) = 4\sqrt{19} + 2\sqrt{37} + 2\sqrt{3}r + 4\sqrt{3}(r-3) + 4\sqrt{7}(r-2).$$

Case 4 $r, s > 2$

$$\begin{aligned}
ESO(P_r \odot P_s) &= r(s-3)\sqrt{3^2 + 3^2 + 3^2} + (r-3)\sqrt{3(s+2)^2} \\
&\quad + 2\sqrt{2^2 + 3^2 + 2 \times 3} + 4\sqrt{(s+1)^2 + 2^2 + 2(s+1)} \\
&\quad + 2(r-1)\sqrt{(s+2)^2 + 2^2 + 2(s+2)} + 2(s-2) \\
&\quad \sqrt{(s+1)^2 + 3^2 + 3(s+1)} + (r-2)(s-2) \\
&\quad \sqrt{(s+2)^2 + 3^2 + 3(s+2)} + 2\sqrt{(s+1)^2 + (s+2)^2 + (s+1)(s+2)}.
\end{aligned}$$

□

Theorem 2.8. *The Elliptic Sombor index and Euler Sombor index of corona product of two cycles $(C_r \odot C_s)$ are given by*

$$ESO(C_r \odot C_s) = 18rs\sqrt{2} + rs(s+5)\sqrt{3^2 + (s+2)^2} + 2\sqrt{2}r(s+2)^2.$$

$$EU(C_r \odot C_s) = 3\sqrt{3} + rs\sqrt{3^2 + (s+2)^2 + 3(s+2)} + \sqrt{3}r(s+2).$$

Proof. The graph $C_r \odot C_s$ with order $rs + r$ and size $2rs + r$ has two types of vertices, such as 3-degree vertex and $s+2$ degree vertex. The edges of $C_r \odot C_s$ can be partitioned into three types.

$$\begin{aligned}
ESO(C_r \odot C_s) &= rs(3+3)\sqrt{3^2 + 3^2} + rs(s+2+3)\sqrt{(s+2)^2 + 3^2} \\
&\quad + r(s+2+s+2)\sqrt{2(s+2)^2},
\end{aligned}$$

$$ESO(C_r \odot C_s) = 18\sqrt{2}rs + rs(s+5)\sqrt{3^2 + (s+2)^2} + 2\sqrt{2}r(s+2)^2.$$

$$EU(C_r \odot C_s) = rs\sqrt{3^2 + 3^2 + 3^2} + rs\sqrt{(s+2)^2 + 3^2 + 3(s+2)} + r\sqrt{3(s+2)^2},$$

$$EU(C_r \odot C_s) = 3\sqrt{3}rs + rs\sqrt{3^2 + (s+2)^2 + 3(s+2)} + \sqrt{3}r(s+2).$$

3 Conclusion

In this article, elliptic Sombor and Euler Sombor indices of join and corona product of certain graphs are obtained for standard graphs such as path, cycle and complete graphs. These results can be further extend for corona product and join of other graphs.

References

- [1] Harary, F. (1969) Graph Theory. Addison-Wesley Publishing Company, New York.
- [2] N. Trinajstić, Chemical Graph Theory, CRC Press, Boca Raton, FL. 1992.
- [3] Gutman, I. (2013). Degree-based topological indices. *Croatica chemica acta*, 86(4), 351-361.
- [4] Milovanović, I., Milovanović, E., & Matejić, M. (2021). On some mathematical properties of Sombor indices. *Bull. Int. Math. Virtual Inst*, 11(2), 341-353.
- [5] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, *MATCH Commun. Math. Comput. Chem.*, 86(2021) 11-16.
- [6] Gutman, I., Furtula, B., & Oz, M. S. (2024). Geometric approach to vertex-degree-based topological indices—Elliptic Sombor index, theory and application. *International Journal of Quantum Chemistry*, 124(2), e27346.
- [7] Ivan, G. (2024). Relating Sombor and Euler indices. *Vojnotehnički glasnik*, 72(1), 1-12.
- [8] Adhikari, B., Singh, A., & Yadav, S. K. (2023). Corona product of signed graphs and its application to modeling signed networks. *Discrete Mathematics, Algorithms and Applications*, 15(01), 2250062.
- [9] Agnes, V. S., & Kannadasan, C. (2023). Y-index of Different Corona Products of Graphs. *Communications in Mathematics and Applications*, 14(1), 131.

- [10] Khalid, A., Kausar, N., Munir, M., Gulistan, M., Al-Shamiri, M. M., & Lamoudan, T. (2022). Topological indices of families of bistar and corona product of graphs. *Journal of Mathematics*, 2022(1), 3567824.
- [11] Sheeja, P. G., Ranjini, P. S., Lokesha, V., & Cevik, A. S. (2021). Computation of the sk index over different corona products of graphs. *Palestine Journal of Mathematics*, 10(1), 8-16.
- [12] Iswadi, H., Baskoro, E. T., & Simanjuntak, R. (2011). On the metric dimension of corona product of graphs. *Far East Journal of Mathematical Sciences (FJMS)*, 52(2), 155-170.
- [13] Kumara, D., Mahasinghe, A., & Erandi, H. (2021). On the ABC and GA indices of the corona products of some graphs.
- [14] Arif, N. E., Karim, A. H., & Hasni, R. (2022). Sombor index of some graph operations. *International Journal of Nonlinear Analysis and Applications*, 13(1), 2561-2571.
- [15] Frucht, R., & Harary, F. (1970). On the corona of two graphs.
- [16] Mishra, V. N., Delen, S., & Cangul, I. N. (2019). Degree sequences of join and corona products of graphs. *Electron J Math Anal Appl*, 7(1), 5-13.
- [17] Gao, W., Iqbal, Z., Ishaq, M., Aslam, A., Aamir, M., & Binyamin, M. A. (2019). Bounds on Topological Descriptors of the Corona Product of F -Sum of Connected Graphs. *IEEE Access*, 7, 26788-26796.
- [18] Ali, N., Kousar, Z., Safdar, M., Tolasa, F. T., & Suleiman, E. (2023). Mapping Connectivity Patterns: Degree-Based Topological Indices of Corona Product Graphs. *Journal of Applied Mathematics*, 2023(1), 8975497.