

A LINEAR PROGRAMMING BOUND FOR SUM-RANK METRIC CODES

AIDA ABIAD, ALEXANDER L. GAVRILYUK, ANTONINA P. KHRAMOVA,
AND ILIA PONOMARENKO

ABSTRACT. We derive a linear programming bound on the maximum cardinality of error-correcting codes in the sum-rank metric. Based on computational experiments on relatively small instances, we observe that the obtained bounds outperform all previously known bounds.

Keywords: sum-rank metric code, linear programming bound, sum-rank metric graph, association scheme, eigenvalues

1. INTRODUCTION

Sum-rank metric codes have received a considerable amount of attention over the last years, especially because of their performance of multi-shot network coding; see [2, 5, 7, 9, 10, 13, 26–35]. A sum-rank metric code is a subset of matrix tuples of fixed dimensions, with every matrix being defined over the same finite field \mathbb{F}_q (see Definition 2.1). The distance between two such tuples is the sum of ranks of differences between respective elements of the tuples. As such, it is a generalization of both rank-metric codes and Hamming codes.

A central question in coding theory, concerning the maximum cardinality of a code with a given minimum distance, has recently been addressed in the context of the sum-rank metric. In [9], properties of sum-rank metric codes are investigated, and several bounds based on classical coding theory approaches are shown. In [2], the authors consider the sum-rank metric space from an algebraic graph theoretical point of view, introduce a sum-rank metric graph, and propose an eigenvalue bound, referred to as Ratio-type bound, by using a connection between the cardinality of a code with a minimum distance $k + 1$ and the k -independence number of the graph. This bound is calculated using the eigenvalues of the adjacency matrix of the sum-rank metric graph. An explicit closed formula is known for the case when the minimum distance is 3 [1] and 4 [23], and otherwise it can be calculated via a linear programming (LP) method which uses the so-called minor polynomials [19]. A geometrical approach to sum-rank metric codes using subspace designs is investigated in [37, Section 5].

While a classical approach to estimate the maximum cardinality of a code is to use Delsarte’s LP method (which was first introduced in the context of the Hamming metric [14]), this has not yet been developed for sum-rank metric codes. The idea behind this method is to consider a code as a subset of points of an association scheme, and then formulate an optimization problem where the objective is to maximize the size of a code subject to linear constraints derived by leveraging the properties of the association scheme. By solving this linear program, one can obtain upper bounds on the number of codewords. This, combined with the duality

of linear programming, provides one of the most powerful methods for bounding the size of codes arising from association schemes. Delsarte's LP approach has already been successfully applied to several metrics, *e.g.*, for Hamming codes [14], rank-metric codes [15], bilinear alternating forms [16], permutation codes [17], and Lee codes [3, 39], but also for newer metrics like subspace codes [36].

Despite its previous success, the LP method has not yet been applied to the sum-rank metric. This is perhaps due to the fact that, unlike for the Hamming or rank metric, the sum-rank metric does not give rise to an association scheme in a straightforward way by taking the distance relations between the codewords. In particular, except some special cases, the sum-rank metric graphs are not distance-regular [2, Proposition 11], unlike the Hamming graphs and the bilinear forms graphs corresponding to the rank-metric space.

In this paper, we develop an approach to utilizing the Delsarte's LP method to estimate the maximum cardinality of a sum-rank metric code with a given minimum distance. From computational experiments for small values of the parameters, we observe that the new LP bound outperforms all previously known bounds for sum-rank metric codes given in [9] and [2]. To prove our results, we describe a way to construct an association scheme for a sum-rank metric space. We do so by making use of the structure of the sum-rank metric graph as a Cartesian product of smaller rank-metric graphs [2, Proposition 9], and considering a direct product of association schemes of the Cartesian factors. Moreover, we investigate the coherent closure of a sum-rank metric graph and discuss conditions under which it is equivalent to the association scheme we construct.

The structure of this paper is as follows. Section 2 provides the preliminaries for the sum-rank metric space, known bounds on the size of sum-rank metric codes, as well as reminds some necessary background on association schemes and Delsarte's LP method. Section 3 introduces a way to construct a suitable association scheme for a sum-rank metric graph using the notion of a direct product of association schemes. Such association scheme is then used to derive the LP bound. In addition, particular attention is given to the coherent closure of the graph, and the conditions under which it coincides with the direct product of the coherent closures of the Cartesian factors of the graph. These conditions are also discussed in the more general context of graphs that can be seen as a Cartesian product. Finally, Section 4 contains a computational comparison of the new LP bound against all the previously known bounds for sum-rank codes.

2. PRELIMINARIES

In this section, we will recall some definitions and results on sum-rank metric codes and association schemes.

2.1. Basic notation. For a positive integer n , let $[n]$ denote the set $\{1, \dots, n\}$ and put $\llbracket n \rrbracket := \{0\} \cup [n]$.

Let Ω be a finite set and R a binary relation on Ω . The **adjacency matrix** of R is a matrix $A \in \mathbb{R}^{\Omega \times \Omega}$ defined as follows: $(A)_{\alpha, \beta} = 1$ if $(\alpha, \beta) \in R$ and 0 otherwise.

By a **graph** we mean a finite simple undirected graph, *i.e.*, a pair $\Gamma = (\Omega, E)$ of a finite set Ω of vertices and a set E of unordered pairs from Ω , $E \subseteq \binom{\Omega}{2}$, called edges. Note that the edge set E can also be identified with an irreflexive symmetric (**adjacency**) relation on Ω . Then the adjacency matrix of a graph Γ is the adjacency matrix of E .

Recall that the **Cartesian product** of graphs $\Gamma_1 = (\Omega_1, E_1)$ and $\Gamma_2 = (\Omega_2, E_2)$ is the graph¹ $\Gamma_1 \square \Gamma_2$ whose vertex set is $\Omega_1 \times \Omega_2$, where vertices (α_1, α_2) and (β_1, β_2) are adjacent if either $\alpha_1 = \beta_1$ and $(\alpha_2, \beta_2) \in E_2$, or $\alpha_2 = \beta_2$ and $(\alpha_1, \beta_1) \in E_1$. If A_i , $i = 1, 2$, denotes the adjacency matrix of a graph Γ_i , and A is the adjacency matrix of $\Gamma_1 \square \Gamma_2$, then

$$A = A_1 \otimes I_{\Omega_2} + I_{\Omega_1} \otimes A_2, \quad (1)$$

where \otimes means the Kronecker product of matrices, and I_{Ω} is the identity matrix of $\mathbb{R}^{\Omega \times \Omega}$. This definition can be inductively extended to a Cartesian product of any finite number of graphs.

2.2. Sum-rank metric and sum-rank metric graphs. Let t be a positive integer and let $\mathbf{n} = (n_1, \dots, n_t)$, $\mathbf{m} = (m_1, \dots, m_t)$ be tuples of positive integers with $m_1 \geq m_2 \geq \dots \geq m_t$, and $m_i \geq n_i$ for all $i \in [t]$.

For a prime power q and positive integers $m \geq n$, let $\mathbb{F}_q^{n \times m}$ denote the vector space of all $n \times m$ matrices over the finite field \mathbb{F}_q . Denote by $\text{rk}(M)$ the rank of a matrix $M \in \mathbb{F}_q^{n \times m}$.

Definition 2.1. *The **sum-rank metric space** is an \mathbb{F}_q -linear vector space $\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ defined as follows:*

$$\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}} := \mathbb{F}_q^{n_1 \times m_1} \times \dots \times \mathbb{F}_q^{n_t \times m_t},$$

where \times stands for the direct product of vector spaces. The **sum-rank** of an element $X = (X_1, \dots, X_t) \in \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ is $\text{srk}(X) = \sum_{i=1}^t \text{rk}(X_i)$. The **sum-rank distance (metric)** between $X, Y \in \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ is $\text{srk}(X - Y)$. In case $t = 1$ we sometimes refer to the sum-rank distance as simply the **rank distance**.

It is easy to see that the sum-rank distance is indeed a distance on $\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$.

Definition 2.2. *A **sum-rank metric code** is a non-empty subset $\mathcal{C} \subseteq \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$. The **minimum sum-rank distance** of a code \mathcal{C} with $|\mathcal{C}| \geq 2$ is defined by*

$$\text{srk}(\mathcal{C}) := \min \{ \text{srk}(X - Y) : X, Y \in \mathcal{C}, X \neq Y \}.$$

The following definition, which is introduced in [2], allows one to study sum-rank metric codes from the graph-theoretical perspective.

Definition 2.3. *The **sum-rank metric graph** $\Gamma(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}})$ is a graph whose vertex set is $\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$, and two vertices X, Y are adjacent if and only if $\text{srk}(X - Y) = 1$.*

We sometimes omit $\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ and simply write Γ when it is clear which sum-rank-metric space the graph corresponds to.

A simple yet important observation is that, for any two vertices X, Y of the graph $\Gamma(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}})$, the geodesic distance between them coincides with the sum-rank distance $\text{srk}(X - Y)$ [10, Proposition 4.3].

The following two special cases covered by Definition 2.1 have been long known and well-studied in the coding theory literature: in case $n_i = m_i = 1$ for all $i \in [t]$, it is equivalent to the Hamming metric on the Hamming space \mathbb{F}_q^t , while in case $t = 1$, the sum-rank metric reduces to the rank metric on the space $\mathbb{F}_q^{n \times m}$. Hence the sum-rank metric is a generalization of both the Hamming metric and the rank metric. In the former case, the corresponding sum-rank metric graphs are well known in the literature as Hamming graphs, denoted $H(t, q)$, while in the latter case those are

¹Note that in [2], the notation $\Gamma_1 \times \Gamma_2$ was used to denote the Cartesian product. However, \square appears to be more common, e.g., see [22].

bilinear forms graphs. Recall that the **bilinear forms graph**, sometimes denoted by $\text{Bil}_q(n, m)$, is a graph with $\mathbb{F}_q^{n \times m}$ as the vertex set, where two vertices (matrices) are adjacent if and only if the rank distance between them is exactly one.

A graph Γ with diameter D is said to be **distance-regular** if for all $i, j, k \in \llbracket D \rrbracket$ and for each pair of vertices x and y at distance k from each other the number of vertices that are at distance i from x and at distance j from y is equal to a constant $p_{i,j}^k$ that does not depend on the choice of vertices x and y .

Note that both Hamming graphs and bilinear forms graphs are distance-regular, see [8].

Proposition 2.4. [2, Proposition 9] *Let $\Gamma_i = \Gamma(\mathbb{F}_q^{n_i \times m_i})$ for $i \in [t]$. Then Γ_i is isomorphic to the bilinear forms graph $\text{Bil}_q(n_i, m_i)$ and the sum-rank metric graph $\Gamma(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}})$ is the Cartesian product $\Gamma_1 \square \dots \square \Gamma_t$.*

Graphs that are Cartesian products of t distance-regular graphs are examples of t -distance-regular graphs considered, e.g., in [6].

Note that in case $n_i = m_i = 1$ for all $i \in [t]$, the graph $\text{Bil}_q(1, 1)$ is a complete graph on q vertices, and $\Gamma(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}})$ is the Cartesian product of t complete graphs on q vertices, which is the Hamming graph $H(t, q)$.

2.3. Coding theory bounds. Several bounds on the maximum size of a sum-rank metric code $\mathcal{C} \subseteq \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ with $|\mathcal{C}| \geq 2$ and a given minimum distance $\text{srk}(\mathcal{C}) \geq d$ were introduced in [9]. The first four bounds are induced by the following connection to the Hamming-metric case: $|\mathcal{C}|$ is upper-bounded by the size of a Hamming code over \mathbb{F}_{q^m} of length N and minimum distance at least d , where $m = \max_{i \in [t]} m_i$ and $N = \sum_{i \in [t]} n_i$.

Theorem 2.5 (see [9, Theorem III.1]). *Let $m = \max_{i \in [t]} m_i$ and let $\mathcal{C} \subseteq \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ be a sum-rank metric code with $|\mathcal{C}| \geq 2$ and $\text{srk}(\mathcal{C}) \geq d$. The following hold:*

Induced Singleton bound: $|\mathcal{C}| \leq q^{m(N-d+1)},$

Induced Hamming bound: $|\mathcal{C}| \leq \left\lfloor \frac{q^{mN}}{\sum_{s=0}^{\lfloor (d-1)/2 \rfloor} \binom{N}{s} (q^m - 1)^s} \right\rfloor,$

Induced Plotkin bound: $|\mathcal{C}| \leq \left\lfloor \frac{q^m d}{q^m d - (q^m - 1)N} \right\rfloor$ if $d > (q^m - 1)N/q^m,$

Induced Elias bound: $|\mathcal{C}| \leq \left\lfloor \frac{Nd(q^m - 1)}{q^m w^2 - 2Nw(q^m - 1) + (q^m - 1)Nd} \cdot \frac{q^{mN}}{V_w(\mathbb{F}_{q^m}^N)} \right\rfloor.$

In the Induced Elias bound, w is any integer between 0 and $N(q^m - 1)/q^m$ such that the denominator is positive, and $V_w(\mathbb{F}_{q^m}^N) = \sum_{i=0}^w \binom{N}{i} (q^m - 1)^i$, i.e., the cardinality of any ball of radius w in $\mathbb{F}_{q^m}^N$ with respect to the Hamming distance.

The other four bounds presented in [9] are not induced by the Hamming-metric case and are specific to the sum-rank metric.

Theorem 2.6 (see [9, Theorems III.2, III.6–III.8]). *Let $\mathcal{C} \subseteq \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}$ be a sum-rank metric code with $|\mathcal{C}| \geq 2$ and $\text{srk}(\mathcal{C}) \geq d$. Let j and δ be the unique integers satisfying $d - 1 = \sum_{i=1}^{j-1} n_i + \delta$ and $0 \leq \delta \leq n_j - 1$. Let $\ell \leq t - 1$ and $\delta' \leq n_{\ell+1} - 1$ be the unique positive integers such that $d - 3 = \sum_{j=1}^{\ell} n_j + \delta'$. Define*

$\mathbf{n}' = (n_{\ell+1} - \delta', n_{\ell+2}, \dots, n_t)$ and $\mathbf{m}' = (m_{\ell+1}, m_{\ell+2}, \dots, m_t)$. Finally, let $Q = \sum_{i=1}^t q^{-m_i}$. The following hold:

Singleton bound: $|\mathcal{C}| \leq q^{\sum_{i=j}^t m_i n_i - m_j \delta},$

Sphere-Packing bound: $|\mathcal{C}| \leq \left\lfloor \frac{|\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}|}{V_r(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}})} \right\rfloor,$ where $r = \lfloor (d-1)/2 \rfloor,$

Projective Sphere-Packing bound: $|\mathcal{C}| \leq \left\lfloor \frac{|\mathbb{F}_q^{\mathbf{n}' \times \mathbf{m}'}|}{V_1(\mathbb{F}_q^{\mathbf{n}' \times \mathbf{m}'})} \right\rfloor$ if $3 \leq d \leq N,$

Total Distance bound: $|\mathcal{C}| \leq \frac{d-N+t}{d-N+Q}$ if $d > N - Q.$

In the Sphere-Packing and Projective Sphere-Packing bounds, the denominator denotes the cardinality of any ball in the sum-rank metric of radius r . For example,

$$V_r(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}}) = |\{(X_1, \dots, X_t) \in \mathbb{F}_q^{\mathbf{n} \times \mathbf{m}} \mid \text{srk}(X_1, \dots, X_t) \leq r\}|,$$

and a closed formula is provided in [9].

2.4. Eigenvalue bounds. In [2, Corollary 16] it was shown that any upper bound on the k -independence number of a graph yields an upper bound on the size of a sum-rank metric code with minimum distance $k + 1$, and vice versa. The next result provides the main bound we will be using to compare our new LP bound with; the so-called Ratio-type LP bound, which was recently proposed to be used for sum-rank metric codes in [2].

From here on, let $\mathbb{R}_k[x]$ denote the set of polynomials in variable x with coefficients in \mathbb{R} of degree at most k .

Theorem 2.7 (Ratio-type bound; see [1]). *Let $\Gamma = (\Omega, E)$ be a regular graph with n vertices and adjacency matrix A with eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$. Let $p \in \mathbb{R}_k[x]$. Define $W(p) = \max_{u \in \Omega} \{p(A)_{uu}\}$ and $\lambda(p) = \min_{i=2, \dots, n} \{p(\lambda_i)\}$. Then*

$$\alpha_k(\Gamma) \leq n \frac{W(p) - \lambda(p)}{p(\lambda_1) - \lambda(p)}. \quad (2)$$

In cases $k = 2$ and $k = 3$, the best polynomial $p \in \mathbb{R}_k[x]$ that minimizes the bound in Theorem 2.7 was shown in [1] and [23], respectively.

Theorem 2.8 (Ratio-Type bound, $k = 2$; see [1]). *Let Γ be a regular graph with n vertices and $r \geq 2$ distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_r$ of the adjacency matrix. Let θ_i be the largest eigenvalue such that $\theta_i \leq -1$. Then*

$$\alpha_2(\Gamma) \leq n \frac{\theta_0 + \theta_i \theta_{i-1}}{(\theta_0 - \theta_i)(\theta_0 - \theta_{i-1})}.$$

Moreover, this is the best possible bound that can be obtained by choosing a polynomial via Theorem 2.7.

Theorem 2.9 (Ratio-Type bound, $k = 3$; see [23]). *Let Γ be a regular graph with the vertex set Ω , $|\Omega| = n$, and with $r \geq 3$ distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_r$ of the adjacency matrix. Let s be the largest index such that $\theta_s \geq -\frac{\theta_0^2 + \theta_0 \theta_r - \Delta}{\theta_0(\theta_r + 1)}$, where $\Delta = \max_{u \in \Omega} \{(A^3)_{uu}\}$. Then*

$$\alpha_3(\Gamma) \leq n \frac{\Delta - \theta_0(\theta_s + \theta_{s+1} + \theta_r) - \theta_s \theta_{s+1} \theta_r}{(\theta_0 - \theta_s)(\theta_0 - \theta_{s+1})(\theta_0 - \theta_r)}.$$

Moreover, this is the best possible bound that can be obtained by choosing a polynomial via Theorem 2.7.

The following lemma is a new contribution to the mentioned eigenvalue bound; it provides conditions under which the Ratio-type bound of Theorem 2.7 is tight.

Lemma 2.10. *Let Γ , $W(p)$, $\lambda(p)$ be as in Theorem 2.7. If the Ratio-type bound of Theorem 2.7 is met by a k -independent set U of the graph Γ , then the following conditions hold:*

- (i) $W(p) = \frac{1}{|U|} \sum_{u \in U} (p(A))_{uu}$,
- (ii) Any vertex $u \in U$ has $W(p)$ neighbors in U ; in particular, the induced subgraph on vertices of U is regular,
- (iii) Any vertex $v \notin U$ has $W(p) - \lambda(p)$ neighbors in U .

Proof. The proof follows directly from analyzing the inequality that arises as a part of the proof of [1, Theorem 3.2], which we now give. Let A denote the adjacency matrix of Γ with adjacency eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$. Let $\mu_1 \geq \mu_2$ denote the eigenvalues of the quotient matrix of the partition of Ω into U and $\Omega \setminus U$:

$$\begin{pmatrix} \frac{1}{r} \sum_{u \in U} (p(A))_{uu} & p(\lambda_1) - \frac{1}{r} \sum_{u \in U} (p(A))_{uu} \\ \frac{rp(\lambda_1) - \sum_{u \in U} (p(A))_{uu}}{n-r} & p(\lambda_1) - \frac{rp(\lambda_1) - \sum_{u \in U} (p(A))_{uu}}{n-r} \end{pmatrix}.$$

Finally, let $\mu_1 = p(\lambda_1)$, and $r = |U| = \alpha_k(\Gamma)$. Then,

$$\lambda(p) \leq \mu_2 \tag{3}$$

$$\begin{aligned} &= \frac{1}{r} \sum_{u \in U} (p(A))_{uu} - \frac{rp(\lambda_1) - \sum_{u \in U} (p(A))_{uu}}{n-r} \\ &\leq W(p) - \frac{rp(\lambda_1) - rW(p)}{n-r}. \end{aligned} \tag{4}$$

The explicit bound on r is then deduced from the inequality between $\lambda(p)$ and the final expression.

In case the Ratio-type bound is tight, the inequalities (3) and (4) must be equalities. In particular, it means that

$$W(p) = \frac{1}{r} \sum_{u \in U} (p(A))_{uu},$$

which gives (i). The inequality (3) being an equality means that the interlacing of eigenvalues is tight, and by [21, Corollary 2.3] it follows that the partition of Γ into U and $\Omega \setminus U$ is equitable (regular). It follows that each vertex from $\Omega \setminus U$ is adjacent to

$$\frac{rp(\lambda_1) - \sum_{u \in U} (p(A))_{uu}}{n-r} = \frac{rp(\lambda_1) - rW(p)}{n-r} = W(p) - \lambda(p)$$

vertices from U , which gives (iii), and each vertex from U is adjacent to

$$\frac{1}{r} \sum_{u \in U} (p(A))_{uu} = W(p)$$

vertices from U , which gives (ii). □

A graph Γ is **l -partially walk-regular** if for any vertex v and any positive integer $i \leq l$ the number of closed walks of length i does not depend on the choice of v . We also say that Γ is **partially walk-regular** if it is l -partially walk-regular for some l , and that Γ is **walk-regular** if it is l -partially walk-regular for any l .

Remark 2.11. *The condition (i) of Lemma 2.10 always holds in a partially walk-regular graph, since all entries of the diagonal of $p(A)$ have the same value. In particular, this condition holds if Γ is a sum-rank metric graph, see [2, Proposition 11].*

For a fixed k , the challenge behind applying the Ratio-type bound is to find a polynomial of degree k which minimizes the right-hand of Eq. (2). This was resolved for $k \in \{2, 3\}$ resulting in closed formulæ presented in Theorems 2.8 and 2.9. However, even in the case $k = 3$ finding the polynomial in general turns out to be an involved problem, with an entire paper devoted to it [23]. In [19], an LP implementation using the so-called minor polynomials was proposed for finding, for a given k and a k -partially walk-regular graph Γ , the polynomial p that optimizes the Ratio-type bound from Theorem 2.7. We will use such LP to compute the Ratio-type bound and compare it with the new Delsarte's LP bound. In the LP for the Ratio-type bound, the inputs are the distinct adjacency eigenvalues of the graph Γ , denoted $\theta_0 > \dots > \theta_r$, with respective multiplicities $m(\theta_i)$, $i \in \{0, \dots, r\}$. The **minor polynomial** $f_k \in \mathbb{R}_k[x]$ is a polynomial that minimizes $\sum_{i=0}^r m(\theta_i) f_k(\theta_i)$. Let $p = f_k$ be defined by $f_k(\theta_0) = x_0 = 1$ and $f_k(\theta_i) = x_i$ for $i \in \{1, \dots, r\}$, where the vector (x_1, \dots, x_r) is a solution of the following linear program:

$$\begin{array}{ll} \text{minimize} & \sum_{i \in [r]} m(\theta_i) x_i \\ \text{subject to} & f[\theta_0, \dots, \theta_s] = 0, \quad s = k+1, \dots, r, \\ & x_i \geq 0, \quad i \in [r]. \end{array} \quad (5)$$

Here, $f[\theta_0, \dots, \theta_m]$ denote m -th divided differences of Newton interpolation, recursively defined by

$$f[\theta_i, \dots, \theta_j] = \frac{f[\theta_{i+1}, \dots, \theta_j] - f[\theta_i, \dots, \theta_{j-1}]}{\theta_j - \theta_i},$$

where $j > i$, starting with $f[\theta_i] = x_i$ for $i \in \{0, \dots, r\}$. In [19], it was shown that, for $k = 3$, using the LP (5) it is possible to obtain tight bounds for every Hamming graph $H(r, 2)$, and it was also shown that for these graphs the Ratio-type bound coincides with Delsarte's LP bound [14]; see also Section 5.

2.5. Association schemes. The basic theory of association schemes and their Bose-Mesner algebras, given in this subsection for the sake of completeness, are standard and can be found in more detail, *e.g.*, in [8, Chapter 2], [14], or [20].

Definition 2.12. *A (symmetric) D -class association scheme $\mathcal{X} = (\Omega, \mathbf{R})$ is a finite set Ω (of points) together with a collection $\mathbf{R} = \{R_i \mid i \in [D]\}$ of non-empty binary relations on Ω , satisfying the following four conditions:*

- (i) \mathbf{R} is a partition of $\Omega \times \Omega$.
- (ii) R_0 is the diagonal of $\Omega \times \Omega$.
- (iii) Each binary relation R_i equals its transpose (converse).
- (iv) For all $i, j, k \in [D]$ and $(\alpha, \beta) \in R_k$, the number of $\gamma \in \Omega$ such that $(\alpha, \gamma) \in R_i$ and $(\gamma, \beta) \in R_j$ is a constant denoted by $p_{i,j}^k$ that does not depend on the choice of (α, β) .

Throughout this paper, we refer to a symmetric D -class association scheme simply as association scheme (informally, we will occasionally call an association scheme just a scheme). By the relations of \mathcal{X} we mean the elements of \mathbb{R} . The number of relations $D + 1$ and the constants $p_{i,j}^k$ are called the **rank** and the **intersection numbers** of a scheme, respectively.

The following are two well-known examples of association schemes, which provide an algebraic framework to study codes in the Hamming and rank metrics.

Definition 2.13. *Let n and q be positive integers. The **Hamming scheme** is an association scheme $([n]^q, \mathbb{R})$ of rank $n + 1$, where all pairs $x, y \in [n]^q$ with Hamming distance equal to i are in the relation $R_i \in \mathbb{R}$, $i \in \llbracket n \rrbracket$.*

Definition 2.14. *Let n, m be positive integers with $n \leq m$ and q be a prime power. The **bilinear forms scheme** is an association scheme $(\mathbb{F}_q^{n \times m}, \mathbb{R})$ of rank $n + 1$, where all pairs $A, B \in \mathbb{F}_q^{n \times m}$ with $\text{rk}(A - B) = i$ are in the relation $R_i \in \mathbb{R}$, $i \in \llbracket n \rrbracket$.*

Let $\mathcal{X} = (\Omega, \mathbb{R})$ be an association scheme, and $A_i \in \mathbb{R}^{\Omega \times \Omega}$ denote the adjacency matrix of $R_i \in \mathbb{R}$. Definition 2.12, which is stated in terms of binary relations on Ω , can also be rewritten in terms of the adjacency matrices that they represent:

- (i) $\sum_{i \in I} A_i = J_\Omega$ (the all-ones matrix).
- (ii) $A_0 = I_\Omega$ (the identity matrix).
- (iii) $A_i = A_i^\top$ for all $i \in \llbracket D \rrbracket$.
- (iv) $A_i A_j = \sum_{k \in \llbracket D \rrbracket} p_{i,j}^k A_k$ for all $i, j \in \llbracket D \rrbracket$.

By (iv) above, the matrices A_i form a basis for a $(D + 1)$ -dimensional matrix algebra over \mathbb{C} , called the **Bose-Mesner algebra** of \mathcal{X} . This algebra is closed under matrix multiplication, Schur (entrywise) multiplication, and transposition, which thus makes it an example of a **coherent algebra**. Since the matrices A_i , $i \in \llbracket D \rrbracket$, pairwise commute, it follows that \mathbb{R}^Ω decomposes into the orthogonal direct sum of their common maximal eigenspaces.

The Bose-Mesner algebra of \mathcal{X} has two distinctive linear bases, namely, the one consisting of the adjacency matrices A_i and another one of the so-called **primitive idempotents** E_j , which are the orthogonal projection matrices onto maximal common eigenspaces of the adjacency matrices. In particular, $E_i E_j$ is the zero matrix if $i \neq j$ and $E_i^2 = E_i$ for all $i, j \in \llbracket D \rrbracket$. For expressing each basis in terms of the other, define the constants P_{ji} and Q_{ij} as

$$A_i = \sum_{j \in \llbracket D \rrbracket} P_{ji} E_j \quad \text{and} \quad E_i = \frac{1}{|\Omega|} \sum_{j \in \llbracket D \rrbracket} Q_{ij} A_j.$$

Now, the corresponding change-of-basis matrices $P = (P_{ji})$ and $Q = (Q_{ij})$ of order $D + 1$ are respectively called the **first and second eigenmatrices** (or character tables) of the Bose-Mesner algebra (or of the scheme). In particular, the i -th column of P consists of the eigenvalues of A_i , i.e., P_{ji} is the eigenvalue of A_i on the j -th maximal common eigenspace. Furthermore, note that $Q = |\Omega| \cdot P^{-1}$.

Next, we briefly (see [8, Chapter 9] for further details) recall how to compute the P and Q matrices for the bilinear forms scheme from Definition 2.14 with $D = n$. First, observe that the adjacency matrix A_1 is the adjacency matrix of the bilinear forms graph $\text{Bil}_q(n, m)$ and has $D + 1$ distinct eigenvalues given by

$$\theta_j = \frac{1}{q-1} \left((q^{n-j} - 1)(q^m - q^j) - q^j + 1 \right), \quad j \in \llbracket D \rrbracket.$$

Second, as the bilinear forms scheme is P -polynomial (see [8, 14]), for every $i \in \llbracket D \rrbracket$, the adjacency matrix A_i equals $p_i(A_1)$, where p_i is a polynomial of degree i defined by the following recursive equation:

$$p_i = \frac{1}{c_i} ((x - a_{i-1})p_{i-1} - b_{i-2}p_{i-2}), \quad i \in \llbracket D \rrbracket,$$

where $p_0 = 1$, $p_1(x) = x$, and

$$\begin{aligned} b_i &:= p_{1,i+1}^i = \frac{q^{2i}(q^{m-i} - 1)(q^{n-i} - 1)}{q - 1}, \\ c_i &:= p_{1,i-1}^i = \frac{q^{i-1}(q^i - 1)}{q - 1}, \\ a_i &:= p_{1,i}^i = b_0 - b_i - c_i. \end{aligned}$$

Furthermore, the bilinear forms scheme is self-dual (see [8, Section 6.1]), which, in particular, means that the matrices P and Q are equal. Therefore,

$$P_{ji} = Q_{ji} = p_i(\theta_j), \quad i, j \in \llbracket D \rrbracket. \quad (6)$$

For more information on bilinear forms scheme, the reader is referred to [14, 15].

2.6. Delsarte's linear programming bound. In this subsection we briefly recall Delsarte's linear programming method [14].

Let $\mathcal{X} = (\Omega, \mathbb{R})$ be an association scheme of rank $D + 1$. Let Δ be a non-empty subset of Ω . The (inner) **distribution vector** of Δ is a vector $\mathbf{a} = (a_0, \dots, a_D)$ with entries

$$a_i = \frac{|(\Delta \times \Delta) \cap R_i|}{|\Delta|}$$

for every $R_i \in \mathbb{R}$. It is clear that $a_i \geq 0$ for all $i \in \llbracket D \rrbracket$, $\sum_{i \in \llbracket D \rrbracket} a_i = |\Delta|$, and the distribution vector is normalized so that $a_0 = 1$.

A key observation for Delsarte's LP bound is the following result.

Theorem 2.15. [14, Theorem 3.3] *With the above notation, the distribution vector \mathbf{a} satisfies*

$$\mathbf{a}Q \geq \mathbf{0},$$

where Q is the second eigenmatrix of \mathcal{X} .

Then a linear programming bound for a code in \mathcal{X} is carried by Theorem 2.15 as follows. For a subset of indices $M \subseteq \llbracket D \rrbracket$, an M -**code** is a subset $\Delta \subseteq \Omega$ such that any two distinct points of Δ are in R_i for some $i \in M$. In other words, Δ is an M -code if $a_i = 0$ for all $i \notin M$. Consider the following LP formulation:

$$\boxed{\begin{array}{ll} \text{maximize} & \sum_{i \in \llbracket D \rrbracket} a_i \\ \text{subject to} & \mathbf{a}Q \geq \mathbf{0}, \\ & \mathbf{a} \geq \mathbf{0}, \\ & a_0 = 1, \\ & a_i = 0, \quad i \notin M. \end{array}} \quad (7)$$

It follows from Theorem 2.15 that the cardinality $|\Delta|$ of such an M -code is upper bounded by the solution of the LP (7).

3. AN ASSOCIATION SCHEME FOR SUM-RANK METRIC GRAPHS

In this section, we define and study an association scheme related to sum-rank metric graphs. Recall that, in the case of the Hamming or bilinear forms association schemes (see Definitions 2.13, 2.14), the adjacency matrix A_1 of the corresponding graph generates the respective Bose-Mesner algebra. In general, unlike these two partial cases, the sum-rank distance itself does not directly determine an association scheme. However, one can define an association scheme based on the fact that the sum-rank metric graph is constructed as the Cartesian product of bilinear forms graphs [2, Proposition 9]. This scheme may possibly be larger (in the sense of Definition 3.3 below, see the discussion in Section 3.3) than we need, but it suffices to estimate Delsarte's LP bound on the size of sum-rank metric codes.

3.1. Direct product of association schemes. The concept of a direct product of association schemes is well known; here we follow the definition proposed in [18].

Definition 3.1. *Let $\mathcal{X}_i = (\Omega_i, \mathbf{R}_i)$ be an association scheme of rank D_i with relations $\mathbf{R}_i = \{R_j^i \mid j \in \llbracket D_i \rrbracket\}$, $i \in [t]$. Then the **direct product** $\mathcal{X}_1 \otimes \cdots \otimes \mathcal{X}_t$ is an association scheme $\mathcal{X} = (\Omega, \mathbf{R})$ where:*

- (i) $\Omega = \Omega_1 \times \cdots \times \Omega_t$,
- (ii) $\mathbf{R} = \{R_{(j_1, \dots, j_t)} \mid (j_1, \dots, j_t) \in \llbracket D_1 \rrbracket \times \cdots \times \llbracket D_t \rrbracket\}$, where

$$R_{(j_1, \dots, j_t)} = \{((\alpha_1, \dots, \alpha_t), (\beta_1, \dots, \beta_t)) \in \Omega^2 : (\alpha_i, \beta_i) \in R_{j_i}^i \in \mathbf{R}_i, i \in [t]\}.$$

The fact that the relations in \mathbf{R} indeed give an association scheme can be easily verified (see [25, Section 2.2]). In the more general context of coherent configurations [12, Section 3.2.2], which are not the focus of this paper (see Section 3.3), the direct product is called the **tensor product**.

Note that the adjacency matrix $A_{(j_1, \dots, j_t)}$ of the relation $R_{(j_1, \dots, j_t)} \in \mathbf{R}$ is the Kronecker product $A_{j_1}^1 \otimes \cdots \otimes A_{j_t}^t$ of the adjacency matrices $A_{j_i}^i$ with $j_i \in \llbracket D_i \rrbracket$, $i \in [t]$.

Lemma 3.2. [25] *In the notation of Definition 3.1, the second eigenmatrix Q of \mathcal{X} is the Kronecker product $Q_1 \otimes Q_2 \otimes \cdots \otimes Q_t$, where Q_i is the second eigenmatrix of \mathcal{X}_i , $i \in [t]$.*

Using Definition 3.1, we can consider an association scheme that contains the edge set of a sum-rank metric graph as a union of some of its relations.

Definition 3.3. *Let \mathcal{X} and \mathcal{Y} be association schemes on the same set of points. We say that $\mathcal{X} \leq \mathcal{Y}$ if and only if every relation of \mathcal{X} is a union of some relations of \mathcal{Y} . If this is the case, then \mathcal{X} is said to be a **fusion** (scheme) of \mathcal{Y} and \mathcal{Y} is a **fission** (scheme) of \mathcal{X} .*

The Bose-Mesner algebra of a fusion scheme \mathcal{X} is a linear subspace of the Bose-Mesner algebra of \mathcal{Y} . This may serve as an alternative definition of a fusion [20, Section 5.2]; note that what we call a fusion scheme is sometimes called a subscheme [20].

Note that the relation \leq is a partial order on the set of association schemes defined on the same point set Ω . For example, the association scheme of the complete graph on Ω is a fusion of any scheme on Ω ; in fact, it is the smallest scheme on Ω .

Lemma 3.4. *Let Γ denote the sum-rank metric graph $\Gamma(\mathbb{F}_q^{n \times m})$ and suppose that \mathcal{X}_i is the bilinear forms scheme on $\mathbb{F}_q^{n_i \times m_i}$, $i \in [t]$. Then the adjacency relation of Γ is a union of some relations of the direct product $\mathcal{X}_1 \otimes \cdots \otimes \mathcal{X}_t$.*

Proof. It follows directly from Proposition 2.4, Eq. (1), and Definition 3.1. \square

3.2. An association scheme for sum-rank metric graphs. We now define the association scheme \mathcal{X} of the sum-rank metric graph Γ as the smallest association scheme such that a union of some of its relations coincides with the adjacency relation of Γ (indeed, such a scheme exists by Lemma 3.4). Again, in the more general context of coherent configurations [12, Section 2.6.1], \mathcal{X} is the **coherent closure** (or **Weisfeiler-Leman closure**) of the graph Γ , denoted $\text{WL}(\Gamma)$, which can be computed by using the Weisfeiler-Leman algorithm [40]. From the above discussion and Lemma 3.4, one has the following immediate consequence.

Corollary 3.5. *Let Γ be a sum-rank metric graph, and let Γ_i , $i \in [t]$, be the t bilinear forms graphs that are the Cartesian factors of Γ . Then*

$$\mathcal{X} := \text{WL}(\Gamma) = \text{WL}(\Gamma_1 \square \cdots \square \Gamma_t) \leq \text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2) \otimes \cdots \otimes \text{WL}(\Gamma_t), \quad (8)$$

moreover, $\text{WL}(\Gamma_i)$ is a bilinear forms scheme, $i \in [t]$.

The relation “to be at distance ℓ in Γ ” is the union of all relations $R_{(j_1, \dots, j_t)}$ such that $j_1 + j_2 + \cdots + j_t = \ell$. Therefore, we can get an upper bound on the size of a sum-rank metric code with prescribed minimum distance d by applying Delsarte’s LP bound to a code in the direct product of bilinear forms schemes. Namely, we solve the LP (7) with the matrix Q given by Lemma 3.2 and Eq. (6) and the distribution vector $\mathbf{a} = (a_{(j_1, \dots, j_t)})$ indexed by $\llbracket D_1 \rrbracket \times \cdots \times \llbracket D_t \rrbracket$:

$$\begin{array}{ll} \text{maximize} & \sum_{i \in \llbracket D_1 \rrbracket \times \cdots \times \llbracket D_t \rrbracket} a_i \\ \text{subject to} & \mathbf{a}Q \geq \mathbf{0}, \\ & \mathbf{a} \geq \mathbf{0}, \\ & a_{(0, \dots, 0)} = 1, \\ & a_{(j_1, \dots, j_t)} = 0, \quad j_1 + j_2 + \cdots + j_t < d. \end{array} \quad (9)$$

It may happen that the scheme on the right-hand side of Eq. (8) is strictly larger than the one on the left-hand side, which results in the LP problem of larger size and brings us to the next subsection.

3.3. On the coherent closure of a Cartesian product of graphs. The question when the inequality in Eq. (8) is strict is a subtle one. This happens precisely when the direct product on the right-hand side of the formula admits a non-trivial fusion containing the edge set of the graph in question as a relation. In the general case, not restricted to bilinear forms graphs, the situation is studied in [11] using the theory of coherent configurations. In this section, we give a partial result in this direction using spectral graph theory.

First, we recall a simple example which shows that the inequality in Eq. (8) can be strict. Indeed, the Hamming association scheme on \mathbb{F}_q^n is a non-trivial fusion of the direct product of the (trivial) schemes of complete graphs. One can easily see this, as the former scheme has rank $n + 1$, while the latter one has rank 2^n . To explain this phenomenon, we need to recall the following definition.

Let $\mathcal{X} = (\Omega, \mathbf{R})$, $\mathbf{R} = \{R_i \mid i \in \llbracket D \rrbracket\}$, and $\mathcal{Y} = (\Omega, \mathbf{T})$, $\mathbf{T} = \{T_i \mid i \in \llbracket D \rrbracket\}$, be two association schemes. Any permutation $\varphi \in \text{Sym}(\llbracket D \rrbracket)$, mapping $i \mapsto i^\varphi$, naturally

induces a bijection from \mathbf{R} to \mathbf{T} , which sends R_i to $T_{i\varphi}$. With a slight abuse of notation, we also denote the latter bijection by φ . Now, if $\varphi \in \text{Sym}(\llbracket D \rrbracket)$ preserves the numbers from the condition (iv) of Definition 2.12 (namely, the intersection numbers $p_{i,j}^k$ of \mathcal{X} and $p_{i\varphi,j\varphi}^{k\varphi}$ of \mathcal{Y} are equal for all $i, j, k \in \llbracket D \rrbracket$), then the induced bijection $\varphi: \mathbf{R} \rightarrow \mathbf{T}$ is called an **algebraic isomorphism** from \mathcal{X} to \mathcal{Y} . If $\mathcal{X} = \mathcal{Y}$, then φ is an **algebraic automorphism** of \mathcal{X} . Clearly, algebraic automorphisms form a group under composition.

Let Φ be a group of algebraic automorphisms of \mathcal{X} . Given $i \in \llbracket D \rrbracket$, put

$$R_i^\Phi = \cup_{\varphi \in \Phi} R_{i\varphi}.$$

One can see that the set $\mathbf{R}^\Phi = \{R_i^\Phi \mid i \in \llbracket D \rrbracket\}$ forms a partition of Ω^2 . Furthermore, it follows (see [12, Lemma 2.3.26], [20, Lemma 9.1.1]) that the pair $\mathcal{X}^\Phi = (\Omega, \mathbf{R}^\Phi)$ is an association scheme, called the **algebraic fusion** of \mathcal{X} with respect to Φ .

Consider now the t -fold direct product:

$$\mathcal{X}^t = \underbrace{\mathcal{X} \otimes \mathcal{X} \otimes \cdots \otimes \mathcal{X}}_t, \quad t \geq 2.$$

One can show (see [12, Section 2.3]) that any permutation on t symbols induces an algebraic automorphism of \mathcal{X}^t ; thus, there exists a non-trivial algebraic fusion of \mathcal{X}^t , called the **exponentiation** of \mathcal{X} by the symmetric group $\text{Sym}(t)$. Furthermore, when an association scheme \mathcal{X} is the coherent closure of a graph Γ , these algebraic automorphisms of \mathcal{X}^t leave invariant the adjacency relation of the Cartesian t -th power $\Gamma^{\square t} := \Gamma \square \cdots \square \Gamma$ of the graph Γ . Thus, the coherent closure of $\Gamma^{\square t}$ is strictly less than \mathcal{X}^t . For example, the Hamming scheme on \mathbb{F}_q^n is the exponentiation of the trivial scheme on q points by the symmetric group $\text{Sym}(n)$.

Therefore, when a Cartesian product of graphs contains isomorphic factors, its coherent closure is definitely smaller than the direct product of the coherent closures of the factors. In particular, this applies to a sum-rank metric graph in Eq. (8). Otherwise, we only have the following conjecture.

Problem 3.6. *Let Γ be a Cartesian product of t bilinear forms graphs Γ_i , $i \in [t]$. Suppose that the coherent closures $\text{WL}(\Gamma_i)$, $\text{WL}(\Gamma_j)$ are not algebraically isomorphic² for all $i \neq j \in [t]$. Is it true that then*

$$\text{WL}(\Gamma_1 \square \cdots \square \Gamma_t) = \text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2) \otimes \cdots \otimes \text{WL}(\Gamma_t)? \quad (10)$$

In other words, Eq. (10) states that the direct product on the right-hand side does not admit a non-trivial fusion containing the edge set of the graph in question as a relation. Note that a general necessary and sufficient existence condition of fusions is given by the Bannai-Muzychuk criterion [4, Lemma 2.48]: see also Remark 3.8. In Theorem 3.7 below, we prove a result which shows the validity of Eq. (10) under a stronger assumption than that in Problem 3.6.

Note that the above discussion applies to the general case when the coherent closure of a graph is not an association scheme. Moreover, instead of the t -fold direct product one can consider a direct product of algebraically isomorphic schemes [11]. For example, the Doob and Hamming association schemes on \mathbb{F}_4^{2s+t} are algebraically isomorphic (for all natural s, t); both are the coherent closures of the Cartesian products of graphs: a Doob graph $Sh^{\square s} \square K_4^{\square t}$ in the former case (here Sh stands

²Two bilinear forms schemes on $\mathbb{F}_q^{n_1 \times m_1}$ and $\mathbb{F}_q^{n_2 \times m_2}$ are (algebraically) isomorphic if and only if $\{n_1, m_1\} = \{n_2, m_2\}$.

for the Shrikhande graph), and a Hamming graph $K_4^{\square(2s+t)}$ in the latter case. Note also that, except for the obvious trivial cases, both association schemes are strictly less than the direct products of the schemes of factors in the corresponding Cartesian products.

Next, we prove a sufficient condition under which the coherent closure of a Cartesian product of two graphs equals the direct product of their coherent closures. The condition is formulated in terms of the eigenvalues of the adjacency matrices of the graphs. Recall that the **adjacency algebra** of a graph is a coherent algebra (see Section 2.5) generated by its adjacency matrix. In the proof of Theorem 3.7 below, the reader may assume that the coherent closures $\text{WL}(\Gamma_1 \square \Gamma_2)$, $\text{WL}(\Gamma_1)$, $\text{WL}(\Gamma_2)$ are association schemes, in which case the adjacency algebras of the graphs $\Gamma_1 \square \Gamma_2$, Γ_1 , Γ_2 are simply the Bose-Mesner algebras (see Section 2.5) of the corresponding schemes. However, the proof remains valid in the general case when these coherent closures are coherent configurations; the corresponding adjacency algebras are coherent algebras associated with coherent configurations [12, Section 2.3.1].

Theorem 3.7. *Let Γ_i , $i \in \{1, 2\}$, be a graph with precisely s_i pairwise distinct eigenvalues θ_{ij} , $j \in [s_i]$. Then*

$$\text{WL}(\Gamma_1 \square \Gamma_2) = \text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2)$$

if the set $S := \{\theta_{1j} + \theta_{2k} \mid j \in [s_1], k \in [s_2]\}$ is of cardinality $s_1 s_2$.

Proof. As in Corollary 3.5, one can see that $\text{WL}(\Gamma_1 \square \Gamma_2) \leq \text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2)$. Therefore, to show the result, we need to prove $\text{WL}(\Gamma_1 \square \Gamma_2) \geq \text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2)$. Note that the fact that $\text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2)$ is a fusion of $\text{WL}(\Gamma_1 \square \Gamma_2)$ implies that the adjacency algebra \mathcal{A} of $\Gamma_1 \square \Gamma_2$ contains the coherent (Bose-Mesner) algebra of $\text{WL}(\Gamma_1) \otimes \text{WL}(\Gamma_2)$. Combining this and [12, Theorem 3.2.25], we conclude that it is sufficient to prove that \mathcal{A} contains the tensor product of the adjacency algebras \mathcal{A}_1 and \mathcal{A}_2 of Γ_1 and Γ_2 , respectively.

To prove the latter statement, it in turn suffices to prove that

$$A_1 \otimes I_{n_2} \in \mathcal{A} \quad \text{or} \quad I_{n_1} \otimes A_2 \in \mathcal{A}, \quad (11)$$

where A_1, A_2 denote the adjacency matrices of the graphs Γ_1, Γ_2 on n_1, n_2 vertices, respectively. Indeed, it is sufficient to prove only one of the two inclusions in Eq. (11): the adjacency algebras $\mathcal{A}, \mathcal{A}_1, \mathcal{A}_2$ are generated (as coherent algebras) by the matrices $A = A_1 \otimes I_{n_2} + I_{n_1} \otimes A_2$ (see Eq. (1)), A_1, A_2 , respectively. Assume, for example, that \mathcal{A} contains $A_1 \otimes I_{n_2}$ and hence \mathcal{A} contains the subalgebra $\mathcal{A}_1 \otimes I_{n_2} = \{X \otimes I_{n_2} : X \in \mathcal{A}_1\}$. Since $A \in \mathcal{A}$, it follows that \mathcal{A} contains $A - A_1 \otimes I_{n_2} = I_{n_1} \otimes A_2$ and hence the subalgebra $I_{n_1} \otimes \mathcal{A}_2 = \{I_{n_1} \otimes X : X \in \mathcal{A}_2\}$. Thus, \mathcal{A} contains both $\mathcal{A}_1 \otimes I_{n_2}$ and $I_{n_1} \otimes \mathcal{A}_2$, and hence

$$\mathcal{A} \supseteq (\mathcal{A}_1 \otimes I_{n_2}) \cdot (I_{n_1} \otimes \mathcal{A}_2) = \mathcal{A}_1 \otimes \mathcal{A}_2,$$

as required.

Next, to prove Eq. (11), we first note that

$$A^k = \sum_{i=0}^k \binom{k}{i} A_1^i \otimes A_2^{k-i},$$

which follows from Eq. (1) and the standard properties of Kronecker product of matrices.

Let $f_i, i \in \{1, 2\}$, be the minimal polynomial of A_i and let I denote the ideal of the ring of polynomials $\mathbb{R}[x, y]$, generated by $f_1(x)$ and $f_2(y)$. Then the mappings $A_1 \otimes I_{n_2} \mapsto x$ and $I_{n_1} \otimes A_2 \mapsto y$, define an isomorphism, say ι , between \mathcal{A} and the subalgebra \mathcal{K} of $\mathbb{R}[x, y]/I$ generated by the polynomial $x + y$,

For $i \in \{1, 2\}$, since f_i is the minimal polynomial of A_i , its degree is s_i . It follows that the algebra $\mathbb{R}[x, y]/I$ is of dimension $N = s_1 s_2$ and has a basis consisting of $x^i y^j, i \in \llbracket s_1 - 1 \rrbracket, j \in \llbracket s_2 - 1 \rrbracket$. In particular, the dimension of \mathcal{K} is at most N .

Consider a linear mapping $\epsilon: \mathcal{K} \rightarrow \mathbb{R}^N$ defined for $p \in \mathcal{K}$ as follows:

$$\epsilon: p(x, y) \mapsto (p(\theta_{11}, \theta_{21}), p(\theta_{11}, \theta_{22}), \dots, p(\theta_{1s_1}, \theta_{2s_2})).$$

Let $K = \langle \epsilon(1), \epsilon(x + y), \epsilon((x + y)^2), \dots, \epsilon((x + y)^{N-1}) \rangle$ be a subspace of \mathbb{R}^N , the image of ϵ . Due to the Rank-Nullity theorem, the dimension of \mathcal{K} is lower-bounded by the dimension of the image of ϵ . Moreover, $\dim K$ equals the rank r of the following $N \times N$ matrix:

$$\begin{bmatrix} 1 & \dots & 1 \\ \theta_{11} + \theta_{21} & \dots & \theta_{1s_1} + \theta_{2s_2} \\ (\theta_{11} + \theta_{21})^2 & \dots & (\theta_{1s_1} + \theta_{2s_2})^2 \\ \vdots & \vdots & \vdots \end{bmatrix},$$

which is a Vandermonde matrix. Hence r equals the number of pairwise distinct numbers in the set S . Therefore, if $|S| = N$, then the dimension of \mathcal{K} is exactly N , and the elements $1, (x + y), \dots, (x + y)^{N-1}$ constitute another basis of \mathcal{K} . Thus, the elements of one basis $x^i y^j, i \in \llbracket s_1 - 1 \rrbracket, j \in \llbracket s_2 - 1 \rrbracket$, can be expressed through the elements of another basis $(x + y)^i, i \in \llbracket N - 1 \rrbracket$. In particular,

$$x = \sum_{i \in \llbracket N - 1 \rrbracket} \alpha_i (x + y)^i$$

for some $\alpha_i \in \mathbb{R}, i \in \llbracket N - 1 \rrbracket$. By applying ι^{-1} , we get

$$A_1 \otimes I_{n_2} = \sum_{i \in \llbracket N - 1 \rrbracket} \alpha_i A^i,$$

which implies $A_1 \otimes I_{n_2} \in \mathcal{A}$, as required by Eq. (11). \square

Remark 3.8. For a sum-rank metric graph $\Gamma = \Gamma(\mathbb{F}_q^{\mathbf{n} \times \mathbf{m}})$, which is the Cartesian product of t bilinear forms graphs $\Gamma_i = \text{Bil}_q(n_i, m_i), i \in [t]$, the set of distinct eigenvalues of Γ is contained in the following set:

$$S = \left\{ \sum_{i=1}^t \frac{1}{q-1} ((q^{n_i - j_i} - 1)(q^{m_i} - q^{j_i}) - q^{j_i} + 1) : 0 \leq j_i \leq n_i, i \in [t] \right\}.$$

Theorem 3.7 states that $\text{WL}(\Gamma)$ is equal to the direct product of $\text{WL}(\Gamma_i), i \in [t]$, if S has $(n_1 + 1) \cdots (n_t + 1)$ distinct elements. One can easily find instances when this condition fails. For example, in case $t = 2$, two eigenvalues corresponding to distinct pairs (j_1, j_2) and $(j'_1, j'_2), 0 \leq j_i, j'_i \leq n_i, i = 1, 2$, coincide if and only if $n_2 - n_1 + m_2 - m_1 = j_2 - j'_1 = j'_2 - j_1$.

Despite this, to support the affirmative answer to Problem 3.6, we checked the Bannai-Muzychuk criterion for the direct product on the right-hand side of Eq. (10) for all sum-rank metric graphs over \mathbb{F}_2 with $t = 2, \sum_{i \in [t]} m_i \leq 6$, and $t = 3, \sum_{i \in [t]} m_i \leq 11$, and found no counterexamples.

4. BOUNDS PERFORMANCE

Tables 1, 2, and 3 list the non-distance regular sum-rank metric graphs (restricted in the number of vertices $|\Omega|$ and the number of blocks t) for which we computed the Delsarte's LP bound on α_{d-1} by solving the LP (9), as well as the other previously known bounds from [9] and the Ratio-type bound introduced in [2]. Based on these computational experiments on relatively small instances, it is observed that the obtained bounds outperform all previously known bounds. While we omit the cases when the result of the Delsarte's LP bound coincides with one of the previously achieved bounds, there are no examples of sum-rank metric graphs under the specified restrictions on $|\Omega|$ and t for which the Delsarte's LP bound returns a result that is strictly larger than the result of one of the other upper bounds.

The column "RT $_{d-1}$ " gives the values of the Ratio-type bound from Theorem 2.7, which is calculated either explicitly in case $d \in \{3, 4\}$ using the results of Theorem 2.8 [1] and Theorem 2.9 [23], or using the LP (5) [19]. The columns "D $_d$ ", "iS $_d$ ", "iH $_d$ ", "iE $_d$ ", "S $_d$ ", "SP $_d$ ", "PSP $_d$ " give the values of the Delsarte's LP (5), Induced Singleton, Induced Hamming, Induced Elias, Singleton, Sphere-Packing, and Projective Sphere-Packing bounds, respectively (see Theorems 2.5 and 2.6).

Since the code cardinality is always an integer, the columns only contain integer values, which are sometimes obtained by taking the floor of the real value given by the bound.

Note that the sum-rank metric graphs considered in Tables 1, 2, and 3 are restricted not only in the size of the vertex set of the graph, but also in t , the number of blocks. Since calculating the Delsarte's LP bound requires taking a Kronecker product of t matrices, the computation speed of the bound changes significantly with the increase of t . This obstacle is avoided when using the other bounds presented, particularly the Ratio-type bound, which, unlike the Delsarte's LP bound, does not depend as much on the structure of the graph in its design. Hence the Ratio-type bound can be useful in case t is large, despite it being often outperformed by Delsarte's LP bound for smaller values of t . The Ratio-type bound can also be computed without the use of LP for $d = 3, 4$ from explicit formulæ [1, 2, 23], which also puts it at advantage in this case.

To conclude this section, we discuss the relationship between the Delsarte's LP bound and the Lovász theta number. For a graph Γ , we define Γ^k to be a **k -th power** of Γ if the vertex set of Γ^k is the same as the vertex set of Γ , and two vertices are adjacent in Γ^k if and only if they are at distance at most k in Γ . It is then clear that the k -independence number of Γ is exactly the independence number of Γ^k . It is well-known that the independence number is upper bounded by the Lovász theta number, denoted by ϑ [24]. Thus we use ϑ_k to denote the Lovász theta number of Γ^k , which is an upper bound on α_k . The number ϑ_k can be estimated using Semidefinite Programming (SDP) as follows [24]: Let $\Gamma = (\Omega, E)$ be a graph on n vertices, and let \mathcal{S}_+^n denote the set of all $n \times n$ symmetric positive semidefinite matrices. Then ϑ is the solution of:

$$\begin{array}{ll} \text{maximize} & \text{tr}(BJ) \\ \text{subject to} & \text{tr } B = 1, \\ & b_{ij} = 0, \quad (i, j) \in E, i, j \in [n], \\ & B \in \mathcal{S}_+^n, \end{array} \tag{12}$$

where J is an all-ones matrix of size $n \times n$. Note that $\text{tr}(BJ)$ is the sum of the entries in B .

Remark 4.1. In [38], it was shown that, for graphs derived from symmetric association schemes, the bound obtained through Delsarte’s LP method [14] coincides with the number ϑ' , which is calculated similarly to ϑ using SDP (12), with the additional constraint that B must be a non-negative matrix. In the same paper [38] it was proven that $\alpha \leq \vartheta' \leq \vartheta$, meaning that the Delsarte’s LP bound performs no worse than the Lovász theta bound. This is confirmed by computations on sum-rank metric graphs given in Table 4 in Appendix.

5. CONCLUDING REMARKS

In this work we show how to implement Delsarte’s LP method to obtain an upper bound on the maximum size of a code with a given minimum distance in sum-rank metric, which does not correspond to a distance-regular graph (unlike the Hamming metric or rank metric). We do so by considering the direct product of the association schemes of the Cartesian factors of the corresponding sum-rank metric graph. Our results mostly make use of the structure of the space itself and not of the properties of matrix ranks that define the distance between the elements. Thus, we expect that this approach for constructing association schemes could also be applied to other metrics that give rise to graphs with a similar structure. In particular, most results of Section 3.3 are provided in a general form and could be applied to any Cartesian product of graphs instead of a sum-rank metric graph. For instance, codes in the Lee-metric can be viewed as subsets of vertices in the standard path-length metric of the Cartesian products of cycles; note that in [3] the linear programming bound for Lee-codes was obtained by applying Delsarte’s LP method to the explicitly defined Lee association scheme [39].

The application of Delsarte’s LP bound in case the graph corresponding to the distance metric is distance-regular also gives rise to some open problems. Consider the rank-metric case, *i.e.* the case of sum-rank metric graphs with $t = 1$. The computational experiments on small graphs (up to 10^7 vertices with diameter at least 3) shown in Table 5 suggest that for such graphs, both Delsarte’s LP bound computed with the LP (9) and the Ratio-type bound of Theorem 2.7 computed with the LP (5) return the same result (this concerns the bounds on the size of the maximum code with minimum distance $d \in [3, n]$). Such output also coincides with the Singleton bound, which is known to be always met in the sum-rank metric case [15]. Thus, we suspect that the first two bounds might be equivalent in the case of rank-metric graphs.

Acknowledgements. Aida Abiad is supported by NWO (Dutch Research Council) through the grant VI.Vidi.213.085. The research of Alexander Gavrilyuk is supported by JSPS KAKENHI Grant Number 22K03403. Antonina P. Khramova is supported by NWO via the grant OCENW.KLEIN.475. This work was initiated during the RICOTA conference, Croatia, in July 2023; the authors would like to thank the organizers of the event.

REFERENCES

- [1] A. Abiad, G. Coutinho, and M. A. Fiol. On the k -independence number of graphs. *Discrete Mathematics*, 342(10):2875–2885, 2019.

- [2] A. Abiad, A. P. Khamova, and A. Ravagnani. Eigenvalue bounds for sum-rank-metric codes. *IEEE Transactions on Information Theory*, 2024.
- [3] J. Astola. The Lee-scheme and bounds for Lee-codes. *Cybernetics and Systems Analysis*, 13:331–343, 1982.
- [4] E. Bannai, E. Bannai, T. Ito, and R. Tanaka. *Algebraic Combinatorics*. De Gruyter, Berlin, Boston, 2021.
- [5] H. Bartz and S. Puchinger. Decoding of interleaved linearized Reed-Solomon codes with applications to network coding. In *2021 IEEE International Symposium on Information Theory*, pages 160–165, 2021.
- [6] P. A. Bernard, L. Vinet, M. Zaimi, and X. Zhang. m -distance-regular graphs and their relation to multivariate P -polynomial association schemes. arXiv:2309.1601, 2023.
- [7] M. Borello and F. Zullo. Geometric dual and sum-rank minimal codes. *Journal of Combinatorial Designs*, 32(5):238–273, 2024.
- [8] A. E. Brouwer, A. M. Cohen, and A. Neumaier. *Distance-Regular Graphs*. Springer, Berlin, Heidelberg, 1989.
- [9] E. Byrne, H. Gluesing-Luerssen, and A. Ravagnani. Fundamental properties of sum-rank-metric codes. *IEEE Transactions on Information Theory*, 67(10):6456–6475, 2021.
- [10] E. Byrne, H. Gluesing-Luerssen, and A. Ravagnani. Anticodes in the sum-rank metric. *Linear Algebra and its Applications*, 643:80–98, 2022.
- [11] J. Cai, J. Guo, A. L. Gavriluk, and I. Ponomarenko. Cartesian product of graphs and their coherent configurations. Preprint, 2024.
- [12] G. Chen and I. Ponomarenko. *Coherent configurations*. Central China Normal University Press, 2019. Updated version available at <http://www.pdmi.ras.ru/~inp/ccNOTES.pdf>.
- [13] H. Chen. New explicit good linear sum-rank-metric codes. *IEEE Transactions on Information Theory*, 69(10):6303–6313, 2023.
- [14] P. Delsarte. An algebraic approach to the association schemes of coding theory. *Philips Research Reports: Supplements*, 10:vi+–97, 1973.
- [15] P. Delsarte. Bilinear forms over a finite field, with applications to coding theory. *Journal of Combinatorial Theory, Series A*, 25(3):226–241, 1978.
- [16] P. Delsarte and J.-M. Goethals. Alternating bilinear forms over $\text{GF}(q)$. *Journal of Combinatorial Theory, Series A*, 19(1):26–50, 1975.
- [17] P. J. Dukes, F. Ihringer, and N. Lindzey. On the algebraic combinatorics of injections and its applications to injection codes. *IEEE Transactions on Information Theory*, 66(11):6898–6907, 2020.
- [18] P. A. Ferguson and A. Turull. Algebraic decompositions of commutative association schemes. *Journal of Algebra*, 96(1):211–229, 1985.
- [19] M. A. Fiol. A new class of polynomials from the spectrum of a graph, and its application to bound the k -independence number. *Linear Algebra and its Applications*, 605:1–20, 2020.
- [20] C. D. Godsil. Association Schemes. <https://www.math.uwaterloo.ca/~cgodsil/pdfs/assoc2.pdf>.
- [21] W. H. Haemers. Interlacing eigenvalues and graphs. *Linear Algebra and its Applications*, 226–228:593–616, 1995.
- [22] R. Hammack, W. Imrich, and S. Klavžar. *Handbook of product graphs*. CRC Press, Boca Raton, FL, 2011.
- [23] L. Kavi and M. Newman. The optimal bound on the 3-independence number obtainable from a polynomial-type method. *Discrete Mathematics*, 346:113471, 2023.
- [24] L. Lovász. On the Shannon capacity of a graph. *IEEE Transactions on Information Theory*, 25(1):1–7, 1979.
- [25] W. Martin. Designs in product association schemes. *Designs, Codes and Cryptography*, 16:271–289, 1999.
- [26] U. Martínez-Peñas. Skew and linearized Reed-Solomon codes and maximum sum rank distance codes over any division ring. *Journal of Algebra*, 504:587–612, 2018.
- [27] U. Martínez-Peñas. Theory of supports for linear codes endowed with the sum-rank metric. *Designs, Codes and Cryptography*, 87:2295–2320, 2019.
- [28] U. Martínez-Peñas. Hamming and simplex codes for the sum-rank metric. *Designs, Codes and Cryptography*, 88:1521–1539, 2020.

- [29] U. Martínez-Peñas and F. R. Kschischang. Reliable and secure multishot network coding using linearized Reed-Solomon codes. *IEEE Transactions on Information Theory*, 65:4785–4803, 2019.
- [30] U. Martínez-Peñas and F. R. Kschischang. Universal and dynamic locally repairable codes with maximal recoverability via sum-rank codes. *IEEE Transactions on Information Theory*, 65:7790–7805, 2019.
- [31] U. Martínez-Peñas, M. Shehadeh, and F. R. Kschischang. Codes in the sum-rank metric: Fundamentals and applications. *Foundations and Trends in Communications and Information Theory*, 19(5):814–1031, 2022.
- [32] U. Martínez-Peñas. Sum-rank BCH codes and cyclic-skew-cyclic codes. *IEEE Transactions on Information Theory*, 67(8):5149–5167, 2021.
- [33] A. Neri, P. Santonastaso, and F. Zullo. The geometry of one-weight codes in the sum-rank metric. *Journal of Combinatorial Theory, Series A*, 194, 2023.
- [34] R. Nóbrega and B. Uchôa-Filho. Multishot codes for network coding using rank-metric codes. In *2010 Third IEEE International Workshop on Wireless Network Coding*, pages 1–6, 2010.
- [35] C. Ott, H. Liu, and A. Wachter-Zeh. Covering properties of sum-rank metric codes. In *2022 58th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 1–7, 2022.
- [36] A. Roy. Bounds for codes and designs in complex subspaces. *Journal of Algebraic Combinatorics*, 31:1–32, 2010.
- [37] P. Santonastaso and F. Zullo. On subspace designs. *EMS Surveys in Mathematical Sciences*, 11(1):1–62, 2023.
- [38] A. Schrijver. A comparison of the Delsarte and Lovász bounds. *IEEE Transactions on Information Theory*, 25(4):425–429, 1979.
- [39] P. Sole. The Lee association scheme. In *International Colloquium on Coding Theory and Applications*, pages 45–55. Springer, 1986.
- [40] B. Weisfeiler and A. Leman. The reduction of a graph to canonical form and the algebra which appears therein. *Nauchno-Tekhnicheskaya Informatsia*, 2(9):12–16, 1968. (Russian), English translation by G. Ryabov available at https://www.iti.zcu.cz/wl2018/pdf/wl_paper_translation.pdf.

APPENDIX

t	q	\mathbf{n}	\mathbf{m}	d	$ \Omega $	RT_{d-1}	D_d	iS_d	iH_d	iE_d	S_d	SP_d	PSP_d
2	2	[2, 2]	[2, 2]	3	256	11 [1]	10	16	19	34	16	13	13
3	2	[2, 2, 1]	[2, 2, 1]	3	512	25 [1]	20	64	64	151	32	25	25
3	2	[2, 2, 1]	[2, 2, 1]	4	512	10 [23]	6	16	64	27	8	25	18
3	2	[2, 2, 1]	[2, 2, 2]	3	1024	38 [1]	34	64	64	151	64	46	46
3	2	[2, 2, 1]	[2, 2, 2]	4	1024	15 [23]	8	16	64	27	16	46	36
3	2	[2, 2, 2]	[2, 2, 2]	3	4096	146 [1]	133	256	215	529	256	146	146
3	2	[2, 2, 2]	[2, 2, 2]	4	4096	45 [23]	32	64	215	119	64	146	102
3	2	[2, 2, 2]	[2, 2, 2]	5	4096	14 [19]	6	16	26	19	16	14	13
4	2	[2, 1, 1, 1]	[2, 2, 2, 1]	3	512	28 [1]	24	64	64	151	32	30	30
4	2	[2, 1, 1, 1]	[2, 2, 2, 1]	4	512	11 [23]	6	16	64	27	8	30	32
4	2	[2, 1, 1, 1]	[2, 2, 2, 2]	3	1024	44 [1]	42	64	64	151	64	53	53
4	2	[2, 1, 1, 1]	[2, 2, 2, 2]	4	1024	18 [23]	10	16	64	27	16	53	64
4	2	[2, 2, 1, 1]	[2, 2, 1, 1]	3	1024	46 [1]	40	256	215	529	64	48	48
4	2	[2, 2, 1, 1]	[2, 2, 1, 1]	4	1024	19 [23]	12	64	215	119	16	48	36
4	2	[2, 2, 1, 1]	[2, 2, 2, 1]	3	2048	85 [1]	68	256	215	529	128	89	89
4	2	[2, 2, 1, 1]	[2, 2, 2, 1]	4	2048	32 [23]	16	64	215	119	32	89	73
4	2	[2, 2, 1, 1]	[2, 2, 2, 1]	5	2048	12 [19]	5	16	26	19	8	10	9
4	2	[2, 2, 1, 1]	[2, 2, 2, 2]	3	4096	146 [1]	134	256	215	529	256	163	163
4	2	[2, 2, 1, 1]	[2, 2, 2, 2]	4	4096	48 [23]	32	64	215	119	64	163	146
4	2	[2, 2, 1, 1]	[2, 2, 2, 2]	5	4096	17 [19]	7	16	26	19	16	17	16
5	2	[2, 1, 1, 1, 1]	[2, 1, 1, 1, 1]	5	256	5 [19]	2	16	26	19	4	4	3
5	2	[2, 1, 1, 1, 1]	[3, 1, 1, 1, 1]	5	1024	8 [19]	2	64	336	240	4	6	3
5	2	[2, 1, 1, 1, 1]	[4, 1, 1, 1, 1]	5	4096	11 [19]	2	256	4096	1970	4	9	3
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 1, 1]	3	1024	56 [1]	49	256	215	529	64	56	56
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 1, 1]	4	1024	22 [23]	13	64	215	119	16	56	64
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 2, 1]	3	2048	102 [1]	85	256	215	529	128	102	102
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 2, 1]	4	2048	36 [23]	20	64	215	119	32	102	128
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 2, 1]	5	2048	14 [19]	6	16	26	19	8	13	11
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 2, 2]	3	4096	153 [1]	146	256	215	529	256	186	186
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 2, 2]	4	4096	53 [23]	40	64	215	119	64	186	256
5	2	[2, 1, 1, 1, 1]	[2, 2, 2, 2, 2]	5	4096	20 [19]	8	16	26	19	16	21	19
5	2	[2, 2, 1, 1, 1]	[2, 2, 1, 1, 1]	3	2048	93 [1]	80	1024	744	1876	128	93	93
5	2	[2, 2, 1, 1, 1]	[2, 2, 1, 1, 1]	4	2048	38 [23]	24	256	744	407	32	93	73
5	2	[2, 2, 1, 1, 1]	[2, 2, 1, 1, 1]	5	2048	13 [19]	7	64	77	99	8	11	9
5	2	[2, 2, 1, 1, 1]	[2, 2, 2, 1, 1]	3	4096	170 [1]	137	1024	744	1876	256	170	170
5	2	[2, 2, 1, 1, 1]	[2, 2, 2, 1, 1]	4	4096	64 [23]	32	256	744	407	64	170	146
5	2	[2, 2, 1, 1, 1]	[2, 2, 2, 1, 1]	5	4096	22 [19]	11	64	77	99	16	19	17
6	2	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	4	512	16 [23]	12	256	512	407	16	34	32
6	2	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	5	512	8 [19]	4	64	77	99	8	6	5
6	2	[2, 1, 1, 1, 1, 1]	[3, 1, 1, 1, 1, 1]	5	2048	15 [19]	4	512	1943	1707	8	11	5
6	2	[2, 1, 1, 1, 1, 1]	[3, 1, 1, 1, 1, 1]	6	2048	11 [19]	2	64	1943	211	4	11	3
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	5	1024	11 [19]	6	64	77	99	8	9	8
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	6	1024	7 [19]	2	16	77	14	4	9	3
6	2	[2, 1, 1, 1, 1, 1]	[3, 2, 1, 1, 1, 1]	6	4096	16 [19]	2	64	1943	211	4	17	3
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 2, 1, 1, 1]	3	2048	102 [1]	99	1024	744	1876	128	107	107
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 2, 1, 1, 1]	4	2048	42 [23]	26	256	744	407	32	107	128
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 2, 2, 1, 1]	3	4096	186 [1]	170	1024	744	1876	256	195	195
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 2, 2, 1, 1]	4	4096	70 [23]	40	256	744	407	64	195	256
6	2	[2, 1, 1, 1, 1, 1]	[2, 2, 2, 2, 1, 1]	5	4096	26 [19]	12	64	77	99	16	23	21
6	2	[2, 2, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	3	4096	170 [1]	160	4096	2621	4096	256	178	178
6	2	[2, 2, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	4	4096	72 [23]	48	1024	2621	1419	64	178	146
6	2	[2, 2, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	5	4096	24 [19]	14	256	236	366	16	21	18
6	2	[2, 2, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	6	4096	12 [19]	6	64	236	79	8	21	16
6	2	[2, 2, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	7	4096	6 [19]	2	16	36	11	4	5	3

TABLE 1. Some of the non-distance-regular sum-rank metric graphs with ≤ 5000 vertices for which the LP (9) outperforms previously known bounds for $A_q(\mathbf{n}, \mathbf{m}, d)$ with $t \leq 6$.

t	q	\mathbf{n}	\mathbf{m}	d	$ \Omega $	RT_{d-1}	D_d	iS_d	iH_d	iE_d	S_d	SP_d	PSP_d
3	3	[2, 2, 2]	[2, 2, 2]	3	531441	4652 [1]	4278	6561	10845	273399	6561	5478	5478
3	3	[2, 2, 2]	[2, 2, 2]	5	531441	198 [19]	33	81	526	343	81	160	100
6	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	5	19683	86 [19]	18	729	3413	2772	27	43	22
6	3	[2, 1, 1, 1, 1, 1]	[3, 1, 1, 1, 1, 1]	5	177147	170 [19]	18	19683	177147	177147	27	97	22
6	3	[2, 1, 1, 1, 1, 1]	[3, 1, 1, 1, 1, 1]	6	177147	148 [19]	6	729	177147	10104	9	97	9
7	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	5	59049	223 [19]	48	6561	23180	26541	81	109	56
7	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	6	59049	142 [19]	18	729	23180	3256	27	109	22
7	3	[2, 1, 1, 1, 1, 1]	[3, 1, 1, 1, 1, 1]	5	531441	482 [19]	48	531441	531441	531441	81	259	56
7	3	[2, 1, 1, 1, 1, 1]	[3, 1, 1, 1, 1, 1]	6	531441	406 [19]	18	19683	531441	531441	27	259	22
7	3	[2, 1, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	6	177147	222 [19]	18	729	23180	3256	27	222	22
8	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	6	177147	346 [19]	48	6561	162987	45526	81	282	56
8	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	7	177147	158 [19]	12	729	8536	4625	27	41	22
8	3	[2, 1, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	6	531441	562 [19]	48	6561	162987	45526	81	592	56
8	3	[2, 1, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	7	531441	233 [19]	18	729	8536	4625	27	70	22
9	3	[2, 1, 1, 1, 1, 1]	[2, 2, 1, 1, 1, 1]	5	531441	1471 [19]	340	531441	531441	531441	729	737	385
9	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	6	531441	863 [19]	139	59049	531441	320840	243	737	145
9	3	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	7	531441	362 [19]	37	6561	54141	32938	81	96	56

TABLE 2. Some of the non-distance-regular sum-rank metric graphs with $\leq 10^6$ vertices for which the LP (9) outperforms previously known bounds for $A_q(\mathbf{n}, \mathbf{m}, d)$ with $q \neq 2$.

t	q	\mathbf{n}	\mathbf{m}	d	$ \Omega $	RT_{d-1}	D_d	iS_d	iH_d	iE_d	S_d	SP_d	PSP_d
7	4	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	5	1048576	1740 [19]	179	65536	668893	549450	256	596	215
8	4	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	5	4194304	6156 [19]	614	1048576	4194304	4194304	1024	2055	744
8	4	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	6	4194304	4274 [19]	179	65536	4194304	1324989	256	2055	215
8	4	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	7	4194304	1537 [19]	40	4096	235553	80457	64	200	64

TABLE 3. Some of the non-distance-regular sum-rank metric graphs with $\leq 10^7$ vertices for which the LP (9) outperforms previously known bounds for $A_q(\mathbf{n}, \mathbf{m}, d)$ with $q > 3$.

t	q	\mathbf{n}	\mathbf{m}	d	$ \Omega $	α_{d-1}	ϑ_{d-1}	RT_{d-1}	Delsarte's LP
2	2	[2, 1]	[2, 1]	3	32	2	2	2	2
2	2	[2, 1]	[2, 2]	3	64	4	4	4	4
2	2	[2, 1]	[3, 3]	3	512	8	8	8	8
2	2	[2, 2]	[2, 2]	3	256	9	10	11	10
2	3	[2, 1]	[2, 2]	3	729	9	9	9	9
3	2	[1, 1, 1]	[2, 1, 1]	3	16	2	2	2	2
3	2	[2, 1, 1]	[2, 2, 2]	3	256	16	16	16	16
3	2	[2, 2, 1]	[2, 2, 1]	3	512	18	20	25	20
4	2	[1, 1, 1, 1]	[2, 1, 1, 1]	3	32	4	4	4	4
4	2	[1, 1, 1, 1]	[2, 1, 1, 1]	4	32	2	2	2	2
4	2	[2, 1, 1, 1]	[2, 1, 1, 1]	4	128	4	4	4	4
4	2	[2, 1, 1, 1]	[2, 2, 1, 1]	3	256	16	16	16	16
4	2	[2, 1, 1, 1]	[2, 2, 2, 1]	3	512	time	24	28	24
5	2	[1, 1, 1, 1, 1]	[2, 1, 1, 1, 1]	3	64	8	8	8	8
5	2	[2, 1, 1, 1, 1]	[2, 2, 1, 1, 1]	3	512	32	32	32	32
6	2	[1, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	3	128	9	12	12	12
6	2	[2, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1]	3	512	time	32	32	32
7	2	[1, 1, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1, 1]	3	256	time	25	25	21
8	2	[1, 1, 1, 1, 1, 1, 1, 1]	[2, 1, 1, 1, 1, 1, 1, 1]	3	512	time	42	42	42

TABLE 4. Some sum-rank metric graphs on up to 1000 vertices with the exact $(d - 1)$ -independence number α_{d-1} , ϑ_{d-1} , Ratio-type bound, and Delsarte's LP bound computed. We write 'time' whenever the computation time exceeds 2 hours on a standard laptop.

Γ	d	$ \Omega $	Ratio-type	Delsarte's LP	Singleton
$\text{Bil}_2(3, 3)$	3	512	8	8	8
$\text{Bil}_2(3, 4)$	3	4096	16	16	16
$\text{Bil}_2(3, 5)$	3	32768	32	32	32
$\text{Bil}_2(3, 6)$	3	262144	64	64	64
$\text{Bil}_2(3, 7)$	3	2097152	128	128	128
$\text{Bil}_2(4, 4)$	3	65536	256	256	256
$\text{Bil}_2(4, 4)$	4	65536	16	16	16
$\text{Bil}_2(4, 5)$	3	1048576	1024	1024	1024
$\text{Bil}_2(4, 5)$	4	1048576	32	32	32
$\text{Bil}_3(3, 3)$	3	19683	27	27	27
$\text{Bil}_3(3, 4)$	3	531441	81	81	81
$\text{Bil}_4(3, 3)$	3	262144	64	64	64
$\text{Bil}_5(3, 3)$	3	1953125	125	125	125

TABLE 5. The list of rank-metric graphs, *i.e.* sum-rank metric graphs with $t = 1$, and the respective values of Delsarte's LP, Ratio-type, and Singleton bounds.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EINDHOVEN UNIVERSITY OF TECHNOLOGY, THE NETHERLANDS

Email address: a.abiad.monge@tue.nl

INTERDISCIPLINARY FACULTY OF SCIENCE AND ENGINEERING, SHIMANE UNIVERSITY, MATSUE, JAPAN

Email address: gavriilyuk@riko.shimane-u.ac.jp

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EINDHOVEN UNIVERSITY OF TECHNOLOGY, THE NETHERLANDS

Email address: a.khramova@tue.nl

HAINAN UNIVERSITY, HAIKOU, CHINA

Email address: iliapon@gmail.com