

# Some Representations of the Deformed Euler Numbers $e_{s,t,u}$ and Proof for Irrationality

Ronald Orozco López

June 25, 2024

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Preliminaries</b>	<b>3</b>
2.1	Generalized Fibonacci numbers for $s^2 + 4t \geq 0$	3
2.2	Deformed $(s, t)$ -exponential function	6
2.3	Deformed binomial formula	7
<b>3</b>	<b>Deformed Euler <math>(s, t)</math>-numbers</b>	<b>8</b>
3.1	Definition	8
3.2	Estimating the numbers $e_{s,t}$	10
<b>4</b>	<b>Some representations of the <math>e_{s,t,u}</math>, <math>s \neq 0</math>, <math>t \neq 0</math></b>	<b>10</b>
4.1	As series compression	11
4.2	As Engel series expansion	12
4.3	As a series of partial Theta functions	15
4.4	As a continued fraction	16
4.5	As the limit of a sequence	17
<b>5</b>	<b><math>e_{s,t,u^{-1}}</math> and <math>e_{s,t,u^{-1}}^{-1}</math>, <math>u \in \mathbb{Q}</math>, <math> u  &gt; 1</math>, are irrationals</b>	<b>19</b>
<b>6</b>	<b><math>\varphi</math>-Euler and <math>\varphi'</math>-Euler numbers</b>	<b>21</b>

## Abstract

In this paper, representations of the Euler  $(s, t)$ -numbers  $e_{s,t,u}$  are given: as series compression, as Engel series expansion, as a series of partial Theta functions, as a continued fraction and as the limit of a sequence. Furthermore, it is proved that  $e_{as,a^2t,u^{-1}}$  and  $e_{as,a^2t,u^{-1}}^{-1}$  are irrational numbers when  $a, u \in \mathbb{Q}$  and  $|au| > 1$ . This is the first step in a program to study the irrationality of  $(s, t)$ -analog of known numbers.

2020 *Mathematics Subject Classification*: Primary 11J72. Secondary 11A67; 11J70; 11B39; 33E20.

*Keywords*: Generalized Fibonacci numbers, Euler  $(s, t)$ -numbers, Engel series, partial Theta functions, deformed  $(s, t)$ -binomial formula, irrationality.

# 1 Introduction

The constant  $e$  can be represented in many ways: as an infinite series

$$e = \sum_{n=0}^{\infty} \frac{1}{n!},$$

as a continued fraction [4]

$$e = \frac{1}{1 - \frac{1}{2 - \frac{1}{3 - \frac{2}{4 - \ddots}}}},$$

as an Engel series expansion [3]

$$e = 1 + \frac{1}{1} \left( 1 + \frac{1}{2} \left( 1 + \frac{1}{3} (1 + \dots) \right) \right),$$

as a limit of a sequence

$$e = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right)^n,$$

among others. In this paper, we aim to give several representations, analogous to the above, of the deformed Euler numbers  $\exp_{s,t}(1, u)$  defined on generalized Fibonacci numbers. The deformed exponential function  $\exp_{s,t}(x, u)$  on generalized Fibonacci numbers was defined by Orozco [11], which is introduced as a solution of the equation

$$\mathbf{D}_{s,t}f(x) = f(ux), \quad f(0) = 1. \quad (1)$$

A solution of Eq.(1) is of the form

$$\exp_{s,t}(x, u) = \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{x^n}{\{n\}_{s,t}!} \quad (2)$$

where  $\{n\}_{s,t}! = \{1\}_{s,t} \{2\}_{s,t} \cdots \{n\}_{s,t}$  is the  $(s, t)$ -analogue of the  $n!$ -factorial.

It is well known that  $e$  is irrational and many proofs of this fact have been given [4, 6, 7, 8, 13, 14]. Another aim of this paper is to prove the irrationality of the numbers  $\exp_{s,t}(1, u)$ . To achieve all the above results, the parameters  $s, t$  will be required to satisfy the condition  $s^2 + 4t \geq 0$ .

The paper is divided as follows. In Section 2 we deal with generalized Fibonacci numbers  $\{n\}_{s,t}$ , where  $s, t \in \mathbb{R} - \{0\}$  and  $s^2 + 4t \geq 0$ . The reason for stating the above condition is because under it the sequence of numbers  $|\{n\}_{s,t}|$  is increasing, which is the natural property of the sequence  $(0, 1, 2, 3, 4, 5, \dots)$ . Next, the  $(s, t)$ -deformed exponential functions and the  $(s, t)$ -deformed binomial formula are introduced. In Section 3 we introduce the deformed Euler  $(s, t)$ -numbers, and an estimate of them is given. Section 4 deals with representations of the deformed Euler  $(s, t)$ -numbers: as series compression, as Engel series expansion, as a series of partial Theta functions, as a continued fraction, and as the limit of a sequence. In Section 5 it is proven that  $e_{as, a^2t, u^{-1}}$  and  $e_{as, a^2t, u^{-1}}^{-1}$  are irrational numbers when  $a, u \in \mathbb{Q}$  and  $|au| > 1$ . Finally, in the last section, we deal with the representation as an infinite product for the numbers  $\varphi$ -Euler and  $\varphi'$ -Euler.

## 2 Preliminaries

### 2.1 Generalized Fibonacci numbers for $s^2 + 4t \geq 0$

The generalized Fibonacci numbers on the parameters  $s, t$  are defined by

$$\{n+2\}_{s,t} = s\{n+1\}_{s,t} + t\{n\}_{s,t} \quad (3)$$

with initial values  $\{0\}_{s,t} = 0$  and  $\{1\}_{s,t} = 1$ , where  $s \neq 0$  and  $t \neq 0$ .

The  $(s, t)$ -Fibonacci constant is the ratio toward which adjacent  $(s, t)$ -Fibonacci numbers tend. This is the only positive root of  $x^2 - sx - t = 0$ . We will let  $\varphi_{s,t}$  denote this constant, where

$$\varphi_{s,t} = \frac{s + \sqrt{s^2 + 4t}}{2}$$

and its conjugate is

$$\varphi'_{s,t} = s - \varphi_{s,t} = -\frac{t}{\varphi_{s,t}} = \frac{s - \sqrt{s^2 + 4t}}{2}.$$

Three cases arise from the discriminant  $\Delta = s^2 + 4t$ :  $\Delta > 0$ , which produces an increasing sequence of the absolute values of its terms.  $\Delta = 0$ , which yields sequences of the form  $n(\pm i\sqrt{t})^{n-1}$ . Finally, when  $\Delta < 0$ , then  $s \neq 0$  and  $t < 0$  and we obtain sequences of the form

$$\{n\}_{\sqrt{a}, -b/4} = \frac{2}{\sqrt{b-a}} \left(\frac{b}{4}\right)^n \sin(\theta n)$$

where  $\theta = \arctan\left(\sqrt{\frac{b}{a} - 1}\right)$  and  $a = s^2$ ,  $b = -4t$ .

Below are some important specializations of Fibonacci numbers when  $\Delta > 0$ .

1. When  $s = 1, t = 1$ , then  $\{n\}_{1,1} = F_n$ , the Fibonacci numbers.
2. When  $s = 2, t = 1$ , then  $\{n\}_{2,1} = P_n$ , where  $P_n$  are the Pell numbers

$$P_n = (0, 1, 2, 5, 12, 29, 70, 169, 408, \dots).$$

3. When  $s = 1, t = 2$ , then  $\{n\}_{1,2} = J_n$ , where  $J_n$  are the Jacobsthal numbers

$$J_n = (0, 1, 1, 3, 5, 11, 21, 43, 85, 171, \dots).$$

4. When  $s = 3, t = -2$ , then  $\{n\}_{3,-2} = M_n$ , where  $M_n = 2^n - 1$  are the Mersenne numbers

$$M_n = (0, 1, 3, 7, 15, 31, 63, 127, 255, \dots).$$

Some specializations of the constants  $\varphi_{s,t}$  and  $\varphi'_{s,t}$  are:

1. When  $s = 1$  and  $t = 1$ , then  $\varphi_{1,1} = \varphi = \frac{1+\sqrt{5}}{2}$  and  $\varphi'_{1,1} = \varphi' = \frac{1-\sqrt{5}}{2}$ .
2. When  $s = 2$  and  $t = 1$ , then  $\varphi_{2,1} = 1 + \sqrt{2}$  and  $\varphi'_{2,1} = 1 - \sqrt{2}$ .
3. When  $s = 1$  and  $t = 2$ , then  $\varphi_{1,2} = 2$  and  $\varphi'_{1,2} = -1$ .
4. When  $s = 3$  and  $t = -2$ , then  $\varphi_{3,-2} = 2$  and  $\varphi'_{3,-2} = 1$ .

The Binet's  $(s, t)$ -identity is

$$\{n\}_{s,t} = \frac{\varphi_{s,t}^n - \varphi'_{s,t}{}^n}{\varphi_{s,t} - \varphi'_{s,t}} \quad (4)$$

As  $\varphi_{us,u^2t} = u\varphi_{s,t}$  and  $\varphi'_{us,u^2t} = u\varphi'_{s,t}$ , then follows that  $\{n\}_{us,u^2t} = u^{n-1}\{n\}_{s,t}$ . Then for a non-zero complex number  $u$  we will say that  $\{n\}_{us,u^2t}$  is an  $u$ -deformation of  $\{n\}_{s,t}$ . We define the *alternating generalized Fibonacci numbers* as the  $-1$ -deformation of sequence  $\{n\}_{s,t}$ , thus,

$$\{n\}_{-s,t} = (-1)^{n-1}\{n\}_{s,t}. \quad (5)$$

For example, the alternating Fibonacci numbers are:  $0, 1, -1, 2, -3, 5, -8, \dots$  and the alternating Pell numbers are:  $0, 1, -2, 5, -12, 29, -70, 169, -408, \dots$ . Another important  $u$ -deformation is a  $\varphi_{s,t}$ -deformation, so  $\{n\}_{s,t} = \varphi_{s,t}^{n-1}[n]_q$  with  $q = \varphi'_{s,t}/\varphi_{s,t}$ .

On the other hand, when  $s^2 + 4t = 0$  we obtain the degenerate case of the  $(s, t)$ -Fibonacci numbers. When  $\varphi_{s,t} \rightarrow \varphi'_{s,t}$ , we obtain

$$\lim_{\varphi_{s,t} \rightarrow \varphi'_{s,t}} \frac{\varphi_{s,t}^n - \varphi'_{s,t}{}^n}{\varphi_{s,t} - \varphi'_{s,t}} = n\varphi_{s,t}^{(n-1)}.$$

Likewise, when  $\varphi'_{s,t} \rightarrow \varphi_{s,t}$ , then  $\{n\}_{s,t} \rightarrow n\varphi_{s,t}^{n-1}$ . Therefore, this implies that  $s \rightarrow \pm 2i\sqrt{t}$ , and that  $\varphi_{s,t} = \varphi'_{s,t} = \pm i\sqrt{t}$ . In this way we obtain the  $(\pm 2i\sqrt{t}, t)$ -Fibonacci function

$$\{n\}_{\pm 2i\sqrt{t}, t} = n(\pm i\sqrt{t})^{n-1} \quad (6)$$

for all  $t \in \mathbb{R}$ ,  $t \neq 0$ . When  $t = -1$ , then  $\{n\}_{\pm 2i\sqrt{t}, t} = \{n\}_{\mp 2, -1} = n(\mp 1)^{n-1}$ . On the other hand, in the  $q$ -calculus the degenerate case is obtained when  $q \mapsto 1$ . In this situation, the  $q$ -numbers  $[n]_q$  tend to the integers  $n$ . Then  $\frac{\varphi'_{s,t}}{\varphi_{s,t}} \mapsto 1$  implies that  $\varphi_{s,t} \mapsto \sqrt{-t}$  and  $\varphi'_{s,t} \mapsto \sqrt{-t}$ . Therefore, if  $t = -1$ , then

$$\lim_{\varphi_{s,-1} \mapsto 1} \frac{\varphi_{s,-1}^n - \varphi'_{s,-1}{}^n}{\varphi_{s,-1} - \varphi'_{s,-1}} = n.$$

The sequence  $\{0, 1, 2, 3, 4, 5, 6, \dots\}$ , the basis of all classical calculus, is strictly increasing, and clearly, none of its elements is 0 except itself. This fact is important for defining the factorial of a number  $n$ . Then, we want to establish the range of values that the parameters  $s$  and  $t$  can take such that  $\{n\}_{s,t}$  is an increasing sequence and that  $\{n\}_{s,t} \neq 0$  para  $n \neq 0$ . Orozco[11] showed that

$$|\{n\}_{s,t}| = \{n\}_{|s|,t} \quad (7)$$

for all  $s, t \in \mathbb{Z}$  such that  $s \neq 0$ ,  $t \neq 0$ . The following Lemma exhibits conditions for  $s$  and  $t$  to achieve the aims of this paper.

**Lemma 1.** *Set  $s, t \in \mathbb{Z}$  such that  $s \neq 0$  and  $\Delta > 0$  and suppose  $|q| \neq 1$ . Then*

1.  $\{n\}_{|s|,t}$  is strictly increasing.
2. If  $|s| + t > 1$  with  $|s| \geq 1$ , then  $\{n+1\}_{|s|,t} > \{n\}_{|s|,t} + 1$ , for all  $n \geq 2$ .
3. If  $|s| + t > 1$  with  $|s| \geq 1$ , then  $n \leq \{n\}_{|s|,t}$ .

*Proof.* 1. Take  $s, t \in \mathbb{Z}$ . As  $s \neq 0$ ,  $t > 0$  and  $s^2 + 4t > 0$ , then every  $\{n\}_{|s|,t} \in \mathbb{N}$  for all  $n \in \mathbb{N}$ . Now, suppose  $0 < |q| < 1$ . Then

$$\lim_{n \rightarrow \infty} \{n\}_{s,t} = \lim_{n \rightarrow \infty} \frac{\varphi_{|s|,t}^n - \varphi_{|s|,t}^m}{\varphi_{|s|,t} - \varphi'_{|s|,t}} = \lim_{n \rightarrow \infty} \varphi_{|s|,t}^{n-1} \frac{1 - q^n}{1 - q} = \infty$$

and  $\{n\}_{|s|,t}$  is increasing. From Eq.(3),  $\{n+2\}_{s,t} > \{n+1\}_{s,t}$  and  $\{n+2\}_{s,t} > \{n\}_{s,t}$ . Then  $\{n\}_{s,t}$  is strictly increasing. It is also proven to  $|q| > 1$ . On the other side, suppose that  $t < 0$ . Therefore must be  $s^2 - 4t > 0$  and thus  $|s| > 2\sqrt{-t}$ . We will prove using induction on  $n$  that  $\{n\}_{|s|,-t} \in \mathbb{N}$ . It is easy to notice that  $\{2\}_{|s|,-t} > 2t^{1/2}$ ,  $\{3\}_{|s|,-t} > 3t^{2/2}$ ,  $\{4\}_{s,-t} > 4t^{3/2}$ . We can therefore assume that  $\{n\}_{|s|,-t} > nt^{(n-1)/2}$  and  $\{n+1\}_{|s|,-t} > (n+1)t^{n/2}$ . Thus

$$\begin{aligned} \{n+2\}_{|s|,-t} &= |s|\{n+1\}_{|s|,-t} - t\{n\}_{|s|,-t} \\ &> 2t^{1/2}(n+1)t^{n/2} - tnt^{(n-1)/2} = (n+2)t^{(n+1)/2} \end{aligned}$$

and  $\{n\}_{|s|,-t} \in \mathbb{N}$  for all  $n \in \mathbb{N}$ . And it immediately follows that  $\{n\}_{|s|,-t} < \{n+1\}_{|s|,-t}$  for all  $n \in \mathbb{N}$ .

2. Take  $|s| \geq 2$  and  $t > 0$  such that  $|s| + t > 1$ . We will show by induction that  $\{n+1\}_{|s|,t} > \{n\}_{|s|,t} + 1$  for all  $n \geq 2$ . When  $n = 2$ ,  $\{2\}_{|s|,t} = |s| \geq 2$  and as it is assumed that  $|s| + t \geq 1$ , then  $\{3\}_{|s|,t} = s^2 + t > (1-t)^2 + t = 1 - t + t^2 > \{2\}_{|s|,t} + 1$ . Suppose it is true that  $\{n\}_{|s|,t} > \{n-1\}_{|s|,t} + 1$  and  $\{n+1\}_{|s|,t} > \{n\}_{|s|,t} + 1$ . Then

$$\begin{aligned} \{n+2\}_{|s|,t} &= |s|\{n+1\}_{|s|,t} + t\{n\}_{|s|,t} \\ &> |s|(\{n\}_{|s|,t} + 1) + t(\{n-1\}_{|s|,t} + 1) \\ &= |s|\{n\}_{|s|,t} + t\{n-1\}_{|s|,t} + |s| + t \\ &\geq \{n+1\}_{|s|,t} + 1 \end{aligned}$$

and the statement is true for all  $n \geq 2$ . Now, take  $|s| = 1$  and  $t \geq 1$ . Then  $\{5\}_{1,t} = 1 + 3t + t^2 > (1 + 2t) + 1 = \{4\}_{1,t} + 1$  and it follows by induction that  $\{n+1\}_{1,t} > \{n\}_{1,t} + 1$ .

3. If  $t \geq 1$ , then  $\{n\}_{|s|,-t} > nt^{(n-1)/2} > n$ , for all  $n \in \mathbb{N}$  and  $|s| > 2\sqrt{-t}$ . Now suppose that  $|s| + t \geq 1$  con  $|s| > 2$ . Then  $\{2\}_{|s|,t} > 2$  and  $\{3\}_{|s|,t} > 3$ . Assume by induction hypothesis that  $\{n\}_{|s|,t} > n$ . Then, according to statement 2,

$$\{n+1\}_{|s|,t} > \{n\}_{|s|,t} + 1 > n + 1.$$

Then the statement is true for all  $n \geq 2$ . Now take  $|s| = 1$  and  $t \geq 1$ . Then  $\{6\}_{1,t} = 1 + 4t + 3t^2 > 6$ . Then, using statement 2 and proving by induction we arrive at the truth that  $\{n\}_{1,t} > n$  for all  $n \geq 6$ . □

Take  $t < 0$  in Eq.(6). Then  $|\{n\}_{\pm 2\sqrt{-a}, -a}|$ , with  $a = -t$ , is strictly increasing. Suppose  $\Delta < 0$ . If  $\arctan \sqrt{\frac{b}{a} - 1} \neq \frac{k\pi}{n}$ , for all  $n \geq 1$ , then  $\{n\}_{\sqrt{a}, -b/4} \neq 0$ , for  $n \neq 0$ . However, it is not an increasing sequence. For example,

$$\{n\}_{1,-2} = (0, 1, 1, -1, -3, -1, 5, 7, -3, \dots).$$

For  $s, t \in \mathbb{Z}$  such that  $\Delta > 0$ , the  $(s, t)$ -Fibonomial coefficients are define by

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{s,t} = \frac{\{n\}_{s,t}!}{\{k\}_{s,t}! \{n-k\}_{s,t}!},$$

where  $\{n\}_{s,t}! = \{1\}_{s,t} \{2\}_{s,t} \cdots \{n\}_{s,t}$  is the  $(s, t)$ -factorial or generalized fibotorial. For  $\Delta = 0$ , we define the  $(\pm 2i\sqrt{t}, t)$ -factorial and the  $(\pm 2i\sqrt{t}, t)$ -Fibonomial functions as

$$\{n\}_{\pm 2i\sqrt{t}, t}! = (\pm i\sqrt{t})^{\binom{n}{2}} n!$$

and

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{\pm 2i\sqrt{t}, t} = (\pm i\sqrt{t})^{k(n-k)} \binom{n}{k},$$

respectively. For  $\Delta < 0$  with  $\theta = \arctan \sqrt{\frac{b}{a} - 1} \neq \frac{k\pi}{n}$ ,  $k \in \mathbb{Z}$ ,  $n \geq 1$ , the  $(s, t)$ -factorial is

$$\{n\}_{\sqrt{a}, -b/4}! = \left( \frac{2}{\sqrt{b-a}} \right)^n \left( \frac{b}{4} \right)^{\binom{n+1}{2}} \prod_{k=1}^n \sin(\theta k) \quad (8)$$

and the  $(s, t)$ -Fibonomials coefficients are

$$\left( \frac{b}{4} \right)^{k(n-k)} \frac{\prod_{h=k+1}^n \sin(\theta h)}{\prod_{h=1}^{n-k} \sin(\theta h)}. \quad (9)$$

## 2.2 Deformed $(s, t)$ -exponential function

Set  $s, t \in \mathbb{R}/\{0\}$ . We will set  $q = \varphi'_{s,t}/\varphi_{s,t}$  in the remainder of the paper. For all  $u \in \mathbb{C}$ , we define formally the deformed  $(s, t)$ -exponential function as

$$\exp_{s,t}(z, u) = \begin{cases} \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{z^n}{\{n\}_{s,t}!} & \text{if } u \neq 0; \\ 1 + z & \text{if } u = 0. \end{cases}$$

Also, we define

$$\begin{aligned} \exp_{s,t}(z) &= \exp_{s,t}(z, 1), \\ \text{Exp}_{s,t}(z) &= \exp_{s,t}(z, \varphi_{s,t}), \\ \text{Exp}'_{s,t}(z) &= \exp_{s,t}(z, \varphi'_{s,t}). \end{aligned}$$

If  $|q| < 1$ , the function  $\exp_{s,t}(z, u)$  is

1. an entire function if either  $(q, u) \in E_1$  or  $(q, u) \in E_2$ , where

$$\begin{aligned} E_1 &= \{(q, u) : 0 < |q| < 1, 0 < u < |\varphi_{s,t}|\}, \\ E_2 &= \{(q, u) : |q| > 1, 0 < u < |\varphi'_{s,t}|\}, \end{aligned}$$

2. convergent in the disks

$$\begin{aligned} \mathbb{D}_1 &= \{z \in \mathbb{C} : |z| < |\varphi_{s,t}|/\sqrt{s^2 + 4t}\}, \text{ when } u = |\varphi_{s,t}| \text{ and } 0 < |q| < 1, \\ \mathbb{D}_2 &= \{z \in \mathbb{C} : |z| < |\varphi'_{s,t}|/\sqrt{s^2 + 4t}\}, \text{ when } u = |\varphi'_{s,t}| \text{ and } |q| > 1, \end{aligned}$$

3. convergent in  $z = 0$  when either  $u > |\varphi_{s,t}|$  or  $u > |\varphi'_{s,t}|$ .

Finally, taking  $s \rightarrow \pm 2i\sqrt{t}$ , then the deformed  $(s, t)$ -exponential functions reduce to the following deformed  $(\pm 2i\sqrt{t}, t)$ -exponential functions:

$$\begin{aligned}\exp_{\pm 2i\sqrt{t}, t}(z, u) &= \sum_{n=0}^{\infty} (u/\pm i\sqrt{t})^{(n)} \frac{z^n}{n!}, \\ \exp_{\pm 2i\sqrt{t}, t}(z) &= \sum_{n=0}^{\infty} (\pm i\sqrt{t})^{-\binom{n}{2}} \frac{z^n}{n!}, \\ \exp'_{\pm 2i\sqrt{t}, t}(z) &= \sum_{n=0}^{\infty} (\pm i\sqrt{t})^{\binom{n}{2}} \frac{z^n}{n!}, \\ \text{Exp}_{\pm 2i\sqrt{t}, t}(z) &= \text{Exp}'_{F_{\pm 2i\sqrt{t}, t}}(z) = e^z.\end{aligned}$$

When  $t = -1$ , then

$$\begin{aligned}\exp_{\mp 2, -1}(z, u) &= \sum_{n=0}^{\infty} (\mp u)^{\binom{n}{2}} \frac{z^n}{n!}, \\ \exp_{\mp 2, -1}(z) &= \exp'_{\mp 2, -1}(z) = \sum_{n=0}^{\infty} (\mp)^{\binom{n}{2}} \frac{z^n}{n!}, \\ \text{Exp}_{2, -1}(z) &= \text{Exp}'_{2, -1}(z) = e^z.\end{aligned}$$

Thus

$$\exp_{2, -1}(x, u) = \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{x^n}{n!}$$

and therefore  $\exp_{2, -1}(x, u)$  is the deformed exponential function  $\text{Exp}(x, u)$  and

$$\exp_{2, -1}(x) = \exp'_{2, -1}(x) = \text{Exp}_{2, -1}(x) = \text{Exp}'_{2, -1}(x) = e^x.$$

**Theorem 1.** For all non-zero complex number  $t$  the function  $\exp_{\pm 2i\sqrt{t}, t}(z, u)$

1. is entire if  $|u| \leq |\sqrt{t}|$ .
2. Converge in  $z = 0$  when  $|u| > |\sqrt{t}|$ .

## 2.3 Deformed binomial formula

Orozco [12] defined the  $(u, v)$ -deformed  $(s, t)$ -binomial formula, for  $x, y$  commuting and for non-zero complex numbers  $u, v$ , as

$$(x \oplus_{u,v} y)_{s,t}^{(n)} = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{s,t} u^{\binom{n-k}{2}} v^{\binom{k}{2}} x^{n-k} y^k \quad n \geq 0. \quad (10)$$

He defined the  $(u, v)$ -deformed  $(s, t)$ -analogue of  $(x - y)^n$  by

$$(x \ominus_{u,v} y)_{s,t}^{(n)} \equiv (x \oplus_{u,v} (-y))_{s,t}^{(n)}.$$

When  $u = \varphi$  and  $v = \varphi'$ , then set  $(x \oplus_{\varphi, \varphi'} y)_{s,t}^{(n)} = (x \oplus y)_{\varphi, \varphi'}^n = (x \ominus (-y))_{\varphi, \varphi'}^n$ , where

$$(x \oplus y)_{\varphi, \varphi'}^n = \begin{cases} \prod_{k=0}^{n-1} (\varphi_{s,t}^k x + \varphi_{s,t}'^k y), & \text{if } n \geq 1; \\ 1, & \text{if } n = 1. \end{cases} \quad (11)$$

We can then use the  $(u, v)$ -deformed  $(s, t)$ -binomial formula to establish a formula for the product of deformed exponential functions

$$\exp_{s,t}(x, u) \exp_{s,t}(y, v) = \sum_{n=0}^{\infty} \frac{(x \oplus_{u,v} y)_{s,t}^{(n)}}{\{n\}_{s,t}!}. \quad (12)$$

### 3 Deformed Euler $(s, t)$ -numbers

#### 3.1 Definition

We want that  $|\varphi_{s,t}| > 1$  and that  $|\varphi'_{s,t}| > 1$ , since in this way  $\exp_{s,t}(1, 1)$  makes sense. With this in mind, the following sets are defined

$$\begin{aligned} E_{11}^* &= \{(q, u) \in E_1 : (t, s) \in [-1, \infty) \times (1-t, \infty) \text{ or } (t, s) \in (-\infty, -1) \times [2\sqrt{-t}, \infty)\}, \\ E_{12}^* &= \{(q, u) \in E_1 : (t, s) \in (-\infty, -1) \times (t-1, -2\sqrt{-t}]\}, \\ E_{21}^* &= \{(q, u) \in E_2 : (t, s) \in (-\infty, -1) \times [2\sqrt{-t}, 1-t)\}, \\ E_{22}^* &= \{(q, u) \in E_2 : (t, s) \in [-1, \infty) \times (-\infty, t-1) \text{ or } (t, s) \in (-\infty, -1) \times (-\infty, -2\sqrt{-t}]\}. \end{aligned}$$

**Definition 1.** Set  $s \neq 0$ ,  $t \neq 0$ . If  $(q, u) \in E_{11}^*, E_{12}^*, E_{21}^*, E_{22}^*$ , we define the  $u$ -deformed Euler  $(s, t)$ -numbers as

$$e_{s,t,u} \equiv \exp_{s,t}(1, u) = \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{\{n\}_{s,t}!}. \quad (13)$$

Also, we denote

$$e_{s,t} \equiv \exp_{s,t}(1, 1) \quad (14)$$

and

$$e_{s,t,u}^{-1} \equiv \sum_{n=0}^{\infty} (-1)^n \frac{u^{\binom{n}{2}}}{\{n\}_{s,t}!}. \quad (15)$$

From Eq.(12),

$$e_{s,t,u}^{-1} = \frac{\exp_{s,t}((1 \ominus_{u,u} 1))}{e_{s,t,u}} = \frac{\sum_{n=0}^{\infty} \frac{(1 \ominus_{u,u} 1)_{s,t}^{(n)}}{\{n\}_{s,t}!}}{\sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{\{n\}_{s,t}!}}.$$

**Remark 1.** In reality the  $e_{s,t,u}$  are not numbers but functions in the variable  $u$  for each  $s, t \in \mathbb{R}$  non-zeros.

Some important specializations are

1. Deformed Euler  $(2, -1)$ -number or deformed Euler number

$$e_u \equiv e_{2,-1,u} = \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{n!}$$



2. Deformed Euler (1, 1)-number or deformed Fibonacci-Euler number

$$e_{F,u} \equiv e_{1,1,u} = \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{F_n!}$$

3. Deformed Euler (2, 1)-number or deformed Pell-Euler number

$$e_{P,u} \equiv e_{2,1,u} = \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{P_n!}$$

4. Deformed Euler (1, 2)-number or deformed Jacobsthal-Euler number

$$e_{J,u} \equiv e_{1,2,u} = \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{J_n!}$$

5. Deformed Euler (3, -2)-number or deformed Mersenne-Euler number

$$e_{M,u} \equiv e_{3,-2,u} = \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{M_n!}$$

Likewise, we want  $\exp_{s,t}(1, u)$  to make sense either when  $u = |\varphi_{s,t}|$  or when  $u = |\varphi'_{s,t}|$ . So we define the following sets

$$\begin{aligned} D_{11}^* &= \{(t, s) \in (-\infty, 0) \times (2\sqrt{-t}, \infty)\}, \\ D_{12}^* &= \{(t, s) \in (-\infty, 0) \times (-3\sqrt{-2t}/2, -2\sqrt{-t})\}, \\ D_{21}^* &= \{(t, s) \in (-\infty, 0) \times (2\sqrt{-t}, 3\sqrt{-2t}/2)\}, \\ D_{22}^* &= \{(t, s) \in (-\infty, 0) \times (-\infty, -2\sqrt{-t})\}. \end{aligned}$$

**Definition 2.** Set  $s \neq 0$ ,  $t \neq 0$ . If  $(t, s) \in D_{11}^*, D_{12}^*$ , we define the  $\varphi$ -deformed Euler  $(s, t)$ -numbers as

$$e_{s,t,\varphi} \equiv \exp_{s,t}(1, \varphi_{s,t}) = \sum_{n=0}^{\infty} \frac{\varphi_{s,t}^{\binom{n}{2}}}{\{n\}_{s,t}!}. \quad (16)$$

Also, we denote

$$e_{s,t,\varphi_{s,t}}^{-1} \equiv \sum_{n=0}^{\infty} (-1)^n \frac{\varphi_{s,t}^{\binom{n}{2}}}{\{n\}_{s,t}!}. \quad (17)$$

Equally, if  $(t, s) \in D_{21}^*, D_{22}^*$ , we define the  $\varphi'$ -deformed Euler  $(s, t)$ -numbers as

$$e_{s,t,\varphi'} \equiv \exp_{s,t}(1, \varphi'_{s,t}) = \sum_{n=0}^{\infty} \frac{\varphi'_{s,t}{}^{\binom{n}{2}}}{\{n\}_{s,t}!}. \quad (18)$$

Also, we denote

$$e_{s,t,\varphi'_{s,t}}^{-1} \equiv \sum_{n=0}^{\infty} (-1)^n \frac{\varphi'_{s,t}{}^{\binom{n}{2}}}{\{n\}_{s,t}!}. \quad (19)$$

From Eq.(11),  $(1 \ominus 1)_{\varphi,\varphi'}^n = 0$  for all  $n \geq 1$ . Then

$$e_{s,t,\varphi} e_{s,t,\varphi'}^{-1} = 1, \quad (t, s) \in D_{1,2}^*, D_{2,1}^*$$

and

$$e_{s,t,\varphi'} e_{s,t,\varphi}^{-1} = 1, \quad (t, s) \in D_{1,2}^*, D_{2,1}^*.$$

### 3.2 Estimating the numbers $e_{s,t}$

**Theorem 2.** *Suppose that  $\varphi_{s,t} > 1$ . Then*

$$2 + \frac{1}{s} + h(s, t) < e_{s,t} < 2 + \frac{1}{s} + h(s, t) + \frac{1}{\{7\}_{s,t}! \{8\}_{s,t} - 1} \quad (20)$$

where  $h(s, t) = \frac{1}{\{3\}_{s,t}!} + \frac{1}{\{4\}_{s,t}} + \frac{1}{\{5\}_{s,t}} + \frac{1}{\{6\}_{s,t}!}$ .

*Proof.* To estimate  $e_{s,t}$  we have the sum

$$\begin{aligned} e_{s,t} &= \sum_{n=0}^{\infty} \frac{1}{\{n\}_{s,t}!} = \frac{1}{\{0\}_{s,t}!} + \frac{1}{\{1\}_{s,t}!} + \frac{1}{\{2\}_{s,t}!} + h(s, t) + \sum_{n=7}^{\infty} \frac{1}{\{n\}_{s,t}!} \\ &= 2 + \frac{1}{s} + h(s, t) + \sum_{n=7}^{\infty} \frac{1}{\{n\}_{s,t}!} \end{aligned}$$

This gives the lower bound

$$2 + \frac{1}{s} + h(s, t) < e_{s,t}.$$

To get the upper bound, we combine

$$e_{s,t} = 2 + \frac{1}{s} + h(s, t) + \frac{1}{\{7\}_{s,t}!} \left( 1 + \frac{1}{\{8\}_{s,t}} + \frac{1}{\{8\}_{s,t}\{9\}_{s,t}} + \cdots \right).$$

Since,

$$\{n\}_{s,t} > \{8\}_{s,t} \text{ for all } n > 8, \implies \frac{1}{\{n\}_{s,t}} < \frac{1}{\{8\}_{s,t}},$$

then

$$\begin{aligned} e_{s,t} &< 2 + \frac{1}{s} + h(s, t) + \frac{1}{\{8\}_{s,t}!} \left( 1 + \frac{1}{\{8\}_{s,t}} + \frac{1}{\{8\}_{s,t}^2} + \frac{1}{\{8\}_{s,t}^3} + \cdots \right) \\ &< 2 + \frac{1}{s} + h(s, t) + \frac{1}{\{8\}_{s,t}!} \frac{1}{1 - \frac{1}{\{8\}_{s,t}}} \\ &< 2 + \frac{1}{s} + h(s, t) + \frac{1}{\{7\}_{s,t}! \{8\}_{s,t} - 1}. \end{aligned}$$

The proof is reached. □

The following approaches follow

$$\begin{aligned} 3.70416 &< e_F < 3.70418, \\ 2.6086247947 &< e_P < 2.6086247948, \\ 3.406355917 &< e_J < 3.406355918, \\ 2.3842310161 &< e_M < 2.3842310162. \end{aligned}$$

## 4 Some representations of the $e_{s,t,u}$ , $s \neq 0$ , $t \neq 0$

In this section, we will assume that either  $0 < u \leq |\varphi_{s,t}|$  or  $0 < u \leq |\varphi'_{s,t}|$ , where  $(t, s)$  belongs to some of the sets  $E_{1,1}^*$ ,  $E_{12}^*$ ,  $E_{21}^*$ ,  $E_{22}^*$ ,  $D_{11}^*$ ,  $D_{12}^*$ ,  $D_{21}^*$ ,  $D_{22}^*$ ,  $s \neq 0$ ,  $t \neq 0$ .

## 4.1 As series compression

For  $u \neq 0$  it is easy to show that  $e_{\frac{s}{u}, \frac{t}{u^2}} = e_{s,t,u}$ . Then set  $a = \frac{s}{u}$  and  $b = \frac{t}{u^2}$  and we will prove the following results for  $e_{a,b}$ . The deformed Euler  $(s, t)$ -numbers can be expressed as the sum of the following infinite series. We start by separating the series into sums of even and odd-term fractions and then amplify the fractions of even terms,

$$\begin{aligned} e_{s,t,u} = e_{a,b} &= \sum_{n=0}^{\infty} \frac{1}{\{2n+1\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} \frac{1}{\{2n+1\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{\{2n+1\}_{a,b}}{\{2n+1\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} \frac{\{2n+1\}_{a,b} + 1}{\{2n+1\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} u^{\binom{2n}{2}} \frac{\{2n+1\}_{s,t} + u^{2n+1}}{\{2n+1\}_{s,t}!}. \end{aligned}$$

Then

$$e_{a,b} = \frac{\{1\}_{s,t} + 1}{\{1\}_{a,b}!} + \frac{\{3\}_{a,b} + 1}{\{3\}_{a,b}!} + \frac{\{5\}_{a,b} + 1}{\{5\}_{a,b}!} + \frac{\{7\}_{a,b} + 1}{\{7\}_{a,b}!} + \dots \quad (21)$$

Similarly, we separate the series into sums of odd and even-term fractions and then amplify the fractions of odd terms,

$$\begin{aligned} e_{s,t,u} &= \sum_{n=0}^{\infty} \frac{1}{\{2n+1\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} \\ &= \sum_{n=1}^{\infty} \frac{1}{\{2n-1\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} \frac{\{2n\}_{a,b}}{\{2n\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} \frac{\{2n\}_{a,b} + 1}{\{2n\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} u^{\binom{2n-1}{2}} \frac{\{2n\}_{s,t} + u^{2n-1}}{\{2n\}_{s,t}!}. \end{aligned}$$

Then

$$e_{a,b} = \frac{\{0\}_{s,t} + 1}{\{0\}_{a,b}!} + \frac{\{2\}_{a,b} + 1}{\{2\}_{a,b}!} + \frac{\{4\}_{a,b} + 1}{\{4\}_{a,b}!} + \frac{\{6\}_{a,b} + 1}{\{6\}_{a,b}!} + \dots \quad (22)$$

Adding Eq.(21) and Eq.(22) leads to another beautiful series representing  $e_{s,t,u}$ ,

$$\begin{aligned} 2e_{s,t,u} &= \frac{\{0\}_{s,t} + 1}{\{0\}_{a,b}!} + \frac{\{1\}_{s,t} + 1}{\{1\}_{a,b}!} + \frac{\{2\}_{a,b} + 1}{\{2\}_{a,b}!} + \frac{\{3\}_{a,b} + 1}{\{3\}_{a,b}!} + \frac{\{4\}_{a,b} + 1}{\{4\}_{a,b}!} + \dots \\ &= \sum_{n=0}^{\infty} \frac{\{n\}_{a,b} + 1}{\{n\}_{a,b}!} \\ &= \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}}}{\{n\}_{s,t}!} \left( \frac{\{n\}_{s,t} + u^{n-1}}{u^{n-1}} \right). \end{aligned}$$

The same techniques can be applied to the series of  $e_{s,t,u}^{-1}$ ,

$$\begin{aligned}
e_{s,t,u}^{-1} &= \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} - \sum_{n=0}^{\infty} \frac{1}{\{2n+1\}_{a,b}!} \\
&= \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} - \sum_{n=1}^{\infty} \frac{1}{\{2n-1\}_{a,b}!} \\
&= \sum_{n=0}^{\infty} \frac{1}{\{2n\}_{a,b}!} - \sum_{n=0}^{\infty} \frac{\{2n\}_{a,b}}{\{2n\}_{a,b}!} \\
&= \sum_{n=0}^{\infty} \frac{1 - \{2n\}_{a,b}}{\{2n\}_{a,b}!} = \sum_{n=0}^{\infty} \frac{u^{\binom{2n}{2}}}{\{2n\}_{s,t}!} \left( \frac{u^{2n-1} - \{2n\}_{s,t}}{u^{2n-1}} \right).
\end{aligned}$$

Equally is proven that

$$e_{s,t,u}^{-1} = \sum_{n=0}^{\infty} \frac{\{2n+1\}_{a,b} - 1}{\{2n+1\}_{a,b}!} = \sum_{n=0}^{\infty} \frac{u^{\binom{2n+1}{2}}}{\{2n+1\}_{s,t}!} \left( \frac{\{2n+1\}_{s,t} - u^{2n+1}}{u^{2n+1}} \right). \quad (23)$$

Finally, the  $(s, t)$ -deformed analogue of the representation

$$e = \sum_{n=0}^{\infty} \frac{(3n)^2 + 1}{(3n)!}$$

is

$$\begin{aligned}
e_{s,t,u} &= \sum_{n=0}^{\infty} \frac{1}{\{3n\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{1}{\{3n+1\}_{a,b}!} + \sum_{n=0}^{\infty} \frac{1}{\{3n+2\}_{a,b}!} \\
&= \sum_{n=0}^{\infty} \frac{1 + \{3n\}_{a,b}(\{3n-1\}_{a,b} + 1)}{\{3n\}_{a,b}!} \\
&= \sum_{n=0}^{\infty} \frac{u^{\binom{3n}{2}}}{\{3n\}_{s,t}!} \left( \frac{u^{6n-3} + \{3n\}_{s,t}(\{3n-1\}_{s,t} + u^{6n-3})}{u^{6n-3}} \right).
\end{aligned}$$

The above results were based on [2].

## 4.2 As Engel series expansion

It is known that any positive real number  $x$  can be represented using a non-decreasing sequence of positive integers  $(a_1, a_2, a_3, a_3, \dots)$  such that

$$x = \frac{1}{a_1} + \frac{1}{a_1 a_2} + \frac{1}{a_1 a_2 a_3} + \dots = \frac{1}{a_1} \left( 1 + \frac{1}{a_2} \left( 1 + \frac{1}{a_3} (1 + \dots) \right) \right). \quad (24)$$

Such a representation is called the Engel series expansion of  $x$  [3]. For example, the Engel expansion of  $e$  is  $(1, 1, 2, 3, 4, 5, \dots)$  corresponding to infinite series

$$e = \frac{1}{1} + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} + \dots$$

From Eq.(13), we have the following definition

**Definition 3.** A  $u$ -deformed Engel series expansion of a positive real number  $x$  is the sequence

$$\left( a_0, a_1, \frac{1}{u}a_2, \frac{1}{u^2}a_3, \frac{1}{u^3}a_4, \frac{1}{u^4}a_5, \dots \right), \quad (25)$$

where the  $a_n$  are positive integer, such that

$$x(u) = a_0 + \sum_{n=1}^{\infty} \frac{u^{\binom{n}{2}}}{a_1 a_2 \cdots a_n} \quad (26)$$

is a convergent series. When  $u = 1$ , we obtain the ordinary Engel series expansion of  $x$ .

Then the deformed Engel series expansion for  $e_{s,t,u}$  is

$$\begin{aligned} e_{s,t,u} &= \frac{1}{\{0\}_{s,t}!} + \frac{1}{\{1\}_{s,t}} + \frac{u}{\{1\}_{s,t}\{2\}_{s,t}} + \frac{u^3}{\{1\}_{s,t}\{2\}_{s,t}\{3\}_{s,t}} + \frac{u^6}{\{1\}_{s,t}\{2\}_{s,t}\{3\}_{s,t}\{4\}_{s,t}} + \cdots \\ &= 1 + \frac{1}{\{1\}_{s,t}} \left( + \frac{u}{\{2\}_{s,t}} \left( 1 + \frac{u^2}{\{3\}_{s,t}} \left( 1 + \frac{u^3}{\{4\}_{s,t}} (1 + \cdots) \right) \right) \right) \end{aligned}$$

with sequence

$$\left( 1, 1, \frac{1}{u}\{2\}_{s,t}, \frac{1}{u^2}\{3\}_{s,t}, \frac{1}{u^3}\{4\}_{s,t}, \frac{1}{u^4}\{5\}_{s,t}, \frac{1}{u^5}\{6\}_{s,t}, \dots \right). \quad (27)$$

Some specializations of Eq.(27) are:

$$\begin{aligned} e_{F,u} &= \left( 1, 1, \frac{1}{u}2, \frac{1}{u^2}3, \frac{1}{u^3}5, \frac{1}{u^4}8, \dots \right), \\ e_{P,u} &= \left( 1, 1, \frac{1}{u}2, \frac{1}{u^2}5, \frac{1}{u^3}12, \frac{1}{u^4}29, \dots \right), \\ e_{J,u} &= \left( 1, 1, \frac{1}{u}3, \frac{1}{u^2}5, \frac{1}{u^3}11, \frac{1}{u^4}21, \dots \right), \\ e_{M,u} &= \left( 1, 1, \frac{1}{u}3, \frac{1}{u^2}7, \frac{1}{u^3}15, \frac{1}{u^4}31, \dots \right). \end{aligned}$$

The deformed Engel expansion of  $e_{s,t,u}$  can also be written as an ascending continued fraction [16].

**Theorem 3.** The ascending continued fraction of  $e_{s,t,u}$  is

$$e_{s,t,u} = 2 + \frac{u + u \frac{u^2 + u^2 \frac{u^3 (1 + \cdots)}{\{4\}_{s,t}}}{\{3\}_{s,t}}}{\{2\}_{s,t}}. \quad (28)$$

The fraction Eq.(28) is of form

$$f(n) = 1 + \frac{u^{n-1} f(n+1)}{\{n\}_{s,t}},$$

where  $f(1)$  computes the sum of the terms of 1 to  $\infty$ . Some specializations of Eq.(28) are:

$$\begin{aligned}
e_u &= 2 + \frac{u + u \frac{u^2 + u^2 \frac{u^3(1+\dots)}{4}}{3}}{2}, \\
e_{F,u} &= 2 + \frac{u + u \frac{u^2 + u^2 \frac{u^3(1+\dots)}{3}}{2}}{1}, \\
e_{P,u} &= 2 + \frac{u + u \frac{u^2 + u^2 \frac{u^3(1+\dots)}{12}}{5}}{2}, \\
e_{J,u} &= 2 + \frac{u + u \frac{u^2 + u^2 \frac{u^3(1+\dots)}{5}}{3}}{1}, \\
e_{M,u} &= 2 + \frac{u + u \frac{u^2 + u^2 \frac{u^3(1+\dots)}{15}}{7}}{3}.
\end{aligned}$$

Now, we obtain a pretty nested radical result when  $x$  is taken to the power of both sides of Eq.(28).

**Theorem 4.** *A representation of  $e_{s,t,u}$  using a nested radical is*

$$x^{e_{s,t,u}-2} = \sqrt[{}_{s,t}^{(2)}]{x^u} \sqrt[{}_{s,t}^{(3)}]{x^{u^2}} \sqrt[{}_{s,t}^{(4)}]{x^{u^3}} \sqrt[{}_{s,t}^{(5)}]{x^{u^4}} \dots \quad (29)$$

Some specializations of Eq.(29) are:

$$\begin{aligned}
x^{e_u-2} &= \sqrt{x^u} \sqrt[3]{x^{u^3}} \sqrt[4]{x^{u^6}} \sqrt[5]{x^{u^{10}}} \dots, \\
x^{e_{F,u}-2-u} &= \sqrt{x^{u^3}} \sqrt[3]{x^{u^6}} \sqrt[5]{x^{u^{10}}} \sqrt[8]{x^{u^{15}}} \dots, \\
x^{e_{P,u}-2} &= \sqrt{x^u} \sqrt[5]{x^{u^3}} \sqrt[12]{x^{u^6}} \sqrt[29]{x^{u^{10}}} \dots, \\
x^{e_{J,u}-2-u} &= \sqrt[3]{x^{u^3}} \sqrt[5]{x^{u^6}} \sqrt[11]{x^{u^{10}}} \sqrt[21]{x^{u^{15}}} \dots, \\
x^{e_{M,u}-2} &= \sqrt[3]{x^u} \sqrt[7]{x^{u^3}} \sqrt[15]{x^{u^6}} \sqrt[31]{x^{u^{10}}} \dots.
\end{aligned}$$

### 4.3 As a series of partial Theta functions

Sokal [15] defined the three-variable Rogers-Ramanujan function

$$R(x, y, q) = \sum_{n=0}^{\infty} \frac{x^n y^{\binom{n}{2}}}{(q; q)_n} \quad (30)$$

as a generalization of the theta function

$$\Theta_0(x, y) = \sum_{n=0}^{\infty} x^n y^{\binom{n}{2}}$$

and the deformed exponential function

$$\exp(x, y) = \sum_{n=0}^{\infty} y^{\binom{n}{2}} \frac{x^n}{n!},$$

which are obtained when  $q = 0$  and  $q = 1$ , respectively. It is easy to notice that the deformed  $(s, t)$ -exponential functions are Rogers-Ramanujan functions; we note

$$\begin{aligned} \exp_{s,t}(x, u) &= \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{x^n}{\{n\}_{s,t}} \\ &= \sum_{n=0}^{\infty} \left( \frac{u}{\varphi_{s,t}} \right)^{\binom{n}{2}} \frac{x^n}{[n]_q!} \\ &= \sum_{n=0}^{\infty} \left( \frac{u}{\varphi_{s,t}} \right)^{\binom{n}{2}} \frac{(1-q)^n x^n}{(q; q)_n} \\ &= R((1-q)x, u/\varphi_{s,t}, q) \end{aligned}$$

where  $q = \varphi'_{s,t}/\varphi_{s,t}$ . Following Corollary 6.2 in [15] we get an identity that expresses  $\exp_{s,t}(x, u)$  in terms of the function  $\Theta_0(x, u)$ :

$$\exp_{s,t}(x, u) = \frac{1}{(q; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n+1}{2}}}{(q; q)_n} \Theta_0((1-q)xq^n, u/\varphi_{s,t}),$$

for  $x \in \mathbb{C}$ ,  $0 < u < |\varphi_{s,t}|$ , and  $|q| < 1$ . Then

$$e_{s,t,u} = \frac{1}{(q; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n+1}{2}}}{(q; q)_n} \Theta_0((1-q)q^n, u/\varphi_{s,t})$$

is a representation of  $e_{s,t,u}$  as a series of partial Theta functions. Equally, for all  $x \in \mathbb{C}$  and  $0 < u < |\varphi'_{s,t}|$ , then

$$\exp_{s,t}(x, u) = \frac{1}{(q^{-1}; q^{-1})_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{-\binom{n+1}{2}}}{(q^{-1}; q^{-1})_n} \Theta_0((1-q^{-1})xq^{-n}, u/\varphi'_{s,t})$$

and

$$e_{s,t,u} = \frac{1}{(q^{-1}; q^{-1})_{\infty}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{-\binom{n+1}{2}}}{(q^{-1}; q^{-1})_n} \Theta_0((1-q^{-1})q^{-n}, u/\varphi_{s,t}).$$

#### 4.4 As a continued fraction

It is a known fact [4] that if

$$x = 1 + \sum_{n=1}^{\infty} r_1 r_2 \cdots r_n = 1 + \sum_{n=1}^{\infty} \left( \prod_{k=1}^n r_k \right), \quad (31)$$

then  $x$  has the infinite generalized continued fraction

$$x = \frac{1}{1 - \frac{r_1}{1 + r_1 - \frac{r_2}{1 + r_2 - \frac{r_3}{1 + r_3 - \ddots}}}}. \quad (32)$$

As

$$e_{s,t,u} = 1 + \sum_{n=1}^{\infty} \left( \prod_{k=1}^n \frac{u^{k-1}}{\{k\}_{s,t}} \right)$$

then

$$e_{s,t,u} = \frac{1}{1 - \frac{1}{2 - \frac{\frac{u}{\{2\}_{s,t}}}{1 + \frac{u}{\{2\}_{s,t}} - \frac{\frac{u^2}{\{3\}_{s,t}}}{1 + \frac{u^2}{\{3\}_{s,t}} - \ddots}}}}. \quad (33)$$

Applying an equivalence transformation,

$$e_{s,t,u} = \frac{1}{1 - \frac{1}{2 - \frac{u}{\{2\}_{s,t} + u - \frac{\{2\}_{s,t} u^2}{\{3\}_{s,t} + u^2 - \ddots}}}}. \quad (34)$$

Some specializations of Eq.(34) are:

$$e_{F,u} = \frac{1}{1 - \frac{1}{2 - \frac{u}{1 + u - \frac{u^2}{3 + u^2 - \ddots}}}},$$

$$e_{P,u} = \frac{1}{1 - \frac{1}{2 - \frac{u}{2 + u - \frac{2u^2}{5 + u^2 - \ddots}}}},$$



$$e_{J,u} = \frac{1}{1 - \frac{1}{2 - \frac{u}{1 + u - \frac{u^2}{3 + u^2 - \dots}}}}$$

$$e_{M,u} = \frac{1}{1 - \frac{1}{2 - \frac{u}{3 + u - \frac{3u^2}{7 + u^2 - \dots}}}}$$

## 4.5 As the limit of a sequence

In the following theorems, we obtain the  $(s, t)$ -analog of the remarkable limit

$$e^x = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n.$$

**Theorem 5.** *Set  $s^2 + 4t > 0$ . If  $|q| < 1$ , then*

$$\exp_{s,t}(x, u) = \lim_{n \rightarrow \infty} \left(1 \oplus_{1,u\varphi} \frac{\sqrt{s^2 + 4t}}{\varphi_{s,t}^n} x\right)^{(n)} \quad (35)$$

and

$$e_{s,t,u} = \lim_{n \rightarrow \infty} \left(1 \oplus_{1,u\varphi} \frac{\sqrt{s^2 + 4t}}{\varphi_{s,t}^n}\right)^{(n)}. \quad (36)$$

If  $|q| > 1$ , then

$$\exp_{s,t}(x, u) = \lim_{n \rightarrow \infty} \left(1 \ominus_{1,u\varphi'} \frac{\sqrt{s^2 + 4t}}{\varphi_{s,t}^n} x\right)^{(n)} \quad (37)$$

and

$$e_{s,t,u} = \lim_{n \rightarrow \infty} \left(1 \ominus_{1,u\varphi'} \frac{\sqrt{s^2 + 4t}}{\varphi_{s,t}^n}\right)^{(n)}. \quad (38)$$

*Proof.* Fixing  $x = 1$ ,  $y \mapsto x/\varphi_{s,t}^n$ ,  $v \mapsto u\varphi_{s,t}$  and  $u = 1$  in Eq.(10), we obtain

$$\left(1 \oplus_{1,u\varphi} \frac{x}{\varphi_{s,t}^n}\right)^{(n)} = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_{s,t} (u\varphi_{s,t})^{\binom{k}{2}} \frac{x^k}{\varphi_{s,t}^{nk}}.$$

As  $|q| < 1$ , then

$$\begin{aligned}
\lim_{n \rightarrow \infty} \binom{n}{k}_{s,t} \frac{1}{\varphi_{s,t}^{nk}} &= \lim_{n \rightarrow \infty} \frac{\{n\}_{s,t} \{n-1\}_{s,t} \cdots \{n-k+1\}_{s,t}}{\{k\}_{s,t}! \varphi_{s,t}^{nk}} \\
&= \frac{1}{\{k\}_{s,t}! (\varphi_{s,t} - \varphi'_{s,t})^k} \\
&\quad \times \lim_{n \rightarrow \infty} \left( 1 - \frac{\varphi_{s,t}^n}{\varphi_{s,t}^n} \right) \left( \frac{1}{\varphi_{s,t}} - \frac{\varphi_{s,t}^{(n-1)}}{\varphi_{s,t}^n} \right) \cdots \left( \frac{1}{\varphi_{s,t}^{k-1}} - \frac{\varphi_{s,t}^{(n-k+1)}}{\varphi_{s,t}^n} \right) \\
&= \frac{1}{\{k\}_{s,t}! (\varphi_{s,t} - \varphi'_{s,t})^k \varphi_{s,t}^{0+1+2+\cdots+k-1}} \\
&= \frac{1}{\{k\}_{s,t}! (\varphi_{s,t} - \varphi'_{s,t})^k \varphi_{s,t}^{\binom{k}{2}}}.
\end{aligned}$$

Thus

$$\begin{aligned}
\lim_{n \rightarrow \infty} \left( 1 \oplus_{1,u\varphi} \frac{x}{\varphi_{s,t}^n} \right)^{(n)} &= \lim_{n \rightarrow \infty} \sum_{k=0}^n \binom{n}{k}_{s,t} (u\varphi_{s,t})^{\binom{k}{2}} \frac{x^k}{\varphi_{s,t}^{nk}} \\
&= \sum_{k=0}^{\infty} (u\varphi_{s,t})^{\binom{k}{2}} \frac{x^k}{\{k\}_{s,t}! (\varphi_{s,t} - \varphi'_{s,t})^k \varphi_{s,t}^{\binom{k}{2}}} \\
&= \exp_{s,t} \left( \frac{x}{\varphi_{s,t} - \varphi'_{s,t}}, u \right)
\end{aligned}$$

and the proof is complete. For  $|q| > 1$  the proof is identical.  $\square$

**Theorem 6.** For all  $s, t \in \mathbb{R}$ ,  $s \neq 0$ ,  $t \neq 0$ . If  $|q| < 1$ , then

$$\exp_{s,t}(x, u) = \lim_{n \rightarrow \infty} \left( 1 \oplus_{1,u\varphi} \frac{x}{\{n\}_{s,t}} \right)^{(n)} \quad (39)$$

and

$$e_{s,t,u} = \lim_{n \rightarrow \infty} \left( 1 \oplus_{1,u\varphi} \frac{1}{\{n\}_{s,t}} \right)^{(n)}. \quad (40)$$

If  $|q| > 1$ , then

$$\exp_{s,t}(x, u) = \lim_{n \rightarrow \infty} \left( 1 \oplus_{1,u\varphi'} \frac{x}{\{n\}_{s,t}} \right)^{(n)} \quad (41)$$

and

$$e_{s,t,u} = \lim_{n \rightarrow \infty} \left( 1 \oplus_{1,u\varphi'} \frac{1}{\{n\}_{s,t}} \right)^{(n)}. \quad (42)$$

*Proof.* It is easy to show that

$$\lim_{n \rightarrow \infty} \binom{n}{k}_{s,t} \frac{1}{\{n\}_{s,t}^k} = \frac{1}{\{k\}_{s,t}! \varphi_{s,t}^{\binom{k}{2}}}$$

when  $|q| < 1$ . It follows that

$$\begin{aligned} \lim_{n \rightarrow \infty} \left( 1 \oplus_{1, u\varphi} \frac{x}{\{n\}_{s,t}} \right)^{(n)} &= \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} \binom{n}{k}_{s,t} (u\varphi_{s,t})^{(k)} \frac{x^k}{\{n\}_{s,t}^k} \\ &= \sum_{k=0}^{\infty} (u\varphi_{s,t})^{(k)} \frac{x^k}{\{k\}_{s,t}! \varphi_{s,t}^{(k)}} = \exp_{s,t}(x, u). \end{aligned}$$

For  $|q| > 1$  the proof is analogous. Thus, the proof is achieved.  $\square$

## 5 $e_{s,t,u^{-1}}$ and $e_{s,t,u^{-1}}^{-1}$ , $u \in \mathbb{Q}$ , $|u| > 1$ , are irrationals

**Theorem 7.** *Set  $s, t \in \mathbb{Z}$  such that  $s \neq 0$ ,  $t \neq 0$  and  $s^2 + 4t > 0$  and set  $u \in \mathbb{Q}$ ,  $u > 1$ . Then  $e_{s,t,u^{-1}}$  is irrational.*

*Proof.* Set  $s > 0$ . First, we prove the following inequality

$$e_{s,t,u^{-1}} - s_n < \frac{1}{u^{\binom{n+1}{2}} \{n\}_{s,t}! \{n\}_{s,t}}, \quad (43)$$

where  $s_n$  is the partial sum  $\sum_{k=0}^n (1/u^{\binom{k}{2}} \{k\}_{s,t}!)$ . Set  $u \in \mathbb{N}$  and denote  $s_n$  the partial sum of  $e_{s,t,u^{-1}}$

$$s_n = \sum_{k=0}^n \frac{1}{u^{\binom{k}{2}} \{k\}_{s,t}!}. \quad (44)$$

Then from Lemma 1

$$\begin{aligned} e_{s,t,u^{-1}} - s_n &= \frac{1}{u^{\binom{n+1}{2}} \{n+1\}_{s,t}!} + \frac{1}{u^{\binom{n+2}{2}} \{n+2\}_{s,t}!} + \frac{1}{u^{\binom{n+3}{2}} \{n+3\}_{s,t}!} + \dots \\ &= \frac{1}{u^{\binom{n+1}{2}} \{n+1\}_{s,t}!} \left( 1 + \frac{1}{u^{n+1} \{n+2\}_{s,t}} + \frac{1}{u^{2n+3} \{n+2\}_{s,t} \{n+3\}_{s,t}} + \dots \right) \\ &< \frac{1}{u^{\binom{n+1}{2}} \{n+1\}_{s,t}} \left( 1 + \frac{1}{\{n+1\}_{s,t}} + \frac{1}{\{n+1\}_{s,t}^2} + \dots \right) \\ &= \frac{1}{u^{\binom{n+1}{2}} \{n+1\}_{s,t}!} \frac{\{n+1\}_{s,t}}{\{n+1\}_{s,t} - 1} \\ &< \frac{1}{u^{\binom{n+1}{2}} \{n\}_{s,t}! \{n\}_{s,t}}. \end{aligned}$$

Therefore

$$0 < e_{s,t,u^{-1}} - s_n < \frac{1}{u^{\binom{n+1}{2}} \{n\}_{s,t}! \{n\}_{s,t}}. \quad (45)$$

Suppose that  $e_{s,t,u^{-1}}$  is a rational number, that is,  $e_{s,t,u^{-1}} = p/q$ , where  $p, q$  are positive integers. From Eq.(45) and Lemma 1 statement 3,

$$0 < u^{\binom{q+1}{2}} \{q\}_{s,t}! q (e_{s,t,u^{-1}} - s_q) < \frac{q}{\{q\}_{s,t}} < 1 \quad (46)$$

Set  $u = \frac{n}{m}$ ,  $n, m \in \mathbb{N}$ . If  $u$  is an integer, then both  $u^{\binom{q+1}{2}} \{q\}_{s,t}! q e_{s,t,u-1}$  and

$$u^{\binom{q+1}{2}} \{q\}_{s,t}! q s_q = u^{\binom{q+1}{2}} \{q\}_{s,t}! q \left( 1 + \frac{1}{u^{\binom{1}{2}} \{1\}_{a,b}!} + \frac{1}{u^{\binom{2}{2}} \{2\}_{a,b}!} + \cdots + \frac{1}{u^{\binom{q}{2}} \{q\}_{a,b}!} \right)$$

are integers too. If  $n$  and  $m$  are co-prime and  $m^{\binom{q+1}{2}}$  divide  $\{q\}_{s,t}!(e_{s,t,u-1} - s_q)$ , then  $u^{\binom{q+1}{2}} \{q\}_{s,t}! q (e_{s,t,u-1} - s_q)$  is an integer between 0 and 1. If  $m^{\binom{q+1}{2}}$  does not divide  $\{q\}_{s,t}!(e_{s,t,u-1} - s_q)$ , then  $u^{\binom{q+1}{2}} \{q\}_{s,t}!(e_{s,t,u-1} - s_q)$  is a fraction larger than 1. Therefore the assumption that  $e_{s,t,u-1}$  is a rational is false and for that reason  $e_{s,t,u-1}$  is irrational. If  $s < 0$ , then

$$0 < |e_{-s,t,u-1} - s_n| < \frac{1}{u^{\binom{n+1}{2}} \{n\}_{|s|,t}! \{n\}_{|s|,t}}$$

Now use the above argument. The proof is reached.  $\square$

**Theorem 8.** Set  $s, t \in \mathbb{Z}$  such that  $s \neq 0$ ,  $t \neq 0$  and  $s^2 + 4t \geq 0$  and set  $u \in \mathbb{Q}$ ,  $u > 1$ . Then  $e_{s,t,u-1}^{-1}$  is irrational.

*Proof.* Set  $s > 0$ . First, we prove the following inequality

$$e_{s,t,u-1}^{-1} - s_{2n-1} < \frac{1}{u^{\binom{n}{2}} \{2n\}_{s,t}!}, \quad (47)$$

where  $s_{2n-1}$  is the partial sum  $\sum_{k=0}^{2n-1} ((-1)^k / u^{\binom{k}{2}} \{k\}_{s,t}!)$ . From Theorem 8.16 of [1]

$$e_{s,t,u-1}^{-1} - s_{2n-1} = \sum_{k=0}^{\infty} (-1)^k \frac{1}{u^{\binom{k}{2}} \{k\}_{s,t}!} - \sum_{k=0}^{2n-1} (-1)^k \frac{1}{u^{\binom{k}{2}} \{k\}_{s,t}!} < \frac{1}{u^{\binom{2n}{2}} \{2n\}_{s,t}!}.$$

Note that if  $u$  is an integer, then  $u^{\binom{2n-1}{2}} \{2n-1\}_{s,t}! s_{2n-1}$  is always an integer. Assume that  $e_{s,t,u-1}^{-1}$  is rational, so  $p/q$ , where  $p$  and  $q$  are co-prime, and  $q \neq 0$ . Choose  $n$  such that  $n \geq (q+1)/2$ . Then  $u^{\binom{2n-1}{2}} \{2n-1\}_{s,t}! q e_{s,t,u-1}^{-1}$  is an integer too. Therefore,  $u^{\binom{2n-1}{2}} \{2n-1\}_{s,t}! q (e_{s,t,u-1}^{-1} - s_{2n-1})$  is an integer less than  $q/\{2n\}_{s,t}$ , which is not possible because of the Lemma 1. Suppose that  $u$  is a fraction of the form  $n/m$ , with  $n > m$ . If  $m^{\binom{2n-1}{2}}$  divide to  $\{2n-1\}_{s,t}! q (e_{s,t,u-1}^{-1} - s_{2n-1})$ , then

$$u^{\binom{2n-1}{2}} \{2n-1\}_{s,t}! q (e_{s,t,u-1}^{-1} - s_{2n-1}) \quad (48)$$

is an integer. If  $m^{\binom{2n-1}{2}}$  does not divide  $\{2n-1\}_{s,t}! q (e_{s,t,u-1}^{-1} - s_{2n-1})$ , then Eq(48) is a fraction larger than 1. Therefore, the assumption that  $e_{s,t,u-1}^{-1}$  is a rational is false and thus  $e_{s,t,u-1}^{-1}$  is irrational. If  $s < 0$ , then use

$$0 < |e_{s,t,u-1}^{-1} - s_{2n-1}| < \frac{1}{u^{\binom{n}{2}} \{2n\}_{|s|,t}!}$$

and the previous argument. The proof is achieved.  $\square$

**Proposition 1.** For all  $a \in \mathbb{R}$  it holds that  $\{n\}_{as,a^2t} = a^{n-1}\{n\}_{s,t}$ , which implies that  $\{n\}_{as,a^2t}! = a^{\binom{n}{2}}\{n\}_{s,t}!$ .

*Proof.* It is easy to notice that if  $a > 0$ , then  $\varphi_{as,a^2t} = a\varphi_{s,t}$  and  $\varphi'_{as,a^2t} = a\varphi'_{s,t}$ . Then

$$\{n\}_{as,a^2t} = \frac{(a\varphi_{s,t})^n - (a\varphi'_{s,t})^n}{a(\varphi_{s,t} - \varphi'_{s,t})} = a^{n-1}\{n\}_{s,t}.$$

If  $a < 0$ , then  $\varphi_{as,a^2t} = a\varphi'_{s,t}$  and  $\varphi'_{as,a^2t} = a\varphi_{s,t}$ . Therefore,  $\{n\}_{as,a^2t} = a^{n-1}\{n\}_{a,t}$  for all  $a \in \mathbb{R}$ ,  $a \neq 0$ .  $\square$

**Corollary 1.** Set  $s, t \in \mathbb{Z}$  such that  $s > 0$ ,  $t \neq 0$  and  $s^2 + 4t \geq 0$  and set  $u \in \mathbb{Q}$ ,  $u > 1$ . Then  $e_{as,a^2t,u^{-1}}$  and  $e_{as,a^2t,u^{-1}}^{-1}$  are irrational, for all  $a \in \mathbb{Q}$  such that  $|au| > 1$ .

*Proof.* From Proposition 1

$$e_{as,a^2t,u^{-1}} = \sum_{n=0}^{\infty} \frac{u^{-\binom{n}{2}}}{\{n\}_{as,a^2t}!} = \sum_{n=0}^{\infty} \frac{u^{-\binom{n}{2}}}{a^{\binom{n}{2}}\{n\}_{s,t}!} = \sum_{n=0}^{\infty} \frac{(au)^{-\binom{n}{2}}}{\{n\}_{s,t}} = e_{s,t,(au)^{-1}}.$$

It is also shown that  $e_{as,a^2t,u^{-1}}^{-1} = e_{s,t,(au)^{-1}}$ . Finally, the Theorems 7 and 8 must be applied.  $\square$

## 6 $\varphi$ -Euler and $\varphi'$ -Euler numbers

In the following results, we will give the infinite product representation of the functions  $\text{Exp}_{s,t}(z)$  and  $\text{Exp}'_{s,t}(z)$ .

**Theorem 9.** Set  $s \neq 0$  and  $t \neq 0$ . If  $|q| < 1$ , then the meromorphic continuation of  $\text{Exp}_{s,t}(z)$  is given by

$$\text{Exp}_{s,t}(z) = \prod_{k=0}^{\infty} \frac{\varphi_{s,t}^{k+1}}{\varphi_{s,t}^{k+1} - (\varphi_{s,t} - \varphi'_{s,t})\varphi_{s,t}^k z}.$$

If  $|q| > 1$ , then  $\text{Exp}_{s,t}(z)$  is given by

$$\text{Exp}_{s,t}(z) = \prod_{k=0}^{\infty} \left( 1 - (\varphi_{s,t} - \varphi'_{s,t}) \frac{\varphi_{s,t}^k}{\varphi_{s,t}^{k+1}} z \right).$$

*Proof.* As  $\mathbf{D}_{s,t} \text{Exp}_{s,t}(z) = \text{Exp}_{s,t}(\varphi_{s,t}z)$ , then

$$\frac{\text{Exp}_{s,t}(\varphi_{s,t}z) - \text{Exp}_{s,t}(\varphi'_{s,t}z)}{(\varphi_{s,t} - \varphi'_{s,t})z} = \text{Exp}_{s,t}(\varphi_{s,t}z).$$

Suppose  $|q| < 1$ . Then

$$\text{Exp}_{s,t}(\varphi_{s,t}z) = [1 - (\varphi_{s,t} - \varphi'_{s,t})z]^{-1} \text{Exp}_{s,t}(\varphi'_{s,t}z)$$

and by setting  $w = \varphi_{s,t}z$ ,

$$\text{Exp}_{s,t}(w) = (1 - (1 - q)w)^{-1} \text{Exp}_{s,t}(qw). \quad (49)$$

Now, suppose  $|q| > 1$ . Then

$$\text{Exp}_{s,t}(\varphi'_{s,t}z) = [1 - (\varphi_{s,t} - \varphi'_{s,t})z] \text{Exp}_{s,t}(\varphi_{s,t}z).$$

By setting  $w = \varphi'_{s,t}z$ ,

$$\text{Exp}_{s,t}(w) = [1 - (q^{-1} - 1)w] \text{Exp}_{s,t}(q^{-1}w). \quad (50)$$

Iterating Eqs. (49) and (50) yields desired results.  $\square$

Thus the meromorphic function  $\prod_{k=0}^{\infty} \frac{1}{1 - (1-q)q^k z}$ , with simple poles at  $\frac{1}{(1-q)q^k}$ ,  $k \in \mathbb{N}_0$ , is a good substitute for  $\text{Exp}_{s,t}(z)$ . Analogously, the following theorem is proved.

**Theorem 10.** *Set  $s \neq 0$  and  $t \neq 0$ . If  $|q| < 1$ , then*

$$\text{Exp}'_{s,t}(z) = \prod_{k=0}^{\infty} \left( 1 + (\varphi_{s,t} - \varphi'_{s,t}) \frac{\varphi_{s,t}^k}{\varphi_{s,t}^{k+1}} z \right).$$

If  $|q| > 1$ , then the meromorphic continuation of  $\text{Exp}'_{s,t}(z)$  is given by

$$\text{Exp}'_{s,t}(z) = \prod_{k=0}^{\infty} \frac{\varphi_{s,t}^{(k+1)'}}{\varphi_{s,t}^{(k+1)'} + (\varphi_{s,t} - \varphi'_{s,t})\varphi_{s,t}^k z}.$$

From the Theorems 9 and 10 we obtain the following results.

**Corollary 2.** *Set  $s \neq 0$ . For all  $q$  with  $q \neq 1$ ,*

$$\text{Exp}_{s,t}(z) \text{Exp}'_{s,t}(-z) = \text{Exp}_{s,t}(-z) \text{Exp}'_{s,t}(z) = 1.$$

**Definition 4.** *Define the  $\varphi$ -Euler number  $E_{s,t}$  as  $E_{s,t} = \text{Exp}_{s,t}(1)$  and the  $\varphi'$ -Euler number  $E'_{s,t}$  as  $E'_{s,t} = \text{Exp}'_{s,t}(1)$ .*

The following are the specializations for the infinite product representation of the  $\varphi$ -Euler number:

**Example 1.**

$$\begin{aligned} E_{1,1} &= \prod_{k=0}^{\infty} \frac{(1 + \sqrt{5})^{k+1}}{(1 + \sqrt{5})^{k+1} - \sqrt{5}(1 - \sqrt{5})^k} \\ &= \frac{1 + \sqrt{5}}{1} \cdot \frac{6 + 2\sqrt{5}}{11 + \sqrt{5}} \cdot \frac{16 + 8\sqrt{5}}{26 + 2\sqrt{5}} \cdot \frac{56 + 24\sqrt{5}}{96 + 8\sqrt{5}} \cdot \frac{176 + 80\sqrt{5}}{296 + 24\sqrt{5}} \cdots \\ &= 2.77148239154444838745613 \cdots \end{aligned}$$

**Example 2.**

$$\begin{aligned} E_{2,1} &= \prod_{k=0}^{\infty} \frac{(1 + \sqrt{2})^{k+1}}{(1 + \sqrt{2})^{k+1} - \sqrt{2}(1 - \sqrt{2})^k} \\ &= \frac{1 + \sqrt{2}}{1} \cdot \frac{3 + 2\sqrt{2}}{5 + \sqrt{2}} \cdot \frac{7 + 5\sqrt{2}}{11 + 2\sqrt{2}} \cdot \frac{17 + 2\sqrt{2}}{27 + 5\sqrt{2}} \cdot \frac{41 + 29\sqrt{2}}{65 + 12\sqrt{2}} \cdots \\ &= 2.2266047604700180108 \cdots \end{aligned}$$

### Example 3.

$$\begin{aligned} E_{1,2} &= \prod_{k=0}^{\infty} \frac{2^{k+1}}{2^{k+1} - 3(-1)^k} \\ &= \frac{2}{1} \cdot \frac{4}{7} \cdot \frac{8}{5} \cdot \frac{16}{19} \cdot \frac{32}{29} \cdot \frac{64}{67} \cdot \frac{128}{125} \cdot \frac{256}{259} \cdot \frac{512}{509} \cdot \frac{1024}{1027} \cdot \frac{2048}{2045} \cdots \\ &= -1.649223312843355358 \cdots \end{aligned}$$

### Example 4.

$$\begin{aligned} E_{3,-2} &= \prod_{k=0}^{\infty} \frac{2^{k+1}}{2^{k+1} - 1} \\ &= \frac{2}{1} \cdot \frac{4}{3} \cdot \frac{8}{7} \cdot \frac{16}{15} \cdot \frac{32}{31} \cdot \frac{64}{63} \cdot \frac{128}{127} \cdot \frac{256}{255} \cdot \frac{512}{511} \cdot \frac{1024}{1023} \cdot \frac{2048}{2047} \cdots \\ &= 3.4627466194550636 \cdots \end{aligned}$$

## References

- [1] Apostol, T. *Mathematical analysis* (2nd ed., Addison-Wesley series in mathematics). Reading, Mass.: Addison-Wesley. (1974).
- [2] Brothers, H.J. Improving the convergence of Newton's series approximation for  $e$ . *The College Mathematics Journal*, **35**(1), (2004), pp. 34–39.
- [3] Engel, F. Entwicklung der Zahlen nach Stammbruechen, *Verhandlungen der 52. Versammlung deutscher Philologen und Schulmaenner in Marburg*, (1913), pp. 190–191.
- [4] Euler, L. De fractionibus continuis dissertation. *Commentarii Academiae Scientiarum Petropolitanae*. **9**, (1744), pp. 98–137.
- [5] Euler, L. *Introductio in Analysin Infinitorum*, T. 1, Chapter XVI, p.259, Lausanne, (1748).
- [6] Hermite, C. Sur la fonction exponentielle. *Comptes rendus de l'Académie des Sciences de Paris* (in French). **77**, (1873), pp. 18–24
- [7] Hurwitz, A. Über die Kettenbruchentwicklung der Zahl  $e$ . *Mathematische Werke* (in German). **2**. Basel: Birkhäuser. (1933) [1891], pp. 129–133.
- [8] Liouville, J. Sur l'irrationalité du nombre  $e = 2,718\dots$ . *Journal de Mathématiques Pures et Appliquées*. 1 (in French). **5**(192), (1840)
- [9] MacDivitt, A. R. G., Yanagisawa, Y. An elementary proof that  $e$  is irrational. *The Mathematical Gazette*. **71**(457), (1987)
- [10] Özvatan, M. Generalized Golden-Fibonacci calculus and applications. Ph.D. thesis, Izmir Institute of Technology (2018).
- [11] Orozco, R. Existence of Solutions of Functional-Difference Equations with Proportional Delay on Deformed Generalized Fibonacci Polynomials via Successive Approximation and Bell Polynomials (under revision), arXiv:2211.04450v3 (2024).

- [12] Orozco, R. Deformed Newton's  $(s, t)$ -Binomial Series, Lehmer Series, and Generating Functions of Generalized Fibonomial Coefficients, arXiv:2306.07431v2 (2024).
- [13] Penesi, L. L. Elementary proof that  $e$  is irrational. *American Mathematical Monthly*. **60(7)**, (1953)
- [14] Sondow, J. A Geometric Proof that  $e$  Is Irrational and a New Measure of Its Irrationality. *Amer. Math. Monthly* **113**, (2006), pp. 637–641.
- [15] Sokal, A. The leading root of the partial theta function. *Advances in Mathematics*. **229**, (2011), pp. 2603–2621.
- [16] Kraaikamp, C., Wu, J. On a new continued fraction expansion with non-decreasing partial quotients, *Monatshefte für Mathematik*, **143(4)**, (2004), pp. 285–298, doi:10.1007/s00605-004-0246-3