

MORE ON YET ANOTHER IDEAL VERSION OF THE BOUNDING NUMBER

ADAM KWELA

ABSTRACT. This is a continuation of the paper [J. Symb. Log. 87 (2022), 1065–1092]. For an ideal \mathcal{I} on ω we denote $\mathcal{D}_{\mathcal{I}} = \{f \in \omega^\omega : f^{-1}[\{n\}] \in \mathcal{I} \text{ for every } n \in \omega\}$ and write $f \leq_{\mathcal{I}} g$ if $\{n \in \omega : f(n) > g(n)\} \in \mathcal{I}$, where $f, g \in \omega^\omega$.

We study the cardinal numbers $\mathfrak{b}(\geq_{\mathcal{I}} \cap (\mathcal{D}_{\mathcal{I}} \times \mathcal{D}_{\mathcal{I}}))$ describing the smallest sizes of subsets of $\mathcal{D}_{\mathcal{I}}$ that are unbounded from below with respect to $\leq_{\mathcal{I}}$.

In particular, we examine the relationships of $\mathfrak{b}(\geq_{\mathcal{I}} \cap (\mathcal{D}_{\mathcal{I}} \times \mathcal{D}_{\mathcal{I}}))$ with the dominating number \mathfrak{d} . We show that, consistently, $\mathfrak{b}(\geq_{\mathcal{I}} \cap (\mathcal{D}_{\mathcal{I}} \times \mathcal{D}_{\mathcal{I}})) > \mathfrak{d}$ for some ideal \mathcal{I} , however $\mathfrak{b}(\geq_{\mathcal{I}} \cap (\mathcal{D}_{\mathcal{I}} \times \mathcal{D}_{\mathcal{I}})) \leq \mathfrak{d}$ for all analytic ideals \mathcal{I} . Moreover, we give example of a Borel ideal with $\mathfrak{b}(\geq_{\mathcal{I}} \cap (\mathcal{D}_{\mathcal{I}} \times \mathcal{D}_{\mathcal{I}})) = \text{add}(\mathcal{M})$.

1. INTRODUCTION

Let \mathcal{I} be an ideal on ω . For $f, g \in \omega^\omega$ write $f \leq_{\mathcal{I}} g$ if $\{n \in \omega : f(n) > g(n)\} \in \mathcal{I}$. Moreover, denote $\mathcal{D}_{\mathcal{I}} = \{f \in \omega^\omega : f^{-1}[\{n\}] \in \mathcal{I} \text{ for every } n \in \omega\}$. Let $\mathfrak{b}(\geq_{\mathcal{I}} \cap (\mathcal{D}_{\mathcal{I}} \times \mathcal{D}_{\mathcal{I}}))$ (for simplicity denoted also by $\mathfrak{b}(\mathcal{I})$) be the smallest size of a set in $\mathcal{D}_{\mathcal{I}}$ not bounded from below with respect to the order $\leq_{\mathcal{I}}$ by any member of $\mathcal{D}_{\mathcal{I}}$:

$$\mathfrak{b}(\mathcal{I}) = \min \{|\mathcal{F}| : \mathcal{F} \subseteq \mathcal{D}_{\mathcal{I}} \wedge \neg(\exists g \in \mathcal{D}_{\mathcal{I}} \forall f \in \mathcal{F} g \leq_{\mathcal{I}} f)\}.$$

The cardinal $\mathfrak{b}(\mathcal{I})$ in the case of maximal ideals has been deeply studied by Canjar in 1980s in the context of smallest sizes of cofinal and coinital subsets in ultrapowers $\omega^\omega/\mathcal{U}$ ordered by $\leq_{\mathcal{I}}$, where $\mathcal{U} = \mathcal{I}^*$ (see [11, 12, 13]). In the case of Borel ideals, we have extensively examined $\mathfrak{b}(\mathcal{I})$ in [17].

This research is partially motivated by the study of ideal-QN-spaces. A topological space X is a QN-space if it does not distinguish pointwise and quasi-normal convergence of sequences of real-valued continuous functions defined on X . The research on QN-spaces was initiated by Bukovský, Reclaw and Repický in [9], who proved that the smallest size of non-QN-space is equal to the bounding number \mathfrak{b} . Studies of QN-spaces have been continued in papers [5, 6, 8, 9, 10, 28, 29, 33, 35]. Ideal variants of QN-spaces were introduced in [14] and studied in [7, 26, 36, 37, 30, 31]. For a given ideal \mathcal{I} , the cardinal number $\mathfrak{b}(\mathcal{I})$ characterizes the smallest size of a space which is not \mathcal{I} QN (see [30]).

The paper is organized as follows.

In Section 2 we collect basics about ideals on ω and some known facts about the cardinal numbers $\mathfrak{b}(\mathcal{I})$. In particular, we recall a very useful combinatorial

Date: October 10, 2024.

2010 Mathematics Subject Classification. Primary: 03E05. Secondary: 03E15, 03E17, 03E35.

Key words and phrases. ideal, filter, Borel ideal, analytic ideal, dominating number, ideal with topological representation, bounding number, quasi-normal convergence, quasi-normal space, QN-space,

characterization of $\mathfrak{b}(\mathcal{I})$ from [17], which we use almost exclusively in the rest of the paper.

In the remaining part of the paper we answer two natural questions concerning $\mathfrak{b}(\mathcal{I})$.

The first question is about possible values of $\mathfrak{b}(\mathcal{I})$ for Borel ideals: in [17] it is shown that there are Σ_2^0 ideals \mathcal{I} with $\mathfrak{b}(\mathcal{I}) = \aleph_1$ provable in ZFC ([17, Corollary 7.3 and Theorem 7.4]) as well as Σ_2^0 ideals \mathcal{I} with $\mathfrak{b}(\mathcal{I}) = \mathfrak{b}$ provable in ZFC ([17, Example 5.15]). However, it was unknown whether \aleph_1 and \mathfrak{b} in the above can be replaced by other cardinals. In Section 3 we study $\mathfrak{b}(\mathcal{I})$ in the case of ideals with topological representation (such ideals were introduced by Sabok and Zapletal in [32]) obtaining a Π_3^0 ideal with $\mathfrak{b}(\mathcal{I}) = \text{add}(\mathcal{M})$ provable in ZFC.

The second question concerns comparison of $\mathfrak{b}(\mathcal{I})$ with the dominating number \mathfrak{d} . In general, $\aleph_1 \leq \mathfrak{b}(\mathcal{I}) \leq \mathfrak{c}$, for every ideal \mathcal{I} ([17, Theorem 4.2]). Moreover, the upper bound can be improved to $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{b}$ in the case of Π_4^0 ideals \mathcal{I} ([17, Corollary 6.9]; in fact, as shown in [17, Theorem 6.8], this holds even for a larger class of all so-called co-analytic weak P-ideals). On the other hand, Canjar proved in [12, 11, 13] that under $\mathfrak{d} = \mathfrak{c}$ there is always a maximal ideal \mathcal{I} with $\mathfrak{b}(\mathcal{I}) = \text{cf}(\mathfrak{d})$ and that in the model obtained by adding λ Cohen reals to a model of GCH, $\mathfrak{b}(\mathcal{I})$ can be equal to any regular cardinal between \aleph_1 and $\lambda = \mathfrak{d}$, for some maximal ideal \mathcal{I} . However, it was unknown if it is consistent to have an ideal \mathcal{I} with $\mathfrak{b}(\mathcal{I}) > \mathfrak{d}$. We show that the answer is affirmative (Section 4), however $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$ for every analytic ideal \mathcal{I} (Section 5). The latter result uses ideas developed by Debs and Saint Raymond in [15], providing a new method of proving statements about analytic ideals.

2. PRELIMINARIES

By ω we denote the set of all natural numbers. We identify a natural number n with the set $\{0, 1, \dots, n-1\}$. We write $A \subseteq^* B$ if $A \setminus B$ is finite. For a set A and a finite or infinite cardinal number κ , we write $[A]^\kappa = \{B \subseteq A : |B| = \kappa\}$ and $[A]^{<\kappa} = \{B \subseteq A : |B| < \kappa\}$. Moreover, by $\text{cf}(\kappa)$ we denote the cofinality of κ .

2.1. Ideals and σ -ideals. An *ideal on a set X* is a family $\mathcal{I} \subseteq \mathcal{P}(X)$ satisfying the following properties:

- (i) if $A, B \in \mathcal{I}$ then $A \cup B \in \mathcal{I}$;
- (ii) if $A \subseteq B$ and $B \in \mathcal{I}$ then $A \in \mathcal{I}$;
- (iii) \mathcal{I} contains all finite subsets of X ;
- (iv) $X \notin \mathcal{I}$.

Note that, thanks to item (iii), $\bigcup \mathcal{I} = X$ – we will use this observation several times in Section 5. A σ -*ideal on X* is an ideal satisfying one additional property:

- (v) if $\langle A_n : n \in \omega \rangle \in \mathcal{I}^\omega$ then $\bigcup_{n \in \omega} A_n \in \mathcal{I}$.

Note that, unlike most Authors, in our paper every σ -ideal on X already contains all countable subsets of X and has to be a proper subset of $\mathcal{P}(X)$ (i.e., cannot be equal to $\mathcal{P}(X)$). In this paper, by an *ideal* we mean ideal on some countable set, while all σ -ideals will be subsets of $\mathcal{P}(2^\omega)$.

The ideal of all finite subsets of ω is denoted by Fin . We say that an ideal \mathcal{I} on X is *maximal* if $\mathcal{I} \subseteq \mathcal{J}$ implies $\mathcal{I} = \mathcal{J}$, for every ideal \mathcal{J} on X .

If \mathcal{I} and \mathcal{J} are ideals on X and Y , respectively, then we say that \mathcal{I} and \mathcal{J} are *isomorphic*, if there is a bijection $f : Y \rightarrow X$ such that:

$$A \in \mathcal{I} \Leftrightarrow f^{-1}[A] \in \mathcal{J},$$

for every $A \subseteq X$. It is easy to see that many properties of ideals are preserved under isomorphisms of ideals.

For $\mathcal{A} \subseteq \mathcal{P}(X)$ we write:

$$\mathcal{I}(\mathcal{A}) = \left\{ B \subseteq X : B \subseteq^* \bigcup \mathcal{A}' \text{ for some } \mathcal{A}' \in [\mathcal{A}]^{<\omega} \right\}.$$

If $X \notin \mathcal{I}(\mathcal{A})$, then $\mathcal{I}(\mathcal{A})$ is an ideal, which we call *the ideal generated by \mathcal{A}* . Note that in this case $\mathcal{I}(\mathcal{A})$ is the smallest ideal containing \mathcal{A} .

By identifying subsets of a countable set X with their characteristic functions, we can equip $\mathcal{P}(X)$ with the topology of the Cantor space $\{0, 1\}^X$ and therefore assign descriptive complexity to ideals on X . In particular, an ideal \mathcal{I} is analytic if \mathcal{I} is analytic as a subset of the space $\{0, 1\}^X$.

By \mathcal{M} (\mathcal{N}) we denote the σ -ideals of meager (null, respectively) subsets of 2^ω . If $\mathcal{A} \subseteq \mathcal{P}(2^\omega)$ and 2^ω cannot be covered by countably many members of \mathcal{A} , then by $\sigma\mathcal{A}$ we denote the σ -ideal generated by \mathcal{A} , i.e.:

$$\sigma\mathcal{A} = \left\{ B \subseteq 2^\omega : \exists \langle A_n \rangle_{n \in \omega} \in \mathcal{A}^\omega \ B \subseteq \bigcup_{n \in \omega} A_n \right\}.$$

For a σ -ideal I on 2^ω let \bar{I} be the family of all compact subsets of 2^ω belonging to I . We will say that I is generated by compact sets if $I = \sigma\bar{I}$. In particular, $[2^\omega]^{<\omega}$, \mathcal{M} and $\sigma\bar{\mathcal{N}}$ (i.e., the σ -ideal on 2^ω generated by compact null subsets of 2^ω) are generated by compact sets.

2.2. Some cardinal invariants. In our paper we will need the following cardinal invariants:

- the *pseudointersection number*:

$$\mathfrak{p} = \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq [\omega]^\omega \wedge \left(\forall \mathcal{A}_0 \in [\mathcal{A}]^{<\omega} \bigcap \mathcal{A}_0 \neq \emptyset \right) \wedge \left(\forall S \in [\omega]^\omega \exists A \in \mathcal{A} |S \setminus A| = \omega \right) \right\};$$

- the *additivity of the σ -ideal of meager subsets of 2^ω* is given by:

$$\text{add}(\mathcal{M}) = \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq \mathcal{M} \wedge \bigcup \mathcal{A} \notin \mathcal{M} \right\};$$

- the *bounding number \mathfrak{b}* , which is the smallest size of an \leq_{Fin} -unbounded subset of ω^ω , that is:

$$\mathfrak{b} = \min \left\{ |\mathcal{F}| : \mathcal{F} \subseteq \omega^\omega \wedge \neg(\exists g \in \omega^\omega \forall f \in \mathcal{F} f \leq_{\text{Fin}} g) \right\};$$

- the *dominating number \mathfrak{d}* , which is the smallest size of a \leq_{Fin} -dominating subset of ω^ω , that is:

$$\mathfrak{d} = \min \left\{ |\mathcal{F}| : \mathcal{F} \subseteq \omega^\omega \wedge \forall g \in \omega^\omega \exists f \in \mathcal{F} g \leq_{\text{Fin}} f \right\};$$

- the *ultrafilter number \mathfrak{u}* , which is the smallest size of a family generating a maximal ideal.

It is known that:

$$\aleph_1 \leq \mathfrak{p} \leq \text{add}(\mathcal{M}) \leq \mathfrak{b} \leq \mathfrak{d} \leq \mathfrak{c}$$

(see [6, Subsection 9.2]). Moreover, $\mathfrak{b} \leq \mathfrak{u} \leq \mathfrak{c}$, however both $\mathfrak{d} < \mathfrak{u}$ and $\mathfrak{u} \leq \mathfrak{d}$ are consistent (see [4]).

The cardinal $\text{add}(\mathcal{M})$ is a particular case of a more general invariant: if I is a σ -ideal on 2^ω , then:

$$\text{add}(I) = \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq I \wedge \bigcup \mathcal{A} \notin I \right\}.$$

Note that $\text{add}(I) \geq \aleph_1$. Moreover, if I is generated by compact sets, then:

$$\text{add}(I) = \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq \bar{I} \wedge \bigcup \mathcal{A} \notin I \right\}.$$

2.3. A characterization of $\mathfrak{b}(\mathcal{I})$. Let \mathcal{I} be an ideal on a set X . By $\widehat{\mathcal{P}}_{\mathcal{I}}$ we will denote the family of all sequences $\langle A_n : n \in \omega \rangle \in [\mathcal{P}(X)]^\omega$ such that $A_n \in \mathcal{I}$ for all $n \in \omega$ and $A_n \cap A_k = \emptyset$ whenever $n \neq k$. By $\mathcal{P}_{\mathcal{I}}$ we will denote the family of all sequences $\langle A_n : n \in \omega \rangle \in \widehat{\mathcal{P}}_{\mathcal{I}}$ such that $\bigcup \{A_n : n \in \omega\} = X$.

By [17, Theorem 3.10], the studied cardinal number $\mathfrak{b}(\mathcal{I})$ has a useful combinatorial characterization, which we will use almost exclusively in the rest of the paper without any reference:

$$\mathfrak{b}(\mathcal{I}) = \min \left\{ |\mathcal{E}| : \mathcal{E} \subseteq \widehat{\mathcal{P}}_{\mathcal{I}} \wedge \forall \langle A_n \rangle \in \mathcal{P}_{\mathcal{I}} \exists \langle E_n \rangle \in \mathcal{E} \bigcup_{n \in \omega} \left(A_n \cap \bigcup_{i \leq n} E_i \right) \notin \mathcal{I} \right\}.$$

The following immediate observation will simplify some of our considerations.

Remark 2.1. *If $\langle B_n : n \in \omega \rangle$ is a partition of a set X (that is $\bigcup_{n \in \omega} B_n = X$ and $B_n \cap B_m = \emptyset$ for all $n \neq m$) and $\langle A_i : i \in \omega \rangle$ is any sequence of subsets of X , then:*

$$\bigcup_{i \in \omega} \left(A_i \setminus \bigcup_{n < i} B_n \right) = \bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} A_i \right)$$

3. IDEALS WITH TOPOLOGICAL REPRESENTATION

In this Section we will deal with ideals having a topological representation in the sense of Sabok and Zapletal (see [32]).

Let X be a separable metrizable space with a countable dense set D and I be a σ -ideal on X (recall that in our paper every σ -ideal on X contains all singletons and is a proper subset of $\mathcal{P}(X)$). Following [32], we define an ideal on D by:

$$\mathcal{J}_I = \{A \subseteq D : \text{cl}_X(A) \in I\}$$

(here $\text{cl}_X(A)$ denotes the closure of the set A in the space X). We say that an ideal *has a topological representation* if it is isomorphic to some \mathcal{J}_I as above. In such case, we say that it is *represented on X by I* .

Note that $\mathcal{J}_I = \mathcal{J}_{\sigma \bar{I}}$, since \mathcal{J}_I depends only on closed members of I . By [27, Proposition 2.1], two ideals represented by the same σ -ideal I , but defined on different countable dense subsets of X , are isomorphic. Moreover, each ideal with topological representation can be represented on the Cantor space 2^ω by a σ -ideal generated by some family of compact nowhere dense sets ([27, Corollary 1.3]). Finally, every analytic ideal with topological representation is $\mathbf{\Pi}_3^0$, but not $\mathbf{\Sigma}_2^0$ ([27, Theorem 1.4]).

The basic examples of ideals having topological representation are $\text{NWD} = \mathcal{J}_{\mathcal{M}}$, $\text{NULL} = \mathcal{J}_{\sigma \bar{N}}$ and $\text{CTBL} = \mathcal{J}_{[2^\omega] \leq \omega}$. First two of them were introduced by Farah and Solecki in [16].

Before calculating the invariant $\mathfrak{b}(I)$ for ideals with topological representation, we need to introduce two notions.

Definition 3.1. For a σ -ideal I on 2^ω we define:

$$\text{add}'(I) = \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq \bar{I} \wedge \forall_{\langle B_n \rangle \in \bar{I}^\omega} \exists_{A \in \mathcal{A}} \forall_{n \in \omega} A \setminus B_n \neq \emptyset \right\}.$$

The coefficient $\text{add}'(I)$ has been studied implicitly by Fremlin in [19, Definition 22A] and by Kankaanpää in [21]. Note that $\text{add}'(I) = \text{add}'(\sigma\bar{I})$, as $\text{add}'(I)$ depends only on compact members of I .

Proposition 3.2. *Let I be a σ -ideal on 2^ω generated by some family of compact sets. Then*

$$\aleph_1 \leq \text{add}'(I) \leq \text{add}(I).$$

Proof. To show that $\aleph_1 \leq \text{add}'(I)$, fix any countable $\mathcal{A} \subseteq \bar{I}$. Then $\mathcal{A} = \{B_n : n \in \omega\}$. Consider now the sequence $\langle B_n : n \in \omega \rangle \in \bar{I}^\omega$. Clearly, for each $A \in \mathcal{A}$ there is $n \in \omega$ with $A = B_n$, so $A \setminus B_n = \emptyset$. Thus, $\aleph_1 \leq \text{add}'(I)$.

Now we show that $\text{add}'(I) \leq \text{add}(I)$. Since I is generated by some family of compact sets, we have:

$$\begin{aligned} \text{add}(I) &= \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq I \wedge \bigcup \mathcal{A} \notin I \right\} \\ &= \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq \bar{I} \wedge \bigcup \mathcal{A} \notin I \right\} \\ &= \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq \bar{I} \wedge \forall_{\langle B_n \rangle \in \bar{I}^\omega} \bigcup \mathcal{A} \setminus \bigcup_n B_n \neq \emptyset \right\} \\ &= \min \left\{ |\mathcal{A}| : \mathcal{A} \subseteq \bar{I} \wedge \forall_{\langle B_n \rangle \in \bar{I}^\omega} \exists_{A \in \mathcal{A}} A \setminus \bigcup_n B_n \neq \emptyset \right\}. \end{aligned}$$

Now it is easy to see that each family $\mathcal{A} \subseteq \bar{I}$ witnessing $\text{add}(I)$ is also a witness for $\text{add}'(I)$. Hence, $\text{add}'(I) \leq \text{add}(I)$. \square

Proposition 3.3.

- (a) $\text{add}'(\mathcal{M}) = \text{add}(\mathcal{M})$;
- (b) $\mathfrak{p} \leq \text{add}'(\mathcal{N}) = \text{add}'(\sigma\bar{\mathcal{N}}) \leq \text{add}(\mathcal{M})$;
- (c) $\text{add}'([2^\omega]^{\leq \omega}) = \aleph_1$.

Proof. (a): The inequality $\text{add}'(\mathcal{M}) \leq \text{add}(\mathcal{M})$ follows from Proposition 3.2, while $\text{add}'(\mathcal{M}) \geq \text{add}(\mathcal{M})$ is [21, Lemma 3.5].

(b): Since $\text{add}'(I)$ depends only on compact members of I , $\text{add}'(\mathcal{N}) = \text{add}'(\sigma\bar{\mathcal{N}})$. From Proposition 3.2 and [3, Theorem 3.1] we get $\text{add}'(\sigma\bar{\mathcal{N}}) \leq \text{add}(\sigma\bar{\mathcal{N}}) = \text{add}(\mathcal{M})$, while $\mathfrak{p} \leq \text{add}'(\mathcal{N})$ is shown in [19, Theorem 22G].

(c): It follows from Proposition 3.2 and the fact that $\text{add}([2^\omega]^{\leq \omega}) = \omega_1$. \square

Definition 3.4. We say that a σ -ideal I on 2^ω is invariant over basic sets if given any $s \in 2^{<\omega}$ we have:

$$A \in I \Leftrightarrow f_s[A] \in I,$$

where $f_s : 2^\omega \rightarrow V_s$ is given by $f_s(x) = s \cap x$ and $V_s = \{x \in 2^\omega : x \upharpoonright \text{lh}(s) = s\}$ is the basic clopen set associated to s (here $\text{lh}(s)$ is the length of s).

Proposition 3.5. *The σ -ideals \mathcal{M} , \mathcal{N} , $\sigma\bar{\mathcal{N}}$ and $[2^\omega]^{<\omega}$ are invariant over basic sets.*

Proof. In the case of $[2^\omega]^{<\omega}$, it suffices to observe that each f_s is a bijection (so $|A| = |f_s[A]|$). Since each f_s is also a homeomorphism, \mathcal{M} is invariant over basic sets. Finally, since for each measurable $A \subseteq 2^\omega$ the measure of $f_s[A]$ is equal to the measure of A multiplied by $\frac{1}{2^{\ln(s)}}$, \mathcal{N} and $\sigma\overline{\mathcal{N}}$ are invariant over basic sets. \square

Finally, we are ready to calculate the invariants $\mathfrak{b}(\mathcal{I})$ in the case of ideals with topological representation.

Theorem 3.6. *Let I be an invariant over basic sets σ -ideal on 2^ω generated by some family of compact sets. Then:*

$$\min\{\text{add}'(I), \mathfrak{b}\} \leq \mathfrak{b}(\mathcal{J}_I) \leq \text{add}'(I).$$

Moreover, if \mathcal{J}_I is analytic, then:

$$\mathfrak{b}(\mathcal{J}_I) = \min\{\text{add}'(I), \mathfrak{b}\}.$$

Proof. The case of analytic ideals will follow from the general case. Indeed, each analytic ideal with topological representation \mathcal{J}_I is $\mathbf{\Pi}_3^0$ (by [27, Theorem 1.4]), so using [17, Corollary 6.9] we get $\mathfrak{b}(\mathcal{J}_I) \leq \mathfrak{b}$.

By [27, Proposition 2.1], without loss of generality we may assume that \mathcal{J}_I is an ideal on $D = \{x \in 2^\omega : \exists k \in \omega \forall n \geq k x(n) = 0\}$. Note that $f_s[D] = D \cap V_s$, where V_s and f_s are as in Definition 3.4.

We start with $\mathfrak{b}(\mathcal{J}_I) \geq \min\{\text{add}'(I), \mathfrak{b}\}$. Fix any $\kappa < \min\{\text{add}'(I), \mathfrak{b}\}$. We need to show that $\kappa < \mathfrak{b}(\mathcal{J}_I)$. Let $\{\langle E_n^\alpha : n \in \omega \rangle : \alpha < \kappa\} \subseteq \widehat{\mathcal{P}}_{\mathcal{J}_I}$ be arbitrary. Since $\omega \cdot \kappa < \text{add}'(I)$ (by Proposition 3.2), there is $\langle B_n : n \in \omega \rangle \in \overline{I}^\omega$ such that for each $n \in \omega$ and $\alpha < \kappa$ there exists $f_\alpha(n) \in \omega$ with $\text{cl}_{2^\omega}(E_n^\alpha) \subseteq B_{f_\alpha(n)}$.

Since $\kappa < \mathfrak{b}$, there exists $g \in \omega^\omega$ such that $f_\alpha \leq^* g$, for all $\alpha < \kappa$. Without loss of generality, we may assume that g is strictly increasing.

Let $\{q_n : n \in \omega\}$ be an enumeration of D and define $C_0 = \bigcup_{i \leq g(1)} (B_i \cap D) \cup \{q_0\}$ and $C_n = (\bigcup_{i \leq g(n+1)} (B_i \cap D) \cup \{q_n\}) \setminus \bigcup_{i < n} C_i$. Then $\langle C_n : n \in \omega \rangle \in \mathcal{P}_{\mathcal{J}_I}$, since:

$$\text{cl}_{2^\omega}(C_n) \subseteq \text{cl}_{2^\omega} \left(\bigcup_{i \leq g(n+1)} B_i \cup \{q_n\} \right) = \bigcup_{i \leq g(n+1)} B_i \cup \{q_n\} \in I.$$

Fix now $\alpha < \kappa$. We need to show that $\bigcup_{n \in \omega} (C_n \cap \bigcup_{i \leq n} E_i^\alpha) \in \mathcal{J}_I$.

Let $k_\alpha \in \omega$ be such that $f_\alpha(m) \leq g(m)$, for every $m > k_\alpha$. Observe that for each $m > k_\alpha$ we get:

$$E_m^\alpha \subseteq \text{cl}_{2^\omega}(E_m^\alpha) \cap D \subseteq B_{f_\alpha(m)} \cap D \subseteq \bigcup_{n \leq g(m)} B_n \cap D \subseteq \bigcup_{n \leq m-1} C_n.$$

and consequently:

$$E_m^\alpha \cap \bigcup_{n \in \omega} \left(C_n \cap \bigcup_{i \leq n} E_i^\alpha \right) \subseteq E_m^\alpha \cap \bigcup_{n \geq m} C_n = \emptyset,$$

since $\langle E_n^\alpha : n \in \omega \rangle \in \widehat{\mathcal{P}}_{\mathcal{J}_I}$. Hence,

$$\bigcup_{n \in \omega} \left(C_n \cap \bigcup_{i \leq n} E_i^\alpha \right) \subseteq \left(\bigcup_{m \in \omega} E_m^\alpha \right) \cap \left(\bigcup_{n \in \omega} \left(C_n \cap \bigcup_{i \leq n} E_i^\alpha \right) \right) \subseteq \bigcup_{m \leq k_\alpha} E_m^\alpha \in \mathcal{J}_I.$$

Now we move to $\mathfrak{b}(\mathcal{J}_I) \leq \text{add}'(I)$. Let $\mathcal{A} \subseteq \bar{I}$ be the family of cardinality $\text{add}'(I)$ such that for each $\langle B_n : n \in \omega \rangle \in \bar{I}^\omega$ one can find $A \in \mathcal{A}$ with $A \setminus B_n \neq \emptyset$, for all n .

Without loss of generality we may assume that $\text{cl}_{2^\omega}(A \cap D) = A$ for each $A \in \mathcal{A}$. Indeed, given $A \in \mathcal{A}$, fix any countable $E \subseteq A$ such that $\text{cl}_{2^\omega}(E) = A$ (this is possible as A is a closed subset of 2^ω) and enumerate $E = \{e_i : i \in \omega\}$. For each $i \in \omega$ find a sequence $\langle x_{i,j} : j \in \omega \rangle \in [D \cap B(e_i, \frac{1}{2^i})]^\omega$ converging to e_i (here $B(e_i, \frac{1}{2^i})$ denotes the open ball in 2^ω of radius $\frac{1}{2^i}$ centered at e_i). By defining $\hat{A} = A \cup \{x_{i,j} : i, j \in \omega\}$ we get a closed set such that $\hat{A} = \text{cl}_{2^\omega}(\hat{A} \cap D)$. Moreover, as $A \subseteq \hat{A}$, the family $\{\hat{A} : A \in \mathcal{A}\}$ satisfies the same property as \mathcal{A} . Thus, we will assume from now on that $\text{cl}_{2^\omega}(A \cap D) = A$, for every $A \in \mathcal{A}$.

Fix any bijection $h : \omega \rightarrow 2^{<\omega}$ and define $E_n^A = f_{h(n)}[A \cap D]$ for each $n \in \omega$ and $A \in \mathcal{A}$. Observe that:

$$\text{cl}_{2^\omega}(E_n^A) = \text{cl}_{2^\omega}(f_{h(n)}[A \cap D]) \subseteq f_{h(n)}[\text{cl}_{2^\omega}(A \cap D)] = f_{h(n)}[A] \in I,$$

since $f_{h(n)}$ is a homeomorphism and I is invariant over basic sets. Hence, each E_n^A belongs to \mathcal{J}_I . We claim that the family $\{\langle \hat{E}_n^A : n \in \omega \rangle : A \in \mathcal{A}\} \subseteq \hat{\mathcal{P}}_{\mathcal{J}_I}$, where $\hat{E}_n^A = E_n^A \setminus \bigcup_{i < n} E_i^A$ for all $n \in \omega$ and $A \in \mathcal{A}$, witnesses $\mathfrak{b}(\mathcal{J}_I) \leq \text{add}'(I)$.

Fix any partition $\langle B_n : n \in \omega \rangle \in \mathcal{P}_{\mathcal{J}_I}$ of D .

Note that given any $n \in \omega$, we have $V_{h(n)} \cap \bigcup_{i < n} \text{cl}_{2^\omega}(B_i) \subseteq \bigcup_{i < n} \text{cl}_{2^\omega}(B_i) \in I$. Since I is invariant over basic sets and $f_{h(n)}$ is a bijection, the set $f_{h(n)}^{-1}[V_{h(n)} \cap \bigcup_{i < n} \text{cl}_{2^\omega}(B_i)]$ belongs to \bar{I} . Hence, there is $A \in \mathcal{A}$ such that:

$$A \setminus \left(f_{h(n)}^{-1} \left[V_{h(n)} \cap \bigcup_{i < n} \text{cl}_{2^\omega}(B_i) \right] \right) \neq \emptyset$$

for all $n \in \omega$. To finish the proof, we will show that $B = \bigcup_{n \in \omega} (B_n \cap \bigcup_{i \leq n} \hat{E}_n^A)$ is dense in 2^ω (hence, $\text{cl}_{2^\omega}(B) = 2^\omega \notin I$ and $B \notin \mathcal{J}_I$).

Fix any $s \in 2^{<\omega}$. We need to find an element of B belonging to V_s . Since

$$\text{cl}_{2^\omega}(A \cap D) \setminus f_s^{-1} \left[V_s \cap \bigcup_{i < h^{-1}(s)} \text{cl}_{2^\omega}(B_i) \right] = A \setminus \left(f_s^{-1} \left[V_s \cap \bigcup_{i < h^{-1}(s)} \text{cl}_{2^\omega}(B_i) \right] \right) \neq \emptyset,$$

and $f_s^{-1} \left[V_s \cap \bigcup_{i < h^{-1}(s)} \text{cl}_{2^\omega}(B_i) \right]$ is closed (as f_s is a homeomorphism), we can find $x_s \in (A \cap D) \setminus \left(f_s^{-1} [V_s \cap \bigcup_{i < h^{-1}(s)} B_i] \right)$. Then, using Remark 2.1, we get:

$$\begin{aligned} f_s(x_s) &\in V_s \cap \left(E_{h^{-1}(s)}^A \setminus \bigcup_{i < h^{-1}(s)} B_i \right) \subseteq V_s \cap \bigcup_{j \leq h^{-1}(s)} \left(\hat{E}_j^A \setminus \bigcup_{i < h^{-1}(s)} B_i \right) \\ &\subseteq V_s \cap \bigcup_{j \leq h^{-1}(s)} \left(\hat{E}_j^A \setminus \bigcup_{i < j} B_i \right) \subseteq V_s \cap B. \end{aligned}$$

This finishes the proof. □

Next result gives some bounds for three well-known ideals having topological representation: $\text{NWD} = \mathcal{J}_{\mathcal{M}}$, $\text{NULL} = \mathcal{J}_{\sigma\bar{\mathcal{N}}}$ and $\text{CTBL} = \mathcal{J}_{[2^\omega] \leq \omega}$.

Corollary 3.7.

- (a) $\mathfrak{b}(\text{NWD}) = \text{add}(\mathcal{M})$.

- (b) $\mathfrak{p} \leq \mathfrak{b}(\text{NULL}) \leq \text{add}(\mathcal{M})$.
- (c) $\mathfrak{b}(\text{CTBL}) = \aleph_1$.

Proof. Follows from Theorem 3.6, Proposition 3.3 and the fact that $\aleph_1 \leq \text{add}(\mathcal{M}) \leq \mathfrak{b}$ (by [2, Theorem 3.11]). \square

We end this Section with calculation of $\mathfrak{b}(\mathcal{I})$ for one more well-known ideal defined on the rationals.

Recall that CONV is the ideal on $\mathbb{Q} \cap [0, 1]$ generated by sequences in $\mathbb{Q} \cap [0, 1]$ that are convergent in $[0, 1]$. This is a Σ_4^0 ideal contained in NWD , NULL and CTBL (see [20, Subsection 3.4]).

Proposition 3.8. $\mathfrak{b}(\text{CONV}) = \aleph_1$.

Proof. By [18, Proposition 4.1], $\mathfrak{b}(\mathcal{I}) \geq \aleph_1$ for every ideal \mathcal{I} , so we only need to show that $\mathfrak{b}(\mathcal{I}) \leq \aleph_1$.

Fix any family $\{A_n^\alpha : n \in \omega, \alpha < \aleph_1\} \subseteq \text{CONV}$ such that:

- each A_n^α is a sequence in $\mathbb{Q} \cap [0, 1]$ convergent in $[0, 1]$;
- $\lim A_n^\alpha \neq \lim A_m^\beta$, for all $(\alpha, n), (\beta, m) \in \aleph_1 \times \omega$ such that $(\alpha, n) \neq (\beta, m)$;
- $A_n^\alpha \cap A_m^\alpha = \emptyset$ for all $\alpha < \aleph_1$ and $n, m \in \omega$ such that $n \neq m$.

We will show that this family witnesses $\mathfrak{b}(\text{CONV}) \leq \aleph_1$.

Let $\langle B_n : n \in \omega \rangle \in \mathcal{P}_{\text{CONV}}$. Since $\bigcup_{n \in \omega} \text{cl}_{[0,1]}(B_n)$ is countable, there is $\alpha < \aleph_1$ such that $\lim A_m^\alpha \notin \bigcup_{n \in \omega} \text{cl}_{[0,1]}(B_n)$, for all $m \in \omega$. In particular, $A_m^\alpha \cap B_n$ is finite for all $n, m \in \omega$.

Suppose to the contrary that $B = \bigcup_{n \in \omega} (B_n \cap \bigcup_{i \leq n} A_i^\alpha) \in \mathcal{I}$, i.e., there are sequences $D_0, \dots, D_m \subseteq \mathbb{Q} \cap [0, 1]$ convergent in $[0, 1]$ and such that $B \subseteq^* \bigcup_{i \leq m} D_i$. Find $j \in \omega$ such that $\lim A_j^\alpha \neq \lim D_i$ for all $i \leq m$. Observe that:

$$A_j^\alpha \setminus \bigcup_{n < j} B_n = A_j^\alpha \cap \bigcup_{n \geq j} B_n \subseteq \bigcup_{n \geq j} \left(B_n \cap \bigcup_{i \leq n} A_i^\alpha \right) \subseteq B.$$

Now, since $A_j^\alpha \cap \bigcup_{n < j} B_n$ is finite, we get $A_j^\alpha \subseteq^* B \subseteq^* \bigcup_{i \leq m} D_i$. Thus, there is $i \leq m$ with $A_j^\alpha \cap D_i$ infinite. This contradicts $\lim A_j^\alpha \neq \lim D_i$ and finishes the proof. \square

4. AN IDEAL WITH $\mathfrak{b}(\mathcal{I}) > \mathfrak{d}$

In this Section we will show that, consistently, there is an ideal \mathcal{I} with $\mathfrak{b}(\mathcal{I}) > \mathfrak{d}$. It will follow from the following more general result.

Lemma 4.1. *If $\kappa < \text{cf}(\mathfrak{c}) \leq \mathfrak{u} = \mathfrak{c} = \mathfrak{c}^\kappa$, then there exists an ideal \mathcal{I} such that $\mathfrak{b}(\mathcal{I}) > \kappa$.*

Proof. Without loss of generality we may assume that κ is infinite (as for finite κ this is trivial). Recall that for $\mathcal{A} \subseteq \mathcal{P}(\omega)$ we write:

$$\mathcal{I}(\mathcal{A}) = \left\{ B \subseteq \omega : B \subseteq^* \bigcup \mathcal{A}' \text{ for some } \mathcal{A}' \in [\mathcal{A}]^{<\omega} \right\}.$$

In particular, $\mathcal{I}(\mathcal{A})$ is an ideal if and only if $\omega \notin \mathcal{I}(\mathcal{A})$.

Since $|\mathcal{P}(\omega)^{\kappa \times \omega}| = \mathfrak{c}^\kappa = \mathfrak{c}$, we can fix an enumeration $\{h_\alpha : \alpha < \mathfrak{c}\}$ of $\mathcal{P}(\omega)^{\kappa \times \omega}$ such that for each $h \in \mathcal{P}(\omega)^{\kappa \times \omega}$ and $\alpha < \mathfrak{c}$ there is $\alpha < \beta < \mathfrak{c}$ with $h_\beta = h$.

We will recursively define a sequence $\{\mathcal{I}_\alpha : \alpha < \mathfrak{c}\}$ of subsets of $\mathcal{P}(\omega)$ such that for each $\alpha < \mathfrak{c}$:

- (i) \mathcal{I}_α is an ideal;
- (ii) \mathcal{I}_α is generated by at most $|\alpha \cdot \kappa|$ sets;
- (iii) if $\beta < \alpha$, then $\mathcal{I}_\beta \subseteq \mathcal{I}_\alpha$;
- (iv) if $h_\alpha[\kappa \times \omega] = \{h_\alpha(\gamma, n) : \gamma < \kappa, n \in \omega\} \subseteq \bigcup_{\delta < \alpha} \mathcal{I}_\delta$, then there is $\langle B_n : n \in \omega \rangle \in \mathcal{P}_{\mathcal{I}_\alpha}$ such that $\bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} h_\alpha(\gamma, i) \right) \in \mathcal{I}_\alpha$, for all $\gamma < \kappa$.

Assume that $\alpha < \mathfrak{c}$ and \mathcal{I}_β , for all $\beta < \alpha$, are already defined. Observe that $\hat{\mathcal{I}}_\alpha = \bigcup_{\delta < \alpha} \mathcal{I}_\delta$ is an ideal (as a union of an increasing sequence of ideals) generated by at most $|\alpha \cdot (\alpha \cdot \kappa)| = |\alpha \cdot \kappa|$ sets (since κ is infinite). Hence, if $h_\alpha(\gamma, n) \notin \hat{\mathcal{I}}_\alpha$ for some $(\gamma, n) \in \kappa \times \omega$, then just put $\mathcal{I}_\alpha = \hat{\mathcal{I}}_\alpha$ and observe that \mathcal{I}_α is as needed.

Assume now that $h_\alpha[\kappa \times \omega] \subseteq \hat{\mathcal{I}}_\alpha$. Recursively define a sequence $\langle C_n : n \in \omega \rangle$ of subsets of ω such that $C_n \notin \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{C_j : j < n\})$ and $\mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{C_j : j \leq n\})$ is an ideal. This is possible as $|\alpha \cdot (\alpha \cdot \kappa) + n| < \mathfrak{c} = \mathfrak{u}$ guarantees that $\mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{C_j : j < n\})$ is not maximal. Next, for each $n \in \omega$ put $B_n = (\{n\} \cup C_n) \setminus \bigcup_{j < n} B_j$ and note that $\langle B_n : n \in \omega \rangle$ is a partition of ω and $B_n \notin \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{B_j : j < n\}) = \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{C_j : j < n\})$, for each n . Moreover, $\mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{B_k : k \in \omega\})$ is an ideal, as otherwise we would have:

$$\omega \in \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{B_k : k \in \omega\}) = \bigcup_{n \in \omega} \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{B_k : k < n\}) = \bigcup_{n \in \omega} \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{C_k : k < n\}),$$

which contradicts the fact that $\omega \notin \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{C_k : k \leq n\})$ for each n .

Define:

$$\mathcal{I}_\alpha = \mathcal{I} \left(\hat{\mathcal{I}}_\alpha \cup \{B_n : n \in \omega\} \cup \left\{ \bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} h_\alpha(\gamma, i) \right) : \gamma < \kappa \right\} \right).$$

Obviously, items (iii) and (iv) are satisfied. What is more, item (ii) is also satisfied as $|\alpha \cdot \kappa + \omega + \kappa| = |\alpha \cdot \kappa|$. Thus, we only need to check that $\omega \notin \mathcal{I}_\alpha$.

Assume to the contrary that $\omega \in \mathcal{I}_\alpha$. Then there are $k, m \in \omega$, $\gamma_0, \dots, \gamma_m < \kappa$ and $A \in \hat{\mathcal{I}}_\alpha$ such that:

$$\omega = A \cup \bigcup_{j < k} B_j \cup \bigcup_{j \leq m} \left(\bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} h_\alpha(\gamma_j, i) \right) \right).$$

However, since $\langle B_n : n \in \omega \rangle$ is a partition of ω , we have:

$$\begin{aligned} B_k &= B_k \cap \omega = B_k \cap \left(A \cup \bigcup_{j < k} B_j \cup \bigcup_{j \leq m} \left(\bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} h_\alpha(\gamma_j, i) \right) \right) \right) \\ &\subseteq A \cup \bigcup_{j \leq m} \bigcup_{i \leq k} h_\alpha(\gamma_j, i) \in \hat{\mathcal{I}}_\alpha, \end{aligned}$$

which contradicts $B_k \notin \mathcal{I}(\hat{\mathcal{I}}_\alpha \cup \{B_j : j < k\})$. Thus, $\omega \notin \mathcal{I}_\alpha$.

Once the recursive construction of $\{\mathcal{I}_\alpha : \alpha < \mathfrak{c}\}$ is completed, define $\mathcal{I} = \bigcup_{\alpha < \mathfrak{c}} \mathcal{I}_\alpha$. Clearly, \mathcal{I} is an ideal (by items (i) and (iii)). To finish the proof we need to show that $\mathfrak{b}(\mathcal{I}) > \kappa$. Fix any $\{\langle E_n^\gamma : n \in \omega \rangle : \gamma < \kappa\} \subseteq \hat{\mathcal{P}}_{\mathcal{I}}$. Then for each $(\gamma, n) \in \kappa \times \omega$ there is $\beta(\gamma, n) < \mathfrak{c}$ such that $E_n^\gamma \in \mathcal{I}_{\beta(\gamma, n)}$. Denote $\beta = \sup\{\beta(\gamma, n) : (\gamma, n) \in \kappa \times \omega\}$. Since $|\kappa \times \omega| = \kappa < \text{cf}(\mathfrak{c})$, we get that $\beta < \mathfrak{c}$. Thus, we can find $\beta < \alpha < \mathfrak{c}$ with $h_\alpha(\gamma, n) = E_n^\gamma$, for all $(\gamma, n) \in \kappa \times \omega$. By item

(iii), $h_\alpha[\kappa \times \omega] \subseteq \bigcup_{\delta < \alpha} \mathcal{I}_\delta$. Using item (iv) for α we get $\langle B_n : n \in \omega \rangle \in \mathcal{P}_{\mathcal{I}_\alpha} \subseteq \mathcal{P}_{\mathcal{I}}$ such that $\bigcup_{n \in \omega} (B_n \cap \bigcup_{i \leq n} E_i^\gamma) \in \mathcal{I}_\alpha \subseteq \mathcal{I}$, for all $\gamma < \kappa$. Therefore, $\mathfrak{b}(\mathcal{I}) > \kappa$ and the proof is completed. \square

Theorem 4.2. *If $\mathfrak{d} < \text{cf}(\mathfrak{c}) \leq \mathfrak{u} = \mathfrak{c} = \mathfrak{c}^\mathfrak{d}$, then there exists an ideal \mathcal{I} with $\mathfrak{b}(\mathcal{I}) > \mathfrak{d}$. In particular, it is consistent that there exists an ideal \mathcal{I} such that $\mathfrak{b}(\mathcal{I}) > \mathfrak{d}$.*

Proof. The first part follows directly from Lemma 4.1.

We will show that the assumptions $\mathfrak{d} < \text{cf}(\mathfrak{c}) \leq \mathfrak{u} = \mathfrak{c} = \mathfrak{c}^\mathfrak{d}$ are consistent (specifically, that they hold in the generic extension obtained by adding \aleph_2 random reals to the model of GCH).

Suppose that GCH is satisfied in the ground model V . Let

$$M(\aleph_2) = \{C \subseteq 2^{\aleph_2} : C \text{ is a closed set of positive measure}\}$$

(ordered by the inclusion modulo null sets) be a forcing which adds \aleph_2 random reals (here we consider the product measure on 2^{\aleph_2}). Let G denote a $M(\aleph_2)$ -generic filter.

It is known that $\mathfrak{d} = \aleph_1$ and $\mathfrak{u} = \mathfrak{c} = \aleph_2$ in $V[G]$ (see e.g. [4, p. 474]), so $\mathfrak{d} < \text{cf}(\mathfrak{c}) = \mathfrak{u} = \mathfrak{c}$. It remains to show that $\mathfrak{c} = \mathfrak{c}^\mathfrak{d}$ in $V[G]$.

Denote $\kappa = |M(\aleph_2)|$, $\lambda = \aleph_1$ and $\delta = \kappa^\lambda$. Since GCH holds in V , we have $\aleph_2 = 2^{\aleph_1}$, and consequently

$$\kappa = |M(\aleph_2)| = |[\aleph_2]^{< \aleph_0}|^{\aleph_0} = \aleph_2^{\aleph_0} = (2^{\aleph_1})^{\aleph_0} = 2^{\aleph_1} = \aleph_2$$

and

$$\delta = \aleph_2^{\aleph_1} = (2^{\aleph_1})^{\aleph_1} = \aleph_2.$$

Hence, κ, λ, δ are infinite cardinals. As $M(\aleph_2)$ is ccc, using [25, Lemma IV.3.11 at p. 267] we obtain in $V[G]$ the inequality $2^\lambda \leq \delta$ which yields:

$$\mathfrak{c}^\mathfrak{d} = (2^{\aleph_0})^\mathfrak{d} = 2^\mathfrak{d} = 2^{\aleph_1} = 2^\lambda \leq \delta = \aleph_2 = \mathfrak{c}.$$

Since $\mathfrak{c} \leq \mathfrak{c}^\mathfrak{d}$, we obtain $\mathfrak{c} = \mathfrak{c}^\mathfrak{d}$, and the proof is finished. \square

5. DOMINATING NUMBER AND ANALYTIC IDEALS

In this Section we show that $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$ for every analytic ideal \mathcal{I} . We will apply ideas developed by Katětov in [22, 24, 23] and by Debs and Saint Raymond in [15]. We start with recalling several technical notions.

If \mathcal{I} and \mathcal{J} are ideals on X and Y , respectively, then we say that \mathcal{I} is below \mathcal{J} in the Katětov order (in short: $\mathcal{I} \leq_K \mathcal{J}$), if there is a function $f : Y \rightarrow X$ such that $f^{-1}[A] \in \mathcal{J}$ for every $A \in \mathcal{I}$. If we can find a bijection with the above property, then we say that \mathcal{J} contains an isomorphic copy of \mathcal{I} and write $\mathcal{I} \sqsubseteq \mathcal{J}$.

If $\{X_t : t \in T\}$ is a family of sets, then $\sum_{t \in T} X_t = \{(t, x) : t \in T, x \in X_t\}$ is its disjoint sum. The vertical section of a set $A \subseteq \sum_{t \in T} X_t$ at a point $t \in T$ is defined by $A_{(t)} = \{x \in X_t : (t, x) \in A\}$.

Let \mathcal{I} be an ideal on a countable set T and $\{\mathcal{I}_t : t \in T\}$ be a family of ideals. Following [22], we define the ideal:

$$\mathcal{I} - \sum_{t \in T} \mathcal{I}_t = \{A \subseteq \sum_{t \in T} \left(\bigcup \mathcal{I}_t \right) : \{t \in T : A_{(t)} \notin \mathcal{I}_t\} \in \mathcal{I}\}.$$

In particular, if \mathcal{J} is some ideal on S and $\mathcal{I}_t = \mathcal{J}$ for all $t \in T$, then we denote:

$$\mathcal{I} \otimes \mathcal{J} = \mathcal{I} - \sum_{t \in T} \mathcal{I}_t = \{A \subseteq T \times S : \{t \in T : A_{(t)} \notin \mathcal{J}\} \in \mathcal{I}\}.$$

Following [23, 24], let $\{\text{Fin}^\alpha : 1 \leq \alpha < \omega_1\}$ be the family of ideals given by:

$$\text{Fin}^\alpha = \begin{cases} \text{Fin}, & \text{if } \alpha = 1, \\ \text{Fin} \otimes \text{Fin}^\beta, & \text{if } \alpha = \beta + 1, \\ \mathcal{I}_\alpha - \sum_{\beta < \alpha} \text{Fin}^\beta, & \text{if } \alpha \text{ is a limit ordinal,} \end{cases}$$

where \mathcal{I}_α is the ideal on the ordinal α generated by the family $\{\beta : \beta < \alpha\}$ (in particular, $\mathcal{I}_\omega = \text{Fin}$).

Finally, for ideals \mathcal{I} , \mathcal{J} and \mathcal{K} we define:

$$\mathfrak{b}(\mathcal{I}, \mathcal{J}, \mathcal{K}) = \min \left\{ |\mathcal{E}| : \mathcal{E} \subseteq \widehat{\mathcal{P}}_{\mathcal{K}} \wedge \forall \langle A_n \rangle \in \mathcal{P}_{\mathcal{J}} \exists \langle E_n \rangle \in \mathcal{E} \bigcup_{n \in \omega} \left(A_n \cap \bigcup_{i \leq n} E_i \right) \notin \mathcal{I} \right\}$$

(the above is an equivalent form of the coefficient defined in [34] – the equivalence is shown in [17, Proposition 3.9]). Note that $\mathfrak{b}(\mathcal{I}, \mathcal{I}, \mathcal{I}) = \mathfrak{b}(\mathcal{I})$. Moreover, it is easy to see that if $\mathcal{K} \subseteq \mathcal{K}'$, then $\mathfrak{b}(\mathcal{I}, \mathcal{J}, \mathcal{K}) \geq \mathfrak{b}(\mathcal{I}, \mathcal{J}, \mathcal{K}')$ (see [17, Proposition 3.8]).

We are ready to prove some lemmas, which will imply that $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$ for every analytic ideal \mathcal{I} .

Lemma 5.1. *If $1 \leq \alpha < \omega_1$ is an ordinal, $\text{Fin}^{\alpha+1} \not\sqsubseteq \mathcal{I}$, but $\text{Fin}^\alpha \sqsubseteq \mathcal{I}$, then $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$.*

Proof. If $\alpha = 1$ then this is true by [17, Theorem 4.2(4) and Theorem 4.2(9)].

Let now $1 < \alpha < \omega_1$ be any ordinal. We need to standardize the notation so that we will be able to proceed with the proof for both successor and limit ordinals α .

If $\alpha = \beta + 1$ is a successor ordinal, put $\gamma = \omega$ and write $\beta_\delta = \beta$ for all $\delta < \gamma$. Note that $\text{Fin}^\alpha = \text{Fin} \otimes \text{Fin}^\beta = \mathcal{I}_\gamma - \sum_{\delta < \gamma} \text{Fin}^{\beta_\delta}$.

On the other hand, if α is a limit ordinal, put $\gamma = \alpha$ and write $\beta_\delta = \delta$ for all $\delta < \gamma$. Note that $\text{Fin}^\alpha = \mathcal{I}_\alpha - \sum_{\delta < \gamma} \text{Fin}^\delta = \mathcal{I}_\gamma - \sum_{\delta < \gamma} \text{Fin}^{\beta_\delta}$.

Now we can proceed with the proof not caring if α is a successor or a limit ordinal.

For each $\delta < \gamma$ let $\text{Fin}^{\beta_\delta}(\omega)$ be any isomorphic copy of $\text{Fin}^{\beta_\delta}$ on ω . Then Fin^α and $\mathcal{J} = \mathcal{I}_\gamma - \sum_{\delta < \gamma} \text{Fin}^{\beta_\delta}(\omega)$ are isomorphic, so $\mathcal{J} \sqsubseteq \mathcal{I}$. Thus, without loss of generality we may assume that \mathcal{I} is an ideal on $\gamma \times \omega$ such that $\mathcal{J} \sqsubseteq \mathcal{I}$ (this can be done by considering the ideal $\{f[A] : A \in \mathcal{I}\} \supseteq \mathcal{J}$, instead of \mathcal{I} , where f is the bijection witnessing $\mathcal{J} \sqsubseteq \mathcal{I}$).

We will show that $\mathfrak{b}(\mathcal{I}, \mathcal{I}, \mathcal{J}) \leq \mathfrak{d}$. This will finish the proof as $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{b}(\mathcal{I}, \mathcal{I}, \mathcal{J})$ (by $\mathcal{J} \sqsubseteq \mathcal{I}$).

Fix a strictly increasing sequence $\langle \gamma_i : i \in \omega \rangle \in \gamma^\omega$ converging to γ (in the order topology) such that $\gamma_0 = 0$. Let $\{f_\alpha : \alpha < \mathfrak{d}\} \subseteq \omega^\omega$ be a \leq_{Fin} -dominating family. Without loss of generality we may assume that each f_α is strictly increasing and satisfies $f_\alpha(0) = 0$. Define $A_n^\alpha = (\gamma_{f_\alpha(n+1)} \setminus \gamma_{f_\alpha(n)}) \times \omega$ for all $\alpha < \mathfrak{d}$ and $n \in \omega$. Then $\{\langle A_n^\alpha : n \in \omega \rangle : \alpha < \mathfrak{d}\} \subseteq \widehat{\mathcal{P}}_{\mathcal{J}}$. We claim that this family witnesses $\mathfrak{b}(\mathcal{I}, \mathcal{I}, \mathcal{J}) \leq \mathfrak{d}$.

Fix any $\langle B_n : n \in \omega \rangle \in \mathcal{P}_{\mathcal{I}}$. There are two possibilities.

Assume first that there is $X \notin \mathcal{I}$ such that $(X \cap B_n)_{(\delta)} \in \text{Fin}^{\beta_\delta}(\omega)$, for all $n \in \omega$ and $\delta < \gamma$. For $\delta < \gamma$ let $m_\delta \in \omega$ be such that $\gamma_{f_0(m_\delta)} \leq \delta < \gamma_{f_0(m_\delta+1)}$ (i.e.,

$\{\delta\} \times \omega \subseteq A_{m_\delta}^0$). Observe that:

$$\begin{aligned} \left(\bigcup_{i \in \omega} \left(A_i^0 \setminus \bigcup_{n < i} B_n \right) \right)_{(\delta)} &\supseteq \left(A_{m_\delta}^0 \setminus \bigcup_{n < m_\delta} B_n \right)_{(\delta)} \\ &= \omega \setminus \bigcup_{n < m_\delta} (B_n)_{(\delta)} \\ &\supseteq X_{(\delta)} \setminus \bigcup_{n < m_\delta} (X \cap B_n)_{(\delta)}. \end{aligned}$$

Define $Y_\delta = \bigcup_{n < m_\delta} (X \cap B_n)_{(\delta)} \in \text{Fin}^{\beta_\delta}(\omega)$ and $Y = \bigcup_{\delta < \gamma} \{\delta\} \times Y_\delta \in \mathcal{J} \subseteq \mathcal{I}$. Then we have:

$$\mathcal{I} \not\# X \setminus Y \subseteq \bigcup_{i \in \omega} \left(A_i^0 \setminus \bigcup_{n < i} B_n \right) \subseteq \bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} A_i^0 \right)$$

(the last inclusion follows from Remark 2.1).

Assume now that if $X \subseteq \gamma \times \omega$ satisfies $(X \cap B_n)_{(\delta)} \in \text{Fin}^{\beta_\delta}(\omega)$ for all $n \in \omega$ and $\delta < \gamma$ then $X \in \mathcal{I}$. Recall that the ideal $\text{Fin} \otimes \mathcal{J}$ (which is isomorphic to $\text{Fin}^{\alpha+1}$) has three kinds of generators:

- sets of the form $\{n\} \times \gamma \times \omega$, for $n \in \omega$ (*generators of the first type*);
- sets of the form $\bigcup_{n \in \omega} \{n\} \times \gamma_{g(n)} \times \omega$, for $g \in \omega^\omega$ (*generators of the second type*);
- sets $G \subseteq \omega \times \gamma \times \omega$ such that $G_{(n,\delta)} \in \text{Fin}^{\beta_\delta}(\omega)$, for all $(n,\delta) \in \omega \times \gamma$ (*generators of the third type*).

Consider the function $h : \gamma \times \omega \rightarrow \omega \times \gamma \times \omega$ defined by the formula $h(\delta, j) = (n(\delta, j), \delta, j)$, where $n(\delta, j) \in \omega$ is given by $(\delta, j) \in B_{n(\delta, j)}$. Observe that:

- if $G = \{n\} \times \gamma \times \omega$ is a generator of the first type, then $h^{-1}[G] \subseteq B_n \in \mathcal{I}$, for every $n \in \omega$;
- if $G \subseteq \omega \times \gamma \times \omega$ is a generator of the third type, then $(h^{-1}[G] \cap B_n)_{(\delta)} \subseteq G_{(n,\delta)} \in \text{Fin}^{\beta_\delta}(\omega)$, for all $(n,\delta) \in \omega \times \gamma$, so $h^{-1}[G] \in \mathcal{I}$ by our assumption.

On the other hand, $\text{Fin}^{\alpha+1} \not\subseteq_K \mathcal{I}$ (by $\text{Fin}^{\alpha+1} \not\subseteq \mathcal{I}$ and [1, Example 4.1]), so there has to be a generator of the second type, $G = \bigcup_{n \in \omega} \{n\} \times \gamma_{g(n)} \times \omega$ for some $g \in \omega^\omega$, such that:

$$\mathcal{I} \not\# h^{-1}[G] \subseteq \bigcup_{n \in \omega} (B_n \cap (\gamma_{g(n)} \times \omega)).$$

Find $\alpha < \mathfrak{d}$ with $g \leq^* f_\alpha$ and let $m \in \omega$ be such that $g(n) \leq f_\alpha(n)$ for all $n > m$. Note that:

$$\begin{aligned} \bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} A_i^\alpha \right) &\supseteq \bigcup_{n \in \omega} (B_n \cap (\gamma_{f_\alpha(n)} \times \omega)) \\ &\supseteq \bigcup_{n > m} (B_n \cap (\gamma_{g(n)} \times \omega)) \supseteq h^{-1}[G] \setminus \bigcup_{n \leq m} B_n \notin \mathcal{I} \end{aligned}$$

(as $\bigcup_{n \leq m} B_n \in \mathcal{I}$). This finishes the proof. \square

Lemma 5.2. *For each $1 < \alpha < \omega_1$ there is a family $\{A_f : f \in \omega^\omega\} \subseteq \text{Fin}^\alpha$ satisfying:*

- for every $A \in \text{Fin}^\alpha$ there is $f \in \omega^\omega$ with $A \subseteq A_f$;

- $A_f \subseteq A_g$ whenever $f \leq g$ (i.e., $f(n) \leq g(n)$ for all $n \in \omega$).

Proof. We will show it inductively.

This is clear for $\alpha = 2$ as witnessed by the family:

$$\left\{ (f(0) \times \omega) \cup \bigcup_{i \in \omega} \{i\} \times f(i) : f \in \omega^\omega \right\} \subseteq \text{Fin}^2.$$

Fix now any partition $\langle C_n : n \in \omega \rangle$ of ω into infinite sets and for each n let $h_n : \omega \rightarrow C_n$ be the increasing enumeration of C_n .

Assume that for some $1 < \alpha < \omega_1$ we have the needed family $\{A_f : f \in \omega^\omega\} \subseteq \text{Fin}^\alpha$. We need to find the required family for $\alpha + 1$. Given $f \in \omega^\omega$, define:

$$B_f = (f(0) \times \bigcup \text{Fin}^\alpha) \cup \bigcup_{n \in \omega} \{n\} \times A_{f \circ h_n}.$$

Again it is clear that $\{B_f : f \in \omega^\omega\} \subseteq \text{Fin}^{\alpha+1}$ is as needed.

Finally, assume that $1 < \alpha < \omega_1$ is a limit ordinal and for each $\beta < \alpha$ we have the required family $\{A_f^\beta : f \in \omega^\omega\} \subseteq \text{Fin}^\beta$. Find a bijection $g : \omega \rightarrow \alpha$ and a strictly increasing sequence $\langle \beta_i : i \in \omega \rangle$ converging to α (in the order topology). For each $f \in \omega^\omega$, define:

$$B_f = \left(\sum_{\beta < \beta_f(0)} \bigcup \text{Fin}^\beta \right) \cup \left(\bigcup_{n \in \omega} \{g(n)\} \times A_{f \circ h_n}^{g(n)} \right).$$

Once more it is clear that $\{B_f : f \in \omega^\omega\} \subseteq \text{Fin}^\alpha$ is as needed. □

Lemma 5.3. *If $\alpha < \omega_1$ is a limit ordinal, $\text{Fin}^\alpha \not\sqsubseteq \mathcal{I}$, but $\text{Fin}^\beta \sqsubseteq \mathcal{I}$ for all $\beta < \alpha$, then $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$.*

Proof. Find an increasing sequence $\langle \beta_i : i \in \omega \rangle$ converging to α (in the order topology). Since $\text{Fin}^\beta \sqsubseteq \mathcal{I}$ for all $\beta < \alpha$, we can assume that \mathcal{I} is an ideal on ω and $\text{Fin}^{\beta_i}(\omega) \subseteq \mathcal{I}$ for all i , where $\text{Fin}^{\beta_i}(\omega)$ is some copy of Fin^{β_i} on ω . For each $i \in \omega$ let $\{B_f^i : f \in \omega^\omega\} \subseteq \text{Fin}^{\beta_i}(\omega)$ be the family from Lemma 5.2.

Fix any partition $\langle C_n : n \in \omega \rangle$ of ω into infinite sets and for each n let $h_n : \omega \rightarrow C_n$ be the increasing enumeration of C_n . Let $\{f_\alpha : \alpha < \mathfrak{d}\} \subseteq \omega^\omega$ be a \leq_{Fin} -dominating family of strictly increasing functions and recursively define $A_n^\alpha = B_{f_\alpha \circ h_n}^n \setminus \bigcup_{i < n} A_i^\alpha$, for all $\alpha < \mathfrak{d}$ and $n \in \omega$. Then $\{\langle A_n^\alpha : n \in \omega \rangle : \alpha < \mathfrak{d}\} \subseteq \widehat{\mathcal{P}}_{\mathcal{I}}$ as $\text{Fin}^{\beta_i}(\omega) \subseteq \mathcal{I}$ for all i . We will show that $\{A_n^\alpha : n \in \omega, \alpha < \mathfrak{d}\}$ witnesses $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$.

Fix any $\langle B_n : n \in \omega \rangle \in \mathcal{P}_{\mathcal{I}}$ and consider the function $h : \omega \rightarrow \alpha \times \omega$ defined by $h(i) = (\beta_n, i)$, where $n \in \omega$ is given by $i \in B_n$. Since $\mathcal{I}_\alpha - \sum_{\beta < \alpha} \text{Fin}^\beta(\omega)$ is isomorphic to Fin^α and $\text{Fin}^\alpha \not\leq_K \mathcal{I}$ (by $\text{Fin}^\alpha \not\sqsubseteq \mathcal{I}$ and [1, Example 4.1]), there is $Y \in \mathcal{I}_\alpha - \sum_{\beta < \alpha} \text{Fin}^\beta(\omega)$ such that $h^{-1}[Y] \notin \mathcal{I}$. Actually, for each $\beta < \alpha$ either $h^{-1}[\{\beta\} \times \omega] = \emptyset \in \mathcal{I}$ (if $\beta \neq \beta_i$ for all $i \in \omega$) or $h^{-1}[\{\beta\} \times \omega] \subseteq B_n \in \mathcal{I}$ for some $n \in \omega$ (if $\beta = \beta_n$). Hence, the set:

$$X = h^{-1} \left[\bigcup \left\{ \{\delta\} \times Y_{(\delta)} : Y_{(\delta)} \in \text{Fin}^\delta(\omega) \right\} \right]$$

does not belong to \mathcal{I} , while $h[X] \subseteq h[h^{-1}[Y]] \subseteq Y \in \mathcal{I}_\alpha - \sum_{\beta < \alpha} \text{Fin}^\beta(\omega)$. Observe that $X \cap B_n \subseteq (h[X])_{(\beta_n)} \in \text{Fin}^{\beta_n}$ for all $n \in \omega$.

Then for each n there is $g_n \in \omega^\omega$ such that $X \cap B_n \subseteq B_{g_n}^n$. Define $g \in \omega^\omega$ by $g \upharpoonright C_n = g_n \circ h_n^{-1}$. Find $\alpha < \mathfrak{d}$ with $g \leq^* f_\alpha$ and let $m \in \omega$ be such that $g(i) \leq f_\alpha(i)$

whenever $i \geq m$. Let $k \in \omega$ be maximal such that $C_k \cap m \neq \emptyset$. Observe that for $n > k$ and any $j \in \omega$ we have:

$$(f_\alpha \circ h_n)(j) = f_\alpha(h_n(j)) \geq g(h_n(j)) = g_n(j).$$

Hence, $X \cap B_n \subseteq B_{g_n}^n \subseteq B_{f_\alpha \circ h_n}^n \subseteq \bigcup_{i \leq n} A_i^\alpha$, whenever $n > k$. Thus,

$$\bigcup_{n \in \omega} \left(B_n \cap \bigcup_{i \leq n} A_i^\alpha \right) \supseteq \bigcup_{n > k} \left(B_n \cap \bigcup_{i \leq n} A_i^\alpha \right) \supseteq \bigcup_{n > k} (X \cap B_n) \notin \mathcal{I}$$

as $\mathcal{I} \not\supseteq X = \bigcup_{n \in \omega} (X \cap B_n)$ and $\bigcup_{n \leq k} B_n \in \mathcal{I}$. \square

Theorem 5.4. *If there is $\alpha < \omega_1$ such that $\text{Fin}^\alpha \not\sqsubseteq \mathcal{I}$, then $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$.*

Proof. Let $\alpha < \omega_1$ be minimal such that $\text{Fin}^\alpha \not\sqsubseteq \mathcal{I}$. If α is a successor ordinal, use Lemma 5.1. On the other hand, if α is a limit ordinal, we can apply Lemma 5.3. \square

Recall that for an ideal \mathcal{I} on X we write $\mathcal{I}^* = \{A \subseteq X : X \setminus A \in \mathcal{I}\}$ and call it the *dual filter of \mathcal{I}* .

Corollary 5.5. *If \mathcal{I} is Borel-separated from \mathcal{I}^* (i.e., there is a Borel set $S \subseteq 2^\omega$ such that $\mathcal{I} \subseteq S$ and $\mathcal{I}^* \cap S = \emptyset$), then $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$. In particular, $\mathfrak{b}(\mathcal{I}) \leq \mathfrak{d}$ for every analytic ideal \mathcal{I} .*

Proof. We claim that there is $\alpha < \omega_1$ such that $\text{Fin}^\alpha \not\sqsubseteq \mathcal{I}$. Indeed, by [15, Theorem 3.2, Theorem 6.5 and Lemma 7.2], if $\text{Fin}^\alpha \sqsubseteq \mathcal{I}$, then \mathcal{I} is not $\Delta_{1+\alpha}^0$ -separated from \mathcal{I}^* . Thus, the assumption that $\text{Fin}^\alpha \sqsubseteq \mathcal{I}$, for all $\alpha < \omega_1$, would contradict the fact that \mathcal{I} is Borel-separated from \mathcal{I}^* . Now it suffices to apply Theorem 5.4.

For each analytic ideal, its dual filter is also analytic (by [15, Subsection 1.1]), so they can be Borel-separated (by the Lusin separation theorem). This proves the "In particular" part. \square

Actually, we do not know whether Corollary 5.5 can be strengthened by replacing \mathfrak{d} with \mathfrak{b} . In particular, we do not know the answer to the following problem.

Problem 1. Does there consistently exist an analytic ideal \mathcal{I} with $\mathfrak{b}(\mathcal{I}) > \mathfrak{b}$?

REFERENCES

1. Paweł Barbarski, Rafał Filipów, Nikodem Mrozek, and Piotr Szuca, *When does the katětov order imply that one ideal extends the other?*, Colloq. Math. **130** (2013), 91–102.
2. Tomek Bartoszyński, *Invariants of measure and category*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 491–555. MR 2768686
3. Tomek Bartoszyński and Saharon Shelah, *Closed measure zero sets*, Ann. Pure Appl. Logic **58** (1992), no. 2, 93–110. MR 1186905
4. Andreas Blass, *Combinatorial cardinal characteristics of the continuum*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 395–489. MR 2768685
5. Lev Bukovský, *On wQN_* and wQN^* spaces*, Topology Appl. **156** (2008), no. 1, 24–27. MR 2463820
6. ———, *The structure of the real line*, Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) [Mathematics Institute of the Polish Academy of Sciences. Mathematical Monographs (New Series)], vol. 71, Birkhäuser/Springer Basel AG, Basel, 2011. MR 2778559
7. Lev Bukovský, Pratulananda Das, and Jaroslav Šupina, *Ideal quasi-normal convergence and related notions*, Colloq. Math. **146** (2017), no. 2, 265–281. MR 3622377

8. Lev Bukovský and Jozef Haleš, *QN-spaces, wQN-spaces and covering properties*, Topology Appl. **154** (2007), no. 4, 848–858. MR 2294632
9. Lev Bukovský, Ireneusz Reclaw, and Miroslav Repický, *Spaces not distinguishing pointwise and quasinormal convergence of real functions*, Topology Appl. **41** (1991), no. 1-2, 25–40. MR 1129696
10. ———, *Spaces not distinguishing convergences of real-valued functions*, Topology Appl. **112** (2001), no. 1, 13–40. MR 1815270
11. Michael Canjar, *Countable ultraproducts without CH^** , Ann. Pure Appl. Logic **37** (1988), no. 1, 1–79. MR 924678
12. R. Michael Canjar, *Cofinalities of countable ultraproducts: the existence theorem*, Notre Dame J. Formal Logic **30** (1989), no. 4, 539–542. MR 1036675
13. Robert Michael Canjar, *Model-theoretic properties of countable ultraproducts without the Continuum Hypothesis*, ProQuest LLC, Ann Arbor, MI, 1982, Thesis (Ph.D.)–University of Michigan. MR 2632174
14. Pratulananda Das and Debraj Chandra, *Spaces not distinguishing pointwise and \mathcal{I} -quasinormal convergence*, Comment. Math. Univ. Carolin. **54** (2013), no. 1, 83–96. MR 3038073
15. Gabriel Debs and Jean Saint Raymond, *Filter descriptive classes of Borel functions*, Fund. Math. **204** (2009), no. 3, 189–213. MR 2520152
16. Ilijas Farah and Sławomir Solecki, *Two $F_{\sigma\delta}$ ideals*, Proc. Amer. Math. Soc. **131** (2003), no. 6, 1971–1975. MR 1955288
17. Rafał Filipów and Adam Kwela, *Yet another ideal version of the bounding number*, J. Symb. Log. **87** (2022), no. 3, 1065–1092. MR 4472525
18. Rafał Filipów and Marcin Staniszewski, *Pointwise versus equal (quasi-normal) convergence via ideals*, J. Math. Anal. Appl. **422** (2015), no. 2, 995–1006. MR 3269495
19. D. H. Fremlin, *Consequences of Martin's axiom*, Cambridge Tracts in Mathematics, vol. 84, Cambridge University Press, Cambridge, 1984. MR 780933
20. Michael Hrušák, *Combinatorics of filters and ideals*, Set theory and its applications, Contemp. Math., vol. 533, Amer. Math. Soc., Providence, RI, 2011, pp. 29–69. MR 2777744
21. Teppo Kankaanpää, *Remarks on gaps in $Dense(\mathbb{Q})/nwd$* , MLQ Math. Log. Q. **59** (2013), no. 1-2, 51–61. MR 3032424
22. Miroslav Katětov, *Products of filters*, Comment. Math. Univ. Carolinae **9** (1968), 173–189. MR 250257
23. ———, *On descriptive classes of functions*, Theory of sets and topology (in honour of Felix Hausdorff, 1868–1942), VEB Deutsch. Verlag Wissensch., Berlin, 1972, pp. 265–278. MR 0345060
24. ———, *On descriptive classification of functions*, General topology and its relations to modern analysis and algebra, III (Proc. Third Prague Topological Sympos., 1971), Academia, Prague, 1972, pp. 235–242. MR 0355956
25. Kenneth Kunen, *Set theory*, Studies in Logic (London), vol. 34, College Publications, London, 2011. MR 2905394
26. Adam Kwela, *Ideal weak QN-spaces*, Topology Appl. **240** (2018), 98–115. MR 3784399
27. Adam Kwela and Marcin Sabok, *Topological representations*, J. Math. Anal. Appl. **422** (2015), no. 2, 1434–1446. MR 3269521
28. Ireneusz Reclaw, *Metric spaces not distinguishing pointwise and quasinormal convergence of real functions*, Bull. Polish Acad. Sci. Math. **45** (1997), no. 3, 287–289. MR 1477547
29. Miroslav Repický, *Spaces not distinguishing convergences*, Comment. Math. Univ. Carolin. **41** (2000), no. 4, 829–842. MR 1800160
30. ———, *Spaces not distinguishing ideal convergences of real-valued functions*, Real Anal. Exchange **46** (2021), no. 2, 367–394. MR 4336563
31. ———, *Spaces not distinguishing ideal convergences of real-valued functions, II*, Real Anal. Exchange **46** (2021), no. 2, 395–421. MR 4336564
32. Marcin Sabok and Jindřich Zapletal, *Forcing properties of ideals of closed sets*, Journal of Symbolic Logic **76** (2010), 4011–4039.
33. Masami Sakai, *The sequence selection properties of $C_p(X)$* , Topology Appl. **154** (2007), no. 3, 552–560. MR 2280899
34. Marcin Staniszewski, *On ideal equal convergence II*, J. Math. Anal. Appl. **451** (2017), no. 2, 1179–1197. MR 3624786

35. Boaz Tsaban and Lyubomyr Zdomskyy, *Hereditarily Hurewicz spaces and Arhangel'skiĭ sheaf amalgamations*, J. Eur. Math. Soc. (JEMS) **14** (2012), no. 2, 353–372. MR 2881299
36. Viera Šottová and Jaroslav Šupina, *Principle $S_1(\mathcal{P}, \mathcal{R})$: ideals and functions*, Topology Appl. **258** (2019), 282–304. MR 3924519
37. Jaroslav Šupina, *Ideal QN -spaces*, J. Math. Anal. Appl. **435** (2016), no. 1, 477–491. MR 3423409

(Adam Kwela) INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS, UNIVERSITY OF GDAŃSK, UL. WITA STWOSZA 57, 80-308 GDAŃSK, POLAND

Email address: `Adam.Kwela@ug.edu.pl`

URL: `https://mat.ug.edu.pl/~akwela`