

Weak recovery, hypothesis testing, and mutual information in stochastic block models and planted factor graphs

Elchanan Mossel*

Allan Sly†

Youngtak Sohn ‡

June 25, 2024

Abstract

The stochastic block model is a canonical model of communities in random graphs. It was introduced in the social sciences and statistics as a model of communities, and in theoretical computer science as an average case model for graph partitioning problems under the name of the “planted partition model.” Given a sparse stochastic block model, the two standard inference tasks are: (i) Weak recovery: can we estimate the communities with non trivial overlap with the true communities? (ii) Detection/Hypothesis testing: can we distinguish if the sample was drawn from the block model or from a random graph with no community structure with probability tending to 1 as the graph size tends to infinity?

In this work, we show that for sparse stochastic block models, the two inference tasks are equivalent except at a critical point. That is, weak recovery is information theoretically possible if and only if detection is possible. We thus find a strong connection between these two notions of inference for the model. We further prove that when detection is impossible, an explicit hypothesis test based on low degree polynomials in the adjacency matrix of the observed graph achieves the optimal statistical power. This low degree test is efficient as opposed to the likelihood ratio test, which is not known to be efficient. Moreover, we prove that the asymptotic mutual information between the observed network and the community structure exhibits a phase transition at the weak recovery threshold.

Our results are proven in much broader settings including the hypergraph stochastic block models and general planted factor graphs. In these settings we prove that the impossibility of weak recovery implies contiguity and provide a condition which guarantees the equivalence of weak recovery and detection.

1 Introduction

The stochastic block model, or simply a block model, is a random graph model generalizing the famous Erdos-Renyi random graph [ER+60] and is a special case of inhomogeneous random graphs [BJR07]. It has been studied extensively in statistics and the social sciences as a model of communities [HLL83, SN97, BC09, RCY11] and in computer science as a model to study the average case behavior of clustering algorithms [DF89, JS98, CK01, McS01, CO10]. In the last decade starting with the work of Decelle et al. [DKMZ11], the sparse block model has been extensively studied with fascinating connections to belief propagation and non-backtracking random

*Department of Mathematics, Massachusetts Institute of Technology. Email: elmos@mit.edu

†Department of Mathematics, Princeton University. Email: asly@math.princeton.edu

‡Department of Mathematics, Massachusetts Institute of Technology. Email: youngtak@mit.edu

walks [KMM⁺13, BLM15]. Furthermore, the block model is one of the canonical models in high-dimensional statistics which is believed to exhibit a statistical-computational gap, meaning a gap between what is achievable information theoretically and what is achievable with known computationally efficient algorithms.

The two central inference tasks related to the sparse block models are *weak recovery* and *detection*. Weak recovery means estimating communities with non trivial overlap with the true communities. Detection refers to a statistical test that achieves vanishing Type 1 and Type 2 errors from a single sample in determining if it is drawn from the block model or from the “null model” with no community structure. Here, a null model refers to a standard random graph model with the same average degree. A number of papers including [MNS15, BMNN16, COKPZ18, COEJ⁺18, COKM20] have studied the information theoretic thresholds for both weak recovery and detection. In particular it was shown that for the two community case [MNS15] and for the disassortative case [COEJ⁺18] the two thresholds coincide. It is thus natural to ask

Is weak recovery equivalent to the detection in general?

The first objective of this paper is to answer this question for sparse block models, and in greater generality, for *planted factor models* [COKPZ18] which encompasses labeled/weighted stochastic block models [SH12, LMX15, XJL20] and hypergraph stochastic block models [ACKZ15, GD17a, GD17b, PZ21, SZ22]. We prove that the information theoretic possibility of detection implies the possibility of weak recovery, and find a condition under which the converse holds. The second objective of this paper is to study *hypothesis testing*, which is a more classical task than detection as it involves testing between two simple hypothesis. Notably, the impossibility of detection merely implies that there is no sequence of statistical tests that distinguishes the block model from the null model with vanishing Type 1 and Type 2 errors. Then, we are led to the question

Can we characterize the asymptotic power of the optimal (likelihood ratio) test at significance level $\alpha \in (0, 1)$? If so, can the optimal power be achieved efficiently?

In this paper, we explicitly characterize the asymptotic power of the likelihood ratio test. Moreover, we prove that it can be achieved efficiently by a test based on low-degree polynomials of the adjacency matrix when the weak recovery, or equivalently detection, is impossible. We also investigate the relationship between these statistical tasks and the mutual information between the observed graph and the community structure.

1.1 Our results for symmetric block models

To illustrate our general results, we first consider the well-studied setup of sparse symmetric block models with $q \geq 2$ communities defined as follows. Let V be a vertex set of size $n \geq 1$. For parameters $a, b > 0$, let $\mathbf{G}^* \sim \mathcal{G}(n, q, \frac{a}{n}, \frac{b}{n})$ denote the model of random graphs in which each vertex $u \in V$ is assigned a label $\sigma_u^* \in \{1, 2, \dots, q\}$ uniformly at random, and then each possible edge (u, v) is included with probability $\frac{a}{n}$ if $\sigma_u^* = \sigma_v^*$ and with probability $\frac{b}{n}$ if $\sigma_u^* \neq \sigma_v^*$. Then, the average degree of \mathbf{G}^* is given by $d \equiv \frac{a+(q-1)b}{q}$. The corresponding random graph with no community structure is $\mathbf{G} \sim \mathcal{G}_{\text{ER}}(n, \frac{d}{n})$, where $\mathcal{G}_{\text{ER}}(n, \frac{d}{n}) \equiv \mathcal{G}(n, q, \frac{d}{n}, \frac{d}{n})$ is the sparse Erdos-Renyi graph with average degree d . Observe that the parameters (a, b) is in one-to-one correspondence with the average degree d and the parameter

$$\lambda \equiv \frac{a-b}{a+(q-1)b} \in \left(-\frac{1}{q-1}, 1 \right).$$

Here, λ corresponds to the second eigenvalue of a certain stochastic matrix, namely the transition matrix from the color of a vertex to its neighbor or child (see Eq. (1)). This parametrization is emphasized in several works on the block model [Mas14, MNS18, BLM15, AS15, AS18] and the phylogeny reconstruction problem [Mos04, DMR11, MRS11, RS17]. We are interested in high-dimensional setting, where $n \rightarrow \infty$ as q, d, λ are fixed.

A first observation is that by fixing $\lambda \in (-\frac{1}{q-1}, 1)$ and increasing d from 0 to ∞ , the task of weak recovery is monotone in d since larger d amounts to more information. Indeed given a sample from the model with a certain d by deleting each edge independently with probability $t \in [0, 1]$ we obtain a perfect sample of the model with the same λ and the new degree $t \cdot d$. Thus, there exists a critical point $d_*(q, \lambda)$, which we call the *weak recovery threshold*, at which the weak recovery becomes information theoretically possible (see Definition 3.3). Our result for the block model shows that the weak recovery and detection is equivalent possibly except for the critical point $d_*(q, \lambda)$.

Theorem 1.1. (Special case of Theorem 3.13) *Consider the sparse symmetric block model $\mathbf{G}^* \sim \mathcal{G}(n, q, \frac{a}{n}, \frac{b}{n})$ with $q \geq 2$ communities and parameters $a, b > 0$. For average degree $d \equiv \frac{a+(q-1)b}{q}$ below the weak recovery threshold $d < d_*(q, \lambda)$, the following holds.*

- (1) $\mathbf{G}^* \sim \mathcal{G}(n, q, \frac{a}{n}, \frac{b}{n})$ is mutually contiguous with the Erdos-Renyi graph $\mathbf{G} \sim \mathcal{G}_{\text{ER}}(n, \frac{d}{n})$. That is, for a sequence of events $(\mathcal{A}_n)_{n \geq 1}$, $\mathbb{P}(\mathbf{G}^* \in \mathcal{A}_n) \rightarrow 0$ as $n \rightarrow \infty$ if and only if $\mathbb{P}(\mathbf{G} \in \mathcal{A}_n) \rightarrow 0$ as $n \rightarrow \infty$. Thus, detection is impossible, i.e. there is no statistical test that distinguishes \mathbf{G}^* from \mathbf{G} with power tending to 1 as $n \rightarrow \infty$. Moreover, there is no consistent estimator for $a, b > 0$.
- (2) The mutual information between \mathbf{G}^* and the community structure $\underline{\sigma}^*$ per-vertex $I(\mathbf{G}^*, \underline{\sigma}^*)/n$ converges to $\frac{d}{2}I_0(q, \lambda)$ as $n \rightarrow \infty$, where $I_0(q, \lambda)$ is the mutual information of $\sigma_\lambda \in [q]$ and $\sigma \in [q]$ defined in Eq. (1) below.

On the contrary, above the weak recovery threshold $d > d_*(q, \lambda)$,

- (3) $\mathbf{G}^* \sim \mathcal{G}(n, q, \frac{a}{n}, \frac{b}{n})$ is not mutually contiguous to the Erdos-Renyi graph $\mathbf{G} \sim \mathcal{G}_{\text{ER}}(n, \frac{d}{n})$. Moreover, there exists a sequence of events $(\mathcal{A}_n^*)_{n \geq 1}$ such that after passing to a subsequence, $\mathbb{P}(\mathbf{G} \in \mathcal{A}_n^*) \leq 2e^{-\eta n}$ and $\mathbb{P}(\mathbf{G}^* \in \mathcal{A}_n^*) \geq 1 - 2e^{-\eta n}$ hold for $\eta > 0$ not depending on n . Thus, detection is possible, and along a subsequence, there exists a statistical test that distinguishes \mathbf{G}^* from \mathbf{G} with power tending to 1 exponentially fast as $n \rightarrow \infty$.
- (4) $\liminf_{n \rightarrow \infty} \frac{1}{n}I(\mathbf{G}^*, \underline{\sigma}^*) < \frac{d}{2}I_0(q, \lambda)$ holds.

The results regarding the mutual contiguity in Theorem 1.1-(1), (3) were previously known for various regimes of parameter: the disassortative case where $a < b$ follows from combining results of a sequence of papers [COKPZ18, COEJ+18, COKM20], the $q = 2$ case follows by [MNS15], and if the average degree d is below the so-called ‘second moment threshold’, Theorem 1.1-(1) follows from [BMNN16] (see Section 2 for further literature). However, both the proofs and the results of all of the previous works have different sets of limitations, which relied on either the ‘interpolation technique’ failing for assortative case $a > b$ or the ‘second moment method’ failing for d slightly below $d_*(q, \lambda)$. Our Theorem 1.1-(1), (3) unifies and extends these results and covers the maximum range of parameters. Furthermore, our results apply in a greater generality to hypergraph stochastic block model, where we allow $k \geq 2$ order interaction of communities and non-symmetric connection probabilities between different communities. We refer to Section 3 for our most general results in the planted factor models.

The proof of Theorem 1.1 is based on a novel equivalence between the impossibility of weak recovery and the *near-orthogonality* of the samples drawn from the posterior (see Theorem 3.10). In particular, the new key insight of the proof of Theorem 1.1-(1) is that below the weak recovery threshold $d < d_*(q, \lambda)$, a suitable truncation of the likelihood ratio function based on *overlaps* of the samples drawn from the posterior reduces the second moment by a multiplicative $e^{\Omega(n)}$ factor while only affecting the first moment by $1 - o(1)$. We combine this insight with other techniques such as small subgraph conditioning from combinatorics [Jan95, Wor99] and I-MMSE relation from information theory [GWSV11]. We refer to Section 4 for a proof overview.

Note that Theorem 1.1 does not cover the critical case $d = d_*(q, \lambda)$. This is not a coincidence: there are cases where the equivalence of the detection and weak recovery fails for $d = d_*(q, \lambda)$! For example, it is known that the weak recovery threshold $d_*(q, \lambda)$ matches the *Kesten-Stigum threshold* $d_{KS} = \lambda^{-2}$ when $q = 2$ [MNS15] or when $q \in \{3, 4\}$ and d is large enough [MSS23]. In these cases, the weak recovery is impossible at the Kesten-Stigum threshold [MNS15, MSS23]. In contrast, it is possible to distinguish the block model from the Erdos Renyi graph with probability approaching 1 by counting short cycles as the asymptotic distribution of the number of cycles of fixed length is Poisson with known parameters [BJR07].

The quantity $I_0(q, \lambda)$ in Theorem 1.1 can be characterized as follows. Let $\sigma \in \text{Unif}(\{1, \dots, q\})$ be a uniform community assignment. Given σ , let σ_λ a sample from the conditional distribution

$$\mathbb{P}(\sigma_\lambda = j \mid \sigma = i) = \begin{cases} \lambda + \frac{1-\lambda}{q} & \text{if } i = j \\ \frac{1-\lambda}{q} & \text{otherwise.} \end{cases} \quad (1)$$

Then, $I_0(q, \lambda) \equiv I(\sigma_\lambda, \sigma)$ is the mutual information between σ_λ and σ :

$$I_0(q, \lambda) \equiv I(\sigma_\lambda, \sigma) = \frac{1 + (q-1)\lambda}{q} \log(1 + (q-1)\lambda) + \frac{(q-1)(1-\lambda)}{q} \log(1-\lambda), \quad (2)$$

Thus, Theorem 1.1-(2) establishes that for all $q \geq 2$ and $\lambda \in (-\frac{1}{q-1}, 1)$, below the weak recovery threshold $d < d_*(q, \lambda)$, the asymptotic per-vertex mutual information between the observed network and the community structure has a “single-letter” characterization. Informally, the mutual information can all be explained by local interactions between neighbors. On the other hand, when $d > d_*(q, \lambda)$ there is global correlation in σ given G and therefore there is an additional term in the mutual information. We refer to Lemma 4.18 and Proposition 4.21 below for a more formal statement along these lines. In particular, in the simplest case of $q = 2$ communities and $d\lambda^2 < 1$, this improves the result by [DAM16], where their proof method requires that the average degree diverges $d \rightarrow \infty$. We further prove in Theorem 3.13 that such single letter characterization holds for (possibly non-symmetric) hypergraph stochastic block model below the weak recovery threshold. Moreover, we provide a general condition under which the phase transition of per-vertex mutual information occurs at the weak recovery threshold.

1.1.1 Hypothesis testing for the existence of communities

Theorem 1.1 establishes that the weak recovery threshold $d_*(q, \lambda)$ marks the sharp phase transition of the following hypothesis testing from a single observation G

$$H_0 : G \sim \mathcal{G}_{\text{ER}}\left(n, \frac{d}{n}\right) \quad \text{vs.} \quad H_1 : G \sim \mathcal{G}\left(n, q, \frac{a}{n}, \frac{b}{n}\right) \quad (3)$$

In particular, if $d < d_*(q, \lambda)$, then there is no perfect sequence of statistical tests, whereas if $d > d_*(q, \lambda)$ there exists a test which achieves vanishing Type I and Type II error exponentially fast at least along a subsequence.

However, for $d < d_*(q, \lambda)$, Theorem 1.1 does not provide any information on the power achieved by the optimal likelihood ratio (LR) test. We next characterize the asymptotic power obtained by LR test below the weak recovery threshold. Let us denote the likelihood ratio of the block model $\mathbf{G}^* \sim \mathcal{G}(n, q, \frac{a}{n}, \frac{b}{n})$ with respect to Erdos Renyi graph $\mathbf{G} \sim \mathcal{G}_{\text{ER}}(n, \frac{d}{n})$ by

$$\mathcal{L}_n(G) = \frac{\mathbb{P}(\mathbf{G}^* = G)}{\mathbb{P}(\mathbf{G} = G)}.$$

Theorem 1.2. (Special case of Theorem 3.14) *For average degree $d \equiv \frac{a+(q-1)b}{q}$ below the weak recovery threshold $d < d_*(q, \lambda)$, the likelihood ratio under null $\mathbf{G} \sim \mathcal{G}_{\text{ER}}(n, \frac{d}{n})$ converges in distribution to*

$$\mathcal{L}_n(\mathbf{G}) \xrightarrow{d} \mathcal{L}_\infty := \prod_{\ell \geq 3} \left\{ (1 + (q-1)\lambda^\ell)^{\mathbf{X}_\ell} e^{-\frac{(q-1)(d\lambda)^\ell}{2\ell}} \right\},$$

where $(\mathbf{X}_\ell)_{\ell \geq 3}$ are independent Poisson random variables with mean $\mathbb{E}\mathbf{X}_\ell = \frac{1}{2\ell}d^\ell$. Here, the random variable $\mathcal{L}_\infty \equiv \mathcal{L}_\infty(d, \lambda)$ satisfies the following.

- (1) For $d \in (0, \lambda^{-2})$, which is a superset of $(0, d_*(q, \lambda))$, \mathcal{L}_∞ is well-defined (i.e. the infinite product converges a.s.) and has finite second moment $\mathbb{E}\mathcal{L}_\infty^2 < \infty$.
- (2) If $1 \leq d < \lambda^{-2}$ holds, then the random variable \mathcal{L}_∞ does not have a point mass. In particular, in the regime $1 \leq d < d_*(q, \lambda)$, the likelihood ratio test for the hypothesis testing task (3) at significance level $\alpha \in (0, 1)$ has the asymptotic power $\beta_*(\alpha) \equiv \beta_*(\alpha; q, d, \lambda) \in (0, 1)$ defined by

$$\beta_*(\alpha) = \mathbb{E}[\mathcal{L}_\infty \mathbf{1}\{\mathcal{L}_\infty \geq C_\alpha\}]. \quad (4)$$

Here, $C_\alpha > 0$ is an arbitrary constant satisfying $\mathbb{P}(\mathcal{L}_\infty \geq C_\alpha) = \alpha$, whose existence is guaranteed.

The threshold $d_{\text{KS}} := \lambda^{-2}$ is called the *Kesten-Stigum* (KS) threshold, which was first discovered in the context of the Markov processes on trees in the seminal work of Kesten and Stigum [KS66]. In the context of sparse block model, it was first conjectured in [DKMZ11] that above the KS threshold $d\lambda^2 > 1$, the weak recovery can be done in a computationally efficient way. This conjecture was later proved for symmetric block model with $q = 2$ communities by [Mas14, MNS18, BLM15] and for general block model by [AS18]. In particular, the fact that the weak recovery threshold $d_*(q, \lambda)$ is at most the KS threshold d_{KS} for any $q \geq 2$ and $\lambda \in (-\frac{1}{q-1}, 1)$ follows from the result [AS18].

In the subcritical regime $d < 1$, the random variable \mathcal{L}_∞ is a discrete random variable supported on the countable set

$$\left\{ z : z = (1-d\lambda)^{\frac{q-1}{2}} \prod_{\ell \geq 1} (1+(q-1)\lambda^\ell)^{x_\ell} \text{ for some } x = (x_\ell)_{\ell \geq 1} \text{ with finitely many non-zero elements.} \right\}.$$

Thus, for $d < 1$ the weak convergence of the likelihood ratio obtained in Theorem 1.2 only characterizes the asymptotic power of the *non-randomized* LR test for the hypothesis test (3) (see also Corollary 4.14). This is because the LR test might require randomization since the law of \mathcal{L}_∞ is atomic for $d < 1$. We refer to [LR05, Section 3.2] for the description of randomization in LR tests.

1.1.2 Computational aspects of hypothesis testing

We next discuss the computational aspects in the hypothesis testing (3). Although the LR test achieves optimal power for hypothesis test (3) at any significance level by the Neyman-Pearson lemma, it is far from clear how to efficiently approximate the likelihood ratio $\mathcal{L}_n(G)$. In fact, $\mathcal{L}_n(G)$ can be interpreted as a certain *partition function* from statistical physics, and it is often NP-hard (or even #P-hard) to approximate partition functions, see e.g. [JS93, Ist00, GJ07]. Note that given an assignment of the communities $\underline{\sigma} = (\sigma_v)_{v \in V} \in [q]^V$ and $G = (V, E)$, respectively denote the number of monochromatic edges and non-edges by

$$e_0(G, \underline{\sigma}) := |\{(u, v) \in E : \sigma_u = \sigma_v\}|, \quad \tilde{e}_0(G, \underline{\sigma}) := |\{(u, v) \notin E : \sigma_u = \sigma_v\}|.$$

Then, it is straightforward to calculate

$$\mathcal{L}_n(G) = q^{-n} \sum_{\underline{\sigma} \in [q]^V} \left(\frac{n-a}{n-(a+(q-1)b/q)} \right)^{\tilde{e}_0(G, \underline{\sigma})} \left(\frac{n-b}{n-(a+(q-1)b/q)} \right)^{\binom{n}{2} - |E| - \tilde{e}_0(G, \underline{\sigma})} \left(\frac{qa}{a+(q-1)b} \right)^{e_0(G, \underline{\sigma})} \left(\frac{qb}{a+(q-1)b} \right)^{|E| - e_0(G, \underline{\sigma})},$$

and in particular, summing over $\underline{\sigma} \in [q]^V$ makes the naive computation of $\mathcal{L}_n(G)$ intractable.

On the other hand, we now construct a *computationally efficient* test for which it achieves the power defined in Eq. (4). Given a graph G and an integer $\ell \geq 3$, let $X_\ell(G)$ denote the number of cycles of length ℓ . Then, consider the statistic

$$\mathcal{T}_n(G) := \prod_{3 \leq \ell \leq K_n} (1 + (q-1)\lambda^\ell)^{X_\ell(G)},$$

where the truncation parameter K_n is chosen to diverge slowly to infinity, say $1 \ll K_n = O(\log \log n)$.

We note that if G is drawn from the sparse block model or the sparse Erdos Renyi graph, then the condition $K_n = O(\log \log n)$ guarantees that the computation of $\mathcal{T}_n(G)$ can be done in nearly linear time (up to log factors) with high probability. This is because when G is sparse, the number of cycles $X_\ell(G)$ up to length $\log \log n$ can be counted by looking at depth $\log \log n$ neighborhood of each nodes, each of which has at most $(Cd)^{\log \log n}$ neighbors with high probability.

In addition, note that $\mathcal{T}_n(G)$ is a degree K_n polynomial in the indicator of the edges of G . Our next theorem establishes that a test based on this low-degree polynomial $\mathcal{T}_n(G)$ is asymptotically most powerful below the weak recovery threshold.

Theorem 1.3. (Special case of Theorem 3.15) *Let $K_n = O(\log \log n)$ and $K_n \rightarrow \infty$ as $n \rightarrow \infty$. For a significance level $\alpha \in (0, 1)$, consider the test $\phi_{n,\alpha}(\cdot)$ which rejects the null \mathbf{H}_0 in Eq. (3) with probability*

$$\phi_{n,\alpha}(G) := \begin{cases} 1 & \text{if } \mathcal{T}_n(G) > C'_{n,\alpha}; \\ 0 & \text{otherwise.} \end{cases}$$

Here, the constants $C'_{n,\alpha} > 0$ is chosen so that we have $\mathbb{P}(\mathcal{T}_n(\mathbf{G}) > C'_{n,\alpha}) \leq \alpha \leq \mathbb{P}(\mathcal{T}_n(\mathbf{G}) \geq C'_{n,\alpha})$, where $\mathbf{G} \sim \mathcal{G}_{\text{ER}}(n, \frac{d}{n})$. Then, for $1 \leq d < \lambda^{-2}$, the test $\phi_{n,\alpha}$ achieves the power $\beta_*(\alpha)$ defined in Eq. (4). That is, under the alternative $\mathbf{G}^* \sim \mathcal{G}(n, q, \frac{a}{n}, \frac{b}{n})$, we have

$$\mathbb{E}\phi_{n,\alpha}(\mathbf{G}^*) \rightarrow \beta_*(\alpha) \quad \text{as } n \rightarrow \infty.$$

In particular, in the regime $1 \leq d < d_*(q, \lambda)$, the test $\phi_{n,\alpha}(\cdot)$ is asymptotically most powerful for the hypothesis test (3).

For $q \geq 5$ communities, it is known [BMNN16, AS16] that at least for small enough λ , there is a gap between the weak recovery threshold $d_*(q, \lambda)$ and the KS threshold $d_{\text{KS}} \equiv \lambda^{-2}$. In this case, Theorems 1.1 and 1.3 uncover an important phenomenon: the weak recovery threshold $d_*(q, \lambda)$ marks the *local* to *global* phase transition of the block model. Specifically, for average degree d below the weak recovery threshold $d_*(q, \lambda)$, the existence of the community structure is fully captured by the *local* structure of the graph, namely the number of short cycles. On the contrary, in the regime $d_*(q, \lambda) < d < d_{\text{KS}}$, a *global* structure emerges, which cannot be described by the number of short cycles. Notably, Theorem 1.1-(3) shows that in the latter regime, the asymptotic power of the LR test must equal 1 (cf. Neyman-Pearson lemma) at least along a subsequence. However, Theorem 1.3 implies that the test $\phi_{n,\alpha}$ based on a low-degree polynomial achieves limited power $\beta_*(\alpha) < 1$ in the intermediate regime $d \in (d_*(q, \lambda), d_{\text{KS}})$. This discrepancy thus demonstrates *statistical-computational gap* in hypothesis testing. A natural question would be to see if the low-degree test $\phi_{n,\alpha}$ achieves the optimal power among all computationally efficient tests in the regime $d \in (d_*(q, \lambda), d_{\text{KS}})$ under a suitable complexity-theoretic conjecture, see e.g. [MW23] for recent related work.

On the contrary, for $q = 2$ communities and $q \in \{3, 4\}$ communities with large enough average degree d , it is known that $d_*(q, \lambda) = d_{\text{KS}}$ holds [MNS15, MSS23]. Thus, in these cases, our results yield that there is no statistical-computational gap in hypothesis testing at any significance level. Indeed, if $d < d_*(q, \lambda) = d_{\text{KS}}$ holds, then the low-degree-polynomial test $\phi_{n,\alpha}(\cdot)$ is most powerful by Theorem 1.1. Moreover, it is known that above the KS threshold $d > d_{\text{KS}}$, the detection is solvable by efficient algorithms (e.g. by counting short cycles [Mas13, MNS18, Abb18] or by a semidefinite programming relaxation [BMR21]). Specifically, we obtain the following corollary by combining our results with the results of [MNS15, MSS23].

Corollary 1.4. *For $q = 2$ communities or $q \in \{3, 4\}$ and small enough $|\lambda| \leq \lambda_0$, where $\lambda_0 > 0$ is a universal constant, the results of Theorems 1.1, 1.2, 1.3 hold with $d_*(q, \lambda) \equiv \lambda^{-2}$. Thus, in this case, for any average degree d such that $d \geq 1$ and $d \neq \lambda^{-2}$ hold and any significance level $\alpha \in (0, 1)$, there exists an efficient sequence of statistical tests $\varphi_{n,\alpha}(\cdot)$ that is asymptotically most powerful at level α .*

2 Further related work

Stochastic block models and their hypergraph analogs The stochastic block model, first introduced in [HLL83], has been studied extensively in statistics [SN97, BC09, RCY11] and in computer science [DF89, JS98, CK01, McS01, CO10]. The block model in the sparse regime became a major object of research due to the landmark paper [DKMZ11]. In particular, [DKMZ11] conjectured a different set of phase transitions within the sparse block model, where the Kesten-Stigum (KS) threshold [KS66] plays a crucial role. Later, it was shown in a series of works [Mas14, MNS18, BLM15, AS15, AS18] that above the KS threshold, there is an efficient algorithm that achieves both weak recovery and detection. The tightness/non-tightness of KS threshold for weak recovery was also studied by a number of papers [MNS15, AS16, BMNN16, RTSZ19, MSS23]. We refer to the survey by Abbe [Abb18] for the developments of the block model for more references.

More recently, community detection in sparse hypergraph has also gained significant interest. When the average degree is at least of order $\Omega(\log^2 n)$, Ghoshdastidar and Dukkipati [GD17a] proved that a spectral algorithm achieves near-perfect recovery of the community structure. When the average degree is of order constant, [ACKZ15] conjectured that a phase transition occurs at the KS threshold - above this threshold, efficient weak recovery is possible, and impossible below the threshold. Efficient recovery above the KS threshold was proved for hypergraph models for 2 communities by Pal and Zhu [PZ21] and general case by Stephan and Zhu [SZ22]. Gu and Polyanskiy [GP23] recently showed that the information theoretic threshold for weak recovery is the KS threshold within the r uniform hypergraph stochastic block model for $r = 3, 4$ while the same is not true for $r \geq 7$.

General factor models The framework for factor models capture many standard models in theoretical computer science such as k-SAT models and in statistical physics such as Potts models (see e.g. Chapter 9 in [MM09]). The planted factor models was initially introduced to study the phase transitions of random constraint satisfaction problems [ACO08, KZ09]. Later, Coja-Oghlan, Krzakala, Perkins, Zdeborová [COKPZ18] studied the information theoretic thresholds for planted factor models that satisfy certain convexity and balanced conditions. In particular, [COKPZ18] obtained an explicit variational principle for the weak recovery threshold for the disassortative block models. Subsequently, Coja-Oghlan, Efthymiou, Jaafari, Kang, Kapetanopoulos [COEJ+18] proved that in the disassortative case, the planted and the null models are mutually contiguous below the weak recovery threshold. The work [COEJ+18] also introduced the Kesten-Stigum bound for planted factor models in the uniform prior $\pi = \text{Unif}([q])$ case. The analysis of [COKPZ18, COEJ+18] was generalized to factor models with hard constraints by [COKM20]. The planted models also encompass certain high-dimensional Bayesian inference problems as studied in [BPS21, BP22].

Statistical Computational Gaps In studying the computational complexity of statistical tasks such as weak recovery and detection, it is natural to question the average case complexity of the task. Many recent works attempt to answer this question using a number of different perspectives. These include average case reduction to a widely-believed statistically hard problems such as the hidden-clique problem [Jer92, Kuč95], see e.g. [BB20]. A different approach is to study a restricted class of algorithms such as local algorithms (see e.g. [GS14] and follow up work), SQ algorithms [Kea98] or low-degree polynomials. In particular, a recent line of work (see e.g. [HS17, Hop18, KWB19, BKW20, GJW20, HW20, BH21, Wei20]) uses a “low-degree heuristic” to predict computational-statistical gaps for a variety of problems. In this context, Hopkins and Steurer [HS17] showed that below the KS threshold, functions that can determine if two vertices are in the same community better than random have to be of degree at least $\Omega(n)$.

3 Main results in the general case

Our results in the most general setting are stated in the framework of planted factor models, which is a generalization of block models. In Sections 3.1,3.2, we define and state our main results for the planted factor model and the associated inference tasks: weak recovery and detection. In Section 3.3, we apply these results to hypergraph stochastic block models (HSBM).

3.1 Planted factor models

Recall that the sparse symmetric block model is defined by first “planting” a community structure $\underline{\sigma}^* \equiv (\sigma_v^*)_{v \in V} \stackrel{i.i.d.}{\sim} \text{Unif}(\{1, \dots, q\})$. Subsequently, based on this planted structure, a linear number of edges are drawn. Likewise, in the planted factor models, communities (also termed as spins) are represented as $\underline{\sigma}^* \equiv (\sigma_v^*)_{v \in V} \stackrel{i.i.d.}{\sim} \pi$, where π encodes the community’s prior. Using $\underline{\sigma}^*$, we create a linear number of clauses, with each clause connecting k nodes. When $k = 2$, these clauses are analogous to the edges of the block model. However, allowing $k \geq 3$ enables us to consider hypergraph analogs of the block models called the hypergraph stochastic block models (HSBM) considered in [GD17a, GD17b, ACKZ15]. In addition, a notable extension in planted factor models is the consideration of clause connectivity probabilities as random variables. Such extension ensures that the planted factor models encompass planted constraint satisfaction problems studied in statistical physics and computer science [AJM05, JMS07, KMZ14, FPV15].

We first define the necessary notations that will be used throughout the paper. Given $q \geq 2$, we denote by $[q] \equiv \{1, \dots, q\}$ the finite set of spins or communities. We let $\pi \in \mathcal{P}([q])$ be a probability measure on $[q]$ which represents the prior of different communities.

We let $k \geq 2$ denote the number of interaction between different nodes. Further, we let Ψ be a finite set of *weight functions* described by

$$\psi : \{1, \dots, q\}^k \rightarrow \mathbb{R}_{>0}.$$

Here, we assumed the weight functions to have positive values, which corresponds to *positive temperature* models in statistical physics language. Let $p \in \mathcal{P}(\Psi)$ be a prior distribution of the weight functions. Without loss of generality we let $\Psi = \text{supp}(p)$ since otherwise we can reduce the set Ψ .

Throughout, we assume the following: for any permutation $\theta \in S_k$, let $\psi^\theta : [q]^k \rightarrow \mathbb{R}_{>0}$ defined by $\psi^\theta(\sigma_1, \dots, \sigma_k) = \psi(\sigma_{\theta(1)}, \dots, \sigma_{\theta(k)})$. Then, for any $\psi \in \Psi$ and $\theta \in S_k$, we assume that

$$\psi^\theta \in \Psi \quad \text{and} \quad p(\psi^\theta) = p(\psi) > 0, \tag{5}$$

which reflects the exchangeability of the model with respect to the different variables.

A (bipartite) factor graph $G = (V, F, E, (\psi_a)_{a \in F})$ consists of the following.

- The set of variables $V := \{v_1, \dots, v_n\}$ and the set of clauses (a.k.a. function nodes) $F := \{a_1, \dots, a_m\}$.
- The set of edges E , where $e = (av) \in E$ connects $a \in F$ and $v \in V$. Here, each clause $a \in F$ is connected to k variables, and we denote its neighborhood by $\delta a := \{v \in V : a \sim v\}$ with the convention that $\delta a = (v_1, \dots, v_k)$ is ordered. We let $\delta_i a \equiv v_i$ be the i 'th variable adjacent to a for $1 \leq i \leq k$.
- The set of weight function $(\psi_a)_{a \in F}$. Here, $\psi_a \in \Psi$ is the weight function assigned to a clause $a \in F$. Since δa is ordered, for an assignment of communities $\underline{x} = (x_v)_{v \in V} \in [q]^V$, the expression $\psi_a(\underline{x}_{\delta a}) \equiv \psi_a(x_{v_1}, \dots, x_{v_k})$ is well-defined.

Given the prior p on Ψ , the random graph model with no community (a.k.a. planted) structure corresponds to the *null model* defined below.

Definition 3.1. (*The null model*) Given $V = \{v_1, \dots, v_n\}$, $F = \{a_1, \dots, a_m\}$, and $p \in \mathcal{P}(\Psi)$, the null model $\mathbf{G}(n, m) \equiv \mathbf{G}(n, m, p)$ is a factor graph $(V, F, E, (\psi_a)_{a \in F})$ defined as follows. For each $a \in F$, its neighborhood $\delta a \in V^k$ is drawn independently and uniformly at random from V^k . The weight function ψ_a is drawn i.i.d. from p , i.e. $(\psi_a)_{a \in F} \stackrel{i.i.d.}{\sim} p$.

Given a factor graph $G = (V, F, E, (\psi_a)_{a \in F})$ and $\underline{\sigma} \in [q]^V$, let

$$\psi_G(\underline{\sigma}) := \prod_{a \in F} \psi_a(\underline{\sigma}_{\delta a}). \quad (6)$$

Subsequently, the random graph model with a community structure corresponds to the *planted model*.

Definition 3.2. (*The planted model*) Given $V = \{v_1, \dots, v_n\}$, $F = \{a_1, \dots, a_m\}$, and priors $\pi \in \mathcal{P}([q])$ and $p \in \mathcal{P}(\Psi)$, the planted model $\mathbf{G}^*(n, m, \underline{\sigma}^*) \equiv \mathbf{G}^*(n, m, \underline{\sigma}^*, p, \pi)$ is a factor graph defined as follows. First draw a planted/community structure $\underline{\sigma}^* \equiv (\sigma_1^*, \dots, \sigma_n^*)$ by $\sigma_i^* \stackrel{i.i.d.}{\sim} \pi$. Given $\underline{\sigma}^*$, draw $\mathbf{G}^*(n, m, \underline{\sigma}^*)$ from the distribution

$$\mathbb{P}\left(\mathbf{G}^*(n, m, \underline{\sigma}^*) = G \mid \underline{\sigma}^* = \underline{\sigma}\right) = \mathbb{P}(\mathbf{G}(n, m) = G) \cdot \frac{\psi_G(\underline{\sigma})}{\mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\sigma})]},$$

where \mathbb{E} denotes the expectation with respect to the null model $\mathbf{G}(n, m)$. Equivalently, given $\underline{\sigma}^* = (\sigma_v)_{v \in V}$, independently draw for each clause $a \in F$ the neighborhood δa and the weight function ψ_a from the distribution

$$\mathbb{P}(\delta a = (v_1, \dots, v_k), \psi_a = \psi) = \frac{p(\psi)\psi(\sigma_{v_1}, \dots, \sigma_{v_k})}{\sum_{\psi' \in \Psi} p(\psi') \sum_{v'_1, \dots, v'_k \in V} \psi'(\sigma_{v'_1}, \dots, \sigma_{v'_k})} = \frac{1}{n^k} \cdot \frac{p(\psi)\psi(\sigma_{v_1}, \dots, \sigma_{v_k})}{\mathbb{E}_{p, u}[\psi(\underline{\sigma}_\omega)]}, \quad (7)$$

where $\mathbb{E}_{p, u}$ denotes the expectation with respect to $\psi \sim p$ and $\omega \sim u := \text{Unif}(V^k)$.

The planted model can be specialized to sparse symmetric block model by taking $k = 2$, $\pi = \text{Unif}([q])$, and p to put all of its mass on a specific weight function (see Eq. (116)). Within this context, a clause $a \in F$ corresponds to an edge connecting $\delta_1 a$ and $\delta_2 a$ in the block model. Thus m corresponds to the number of edges in the block model, which is approximately Poisson with mean $dn/2$. Therefore, in the broader planted factor model, we let the clause count m to follow a Poisson distribution. Using a concise notation, we let

$$\mathbf{G}^* \equiv \mathbf{G}^*(n, \mathbf{m}, \underline{\sigma}^*, p, \pi), \quad \mathbf{G} \equiv \mathbf{G}(n, \mathbf{m}, p), \quad \text{where } \mathbf{m} \sim \text{Poi}(dn/k). \quad (8)$$

Further, we respectively denote the probability distribution of \mathbf{G}^* and \mathbf{G} by

$$\mathcal{G}_{\text{plant}}(n, d, p, \pi) := \text{Law}(\mathbf{G}^*), \quad \mathcal{G}_{\text{null}}(n, d, p) := \text{Law}(\mathbf{G}). \quad (9)$$

Given a single observation G , the two central tasks in statistical inference are *weak recovery* and *detection*.

Definition 3.3. (*Weak recovery*) Consider the planted model $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$. We say that *weak recovery is possible* at d , if there exists an $\varepsilon > 0$ and (sequence of) estimators $\hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}_n(\mathbf{G}^*)$ that takes as an input the factor graph \mathbf{G}^* and returns $\hat{\underline{\sigma}} \equiv (\hat{\sigma}_v)_{v \in V} \in [q]^V$, such that

$$\limsup_{n \rightarrow \infty} \mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon, \quad (10)$$

where the *overlap* between $\underline{\sigma}^1 \equiv (\sigma_v^1)_{v \in V} \in [q]^V$ and $\underline{\sigma}^2 \equiv (\sigma_v^2)_{v \in V} \in [q]^V$ is defined by

$$A(\underline{\sigma}^1, \underline{\sigma}^2) := \max_{\Gamma \in S_q} \left\{ \frac{1}{q} \sum_{i=1}^q \frac{|\{v \in V : \sigma_v^1 = i, \sigma_v^2 = \Gamma(i)\}|}{|\{v \in V : \sigma_v^1 = i\}|} \right\}. \quad (11)$$

Here, the maximum is taken with respect to permutations Γ among the set of communities $[q]$. By fixing p, π and varying $d > 0$, the weak recovery threshold $d_* \equiv d_*(p, \pi)$ is defined by

$$d_* \equiv d_*(p, \pi) := \inf \left\{ d > 0 : \text{weak recovery is possible at } d \text{ for } \mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi) \right\}.$$

We remark that Definition 3.3 appeared previously in [AS18, MSS23]. In particular, the maximum over $\Gamma \in S_q$ is taken in the definition of the overlap $A(\underline{\sigma}^1, \underline{\sigma}^2)$ to account for symmetries, if any, of the labels of the communities. For example, if $\pi = \text{Unif}([q])$, inherent symmetries between communities imply that communities can only be recovered up to a permutation. For an equivalent, but slightly different formulation, see [AS18, Definition 2.3].

We further remark that in Theorem 3.10 below, we give equivalent descriptions of the weak recovery that involve two-point correlations and orthogonality of the samples drawn from the posterior. Furthermore, it is crucial to note that in Proposition 4.1, we characterize a simple condition (SYM) which ensures that the weak recovery threshold $d_*(p, \pi)$ is non-trivial, lying in the interval $d_*(p, \pi) \in [\frac{1}{k-1}, \infty)$.

Before proceeding further, we make two elementary observations. First, note that the task of weak recovery is monotone in $d > 0$. Specifically, if weak recovery is impossible at d , then weak recovery is impossible at d' for all $0 < d' < d$. This is validated by deleting the clauses of $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$ independently with probability d'/d yields a sample drawn from $\mathcal{G}_{\text{plant}}(n, d', p, \pi)$. In particular, for all $d > d_*(p, \pi)$, weak recovery is possible at $d > 0$. Second, for arbitrary $\underline{\sigma}^1, \underline{\sigma}^2 \in [q]^V$, the overlap satisfies $A(\underline{\sigma}^1, \underline{\sigma}^2) \geq \frac{1}{q}$ by taking average over $\Gamma \in S_q$ instead of taking maximum as in (11). Thus, the impossibility of weak recovery is equivalent to

$$\lim_{n \rightarrow \infty} \mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] = \frac{1}{q}, \quad \text{for any (sequence of) estimators } \hat{\sigma} \equiv \hat{\sigma}_n(\mathbf{G}^*). \quad (12)$$

Definition 3.4. (*Detection*) Consider the following hypothesis testing task given a single observation G :

$$\mathbf{H}_0 : G \sim \mathcal{G}_{\text{null}}(n, d, p) \quad \text{vs.} \quad \mathbf{H}_1 : G \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi) \quad (13)$$

We say that *detection is possible* at d , if there exists a sequence of test $(\phi_n(G))_{n \geq 1}$, which rejects the null \mathbf{H}_0 with probability $\phi_n(G) \in [0, 1]$, such that it achieves vanishing Type 1 and Type 2 errors. That is,

$$\mathbb{E}[\phi_n(\mathbf{G})] + \mathbb{E}[1 - \phi_n(\mathbf{G}^*)] \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (14)$$

where $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$ and $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$. If there does not exist such sequence of tests, we say *detection is impossible* at d .

3.2 Main results for the factor models

In our first result for the planted factor model, we prove that below the weak recovery threshold d_* and the (generalized) Kesten-Stigum (KS) threshold $d_{\text{KS}} \equiv d_{\text{KS}}(\pi, p)$ defined in Section 4.1, the detection is impossible.

Theorem 3.5. *Consider the planted factor model $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$ in the regime $d < d_* \wedge d_{\text{KS}}$. Then, the planted model \mathbf{G}^* is mutually contiguous with the null model $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$. That is, for any sequence of events $(\mathcal{A}_n)_{n \geq 1}$, $\mathbb{P}_{\mathbf{G}^*}(\mathcal{A}_n) \rightarrow 0$ if and only if $\mathbb{P}_{\mathbf{G}}(\mathcal{A}_n) \rightarrow 0$. In particular, the detection is impossible at d .*

In some special cases, it is known that the weak recovery threshold $d_* \equiv d_*(p, \pi)$ is at most the KS threshold $d_{\text{KS}} \equiv d_{\text{KS}}(p, \pi)$ in which case the condition $d < d_* \wedge d_{\text{KS}}$ in Theorem 3.5 can be simplified to $d < d_*$. This is the case for sparse (non-symmetric) block model, which corresponds to $k = 2$ and p being Dirac measure, by [AS18]. The same holds when $k \geq 3$, p is a Dirac measure, and π is the uniform measure $\text{Unif}([q])$ by [SZ22]. It would be interesting to show that for general planted factor models $d_* \leq d_{\text{KS}}$ holds.

In order to prove Theorem 3.5, we will prove a stronger result (cf. Theorem 4.12) which characterizes the asymptotic power of the likelihood ratio test. Since such characterization requires extra technical notations regarding ζ -cycles, we defer the statement to Section 4. The proof of Theorem 3.5 is in Section 4.1.

3.2.1 Consequences of contiguity in point estimation

As a consequence of the mutual contiguity in Theorem 3.5, we establish that it is impossible to consistently estimate the set of weight functions $\Psi \equiv \text{supp}(p)$. Specifically, we consider the following notion of consistency.

Definition 3.6. Given $T \geq 1$, consider the parametric family of planted factor models $\mathcal{G}_{\text{plant}}(n, d, p, \pi)$ with fixed size of the support $T \equiv |\text{supp}(p)|$. Let $\widehat{\Psi}_n(G)$ be an estimator that takes as an input a factor graph G with n variables and outputs a set of weight functions $\{\widehat{\psi}_1, \dots, \widehat{\psi}_T\}$, where $\widehat{\psi}_t : [q]^k \rightarrow \mathbb{R}_{>0}$ for $1 \leq t \leq T$. For a fixed $\varepsilon > 0$, we say that the sequence of estimators $(\widehat{\Psi}_n(G))_{n \geq 1}$ is ε -locally consistent at (d, p, π) if the following holds. Let $\Psi = \{\psi_1, \dots, \psi_T\}$. Then, for any $\Psi' = \{\psi'_1, \dots, \psi'_T\}$ such that $\|\psi'_t - \psi_t\|_\infty \leq \varepsilon$ holds for all $1 \leq t \leq T$, we have as $n \rightarrow \infty$

$$\widehat{\psi}_t(\mathbf{G}^*) \xrightarrow{p} \psi', \quad \text{for } \mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p', \pi) \text{ and } 1 \leq t \leq T.$$

Here, $p' \sim \mathcal{P}(\Psi')$ is defined by letting $p'(\psi'_t) = p(\psi_t)$ for $1 \leq t \leq T$.

To clarify the definition, the non-existence of ε -locally consistent estimator describes circumstances in which, despite having access to the information of the prior of the communities $\pi \in \mathcal{P}([q])$ and the mass of each weight functions $(p(\psi))_{\psi \in \Psi} \in \mathbb{R}^T$, it remains infeasible to estimate the set of weight functions $\Psi \equiv \text{supp}(p)$ even within its ε neighborhood. Consequently, our definition of an ε -locally consistent estimator is notably weaker than the usual definition of consistent estimator, which imposes no parameter restriction. Our subsequent corollary whose proof is in Section 4.1 demonstrate that, even within this weaker notion of consistency, estimating Ψ below the weak recovery threshold and the KS threshold is impossible.

Corollary 3.7. *If $d < d_* \wedge d_{\text{KS}}$ holds, then for any $\varepsilon > 0$, there does not exist ε -locally consistent estimator of Ψ at (d, p, π) .*

3.2.2 Asymptotic mutual information

We next consider the mutual information $I(\mathbf{G}^*, \underline{\sigma}^*)$ between the planted factor model $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$ and the community structure $\underline{\sigma}^* \in [q]^V$, and the Kuller-Leibler divergence $D_{\text{KL}}(\mathbf{G}^* \parallel \mathbf{G})$

between the planted model \mathbf{G}^* and the null model $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$. In the regime $d > d_*$, our results are stated under the following assumption:

- (MIN): For a probability vector $\pi \in \mathbb{R}^q$, define the set

$$\mathcal{R}_\pi := \left\{ R \in [0, 1]^{q \times q} : \sum_{i \in [q]} R(i, j) = \pi_j \text{ for } j \in [q], \text{ and } \sum_{j \in [q]} R(i, j) = \pi_i \text{ for } i \in [q] \right\}. \quad (15)$$

Let the function $\mathcal{F} : \mathcal{R}_\pi \rightarrow \mathbb{R}_{\geq 0}$ be defined by

$$\mathcal{F}(R) := \sum_{\underline{\sigma}, \underline{\tau} \in [q]^k} \mathbb{E}_p[\psi(\underline{\sigma})\psi(\underline{\tau})] \prod_{s=1}^k R(\sigma_s, \tau_s). \quad (16)$$

Then, the $R \mapsto \mathcal{F}(R)$ is uniquely minimized at $R = \pi\pi^\top$.

We remark that in the case $\pi = \text{Unif}([q])$, the assumption (MIN) was first considered in [COEJ⁺18]. In addition, it is straightforward to verify that the sparse symmetric block model satisfies such assumption (see Eq. (28) below). The following result shows that under such condition, the normalized mutual information $I(\mathbf{G}^*, \underline{\sigma}^*)/n$ exhibits a phase transition at the weak recovery threshold $d_*(p, \pi)$ defined in Definition 3.3.

Theorem 3.8. *We have the following.*

- (1) *Suppose that $d < d_*$ holds. Then as $n \rightarrow \infty$, we have*

$$\frac{1}{n} D_{\text{KL}}(\mathbf{G}^* \parallel \mathbf{G}) \rightarrow 0, \quad \text{and} \quad \frac{1}{n} I(\mathbf{G}^*, \underline{\sigma}^*) \rightarrow \frac{d}{k} \cdot \mathbb{E}_{p, \pi} \left[\frac{\psi(\underline{\sigma})}{\xi} \log \left(\frac{\psi(\underline{\sigma})}{\xi} \right) \right], \quad (17)$$

where $\mathbb{E}_{p, \pi}$ denotes the expectation with respect to $\psi \sim p$ and $\underline{\sigma} \sim \pi^{\otimes k}$, and we denoted $\xi := \mathbb{E}_{p, \pi}[\psi(\underline{\sigma})]$.

- (2) *Conversely, suppose that $d > d_*$ and assume that the condition (MIN) holds. Then, we have*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} D_{\text{KL}}(\mathbf{G}^* \parallel \mathbf{G}) > 0, \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{1}{n} I(\mathbf{G}^*, \underline{\sigma}^*) < \frac{d}{k} \cdot \mathbb{E}_{p, \pi} \left[\frac{\psi(\underline{\sigma})}{\xi} \log \left(\frac{\psi(\underline{\sigma})}{\xi} \right) \right].$$

- (3) *Assuming the condition (MIN), the following holds for any $d_0 > 0$. For any $\varepsilon > 0$, there exists $\eta \equiv \eta(\varepsilon) > 0$ not depending on n such that if $\mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds for some estimator $\hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}_n(\mathbf{G}_0^*)$ where $\mathbf{G}_0^* \sim \mathcal{G}_{\text{plant}}(n, d_0, p, \pi)$, then for all $d > d_0$, there is an event \mathcal{A}_n^* satisfying*

$$\mathbb{P}(\mathbf{G} \in \mathcal{A}_n^*) \leq 2e^{-\eta m}, \quad \text{and} \quad \mathbb{P}(\mathbf{G}^* \in \mathcal{A}_n^*) \geq 1 - 2e^{-\eta m}, \quad (18)$$

where $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$ and $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$. In particular, if $d > d_*$ and the condition (MIN) holds, then there exists a sequence of events $(\mathcal{A}_n^*)_{n \geq 1}$, a subsequence $(n_\ell)_{\ell \geq 1}$, and a constant $\eta > 0$ such that $\mathbb{P}(\mathbf{G} \in \mathcal{A}_{n_\ell}^*) \leq 2e^{-\eta m_\ell}$ while $\mathbb{P}(\mathbf{G}^* \in \mathcal{A}_{n_\ell}^*) \geq 1 - 2e^{-\eta m_\ell}$. Thus, \mathbf{G} and \mathbf{G}^* are mutually orthogonal along a subsequence.

Remark 3.9. We note that in general it is not known that if there exists a subsequence $(n_\ell)_{\ell \geq 1}$ such that the weak recovery is possible at d along $(n_\ell)_{\ell \geq 1}$, i.e. there exists $\varepsilon > 0$ such that for all $\ell \geq 1$, $\mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds for $\hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}_{n_\ell}(\mathbf{G}^*)$, $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n_\ell, d, p, \pi)$, then the weak recovery is possible at d (or even for $d' > d$) along every subsequence. Thus, if we replace \limsup with \liminf in Eq. (10) in Definition 3.3 we may in principle obtain a different threshold for weak recovery. The fact that we do not know that the ‘ \limsup ’ and ‘ \liminf ’ thresholds are the same is the reason that the statements in Theorem 1.1-(3) and Theorem 3.8-(3) use subsequences. However, like in many other random graph and spin-glass models [COKPZ18] we believe that these thresholds should be the same for weak recovery, and similarly for detection. Moreover, as seen in Theorem 3.8-(3), if we let d'_* be the ‘ \liminf ’ threshold where we replace \limsup with \liminf in Eq. (10), then we have mutual orthogonality for the entire sequence above this (possibly different) threshold $d > d'_*$.

3.2.3 Equivalent notions of weak recovery

The proof of Theorem 3.5 is based on a novel equivalence between the impossibility of weak recovery and near-orthogonality of the samples drawn from the posterior. To state the equivalence, we introduce more notations. Given a factor graph G with n variables and m clauses, we denote the posterior $\mu_G \in \mathcal{P}([q]^V)$ by

$$\mu_G(\underline{\sigma}) = \mathbb{P}(\underline{\sigma}^* = \underline{\sigma} \mid \mathbf{G}^*(n, m, \underline{\sigma}^*) = G), \quad \underline{\sigma} \in [q]^V. \quad (19)$$

We denote by $\langle \cdot \rangle_G$ the expectation with respect to samples $(\sigma^\ell)_{\ell \geq 1} \stackrel{i.i.d.}{\sim} \mu_G$ from the posterior. That is, for any $L \geq 1$ and $f : ([q]^V)^L \rightarrow \mathbb{R}$, we let

$$\langle f(\underline{\sigma}^1, \dots, \underline{\sigma}^L) \rangle_G := \sum_{\underline{\sigma}^1, \dots, \underline{\sigma}^L \in [q]^V} f(\underline{\sigma}^1, \dots, \underline{\sigma}^L) \prod_{\ell=1}^L \mu_G(\underline{\sigma}^\ell). \quad (20)$$

Given two samples $\underline{\sigma}^\ell \equiv (\sigma_v^\ell)_{v \in V} \in [q]^V$, $\ell = 1, 2$, the *overlap matrix* $R_{\underline{\sigma}^1, \underline{\sigma}^2} \equiv (R_{\underline{\sigma}^1, \underline{\sigma}^2}(i, j))_{i, j \leq q} \in \mathbb{R}^{q \times q}$ is defined by

$$R_{\underline{\sigma}^1, \underline{\sigma}^2}(i, j) = \frac{1}{n} \sum_{v \in V} \mathbb{1}\{\sigma_v^1 = i, \sigma_v^2 = j\}. \quad (21)$$

Concisely, the overlap matrix $R_{\underline{\sigma}^1, \underline{\sigma}^2}$ represents the empirical distribution of $(\sigma_v^1, \sigma_v^2)_{v \in V}$. In particular, the overlap matrix $R_{\underline{\sigma}^1, \underline{\sigma}^2}$ being close to $\pi \pi^\top$ indicates that the two samples $\underline{\sigma}^1, \underline{\sigma}^2$ are *near-orthogonal*, i.e. $\underline{\sigma}^1$ and $\underline{\sigma}^2$ appear as if they are drawn independently from $\pi^{\otimes V}$. In our subsequent result, we establish that the impossibility of weak recovery is *equivalent* to the near-orthogonality of two samples drawn from the posterior. It plays a crucial role in deriving Theorem 3.5, but it might also be of independent interest.

Theorem 3.10. *Given $k \geq 2, d > 0, \pi, p$, the following are equivalent:*

- (a) *Weak recovery is impossible at d . That is, (12) holds.*
- (b) *For any 2 distinct variables $u \neq v$ and $i, j \in [q]$, we have that as $n \rightarrow \infty$,*

$$\mathbb{P}(\sigma_u^* = i, \sigma_v^* = j \mid \mathbf{G}^*) \xrightarrow{p} \pi_i \pi_j.$$

(c) For any 2 distinct variables $u \neq v$, we have

$$\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{G}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2, \quad \text{and} \quad \sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{G}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2.$$

(d) The overlap matrix $R_{\underline{\sigma}^1, \underline{\sigma}^2}$ of the samples $\underline{\sigma}^1, \underline{\sigma}^2$ drawn from the posterior is trivial. That is, as $n \rightarrow \infty$,

$$\mathbb{E} \left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_1 \right\rangle_{\mathbf{G}^*} \rightarrow 0.$$

Theorem 3.10 is actually established within a more general framework, where \mathbf{G}^* can be replaced by any random variable $\mathbf{X}^* \equiv \mathbf{X}^*(\boldsymbol{\sigma}^*)$ that satisfy a certain condition (EXG) (cf. Proposition 5.3).

3.3 Main results for hypergraph stochastic block models

In this section, we apply Theorems 3.5, 3.8, 4.12 in the setting of hypergraph stochastic block models (HSBM). Notably, HSBM significantly generalizes the symmetric block model considered in Section 1. In particular, the results stated in this section immediately imply Theorems 1.1, 1.2, 1.3 whose proofs are proved at the end of this section. To begin with, we define HSBM and the associated inference tasks.

Definition 3.11. (*Hypergraph Stochastic Block Model*) Given $k, q \geq 2$, let $M \equiv (M(i_1, \dots, i_k))_{1 \leq i_1, \dots, i_k \leq q}$ be a symmetric tensor of order $k \geq 2$ with positive entries. That is, for any permutation $\omega \in \overline{S}_q$, $M(i_1, \dots, i_k) = M(\omega(i_1), \dots, \omega(i_k)) > 0$ holds. Also, let $\pi = (\pi_i)_{i \leq q} \in [0, 1]^q$ be a probability vector, which encodes the prior of different communities. Given $n \geq 1$, let $\mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, M, \pi)$ be a random k -uniform hypergraph model with n vertices defined as follows. Every vertex $v \in V$ is assigned a community $\boldsymbol{\sigma}_v^* \in \{1, 2, \dots, q\}$ independently according to $\boldsymbol{\sigma}_v^* \stackrel{i.i.d.}{\sim} \pi$. Given the community structure $\underline{\boldsymbol{\sigma}}^*$, each possible hyperedge (v_1, \dots, v_k) consisting of distinct vertices $v_1, \dots, v_k \in V$ is included independently with probability $M(\boldsymbol{\sigma}_{v_1}^*, \dots, \boldsymbol{\sigma}_{v_k}^*) / \binom{n}{k-1}$.

Note that by restricting to $k = 2$, HSBM specializes to stochastic block models with prior $\pi \in \mathcal{P}([q])$ and connection probabilities encoded by an arbitrary symmetric matrix $M \in \mathbb{R}_{>0}^{q \times q}$. Let the average degree of HSBM denoted by

$$d := \sum_{i_1, \dots, i_k \in [q]} M(i_1, \dots, i_k) \prod_{s=1}^k \pi_{i_s}. \quad (22)$$

Observe that the normalized tensor $M_0 \equiv (M_0(i_1, \dots, i_k))_{1 \leq i_1, \dots, i_k \leq q} := M/d$ must satisfy

$$1 = \sum_{i_1, \dots, i_k \in [q]} M_0(i_1, \dots, i_k) \prod_{s=1}^k \pi_{i_s}. \quad (23)$$

By fixing such M_0 and varying d , the weak recovery threshold $d_*^{\text{H}} \equiv d_*^{\text{H}}(M_0, \pi)$ is defined by

$$d_*^{\text{H}} \equiv d_*^{\text{H}}(M_0, \pi) := \inf \left\{ d > 0 : \text{weak recovery is possible at } d \text{ for } \mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, dM_0, \pi) \right\}.$$

Here, the weak recovery in HSBM is defined analogously to Definition 3.3. That is, we say that the weak recovery is possible at d for $\mathcal{G}^{\text{H}}(n, dM_0, \pi)$ if there exists an $\varepsilon > 0$ and (sequence of) estimators $\hat{\underline{c}} \equiv \hat{\underline{c}}_n(\mathbf{G}_{\text{HSBM}}^*)$ such that Eq. (10) is satisfied.

It is crucial to observe that the weak recovery threshold is trivial if the average degree around every node is not the same. That is, $d_*^{\text{H}}(M_0, \pi) > 0$ holds only if the following is satisfied:

$$1 = \sum_{i_1, \dots, i_{k-1} \in [q]} M_0(i_1, \dots, i_{k-1}, i) \prod_{s=1}^{k-1} \pi_{i_s}, \quad \text{for any } 1 \leq i \leq q. \quad (24)$$

To see this, note that weak recovery is possible at d just by assigning the communities to each vertex based on their degree. Thus, statements such as “For $d < d_*^{\text{H}}(M_0, \pi)$, ...” are not vacant only if Eq. (24) is satisfied. We further note that if Eq. (24) is satisfied, Proposition 4.1 below shows that $d_*^{\text{H}}(M_0, \pi) \geq \frac{1}{k-1}$ holds.

In the case $M_0 = \mathbf{1}_{k,q}$, the all-1-tensor, the random graph model $\mathcal{G}_{\text{ER}}^{\text{H}}(n, d, k) \equiv \mathcal{G}^{\text{H}}(n, d \cdot \mathbf{1}_{k,q}, \pi)$ does not have community structure, and does not depend on π . This is the hypergraph analog of sparse Erdos-Renyi graphs with average degree d . Thus, the hypothesis test corresponding to Eq. (3) which determines the existence of a community structure for a k -uniform hypergraph G is given by

$$H_0 : G \sim \mathcal{G}_{\text{ER}}^{\text{H}}(n, d, k) \quad \text{vs.} \quad H_1 : G \sim \mathcal{G}^{\text{H}}(n, dM_0, \pi) \quad (25)$$

Analogously to Definition 3.4, we say that detection is possible at d if there exists a sequence of tests $(\phi_n(G))_{n \geq 1}$ which achieves vanishing Type 1 and Type 2 errors (cf. Eq. (14)).

3.3.1 Contiguity, point estimation, and mutual information in HSBM

We now specialize Theorems 3.5, 3.8, and Corollary 3.7 to HSBM. To this end, we first define the KS threshold for HSBM, which is simple to state. Given M_0 and π , define the matrix $B \equiv (B(i, j))_{i, j \leq q} \in \mathbb{R}^{q \times q}$ by

$$B(i, j) = \sum_{i_1, \dots, i_{k-2} \in [q]} M_0(i_1, \dots, i_{k-2}, i, j) \pi_j \prod_{s=1}^{k-2} \pi_{i_s}. \quad (26)$$

Note that under the degree condition in Eq. (24), the matrix B is a stochastic matrix. Thus, if we let $\lambda_1, \dots, \lambda_q$ be the eigenvalues of B ordered in decreasing absolute values, then by Perron-Frobenius theorem, we have

$$1 = \lambda_1 > |\lambda_2| \geq \dots \geq |\lambda_q|.$$

Then, the KS threshold for HSBM is defined by

$$d_{\text{KS}}^{\text{H}} \equiv d_{\text{KS}}^{\text{H}}(M_0, \pi) := |\lambda_2|^{-2}. \quad (27)$$

The following definition is the translation of local consistent estimators (cf. Definition 3.6) within HSBM.

Definition 3.12. Given k, q, π, M , consider an estimator $\widehat{M}(G)$, which takes as an input a k -uniform hypergraph G , and outputs a symmetric tensor of order $k \geq 2$ with positive entries. For $\varepsilon > 0$, we say that $\widehat{M}(\cdot)$ is ε -locally-consistent for M at (d, π, M) if for all symmetric tensor M' of order k such that $\|M' - M\|_1 \leq \varepsilon$ holds, we have that as $n \rightarrow \infty$,

$$\widehat{M}(\mathbf{G}_{\text{HSBM}}^*) \xrightarrow{p} M', \quad \text{for } \mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, M', \pi).$$

We reiterate that ε -locally consistency is a weaker notion than standard consistency since the parameter space is restricted to be the ε neighborhood of M . Our results establish that below the weak recovery threshold, even this weaker notion of parameter estimation cannot be achieved.

Theorem 3.13. *Let M_0 be a symmetric tensor of order $k \geq 2$ with positive entries, and π be a q -dimensional probability vector for $q \geq 2$. For average degree d below the weak recovery threshold and the KS threshold $d < d_*^{\text{H}}(M_0, \pi) \wedge d_{\text{KS}}^{\text{H}}(M_0, \pi)$, the following holds.*

- (1) $\mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, dM_0, \pi)$ is mutually contiguous with the Erdos-Renyi hypergraph $\mathbf{G}_{\text{ER}} \sim \mathcal{G}_{\text{ER}}^{\text{H}}(n, d, k)$. Thus, detection is impossible. Moreover, for any fixed $\varepsilon > 0$, there exists no ε -locally-consistent estimator for M .
- (2) The asymptotic per-vertex mutual information between $\mathbf{G}_{\text{HSBM}}^*$ and the community structure $\underline{\sigma}^*$ is given by

$$\lim_{n \rightarrow \infty} \frac{1}{n} I(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) = \frac{d}{k} \cdot \sum_{i_1, \dots, i_k=1}^q M_0(i_1, \dots, i_k) \log(M_0(i_1, \dots, i_k)) \prod_{s=1}^k \pi_{i_s}.$$

On the other hand, suppose the average degree is above the weak recovery threshold $d > d_*^{\text{H}}(M_0, \pi)$ and the condition (MIN) holds. Then, we have

- (3) There exists a sequence of events $(\mathcal{A}_n^*)_{n \geq 1}$, a subsequence $(n_\ell)_{\ell \geq 1}$, and a positive constant $\eta > 0$ not depending on ℓ such that for any $\ell \geq 1$,

$$\mathbb{P}(\mathbf{G}_{\text{ER}} \in \mathcal{A}_{n_\ell}^*) \leq 2e^{-\eta n_\ell}, \quad \text{and} \quad \mathbb{P}(\mathbf{G}_{\text{HSBM}}^* \in \mathcal{A}_{n_\ell}^*) \geq 1 - 2e^{-\eta n_\ell},$$

where $\mathbf{G}_{\text{ER}} \sim \mathcal{G}_{\text{ER}}^{\text{H}}(n_\ell, d, k)$ and $\mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n_\ell, dM_0, \pi)$. Thus, detection is possible along the subsequence $(n_\ell)_{\ell \geq 1}$.

- (4) We have that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} I(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) < \frac{d}{k} \cdot \sum_{i_1, \dots, i_k=1}^q M_0(i_1, \dots, i_k) \log(M_0(i_1, \dots, i_k)) \prod_{s=1}^k \pi_{i_s}.$$

We remark the condition (MIN) within HSBM is translated to the following.

- (MIN): Recall the set \mathcal{R}_π in (15). Consider the function $\mathcal{F}_{M_0} : \mathcal{R}_\pi \rightarrow \mathbb{R}_{\geq 0}$ defined by

$$\mathcal{F}_{M_0}(R) := \sum_{\underline{\sigma}, \underline{\tau} \in [q]^k} M_0(\underline{\sigma}) M_0(\underline{\tau}) \prod_{s=1}^k R(\sigma_s, \tau_s).$$

Then, the $\mathcal{F}_{M_0}(R)$ for $R \in \mathcal{R}_\pi$ is uniquely minimized at $R = \pi \pi^\top$.

Symmetric HSBM corresponds to the uniform prior $\pi = \text{Unif}([q])$ and the tensor M_0 taking at most two values $a, b > 0$, where $M_0(i_1, \dots, i_k) = a$ if $i_1 = \dots = i_k$ and $M_0(i_1, \dots, i_k) = b$, otherwise. The symmetric HSBM clearly satisfies (MIN) since for $R \in \mathcal{R}_{q^{-1}\mathbf{1}}$, it is straightforward to compute

$$\mathcal{F}_{M_0}(R) = 1 + (a - b)^2 \sum_{i=1}^q R(i, i)^k, \quad (28)$$

which is uniquely minimized at $R = q^{-2}\mathbf{1}\mathbf{1}^\top$.

3.3.2 Hypothesis testing in HSBM

Our next result concerns the asymptotic power of the likelihood ratio of the hypothesis test (25). To this end, consider the likelihood ratio

$$\mathcal{L}_n(G) = \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G)}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)},$$

where $\mathbf{G}_{\text{ER}} \sim \mathcal{G}_{\text{ER}}^{\text{H}}(n, d, k)$ and $\mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, dM_0, \pi)$.

Theorem 3.14. *Let M_0 be a symmetric tensor of order $k \geq 2$ with positive entries, and π be a q -dimensional probability vector for $q \geq 2$. For average degree d below the weak recovery threshold and the KS threshold $d < d_*^{\text{H}}(M_0, \pi) \wedge d_{\text{KS}}^{\text{H}}(M_0, \pi)$, the likelihood ratio under the null $\mathbf{G}_{\text{ER}} \sim \mathcal{G}_{\text{ER}}^{\text{H}}(n, d, k)$ converges in distribution to*

$$\mathcal{L}_n(\mathbf{G}_{\text{ER}}) \xrightarrow{d} \mathcal{L}_{\infty} := \prod_{\ell=2+\mathbf{1}\{k=2\}}^{\infty} \frac{(1 + \alpha_{\ell})^{\mathbf{X}_{\ell}}}{\mathbb{E}(1 + \alpha_{\ell})^{\mathbf{X}_{\ell}}}.$$

Here, $(\mathbf{X}_{\ell})_{\ell \geq 2}$ are independent Poisson random variables with mean $\mathbb{E}\mathbf{X}_{\ell} = \frac{1}{2\ell}((k-1)d)^{\ell}$, and the constants $(\alpha_{\ell})_{\ell \geq 2}$ are defined by

$$\alpha_{\ell} := \text{tr}(B^{\ell}) - 1 = \sum_{i=2}^q \lambda_i^{\ell}, \quad (29)$$

where $B \in \mathbb{R}^{q \times q}$ is defined in Eq. (26). Moreover, the random variable $\mathcal{L}_{\infty} \equiv \mathcal{L}_{\infty}(d, M_0, \pi)$ satisfies the following.

- (1) *Below the Kesten-Stigum threshold $d < d_{\text{KS}}^{\text{H}}(M_0, \pi)$, \mathcal{L}_{∞} is well-defined (i.e. the infinite product converges a.s.) and has finite second moment $\mathbb{E}\mathcal{L}_{\infty}^2 < \infty$.*
- (2) *For $\frac{1}{k-1} \leq d < d_{\text{KS}}^{\text{H}}(M_0, \pi)$, the random variable \mathcal{L}_{∞} does not have a point mass. In particular, in the regime $\frac{1}{k-1} \leq d < d_*^{\text{H}}(M_0, \pi) \wedge d_{\text{KS}}^{\text{H}}(M_0, \pi)$, the asymptotic power of the likelihood ratio test for the hypothesis test (3) at significance level $\alpha \in (0, 1)$ is given by $\beta_*(\alpha) \in (0, 1)$, where*

$$\beta_*(\alpha) = \mathbb{E}[\mathcal{L}_{\infty} \mathbf{1}\{\mathcal{L}_{\infty} \geq C_{\alpha}\}]. \quad (30)$$

Here, $C_{\alpha} > 0$ is an arbitrary constant satisfying $\mathbb{P}(\mathcal{L}_{\infty} \geq C_{\alpha}) = \alpha$, whose existence is guaranteed.

We next construct a computationally efficient and most powerful test based on a low-degree polynomial of the adjacency matrix. Given a k -uniform hypergraph G with n nodes, let $X_{\ell}(G)$ count the number of cycles of length ℓ . Here, a cycle of length ℓ is a set of vertices $\{v_1, \dots, v_{\ell}, v_{\ell+1} \equiv v_1\}$, where v_i and v_{i+1} are connected by an hyperedge for all $1 \leq i \leq \ell$. Then, consider the following statistic based on the cycle counts of G :

$$\mathcal{T}_n(G) := \prod_{\ell=2+\mathbf{1}\{k=2\}}^{K_n} (1 + \alpha_{\ell})^{X_{\ell}(G)},$$

where the truncation parameter is chosen so that $1 \ll K_n = O(\log \log n)$.

Theorem 3.15. *Let M_0 be a symmetric tensor of order $k \geq 2$ with positive entries, and π be a q -dimensional probability vector for $q \geq 2$. Also, let $K_n = O(\log \log n)$ and $K_n \rightarrow \infty$ as $n \rightarrow \infty$. For a significance level $\alpha \in (0, 1)$, consider the test $\phi_{n,\alpha}(\cdot)$ which rejects the null H_0 in (25) with probability*

$$\phi_{n,\alpha}(G) := \begin{cases} 1 & \text{if } \mathcal{T}_n(G) > C'_{n,\alpha}; \\ 0 & \text{otherwise.} \end{cases}$$

Here, the constants $C'_{n,\alpha} > 0$ is chosen so that we have

$$\mathbb{P}(\mathcal{T}_n(\mathbf{G}_{\text{ER}}) > C'_{n,\alpha}) \leq \alpha \leq \mathbb{P}(\mathcal{T}_n(\mathbf{G}_{\text{ER}}) \geq C'_{n,\alpha}),$$

where $\mathbf{G}_{\text{ER}} \sim \mathcal{G}_{\text{ER}}^{\text{H}}(n, d, k)$. Then, for $\frac{1}{k-1} \leq d < d_{\text{KS}}^{\text{H}}(M_0, \pi)$, the test $\phi_{n,\alpha}$ achieves the power $\beta_*(\alpha)$ in Eq. (30). That is, under the alternative $\mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, dM_0, \pi)$, we have

$$\mathbb{E}\phi_{n,\alpha}(\mathbf{G}_{\text{HSBM}}^*) \rightarrow \beta_*(\alpha) \quad \text{as } n \rightarrow \infty.$$

In particular, in the regime $\frac{1}{k-1} \leq d < d_*^{\text{H}}(M_0, \pi) \wedge d_{\text{KS}}^{\text{H}}(M_0, \pi)$, the test $\phi_{n,\alpha}(\cdot)$ is asymptotically most powerful for the hypothesis test (3).

For symmetric HSBM, it was shown in [GP23] that the weak recovery threshold equals the KS threshold for $q = 2$ and $k \in \{3, 4\}$. Thus, combining this with our results give the following corollary.

Corollary 3.16. *Consider symmetric HSBM with $q = 2$ communities and $k \in \{3, 4\}$ interactions. Then, Theorems 3.13, 3.14, 3.15 hold with $d_*^{\text{H}}(M_0, \pi) \equiv d_{\text{KS}}^{\text{H}}(M_0, \pi)$. Thus, in these cases, for any average degree d such that $d \geq \frac{1}{k-1}$ and $d \neq d_{\text{KS}}^{\text{H}}(M_0, \pi)$ hold, and any significance level $\alpha \in (0, 1)$, there exists an efficient sequence of statistical tests $\varphi_{n,\alpha}(\cdot)$ that is asymptotically most powerful at level α .*

4 Proof overview

This section provides the proof of Theorems 3.5, 3.8, and Corollary 3.7 while deferring the technical steps to the appendices. Theorems 3.10, 3.13, 3.14, and 3.15 are derived from our results for planted factor models and their proof is deferred to Appendix C.

Notations: The notation \mathbb{E}_π is used for the expectation with respect to $\underline{\sigma} \equiv (\sigma_1, \dots, \sigma_k) \sim \pi^{\otimes k}$, while \mathbb{E}_p is used for the expectation with respect to $\psi \sim p$. Additionally $u = \text{Unif}(V^k)$ denotes the uniform distribution among k tuples of variables, and \mathbb{E}_u denotes the expectation with respect to $\omega \sim u$. These notations can also be used jointly. For instance, $\mathbb{E}_{p,\pi}$ denotes the expectation with respect to $\psi \sim \pi$ and $\underline{\sigma} \sim \pi^{\otimes k}$. We reserve the greek letter ξ to denote

$$\xi := \mathbb{E}_{p,\pi}[\psi(\underline{\sigma})].$$

We denote by $\mathcal{P}(\Omega)$ the set of probability measures on a finite set Ω .

4.1 Proof of Theorem 3.5 and Corollary 3.7

In this section, we prove Theorem 3.5 and its consequences in hypothesis testing (see Corollaries 4.14, 4.16) and point estimation in Corollary 3.7. The proof of Theorem 3.5 and Corollary 3.7 is at the end of Section 4.1.3 and 4.1.5 respectively. To begin with, we consider the following property, which guarantees that the task of weak recovery is non-trivial.

- (SYM) : For any $\psi \in \Psi$, $1 \leq s \leq k$, and $\tau \in [q]$, $\mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_s = \tau] = \mathbb{E}_{p,\pi}[\psi(\underline{\sigma})]$ holds.

Indeed, we prove that if (SYM) is violated, then weak recovery is possible for any $d > 0$ based on estimating communities based on the degree of each variables. On the contrary, if (SYM) is satisfied, then we prove that the task of weak recovery is non-trivial. Specifically, the following result is proven in Section A.

Proposition 4.1. *If (SYM) does not hold for p, π , then weak recovery is possible for any $d > 0$. That is, the violation of (SYM) implies $d_*(p, \pi) = 0$. On the other hand, if (SYM) is satisfied for p, π and at least one of the weight function $\psi \in \Psi$ are not constant functions, then we have $\frac{1}{k-1} \leq d_*(p, \pi) < \infty$.*

Note that Theorem 3.5 and Corollary 3.7 is trivial if $d_*(p, \pi) = 0$. Therefore, by Proposition 4.1, we may safely assume (SYM) throughout the proof.

4.1.1 Generalized Kesten-Stigum threshold

This section defines the KS threshold for the planted factor model, which is a generalization of the uniform prior case $\pi = \text{Unif}([q])$ considered in [COEJ⁺18] and sparse (non-symmetric) block model case considered in [AS18]. For $\psi \in \Psi$, let the matrix $\Phi_\psi \equiv \Phi_\psi(i, j)_{i, j \in [q]} \in \mathbb{R}^{q \times q}$ be defined by

$$\Phi_\psi(i, j) := \xi^{-1} \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_1 = i, \sigma_2 = j] \cdot \pi_j. \quad (31)$$

With a slight abuse of notation, we often identify a matrix $A \equiv (A(\tau_1, \tau_2))_{\tau_1, \tau_2 \in [q]}$ with the linear map $x \in \mathbb{R}^q \rightarrow Ax \in \mathbb{R}^q$. With this convention, let the linear operator $\Xi \equiv \Xi_{p, \pi}$ on $\mathbb{R}^q \otimes \mathbb{R}^q$ defined by

$$\Xi \equiv \Xi_p := \mathbb{E}_p[\Phi_\psi \otimes \Phi_\psi],$$

where \otimes denotes the tensor product. In addition, define the linear operator $\Xi_* \equiv \Xi_{*, p, \pi}$ on $\mathbb{R}^q \otimes \mathbb{R}^q$ by

$$\Xi_* \equiv \Xi_{*, p, \pi} := \mathbb{E}_p \left[\left(\Phi_\psi - \mathbf{1}\pi^\top \right) \otimes \left(\Phi_\psi - \mathbf{1}\pi^\top \right) \right],$$

where $\mathbf{1} \in \mathbb{R}^q$ denote the all-1-vector. Here, we identified the matrix $\Phi_\psi - \mathbf{1}\pi^\top \in \mathbb{R}^q$ with the corresponding linear map on \mathbb{R}^q . Let $\langle \cdot, \cdot \rangle_\pi$ denote the unique inner product on $\mathbb{R}^q \otimes \mathbb{R}^q$ that satisfies

$$\langle x_1 \otimes x_2, y_1 \otimes y_2 \rangle_\pi := x_1^\top \text{diag}(\pi) y_1 \cdot x_2^\top \text{diag}(\pi) y_2, \quad (32)$$

where $x_1, x_2, y_1, y_2 \in \mathbb{R}^q$ and $\text{diag}(\pi) \equiv \text{diag}((\pi)_{i \leq q})$. Further, let $\mathcal{S} \subset \mathbb{R}^q \otimes \mathbb{R}^q$ be the subspace defined by

$$\mathcal{S} := \{x \in \mathbb{R}^q \otimes \mathbb{R}^q : \langle x, w \otimes \mathbf{1} \rangle_\pi = \langle x, \mathbf{1} \otimes w \rangle_\pi = 0, \text{ for all } w \in \mathbb{R}^q\}.$$

We denote the projection operator onto the subspace \mathcal{S} by $\text{proj}_{\mathcal{S}}$. In order to define the KS threshold, we need the following lemma whose proof is deferred to Section 6.

Lemma 4.2. *Assuming (SYM), the following holds for the linear operator Ξ on $\mathbb{R}^q \otimes \mathbb{R}^q$.*

- (1) Ξ is self-adjoint on the inner product space $(\mathbb{R}^q \otimes \mathbb{R}^q, \langle \cdot, \cdot \rangle_\pi)$.
- (2) $\Xi(\mathcal{S}) \subset \mathcal{S}$ and $\Xi(\mathcal{S}^\perp) \subset \mathcal{S}^\perp$ hold, where \mathcal{S}^\perp is the orthogonal subspace of \mathcal{S} with respect to the inner product $\langle \cdot, \cdot \rangle_\pi$.
- (3) The composition of Ξ and $\text{proj}_{\mathcal{S}}$ is given by $\Xi \circ \text{proj}_{\mathcal{S}} = \Xi_*$.

Having Lemma 4.2 in hand, we denote the eigenspace of Ξ restricted to the subspace \mathcal{S} by

$$\text{Eig}_{\mathcal{S}}(\Xi) := \{ \lambda \in \mathbb{R} : \Xi x = \lambda x \text{ for some } x \in \mathcal{S} \setminus \{0\} \}. \quad (33)$$

By Lemma 4.2 and the spectral theorem, $\text{Eig}_{\mathcal{S}}(\Xi)$ is non-empty whose non-zero elements equal the non-zero elements of $\text{Eig}(\Xi_*)$, the set of eigenvalues of Ξ_* . The KS threshold is defined as follows.

Definition 4.3. (*Kesten-Stigum threshold*) Let $\lambda_{\text{KS}} \equiv \lambda_{\text{KS}}(p, \pi) := \max_{\lambda \in \text{Eig}_{\mathcal{S}}(\Xi)} |\lambda| = \max_{\lambda \in \text{Eig}(\Xi_*)} |\lambda|$. The Kesten-Stigum (KS) threshold is defined by $d_{\text{KS}} \equiv ((k-1)\lambda_{\text{KS}})^{-1}$.

Notably, the KS threshold for HSBM in Eq. (27) is a special case of Definition 4.3, where the prior p puts all of its mass on a single weight function (or equivalently a tensor) on M .

4.1.2 Second moment of the truncated likelihood ratio

For a factor graph G with n variables and m clauses, the likelihood ratio (LR) for the hypothesis test (13) evaluated at G is given by

$$L(G) \equiv L(G; p, \pi) := \frac{\mathbb{P}(\mathbf{G}^*(n, m, \underline{\sigma}^*) = G)}{\mathbb{P}(\mathbf{G}(n, m) = G)}, \quad (34)$$

where $\mathbf{G}(n, m)$ is the null model in Definition 3.1 and $\mathbf{G}^*(n, m, \underline{\sigma}^*)$ is the planted factor model in Definition 3.2. To show that $\mathbf{G} \equiv \mathbf{G}(n, m)$ and $\mathbf{G}^* \equiv \mathbf{G}^*(n, m)$ are mutually contiguous for $d < d_* \wedge d_{\text{KS}}$, we apply the second moment method to a suitable truncation of the likelihood ratio supplemented with the small subgraph conditioning method [Jan95, Wor99].

A standard observation is that because the randomness of the number of clauses $\mathbf{m} \sim \text{Poi}(dn/k)$ causes the second moment $\mathbb{E}L(\mathbf{G})^2$ to diverge, it is necessary to first condition on $\mathbf{m} = m_n$ for typical values of $(m_n)_{n \geq 1}$. In addition, note that $\mathbb{E}L(\mathbf{G}(n, m)) = 1$ holds for $n, m \geq 1$ by a simple change of measure, thus the second moment method requires to prove that $\mathbb{E}L(\mathbf{G}(n, m_n))^2 = O(1)$ holds.

However, even after conditioning on $\mathbf{m} = m_n$, $\mathbb{E}L(\mathbf{G}(n, m_n))^2 = O(1)$ fails to hold in general for all range of $d < d_* \wedge d_{\text{KS}}$. Indeed, by the results of [BMNN16], even restricted to symmetric block model with 3 or more communities, there is a non-trivial regime below d_* such that the second moment is exponentially large, i.e. $\mathbb{E}L(\mathbf{G}(n, m_n))^2 = e^{\Omega(n)}$.

To overcome such difficulty, we truncate the likelihood ratio $L(\mathbf{G}(n, m_n))$ to ensure that the two samples drawn from the posterior is near-orthogonal. Such truncation is motivated by Theorem 3.10, which is detailed as follows. For a sequence of truncation parameters $(\varepsilon_n)_{n \geq 1}$, let the truncated LR be defined by

$$L^*(G) \equiv L^*(G; \varepsilon_n) := \frac{\mathbb{P}(\mathbf{G}^*(n, m, \underline{\sigma}^*) = G)}{\mathbb{P}(\mathbf{G}(n, m) = G)} \mathbf{1} \left\{ \langle \|R_{\underline{q}^1, \underline{q}^2} - \pi \pi^\top\|_1 \rangle_G \leq \varepsilon_n \right\}, \quad (35)$$

where we recall that $\langle \cdot \rangle_G$ denotes the expectation taken w.r.t. the sampled $\underline{\sigma}^1, \underline{\sigma}^2$ drawn from the posterior (see Eq. (20)). A crucial observation is that by a change of measure, the truncation affects the first moment by

$$\begin{aligned} \mathbb{E}|L(\mathbf{G}(n, m_n)) - L^*(\mathbf{G}(n, m_n))| &= \mathbb{E}L(\mathbf{G}(n, m_n)) \mathbb{1}\{\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 \rangle_{\mathbf{G}(n, m_n)} > \varepsilon_n\} \\ &= \mathbb{P}\left(\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 \rangle_{\mathbf{G}^*(n, m_n)} > \varepsilon_n\right). \end{aligned} \quad (36)$$

Note that if $d < d_*(p, \pi)$ and m_n is sufficiently close to dn/k , Theorem 3.10 suggests that the right hand side tends to 0 by choosing $\varepsilon_n = o_n(1)$ appropriately. Indeed, the following lemma guarantees that the truncation does not affect the first moment below the weak recovery threshold for some $\varepsilon_n = o_n(1)$. Its proof is deferred to Section 6.

Lemma 4.4. *Let $d < d_*$ and $|m_n - dn/k| \leq n^{2/3}$ for $n \geq 1$. Then, there exists a sequence $(\varepsilon_n)_{n \geq 1}$ such that as $n \rightarrow \infty$, $\varepsilon_n \rightarrow 0$ and the following convergence is satisfied.*

$$\mathbb{E}|L(\mathbf{G}(n, m_n)) - L^*(\mathbf{G}(n, m_n))| \rightarrow 0.$$

On the other hand, the truncation in $L^*(G)$ reduces the second moment tremendously. In particular, the following second moment estimate is at the heart of the proof of Theorem 3.5. Its proof is the most technical piece of the paper and it is deferred to Section 6.

Proposition 4.5. *Let $d < d_{\kappa_S}$ and assume the condition (SYM). Then, for any sequences $(m_n)_{n \geq 1}$ and $(\varepsilon_n)_{n \geq 1}$ such that $|m_n - dn/k| \leq n^{2/3}$ and $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$, we have*

$$\mathbb{E}L^*(\mathbf{G}(n, m_n))^2 \leq (1 + o_n(1)) \prod_{\lambda \in \text{Eig}_S(\Xi)} \frac{1}{\sqrt{1 - (k-1)d\lambda}}. \quad (37)$$

4.1.3 Cycles and small subgraph conditioning

Proposition 4.5 guarantees that $L^*(\mathbf{G}(n, m_n))$ is bounded away from zero with uniformly positive probability by the Payley-Zygmund inequality (a.k.a. second moment method). To boost this probability to close to one, we use the second moment method conditioned on the number of small cycles. This is formalized by the *small subgraph conditioning method* developed by [Jan95, Wor99]. In particular, we have the following theorem from [Wor99] (see also [Jan95, Theorem 1] and [JLuR00, Theorem 9.12]).

Theorem 4.6. [Wor99, Theorems 4.3], [JLuR00, Theorem 9.12] *Let $(X_{in})_{i \geq 1}$ be a set of non-negative integer valued random variables indexed by $n \geq 1$, and let Y_n be another non-negative random variable on the same probability space as $(X_{in})_{i \geq 1}$. Suppose that $\mathbb{E}Y_n > 0$ for large enough n , and there exist $\lambda_i > 0$ and $\delta_i \geq -1$ such that the following hold.*

- (A1) $X_{in} \xrightarrow{d} X_{i\infty}$ as $n \rightarrow \infty$, jointly for all i , where $X_{i\infty}$ are independent $\text{Poi}(\lambda_i)$ distributed random variables.
- (A2) For any finite sequence x_1, \dots, x_L of non-negative integers, as $n \rightarrow \infty$,

$$\frac{\mathbb{E}[Y_n \mathbb{1}\{X_{1n} = x_1, \dots, X_{Ln} = x_L\}]}{\mathbb{E}Y_n} \rightarrow \prod_{i=1}^L \frac{((1 + \delta_i)\lambda_i)^{x_i}}{x_i!} e^{-(1+\delta_i)\lambda_i}.$$

$$(A3) \quad \sum_{i \geq 1} \lambda_i \delta_i^2 < \infty.$$

$$(A4) \quad \frac{\mathbb{E}Y_n^2}{(\mathbb{E}Y_n)^2} \leq \exp\left(\sum_{i \geq 1} \lambda_i \delta_i^2\right) + o_n(1) \text{ as } n \rightarrow \infty.$$

Then, we have as $n \rightarrow \infty$ that

$$\frac{Y_n}{\mathbb{E}Y_n} \xrightarrow{d} W \equiv \prod_{i=1}^{\infty} (1 + \delta_i)^{X_{i\infty}} e^{-\lambda_i \delta_i}.$$

Moreover, this and the convergence in (A1) hold jointly.

Remark 4.7. An implicit conclusion of Theorem 4.6 is that W is well-defined as the almost-sure-limit of $W_m \equiv \prod_{i \leq m} (1 + \delta_i)^{X_{i\infty}} e^{-\lambda_i \delta_i}$ as $m \rightarrow \infty$. This can be easily seen by martingale convergence theorem since $(W_m)_{m \geq 1}$ is a martingale with mean 1 and bounded variance by (A3).

In random graph theory, it is common to take $(X_{in})_{i \geq 1}$ in Theorem 4.6 as the number of cycles in the random graph with n nodes. Indeed, the works [COW18, COEJ⁺18, NSS24] applied Theorem 4.6 for various factor graphs by setting $(X_{in})_{i \geq 1}$ as the number of cycles with specific signatures. We consider the following specific notion of cycles with signature ζ , which is essentially the same as the one considered in [COEJ⁺18].

Definition 4.8. (ζ -cycle) A signature of order $\ell \geq 1$ is defined by a family

$$\zeta = (\psi_1, \dots, \psi_\ell, s_1, t_1, \dots, s_\ell, t_\ell),$$

where $\psi_1, \dots, \psi_\ell \in \Psi$, and $s_1, t_1, \dots, s_\ell, t_\ell \in [k]$ that satisfy $s_i \neq t_i$ for $i \leq \ell$. In a factor graph $G = (V, F, E, (\psi_a)_{a \in F})$, for variables $v_{i_1}, \dots, v_{i_\ell} \in V$ and clauses $a_{j_1}, \dots, a_{j_\ell} \in F$, we call $\{v_{i_1}, a_{j_1}, \dots, v_{i_\ell}, a_{j_\ell}\}$ a ζ -cycle if it satisfies the following conditions.

- The indices $i_1, \dots, i_\ell \in [n]$ are distinct such that $i_1 = \min\{i_1, \dots, i_\ell\}$. Similarly, $j_1, \dots, j_\ell \in [m]$ are distinct and $j_1 = \min\{j_1, \dots, j_\ell\}$.
- For $1 \leq h \leq \ell$, the weight function $\psi_{a_{j_h}}$ assigned to the clause a_{j_h} in G is given by $\psi_{a_{j_h}} = \psi_h$.
- For $1 \leq h \leq \ell$, the s_h 'th variable adjacent to the clause a_{j_h} in G is v_{i_h} , i.e. $\delta_{s_h} a_{j_h} = v_{i_h}$. Similarly, for $1 \leq h \leq \ell - 1$, $\delta_{t_h} a_{j_h} = v_{i_{h+1}}$ holds, and $\delta_{t_\ell} a_{j_\ell} = v_{i_1}$.

We let $X_\zeta(G)$ be the number of ζ -cycles in a factor graph G . Also, we let S_ℓ denote the set of signatures ζ of order ℓ , and let $S := \cup_{\ell \geq 1} S_\ell$ be the set of signatures.

The first condition $i_1 = \min\{i_1, \dots, i_\ell\}, j_1 = \min\{j_1, \dots, j_\ell\}$ is to avoid overcounting. Also, note that the set S_ℓ is finite for each $\ell \geq 1$ because we assumed that Ψ is finite. Thus, the set S is countable.

We will take $(X_{in})_{i \geq 1} = (X_\zeta(\mathbf{G}(n, m_n)))_{\zeta \in S}$ and $Y_n = L(\mathbf{G}(n, m_n))$ in Theorem 4.6. To verify the assumptions (A1) and (A2), we describe the asymptotic distribution of $X_\zeta(\mathbf{G}(n, m_n))$ and $X_\zeta(\mathbf{G}^*(n, m_n))$. To this end, we introduce additional notations. Let the matrix $\Phi_{\psi, s, t} \equiv (\Phi_{\psi, s, t}(i, j))_{i, j \in [q]} \in \mathbb{R}^{q \times q}$ for $\psi \in \Psi$ and $s, t \in [k]$ be defined by

$$\Phi_{\psi, s, t}(i, j) := \xi^{-1} \cdot \mathbb{E}_\pi[\psi(\boldsymbol{\sigma}) \mid \boldsymbol{\sigma}_s = i, \boldsymbol{\sigma}_t = j] \cdot \pi_j. \quad (38)$$

Notably, $\Phi_{\psi,1,2} = \Phi_{\psi}$ holds by definition. Moreover, for a signature $\zeta = (\psi_1, \dots, \psi_{\ell}, s_1, t_1, \dots, s_{\ell}, t_{\ell}) \in S_{\ell}$, we let $\Phi_{\zeta} := \prod_{i=1}^{\ell} \Phi_{\psi_i, s_i, t_i} \in \mathbb{R}^{q \times q}$ and define the constants $\lambda_{\zeta}, \lambda_{\zeta}^*$ by

$$\lambda_{\zeta} \equiv \lambda_{\zeta}(k, d, p) := \frac{1}{2\ell} \left(\frac{d}{k}\right)^{\ell} \prod_{i=1}^{\ell} p(\psi_i), \quad \lambda_{\zeta}^* \equiv \lambda_{\zeta}^*(k, d, \pi, p) := \lambda_{\zeta} \cdot \text{tr}(\Phi_{\zeta}). \quad (39)$$

In order to obtain our results for hypergraph stochastic block models, we condition the planted factor model on a specific event \mathcal{G}_n defined as follows. Let the event \mathcal{G}_n consist of factor graphs $G = (V, F, E, (\psi_a)_{a \in F})$ with $|V| = n$ variables such that

- (H1) Every clause is connected to k different variables, i.e. for any $a \in F$, $|\delta a| = k$.
- (H2) For any $a, a' \in F$ with $a \neq a'$, the set of neighbors of a is distinct from a' , i.e. δa does not equal any permutation of $\delta a'$.

By using the moment method, [COEJ⁺18, Proposition 3.12] deduced the following result.

Fact 4.9. [COEJ⁺18, Proposition 3.12] *Assume the condition (SYM). Then, for $|m_n - dn/k| \leq n^{2/3}$, we have the following.*

- (1) $X_{\zeta}(\mathbf{G}(n, m_n)) \xrightarrow{d} X_{\zeta, \infty}$ as $n \rightarrow \infty$, jointly for all $\zeta \in \cup_{\ell \geq 1} S_{\ell}$, where $X_{\zeta, \infty}$ are independent $\text{Poi}(\lambda_{\zeta})$ distributed random variables.
- (2) $X_{\zeta}(\mathbf{G}^*(n, m_n)) \xrightarrow{d} X_{\zeta, \infty}^*$ as $n \rightarrow \infty$, jointly for all $\zeta \in \cup_{\ell \geq 1} S_{\ell}$, where $X_{\zeta, \infty}^*$ are independent $\text{Poi}(\lambda_{\zeta}^*)$ distributed random variables.
- (3) Let $C(G) := \sum_{\zeta \in S_1} X_{\zeta}(G) + \mathbf{1}\{k = 2\} \sum_{\zeta \in S_2} X_{\zeta}(G)$. Then, note that for $k = 2$, the event \mathcal{G}_n equals $\{C(G) = 0\}$, and in general $\mathcal{G}_n \subseteq \{C(G) = 0\}$ holds. For $k \geq 3$, the probability of $\mathbf{G}(n, m_n)$ and $\mathbf{G}^*(n, m_n)$ satisfying (H2) is $O(1/n)$, and we have

$$\begin{aligned} \mathbb{P}(C(\mathbf{G}(n, m_n)) = 0) &= \mathbb{P}(\mathbf{G}(n, m_n) \in \mathcal{G}_n) + O(1/n), \\ \mathbb{P}(C(\mathbf{G}^*(n, m_n)) = 0) &= \mathbb{P}(\mathbf{G}^*(n, m_n) \in \mathcal{G}_n) + O(1/n). \end{aligned}$$

Remark 4.10. We remark that the results of [COEJ⁺18, Proposition 3.12] are only stated for $\widehat{\mathbf{G}}(n, m_n)$ instead of $\mathbf{G}^*(n, m_n)$, $\pi = \text{Unif}([q])$, and when p satisfies an extra condition called (BAL). Here, $\widehat{\mathbf{G}}(n, m_n)$ is a certain ‘tweaked version’ of $\mathbf{G}^*(n, m_n)$ defined in Section 3.2 therein. In particular, the condition (BAL) therein guarantees that $\widehat{\mathbf{G}}(n, m)$ is mutually contiguous with $\mathbf{G}^*(n, m)$ (see [COEJ⁺18, Lemma 3.2]). However, a cursory examination of [COEJ⁺18, Proposition 3.12 and Lemma 8.2] reveals that the claim was proven in two stages: first, they calculate the joint moments of $(X_{\zeta}(\mathbf{G}^*(n, m_n)))_{\zeta \in \cup_{\ell \geq 1} S_{\ell}}$ up to $o(1)$ error (see Eq. (8.10)) to determine its asymptotic distribution and then translate the result to $\widehat{\mathbf{G}}(n, m_n)$ using the condition (BAL). Since the argument of [COEJ⁺18, Proposition 3.12] works verbatim to show Fact 4.9, we omit its proof.

To verify the assumptions (A3) and (A4), we need the following lemma which relates the quantities $\lambda_{\zeta}, \lambda_{\zeta}^*$ in (39) and $\lambda \in \text{Eig}_{\mathcal{S}}(\Xi \circ \Lambda^{\otimes 2})$ in (33). Its proof is deferred to Section 6.

Lemma 4.11. For a signature ζ , define the constant

$$\delta_\zeta := \lambda_\zeta^*/\lambda_\zeta - 1 = \text{tr}(\Phi_\zeta) - 1.$$

Assuming (SYM) and $d < d_{\text{KS}}$, we have that

$$\exp\left(\sum_{\ell \geq 1} \sum_{\zeta \in S_\ell} \lambda_\zeta \delta_\zeta^2\right) = \prod_{\lambda \in \text{Eig}_S(\Xi)} \frac{1}{\sqrt{1 - (k-1)d\lambda}}.$$

By combining our second moment estimate in Proposition 4.5 and small subgraph conditioning method in Theorem 4.6, we obtain the following result.

Theorem 4.12. For average degree d below the weak recovery threshold and the Kesten-Stigum threshold $d < d_* \wedge d_{\text{KS}}$, the likelihood ratio evaluated at the null model $\mathbf{G} \equiv \mathbf{G}(n, \mathbf{m})$ converges in distribution to

$$L(\mathbf{G}) \xrightarrow{d} \mathbf{L}_\infty := \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta, \infty}} e^{-\lambda_\zeta \delta_\zeta} \right\}, \quad (40)$$

where $X_{\zeta, \infty} \sim \text{Poi}(\lambda_\zeta)$ are independent Poisson random variables and $\delta_\zeta \equiv \lambda_\zeta^*/\lambda_\zeta - 1$. Moreover, the convergence in (40) holds jointly with the convergence in Fact 4.9-(1).

Proof. By Proposition 4.1, we assume (SYM) without loss of generality. Fix any sequence $(m_n)_{n \geq 1}$ such that $|m_n - dn/k| \leq n^{2/3}$ holds. We use Theorem 4.6 by setting $Y_n \equiv L^*(\mathbf{G}(n, m_n))$ and $(X_{in})_{i \geq 1} \equiv (X_\zeta(\mathbf{G}(n, m_n)))_{\zeta \in S}$. To this end, we check the assumptions (A1)-(A4). First, Fact 4.9-(1) implies the first assumption (A1). In addition, Fact 4.9-(2) and a simple change of measure guarantee that for a finite sequence of non-negative integers x_1, \dots, x_L , we have

$$\mathbb{E}\left[L(\mathbf{G}(n, m_n)) \mathbb{1}\{X_{\zeta_1} = x_1, \dots, X_{\zeta_L} = x_L\}\right] \longrightarrow \prod_{i=1}^L \mathbb{P}(X_{\zeta_i, \infty}^* = x_i).$$

Meanwhile, Lemma 4.4 shows that the LHS equals $\mathbb{E}[L^*(\mathbf{G}(n, m_n)) \mathbb{1}\{X_{\zeta_1} = x_1, \dots, X_{\zeta_L} = x_L\}]$ up to $o_n(1)$ error. Moreover, $\mathbb{E}L^*(\mathbf{G}(n, m_n)) = 1 - o_n(1)$ holds by Lemma 4.4 since $\mathbb{E}L(\mathbf{G}(n, m_n)) = 1$. Thus, it follows that

$$\frac{\mathbb{E}\left[L^*(\mathbf{G}(n, m_n)) \mathbb{1}\{X_{\zeta_1} = x_1, \dots, X_{\zeta_L} = x_L\}\right]}{\mathbb{E}L^*(\mathbf{G}(n, m_n))} \longrightarrow \prod_{i=1}^L \mathbb{P}(X_{\zeta_i, \infty}^* = x_i),$$

which verifies (A2). (A3) is immediate from Lemma 4.11. Furthermore, for $d < d_{\text{KS}}$, Proposition 4.5 shows that

$$\frac{\mathbb{E}\left(L^*(\mathbf{G}(n, m_n))\right)^2}{\left(\mathbb{E}L^*(\mathbf{G}(n, m_n))\right)^2} \leq (1 + o_n(1)) \mathbb{E}\left(L^*(\mathbf{G}(n, m_n))\right)^2 \leq (1 + o_n(1)) \prod_{\lambda \in \text{Eig}_S(\Xi \circ \Lambda^{\otimes 2})} \frac{1}{\sqrt{1 - (k-1)d\lambda}},$$

where the first inequality holds since $\mathbb{E}L^*(\mathbf{G}(n, m_n)) = 1 - o_n(1)$. Since Lemma 4.11 shows that the final product equals $(1 + o_n(1)) \exp\left(\sum_{\ell \geq 1} \sum_{\zeta \in S_\ell} \lambda_\zeta \delta_\zeta^2\right)$, (A4) also holds. Therefore, by Theorem 4.6, we have for $d < d_{\text{KS}}$ that

$$\frac{L^*(\mathbf{G}(n, m_n))}{\mathbb{E}L^*(\mathbf{G}(n, m_n))} \xrightarrow{d} \mathbf{L}_\infty \equiv \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta, \infty}} e^{-\lambda_\zeta \delta_\zeta} \right\}, \quad (41)$$

and this convergence holds jointly with the convergence of $X_\zeta(\mathbf{G}(n, m_n))$ in Fact 4.9-(1). For $d < d_* \wedge d_{\text{KS}}$, Lemma 4.4 allows us to translate this convergence to the convergence of $L(\mathbf{G}(n, m_n))$:

$$L(\mathbf{G}(n, m_n)) \xrightarrow{d} \mathbf{L}_\infty,$$

where the convergence is also joint with the convergence in Fact 4.9-(1). Finally, since $|\mathbf{m} - dn/k| \leq n^{2/3}$ holds with probability tending to one by a Chernoff bound, this concludes the proof. \square

Proof of Theorem 3.5. Let $d < d_* \wedge d_{\text{KS}}$. First, observe that the likelihood ratio of $\mathbf{G}^* \equiv \mathbf{G}^*(n, \mathbf{m})$ and $\mathbf{G} \equiv \mathbf{G}(n, \mathbf{m})$ is given $L(\cdot)$ since for a factor graph G with n variables and m clauses, we have

$$\frac{\mathbb{P}(\mathbf{G}^* = G)}{\mathbb{P}(\mathbf{G} = G)} = \frac{\mathbb{P}(\mathbf{G}^*(n, m) = G)\mathbb{P}(\mathbf{m} = m)}{\mathbb{P}(\mathbf{G}(n, m) = G)\mathbb{P}(\mathbf{m} = m)} = L(G). \quad (42)$$

Moreover, Theorem 4.12 guarantees that for $d < d_* \wedge d_{\text{KS}}$, the convergence $L(\mathbf{G}) \xrightarrow{d} \mathbf{L}_\infty$ holds as $n \rightarrow \infty$. Note that $\mathbb{E}\mathbf{L}_\infty = 1$ holds since $X_{\zeta, \infty} \sim \text{Poi}(\lambda_\zeta)$ are independent Poisson random variables. In addition, $\mathbf{L}_\infty > 0$ holds a.s.. Therefore, Le Cam's first lemma (see e.g. [Vaa98, Lemma 6.4] or [JLuR00, Proposition 9.49]) implies that $\mathbf{G} \equiv \mathbf{G}(n, \mathbf{m})$ and $\mathbf{G}^* \equiv \mathbf{G}^*(n, \mathbf{m})$ are mutually contiguous. \square

4.1.4 Consequences in hypothesis testing

We next state the consequences of likelihood ratio convergence in hypothesis testing. To do so, we need the the following result, which shows that the distribution of \mathbf{L}_∞ contains a point mass depending on the average degree $d > 0$. Its proof is deferred to Section B.

Lemma 4.13. *In the subcritical regime $d \in (0, \frac{1}{k-1})$, the support of the random variable \mathbf{L}_∞ equals the countable set*

$$\left\{ z : z = \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} (1 + \delta_\zeta)^{x_\zeta} e^{-\lambda_\zeta \delta_\zeta} \text{ for some } x = (x_\zeta)_{\ell \geq 1, \zeta \in S_\ell} \text{ with finitely many non-zero elements.} \right\}. \quad (43)$$

On the contrary, in the critical or subcritical regime $d \in [\frac{1}{k-1}, d_{\text{KS}})$, the random variable \mathbf{L}_∞ does not have a point mass.

As an immediate consequence of Theorem 4.12 and Lemma 4.13, we have the following corollary.

Corollary 4.14. *In the subcritical regime $d \in (0, \frac{1}{k-1})$, the asymptotic power of non-randomized likelihood ratio test for the hypothesis test (13) at significance level $\alpha \in (0, 1)$ is given by*

$$\beta_*(\alpha, d, p, \pi) := \mathbb{E}[\mathbf{L}_\infty \mathbf{1}\{\mathbf{L}_\infty > C_\alpha\}]. \quad (44)$$

Here, $C_\alpha > 0$ is an arbitrary constant that satisfy

$$\mathbb{P}(\mathbf{L}_\infty > C_\alpha) \leq \alpha \leq \mathbb{P}(\mathbf{L}_\infty \geq C_\alpha).$$

In the critical or supercritical regime $d \in [\frac{1}{k-1}, d_{\text{KS}})$, the asymptotic power of (possibly randomized) likelihood ratio test for the hypothesis test (13) at significance level $\alpha \in (0, 1)$ is given by $\beta_(\alpha, d, p, \pi)$.*

Next, we consider computationally efficient hypothesis using the following statistic based on ζ -cycles:

$$T_n(\mathbf{G}) := \prod_{1 \leq \ell \leq K_n} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_\zeta(\mathbf{G})} e^{-\lambda_\zeta \delta_\zeta} \right\},$$

where the truncation parameter is taken so that $1 \ll K_n = O(\log \log n)$. Then, we prove the following lemma in Section B by utilizing Fact 4.9.

Lemma 4.15. *Under the condition (SYM), let $d < d_{\text{KS}}$ and $1 \ll K_n = O(\log \log n)$. Then, under the null model $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$,*

$$T_n(\mathbf{G}) \xrightarrow{d} \mathbf{L}_\infty \equiv \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta, \infty}} e^{-\lambda_\zeta \delta_\zeta} \right\},$$

where $X_{\zeta, \infty} \sim \text{Poi}(\lambda_\zeta)$ are independent Poisson random variables. Under the planted model $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$,

$$T_n(\mathbf{G}^*) \xrightarrow{d} \mathbf{L}_\infty^* \equiv \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta, \infty}^*} e^{-\lambda_\zeta \delta_\zeta} \right\},$$

where $X_{\zeta, \infty}^* \sim \text{Poi}(\lambda_\zeta^*)$ are independent Poisson random variables.

We note that the random variable \mathbf{L}_∞^* is well-defined for $d < d_{\text{KS}}$, i.e. the infinite product over $\ell \geq 1$ converges almost surely, since \mathbf{L}_∞ is well-defined (cf. Remark 4.7) with mean 1 from which \mathbf{L}_∞^* can be obtained by a change of measure (see Eq. (47) below). As a corollary, we have the following result.

Corollary 4.16. *Under the condition (SYM), let $1 \leq d < d_{\text{KS}}$ and $1 \ll K_n = O(\log \log n)$. At significance level $\alpha \in (0, 1)$, consider the non-randomized test $\varphi_{n, \alpha}(\cdot)$ which rejects the null \mathbf{H}_0 in Eq. (13) with probability*

$$\varphi_{n, \alpha}(\mathbf{G}) := \begin{cases} 1 & \text{if } T_n(\mathbf{G}) > C_{n, \alpha}; \\ 0 & \text{otherwise.} \end{cases}$$

Here, the constants $C_{n, \alpha} > 0$ is chosen so that we have

$$\mathbb{P}(T_n(\mathbf{G}) > C_{n, \alpha}) \leq \alpha \leq \mathbb{P}(T_n(\mathbf{G}) \geq C_{n, \alpha}), \quad (45)$$

where $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$. Then, for $1 \leq d < d_{\text{KS}}$, the test $\varphi_{n, \alpha}$ achieves the power $\beta_*(\alpha, d, p, \pi)$ defined in Eq. (44). That is, under the alternative $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$, we have

$$\mathbb{E}\varphi_{n, \alpha}(\mathbf{G}^*) \rightarrow \beta_*(\alpha) \quad \text{as } n \rightarrow \infty.$$

In particular, in the regime $1 \leq d < d_* \wedge d_{\text{KS}}$, the test $\varphi_{n, \alpha}(\cdot)$ is asymptotically most powerful for the hypothesis test (3).

Proof. Let $1 \leq d < d_{\text{KS}}$. By passing to a subsequence, let $\tilde{C}_\alpha \in [-\infty, \infty]$ be the limit of $(C_{n, \alpha})_{n \geq 1}$. We first claim that $\mathbb{P}(\mathbf{L}_\infty > \tilde{C}_\alpha) = \alpha$. To see this, note that by Lemma 4.15 and Skorohod embedding, there exists a coupling such that $T_n(\mathbf{G})$ converges to \mathbf{L}_∞ almost surely. Moreover, $\mathbb{P}(\mathbf{L}_\infty = \tilde{C}_\alpha) = 0$ holds by Lemma 4.13. Thus, it follows that

$$\mathbb{1}\{T_n(\mathbf{G}) > C_{n, \alpha}\} \rightarrow \mathbb{1}\{\mathbf{L}_\infty > \tilde{C}_\alpha\} \quad \text{as } n \rightarrow \infty \text{ almost surely.}$$

Similarly, $\mathbb{1}\{T_n(\mathbf{G}) \geq C_{n,\alpha}\}$ converges to $\mathbb{1}\{\mathbf{L}_\infty > \tilde{C}_\alpha\} = \mathbb{1}\{\mathbf{L}_\infty \geq \tilde{C}_\alpha\}$ a.s.. Hence, by dominated convergence theorem, sending $n \rightarrow \infty$ in (45) yield that

$$\alpha = \lim_{n \rightarrow \infty} \mathbb{P}(T_n(\mathbf{G}) > C_{n,\alpha}) = \lim_{n \rightarrow \infty} \mathbb{P}(T_n(\mathbf{G}) \geq C_{n,\alpha}) = \mathbb{P}(\mathbf{L}_\infty > \tilde{C}_\alpha), \quad (46)$$

which proves our first claim. Next, note that for any Borel measurable set $A \subset \mathbb{R}$, we have

$$\begin{aligned} \mathbb{E}[\mathbf{L}_\infty \mathbb{1}\{\mathbf{L}_\infty \in A\}] &\equiv \sum_{x=(x_\zeta)_\zeta} \mathbb{1}\left\{\prod_{\zeta} (1 + \delta_\zeta)^{x_\zeta} e^{-\lambda_\zeta \delta_\zeta} \in A\right\} \prod_{\zeta} \left\{(1 + \delta_\zeta)^{x_\zeta} e^{-\lambda_\zeta \delta_\zeta} \mathbb{P}(X_{\zeta,\infty} = x_\zeta)\right\} \\ &= \sum_{x=(x_\zeta)_\zeta} \mathbb{1}\left\{\prod_{\zeta} (1 + \delta_\zeta)^{x_\zeta} e^{-\lambda_\zeta \delta_\zeta} \in A\right\} \prod_{\zeta} \frac{(\lambda_\zeta^*)^{x_\zeta}}{x_\zeta!} e^{-\lambda_\zeta^*} \\ &= \mathbb{P}(\mathbf{L}_\infty^* \in A), \end{aligned} \quad (47)$$

where the second equality holds since $1 + \delta_\zeta \equiv \lambda_\zeta^*/\lambda_\zeta$. In particular, the distribution of \mathbf{L}_∞^* does not have a point mass by Lemma 4.13. Moreover, by repeating the argument in deriving (46), Lemma 4.15 shows that

$$\lim_{n \rightarrow \infty} \mathbb{E}\varphi_{n,\alpha}(\mathbf{G}^*) = \lim_{n \rightarrow \infty} \mathbb{P}(T_n(\mathbf{G}^*) > C_{n,\alpha}) = \mathbb{P}(\mathbf{L}_\infty^* > \tilde{C}_\alpha).$$

Meanwhile, the final term equals $\mathbb{P}(\mathbf{L}_\infty^* > \tilde{C}_\alpha) = \mathbb{E}[\mathbf{L}_\infty \mathbb{1}\{\mathbf{L}_\infty > \tilde{C}_\alpha\}]$ by (47). Since $\mathbb{P}(\mathbf{L}_\infty > \tilde{C}_\alpha) = \alpha$ holds by our first claim, the final term further equals $\beta_*(\alpha, d, p, \pi)$, which concludes the proof. \square

4.1.5 Proof of Corollary 3.7

We establish the impossibility of point estimation in Corollary 3.7 based on the mutual contiguity in Theorem 3.5 and a *resampling argument*. Given a factor graph $G = (V, F, E, \{\psi_a\}_{a \in F})$ and $\gamma \in [0, 1]$, consider the γ -resampling procedure as follows. For each clause $a \in F$, toss a coin independently with probability γ and let $(Z_a)_{a \in F} \stackrel{i.i.d.}{\sim} \text{Ber}(\gamma)$. If $Z_a = 1$, then independently resample the neighborhood δa and the factor ψ_a of the clause a from the ‘‘null distribution’’:

$$\mathbb{P}(\delta a = (v_1, \dots, v_k), \psi_a = \psi) = \frac{p(\psi)}{n^k}, \quad (48)$$

and if $Z_a = 0$, keep $(\delta a, \psi_a)$ unchanged. We denote by G_γ the resulting factor graph obtained from G by the γ -resampling procedure. For instance, \mathbf{G}_γ^* denotes the factor graph obtained from the planted model $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$ by the γ -resampling procedure. Note that since γ fraction of clauses are resampled according to the null distribution, the probability distribution of \mathbf{G}_γ^* only gets closer to that of the null model \mathbf{G} as γ becomes large. Indeed, in the extreme case $\gamma = 1$, \mathbf{G}_1^* is distributed the same as \mathbf{G} . Thus, the family $(\mathbf{G}_\gamma^*)_{\gamma \in [0,1]}$ is a continuous interpolation of the planted and the null model.

Our next result, which is used for the proof of Corollary 3.7, shows that \mathbf{G}_γ^* is contiguous to another planted model described as follows. Let $\Psi = \{\psi_1, \dots, \psi_T\}$ and $p \in \mathcal{P}(\Psi)$. Then, let Ψ_γ and $p_\gamma \sim \mathcal{P}(\Psi_\gamma)$ be their γ -modifications defined by

$$\Psi_\gamma := \{\psi_{1,\gamma}, \dots, \psi_{T,\gamma}\}, \quad \text{where} \quad \psi_{t,\gamma}(\cdot) := (1 - \gamma)\psi_t(\cdot) + \gamma\xi, \quad (49)$$

and $p_\gamma(\psi_{t,\gamma}) := p(\psi_t)$ for $1 \leq t \leq T$. Here, we recall the notation $\xi \equiv \mathbb{E}_{p,\pi}[\psi(\underline{\sigma})]$. The proof of the following lemma is deferred to Section B.

Lemma 4.17. *Suppose that the priors $p \in \mathcal{P}(\Psi)$ and $\pi \in \mathcal{P}([q])$ satisfy (SYM), and consider the planted model $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$. Then, for $\gamma \in [0, 1]$, the γ -resampled factor graph \mathbf{G}_γ^* is mutually contiguous to the planted distribution with γ -modified weight functions $\mathcal{G}_{\text{plant}}(n, d, p_\gamma, \pi)$.*

Proof of Corollary 3.7. Fix $\varepsilon > 0$ and $d < d_* \wedge d_{\text{KS}}$. By Proposition 4.1, w.l.o.g. we assume (SYM).

We first argue that the γ -resampled graph \mathbf{G}_γ^* of the planted model $\mathbf{G}^* \equiv \mathbf{G}^*(n, \mathbf{m}, \underline{\sigma}^*) \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$ is mutually contiguous to the null model $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$ for any fixed $\gamma \in [0, 1]$. To see this, recall that \mathbf{G}^* has $\mathbf{m} \sim \text{Poi}(dn/k)$ clauses. By Poisson thinning, \mathbf{G}_γ^* has $\mathbf{m}_1 \sim \text{Poi}((1-\gamma)dn/k)$ clauses distributed as Eq. (7) and \mathbf{m}_2 clauses distributed as Eq. (48). Thus, the likelihood ratio (a.k.a. Radon-Nikodym derivative) of \mathbf{G}_γ^* and $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p)$ is the same as the likelihood ratio of $(\mathbf{G}^*)' \sim \mathcal{G}_{\text{plant}}(n, (1-\gamma)d, p, \pi)$ and $\mathbf{G}' \sim \mathcal{G}_{\text{null}}(n, (1-\gamma)d, p)$. Noting that $(1-\gamma)d < d_*(p, \pi) \wedge d_{\text{KS}}$ holds, combining this equality between likelihood ratios with Theorem 3.5 proves that \mathbf{G}_γ^* is mutually contiguous to \mathbf{G} .

Hence, by Lemma 4.17, the planted distribution with γ -modified weight functions $\mathcal{G}_{\text{plant}}(n, d, p_\gamma, \pi)$ is mutually contiguous to the null model \mathbf{G} for any $\gamma \in [0, 1]$. Note that by taking $\gamma > 0$ small enough compared to ε , the weights $\psi_{t,\gamma}$ defined in (49) satisfy $\|\psi_{t,\gamma} - \psi_t\|_\infty \leq \varepsilon$ for any $1 \leq t \leq T$. Therefore, the mutual contiguity of $\mathcal{G}_{\text{plant}}(n, d, p_\gamma, \pi)$ and $\mathcal{G}_{\text{plant}}(n, d, p_{\gamma'}, \pi)$ for sufficiently small $\gamma \neq \gamma'$ imply the non-existence of ε -locally consistent estimator of Ψ at (d, p, π) (cf. Definition 3.6). \square

4.2 Proof of Theorem 3.8

The key ingredient to prove Theorem 3.8 is to analyze the (normalized) log likelihood ratio evaluated at the planted model $\frac{1}{n} \log L(\mathbf{G}^*)$. Its relationship between the KL divergence and the mutual information is given by the following lemma.

Lemma 4.18. *We have the equality $D_{\text{KL}}(\mathbf{G}^* \parallel \mathbf{G}) = \mathbb{E} \log L(\mathbf{G}^*)$. Moreover, the normalized mutual information can be approximated by*

$$\frac{1}{n} I(\mathbf{G}^*, \underline{\sigma}^*) = -\frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) + \frac{d}{k} \cdot \mathbb{E}_{p,\pi} \left[\frac{\psi(\underline{\sigma})}{\xi} \log \left(\frac{\psi(\underline{\sigma})}{\xi} \right) \right] + o_n(1).$$

Thus, to prove Theorem 3.8, it suffices to analyze $\frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*)$. By borrowing terminology from statistical physics, we call $\frac{1}{n} \log L(\mathbf{G}^*)$ the *free energy* since $L(G)$ can be interpreted as a partition function by the equality

$$L(G) \equiv \frac{\mathbb{P}(\mathbf{G}^*(n, \mathbf{m}, \underline{\sigma}^*) = G)}{\mathbb{P}(\mathbf{G}(n, \mathbf{m}) = G)} = \sum_{\underline{\sigma} \in [q]^V} \frac{\psi_G(\underline{\sigma})}{\mathbb{E}[\psi_{\mathbf{G}(n, \mathbf{m})}(\underline{\sigma})]} \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\sigma}).$$

Because the weight functions $\psi \in \Psi$ are strictly positive, it is well-known that the free energies concentrate tightly around their expectation (see e.g. [COKPZ18, Lemma 3.3]).

Fact 4.19. *There exists a constant $c = c(\Psi) > 0$ that only depends on the set of weight functions Ψ such that the following hold. For any $\varepsilon > 0$, we have*

$$\mathbb{P} \left(\left| \frac{1}{n} \log L(\mathbf{G}^*) - \frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) \right| \geq \varepsilon \right) \leq e^{-c n \varepsilon^2}.$$

Proof. Note that if we have two factor graphs G and G' , which differ only in a single clause, then $|n^{-1} \log L(G) - n^{-1} \log L(G')| \leq C$ holds, where $C \equiv C(\Psi) > 0$. Since conditional on $\underline{\sigma}^*$, the clauses in \mathbf{G}^* are independent, the desired inequality is immediate from Azuma-Hoeffding inequality applied to the Doob martingale w.r.t. the clause revealing filtration. \square

The next step is to calculate derivative of the expected free energy w.r.t. d . Recall that $\langle \cdot \rangle_G$ denotes the expectation with respect to samples $(\underline{\sigma}^\ell)_{\ell \geq 1} \stackrel{i.i.d.}{\sim} \mu_G$ from the posterior. If there are only one sample, i.e. $\ell = 1$, we omit the superscript and write $\langle f(\underline{\sigma}^1) \rangle_G \equiv \langle f(\underline{\sigma}) \rangle_G$ for simplicity.

Proposition 4.20. *For any $d > 0$ and $n \geq 1$, the derivative of the expected free energy is given by*

$$\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) = \frac{1}{k} \cdot \mathbb{E}_{\mathbf{G}^*, p, u} \left[\left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p, u}[\psi'(\underline{\sigma}_{\omega'})]} \right\rangle_{\mathbf{G}^*} \log \left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p, u}[\psi'(\underline{\sigma}_{\omega'})]} \right\rangle_{\mathbf{G}^*} \right], \quad (50)$$

where the expectation $\mathbb{E}_{\mathbf{G}^*, p, u}$ is with respect to \mathbf{G}^* , $\psi \sim p$, and $\omega \sim u \equiv \text{Unif}(V^k)$. The expectation $\mathbb{E}_{p, u}$ in the denominator is with respect to $\psi' \sim p$ and $\omega' \sim u$. Furthermore, we can approximate

$$\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) = \frac{1}{k} \cdot \mathbb{E}_{\mathbf{G}^*, p, u} \left[\frac{\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*}}{\xi} \cdot \log \left(\frac{\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*}}{\xi} \right) \right] + O_{k, q, \Psi}(n^{-1/3}), \quad (51)$$

where $f = O_{k, q, \Psi}(g)$ if there exists a constant C that only depends on k, q, Ψ such that $|f| \leq Cg$.

The first equality (51) can be interpreted as the analog of I-MMSE relation from information theory [GWSV11] within the planted factor model. By appealing to Theorem 3.10, we prove that below the weak recovery threshold $d < d_*$, the RHS of (51) tends to 0 as $n \rightarrow \infty$, whereas for $d > d_*$ it is uniformly positive along a subsequence under the (MIN) condition. As a consequence, we prove the following Proposition, which plays a crucial role in the proof of Theorem 3.8.

Proposition 4.21. *The following holds.*

(1) *For any $n \geq 1$ and $d > 0$, we have $\frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) \geq 0$ and $\mathbb{E} \log L(\mathbf{G}^*) \geq 0$.*

(2) *If $d < d_*$, then we have*

$$\frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

(3) *Assume that the condition (MIN) holds. Then, for any $\varepsilon > 0$, there exists $\eta \equiv \eta(\varepsilon) > 0$ and $n_0 \equiv n_0(\varepsilon)$ not depending on n nor $d > 0$ such that if $\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] \geq \frac{1}{q} + \varepsilon$ holds for some estimator $\hat{\sigma} \equiv \hat{\sigma}_n(\mathbf{G}^*)$ and $n \geq n_0$, then $\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) \geq \eta$ holds.*

(4) *If $d > d_*$ and (MIN) holds, then there exists a constant $\eta > 0$ such that*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) \geq \eta > 0$$

The proofs of Lemma 4.18 and Propositions 4.20, 4.21 are deferred to Section 7. Here, we prove that they imply Theorem 3.8.

Proof of Theorem 3.8. The first statement is an immediate consequence of Lemma 4.18 and Proposition 4.21-(2) while the second statement follows from Lemma 4.18 and Proposition 4.21-(4).

To prove the third statement, assume the condition (MIN). Suppose that for some $d_0 > 0$ and $\varepsilon > 0$, $\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] \geq \frac{1}{q} + \varepsilon$ holds for some estimator $\hat{\sigma} \equiv \hat{\sigma}_n(\check{\mathbf{G}}^*)$ where $\check{\mathbf{G}}^* \sim \mathcal{G}_{\text{plant}}(n, d_0, p, \pi)$. Then, note that for all $d > d_0$, $\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] \geq \frac{1}{q} + \varepsilon$ holds for some estimator $\hat{\sigma} \equiv \hat{\sigma}_n(\mathbf{G}^*)$ where $\mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n, d, p, \pi)$. This is because starting from \mathbf{G}^* , subsampling each clause independently with probability d_0/d yields a sample drawn from $\mathcal{G}_{\text{plant}}(n, d_0, p, \pi)$ by Poisson thinning. Thus, if we let $\eta \equiv \eta(\varepsilon)$ and $n_0 \equiv n_0(\varepsilon)$ be as in Proposition 4.21-(3), then $\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) \geq \eta$ holds for all $d > d_0$ and $n \geq n_0$. Since in general $\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) \geq 0$ holds by Proposition 4.21-(1) and η does not depend on d , it follows that for all $n \geq n_0$,

$$\frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) \geq \eta(d - d_0) \equiv \eta_d > 0.$$

Then, consider the event

$$\mathcal{A}_n^* := \{G : L(G) \geq e^{\eta_d n^2}\}.$$

Then by Fact 4.19, there exists a constant $c > 0$ such that for $n \geq n_0$,

$$\mathbb{P}(\mathbf{G}^* \in \mathcal{A}_n^*) \geq \mathbb{P}\left(\left|\frac{1}{n} \log L(\mathbf{G}^*) - \frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*)\right| \leq \frac{\eta}{2}\right) \geq 1 - e^{-c\eta_d^2}.$$

Meanwhile, note that $\mathbb{E}L(\mathbf{G}) = 1$ holds since $L(\cdot)$ is Radon-Nikodym derivative of the law of \mathbf{G}^* and \mathbf{G} . Thus, we have by Markov's inequality that for any $n \geq 1$,

$$\mathbb{P}(\mathbf{G} \in \mathcal{A}_n^*) = \mathbb{P}(L(\mathbf{G}) \geq e^{\eta_d n^2}) \leq e^{-\eta_d n^2}.$$

Therefore, by adjusting η appropriately to account for small $n < n_0$, the desired claim (18) holds for any $n \geq 1$ under the stated assumptions, concluding the proof. \square

5 Equivalent notions of weak recovery

In this section, we prove Theorem 3.10. We will prove a generalized statement, where $\mathbf{G}^* \equiv \mathbf{G}^*(n, \mathbf{m}, \underline{\sigma}^*, p, \pi)$ is replaced by an arbitrary random variable $\mathbf{X}^* \equiv \mathbf{X}_n^*$ defined on the same probability space as $\underline{\sigma}^*$ and satisfy a certain condition (EXG). This generalization is stated in Proposition 5.3, which will be useful for the proof of Lemma 4.4, but it is interesting in its own right. We first specify the task of weak recovery with respect to \mathbf{X}_n^* , extending Definition 3.3. Recall the definition of $A(\underline{\sigma}^1, \underline{\sigma}^2)$ in (11).

Definition 5.1. Let $(\mathbf{X}_n^*)_{n \geq 1}$ be a sequence of random variables which are defined on the same probability space as $\underline{\sigma}^*$ and take values in a probability space \mathcal{X} . We say that weak recovery is possible for $(\mathbf{X}_n^*)_{n \geq 1}$ if there exists $\varepsilon > 0$ and an estimator (i.e. a measurable function) $\hat{\sigma} \equiv \hat{\sigma}(\mathbf{X}_n^*)$ that takes as an input \mathbf{X}_n^* and returns $\hat{\sigma} \in [q]^V$ such that $\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] \geq \frac{1}{q} + \varepsilon$ for large enough $n \geq 1$. Otherwise, we say that the weak recovery is impossible for $(\mathbf{X}_n^*)_{n \geq 1}$.

Given ‘data’ \mathbf{X}_n^* , we define the ‘posterior’ $\mu_X \in \mathcal{P}([q]^V)$ as

$$\mu_X(\underline{\sigma}) = \mathbb{P}(\underline{\sigma}^* = \underline{\sigma} \mid \mathbf{X}^* = X), \quad \underline{\sigma} \in [q]^V. \quad (52)$$

We denote by $\langle \cdot \rangle_X$ the expectation with respect to samples $(\sigma^\ell)_{\ell \geq 1} \stackrel{i.i.d.}{\sim} \mu_X$ from from the posterior.

The condition we impose on the data $(\mathbf{X}_n^*)_{n \geq 1}$ is

- Exchangeability: for any $n \geq 1$ and any pairs of distinct variables (u_1, v_1) and (u_2, v_2) with $u_i \neq v_i, i = 1, 2$,

$$\text{(EXG)} : \left(\mathbb{P}(\sigma_{u_1}^* = i, \sigma_{v_1}^* = j \mid \mathbf{X}_n^*) \right)_{i,j \in [q]} \stackrel{d}{=} \left(\mathbb{P}(\sigma_{u_2}^* = i, \sigma_{v_2}^* = j \mid \mathbf{X}_n^*) \right)_{i,j \in [q]} \quad (53)$$

By the symmetry of variables, the planted factor model satisfies the property (EXG) as verified by the next lemma.

Lemma 5.2. *For any sequence of $(m_n)_{n \geq 1}$, if we let $\mathbf{X}_n^* = \mathbf{G}^*(n, m_n)$, then the sequence of random variables (\mathbf{X}_n^*) satisfy (EXG). In particular, $\mathbf{X}_n^* = \mathbf{G}^*$ satisfies (EXG).*

Proof. Fix two sets of distinct variables (u_i, v_i) , $i = 1, 2$. We show that for a continuous and bounded test function $f : \mathbb{R}^{q \times q} \rightarrow \mathbb{R}$,

$$\mathbb{E}f\left(\left\{\mathbb{P}(\sigma_{u_1}^* = i, \sigma_{v_1}^* = j \mid \mathbf{G}^*(n, m))\right\}_{i,j \leq q}\right) = \mathbb{E}f\left(\left\{\mathbb{P}(\sigma_{u_2}^* = i, \sigma_{v_2}^* = j \mid \mathbf{G}^*(n, m))\right\}_{i,j \leq q}\right). \quad (54)$$

Since $u_i \neq v_i$, note that there exists a permutation $\kappa : V \rightarrow V$ such that $\kappa(u_1) = u_2$ and $\kappa(v_1) = v_2$. With abuse of notation, denote $\kappa \circ \underline{\sigma}^* \equiv (\sigma_{\kappa(v)}^*)_{v \in V}$ and for a factor graph G , let $\kappa \circ G$ be the factor graph defined by mapping G by a graph isomorphism that permutes the labels of the variables according to κ . That is, $\kappa(v) \sim a$ in $\kappa \circ G$ if and only if $v \sim a$ in G . Then, the key observation is as follows. Recalling the definition of the planted model (cf. Definition 3.2) and our assumption that $p(\psi^\theta) = p(\psi)$ holds for $\theta \in S_\kappa$, we have

$$(\kappa \circ \underline{\sigma}^*, \kappa \circ \mathbf{G}^*(n, m)) \stackrel{d}{=} (\underline{\sigma}^*, \mathbf{G}^*(n, m)).$$

Thus, $\mathbb{P}(\sigma_{u_1}^* = i, \sigma_{v_1}^* = j \mid \mathbf{G}^*(n, m) = G) = \mathbb{P}(\sigma_{u_2}^* = i, \sigma_{v_2}^* = j \mid \kappa \circ \mathbf{G}^*(n, m) = G)$ holds for any factor graph G and $i, j \in [q]$. As a consequence, the LHS of (54) equals

$$\begin{aligned} & \mathbb{E}f\left(\left\{\mathbb{P}(\sigma_{u_1}^* = i, \sigma_{v_1}^* = j \mid \mathbf{G}^*(n, m))\right\}_{i,j \leq q}\right) \\ &= \sum_G f\left(\left\{\mathbb{P}(\sigma_{u_1}^* = i, \sigma_{v_1}^* = j \mid \mathbf{G}^*(n, m) = G)\right\}_{i,j \leq q}\right) \mathbb{P}(\mathbf{G}^*(n, m) = G) \\ &= \sum_G f\left(\left\{\mathbb{P}(\sigma_{u_2}^* = i, \sigma_{v_2}^* = j \mid \kappa \circ \mathbf{G}^*(n, m) = G)\right\}_{i,j \leq q}\right) \mathbb{P}(\kappa \circ \mathbf{G}^*(n, m) = G) \\ &= \mathbb{E}f\left(\left\{\mathbb{P}(\sigma_{u_2}^* = i, \sigma_{v_2}^* = j \mid \mathbf{G}^*(n, m))\right\}_{i,j \leq q}\right), \end{aligned}$$

which concludes the proof of (54). \square

Under the condition (EXG), we then establish the analog of Theorem 3.10.

Proposition 5.3. *Assume that the sequence of random variables $(\mathbf{X}_n^*)_{n \geq 1}$ satisfies the condition (EXG). Then, the impossibility of weak recovery for (\mathbf{X}_n^*) is equivalent to the (analog of) conditions in Theorem 3.10-(b), (c), (d), where one replaces \mathbf{G}^* therein with \mathbf{X}_n^* .*

Proof of Theorem 3.10. This is immediate from Lemma 5.2 and Proposition 5.3. \square

For the rest of this section, we prove Proposition 5.3. In Section 5.1, we show that (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (a) holds. In Section 5.2, we show that (c) \Leftrightarrow (d) holds. Throughout, we abbreviate $\mathbf{X}^* \equiv \mathbf{X}_n^*$. We recall that $\langle \cdot \rangle_X$ denotes the expectation taken w.r.t. the posterior μ_X .

5.1 Triviality of two point correlation

We first substitute the quantity $A(\underline{\sigma}^*, \hat{\sigma})$ in Definition 3.3 by $\tilde{A}(\underline{\sigma}^*, \hat{\sigma})$, which will be more convenient for the proof of (a) \Rightarrow (b) and (c) \Rightarrow (a). For $\underline{\sigma}^\ell \equiv (\sigma_u^\ell)_{u \in V} \in [q]^V$, $\ell = 1, 2$, define

$$\tilde{A}(\underline{\sigma}^1, \underline{\sigma}^2) \equiv \tilde{A}(\underline{\sigma}^1, \underline{\sigma}^2; \pi) := \max_{\Gamma \in S_q} \frac{1}{q} \sum_{i=1}^q \frac{1}{n\pi_i} \sum_{u \in V} \mathbb{1}\{\sigma_u^1 = i, \sigma_u^2 = \Gamma(i)\}. \quad (55)$$

That is $\tilde{A}(\underline{\sigma}^1, \underline{\sigma}^2)$ is defined by replacing $|\{v \in V : \sigma_v^1 = i\}|$ term in (11) by $n\pi_i$. If $\underline{\sigma}^1 = \underline{\sigma}^*$, then such replacement is valid by the concentration of $|\{v \in V : \sigma_v^* = i\}|$ around $n\pi_i$ as shown in the lemma below.

Lemma 5.4. *There exists a constant $C \equiv C_{q,\pi} > 0$ such that for any estimator $\hat{\sigma} \equiv \hat{\sigma}(\mathbf{X}^*)$, we have*

$$|\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] - \mathbb{E}[\tilde{A}(\underline{\sigma}^*, \hat{\sigma})]| \leq Cn^{-1/3}.$$

Proof. Note that we can write $A(\underline{\sigma}^*, \hat{\sigma})$ by

$$A(\underline{\sigma}^*, \hat{\sigma}) = \max_{\Gamma \in S_q} \frac{1}{q} \sum_{i=1}^q \frac{\frac{1}{n\pi_i} \sum_{u \in V} \mathbb{1}\{\sigma_u^* = i, \hat{\sigma}_u = \Gamma(i)\}}{\frac{1}{n\pi_i} \sum_{u \in V} \mathbb{1}\{\sigma_u^* = i\}}.$$

We show that on a w.h.p. event, the denominator above is close to 1: since $\underline{\sigma}^* \equiv (\sigma_v^*)_{v \in V} \stackrel{i.i.d.}{\sim} \pi$, we have by Hoeffding's inequality and a union bound that

$$\mathbb{P}(\mathcal{A}_{\text{bal}}) \geq 1 - q \exp(-2n^{1/3}), \quad \text{where } \mathcal{A}_{\text{bal}} := \left\{ \left| \frac{1}{n} \sum_{u \in V} \mathbb{1}\{\sigma_u^* = i\} - \pi_i \right| \leq n^{-1/3}, \text{ for all } i \in [q] \right\}. \quad (56)$$

Thus, it follows that on the event \mathcal{A}_{bal} , for any estimator $\hat{\sigma}$,

$$1 - (\min_i \pi_i)^{-1} \cdot n^{-1/3} \leq \frac{A(\underline{\sigma}^*, \hat{\sigma})}{\tilde{A}(\underline{\sigma}^*, \hat{\sigma})} \leq 1 + (\min_i \pi_i)^{-1} \cdot n^{-1/3}.$$

Further, $\tilde{A}(\underline{\sigma}^1, \underline{\sigma}^2) \in [0, (\min_i \pi_i)^{-1}]$ holds, so it follows that on the event \mathcal{A}_{bal} ,

$$|A(\underline{\sigma}^*, \hat{\sigma}) - \tilde{A}(\underline{\sigma}^*, \hat{\sigma})| \leq (\min_i \pi_i)^{-2} \cdot n^{-1/3}.$$

Combining with (56) concludes the proof. \square

With Lemma 5.4 in hand, we prove (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (a). We first start with (a) \Rightarrow (b).

Lemma 5.5. *Suppose that weak recovery is impossible for \mathbf{X}^* at d . Then, for any $u \neq v$ and $i, j \in [q]$, $\mathbb{P}(\sigma_u^* = i, \sigma_v^* = j | \mathbf{X}^*) \xrightarrow{p} \pi_i \pi_j$ holds as $n \rightarrow \infty$.*

Proof. We first prove that the impossibility of weak recovery implies $\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*) \xrightarrow{p} \pi_i$ for any $u \in V$ and $i \in [q]$. By (EXG), this implies that for every $u \in V$, $\mathbb{P}(\sigma_u^* = i_0 | \mathbf{X}^*)$ does not converge in probability to π_{i_0} . Now, note that $\max_{i \leq q} \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*)}{\pi_i} \geq 1$, almost surely, since $\sum_{i=1}^q \mathbb{P}(\sigma_u^* = i | \mathbf{X}^*) = 1$ holds. Hence, it follows that

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left[\max_{i \in [q]} \left\{ \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*)}{\pi_i} \right\} \right] > 1, \quad (57)$$

since otherwise, $\max_{i \leq q} \frac{\mathbb{P}(\sigma_u^* = i)}{\pi_i} \xrightarrow{p} 1$ holds, which implies that $\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*) \xrightarrow{p} \pi_i$ for every $i \in [q]$. We now show that (57) implies that weak recovery is possible. To this end, for $X \in \mathcal{X}$, consider the estimator $\hat{\underline{\alpha}}(X) \equiv (\hat{\sigma}_u(X))_{u \in V}$ defined by

$$\hat{\sigma}_u(X) := \arg \max_{i \in [q]} \left\{ \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^* = X)}{\pi_i} \right\}.$$

Then, by taking $\Gamma = \text{Id}$ in the definition of $\tilde{A}(\underline{\sigma}^*, \hat{\underline{\alpha}})$ (cf. (55)), and using (EXG) (cf. (53)), we have

$$\mathbb{E} \left[\tilde{A}(\underline{\sigma}^*, \hat{\underline{\alpha}}(\mathbf{X}^*)) \right] \geq \frac{1}{q} \sum_{i=1}^q \frac{1}{\pi_i} \mathbb{P}(\sigma_u^* = i, \hat{\sigma}_u(\mathbf{X}^*) = i) = \frac{1}{q} \sum_{i=1}^q \mathbb{E} \left[\mathbb{1}(\hat{\sigma}_u(\mathbf{X}^*) = i) \cdot \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*)}{\pi_i} \right], \quad (58)$$

where the last equality is due to tower property and the fact that $\hat{\sigma}_u(\mathbf{X}^*)$ is \mathbf{X}^* -measurable. Note that the RHS equals $q^{-1} \mathbb{E}[\max_{i \leq q} \pi_i^{-1} \mathbb{P}(\sigma_u^* = i | \mathbf{X}^*)]$. Thus, combining with (57), we have

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left[\tilde{A}(\underline{\sigma}^*, \hat{\underline{\alpha}}(\mathbf{X}^*)) \right] \geq \frac{1}{q} \limsup_{n \rightarrow \infty} \mathbb{E} \left[\max_{i \in [q]} \left\{ \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*)}{\pi_i} \right\} \right] > \frac{1}{q}.$$

Therefore, by Lemma 5.4, weak recovery is possible at d , which contradicts our assumption. We thus conclude that $\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*) \xrightarrow{p} \pi_i$ for any $u \in V$ and $i \in [q]$.

Next, we show that the impossibility of weak recovery implies $\mathbb{P}(\sigma_u^* = i, \sigma_v^* = j | \mathbf{X}^*) \xrightarrow{p} \pi_i \pi_j$ for any $u \neq v$ and $i, j \in [q]$ by using a similar argument as above. Assume by contradiction that there exists $u_0 \neq v_0$ and $i_0, j_0 \in [q]$ such that $\mathbb{P}(\sigma_{u_0}^* = i_0, \sigma_{v_0}^* = j_0 | \mathbf{X}^*)$ does not converge in probability to $\pi_{i_0} \pi_{j_0}$. Since we have shown that we must have $\mathbb{P}(\sigma_{v_0}^* = j_0 | \mathbf{X}^*) \xrightarrow{p} \pi_{j_0}$, this implies that $\mathbb{P}(\sigma_{u_0}^* = i_0 | \mathbf{X}^*, \sigma_{v_0}^* = j_0)$ does not converge in probability to π_{i_0} . Then, by the same argument as above, the property (EXG) in (53) shows that for any two district variables $u \neq v$,

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left[\max_{i \in [q]} \left\{ \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*, \sigma_v^* = j_0)}{\pi_i} \right\} \right] > 1, \quad (59)$$

since otherwise, $\max_{i \in [q]} \frac{\mathbb{P}(\sigma_{u_0}^* = i | \mathbf{X}^*, \sigma_{v_0}^* = j_0)}{\pi_i} \xrightarrow{p} 1$ holds, which implies that $\mathbb{P}(\sigma_{u_0}^* = i | \mathbf{X}^*, \sigma_{v_0}^* = j_0) \xrightarrow{p} \pi_i$ holds for any $i \in [q]$. Now, for $X \in \mathcal{X}$, consider the estimator $\hat{\underline{\alpha}}'(X) \equiv (\hat{\sigma}'_u(X))_{u \in V}$ defined by

$$\hat{\sigma}'_u(X) := \arg \max_{i \in [q]} \left\{ \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^* = X, \sigma_v^* = j_0)}{\pi_i} \right\}.$$

Then, taking $\Gamma = \text{Id}$ in the definition of $\tilde{A}(\underline{\sigma}^*, \hat{\underline{\alpha}}'(\mathbf{X}^*))$ and using a tower property as in (58), we have

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left[\tilde{A}(\underline{\sigma}^*, \hat{\underline{\alpha}}'(\mathbf{X}^*)) \right] \geq \frac{1}{q} \limsup_{n \rightarrow \infty} \mathbb{E} \left[\max_{i \in [q]} \left\{ \frac{\mathbb{P}(\sigma_u^* = i | \mathbf{X}^*, \sigma_v^* = j_0)}{\pi_i} \right\} \right] > \frac{1}{q},$$

where the last inequality is by (59). Therefore, by Lemma 5.4, weak recovery is possible at d , which contradicts our assumption. We thus conclude that $\mathbb{P}(\sigma_u^* = i, \sigma_v^* = j | \mathbf{X}^*) \xrightarrow{p} \pi_i \pi_j$ holds for any $u \neq v$ and $i, j \in [q]$. \square

The step (b) \Rightarrow (c) is trivial, thus we next prove (c) \Rightarrow (a)

Lemma 5.6. *Suppose that $\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ and $\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ hold as $n \rightarrow \infty$. Then, weak recovery is impossible at d .*

Proof. By Lemma 5.4, it suffices to show that $\mathbb{E}[\tilde{A}(\underline{\boldsymbol{\sigma}}^*, \hat{\underline{\boldsymbol{\sigma}}}(\mathbf{X}^*))] = q^{-1} + o_n(1)$ holds for any estimator $\hat{\underline{\boldsymbol{\sigma}}}(\mathbf{X}^*)$. For simplicity, denote $\hat{\underline{\boldsymbol{\sigma}}}(\mathbf{X}^*) \equiv (\hat{\sigma}_u)_{u \in V}$. Then, note that we can express

$$\mathbb{E}[\tilde{A}(\underline{\boldsymbol{\sigma}}^*, \hat{\underline{\boldsymbol{\sigma}}}(\mathbf{X}^*))] - \frac{1}{q} = \mathbb{E} \max_{\Gamma \in S_q} \left\{ \frac{1}{q} \sum_{i=1}^q \frac{1}{n\pi_i} \sum_{u \in V} \left(\mathbf{1}(\boldsymbol{\sigma}_u^* = i, \hat{\sigma}_u = \Gamma(i)) - \pi_i \mathbf{1}(\hat{\sigma}_u = \Gamma(i)) \right) \right\},$$

Thus, by triangle inequality, we can bound

$$\begin{aligned} \left| \mathbb{E}[\tilde{A}(\underline{\boldsymbol{\sigma}}^*, \hat{\underline{\boldsymbol{\sigma}}}(\mathbf{X}^*))] - \frac{1}{q} \right| &\leq (q \min_{i \leq q} \pi_i)^{-1} \mathbb{E} \max_{\Gamma \in S_q} \left\{ \sum_{i=1}^q \frac{1}{n} \left| \sum_{u \in V} (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbf{1}(\hat{\sigma}_u = \Gamma(i)) \right| \right\} \\ &\leq (q \min_{i \leq q} \pi_i)^{-1} \frac{1}{n} \mathbb{E} \sum_{i,j=1}^q \left| \sum_{u \in V} (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbf{1}(\hat{\sigma}_u = j) \right|. \end{aligned} \quad (60)$$

By Cauchy-Schwarz, we can bound the right most term by

$$\frac{1}{n} \sum_{i,j=1}^q \mathbb{E} \left| \sum_{u \in V} (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbf{1}(\hat{\sigma}_u = j) \right| \leq \frac{q^2}{n} \left(\sum_{i,j=1}^q \mathbb{E} \left(\sum_{u \in V} (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbf{1}(\hat{\sigma}_u = j) \right)^2 \right)^{1/2}. \quad (61)$$

Note that we can expand the sum inside the square root by

$$\sum_{i,j=1}^q \mathbb{E} \left(\sum_{u \in V} (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbf{1}(\hat{\sigma}_u = j) \right)^2 = \sum_{i=1}^q \mathbb{E} \sum_{u,v \in V} \mathbf{1}(\hat{\sigma}_u = \hat{\sigma}_v) (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) (\mathbf{1}(\boldsymbol{\sigma}_v^* = i) - \pi_i).$$

Since $\hat{\underline{\boldsymbol{\sigma}}}(\mathbf{X}^*) \equiv (\hat{\sigma}_u)_{u \in V}$ is \mathbf{X}^* -measurable, tower property shows that the RHS above equals

$$\begin{aligned} &\mathbb{E} \left[\sum_{u,v \in V} \mathbf{1}(\hat{\sigma}_u = \hat{\sigma}_v) \mathbb{E} \left[\sum_{i=1}^q (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) (\mathbf{1}(\boldsymbol{\sigma}_v^* = i) - \pi_i) \mid \mathbf{X}^* \right] \right] \\ &= \mathbb{E} \sum_{u,v \in V} \mathbf{1}(\hat{\sigma}_u = \hat{\sigma}_v) \left(\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*) - \sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*) - \sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_v^* = i | \mathbf{X}^*) + \sum_{i=1}^q \pi_i^2 \right). \end{aligned}$$

By dividing the sum over $u, v \in V$ into the cases $u = v$ and $u \neq v$, and using triangle inequality, we can bound

$$\begin{aligned} &\sum_{i,j=1}^q \mathbb{E} \left(\sum_{u \in V} (\mathbf{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbf{1}(\hat{\sigma}_u = j) \right)^2 \\ &\leq 2n + n(n-1) \cdot \mathbb{E} \left| \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = \boldsymbol{\sigma}_{v_1}^* | \mathbf{X}^*) - 2 \sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i | \mathbf{X}^*) + \sum_{i=1}^q \pi_i^2 \right|, \end{aligned}$$

where we fixed two distinct variables $u_1 \neq v_1$. Plugging the inequality above into (61), we have

$$\begin{aligned} & \frac{1}{n} \sum_{i,j=1}^q \mathbb{E} \left| \sum_{u \in V} (\mathbb{1}(\boldsymbol{\sigma}_u^* = i) - \pi_i) \cdot \mathbb{1}(\hat{\sigma}_u = j) \right| \\ & \leq q^2 \cdot \left(\mathbb{E} \left| \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = \boldsymbol{\sigma}_{v_1}^* | \mathbf{X}^*) - 2 \sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i | \mathbf{X}^*) + \sum_{i=1}^q \pi_i^2 \right| + \frac{2}{n} \right)^{1/2}. \end{aligned} \quad (62)$$

Since $\mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = \boldsymbol{\sigma}_{v_1}^* | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ and $\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ hold by our assumption, the RHS of (62) converges to 0 as $n \rightarrow \infty$ since the convergent random variables are bounded. This further implies that the RHS of (60) converges to 0. We thus conclude that $\mathbb{E}[\tilde{A}(\hat{\boldsymbol{\sigma}}^*, \hat{\boldsymbol{\sigma}}(\mathbf{X}^*))] = q^{-1} + o_n(1)$ holds, which implies that the weak recovery is impossible at d by Lemma 5.4. \square

5.2 Triviality of the overlap

Next, we establish (c) \Leftrightarrow (d). The proof of (c) \Rightarrow (d) resembles the proof of Lemma 5.6.

Lemma 5.7. *Suppose that $\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ and $\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ hold as $n \rightarrow \infty$. Then, $\mathbb{E} \langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top\|_1 \rangle_{\mathbf{X}^*} \rightarrow 0$ holds as $n \rightarrow \infty$.*

Proof. By Cauchy-Schwarz inequality, we can bound

$$\mathbb{E} \langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top\|_1 \rangle_{\mathbf{X}^*} \leq q \cdot \left(\sum_{i,j=1}^q \mathbb{E} \langle (R_{\underline{\sigma}^1, \underline{\sigma}^2}(i,j) - \pi_i \pi_j)^2 \rangle_{\mathbf{X}^*} \right)^{1/2}. \quad (63)$$

Recalling the definition of $R_{\underline{\sigma}^1, \underline{\sigma}^2}(i,j)$ in (21), we can expand

$$\begin{aligned} \sum_{i,j=1}^q \mathbb{E} \langle (R_{\underline{\sigma}^1, \underline{\sigma}^2}(i,j) - \pi_i \pi_j)^2 \rangle_{\mathbf{X}^*} &= \frac{1}{n^2} \sum_{u,v \in V} \sum_{i,j=1}^q \mathbb{E} \langle \mathbb{1}(\sigma_u^1 = i, \sigma_v^1 = j, \sigma_u^2 = i, \sigma_v^2 = j) \rangle_{\mathbf{X}^*} \\ &\quad - \frac{2}{n} \sum_{u \in V} \sum_{i,j=1}^q \pi_i \pi_j \mathbb{E} \langle \mathbb{1}(\sigma_u^1 = i, \sigma_u^2 = j) \rangle_{\mathbf{X}^*} + \sum_{i,j=1}^q \pi_i^2 \pi_j^2. \end{aligned}$$

Recalling that $\underline{\sigma}^1, \underline{\sigma}^2$ are independent samples from the posterior $\mu_{\mathbf{X}^*}$, we can simplify the RHS above by using the property (EXG) in (53) as

$$\frac{1}{n} + \frac{n-1}{n} \cdot \mathbb{E} \left(\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*) \right)^2 - 2 \cdot \mathbb{E} \left(\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*) \right)^2 + \left(\sum_{i=1}^q \pi_i^2 \right)^2,$$

where $u \neq v$ are distinct variables. By our assumption, $\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*)$ and $\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*)$ converge in probability to $\sum_{i=1}^q \pi_i^2$, which further implies L^2 convergence since they are bounded. Therefore, we have $\sum_{i,j=1}^q \mathbb{E} \langle (R_{\underline{\sigma}^1, \underline{\sigma}^2}(i,j) - \pi_i \pi_j)^2 \rangle_{\mathbf{X}^*} \rightarrow 0$, and combining with (63) concludes the proof. \square

The implication (d) \Rightarrow (c) is a consequence of the following lemma, which is a generalization of [COKPZ18, Lemma 4.8] to non-uniform priors and is built on [BCO16, Corollary 2.2].

Lemma 5.8. For all $\varepsilon > 0$ and $q \geq 2$, there exists $\delta \equiv \delta(\varepsilon, q) > 0$ and $n_0 \equiv n_0(\varepsilon, q)$ such that if $n \geq n_0(\varepsilon, q)$, the following holds. Suppose that a probability measure $\mu \in \mathcal{P}([q]^n)$ and a probability vector on $[q]$, $\pi \in \mathbb{R}^q$, $\sum_{i=1}^q \pi_i = 1$, $\pi_i \geq 0$, satisfy

$$\left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_{\mathbb{F}}^2 \right\rangle_{\mu} < \delta, \quad (64)$$

where $\langle \cdot \rangle_{\mu}$ is the expectation with respect to independent samples $\underline{\sigma}^1, \underline{\sigma}^2 \sim \mu$, $R_{\underline{\sigma}^1, \underline{\sigma}^2}$ is defined in (21), and $\|\cdot\|_{\mathbb{F}}$ denotes the Frobenius norm. Then, we have

$$\frac{1}{n^2} \sum_{u, v \in [n]} \|\mu_{u, v} - \pi^{\otimes 2}\|_{\text{TV}} < \varepsilon,$$

where $\mu_{u, v}$ denotes the marginal distribution of (σ_u, σ_v) when $\underline{\sigma} \equiv (\sigma_u)_{u \in [n]} \sim \mu$.

Proof. The following is a restatement of [BCO16, Corollary 2.2]: for all $\varepsilon_0 > 0$, there exists $\eta \equiv \eta(\varepsilon_0, q)$ and $n_0 = n_0(\varepsilon_0, q)$ such that for every $n \geq n_0$ and $\mu \in \mathcal{P}([q]^n)$, the following holds. There exists a decomposition S_0, S_1, \dots, S_N of $[q]^n$ such that it satisfies

- (C1): $\mu(S_0) \leq \varepsilon_0$ and for any $1 \leq k \leq N$, $\mu(S_k) \geq \eta$.
- (C2): For $1 \leq k \leq N$ and $u, v \in [n]$, denote by $\mu_{u, v}(\cdot | S_k)$ the marginal distribution of (σ_u, σ_v) , where $\underline{\sigma} = (\sigma_u)_{u \in V}$ is drawn from the conditional distribution $\mu(\cdot | S_k) \equiv \mu(\cdot \cap S_k) / \mu(S_k)$. Similarly, denote by $\mu_u(\cdot | S_k)$ the marginal distribution of σ_u , where $\underline{\sigma} \sim \mu(\cdot | S_k)$. Then, for any $1 \leq k \leq N$,

$$\frac{1}{n^2} \sum_{u, v \in [n]} \|\mu_{u, v}(\cdot | S_k) - \mu_u(\cdot | S_k) \otimes \mu_v(\cdot | S_k)\|_{\text{TV}} < \varepsilon_0.$$

We proceed by taking advantage of such decomposition guaranteed by [BCO16, Corollary 2.2]. Note that it suffices to prove the lemma for small enough $\varepsilon > 0$, so we fix $\varepsilon \in (0, 1/100q)$. Given such $\varepsilon > 0$, we take ε_0 and δ small enough depending only on ε and q whose precise value is determined below. Suppose that for a probability measure $\mu \in \mathcal{P}([q]^n)$ satisfies (64). Given μ , consider the decomposition S_0, S_1, \dots, S_N of $[q]^n$, which satisfies (C1), (C2) above, and denote by $\langle \cdot \rangle_k$ the average w.r.t. $\underline{\sigma}^1, \underline{\sigma}^2 \stackrel{i.i.d.}{\sim} \mu(\cdot | S_k)$. Then, by our assumption (64) and since $\mu(S_k) \geq \eta \equiv \eta(\varepsilon_0, q)$ holds by (C1), we have for any $1 \leq k \leq N$,

$$\left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_{\mathbb{F}}^2 \right\rangle_k < \frac{\delta}{\eta^2}. \quad (65)$$

Recalling the definition of $R_{\underline{\sigma}^1, \underline{\sigma}^2}$ in (21), we can expand the LHS by

$$\left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_{\mathbb{F}}^2 \right\rangle_k = \sum_{i, j=1}^q \left(\frac{1}{n^2} \sum_{u, v \in [n]} \mu_{u, v}(i, i | S_k) \mu_{u, v}(j, j | S_k) - \frac{2\pi_i \pi_j}{n} \sum_{u \in [n]} \mu_u(i | S_k) \mu_u(j | S_k) + \pi_i^2 \pi_j^2 \right). \quad (66)$$

Note that by (C2), we can replace $\mu_{u, v}(\cdot | S_k)$ with $\mu_u(\cdot | S_k) \mu_v(\cdot | S_k)$ with error depending on ε_0 :

$$\frac{1}{n^2} \sum_{i, j=1}^q \sum_{u, v \in [n]} \mu_{u, v}(i, i | S_k) \mu_{u, v}(j, j | S_k) \geq \frac{1}{n^2} \sum_{i, j=1}^q \sum_{u, v \in [n]} \mu_u(i | S_k) \mu_v(i | S_k) \mu_u(j | S_k) \mu_v(j | S_k) - 4\varepsilon_0,$$

thus plugging the above bound into (66) and combining with (65) gives

$$\sum_{i,j=1}^q \left(\frac{1}{n} \sum_{u \in [n]} \mu_u(i | S_k) \mu_v(j | S_k) - \pi_i \pi_j \right)^2 < \frac{\delta}{\eta^2} + 4\varepsilon_0.$$

Thus, by Cauchy-Schwarz, we have that

$$\sum_{i,j=1}^q \left| \frac{1}{n} \sum_{u \in [n]} \mu_u(i | S_k) \mu_v(j | S_k) - \pi_i \pi_j \right| \leq q^2 \cdot \left(\frac{\delta}{\eta^2} + 4\varepsilon_0 \right)^{1/2} \leq \varepsilon^3, \quad (67)$$

where in the last inequality we took $\varepsilon_0 \equiv \frac{\varepsilon^6}{8q^4}$ and $\delta \equiv \frac{\eta^2 \varepsilon^6}{2q^4}$ for $\eta \equiv \eta(\varepsilon_0, q)$. Observe that restricting the sum over i, j in (67) to $i = j$, and pushing the sum over j inside the absolute value by triangle inequality shows that

$$\left(\sum_{i=1}^q \left| \frac{1}{n} \sum_{u \in [n]} \mu_u(i | S_k)^2 - \pi_i^2 \right| \right) \vee \left(\sum_{i=1}^q \left| \frac{1}{n} \sum_{u \in [n]} \mu_u(i | S_k) - \pi_i \right| \right) \leq \varepsilon^3.$$

Thus, by triangle inequality,

$$\sum_{i=1}^q \frac{1}{n} \sum_{u \in [n]} (\mu_u(i | S_k) - \pi_i)^2 = \sum_{i=1}^q \left(\frac{1}{n} \sum_{u \in [n]} \mu_u(i | S_k)^2 - \pi_i^2 \right) - 2 \sum_{i=1}^q \pi_i \left(\frac{1}{n} \sum_{u \in [n]} \mu_u(i | S_k) - \pi_i \right) \leq 3\varepsilon^3.$$

Then, it follows from Cauchy-Schwarz that

$$\frac{1}{n} \sum_{i=1}^q \sum_{u \in [n]} \left| \mu_u(i | S_k) - \pi_i \right| \leq (3q\varepsilon^3)^{1/2} < \frac{\varepsilon}{4}, \quad (68)$$

where the last inequality holds since $\varepsilon < (100q)^{-1}$. Consequently, combining with (C2), we have

$$\begin{aligned} \frac{1}{n^2} \sum_{u,v \in [n]} \left\| \mu_{u,v}(\cdot | S_k) - \pi \pi^\top \right\|_{\text{TV}} &\leq \varepsilon_0 + \frac{1}{n^2} \sum_{u,v \in [n]} \left\| \mu_u(\cdot | S_k) \otimes \mu_v(\cdot | S_k) - \pi \pi^\top \right\|_{\text{TV}} \\ &\leq \varepsilon_0 + \frac{2}{n} \sum_{u \in [n]} \left\| \mu_u(\cdot | S_k) - \pi \right\|_{\text{TV}} < \frac{\varepsilon}{2}, \end{aligned} \quad (69)$$

where the last inequality is due to (68). Therefore, by triangle inequality, we have that

$$\begin{aligned} \frac{1}{n^2} \sum_{u,v \in [n]} \left\| \mu_{u,v} - \pi^{\otimes 2} \right\|_{\text{TV}} &\leq \frac{1}{n^2} \sum_{u,v \in [n]} \sum_{k=0}^N \left\| \mu_{u,v}(\cdot | S_k) - \pi^{\otimes 2} \right\|_{\text{TV}} \cdot \mu(S_k) \\ &\stackrel{(a)}{\leq} \mu(S_0) + \frac{\varepsilon}{2} \sum_{k=1}^N \mu(S_k) \stackrel{(b)}{<} \varepsilon_0 + \frac{\varepsilon}{2} < \varepsilon, \end{aligned}$$

where (a) is due to (69) and (b) is due to (C1). This concludes the proof. \square

With Lemma 5.8 in hand, the step (d) \Rightarrow (c) is straightforward.

Lemma 5.9. *Suppose that $\mathbb{E}\langle\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1\rangle_{\mathbf{X}^*} \rightarrow 0$ as $n \rightarrow \infty$. Then, $\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ and $\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ hold as $n \rightarrow \infty$.*

Proof. By Markov's inequality, $\|R_{\underline{\sigma}^1, \underline{\sigma}^2}(i, j) - \pi_i\pi_j\| \xrightarrow{p} 0$ under $\mathbb{E}\mu_{\mathbf{X}^*}^{\otimes 2}$ for any $i, j \in [q]$. Since $\|R_{\underline{\sigma}^1, \underline{\sigma}^2}(i, j) - \pi_i\pi_j\| \leq 1$, convergence in probability further implies L^2 convergence. Summing over $i, j \in [q]$, we have $\mathbb{E}\langle\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_F^2\rangle_{\mathbf{X}^*} \rightarrow 0$. Hence, Lemma 5.8 further implies that

$$\frac{1}{n^2} \sum_{u, v \in V} \left\| \mathbb{P}(\boldsymbol{\sigma}_u^* = \cdot, \boldsymbol{\sigma}_v^* = \cdot | \mathbf{X}^*) - \pi^{\otimes 2} \right\|_{\text{TV}} \xrightarrow{p} 0.$$

Note that the LHS above is at most 1, thus above further implies L^1 convergence. Moreover,

$$\frac{1}{n^2} \sum_{u, v \in V} \mathbb{E} \left\| \mathbb{P}(\boldsymbol{\sigma}_u^* = \cdot, \boldsymbol{\sigma}_v^* = \cdot | \mathbf{X}^*) - \pi^{\otimes 2} \right\|_{\text{TV}} = \sum_{i, j=1}^q \mathbb{E} \left| \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i, \boldsymbol{\sigma}_{v_1}^* = j | \mathbf{X}^*) - \pi_i\pi_j \right| + O(n^{-1}), \quad (70)$$

where we used the property (EXG) in the equality above for distinct variables $u_1 \neq v_1$. Therefore, we have that $\mathbb{P}(\boldsymbol{\sigma}_u^* = \boldsymbol{\sigma}_v^* | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$ and $\sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_u^* = i | \mathbf{X}^*) \xrightarrow{p} \sum_{i=1}^q \pi_i^2$. \square

Finally, we prove a non-asymptotic version of the implication (d) \Rightarrow (a), which is used for the proof of Proposition 4.21 (see Lemma 7.4).

Lemma 5.10. *For any $\varepsilon > 0$, there exists $\eta \equiv \eta(\varepsilon, q, \pi) > 0$ and $n_0 \equiv n_0(\varepsilon, q, \pi)$ such that the following holds. Suppose that $n \geq n_0$ and there exists an estimator $\hat{\boldsymbol{\sigma}} \equiv \hat{\boldsymbol{\sigma}}(\mathbf{X}^*)$ such that $\mathbb{E}[A(\boldsymbol{\sigma}^*, \hat{\boldsymbol{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds. Then, we have $\mathbb{E}\langle\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1\rangle_{\mathbf{X}^*} \geq \eta$.*

Proof. Throughout, we write $C \equiv C(q, \pi) > 0$ for a positive constant that only depends on q, π that may differ from line to line. Also, we fix $\varepsilon > 0$ and write $n_0 \equiv n_0(\varepsilon, q, \pi)$ for an integer that only depends on ε, q, π . Suppose that $\mathbb{E}[A(\boldsymbol{\sigma}^*, \hat{\boldsymbol{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds for some $\hat{\boldsymbol{\sigma}} \equiv \hat{\boldsymbol{\sigma}}(\mathbf{X}^*)$. By Lemma 5.4, we have $\mathbb{E}[\tilde{A}(\boldsymbol{\sigma}^*, \hat{\boldsymbol{\sigma}})] \geq \frac{1}{q} + \frac{\varepsilon}{2}$ for $n \geq n_0$. Subsequently, by the inequalities (60) and (62) in the proof of Lemma 5.6, it follows that for $n \geq n_0$

$$C\varepsilon^2 \leq \mathbb{E} \left| \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = \boldsymbol{\sigma}_{v_1}^* | \mathbf{X}^*) - 2 \sum_{i=1}^q \pi_i \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i | \mathbf{X}^*) + \sum_{i=1}^q \pi_i^2 \right|,$$

where $u_1 \neq v_1$ are two distinct variables. By triangle inequality, the RHS is upper bounded by

$$\begin{aligned} & \mathbb{E} \left| \sum_{i=1}^q (\mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i, \boldsymbol{\sigma}_{v_1}^* = i | \mathbf{X}^*) - \pi_i^2) - 2 \sum_{i=1}^q \pi_i (\mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = i | \mathbf{X}^*) - \pi_i) \right| \\ & \leq 3\mathbb{E} \left\| \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = \cdot, \boldsymbol{\sigma}_{v_1}^* = \cdot | \mathbf{X}^*) - \pi^{\otimes 2} \right\|_{\text{TV}}. \end{aligned}$$

Thus, combining the two inequalities above, $\mathbb{E} \left\| \mathbb{P}(\boldsymbol{\sigma}_{u_1}^* = \cdot, \boldsymbol{\sigma}_{v_1}^* = \cdot | \mathbf{X}^*) - \pi^{\otimes 2} \right\|_{\text{TV}} \geq C\varepsilon^2$ holds for $n \geq n_0$. Recalling the equality (70) and using that the total variation distance is at most 1, we have

$$\mathbb{P} \left(\frac{1}{n^2} \sum_{u, v \in V} \left\| \mathbb{P}(\boldsymbol{\sigma}_u^* = \cdot, \boldsymbol{\sigma}_v^* = \cdot | \mathbf{X}^*) - \pi^{\otimes 2} \right\|_{\text{TV}} \geq C\varepsilon^2 \right) \geq C\varepsilon^2.$$

Then by Lemma 5.8, there exists a constant $\delta \equiv \delta(\varepsilon, q, \pi) > 0$ such that

$$\mathbb{P}\left(\left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top \right\|_{\mathbf{F}}^2 \right\rangle_{\mathbf{X}^*} \geq \delta \right) \geq C\varepsilon^2.$$

Finally, note that $\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_{\mathbf{F}}^2 \leq \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1$ holds since the entries of $R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top$ have absolute values at most 1. Therefore, setting $\eta \equiv C\delta\varepsilon^2$ concludes the proof. \square

Proof of Proposition 5.3. This is immediate from combining Lemmas 5.5, 5.6, 5.7, and 5.9. \square

6 Contiguity below the weak recovery threshold

In this section, we prove Lemma 4.2, Proposition 4.5 and Lemma 4.11, which were crucial for the proof of Theorem 3.5. In Section 6.3, we prove Proposition 4.5 by appealing to a central limit theorem. In Section 6.1, we prove Lemmas 4.2 and 4.11.

6.1 Proof of Lemma 4.2 and Lemma 4.11

Throughout, we assume (SYM). We start with the first statement of Lemma 4.2.

Lemma 6.1. *The linear operator Ξ is self-adjoint on the inner product space $(\mathbb{R}^q \otimes \mathbb{R}^q, \langle \cdot, \cdot \rangle_\pi)$.*

Proof. By linearity, it suffices to check $\langle \Xi(e_i \otimes e_j), e_s \otimes e_t \rangle_\pi = \langle e_i \otimes e_j, \Xi(e_s \otimes e_t) \rangle_\pi$ for $i, j, s, t \in [q]$, where $(e_i)_{i \in [q]}$ are standard basis of \mathbb{R}^q . Note that by definition of Ξ and $\langle \cdot, \cdot \rangle_\pi$, we have

$$\langle \Xi(e_i \otimes e_j), e_s \otimes e_t \rangle_\pi = \sum_{\psi \in \Psi} p(\psi) \cdot \langle \Phi_\psi e_i \otimes \Phi_\psi e_j, e_s \otimes e_t \rangle_\pi = \mathbb{E}_p \left[e_i^\top \Phi_\psi^\top \text{diag}(\pi) e_s \cdot e_j^\top \Phi_\psi^\top \text{diag}(\pi) e_t \right].$$

Recalling the condition (5) and the definition of Φ_ψ in (31), note that we have

$$\text{diag}(\pi) \Phi_\psi \stackrel{d}{=} \Phi_\psi^\top \text{diag}(\pi) \quad \text{for } \psi \sim p. \quad (71)$$

Thus, it follows that $\langle \Xi(e_i \otimes e_j), e_s \otimes e_t \rangle_\pi = \langle e_i \otimes e_j, \Xi(e_s \otimes e_t) \rangle_\pi$ holds for $i, j, s, t \in [q]$. \square

Consider the matrix $\Phi \equiv \Phi_p \in \mathbb{R}^{q \times q}$ defined by the average of $\Phi_\psi, \psi \sim p$.

$$\Phi \equiv \Phi_p := \mathbb{E}_p[\Phi_\psi].$$

As before, we often identify the matrix Φ with the linear map $x \rightarrow \Phi x$. Further, consider the inner product $\langle \cdot, \cdot \rangle_{\pi, 1}$ on \mathbb{R}^q defined by $\langle x, y \rangle_{\pi, 1} := x^\top \text{diag}(\pi) y$. The notation 1 is to distinguish $\langle \cdot, \cdot \rangle_{\pi, 1}$ from the inner product $\langle \cdot, \cdot \rangle_\pi$ in (32), which is the tensor product of $\langle \cdot, \cdot \rangle_{\pi, 1}$ with itself. Then, we have the following lemma.

Lemma 6.2. *The linear operator Φ is self-adjoint on the inner product space $(\mathbb{R}^q, \langle \cdot, \cdot \rangle_{\pi, 1})$ such that $\Phi \mathbf{1} = \mathbf{1}$. Further, for $w \in \mathbb{R}^q$, we have that $\Xi(\mathbf{1} \otimes w) = \mathbf{1} \otimes \Phi w$.*

Proof. Note that since $\text{diag}(\pi)\Phi_\psi \stackrel{d}{=} \Phi_\psi^\top \text{diag}(\pi)$ holds for $\psi \sim p$ (cf. (71)), $\text{diag}(\pi)\Phi = \Phi \text{diag}(\pi)$ holds. Thus, for any $x, y \in \mathbb{R}^q$, we have that

$$\langle \Phi x, y \rangle_{\pi, \mathbf{1}} = x^\top \Phi^\top \text{diag}(\pi) y = x^\top \text{diag}(\pi) \Phi y = \langle x, \Phi y \rangle_{\pi, \mathbf{1}}.$$

Further, note that by (SYM), for any $\psi \in \Psi$, we have the equality $\Phi_\psi \mathbf{1} = \mathbf{1}$. Thus, $\Phi \mathbf{1} = \mathbf{1}$ holds and we have for any $w \in \mathbb{R}^q$ that

$$\Xi(\mathbf{1} \otimes w) = \sum_{\psi \in \Psi} p(\psi) \cdot \Phi_\psi \mathbf{1} \otimes \Phi_\psi w = \sum_{\psi \in \Psi} p(\psi) \cdot \mathbf{1} \otimes \Phi_\psi w = \mathbf{1} \otimes \Phi w,$$

which concludes the proof. \square

Observe that by Lemma 6.2, there exists an orthonormal basis $\{w_1 = \mathbf{1}, w_2, \dots, w_q\}$ of the inner product space $(\mathbb{R}^q, \langle \cdot, \cdot \rangle_{\pi, \mathbf{1}})$ such that w_i 's are eigenvectors of Φ . We denote by λ_i the eigenvalue corresponding to w_i , where $\lambda_1 = 1$. Then, we have the following lemma. Below, $\text{proj}_{\mathcal{S}^\perp}$ and $\text{proj}_{w_i \otimes w_j}$ for $i, j \in [q]$ respectively denotes the projection operator in $(\mathbb{R}^q \otimes \mathbb{R}^q, \langle \cdot, \cdot \rangle_\pi)$ onto the subspace \mathcal{S}^\perp and the subspace spanned by the vector $w_i \otimes w_j$.

Lemma 6.3. *The subspaces \mathcal{S} and \mathcal{S}^\perp is invariant under Ξ . Further, we have that*

$$\Xi \circ \text{proj}_{\mathcal{S}^\perp} = \text{proj}_{\mathbf{1} \otimes \mathbf{1}} + \sum_{i=2}^q \lambda_i \text{proj}_{\mathbf{1} \otimes w_i} + \sum_{i=2}^q \lambda_i \text{proj}_{w_i \otimes \mathbf{1}}.$$

Thus, $\Xi|_{\mathcal{S}^\perp}$ has eigenvalues $\{1, \lambda_2, \lambda_2, \dots, \lambda_q, \lambda_q\}$ counting multiplicities.

Proof. Since $(w_i)_{i \leq q}$ are eigenvectors of Φ , Lemma 6.2 implies that $\mathbf{1} \otimes w_i$ and $w_i \otimes \mathbf{1}$ are eigenvectors of Ξ with λ_i the corresponding eigenvalue. Meanwhile, since $\mathbb{R}^q \otimes \mathbb{R}^q$ is finite dimensional, \mathcal{S}^\perp is given by

$$\mathcal{S}^\perp = \{x \in \mathbb{R}^q \otimes \mathbb{R}^q : x = w \otimes \mathbf{1} \text{ or } x = \mathbf{1} \otimes w \text{ for some } w \in \mathbb{R}^q\}.$$

Thus, the set $\mathcal{W} := \{\mathbf{1} \otimes \mathbf{1}, \mathbf{1} \otimes w_2, \dots, \mathbf{1} \otimes w_q, w_2 \otimes \mathbf{1}, \dots, w_q \otimes \mathbf{1}\}$ is an orthonormal basis of the inner product space $(\mathbb{R}^q \otimes \mathbb{R}^q, \langle \cdot, \cdot \rangle_\pi)$, which are also eigenvectors of Ξ . Hence, the subspace of \mathcal{S}^\perp is invariant under Ξ , which further implies that \mathcal{S} is invariant since Ξ is self-adjoint by Lemma 6.1. Moreover, since \mathcal{W} is the eigenbasis of $\Xi|_{\mathcal{S}^\perp}$, the rest of the claims follow by the spectral theorem. \square

Having Lemmas 6.1, 6.2, and 6.3 in hand, we now prove Lemma 4.2 and Lemma 4.11.

Proof of Lemma 4.2. The first and second statements follow immediately from Lemma 6.1 and Lemma 6.3, thus it remains to prove the third statement. Note that applying the spectral theorem on the self-adjoint operator Φ on $(\mathbb{R}^q, \langle \cdot, \cdot \rangle_{\pi, \mathbf{1}})$ (cf. Lemma 6.2), we have

$$\Phi = \sum_{i=1}^q \lambda_i \text{proj}_{w_i} = \mathbf{1}\pi^\top + \sum_{i=2}^q \lambda_i \text{proj}_{w_i},$$

where proj_{w_i} denotes the projection operator in $(\mathbb{R}^q, \langle \cdot, \cdot \rangle_{\pi, \mathbf{1}})$ onto the subspace spanned by the vector w_i . In particular, the last equality follows since $\text{proj}_{\mathbf{1}} = \mathbf{1}\pi^\top$. Thus, combining with Lemma 6.3, we have

$$\Xi \circ \text{proj}_{\mathcal{S}^\perp} = (\mathbf{1}\pi^\top) \otimes \Phi + \Phi \otimes (\mathbf{1}\pi^\top) - (\mathbf{1}\pi^\top) \otimes (\mathbf{1}\pi^\top).$$

Since \mathcal{S} and \mathcal{S}^\perp are invariant subspaces of Ξ by Lemma 6.3, $\Xi = \Xi \circ \text{proj}_{\mathcal{S}} + \Xi \circ \text{proj}_{\mathcal{S}^\perp}$ holds by the spectral theorem. Thus, it follows that

$$\Xi \circ \text{proj}_{\mathcal{S}} = \Xi - (\mathbf{1}\pi^\top) \otimes \Phi - \Phi \otimes (\mathbf{1}\pi^\top) + (\mathbf{1}\pi^\top) \otimes (\mathbf{1}\pi^\top) = \mathbb{E}_p \left[\left(\Phi - \mathbf{1}\pi^\top \right)^{\otimes 2} \right] \equiv \Xi_*,$$

which concludes the proof. \square

Proof of Lemma 4.11. For each $\ell \geq 1$, we have by definition,

$$\sum_{\zeta \in \mathcal{S}_\ell} \lambda_\zeta \delta_\zeta^2 = \sum_{\psi_1, \dots, \psi_\ell \in \Psi} \sum_{\substack{s_1, t_1, \dots, s_\ell, t_\ell \in [k] \\ s_i \neq t_i, 1 \leq i \leq \ell}} \frac{1}{2^\ell} \left(\frac{d}{k} \right)^\ell \left(\text{tr} \left(\prod_{i=1}^\ell \Phi_{\psi_i, s_i, t_i} \right) - 1 \right)^2 \prod_{i=1}^\ell p(\psi_i).$$

Note that given $s, t \in [k]$ with $s \neq t$, there exists a permutation θ in $[k]$ such that $\theta(1) = s, \theta(2) = t$, and for such θ , we have $\Phi_{\psi, s, t} = \Phi_{\psi^\theta}$, where $\psi^\theta(\sigma_1, \dots, \sigma_k) \equiv \psi(\sigma_{\theta(1)}, \dots, \sigma_{\theta(k)})$. Thus, by our assumption that $p(\psi^\theta) = p(\psi)$ in (5), we can express the equation above by

$$\sum_{\zeta \in \mathcal{S}_\ell} \lambda_\zeta \delta_\zeta^2 = \sum_{\psi_1, \dots, \psi_\ell \in \Psi} \frac{((k-1)d)^\ell}{2^\ell} \left(\text{tr} \left(\prod_{i=1}^\ell \Phi_{\psi_i} \right) - 1 \right)^2 \prod_{i=1}^\ell p(\psi_i) = \frac{((k-1)d)^\ell}{2^\ell} \mathbb{E} \left[\left(\text{tr} \left(\prod_{i=1}^\ell \Phi_{\psi_i} \right) - 1 \right)^2 \right],$$

where $(\psi_i)_{i \leq \ell} \stackrel{i.i.d.}{\sim} p$. Note that we can expand the rightmost term by

$$\mathbb{E} \left[\left(\text{tr} \left(\prod_{i=1}^\ell \Phi_{\psi_i} \right) - 1 \right)^2 \right] = \mathbb{E} \left[\text{tr} \left(\prod_{i=1}^\ell (\Phi_{\psi_i} \otimes \Phi_{\psi_i}) \right) \right] - 2 \mathbb{E} \left[\text{tr} \left(\prod_{i=1}^\ell \Phi_{\psi_i} \right) \right] + 1 = \text{tr}(\Xi^\ell) - 2 \text{tr}(\Phi^\ell) + 1,$$

where the last equality holds since the expectation and trace is exchangeable and ψ_i 's are independent. Since Ξ and Φ are self-adjoint by Lemma 6.1 and Lemma 6.2, we have

$$\text{tr}(\Xi^\ell) - 2 \text{tr}(\Phi^\ell) + 1 = \sum_{\lambda \in \text{Eig}(\Xi)} \lambda^\ell - \sum_{\lambda' \in \text{Eig}(\Phi)} (\lambda')^\ell + 1 = \sum_{\lambda \in \text{Eig}_{\mathcal{S}}(\Xi)} \lambda^\ell,$$

where the last equality follows from Lemma 6.3. Thus, combining the 3 equations in the displays above and summing over $\ell \geq 1$, we have

$$\sum_{\ell \geq 1} \sum_{\zeta \in \mathcal{S}_\ell} \lambda_\zeta \delta_\zeta^2 = \sum_{\ell \geq 1} \sum_{\lambda \in \text{Eig}_{\mathcal{S}}(\Xi)} \frac{((k-1)d)^\ell}{2^\ell} \lambda^\ell = \sum_{\lambda \in \text{Eig}_{\mathcal{S}}(\Xi)} \frac{1}{2} \log(1 - (k-1)d\lambda),$$

where the last equality holds for $d < d_{\text{KS}}$ since $\sum_{\ell \geq 1} \frac{x^\ell}{\ell} = \log(1-x)$ holds for $|x| < 1$. Therefore, exponentiating the equation above concludes the proof. \square

6.2 Proof of Lemma 4.4

Fix $d < d_*$. We first claim that for any $(m_n)_{n \geq 1}$ such that $|m_n - dn/k| \leq n^{2/3}$, weak recovery is impossible for $\mathbf{G}^*(n, m_n)$. We refer to Definition 5.1 for a general definition of weak recovery. To this end, fix such $(m_n)_{n \geq 1}$ and $\eta > 0$ such that $d < d + \eta < d_*$ holds. Then, weak recovery is

impossible at $d + \eta$ by definition of d_* . That is, for any estimator $\hat{\sigma} \equiv \hat{\sigma}(\mathbf{G}^*(n, \mathbf{m})) \in [q]^V$, where $\mathbf{m} \sim \text{Poi}((d + \eta)n/k)$, we have

$$\lim_{n \rightarrow \infty} \mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] = \frac{1}{q}. \quad (72)$$

Note that by definition of the planted model (cf. Definition 3.2), conditional on the event $\mathbf{m} \geq m_n$, the subgraph of $\mathbf{G}^*(n, \mathbf{m})$ formed by excluding $\mathbf{m} - m_n$ clauses is distributed the same as $\mathbf{G}^*(n, m_n)$. Moreover, since $(d + \eta)n/k \geq m_n + \Omega(n)$, it follows from Chernoff bound that $\mathbf{m} > m_n$ holds with high probability. As a consequence, for any estimator $\hat{\sigma} \equiv \hat{\sigma}(\mathbf{G}^*(n, m_n))$ of $\mathbf{G}^*(n, m_n)$, (72) must also hold. Therefore, for any $(m_n)_{n \geq 1}$ such that $|m_n - dn/k| \leq n^{2/3}$, weak recovery is impossible for $\mathbf{G}^*(n, m_n)$.

Note that since $m_n \rightarrow \infty$ as $n \rightarrow \infty$, Lemma 5.2 implies that $(\mathbf{G}^*(n, m_n))_{n \geq 1}$ satisfies the properties (EXG). Thus, it follows from Proposition 5.3 that as $n \rightarrow \infty$,

$$\delta_n := \mathbb{E} \left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_1 \right\rangle_{\mathbf{G}^*(n, m_n)} \rightarrow 0.$$

Let $\varepsilon_n := \delta_n^{1/2}$ and consider $L^*(\mathbf{G}(n, m_n)) \equiv L^*(\mathbf{G}(n, m_n); \varepsilon_n)$. Then, we have by a change of measure (cf. (36)) that

$$\begin{aligned} \mathbb{E} \left[\left| L(\mathbf{G}(n, m_n)) - L^*(\mathbf{G}(n, m_n)) \right| \right] &= \mathbb{P} \left(\left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_1 \right\rangle_{\mathbf{G}^*(n, m_n)} > \varepsilon_n \right) \\ &\leq \varepsilon_n^{-1} \mathbb{E} \left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_1 \right\rangle_{\mathbf{G}^*(n, m_n)} = \delta_n^{1/2}, \end{aligned}$$

where the second equality is due to the change of measure and the inequality is by Markov's inequality. Since $\delta_n \rightarrow 0$ as $n \rightarrow \infty$, this concludes the proof.

6.3 Proof of Proposition 4.5

Throughout this subsection, we fix $d < d_{\text{KS}}$ and assume the condition (SYM). For any factor graph G with n variables and m clauses, note that we can express the square of the truncated likelihood ratio $L^*(G)^2$ as

$$\begin{aligned} L^*(G)^2 &= \sum_{\underline{\sigma}, \underline{\tau} \in [q]^V} L_{\underline{\sigma}, \underline{\tau}}^2(G), \quad \text{where} \\ L_{\underline{\sigma}, \underline{\tau}}^2(G) &:= \frac{\mathbb{P}(\underline{\sigma}^* = \underline{\sigma}, \mathbf{G}^*(n, m) = G) \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\tau}, \mathbf{G}^*(n, m) = G)}{\mathbb{P}(\mathbf{G}(n, m) = G)^2} \mathbb{1} \left\{ \left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_1 \right\rangle_G \leq \varepsilon_n \right\}. \end{aligned} \quad (73)$$

We divide the sum above into the *near-independent* regime where $\underline{\sigma}, \underline{\tau} \in [q]^V$ satisfies $\left\| R_{\underline{\sigma}, \underline{\tau}} - \pi \pi^\top \right\|_1 \leq \sqrt{\varepsilon_n}$, and *correlated* regime where $\left\| R_{\underline{\sigma}, \underline{\tau}} - \pi \pi^\top \right\|_1 > \sqrt{\varepsilon_n}$. Namely, we have $L^*(G)^2 = L_{\text{ind}}^2(G) + L_{\text{cor}}^2(G)$, where

$$L_{\text{ind}}^2(G) = \sum_{\underline{\sigma}, \underline{\tau} \in [q]^V : \left\| R_{\underline{\sigma}, \underline{\tau}} - \pi \pi^\top \right\|_1 \leq \sqrt{\varepsilon_n}} L_{\underline{\sigma}, \underline{\tau}}^2(G), \quad L_{\text{cor}}^2(G) = \sum_{\underline{\sigma}, \underline{\tau} \in [q]^V : \left\| R_{\underline{\sigma}, \underline{\tau}} - \pi \pi^\top \right\|_1 > \sqrt{\varepsilon_n}} L_{\underline{\sigma}, \underline{\tau}}^2(G). \quad (74)$$

An important consequence of the truncation $\mathbb{1} \left\{ \left\langle \left\| R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top \right\|_1 \right\rangle_G \leq \varepsilon_n \right\}$ in $L^*(G)$ is that the contribution to its second moment from the correlated regime is negligible compared to the contribution from the near-independence regime as seen by the following lemma.

Lemma 6.4. For any factor graph G , $L_{\text{cor}}^2(G) \leq \sqrt{\varepsilon_n} \cdot (L^*(G))^2$ holds.

Proof. Recalling the posterior $\mu_G(\underline{\sigma})$ (cf. (19)), note that $L_{\underline{\sigma}, \underline{\tau}}(G) = (L^*(G))^2 \mu_G(\underline{\sigma}) \mu_G(\underline{\tau})$ holds. Thus, it follows that

$$L_{\text{cor}}^2(G) = (L^*(G))^2 \cdot \left\langle \mathbf{1} \left\{ \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top\|_1 > \sqrt{\varepsilon_n} \right\} \right\rangle_G \leq \varepsilon_n^{-1/2} (L^*(G))^2 \left\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top\|_1 \right\rangle_G.$$

where the inequality is due to Markov's inequality. Since $L^*(G) = 0$ holds if $\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top\| \rangle_G > \varepsilon_n$, the RHS can be further bounded by $L_{\text{cor}}^2(G) \leq \sqrt{\varepsilon_n} (L^*(G))^2$, which concludes the proof. \square

Having Lemma 6.4 in hand, we now compute $\mathbb{E} L_{\text{ind}}^2(\mathbf{G}(n, m))$. The first step is to compute $\mathbb{E} L_{\underline{\sigma}, \underline{\tau}}^2(\mathbf{G}(n, m))$.

Lemma 6.5. For any $\underline{\sigma}, \underline{\tau} \in [q]^V$ and $n, m \geq 1$, we have

$$\mathbb{E} L_{\underline{\sigma}, \underline{\tau}}^2(\mathbf{G}(n, m)) \leq \left(\frac{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)]}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] \cdot \mathbb{E}_{p,u}[\psi(\underline{\tau}_\omega)]} \right)^m \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\sigma}) \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\tau}).$$

Proof. Recall that by definition of the planted model (cf. Definition 3.2), $\frac{\mathbb{P}(\mathbf{G}^*(n, m) = G | \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}(n, m) = G)} = \frac{\psi_G(\underline{\sigma})}{\mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\sigma})]}$ holds for a factor graph G with n variables and m clauses. Thus, by dropping the indicator in the definition of $L_{\underline{\sigma}, \underline{\tau}}^2(G)$ in (73), we have

$$\mathbb{E} L_{\underline{\sigma}, \underline{\tau}}^2(\mathbf{G}(n, m)) \leq \frac{\mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\sigma}) \psi_{\mathbf{G}(n, m)}(\underline{\tau})]}{\mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\sigma})] \cdot \mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\tau})]} \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\sigma}) \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\tau}).$$

Meanwhile, for the null model $\mathbf{G}(n, m)$, its weight functions $(\psi_a)_{a \in F}$ and the neighborhoods $(\delta a)_{a \in F}$ are drawn i.i.d. from p and $u := \text{Unif}(V^k)$ respectively. Thus, we have

$$\frac{\mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\sigma}) \psi_{\mathbf{G}(n, m)}(\underline{\tau})]}{\mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\sigma})] \cdot \mathbb{E}[\psi_{\mathbf{G}(n, m)}(\underline{\tau})]} = \left(\frac{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)]}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] \cdot \mathbb{E}_{p,u}[\psi(\underline{\tau}_\omega)]} \right)^m,$$

which concludes the proof. \square

Lemma 6.5 shows that in order to bound $\mathbb{E} L_{\text{ind}}^2(\mathbf{G}(n, m))$, it suffices to compute the expected value of $\left(\frac{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)]}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] \cdot \mathbb{E}_{p,u}[\psi(\underline{\tau}_\omega)]} \right)^m$ under $\underline{\sigma}, \underline{\tau} \stackrel{i.i.d.}{\sim} \pi^{\otimes V}$ conditioned on the event $\|R_{\underline{\sigma}, \underline{\tau}} - \pi \pi^\top\|_1 \leq \sqrt{\varepsilon_n}$, which happens w.h.p. since we assumed $n^{-1} \varepsilon_n \rightarrow \infty$. As we will see next, $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)]$ and $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)]$ can be computed in terms of the normalized overlap matrix $X \equiv X_{\underline{\sigma}, \underline{\tau}} \equiv (X(i, j))_{i, j \leq q} \in \mathbb{R}^{q \times q}$, where

$$X \equiv X_{\underline{\sigma}, \underline{\tau}} := \sqrt{n} \left(R_{\underline{\sigma}, \underline{\tau}} - \pi \pi^\top \right). \quad (75)$$

Note that such normalization guarantees that $\sum_{i, j=1}^q X(i, j) = 0$ and X has $O(1)$ fluctuations: by viewing X as a q^2 dimensional vector and denoting $\mathbf{X}_n \equiv X_{\underline{\sigma}, \underline{\tau}}$ for $\underline{\sigma}, \underline{\tau} \sim \pi^{\otimes n}$, the central limit theorem shows that

$$\mathbf{X}_n \xrightarrow{d} \mathbf{N}(0, \text{diag}(\pi)^{\otimes 2} - (\pi \pi^\top)^{\otimes 2}). \quad (76)$$

Here, for two matrices $A = (A_{i,j})_{i,j \leq q}$ and $B = (B_{i,j})_{i,j \leq q}$, we recall that their tensor product $A \otimes B$ is defined by the $q^2 \times q^2$ matrix with $((i_1, j_1), (i_2, j_2))$ entry $(A \otimes B)_{(i_1, j_1), (i_2, j_2)} \equiv A_{i_1, i_2} B_{j_1, j_2}$.

In addition, for $\psi \in \Psi$, we define the matrix $\widehat{\Phi}_\psi \equiv (\widehat{\Phi}_\psi(i, j))_{i, j \leq q}$ by

$$\widehat{\Phi}_\psi(i, j) := \xi^{-1} \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_1 = i, \sigma_2 = j].$$

Here, we note that $\widehat{\Phi}_\psi \text{diag}(\pi) = \Phi_\psi$ holds by definition. We then define the the matrix $\widehat{\Phi} \in \mathbb{R}^{q \times q}$ by $\widehat{\Phi} := \mathbb{E}_p[\widehat{\Phi}_\psi]$ and define the matrix $\widehat{\Xi} \in \mathbb{R}^{q^2 \times q^2}$ by the averaged tensor product of the matrix $\widehat{\Phi}_\psi$ with itself:

$$\widehat{\Xi} := \mathbb{E}_p[\widehat{\Phi}_\psi \otimes \widehat{\Phi}_\psi].$$

Note that viewing X as a q^2 dimensional vector, quantities such as $\langle \widehat{\Xi} X, X \rangle$ is well-defined, where

$$\langle \widehat{\Xi} X, X \rangle \equiv \sum_{(i_1, j_1), (i_2, j_2) \in [q]^2} \widehat{\Xi}((i_1, j_1), (i_2, j_2)) X(i_1, j_1) X(i_2, j_2).$$

We also recall that $\mathbf{1} \in \mathbb{R}^q$ denoted all 1-vector.

Lemma 6.6. For $\underline{\sigma}, \underline{\tau} \in [q]^V$, we have for $X \equiv X_{\underline{\sigma}, \underline{\tau}}$ that

$$\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)] = \xi^2 \cdot \exp\left(\frac{k(k-1)}{2n} \langle \widehat{\Xi} X, X \rangle + O_{k,q,\Psi}\left(\frac{\|X\|_\infty^3}{n^{3/2}}\right)\right), \quad (77)$$

where we used the notation $f = O_{k,q,\Psi}\left(\frac{\|X\|_\infty^3}{n^{3/2}}\right)$ to indicate that there exists a constant $C \equiv C_{k,q,\Psi}$ which only depends on k, q, Ψ such that $|f| \leq C \frac{\|X\|_\infty^3}{n^{3/2}}$ holds. Further, we have

$$\begin{aligned} \mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] &= \xi \cdot \exp\left(\frac{k(k-1)}{2n} \langle (\widehat{\Phi} \otimes \mathbf{11}^\top) X, X \rangle + O_{k,q,\Psi}\left(\frac{\|X\|_\infty^3}{n^{3/2}}\right)\right), \\ \mathbb{E}_{p,u}[\psi(\underline{\tau}_\omega)] &= \xi \cdot \exp\left(\frac{k(k-1)}{2n} \langle (\mathbf{11}^\top \otimes \widehat{\Phi}) X, X \rangle + O_{k,q,\Psi}\left(\frac{\|X\|_\infty^3}{n^{3/2}}\right)\right). \end{aligned} \quad (78)$$

Proof. We start with the proof of (77). By first calculating the expectation w.r.t. $\omega \sim u := \text{Unif}(V^k)$, we can express $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)]$ by summing the following over $\underline{i} \equiv (i_1, \dots, i_k), \underline{j} \equiv (j_1, \dots, j_k) \in [q]^k$:

$$\begin{aligned} \mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega) \psi(\underline{\tau}_\omega)] &= \sum_{\underline{i}, \underline{j} \in [q]^k} \mathbb{E}_p[\psi(\underline{i}) \psi(\underline{j})] \prod_{s=1}^k (\pi_{i_s} \pi_{j_s} + n^{-1/2} X(i_s, j_s)) \\ &= \sum_{\ell=0}^k \sum_{S \subseteq [k]: |S|=\ell} n^{-\ell/2} \sum_{\underline{i}, \underline{j} \in [q]^k} \mathbb{E}_p[\psi(\underline{i}) \psi(\underline{j})] \prod_{t \in [k] \setminus S} \pi_{i_t} \pi_{j_t} \prod_{s \in S} X(i_s, j_s). \end{aligned}$$

Given ℓ and S with $|S| = \ell$, we divide the sum $\underline{i} \in [q]^k$ into the sum over $\underline{i}_1 \equiv (i_s)_{s \in S} \in [q]^S$ and $\underline{i}_2 \equiv (i_t)_{t \notin S} \in [q]^{[k] \setminus S}$, and similarly for the sum $\underline{j} \in [q]^k$. Note that given $\underline{i}_1, \underline{j}_1 \in [q]^S$, we have

$$\sum_{\underline{i}_2, \underline{j}_2 \in [q]^{[k] \setminus S}} \mathbb{E}_p[\psi(\underline{i}) \psi(\underline{j})] \prod_{t \in [k] \setminus S} \pi_{i_t} \pi_{j_t} = \mathbb{E}_p\left[\mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \underline{\sigma}_S = \underline{i}_1] \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \underline{\sigma}_S = \underline{j}_1]\right],$$

where \mathbb{E}_π is taken w.r.t. $\underline{\sigma} \sim \pi^{\otimes k}$ conditional on ψ and $\underline{\sigma}_S \equiv (\sigma_s)_{s \in S}$. By our assumption that $\psi \stackrel{d}{=} \psi^\theta$ for $\psi \sim p$ and any permutation θ in (5), the RHS above does not depend S given i_1, j_1 . Thus, by fixing $S = [\ell]$, we can express $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)\psi(\underline{\tau}_\omega)]$ by

$$\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)\psi(\underline{\tau}_\omega)] = \sum_{\ell=0}^k \binom{k}{\ell} n^{-\ell/2} \sum_{\underline{i}, \underline{j} \in [q]^\ell} \mathbb{E}_p \left[\mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{[\ell]} = \underline{i}] \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{[\ell]} = \underline{j}] \right] \cdot \prod_{s=1}^{\ell} X(i_s, j_s). \quad (79)$$

We now divide the summand above into the cases $\ell \in \{0, 1, 2\}$ and $\ell \geq 3$. For $\ell = 0$, by (SYM),

$$\mathbb{E}_p \left(\mathbb{E}_\pi[\psi(\underline{\sigma})] \right)^2 = \sum_{\psi \in \Psi} p(\psi) \left(\mathbb{E}_\pi[\psi(\underline{\sigma})] \right)^2 = \xi^2. \quad (80)$$

For $\ell = 1$, again by (SYM), we have

$$kn^{-1/2} \sum_{i,j=1}^q X(i, j) \mathbb{E}_p \left[\mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_1 = i] \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_1 = j] \right] = kn^{-1/2} \sum_{i,j=1}^q X(i, j) \xi^2 = 0, \quad (81)$$

where the last equality follows since $\sum_{i,j=1}^q X(i, j) = 0$. For $\ell = 2$, we have by definition of $\widehat{\Xi}$ that

$$\binom{k}{2} n^{-1} \sum_{\underline{i}, \underline{j} \in [q]^2} \mathbb{E}_p \left[\mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{\{1,2\}} = \underline{i}] \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{\{1,2\}} = \underline{j}] \right] \cdot \prod_{s=1}^2 X(i_s, j_s) = \binom{k}{2} \frac{\xi^2}{n} \langle \widehat{\Xi} X, X \rangle. \quad (82)$$

Finally, for $\ell \geq 3$, we can crudely bound

$$\begin{aligned} & \sum_{\ell \geq 3} \binom{k}{\ell} n^{-\ell/2} \sum_{\underline{i}, \underline{j} \in [q]^\ell} \mathbb{E}_p \left[\mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{[\ell]} = \underline{i}] \cdot \mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{[\ell]} = \underline{j}] \right] \cdot \prod_{s=1}^{\ell} X(i_s, j_s) \\ & \leq \max_{\psi \in \Psi} \|\psi\|_\infty^2 \cdot \sum_{\ell \geq 3} \binom{k}{\ell} n^{-\ell/2} q^{2\ell} \cdot \|X\|_\infty^\ell \leq \max_{\psi \in \Psi} \|\psi\|_\infty^2 \cdot (1+q)^{2k} \cdot n^{-3/2} \|X\|_\infty^3, \end{aligned} \quad (83)$$

where the last inequality follows since $\|X\|_\infty \leq \sqrt{n}$ by definition of X in (75). Therefore, plugging in the equalities (80)-(82) and the bound (83) into (79), we have

$$\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)\psi(\underline{\tau}_\omega)] = \xi^2 \cdot \left(1 + \frac{k(k-1)}{2n} \langle \widehat{\Xi} X, X \rangle + O_{k,q,\Psi} \left(\frac{\|X\|_\infty^3}{n^{3/2}} \right) \right). \quad (84)$$

Note that since $\psi(\cdot) > 0$ for any $\psi \in \Psi$, there exists a constant $c_i \equiv c_{i,\Psi}$ for $i = 1, 2$, such that $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)\psi(\underline{\tau}_\omega)] \in [c_1 \xi^2, c_2 \xi^2]$. Moreover, there exists a constant $C \equiv C_{c_1, c_2} > 0$ such that for $1+x \in [c_1, c_2]$, $e^{x-Cx^2} \leq 1+x \leq e^x$ holds. Applying this inequality for $x = k(k-1) \langle \widehat{\Xi} X, X \rangle / n + O_{k,q,\Psi}(\|X\|_\infty^3 / n^{3/2})$ and using the fact that $\|X\|_\infty \leq \sqrt{n}$, we obtain the estimate (77).

Next, we prove (78) by a similar argument. Denote by $X(i, \cdot) := \sum_{j=1}^q X(i, j)$. Then, proceeding in the same manner as in (79), we can express $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)]$ by

$$\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] = \sum_{\ell=0}^k \binom{k}{\ell} n^{-\ell/2} \sum_{\underline{i} \in [q]^\ell} \mathbb{E}_p \left[\mathbb{E}_\pi[\psi(\underline{\sigma}) | \underline{\sigma}_{[\ell]} = \underline{i}] \right] \prod_{s=1}^{\ell} X(i_s, \cdot) =: \sum_{\ell=0}^k G(\ell). \quad (85)$$

Then, by the same calculations done in (80) and (81), $G(0) = \xi$ and $G(1) = 0$ hold by (SYM). For $\ell = 2$, we have

$$\begin{aligned} G(2) &\equiv \binom{k}{2} n^{-1} \sum_{i \in [q]^2} \mathbb{E}_p \left[\mathbb{E}_\pi [\boldsymbol{\psi}(\boldsymbol{\sigma}) \mid \boldsymbol{\sigma}_{\{1,2\}} = \underline{i}] \right] \prod_{s=1}^2 X(i_s, \cdot) \\ &= \binom{k}{2} \frac{\xi}{n} \sum_{i_1, i_2, j_1, j_2 \in [q]} \Phi(i_1, i_2) \prod_{s=1}^2 X(i_s, j_s) = \binom{k}{2} \frac{\xi}{n} \langle (\widehat{\Phi} \otimes \mathbf{1}\mathbf{1}^\top) X, X \rangle. \end{aligned}$$

For $\ell \geq 3$, proceeding in the same manner as done in (83), we can bound

$$\sum_{\ell \geq 3} G(3) \equiv \sum_{\ell \geq 3} \binom{k}{\ell} n^{-\ell/2} \sum_{i, j \in [q]^\ell} \mathbb{E}_p \left[\mathbb{E}_\pi [\boldsymbol{\psi}(\boldsymbol{\sigma}) \mid \boldsymbol{\sigma}_{[\ell]} = \underline{i}] \right] \prod_{s=1}^{\ell} X(i_s, j_s) \leq \max_{\psi \in \Psi} \|\psi\|_\infty^2 \cdot (1+q)^{2k} \cdot n^{-3/2} \|X\|_\infty^3$$

By plugging in the obtained bounds into (85), and using the inequality $e^{x-O(x^2)} \leq 1+x \leq e^x$ for $1+x$ bounded away from 0 and ∞ as before, we obtain the first inequality of (78). The second inequality of (78) then follows from the first by exchanging the role of $\underline{\sigma}$ and $\underline{\tau}$. \square

Define the matrix $\widehat{\Xi}_* \in \mathbb{R}^{q^2 \times q^2}$ by

$$\widehat{\Xi}_* = \widehat{\Xi} - \widehat{\Phi} \otimes \mathbf{1}\mathbf{1}^\top - \mathbf{1}\mathbf{1}^\top \otimes \widehat{\Phi} + \mathbf{1}\mathbf{1}^\top \otimes \mathbf{1}\mathbf{1}^\top.$$

We note that the matrix $\widehat{\Xi}_*$ is related with the linear operator Ξ_* on $\mathbb{R}^q \otimes \mathbb{R}^q$ as follows. By identifying the vector spaces $\mathbb{R}^q \otimes \mathbb{R}^q$ and \mathbb{R}^{q^2} by the unique isomorphism that maps $e_i \otimes e_j \rightarrow e_{(i,j)}$ for $i, j \in [q]$, where $(e_i)_{i \leq q}$ is the standard basis in \mathbb{R}^q and $(e_{(i,j)})_{i,j \leq q}$ is the standard basis in \mathbb{R}^{q^2} , Ξ can be identified with the $q^2 \times q^2$ matrix $\Xi((i,j), (s,t))_{i,j,s,t \leq q} \in \mathbb{R}^{q^2 \times q^2}$, where

$$\Xi((i,j), (s,t)) \equiv \mathbb{E}_p \left[\Phi_\psi(i,s) \cdot \Phi_\psi(j,t) \right].$$

Similarly, Ξ_* can be identified with a $q^2 \times q^2$ matrix. With such identification, we have that $\Xi = \widehat{\Xi} \text{diag}(\pi)^{\otimes 2}$, thus

$$\widehat{\Xi}_* \text{diag}(\pi)^{\otimes 2} = \Xi - \Phi \otimes \mathbf{1}\pi^\top - \mathbf{1}\pi^\top \otimes \Phi + (\mathbf{1}\pi^\top) \otimes (\mathbf{1}\pi^\top) = \Xi_*. \quad (86)$$

As a consequence of Lemmas 6.4, 6.5, and 6.6, we have the following proposition. We recall the random variable $\mathbf{X}_n \equiv X_{\underline{\sigma}, \underline{\tau}} \in \mathbb{R}^{q \times q} \cong \mathbb{R}^{q^2}$, where $\underline{\sigma}, \underline{\tau} \sim \pi^{\otimes n}$.

Proposition 6.7. *There exists a constant $C \equiv C_{k,q,\Psi} > 0$ which only depends on k, q, Ψ such that the following holds. For $d < d_{\kappa S}$ and any $(m_n)_{n \geq 1}, (\varepsilon_n)_{n \geq 1}$ such that $|m_n - dn/k| \leq n^{2/3}$ and $\varepsilon_n \rightarrow 0, n^{2/3}\varepsilon_n \rightarrow \infty$ as $n \rightarrow \infty$, we have for large enough n that*

$$\mathbb{E} \left(L^*(\mathbf{G}(n, m_n)) \right)^2 \leq (1 - \sqrt{\varepsilon_n})^{-1} \mathbb{E} \left[\exp \left(\frac{(k-1)d}{2} \left(\langle \widehat{\Xi}_* \mathbf{X}_n, \mathbf{X}_n \rangle + C\sqrt{\varepsilon_n} \|\mathbf{X}_n\|_\infty^2 \right) \right) \mathbf{1} \left\{ \|\mathbf{X}_n\|_1 \leq \sqrt{n\varepsilon_n} \right\} \right]$$

Proof. Recall the decomposition $L^*(G)^2 = L_{\text{ind}}^2(G) + L_{\text{cor}}^2(G)$ in (74). By Lemma 6.4, we have

$$\mathbb{E}\left(L^*(\mathbf{G}(n, m_n))\right)^2 \leq (1 - \sqrt{\varepsilon_n})^{-1} \cdot \mathbb{E}L_{\text{ind}}^2(\mathbf{G}(n, m_n)). \quad (87)$$

Meanwhile, by Lemma 6.5, $\mathbb{E}L_{\text{ind}}^2(\mathbf{G}(n, m_n))$ can be upper bounded by

$$\sum_{\underline{\sigma}, \underline{\tau} \in [q]^V: \|R_{\underline{\sigma}, \underline{\tau}} - \pi\pi^\top\|_1 \leq \sqrt{\varepsilon_n}} \left(\frac{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)\psi(\underline{\tau}_\omega)]}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] \cdot \mathbb{E}_{p,u}[\psi(\underline{\tau}_\omega)]} \right)^{m_n} \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\sigma}) \cdot \mathbb{P}(\underline{\sigma}^* = \underline{\tau}).$$

Observe that for $\underline{\sigma}, \underline{\tau} \in [q]^V$, the restriction $\|R_{\underline{\sigma}, \underline{\tau}} - \pi\pi^\top\|_1 \leq \sqrt{\varepsilon_n}$ is equivalent to $\|X_{\underline{\sigma}, \underline{\tau}}\|_1 \leq \sqrt{n\varepsilon_n}$. In particular, $\|X_{\underline{\sigma}, \underline{\tau}}\|_\infty^3/n^{3/2} \leq \sqrt{\varepsilon_n}\|X_{\underline{\sigma}, \underline{\tau}}\|_\infty$ holds under such restriction. Thus, for $\underline{\sigma}, \underline{\tau} \in [q]^V$ such that $\|R_{\underline{\sigma}, \underline{\tau}} - \pi\pi^\top\|_1 \leq \sqrt{\varepsilon_n}$, we have by Lemma 6.6 that

$$\left(\frac{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)\psi(\underline{\tau}_\omega)]}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] \cdot \mathbb{E}_{p,u}[\psi(\underline{\tau}_\omega)]} \right)^{m_n} \leq \exp\left(\frac{m_n}{n} \binom{k}{2} \left(\left\langle \left(\widehat{\Xi} - \widehat{\Phi} \otimes \mathbf{1}\mathbf{1}^\top - \mathbf{1}\mathbf{1}^\top \otimes \widehat{\Phi} \right) X, X \right\rangle + C\sqrt{\varepsilon_n}\|X\|_\infty^2 \right)\right),$$

where $C \equiv C_{k,q,\Psi} > 0$ denotes a constant that only depend on k, q , and the set Ψ . Note that since $\langle \mathbf{1}\mathbf{1}^\top, X \rangle = \sum_{i,j=1}^q X(i, j) = 0$, the inner product above equals $\langle \widehat{\Xi}_* X, X \rangle$. Moreover, since $m_n \leq dn/k + n^{3/2}$ and $n^{-1/3} \ll \sqrt{\varepsilon_n}$ holds, the factor $\frac{m_n}{n} \binom{k}{2}$ in the RHS above can be replaced by $\frac{(k-1)d}{2}$ for large enough n with the modification of the constant C . Therefore,

$$\mathbb{E}L_{\text{ind}}^2(\mathbf{G}(n, m_n)) \leq \mathbb{E}\left[\exp\left(\frac{(k-1)d}{2} \left(\langle \widehat{\Xi}_* \mathbf{X}_n, \mathbf{X}_n \rangle + C'\sqrt{\varepsilon_n}\|\mathbf{X}_n\|_\infty^2 \right) \right) \mathbf{1}\left\{\|\mathbf{X}_n\|_1 \leq \sqrt{n\varepsilon_n}\right\} \right],$$

for some constant $C' \equiv C'_{k,q,\Psi}$. Combining this with (87) concludes the proof. \square

Having Proposition 6.7 in hand, the final ingredient to prove Proposition 4.5 is the uniform integrability of $(\mathbf{Z}_n)_{n \geq 1} \equiv (\mathbf{Z}_n(d, k, \varepsilon, C))_{n \geq 1}$, where $\varepsilon \equiv (\varepsilon_n)_{n \geq 1}$ and

$$\mathbf{Z}_n := \exp\left(\frac{(k-1)d}{2} \left(\langle \widehat{\Xi}_* \mathbf{X}_n, \mathbf{X}_n \rangle + C\sqrt{\varepsilon_n}\|\mathbf{X}_n\|_\infty^2 \right) \right) \mathbf{1}\left\{\|\mathbf{X}_n\|_1 \leq \sqrt{n\varepsilon_n}\right\}.$$

Here, we emphasize that the indicator $\mathbf{1}\left\{\|\mathbf{X}_n\|_1 \leq \sqrt{n\varepsilon_n}\right\}$ due to the truncation $\mathbf{1}\left\{\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top \rangle_G \leq \varepsilon_n\right\}$ in $L_*(G)$ is crucial for uniform integrability stated below.

Lemma 6.8. *For $d < d_{\text{KS}}$ and $\varepsilon = (\varepsilon_n)_{n \geq 1}$, where $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$, the random variables $(\mathbf{Z}_n)_{n \geq 1}$ are uniformly integrable.*

Proof. Denote by $\mathcal{P}_n([q]^2)$ the set of $R \equiv (R(i, j))_{i, j \leq q} \in \mathbb{R}^{q \times q}$ such that $R(i, j) \in \mathbb{Z}_{\geq 0}/n$ for any $i, j \leq q$ and $\sum_{i, j=1}^q R(i, j) = 1$. Further, denote by $\mathbf{R}_n \equiv R_{\underline{\sigma}, \underline{\tau}}$ the overlap matrix under $\underline{\sigma}, \underline{\tau} \stackrel{i.i.d.}{\sim} \pi^{\otimes n}$. Then, by Sanov's theorem (a.k.a. Stirling's approximation), we have for any $R \in \mathcal{P}_n([q]^2)$ that

$$\mathbb{P}(\mathbf{R}_n = R) = n^{O_q(1)} \exp(-n \cdot D_{\text{KL}}(R \| \pi\pi^\top)), \quad (88)$$

where $O_q(1)$ denotes a bounded constant that only depends on q and $D_{\text{KL}}(\mu \parallel \nu) := \sum_{i,j \in [q]} \mu(i,j) \log \left(\frac{\nu(i,j)}{\mu(i,j)} \right)$ for $\mu, \nu \in \mathcal{P}([q]^2)$ is the KL divergence (a.k.a. relative entropy) between μ and ν . By viewing $R \in \mathcal{P}_n([q]^2)$ as a q^2 dimensional vector, we first claim that for $d < d_{\text{KS}}$,

$$\nabla_R^2 D_{\text{KL}}(R \parallel \pi\pi^\top) \Big|_{R=\pi\pi^\top} \succ (k-1)d \cdot \widehat{\Xi}_*, \quad (89)$$

where ∇_R^2 denotes the Hessian with respect to $R \in \mathbb{R}^{q^2}$. Indeed, note that a direct computation gives $\nabla_R^2 D_{\text{KL}}(R \parallel \pi\pi^\top) \Big|_{R=\pi\pi^\top} = \text{diag}((\pi_i^{-1})_{i \leq q})^{\otimes 2}$, thus (89) is equivalent to $I_{q^2 \times q^2} \succ (k-1)d \cdot \widehat{\Xi}_* \text{diag}(\pi)^{\otimes 2}$, which follows for $d < d_{\text{KS}}$ by Lemma 4.2 since the maximum eigenvalue of $\widehat{\Xi}_* \text{diag}(\pi)^{\otimes 2}$ equals $\max_{\lambda \in \text{Eig}(\Xi_*)} |\lambda|$ by (86). Hence, (89) holds for $d < d_{\text{KS}}$.

Now, observe that since $R \rightarrow D_{\text{KL}}(R \parallel \pi\pi^\top)$ is convex with $D_{\text{KL}}(\pi\pi^\top \parallel \pi\pi^\top) = 0$, (89) implies that there exists small enough $\varepsilon, \delta > 0$ such that if $\|R - \pi\pi^\top\|_1 \leq \varepsilon$, then

$$D_{\text{KL}}(R \parallel \pi\pi^\top) \geq (1+\delta) \frac{(k-1)d}{2} \left\langle \widehat{\Xi}_*(R - \pi\pi^\top), R - \pi\pi^\top \right\rangle + \delta \|R - \pi\pi^\top\|_2^2. \quad (90)$$

For such $\delta > 0$, we now claim that $(\mathbb{E}Z_n^{1+\delta})_{n \geq 1}$ is bounded, which is sufficient for uniform integrability. Note that $\mathbf{X}_n = \sqrt{n}(R_n - \pi\pi^\top)$ holds by definition. Thus, by (88), we can bound $\mathbb{E}Z_n^{1+\delta}$ by

$$\sum_{\substack{R \in \mathcal{P}_n([q]^2) \\ \|R - \pi\pi^\top\|_1 \leq \sqrt{\varepsilon_n}}} \exp \left(\frac{(1+\delta)(k-1)dn}{2} \left(\left\langle \widehat{\Xi}_*(R - \pi\pi^\top), R - \pi\pi^\top \right\rangle + C\sqrt{\varepsilon_n} \|R - \pi\pi^\top\|_\infty^2 \right) - nD_{\text{KL}}(R \parallel \pi\pi^\top) \right).$$

Observe that since $\varepsilon_n = o_n(1)$, the above sum is restricted to $\|R - \pi\pi^\top\|_1 \leq \sqrt{\varepsilon_n} < \varepsilon$ for large enough n . Thus, we can use the inequality (90) to further bound

$$\begin{aligned} \mathbb{E}Z_n^{1+\delta} &\leq \sum_{R \in \mathcal{P}_n([q]^2)} \exp \left(-n\delta \|R - \pi\pi^\top\|_2^2 + \frac{C(1+\delta)(k-1)d}{2} n\sqrt{\varepsilon_n} \|R - \pi\pi^\top\|_\infty^2 \right) \\ &\leq \sum_{R \in \mathcal{P}_n([q]^2)} \exp \left(-\frac{n\delta}{2} \|R - \pi\pi^\top\|_2^2 \right) \leq C', \end{aligned}$$

where the second inequality follows for large enough n since $\varepsilon_n = o_n(1)$ and $\|R - \pi\pi^\top\|_\infty \leq \|R - \pi\pi^\top\|_2$, and the last inequality follows by Gaussian integration for some constant $C' \equiv C'(\delta, q) < \infty$ that only depends on δ, q . Therefore, $\sup_{n \geq 1} \mathbb{E}Z_n^{1+\delta} < \infty$ holds, and uniform integrability of $(Z_n)_{n \geq 1}$ follows. \square

Proof of Proposition 4.5. Note that $\mathbb{E}(L^*(\mathbf{G}(n, m_n)))^2$ increases as ε_n increases. Thus, w.l.o.g., we may assume that $n^{2/3}\varepsilon_n \rightarrow \infty$ as $n \rightarrow \infty$. Also, recall that $\mathbf{X}_n \equiv \sqrt{n}(R_{\underline{\sigma}, \underline{\tau}} - \pi\pi^\top)$, where $\underline{\sigma}, \underline{\tau} \sim \pi^{\otimes n}$. Then, $\mathbf{X}_n \xrightarrow{d} \mathbf{X}_\infty \sim \mathbf{N}(\text{diag}(\pi)^{\otimes 2} - (\pi\pi^\top)^{\otimes 2})$ holds by the central limit theorem (cf. (76)). Recalling the random variable $Z_n \equiv \exp \left(\frac{(k-1)d}{2} \left(\left\langle \widehat{\Xi}_* \mathbf{X}_n, \mathbf{X}_n \right\rangle + C\sqrt{\varepsilon_n} \|\mathbf{X}_n\|_\infty^2 \right) \right) \mathbf{1} \left\{ \|\mathbf{X}_n\|_1 \leq \sqrt{n\varepsilon_n} \right\}$, we claim that as $n \rightarrow \infty$,

$$\mathbf{Z}_n \xrightarrow{d} \mathbf{Z}_\infty := \exp \left(\frac{(k-1)d}{2} \left\langle \widehat{\Xi}_* \mathbf{X}_\infty, \mathbf{X}_\infty \right\rangle \right). \quad (91)$$

To see this, note that $\exp\left(\frac{(k-1)d}{2}\langle\widehat{\Xi}_*\mathbf{X}_n,\mathbf{X}_n\rangle\right)\xrightarrow{d}\exp\left(\frac{(k-1)d}{2}\langle\widehat{\Xi}_*\mathbf{X}_\infty,\mathbf{X}_\infty\rangle\right)$ holds by the continuous mapping theorem. Moreover, since $n\varepsilon_n\rightarrow\infty$ and $\varepsilon_n\rightarrow 0$ as $n\rightarrow\infty$, we have that

$$\exp\left(C\sqrt{\varepsilon_n}\|\mathbf{X}_n\|_\infty^3\right)\xrightarrow{p}1\quad\text{and}\quad\mathbb{1}\{\|\mathbf{X}_n\|_1\leq\sqrt{n\varepsilon_n}\}\xrightarrow{p}1$$

Thus, by Slutsky, (91) holds. Meanwhile, $(\mathbf{Z}_n)_{n\geq 1}$ is uniformly integrable by Lemma 6.8, thus

$$\lim_{n\rightarrow\infty}\mathbb{E}\mathbf{Z}_n=\mathbb{E}\mathbf{Z}_\infty.$$

Note that $\mathbb{E}\mathbf{Z}_\infty$ can be calculated explicitly as follows. Recall that if $Z\sim\mathbf{N}(0,\Sigma)$ and the eigenvalues of $A\Sigma$ is bounded by 1 in absolute value for a symmetric matrix A , then $\mathbb{E}\exp(Z^\top AZ/2)=\prod_{\lambda\in\text{Eig}(A\Sigma)}(1-\lambda)^{-1/2}$ holds, where $\text{Eig}(\Sigma^{1/2}A\Sigma)=\text{Eig}(A\Sigma)$ denotes the set of eigenvalues of $A\Sigma$ (see e.g. [MP92, Theorem 3.2a.2]). Also, note that

$$\widehat{\Xi}_*(\pi\pi^\top)^{\otimes 2}=\mathbb{E}_p\left[\widehat{\Phi}_\psi\pi\pi^\top\otimes\widehat{\Phi}_\psi\pi\pi^\top\right]-\widehat{\Phi}\pi\pi^\top\otimes\mathbf{1}\pi^\top-\mathbf{1}\pi^\top\otimes\widehat{\Phi}\pi\pi^\top+(\mathbf{1}\pi^\top)\otimes(\mathbf{1}\pi^\top)=0,$$

where the last equality holds because (SYM) yields $\widehat{\Phi}_\psi\pi=\mathbf{1}$ for any $\psi\in\Psi$. Thus, it follows that $\widehat{\Xi}_*(\text{diag}(\pi)^{\otimes 2}-(\pi\pi^\top)^{\otimes 2})=\Xi_*$ holds. Therefore, by Proposition 6.7, we can bound

$$\mathbb{E}\left(L^*(\mathbf{G}(n,m_n))\right)^2\leq\mathbb{E}\mathbf{Z}_n=\mathbb{E}\mathbf{Z}_\infty+o_n(1)=\prod_{\lambda\in\text{Eig}(\Xi_*)}\frac{1}{\sqrt{1-(k-1)d\lambda}}+o_n(1),$$

which concludes the proof since non-zero elements of $\text{Eig}(\Xi_*)$ equals the non-zero elements of $\text{Eig}_S(\Xi)$ by Lemma 4.2. \square

7 Mutual information between the planted and the null model

In this section, we prove Lemma 4.18 and Propositions 4.20, 4.21. In Section 7.1, we prove Lemma 4.18. In Section 7.2, we prove Proposition 4.20. In Section 7.3, we prove Proposition 4.21.

7.1 Proof of Lemma 4.18

By invoking the identity (42), we have

$$\frac{1}{n}D_{\text{KL}}(\mathbf{G}^*\|\mathbf{G})\equiv\frac{1}{n}\sum_G\mathbb{P}(\mathbf{G}^*=G)\log\frac{\mathbb{P}(\mathbf{G}^*=G)}{\mathbb{P}(\mathbf{G}=G)}=\frac{1}{n}\mathbb{E}\log L(\mathbf{G}^*),$$

which finishes the proof of the first equality. Turning to the mutual information, we have by definition that

$$\begin{aligned}\frac{1}{n}I(\mathbf{G}^*,\underline{\sigma}^*)&\equiv\frac{1}{n}\sum_{G,\underline{\sigma}}\mathbb{P}(\mathbf{G}^*=G,\underline{\sigma}^*=\underline{\sigma})\log\left(\frac{\mathbb{P}(\mathbf{G}=G)}{\mathbb{P}(\mathbf{G}^*=G)}\cdot\frac{\mathbb{P}(\mathbf{G}^*=G|\underline{\sigma}^*=\underline{\sigma})}{\mathbb{P}(\mathbf{G}=G)}\right) \\ &= -\frac{1}{n}\mathbb{E}\log L(\mathbf{G}^*)+\frac{1}{n}\sum_{G,\underline{\sigma}}\mathbb{P}(\mathbf{G}^*=G,\underline{\sigma}^*=\underline{\sigma})\log\frac{\mathbb{P}(\mathbf{G}^*=G|\underline{\sigma}^*=\underline{\sigma})}{\mathbb{P}(\mathbf{G}=G)},\end{aligned}\tag{92}$$

where the last equality is due to (42). We now simplify the last sum above. Recall that by Definition 3.2, if $G = (V, F, E, (\psi_a)_{a \in F})$ is a factor graph with m clauses, then

$$\frac{\mathbb{P}(\mathbf{G}^* = G \mid \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G} = G)} = \frac{\psi_G(\underline{\sigma})}{\mathbb{E}[\psi_{\mathbf{G}(n,m)}(\underline{\sigma})]} = \prod_{a \in F} \frac{\psi_a(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)]}, \quad (93)$$

where $\psi \sim p$ and $\omega \sim \text{Unif}(V^k)$. Thus, if we denote by $(\psi_{a_i}, \delta a_i)_{1 \leq i \leq m}$ the random weight functions and neighborhoods of clauses of \mathbf{G}^* , which are i.i.d. with distribution (7), then it follows that

$$\frac{1}{n} \sum_{G, \underline{\sigma}} \mathbb{P}(\mathbf{G}^* = G, \underline{\sigma}^* = \underline{\sigma}) \log \frac{\mathbb{P}(\mathbf{G}^* = G \mid \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G} = G)} = \frac{1}{n} \mathbb{E} \left[\sum_{i=1}^m \log \frac{\psi_{a_i}(\underline{\sigma}_{\delta a_i}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \right] = \frac{d}{k} \mathbb{E} \left[\log \frac{\psi_{a_1}(\underline{\sigma}_{\delta a_1}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \right], \quad (94)$$

where the last equality holds since $(\psi_{a_i}, \delta a_i)_{1 \leq i \leq m}$ are i.i.d., and $\mathbb{E}[m] = dn/k$. Since $(\psi_{a_1}, \delta a_1)$ is distributed according to (7), the RHS equals

$$\frac{d}{k} \mathbb{E} \left[\log \frac{\psi_{a_1}(\underline{\sigma}_{\delta a_1}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \right] = \frac{d}{k} \mathbb{E} \left[\frac{\psi'(\underline{\sigma}_{\omega'}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \cdot \log \frac{\psi'(\underline{\sigma}_{\omega'}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \right], \quad (95)$$

where the outer expectation in the RHS is w.r.t. $\underline{\sigma}^* \sim \pi^{\otimes V}$, $\psi' \sim p$, and $\omega' \sim \text{Unif}(V^k)$. Observe that since $\underline{\sigma}^* \sim \pi^{\otimes V}$, the total variation distance between the empirical distribution of $\underline{\sigma}^*$ and π is at most $O(n^{-1/3})$ with probability tending to one. Thus, w.h.p., $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)] = \mathbb{E}_{p,\pi}[\psi(\underline{\sigma})] + O(n^{-1/3}) \equiv \xi + O(n^{-1/3})$ holds. In addition, by our assumption that $\psi(\cdot) \in (0, \infty)$ and Ψ is finite, there exists constants $c, C \in (0, \infty)$ such that $\psi(\cdot) \in [c, C]$ holds for any $\psi \in \Psi$. As a consequence, we have that

$$\begin{aligned} \frac{d}{k} \mathbb{E} \left[\frac{\psi'(\underline{\sigma}_{\omega'}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \cdot \log \frac{\psi'(\underline{\sigma}_{\omega'}^*)}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega^*)]} \right] &= \frac{d}{k} \mathbb{E} \left[\frac{\psi'(\underline{\sigma}_{\omega'}^*)}{\xi} \cdot \log \frac{\psi'(\underline{\sigma}_{\omega'}^*)}{\xi} \right] + o_n(1) \\ &= \frac{d}{k} \mathbb{E}_{p,\pi} \left[\frac{\psi(\underline{\sigma})}{\xi} \cdot \log \frac{\psi(\underline{\sigma})}{\xi} \right] + o_n(1), \end{aligned} \quad (96)$$

where in the last equality, we again used the fact that the total variation distance between the empirical distribution of $\underline{\sigma}^*$ and π is at most $O(n^{-1/3})$ with probability tending to one. Combining (94), (95), (96) with (93) concludes the proof of Lemma 4.18.

7.2 Proof of Proposition 4.20

It is well-known that given a function $f : \mathbb{N} \rightarrow \mathbb{R}$ at most of exponential growth and $\alpha > 0$, the map $d \rightarrow \mathbb{E}f(X_d)$ for $X_d \sim \text{Poi}(\alpha d)$ is differentiable w.r.t. d with derivative $\frac{\partial}{\partial d} \mathbb{E}f(X_d) = \alpha(\mathbb{E}f(X_d + 1) - \mathbb{E}f(X_d))$. Thus, it follows that

$$\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) = \frac{1}{k} \left(\mathbb{E} \log L(\mathbf{G}^*(n, \mathbf{m} + 1)) - \mathbb{E} \log L(\mathbf{G}^*(n, \mathbf{m})) \right). \quad (97)$$

We couple $\mathbf{G}^*(n, \mathbf{m})$ and $\mathbf{G}^*(n, \mathbf{m} + 1)$ to show that the RHS can be calculated as follows.

Lemma 7.1. *For any $d > 0$ and $n, m \geq 1$, there exists a coupling between $\mathbf{G}^*(n, m)$ and $\mathbf{G}^*(n, m + 1)$ such that*

$$\mathbb{E} \log \left(\frac{L(\mathbf{G}^*(n, m + 1))}{L(\mathbf{G}^*(n, m))} \right) = \mathbb{E}_{\mathbf{G}^*(n,m), p, u} \left[\left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p,u}[\psi'(\underline{\sigma}_\omega)]} \right\rangle_{\mathbf{G}^*(n,m)} \log \left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p,u}[\psi'(\underline{\sigma}_\omega)]} \right\rangle_{\mathbf{G}^*(n,m)} \right],$$

where the expectation $\mathbb{E}_{\mathbf{G}^*(n,m),p,u}$ is with respect to $\mathbf{G}^*(n,m), \boldsymbol{\psi} \sim p, \boldsymbol{\omega} \sim u := \text{Unif}(V^k)$, and the expectation $\mathbb{E}_{p,u}$ in the denominator is with respect to $\boldsymbol{\psi}' \sim p$ and $\boldsymbol{\omega}' \sim u$.

Proof. Recalling the Definition 3.2 of the planted model $\mathbf{G}^*(n,m) \equiv \mathbf{G}^*(n,m,\boldsymbol{\sigma}^*)$, we can couple $\mathbf{G}^*(n,m)$ and $\mathbf{G}^*(n,m+1)$ by first drawing $\boldsymbol{\sigma}^* \sim \pi^{\otimes n}$ and $\mathbf{G}^*(n,m,\boldsymbol{\sigma}^*)$, and then conditional on $\boldsymbol{\sigma}^*, \mathbf{G}^*(n,m,\boldsymbol{\sigma}^*)$, adding an independent clause a_{m+1} to $\mathbf{G}^*(n,m)$ with neighborhood δa_{m+1} and the weight function $\psi_{a_{m+1}}$ from the distribution

$$\mathbb{P}\left(\delta a_{m+1} = (v_1, \dots, v_k), \psi_{a_{m+1}} = \psi \mid \boldsymbol{\sigma}^*, \mathbf{G}^*(n,m,\boldsymbol{\sigma}^*)\right) = \frac{1}{n^k} \cdot \frac{p(\psi)\psi(\boldsymbol{\sigma}_{v_1}^*, \dots, \boldsymbol{\sigma}_{v_k}^*)}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\boldsymbol{\sigma}_{\boldsymbol{\omega}}^*)]}. \quad (98)$$

Recalling the notation $\psi_G(\underline{\sigma})$ in (6), we can express $L(\mathbf{G}^*(n,m+1))$ under such coupling by

$$L(\mathbf{G}^*(n,m+1)) = \sum_{\underline{\sigma} \in [q]^V} \frac{\psi_{\mathbf{G}^*(n,m+1)}(\underline{\sigma})}{\mathbb{E}[\boldsymbol{\psi}_{\mathbf{G}^*(n,m+1)}(\underline{\sigma})]} \mathbb{P}(\boldsymbol{\sigma}^* = \underline{\sigma}) = \sum_{\underline{\sigma} \in [q]^V} \frac{\psi_{a_{m+1}}(\underline{\sigma}_{\delta a_{m+1}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \cdot \frac{\psi_{\mathbf{G}^*(n,m)}(\underline{\sigma})}{\mathbb{E}[\boldsymbol{\psi}_{\mathbf{G}^*(n,m)}(\underline{\sigma})]} \mathbb{P}(\boldsymbol{\sigma}^* = \underline{\sigma}). \quad (99)$$

Meanwhile, by Bayes rule, the posterior $\mu_G(\underline{\sigma})$ in (19), where G have n variables m clauses, equals

$$\mu_G(\underline{\sigma}) = \frac{\mathbb{P}(\mathbf{G}^*(n,m) = G \mid \boldsymbol{\sigma}^* = \underline{\sigma}) \mathbb{P}(\boldsymbol{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}^*(n,m) = G)} = \frac{\frac{\psi_G(\underline{\sigma})}{\mathbb{E}[\boldsymbol{\psi}_{\mathbf{G}^*(n,m)}(\underline{\sigma})]} \mathbb{P}(\boldsymbol{\sigma}^* = \underline{\sigma})}{L(G)}.$$

By taking $G = \mathbf{G}^*(n,m)$ and combining with (99), it follows that

$$\mathbb{E} \log \left(\frac{L(\mathbf{G}^*(n,m+1))}{L(\mathbf{G}^*(n,m))} \right) = \mathbb{E} \log \left(\sum_{\underline{\sigma} \in [q]^V} \frac{\psi_{a_{m+1}}(\underline{\sigma}_{\delta a_{m+1}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \mu_{\mathbf{G}^*(n,m)}(\underline{\sigma}) \right) = \mathbb{E} \log \left\langle \frac{\psi_{a_{m+1}}(\underline{\sigma}_{\delta a_{m+1}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \right\rangle_{\mathbf{G}^*(n,m)}, \quad (100)$$

where in the last expression, we abbreviated the superscript in $\boldsymbol{\sigma}^1 \sim \mu_{\mathbf{G}^*(n,m)}$ and the outer expectation is w.r.t. $\mathbf{G}^*(n,m)$ and $(\delta a_{m+1}, \psi_{a_{m+1}})$. By the conditional law of $(\delta a_{m+1}, \psi_{a_{m+1}})$ in (98), we have

$$\begin{aligned} \mathbb{E} \left[\log \left\langle \frac{\psi_{a_{m+1}}(\underline{\sigma}_{\delta a_{m+1}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \right\rangle_{\mathbf{G}^*(n,m)} \mid \boldsymbol{\sigma}^*, \mathbf{G}^*(n,m) \right] &= \sum_{\psi \in \Psi} \sum_{\delta a \in V^k} \log \left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \right\rangle_{\mathbf{G}^*(n,m)} \frac{p(\psi)\psi(\boldsymbol{\sigma}_{\delta a}^*)}{n^k \cdot \mathbb{E}_{p,u}[\boldsymbol{\psi}(\boldsymbol{\sigma}_{\boldsymbol{\omega}}^*)]} \\ &= \mathbb{E}_{p,u} \left[\frac{\boldsymbol{\psi}'(\boldsymbol{\sigma}_{\boldsymbol{\omega}'})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\boldsymbol{\sigma}_{\boldsymbol{\omega}}^*)]} \log \left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \right\rangle_{\mathbf{G}^*(n,m)} \right], \end{aligned}$$

where the outer expectation $\mathbb{E}_{p,u}$ is taken w.r.t. $\boldsymbol{\psi}' \sim p$ and $\boldsymbol{\omega}' \sim \text{Unif}(V^k)$ independent of everything else. Thus, by tower property, we have

$$\begin{aligned} &\mathbb{E} \left[\log \left\langle \frac{\psi_{a_{m+1}}(\underline{\sigma}_{\delta a_{m+1}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \right\rangle_{\mathbf{G}^*(n,m)} \mid \mathbf{G}^*(n,m) \right] \\ &= \mathbb{E} \left[\mathbb{E}_{p,u} \left[\frac{\boldsymbol{\psi}'(\boldsymbol{\sigma}_{\boldsymbol{\omega}'})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\boldsymbol{\sigma}_{\boldsymbol{\omega}}^*)]} \log \left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})]} \right\rangle_{\mathbf{G}^*(n,m)} \right] \mid \mathbf{G}^*(n,m) \right] \\ &= \mathbb{E}_{p,u} \left[\left\langle \frac{\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}'(\boldsymbol{\sigma}_{\boldsymbol{\omega}'})]} \right\rangle_{\mathbf{G}^*(n,m)} \log \left\langle \frac{\boldsymbol{\psi}(\underline{\sigma}_{\boldsymbol{\omega}})}{\mathbb{E}_{p,u}[\boldsymbol{\psi}'(\boldsymbol{\sigma}_{\boldsymbol{\omega}'})]} \right\rangle_{\mathbf{G}^*(n,m)} \right], \end{aligned}$$

where the last equality follows since $\mathbb{E}_{p,u}$ and the conditional expectation over \mathbf{G}^* is exchangeable and the expectation w.r.t. to the measure $\mathbb{P}(\underline{\sigma}^* = \cdot \mid \mathbf{G}^*(n, m))$ equals the expectation over $\underline{\sigma} \sim \mu_{\mathbf{G}^*(n, m)}$. Therefore, taking expectation in the equality above and combining with (100) concludes the proof. \square

In order to prove Proposition 4.20, it remains to approximate $\mathbb{E}_{p,u}[\psi'(\underline{\sigma}_{\omega'})]$ under $\underline{\sigma} \sim \mu_{\mathbf{G}^*(n, m)}$. Recalling that $R_{\underline{\sigma}}$ denotes the empirical distribution of $\underline{\sigma} \in [q]^V$ (cf. (115)), consider the event

$$\mathcal{A}_{\text{bal}}(n, m) := \left\{ \mathbb{P}\left(\|R_{\underline{\sigma}^*} - \pi\|_{\infty} \geq n^{-1/3} \mid \mathbf{G}^*(n, m)\right) \leq \exp(-n^{1/4}) \right\}.$$

Then, we have the following lemma.

Lemma 7.2. *There exists a constant $c \equiv c_q$ which only depends on q such that for any $n, m \geq 1$, we have $\mathbb{P}(\mathbf{G}^*(n, m) \in \mathcal{A}_{\text{bal}}(n, m)) \leq \exp(-cn^{1/3})$. Moreover, there exists a constant $C \equiv C_{k, q, \Psi}$ depending only on k, q, Ψ such that on the w.h.p. event \mathcal{A}_{bal} , for any $\psi \in \Psi$ and $\delta a \in V^k$, we have*

$$\left| \left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_{\omega})]} \right\rangle_{\mathbf{G}^*(n, m)} - \xi^{-1} \cdot \left\langle \psi(\underline{\sigma}_{\delta a}) \right\rangle_{\mathbf{G}^*(n, m)} \right| \leq Cn^{-1/3}. \quad (101)$$

Proof. For the first statement, note that by Markov's inequality and tower property, we have

$$\mathbb{P}(\mathbf{G}^*(n, m) \in \mathcal{A}_{\text{bal}}(n, m)) \leq \exp(-n^{1/4}) \cdot \mathbb{P}\left(\|R_{\underline{\sigma}^*} - \pi\|_{\infty} \geq n^{-1/3}\right).$$

Invoking (56), $\mathbb{P}(\|R_{\underline{\sigma}^*} - \pi\|_{\infty} \geq n^{-1/3}) \leq q \exp(-2n^{1/3})$ holds, thus plugging this bound shows that $\mathbb{P}(\mathbf{G}^*(n, m) \in \mathcal{A}_{\text{bal}}(n, m)) \leq \exp(-cn^{1/3})$ holds for some $c \equiv c_q > 0$.

We next prove (101). To this end, assume that $\mathbf{G}^*(n, m) \in \mathcal{A}_{\text{bal}}(n, m)$ happens, and fix any $\psi \in \Psi$ and $\delta a \in V^k$. Throughout, we denote C by a positive constant that depends only on k, q, Ψ and write $f = O_{k, q, \Psi}(g)$ if there exists a constant C such that $|f| \leq Cg$. Note that

$$\left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_{\omega})]} \mathbb{1}\left\{\|R_{\underline{\sigma}} - \pi\|_{\infty} \geq n^{-1/3}\right\} \right\rangle_{\mathbf{G}^*(n, m)} \leq C \left\langle \mathbb{1}\left\{\|R_{\underline{\sigma}} - \pi\|_{\infty} \geq n^{-1/3}\right\} \right\rangle_{\mathbf{G}^*(n, m)} \leq C \exp(-n^{1/4}), \quad (102)$$

where the last inequality holds since $\langle f(\underline{\sigma}) \rangle_G \equiv \mathbb{E}[f(\underline{\sigma}^*) \mid \mathbf{G}^* = G]$ holds by definition for $f : [q]^V \rightarrow \mathbb{R}$. Meanwhile, note that if $\underline{\tau} \in [q]^V$ satisfies $\|R_{\underline{\tau}} - \pi\|_{\infty} \leq n^{-1/3}$, then

$$\mathbb{E}_{p,u}[\psi(\underline{\tau})_{\omega}] = \mathbb{E}_p \left[\sum_{\underline{i}=(i_1, \dots, i_k) \in [q]^k} \psi(\underline{i}) \prod_{s=1}^k R_{\underline{\tau}}(i_s) \right] = \mathbb{E}_p \left[\sum_{\underline{i}=(i_1, \dots, i_k) \in [q]^k} \psi(\underline{i}) \prod_{s=1}^k \pi_{i_s} \right] + O_{k, q, \Psi}(n^{-1/3}),$$

which equals $\xi + O_{k, q, \Psi}(n^{-1/3})$. Thus, it follows that

$$\begin{aligned} \left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_{\omega})]} \mathbb{1}\left\{\|R_{\underline{\sigma}} - \pi\|_{\infty} \leq n^{-1/3}\right\} \right\rangle_{\mathbf{G}^*(n, m)} &= \left\langle \frac{\psi(\underline{\sigma}_{\delta a})}{\xi + O_{k, q, \Psi}(n^{-1/3})} \mathbb{1}\left\{\|R_{\underline{\sigma}} - \pi\|_{\infty} \leq n^{-1/3}\right\} \right\rangle_{\mathbf{G}^*(n, m)} \\ &= \xi^{-1} \cdot \left\langle \psi(\underline{\sigma}_{\delta a}) \right\rangle_{\mathbf{G}^*(n, m)} + O_{k, q, \Psi}(n^{-1/3}), \end{aligned}$$

where the last equality holds since $\langle \psi(\underline{\sigma}_{\delta a}) \mathbb{1}\left\{\|R_{\underline{\sigma}} - \pi\|_{\infty} \geq n^{-1/3}\right\} \rangle_{\mathbf{G}^*(n, m)} \leq C \exp(-n^{1/4})$ holds on the event $\mathcal{A}_{\text{bal}}(n, m)$. Therefore, combining with (102) completes the proof of (101). \square

The proof of Proposition 4.20 is straightforward from Lemma 7.1 and Lemma 7.2.

Proof of Proposition 4.20. The first statement (50) follows immediately from (97) and Lemma 7.1. To establish the second statement (51) from the first statement, note that Lemma 7.2 implies the following. Given a continuous function $F : \mathbb{R}_+ \rightarrow \mathbb{R}$, there exists a constant $C \equiv C_{F,k,q,\Psi}$, which only depends on F, k, q, Ψ , such that

$$\left| \mathbb{E}_{\mathbf{G}^*, p, u} F \left(\left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p, u}[\psi'(\underline{\sigma}_\omega)]} \right\rangle_{\mathbf{G}^*} \right) - \mathbb{E}_{\mathbf{G}^*, p, u} F \left(\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} \right) \right| \leq C n^{-1/3}. \quad (103)$$

Here, we note that the continuity of F suffices (rather than Lipschitz continuity) since there exist constants $c_i \equiv c_{i, \Psi} > 0$ such that for any $\psi \in \Psi$, $\psi(\cdot) \in [c_1, c_2]$ holds. Thus, taking $F(x) = x \log x$, the second statement (51) follows from (50). \square

7.3 Proof of Proposition 4.21

In order to prove Proposition 4.21 from Proposition 4.20, we establish the following lemmas. Recall that $\mathbf{G}^* \equiv \mathbf{G}^*(n, \mathbf{m})$, where $\mathbf{m} \sim \text{Poi}(dn/k)$.

Lemma 7.3. *There exists a constant $C \equiv C_\Psi > 0$ such that for any $m \geq 1$, $|\log L(G)| \leq Cm$ holds for any factor graph G with m clauses. In particular, we have $\mathbb{E}|\log L(\mathbf{G}^*)| \leq Cdn/k$.*

Proof. We have for a factor graph $G = (V, F, E, (\psi_a)_{a \in F})$ that

$$L(G) \equiv \sum_{\underline{\sigma} \in [q]^V} \mathbb{P}(\underline{\sigma}^* = \underline{\sigma}) \prod_{a \in F} \frac{\psi_a(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p, u}[\psi(\underline{\sigma}_\omega)]}.$$

Note that since $\psi(\cdot), \psi \in \Psi$ is bounded away from 0 and ∞ , there exist constants $C_i \equiv C_{i, \Psi} > 0, i = 1, 2$, such that for any $\underline{\sigma} \in [q]^V, \delta a \in V^k, \psi \in \Psi, \frac{\psi(\underline{\sigma}_{\delta a})}{\mathbb{E}_{p, u}[\psi(\underline{\sigma}_\omega)]} \in [C_1, C_2]$ holds. Thus, if G has m clauses, $L(G) \in [C_1^m, C_2^m]$ holds. Therefore, $|\log L(G)| \leq Cm$ for some constant $C \equiv C_\Psi > 0$. In particular, this implies that $\mathbb{E}|\log L(\mathbf{G}^*)| \leq C \cdot \mathbb{E}m = Cdn/k$, which concludes the proof. \square

Lemma 7.4. *For $d < d_*$, we have $\mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 \rightarrow 0$ as $n \rightarrow \infty$. Moreover, assuming (MIN), for any $\varepsilon > 0$, there exists $\delta \equiv \delta(\varepsilon, q, \pi) > 0$ and $n_0 \equiv n_0(\varepsilon, q, \pi)$ which only depends on ε, q, π such that the following holds: if $n \geq n_0$ and there exists an estimator $\hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}(\mathbf{G}^*)$ such that $\mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds, then $\mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 > \delta$ holds.*

Proof. To start with, invoking (103) for $F(x) = x$, we have that

$$\mathbb{E}_{\mathbf{G}^*, p, u} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} = \xi + O_{k, q, \Psi}(n^{-1/3}). \quad (104)$$

Thus, recalling the function $\mathcal{F}(R) \equiv \sum_{\underline{\sigma}, \underline{\tau} \in [q]^k} \mathbb{E}_p[\psi(\underline{\sigma})\psi(\underline{\tau})] \prod_{s=1}^k R(\sigma_s, \tau_s)$, it follows that

$$\mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 = \mathbb{E} \langle \mathcal{F}(R_{\underline{\sigma}^1, \underline{\sigma}^2}) \rangle_{\mathbf{G}^*} - \xi^2 + O_{k, q, \Psi}(n^{-1/3}). \quad (105)$$

Let us now assume $d < d_*$. Then, $\mathbb{E} \langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi \pi^\top\|_1 \rangle_{\mathbf{G}^*} \rightarrow 0$ as $n \rightarrow \infty$ by Theorem 3.10. Thus,

$$\mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 \rightarrow \mathbb{E} \mathcal{F}(\pi \pi^\top) - \xi^2 = 0.$$

To prove the second claim, assume that the condition (MIN) and fix $\varepsilon > 0$. Since \mathbf{G}^* satisfies (EXG) (cf. Lemma 5.2), Lemma 5.10 shows that there exists $\eta \equiv \eta(\varepsilon, q, \pi)$ and $n_0 \equiv n_0(\varepsilon, q, \pi)$ such that if $\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] \geq \frac{1}{q} + \varepsilon$ holds for some $\hat{\sigma} \equiv \hat{\sigma}(\mathbf{G}^*)$, then $\mathbb{E}\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 \rangle_{\mathbf{G}^*} \geq \eta$. Note that since $\|R_{\underline{\sigma}, \underline{\tau}} - \pi\pi^\top\|_1 \leq q$ for $\underline{\sigma}, \underline{\tau} \in [q]^V$, $\mathbb{E}\langle \|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 \rangle_{\mathbf{G}^*} \geq \eta$ implies that

$$\mathbb{E}\left\langle \mathbb{1}\left\{\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 > \frac{\eta}{2}\right\} \right\rangle_{\mathbf{G}^*} \geq \frac{\eta}{2q}.$$

Meanwhile, as shown in the proof of Lemma 7.2, there exists $C \equiv C_q > 0$ such that for $\ell = 1, 2$,

$$\mathbb{E}\left\langle \mathbb{1}\left\{\|R_{\underline{\sigma}^\ell} - \pi\|_1 > n^{-1/3}\right\} \right\rangle_{\mathbf{G}^*} \leq Cn^{-1/3},$$

which can be seen by conditioning on $\mathbf{m} = m$ and decomposing the LHS into the events where $\mathbf{G}^*(n, m) \in \mathcal{A}(n, m)$ or $\mathbf{G}^*(n, m) \notin \mathcal{A}(n, m)$. Thus, there exists a constant $c \equiv c_q > 0$ such that for large enough $n \geq n_0(\varepsilon, q, \pi)$,

$$\mathbb{E}\left\langle \mathbb{1}\left\{\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 > \frac{\eta}{2} \text{ and } \|R_{\underline{\sigma}^\ell} - \pi\|_1 \leq n^{-1/3} \text{ for } \ell = 1, 2\right\} \right\rangle_{\mathbf{G}^*} > c\eta. \quad (106)$$

Now, observe that by the condition (MIN), there exists $\epsilon \equiv \epsilon(\eta) > 0$ such that

$$\min \left\{ \mathcal{F}(R) : R \in [0, 1]^q, \quad R\mathbf{1} = R^\top\mathbf{1} = \pi, \quad \|R - \pi\pi^\top\| \geq \frac{\eta}{2} \right\} > \mathcal{F}(\pi\pi^\top) + \epsilon.$$

Thus, by continuity of $R \rightarrow \mathcal{F}(R)$, there exists small enough $\epsilon' \equiv \epsilon'(\eta) > 0$ such that

$$\min \left\{ \mathcal{F}(R) : R \in [0, 1]^q, \quad \|R\mathbf{1} - \pi\|_1 + \|R^\top\mathbf{1} - \pi\|_1 < \epsilon', \quad \|R - \pi\pi^\top\| \geq \frac{\eta}{2} \right\} > \mathcal{F}(\pi\pi^\top) + \epsilon'.$$

As a consequence, for any $\underline{\sigma}^1, \underline{\sigma}^2 \in [q]^V$, we have

$$\|R_{\underline{\sigma}^1, \underline{\sigma}^2} - \pi\pi^\top\|_1 > \frac{\eta}{2} \text{ and } \|R_{\underline{\sigma}^\ell} - \pi\|_1 \leq n^{-1/3} \text{ for } \ell = 1, 2 \implies \mathcal{F}(R_{\underline{\sigma}^1, \underline{\sigma}^2}) > \xi^2 + \epsilon', \quad (107)$$

where we used the fact that $\mathcal{F}(\pi\pi^\top) = \xi^2$. Finally, recalling the estimate (105), we have

$$\begin{aligned} \mathbb{E}_{\mathbf{G}^*, p, u} \left(\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi \right)^2 + O(n^{-1/3}) &= \mathbb{E}\langle \mathcal{F}(R_{\underline{\sigma}^1, \underline{\sigma}^2}) \rangle_{\mathbf{G}^*} - \xi^2 \\ &\geq \epsilon' \mathbb{E}\left\langle \mathbb{1}\left\{\mathcal{F}(R_{\underline{\sigma}^1, \underline{\sigma}^2}) \geq \xi^2 + \epsilon'\right\} \right\rangle_{\mathbf{G}^*} \\ &> c\epsilon'\eta, \end{aligned}$$

where we used (MIN) in the second inequality and the last inequality is by (106) and (107). Therefore, taking $\delta = c\epsilon'\eta$ completes the proof. \square

Proof of Proposition 4.21. The statements $\frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) \geq 0$ and $\mathbb{E} \log L(\mathbf{G}^*) \geq 0$ follows from Jensen's inequality. Indeed, for ease of notations, let $F_0(x) = x \log x$. Since $F_0(\cdot)$ is convex,

$$\mathbb{E} \log L(\mathbf{G}^*) = \mathbb{E} L(\mathbf{G}) \log L(\mathbf{G}) \geq F_0(\mathbb{E} L(\mathbf{G})) = F_0(1) = 0,$$

where the first equality follows from change of measure (see also (42)). Similarly, by Proposition 4.20,

$$\frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) = \frac{1}{k} \mathbb{E}_{\mathbf{G}^*, p, u} F_0 \left(\left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p, u}[\psi'(\underline{\sigma}_{\omega'})]} \right\rangle_{\mathbf{G}^*} \right) \geq \frac{1}{k} F_0 \left(\mathbb{E}_{\mathbf{G}^*, p, u} \left\langle \frac{\psi(\underline{\sigma}_\omega)}{\mathbb{E}_{p, u}[\psi'(\underline{\sigma}_{\omega'})]} \right\rangle_{\mathbf{G}^*} \right) = 0.$$

To prove the second statement, consider $d_0 < d_*$. Note that for any $0 < d \leq d_0$, we have by Proposition 4.20 that

$$\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) = \frac{1}{k} \cdot \mathbb{E}_{\mathbf{G}^*, p, u} F_0 \left(\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} \right) + o_n(1) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (108)$$

where the convergence holds since $\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} \xrightarrow{p} 0$ by Lemma 7.4 and $(\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*})_{n \geq 1}$ is bounded away from 0 and ∞ . Moreover, note that $\lim_{d \rightarrow 0} \mathbb{E} \log L(\mathbf{G}^*) = 0$ holds for fixed $n \geq 1$ by Lemma 7.3. Thus, by fundamental theorem of calculus, we have

$$\frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) \Big|_{d=d_0} = \int_0^{d_0} \frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) dd \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

where the convergence holds by (108) and dominated convergence theorem, since the equality (50) in Proposition 4.20 shows that $\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*)$ is uniformly bounded for $n \geq 1$ and $d \leq d_0$.

Next, we prove the third statement. Assume that (MIN) holds and that $\mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds for some estimator $\hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}_n(\mathbf{G}^*)$ and $\varepsilon > 0$. Then, Lemma 7.4 yields that under this assumption, $\mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 > \delta$ holds for $n \geq n_0$ for some $n_0 \equiv n_0(\varepsilon, q, \pi)$ and $\delta \equiv \delta(\varepsilon, q, \pi) > 0$. Moreover, since $F_0(x) = x \log x$ is strongly convex with $F_0(1) = 0, F'_0(1) = 1$, there exists a universal constant $c > 0$ such that $F_0(x) \geq x - 1 + C(x - 1)^2$ holds for $x > 0$. Thus, for $n \geq n_0$,

$$\begin{aligned} \mathbb{E}_{\mathbf{G}^*, p, u} F_0 \left(\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} \right) &\geq \mathbb{E}_{\mathbf{G}^*, p, u} [\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - 1] + C\xi^{-2} \cdot \mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 \\ &= C\xi^{-2} \cdot \mathbb{E}_{\mathbf{G}^*, p, u} (\langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} - \xi)^2 + O_{k, q, \Psi}(n^{-1/3}) \\ &\geq C\xi^{-2} \delta + O_{k, q, \Psi}(n^{-1/3}), \end{aligned}$$

where the equality follows from (104). Combining with Proposition 4.20, it follows that

$$\frac{1}{n} \frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) = k^{-1} \mathbb{E}_{\mathbf{G}^*, p, u} F_0 \left(\xi^{-1} \langle \psi(\underline{\sigma}_\omega) \rangle_{\mathbf{G}^*} \right) + O_{k, q, \Psi}(n^{-1/3}) \geq C\xi^{-2} k^{-1} \delta + O_{k, q, \Psi}(n^{-1/3}),$$

thus letting $\eta \equiv C\xi^{-2} k^{-1} \delta / 2$ concludes the proof of the third statement.

Finally, we prove the fourth statement under (MIN) condition. To this end, consider $d > d_*$. Then, there exists $d_0 \in (d_*, d)$, and since weak recovery is possible at d_0 , there exists a subsequence $(n_\ell)_{\ell \geq 1}$ and $\varepsilon > 0$ such that for any $\ell \geq 1$, $\mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon$ holds for some estimator $\hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}_{n_\ell}(\mathbf{G}_0^*)$ where $\mathbf{G}_0^* \sim \mathcal{G}_{\text{plant}}(n_\ell, d_0, p, \pi)$. Observe that this implies that for any fixed $\ell \geq 1$ and $\check{d} \in (d_0, d)$,

$$\mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] \geq \frac{1}{q} + \varepsilon \quad \text{for some } \hat{\underline{\sigma}} \equiv \hat{\underline{\sigma}}_{n_\ell}(\check{\mathbf{G}}^*) \quad \text{where } \check{\mathbf{G}}^* \sim \mathcal{G}_{\text{plant}}(n_\ell, \check{d}, p, \pi)$$

since subsampling the clauses independently with probability \check{d}/d from $\check{\mathbf{G}}^* \sim \mathcal{G}_{\text{plant}}(n_\ell, \check{d}, p, \pi)$ gives a sample drawn from $\mathcal{G}_{\text{plant}}(n_\ell, d_0, p, \pi)$ by Poisson thinning. Hence, by the third statement which

we established in the previous paragraph, there exists $\eta \equiv \eta(\varepsilon)$ and $n_0 \equiv n_0(\varepsilon)$ such that for all $n_\ell \geq n_0$ and $\check{d} \in (d_0, d)$, $\frac{1}{n_\ell} \frac{\partial}{\partial d} \mathbb{E} \log L(\check{\mathbf{G}}^*) \geq \eta$ holds. Since we proved in the first statement that $\frac{\partial}{\partial d} \mathbb{E} \log L(\mathbf{G}^*) \geq 0$ holds in general, it follows that for all ℓ large enough so that $n_\ell \geq n_0$,

$$\frac{1}{n} \mathbb{E} \log L(\mathbf{G}^*) \geq \eta(d - d_0) \quad \text{where} \quad \mathbf{G}^* \sim \mathcal{G}_{\text{plant}}(n_\ell, d, p, \pi),$$

which concludes the proof. □

Acknowledgements

E.M. and Y.S. are supported by Simons-NSF collaboration on deep learning NSF DMS-2031883 and Vannevar Bush Faculty Fellowship award ONR-N00014-20-1-2826. E.M. is also supported by ARO MURI W911NF1910217 and a Simons Investigator Award in Mathematics (622132). A.S. is supported by NSF grants DMS-1855527, DMS-1749103, a Simons Investigator grant, and a MacArthur Fellowship.

References

- [Abb18] Emmanuel Abbe, *Community detection and stochastic block models: Recent developments*, Journal of Machine Learning Research **18** (2018), no. 177, 1–86.
- [ACKZ15] Maria Chiara Angelini, Francesco Caltagirone, Florent Krzakala, and Lenka Zdeborová, *Spectral detection on sparse hypergraphs*, 53rd Annual Allerton Conference on Communication, Control, and Computing, Allerton 2015, Allerton Park & Retreat Center, Monticello, IL, USA, September 29 - October 2, 2015, IEEE, 2015, pp. 66–73.
- [ACO08] Dimitris Achlioptas and Amin Coja-Oghlan, *Algorithmic barriers from phase transitions*, 2008 49th Annual IEEE Symposium on Foundations of Computer Science, 2008, pp. 793–802.
- [AJM05] Dimitris Achlioptas, Haixia Jia, and Cristopher Moore, *Hiding satisfying assignments: Two are better than one*, J. Artif. Int. Res. **24** (2005), no. 1, 623–639.
- [AS15] Emmanuel Abbe and Colin Sandon, *Community detection in general stochastic block models: Fundamental limits and efficient algorithms for recovery*, 2015 IEEE 56th Annual Symposium on Foundations of Computer Science, IEEE, 2015, pp. 670–688.
- [AS16] ———, *Crossing the ks threshold in the stochastic block model with information theory*, 2016 IEEE International Symposium on Information Theory (ISIT), IEEE, 2016, pp. 840–844.
- [AS18] ———, *Proof of the achievability conjectures for the general stochastic block model*, Communications on Pure and Applied Mathematics **71** (2018), no. 7, 1334–1406.
- [BB20] Matthew Brennan and Guy Bresler, *Reducibility and statistical-computational gaps from secret leakage*, Conference on Learning Theory, PMLR, 2020, pp. 648–847.

- [BC09] P.J. Bickel and A. Chen, *A nonparametric view of network models and Newman-Girvan and other modularities*, Proceedings of the National Academy of Sciences **106** (2009), no. 50, 21068–21073.
- [BCO16] Victor Bapst and Amin Coja-Oghlan, *Harnessing the bethe free energy*, Random Structures & Algorithms **49** (2016), no. 4, 694–741.
- [BH21] Guy Bresler and Brice Huang, *The algorithmic phase transition of random k -sat for low degree polynomials*, FOCS, 2021.
- [BJR07] Béla Bollobás, Svante Janson, and Oliver Riordan, *The phase transition in inhomogeneous random graphs*, Random Structures & Algorithms **31** (2007), no. 1, 3–122.
- [BKW20] Afonso S Bandeira, Dmitriy Kunisky, and Alexander S Wein, *Computational hardness of certifying bounds on constrained pca problems*, ITCS, 2020.
- [BLM15] Charles Bordenave, Marc Lelarge, and Laurent Massoulié, *Non-backtracking spectrum of random graphs: community detection and non-regular ramanujan graphs*, Foundations of Computer Science (FOCS), 2015 IEEE 56th Annual Symposium on, IEEE, 2015, pp. 1347–1357.
- [BMNN16] Jess Banks, Cristopher Moore, Joe Neeman, and Praneeth Netrapalli, *Information-theoretic thresholds for community detection in sparse networks*, Conference on Learning Theory, PMLR, 2016, pp. 383–416.
- [BMR21] Jess Banks, Sidhanth Mohanty, and Prasad Raghavendra, *Local statistics, semidefinite programming, and community detection*, pp. 1298–1316, 2021.
- [BP22] Jean Barbier and Dmitry Panchenko, *Strong replica symmetry in high-dimensional optimal bayesian inference*, Communications in Mathematical Physics **393** (2022), no. 3, 1199–1239.
- [BPS21] Jean Barbier, Dmitry Panchenko, and Manuel Sáenz, *Strong replica symmetry for high-dimensional disordered log-concave Gibbs measures*, Information and Inference: A Journal of the IMA **11** (2021), no. 3, 1079–1108.
- [CK01] A. Condon and R.M. Karp, *Algorithms for graph partitioning on the planted partition model*, Random Structures and Algorithms **18** (2001), no. 2, 116–140.
- [CO10] A. Coja-Oghlan, *Graph partitioning via adaptive spectral techniques*, Combinatorics, Probability and Computing **19** (2010), no. 02, 227–284.
- [COEJ⁺18] Amin Coja-Oghlan, Charilaos Efthymiou, Nor Jaafari, Mihyun Kang, and Tobias Kapetanopoulos, *Charting the replica symmetric phase*, Communications in Mathematical Physics **359** (2018), no. 2, 603–698.
- [COKM20] Amin Coja-Oghlan, Tobias Kapetanopoulos, and Noela Müller, *The replica symmetric phase of random constraint satisfaction problems*, Combinatorics, Probability and Computing **29** (2020), no. 3, 346–422.

- [COKPZ18] Amin Coja-Oghlan, Florent Krzakala, Will Perkins, and Lenka Zdeborová, *Information-theoretic thresholds from the cavity method*, *Advances in Mathematics* **333** (2018), 694–795.
- [COW18] Amin Coja-Oghlan and Nick Wormald, *The number of satisfying assignments of random regular k -SAT formulas*, *Combin. Probab. Comput.* **27** (2018), no. 4, 496–530.
- [DAM16] Yash Deshpande, Emmanuel Abbe, and Andrea Montanari, *Asymptotic mutual information for the balanced binary stochastic block model*, *Information and Inference: A Journal of the IMA* **6** (2016), no. 2, 125–170.
- [DF89] M.E. Dyer and A.M. Frieze, *The solution of some random NP-hard problems in polynomial expected time*, *Journal of Algorithms* **10** (1989), no. 4, 451–489.
- [DKMZ11] A. Decelle, F. Krzakala, C. Moore, and L. Zdeborová, *Asymptotic analysis of the stochastic block model for modular networks and its algorithmic applications*, *Physics Review E* **84** (2011), 066106.
- [DMR11] C. Daskalakis, E. Mossel, and S. Roch, *Evolutionary trees and the ising model on the bethe lattice: a proof of steel’s conjecture*, *PTRF* **149** (2011), no. 1-2, 149–189.
- [ER⁺60] Paul Erdos, Alfréd Rényi, et al., *On the evolution of random graphs*, *Publ. Math. Inst. Hung. Acad. Sci* **5** (1960), no. 1, 17–60.
- [FPV15] Vitaly Feldman, Will Perkins, and Santosh Vempala, *On the complexity of random satisfiability problems with planted solutions*, *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing (New York, NY, USA), STOC ’15*, Association for Computing Machinery, 2015, p. 77–86.
- [GD17a] Debarghya Ghoshdastidar and Ambedkar Dukkipati, *Consistency of spectral hypergraph partitioning under planted partition model*, *The Annals of Statistics* **45** (2017), no. 1, 289–315.
- [GD17b] ———, *Uniform hypergraph partitioning: Provable tensor methods and sampling techniques*, *Journal of Machine Learning Research* **18** (2017), no. 50, 1–41.
- [GJ07] Leslie Ann Goldberg and Mark Jerrum, *Inapproximability of the tutte polynomial*, *Proceedings of the thirty-ninth annual ACM symposium on Theory of computing*, 2007, pp. 459–468.
- [GJW20] David Gamarnik, Aukosh Jagannath, and Alexander S Wein, *Low-degree hardness of random optimization problems*, *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, IEEE, 2020, pp. 131–140.
- [GP23] Yuzhou Gu and Yury Polyanskiy, *Weak recovery threshold for the hypergraph stochastic block model*, *arXiv preprint arXiv:2303.14689* (2023).
- [GS14] David Gamarnik and Madhu Sudan, *Limits of local algorithms over sparse random graphs*, *Proceedings of the 5th conference on Innovations in theoretical computer science*, 2014, pp. 369–376.

- [GWSV11] Dongning Guo, Yihong Wu, Shlomo S Shitz, and Sergio Verdú, *Estimation in gaussian noise: Properties of the minimum mean-square error*, IEEE Transactions on Information Theory **57** (2011), no. 4, 2371–2385.
- [HLL83] P.W. Holland, K.B. Laskey, and S. Leinhardt, *Stochastic blockmodels: First steps*, Social Networks **5** (1983), no. 2, 109 – 137.
- [Hop18] Samuel Hopkins, *Statistical inference and the sum of squares method*, Ph.D. thesis, Cornell University, 2018.
- [HS17] Samuel Hopkins and David Steurer, *Efficient bayesian estimation from few samples: community detection and related problems*, 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), IEEE, 2017, pp. 379–390.
- [HW20] Justin Holmgren and Alexander S Wein, *Counterexamples to the low-degree conjecture*, ITCS, 2020.
- [Ist00] Sorin Istrail, *Statistical mechanics, three-dimensionality and np-completeness: I. universality of intracatability for the partition function of the ising model across non-planar surfaces (extended abstract)*, Proceedings of the Thirty-Second Annual ACM Symposium on Theory of Computing (New York, NY, USA), STOC '00, Association for Computing Machinery, 2000, p. 87–96.
- [Jan95] Svante Janson, *Random regular graphs: Asymptotic distributions and contiguity*, Combinatorics, Probability and Computing **4** (1995), no. 4, 369–405.
- [Jer92] Mark Jerrum, *Large cliques elude the metropolis process*, Random Structures & Algorithms **3** (1992), no. 4, 347–359.
- [JLuR00] Svante Janson, Tomasz Łuczak, and Andrzej Ruciński, *Random graphs*, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000. MR 1782847
- [JMS07] Haixia Jia, Cristopher Moore, and Doug Strain, *Generating hard satisfiable formulas by hiding solutions deceptively*, J. Artif. Int. Res. **28** (2007), no. 1, 107–118.
- [JS93] Mark Jerrum and Alistair Sinclair, *Polynomial-time approximation algorithms for the ising model*, SIAM Journal on computing **22** (1993), no. 5, 1087–1116.
- [JS98] M. Jerrum and G.B. Sorkin, *The Metropolis algorithm for graph bisection*, Discrete Applied Mathematics **82** (1998), no. 1-3, 155–175.
- [Kea98] Michael Kearns, *Efficient noise-tolerant learning from statistical queries*, Journal of the ACM (JACM) **45** (1998), no. 6, 983–1006.
- [KMM⁺13] F. Krzakala, C. Moore, E. Mossel, J. Neeman, A. Sly, Zdeborova L, and P. Zhang, *Spectral redemption: clustering sparse networks*, PNAS **100** (2013), no. 52, 20935–20940.

- [KMZ14] Florent Krzakala, Marc Mézard, and Lenka Zdeborová, *Reweighted belief propagation and quiet planting for random K -SAT*, J. Satisf. Boolean Model. Comput. **8** (2014), no. 3/4, 149–171.
- [KS66] H. Kesten and B. P. Stigum, *Additional limit theorems for indecomposable multidimensional Galton-Watson processes*, Ann. Math. Statist. **37** (1966), 1463–1481.
- [Kuč95] Luděk Kučera, *Expected complexity of graph partitioning problems*, Discrete Applied Mathematics **57** (1995), no. 2-3, 193–212.
- [KWB19] Dmitriy Kunisky, Alexander S Wein, and Afonso S Bandeira, *Notes on computational hardness of hypothesis testing: Predictions using the low-degree likelihood ratio*, arXiv preprint arXiv:1907.11636 (2019).
- [KZ09] Florent Krzakala and Lenka Zdeborová, *Hiding quiet solutions in random constraint satisfaction problems*, Phys. Rev. Lett. **102** (2009), 238701.
- [LMX15] Marc Lelarge, Laurent Massoulié, and Jiaming Xu, *Reconstruction in the labelled stochastic block model*, IEEE Transactions on Network Science and Engineering **2** (2015), no. 4, 152–163.
- [LPW08] D.A. Levin, Y. Peres, and E.L. Wilmer, *Markov chains and mixing times*, American Mathematical Soc., 2008.
- [LR05] E. L. Lehmann and Joseph P. Romano, *Testing statistical hypotheses*, third ed., Springer Texts in Statistics, Springer, New York, 2005. MR MR2135927 (2006m:62005)
- [Mas13] Laurent Massoulié, *Community detection thresholds and the weak ramanujan property*, arXiv preprint arXiv:1311.3085, 2013.
- [Mas14] Laurent Massoulié, *Community detection thresholds and the weak ramanujan property*, Proceedings of the forty-sixth annual ACM symposium on Theory of computing, ACM, 2014, pp. 694–703.
- [McS01] Frank McSherry, *Spectral partitioning of random graphs*, Foundations of Computer Science, 2001. Proceedings. 42nd IEEE Symposium on, IEEE, 2001, pp. 529–537.
- [MM09] M. Mézard and A. Montanari, *Information, physics, and computation*, Oxford University Press, USA, 2009.
- [MNS15] E. Mossel, J. Neeman, and A. Sly, *Reconstruction and estimation in the planted partition model*, Probability Theory and Related Fields (2015), no. 3-4, 431–461, The Arxiv version of this paper is titled Stochastic Block Models and Reconstruction.
- [MNS18] Elchanan Mossel, Joe Neeman, and Allan Sly, *A proof of the block model threshold conjecture*, Combinatorica **38** (2018), no. 3, 665–708.
- [Mos04] E. Mossel, *Survey: Information flow on trees*, Graphs, Morphisms and Statistical Physics. DIMACS series in discrete mathematics and theoretical computer science (J. Neštril and P. Winkler, eds.), 2004, pp. 155–170. MR MR2056226

- [MP92] Arak Mathai and Serge Provost, *Quadratic forms in random variables*, Statistics: textbooks and monographs, vol. 126, Marcel Dekker, 01 1992.
- [MRS11] E. Mossel, S. Roch, and A. Sly, *On the inference of large phylogenies with long branches: How long is too long?*, Bull. Math. Bio. **73** (2011), no. 7, 1627–1644.
- [MSS23] Elchanan Mossel, Allan Sly, and Youngtak Sohn, *Exact phase transitions for stochastic block models and reconstruction on trees*, Proceedings of the 55th Annual ACM Symposium on Theory of Computing (New York, NY, USA), STOC 2023, Association for Computing Machinery, 2023, p. 96–102.
- [MW23] Ankur Moitra and Alex Wein, *Precise error rates for computationally efficient testing*, arXiv preprint, arXiv:2311.00289 (2023).
- [NSS24] Danny Nam, Allan Sly, and Youngtak Sohn, *One-step replica symmetry breaking of random regular NAE-SAT II*, Communications in Mathematical Physics **405** (2024), no. 3, 61.
- [PZ21] Soumik Pal and Yizhe Zhu, *Community detection in the sparse hypergraph stochastic block model*, Random Structures & Algorithms **59** (2021), no. 3, 407–463.
- [RCY11] Karl Rohe, Sourav Chatterjee, and Bin Yu, *Spectral clustering and the high-dimensional stochastic blockmodel*, The Annals of Statistics **39** (2011), no. 4, 1878–1915.
- [RS17] Sebastien Roch and Allan Sly, *Phase transition in the sample complexity of likelihood-based phylogeny inference*, Probability Theory and Related Fields **169** (2017), no. 1, 3–62.
- [RTSZ19] Federico Ricci-Tersenghi, Guilhem Semerjian, and Lenka Zdeborová, *Typology of phase transitions in Bayesian inference problems*, Phys. Rev. E **99** (2019), 042109.
- [SH12] Laurent Massoulié Simon Heimlicher, Marc Lelarge, *Community detection in the labelled stochastic block model*, arXiv preprint, arXiv:1209.2910 (2012).
- [SN97] T.A.B. Snijders and K. Nowicki, *Estimation and prediction for stochastic blockmodels for graphs with latent block structure*, Journal of Classification **14** (1997), no. 1, 75–100.
- [SZ22] L. Stephan and Y. Zhu, *Sparse random hypergraphs: Non-backtracking spectra and community detection*, 2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS) (Los Alamitos, CA, USA), IEEE Computer Society, nov 2022, pp. 567–575.
- [Vaa98] A. W. van der Vaart, *Asymptotic statistics*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, 1998.
- [Wei20] Alexander S Wein, *Optimal low-degree hardness of maximum independent set*, arXiv preprint arXiv:2010.06563 (2020).
- [Wor99] N. C. Wormald, *Models of random regular graphs*, London Mathematical Society Lecture Note Series, p. 239–298, Cambridge University Press, 1999.

[XJL20] Min Xu, Varun Jog, and Po-Ling Loh, *Optimal rates for community estimation in the weighted stochastic block model*, The Annals of Statistics **48** (2020), no. 1, 183 – 204.

A Nontriviality of the weak recovery threshold

In this section, we prove Proposition 4.1. We begin with the case where (SYM) does not hold. Then we must have some ψ , s and τ, τ' such that

$$\mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_s = \tau] > \mathbb{E}_{p,\pi}[\psi(\underline{\sigma})] > \mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_s = \tau'].$$

For each vertex v let d_v^ψ be the number of clauses of type ψ containing v . Define

$$x_i := \frac{dp(\psi) \sum_{\tau_1, \dots, \tau_{k-1}} \psi(\tau_1, \dots, \tau_{k-1}, i) \prod_{j=1}^{k-1} \pi_{\tau_j}}{\mathbb{E}_{p,\pi}[\psi(\underline{\sigma})]} = \frac{dp(\psi) \mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_s = i]}{\mathbb{E}_{p,\pi}[\psi(\underline{\sigma})]},$$

When $\sigma_v = i$ we have that d_v^ψ is Poisson with mean $(1 + o(1))x_i$. We take the very simple estimator

$$\hat{\sigma}_v = \begin{cases} \tau & d_v^\psi \geq dp(\psi) \\ \tau' & d_v^\psi < dp(\psi). \end{cases}$$

Then

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] &= \frac{1}{q\pi_\tau} \mathbb{P}[\sigma_v = \tau, d_v^\psi \geq dp(\psi)] + \frac{1}{q\pi_{\tau'}} \mathbb{P}[\sigma_v = \tau', d_v^\psi < dp(\psi)] \\ &= \frac{1}{q} (\mathbb{P}[\text{Pois}(x_\tau) \geq dp(\psi)] - \mathbb{P}[\text{Pois}(x_{\tau'}) < dp(\psi)]) \\ &= \frac{1}{q} + \frac{1}{q} (\mathbb{P}[\text{Pois}(x_\tau) \geq dp(\psi)] - \mathbb{P}[\text{Pois}(x_{\tau'}) \geq dp(\psi)]) \geq \frac{1}{q} + \epsilon, \end{aligned}$$

since $x_\tau > x_{\tau'}$ which establishes weak recovery whenever $d > 0$.

Now suppose that (SYM) holds and $d < \frac{1}{k-1}$. In this case the graph is subcritical and almost all the variables are in trees of size $O(1)$. We will consider the local weak limit of $(G, \underline{\sigma})$ under the planted model. A vertex of type i is connected to a Poisson with mean

$$\sum_{\psi} \frac{dp(\psi) \mathbb{E}_\pi[\psi(\underline{\sigma}) \mid \sigma_s = i]}{\mathbb{E}_{p,\pi}[\psi(\underline{\sigma})]} = dp(\psi)$$

neighbouring clauses. In particular, the distribution does not depend on the type i and so the law of the local neighbourhood is a branching process independent of the state of the root. Among all the trees in the graph with a fixed topology, a $\pi_i + o(1)$ will have state i at the root. Since the estimator cannot distinguish between trees with the same topology, it will be independent of $\underline{\sigma}$ and so $\mathbb{E}[A(\underline{\sigma}^*, \hat{\sigma})] = \frac{1}{q} + o(1)$. Hence weak recovery is impossible when $d < \frac{1}{k-1}$.

Finally, we show that for d large that weak recovery is possible. We define

$$\Psi(\underline{\sigma}) = \sum_a \psi_a(\sigma_{\delta a})$$

and take our estimator to be the maximizer

$$\hat{\underline{\sigma}} = \arg \max \Psi(\hat{\underline{\sigma}}).$$

among $\hat{\sigma}$ with empirical distribution $\pi + o(1)$. We will first show that when d is large enough that the joint distribution of $(\underline{\sigma}^*, \hat{\underline{\sigma}})$ is not close to the product measure.

Recall that we defined the map $\mathcal{F}(R') := \sum_{\underline{\sigma}, \underline{\tau} \in [q]^k} \mathbb{E}_p[\psi(\underline{\sigma})\psi(\underline{\tau})] \prod_{s=1}^k R'(\sigma_s, \tau_s)$ in (16). Let $R = \pi\pi^\top$ and $R_\star = \text{diag}(\pi_1, \dots, \pi_q)$. Then

$$\mathcal{F}(R_\star) - \mathcal{F}(R) = \sum_{\underline{\sigma}, \underline{\tau} \in [q]^k} \mathbb{E}_p[\psi(\underline{\sigma})\psi(\underline{\tau})] \prod_{s=1}^k \pi(\sigma_s)\pi(\tau_s) - \sum_{\underline{\sigma}, \underline{\tau} \in [q]^k} \mathbb{E}_p[\psi(\underline{\sigma})\psi(\underline{\tau})] \prod_{s=1}^k \pi(\sigma_s)\pi(\tau_s)$$

This can be expressed as $\sum_{\psi} p(\psi)\psi(X_1, \dots, X_k)$ where X_i are IID according to π and since at least one of the ψ is not constant we have that this variance is strictly positive and hence $\mathcal{F}(R_\star) > \mathcal{F}(R)$. Let $\underline{\sigma}^*$ be fixed and let $\underline{\tau} \in [q]^V$ be a fixed configuration. Define their joint empirical distribution as

$$W_{ij} = W_{ij}(\underline{\sigma}^*, \underline{\tau}) = \frac{1}{n} |\{v \in V : \sigma_v^* = i, \tau_v = j\}|.$$

The probability that a clause of type ψ connects variables v_1, \dots, v_k is

$$\frac{d}{Z \binom{n}{k}} \psi(\sigma_{v_1}^*, \dots, \sigma_{v_k}^*)$$

for some normalizing constant Z and so

$$\mathbb{E}[\Psi(\underline{\tau})] = \frac{d}{Z \binom{n}{k}} \sum_{v_1, \dots, v_k} \psi(\sigma_{v_1}^*, \dots, \sigma_{v_k}^*) \psi(\tau_{v_1}, \dots, \tau_{v_k}) = \frac{dn}{Z} \mathcal{F}(W) + o(n).$$

and

$$\mathbb{E}[\Psi(\underline{\sigma}^*)] = \frac{dn}{Z} \mathcal{F}(R_\star) + o(n).$$

If we reveal each clause one by one, each step affects the value of $\Psi(\underline{\tau})$ by $O(1)$ so by the Azuma-Hoeffding Inequality,

$$\mathbb{P}[|\Psi(\underline{\tau}) - \mathbb{E}[\Psi(\underline{\tau})]| \geq den] \leq 2 \exp(-C\epsilon^2 dn)$$

By the continuity of \mathcal{F} , for some $\delta > 0$ sufficiently small we have that

$$\inf_{R': \|R' - R\| \leq \delta} \mathcal{F}(R_\star) - \mathcal{F}(R') \geq \delta.$$

Then

$$\begin{aligned} \mathbb{P}[\|W(\underline{\sigma}^*, \hat{\underline{\sigma}}) - R\| < \delta] &\leq \mathbb{P}\left[|\Psi(\underline{\sigma}) - \mathbb{E}[\Psi(\underline{\sigma})]| \geq \frac{d\delta n}{3Z}\right] + \sum_{\underline{\tau}: \|W(\underline{\sigma}^*, \underline{\tau}) - R\| \leq \delta} \mathbb{P}\left[|\Psi(\underline{\tau}) - \mathbb{E}[\Psi(\underline{\tau})]| \geq \frac{d\delta n}{3Z}\right] \\ &\leq 2q^n \exp(-C(\delta/3Z)^2 dn) = o(1), \end{aligned}$$

for large enough d . Hence, we have that with high probability $W(\underline{\sigma}^*, \hat{\underline{\sigma}})$ is at least δ away from $\pi\pi^\top$. The performance of our estimator can be written as

$$\limsup_{n \rightarrow \infty} \mathbb{E}[A(\underline{\sigma}^*, \hat{\underline{\sigma}})] = \limsup_{n \rightarrow \infty} \mathbb{E} \max_{\Gamma \in S_q} \left\{ \frac{1}{q} \sum_{i=1}^q \frac{W_{i, \Gamma(i)}(\underline{\sigma}^*, \hat{\underline{\sigma}})}{\pi_i} \right\}.$$

Let us write

$$\Upsilon(R') = \max_{\Gamma \in S_q} \left\{ \frac{1}{q} \sum_{i=1}^q \frac{R'_{i,\Gamma(i)}}{\pi_i} \right\}$$

Let

$$\mathcal{R}_\pi^\delta = \{R' : \|R' - R\| \geq \delta, \sum_i R_{ij} = \pi_j, \sum_j R_{ij} = \pi_i\}$$

Since w.h.p. we have that $W(\underline{\sigma}^*, \hat{\underline{\sigma}})$ is $o(1)$ distance from \mathcal{R}_π^δ , to prove weak recovery it is enough to show that

$$\inf_{R' \in \mathcal{R}} \Upsilon(R') \geq \frac{1}{q} + \epsilon. \quad (109)$$

Since \mathcal{R} is compact it is enough to show that $\Upsilon(R') > \frac{1}{q}$ for all $R' \in \mathcal{R}_\pi^\delta$. First we have that for any $R' \in \mathcal{R}_\pi$ that

$$\Upsilon(R') \geq \frac{1}{q!} \sum_{\Gamma \in S_q} \frac{1}{q} \sum_{i=1}^q \frac{R'_{i,\Gamma(i)}}{\pi_i} = \frac{1}{q^2} \sum_{i=1}^q \frac{\sum_{i'=1}^q R'_{i,i'}}{\pi_i} = \frac{1}{q}$$

since the max is always at least the average and since every $\Gamma(i)$ is equal to i' for a $\frac{1}{q}$ fraction of Γ . So if $\Upsilon(R') = \frac{1}{q}$ then we must have

$$\forall \Gamma : \frac{1}{q} \sum_{i=1}^q \frac{R'_{i,\Gamma(i)}}{\pi_i} = \frac{1}{q}.$$

In particular, switching two entries of Γ cannot change the above expression so for any i, i', j, j'

$$\frac{R_{ij}}{\pi_i} - \frac{R_{ij'}}{\pi_i} = \frac{R_{i'j}}{\pi_{i'}} - \frac{R_{i'j'}}{\pi_{i'}}.$$

But this can only occur if all the rows of R_{ij}/π_i are the same in which case $R' = \pi\pi^T$. Hence, (109) holds and for large enough d we have weak recovery.

B Hypothesis testing below the weak recovery threshold

In this section, we prove Lemmas 4.13, 4.15, and 4.17.

B.1 Proof of Lemma 4.13

We show that

$$\mathbf{L}_\infty \equiv \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta\infty}} e^{-\lambda_\zeta \delta_\zeta} \right\}.$$

is discrete if and only if $d \in (0, \frac{1}{k-1})$. The case where $d < \frac{1}{k-1}$ is straightforward as in this case

$$\sum_{\ell \geq 1} \sum_{\zeta \in S_\ell} \mathbb{P}[X_{\zeta\infty} \geq 1] \leq \sum_{\ell \geq 1} \sum_{\zeta \in S_\ell} \mathbb{E}[X_{\zeta\infty}] < \infty$$

and so by Borel-Cantelli, only finitely many $X_{\zeta\infty}$ are non-zero and so \mathbf{L}_∞ takes values in the set given in equation (43) and hence is discrete. When $d \geq \frac{1}{k-1}$ we have that \mathbf{L}_∞ is the exponential

of a weighted sum of Poisson random variables the sum of whose means is infinite and so we can write

$$\log(\mathbf{L}_\infty) = \lim_{T \rightarrow \infty} \int_0^T f(t) dN_t - g(T)$$

where $f(t)$ and $g(t)$ are deterministic functions, $f(t)$ is strictly positive and N_t is a standard Poisson process. We will show that any such random variable must be continuous. Define

$$R(K, t) = \max_{x_1 < x_2 < \dots < x_k \in \mathbb{R}} \sum_{i=1}^K \mathbb{P} \left[\int_0^T f(t) dN_t = x_i \right],$$

that is the maximum probability $\int_0^T f(t) dN_t$ puts on k distinct points. We claim that,

$$\left. \frac{d}{dt} R(K, t) \right|_{t=T} \leq -(R(K, T) - R(K-1, T)) + (R(K+1, T) - R(K, T)). \quad (110)$$

To see this, suppose that $\mathcal{X} = \{x_1, x_2, \dots, x_K\}$ is some maximizing set of k atoms at time T . Note that at time t the size of the j -th largest atom is $R(j, t) - R(j-1, t)$. Between time t and $t + \delta t$ there is probability δt of a new Poisson point which increases the integral by $f(t)$. Thus we have that

$$\begin{aligned} \left. \frac{d}{dt} \sum_{x \in \mathcal{X}} \mathbb{P} \left[\int_0^T f(t) dN_t = x \right] \right|_{t=T} &= - \sum_{x \in \mathcal{X}} (\mathbb{P} \left[\int_0^T f(t) dN_t = x \right] - \mathbb{P} \left[\int_0^T f(t) dN_t = x - f(t) \right]) \\ &= - \sum_{x \in \mathcal{X}} \mathbb{P} \left[\int_0^T f(t) dN_t = x \right] + \sum_{y: y+f(T) \in \mathcal{X}} \mathbb{P} \left[\int_0^T f(t) dN_t = y \right] \\ &= - \sum_{x \in \mathcal{X}: x-f(T) \notin \mathcal{X}} \mathbb{P} \left[\int_0^T f(t) dN_t = x \right] + \sum_{y \notin \mathcal{X}: y+f(T) \in \mathcal{X}} \mathbb{P} \left[\int_0^T f(t) dN_t = y \right] \\ &\leq -|\{x \in \mathcal{X} : x - f(T) \notin \mathcal{X}\}|(R(K, T) - R(K-1, T)) \\ &\quad + |\{y \notin \mathcal{X} : y + f(T) \in \mathcal{X}\}|(R(K+1, T) - R(K, T)) \end{aligned}$$

where the third equality holds by canceling out terms that appear in the first and second sums while the inequality holds by noting that every atom in \mathcal{X} is one of the largest K atoms while every atom in \mathcal{X}^c is at most the $(K+1)$ -th biggest. Note that

$$|\{x \in \mathcal{X} : x - f(T) \notin \mathcal{X}\}| = |\{y \notin \mathcal{X} : y + f(T) \in \mathcal{X}\}|$$

and both are at least of size one, since x_1 is not in the former. Hence we have equation (110). Also since $R(j, t) - R(j-1, t)$ is the size of the j -th atoms it is decreasing in j so we have in particular that

$$\left. \frac{d}{dt} R(K, t) \right|_{t=T} \leq 0.$$

Suppose that $R(1, t) \rightarrow q > 0$ as $t \rightarrow \infty$. Then set $K = \lceil \frac{1}{q} \rceil$. Then

$$\begin{aligned} \frac{d}{dt} \sum_{j=1}^K R(j, t) &\leq \sum_{j=1}^K -(R(j, t) - R(j-1, t)) + (R(j+1, t) - R(j, t)) \\ &= -R(1, t) + (R(K+1, t) - R(K, t)) \\ &\leq -q + \frac{1}{K+1} < 0. \end{aligned}$$

But $\sum_{j=1}^K R(j, 0) = K$ we have that $\sum_{j=1}^K R(j, t) \leq K - t(q - (K + 1)^{-1})$ for all t which contradicts the fact that it is non-negative. Hence $R(1, t) \rightarrow 0$ as $t \rightarrow \infty$. Now if X and Y are independent random variables,

$$\max_x \mathbb{P}[X + Y = x] \leq \max_x \mathbb{P}[X = x]$$

and so

$$\max_x \mathbb{P}[\log(\mathbf{L}_\infty) = x] \leq \inf_t R(1, t) = 0.$$

Hence \mathbf{L}_∞ has a continuous distribution.

B.2 Proof of Lemma 4.15

Fix $d < d_{\kappa_S}$ and assume (SYM). We only prove the weak convergence

$$T_n(\mathbf{G}^*) \xrightarrow{d} \mathbf{L}_\infty^* \equiv \prod_{\ell \geq 1} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta, \infty}^*} e^{-\lambda_\zeta \delta_\zeta} \right\}, \quad (111)$$

since the weak convergence for $T_n(\mathbf{G})$ can be established using the same argument. To start with, for $K \geq 1$, let us denote

$$T_{n,K}(G) := \prod_{1 \leq \ell \leq K} \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_\zeta(G)} e^{-\lambda_\zeta \delta_\zeta} \right\}.$$

Thus, $T_n(\mathbf{G}^*) \equiv T_{n,K_n}(\mathbf{G}^*)$ holds by definition. Observe that by considering a fixed $K \geq 1$, we have by Fact 4.9-(2) that as $n \rightarrow \infty$,

$$T_{n,K}(\mathbf{G}^*) \xrightarrow{d} \mathbf{L}_K^* := \prod_{\ell=1}^K \prod_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta)^{X_{\zeta, \infty}^*} e^{-\lambda_\zeta \delta_\zeta} \right\}.$$

Observe that since \mathbf{L}_K^* converges a.s. to \mathbf{L}_∞^* as $K \rightarrow \infty$, we have $\mathbf{L}_K^* \xrightarrow{d} \mathbf{L}_\infty^*$ holds. Thus, combining with the above weak convergence, a diagonal argument shows that there exists an arbitrarily slowly growing sequence $(K'_n)_{n \geq 1}$ such that $1 \ll K'_n \leq K_n$ holds and as $n \rightarrow \infty$,

$$T_{n,K'_n}(\mathbf{G}^*) \xrightarrow{d} \mathbf{L}_\infty^*.$$

Hence, in order to achieve our goal in Eq. (111), it suffices to prove that as $n \rightarrow \infty$,

$$X_n := \sum_{\ell=K'_n+1}^{K_n} \sum_{\zeta \in S_\ell} \left\{ \log(1 + \delta_\zeta) X_\zeta(\mathbf{G}^*) - \lambda_\zeta \delta_\zeta \right\} \xrightarrow{p} 0, \quad (112)$$

since $T_{n,K_n}(\mathbf{G}^*)/T_{n,K'_n}(\mathbf{G}^*) = e^{X_n}$ holds by definition. For the rest of the proof, we prove Eq. (112) by showing that $\mathbb{E}X_n^2 \rightarrow 0$ as $n \rightarrow \infty$. To this end, we calculate the moments of $(X_\zeta(\mathbf{G}^*))_{\zeta \in \cup_{\ell \leq K_n} S_\ell}$. Since the length of ζ is $2\ell \leq K_n \ll \log n$, it is standard to approximate the the first and second moments of $X_\zeta(\mathbf{G}^*)$ (see e.g. [COEJ⁺18, Eq. (8.7)]) as follows. For two distinct signatures $\zeta, \zeta' \in \cup_{\ell \leq K_n} S_\ell$, we have

$$\begin{aligned} \mathbb{E}X_\zeta(\mathbf{G}^*) &= \lambda_\zeta^* + O_{k,q,\Psi} \left(\frac{\log n}{\sqrt{n}} \right), & \text{Var}(X_\zeta(\mathbf{G}^*)) &= \lambda_\zeta^* + O_{k,q,\Psi} \left(\frac{\log n}{\sqrt{n}} \right), \\ \mathbb{E}X_\zeta(\mathbf{G}^*)X_{\zeta'}(\mathbf{G}^*) &= \lambda_\zeta^* \lambda_{\zeta'}^* + O_{k,q,\Psi} \left(\frac{\log n}{\sqrt{n}} \right), \end{aligned}$$

where the dominant error $\frac{\log n}{\sqrt{n}}$ comes from approximating the empirical distribution of $\underline{\sigma}^*$ by π . Moreover, note that the number of signatures $\zeta \in \cup_{\ell \leq K_n} S_\ell$ is at most $e^{O(\log \log n)} \ll n^{1/10}$ since $K_n = O(\log \log n)$. Thus, $\mathbb{E}X_n^2$ can be bounded by

$$\begin{aligned}
\mathbb{E}X_n^2 &\leq \sum_{\ell=K'_n+1}^{K_n} \sum_{\zeta \in S_\ell} \left\{ \log^2(1 + \delta_\zeta) \lambda_\zeta^* + (\log(1 + \delta_\zeta) \lambda_\zeta^* - \lambda_\zeta \delta_\zeta)^2 \right\} \\
&\quad + \sum_{\ell, \ell'=K'_n+1}^{K_n} \sum_{\zeta \in S_\ell, \zeta' \in S_{\ell'}} \left(\log(1 + \delta_\zeta) \lambda_\zeta^* - \lambda_\zeta \delta_\zeta \right) \left(\log(1 + \delta_{\zeta'}) \lambda_{\zeta'}^* - \lambda_{\zeta'} \delta_{\zeta'} \right) + O_{k,q,\Psi}(n^{-1/3}) \\
&\leq \sum_{\ell=K'_n+1}^{\infty} \sum_{\zeta \in S_\ell} \left\{ (1 + \delta_\zeta) \log^2(1 + \delta_\zeta) \lambda_\zeta + ((1 + \delta_\zeta) \log(1 + \delta_\zeta) - \delta_\zeta)^2 \lambda_\zeta^2 \right\} \\
&\quad + \left(\sum_{\ell=K'_n+1}^{\infty} \sum_{\zeta \in S_\ell} ((1 + \delta_\zeta) \log(1 + \delta_\zeta) - \delta_\zeta) \lambda_\zeta \right)^2 + O_{k,q,\Psi}(n^{-1/3}),
\end{aligned} \tag{113}$$

where the final inequality holds since $\lambda_\zeta^* = \lambda_\zeta(1 + \delta_\zeta)$. We upper bound the RHS by using a Taylor approximation w.r.t. δ_ζ . To do so, we first argue that $\sup_{\zeta \in S_\ell} |\delta_\zeta| \rightarrow 0$ as $\ell \rightarrow \infty$, which can be argued using the following lemma.

Lemma B.1. *Consider a triangular array of stochastic matrices $(P_{n,m})_{1 \leq m \leq a_n, n \geq 1} \in \mathbb{R}^{q \times q}$, where $a_n \rightarrow \infty$ as $n \rightarrow \infty$, and it satisfies the following. For every $1 \leq m \leq a_n$, π is the stationary distribution of $P_{n,m}$, and there exists a constant $c > 0$ such that the minimal element of $P_{n,m}$ is bounded below by c , i.e. $P_{n,m}(i, j) \geq c$ holds for $i, j \in [q]$. Then, the product $\prod_{m=1}^{a_n} P_{n,m}$ converges to $\mathbf{1}\pi^\top$ as $n \rightarrow \infty$.*

Proof. For $n \geq 1$ and $1 \leq t \leq a_n$, consider the following distances:

$$\Delta_n(t) := \sup_{i \in [q]} \left\| \left(\prod_{m=1}^t P_{n,m} \right) (i, \cdot) - \pi \right\|_{\text{TV}}, \quad \bar{\Delta}_n(t) := \sup_{i, j \in [q]} \left\| \left(\prod_{m=1}^t P_{n,m} \right) (i, \cdot) - \left(\prod_{m=1}^t P_{n,m} \right) (j, \cdot) \right\|_{\text{TV}}.$$

Then, since π is the stationary distribution for $\prod_{m=1}^t P_{n,m}$, $\Delta_n(t) \leq \bar{\Delta}_n(t)$ holds (see e.g. proof of [LPW08, Lemma 4.10]). Moreover, a standard coupling argument (see e.g. [LPW08, Lemma 4.11]) shows that for any $n \geq 1$ and $1 \leq s \leq s + t \leq a_n$, we have

$$\begin{aligned}
\bar{\Delta}_n(t + s) &\leq \bar{\Delta}_n(t) \bar{\Delta}_n(s; t), \quad \text{where,} \\
\bar{\Delta}_n(s; t) &:= \sup_{i, j \in [q]} \left\| \left(\prod_{m=t+1}^{t+s} P_{n,m} \right) (i, \cdot) - \left(\prod_{m=t+1}^{t+s} P_{n,m} \right) (j, \cdot) \right\|_{\text{TV}}.
\end{aligned}$$

Finally, note that $\bar{\Delta}_n(1; t) \leq 1 - c$ holds for any $0 \leq t \leq a_n - 1$ since every element of $P_{n,t+1}$ is bounded below by $c > 0$. Thus, by the above submultiplicative property, we have that

$$\Delta_n(t) \leq \bar{\Delta}_n(t) \leq (1 - c)^t.$$

By letting $t = a_n$, the inequality above establishes that $\prod_{m=1}^{a_n} P_{n,m}$ converges to $\mathbf{1}\pi^\top$. \square

To use Lemma B.1, observe that under (SYM), the matrix $\Phi_{\zeta,s,t} \in \mathbb{R}^{q \times q}$ defined in (38) is a stochastic matrix with stationary distribution π . Moreover, since we assumed that Φ is finite whose elements are positive weight functions, the elements of $\Phi_{\psi,s,t}$ are lower bounded by a constant $c_\Psi > 0$. Recalling that $\Phi_\zeta := \prod_{i=1}^\ell \Phi_{\psi_i, s_i, t_i}$ for a signature $\zeta = (\psi_1, s_1, t_1, \dots, \psi_\ell, s_\ell, t_\ell)$, it follows from Lemma B.1 that

$$\limsup_{\ell \rightarrow \infty} \sup_{\zeta \in S_\ell} |\delta_\zeta| \equiv \limsup_{\ell \rightarrow \infty} \sup_{\zeta \in S_\ell} |\text{tr}(\Phi_\zeta) - 1| = 0.$$

Hence, the constants $(\delta_\zeta)_{\zeta \in \cup_{\ell \geq 1} S_\ell}$ are bounded from -1 and ∞ . Therefore, by using a Taylor approximation for the term $\log(1 + \delta_\zeta)$ in the RHS of (113), we can further bound

$$\mathbb{E}X_n^2 \leq C \left(\sum_{\ell=K'_n+1}^{\infty} \sum_{\zeta \in S_\ell} \left\{ \lambda_\zeta \delta_\zeta^2 + \lambda_\zeta^2 \delta_\zeta^4 \right\} + \left(\sum_{\ell=K'_n+1}^{\infty} \sum_{\zeta \in S_\ell} \lambda_\zeta \delta_\zeta^2 \right)^2 + n^{-1/3} \right),$$

where $C > 0$ only depends on k, q, Ψ . Since for $d < d_{KS}$, $\sum_{\ell \geq 1} \sum_{\zeta \in S_\ell} \lambda_\zeta \delta_\zeta^2 < \infty$ holds by Lemma 4.11, the RHS tends to 0 as $n \rightarrow \infty$. Therefore, $X_n \xrightarrow{p} 0$, which concludes the proof of (112).

B.3 Proof of Lemma 4.17

Let $\mathbf{G}_\gamma^*(n, m, \underline{\sigma}^*)$ be the factor graph obtained from γ resampling procedure starting from $\mathbf{G}^*(n, m, \underline{\sigma}^*)$. By definitions of the planted model and the resampling procedures, the factor graph $\mathbf{G}_\gamma^*(n, m, \underline{\sigma}^*)$ is distributed as follows. Conditional on $\underline{\sigma}^* = \underline{\sigma}$, independently draw for each clause $a \in F$ the neighborhood δa and the weight function ψ_a from the distribution

$$\mathbb{P}(\delta a = (v_1, \dots, v_k), \psi_a = \psi) = \frac{p(\psi)}{n^k} \left(\frac{(1 - \gamma)\psi(\sigma_{v_1}, \dots, \sigma_{v_k})}{\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)]} + \frac{\gamma}{\xi} \right).$$

On the other hand, let $(\mathbf{G}^*)'(n, m, \underline{\sigma}^*) \equiv (\mathbf{G}^*)'(n, m, \underline{\sigma}^*, p_\gamma, \pi)$ be the planted model with γ -modified weight functions defined in (49). Then, conditional on $\underline{\sigma}^* = \underline{\sigma}$, the neighborhood δa and the weight function ψ_a in $(\mathbf{G}^*)'(n, m, \underline{\sigma})$ are independently drawn from a slightly different distribution

$$\mathbb{P}(\delta a = (v_1, \dots, v_k), \psi_a = \psi) = \frac{p(\psi)}{n^k} \cdot \frac{(1 - \gamma)\psi(\sigma_{v_1}, \dots, \sigma_{v_k}) + \gamma\xi}{\xi}.$$

Hence, given $\underline{\sigma}^* = \underline{\sigma}$, the likelihood ratio of $\mathbf{G}_\gamma^* \equiv \mathbf{G}_\gamma^*(n, m, \underline{\sigma}^*)$ and $(\mathbf{G}^*)' \equiv (\mathbf{G}^*)'(n, m, \underline{\sigma}^*)$ evaluated at a factor graph $G = (V, F, E, (\psi_a)_{a \in F})$ is given by

$$\frac{\mathbb{P}(\mathbf{G}_\gamma^* = G \mid \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}((\mathbf{G}^*)' = G \mid \underline{\sigma}^* = \underline{\sigma})} = \prod_{a \in F} \left(1 + \frac{\gamma(\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] - \xi)}{\gamma\xi + (1 - \gamma)\psi_a(\delta a)} \right). \quad (114)$$

Observe that under (SYM), Lemma 6.6 shows that $\mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] - \xi = O_p(n^{-1})$. More precisely, for $\underline{\sigma} \in [q]^V$, define the vector $R_\sigma \equiv (R_\sigma(i))_{i \leq q} \in \mathbb{R}^q$ by the empirical distribution of $\underline{\sigma}$:

$$R_\sigma(i) := \frac{1}{n} \sum_{v \in V} \mathbf{1}\{\sigma_v = i\}. \quad (115)$$

Then, by Lemma 6.6, for any $\underline{\sigma} \in [q]^V$, we have the bound

$$\left| \mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] - \xi \right| \leq \frac{C}{n} \left\| \sqrt{n}(R_{\underline{\sigma}} - \pi) \right\|_1^2,$$

where $C \equiv C_{k,q,\Psi} > 0$. Thus, on the event $\left\{ \left\| \sqrt{n}(R_{\underline{\sigma}} - \pi) \right\|_1 \leq K \right\}$, which happens with probability $1 - o_K(1)$ under $\underline{\sigma}^* \sim \pi^{\otimes n}$ by the central limit theorem, we have that $\left| \mathbb{E}_{p,u}[\psi(\underline{\sigma}_\omega)] - \xi \right| = O_{k,q,\Psi}\left(\frac{K^2}{n}\right)$ holds. Therefore, using this bound in the RHS of (114) shows that on the event $\left\{ \left\| \sqrt{n}(R_{\underline{\sigma}} - \pi) \right\|_1 \leq K \right\}$, we have

$$\frac{\mathbb{P}(\mathbf{G}_\gamma^* = G \mid \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}((\mathbf{G}^*)' = G \mid \underline{\sigma}^* = \underline{\sigma})} = \exp\left(O_{k,q,\Psi}\left(\frac{K^2|F|}{n}\right)\right).$$

Since the number of clauses m is Poisson with mean dn/k for both \mathbf{G}_γ^* and $(\mathbf{G}^*)'$, we have $|F| \leq Cn$ holds with probability tending to 1 under both models. Therefore, we conclude that $(\mathbf{G}_\gamma^*, \underline{\sigma}^*)$ and $((\mathbf{G}^*)', \underline{\sigma}^*)$ are contiguous (see e.g. [Jan95, Proposition 9.47]).

C Applications to hypergraph stochastic block models

In this section, we prove Theorems 3.13, 3.14, and 3.15 by applying our results for the planted factor models. Throughout, we fix $k, q \geq 2$ and $\pi \in \mathcal{P}([q])$. Also, we recall that \mathcal{G}_n denotes the event consisting of factor graphs that satisfy (H1) and (H2). We denote by $(\tilde{\mathbf{G}}^*, \underline{\sigma}^*)$ the planted model $(\mathbf{G}^*, \underline{\sigma}^*)$ conditioned on the event \mathcal{G}_n . Then, $\tilde{\mathbf{G}}^*$ is a k -uniform hypergraph with no multiple edges by viewing each clause as an hyperedge.

C.1 Proof of Theorem 3.14 and Theorem 3.15

First, we prove that $\mathbf{G}_{\text{HSBM}}^* \sim \mathcal{G}^{\text{H}}(n, M, \pi)$ is mutually contiguous with respect to $\tilde{\mathbf{G}}^*$ with a particular choice of weight function defined as follows. By viewing the symmetric tensor M of order k as a weight function $M : [q]^k \rightarrow \mathbb{R}_+$, let p_M be the prior on the weight functions which puts all of its mass on M

$$p_M = \delta_M. \tag{116}$$

That is, there is a single weight function $\Psi \equiv \{M\}$. Then, we consider the planted model

$$\mathbf{G}_M^* \equiv \mathbf{G}^*(n, m, \underline{\sigma}^*, p_M, \pi), \quad \text{where } m \sim \text{Poi}(dn/k),$$

where d is the average degree of the HSBM defined in (22). We let $\tilde{\mathbf{G}}_M^*$ be the k -uniform hypergraph obtained from \mathbf{G}_M^* by conditioning on the event \mathcal{G}_n .

Lemma C.1. *Let M be a symmetric tensor of order k with positive entries such that the degree condition (24) holds for $M_0 = M/d$. Then, the total variation distance between $(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*)$ and $(\tilde{\mathbf{G}}_M^*, \underline{\sigma}^*)$ tends to 0 as $n \rightarrow \infty$:*

$$\lim_{n \rightarrow \infty} \text{TV}\left((\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*), (\tilde{\mathbf{G}}_M^*, \underline{\sigma}^*)\right) = 0.$$

Proof. Let $Q(G, \underline{\sigma}) := \frac{\mathbb{P}(\tilde{\mathbf{G}}_M^* = G, \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G, \underline{\sigma}^* = \underline{\sigma})}$ be the likelihood ratio between $(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*)$ and $(\tilde{\mathbf{G}}_M^*, \underline{\sigma}^*)$. Then, it is standard to see that

$$\text{TV} \left((\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*), (\tilde{\mathbf{G}}_M^*, \underline{\sigma}^*) \right) = \mathbb{E} \left[(1 - Q(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*)) \mathbb{1} \{ Q(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) \leq 1 \} \right].$$

Since the term inside the expectation in the right hand side is bounded, if we establish that

$$Q(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) \xrightarrow{p} 1, \quad (117)$$

then this implies our goal. Thus, we aim to show (117) for the rest of the proof.

Given a factor graph $G = (V, F, E, (\psi_a)_{a \in F})$ and a k -tuple of variables $\omega \equiv (v_1, \dots, v_k) \in V^k$, let $F_\omega(G)$ be the set of $a \in F$ such that $\delta a = (v_1, \dots, v_k)$ and denote its size by $f_\omega(G) \equiv |F_\omega(G)|$. For $\mathbf{G}_M^* \equiv \mathbf{G}^*(n, \mathbf{m}, \underline{\sigma}^*, p_M, \pi)$, we have by Poisson thinning that conditional on $\underline{\sigma}^*$, $(f_\omega(\mathbf{G}_M^*))_{\omega \in V^k}$ are independent Poisson random variables with mean

$$\mathbb{E} \left[f_\omega(\mathbf{G}_M^*) \mid \underline{\sigma}^* \right] = \frac{dn}{k} \frac{M(\underline{\sigma}_\omega^*)}{\sum_{\omega' \in V^k} M(\underline{\sigma}_{\omega'}^*)}.$$

Observe that $G \in \mathcal{G}_n$ if and only if $f_\omega(G) = 0$ for any $\omega = (v_1, \dots, v_k)$ such that v_1, \dots, v_k are not distinct and $f_\omega(G) \leq 1$ holds for any $\omega \in V^k$. Thus, conditional on $\underline{\sigma}^* = \underline{\sigma}$, each possible hyperedge (v_1, \dots, v_k) in $\tilde{\mathbf{G}}_M^*$, where v_1, \dots, v_k are distinct, is included independently with probability $\frac{\phi(\sigma_{v_1, \dots, v_k})}{1 + \phi(\sigma_{v_1, \dots, v_k})}$. Here, for $\underline{i} = (i_1, \dots, i_k) \in [q]^k$, the function $\phi(\underline{i}) \equiv \phi_\underline{\sigma}(\underline{i})$ is defined by

$$\phi(\underline{i}) := \frac{d(k-1)!M(\underline{i})}{\sum_{\omega' \in V^k} M(\underline{\sigma}_{\omega'}^*)} \cdot n = \frac{d(k-1)!M(\underline{i})}{\sum_{i'_1, \dots, i'_k} M(i'_1, \dots, i'_k) \prod_{s=1}^k R_{\underline{\sigma}}(i'_s)} n^{-(k-1)}, \quad (118)$$

where in the last equality, $R_{\underline{\sigma}} \equiv (R_{\underline{\sigma}}(i))_{1 \leq i \leq q}$ denotes the empirical distribution of $\underline{\sigma}$ (cf. (115)). Thus, if we let $E_{\underline{i}} \equiv E_{\underline{i}}(G)$ denote the number of hyperedges in G such that the communities of the end points of the hyperedges are given by $\underline{i} = (i_1, \dots, i_k)$, then we have that

$$\mathbb{P}(\tilde{\mathbf{G}}_M^* = G \mid \underline{\sigma}^* = \underline{\sigma}) = \prod_{\underline{i} = (i_1, \dots, i_k) \in [q]^k} \left(\frac{\phi(\underline{i})}{1 + \phi(\underline{i})} \right)^{E_{\underline{i}}} \cdot \left(\frac{1}{1 + \phi(\underline{i})} \right)^{n^k \prod_{s=1}^k R_{\underline{\sigma}}(i_s) - E_{\underline{i}}}.$$

Hence, $Q(G, \underline{\sigma})$ can be expressed by

$$\begin{aligned} Q(G, \underline{\sigma}) &= \frac{\mathbb{P}(\tilde{\mathbf{G}}_M^* = G \mid \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G \mid \underline{\sigma}^* = \underline{\sigma})} \\ &= \prod_{\underline{i} \in [q]^k} \left(\frac{\phi(\underline{i}) \cdot \binom{n}{k-1}}{(1 + \phi(\underline{i})) \cdot M(\underline{i})} \right)^{E_{\underline{i}}} \cdot \left((1 + \phi(\underline{i})) \left(1 - \frac{M(\underline{i})}{\binom{n}{k-1}} \right) \right)^{-n^k \prod_{s=1}^k R_{\underline{\sigma}}(i_s) + E_{\underline{i}}} \end{aligned} \quad (119)$$

To this end, we estimate the RHS when $G = \mathbf{G}_{\text{HSBM}}^*$ and $\underline{\sigma} = \underline{\sigma}^*$. Observe that $\phi(i_1, \dots, i_k)$ can be estimated as follows. If we let $Z_{\underline{\sigma}} := \sqrt{n}(R_{\underline{\sigma}} - \pi)$, then $Z_{\underline{\sigma}^*}$ is asymptotically normal by the central limit theorem. In particular, $Z_{\underline{\sigma}^*}$ has $O(1)$ fluctuations. Moreover, since $M_0 = M/d$ satisfies the condition (24), we have

$$\sum_{\underline{i} = (i_1, \dots, i_k) \in [q]^k} M(i_1, \dots, i_k) \prod_{s=1}^k \left(\pi_{i_s} + n^{-1/2} Z_{\underline{\sigma}}(i_s) \right) = d + O \left(\frac{\|Z_{\underline{\sigma}}\|_\infty^2 \cdot \left(\frac{Z_{\underline{\sigma}}}{\sqrt{n}} \vee 1 \right)^{k-2}}{n} \right).$$

Thus, by plugging in the estimate above to (118), we can approximate

$$\phi_{\underline{\sigma}}(\underline{i}) = \left(1 + \frac{h_{\underline{\sigma}}(\underline{i})}{n}\right) \cdot \frac{M(\underline{i})}{\binom{n}{k-1}}, \quad \text{where } h_{\underline{\sigma}^*}(i_1, \dots, i_k) \leq \log n \quad \text{holds w.h.p.} \quad (120)$$

Moreover, note that

$$E_{\underline{i}}(\mathbf{G}_{\text{HSBM}}^*) = \frac{n^k M(\underline{i}) \prod_{s=1}^k \pi_{i_s}}{\binom{n}{k-1}} + o_{\mathbb{P}}(n^{2/3}) = n \cdot (k-1)! M(\underline{i}) \prod_{s=1}^k \pi_{i_s} + o_{\mathbb{P}}(n^{2/3}), \quad (121)$$

where $Y = o_{\mathbb{P}}(n^{2/3})$ denotes a term that $n^{-2/3}Y \xrightarrow{p} 0$ holds under $(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*)$. Hence, by using the estimates (120) and (121), we have the approximation

$$\left(\frac{\phi_{\underline{\sigma}^*}(\underline{i}) \cdot \binom{n}{k-1}}{(1 + \phi_{\underline{\sigma}^*}(\underline{i})) \cdot M(\underline{i})}\right)^{E_{\underline{i}}(\mathbf{G}_{\text{HSBM}}^*)} = \exp\left((k-1)! \left(h_{\underline{\sigma}^*}(\underline{i}) - \mathbb{1}\{k=2\}M(\underline{i})\right) M(\underline{i}) \prod_{s=1}^k \pi_{i_s} + o_{\mathbb{P}}(1)\right).$$

Similarly, the second term of the product in (119) can be approximated by

$$\begin{aligned} & \left(\left(1 + \phi_{\underline{\sigma}^*}(\underline{i})\right) \left(1 - \frac{M(\underline{i})}{\binom{n}{k-1}}\right)\right)^{-n^k \prod_{s=1}^k R_{\underline{\sigma}^*}(i_s) + E_{\underline{i}}(\mathbf{G}_{\text{HSBM}}^*)} \\ &= \exp\left(- (k-1)! \left(h_{\underline{\sigma}^*}(\underline{i}) - \mathbb{1}\{k=2\}M(\underline{i})\right) M(\underline{i}) \prod_{s=1}^k R_{\underline{\sigma}^*}(i_s) + o_{\mathbb{P}}(1)\right) \\ &= \exp\left(- (k-1)! \left(h_{\underline{\sigma}^*}(\underline{i}) - \mathbb{1}\{k=2\}M(\underline{i})\right) M(\underline{i}) \prod_{s=1}^k \pi_{i_s} + o_{\mathbb{P}}(1)\right), \end{aligned}$$

where the last approximation holds since $R_{\underline{\sigma}^*} = \pi + o_{\mathbb{P}}(n^{-1/3})$. Therefore, the RHS in the two displays above exactly cancels out, and combining with (119) yields that $Q(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) \xrightarrow{p} 1$. \square

Let $\tilde{d}_*(p_{M_0}, \pi)$ denote the weak recovery threshold w.r.t. $\tilde{\mathbf{G}}_{d, M_0}^*$. That is, recalling the general definition of weak recovery in Definition 5.1, let

$$\tilde{d}_*(p_{M_0}, \pi) := \inf \left\{ d > 0 : \text{weak recovery is possible at } d \text{ for } \tilde{\mathbf{G}}_{d, M_0}^* \right\}.$$

As a consequence of Lemma C.1, we have

$$d_*^{\text{H}}(M_0, \pi) = \tilde{d}_*(p_{M_0}, \pi). \quad (122)$$

To transfer Theorem 4.12 for planted factor models to HSBM, we need to further prove that the weak recovery threshold is unchanged after conditioning on \mathcal{G}_n .

Lemma C.2. *Let M_0 be a symmetric tensor of order k with positive entries such that the degree condition (24) is satisfied. Then, we have $\tilde{d}_*(p_{M_0}, \pi) = d_*(p_{M_0}, \pi)$. and $d_{\text{KS}}(p_{M_0}, \pi) = d_{\text{KS}}^{\text{H}}(M_0, \pi)$.*

Proof. The final assertion that $d_{\text{KS}}(p_{M_0}, \pi) = d_{\text{KS}}^{\text{H}}(M_0, \pi)$ is immediate from the definition of KS thresholds for HSBM and planted factor models stated in (27) and Definition 4.3 respectively. For the rest of the proof, we aim to prove that $\tilde{d}_*(p_{M_0}, \pi) = d_*(p_{M_0}, \pi)$ holds by showing that

$$\limsup_n \mathbb{E} \left[\sup_{\hat{\sigma}(\tilde{\mathbf{G}}_M^*)} A(\underline{\sigma}^*, \hat{\sigma}) - \frac{1}{q} \right] = 0 \quad \Leftrightarrow \quad \limsup_n \mathbb{E} \left[\sup_{\hat{\sigma}(\mathbf{G}_M^*)} A(\underline{\sigma}^*, \hat{\sigma}) - \frac{1}{q} \right] = 0.$$

Note that $A(\underline{\sigma}^*, \hat{\sigma}) - \frac{1}{q} \in [0, 1]$ and so the above equivalence would follow if $\tilde{\mathbf{G}}_M^*$ and \mathbf{G}_M^* were mutually contiguous since the events $\{A(\underline{\sigma}^*, \hat{\sigma}) - \frac{1}{q} \geq \epsilon\}$ would both either tend to 0 or not. Unfortunately, while $\tilde{\mathbf{G}}_M^*$ is contiguous with respect to \mathbf{G}_M^* (which gives one direction of the equivalence) \mathbf{G}_M^* is not contiguous with respect to $\tilde{\mathbf{G}}_M^*$ because there is a constant probability of clauses that violate (H1) or (H2). Instead, we will add clauses to $\tilde{\mathbf{G}}_M^*$, using no information about $\underline{\sigma}^*$ so that the resulting graph $\check{\mathbf{G}}_M^*$ is mutually contiguous with respect to \mathbf{G}_M^* . Let

$$\mathcal{C} = \{\underline{j} \in \mathbb{Z}^k : 1 \leq j_1 \leq j_2 \leq \dots \leq j_k \leq n\},$$

be the set of increasing integer sequences of length k between 1 and n . For $\underline{j} \in \mathcal{C}$ let $N_{\underline{j}}$ be the number of clauses in \mathbf{G}_M^* with vertex set given by a permutation of $v_{j_1}, v_{j_2}, \dots, v_{j_k}$ (counting multiplicity if there are repeated j_i). By construction of \mathbf{G}_M^* , conditional on $\underline{\sigma}^*$, the $N_{\underline{j}}$ are independent Poisson random variables with means $p_{\underline{j}}$ where

$$C_1^{-1} n^{-(k-1)} \leq p_{\underline{j}} \leq C_1 n^{-(k-1)}$$

for some $C_1 > 1$ which may depend on the model but not on n or \underline{j} . We will let \mathcal{D} denote the set of \underline{j} with all j_i distinct. The $p_{\underline{j}}$ satisfy

$$\sum_{\underline{j} \in \mathcal{C}} p_{\underline{j}} \leq CN, \quad \sum_{\underline{j} \in \mathcal{C} \setminus \mathcal{D}} p_{\underline{j}} \leq C. \quad (123)$$

Letting $\tilde{N}_{\underline{j}}$ denote the number of clauses of $\tilde{\mathbf{G}}_M^*$ with vertices \underline{j} , the $\tilde{N}_{\underline{j}}$ are conditionally independent with distribution Poisson with mean $p_{\underline{j}}$ conditioned to be at most 1 if $\underline{j} \in \mathcal{D}$ and are equal to 0 if $\underline{j} \in \mathcal{C} \setminus \mathcal{D}$.

Next let $N'_{\underline{j}}$ be independent Poisson with mean $n^{-(k-1)}$ and define the transformed graph $\check{\mathbf{G}}_M^*$ by adding clauses such that

$$\check{N}_{\underline{j}} := \begin{cases} N'_{\underline{j}} & \text{if } \tilde{N}_{\underline{j}} \in \mathcal{D}, N'_{\underline{j}} \geq 2 \\ N'_{\underline{j}} & \text{if } \tilde{N}_{\underline{j}} \in \mathcal{C} \setminus \mathcal{D} \\ \tilde{N}_{\underline{j}} & \text{otherwise.} \end{cases}$$

Note that we do not use any information about $\underline{\sigma}^*$ to construct $\check{\mathbf{G}}_M^*$ from $\tilde{\mathbf{G}}_M^*$. For $\underline{j} \in \mathcal{D}$, the distribution of $\check{N}_{\underline{j}}$ satisfies

$$\frac{\mathbb{P}[\check{N}_{\underline{j}} = \ell]}{\mathbb{P}[N_{\underline{j}} = \ell]} = \begin{cases} \frac{\mathbb{P}[N'_{\underline{j}} \leq 2]}{\mathbb{P}[\tilde{N}_{\underline{j}} \leq 2]} & \text{if } \ell \leq 1 \\ \frac{n^{-(k-1)\ell} e^{-n^{-(k-1)}}}{p_{\underline{j}}^\ell e^{-p_{\underline{j}}}} & \text{if } \ell \geq 1 \end{cases} = \begin{cases} 1 + O(n^{-2(k-1)}) & \text{if } \ell \leq 1 \\ e^{O(\ell)} & \text{if } \ell \geq 2 \end{cases}$$

Hence, if we write $\nu_{\underline{j}}$ and $\check{\nu}_{\underline{j}}$ for the law of $N_{\underline{j}}$ and $\check{N}_{\underline{j}}$ respectively then

$$\|\nu_{\underline{j}} - \check{\nu}_{\underline{j}}\|_{L^2(\nu_{\underline{j}})}^2 = \sum_{\ell=0}^{\infty} \left| \frac{\mathbb{P}[\check{N}_{\underline{j}} = \ell]}{\mathbb{P}[N_{\underline{j}} = \ell]} - 1 \right|^2 \mathbb{P}[N_{\underline{j}} = \ell] \leq O(n^{-4(k-1)}) + \sum_{\ell=2}^{\infty} C^{2\ell} \frac{p_{\underline{j}}^{\ell} e^{-p_{\underline{j}}}}{\ell!} = O(n^{-(k-1)} p_{\underline{j}}),$$

and similarly

$$\|\nu_{\underline{j}} - \check{\nu}_{\underline{j}}\|_{L^2(\check{\nu}_{\underline{j}})}^2 = O(n^{-(k-1)} p_{\underline{j}}).$$

For $\underline{j} \in \mathcal{C} \setminus \mathcal{D}$,

$$\begin{aligned} \|\nu_{\underline{j}} - \check{\nu}_{\underline{j}}\|_{L^2(\nu_{\underline{j}})}^2 &= \sum_{\ell=0}^{\infty} \left| \frac{n^{-(k-1)\ell} e^{-n^{-(k-1)}}}{p_{\underline{j}}^{\ell} e^{-p_{\underline{j}}^{\ell}}} - 1 \right|^2 \frac{p_{\underline{j}}^{\ell} e^{-p_{\underline{j}}}}{\ell!} \\ &\leq O(n^{-2(k-1)}) + \sum_{\ell=1}^{\infty} C^{2\ell} \frac{p_{\underline{j}}^{\ell} e^{-p_{\underline{j}}}}{\ell!} \\ &= O(n^{-(k-1)}) = O(p_{\underline{j}}), \end{aligned}$$

and similarly

$$\|\nu_{\underline{j}} - \check{\nu}_{\underline{j}}\|_{L^2(\check{\nu}_{\underline{j}})}^2 = O(p_{\underline{j}}).$$

Setting ν and $\check{\nu}$ for the law of the vectors $\{N_{\underline{j}}\}_{\underline{j} \in \mathcal{C}}$ and $\{\check{N}_{\underline{j}}\}_{\underline{j} \in \mathcal{C}}$ respectively, since both are product measures,

$$\begin{aligned} 1 + \|\nu - \check{\nu}\|_{L^2(\check{\nu})}^2 &= \prod_{\underline{j} \in \mathcal{C}} (1 + \|\nu_{\underline{j}} - \check{\nu}_{\underline{j}}\|_{L^2(\check{\nu}_{\underline{j}})}^2) \\ &\leq \exp \left(C n^{-(k-1)} \sum_{\underline{j} \in \mathcal{D}} p_{\underline{j}} + C \sum_{\underline{j} \in \mathcal{C} \setminus \mathcal{D}} p_{\underline{j}} \right) \\ &\leq O(1), \end{aligned}$$

where the last inequality used equation (123). Similarly

$$\|\tilde{\nu} - \check{\nu}\|_{L^2(\check{\nu})}^2 = O(1).$$

It follows that $\{N_{\underline{j}}\}_{\underline{j} \in \mathcal{C}}$ and $\{\check{N}_{\underline{j}}\}_{\underline{j} \in \mathcal{C}}$ are mutually contiguous and hence $\check{\mathbf{G}}_M^*$ is mutually contiguous with respect to \mathbf{G}_M^* . Hence we have that $\tilde{d}_*(p_{M_0}, \pi) = d_*(p_{M_0}, \pi)$. \square

Let $\tilde{\mathbf{G}}$ denote the null model $\mathbf{G} \sim \mathcal{G}_{\text{null}}(n, d, p_M)$ conditioned on the event \mathcal{G}_n . Here, we drop the subscript M since the factors $(\psi_a)_{a \in F} \equiv M$ in $\tilde{\mathbf{G}}$ (and \mathbf{G}) do not play any role. For a factor graph G , let

$$\tilde{L}(G) := \frac{\mathbb{P}(\tilde{\mathbf{G}}_M^* = G)}{\mathbb{P}(\tilde{\mathbf{G}} = G)} = \frac{L(G) \mathbf{1}\{G \in \mathcal{G}_n\}}{\mathbb{E}[L(\mathbf{G}) | \mathbf{G} \in \mathcal{G}_n]}, \quad (124)$$

where the last equality holds by definition of $L(G)$ (see Eq. (42)). Then, the following proposition plays an important role in the proof of Theorem 3.14. Recall the constants $(\alpha_{\ell})_{\ell \geq 2}$ defined in Eq. (29), which are functions of M_0 and π .

Proposition C.3. Let M_0 be a symmetric tensor of order k with positive entries. Below the weak revery and KS thresholds $d < d_*(p_{M_0}, \pi) \wedge d_{\text{KS}}(p_{M_0}, \pi)$, we have that as $n \rightarrow \infty$,

$$\tilde{L}(\tilde{\mathbf{G}}) \xrightarrow{d} \mathcal{L}_\infty := \prod_{\ell=2+\mathbf{1}\{k=2\}}^{\infty} \frac{(1 + \alpha_\ell)^{\mathbf{X}_\ell}}{\mathbb{E}(1 + \alpha_\ell)^{\mathbf{X}_\ell}}, \quad (125)$$

where $(\mathbf{X}_\ell)_{\ell \geq 2}$ are independent Poisson random variables with mean $\mathbb{E}\mathbf{X}_\ell = \frac{1}{2\ell}((k-1)d)^\ell$.

Proof. The proof follows from a combination of Fact 4.9-(3) Theorem 4.12. For a signature $\zeta \in S_\ell$, if the prior on the weight functions p puts all of its mass on M , then the constants λ_ζ and λ_ζ^* in (39) equal

$$\lambda_\zeta = \frac{1}{2\ell} \left(\frac{d}{k} \right)^\ell, \quad \lambda_\zeta^* = \frac{1}{2\ell} \text{tr}(B^\ell),$$

where B is the matrix defined in (26). Thus, $\lambda_\zeta, \lambda_\zeta^*$ depends on ζ on through its order ℓ and

$$\delta_\zeta \equiv \lambda_\zeta^*/\lambda_\zeta - 1 = \alpha_\ell.$$

Consequently, Theorem 4.12 yields that for $d < d_*(p_{M_0}, \pi) \wedge d_{\text{KS}}(p_{M_0}, \pi)$,

$$L(\mathbf{G}) \xrightarrow{d} \mathbf{L}_\infty = \prod_{\ell \geq 1} \frac{(1 + \alpha_\ell)^{\mathbf{X}_\ell}}{\mathbb{E}(1 + \alpha_\ell)^{\mathbf{X}_\ell}},$$

where this convergence holds jointly with the weak convergence of the number of cycles of given length in Fact 4.9-(1). Meanwhile by Fact 4.9-(3), $\mathbb{P}(\mathbf{G} \in \mathcal{G}_n) = \mathbb{P}(C(\mathbf{G}) = 0) + O(1/n)$ and $\mathbb{P}(\mathbf{G}^* \in \mathcal{G}_n) = \mathbb{P}(C(\mathbf{G}^*) = 0) + O(1/n)$ hold, where $C(G)$ is the number of self-loops if $k \geq 3$ and the sum of the number of self-loops and the number of cycles of length 2 if $k = 2$. Hence, it follows that $L(\tilde{\mathbf{G}})$ weakly converges to the distribution of \mathbf{L}_∞ conditional on the event $\{\mathbf{X}_1 + \mathbf{1}\{k=2\}\mathbf{X}_2 = 0\}$. Moreover, $\tilde{L}(G)$ and $L(G)$ just differs by a multiplicative factor for $G \in \mathcal{G}_n$ by (124) where $\tilde{L}(\tilde{\mathbf{G}})$ satisfies the normalization $\mathbb{E}\tilde{L}(\tilde{\mathbf{G}}) = 1$. Therefore, the desired claim (125) follows. \square

Proof of Theorem 3.14. Consider $d < d_*^{\text{H}}(M_0, \pi) \wedge d_{\text{KS}}^{\text{H}}(M_0, \pi)$. Recalling (122), we have $d_*^{\text{H}}(M_0, \pi) = \tilde{d}_*(p_{M_0}, \pi)$. Moreover, $\tilde{d}_*(p_{M_0}, \pi) = d_*(p_{M_0}, \pi)$ and $d_{\text{KS}}^{\text{H}}(M_0, \pi) = d_{\text{KS}}^{\text{H}}(p_{M_0}, \pi)$ hold by Lemma C.2. Hence, Proposition C.3 yields $\tilde{L}(\tilde{\mathbf{G}}) \xrightarrow{d} \mathcal{L}_\infty$. Note that $\tilde{L}(G)$ and $\mathcal{L}_n(G)$ are related by the chain rule

$$\mathcal{L}_n(G) = \tilde{L}(G) \cdot \frac{\mathbb{P}(\tilde{\mathbf{G}} = G)}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)} \cdot \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G)}{\mathbb{P}(\tilde{\mathbf{G}}_M^* = G)} \equiv \tilde{L}(G) \frac{f_1(G)}{f_2(G)}, \quad (126)$$

where

$$f_1(G) := \frac{\mathbb{P}(\tilde{\mathbf{G}} = G)}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)}, \quad f_2(G) := \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G)}{\mathbb{P}(\tilde{\mathbf{G}}_M^* = G)}.$$

Note that by considering $M_0 = \mathbf{1}_{k,q}$, the all-1-tensor in Lemma C.1, we have $f_1(\tilde{\mathbf{G}}) \xrightarrow{p} 1$. Clearly, $f_2(\tilde{\mathbf{G}}_M^*) \xrightarrow{p} 1$ follows from Lemma C.1. Note that $\tilde{\mathbf{G}}_M^*$ is mutually contiguous with $\tilde{\mathbf{G}}$ by Proposition C.3, thus $f_2(\tilde{\mathbf{G}}) \xrightarrow{p} 1$ also holds. Therefore, by combining the established convergences $\tilde{L}(\tilde{\mathbf{G}}) \xrightarrow{d} \mathcal{L}_\infty$, $f_i(\tilde{\mathbf{G}}) \xrightarrow{p} 1$ for $i = 1, 2$, we have $\mathcal{L}_n(\tilde{\mathbf{G}}) \xrightarrow{d} \mathcal{L}_\infty$ by (126). Finally, the stated properties of \mathcal{L}_∞ are a special case of Lemma 4.13 and Corollary 4.14. \square

Proof of Theorem 3.15. This is a special case of Lemma 4.15 and Corollary 4.16. \square

C.2 Proof of Theorem 3.13

The following lemma relates the mutual information and the free energy for HSBMs, which is the analog of Lemma 4.18 for planted factor models.

Lemma C.4. *Let M_0 be a symmetric tensor of order k with positive entries such that the normalization (23) is satisfied. Then, the normalized mutual information can be approximated by*

$$\frac{1}{n}I(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) + \frac{1}{n}\mathbb{E}\log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) = \frac{d}{k} \sum_{i_1, \dots, i_k=1}^q M_0(i_1, \dots, i_k) \log(M_0(i_1, \dots, i_k)) \prod_{s=1}^k \pi_{i_s} + o_n(1).$$

Proof. By definition of mutual information and the likelihood ratio $\mathcal{L}_n(G)$, we have the identity

$$\begin{aligned} & \frac{1}{n}I(\mathbf{G}_{\text{HSBM}}^*, \underline{\sigma}^*) \\ &= -\frac{1}{n}\mathbb{E}\log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) + \frac{1}{n} \sum_{G, \underline{\sigma}} \mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G, \underline{\sigma}^* = \underline{\sigma}) \log \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G | \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)}. \end{aligned} \quad (127)$$

Note that for any G and $\underline{\sigma}$, we can compute

$$\frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G | \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)} = \prod_{(v_1, \dots, v_k) \in E(G)} \frac{\frac{M(\sigma_{v_1}, \dots, \sigma_{v_k})}{\binom{n}{k-1}}}{\frac{d}{\binom{n}{k-1}}} \prod_{(v_1, \dots, v_k) \notin E(G)} \frac{1 - \frac{M(\sigma_{v_1}, \dots, \sigma_{v_k})}{\binom{n}{k-1}}}{1 - \frac{d}{\binom{n}{k-1}}},$$

where $E(G)$ denote the set of hyperedges in G . Recall that $M = dM_0$, so taking logarithm and conditional expectation w.r.t. $\mathbf{G}_{\text{HSBM}}^*$ conditional on $\underline{\sigma}^* = \underline{\sigma}$ in the RHS gives

$$\begin{aligned} & \sum_G \mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G | \underline{\sigma}^* = \underline{\sigma}) \log \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G | \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)} \\ &= \sum_{(v_1, \dots, v_k)} \frac{dM_0(\sigma_{v_1}, \dots, \sigma_{v_k})}{\binom{n}{k-1}} \log M_0(\sigma_{v_1}, \dots, \sigma_{v_k}) + \left(1 - \frac{dM_0(\sigma_{v_1}, \dots, \sigma_{v_k})}{\binom{n}{k-1}}\right) \log \left(\frac{1 - \frac{dM_0(\sigma_{v_1}, \dots, \sigma_{v_k})}{\binom{n}{k-1}}}{1 - \frac{d}{\binom{n}{k-1}}}\right), \end{aligned}$$

where the sum is over distinct k vertices (v_1, \dots, v_k) . Taking expectation over $\underline{\sigma}^*$ yields

$$\begin{aligned} & \sum_{G, \underline{\sigma}} \mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G, \underline{\sigma}^* = \underline{\sigma}) \log \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* = G | \underline{\sigma}^* = \underline{\sigma})}{\mathbb{P}(\mathbf{G}_{\text{ER}} = G)} \\ &= \sum_{\underline{i}=(i_1, \dots, i_k) \in [q]^k} \binom{n}{k} \cdot \left(\frac{dM_0(\underline{i})}{\binom{n}{k-1}} \log M_0(\underline{i}) + \left(1 - \frac{dM_0(\underline{i})}{\binom{n}{k-1}}\right) \log \left(\frac{1 - \frac{dM_0(\underline{i})}{\binom{n}{k-1}}}{1 - \frac{d}{\binom{n}{k-1}}}\right)\right) \prod_{s=1}^k \pi_{i_s} \\ &= \frac{dn}{k} \left(\sum_{\underline{i} \in [q]^k} M_0(\underline{i}) \log M_0(\underline{i}) \prod_{s=1}^k \pi_{i_s} + \sum_{\underline{i} \in [q]^k} (1 - M_0(\underline{i})) \prod_{s=1}^k \pi_{i_s} + o_n(1) \right) \\ &= \frac{dn}{k} \sum_{\underline{i} \in [q]^k} M_0(\underline{i}) \log M_0(\underline{i}) \prod_{s=1}^k \pi_{i_s} + o_n(1), \end{aligned}$$

where we used $\sum_{\underline{i} \in [q]^k} (1 - M_0(\underline{i})) \prod_{s=1}^k \pi_{i_s} = 0$ (cf. (23)) in the last equality. Combining this with (127) concludes the proof. \square

We next prove that the free energy is bounded away from 0 along a subsequence by transferring Proposition 4.21 for the planted factor models to HSBM.

Proposition C.5. *For any $n \geq 1$ and $d > 0$, we have $\mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) \geq 0$. Moreover, if $d > d_*(p_{M_0}, \pi)$ and the condition (MIN) holds, then there exists $\eta > 0$ such that*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) \geq \eta. \quad (128)$$

Proof. Letting $F_0(x) = x \log x$, we have by a change of measure that

$$\mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) = \mathbb{E} \mathcal{L}_n(\mathbf{G}_{\text{ER}}) \log \mathcal{L}_n(\mathbf{G}_{\text{ER}}) \geq F_0(\mathbb{E} \mathcal{L}_n(\mathbf{G}_{\text{ER}})) = 0,$$

where the inequality follows from Jensen's inequality and the last equality holds by $\mathbb{E} \mathcal{L}_n(\mathbf{G}_{\text{ER}}) = 1$.

Next, suppose that $d > d_*(p_{M_0}, \pi)$ and the condition (MIN) holds. Assume by contraction that (128) does not hold. Then, since $\mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) \geq 0$, we must have that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) = 0.$$

Now, recall the chain rule $\mathcal{L}_n(G) = \tilde{L}(G) f_1(G) / f_2(G)$ in (126), where we defined $f_1(G) \equiv \frac{\mathbb{P}(\tilde{\mathbf{G}}=G)}{\mathbb{P}(\mathbf{G}_{\text{ER}}=G)}$ and $f_2(G) \equiv \frac{\mathbb{P}(\mathbf{G}_{\text{HSBM}}^*=G)}{\mathbb{P}(\tilde{\mathbf{G}}_M^*=G)}$. Then, note that for $i = 1, 2$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log f_i(\mathbf{G}_{\text{HSBM}}^*) = 0.$$

Indeed, $i = 2$ case holds since $\mathbf{G}_{\text{HSBM}}^*$ and $\tilde{\mathbf{G}}_M^*$ are mutually contiguous by Lemma C.1 and $n^{-1} \log f_2(\cdot)$ is bounded. For $i = 1$, $n^{-1} \log f_1(G)$ can be calculated directly since $\tilde{\mathbf{G}}$ and \mathbf{G}_{ER} doesn't have planted structure, from which it can be seen that for any sparse random graph $\tilde{\mathbf{G}}$ with at most linear number of edges in expectation, $n^{-1} \mathbb{E} \log f_1(\tilde{\mathbf{G}}) = o_n(1)$. Therefore, it follows that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log \tilde{L}(\mathbf{G}_{\text{HSBM}}^*) = 0.$$

Since $\text{TV}(\mathbf{G}_{\text{HSBM}}^*, \tilde{\mathbf{G}}_M^*) = o_n(1)$ holds by Lemma C.1 and $n^{-1} \log \tilde{L}(G) = O(1)$ holds for factor graph G with at most $O(n)$ clauses, it follows that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log \tilde{L}(\tilde{\mathbf{G}}_M^*) = 0.$$

Note that by Azuma Hoeffding's inequality, $n^{-1} \log \tilde{L}(\tilde{\mathbf{G}}_M^*)$ and $n^{-1} \log L(\mathbf{G}_M^*)$ concentrate tightly around their expectation with $O(n^{-1/2})$ fluctuation, and that $\mathbf{G}_M^* \in \mathcal{G}_n$ holds with uniformly positive probability. Thus, $n^{-1} \mathbb{E} \log \tilde{L}(\tilde{\mathbf{G}}_M^*) = n^{-1} \mathbb{E} \log L(\mathbf{G}_M^*) + O(n^{-1/2})$ holds. Consequently, we have that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log L(\mathbf{G}_M^*) = 0,$$

which contradicts Proposition 4.21-(3). □

Proof of Theorem 3.13. The first statement regarding the mutual contiguity follows immediately from Theorem 3.14 and Le Cam's first lemma (see e.g. [Vaa98, Lemma 6.4] or [JLuR00, Proposition 9.49]) since $\mathcal{L}_\infty > 0$ a.s. and $\mathbb{E}\mathcal{L}_\infty = 1$. The mutual contiguity between $\mathbf{G}_{\text{HSBM}}^*$ and \mathbf{G}_{ER} implies that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) = 0.$$

Combining with Lemma C.4 finishes the proof of the second statement. For the third statement, recall that $d_*^{\text{H}}(M_0, \pi) = \tilde{d}_*(p_{M_0}, \pi)$ holds (cf. (122)) and that $\tilde{d}_*(p_{M_0}, \pi) = d_*(p_{M_0}, \pi)$ holds by Lemma C.2. Thus, for $d > d_*^{\text{H}}(M_0, \pi)$, Proposition C.5 yields that there exists a constant $\eta > 0$ such that $n^{-1} \mathbb{E} \log \mathcal{L}_n(\mathbf{G}_{\text{HSBM}}^*) \geq \eta > 0$ holds along a subsequence $(n_\ell)_{\ell \geq 1}$. Then, consider the event

$$\mathcal{A}_n^* = \{G : \mathcal{L}_n(G) \geq e^{\eta n/2}\}.$$

Then, the same argument as in the proof of Theorem 3.8 shows that $\mathbb{P}(\mathbf{G}_{\text{ER}} \in \mathcal{A}_{n_\ell}^*) \leq e^{-\eta' n}$ and $\mathbb{P}(\mathbf{G}_{\text{HSBM}}^* \in \mathcal{A}_{n_\ell}^*) \geq 1 - e^{-\eta' n}$ for some constant $\eta' > 0$. Finally, the fourth statement is immediate from Lemma C.4 and Proposition C.5, which concludes the proof. \square