

The local well-posedness, blow-up and global solution of a new integrable system in Besov spaces

Pei Zheng¹*, Zhaoyang Yin^{1,2†}

¹Department of Mathematics, Sun Yat-sen University, Guangzhou 510275, China

²School of Science,

Shenzhen Campus of Sun Yat-sen University, Shenzhen 518107, China

Abstract

In this paper, we first establish the local well-posedness for the Cauchy problem of a N -peakon system in the sense of Hadamard in both critical Besov spaces and supercritical Besov spaces. Second, we gain a blow-up criterion. According to blow-up criterion we reach a precise blow-up criterion. Under a sign condition, we reach the existence of global solution. Finally, based on the first-order difference method, we give a simulation example of the blow-up of the equation and the properties of the global solution.

Contents

1	Introduction	2
2	Preliminaries	3
3	Local well-posedness	7
4	Blow-up criterion	14
5	Blow-up	16
6	Existence of global solution	17
7	Simulation	19
	References	20

*Email: zhengp25@mail2.sysu.edu.cn

†E-mail: mcsyzy@mail.sysu.edu.cn

1 Introduction

Korteweg-de Vries (KdV) equation which was introduced to describe the behavior of long waves on shallow water in the most famous model in soliton theory, because it is integrable and includes the phenomena of soliton interaction. But the KdV equation cannot model the occurrence of breaking waves. In 1993, Camassa and Holm obtain a nonlinear partial differential equation [2]

$$m_t + um_x + 2u_x m = 0, m = u - u_{xx} + \kappa$$

which is named Camassa-Holm (CH) equation.

A particular feature of the CH equation is that when $\kappa = 0$ it admits peaked soliton solutions which are also called peakons. It can be regarded as a shallow water wave equation.[2, 8, 18]. Its complete integrability was discussed in [2, 4, 9]. The CH equation also has a Hamilton structure[3, 12], and admits exact orbitally stable peaked solitons of the form $ce^{|x-ct|}$ with $c > 0$. The local well-posedness of the Cauchy problem of CH equation in Besov spaces and Sobolev spaces was proved in [5, 6, 10, 22].

Since constructing peakon equation are interested in the fields of Physics and Mathematics. Based on an asymptotic integrability approach, Degasperis and Procesi discovered a new equation which has peakon solutions[20, 11]. Similarly, the Cauchy problem of the DP equation is locally well-posed in certain Sobolev and Besov spaces[24, 16, 25]. And Geng and Xue constructed a two component peakon with cubic nonlinearity[13], three components generalization of CH equation[14] and super CH equation with N -peakon[15].

Geng and Xue constructed from the compatibility of a 2×2 matrix spectral problem, a new nonlinear evolution with N -peakon is derived[23]

$$(1.1) \quad \begin{cases} n_t = 4[n(v_x + 2\beta_0 v)]_x \\ n = 4\beta_0^2 v - v_{xx} \\ v|_{t=0} = v_0 \end{cases}$$

It is shown in that system (1.1) admits exact solutions with N -peakon, which takes the form

$$v(x, t) = \sum_{i=1}^N n_i(t) e^{-2\beta_0 |x - x_i(t)|}$$

where n_i and x_i evolve according to a dynamical system.

After some transformations, (1.1) can be changed into a transport-like equation

$$(1.2) \quad \begin{cases} v_t - (8\beta_0 v + 2v_x)v_x = -8\beta_0^2 v^2 + 8\beta_0 P_1(D)(2\beta_0^2 v^2 + v_x^2) + 8\beta_0^2 P_2(D)(4\beta_0^2 v^2 - v_x^2) \\ v|_{t=0} = v_0 \end{cases}$$

with $P_2(D) = (4\beta_0^2 - \partial_x^2)^{-1}$, $P_1(D) = \partial_x P_2(D)$.

In this paper, firstly, applying Littlewood-Paley theory and transport theory, for the initial data in certain Besov spaces of high regularity or of critical regularity, complete the locally well-posedness in Besov space $B_{p,r}^s$ with $\{s \geq \frac{1}{2}, p > 2, 1 \leq r \leq \infty\}$ or $\{s > \frac{1}{p}, 1 \leq p \leq 2, 1 \leq r \leq \infty\}$ or $\{s = \frac{1}{p}, 1 \leq p \leq 2, r = 1\}$, we prove the local solution to equation (1.1) exists uniquely and depends continuously on the initial data. And we will talk about the blow-up criterion and the blow up condition of initial data. In the last part of the paper, according to the structure of equation, we can find a sign-preserved property, and by the virtue of the high-preserved we see that the H^1 -norm of v is non-increasing, then we can finally give the condition of existence of the global solution.

2 Preliminaries

In this section, we will present some propositions about the Littlewood-Paley decomposition and the non homogeneous Besov spaces with their properties.

Proposition 2.1 (Littlewood-Paley decomposition). *[1, 17] There exists a couple of smooth function (χ, φ) valued in $[0, 1]$, such that χ is supported in the ball $B \triangleq \{\xi \in \mathbb{R}^d : \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$. Moreover*

$$\forall \xi \in \mathbb{R}^d, \chi(\xi) + \sum_{j \geq 0} \varphi(2^{-j}\xi) = 1$$

and

$$\begin{aligned} \text{supp } \varphi(2^{-j}\cdot) \cap \text{supp } \varphi(2^{-j'}\cdot) &= \emptyset, \text{ if } |j - j'| \geq 2 \\ \text{supp } \chi(\cdot) \cap \text{supp } \varphi(2^{-j'}\cdot) &= \emptyset, \text{ if } j \geq 1 \end{aligned}$$

Then for all $u \in \mathcal{S}'$, we can define the nonhomogeneous dyadic blocks as follows. Let

$$\begin{aligned} \Delta_j u &\triangleq 0, \text{ if } j \leq -2, \quad \Delta_{-1} u \triangleq \chi(D)u = \mathcal{F}^{-1}(\chi \mathcal{F}u) \\ \Delta_j &\triangleq \varphi(2^{-j}u) = \mathcal{F}^{-1}(\varphi(2^{-j}\cdot)\mathcal{F}u), \text{ if } j \geq 0 \end{aligned}$$

Hence,

$$u = \sum_{j \in \mathbb{Z}} \Delta_j u \text{ in } \mathcal{S}'(\mathbb{R}^d)$$

where the right-hand side is called the nonhomogeneous Littlewood-Paley decomposition of u .

According to Young's inequality, we get

$$\|\Delta_j u\|_{L^p}, \|S_j u\|_{L^p} \leq C \|u\|_{L^p}, \quad \forall 1 \leq p \leq \infty$$

The low frequency cut-off operator S_j is defined by

$$S_j u \triangleq \sum_{j'=-1}^{j-1} \Delta_{j'} u = \chi(2^{-j}D)u = \mathcal{F}^{-1}(\chi(2^{-j}\xi)\mathcal{F}u), \quad \forall j \in \mathbb{N}$$

and the Littlewood-Paley decomposition is quasi-orthogonal in L^2 in the following sense:

$$\begin{aligned} \Delta_j \Delta_k u &\equiv 0, \text{ if } |j - k| \geq 2 \\ \Delta_j (S_{k-1} u \Delta_k u) &\equiv 0, \text{ if } |j - k| \geq 5 \end{aligned}$$

Definition 2.2 (Nonhomogeneous Besov space). *[1] Let $s \in \mathbb{R}$, $1 \leq p, r \leq \infty$. The nonhomogeneous Besov space $B_{p,r}^s(\mathbb{R}^d)$ ($B_{p,r}^s$ for short) is defined by*

$$B_{p,r}^s(\mathbb{R}^d) \triangleq \left\{ f \in \mathcal{S}' : \|f\|_{B_{p,r}^s} = \triangleq \|(2^{js} \|\Delta_j f\|_{L^p})_{j \geq -1}\|_{l^r} < \infty \right\}$$

If $s = \infty$, we have

$$B_{p,r}^\infty \triangleq \cap_{s \in \mathbb{R}} B_{p,r}^s = \left\{ f \in \mathcal{S}' : \sup_{j \geq -1} 2^{js} \|\Delta_j f\|_{L^p} < \infty \right\}$$

In the following lemma, we list some important properties of Besov spaces.

Lemma 2.3. [1, 17] Let $s \in \mathbb{R}$, $1 \leq p, p_1, p_2, r, r_1, r_2 \leq \infty$. We have

(1) $B_{p,r}^s$ is a Banach space, and is continuously embedded in \mathcal{S}' .

(2) If $r < \infty$, then $\lim_{j \rightarrow \infty} \|S_j u - u\|_{B_{p,r}^s} = 0$. If $p, r < \infty$, C_0^∞ is dense in $B_{p,r}^s$.

(3) If $p_1 \leq p_2$ and $r_1 \leq r_2$, then $B_{p_1, r_1}^s \hookrightarrow B_{p_2, r_2}^{s-d(\frac{1}{p_1} - \frac{1}{p_2})}$. If $s_1 < s_2$, the embedding $B_{p, r_2}^{s_2} \hookrightarrow B_{p, r_1}^{s_1}$ is locally compact.

(4) $\forall s > 0$, $B_{p,r}^s \cap L^\infty$ is an algebra. Moreover, $B_{p,r}^s \hookrightarrow L^\infty \Leftrightarrow s > \frac{d}{p}$ or $s = \frac{d}{p}$, $r = 1 \Leftrightarrow B_{p,r}^s$ is an algebra.

(5) Complex interpolation: $\forall f \in B_{p,r}^{s_1} \cap B_{p,r}^{s_2}$, $\forall \theta \in [0, 1]$

$$(2.1) \quad \|f\|_{B_{p,r}^{\theta s_1 + (1-\theta)s_2}} \leq \|f\|_{B_{p,r}^{s_1}}^\theta \|f\|_{B_{p,r}^{s_2}}^{1-\theta}$$

(6) Logarithm interpolation: $\forall s \in \mathbb{R}$, $\varepsilon > 0$, and $1 \leq p \leq \infty$, there exists a constant C such that

$$(2.2) \quad \|u\|_{B_{p,1}^s} \leq C \frac{1+\varepsilon}{\varepsilon} \|u\|_{B_{p,\infty}^s} \left(1 + \log \frac{\|u\|_{B_{p,\infty}^{s+\varepsilon}}}{\|u\|_{B_{p,\infty}^s}} \right)$$

(7) Fatou property: if $(u_n)_{n \in \mathbb{N}}$ is a bounded sequence in $B_{p,r}^s$, then an element $u \in B_{p,r}^s$ and a subsequence $(u_{n_k})_{k \in \mathbb{N}}$ exists such that

$$\lim_{k \rightarrow \infty} u_{n_k} = u \text{ in } \mathcal{S}' \quad \text{and} \quad \|u\|_{B_{p,r}^s} \leq C \liminf_{k \rightarrow \infty} \|u_{n_k}\|_{B_{p,r}^s}.$$

(8) Let $m \in \mathbb{R}$ and f be a S^m -multiplier, (i.e. f is a smooth function and satisfies that $\forall \alpha \in \mathbb{N}^d$, $\exists C = C(\alpha)$, such that $|\partial^\alpha f(\xi)| \leq C(1 + |\xi|)^{m-|\alpha|}$, $\forall \xi \in \mathbb{R}^d$). Then the operator $f(D) = \mathcal{F}^{-1}(f\mathcal{F}\cdot)$ is continuous from $B_{p,r}^s$ to $B_{p,r}^{s-m}$.

Proposition 2.4. $\forall \beta_0 \neq 0$, we have $(4\beta_0^2 - \partial_x^2)^{-1}$ is a S^{-2} -multiplier, and $\partial_x(4\beta_0^2 - \partial_x^2)^{-1}$ is a S^{-1} -multiplier.

Next, we introduce the paradifferential calculus of Besov space and the main continuity properties of the paraproduct and the remainder.

Definition 2.5. [1] The nonhomogeneous paraproduct of v by u is defined by

$$T_u v = \sum_j S_{j-1} u \Delta_j v$$

The nonhomogeneous remainder of u and v is defined by

$$R(u, v) = \sum_{|k-j| \leq 1} \Delta_k u \Delta_j v$$

Lemma 2.6. [1] (1) $\forall t < 0$, $s \in \mathbb{R}$, $u \in B_{p,r_1}^t \cap L^\infty$, $v \in B_{p,r_2}^s$ with $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$, then

$$\|T_u v\|_{B_{p,r_2}^s} \leq C \|u\|_{L^\infty} \|v\|_{B_{p,r_2}^s}$$

or

$$\|T_u v\|_{B_{p,r}^{s+t}} \leq C \|u\|_{B_{\infty,r_1}^t} \|v\|_{B_{p,r_2}^s}$$

(2) $\forall s_1, s_2 \in \mathbb{R}, 1 \leq p_1, p_2, r_1, r_2 \leq \infty$, with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2} \leq 1, \frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2} \leq 1$. Then $\forall (u, v) \in B_{p_1, r_1}^{s_1} \times B_{p_2, r_2}^{s_2}$, if $s_1 + s_2 > 0$

$$\|R(u, v)\|_{B_{p, r}^{s_1+s_2}} \leq C \|u\|_{B_{p_1, r_1}^{s_1}} \|v\|_{B_{p_2, r_2}^{s_2}}$$

If $r = 1$ and $s_1 + s_2 = 0$,

$$\|R(u, v)\|_{B_{p, \infty}^0} \leq C \|u\|_{B_{p_1, r_1}^{s_1}} \|v\|_{B_{p_2, r_2}^{s_2}}$$

We then have the following product laws:

Lemma 2.7. [1, 17] (1) For any $s > 0$ and any (p, r) in $[1, \infty]^2$, the space $L^\infty \cap B_{p, r}^s$ is an algebra, and a constant C exists such that

$$\|uv\|_{B_{p, r}^s} \leq C (\|u\|_{L^\infty} \|v\|_{B_{p, r}^s} + \|u\|_{B_{p, r}^s} \|v\|_{L^\infty}).$$

(2) If $1 \leq p, r \leq \infty, s_1 \leq s_2, s_2 > \frac{d}{p} (s_2 \geq \frac{d}{p} \text{ if } r = 1)$ and $s_1 + s_2 > \max(0, \frac{2d}{p} - d)$, there exists C such that

$$\|uv\|_{B_{p, r}^{s_1}} \leq C \|u\|_{B_{p, r}^{s_1}} \|v\|_{B_{p, r}^{s_2}}.$$

(3) If $1 \leq p \leq 2, \forall u \in B_{p, \infty}^{\frac{d}{p}-d}(\mathbb{R}^d), \forall v \in B_{p, 1}^{\frac{d}{p}}(\mathbb{R}^d)$, there exists C such that

$$\|uv\|_{B_{p, \infty}^{\frac{d}{p}-d}} \leq C \|u\|_{B_{p, \infty}^{\frac{d}{p}-d}} \|v\|_{B_{p, 1}^{\frac{d}{p}}}.$$

Proposition 2.8. [1] Let $s \in \mathbb{R}, 1 \leq p, r \leq \infty$.

$$(2.3) \quad \begin{cases} B_{p, r}^s \times B_{p', r'}^{-s} & \longrightarrow \mathbb{R}, \\ (u, \phi) & \longmapsto \sum_{|j-j'| \leq 1} \langle \Delta_j u, \Delta_{j'} \phi \rangle, \end{cases}$$

defines a continuous bilinear functional on $B_{p, r}^s \times B_{p', r'}^{-s}$. Denoted by $Q_{p', r'}^{-s}$ the set of functions ϕ in \mathcal{S}' such that $\|\phi\|_{B_{p', r'}^{-s}} \leq 1$. If u is in \mathcal{S}' , then we have

$$\|u\|_{B_{p, r}^s} \leq C \sup_{\phi \in Q_{p', r'}^{-s}} \langle u, \phi \rangle.$$

We now state the definition of Osgood modulus of continuity and Osgood lemma.

Definition 2.9. [1] Let $a \in (0, 1]$. Nondecreasing nonzero continuous function μ is defined on $[0, a]$ and $\mu : [0, a] \rightarrow \mathbb{R}^+$ s.t. $\mu(0) = 0$, then μ is called a modulus of continuity. And if μ is called an Osgood modulus of continuity if

$$\int_0^a \frac{dr}{\mu(r)} = \infty$$

The Osgood lemma are useful to prove the uniqueness of some ordinary differential equations and one can utilize a procedure to construct a large families of Osgood functions.

Lemma 2.10 (Osgood lemma). [1] Let ρ be a measurable function mapping $[t_0, T]$ to $[0, a]$, γ is a locally integrable function from $[t_0, T]$ to \mathbb{R}^+ , and μ is a nondecreasing function mapping $[0, a]$ to \mathbb{R}^+ . Assume that, for some nonnegative real number c , the function ρ satisfies

$$\rho(t) \leq c + \int_{t_0}^t \gamma(t') \mu(\rho(t')) dt' \quad \text{for a.e. } t \in [t_0, T]$$

- If c is positive

$$-\mathcal{M}(\rho(t)) + \mathcal{M}(c) \leq \int_{t_0}^t \gamma(t') dt' \quad \text{with} \quad \mathcal{M}(x) = \int_x^a \frac{dr}{\mu(r)} \quad \text{for a.e. } t \in [t_0, T]$$

- If $c = 0$ and μ is an Osgood modulus of continuity, then $\rho = 0$, a.e. $t \in [t_0, T]$.

Next, we state some useful result in the transport equation theory, which are crucial to the proofs of our main theorems later. Let us consider the linear transport equation:

$$(2.4) \quad \begin{cases} \partial_t f + v \cdot \nabla f = g \\ f|_{t=0} = f_0 \end{cases}$$

Lemma 2.11 (A priori estimates in Besov spaces). [1] Let $s \in \mathbb{R}$, $1 \leq p, r \leq \infty$. Assume that

$$s \geq -d \min\left(\frac{1}{p_1}, \frac{1}{p'}\right)$$

There exists a constant C such that for all solutions $f \in L^\infty([0, T]; B_{p,r}^s)$ of (2.4) in d -dimensional space with initial data $f_0 \in B_{p,r}^s$, and $g \in L^1([0, T]; B_{p,r}^s)$, we have for a.e. $t \in [0, T]$,

$$\|f(t)\|_{B_{p,r}^s} \leq \|f_0\|_{B_{p,r}^s} + \int_0^t \|g(t')\|_{B_{p,r}^s} dt' + \int_0^t V'_{p_1}(t') \|f(t)\|_{B_{p,r}^s} dt'$$

or

$$\|f(t)\|_{B_{p,r}^s} \leq e^{CV_{p_1}(t)} \left(\|f_0\|_{B_{p,r}^s} + \int_0^t e^{-CV_{p_1}(t')} \|g(t')\|_{B_{p,r}^s} dt' \right)$$

with

$$V'_{p_1}(t) = \begin{cases} \|\nabla v\|_{B_{p_1, \infty}^{\frac{d}{p_1}} \cap L^\infty}, & \text{if } s < 1 + \frac{d}{p_1} \\ \|\nabla v\|_{B_{p_1, r}^{s-1}}, & \text{if } s > 1 + \frac{d}{p_1} \text{ or } \left\{ s = 1 + \frac{d}{p_1} \text{ and } r = 1 \right\} \end{cases}$$

If $f = v$, for all $s > 0$

$$V'_{p_1}(t) = \|\nabla v(t)\|_{L^\infty}.$$

Lemma 2.12. [21, 17] If

$$\left\{ s \geq \frac{d}{2}, p > 2, 1 \leq r \leq \infty \right\} \text{ or } \left\{ s > \frac{d}{p}, 1 \leq p \leq 2, 1 \leq r \leq \infty \right\} \text{ or } \left\{ s = \frac{d}{p}, 1 \leq p \leq 2, r = 1 \right\}$$

then for the solution $f \in L^\infty([0, T]; B_{p,r}^s)$ of (2.4) with $v \in L^1([0, T]; B_{p,r}^{s+1})$, then the initial data $f_0 \in B_{p,r}^s$ and $g \in L^1([0, T]; B_{p,r}^s)$, we then have

$$\|f(t)\|_{B_{p,r}^s} \leq e^{CV(t)} (\|f_0\|_{B_{p,r}^s} + \int_0^t e^{-CV(\tau)} \|g(\tau)\|_{B_{p,r}^s} d\tau)$$

with $V(t) = \int_0^t \|v(\tau)\|_{B_{p,r}^{s+1}} d\tau$ and $C = C(p, r, s)$.

Lemma 2.13. [1] Let $1 \leq p \leq p_1 \leq \infty$, $1 \leq r \leq \infty$, $s > -d \min(\frac{1}{p_1}, \frac{1}{p'})$. Let $f_0 \in B_{p,r}^s$, $g \in L^1([0, T]; B_{p,r}^s)$, and let v be a time-dependent vector field such that $v \in L^\rho([0, T]; B_{\infty, \infty}^{-M})$ for some $\rho > 1$ and $M > 0$, and

$$\begin{aligned} \nabla v &\in L^1([0, T]; B_{p_1, \infty}^{\frac{d}{p_1}}), & \text{if } s < 1 + \frac{d}{p_1}, \\ \nabla v &\in L^1([0, T]; B_{p,r}^{s-1}), & \text{if } s > 1 + \frac{d}{p_1} \text{ or } (s = 1 + \frac{d}{p_1} \text{ and } r = 1). \end{aligned}$$

Then the equation (2.4) has a unique solution f in

- the space $C([0, T]; B_{p,r}^s)$, if $r < \infty$;
- the space $\left(\bigcap_{s' < s} C([0, T]; B_{p,\infty}^{s'})\right) \cap C_w([0, T]; B_{p,\infty}^s)$, if $r = \infty$.

Lemma 2.14. [19] Let $1 \leq p \leq \infty, 1 \leq r \leq \infty, s > 1 + \frac{d}{p}$ (or $s = 1 + \frac{d}{p}, r = 1, 1 \leq p < \infty$). Denote $\bar{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$. Let $\{v^n\}_{n \in \bar{\mathbb{N}}} \subset C([0, T]; B_{p,r}^{s-1})$. Assume that v^n is the solution to

$$\begin{cases} \partial_t v^n + a^n \partial_x v^n = f \\ v^n|_{t=0} = v_0 \end{cases}$$

with $v_0 \in B_{p,r}^{s-1}$, $f \in L^1([0, T]; B_{p,r}^{s-1})$ and for some $\alpha \in L^1([0, T])$

$$\sup_{n \in \bar{\mathbb{N}}} \|a^n\|_{B_{p,r}^s} \leq \alpha(t)$$

If $a^n \rightarrow a^\infty$ in $L^1([0, T]; B_{p,r}^{s-1})$ as n tends to ∞ , then $v^n \rightarrow v^\infty$ in $C([0, T]; B_{p,r}^{s-1})$ as n tends to ∞ .

3 Local well-posedness

In this section, we will establish the local well-posedness for the Cauchy problem (1.1) or (1.2) in Besov spaces. We want to solve the Cauchy problem (1.2) directly by general Picard scheme and transport equation theory, but we will fail to do that due to the loss of regularity of the transport term. By Prop 2.4, we can overcome the difficulty by solving the Cauchy problem (1.1).

Firstly, we give a definition as follows

Definition 3.1. Let $T > 0, s \in \mathbb{R}$ and $1 \leq p, r \leq \infty$. Denote

$$E_{p,r}^s(T) \triangleq \begin{cases} C([0, T]; B_{p,r}^s) \cap C^1([0, T]; B_{p,r}^{s-1}) & \text{if } r < \infty \\ C_w([0, T]; B_{p,r}^s) \cap C^{0,1}([0, T]; B_{p,r}^{s-1}) & \text{if } r = \infty \end{cases}$$

$$E_{p,r}^s(T_-) \triangleq \begin{cases} C([0, T]; B_{p,r}^s) \cap C^1([0, T]; B_{p,r}^{s-1}) & \text{if } r < \infty \\ C_w([0, T]; B_{p,r}^s) \cap C^{0,1}([0, T]; B_{p,r}^{s-1}) & \text{if } r = \infty \end{cases}$$

Theorem 3.2. Let $1 \leq p, r \leq \infty, s \in \mathbb{R}$ and (s, p, r) satisfies the condition

$$(3.1) \quad \left\{s \geq \frac{1}{2}, p > 2, 1 \leq r \leq \infty\right\} \text{ or } \left\{s > \frac{1}{p}, 1 \leq p \leq 2, 1 \leq r \leq \infty\right\} \text{ or } \left\{s = \frac{1}{p}, 1 \leq p \leq 2, r = 1\right\}$$

Let $n_0 \triangleq 4\beta_0^2 v_0 - v_{0xx}$ with $\beta_0 \neq 0$ be in $B_{p,r}^s$. Then (1.1) has a unique maximal solution n in $E_{p,r}^s(T_-^*)$ and n depends continuously on the initial data n_0 .

Moreover, there exists a positive constant C , depending only on s, p, r, β_0 s.t

$$T^* \geq \frac{C}{\|n_0\|_{B_{p,r}^s}}$$

Proof. Now we prove Theorem 3.2 as the following steps.

Step 1: Constrcuting Approximate Solitions and Uniform Bounds.

Denoting $v^0 = 0$, we define a sequence $(n^k)_{k \in \mathbb{N}}$ of smooth functions by solving the following linear transport equation:

$$(T_k) \begin{cases} n_t^{k+1} - 4(v_x^k + 2\beta_0 v^k) n_x^{k+1} = 16\beta_0^2 v^k n^k - 4(n^k)^2 + 8\beta_0 v_x^k n^k \\ v^{k+1} = (4\beta_0^2 - \partial_x^2)^{-1} n^{k+1} \\ v^{k+1}|_{t=0} = S_{k+1} v_0 \end{cases}$$

By induction, we assume that $(n^k)_{k \in \mathbb{N}}, \forall T > 0$. Since $(4\beta_0^2 - \partial_x^2)^{-1}$ is an S^{-2} -multiplier and

$$v^{k+1} = (4\beta_0^2 - \partial_x^2)^{-1} n^{k+1} = \frac{1}{4|\beta_0|} e^{-|2\beta_0 \cdot| * n^{k+1}}$$

also noticing that $B_{p,r}^s$ is an algebra when (s, p, r) satisfies the condition (3.1) and the embedding $B_{p,r}^s \hookrightarrow B_{p,r}^{s-1} \hookrightarrow B_{p,r}^{s-2}$, we obtain

$$\|16\beta_0^2 v^k n^k - 4(n^k)^2 + 8\beta_0 v_x^k n^k\|_{B_{p,r}^s} \leq C \|n^k\|_{B_{p,r}^s}^2$$

At the same time $\|S_{k+1} n_0\|_{B_{p,r}^s} \leq C \|n_0\|_{B_{p,r}^s}$, and denote that

$$\int_0^t \|4(v_x^k + 2\beta_0 v^k)_x(\tau)\|_{B_{p,r}^s} d\tau \leq C N^k(t) \triangleq \int_0^t \|n^k(\tau)\|_{B_{p,r}^s} d\tau$$

Applying Thm2.11, we have

$$(3.2) \quad \|n^{k+1}(t)\|_{B_{p,r}^s} \leq e^{CN^k(t)} (\|n_0\|_{B_{p,r}^s} + C \int_0^t e^{-CN^k(\tau)} \|n^k(\tau)\|_{B_{p,r}^s}^2 d\tau)$$

We may assume $C \geq 1$ and fix a $T > 0$ s.t.

$$(3.3) \quad 2C^2 \|n_0\|_{B_{p,r}^s} T < 1$$

Suppose that $\forall t \in [0, T]$, n^k satisfies

$$(3.4) \quad \|n^k(t)\|_{B_{p,r}^s} \leq \frac{C \|n_0\|_{B_{p,r}^s}}{1 - 2C^2 \|n_0\|_{B_{p,r}^s} t}$$

Note that

$$e^{CN^k(t_2) - CN^k(t_1)} = e^{C \int_{t_1}^{t_2} \|v^k(\tau)\|_{B_{p,r}^s} d\tau} \leq e^{\int_{t_1}^{t_2} \frac{C^2 \|n_0\|_{B_{p,r}^s}}{1 - 2C^2 \tau \|n_0\|_{B_{p,r}^s}} d\tau} = \left(\frac{1 - 2C^2 \|n_0\|_{B_{p,r}^s} t_1}{1 - 2C^2 \|n_0\|_{B_{p,r}^s} t_2} \right)^{1/2}$$

Plugging (3.4) into (3.2) yields

$$(3.5) \quad \begin{aligned} & \|n^{k+1}(t)\|_{B_{p,r}^s} \\ & \leq (1 - 2C^2 \|n_0\|_{B_{p,r}^s} t)^{1/2} \left(C \|n_0\|_{B_{p,r}^s} + \int_0^t C (1 - 2C^2 \|n_0\|_{B_{p,r}^s} \tau)^{1/2} \frac{C^2 \|n_0\|_{B_{p,r}^s}}{(1 - 2C^2 \|n_0\|_{B_{p,r}^s} \tau)^2} d\tau \right) \\ & \leq \frac{C \|n_0\|_{B_{p,r}^s}}{1 - 2C^2 \|n_0\|_{B_{p,r}^s} t} \end{aligned}$$

Therefore, $(n^k)_{k \in \mathbb{N}}$ is uniformly bounded in $C([0, T]; B_{p,r}^s)$. According to the system (T_k) and Lemma2.11 and Lemma2.12, we can deduce that $\{n^k\}_{k \in \mathbb{N}}$ is uniformly bounded in $E_{p,r}^s(T)$.

Step 2: Convergence and Regularity.

We will claim that $\{n^k\}_{k \in \mathbb{N}}$ is a Cauchy sequence in $C([0, T]; B_{p,r}^{s-1})$, if $\{s \geq \frac{1}{2}, p > 2, 1 \leq r \leq \infty\}$ or $\{s > \frac{1}{p}, 1 \leq p \leq 2, 1 \leq r \leq \infty\}$; or of $C([0, T]; B_{p,\infty}^{\frac{1}{p}-1})$ if $\{1 \leq p \leq 2, s = \frac{1}{p}, r = 1\}$.

Indeed, for all $k, p \in \mathbb{R}$, taking the difference between the system (T_{k+p}) and T_k , and denote $w^{k,p} = n^{k+p} - n^k$, we have

$$(3.6) \quad \begin{cases} (\partial_t - 4(v_x^{k+p} + 2\beta_0 v^{k+p})) \partial_x w^{k+1,p} \\ = -4(v_x^{k+p} - v_x^k) n_x^{k+1} - 8\beta_0 (v^{k+p} - v^k) n_x^{k+1} - 4(n^{k+p} - n^k)(n^{k+p} + n^k) \\ \quad - 16\beta_0^2 ((v^{k+p} - v^k) n^{k+p} + v^k (n^{k+p} - n^k)) + 8\beta ((v_x^{k+p} - v_x^k) n_x^{k+p} + (n^{k+p} - n^k) v_x^k) \\ w^{k+1,p}|_{t=0} = S_{k+p+1} n_0 - S_{k+1} n_0 \end{cases}$$

Case1: $s \geq \frac{1}{2}, p > 2, 1 \leq r \leq \infty$ or $s > \frac{1}{p}, 1 \leq p \leq 2, 1 \leq r \leq \infty$

First, estimate the left-hand side of system (3.6) in $B_{p,r}^{s-1}$. In fact, by Lemma2.6 and Lemma2.4, we can see

$$\begin{aligned} \|(v_x^{k+p} - v_x^k) n_x^{k+1}\|_{B_{p,r}^{s-1}} &\leq C \|v_x^{k+p} - v_x^k\|_{L^\infty} \|n_x^{k+1}\|_{B_{p,r}^{s-1}} + C \|v_x^{k+p} - v_x^k\|_{B_{p,r}^s} \|n_x^{k+1}\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C \|v_x^{k+p} - v_x^k\|_{B_{p,r}^s} \|n_x^{k+1}\|_{B_{p,r}^{\frac{1}{p}-1}} \leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} \|n^{k+1}\|_{B_{p,r}^s} \\ \|(v^{k+p} - v^k) n_x^{k+1}\|_{B_{p,r}^{s-1}} &\leq C \|v^{k+p} - v^k\|_{L^\infty} \|n_x^{k+1}\|_{B_{p,r}^{s-1}} + C \|v^{k+p} - v^k\|_{B_{p,r}^s} \|n_x^{k+1}\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C \|v^{k+p} - v^k\|_{B_{p,r}^s} \|n_x^{k+1}\|_{B_{p,r}^{\frac{1}{p}-1}} \leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} \|n^{k+1}\|_{B_{p,r}^s} \\ \|(n^{k+p} - n^k)(n^{k+p} + n^k)\|_{B_{p,r}^{s-1}} &\leq C \|n^{k+p} + n^k\|_{L^\infty} \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} + C \|n^{k+p} + n^k\|_{B_{p,r}^s} \|n^{k+p} - n^k\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C \|n^{k+p} + n^k\|_{B_{p,r}^s} \|n^{k+p} - n^k\|_{B_{p,r}^{\frac{1}{p}-1}} \\ &\leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} (\|n^{k+p}\|_{B_{p,r}^s} + \|n^k\|_{B_{p,r}^s}) \\ \|(v^{k+p} - v^k) n^{k+p}\|_{B_{p,r}^{s-1}} &\leq C \|v^{k+p} - v^k\|_{L^\infty} \|n^{k+p}\|_{B_{p,r}^{s-1}} + C \|v^{k+p} - v^k\|_{B_{p,r}^s} \|n^{k+p}\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C \|v^{k+p} - v^k\|_{B_{p,r}^s} \|n^{k+p}\|_{B_{p,r}^{\frac{1}{p}-1}} \leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} \|n^k\|_{B_{p,r}^s} \\ \|v^k (n^{k+p} - n^k)\|_{B_{p,r}^{s-1}} &\leq C \|v^k\|_{L^\infty} \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} + C \|v^k\|_{B_{p,r}^s} \|n^{k+p} - n^k\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C \|v^k\|_{B_{p,r}^s} \|n^{k+p} - n^k\|_{B_{p,r}^{\frac{1}{p}-1}} \leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} \|n^k\|_{B_{p,r}^s} \\ \|(v_x^{k+p} - v_x^k) n^{k+p}\|_{B_{p,r}^{s-1}} &\leq C \|v_x^{k+p} - v_x^k\|_{B_{p,r}^s} \|n^{k+p}\|_{B_{p,r}^s} \leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} \|n^{k+p}\|_{B_{p,r}^s} \\ \|(n^{k+p} - n^k) v_x^k\|_{B_{p,r}^{s-1}} &\leq C \|v_x^k\|_{L^\infty} \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} + C \|v_x^k\|_{B_{p,r}^s} \|n^{k+p} - n^k\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C \|v_x^k\|_{B_{p,r}^s} \|n^{k+p} - n^k\|_{B_{p,r}^{\frac{1}{p}-1}} \leq C \|n^{k+p} - n^k\|_{B_{p,r}^{s-1}} \|n^k\|_{B_{p,r}^s} \end{aligned}$$

Also we can obtain

$$\|S_{k+p+1} n_0 - S_{k+1} n_0\|_{B_{p,r}^{s-1}} \leq C \left(\sum_{l=k}^{k+p} 2^{-lr} 2^{lsr} \|\Delta_l n_0\|_{L^p}^r \right)^{\frac{1}{r}} \rightarrow 0 (k \rightarrow \infty)$$

Now, recalling (3.4) and the uniform bound estimates of $\|n^k\|_{B_{p,r}^s}$, and making use of Lemma2.11 or Lemma2.12 to the system (3.6), we have

$$\|w^{k+1,p}\|_{B_{p,r}^{s-1}} \leq C_T (2^{-k} + \int_0^t \|w^{k,p}(\tau)\|_{B_{p,r}^{s-1}} d\tau)$$

where $C_T = C(s, p, r, T, \beta_0, \|v_0\|_{B_{p,r}^s})$ is a constant, then by induction as k tends to ∞

$$\begin{aligned} \|w^{k+1,p}\|_{B_{p,r}^{s-1}} &\leq \sum_{l=0}^p \|w^{k+l+1,1}\|_{B_{p,r}^{s-1}} \leq \sum_{l=0}^p C_T(2^{-(k+l)}) + \int_0^t C_T(2^{-(k+l-1)}) + \int_0^{t_1} \|w^{k+l,1}\|_{B_{p,r}^{s-1}} dt_2 dt_1 \\ &\leq \sum_{l=0}^p C_t(2^{-(k+l)}) \sum_{m=0}^{k+l} \frac{(2TC_T)^m}{m!} + (C_T)^{k+l+1} \int_0^t \frac{(t-\tau)^{k+l+1}}{(k+l+1)!} d\tau \\ &\leq \sum_{l=0}^p \left(C_T 2^{-(k+l)} e^{2TC_T} + C_T \frac{(TC_T)^{k+l+1}}{(k+l+1)!} \right) \leq C_T 2^{-k+1} e^{2TC_T} + \sum_{m=k+1}^{k+p+1} C_T \frac{(TC_T)^m}{m!} \rightarrow 0 \end{aligned}$$

which means $(n^k)_{k \in \mathbb{N}} \subset E_{p,r}^s(T)$, and by Fatou's property, we obtain that $n \in L^\infty([0, T]; B_{p,r}^s)$.

Now for any test function $\phi \in C([0, T]; \mathcal{S})$ in the system (T_k) , applying Prop2.8, and taking the limits, it is not hard to check that n is a solution to the system (1.1).

According to Lemma2.13, $n \in C([0, T]; B_{p,r}^s)$, and by system(1.1), we can get that $n_t \in C([0, T]; B_{p,r}^{s-1})$, thus $n \in E_{p,r}^s(T)$.

Case2: $s = 1/p$, $1 \leq p \leq 2$ and $r = 1$

Since the critical index $s - 1 = -(1 - \frac{1}{p}) = -\min(\frac{1p}{p}, \frac{1p'}{p'})$, we can only apply Lemma2.11 in the stage space $B_{p,1}^{\frac{1}{p}-1}$. And noticing that $B_{p,1}^{\frac{1}{p}-1} \hookrightarrow B_{p,1}^{\frac{1}{p}-1} \hookrightarrow B_{p,\infty}^{\frac{1}{p}-1}$, and applying Lemma2.6 to the system (3.6)

$$\begin{aligned} \|(v_x^{k+p} - v_x^k)n_x^{k+1}\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|v_x^{k+p} - v_x^k\|_{L^\infty} \|n_x^{k+1}\|_{B_{p,\infty}^{\frac{1}{p}-1}} + C\|v_x^{k+p} - v_x^k\|_{B_{p,\infty}^{\frac{1}{p}}} \|n_x^{k+1}\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C\|v_x^{k+p} - v_x^k\|_{B_{p,1}^{\frac{1}{p}}} \|n_x^{k+1}\|_{B_{p',\infty}^{-\frac{1}{p}}} \leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \|n^{k+1}\|_{B_{p,1}^{\frac{1}{p}}} \\ \|(v^{k+p} - v^k)n_x^{k+1}\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|v^{k+p} - v^k\|_{L^\infty} \|n_x^{k+1}\|_{B_{p,\infty}^{\frac{1}{p}-1}} + C\|v^{k+p} - v^k\|_{B_{p,\infty}^{\frac{1}{p}}} \|n_x^{k+1}\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C\|v^{k+p} - v^k\|_{B_{p,1}^{\frac{1}{p}}} \|n_x^{k+1}\|_{B_{p',\infty}^{-\frac{1}{p}}} \leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \|n^{k+1}\|_{B_{p,1}^{\frac{1}{p}}} \\ \|(n^{k+p} - n^k)(n^{k+p} + n^k)\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|n^{k+p} + n^k\|_{L^\infty} \|n^{k+p} - n^k\|_{B_{p,\infty}^{\frac{1}{p}}} + C\|n^{k+p} + n^k\|_{B_{p,\infty}^{\frac{1}{p}-1}} \|n^{k+p} - n^k\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C\|n^{k+p} + n^k\|_{B_{p',\infty}^{-\frac{1}{p}+1}} \|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \\ &\leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} (\|n^{k+p}\|_{B_{p,1}^{\frac{1}{p}}} + \|n^k\|_{B_{p,1}^{\frac{1}{p}}}) \\ \|(v^{k+p} - v^k)n^{k+p}\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|v^{k+p} - v^k\|_{L^\infty} \|n^{k+p}\|_{B_{p,\infty}^{\frac{1}{p}-1}} + C\|v^{k+p} - v^k\|_{B_{p,\infty}^{\frac{1}{p}}} \|n^{k+p}\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C\|v^{k+p} - v^k\|_{B_{p,1}^{\frac{1}{p}}} \|n^{k+p}\|_{B_{p',\infty}^{-\frac{1}{p}}} \leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \|n^k\|_{B_{p,1}^{\frac{1}{p}}} \\ \|v^k(n^{k+p} - n^k)\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|v^k\|_{L^\infty} \|n^{k+p} - n^k\|_{B_{p,\infty}^{\frac{1}{p}-1}} + C\|v^k\|_{B_{p,\infty}^{\frac{1}{p}}} \|n^{k+p} - n^k\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C\|v^k\|_{B_{p',\infty}^{-\frac{1}{p}+1}} \|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \|n^k\|_{B_{p,1}^{\frac{1}{p}}} \\ \|(v_x^{k+p} - v_x^k)n^{k+p}\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|v_x^{k+p} - v_x^k\|_{B_{p,1}^{\frac{1}{p}}} \|n^{k+p}\|_{B_{p,1}^{\frac{1}{p}}} \leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \|n^{k+p}\|_{B_{p,1}^{\frac{1}{p}}} \\ \|(n^{k+p} - n^k)v_x^k\|_{B_{p,\infty}^{\frac{1}{p}-1}} &\leq C\|v_x^k\|_{B_{p,1}^{\frac{1}{p}}} \|n^{k+p} - n^k\|_{B_{p,\infty}^{\frac{1}{p}-1}} + C\|v_x^k\|_{B_{p,\infty}^{\frac{1}{p}}} \|n^{k+p} - n^k\|_{B_{\infty,\infty}^{-1}} \\ &\quad + C\|v_x^k\|_{B_{p',\infty}^{\frac{1}{p}}} \|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \leq C\|n^{k+p} - n^k\|_{B_{p,1}^{\frac{1}{p}-1}} \|n^k\|_{B_{p,1}^{\frac{1}{p}}} \end{aligned}$$

Also we can obtain the initial data of system (3.6) satisfies

$$\|S_{k+p+1}n_0 - S_{k+1}n_0\|_{B_{p,\infty}^{\frac{1}{p-1}}} \leq C \sum_{l=k}^{k+p+1} 2^{l(1/p-1)} \|\Delta_l n_0\|_{L^p} \leq C 2^{-k} \|n_0\|_{B_{p,1}^{\frac{1}{p-1}}} \longrightarrow 0 (k \rightarrow \infty)$$

Since $B_{p,1}^{\frac{1}{p}} \hookrightarrow B_{p,\infty}^{\frac{1}{p}}$ and $B_{p,1}^{\frac{1}{p}} \hookrightarrow L^\infty$, we obtain the estimate of the flow

$$\int_0^t \|4\partial_x(v_x^{k+p} + 2\beta_0 v^{k+p})\|_{B_{p,\infty}^{1/p} \cap L^\infty} d\tau \leq C \int_0^t \|n^{k+p}\|_{B_{p,1}^{1/p}} d\tau$$

According to the logarithm interpolation of Lemma 2.3, and the uniform bound of $\{n^k\}_{k \in \mathbb{N}}$, we obtain

$$\|w^{k+1,p}\|_{B_{p,\infty}^{\frac{1}{p-1}}} \leq C_T \left(2^{-k} + \int_0^t \|w^{k,p}\|_{B_{p,\infty}^{\frac{1}{p-1}}} \log \left(e + \frac{1}{\|w^{k,p}\|_{B_{p,\infty}^{\frac{1}{p-1}}}} \right) d\tau \right)$$

denote that $w_k(t) = \sup_{\tau \in [0,t], p \in \mathbb{N}} \|(n^{k+p} - n^k)(\tau)\|_{B_{p,\infty}^{\frac{1}{p-1}}}$, then we have

$$w_{k+1}(t) \leq C_T \left(2^{-k} + \int_0^t w_k(\tau) \log \left(e + \frac{1}{w_k(\tau)} \right) d\tau \right)$$

Note that $\{w_k\}_{k \in \mathbb{N}}$, thus it is obvious that $\mu(x) = C_T x \log(e + \frac{1}{x})$ is an Osgood modulus of continuity. Applying Lesbesgue-Fatou's lemma in real analysis, we conclude that, for $\forall t \in [0, T]$

$$\limsup_{k \rightarrow \infty} w_k(t) \leq \int_0^t \mu(\limsup_{k \rightarrow \infty} w_k(\tau)) d\tau$$

Hence according to the Osgood lemma, we have

$$\limsup_{k \rightarrow \infty} w_k(t) = 0, \quad t \in [0, T]$$

which implies that $n^k_{k \in \mathbb{N}}$ is a Cauchy sequence in the space $C([0, T]; B_{p,\infty}^{1/p-1})$ and converges to the limit function $n \in C([0, T]; B_{p,\infty}^{1/p})$. Along the same line of Case 1, similarly we deduce that n is a solution to (1.1) and $n \in E_{p,1}^{\frac{1}{p}}(T)$.

Step 3: Uniqueness.

Suppose n_1, n_2 are two solutions of (1.1) with initial data n_{10} and n_{20} respectively. Denoting $w \triangleq n_1 - n_2$, we have

$$(3.7) \quad \begin{cases} (\partial_t - 4(v_{1x} + 2\beta_0 v_1))\partial_x w \\ = -4(v_{1x} - v_{2x})n_{2x} - 8\beta_0(v_1 - v_2)n_{2x} - 4(n_1 - n_2)(n_1 + n_2) \\ \quad - 16\beta_0^2((v_1 - v_2)n_1 + v_2(n_1 - n_2)) + 8\beta((v_{1x} - v_{2x})n_1 + (n_1 - n_2)v_{1x}) \\ v_1 = (4\beta_0^2 - \partial_x^2)^{-1}n_1, \quad v_2 = (4\beta_0^2 - \partial_x^2)^{-1}n_2 \\ w|_{t=0} = n_{10} - n_{20} \end{cases}$$

If $s \geq 1/2, p > 2$ or $s > 1/p, 1 \leq p \leq 2$, along the same computation in step 2, $\forall t \in [0, T]$, we have inequality

$$e^{-C \int_0^t \|n_1\|_{B_{p,r}^s} d\tau} \|w(t)\|_{B_{p,r}^{s-1}} \leq \|w_0\|_{B_{p,r}^{s-1}} + C \int_0^t e^{-C \int_0^{t'} \|n_{10}\|_{B_{p,r}^s} d\tau} \|w(t')\|_{B_{p,r}^{s-1}} (\|n_1\|_{B_{p,r}^s} + \|n_2\|_{B_{p,r}^s}) dt'$$

By Gronwall's inequality

$$(3.8) \quad \|w(t)\|_{B_{p,r}^{s-1}} \leq \|w_0\|_{B_{p,r}^{s-1}} \exp\left(C \int_0^t (\|n_1\|_{B_{p,r}^s} + \|n_2\|_{B_{p,r}^s}) d\tau\right)$$

If $s = 1/p, 1 \leq p \leq 2, r = 1$, similar to step 2, we have inequality

$$\begin{aligned} & e^{-C \int_0^t \|n_1\|_{B_{p,1}^{1/p}} d\tau} \|w(t)\|_{B_{p,\infty}^{\frac{1}{p}-1}} \\ & \leq \|w_0\|_{B_{p,\infty}^{\frac{1}{p}-1}} + C \int_0^t e^{-C \int_0^{\tau'} \|n_1\|_{B_{p,1}^{1/p}} d\tau} \|w(\tau')\|_{B_{p,\infty}^{\frac{1}{p}-1}} (\|n_1\|_{B_{p,1}^{1/p}} + \|n_2\|_{B_{p,1}^{1/p}}) \log\left(e + \frac{\|n_1\|_{B_{p,1}^{1/p}} + \|n_2\|_{B_{p,1}^{1/p}}}{\|w(\tau')\|_{B_{p,\infty}^{\frac{1}{p}-1}}}\right) d\tau' \end{aligned}$$

denote $h(t) \triangleq \|w(t)\|_{B_{p,\infty}^{\frac{1}{p}-1}} e^{-C \int_0^t \|n_1\|_{B_{p,1}^{1/p}} d\tau} / (\|n_1\|_{L_T^\infty(B_{p,1}^{1/p})} + \|n_2\|_{L_T^\infty(B_{p,1}^{1/p})})$ and $|h(t)| \leq 1$, obviously, since

$$x \log\left(e + \frac{1}{x}\right) \leq x(2 - \log x), \quad \forall x \in [0, 1]$$

then we have

$$h(t) \leq h(0) + C \int_0^t h(\tau) (\|n_1\|_{L_T^\infty(B_{p,1}^{1/p})} + \|n_2\|_{L_T^\infty(B_{p,1}^{1/p})})^2 (2 - \log h(\tau)) d\tau$$

Since $\mu(x) = x(2 - \log x)$ is an Osgood modulus of continuity on $[0, 1]$, then by Osgood lemma, we get inequality

$$\int_{h(0)}^{h(t)} \frac{dr}{r(2 - \log r)} \leq C \int_0^t (\|n_1\|_{L_T^\infty(B_{p,1}^{1/p})} + \|n_2\|_{L_T^\infty(B_{p,1}^{1/p})})^2 d\tau$$

then by the definition of h , we have

$$(3.9) \quad \begin{aligned} & \|n_1 - n_2\|_{L_T^\infty(B_{p,1}^{1/p-1})} \\ & \leq e^2 (\|n_1\|_{L_T^\infty(B_{p,1}^{1/p})} + \|n_2\|_{L_T^\infty(B_{p,1}^{1/p})}) \exp\left(C \int_0^t (\|n_1\|_{L_T^\infty(B_{p,1}^{1/p})} + \|n_2\|_{L_T^\infty(B_{p,1}^{1/p})}) d\tau\right) \\ & \quad \times (\|n_{10} - n_{20}\|_{(B_{p,\infty}^{1/p-1})})^{\exp(-C \int_0^t (\|n_1\|_{L_T^\infty(B_{p,1}^{1/p})} + \|n_2\|_{L_T^\infty(B_{p,1}^{1/p})}) d\tau)} \end{aligned}$$

Then the uniqueness of solution n to the system (1.1) in the space $E_{p,r}^p(T)$ follows from (3.8) and (3.9).

Step 4: Continuous Dependence.

Now according to Lemma 2.14, denote $\bar{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$. suppose $n^k \in C([0, T]; B_{p,r}^s)$ is the solution to (1.1) with initial data $n_0^k \in B_{p,r}^s$, then we have a sequence of systems

$$\begin{cases} \partial_t n^k - 4(v_x^k + 2\beta_0 v^k) \partial_x n^k = f^k \\ n^k = 4\beta_0^2 v^k - v_{xx}^k \\ n^k|_{t=0} = n_0^k \end{cases}$$

with $f^k \triangleq 16\beta_0^2 v^k n^k - 4(n^k)^2 + 8\beta v_x^k n^k$.

Next we claim that if $n_0^k \rightarrow n_0^\infty$ in $B_{p,r}^s$ when k tends to ∞ , then $n^k \rightarrow n^\infty$ in $C([0, T]; B_{p,r}^s)$ with T satisfying $2C^2 \sup_{k \in \bar{\mathbb{N}}} \|n_0^k\|_{B_{p,r}^s} T < 1$.

(1) $r|_\infty$. Decompose n^k into $n^k = y^k + z^k$, and satisfy

$$\begin{cases} \partial_t y^k - 4(v_x^k + 2\beta_0 v^k) \partial_x y^k = f^\infty \\ y^k|_{t=0} = n_0^k \end{cases} \quad \text{and} \quad \begin{cases} \partial_t z^k - 4(v_x^k + 2\beta_0 v^k) \partial_x z^k = f^k - f^\infty \\ z^k|_{t=0} = n_0^k - n_0^\infty \end{cases}$$

Denote $m = \sup_{k \in \mathbb{N}} \|n_0^k\|_{B_{p,r}^s}$, and letting $M = \frac{Cm}{1-2C^2mT}$, according to the proof of the existence, we get that

$$\|n^k(t)\|_{B_{p,r}^s} \leq M \quad \forall t \in [0, T]$$

Hence, $(f^k)_{k \in \mathbb{N}}$ is uniformly bounded in $C([0, T]; B_{p,r}^s)$ and

$$\|v_x^k + 2\beta_0 v^k\|_{B_{p,r}^{s+1}} \leq C \|v^k\|_{B_{p,r}^{s+2}} \leq C \|n^k\|_{B_{p,r}^s} \leq CM$$

also it is obvious that

$$\|(v^k - v^\infty)_x + 2\beta_0(v^k - v^\infty)\|_{B_{p,r}^{s+1}} \leq C \|n^k - n^\infty\|_{B_{p,r}^s}$$

similar to the proof of the uniqueness, since $n_0^k \rightarrow n_0^\infty$ in $B_{p,r}^s$, then we have $n^k \rightarrow n^\infty$ in $L^1([0, T]; B_{p,r}^{s-1}) \Rightarrow v_x^k + 2\beta_0 v^k \rightarrow v_x^\infty + 2\beta_0 v^\infty$ in $L^1([0, T]; B_{p,r}^s)$. Then by Lemma2.14

$$y^k \rightarrow n^\infty \text{ in } C([0, T]; B_{p,r}^s) \text{ as } k \rightarrow \infty$$

To control z^k , we need to estimate $f^k - f^\infty$, by Lemma2.6, we have

$$\|f^k - f^\infty\|_{B_{p,r}^s} \leq C(\|n^k\|_{B_{p,r}^s} + \|n^\infty\|_{B_{p,r}^s}) \|n^k - n^\infty\|_{B_{p,r}^s}$$

then according to Lemma2.11

$$\|z^k\|_{B_{p,r}^s} \leq Ce^{CMT} (\|n_0^k - n_0^\infty\|_{B_{p,r}^s} + \int_0^t \|(n^k - n^\infty)(\tau)\|_{B_{p,r}^s} d\tau)$$

Therefore, we have $\forall \varepsilon > 0, \exists N_1$ large enough, when $k > N_1, \|y^k - n^\infty\|_{B_{p,r}^s} < \varepsilon$, then

$$\|n^k - n^\infty\|_{B_{p,r}^s} \leq \|z^k\|_{B_{p,r}^s} + \|y^k - n^\infty\|_{B_{p,r}^s} \leq \varepsilon + Ce^{CMT} (\|n_0^k - n_0^\infty\|_{B_{p,r}^s} + \int_0^t \|(n^k - n^\infty)(\tau)\|_{B_{p,r}^s} d\tau)$$

by Gronwall's inequality, $\forall t \in [0, T]$

$$\|n^k - n^\infty\|_{B_{p,r}^s} \leq C(\varepsilon + \|n_0^k - n_0^\infty\|_{B_{p,r}^s})$$

Hence we gain the continuity of the system (1.1) in $C([0, T]; B_{p,r}^s)$ with respect to the initial data in $B_{p,r}^s$.

(2) $r = \infty$. Similar to Step2, we have $\|n^k - n^\infty\|_{L^\infty([0, T]; B_{p,\infty}^{s-1})} \rightarrow 0 (k \rightarrow \infty)$. Then $\forall \phi \in B_{p',1}^{-s}$, we have

$$\langle n^k - n^\infty \rangle = \langle S_j(n^k - n^\infty), \phi \rangle + \langle (Id - S_j)(n^k - n^\infty), \phi \rangle = \langle n^k - n^\infty, S_j \phi \rangle + \langle n^k - n^\infty, (Id - S_j) \phi \rangle$$

and

$$|\langle n^k - n^\infty, (Id - S_j) \phi \rangle| \leq CM \|\phi - S_j \phi\|_{B_{p',1}^{-s}} \quad \text{and} \quad |\langle n^k - n^\infty, S_j \phi \rangle| \leq \|n^k - n^\infty\|_{B_{p,\infty}^{s-1}} \|S_j \phi\|_{B_{p',1}^{1-s}}$$

by Lemma2.3, we can conclude that $\langle n^k - n^\infty, \phi \rangle \rightarrow 0$ as $k \rightarrow \infty$ for $r = \infty$. Thus we gain the continuity of the system (1.1) in $C_w([0, T]; B_{p,\infty}^s)$ with respect to the initial data in $B_{p,\infty}^s$. \square

4 Blow-up criterion

We can use the method introduced by Zhang and Yin[?] to obtain a blow-up criterion as follows.

Theorem 4.1. *Let $n_0 \in B_{p,r}^s$ with $1 \leq p, r \leq \infty$, and (s, p, r) satisfy condition (3.1) and let $T > 0$ be the maximal existence time of the corresponding solution n to (1.1). If $T < \infty$, then we have*

$$\int_0^T \|n(t)\|_{L^\infty} dt$$

Proof. For simplicity, consider the case $1 < p < \infty$. For all σ satisfies $0 < \sigma \leq s$, according to the product laws, since $\|v\|_{L^\infty} \leq C\|n\|_{L^\infty}$ and $\|v\|_{B_{p,r}^\sigma} \leq C\|n\|_{B_{p,r}^\sigma}$, we have

$$(4.1) \quad \|16\beta_0^2 vn - 4n^2 + 8\beta_0 v_x n\|_{B_{p,r}^\sigma} \leq C\|n\|_{L^\infty} \|n\|_{B_{p,r}^\sigma}$$

(1) $\sigma \in (0, 1)$.

Applying Lemma2.11 with $p_1 = \infty$, then

$$\|n(t)\|_{B_{p,r}^\sigma} \leq \|n_0\|_{B_{p,r}^\sigma} + C \int_0^t \|n(\tau)\|_{B_{p,r}^\sigma} \|n(\tau)\|_{L^\infty} d\tau + \int_0^t CV'_{p_1} \|n(\tau)\|_{B_{p,r}^\sigma} d\tau$$

with

$$V'_{p_1}(t) = \|4(v_x + 2\beta_0 v)_x\|_{B_{\infty,\infty}^0 \cap L^\infty} \leq C\|n\|_{L^\infty}$$

by Gronwall's inequality

$$\|n(t)\|_{B_{p,r}^\sigma} \leq \|n_0\|_{B_{p,r}^\sigma} \exp\left(C \int_0^t \|n(\tau)\|_{L^\infty} d\tau\right)$$

which implies

$$(4.2) \quad \int_0^T \|n(\tau)\|_{L^\infty} d\tau < \infty \implies \|n\|_{L_T^\infty(B_{p,r}^\sigma)} < \infty$$

(2) $\sigma = 1$

Applying Lemma2.11 with $p = p_1$, and

$$V'_{p_1} = \|4(v_x + 2\beta_0 v_x)\|_{B_{p,\infty}^{1/p} \cap L^\infty} \leq C(\|v\|_{L^\infty} + \|v\|_{B_{p,\infty}^{1/p+2}}) \leq C(\|n\|_{L^\infty} + \|n\|_{B_{p,r}^{1/p}})$$

then we have

$$\|n(t)\|_{B_{p,r}^1} \leq \|n_0\|_{B_{p,r}^1} + C \int_0^t \|n(\tau)\|_{B_{p,r}^1} \|n(\tau)\|_{L^\infty} d\tau + C \int_0^t (\|n(\tau)\|_{L^\infty} + \|n(\tau)\|_{B_{p,r}^{1/p}}) \|m(\tau)\|_{B_{p,r}^1} d\tau$$

by Gronwall's inequality

$$(4.3) \quad \|n(t)\|_{B_{p,r}^1} \leq C\|n_0\|_{B_{p,r}^1} \exp\left(C \int_0^t (\|n(\tau)\|_{L^\infty} + \|n(\tau)\|_{B_{p,r}^{1/p}}) d\tau\right)$$

which implies

$$(4.4) \quad \|n\|_{L_T^\infty(B_{p,r}^{1/p})} < \infty \quad \text{and} \quad \int_0^T \|n(\tau)\|_{L^\infty} d\tau < \infty \implies \|n\|_{L_T^\infty(B_{p,r}^\sigma)} < \infty$$

(3) $\sigma > 1$

According to (4.1) and Lemma2.11 with $p_1 = \infty$, and

$$V'_{p_1}(t) = \|4(v_x + 2\beta_0 v_x)\|_{B_{\infty,r}^{\sigma-1}} \leq C\|v\|_{B_{\infty,r}^{\sigma+1}} \leq C\|n\|_{B_{p,r}^{\sigma-1+1/p}}$$

then we have

$$\|n(t)\|_{B_{p,r}^{\sigma}} \leq \|n_0\|_{B_{p,r}^{\sigma}} + C \int_0^t \|n(\tau)\|_{B_{p,r}^{\sigma}} \|n(\tau)\|_{L^{\infty}} d\tau + C \int_0^t \|n(\tau)\|_{B_{p,r}^{\sigma-1+1/p}} \|n(\tau)\|_{B_{p,r}^{\sigma}} d\tau$$

by Gronwall's inequality

$$\|n(t)\|_{B_{p,r}^{\sigma}} \leq \|n_0\|_{B_{p,r}^{\sigma}} \exp\left(C \int_0^t (\|n(\tau)\|_{L^{\infty}} + \|n(\tau)\|_{B_{p,r}^{\sigma-1+1/p}}) d\tau\right)$$

which implies

$$\|n\|_{L_T^{\infty}(B_{p,r}^{\sigma-1+1/p})} < \infty \quad \text{and} \quad \int_0^T \|n(\tau)\|_{L^{\infty}} d\tau < \infty \implies \|n\|_{L_T^{\infty}(B_{p,r}^{\sigma})} < \infty$$

If $\sigma - 1 + \frac{1}{p} > 1$, we can repeat the above process. Indeed, since

$$\|4(v_x + 2\beta_0 v)_x\|_{B_{\infty,r}^{\sigma-1+1/p-1}} \leq C\|v\|_{B_{\infty,r}^{\sigma+1/p}} \leq C\|n\|_{B_{p,r}^{\sigma-2+2/p}}$$

we similar have

$$\|n\|_{L_T^{\infty}(B_{p,r}^{\sigma-2+1/p})} < \infty \quad \text{and} \quad \int_0^T \|n(\tau)\|_{L^{\infty}} d\tau < \infty \implies \|n\|_{L_T^{\infty}(B_{p,r}^{\sigma-1+1/p})} < \infty$$

By induction, we can choose $k \in \mathbb{N}^+$ such that $0 < \sigma - k(1 - \frac{1}{p}) \leq 1$, and $B_{p,r}^1 \hookrightarrow B_{p,r}^{\sigma-k(1-\frac{1}{p})}$, gives that

$$\|n\|_{L_T^{\infty}(B_{p,r}^1)} \leq \infty \quad \text{and} \quad \int_0^T \|n\|_{L^{\infty}} < \infty \implies \|n\|_{L_T^{\infty}(B_{p,r}^{\sigma-(k-1)(1-\frac{1}{p})})} < \infty \implies \dots \implies \|n\|_{L_T^{\infty}(B_{p,r}^{\sigma})} < \infty$$

Therefore we can conclude that

$$\|n\|_{L_T^{\infty}(B_{p,r}^1)} < \infty \quad \text{and} \quad \int_0^T \|n\|_{L^{\infty}} < \infty \implies \|n\|_{L_T^{\infty}(B_{p,r}^{\sigma})} < \infty$$

Return to $L_T^{\infty}(B_{p,r}^{\sigma})$ and suppose $\int_0^T \|n\|_{L^{\infty}} d\tau < \infty$, we have:

Case1: $s \geq 1$ By (1), we have $\|n\|_{L_T^{\infty}(B_{p,r}^s)} < \infty$.

Case2: $s=1$ By (1), we have $\|n\|_{L_T^{\infty}(B_{p,r}^{1/p})} < \infty$ and by (2) we get that $\|n\|_{L_T^{\infty}(B_{p,r}^1)} < \infty$.

Case3: $s \leq 1$ By (1), we have $\|n\|_{L_T^{\infty}(B_{p,r}^{1/p})} < \infty$ and by (2) we get that $\|n\|_{L_T^{\infty}(B_{p,r}^1)} < \infty$, lastly by the induction in (3), we can deduce that $\|n\|_{L_T^{\infty}(B_{p,r}^1)} < \infty$.

Therefore $\int_0^T \|n\|_{L^{\infty}} d\tau < \infty \implies \|n\|_{L^{\infty}(B_{p,r}^s)} < \infty$. Then by what we proved in Thm3.2, $\exists T_0$ small enough such that for some suitable $\varepsilon > 0$, $2C^2\|n(T-\varepsilon)\|_{B_{p,r}^s} T_0 < 1$, then follow then same line of Step1 in the proof of Thm3.2, we can extend the solution n on $[0, T) \times \mathbb{R}$ to \tilde{n} on $[0, T-\varepsilon+T_0) \times \mathbb{R}$ and by the uniqueness $n \equiv \tilde{n}$ for any $t \in [0, T]$. □

Corollary 4.2. *Let $n_0 \in B_{p,r}^s$ with (s, p, r) satisfying (3.1), $T > 0$ be the maximal existence time of the corresponding solution $n \in E_{p,r}^s(T_-)$ to system (1.1). Then n blows up in finite time if and only if*

$$\limsup_{t \uparrow T} \|n\|_{L^{\infty}} = \infty$$

5 Blow-up

As we have the blow-up criterion, consider the following initial value problem

$$(5.1) \quad \begin{cases} \psi_t(t, x) = -4(v_x + 2\beta_0 v)(t, \psi(t, x)), & t \in [0, T] \\ \psi(0, x) = x \end{cases}$$

then the system (1.1) in Lagrangian coordinates can be written as

$$(5.2) \quad \begin{cases} \partial_t n(t, \psi(t, x)) = (16\beta_0^2 v n - 4n^2 + 8\beta_0 v_x n)(t, \psi(t, x)) \\ n(0, \psi(0, x)) = n_0(x) \end{cases}$$

Theorem 5.1. *Let $n_0 \in B_{p,r}^s$ with (s, p, r) satisfying (3.1). There exists time $T_1 = \frac{1}{32 \max(\beta_0^2, \beta_0^{-2}) \|v_0\|_{W^{1,\infty}}}$, if $\exists x_0 \in \mathbb{R}$ such that*

$$(5.3) \quad n_0(x_0) < -\sqrt{\frac{b}{2}} + \frac{\sqrt{2b}}{1 - \exp(2\sqrt{2b}T_1)} \quad \text{with} \quad b = (16\beta_0^2 \|v_0\|_{L^\infty})^2 + (8\beta_0 \|v_{0x}\|_{L^\infty})^2$$

then n will blow up within T_1 .

More precisely, n will blow up at time $T_2 = \frac{1}{2\sqrt{2b}} \log\left(\frac{\sqrt{2n_0(x_0)} - \sqrt{b}}{\sqrt{2n_0(x_0)} + \sqrt{b}}\right)$.

Proof. Since v is the solution of system (1.2), and v_x satisfies

$$\begin{cases} \partial_t(v_x) - (4v_x + 8\beta_0 v)\partial_x(v_x) = 8\beta_0(v_x)^2 - 16\beta_0^2 v v_x + 8\beta_0 \partial_x P_1(D)(2\beta_0^2 v^2 + v_x^2) + 8\beta_0^2 P_1(D)(4\beta_0^2 v^2 - v_x^2) \\ v_x|_{t=0} = v_{0x} \end{cases}$$

then we have estimates

$$(5.4) \quad \|v\|_{W^{1,\infty}} \leq \|v_0\|_{W^{1,\infty}} + 8 \max(\beta_0^2, \beta_0^{-2}) \int_0^t \|v(\tau)\|_{W^{1,\infty}}^2 d\tau$$

Assume that $\|v(t)\|_{W^{1,\infty}} \leq 2\|v_0\|_{W^{1,\infty}}$ is satisfied for all $t \in [0, T_1]$, then by inequality (5.4), we have

$$\|v(t)\|_{W^{1,\infty}} \leq \|v_0\|_{W^{1,\infty}} + 32 \max(\beta_0^2, \beta_0^{-2}) t \|v_0\|_{W^{1,\infty}}^2 \leq 2\|v_0\|_{W^{1,\infty}}$$

Thus we have $\|v(t)\|_{W^{1,\infty}} \leq 2\|v_0\|_{W^{1,\infty}}$ for all $t \in [0, T_1]$.

Consider the system (5.2) for all $t \in [0, T_1]$, then it is obvious that for all $(t, x_0) \in [0, T_1] \times \mathbb{R}$

$$|16\beta_0^2 v \cdot n| \leq 32\beta_0^2 \|v_0\|_{L^\infty} |n| \leq (16\beta_0^2 \|v_0\|_{L^\infty})^2 + n^2$$

and

$$|8\beta_0 v_x \cdot n| \leq 16|\beta_0| \|v_{0x}\|_{L^\infty} |n| \leq (8\beta_0 \|v_{0x}\|_{L^\infty})^2 + n^2$$

denote $b = (16\beta_0^2 \|v_0\|_{L^\infty})^2 + (8\beta_0 \|v_{0x}\|_{L^\infty})^2$, then according to (5.2), we have

$$\partial_t n(t, \psi(t, x_0)) \leq -2n^2 + b$$

Consider the ordinary differential equation

$$f'(t) = -2f^2 + b, f(0) = n_0(x_0)$$

then we have

$$n(t, \psi(t, x_0)) \leq \int_0^t f'(\tau) d\tau + n_0(x_0) = \sqrt{\frac{b}{2}} \cdot \frac{1 + Ce^{-2\sqrt{2b}t}}{1 - Ce^{-2\sqrt{2b}t}} \quad \text{with} \quad C = \frac{\sqrt{2}n_0(x_0) - \sqrt{b}}{\sqrt{2}n_0(x_0) + \sqrt{b}}$$

It is obvious that $1 - Ce^{-2\sqrt{2b}t}$ is an increasing function, and by (5.3)

$$1 - C < 0 \quad \text{and} \quad 1 - Ce^{-2\sqrt{2b}T_1} > 0 \quad \text{and} \quad 1 - Ce^{-2\sqrt{2b}T_2} = 0$$

then we can easily get that $n(T_2, \psi(T_2, x_0)) = -\infty$ and $T_2 < T_1$.

Therefore n will blow up at time $T_2 = \frac{1}{2\sqrt{2b}} \log\left(\frac{\sqrt{2}n_0(x_0) - \sqrt{b}}{\sqrt{2}n_0(x_0) + \sqrt{b}}\right)$. \square

6 Existence of global solution

Similarly, we consider the initial value problem (5.1), applying classical results in the theory of ODEs, one can obtain the following results of ψ .

Lemma 6.1. *[7] Let $v_x + 2\beta_0 v \in C([0, T]; H^s(\mathbb{R})) \cap C^1([0, T]; H^{s-1}(\mathbb{R}))$, $s \geq 2$. Then the problem (5.1) has a unique solution $\psi \in C^1([0, T] \times \mathbb{R})$. Moreover, the map $\psi(t, \cdot)$ is an increasing diffeomorphism of \mathbb{R} with $\psi_x(0, x) = 1$ and*

$$\psi_x(t, x) = \exp\left(\int_0^t -4(v_x + 2\beta_0 v)_x(\tau, \psi(\tau, x)) d\tau\right) > 0, \quad \forall (t, x) \in [0, T] \times \mathbb{R}$$

We then have the following sign-preserved result.

Lemma 6.2. *Let $n_0 \in H^s(\mathbb{R})$, $s > \frac{1}{2}$ and $n_0(x_0) \geq 0$, $\forall x \in \mathbb{R}$. Assume T_{n_0} is the maximal existence time of the solution $n(t, x)$ to (1.1), then*

$$n(t, x) \geq 0, \quad \forall (t, x) \in [0, T_{n_0}] \times \mathbb{R}$$

Proof. From (1.1) we see that $\forall x \in \mathbb{R}$

$$\begin{aligned} \frac{d}{dt}(n(t, \psi(t, x)) \cdot \psi_x(t, x)) &= n_t(t, \psi(t, x)) \cdot \psi_x + n_x(t, \psi(t, x)) \cdot \psi_t \psi_x + n(t, \psi(t, x)) \cdot \psi_{tx} \\ &= \psi_x \cdot (n_t - 4(n(v_x + 2\beta_0 v))_x) = 0 \end{aligned}$$

solving the ODE, we have

$$n(t, \psi(t, x)) \cdot \psi_x(t, x) = n_0(x) \cdot \psi_x(0, x)$$

Noticing $n_0 \geq 0$ and $\psi_x > 0$, we get $n(t, \psi(t, x)) \geq 0$. Since $\psi(t, x)$ is a diffeomorphism on \mathbb{R} , we completes the proof. \square

Lemma 6.3. *Let $n_0 \in H^s(\mathbb{R})$, $s > \frac{1}{2}$ and $n_0(x) \geq 0$, $\forall x \in \mathbb{R}$. Assume $T_{n_0} > 0$ is the maximal existence time of the solution $n(t, x)$ to (1.1) with initial data n_0 . Then*

$$v(t, x) \geq 0 \quad \text{and} \quad |v_x(t, x)| \leq 2|\beta_0|v(t, x) \leq \sqrt{2}|\beta_0|\|v\|_{H^1}, \quad \forall (t, x) \in [0, T_{n_0}] \times \mathbb{R}$$

Proof. Since we have

$$v(t, x) = \frac{1}{4|\beta_0|} e^{-|2\beta_0 \cdot|} * n(t, x) \geq 0$$

and

$$|v_x(t, x)| = \left| \frac{1}{2} (-\text{sgn}(\cdot) e^{-|2\beta_0 \cdot|} * n)(t, x) \right| \leq 2|\beta_0| v(t, x)$$

On the other hand, for almost $\forall x \in \mathbb{R}$, we have

$$\begin{aligned} 2v^2(t, x) &= \int_{-\infty}^x \frac{d}{dy} (v^2(t, y)) dy - \int_x^{\infty} \frac{d}{dy} (v^2(t, y)) dy = \int_{-\infty}^x 2v \cdot v_y(t, y) dy - \int_x^{\infty} 2v \cdot v_y(t, y) dy \\ &\leq \int_{-\infty}^x v^2 + v_y^2 dy + \int_x^{\infty} v^2 + v_y^2 dy \leq \int_{-\infty}^{\infty} v^2 + v_y^2 dy = \|v(t, \cdot)\|_{H^1}^2 \end{aligned}$$

□

Lemma 6.4. *Let $n_0 \in H^s(\mathbb{R})$, $s > \frac{1}{2}$ and $n_0(x) \geq 0$, $\forall x \in \mathbb{R}$. Define an equivalent norm $H_{\beta_0}^1$ of H^1 -norm:*

$$\|v\|_{H_{\beta_0}^1} \triangleq \left(\int_{\mathbb{R}} (4\beta_0^2 + |\xi|^2) |\hat{v}(\xi)|^2 d\xi \right)^{1/2}$$

Then the $H_{\beta_0}^1$ -norm of v in $[0, T_{n_0})$ is non-increasing, namely, if $0 \leq t_1 \leq t_2 < T_{n_0}$

$$\|v(t_2)\|_{H_{\beta_0}^1} \leq \|v(t_1)\|_{H_{\beta_0}^1} \leq \|v_0\|_{H_{\beta_0}^1}$$

Proof. Since $n_0 \geq 0$, according to Lemma6.3, it follows that $v \geq 0$ and $|v_x(t, x)| \leq 2|\beta_0|v(t, x)$. Hence for any $t \in [0, T_{n_0})$, we have

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} 4\beta_0^2 v^2 + v_x^2 dx &= 2 \int_{\mathbb{R}} 4\beta_0^2 v \cdot v_t + v_x \cdot v_{xt} dx = 2 \int_{\mathbb{R}} v(4\beta_0^2 v_t - v_{xxt}) dx \\ &= 8 \int_{\mathbb{R}} v[n(v_x + 2\beta_0 v)]_x dx = -8 \int_{\mathbb{R}} v_x(4\beta_0^2 v - v_{xx})(v_x + 2\beta_0 v) dx \\ &= -8 \int_{\mathbb{R}} 4\beta_0^2 v \cdot v_x^2 + 8\beta_0^3 v^2 \cdot v_x - v_x^2 v_{xx} - 2\beta_0 v \cdot v_x \cdot v_{xx} dx \\ &= -8 \int_{\mathbb{R}} v_x^2 (4\beta_0^2 v + \beta_0 v_x) dx = -8 \int_{\mathbb{R}} v_x^2 (|\beta_0| + \frac{\beta_0}{2} \text{sgn}(\cdot)) e^{-|2\beta_0 \cdot|} * n dx \leq 0 \end{aligned}$$

Integrating the above inequality on $[0, t_1]$ and $[t_1, t_2]$ we have

$$\|v(t_2)\|_{H_{\beta_0}^1} \leq \|v(t_1)\|_{H_{\beta_0}^1} \leq \|v_0\|_{H_{\beta_0}^1}$$

□

Theorem 6.5. *Let $n_0 \in B_{2,2}^s$ with $s > 1/2$ and $n_0 \geq 0$, the system of (1.1) has a unique global solution, i.e. $\forall T > 0$, $n \in E_{2,2}^s(T)$.*

Proof. Let $\psi(t, x)$ satisfies the flow equation (5.1). Then by Lemma6.3 and Sobolev embedding

$$\frac{d}{dt} (n(t, \psi(t, x))) \leq (16\beta_0^2 v n - 4n^2 + 8\beta_0 v_x n)(t, \psi(t, x)) \leq 32\beta_0^2 v \cdot n(t, \psi(t, x)) \leq C \|v_0\|_{H^1} \cdot n(t, \psi(t, x))$$

by Gronwall's inequality we have

$$n(t, \psi(t, x)) \leq n_0(x) \exp(C \|v_0\|_{H^1})$$

thus $\forall T > 0$, we have

$$\int_0^T \|n(t, \cdot)\|_{L^\infty} dt \leq \int_0^T \|n_0\|_{L^\infty} \exp(C\|v_0\|_{H^1}) dt$$

Therefore by well-posedness and blow-up criterion of the system (1.1), we prove that

$$n \in E_{2,2}^s(T), \quad \forall T > 0$$

□

7 Simulation

According to our blow up condition, we can construct an exact example of the system (1.1).

Let $\beta_0 = 1$ and denote

$$f(x) = \begin{cases} \exp(-\frac{1}{1-x^2}) & |x| \leq 1 \\ 0 & |x| > 1 \end{cases}$$

taking $n_0(x) = -20e \cdot f(20x)$, it is obviously that $\|n_0\|_{L^\infty} = 20$. Since $\|n_0\|_{L^1} \leq \frac{e}{2}$, by Young's inequality we can get the rough estimate of v_0 and v_{0x}

$$\|v_0\|_{L^\infty} \leq \frac{e}{8}, \quad \|v_{0x}\|_{L^\infty} \leq \frac{e}{4}$$

then we can denote $T_1 = \frac{1}{12e}$ and $b = 8e^2$.

It is obvious that

$$-\sqrt{\frac{b}{2}} + \frac{\sqrt{2b}}{1 - \exp(2\sqrt{2b}T_1)} \geq -2e - \frac{4e}{1 - \exp(3/4)} \geq -16 \geq n_0(0) = -20$$

then we have n_0 will blow up within $T_1 \approx 0.0307$.

The solution of system (1.1) with the initial data n_0 can be roughly expressed as Figure 1:




Figure 1: blow up

It is consistent with the approximate numerical calculation of blow up condition we have given.

And for $n_0(x) = f(x)$, it is easy to check that $n_0 \in B_{2,2}^s$ for all $s > 1/2$ and $n_0(x) \geq 0$, the solution of system (1.1) with the initial data n_0 can be roughly expressed as Figure 2:




Figure 2: global solution

It is consistent with the existence of the global solution. And the above figure of the solution $v(t, x)$ is similar to the exact single peakon of the form[23]

$$v(t, x) = a_1 e^{-2\beta_0 |x + 8\beta_0 a_1 t - a_2|}$$




Figure 3: single peakon solution with $a_1 = a_2 = 1, \beta_0 = 1$

References

- [1] H. Bahouri, J.-Y. Chemin, and R. Danchin. *Fourier analysis and nonlinear partial differential equations*, volume 343 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles]*

-
- ples of Mathematical Sciences*]. Springer, Heidelberg, 2011.
- [2] R. Camassa and D. D. Holm. An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.*, 71(11):1661–1664, 1993.
- [3] A. Constantin. The Hamiltonian structure of the Camassa-Holm equation. *Exposition. Math.*, 15(1):53–85, 1997.
- [4] A. Constantin. On the scattering problem for the Camassa-Holm equation. *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.*, 457(2008):953–970, 2001.
- [5] A. Constantin and J. Escher. Global existence and blow-up for a shallow water equation. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 26(2):303–328, 1998.
- [6] A. Constantin and J. Escher. Well-posedness, global existence, and blowup phenomena for a periodic quasi-linear hyperbolic equation. *Comm. Pure Appl. Math.*, 51(5):475–504, 1998.
- [7] A. Constantin and R. I. Ivanov. On an integrable two-component Camassa-Holm shallow water system. *Phys. Lett. A*, 372(48):7129–7132, 2008.
- [8] A. Constantin and D. Lannes. The hydrodynamical relevance of the Camassa-Holm and Degasperis-Procesi equations. *Arch. Ration. Mech. Anal.*, 192(1):165–186, 2009.
- [9] A. Constantin and H. P. McKean. A shallow water equation on the circle. *Comm. Pure Appl. Math.*, 52(8):949–982, 1999.
- [10] R. Danchin. A few remarks on the Camassa-Holm equation. *Differential Integral Equations*, 14(8):953–988, 2001.
- [11] A. Degasperis, D. D. Holm, and A. N. W. Hone. A new integrable equation with peakon solutions. (2), 2002.
- [12] B. Fuchssteiner and A. S. Fokas. Symplectic structures, their Bäcklund transformations and hereditary symmetries. *Phys. D*, 4(1):47–66, 1981/82.
- [13] X. Geng and B. Xue. An extension of integrable peakon equations with cubic nonlinearity. *Nonlinearity*, 22(8):1847–1856, 2009.
- [14] X. Geng and B. Xue. A three-component generalization of Camassa-Holm equation with N -peakon solutions. *Adv. Math.*, 226(1):827–839, 2011.
- [15] X. Geng, B. Xue, and L. Wu. A super Camassa-Holm equation with N -peakon solutions. *Stud. Appl. Math.*, 130(1):1–16, 2013.
- [16] G. Gui and Y. Liu. On the Cauchy problem for the Degasperis-Procesi equation. *Quart. Appl. Math.*, 69(3):445–464, 2011.
- [17] H. He and Z. Yin. On a generalized Camassa-Holm equation with the flow generated by velocity and its gradient. *Appl. Anal.*, 96(4):679–701, 2017.
- [18] D. Ionescu-Kruse. Variational derivation of two-component Camassa-Holm shallow water system. *Appl. Anal.*, 92(6):1241–1253, 2013.

-
- [19] J. Li and Z. Yin. Well-posedness and global existence for a generalized Degasperis-Procesi equation. *Nonlinear Anal. Real World Appl.*, 28:72–90, 2016.
- [20] H. Lundmark and J. Szmigielski. Multi-peakon solutions of the degasperis-procesi equation. *Inverse Problems*, 19(6), 2005.
- [21] W. Luo and Z. Yin. Local well-posedness and blow-up criteria for a two-component Novikov system in the critical Besov space. *Nonlinear Anal.*, 122:1–22, 2015.
- [22] G. Rodríguez-Blanco. On the Cauchy problem for the Camassa-Holm equation. *Nonlinear Anal.*, 46(3):309–327, 2001.
- [23] B. Xue, H. Du, and X. Geng. New integrable peakon equation and its dynamic system. *Appl. Math. Lett.*, 146:Paper No. 108795, 7, 2023.
- [24] K. Yan and Z. Yin. On the Cauchy problem for a two-component Degasperis-Procesi system. *J. Differential Equations*, 252(3):2131–2159, 2012.
- [25] Z. Yin. Global existence for a new periodic integrable equation. *J. Math. Anal. Appl.*, 283(1):129–139, 2003.