Quantum origin of Ohm's reciprocity relation and its violation: conductivity as inverse resistivity

Giorgio Frangi¹ and Sašo Grozdanov^{1, 2}

¹ Higgs Centre for Theoretical Physics, University of Edinburgh, Edinburgh EH9 3FD, Scotland

²Faculty of Mathematics and Physics, University of Ljubljana, Jadranska ulica 19, SI-1000 Ljubljana, Slovenia

Conventional wisdom teaches us that the electrical conductivity in a material is the inverse of its resistivity. In this work, we show that when both of these transport coefficients are defined in linear response through the Kubo formulae as two-point correlators of quantum fields, this Ohm's reciprocity relation is generically violated in theories with dynamical electromagnetism. We also elucidate how in special limits (e.g., in the DC limit in the presence of thermal effects, in certain $2+1d$ conformal theories, and in holographic supersymmetric theories) the reciprocity is again reinstated, and show that if the response of a material is measured with respect to the total electric field that includes quantum corrections, then the reciprocity relation is satisfied by definition. However, in that case, the transport coefficients are given by the photon vacuum polarisation and not the correlators of conserved currents that dominate macroscopic late-time transport.

Introduction.—Ohm's law is a cornerstone of physics. It provides a simple framework to characterise materials that conduct electric currents. It is a phenomenological statement that stipulates that in a piece of a conductor, the voltage V is linearly related to the total current I flowing through the material: $V = RI$. The coefficient of proportionality R is the material's resistance. A more fine-grained way to express Ohm's law is in the local form:

$$
\mathbf{j}(\omega) = \sigma(\omega)\mathbf{E}(\omega),\tag{1}
$$

where $\mathbf{j}(\omega)$ is the current density, $\sigma(\omega)$ the AC frequencydependent electrical *conductivity* (matrix) and $\mathbf{E}(\omega)$ the electric field. To define the electrical resistivity, it is standard to invert Eq. (1) and write

$$
\mathbf{E}(\omega) = \rho(\omega)\mathbf{j}(\omega),\tag{2}
$$

where $\rho(\omega)$ is the electrical *resistivity* (matrix), simply defined as: $\rho(\omega) = \sigma^{-1}(\omega)$ (see Refs. [\[1,](#page-4-0) [2\]](#page-4-1)). In rotationally (and in $2+1d$ also parity) invariant systems, $\sigma_{ij} = \sigma \delta_{ij}$ and $\rho_{ij} = \rho \delta_{ij}$. Hence,

$$
\sigma(\omega)\rho(\omega) = 1. \tag{3}
$$

We will refer to Eq. (3) as *Ohm's reciprocity relation*.

The simplest microscopic model of a conductor is the well-known Drude model (see e.g. Ref. [\[2\]](#page-4-1)). Despite its qualitative and quantitative value, it is a phenomenological setup that cannot be extended to the breadth of conductors of interest. Instead, more powerful techniques are required, which must ultimately account for the effects of the underlying quantum field theory (QFT). This can be done by using linear response theory, which connects a microscopic calculation of correlators from either kinetic theory, perturbative QFT or holography to the appropriate low-energy effective theory, such as hydrodynamics (see Refs. $[3-8]$ $[3-8]$). The macroscopic quantities such as σ and ρ are then identified with the infra-red (IR) limit of QFT correlators. In the process, a system is perturbed away from equilibrium by coupling an external field to a QFT operator. In a conductor, an external

electric field \mathbf{E}_{ext} couples to the electric current, giving it a finite expectation value: $\langle \mathbf{j} \rangle \neq 0$. If the theory has dynamical electromagnetism, $\mathbf{E}_{\rm ext}$ couples to the dynamical electromagnetic field as well, giving $\langle \mathbf{E} \rangle \neq 0$. We write

$$
\langle \mathbf{j}(\omega) \rangle = \sigma(\omega) \mathbf{E}_{\text{ext}}(\omega), \quad \langle \mathbf{E}(\omega) \rangle = \rho(\omega) \mathbf{j}_{\text{ext}}(\omega), \quad (4)
$$

where j_{ext} is an external current related to E_{ext} via Maxwell's equations, and the following Kubo formulae define the conductive transport coefficients:

$$
\sigma(\omega) = \lim_{k \to 0} \frac{G_R^{jj}(\omega, k)}{i\omega}, \quad \rho(\omega) = \lim_{k \to 0} \frac{G_R^{EE}(\omega, k)}{i\omega}.
$$
 (5)

Here, $k \equiv |\mathbf{k}|$ is the magnitude of the wavevector and G_R denote the retarded two-point functions of j and E, chosen to be longitudinal (parallel to k). The Kubo formula for σ is extremely well known. The formula for ρ , however, was to the best of our knowledge first derived in the context of magnetohydrodynamics formulated with higher-form symmetries in Ref. $[9]$ (see also $[10-12]$ $[10-12]$).

An alternative definition of the conductivity that is sometimes used in the literature relates the expectation value of j and the total electromagnetic field $E_{\text{tot}} =$ $\mathbf{E}_{\text{ext}} + \langle \mathbf{E} \rangle$ consisting of the external source and the induced quantum field. In such case,

$$
\langle \mathbf{j}(\omega) \rangle = \tilde{\sigma}(\omega) \mathbf{E}_{\text{tot}}(\omega), \quad \mathbf{E}_{\text{tot}}(\omega) = \tilde{\rho}(\omega) \langle \mathbf{j}(\omega) \rangle. \tag{6}
$$

The appropriate analogue Kubo formula for $\tilde{\sigma}$ is not given in terms of the conserved $U(1)$ current correlator but the retarded vacuum polarisation Π_R (see e.g. Ref. [\[2,](#page-4-1) [7\]](#page-4-6)):

$$
\tilde{\sigma}(\omega) = \lim_{k \to 0} \frac{\Pi_R(\omega, k)}{i\omega} = \frac{1}{\tilde{\rho}(\omega)}.
$$
\n(7)

The validity of Eq. [\(3\)](#page-0-1) for $\tilde{\sigma}$ and $\tilde{\rho}$ follows from the definition [\(6\)](#page-0-2). The vacuum polarisation Π is the oneparticle-irreducible (1PI) part of the photon Feynman (time-ordered) correlator G_F^{AA} , with A^{μ} parallel to **k**. Π_R is found from Π after an analytic continuation.

It should be clear that, generically, $\sigma \neq \tilde{\sigma}$ and $\rho \neq \tilde{\rho}$ and that the use of either Eq. (4) or (6) must depend

FIG. 1. Diagrammatic expansion of G^{jj} in QED₄ to two loops. The diagrams in the first line contribute to the 1PI computation of Π. The 1PR diagram in a blue box does not.

on the context of the physical scenario that the theory is attempting to describe. While the (intrinsically) defined conductive transport coefficients in terms of Π_R can be computed in pertubative QFT, this is less clear nonperturbatively. Moreover, such coefficients are harder to discuss in an IR effective field theory context or an experimental setting, where it is the long-lived, conserved currents that play the central role in determining the dynamics of the low-energy degrees of freedom. Note that both expressions in Eq. (5) can be understood as twopoint correlators of conserved currents: the usual j^{μ} and the two-form current $J^{\mu\nu}$ related to **E** (see Ref. [\[9\]](#page-4-4)).

In this paper, we argue that, generically, Ohm's reciprocity relation [\(3\)](#page-0-1), as applied to the definitions using conserved currents in [\(5\)](#page-0-4), does not hold in theories with dynamical electromagnetism (dynamical photons):

$$
\sigma(\omega)\rho(\omega) \neq 1. \tag{8}
$$

Exceptions are theories in which at low energies, either photons cease to propagate, or another mechanism makes G^{jj} and Π indistinguishable. In fact, Ref. [\[9\]](#page-4-4) already argued for the inequality (8) . Here, we explore this claim in a more general and comprehensive manner to establish the difference between σ and ρ precisely and to demonstrate it with explicit examples. We begin by placing Eq. [\(8\)](#page-1-0) within the context of the theory of electrodynamics in our universe: quantum electrodynamics or QED4. We then state a set of special conditions under which Ohm's reciprocity [\(3\)](#page-0-1) is recovered and show examples of simpler QFTs in which the violation (or the absence of violation) of Ohm's reciprocity can be seen analytically.

General considerations and QED₄.—The Kubo formula [\(5\)](#page-0-4) for $\sigma(\omega)$ is usually derived in a setting with a global $U(1)$ symmetry (see Refs. [\[6,](#page-4-7) [7\]](#page-4-6)). The absence of dynamical photons (and of photon propagators in a diagrammatic expansion) then effectively identifies $G^{jj} =$ Π. Hence, $\sigma = \tilde{\sigma}$, and the resistivity can only be given by $\tilde{\rho}$ from Eq. [\(7\)](#page-0-5), which is by definition the inverse of $\tilde{\sigma}$.

Gauging the global $U(1)$ symmetry then introduces one-particle-reducible (1PR) diagrams from two loops on, distinguishing G^{jj} and Π . One may thus expect that, at least, $\sigma \neq \tilde{\sigma}$. However, as we argue below, this depends on whether the state has $T = 0$ or $T \neq 0$.

We start by considering a $T = 0$ state in QED₄ with a single massive Dirac fermion. The diagrams up to two

FIG. 2. Two types of diagrams contributing to the leadingorder perturbative expansion of G^{jj} at $T \neq 0$ [\[15\]](#page-5-1). Crossed ladder diagrams and mixed-type diagrams are subleading.

loops contributing to G^{jj} are depicted in Fig. [1](#page-1-1) and we show in [A](#page-4-8)ppendix A that, indeed, $G^{jj} \neq \Pi$. While σ and ρ , as defined in [\(5\)](#page-0-4), have little physical meaning in 4d at $T = 0$, it is nevertheless instructive to formally compute $\sigma \rho$ and investigate Eq. [\(8\)](#page-1-0). First, we express $G^{E\overline{E}}$ in terms of G^{AA} to write the Kubo formula [\(5\)](#page-0-4) for ρ as

$$
\rho(\omega) = \lim_{k \to 0} i\omega \left(G_R^{AA}(\omega, k) - \Box^{-1}(\omega, k) \right), \tag{9}
$$

where \square is the classical photon kinetic operator. For purposes of perturbative calculations, we can use $\Box^{-1}(\omega, 0) = G_{\text{free}}^{A\overline{A}}(\omega, 0) = 1/\omega^2$. We then observe that G_F^{jj} is the bubble containing all loop corrections to the photon propagator, resumming diagrams beyond Fig. [1.](#page-1-1) It can be obtained by removing its free part and amputating the remaining external legs (see Ref. [\[13\]](#page-5-2)):

$$
G_F^{jj} = \Box \cdot (G_F^{AA} - G_{\text{free}}^{AA}) \cdot \Box, \tag{10}
$$

where the arguments (ω, k) have been removed for brevity. Analytically continuing [\(10\)](#page-1-2) to G_R^{jj} and plug-ging it into [\(4\)](#page-0-3) gives an alternative expression for $\sigma(\omega)$, which allows us to formally express the reciprocity relation as

$$
\sigma(\omega)\rho(\omega) = \lim_{k \to 0} \left(\frac{\pi(\omega, k)}{\pi(\omega, k) - 1}\right)^2, \quad (11)
$$

where π is the scalar dimensionless vacuum polarisation that enters G_F^{AA} . Eq. [\(11\)](#page-1-3) therefore shows that, generically, Ohm's reciprocity relation is violated at $T = 0$, confirming Eq. [\(8\)](#page-1-0). To one loop in $\alpha = e^2/4\pi$, $\pi(\omega, k =$ $(0) = \alpha \omega^2 / 15 \pi m^2 + O(\omega^4)$ [\[14\]](#page-5-3) and therefore

$$
\sigma(\omega)\rho(\omega) = \frac{\alpha^2 \omega^4}{225\pi^2 m^4} + O(\omega^6). \tag{12}
$$

Eq. [\(11\)](#page-1-3) also holds at $T \neq 0$, but for finite (non-zero) ω, with π computed e.g. in Ref. [\[7\]](#page-4-6). In the DC limit, however, the situation is dramatically different because the loop expansion of Fig. [1](#page-1-1) breaks down due to the appearance of pinching-pole singularities in thermal propagators that enhance higher-loop diagrams $[15]$. In QED_4 , beyond the simple fermion loop, two types of diagrams contribute to G^{jj} . They are shown in Fig. [2.](#page-1-4) One is the 1PR diagram relevant at $T = 0$ (cf. Fig. [1\)](#page-1-1). The second represents an infinite family of 1PI ladder diagrams that require a resummation. If the ladder diagrams dominate the DC limit, then the 1PR part of G^{jj} is effectively suppressed, and one recovers $\sigma = \tilde{\sigma}$. From the viewpoint

of the $2 \rightarrow 2$ scattering, this corresponds to a regime in which the t - and u -channel photon exchanges between on-shell fermions dominate over the s-channel exchanges, which is consistent with the effective kinetic theory of [\[16\]](#page-5-4), at least at the order of leading logarithms. On the other hand, the thermal dressing of photons implies that, due to the definition of Π , $G^{AA} \simeq \Pi^{-1}$ in the DC limit. As a consequence, the thermal DC limit reinstates the reciprocity relation [\(3\)](#page-0-1), at least up to next-to-leading-log corrections. We note that, qualitatively, the same result is obtained by simply using the small- ω limit of the oneloop, hard thermal loop expression for π given in [\[17\]](#page-5-5). Using $\pi(\omega, k = 0) = 4\pi\alpha T^2/9\omega^2 + O(\omega^0)$, one finds

$$
\sigma(\omega)\rho(\omega) = 1 + \frac{9\omega^2}{2\pi\alpha T^2} + O(\omega^4). \tag{13}
$$

While informative, this is a heuristic result as π used here is computed outside the strict hydrodynamic regime.

In summary, while, in general, Ohm's reciprocity is violated (Eq. (8)) for AC transport in QED₄, thermal effects can reestablish it in the DC limit. Importantly, it is also clear that, generically, $\sigma \neq \tilde{\sigma}$, $\rho \neq \tilde{\rho}$, and that context needs to be provided to be able to choose the physically relevant definition. It should, however, be born in mind that the analysis presented here is perturbative and therefore our claims may not apply to the non-perturbative regime studied in Ref. [\[18\]](#page-5-6), which found that $\sigma \rho \neq 1$ even in the DC limit of scalar QED₄.

Suppression of dynamical photons in the hydrodynamic regime can occur via a number of mechanisms, either only for DC transport or even for AC transport. Bellow, we study two simpler classes of QFTs to further elucidate the conditions needed to reinstate Eq. [\(3\)](#page-0-1):

• 3d CFTs in which the kinetic photon terms are irrelevant in the IR. Such theories typically exhibit a particle-vortex duality and Ohm's reciprocity [\(3\)](#page-0-1) even at $T = 0$. Fascinatingly, however, special 3d CFTs exist that evade these conditions and violate the reciprocity relation.

 \bullet Holographic supersymmetric large- N QFTs with a one-loop exact electromagnetic beta function.

Three-dimensional CFTs and particle-vortex duality.—Three-dimensional QFTs frequently possess more symmetries than those in 4d and exhibit dualities, such as the prominent particle-vortex duality that was long known in bosonic theories and has recently been extended to theories with fermions $[19-22]$ $[19-22]$. Many 3d theories also exhibit fixed points and can be understood as CFTs. In fact, it was argued in Ref. [\[23\]](#page-5-9) that a generic CFT_3 with a global $U(1)$ current has a particle-vortex dual (an S-dual) theory (see also [\[4\]](#page-4-9)).

We begin our analysis of such theories by studying the so-called $QED_{3,4}$ (see Refs. [\[20,](#page-5-10) [24–](#page-5-11)[26\]](#page-5-12)), which is a theory of a Dirac fermion Ψ (one can also consider a theory with N flavours) confined to a 3d brane that interacts with a photon propagating in 4d half-space:

$$
S_{\Psi} = \int d^3x \, i\bar{\Psi}\gamma^b \left(\partial_b - iA_b\right) \Psi - \frac{1}{4e^2} \int d^4x \, F_{\mu\nu} F^{\mu\nu}.\tag{14}
$$

At weak coupling e, the theory is scale invariant. At strong coupling, it is known to be scale invariant for large $N > N_{\text{crit}}$. For small N, the theory may undergo spontaneous chiral symmetry breaking and generate a fermion mass [\[24\]](#page-5-11), thereby remaining scale invariant only for some $e < e(N_{\text{crit}})$. Whether a genuine CFT₃ for $N = 1$ or not, $QED_{3,4}$ in (14) is conjectured to be particle-vortex dual, at least in the IR, to a theory of a neutral composite Dirac fermion ψ coupled to an emergent gauge field a_b :

$$
\bar{S}_{\psi} = \int d^3x \left[i \bar{\psi} \gamma^b \left(\partial_b - i a_b \right) \psi - \frac{1}{4\pi} \varepsilon^{abc} A_a \partial_b a_c \right] - \frac{1}{4e^2} \int d^4x F_{\mu\nu} F^{\mu\nu}.
$$
\n(15)

The weak-strong duality relates the fermionic $U(1)$ currents to the topological currents (see Ref. [\[26\]](#page-5-12)):

$$
j^{\mu}_{\Psi} = \bar{\Psi}\gamma^{\mu}\Psi \leftrightarrow -\frac{1}{4\pi}\varepsilon^{\mu\nu\lambda}\partial_{\nu}a_{\lambda},\tag{16}
$$

$$
j^{\mu}_{\psi} = \bar{\psi}\gamma^{\mu}\psi \leftrightarrow \frac{1}{4\pi} \varepsilon^{\mu\nu\lambda} \partial_{\nu} A_{\lambda}, \tag{17}
$$

or, in terms of spatial components of currents and electric fields in two dual theories (E^j is defined in terms of A^{μ} and e^j in terms of a^{μ}),

$$
j^i_{\Psi} \leftrightarrow -\frac{1}{4\pi} \varepsilon^{ij} e_j, \qquad j^i_{\psi} \leftrightarrow \frac{1}{4\pi} \varepsilon^{ij} E_j, \tag{18}
$$

with ε^{ij} the 2d Levi-Civita symbol. The couplings are dualised as $e \leftrightarrow \overline{e} = 8\pi/e$. All quantities computed in \bar{S}_{ψ} will be given overhead bars. In each of the theories, we now assume (anisotropic) Ohm's law: $j_{\Psi}^i = \sigma^{ij} E_j$ and $j_{\psi}^{i} = \bar{\sigma}^{ij} e_j$. Using the duality [\(18\)](#page-2-1), it then follows that

$$
\sigma^{il} = -\frac{1}{(4\pi)^2} \varepsilon^{ij} \bar{\sigma}_{jk}^{-1} \varepsilon^{kl}.
$$
 (19)

Moreover, the correlators that enter the Kubo formulae [\(5\)](#page-0-4) (for any type of G^{jj} and G^{EE}) are dualised as

$$
\langle j^i_{\Psi}, j^j_{\Psi} \rangle \leftrightarrow \frac{1}{(4\pi)^2} \varepsilon^{ik} \varepsilon^{jl} \langle e_k, e_l \rangle, \tag{20}
$$

$$
\langle j_{\psi}^{i}, j_{\psi}^{j} \rangle \leftrightarrow \frac{1}{(4\pi)^{2}} \varepsilon^{ik} \varepsilon^{jl} \langle E_{k}, E_{l} \rangle, \tag{21}
$$

which relates $\sigma(e)$ to $\bar{\rho}(\bar{e})$ and $\sigma(\bar{e})$ to $\rho(e)$ as

$$
\sigma^{ij}(e) = \frac{1}{(4\pi)^2} \varepsilon^{ik} \varepsilon^{jl} \bar{\rho}_{kl}(\bar{e}), \qquad (22)
$$

$$
\bar{\sigma}^{ij}(\bar{e}) = \frac{1}{(4\pi)^2} \varepsilon^{ik} \varepsilon^{jl} \rho_{kl}(e). \tag{23}
$$

Thus, we recover Ohm's reciprocity relations [\(3\)](#page-0-1) (in matrix form) in both dual theories: $\sigma \cdot \rho = 1$ and $\bar{\sigma} \cdot \bar{\rho} = 1$, at finite temperature T, and for all ω and couplings e.

We continue with a discussion of general 3d CFTs at zero temperature. In such theories, \tilde{G}_R^{jj} is constrained by the relativistic symmetries to have the form

$$
G_R^{j^{\mu}j^{\nu}}(p) = \sqrt{p^2} \left(\eta^{\mu\nu} - \frac{p^{\mu}p^{\nu}}{p^2} \right) C_j, \tag{24}
$$

where p^{μ} is the relativistic three-momentum and C_j is momentum-independent constant. If the $CFT₃$ has a number of conserved (non-commuting) currents j_I^{μ} labeled by I, then we have a tensor C_j^{IJ} of constants. The form [\(24\)](#page-2-2) is sufficient to determine the conductivity at $T = 0$, and, equivalently, in the $\omega \gg T$ limit of the AC conductivity in a thermal state (see Refs. [\[4,](#page-4-9) [27](#page-5-13)[–29\]](#page-5-14)):

$$
\sigma(\omega \gg T) \approx \sigma(T = 0) = C_j. \tag{25}
$$

The resistivity [\(5\)](#page-0-4) can be computed from the photon correlator $G_R^{\tilde{A}^{\mu} A^{\nu}}$ (cf. Eq. [\(9\)](#page-1-5)) that has the same tensor structure as Eq. (24) (in the appropriate gauge) but a generically independent constant C_A . We find

$$
\rho(\omega \gg T) \approx \rho(T = 0) = C_A. \tag{26}
$$

Evaluation of the violation of Ohm's reciprocity [\(8\)](#page-1-0) is therefore reduced to computing C_j and C_A , which can be done diagrammatically. This is, unsurprisingly, simplified in CFTs with a large number of flavours N . In the rest of this section, we will focus only on such theories.

We first consider QED₃ with N flavours of fermions ψ_I at $T = 0$ and in the $1/N$ expansion (see Ref. [\[30\]](#page-5-15)):

$$
S = \int d^3x \left[i \sum_{I=1}^{N} \bar{\psi}_I \gamma^{\mu} \left(\partial_{\mu} - i A_{\mu} \right) \Psi_I - \frac{1}{4e^2} F_{\mu\nu} F^{\mu\nu} \right].
$$
\n(27)

In this theory, the gauged 'electric' current that couples to A_{μ} is $j^{\mu} = \sum_{I=1}^{N} \bar{\psi}_{I} \gamma^{\mu} \psi_{I}$, which is a commuting 'Casimir operator'. For a detailed analysis of all relevant aspects of this theory, see Ref. [\[31\]](#page-5-16). Importantly, after one integrates out the fermions, it can be seen that the photon propagator is an irrelevant operator, so that $G^{AA} = \Pi$. Moreover, in this theory, A^{μ} and j^{μ} are related by a Legendre transform [\[23,](#page-5-9) [32\]](#page-5-17). The computation of C_j and C_A was also performed in [\[31\]](#page-5-16), giving

$$
\sigma = C_j = \frac{N}{16} \left[1 + \frac{1}{N} \left(\frac{736}{9\pi^2} - 8 \right) + O(1/N^2) \right], \quad (28)
$$

$$
\rho = C_A = \frac{16}{N} \left[1 - \frac{1}{N} \left(\frac{736}{9\pi^2} - 8 \right) + O(1/N^2) \right].
$$
 (29)

Hence, as expected in such 3d CFTs, $\sigma \rho = 1$.

Finally, we turn our attention to a very special but instructive CFT_3 : the $\mathbb{C}P^{N-1}$ model with the action

$$
S = \frac{1}{e^2} \int d^3x \left| (\partial_\mu - iA_\mu) \varphi \right|^2, \tag{30}
$$

where the N complex scalars φ_I satisfy the constraint $|\varphi|^2 = \sum_{I=1}^{N} |\varphi_I|^2 = 1$. For details, see Refs. [\[33,](#page-5-18) [34\]](#page-5-19). Remarkably, its unique properties allow us to evade the above-discussed, 'generally expected' properties of a 3d CFT with dynamical electromagnetism. The reason is that the action (30) has no kinetic term for the photon. Nevertheless, such a term is dynamically generated by quantum corrections in the $1/N$ expansion. Crucially, it is now not irrelevant. The QFT therefore has

dynamical photons in the IR that gauge the current $j^{\mu} = -i \sum_{I=1}^{N} \left[\varphi_{I}^{\dagger} D_{\mu} \varphi_{I} - \varphi_{I} (D_{\mu} \varphi^{I})^{\dagger} \right]$ (see Ref. [\[29\]](#page-5-14)). The theory at $\dot{T} = 0$ offers an explicit example of the Ohm's reciprocity relation violation, i.e., of Eq. [\(8\)](#page-1-0). This again follows from the coefficients C_j and C_A that were computed in Ref. [\[29\]](#page-5-14), giving us

$$
\sigma = C_j = \frac{N}{16} \left[1 - \frac{2.74}{N} + O(1/N^2) \right],\tag{31}
$$

$$
\rho = C_A = \frac{16}{N} \left[1 + \frac{0.578}{N} + O(1/N^2) \right].
$$
 (32)

The leading term still preserves $\sigma \rho = 1$. However, the $1/N$ corrections, which introduce dynamical photons, violate the reciprocity relation:

$$
\sigma \rho = 1 - \frac{2.16}{N} + O(1/N^2). \tag{33}
$$

As in QED4, this example elucidates the essential role that dynamical IR photons play in the violation of Ohm's reciprocity relation [\(8\)](#page-1-0). It is also important to note that since this is a $T = 0$ calculation, no screening protects Ohm's reciprocity even for DC transport coefficients.

Supersymmetric field theories and holographic duality.—Holographic duality is a powerful tool for computing correlation functions in certain large- N QFTs [\[4,](#page-4-9) [5\]](#page-4-10). Consider the $\mathcal{N} = 4$ supersymmetric Yang-Mills theory with $N \to \infty$ number of colours and an infinite 't Hooft coupling. We are interested in a scenario in which a current j^{μ} that corresponds to a $U(1)$ subgroup of the R-symmetry is gauged so that the QFT has dynamical electromagnetism with a one-loop exact beta function of the running electromagnetic coupling and a UV Landau pole. For details, see Refs. [\[35](#page-5-20)[–37\]](#page-5-21). Correlators of j (giving σ) and **E** (giving ρ) can be computed by using two different 5d bulk actions with mixed boundary conditions:

$$
S_{\sigma} = \int d^5 x \sqrt{-g} \left[R - 2\Lambda - \frac{1}{4e^2} F_{\mu\nu} F^{\mu\nu} \right],\tag{34}
$$

$$
S_{\rho} = \int d^5 x \sqrt{-g} \left[R - 2\Lambda - \frac{1}{12e^2} H_{\mu\nu\rho} H^{\mu\nu\rho} \right], \quad (35)
$$

where $\Lambda = -6/L^2$. L is the AdS scale, which we set to one. Eq. [\(34\)](#page-3-1) is the 5d Einstein-Maxwell theory and Eq. [\(35\)](#page-3-2) the theory of a two-form field $B_{\mu\nu}$ with a threeform field strength $H = dB$ developed in [\[36,](#page-5-22) [37\]](#page-5-21) (see also [\[38,](#page-5-23) [39\]](#page-5-24)) in constructing a holographic dual to a theory with a one-form symmetry and magnetohydrodynamics with dynamical electromagnetism. The actions [\(34\)](#page-3-1) and [\(35\)](#page-3-2) are related by the 5d Hodge dualisation $H = \ast_5 F$.

We assume electric charge neutrality and neglect the energy-momentum fluctuations (the probe limit). The relevant fluctuations of the bulk fields in the background of the Schwarzschild black brane in the two cases are $\delta A = \delta A(u) e^{-i\omega t} dx$ and $\delta B = \delta B(u) e^{-i\omega t} dy \wedge dz$, where u is the radial Fefferman-Graham (FG) coordinate (see Refs. [\[40,](#page-5-25) [41\]](#page-5-26)). Following the procedure of Ref. [\[36\]](#page-5-22) to evaluate the Kubo formulae [\(5\)](#page-0-4) then gives

$$
\sigma(\omega) = \frac{2\delta A^{(2)}}{i\omega \delta A^{(0)}} + i\omega \left(\ln \bar{u} + \frac{1}{2}\right),\tag{36}
$$

$$
\rho(\omega) = \frac{\delta B^{(1)}}{i\omega \left(\delta B^{(0)} + \delta B^{(1)} \ln \bar{u}\right)},\tag{37}
$$

where $\delta A^{(i)}$ and $\delta B^{(i)}$ are the *i*-th order coefficients in the FG expansion, and \bar{u} is the radial position of a finite cutoff brane introduced to regularise the theory. As a result of the dualisation $H = *_{5}F$, we have $\omega \delta A^{(0)} = i \delta B^{(1)}$ and $2\delta A^{(2)} = i\omega(\delta B^{(0)} - \delta B^{(1)}/2)$. Inserting these relations into Eqs. [\(36\)](#page-4-11) and [\(37\)](#page-4-12) then immediately establishes that Ohm's reciprocity relation [\(3\)](#page-0-1) is retained for the AC transport coefficients: $\sigma(\omega)\rho(\omega) = 1$. In fact, Eq. [\(3\)](#page-0-1) holds in a large class of holographic models, discussed in detail in Ref. $[42]$. The large-N and supersymmetric nature of this holographic QFT therefore ensures that Ohm's reciprocity [\(3\)](#page-0-1) is preserved for all frequencies.

Discussion.—Our work shows that one should approach conventional wisdom with care. In particular, when it comes to the measurement of electric response, one needs to carefully and precisely specify the nature of 'external' perturbations, as per Eqs. (4) and (6) . When this is done in the usual sense of Eq. (4) , then one arrives at the conclusion that Ohm's reciprocity is generically violated $(Eq. (8))$ $(Eq. (8))$ $(Eq. (8))$, except in special circumstances when the dynamics of electromagnetism is effectively suppressed, most notably in the perturbative DC limit with thermal effects. Phenomenologically, Eq. [\(8\)](#page-1-0) is important as it refers to transport given in terms of the IR conserved operators that dominate late-time dynamics, and not in terms of the 1PI vacuum polarisation of the photons. We believe it should be possible to devise simpler (e.g., nonrelativistic lattice) models to show this behaviour and potentially even measure the differences between AC σ and $1/\rho$ in experiments. Theories that will exhibit DC violation seem harder to access in real world at least in states with hydrodynamic behaviour. This is because the DC relation $\sigma = 1/\rho$ appears naturally in the derivation of 5

magnetohydrodynamics $[9-11]$ $[9-11]$. Finally, to develop better understanding of the differences between conductive and resistive transport in QFTs beyond linear response theory, it should be most transparent to further develop and employ the Schwinger-Keldysh effective field theory methods $[43-45]$ $[43-45]$, already used in QED₄ in Ref. [\[46\]](#page-5-31).

Acknowledgements.—We would like to thank Arpit Das, Sean Hartnoll, Nabil Iqbal, Janos Polonyi, Nick Poovuttikul, Tomaž Prosen, Paul Romatschke, Alexander Soloviev and Mile Vrbica for illuminating discussions on related topics. The work of G.F. is supported by an Edinburgh Doctoral College Scholarship (ECDS). The work of S.G. was supported by the STFC Ernest Rutherford Fellowship ST/T00388X/1. The work is also supported by the research programme P1-0402 and the project N1-0245 of Slovenian Research Agency (ARIS).

Appendix A: QED⁴ at zero temperature

In this appendix, we briefly review some relevant results in QED_4 computed to two loops at $T = 0$ (see Ref. [\[14\]](#page-5-3)). In particular, the current correlator $G_F^{j\bar{j}}$ is given in the low frequency limit and at $k = 0$ by

$$
G_F^{jj}(\omega) = \omega^2 \left[\left(\frac{\alpha}{15\pi} + \frac{41\alpha^2}{162\pi^2} \right) \frac{\omega^2}{m^2} + \left(\frac{\alpha}{140\pi} + \frac{497\alpha^2}{10800\pi^2} \right) \frac{\omega^4}{m^4} \right] + O\left(\alpha^3, \frac{\omega^6}{m^6}\right), \quad \text{(A1)}
$$

while the 1PI vacuum polarisation is given by

$$
\Pi(\omega) = \omega^2 \left[\left(\frac{\alpha}{15\pi} + \frac{41\alpha^2}{162\pi^2} \right) \frac{\omega^2}{m^2} + \left(\frac{\alpha}{140\pi} + \frac{449\alpha^2}{10800\pi^2} \right) \frac{\omega^4}{m^4} \right] + O\left(\alpha^3, \frac{\omega^6}{m^6}\right). \tag{A2}
$$

The difference between Eqs. $(A1)$ and $(A2)$ can be traced back precisely to the 1PR diagram (shaded in blue) in Fig. [1.](#page-1-1) These results are sufficient to formally compute σ , $\tilde{\sigma}$, ρ , $\tilde{\rho}$, and, thereby, show that $\sigma \rho \neq 1$, as stated in Eq. (12) .

- [1] C. Kittel, *Introduction to Solid State Physics* (Wiley, 2004).
- [2] G. D. Mahan, Many Particle Physics, Third Edition (Plenum, New York, 2000).
- [3] P. Kovtun, Lectures on hydrodynamic fluctuations in relativistic theories, J. Phys. A 45[, 473001 \(2012\),](https://doi.org/10.1088/1751-8113/45/47/473001) [arXiv:1205.5040 \[hep-th\].](https://arxiv.org/abs/1205.5040)
- [4] S. Hartnoll, A. Lucas, and S. Sachdev, *Holographic Quan*[tum Matter](https://books.google.si/books?id=DW9RDwAAQBAJ), The MIT Press (MIT Press, 2018).
- [5] J. Zaanen, Y. Liu, Y. Sun, and K. Schalm, [Holographic](https://books.google.nl/books?id=vNfECgAAQBAJ) [Duality in Condensed Matter Physics](https://books.google.nl/books?id=vNfECgAAQBAJ) (Cambridge University Press, 2015).
- [6] J. I. Kapusta and C. Gale, [Finite-temperature field](https://doi.org/10.1017/CBO9780511535130) [theory: Principles and applications](https://doi.org/10.1017/CBO9780511535130), Cambridge Mono-

graphs on Mathematical Physics (Cambridge University Press, 2011).

- [7] M. L. Bellac, *[Thermal Field Theory](https://doi.org/10.1017/CBO9780511721700)*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2011).
- [8] P. M. Chaikin and T. C. Lubensky, Principles of Condensed Matter Physics (Cambridge University Press, Cambridge, 1995).
- [9] S. Grozdanov, D. M. Hofman, and N. Iqbal, Generalized global symmetries and dissipative magnetohydrodynamics, Phys. Rev. D 95[, 096003 \(2017\),](https://doi.org/10.1103/PhysRevD.95.096003) [arXiv:1610.07392](https://arxiv.org/abs/1610.07392) [\[hep-th\].](https://arxiv.org/abs/1610.07392)
- [10] J. Hernandez and P. Kovtun, Relativistic magnetohydrodynamics, [JHEP](https://doi.org/10.1007/JHEP05(2017)001) 05, 001, [arXiv:1703.08757 \[hep-th\].](https://arxiv.org/abs/1703.08757)
- [11] J. Armas and A. Jain, Magnetohydrodynamics as superfluidity, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.122.141603) 122, 141603 (2019), [arXiv:1808.01939 \[hep-th\].](https://arxiv.org/abs/1808.01939)
- [12] S. Vardhan, S. Grozdanov, S. Leutheusser, and H. Liu, A new formulation of strong-field magnetohydrodynamics for neutron stars, (2022), [arXiv:2207.01636 \[astro](https://arxiv.org/abs/2207.01636)[ph.HE\].](https://arxiv.org/abs/2207.01636)
- [13] S. Weinberg, *[The Quantum theory of fields. Vol. 1: Foun](https://doi.org/10.1017/CBO9781139644167)*[dations](https://doi.org/10.1017/CBO9781139644167) (Cambridge University Press, 2005).
- [14] S. Laporta and U. D. Jentschura, Dimensional regularization and two-loop vacuum polarization operator: Master integrals, analytic results and energy shifts (2024), [arXiv:2403.07127 \[hep-ph\].](https://arxiv.org/abs/2403.07127)
- [15] S. Jeon, Hydrodynamic transport coefficients in relativistic scalar field theory, Phys. Rev. D 52[, 3591 \(1995\),](https://doi.org/10.1103/PhysRevD.52.3591) [arXiv:hep-ph/9409250.](https://arxiv.org/abs/hep-ph/9409250)
- [16] P. Arnold, G. D. Moore, and L. G. Yaffe, Transport coefficients in high temperature gauge theories (i): leadinglog results, [Journal of High Energy Physics](https://doi.org/10.1088/1126-6708/2000/11/001) 2000, 001 $(2000).$
- [17] H. A. Weldon, Covariant calculations at finite temperature: The relativistic plasma, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.26.1394) 26, 1394 [\(1982\).](https://doi.org/10.1103/PhysRevD.26.1394)
- [18] A. Das, A. Florio, N. Iqbal, and N. Poovuttikul, Higherform symmetry and chiral transport in real-time lattice $u(1)$ gauge theory (2023), $a\overline{x}$ Xiv:2309.14438 [hep-th].
- [19] M. A. Metlitski and A. Vishwanath, Particle-vortex duality of two-dimensional Dirac fermion from electricmagnetic duality of three-dimensional topological insulators, Phys. Rev. B 93[, 245151 \(2016\),](https://doi.org/10.1103/PhysRevB.93.245151) [arXiv:1505.05142](https://arxiv.org/abs/1505.05142) [\[cond-mat.str-el\].](https://arxiv.org/abs/1505.05142)
- [20] D. T. Son, Is the Composite Fermion a Dirac Particle?. Phys. Rev. X 5[, 031027 \(2015\),](https://doi.org/10.1103/PhysRevX.5.031027) [arXiv:1502.03446 \[cond](https://arxiv.org/abs/1502.03446)[mat.mes-hall\].](https://arxiv.org/abs/1502.03446)
- [21] N. Seiberg, T. Senthil, C. Wang, and E. Witten, A Duality Web in 2+1 Dimensions and Condensed Matter Physics, [Annals Phys.](https://doi.org/10.1016/j.aop.2016.08.007) 374, 395 (2016), [arXiv:1606.01989](https://arxiv.org/abs/1606.01989) [\[hep-th\].](https://arxiv.org/abs/1606.01989)
- [22] A. Karch and D. Tong, Particle-Vortex Duality from 3d Bosonization, Phys. Rev. X 6[, 031043 \(2016\),](https://doi.org/10.1103/PhysRevX.6.031043) [arXiv:1606.01893 \[hep-th\].](https://arxiv.org/abs/1606.01893)
- [23] E. Witten, $SL(2,Z)$ action on three-dimensional conformal field theories with Abelian symmetry, in From Fields to Strings: Circumnavigating Theoretical Physics: A Conference in Tribute to Ian Kogan (2003) pp. 1173– 1200, [arXiv:hep-th/0307041.](https://arxiv.org/abs/hep-th/0307041)
- [24] E. V. Gorbar, V. P. Gusynin, and V. A. Miransky, Dynamical chiral symmetry breaking on a brane in reduced QED, Phys. Rev. D 64[, 105028 \(2001\),](https://doi.org/10.1103/PhysRevD.64.105028) [arXiv:hep](https://arxiv.org/abs/hep-ph/0105059)[ph/0105059.](https://arxiv.org/abs/hep-ph/0105059)
- [25] S. Teber, Electromagnetic current correlations in reduced quantum electrodynamics, Physical Review D 86, [10.1103/physrevd.86.025005](https://doi.org/10.1103/physrevd.86.025005) (2012).
- [26] W.-H. Hsiao and D. T. Son, Duality and universal transport in mixed-dimension electrodynamics, [Phys. Rev.](https://doi.org/10.1103/PhysRevB.96.075127) B 96[, 075127 \(2017\),](https://doi.org/10.1103/PhysRevB.96.075127) [arXiv:1705.01102 \[cond-mat.mes](https://arxiv.org/abs/1705.01102)[hall\].](https://arxiv.org/abs/1705.01102)
- [27] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, Integer quantum hall transition: An alter-

native approach and exact results, [Phys. Rev. B](https://doi.org/10.1103/PhysRevB.50.7526) 50, 7526 [\(1994\).](https://doi.org/10.1103/PhysRevB.50.7526)

- [28] C. P. Herzog, P. Kovtun, S. Sachdev, and D. T. Son, Quantum critical transport, duality, and m theory, Physical Review D 75, [10.1103/physrevd.75.085020](https://doi.org/10.1103/physrevd.75.085020) (2007).
- [29] Y. Huh, P. Strack, and S. Sachdev, Conserved current correlators of conformal field theories in 2+1 dimensions, Phys. Rev. B 88[, 155109 \(2013\),](https://doi.org/10.1103/PhysRevB.88.155109) [Erratum: Phys.Rev.B 90, 199902 (2014)], [arXiv:1307.6863 \[cond-mat.str-el\].](https://arxiv.org/abs/1307.6863)
- [30] P. Romatschke and S. Säppi, Thermal free energy of large Nf QED in 2+1 dimensions from weak to strong coupling, Phys. Rev. D 100[, 073009 \(2019\),](https://doi.org/10.1103/PhysRevD.100.073009) [arXiv:1908.09835 \[hep](https://arxiv.org/abs/1908.09835)[th\].](https://arxiv.org/abs/1908.09835)
- [31] S. Giombi, G. Tarnopolsky, and I. R. Klebanov, On C_J and C_T in Conformal QED, JHEP 08[, 156,](https://doi.org/10.1007/JHEP08(2016)156) [arXiv:1602.01076 \[hep-th\].](https://arxiv.org/abs/1602.01076)
- [32] R. G. Leigh and A. C. Petkou, $SL(2,Z)$ action on three-dimensional CFTs and holography, [JHEP](https://doi.org/10.1088/1126-6708/2003/12/020) 12, 020, [arXiv:hep-th/0309177.](https://arxiv.org/abs/hep-th/0309177)
- [33] S. Coleman, [Aspects of Symmetry: Selected Erice Lec](http://www.amazon.com/Aspects-Symmetry-Selected-Erice-Lectures/dp/0521318270%3FSubscriptionId%3D13CT5CVB80YFWJEPWS02%26tag%3Dws%26linkCode%3Dxm2%26camp%3D2025%26creative%3D165953%26creativeASIN%3D0521318270)[tures](http://www.amazon.com/Aspects-Symmetry-Selected-Erice-Lectures/dp/0521318270%3FSubscriptionId%3D13CT5CVB80YFWJEPWS02%26tag%3Dws%26linkCode%3Dxm2%26camp%3D2025%26creative%3D165953%26creativeASIN%3D0521318270) (Cambridge University Press, 1988).
- [34] A. M. Polyakov, *Gauge Fields and Strings*, Vol. 3 (1987).
- [35] J. F. Fuini and L. G. Yaffe, Far-from-equilibrium dynamics of a strongly coupled non-Abelian plasma with nonzero charge density or external magnetic field, [JHEP](https://doi.org/10.1007/JHEP07(2015)116) 07, [116,](https://doi.org/10.1007/JHEP07(2015)116) [arXiv:1503.07148 \[hep-th\].](https://arxiv.org/abs/1503.07148)
- [36] S. Grozdanov and N. Poovuttikul, Generalised global symmetries in holography: magnetohydrodynamic waves in a strongly interacting plasma, JHEP 04[, 141,](https://doi.org/10.1007/JHEP04(2019)141) [arXiv:1707.04182 \[hep-th\].](https://arxiv.org/abs/1707.04182)
- [37] D. M. Hofman and N. Iqbal, Generalized global symmetries and holography, [SciPost Phys.](https://doi.org/10.21468/SciPostPhys.4.1.005) 4, 005 (2018), [arXiv:1707.08577 \[hep-th\].](https://arxiv.org/abs/1707.08577)
- [38] S. Grozdanov, A. Lucas, and N. Poovuttikul, Holography and hydrodynamics with weakly broken symmetries, Phys. Rev. D 99[, 086012 \(2019\),](https://doi.org/10.1103/PhysRevD.99.086012) [arXiv:1810.10016 \[hep](https://arxiv.org/abs/1810.10016)[th\].](https://arxiv.org/abs/1810.10016)
- [39] O. DeWolfe and K. Higginbotham, Generalized symmetries and 2-groups via electromagnetic duality in AdS/CFT , Phys. Rev. D 103[, 026011 \(2021\),](https://doi.org/10.1103/PhysRevD.103.026011) [arXiv:2010.06594 \[hep-th\].](https://arxiv.org/abs/2010.06594)
- [40] K. Skenderis, Lecture notes on holographic renormalization, [Classical and Quantum Gravity](https://doi.org/10.1088/0264-9381/19/22/306) 19, 5849 (2002).
- [41] M. Taylor, More on counterterms in the gravitational action and anomalies, (2000), [arXiv:hep-th/0002125.](https://arxiv.org/abs/hep-th/0002125)
- [42] G. Frangi, To appear, (2024).
- [43] J. Polonyi, Classical and quantum effective theories, Phys. Rev. D 90[, 065010 \(2014\),](https://doi.org/10.1103/PhysRevD.90.065010) [arXiv:1407.6526 \[hep](https://arxiv.org/abs/1407.6526)[th\].](https://arxiv.org/abs/1407.6526)
- [44] J. Polonyi, Dissipation and decoherence by a homogeneous ideal gas, Phys. Rev. A 92[, 042111 \(2015\),](https://doi.org/10.1103/PhysRevA.92.042111) [arXiv:1502.02540 \[cond-mat.stat-mech\].](https://arxiv.org/abs/1502.02540)
- [45] H. Liu and P. Glorioso, Lectures on non-equilibrium effective field theories and fluctuating hydrodynamics, [PoS](https://doi.org/10.22323/1.305.0008) TASI2017[, 008 \(2018\),](https://doi.org/10.22323/1.305.0008) [arXiv:1805.09331 \[hep-th\].](https://arxiv.org/abs/1805.09331)
- [46] S. Grozdanov and J. Polonyi, Dynamics of the electric current in an ideal electron gas: A sound mode inside the quasiparticles, Phys. Rev. D **92**[, 065009 \(2015\),](https://doi.org/10.1103/PhysRevD.92.065009) [arXiv:1501.06620 \[hep-th\].](https://arxiv.org/abs/1501.06620)