Short-range correlation and Urca process in neutron stars

Armen Sedrakian

Frankfurt Institute for Advanced Studies, D-60438 Frankfurt am Main, Germany

and Institute of Theoretical Physics, University of Wroclaw, 50-204 Wroclaw, Poland*

(Dated: 20 June 2014)

Recent measurements of high-momentum correlated neutron-proton pairs at JLab suggest that the dense nucleonic component of the compact stars contain a fraction of high-momentum neutronproton pairs that is not accounted in the familiar Fermi-liquid theory of the neutron-proton fluid mixture. We compute the rate of the Urca process in compact stars taking into the non-Fermi liquid contributions to the proton's spectral widths induced by short-range correlations. The Urca rate differs strongly from the Fermi-liquid prediction at low temperatures, in particular, the high threshold on the proton fraction precluding the Urca process in neutron stars is lifted. This observation may have a profound impact on the theories of cooling of compact stars.

PACS numbers: 97.60.Jd

Introduction. The Landau Fermi-liquid theory forms the basis of semi-phenomenological description of many low-temperature strongly correlated fermionic systems, including liquid ³He in its normal state, electrons in metals, nuclear matter, etc. It assumes that slow enough switching of interaction adiabatically evolves the lowlying states, therefore the strongly interacting regime features a sharp Fermi surface, whereas the excitations are weakly interacting quasiparticles [1]. More recent research discovered that the low-energy thermodynamics and transport of a wide class of condensed matter systems, including doped metals, heavy fermions, d-f electrons in metals, quantum dots etc. do not agree with the predictions of Fermi-liquid theory [2, 3]. The non-Fermiliquid (nFL) behavior of these systems is often associated with the breakdown of the concepts of well-defined quasiparticles, Fermi-surfaces, or both.

There exists now sufficient evidence that the proton component of the high density matter in neutron stars is an example of nuclear nFL. Recent high-energy electronscattering measurements at JLab [4] which used medium to heavy nuclear targets show that in neutron-rich nuclei, short-range interactions between the fermions form correlated high-momentum neutron-proton pairs. This, in turn, implies that protons can occupy states that are much greater than their Fermi momentum, i. e. the concept of sharp Fermi-surface does not apply. Theoretical interpretations are based on short-range-correlations (SRC) in the isospin singlet channel and led to phenomenological modeling of the proton momentum distribution tails [5]. Microscopic calculations of isospin asymmetrical nuclear systems confirm, at least qualitatively, the presence and magnitude of SRC in the momentum distributions of infinite nuclear matter and in nuclei [6, 7].

Neutrino emissivities of neutron stars play a central role in the modeling their thermal evolution. Neutron star's cores that are composed of neutrons (n), protons (p) and electrons (e) and cool predominantly by emission of neutrinos ν and anti-neutrinos $\bar{\nu}$ via the direct Urca (dUrca) [8] and and modified Urca (mUrca) processes [9, 10]:

$$n \rightarrow p + e + \bar{\nu}_e,$$
 (1)

$$N+n \rightarrow N+p+e+\bar{\nu}_e, \quad N \in p, n$$
 (2)

up to the onset of the superfluidity of the nucleonic components, after which these processes are suppressed by the pairing gap in the spectra of neutrons and protons, for a review see [11]. The small fraction of μ -ons is neglected here.

The mUrca process has been studied in the pioneering work [9, 10] as the dominant cooling process in neutron stars. (We do not consider here the interesting possibility that the neutron star cores consist of hypernuclear or deconfined quark matter.) The more efficient Urca process was later discovered by Lattimer et al. [8] to occurs in the neutron-proton-electron Fermi-liquid mixture above the threshold $x_p = n_p/n \ge 0.11$, where $n_i \ i \in n, p$ denote the particle numbers of neutrons and protons and $n \equiv n_n + n_p$ [8]. This threshold is reflected in the stepfunction in the Fermi-liquid Urca neutrino emissivity

$$\epsilon_{\text{Urca}} = \epsilon_0 \,\theta(|p_{Fe} + p_{Fp} - p_{Fn}|),\tag{3}$$

 p_{Fi} being the Fermi momentum of particle $i \in n, p, e$. If kinematically allowed the emissivity of the dUrca process is given by [8]

$$\epsilon_0 = \frac{457\pi}{10080} g_A^2 G^2 p_{Fe} m_p^* m_n^* T^6, \qquad (4)$$

where m_i^* is the effective mass of particle *i*, *G* is the weak coupling constant, g_A is the axial vector coupling constant, *T* is the temperature [24]. The "triangle relation" enforced by this step-function requires proton concentration $x_p \ge 0.11$ to fulfill the momentum and energy conservation. The nFL nature of the proton component argued above may mitigate this constrained [12, 13]. To date, the SRC-induced modification of the dUrca process was accessed in terms of correction factors built from

the momentum distribution of protons in dense matter [12, 13].

In this Letter we compute the Urca process microscopically in terms of the nFL spectral function of the proton component. By doing so we establish direct contact with the fundamental microscopic quantities - the selfenergies of the system, which can be computed in a controlled many-body theory. Physically, the tensor interactions between the minority component of protons and the majority component of neutrons nFL behavior. We show that in the quasiparticle limit for proton spectral function we naturally recover the standard Fermi-liquid result [8]. We also argue that the dichotomy of the cooling scenarios of neutron stars (fast vs slow) triggered by the dUrca process in massive stars will be lifted if the nFL rates are used, i. e., the paradigm of fast cooling of (only) massive neutron stars via the dUrca process should be abandoned and replaced by a smooth transition from slow to fast cooling scenarios. Before proceeding, let us mention that the finite width effects have been considered in the context of neutrino bremsstrahlung in the pioneering work by Raffelt et al. [14, 15] and have been later put on more formal grounds in the real-time Green's functions formalism in Refs. [16, 17].

Formalism. The anti-neutrino emissivity is given by [18]

$$\epsilon_{\nu} = -\left(\frac{2G}{\sqrt{2}}\right)^{2} \int \frac{d^{3}k}{(2\pi)^{3}2\omega_{\bar{\nu}}(\vec{k})} \int \frac{d^{3}k'}{(2\pi)^{3}2\omega_{e}(\vec{k}')} \\ \int d^{4}q \,\delta(k+k'-q)g(q_{0})\bar{f}_{e}(\omega_{e})\Im[\Lambda^{\mu\nu}(k,k')\Pi^{R}_{\mu\nu}(q)],$$
(5)

where k and k' to the four-momenta of anti-neutrino and electron, ω_{ν} and ω_e are their on-mass-shell energies, $\bar{f}(p) = 1 - f(p)$ with f(p) being the Fermi-Dirac distribution function and index e refers to electrons, $\Lambda^{\mu\nu} = \text{Tr} \left[\gamma^{\mu}(1-\gamma^5) \not{k}\gamma^{\nu}(1-\gamma^5) \not{k'}\right], g(q_0)$ is the Bose distribution function and $\Pi^R_{\mu\nu}(q)$ is the retarded polarization. We use the spectral representation of the retarded Green's functions of neutrons and protons and compute the polarization tensor in the one-loop approximation. After contracting the leptonic and hadronic traces we obtain

$$\epsilon_{\nu} = 2G^2 \int d^4 q L_1(q) L_2(q),$$
 (6)

where the loop integrals are given by

$$L_1(q) = \int \frac{d^4 p' d^4 p}{(2\pi)^2} \delta^{(4)}(p - p' + q) A_n(p) A_p(p') f(p') \bar{f}(p),$$
(7)

$$L_2(q) = \int \frac{d^3k d^3k'}{(2\pi)^6} \bar{f}(k') \bar{f}(k) \omega_{\bar{\nu}} \delta^4(k+k'-q), \qquad (8)$$

where p' and p refer to the four-momenta of the neutron and proton. The spectral functions of the neutrons with $\tau \in n, p$ are given by

$$A_{\tau}(p) = \frac{\gamma_{\tau}}{[\epsilon - \epsilon_{\tau}(p)]^2 + \gamma_{\tau}^2/4},\tag{9}$$

where its width and quasiparticle energy are given by $\gamma_{\tau} = -2\Im m \Sigma_{\tau}^{R}(p)$ and $\epsilon_{\tau}(p) = \epsilon_{\tau}^{\text{kin}} + \text{Re} \Sigma_{\tau}^{R}(p) - \mu_{\tau}$, respectively. Here $\Sigma_{\tau}^{R}(p)$ is the retarded self-energy, μ_{τ} is the chemical potential and $\epsilon_{\tau}^{\text{kin}}(p)$ is the kinetic energy. In the small γ_{τ} limit [19, 20]

$$A_{\tau}(p) = 2\pi\delta(\epsilon_{\tau} - \epsilon_{\tau}(p)) - \frac{\gamma_{\tau}}{(\epsilon_{\tau} - \epsilon_{\tau}(p))^2} + O(\gamma_{\tau}^2), (10)$$

where any integration over the second term is understood as Cauchy principal value, $\epsilon_{\tau}(p) = p^2/2m_{\tau}^* - \mu_{\tau}$ is the quasiparticle spectrum, m_{τ}^* is the quasiparticle mass which is determined by the on-shell derivative of Re $\Sigma_{\tau}^R(p)$ with respect to the momentum:

$$\frac{m_{\tau}}{m_{\tau}^*} = 1 + \frac{m_{\tau}}{p_{F\tau}} \frac{\partial \Sigma_{\tau}^R(p)}{\partial p} \bigg|_{p=p_{F\tau}}.$$
(11)

The first term in Eq. (10) is the quasiparticle spectral function corresponding to $\gamma_{\tau} = 0$. The Fermi Golden Rule expression for the emissivity can be easily recovered in this limit (i. e. $\gamma_{\tau} \to 0$). Indeed one finds the standard expression by doing first integrations over the energy components p_0 and p'_0 in Eq. (7) using Eq. (10) and subsequently taking the *q*-integral in Eq. (7) using delta functions. Note that the angle integrations eliminate the spatial parts of the four-vector products in the matrix element and it can be approximated as

$$\sum |\mathcal{M}_{\text{Urca}}|^2 = 64(k \cdot p')(p \cdot k') \simeq 64p_0 p'_0 k_0 k'_0, \quad (12)$$

where the sum stands for the average over the initial and summation over the final spins of baryons. The L_2 integral is computed by neglecting in the δ -function the neutrino energy-momentum compared to that of the electron, which fixes the momentum transfer to the Fermi momentum of the electron. In this manner, we obtain the neutrino emissivity

$$\epsilon_{\nu} = G^2 \frac{p_{Fe} m_p^* m_n^*}{4\pi^5} \int_0^\infty d\omega_{\nu} \omega_{\nu}^3 \\ \times \int_{-\infty}^\infty d\omega g(\omega) f_e(\omega_{\nu} - \omega) L_1(\omega, q = p_{Fe}).$$
(13)

The three-dimensional integral in Eq. (13) must be computed numerically in general. If the spectral function of protons is taken in the limit $\gamma_p \to 0$ the integral can be computed analytically and we find

$$L_1(q) = cT \ln \left| \frac{1 + \exp\left(-\frac{\epsilon_{\min} - \mu_p}{T}\right)}{1 + \exp\left(-\frac{\epsilon_{\min} + \omega - \mu_p}{T}\right)} \right|, \quad (14)$$

where $c \equiv m_n^* m_p^* / 2\pi q$ and the minimal energy ϵ_{\min} arises from the limits on the azimuthal angular integration and is given by

$$\epsilon_{\min} = (m_n^*/2q^2) \left(\mu_n + \omega - \mu_p + q^2/2m_p^*\right)^2.$$

Substituting Eq. (14) in Eq. (13) we obtain the general form of Urca emissivity at finite temperatures for $\gamma_{\tau} = 0$. Further simplifications arise at low temperatures, where the small ω expansion leads to

$$L_1(q)/cT\omega = [\exp(\epsilon_{\min} - \mu_p) + 1]^{-1}$$
(15)

after which the remaining integrals can be computed analytically. In the limit $T \to 0$ one finds then $L_1(q)/cT\omega = \theta(\mu_p - \epsilon_{\min}) = \theta(|p_{Fe} + p_{Fp} - p_{Fn}|)$, i. e., the quasiparticle result of Ref. [8] is recovered from Eq. (13).

We now turn to the finite width effects in the proton spectral function, while keeping the neutron spectral function in the quasiparticle approximation, given by the δ -function term in Eq. (10). With the full spectral function for protons given by Eq. (9) we find

$$L_1(q) = c \int d\epsilon [f(\epsilon) - f(\omega + \epsilon)] \left\{ \frac{1}{2} - \frac{1}{\pi} \operatorname{atan} \left[\frac{\epsilon_{\min} - \epsilon}{\gamma_p / 2} \right] \right\}.$$
(16)

For concrete calculations, we need to specify the spectrum of protons and neutrons $\epsilon_{\tau}(p)$ and the width of the proton spectral function. We adopt for neutrons and protons $\epsilon_{\tau}(p) = p^2/2m_{\tau}^* - \mu_{\tau}$, where $m_{\tau}^*/m\tau = 0.7$ and μ_{τ} is taken to be equal to the Fermi energy computed from the partial density $n_{\tau} = k_{F\tau}^3/3\pi^2$, where $k_{F\tau}$ is the Fermi wave-vector. We model the width of protons by

$$\gamma_p = a_p T^2 \left[1 + \left(\frac{\epsilon - \mu_p}{\pi T}\right)^2 \right], \qquad (17)$$

where $a_p = (\hbar v_{Fp} \mu_p^{-1}) \sum_{\tau=n,p} \bar{\sigma}_{p\tau} n_{\tau} \mu_{\tau}^{-1}$, where n_{τ} are the number densities and $\bar{\sigma}_{np} = 7 \text{ fm}^2$ and $\bar{\sigma}_{pp} = 4 \text{ fm}^2$ are the angle averaged cross-sections for neutron-proton and proton-proton scattering.

Results. The neutrino emissivity was computed using Eq. (13) and the spectral width is given by Eq. (17). In Fig. 1 we show the nFL dUrca rate normalized by the same quantity in the quasiparticle approximation as a function of proton fraction $x_p = \rho_p/\rho$, where ρ_p and ρ are the proton and total densities for two fixed values of temperature T = 0.2 and 1 MeV. In the quasiparticle limit we observe the thermal "activation" of the dUrca process due to the opening of the phase space when using the finite temperature expression (15) instead of zerotemperature analogue containing θ -function. At temperature 1 MeV ~ 10¹⁰ K the dUrca process is comparable to the mUrca already for proton fraction $x_p = 0.09$ instead of 0.11 in the zero temperature limit. Next, including the width of protons, which implies L_1 given by (16), we see




FIG. 1: The nFL-Urca emissivity in units of ϵ_0 for T = 1 MeV and T = 0.2 MeV for density $\rho = 1.5\rho_0$ with $\rho_0 = 2.7 \times 10^{14}$ g cm⁻³ as a function of proton fraction x_p . The dashed lines show the quasiparticle limit corresponding to $\gamma_p = 0$.

that the sharp threshold is replaced by a smooth increase of the Urca rate as the proton fraction increases. The fundamental difference to the quasiparticle limit is that the dUrca process dominates the mUrca processes even for rather small values of x_P that are compatible with the symmetry energy for the nuclear equation of states, i.e., practically all the EoS of neutron star matter allow for dUrca process, if the width of protons induced by the short-range correlations are included.

Figure 2 illustrates the temperature dependence of the dUrca process for fixed proton fractions (x_n) under both the quasiparticle and finite-width approximations. Starting with the quasiparticle limit, we observe the wellknown behavior of the dUrca process: for sufficiently large proton fractions (here $x_p = 0.125$), the dUrca process is kinematically allowed and finite width and quasiparticle results nearly coincide. In contrast, for lower proton fractions (here $x_p = 0.1, 0.075$, and 0.05), the process is strongly suppressed in the quasiparticle limit at low temperatures. However, for temperatures around 1 MeV and higher, thermal "activation" of the dUrca process becomes evident for these smaller values of x_p . Including the proton's spectral width leads to significant differences in the emissivity at low temperatures $T \leq 1$ MeV. Instead of vanishingly small results, the emissivity decays much more slowly. Thus, it is evident that the finite width of protons induced by short-range correlations lift the sharp threshold on the dUrca processes in neutron star matter. The asymptotics of the low-temperature Urca process appears to be weakly dependent on the temperature and are comparable to the rate of the mUrca process.

Implications for neutron star cooling. The common picture of neutron star cooling assumes that for $x_p \geq$




FIG. 2: The nFL-Urca emissivity in units of ϵ_0 for $x_p = 0.05$, 0.10 and 0.125 and $\rho = 1.5\rho_0$ as a function of temperature. The solid lines show the full result, whereas the dashed lines – the limit $\gamma_p = 0$.

0.11 the dUrca process is operative, otherwise the mUrca process (which is permitted for any value of x_p) is the dominant cooling mechanism. Such large values of x_p are achieved only in massive stars $M \ge 1.7 \ M_{\odot}$, the exact value depending on the symmetry energy behavior of the underlying equation of state at supranuclear densities. As a consequence, modern cooling simulations display a sharp drop in temperatures of stars that have masses lying beyond the threshold mass for the dUrca process. As a consequence the massive stars featuring the dUrca process cool much faster than less massive stars that cool by the mUrca process.

As we have shown above, the inclusion of the proton spectral width, which is one of the manifestations of SRC in asymmetrical nuclear matter leads to the dUrca process which is operative far below the threshold proton fraction. This implies that cooling simulations including short-range correlation will be in stark contrast to the standard cooling scenarios where there is a sharp drop in the stellar temperatures for massive stars once the mass threshold is crossed.

Conclusions. In conclusion, there is strong experimental evidence for short-range correlations in asymmetrical nuclear systems, which manifest as tails in the momentum distribution of protons. These correlations have been demonstrated in numerous experiments, particularly those conducted at JLab [4]. Consequently, the Fermi-liquid approximation is likely invalid for protons in dense matter. In this work, we obtained the emissivity of dense n - p - e matter via the dUrca process, accounting for the finite width of protons. In the zero-width approximation, we recover the pioneering result of Ref. [8]. We have shown that the dUrca process remains effective even for low proton fractions, replacing the threshold behavior of the dUrca process with a smooth dependence on x_p . Finally, it should be noted that from the perspective of many-body nuclear theory, the distinction between the dUrca and mUrca processes is strictly possible only in the quasiparticle approximation. Once the width of the proton propagator is taken into account, the higher-order interactions (higher-order loop diagrams) are automatically taken into account. This has been extensively discussed in the case of the neutrino bremsstrahlung process in Refs. [14–17].

In the case of Urca processes, for example, the mUrca rate computed in Ref. [10] can be viewed as the dUrca process with the width computed in the second Born approximation to the nucleon (neutron or proton) propagator in the one-pion (eventually plus ρ -meson) exchange approximation. Nevertheless, the description of dUrca processes in terms of proton spectral functions allow one to establish contact with the study of SRC in nuclear systems, as well as many-body formulations of nuclear theory that are based on spectral functions. It allows also us to trace the smooth physical dependence of neutrino emission rates on the proton fraction and temperature. This last point may have important ramifications for the cooling studies of neutron stars. Finally, it is important to note that SRC can significantly impact the low-density physics of neutron stars. This includes influencing pairing-type correlations such as isospin-singlet neutron-proton pairing and deuteronization [21, 22], as well as isospin-triplet proton polarons [23].

Acknowledgements. The author acknowledges support from Deutsche Forschungsgemeinschaft Grant No. SE 1836/5-3 and the Polish NCN Grant No. 2020/37/B/ST9/01937. Gratitude is extended to the Institute for Nuclear Theory at the University of Washington for its kind hospitality, stimulating research environment, and partial support through the INT's U.S. Department of Energy Grant No. DE-FG02-00ER41132 during INT workshop "Astrophysical neutrinos and the origin of the elements". The author also gratefully acknowledges discussions on related topics with M. Alford, A. Haber, A. Harutyunyan, and S. Reddy.

- * Electronic address: sedrakian@fias.uni-frankfurt.de
- A. Leggett, Quantum Liquids. Oxford University Press, Oxford, 2006.
- [2] G. R. Stewart, Non-Fermi-liquid behavior in d- and f-electron metals, Reviews of Modern Physics 73 (Oct., 2001) 797–855.
- [3] C. M. Varma, Z. Nussinov and W. van Saarloos, Singular or non-Fermi liquids, Physics Reports 361 (May, 2002) 267-417, [cond-mat/0103393].
- [4] O. Hen, M. Sargsian, L. B. Weinstein, E. Piasetzky and et al, Momentum sharing in imbalanced Fermi systems, Science 346 (Oct., 2014) 614–617, [1412.0138].

- [5] M. M. Sargsian, New properties of the high-momentum distribution of nucleons in asymmetric nuclei, Phys. Rev. C 89 (Mar, 2014) 034305.
- [6] T. Frick, H. Müther, A. Rios, A. Polls and A. Ramos, Correlations in hot asymmetric nuclear matter, Phys. Rev. C 71 (Jan., 2005) 014313, [nucl-th/0409067].
- [7] A. Rios, A. Polls and W. H. Dickhoff, Density and isospin-asymmetry dependence of high-momentum components, Phys. Rev. C 89 (Apr., 2014) 044303, [1312.7307].
- [8] J. M. Lattimer, M. Prakash, C. J. Pethick and P. Haensel, Direct URCA process in neutron stars, Physical Review Letters 66 (May, 1991) 2701–2704.
- H.-Y. Chiu and E. E. Salpeter, Surface X-Ray Emission from Neutron Stars, Phys. Rev. Lett. 12 (Apr., 1964) 413–415.
- [10] B. L. Friman and O. V. Maxwell, Neutrino emissivities of neutron stars., Astrophys. J. 232 (Sept., 1979) 541–557.
- [11] A. Sedrakian and J. W. Clark, Superfluidity in nuclear systems and neutron stars, European Physical Journal A 55 (Sept., 2019) 167, [1802.00017].
- [12] L. Frankfurt, M. Sargsian and M. Strikman, Recent Observation of Short-Range Nucleon Correlations in Nuclei and Their Implications for the Structure of Nuclei and Neutron Stars, International Journal of Modern Physics A 23 (2008) 2991–3055, [0806.4412].
- [13] M. McGauley and M. M. Sargsian, Off-Fermi Shell Nucleons in Superdense Asymmetric Nuclear Matter, arXiv e-prints (Feb., 2011) arXiv:1102.3973, [1102.3973].
- [14] G. Raffelt, D. Seckel and G. Sigl, Supernova neutrino scattering rates reduced by nucleon spin fluctuations: Perturbative limit, Phys. Rev. D 54 (Aug., 1996) 2784-2792, [astro-ph/9603044].
- [15] G. Raffelt and G. Sigl, Numerical toy-model calculation

of the nucleon spin autocorrelation function in a supernova core, Phys. Rev. D **60** (July, 1999) 023001, [hep-ph/9808476].

- [16] A. Sedrakian and A. Dieperink, Coherence effects and neutrino pair bremsstrahlung in neutron stars, Physics Letters B 463 (Sept., 1999) 145–152, [nucl-th/9905039].
- [17] A. Sedrakian and A. E. L. Dieperink, Coherent neutrino radiation in supernovae at two loops, Phys. Rev. D 62 (Oct., 2000) 083002, [astro-ph/0002228].
- [18] A. Sedrakian, The physics of dense hadronic matter and compact stars, Progress in Particle and Nuclear Physics 58 (Jan., 2007) 168-246, [nucl-th/0601086].
- [19] T. Alm, G. Röpke, A. Schnell and H. Stein, Two-nucleon correlations and the nucleon spectral function in nuclear matter at finite temperature: low-density region, Physics Letters B 346 (Feb., 1995) 233–239.
- [20] A. Schnell, T. Alm and G. Röpke, Nucleon self-energies and spectral functions in nuclear matter at finite temperature with a realistic potential, Physics Letters B 387 (Feb., 1996) 443–448.
- [21] A. Sedrakian and J. W. Clark, Pair condensation and bound states in fermionic systems, Phys. Rev. C 73 (Mar., 2006) 035803, [nucl-th/0511076].
- [22] M. Stein, A. Sedrakian, X.-G. Huang and J. W. Clark, BCS-BEC crossovers and unconventional phases in dilute nuclear matter, Phys. Rev. C 90 (Dec., 2014) 065804, [1410.1053].
- [23] H. Tajima, H. Moriya, W. Horiuchi, E. Nakano and K. Iida, *Polaronic proton and diproton clustering in neutron-rich matter*, *Physics Letters B* 851 (Apr., 2024) 138567, [2304.00535].
- [24] For the sake of simplicity we drop the numerically subdominant vector current neutrino emission.