

# Deformation Families of Quasi-Projective Varieties and Symmetric Projective K3 Surfaces

Fan Xu

## ABSTRACT

The main aim of this paper is to construct a deformation family of symmetric projective K3 surfaces through a deformation family of quasi-projective varieties together with construction of symmetric Kahler metrics on symmetric projective K3 surfaces. Firstly, if the normal bundle satisfies the Diophantine condition, the tubular neighborhood of an elliptic curve embedded in  $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$  can be identified through the toroidal group. Secondly, with ample line bundles fixed on  $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$ , complete Kahler metrics are constructed on the quasi-projective variety from  $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$ . Finally, a deformation family of symmetric projective K3 surfaces over a 10-dimensional complex manifold is constructed through a smooth compactifiable deformation of complete quasi-projective varieties from  $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$  over a 9-dimensional complex manifold and an analogue deformation family.

## 1. Introduction

This paper is motivated by gluing construction of K3 surfaces through two open complements of tubular neighborhoods of elliptic curves embedded in  $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P^2}$  in [1]. What's more, a gluing construction of projective K3 surface was present in [2].

For a rational elliptic surface  $X$  defined as the blow-up of a projective plane at nine base points for a pencil of cubics, Hans-Joachim Hein present some complete Calabi-Yau metrics on  $X \setminus D$  with  $D$  being a fiber on  $X$  in [3]. However, blowing up arbitrary nine points of a projective plane, the result manifold may not be a rational elliptic surface.

Let  $C_0(\tau) = \mathbb{C} / \langle 1, \tau \rangle$  with  $\tau \in Y := \{\tau | \tau \in \mathbb{C} \ \& \ \text{Im}\tau > 0\}$ . What's more, let  $C_0 = C_0(\tau_0) = \mathbb{C} / \langle 1, \tau_0 \rangle$  with  $\tau_0 \in Y$ . In addition, for a smooth elliptic curve  $C_0(\tau) = \mathbb{C} / \langle 1, \tau \rangle$  with  $\tau \in Y$  embedded in  $\mathbb{C}P^2$ . The embedding map from  $C_0(\tau)$  to  $\mathbb{C}P^2$  can be induced by the Weierstrass  $\wp$ -function equation:

$$\wp = \sum_{\lambda \in \langle 1, \tau \rangle \setminus \{0\}} \left( \frac{1}{(z-\lambda)^2} - \frac{1}{\lambda^2} \right) + \frac{1}{z^2}.$$

The Eisenstein series are defined as

$$G_{2k}(\langle 1, \tau \rangle) = \sum_{\lambda \in \langle 1, \tau \rangle \setminus \{0\}} \lambda^{-2k}.$$

Here,  $\langle 1, \tau \rangle$  is a lattice and  $\mathbb{C}$  represents the set consisting of complex numbers. The holomorphic embedding map  $f_\tau : C_0(\tau) = \mathbb{C} / \langle 1, \tau \rangle \rightarrow \mathbb{C}P^2$  defined by  $f_\tau([z]) = [\wp(z) : \wp'(z) : 1] = [z_1 : z_2 : z_3]$  for  $[z] \neq [0]$  and  $f_\tau([0]) = [0 : 1 : 0]$  for  $\tau \in Y$ . What's more,  $f_\tau(C_0(\tau)) = \{-4z_1^3 + 60G_4(\langle 1, \tau \rangle)z_2^2z_1 + 140G_6(\langle 1, \tau \rangle)z_3^3 + z_2^2z_3 = 0\} \subset \mathbb{C}P^2$  for  $\tau \in Y$ .

So  $f_\tau(C_0(\tau))$  is a submanifold of  $S_0 = \mathbb{C}P^2$ . Blowing up nine points  $Z = \{p_1, p_2, \dots, p_9\}$  on  $f_\tau(C_0(\tau))$  such that the normal bundle of  $C(\tau)$  satisfies the Diophantine condition proposed in [2],  $S_0 = \mathbb{C}P^2$  will be transferred to  $S(\tau) \cong \mathbb{C}P^2 \# 9\overline{\mathbb{C}P}^2$ . In addition, let  $C(\tau)$  be the strict transform of  $f_\tau(C_0(\tau))$  in  $S(\tau)$ . Then  $S(\tau) \setminus C(\tau)$  is a quasi-projective variety for  $\tau \in Y$ .

Furthermore, under the condition that the normal bundle of  $C(\tau)$  satisfies the Diophantine condition, there is a deformation family of  $S(\tau_0) \cong \mathbb{C}P^2 \# 9\overline{\mathbb{C}P}^2$  over a 9-dimensional complex manifold.

Elizabeth Gasparim and Francisco Rubilar present some new definitions for deformation family in [4], especially for the deformation family of open manifolds. Taking use of the new definitions, the first main theorem is as below.

**Theorem 1.1:** *There is a smooth compactifiable deformation of  $S(\tau_0) \setminus C(\tau_0)$  over a 9-dimensional complex manifold  $Y_1$ . In addition, the deformation is differential trivial along  $Y_1$ .*

Firstly, there are ample line bundles properly defined on the fibers of the deformation family of  $S(\tau_0) \setminus C(\tau_0)$ .

Secondly, in [2], there was description for construction of the complete Kahler metric on  $S(\tau) \setminus C(\tau)$  corresponding to an ample line bundle on  $S(\tau)$ . So, in this paper, the analogue complete Kahler metrics corresponding to the ample line bundles selected will be constructed in more detail on the fibers.

Then the deformation family of quasi-projective varieties above tends to be a deformation family of non-compact complete Kahler manifolds.

Thirdly, a symmetric projective K3 surface can be constructed through gluing the open complement of the tubular neighborhood of  $C(\tau)$  in  $S(\tau)$  and an analogue manifold.

Then a deformation family of symmetric projective K3 surfaces can be constructed the compactifiable deformation of  $S(\tau_0) \setminus C(\tau_0)$  and an analogue deformation family. That is the second main theorem as below.

**Theorem 1.2:** *There is a deformation family of symmetric projective K3 surfaces over a 10-dimensional complex manifold along with a symmetric Kahler metric on each fiber.*

The following is to introduce the main contents in following sections of this paper.

In section 2, the main goal is to present the gluing construction of the symmetric projective K3 surface and the corresponding toroidal group for the tubular neighborhood.

Firstly, the Diophantine condition will be introduced for the construction of tubular neighborhood of an elliptic curve embedded in  $\mathbb{C}P^2 \# 9\overline{\mathbb{C}P}^2$ .

Secondly, fixing a suitable ample line bundle on  $S(\tau) \setminus C(\tau)$ , the open complement of the tubular neighborhood of  $C(\tau)$  in  $S(\tau)$  and an analogue open manifold can be glued to a symmetric projective K3 surface.

Finally, the specific defined toroidal groups satisfying the Diophantine condition is just properly defined in the special class of the toroidal groups defined in [11, Theorem].

In section 3, the main goal is to prove theorem 1.1 and construct complete Kahler metrics on the fibers of the deformation family through fixing an ample line bundle on  $S(\tau)$ .

What's more, through the process of constructing the complete Kahler metric on  $S(\tau)\setminus C(\tau)$ , the condition for the gluing construction of symmetric K3 surface can be confirmed.

In section 4, the main goal is to prove theorem 1.2.

## 2. Symmetric Projective K3 Surfaces and Toroidal Groups

### 2.1. A Gluing Construction of Symmetric Projective K3 Surfaces

Firstly, it is to give the details for the gluing construction of symmetric projective K3 surfaces.

Taking use of the notations from the introduction, Julius Ross and David Witt Nyström stated that the holomorphic tubular neighborhood of  $C(\tau)$  in  $S(\tau)$  might not exist for  $\tau \in Y$  in [5].

V. I. Arnold proposed a theorem that in most situations an elliptic curve smoothly embedded in a complex surface with self-intersection index 0 should have a small neighborhood biholomorphically equivalent to a neighborhood of zero section of a normal bundle in [6]. Ueda also mentioned a corresponding theorem in [7].

Another fact is that any topologically trivial holomorphic line bundle on an elliptic curve is flat [6]. The following is a smooth version definition for the monodromy of a complex line bundle over a Riemann surface.

**Definition 2.1.1:** For a flat connection  $\nabla$  on a complex line bundle  $L$  representing an isomorphic class of complex line bundle over a Riemann surface  $M$ . What's more, define  $g = e^{\sqrt{-1}t} \in S^1$  for  $t \in [0, 2\pi]$  and let  $\gamma : S^1 \rightarrow M$  be a smooth closed curve. In addition, let  $\psi$  denote a parallel section of  $\hat{\gamma}^*L$  with  $\hat{\gamma} = \gamma \circ g$ . Then  $h(\gamma) \in S^1$  with  $\psi_{2\pi} = h(\gamma)\psi_0$  is defined to be the monodromy of  $L$  along  $\gamma$ .

Let  $R \subset \mathbb{C}$  be the set consisting of real numbers and  $\mathbb{Z} \subset R$  be the set consisting of integers. In addition, let  $\mathbb{N} \subset \mathbb{Z}$  be the set consisting of positive integers.

The following is the Diophantine condition for a pair of real numbers from [2].

**Definition 2.1.2:** [2, Definition 2.1] A pair of numbers  $(p, q) \in \mathbb{R}^2$  is said to satisfy the Diophantine condition if there exists  $\alpha > 0$  and  $A > 0$  such that

$$\min_{\mu, \nu \in \mathbb{Z}} |n(p + q\sqrt{-1}) - (\mu + \nu\sqrt{-1})| \geq A \cdot n^{-\alpha}$$

for any  $n \in \mathbb{N}$ .

With the definitions above, the following is the Diophantine condition for a flat line bundle on an elliptic curve embedded in a complex surface to construct the holomorphic tubular neighborhood of the elliptic curve.

**Definition 2.1.3** [2, Definition 2.2] For  $\tau \in Y$ , let  $C(\tau) = \mathbb{C}/\langle 1, \tau \rangle$  be a smooth elliptic curve with  $\alpha$  and  $\beta$  being smooth loops corresponding to the line segments  $[0, 1]$  and  $[0, \tau]$  respectively. For a topologically trivial line bundle  $L \in \text{Pic}^0(C)$ , let  $e^{p \cdot 2\pi\sqrt{-1}}$  and  $e^{q \cdot 2\pi\sqrt{-1}}$  be the monodromies of  $L$  along the loops  $\alpha$  and  $\beta$  respectively with  $(p, q) \in \mathbb{R}^2$ . Then  $L$  is said to satisfy the Diophantion condition if  $(p, q)$  satisfies the Diophantine condition.

Let  $(p, q)$  be a number pair satisfying the Diophantine condition in definition 2.1.2. According to [1, Theorem 6.2], if  $N_{C/S} \in \text{Pic}^0(C)$  satisfies the Diophantine condition in definition 2.1.3,  $-\log d(\mathbb{I}_C, L^n) = O(\log n)$  as  $n \rightarrow \infty$  with  $d$  being an invariant distance on  $\text{Pic}^0(C)$  and  $\mathbb{I}_C$  being the holomorphically trivial line bundle on  $C$ . So from theorem 3 in [7] which is a generalized result originating from [6] by V.I. Anorld, the following theorem is true.

**Theorem 2.1.4:** [2] If the normal bundle  $N_{C/S} \in \text{Pic}^0(C)$  satisfies the Diophantine condition, then  $C$  admits a holomorphic tubular neighborhood in  $S \cong \mathbb{C}P^2 \# 9\mathbb{C}P^2$  biholomorphic to a neighborhood of the zero section in  $N_{C/S}$ .

Now assume that  $N_{C(\tau)/S(\tau)}$  satisfies the Diophantine condition. Let  $\alpha$  and  $\beta$  be smooth loops corresponding to the line segments  $[0, 1]$  and  $[0, \tau]$  on  $C(\tau)$ . Then let  $e^{p \cdot 2\pi\sqrt{-1}}$  and  $e^{q \cdot 2\pi\sqrt{-1}}$  be the monodromies of  $N_{C(\tau)/S(\tau)}$  along the loops  $\alpha$  and  $\beta$  respectively with  $(p, q) \in \mathbb{R}^2$ . So  $(p, q)$  should satisfy the Diophantine condition. Taking use of theorem 2.1.4, the following statement is true.

There is a tubular neighborhood  $W$  of  $C(\tau)$  biholomorphic to a neighborhood of the zero section in  $N_{C(\tau)/S(\tau)}$  which can be expressed as

$$W := \{(z, w) \in \mathbb{C}^2 \mid |w| < r\} / \sim$$

for a real number  $r > 1$ , where  $\sim$  is the equivalence relationship generated by

$$(z, w) \sim (z + 1, \exp(p \cdot 2\pi\sqrt{-1}) \cdot w) \sim (z + \tau, \exp(q \cdot 2\pi\sqrt{-1}) \cdot w)$$

for  $\tau \in Y$ .

Then for  $s \in \Delta \setminus \{0\}$  with  $\Delta := \{s \in \mathbb{C} \mid |s| < 1\}$ , let

$$M_s := S \setminus \left\{ [(z, w)] \in W \mid |w| \leq \sqrt{|s|/r} \right\}$$

and

$$V_s := \left\{ [(z, w)] \in W \mid \sqrt{|s|/r} < |w| < \sqrt{|s|r} \right\}.$$

The following is to describe the gluing construction of symmetric projective K3 surfaces.

Let  $E_1, E_2, \dots, E_8$  and  $E_9$  being the exceptional divisors corresponding to the points in  $Z := \{p_1, p_2, \dots, p_9\}$  respectively. What's more, let  $\pi : S(\tau) \rightarrow \mathbb{C}P^2$  be the blow up of  $\mathbb{C}P^2$  at nine points  $Z := \{p_1, p_2, \dots, p_9\} \subset f_\tau(C_0(\tau))$  and  $H = (\pi)^* \mathcal{O}_{\mathbb{C}P^2}(1)$  be the pull back bundles. Then the Nakai-Moishezon criterion [8, Appendix A] could be used to prove the following simplified theorem.

**Lemma 2.1.5:** [9, Theorem 3] Let  $\pi : S(\tau) \rightarrow \mathbb{C}P^2$  be the blowing up of at nine points defined above. For the line bundles of the form  $L = d \cdot H - k \cdot \sum_{i=1}^9 E_i$  with  $k \geq 2$  and  $d \geq 3k + 1$  being integers, if  $9 \leq \frac{d^2}{k^2} - 1$ , then the line bundles are ample.

Now let the  $C_0^\pm(\tau)$  be two copies of  $C_0(\tau)$  embedded in  $S_0^\pm = CP^2$  with the map defined in the introduction for  $\tau \in H$ . Then the map  $\ell_{CP^2}([a_1 : a_2 : a_3]) = [a_1 : -a_2 : a_3]$  for  $[a_1 : a_2 : a_3] \in CP^2$  is a holomorphic involution on  $CP^2$  and induces a holomorphic involution on  $C_0(\tau)$  as well as  $C_0^\pm(\tau)$  (i.e.  $(\ell_{CP^2})^{-1} = \ell_{CP^2}$ ). So there is a biholomorphic map  $\ell_{S_0^\pm}$  induced by  $\ell_{CP^2}$  from  $S_0^+$  to  $S_0^-$ . Since  $C_0^\pm(\tau)$  are two copies of  $C_0(\tau)$ , there exists an identity map  $g_\tau$  from  $C_0^+(\tau)$  to  $C_0^-(\tau)$ . Let  $Z^\pm = \{p_1^\pm, p_2^\pm, \dots, p_9^\pm\}$  with  $p_i^- = \ell_{CP^2}(p_i^+)$  for  $i \in \{1, 2, \dots, 9\}$  be the sets collecting the nine blow up points on  $C_0^+(\tau)$  and  $C_0^-(\tau)$  respectively. What's more, let  $p_0^+$  and  $p_0^- = \ell_{CP^2}(p_0^+)$  being inflection points of  $C_0^\pm(\tau)$  respectively.

In addition, let  $C^\pm(\tau)$  be the strict transforms of  $C_0^\pm(\tau)$  after blowing up nine points in the sets  $Z^\pm = \{p_1^\pm, p_2^\pm, \dots, p_9^\pm\}$  on  $C_0^\pm(\tau)$  and  $S^\pm(\tau) \cong CP^2 \# 9\overline{CP}^2$  be the manifolds derived from  $S_0^\pm = CP^2$  after blowing up respectively. Furthermore, let  $\ell_{S^+(\tau)} : S^+(\tau) \rightarrow S^-(\tau)$  be the biholomorphic map defined on  $S^+(\tau)$  induced by the biholomorphic map  $\ell_{S_0^\pm} : S_0^+ \rightarrow S_0^-$  defined on  $S_0^\pm$ .

Then the biholomorphic map  $\tilde{g}_\tau : C^+(\tau) \rightarrow C^-(\tau)$  induced by  $g_\tau$  will map the normal bundle  $N_+ \in Pic^0(C^+(\tau))$  to the normal bundle  $N_- \in Pic^0(C^-(\tau))$  through  $(\tilde{g}_\tau)^*N_- \cong (N_+)^{-1}$ .

Taking use of lemma 2.1.5, let  $E_1^\pm, E_2^\pm, \dots, E_8^\pm$  and  $E_9^\pm$  be the exceptional divisors corresponding to the points in  $Z^\pm = \{p_1^\pm, p_2^\pm, \dots, p_9^\pm\}$  respectively. The following is to give the ample line bundles  $L^\pm$  on  $S^\pm(\tau)$  such that  $(L^+.C^+(\tau)) = (L^-.C^-(\tau))$  with  $(\ell_{S^+(\tau)})^*L^- = L^+$ .

**Theorem 2.1.6:** *There exist ample line bundles  $L^\pm$  on  $S^\pm(\tau)$  such that  $(L^+.C^+(\tau)) = (L^-.C^-(\tau))$  with  $(\ell_{S^+(\tau)})^*L^- = L^+$ .*

**Proof of theorem 2.1.6:** Since  $-K_{S^+(\tau)} = [C^+(\tau)] = 3H^+ - \sum_{i=1}^9 E_i^+$  and  $-K_{S^-(\tau)} = [C^-(\tau)] = 3H^- - \sum_{i=1}^9 E_i^-$ ,  $(L^+.C^+(\tau)) = 3d^+ - 9k^+$  and  $(L^-.C^-(\tau)) = 3d^- - 9k^-$ . Therefore, for  $L^+ = d^+ \cdot H^+ - k^+ \cdot \sum_{i=1}^9 E_i^+$  and  $L^- = d^- \cdot H^- - k^- \cdot \sum_{i=1}^9 E_i^-$  with  $k^\pm \geq 2$ ,  $d^+ \geq 3k^+ + 1$  and  $d^- \geq 3k^- + 1$  being integers,  $(L^+.C^+(\tau)) = (L^-.C^-(\tau))$  if and only if  $d^+ - 3k^+ = d^- - 3k^-$ .

Then for  $d^+ = d^-$  and  $k^+ = k^-$ ,  $10 \leq \frac{(d^+)^2}{(k^+)^2} \leq \frac{(d^-)^2}{(k^-)^2}$  with  $k^+ = k^- \geq 2$  and  $d^+ = d^- \geq 3k^+ + 1 = 3k^- + 1$  being integers,  $L^+ = d^+ \cdot H^+ - k^+ \cdot \sum_{i=1}^9 E_i^+$  and  $L^- = d^- \cdot H^- - k^- \cdot \sum_{i=1}^9 E_i^-$  will be ample line bundles on  $S^+(\tau)$  and  $S^-(\tau)$  respectively such that  $(L^+.C^+(\tau)) = (L^-.C^-(\tau))$ . In addition,  $(\ell_{S^+(\tau)})^*L^- = L^+$ . So theorem 2.1.6 is proved.

Now let  $\alpha_\pm$  and  $\beta_\pm$  be smooth loops corresponding to the line segments  $[0, 1]$  and  $[0, \tau]$  on  $C^\pm(\tau)$  respectively. Then let  $e^{\pm p \cdot 2\pi\sqrt{-1}}$  and  $e^{\pm q \cdot 2\pi\sqrt{-1}}$  be the monodromies of  $N_\pm$  along the loops  $\alpha_\pm$  and  $\beta_\pm$  respectively with  $(p, q) \in R^2$ . Taking use of theorem 2.1.4, the following statement is true.

There are tubular neighborhoods  $W^\pm$  of  $C^\pm(\tau)$  biholomorphic to neighborhoods of the zero section in  $N_{C^\pm(\tau)/S^\pm(\tau)}$  such that  $W^+$  is just a copy of  $W$  and  $W^- = \ell_{S^+(\tau)}(W^+)$  for  $\tau \in Y$ .

Then for arbitrary ample line bundles  $L^\pm \rightarrow S^\pm(\tau)$  satisfying  $(L^+.C^+(\tau)) = (L^-.C^-(\tau))$  with  $(\ell_{S^+(\tau)})^*L^- = L^+$ , there exists a unique  $\xi \in \mathbb{C}$  up to modulo  $\langle 1, \tau \rangle$  such that  $g_\xi = \ell_\xi \circ \tilde{g}_\tau$  with  $\ell_\xi$  being a translation induced from  $\mathbb{C} \ni z \rightarrow z + \xi \in \mathbb{C}$  gives the following gluing construction. Here,  $\xi = \frac{1}{b}((p^+ \cdot L^+) - (p^- \cdot L^-))$  with  $b = (L^+.C^+(\tau)) = (L^-.C^-(\tau))$  and  $p^\pm = 3p_0^\pm H^\pm - \sum_{i=1}^9 p_j^\pm E_j^\pm$  as it was computed in [2, Proposition 5.1].

For  $s \in \Delta \setminus \{0\}$  with  $\Delta := \{s \in \mathbb{C} \mid |s| < 1\}$ , let  $M_s^+ \subset S^+(\tau)$  be a copy of  $M_s$ ,  $M_s^- = \ell_{S^+(\tau)}(M_s^+) \subset S^-(\tau)$ ,  $V_s^+ \subset S^+(\tau)$  be a copy of  $V_s$  and  $V_s^- = \ell_{S^+(\tau)}(V_s^+) \subset S^-(\tau)$  with  $f_s : V_s^+ \rightarrow V_s^-$  defined by  $f_s([(z^+, w^+)]) = [(g_\xi(z^+), s/w^+)]$  for  $[(z^+, w^+)] \in V_s^+$ .

The following is the definition for the symmetric K3 surfaces here.

**Definition 2.1.7:** For a K3 surface  $\mathcal{K}$  with an ample line bundle  $L_{\mathcal{K}}$ , if there exist a holomorphic involution  $f_{\mathcal{K}}$  defined on  $\mathcal{K}$  such that  $f_{\mathcal{K}}^* L_{\mathcal{K}} = L_{\mathcal{K}}$ , then  $\mathcal{K}$  is said to be symmetric.

So the gluing construction of symmetric K3 surfaces is as below.

**Theorem 2.1.8:**  $\exists$  sufficient small  $\varepsilon_0 > 0$  such that for any  $s \in \Delta \setminus \{0\}$  and  $0 < |s| < \varepsilon_0$ , through identifying  $V_s^+$  and  $V_s^-$  through the biholomorphic map  $f_s$  defined above, the two open complements  $M_s^\pm$  of the tubular neighborhoods  $W^\pm$  of  $C^\pm(\tau)$  are glued to be a symmetric projective K3 surface  $X_s$ .

**Proof of theorem 2.1.8:** Let  $\ell_{S^+(\tau) \setminus C^+(\tau)}$  be defined as  $\ell_{S^+(\tau) \setminus C^+(\tau)} = \ell_{S^+(\tau)}|_{S^+(\tau) \setminus C^+(\tau)}$  for  $\tau \in Y$ . Since  $\ell_{S^+(\tau)} : S^+(\tau) \rightarrow S^-(\tau)$  is a biholomorphic map with  $\ell_{S^+(\tau)}|_{C^+(\tau)} : C^+(\tau) \rightarrow C^-(\tau)$  being biholomorphic, taking use of [1, Propostion 2.1], there are global holomorphic non-vanishing 2-forms  $\eta_s^\pm$  defined on  $S^\pm(\tau) \setminus C^\pm(\tau)$  with  $\ell_{S^+(\tau) \setminus C^+(\tau)}^*(\eta_s^+) = \eta_s^-$  such that the restrictions of  $\eta_s^\pm$  on two open complements of the tubular neighborhoods of  $C^\pm(\tau)$  can be glued through  $f_s$  to be a global holomorphic non-vanishing 2-form  $\sigma_s$  on  $X_s$ . In addition,  $\sigma_s|_{V_s} = a \cdot \frac{dz \wedge dw}{w}$  for  $a \in \mathbb{C} \setminus \{0\}$ . So  $\sigma_s$  can be assumed to be normalized in this paper. That is to say,  $a = 1$ .

**Lemma 2.1.9:** [2, Theorem 1.2] Let  $L_s$  be the line bundle on  $X_s$  through gluing  $L^\pm$  by  $f_s$  defined above. Then  $\exists$  sufficient small  $\varepsilon_0 > 0$  such that  $L_s$  is ample for any  $s \in \Delta \setminus \{0\}$  and  $0 < |s| < \varepsilon_0$ .

Taking use of lemma 2.1.9, for any  $s \in \Delta \setminus \{0\}$  and  $0 < |s| < \varepsilon_0$ ,  $X_s$  is a K3 surface with an ample line bundle on it. So  $X_s$  is projective for  $s \in \Delta \setminus \{0\}$  and  $0 < |s| < \varepsilon_0$ .

Let  $F_s : X_s \rightarrow X_s$  defined by

$$F_{X_s} = \begin{cases} \ell_{M_s^+}(x), & x \in M_s^+ \\ \ell_{M_s^+}^{-1}(x), & x \in M_s^- \end{cases}$$

with  $\ell_{M_s^+}$  being the restriction of  $\ell_{S^+(\tau)}$  on  $M_s^+$ . Then  $F_{X_s}$  is a holomorphic involution on the K3 surface  $X_s$ . In addition, since  $(\ell_{S^+(\tau)})^* L^- = L^+$ ,  $F_{X_s}^* L_s = L_s$  for  $s \in \Delta \setminus \{0\}$  and  $0 < |s| < \varepsilon_0$ . So  $X_s$  is a symmetric projective K3 surface for  $s \in \Delta \setminus \{0\}$  and  $0 < |s| < \varepsilon_0$ . Then theorem 2.1.8 is proved.

Here, the existence of  $\varepsilon_0$  will also be confirmed in section 3 during the process of the construction of complete Kahler metrics on the quasi-projective varieties  $S(\tau) \setminus C(\tau)$  for  $\tau \in Y$ .

Then the next part is to introduce the corresponding toroidal groups  $U = \mathbb{C}^2/\Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  provided in [2] for the Diphantine pair  $(p, q)$ .

## 2.2. The Toroidal Group and the Tubular Neighborhood.

The following is to show that  $U = \mathbb{C}^2/\Lambda_0$  is a quasi-abelian variety.

**Definition 2.2.1:** [10, Definition 1.1.1] Let  $\Lambda$  be a lattice which is a discrete subgroup of  $\mathbb{C}^n$  for  $n \in \mathbb{N}$ . Then  $\mathbb{C}^n/\Lambda$  is called a toroidal group if there does not exist any non-constant holomorphic function on  $\mathbb{C}^n/\Lambda$ .

The toroidal group is a topological group. What's more, it is an abelian complex Lie group [10].

**Lemma 2.2.2:** [10, Theorem 1.1.4] For  $\mathbb{C}^2/\Lambda$  with  $\Lambda$  a discrete subgroup of  $\mathbb{C}^2$ ,  $\mathbb{C}^2/\Lambda$  is toroidal if and only if there does not exist any  $\sigma \in \mathbb{C}^2 \setminus \{0\}$  such that the scalar product  $\langle \sigma, \lambda \rangle \in \mathbb{Z}$  is integral for all  $\lambda \in \Lambda$ .

Taking use of lemma 2.2.2, the following lemma can be derived. Here, let  $\mathbb{Q}$  represent the set consisting of rational numbers.

**Lemma 2.2.3:** For any real number pair  $(p, q)$  satisfying the Diophantine condition and complex number  $\tau$  with  $Im\tau > 0$ ,  $U = \mathbb{C}^2/\Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  is a toroidal group.

**Proof of lemma 2.2.3:** For  $\sigma = \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix} \in \mathbb{C}^2 \setminus \{0\}$ ,  $\langle \sigma, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rangle = \langle \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rangle = \sigma_2$ ,  
 $\langle \sigma, \begin{pmatrix} 1 \\ p \end{pmatrix} \rangle = \langle \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix} \rangle = \sigma_1 + p\sigma_2$  and  $\langle \sigma, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle = \langle \begin{pmatrix} \sigma_1 \\ \sigma_2 \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle = \sigma_1 + q\sigma_2$ .

If  $q \in \mathbb{Q}$ , since  $(p, q) \in R^2$  satisfies the Diophantine condition,  $q \in \mathbb{Q}$ . Under the condition that  $\sigma_2 \in \mathbb{Z}$  and  $\sigma_1\tau + q\sigma_2 \in \mathbb{Z}$ , then  $\sigma_1 = 0$  or  $\sigma_1 \in \mathbb{C} \setminus R$ . So  $\sigma_1\tau + q\sigma_2 \notin \mathbb{Z}$ .

If  $q \notin \mathbb{Q}$ , under the condition that  $\sigma_2 \in \mathbb{Z}$  and  $\sigma_1\tau + q\sigma_2 \in \mathbb{Z}$ , then  $\sigma_1 \in \mathbb{C} \setminus R$ . So  $\sigma_1\tau + q\sigma_2 \notin \mathbb{Z}$ .

Therefore, there does not exist any  $\sigma \in \mathbb{C}^2 \setminus \{0\}$  so that the scalar product  $\langle \sigma, \lambda \rangle \in \mathbb{Z}$  is integral for all  $\lambda \in \Lambda_0$ .

Taking use of lemma 2.2.2,  $U = \mathbb{C}^2/\Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  is a toroidal group for any satisfying the Diophantine condition. So lemma 2.2.3 is proved.

**Theorem 2.2.4:** [2, Proposition 2.4] For  $(p, q) \in R^2$  satisfying the Diophantine condition and  $\tau \in \mathbb{C}$  with  $\text{Im}\tau > 0$ , every topologically trivial line bundle on the toroidal group defined as  $U = \mathbb{C}^2/\Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  is homogeneous.

**Proof of theorem 2.2.4:** Taking use of the result from [11], there are nine equivalent conditions for the special class of toroidal groups with homogeneous topologically trivial line bundles [11, Theorem]. The following lemma is to show the rewritten condition 8.

**Lemma 2.2.5:** [11] For  $(p, q) \in R^2$  satisfying the Diophantine condition and  $\tau \in \mathbb{C}$  with  $\text{Im}\tau > 0$ , let  $D_0 = (p \ q)$ . Then every topologically trivial line bundle on the toroidal group defined as  $U = \mathbb{C}^2/\Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  is homogeneous if and only if  $\exists c > 0$  &  $a \geq 0$  such that  $\|\sigma^t \cdot D_0 + \delta^t\| \geq ce^{-a|\sigma|}$  for all  $\sigma \in \mathbb{Z} \setminus \{0\}$  and  $\delta \in \mathbb{Z}^2$ .

Since  $(p, q) \in R^2$  satisfying the Diophantine condition and  $\tau \in \mathbb{C}$  with  $\text{Im}\tau > 0$ , there exists  $\alpha > 0$  and  $A > 0$  such that

$$\min_{\mu, \nu \in \mathbb{Z}} |n(p + q\sqrt{-1}) - (\mu + \nu\sqrt{-1})| \geq A \cdot n^{-\alpha}$$

for any  $n \in \mathbb{N}$ .

That is to say,  $\|\sigma^t \cdot D_0 + \delta^t\| = \|\sigma \cdot (p \ q) + \delta^t\| \geq A \cdot e^{-\alpha|\sigma|}$  for all  $\sigma \in \mathbb{Z} \setminus \{0\}$  and  $\delta \in \mathbb{Z}^2$ . Therefore, the nine equivalent conditions for the special class of toroidal groups [11, Theorem] are all satisfied here. That is to say, every topologically trivial line bundle on the toroidal group defined as  $U = \mathbb{C}^2/\Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  is homogeneous. So theorem 2.2.4 is proved.

With the above result, from [11], the complex vector space  $H^1(U, \mathcal{O})$  is finite-dimensional. The following is to show that  $U = \mathbb{C}^2/\Lambda_0$  defined above are quasi-abelian varieties.

Let  $R_{\Lambda_0} := \left\{ x_1 \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} + x_2 \cdot \begin{pmatrix} 1 \\ p \end{pmatrix} + x_3 \cdot \begin{pmatrix} \tau \\ q \end{pmatrix} \mid x_1 \in R, x_2 \in R, x_3 \in R \right\}$  be the  $R$ -span of  $\Lambda_0$ .

What's more, let  $MC_{\Lambda_0} := R_{\Lambda_0} \cap \sqrt{-1}R_{\Lambda_0} = \left\{ \begin{pmatrix} x \\ 0 \end{pmatrix} \mid x \in \mathbb{C} \right\}$  be the maximal  $\mathbb{C}$ -linear subspace of  $R_{\Lambda_0}$ .

**Definition 2.2.6:** [10, Definition 3.1.6] A quasi-abelian variety is a toroidal group  $\mathbb{C}^n/\Lambda$  with an ample Riemann form for the lattice  $\Lambda$ .

In addition, the following is the definition of the ample Riemann form.

**Definition 2.2.7:** [10, Definition 3.1.6] An ample Riemann form  $H_0$  for the discrete subgroup  $\Lambda \subset \mathbb{C}^n$  of complex rank  $n$  is a Hermitian form  $H_0$  on  $\mathbb{C}^n$  such that



1.  $ImH_0|_{\Lambda \times \Lambda}$  is  $\mathbb{Z}$ -valued,

2.  $H_0$  is positive definite on the maximal  $\mathbb{C}$ -linear subspace  $MC_\Lambda$  of  $R_\Lambda$ .

Here,  $R_\Lambda$  is the  $R$ -span of  $\Lambda$ . What's more,  $MC_\Lambda$  is the maximal  $\mathbb{C}$ -linear subspace of  $R_\Lambda$ .

**Theorem 2.2.8:** For  $(p, q) \in R^2$  satisfying the Diophantine condition and  $\tau \in \mathbb{C}$  with  $Im\tau > 0$ ,  $U = \mathbb{C}_{(z, \eta)}^2 / \Lambda_0$  with  $\Lambda_0 = \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ p \end{pmatrix}, \begin{pmatrix} \tau \\ q \end{pmatrix} \rangle$  is a quasi-abelian variety.

**Proof of theorem 2.2.8:** Let  $G = \begin{pmatrix} \frac{1}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$ , then  $G$  defines a hermitian form on  $\mathbb{C}^2$ .  $ImG|_{\Lambda_0 \times \Lambda_0}$  is  $\mathbb{Z}$ -valued and  $G$  is positive definite on the maximal  $\mathbb{C}$ -linear subspace  $MC_{\Lambda_0}$  of  $R_{\Lambda_0}$ . So  $G = \begin{pmatrix} \frac{1}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$  is an ample Riemann form for the lattice  $\Lambda_0$ . Therefore, according to definition 2.2.6,  $U = \mathbb{C}^2 / \Lambda_0$  is a quasi-abelian variety. Then theorem 2.2.8 is proved.

Since  $MC_{\Lambda_0} := R_{\Lambda_0} \cap \sqrt{-1}R_{\Lambda_0} = \left\{ \begin{pmatrix} x \\ 0 \end{pmatrix} \mid x \in \mathbb{C} \right\}$  is the maximal  $\mathbb{C}$ -linear subspace of  $R_{\Lambda_0}$  and  $MC_{\Lambda_0}$  have the complex dimension 1, the toroidal group  $U = \mathbb{C}^2 / \Lambda_0$  is of type 1. [10, Definition 1.2.1]

For  $G = \begin{pmatrix} \frac{1}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$ ,  $ImG$  has rank 2 on  $R_{\Lambda_0}$ . What's more, since the toroidal group  $U = \mathbb{C}^2 / \Lambda_0$  is of type 1, the ample Riemann form  $G = \begin{pmatrix} \frac{1}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$  for  $\Lambda_0$  is said to be of kind  $\frac{2-2 \times 1}{2} = 0$  (see [10, Definition 3.1.12]).

The following theorem will give a fibration of the toroidal group  $U = \mathbb{C}^2 / \Lambda_0$ .

**Theorem 2.2.9:** [10, Theorem 3.1.16] For a toroidal group  $\mathbb{C}^n / \Lambda$  of type  $q$  with  $\Lambda$  being a lattice,  $\mathbb{C}^n / \Lambda$  is a quasi-abelian variety with an ample Riemann form of kind  $k$  if and only if  $\mathbb{C}^n / \Lambda$  has a maximal closed Stein subgroup  $K \cong \mathbb{C}^k \times \mathbb{C}^{*m}$  with  $2k + m = n - q$  and  $(\mathbb{C}^n / \Lambda) / K$  is an abelian variety of dimension  $q + k$ .

Taking use of theorem 2.2.9, the toroidal group  $U = \mathbb{C}^2 / \Lambda_0$  has a maximal closed stein subgroup  $K \cong \mathbb{C}^*$  and  $U / K$  is an abelian variety of dimension 1.

Furthermore, from [2], through the map  $f_U : U = \mathbb{C}_{(z, \eta)}^2 / \Lambda_0 \rightarrow V_{0, \infty}$  defined by sending  $[(z, \eta)] \in U = \mathbb{C}_{(z, \eta)}^2 / \Lambda_0$  to  $[(z, e^{2\pi\sqrt{-1}\eta})] \in V_{0, \infty}$ ,  $U = \mathbb{C}_{(z, \eta)}^2 / \Lambda_0 \cong V_{0, \infty}$  with  $V_{0, \infty}$  defined by

$$V_{b_1, b_2} := \{(z, w) \in \mathbb{C}^2 \mid b_1 < |w| < b_2\} / \sim$$

for  $0 \leq b_1 < b_2 \leq \infty$ , where  $\sim$  is the equivalence relationship generated by

$$(z, w) \sim (z + 1, \exp(p \cdot 2\pi\sqrt{-1}) \cdot w) \sim (z + \tau, \exp(q \cdot 2\pi\sqrt{-1}) \cdot w)$$

with  $\tau \in \mathbb{C}$  and  $Im\tau > 0$ .

From the definitions above,  $V_{0, r} \cup C = W$ . Let  $\pi_W : W \rightarrow C$  be the natural projection defined as  $\pi_W([z, w]) = [z]$ . The next step is to give a theorem for the line bundles on the tubular neighborhoods defined above.

**Theorem 2.2.10:** [2, Proposition 3.3] For any  $L \in \text{Pic}(W)$ ,  $L$  is the pull back line bundle  $(\pi_W)^*(L|_C)$  of the restricted line bundle  $L|_C$  on  $C$ . Here, the assumption is as above.

This theorem is proved in [2] through the toroidal group defined above. The next part is to prove theorem 1.1.

### 3. A Deformation Family of Complete Non-Compact Kahler Manifolds

#### 3.1. A Deformation Family of Quasi-Projective Varieties from $CP^2 \# 9\overline{CP^2}$ .

First of all, the following is the definition for complex analytic family of compact complex manifolds.

**Definition 3.1.1:**[12, Definition 2.8] For a collection of compact complex manifolds  $\{M_t|t \in B\}$  with  $B$  being a domain of  $\mathbb{C}^m$  with  $m \in \mathbb{N}$ ,  $\{M_t|t \in B\}$  is called a complex analytic family of compact complex manifolds if there is a complex manifold  $\mathcal{M}$  and a holomorphic map  $\pi$  from  $\mathcal{M}$  onto  $B$  satisfying the following conditions:

- (1) The rank of the Jacobian matrix of  $\pi$  is equal to  $m$  at every point of  $\mathcal{M}$ .
- (2) For each  $t \in B$ ,  $\pi^{-1}(t) = M_t$  is a compact complex submanifold of  $\mathcal{M}$ .

Here, a deformation family of  $CP^2 \# 9\overline{CP^2}$  over a 9-dimensional complex manifold will be present before the construction of deformation family of open quasi-projective varieties.

However, a deformation family of elliptic curves over 1-dimensional complex manifold should be present at first. The following is to describe the deformation family of elliptic curves.

Taking use of the notations from the Introduction and section 2. Then the following lemma is true.

**Lemma 3.1.2:** Through changing the parameter  $\tau$ , there is a complex analytic family of elliptic curves over a 1-dimensional complex manifold together a complex analytic family of  $CP^2$  over 1-dimensional complex manifolds.

**Proof of lemma 3.1.2:** For  $\tau_0 \in Y := \{\tau \in \mathbb{C} \ \& \ \text{Im}\tau > 0\}$ , let  $U$  be a sufficiently small disc neighborhood of  $\tau_0$  in  $Y$ . Then  $CP^2 \times U$  is a complex manifold. Let  $\pi_{CP^2 \times U} : CP^2 \times U \rightarrow U$  be a projective map.  $CP^2 \times U$  and  $U$  are smooth complex manifolds.  $\pi_{CP^2 \times U}$  is a holomorphic map. The rank of the Jacobian matrix of  $\pi_{CP^2 \times U}$  is equal to 1 at every point of  $CP^2 \times U$ . For each  $t \in U$ ,  $\pi_{CP^2 \times U}^{-1}(t) = CP^2 \times \{t\}$  is a compact complex submanifold of  $CP^2 \times U$ . So  $\pi_{CP^2 \times U} : CP^2 \times U \rightarrow U$  is a complex analytic family of  $CP^2$  over a 1-dimensional complex manifold  $U$ .

Let  $C_0(\tau) = \mathbb{C}/\langle 1, \tau \rangle$  be a smooth elliptic curve embedded in  $\pi_{CP^2 \times U}^{-1}(\tau) = CP^2 \times \{\tau\}$  as  $f_\tau(C_0(\tau)) \times \{\tau\}$  with  $C_0(\tau_0) = C_0$ . Then let  $\tilde{\mathcal{S}} = \{(x, \tau) \in CP^2 \times U | x \in f_\tau(C_0(\tau))\}$ . So  $\tilde{\mathcal{S}}$  is a complex submanifold of  $CP^2 \times U$ . The rank of the Jacobian matrix of  $\pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}}$  is equal to 1 at every point of  $CP^2 \times U$ . For each  $t \in U$ ,  $\pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}}^{-1}(t) = f_t(C_0(t)) \times \{t\}$  is a compact

complex submanifold of  $\tilde{\mathcal{S}}$ . So  $\pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}} : \tilde{\mathcal{S}} \rightarrow U$  is a complex analytic family of elliptic curves over a 1-dimensional complex manifold  $U$ .

Therefore, through changing the parameter  $\tau$  above, there is a deformation family of elliptic curves,  $\pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}} : \tilde{\mathcal{S}} \rightarrow U$ , over a 1-dimensional complex manifold  $U$  embedded in a complex analytic of  $CP^2$ ,  $\pi_{CP^2 \times U} : CP^2 \times U \rightarrow U$ , over a 1-dimensional complex manifold  $U$ . So lemma 3.1.2 is proved.

Through blowing up points on the complex analytic family of  $CP^2$  defined above, the following is to describe a complex analytic family of  $CP^2 \# 9\overline{CP^2}$ .

**Lemma 3.1.3:** [1] For a fixed pair  $(p, q) \in \mathbb{R}^2$  satisfying the Diophantine condition, through changing the positions of the nine blowing up points on  $\pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}}^{-1}(\tau) = f_\tau(C_0(\tau)) \times \{\tau\} \cong \mathbb{C}/\langle 1, \tau \rangle$  with  $\mathbb{C}/\langle 1, \tau_0 \rangle = C_0$  and the complex number  $\tau \in Y$ , there will be a complex analytic family of  $CP^2 \# 9\overline{CP^2}$  over a 9-dimensional complex manifold.

**Proof of lemma 3.1.3:** Taking use of the notation in the proof of lemma 3.1.2, let  $U_\nu$  be sufficient small disc neighborhoods of  $q_\nu$  in the universal cover  $\mathbb{C}$  of  $\mathbb{C}/\langle 1, \tau \rangle = C_0(\tau)$  for  $\nu \in \{1, 2, 3, 4, 5, 6, 7, 8\}$  with  $C_0(\tau_0) = C_0$  and  $\tau \in U$  (a sufficiently small disc neighborhood of  $\tau_0$  in  $Y$ ). Let  $f_{\tilde{\tau}, \nu} : U_\nu \rightarrow U_\nu / \langle 1, \tilde{\tau} \rangle$  be the natural projection for  $\nu \in \{1, 2, 3, 4, 5, 6, 7, 8\}$  and  $\tilde{\tau} \in U$ .

Let  $N_{\tilde{\tau}}$  be the normal bundle of the strict transform for the elliptic curve  $f_{\tilde{\tau}}(C_0(\tilde{\tau})) \times \{\tilde{\tau}\} \cong \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  after blowing up at nine points  $\{\tilde{p}_1 = (f_{\tilde{\tau}}(p_1), \tilde{\tau}), \tilde{p}_2 = (f_{\tilde{\tau}}(p_2), \tilde{\tau}), \dots, \tilde{p}_9 = (f_{\tilde{\tau}}(p_9), \tilde{\tau})\} \subset f_{\tilde{\tau}}(C_0(\tilde{\tau})) \times \{\tilde{\tau}\}$  with  $\{p_1 = f_{\tilde{\tau}, 1}(\hat{p}_1), p_2 = f_{\tilde{\tau}, 2}(\hat{p}_2), \dots, p_8 = f_{\tilde{\tau}, 8}(\hat{p}_8), p_9\} \subset C_0(\tilde{\tau}) = \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  and  $(\hat{p}_1, \dots, \hat{p}_8) \in U_1 \times \dots \times U_8$ . In addition, let  $C(\tilde{\tau}) \times \{\tilde{\tau}\}$  be the strict transform of  $f_{\tilde{\tau}}(C_0(\tilde{\tau})) \times \{\tilde{\tau}\}$ . Here,  $N_{\tilde{\tau}}$  is assumed to satisfy the Diophantine condition.

Then  $N_{\tilde{\tau}}$  is isomorphic to  $\mathcal{O}_{CP^2}(3)|_{C_0(\tilde{\tau})} \otimes \mathcal{O}_{C_0(\tilde{\tau})}(-p_1 - p_2 - \dots - p_8 - p_9) \in Pic^0(C_0(\tilde{\tau}))$ . Furthermore, taking use of notations from section 2, since  $V_s$  is biholomorphic to a topologically trivial annulus bundle over the strict transform of  $f_{\tilde{\tau}}(C_0(\tilde{\tau}))$ ,  $V_s$  is homotopic to  $S_1 \times S_2 \times S_3$  with  $\{S_1, S_2, S_3\}$  being circles and  $S_1 \times S_2$  being  $C^\infty$  sections for  $s \in \Delta \setminus \{0\}$  and sufficient small [2] (also see theorem 2.1.8 in section 2). Then through integrating the global holomorphic form  $\sigma_s$  defined in section 2 on  $S_1 \times S_2$ , the result value is  $q - p \cdot \tilde{\tau}$  [1]. So there is an equation  $9p_0 - \sum_{j=1}^8 p_j - p_9 = q - p \cdot \tilde{\tau} \pmod{\langle 1, \tilde{\tau} \rangle}$  with  $p_0$  being an inflection point on  $C_0(\tilde{\tau})$  derived in [2].

Then for  $\hat{t} = (\tilde{\tau}, \hat{p}_1, \dots, \hat{p}_8) \in T = U \times U_1 \times \dots \times U_8$ ,  $p_9$  can be regarded as a function of  $\hat{t}$ . Let  $p_9 = f_9(\hat{t}) \in C_0(\tilde{\tau}) = \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  for  $\hat{t} \in T$ .

In addition, let  $\pi : CP^2 \times T = CP^2 \times U \times U_1 \times \dots \times U_8 \rightarrow T$  be a projection from  $CP^2 \times T$  to  $T$ . Then  $\mathcal{S}_0 = \tilde{\mathcal{S}} \times U_1 \times \dots \times U_8$  is a submanifold of  $CP^2 \times T$ . Furthermore,  $C_0(\tilde{\tau}) \cong \pi|_{\mathcal{S}_0}^{-1}(\hat{t}) = \pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}}^{-1}(\tilde{\tau}) \times \{(\hat{p}_1, \hat{p}_2, \dots, \hat{p}_8)\} \subset (\pi)^{-1}(\hat{t}) = CP^2 \times \{(\tilde{\tau}, \hat{p}_1, \dots, \hat{p}_8)\}$  for  $\hat{t} = (\tilde{\tau}, \hat{p}_1, \dots, \hat{p}_8) \in T$ .

Through blowing up nine points  $\{f_{\tilde{\tau}}(p_1), f_{\tilde{\tau}}(p_2), \dots, f_{\tilde{\tau}}(p_9)\} \times \{(\hat{\tau}, \hat{p}_1, \dots, \hat{p}_8)\}$  on  $\pi|_{\mathcal{S}_0}^{-1}(\hat{t}) = \pi_{CP^2 \times U}|_{\tilde{\mathcal{S}}}^{-1}(\tilde{\tau}) \times \{(\hat{p}_1, \dots, \hat{p}_8)\}$  with  $\{p_1 = f_{\tilde{\tau}, 1}(\hat{p}_1), p_2 = f_{\tilde{\tau}, 2}(\hat{p}_2), \dots, p_8 = f_{\tilde{\tau}, 8}(\hat{p}_8), p_9\} \subset \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  and  $p_9 = f_9(\hat{t})$  for  $\hat{t} = (\hat{\tau}, \hat{p}_1, \dots, \hat{p}_8) \in T$ , there will be a 11-dimensional complex manifold  $\mathcal{F}$  derived from  $CP^2 \times T$ . Then there exist a proper holomorphic submersion  $\tilde{\pi} : \mathcal{F} \rightarrow T$  from  $\mathcal{F}$  to  $T$  such that  $(\tilde{\pi})^{-1}(\hat{t}) \cong CP^2 \# 9\overline{CP^2}$  is the manifold derived

from blowing up  $\pi^{-1}(\hat{t})$  on the nine points  $\{f_{\tilde{\tau}}(p_1), f_{\tilde{\tau}}(p_2), \dots, f_{\tilde{\tau}}(p_9)\} \times \{(\hat{\tau}, \hat{p}_1, \dots, \hat{p}_8)\}$  with  $\{p_1 = f_{\tilde{\tau},1}(\hat{p}_1), p_2 = f_{\tilde{\tau},2}(\hat{p}_2), \dots, p_8 = f_{\tilde{\tau},8}(\hat{p}_8), p_9\} \subset \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  and  $p_9 = f_9(\hat{t})$  for  $\hat{t} = (\hat{\tau}, \hat{p}_1, \dots, \hat{p}_8) \in T$ .

What's more, let  $\mathcal{S} \subset \mathcal{F}$  be the strict transforms of  $\mathcal{S}_0 \subset CP^2 \times T$ . Then  $\mathcal{S}$  is a complex codimension one submanifold of  $\mathcal{F}$ . In addition,  $(\tilde{\pi}|_{\mathcal{S}})^{-1}(\hat{t}) \cong \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  is the strict transform of  $\pi|_{\mathcal{S}_0}^{-1}(\hat{t})$  after blowing up on the nine points  $\{f_{\tilde{\tau}}(p_1), f_{\tilde{\tau}}(p_2), \dots, f_{\tilde{\tau}}(p_9)\} \times \{(\hat{\tau}, \hat{p}_1, \dots, \hat{p}_8)\}$  with  $\{p_1 = f_{\tilde{\tau},1}(\hat{p}_1), p_2 = f_{\tilde{\tau},2}(\hat{p}_2), \dots, p_8 = f_{\tilde{\tau},8}(\hat{p}_8), p_9\} \subset \mathbb{C}/\langle 1, \tilde{\tau} \rangle$  and  $p_9 = f_9(\hat{t})$  for  $\hat{t} = (\hat{\tau}, \hat{p}_1, \dots, \hat{p}_8) \in T$ .

In conclusion,  $\tilde{\pi} : \mathcal{F} \rightarrow T$  is a complex analytic family of  $CP^2 \# 9\overline{CP}^2$  over a 9-dimensional complex manifold  $T$ . So lemma 3.1.3 is proved.

Secondly, the following is to give a deformation family of quasi-projective varieties.

Here, the fibers of deformation families defined in definition 3.1.1 should be compact. Since there will be a deformation family of open complex manifolds to be confirmed, the definitions of deformation in [4] will be used. Then the corresponding definition for the category of smooth manifolds is as below.

**Definition 3.1.4:** (*The Category of Smooth Manifolds*). *The objects for the category of smooth manifolds are the smooth manifolds. For any objects  $M$  and  $N$ , the morphisms from  $M$  to  $N$  are the maps in the set  $C^\infty(M, N)$ .*

So the definition of deformation family from [4] is as the following.

**Definition 3.1.5:**[4, Definition 2.3] *A deformation family of a complex manifold  $X$  is a holomorphic surjective submersion  $\mathcal{X} \rightarrow D_1$ , where  $D_1$  is a complex disc centred at 0, satisfying:*

(1)  $\pi^{-1}(0) = X$ ,

(2)  $\mathcal{X}$  is locally trivial in the  $C^\infty$  category.

For any  $t \in D_1$ , the fibre  $\pi^{-1}(t)$  is called a deformation of  $X$ .

What's more, the discussion about definition 3.1.5 can be found in [13]. Here, there is a simple corresponding lemma.

**Lemma 3.1.6:** For a compact analytic family  $(\mathcal{M}, B, \varpi)$  with  $B$  being a complex disc centred at 0,  $(\mathcal{M}, B, \varpi)$  is also a deformation family of the complex manifold  $\varpi^{-1}(0)$  satisfying the conditions in definition 3.1.5.

**Proof of lemma 3.1.6:** From the definition 3.1.1, since the rank of the Jacobian matrix of  $\varpi$  is equal to 1 at every point of  $\mathcal{M}$ ,  $\varpi$  is a holomorphic surjective submersion  $\mathcal{M} \rightarrow B$ . What's more,  $\varpi^{-1}(0)$  is a compact complex submanifold of  $\mathcal{M}$ . In addition, the following lemma is from [12].

**Lemma 3.1.7:** [12, Theorem 2.5]  $\forall x \in B, \exists$  a polydisc  $U_x \subset B$  centered at  $x$  with a diffeomorphism  $\Psi_x : \varpi^{-1}(x) \times U_x \rightarrow \varpi^{-1}(U_x)$  such that  $\varpi \circ \Psi_x : \varpi^{-1}(x) \times U_x \rightarrow U_x$  is a projection.

Taking use of lemma 3.1.7,  $\varpi$  is  $C^\infty$  locally trivial on  $B$ . That is to say,  $\mathcal{M}$  is locally trivial in the  $C^\infty$  category. Therefore,  $(\mathcal{M}, B, \varpi)$  is a deformation family of the complex manifold  $\varpi^{-1}(0)$  satisfying the conditions in definition 3.1.5. So for a compact analytic family  $(\mathcal{M}, B, \varpi)$  with  $B$  being a complex disc centred at 0,  $(\mathcal{M}, B, \varpi)$  is also a deformation family of the complex manifold  $\varpi^{-1}(0)$  satisfying the conditions in definition 3.1.5. Then lemma 3.1.6 is proved.

In addition, the following is to introduce the definition of the smooth compactifiable deformation family. The complex model space is defined as below.

**Definition 3.1.8:** [13, Definition 13]  $\forall f_1, \dots, f_k \in \mathcal{O}(D)$  with  $D \subset \mathbb{C}^n$  being a domain and  $n \in \mathbb{N}$  with  $\mathbb{N}$  being the set collecting positive integers, then  $I_D = \mathcal{O}_D f_1 + \mathcal{O}_D f_2 + \dots + \mathcal{O}_D f_k$  is an ideal sheaf. Let  $X := N(f_1, \dots, f_k)$  and  $\mathcal{O}_X = (\mathcal{O}_D / I_D)|_X$ . So the  $\mathbb{C}$ -ringed space  $(X, \mathcal{O}_X)$  is called a complex model space.

The complex space is defined as the following.

**Definition 3.1.9:** [13, Definition 14] For a  $\mathbb{C}$ -ringed space  $(X, \mathcal{O}_X)$ , if it is a Hausdorff space and  $\forall x \in X, \exists$  an open neighborhood  $U_x$  such that the open  $\mathbb{C}$ -ringed subspace  $(U_x, \mathcal{O}_{U_x})$  is isomorphic to a complex model space. Then  $(X, \mathcal{O}_X)$  is called a complex space.

So the following is the definition of the smooth compactifiable deformation family.

**Definition 3.1.10:** [13, Definition 16] For a complex space  $X$ ,  $X$  is compactifiable if there is an open embedding  $\rho : X \rightarrow \tilde{X}$  such that  $\rho(X) = \tilde{X} \setminus D_2$  with  $\tilde{X}$  being a compact complex space and  $D_2$  being a closed analytic subset of  $\tilde{X}$ . Then a compactifiable deformation of  $X$  is a sextuple  $(\mathcal{X}, D_2, Y_1, f, o, j)$  such that

- 1)  $\mathcal{X}$  and  $Y_1$  are complex spaces,
- 2)  $D_2$  is a closed analytic subset of  $\mathcal{X}$  and  $o$  is a point in  $Y_1$ ,
- 3)  $f$  is a proper holomorphic map from  $\mathcal{X}$  to  $Y_1$  such that  $f|_{\mathcal{X} \setminus D_2}$  is flat,
- 4)  $j$  is an open embedding from  $X$  to  $f^{-1}(o)$  with  $j(X) = f^{-1}(o) \setminus D_2$ .

Furthermore, the differentially local triviality can be defined as the following.

**Definition 3.1.11:** [13, Definition 21] For a holomorphic submersion  $g : X \rightarrow \hat{Y}$  with  $X$  and  $\hat{Y}$  being complex manifolds,  $g$  is differentially locally trivial over  $\hat{Y}$  if  $\forall y \in \hat{Y}, \exists$  an open neighborhood  $U_y$  such that  $g|_{g^{-1}(U_y)} \rightarrow U_y$  is differentially isomorphic to the projection  $g^{-1}(y) \times U_y \rightarrow U_y$ .

Then for the compactifiable deformation of  $X$  defined above, if  $f|_{\mathcal{X}\setminus D_2} : \mathcal{X}\setminus D_2 \rightarrow Y_1$  is differentially locally trivial over  $Y_1$ , then  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is said to be differentially trivial along  $Y_1$ .

**Definition 3.1.12:** For a compactifiable complex space  $X$  defined above, assuming that  $X$  is smooth, if  $f|_{\mathcal{X}\setminus D_2}$  is a smooth morphism of complex spaces and each fibre of  $f$  is a complex manifold, then  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a smooth compactifiable deformation.

Taking use of the notations from the proof of lemma 3.1.3, let  $t_0 = (\tau_0, q_1, q_2, \dots, q_8) \in T$ , then  $\{p_1 = f_{\tau_0,1}(q_1), \dots, p_8 = f_{\tau_0,8}(q_8), p_9 = f_9(t_0)\} \subset \mathbb{C}/\langle 1, \tau_0 \rangle = C_0(\tau_0) = C_0$ . Through blowing up the nine points  $\{f_{\tau_0}(p_1), f_{\tau_0}(p_2), \dots, f_{\tau_0}(p_9)\} \subset f_{\tau_0}(C_0) \subset S_0 = CP^2$ ,  $S_0$  can be transformed to  $S \cong CP^2 \# 9\overline{CP}^2$ . In addition, let  $C \subset S$  be the strict transform of  $f_{\tau_0}(C_0)$ . The following is to give a smooth compactifiable deformation of a quasi-projective variety  $X = S \setminus C \subset S \cong CP^2 \# 9\overline{CP}^2$  differentially trivial along  $Y_1 = T$ .

**Theorem 3.1.13:** Let  $X = S \setminus C$ . What's more, let  $\mathcal{X} = \mathcal{F}$ ,  $D_2 = \mathcal{S}$ ,  $Y_1 = T$ ,  $f = \tilde{\pi}$  and  $o = (\tau_0, q_1, \dots, q_8)$ . In addition, define  $j : X \rightarrow f^{-1}(o)$  by  $j(x) = (x, \tau_0, q_1, q_2, \dots, q_8)$  for  $x \in X$ . Then  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a compactifiable deformation of  $X$ .

**Proof of theorem 3.1.13:** Firstly, it is to prove that  $X = S \setminus C$  is compactifiable.

$X$  is a quasi-projective variety.  $f^{-1}(o) \cong S$  is a compact complex manifold. So  $f^{-1}(o)$  is a complex space. What's more,  $f^{-1}(o) \cap D_2 \cong \mathbb{C}/\langle 1, \tau_0 \rangle$  is a closed analytic submanifold of  $f^{-1}(o)$ . Since  $f^{-1}(o) \setminus j(X) = f^{-1}(o) \cap D_2 \cong \mathbb{C}/\langle 1, \tau_0 \rangle$  is closed,  $j(X)$  is an open submanifold of  $f^{-1}(o) \cong S$ . Then  $j$  is an open embedding with  $j(X) = f^{-1}(o) \setminus D_2$ . So  $(f^{-1}(o), f^{-1}(o) \cap D_2, j)$  is a compactification of  $X$ . That is to say,  $X$  is compactifiable.

Secondly, it is to prove that  $D_2$  is a closed analytic subset of  $\mathcal{X}$  and  $f$  is a proper holomorphic map from  $\mathcal{X}$  to  $Y_1$  such that  $f|_{\mathcal{X}\setminus D_2}$  is flat.

Since  $\mathcal{X}$  and  $Y_1$  are complex manifolds,  $\mathcal{X}$  and  $Y_1$  are complex spaces. From the proof of lemma 3.1.3,  $D_2 = \mathcal{S}$  is a codimension 1 complex submanifold of  $\mathcal{X} = \mathcal{F}$ . In addition, since the blow up points are all on  $S_0$ , it is easy to get an open neighborhood for every point in  $\mathcal{F} \setminus \mathcal{S}$ . Therefore,  $\mathcal{X} \setminus D_2 = \mathcal{F} \setminus \mathcal{S}$  should be an open subset of  $\mathcal{F}$ . Then  $D_2 = \mathcal{S}$  is a closed subset of  $\mathcal{X} = \mathcal{F}$ . So  $D_2 = \mathcal{S}$  is a closed analytic subset of  $\mathcal{X} = \mathcal{F}$ .

Furthermore, since  $\tilde{\pi} : \mathcal{F} \rightarrow T$  is the complex analytic family of compact complex manifold  $CP^2 \# 9\overline{CP}^2$ ,  $f = \tilde{\pi}$  is a proper holomorphic map from  $\mathcal{X} = \mathcal{F}$  to  $Y_1 = T$ . So  $f|_{\mathcal{X}\setminus D_2}$  is also proper.

Since  $f|_{\mathcal{X}\setminus D_2}$  is a proper submersion from the complex manifold  $\mathcal{X}\setminus D_2 = \mathcal{F}\setminus \mathcal{S}$  to  $Y_1 = T$ ,  $f|_{\mathcal{X}\setminus D_2}$  is flat.

Finally, it is to prove that  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a compactifiable deformation of  $X$ .

From above  $j : X \rightarrow f^{-1}(o)$  is an open embedding with  $j(X) = f^{-1}(o) \setminus D_2$ . Therefore, with all the proofs above,  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a compactifiable deformation of  $X$ .

In addition, the following statement is true.

**Proposition 3.1.14:**  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a smooth compactifiable deformation differentially trivial along  $Y_1$ .

**Proof of proposition 3.1.14:** Since  $f|_{\mathcal{X} \setminus D_2}$  is a proper submersion from the complex manifold  $\mathcal{X} \setminus D_2 = \mathcal{F} \setminus \mathcal{S}$  to the complex manifold  $Y_1 = T$  and  $f^{-1}(\hat{t}) = (\tilde{\pi})^{-1}(\hat{t}) \cong \mathbb{C}\mathbb{P}^2 \# 9\overline{\mathbb{C}\mathbb{P}^2}$  is a complex manifold for  $\hat{t} \in Y_1 = T$ ,  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a smooth compactifiable deformation.

What's more, the following lemma from Ehresmann indicates that  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is differentially trivial along  $Y_1$ .

**Lemma 3.1.15:** [14, Theorem 9.3] For any proper holomorphic submersion  $g : M \rightarrow N$  with  $M$  and  $N$  being complex manifolds,  $g$  is a locally trivial fibration.

Taking use of lemma 3.1.16, since  $f|_{\mathcal{X} \setminus D_2}$  is a proper submersion from the complex manifold  $\mathcal{X} \setminus D_2 = \mathcal{F} \setminus \mathcal{S}$  to the complex manifold  $Y_1 = T$ ,  $f|_{\mathcal{X} \setminus D_2}$  is a locally trivial fibration. So  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is differentially trivial along  $Y_1$ .

In conclusion,  $(\mathcal{X}, D_2, Y_1, f, o, j)$  is a smooth compactifiable deformation differentially trivial along  $Y_1$ . So proposition 3.1.14 is proved.

Therefore,  $(\mathcal{F}, \mathcal{S}, T, \tilde{\pi}, o, j)$  is a compactifiable deformation of the quasi-projective variety  $S \setminus C$  with  $o = (\tau_0, q_1, q_2, \dots, q_8)$  and  $j : S \setminus C \rightarrow (\tilde{\pi}^{-1}(o))$  defined by  $j(x) = (x, \tau_0, q_1, q_2, \dots, q_8)$  for  $x \in S \setminus C$ .

From theorem 3.1.13 and proposition 3.1.14, the proof of theorem 1.1 is completed. The next part is to construct the complete Kahler metrics on the fibers of the deformation family.

### 3.2. Complete Kahler Metrics on the Fibers of the Deformation Family of Quasi-Projective Varieties.

The following is to construct the complete Kahler metrics on the fibers of the deformation family of quasi-projective varieties.

Taking use of the notations from the introduction and section 2, the complete Kahler metrics will be constructed on  $S(\tau) \setminus C(\tau)$  for  $\tau \in Y$ .

Let  $L = d \cdot H - k \cdot \sum_{i=1}^9 E_i$  with  $k \geq 2$  and  $d \geq 3k + 1$  being integers together with  $10 \leq d^2/k^2$ . Taking use of lemma 2.1.5,  $L$  is an ample line bundle on  $S(\tau)$ . Then there exists  $n \in \mathbb{N}$  such that  $L^n$  and  $L^n \otimes [-C(\tau)]$  are very ample [15]. So let  $\{g_0(x), g_1(x), \dots, g_\alpha(x)\}$  be a basis for the space of holomorphic sections on  $L^n \otimes [-C(\tau)]$  with  $n \in \mathbb{N}$  and  $x \in S(\tau)$ . What's more, let  $s$  be the defining section of  $C(\tau)$  and  $\{\zeta\}$  be a frame for  $L$ . Then  $\{g_0(x)s(x), g_1(x)s(x), \dots, g_\alpha(x)s(x)\}$  are the holomorphic sections on  $L^n$  with value zero on  $C(\tau)$ . Then the hermitian metric  $h_L$  defined on  $L|_{S(\tau) \setminus C(\tau)}$  through the  $C^\infty$  function

$$h_L(x) = \frac{\zeta(x) \cdot \overline{\zeta(x)}}{(|g_0(x) \cdot s(x)|^2 + |g_1(x) \cdot s(x)|^2 + \dots + |g_\alpha(x) \cdot s(x)|^2)^{\frac{1}{n}}}$$

for  $x \in S(\tau) \setminus C(\tau)$  is smooth.

**Lemma 3.2.1:**  $\sqrt{-1}\Theta_{h_L}$  is positive with  $\Theta_{h_L}$  being the curvature form of  $h_L$ .

**Proof of lemma 3.2.1:** Since  $L^n \otimes [-C(\tau)]$  are very ample, taking use of the Kodaira embedding theorem [15], the evaluation map  $\Phi_{L^n \otimes [-C(\tau)]}(x) = [g_0(x) : g_1(x) : \dots : g_\alpha(x)] \in CP^\alpha$  gives a holomorphic embedding of  $S(\tau) \cong \mathbb{C}P^2 \# 9\mathbb{C}P^2$  into  $CP^\alpha$ . In addition, let  $[\xi_0, \xi_1, \dots, \xi_\alpha]$  be the coordinate system of  $CP^\alpha$  coinciding with the image of the evaluation map  $\Phi_{L^n \otimes [-C(\tau)]}$ .

The function  $g([\xi_0 : \xi_1 : \dots : \xi_\alpha]) = \frac{|\xi_i|^2}{|\xi_0|^2 + |\xi_1|^2 + \dots + |\xi_\alpha|^2}$  on  $\{[\xi_0 : \xi_1 : \dots : \xi_\alpha] \in CP^\alpha \mid \xi_i \neq 0\}$  for  $i \in \{0, 1, \dots, \alpha\}$  defines a hermitian metric on the hyperplane line bundle  $\mathcal{O}(1)$ . Then the curvature form for the hermitian metric  $g$  is  $\Theta_g = \partial\bar{\partial}\log(|\xi_0|^2 + |\xi_1|^2 + \dots + |\xi_\alpha|^2)$  for  $[\xi_0 : \xi_1 : \dots : \xi_\alpha] \in CP^\alpha$ . Furthermore, the associate  $(1, 1)$ -form for the Fubini-Study metrics on  $CP^\alpha$  is the positive Kahler form  $\frac{\sqrt{-1}}{2\pi}\partial\bar{\partial}\log(|\xi_0|^2 + |\xi_1|^2 + \dots + |\xi_\alpha|^2)$  for  $[\xi_0 : \xi_1 : \dots : \xi_\alpha] \in CP^\alpha$ . So  $\sqrt{-1}\Theta_g = \sqrt{-1}\partial\bar{\partial}\log(|\xi_0|^2 + |\xi_1|^2 + \dots + |\xi_\alpha|^2)$  is a positive form for  $[\xi_0 : \xi_1 : \dots : \xi_\alpha] \in CP^\alpha$ .

Therefore,  $g$  is a smooth positive hermitian metric on the hyperplane line bundle  $\mathcal{O}(1)$  for the coordinate system  $[\xi_0 : \xi_1 : \dots : \xi_\alpha]$  on  $CP^\alpha$ . Then the pullback metric  $\hat{g} = (\Phi_{L^n \otimes [-C(\tau)]})^*g$  is a smooth positive hermitian metric on the pullback bundle  $(\Phi_{L^n \otimes [-C(\tau)]})^*\mathcal{O}(1) = L^n \otimes [-C(\tau)]$ . So  $\sqrt{-1}\Theta_{\hat{g}}$  is positive with  $\Theta_{\hat{g}}$  being the curvature form of  $\hat{g}$ .

What's more, in [16], Macro Brunella stated that under the condition that the normal bundle  $N_{C(\tau)/S(\tau)}$  satisfied the Diophantine condition,  $K_{S(\tau)}^{-1} = [C(\tau)]$  admits a smooth hermitian metric with semi-positive curvature. That is to say,  $\sqrt{-1}\Theta_{[C(\tau)]}$  is non-negative. Therefore,  $\sqrt{-1}\Theta_{h_L} = \frac{1}{n}(\sqrt{-1}\Theta_{\hat{g}}|_{S(\tau) \setminus C(\tau)} + \sqrt{-1}\Theta_{[C(\tau)]}|_{S(\tau) \setminus C(\tau)})$  is positive. Then lemma 3.2.1 is proved.

In addition, let  $L|_{C(\tau)}$  be the restricted line bundle of  $L$  on  $C(\tau)$ . Furthermore, let  $L^*$  be the pull back bundle of  $L|_{C(\tau)}$  the restricted line bundle on  $V_{0,\infty}$ . Then  $L^*|_W = L|_W$ . Taking use of theorem 2.2.10 and the equivalent condition in [11, Theorem](also see [2]), there exists a corresponding theta line bundle on  $U = \mathbb{C}^2/\Lambda_0$  holomorphically isomorphic to the pull back line bundle  $(f_U)^*(L^*)$ .

Let  $H = \begin{pmatrix} \frac{(L.C(\tau))}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$ . Defining  $\rho : \Lambda_0 \rightarrow U(1)$  through  $\rho(\lambda + \mu) = \rho(\lambda)\rho(\mu)e^{\pi\sqrt{-1}\cdot ImH(\lambda,\mu)}$  for  $\lambda, \mu \in \Lambda_0$ , let  $\alpha_\lambda(x) = \rho(\lambda)e^{nH(x,\lambda) + (\pi/2)\cdot H(\lambda,\lambda)}$  for  $\lambda \in \Lambda_0$  and  $x \in \mathbb{C}^2$ . The theta line bundle on  $U = \mathbb{C}^2/\Lambda_0$  corresponding to  $H = \begin{pmatrix} \frac{(L.C(\tau))}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$  can be defined as  $L_{H,\rho} = (\mathbb{C}_\zeta \times \mathbb{C}^2/\Lambda_0)$  with  $\lambda \cdot (\zeta, x) = (\alpha_\lambda(x) \cdot \zeta, x + \lambda)$  for  $\lambda \in \Lambda_0, \zeta \in \mathbb{C}_\zeta$  and  $x \in \mathbb{C}^2$ .

**Lemma 3.2.2:** The theta line bundles  $L_{H,\rho}$  on  $U = \mathbb{C}^2/\Lambda_0$  is holomorphically isomorphic to the pull back bundle  $(f_U)^*(L^*)$  of line bundle  $L^*$  on  $V_{0,\infty}$  with  $H = \begin{pmatrix} \frac{(L.C(\tau))}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$ .

**Proof of lemma 3.2.2:** For the pull back bundle  $L_{H,\rho}^* = (f_U^{-1})^*L_{H,\rho}$  of the line bundle  $L_{H,\rho}$  on  $V_{0,\infty}$ ,  $(L_{H,\rho}^*.C(\tau)) = (L.C(\tau))$  (see, [2, lemma 3.4]).



What's more, according to [2, lemma 3.4] and [2, proposition 3.5],  $L_{H,\rho}^*$  with  $H = \begin{pmatrix} \frac{(L.C(\tau))}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$  is the unique theta line bundle defined in the formula as above having the same intersection numbers with  $C(\tau)$  as  $L$ . Therefore, the theta line bundles  $L_{H,\rho}$  on  $U = \mathbb{C}_{(z,\eta)}^2/\Lambda_0$  are holomorphically isomorphic to the pull back bundle  $(f_U)^*(L^*)$  of the line bundle  $L^*$  on  $V_{0,\infty}$  with  $H = \begin{pmatrix} \frac{(L.C(\tau))}{Im\tau} & 0 \\ 0 & 0 \end{pmatrix}$ . So lemma 3.2.2 is proved.

Taking use of lemma 3.2.2, the first chern class of  $L^*$  is  $[\frac{(L.C(\tau))}{2Im\tau} \cdot \sqrt{-1}dz \wedge d\bar{z}]$ . That is to say,  $c_1(L|_{C(\tau)}) = [\frac{(L.C(\tau))}{2Im\tau} \cdot \sqrt{-1}dz \wedge d\bar{z}]$ .

According to [17, Lemma 7.31] ( $\partial\bar{\partial}$ -lemma) and section 8.2 in [17], there exist a smooth hermitian metric  $h_{C(\tau)}$  on  $L|_{C(\tau)}$  such that  $\Theta_{h_{C(\tau)}} = \frac{\pi(L.C(\tau))}{Im\tau} dz \wedge d\bar{z}$ . So  $(\pi_W)^*h_{C(\tau)}$  is a smooth hermitian metric on  $L|_W$  such that  $\Theta_{(\pi_W)^*h_{C(\tau)}} = \frac{\pi(L.C(\tau))}{Im\tau} dz \wedge d\bar{z}$ .

For the line bundle  $L|_{C(\tau)}$ , there exist open finite covers  $\{\mathcal{U}_\alpha\}_{\alpha \in I}$  of  $C(\tau)$  and local trivializations  $\Phi : \pi_{L|_{C(\tau)}}^{-1}(\mathcal{U}_\alpha) \rightarrow \mathcal{U}_\alpha \times \mathbb{C}$  mapping  $\pi_{L|_{C(\tau)}}^{-1}(x)$  isomorphically onto  $\{x\} \times \mathbb{C}$  for  $x \in \mathbb{C}$  and  $\alpha \in I$  with  $I$  being an index set. What's more, let  $g_{\alpha\beta}(x) = [\Phi_\alpha \circ (\Phi_\beta)^{-1}]|_{\{x\} \times \mathbb{C}}$  for  $x \in \mathcal{U}_\alpha \cap \mathcal{U}_\beta$  with  $\alpha, \beta \in I$ .

Since the line bundle  $L|_W = (\pi_W)^*L|_{C(\tau)}$ , the pull back open sets  $\{W_\alpha = \pi_W^{-1}(\mathcal{U}_\alpha)\}_{\alpha \in I}$  is an open cover of  $W$ . What's more, there exist local trivializations  $\Psi_\alpha : \pi_{L|_W}^{-1}(W_\alpha) \rightarrow W_\alpha \times \mathbb{C}$  mapping  $\pi_{L|_W}^{-1}(x)$  isomorphically onto  $\{x\} \times \mathbb{C}$  for  $x \in W$  such that  $\Psi_\alpha|_{\pi_{L|_W}^{-1}(\mathcal{U}_\alpha)} = \Phi_\alpha$  for  $\alpha \in I$ . Let  $\tilde{g}_{\alpha\beta}(x) = [\Psi_\alpha \circ (\Psi_\beta)^{-1}]|_{\{x\} \times \mathbb{C}}$  for  $x \in W_\alpha \cap W_\beta$  with  $\alpha, \beta \in I$ .

A singular hermitian metric for a line bundle is defined through local weights [17, Definition 8.7]. For  $\varepsilon > 0$ , let  $\varphi_L^\alpha \in Psh(W_\alpha \setminus \mathcal{U}_\alpha) \cap C^\infty(W_\alpha \setminus \mathcal{U}_\alpha)$  and  $\varphi_{C(\tau)}^\alpha - \log\varepsilon \in Psh(W_\alpha) \cap C^\infty(W_\alpha)$  be the local weights of the metrics  $h_L$  and  $\varepsilon^{-1}(\pi_W)^*h_{C(\tau)}$  with  $Psh(W_\alpha \setminus \mathcal{U}_\alpha)$  and  $Psh(W_\alpha)$  being the set consisting of plurisubharmonic functions on  $W_\alpha \setminus \mathcal{U}_\alpha$  and  $W_\alpha$  for  $\alpha \in I$  respectively.

Through the following theorem,  $\varphi_L^\alpha$  and  $\varphi_{C(\tau)}^\alpha - \log\varepsilon$  can be patched to smooth plurisubharmonic functions on  $W_\alpha$  for  $\alpha \in I$ .

Let  $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_p)$  with  $\gamma_i > 0$  for  $i \in \{1, \dots, p\}$  and  $X$  be a compact complex manifold. What's more, let  $\theta$  be real non-negative smooth function defined on  $R$  supported on  $[-1, 1]$  with  $\int_R \theta(h)dh = 1$  and  $\int_R h\theta(h)dh = 0$ . In addition, let  $M_\gamma(t_1, t_2, \dots, t_p) = \int_{R^p} \max\{t_1 + h_1, \dots, t_p + h_p\} \prod_{1 \leq j \leq p} \gamma_j^{-1} \theta(h_j/\gamma_j) dh_1 dh_2 \dots dh_p$  for  $t_1, t_2, \dots, t_p \in R$  which is called a regularized max function [18].

**Lemma 3.2.3:** [19, Lemma 5.18]  $M_\gamma(t_1, t_2, \dots, t_p)$  defined above satisfying the following properties:

- 1)  $M_\gamma(t_1, t_2, \dots, t_p)$  is smooth and convex in  $R^p$ ;
- 2)  $M_\gamma(t_1 + a, t_2 + a, \dots, t_p + a) = M_\gamma(t_1, t_2, \dots, t_p) + a, \forall a \in R$ .

Let  $h = h_L$  on  $S(\tau) \setminus W$ . What's more, let  $M_{(1,1)}(\varphi_L^\alpha, \varphi_{C(\tau)}^\alpha - \log\varepsilon) = \int_{R^2} \max\{\varphi_L^\alpha + h_1, \varphi_{C(\tau)}^\alpha - \log\varepsilon + h_2\} \prod_{1 \leq j \leq 2} \theta(h_j) dh_1 dh_2$  be the local weights of the metric  $h$  on  $W_\alpha$  for  $\alpha \in I$ .

Let  $r_1$  and  $r_2$  be positive numbers with  $r_1 < r_2 < r$ . Choosing  $\varepsilon$  small enough,  $M_{(1,1)}(\varphi_L^\alpha, \varphi_{C(\tau)}^\alpha - \log\varepsilon) = \int_{R^2} \max\{\varphi_L^\alpha + h_1, \varphi_{C(\tau)}^\alpha - \log\varepsilon + h_2\} \prod_{1 \leq j \leq 2} \theta(h_j) dh_1 dh_2 = \varphi_L^\alpha$  on  $W_\alpha \cap \{(z, w) \in W | r_1 < |w| \}$  for  $\alpha \in I$ . That is to say,  $h = h_L$  on  $\{(z, w) \in W | r_1 < |w| \}$ .

Taking use of condition 1 of theorem 3.1.5,  $M_{(1,1)}(\varphi_L^\alpha, \varphi_{C(\tau)}^\alpha - \log \varepsilon)$  is plurisubharmonic for  $\alpha \in I$ . Furthermore, from condition 2 of theorem 3.2,3,  $h$  is a smooth hermitian metric on  $L$ . Therefore,  $h$  is a smooth hermitian metric on  $L$  with  $\sqrt{-1}\Theta_h \geq 0$  (see section 8.2 in [17] for the definition of singular hermitian metrics).

Furthermore, there exists  $\varepsilon_0 > 0$  satisfying  $\sqrt{\varepsilon_0}r < r_1$  such that  $M_{(1,1)}(\varphi_L^\alpha, \varphi_{C(\tau)}^\alpha - \log \varepsilon) = \int_{\mathbb{R}^2} \max \left\{ \varphi_L^\alpha + h_1, \varphi_{C(\tau)}^\alpha - \log \varepsilon + h_2 \right\} \prod_{1 \leq j \leq 2} \theta(h_j) dh_1 dh_2 = \varphi_L^\alpha$  on  $W_\alpha \cap \{(z, w) \in W \mid |w| < \sqrt{\varepsilon_0}r\}$  for  $\alpha \in I$ . So  $h = \varepsilon^{-1}(\pi_W)^* h_{C(\tau)}$  on  $\{(z, w) \in W \mid |w| < \sqrt{\varepsilon_0}r\}$ .

Let  $\phi_s(t) = (\log(t^2/|s|))^2$  be a smooth function for  $t > 0$  and  $s \in \{s \mid s \in \Delta \setminus \{0\} \text{ \& } |s| < \varepsilon_0\}$  with  $\Delta = \{s \in \mathbb{C} \mid |s| < 1\}$ .

Let  $f(x) = 1$  for  $|x| < r - \frac{r-r_2}{2}$  and  $f(x) = 0$  for  $r - \frac{r-r_2}{2} \leq |x|$ . The standard mollifier  $\eta \in C^\infty(\mathbb{R})$  is defined as

$$\eta(x) := \begin{cases} a \cdot e^{1/(|x|^2-1)} & \text{if } |x| < 1 \\ 0 & \text{if } |x| \geq 1, \end{cases}$$

with  $a = 1/\int_{-1}^1 e^{1/(|x|^2-1)} dx > 0$  [20].

What's more, let  $\eta_\delta(x) := \frac{1}{\delta} \eta(\frac{x}{\delta})$ . Then for  $\delta = \frac{r-r_2}{4}$ , let  $\tilde{f}(x) = \eta_\delta * f(x) = \int_{-\infty}^{\infty} \eta_\delta(x-y)f(y)dy$ . Then  $\tilde{f}$  is a smooth function on  $\mathbb{R}$  with  $\tilde{f}(x) = 1$  for  $|x| < r_2$  and  $\tilde{f}(x) = 0$  for  $r < |x|$ .

Let  $\Psi_s : S(\tau) \setminus C(\tau) \rightarrow \mathbb{R}$  be the smooth function defined by

$$\Psi_s(p) := \begin{cases} \tilde{\phi}_s(|w|), & p = [(z, w)] \in W \setminus C(\tau) \\ \tilde{\phi}_s(r), & p \notin W \end{cases}$$

with  $\tilde{\phi}_s(x) = \tilde{f}(x) \cdot \phi_s(x)$  for  $x > 0$ .

Then following is to give the complete Kahler metrics on  $S(\tau) \setminus C(\tau)$  originating from the complete Kahler metrics described in [2].

**Theorem 3.2.4:** [2] *There exists  $b > 0$  such that the metric is a smooth hermitian metric on  $L|_{S(\tau) \setminus C(\tau)}$  with  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}} > 0$ . What's more, the Kahler form  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}}$  is of the form*

$$\omega|_{W_{\varepsilon_0}} = \frac{\pi(L \cdot C(\tau))}{Im\tau} \cdot \sqrt{-1}dz \wedge d\bar{z} + 2b \cdot \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2}$$

on the set  $W_{\varepsilon_0} = \{(z, w) \in W \setminus C(\tau) \mid |w| < \sqrt{\varepsilon_0}r\}$ . In addition,  $\omega|_{W_{\varepsilon_0}}$  is Ricci-flat. Then the Kahler form  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}}$  will give a complete Kahler metric on  $S(\tau) \setminus C(\tau)$ .

**Proof of theorem 3.2.4:** Firstly, it is to prove that there exists  $b > 0$  such that the metric  $h \cdot e^{-b\Psi_s}$  is a smooth hermitian metric on  $L|_{S(\tau) \setminus C(\tau)}$  with  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}} > 0$ .

On the set  $S(\tau) \setminus W$ , since  $\Psi_s$  are constants,  $\partial\bar{\partial}\Psi_s = 0$ . Then  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}} = \sqrt{-1}\Theta_h + b\sqrt{-1}\partial\bar{\partial}\Psi_s = \sqrt{-1}\Theta_h = \sqrt{-1}\Theta_{h_L} > 0$ .

On the set  $\{(z, w) \in W \setminus C(\tau) \mid r_2 \leq |w| < r\}$ , from the construction of  $\Psi_s$  above, since  $\tilde{\phi}_s(x)$  is smooth for  $s > 0$ , the coefficients of  $\partial\bar{\partial}\Psi_s$  is bounded. What's more, since  $\sqrt{-1}\Theta_h > 0$  on  $S(\tau) \setminus C(\tau)$ , the coefficients of  $\Theta_h$  have a positive lower bound on  $\{(z, w) \in W \setminus C(\tau) \mid r_2 \leq |w| < r\}$ . Then  $\exists$  sufficient small  $b > 0$  such that  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}} = \sqrt{-1}\Theta_h + b\sqrt{-1}\partial\bar{\partial}\Psi_s > 0$ .

On the set  $\{(z, w) \in W \setminus C(\tau) \mid |w| < r_2\}$ ,  $\sqrt{-1}\partial\bar{\partial}\Psi_s = 2\sqrt{-1} \cdot dw \wedge d\bar{w}/|w|^2 > 0$ . So  $\sqrt{-1}\Theta_{h \cdot e^{-b\Psi_s}} = \sqrt{-1}\Theta_h + b\sqrt{-1}\partial\bar{\partial}\Psi_s > 0$ .

Secondly, it is to prove that  $\sqrt{-1}\Theta_{h,e^{-b\Psi_s}}$  is of the form

$$\omega|_{W_{\varepsilon_0}} = \frac{\pi(L.C(\tau))}{Im\tau} \cdot \sqrt{-1}dz \wedge d\bar{z} + 2b \cdot \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2}$$

on the set  $W_{\varepsilon_0} = \{(z, w) \in W \setminus C(\tau) \mid |w| < \sqrt{\varepsilon_0}r\}$ .

On the set  $W_{\varepsilon_0} = \{(z, w) \in W \setminus C(\tau) \mid |w| < \sqrt{\varepsilon_0}r\}$ ,  $\sqrt{-1}\Theta_{h,e^{-b\Psi_s}} = \sqrt{-1}\Theta_h + b\sqrt{-1}\partial\bar{\partial}\Psi_s = \sqrt{-1}\Theta_{\varepsilon^{-1}(\pi_W)^*h_{C(\tau)}} + b\sqrt{-1}\partial\bar{\partial}\Psi_s = \sqrt{-1}\Theta_{(\pi_W)^*h_{C(\tau)}} + 2b \cdot \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2} = \frac{\pi(L.C(\tau))}{Im\tau} \cdot \sqrt{-1}dz \wedge d\bar{z} + 2b \cdot \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2}$ . So the curvature form  $\sqrt{-1}\Theta_{h,e^{-b\Psi_s}}$  gives a complete Kahler metric on  $S(\tau) \setminus C(\tau)$ .

Finally, it is to prove that  $\omega|_{W_{\varepsilon_0}}$  is Ricci-flat on the set  $W_{\varepsilon_0} = \{(z, w) \in W \setminus C(\tau) \mid |w| < \sqrt{\varepsilon_0}r\}$ .

The coefficient matrix of  $\omega|_{W_{\varepsilon_0}}$  on the set  $W_{\varepsilon_0} = \{(z, w) \in W \setminus C(\tau) \mid |w| < \sqrt{\varepsilon_0}r\}$  is  $(g_{ij})_{2 \times 2} = \begin{pmatrix} \frac{\pi(L.C(\tau))}{Im\tau} & 0 \\ 0 & \frac{2b}{|w|^2} \end{pmatrix}$ . Then the determinate of the coefficient matrix is  $\det((g_{ij})_{2 \times 2}) = \frac{2\pi b(L.C(\tau))}{Im\tau \cdot |w|^2}$ . So  $Ric(\omega|_{W_{\varepsilon_0}}) = 0$ . Therefore,  $\omega|_{W_{\varepsilon_0}}$  is Ricci-flat.

In conclusion,  $\exists b > 0$  such that  $\sqrt{-1}\Theta_{h,e^{-b\Psi_s}}$  defines a complete Kahler metric on  $S(\tau) \setminus C(\tau)$ . In addition,  $\sqrt{-1}\Theta_{h,e^{-b\Psi_s}}$  is of the form

$$\omega|_{W_{\varepsilon_0}} = \frac{\pi(L.C(\tau))}{Im\tau} \cdot \sqrt{-1}dz \wedge d\bar{z} + 2b \cdot \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2}$$

on the set  $W_{\varepsilon_0} = \{(z, w) \in W \setminus C(\tau) \mid |w| < \sqrt{\varepsilon_0}r\}$ . Furthermore,  $\omega|_{W_{\varepsilon_0}}$  is Ricci-flat. That is to say,  $\omega|_{W_{\varepsilon_0}}$  corresponds to a Ricci-flat Kahler metric. So theorem 3.2.4 is proved.

Therefore, the quasi-projective variety above tends to be a non-compact complete Kahler manifold. The following is to show that the deformation family of quasi-projective varieties can be easily changed to a deformation family satisfying the conditions in definition 3.1.5.

### 3.3. A Deformation Family of Non-Compact Complete Kahler Manifolds.

Since the fibers are now proved to be non-compact complete Kahler manifolds, the deformation family above tends to be a deformation family of non-compact complete Kahler manifolds over a 9-dimensional complex manifold.

What's more, for  $T$  defined above, there exists a complex disc  $D_3 \subset T$  centred at  $o = (\tau_0, q_1, q_2, q_3, q_4, q_5, q_6, q_7, q_8)$ .

**Corollary 3.3.1:** *Let  $f : T \rightarrow \mathbb{C}^9$  defined by  $f(x) = x - (\tau_0, q_1, q_2, q_3, q_4, q_5, q_6, q_7, q_8)$ . Then  $f \circ \tilde{\pi}|_{\mathcal{F} \setminus S} : \mathcal{F} \rightarrow f(D_3)$  is a deformation family of the non-compact complete Kahler manifold  $j(S \setminus C)$  over a 9-dimensional complex manifold satisfying the conditions in definition 3.1.5.*

**Proof of corollary 3.3.1:** Since  $\tilde{\pi}|_{\mathcal{F} \setminus S}$  is a holomorphic surjective submersion and  $f$  is a bihomorphic map from  $D_3$  to  $f(D_3)$ , then  $f \circ \tilde{\pi}|_{\mathcal{F} \setminus S}$  is also a holomorphic surjective submersion.  $(f \circ \tilde{\pi}|_{\mathcal{F} \setminus S})^{-1}(0) = j(S \setminus C)$ .

What's more, since  $\tilde{\pi}|_{\mathcal{F} \setminus S}$  is locally trivial in the  $C^\infty$  category and  $f$  is a diffeomorphism, then  $f \circ \tilde{\pi}|_{\mathcal{F} \setminus S}$  is also locally trivial in the  $C^\infty$  category.

In conclusion,  $f \circ \tilde{\pi}|_{\mathcal{F} \setminus \mathcal{S}} : \mathcal{F} \rightarrow f(D_3)$  is a deformation family of the non-compact complete Kahler manifold  $j(S \setminus C)$  over a 9-dimensional complex manifold  $f(D_3)$  satisfying the conditions in definition 3.1.5. So corollary 3.3.1 is proved.

Completing the construction of the deformation family of the non-compact complete Kahler manifold, the next section is to give a deformation family of symmetric projective K3 surfaces over a 10 dimensional complex manifold together with symmetric Calabi-Yau metrics on the fibers to complete the proof of theorem 1.2.

#### 4. A Deformation Family of Symmetric Projective K3 Surfaces

##### 4.1. Symmetric Kahler Metrics on the Symmetric Projective K3 Surfaces.

The following is to give symmetric Kahler metrics of the symmetric projective K3 surfaces.

In section 3, the complete Kahler metric is constructed on  $S(\tau) \setminus C(\tau)$  for  $\tau \in Y$ . That is to say, the complete Kahler metrics are constructed on the fibers of a deformation family of  $S(\tau_0) \setminus C(\tau_0)$ .

Taking use of the notations from section 2, the symmetric projective K3 surface is constructed in theorem 2.1.8 through  $S^+(\tau) \setminus C^+(\tau)$  and  $S^-(\tau) \setminus C^-(\tau)$ .

Since  $S^+(\tau) \setminus C^+(\tau)$  is just a copy of  $S(\tau) \setminus C(\tau)$ , let  $\omega^+$  be a copy of  $\omega$  on  $S(\tau) \setminus C(\tau)$  constructed above. What's more, let  $L^+ = d^+ \cdot H^+ - k^+ \cdot \sum_{i=1}^9 E_i^+$  be defined on  $S^+(\tau)$  as a copy of  $L$  in section 3. Then  $L^- = d^- \cdot H^- - k^- \cdot \sum_{i=1}^9 E_i^-$  is an ample line bundle on  $S^-(\tau)$  and  $(\ell_{S^+(\tau)})^* L^- = L^+$ . In addition, let  $h^+$  be defined on  $L^+$  as a copy of  $h \cdot e^{-b\psi_s}$  on  $L$ . Then there is a hermitian metric  $h^-$  defined on  $L^-$  satisfying  $h^+ = (\ell_{S^+(\tau) \setminus C^+(\tau)})^* h^-$  such that  $\omega^- = \sqrt{-1} \Theta_{h^-} = (\ell_{S^+(\tau) \setminus C^+(\tau)})^* \omega^+$  is a Kahler form on  $S^-(\tau) \setminus C^-(\tau)$ .

Through gluing  $V_s^+$  and  $V_s^-$  through  $f_s$ ,  $M_s^-$  and  $M_s^+$  can be glued to a K3 surface  $X_s$  together with a line bundle  $L_s$  glued from the line bundles  $L^+$  and  $L^-$ . Taking use of the result from section 3,  $\exists \varepsilon_0 > 0$ ,  $\omega$  is well defined on  $S(\tau) \setminus C(\tau)$  with the results of theorem 3.2.4 holding such that for  $s \in \{s | s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$  with  $\Delta = \{s \in \mathbb{C} | |s| < 1\}$ ,  $\omega^+$  and  $\omega^-$  can be glued to be a Kahler form  $\omega_s$  on  $X_s$ . That is to say,  $L_s$  is ample for  $s \in \{s | s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$ . So  $X_s$  is projective for  $s \in \{s | s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$  (also see [2, Theorem 1.2]).

**Definition 4.1.1:** Let  $\mathcal{K}$  be a K3 surface with an ample line bundle  $L_{\mathcal{K}}$  and a holomorphic involution  $f_{\mathcal{K}}$  satisfying  $f_{\mathcal{K}}^* L_{\mathcal{K}} = L_{\mathcal{K}}$ . Then if  $\tilde{\omega}$  is a Kahler metric on  $\mathcal{K}$  such that  $f_{\mathcal{K}}^* \tilde{\omega} = \tilde{\omega}$ , then  $\tilde{\omega}$  is said to be a symmetric Kahler form on  $\mathcal{K}$  corresponding to a symmetric Kahler metric.

With the discussion above, the following lemma is true.

**Lemma 4.1.2:** There exists a symmetric Kahler metric on the projective symmetric K3 surface  $X_s$  for  $s \in \{s | s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$ .

**Proof of lemma 4.1.2:** Since  $\omega^- = (\ell_{S^+(\tau) \setminus C^+(\tau)})^* \omega^+$ ,  $\omega_s = F_{X_s}^* \omega_s$  for  $s \in \{s | s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$ .

Then  $\omega_s$  is the symmetric Kahler form needed for  $s \in \{s|s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$ . So there exists a symmetric corresponding Kahler metric on the projective symmetric K3 surface  $X_s$  for  $s \in \{s|s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$ . Then lemma 4.1.2 is proved.

The following is to complete the proof of the theorem 1.2.

#### 4.2. A Deformation Family of Symmetric Projective K3 Surfaces.

Let  $\hat{\Delta} = \{s|s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$ . Then there are manifolds

$$\mathcal{M}^+ = (S(\tau) \times \hat{\Delta}) \setminus \{([z, w]), s \in W \times \hat{\Delta} | |w| \leq \sqrt{|s|}r\} = \{(x, s) \in M_s^+ \times \hat{\Delta}\}$$

and

$$\mathcal{M}^- = \{(\ell_{S+(\tau)}(x), s) | (x, s) \in M_s^+ \times \hat{\Delta}\}.$$

In addition, let

$$\mathcal{V}^+ = \{([z, w]), s \in W \times \hat{\Delta} | \sqrt{|s|}r < |w| < \sqrt{|s|}r\} = \{(x, s) \in V_s^+ \times \hat{\Delta}\}$$

and

$$\mathcal{V}^- = \{(\ell_{S+(\tau)}(x), s) | (x, s) \in V_s^+ \times \hat{\Delta}\}$$

be the submanifolds of  $\mathcal{M}^+$  and  $\mathcal{M}^-$  respectively.

Define  $f : \mathcal{V}^+ \rightarrow \mathcal{V}^-$  by  $f([z, w], s) = ([g_\xi(z), s/w], s) = (f_s([z, w]), s)$  for  $([z, w], s) \in \mathcal{V}^+$ . Then through the map  $f$ ,  $\mathcal{M}^+$  and  $\mathcal{M}^-$  can be glued to a deformation family of symmetric projective K3 surfaces over one dimensional complex manifold  $\hat{\Delta}$  (also see [2] for a more general result).

For each fiber of the smooth compactifiable deformation of  $S(\tau_0) \setminus C(\tau_0)$  over a 9-dimensional complex manifold constructed in theorem 1.1, there is a deformation family of symmetric projective K3 surfaces over one dimensional complex manifold  $\hat{\Delta} = \{s|s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$  can be constructed.

In fact, there is an analogue deformation family of quasi-projective varieties constructed here. Let  $\mathcal{F}^-$  be the result manifold after blowing up  $CP^2 \times T$  at the points  $\bigcup_{(\tilde{\tau}, \hat{p}_1, \hat{p}_2, \dots, \hat{p}_8) \in T} \{(\ell_{CP^2} \circ f_{\tilde{\tau}} \circ f_{\tilde{\tau}, 1}(\hat{p}_1), \tilde{\tau}, \hat{p}_1, \hat{p}_2, \dots, \hat{p}_8), \dots, (\ell_{CP^2} \circ f_{\tilde{\tau}} \circ f_{\tilde{\tau}, 8}(\hat{p}_8), \tilde{\tau}, \hat{p}_1, \hat{p}_2, \dots, \hat{p}_8), (\ell_{CP^2} \circ f_{\tilde{\tau}} \circ f_9(\tilde{\tau}, \hat{p}_1, \hat{p}_2, \dots, \hat{p}_8), \tilde{\tau}, \hat{p}_1, \hat{p}_2, \dots, \hat{p}_8)\}$ .

What's more, let  $\mathcal{S}^-$  be the analogue complex manifold of  $\mathcal{S}$  and  $\tilde{\pi}^- : \mathcal{F}^- \rightarrow T^- = T$  be the map induced by  $\tilde{\pi}$ . Then  $\tilde{\pi}^- : \mathcal{F}^- \rightarrow T^-$  gives a deformation family of  $CP^2 \# 9\overline{CP}^2$  analogue to the deformation family  $\tilde{\pi} : \mathcal{F} \rightarrow T$ . What's more,  $\tilde{\pi}^-|_{\mathcal{F}^- \setminus \mathcal{S}^-} : \mathcal{F}^- \setminus \mathcal{S}^- \rightarrow T^-$  is a deformation family of quasi-projective varieties analogue to  $\tilde{\pi}|_{\mathcal{F} \setminus \mathcal{S}} : \mathcal{F} \setminus \mathcal{S} \rightarrow T$ .

In addition, let  $\tilde{\pi}^+ : \mathcal{F}^+ \rightarrow T^+ = T$  be a copy of the deformation family  $\tilde{\pi} : \mathcal{F} \rightarrow T$  and  $\mathcal{S}^+$  be a copy of  $\mathcal{S}$ . Then the following theorem is true.

**Theorem 4.2.1:** *There is a deformation family of symmetric projective K3 surfaces over a 10-dimensional complex manifold constructed from a smooth compactifiable deformation of  $S(\tau_0) \setminus C(\tau_0)$  over a 9-dimensional complex manifold.*

**Proof of theorem 4.2.1:** Here,  $\tilde{\pi}|_{\mathcal{F} \setminus \mathcal{S}} : \mathcal{F} \setminus \mathcal{S} \rightarrow T$  is a smooth compactifiable deformation of  $S(\tau_0) \setminus C(\tau_0)$  over a 9-dimensional complex manifold.  $\tilde{\pi}^-|_{\mathcal{F}^- \setminus \mathcal{S}^-} : \mathcal{F}^- \setminus \mathcal{S}^- \rightarrow T^-$  is a deformation family of quasi-projective varieties analogue to  $\tilde{\pi}|_{\mathcal{F} \setminus \mathcal{S}} : \mathcal{F} \setminus \mathcal{S} \rightarrow T$  and  $\tilde{\pi}^+|_{\mathcal{F}^+ \setminus \mathcal{S}^+} : \mathcal{F}^+ \setminus \mathcal{S}^+ \rightarrow T^+$  is a copy of  $\tilde{\pi}|_{\mathcal{F} \setminus \mathcal{S}} : \mathcal{F} \setminus \mathcal{S} \rightarrow T$ .

From the discussion above, choosing two corresponding fibers from  $\tilde{\pi}^+ : \mathcal{F}^+ \rightarrow T^+$  and  $\tilde{\pi}^- : \mathcal{F}^- \rightarrow T^-$  respectively, a deformation family of symmetric projective K3 surface over one dimensional complex manifold  $\hat{\Delta} = \{s | s \in \Delta \setminus \{0\} \ \& \ |s| < \varepsilon_0\}$  can be constructed. Therefore, a deformation family of symmetric projective K3 surfaces over a 10-dimensional complex manifold can be constructed through  $\tilde{\pi}|_{\mathcal{F} \setminus \mathcal{S}} : \mathcal{F} \setminus \mathcal{S} \rightarrow T$  and an analogue deformation family. So theorem 4.2.1 is proved.

Therefore, combining the results from lemma 4.1.2 and theorem 4.2.1, the proof of theorem 1.2 is completed.

## References

1. TAKAYUKI KOIKE and TAKATO UEHARA, ‘A gluing construction of K3 surfaces’, *arXiv preprint arXiv:1903.01444*. (2019).
2. TAKAYUKI KOIKE and TAKATO UEHARA, ‘A gluing construction of projective K3 surfaces’, *Épjournal de Géométrie Algébrique*. 6 (2022).
3. HANS-JOACHIM HEIN, ‘Complete Calabi-Yau metrics from  $\mathbb{C}P^2\#9\overline{\mathbb{C}P^2}$ ’, *arXiv preprint arXiv:1003.2646*. (2010).
4. ELIZABETH GASPARIM and FRANCISCO RUBILAR, ‘Deformations of noncompact Calabi–Yau manifolds, families and diamonds’, *arXiv preprint arXiv:1908.09045*. (2019).
5. JULIUS ROSS and DAVID WITT NYSTRÖM, ‘Homogeneous Monge–Ampère equations and canonical tubular neighbourhoods in Kähler geometry, preprint (2014)’, *International Mathematics Research Notices* 2017.23. (2017): 7069–7108.
6. VLADIMIR IGOREVICH ARNODL’D, ‘Bifurcations of invariant manifolds of differential equations and normal forms in neighborhoods of elliptic curves’, *Funktsional’nyi Analiz i ego Prilozheniya*. 10.4 (1976): 1–12.
7. TETSUO UEDA, ‘On the neighborhood of a compact complex curve with topologically trivial normal bundle’, *Journal of Mathematics of Kyoto University*. 22.4 (1982): 583–607.
8. ROBIN HARTSHORNE, ‘Graduate texts in mathematics’, *Algebraic Geometry*. 52 (1977).
9. TOMASZ SZEMBERG and HALSZKA TUTAJ-GASIŃSKA, ‘General blow-ups of the projective plane’, *Proceedings of the American Mathematical Society*. 130.9 (2002): 2515–2524.
10. YUKITAKA ABE and KLAUS KOPFERMANN, eds, ‘Toroidal groups: line bundles, cohomology and quasi-Abelian varieties’, *Berlin, Heidelberg: Springer Berlin Heidelberg*. 2001.
11. CHRISTIAN VOGT, ‘Line bundles on toroidal groups’, (1982): 197–215.
12. KUNIHICO KODAIRA, ‘Complex manifolds and deformation of complex structures’, *New York, NY: Springer New York*. 1986. 28–75.
13. EDOARDO BALLLICO, ELIZABETH GASPARIM and FRANCISCO RUBILAR, ‘20 Open questions about deformations of compactifiable manifolds’, *São Paulo Journal of Mathematical Sciences*. (2020): 1–21.
14. CLAIRE VOISIN, ‘Hodge theory and complex algebraic geometry. I, Translated from the French original by Leila Schneps’, *Cambridge Studies in Advanced Mathematics*. 76.11 (2002): 3.
15. PHILLIP GRIFFITHS and JOSEPH HARRIS, ‘Principles of algebraic geometry’, *John Wiley & Sons*. 2014.
16. MARCO BRUNELLA, ‘On Kähler surfaces with semipositive Ricci curvature’, *Riv. Mat. Univ. Parma*. 1.2 (2010): 441.
17. VINCENT GUEDJ and AHMED ZERIAHI, ‘Degenerate complex Monge–Ampère equations’, 2017.
18. ALEXANDRE SUKHOV, ‘Regularized maximum of strictly plurisubharmonic functions on an almost complex manifold’, *arXiv preprint arXiv:1303.5312*. 2013.
19. JEAN-PIERRE DEMAILLY, ‘Complex analytic and differential geometry’, *Grenoble: Université de Grenoble I*. 1997.
20. LAWRENCE C. EVANS, ‘Partial differential equations’, *Vol. 19. American Mathematical Society*. 2022.

Fan Xu

Graduate School of Mathematics,

Nagoya University.

Furocho, Chikusaku, Nagoya, Japan,

464-8602

fan.xu.e6@math.nagoya-u.ac.jp