THE MICROLOCAL RIEMANN-HILBERT CORRESPONDENCE FOR COMPLEX CONTACT MANIFOLDS

LAURENT CÔTÉ, CHRISTOPHER KUO, DAVID NADLER, AND VIVEK SHENDE

ABSTRACT. Kashiwara showed in 1996 that the categories of microlocalized D-modules can be canonically glued to give a sheaf of categories over a complex contact manifold. Much more recently, and by rather different considerations, we constructed a canonical notion of perverse microsheaves on the same class of spaces. Here we provide a Riemann-Hilbert correspondence.

1.	Intro	oduct	ion	2
2.	Twis 2.1. 2.2. 2.3. 2.4. 2.5.	ted sl Line Twis Twis Mic Mic	heaves and microlocalization earity sting sted sheaves rolocalization for twisted sheaves rolocalizing integral transforms	4 4 5 7 9
3.	Integ 3.1. 3.2. 3.3. 3.4.	gral t Mas Pola Som Fror	ransforms and Maslov dataslov dataslov dataurization-to-kernelne remarks on principal bundlesn polarization-to-kernel to Maslov-to-kernel	12 13 14 18 19
4.	Verd	ier d	uality on contact manifolds	21
5.	Micr	oshe	aves on complex contact manifolds	23
6.	Mod 6.1. 6.2. 6.3. 6.4. 6.5.	ules Alge Line Twis Twis The	over twisted differential operators ebroids $ebroids$ <	25 26 26 27 28 29
7.	The 7.1. 7.2. 7.3. 7.4.	canor Mic Fror \mathcal{E} -m The	nical sheaf of microdifferential operators rolocal operators	30 30 31 33 35
Ap	pendix	κ Α.	Recollections from Morita theory	39
Ap	pendix	B.	Recollections from algebraic topology	40
Ap	pendix	к С.	Recollections from symplectic and contact geometry	43
			References	46

1. INTRODUCTION

The Riemann-Hilbert correspondence, originating in the question of the existence of Fuchsian differential equations whose solutions have prescribed monodromy, takes as its modern form the assertions that (1) the derived solution functor carries holonomic \mathcal{D} -modules on an algebraic variety to perverse sheaves [19] and (2) after restricting to regular holonomic \mathcal{D} -modules, this is an equivalence of categories [20, 32].

From the beginning, this correspondence was understood to have a microlocal character. In particular, Kashiwara had previously introduced the characteristic variety of a \mathcal{D} -module and generalized the Cauchy-Kowalevskya theorem: the characteristic variety of the \mathcal{D} -module records the obstructions to propagation of its solutions [23]. That is, in the later terminology of Kashiwara and Schapira, the solution functor carries the characteristic variety of a holonomic \mathcal{D} -module to the microsupport of its sheaf of solutions [28, Theorem 11.3.3].

Both the notion of \mathcal{D} -module and sheaf 'microlocalize' over the cotangent bundle.¹ On the one hand, for \mathcal{D} -modules, there is a sheaf \mathcal{E} of 'microlocal differential operators' over the cotangent bundle, whose pushforward to the base recovers \mathcal{D} . One can then introduce the sheaf of \mathcal{E} -modules [38, 25], whose pushforward to the base likewise recovers the sheaf of \mathcal{D} -modules. Moreover, the notions of support coincide: the support of an \mathcal{E} -module is the characteristic variety of the corresponding \mathcal{D} -module [25, Theorem 7.27].

On the other hand, for sheaves, the notion of microsupport gives a formal microlocalization as developed by Kashiwara and Schapira [28]. Namely, given the notion of microsupport $ss(F) \subset T^*X$ for a sheaf F on a manifold X, one can form the sheaf of categories μ sh over T^*X by sheafifying the presheaf of categories

$$\mu \mathrm{sh}^{pre}(U) := \mathrm{sh}(X) / \{F \,|\, ss(F) \cap U = \emptyset\}$$

It is not difficult to show that $\mu sh(T^*X) = sh(X)$, and more generally, the pushforward of the sheaf of categories μsh recovers the sheaf of categories sh. Moreover, the notions of support coincide: the support of a microsheaf matches the microsupport of the corresponding sheaf.

The work of Andronikov [1] and Waschkies [43] establishes a compatible microlocalization of the Riemann-Hilbert correspondence: the derived solution functor induces an equivalence between regular holonomic \mathcal{E} -modules with perverse microsheaves [43, Theorem 3.6.5]. The new content in this result occurs away from the zero-section, hence on the projectivized cotangent bundle \mathbb{P}^*X viewed as a complex contact manifold.

The main theorem of this paper is a globalization of this microlocal equivalence.

Theorem 1.1 (7.15). Let V be a complex contact manifold. There is a canonical equivalence

$$\mu \mathrm{RH}_V : \mathbb{P}erv_V \xrightarrow{\sim} \mathcal{E}_V - \mathfrak{M}od_{rh}$$

between perverse microsheaves on V (in the sense of [8]) and regular holonomic \mathcal{E}_V -modules (in the sense of [24]) extending the microlocal Riemann–Hilbert correspondence.

Here the domain category $\mathbb{P}erv_V$ of perverse microsheaves on V was introduced in our prior work [8]. It builds upon the works [41, 33] which construct a sheaf of microsheaves $\mu sh_{V,\xi}$ on any *real* contact manifold V equipped with 'Maslov data' ξ . (A couple of remarks: this construction did not proceed via gluing, but rather via high-codimensional embeddings; as a major application, it allows for the expression of Fukaya categories of Weinstein manifolds in terms of microlocal sheaf

¹As a default, we will use \mathcal{D} -module to mean an object of the abelian category, but use sheaf to mean an object of the derived category.

theory [12, 14, 13].) We subsequently showed in [8] that on a complex contact manifold V, there is a canonical sheaf of perverse microsheaves $\mathbb{P}erv_V$, locally agreeing with the perverse microsheaves of Andronikov [3] and Waschkies [42]. This ultimately relies on the core observation: the simpleconnectedness of the *complex* symplectic group provides canonical Maslov data ξ for which we can define a canonical abelian subsheaf $\mathbb{P}erv_V \subset \mu \operatorname{sh}_{V\xi}$.

The target category $\mathcal{E}_V - \mathfrak{M}od_{rh}$ of of regular holonomic \mathcal{E}_V -modules was introduced by Kashiwara [24]. It is based on the key insight: while there are many possible choices for a global theory of \mathcal{E} -modules, requiring compatibility with Verdier duality leads to a canonical choice. Regular holonomic modules are characterized microlocally, so within all such \mathcal{E}_V -modules, there is a canonical abelian subsheaf $\mathcal{E}_V - mod_{rh}$ of regular holonomic \mathcal{E}_V -modules. Already, Kashiwara asked for a description of the "stack of 'perverse sheaves on Z', which is equivalent to the stack $\operatorname{Reg}(Z)$ of regular holonomic systems on Z" [24] (here Kashiwara's Z is our complex contact manifold V). Our Theorem 1.1 provides an answer.

Remark 1.2. Theorem 1.1 bears directly on geometric representation theory. For symplectic resolutions and other complex exact symplectic manifolds, one can identify \mathcal{E} -modules on their contactizations with modules over their deformation quantization [26, 7, 6]. Thus Theorem 1.1 says these categories are equivalent to microlocal sheaves, and hence amenable to topological arguments. Conversely, since [40], there has been some interest in the Fukaya categories of such target spaces. After [13], said categories admit microsheaf descriptions, and, after the present article, comparisons to \mathcal{E} -module categories.

In the rest of the introduction, we provide an overview of the strategy and steps of the proof of Theorem 1.1.

To pursue gluing local categories of microsheaves and \mathcal{E} -modules, recall for both sheaves and \mathcal{D} -modules, functors are given by integral transforms. For example, given manifolds M, N and a sheaf $K \in sh(M \times N)$, there is a functor $sh(M) \to sh(N)$ given by $F \mapsto K_!F := \pi_{N!}(K \otimes \pi_M^*F)$. Moreover, microsupport is naturally compatible with such operations so they induce a functor on microsheaves. In particular, when $K_!$ is invertible, notably with ss(K) (away from the zero section) the graph of a conic symplectomorphism ϕ , one obtains an equivalence $K_! : \phi^* \mu sh \simeq \mu sh$. In this case, one says K is a sheaf quantization of ϕ . One similarly arrives at the notion of \mathcal{D} -module quantization.

The gluing data between local categories of microsheaves and \mathcal{E} -modules are such sheaf and \mathcal{D} -module quantizations. Suppose (falsely) that there were *unique* such quantizations of any given conic symplectomorphism.² Then on any contact manifold V, one could obtain a canonical sheaf of microsheaves μsh_V as follows: choose any cover of V by Darboux charts, identify these with balls in cosphere bundles, pull back microsheaves from the cosphere bundles, and glue these together via the putatively unique quantizations of the change-of-charts contactomorphisms. One could similarly construct a canonical sheaf of \mathcal{E} -modules.

But in fact, the quantization of a conic symplectomorphism ϕ is not unique. For sheaves, the freedom was classified in [17]; and for \mathcal{D} -modules, in [10]. In general, a quantization is given by the choice of 'brane structure' on the graph of ϕ . As a first step in the proof of Theorem 1.1, we explain in Section 3 how the global 'Maslov data' of of [33] translates into the local gluing data of sheaf quantizations for overlaps of charts. This turns out to be a delicate exercise, and moreover, must be done in the general setting of twisted sheaves as recalled in Section 2. This generality

²This generally false supposition does in fact hold for sheaves with coefficients in the 1-periodicization of $\mathbb{Z}/2\mathbb{Z}$ -mod.

is necessary to match the canonical orientation datum in [8] with the corresponding twisting by half-forms in [24].

After detailing the above global-to-local translation, we arrive at the key calculation which motivates the rest of the paper: the verification that the resulting microsheaf kernels for gluing $\mathbb{P}erv_V$ commute appropriately with Verdier duality (Proposition 4.3). Once this is established, we are able to proceed directly as follows. First, we apply the existing microlocal Riemann-Hilbert [1, 43] to the kernels to obtain kernels for gluing \mathcal{E} -modules, in particular regular holonomic \mathcal{E} -modules. Then we recall a characterization of Polesello [34, Theorem 3.3], recalled in Theorem 7.9 in the text, that uniquely distinguishes Kashiwara's quantization [24, Theorem 2] based on its self-duality. Finally, we match this criterion with the Verdier duality we have established for the microsheaf kernels, and thus conclude the gluing data coincide.

Acknowledgements. We thank David Ayala, Sam Gunningham, Mikhail Kapranov, Motohico Mulase, Kevin Sackel, Pierre Schapira, and Filip Živanović for helpful conversations.

Part of this work was conducted while L.C. was supported by NSF grant DMS-2305257, and while LC was a member of the Max Planck Institute for Mathematics. C.K. was partially supported by VILLUM FONDEN grant 37814, and also NSF grant DMS-1928930 when in residence at the SLMath during Spring 2024. D.N. was supported by NSF grant DMS-2101466. V.S. was supported by Villum Fonden Villum Investigator grant 37814, Novo Nordisk Foundation grant NNF20OC0066298, and Danish National Research Foundation grant DNRF157.

Notation and conventions. Throughout the paper, C shall denote a stable presentable symmetric monoidal category and $Pic(\mathcal{C})$ shall denote the subcategory of \mathcal{C} consisting of invertible objects with respect to the symmetric monoidal product. Whenever we discuss microlocal sheaf theory, we always implicitly fix coefficients C, which will often be left unspecified. For Riemann-Hilbert, the relevant choice is $C = \mathbb{C} - mod$. Given a manifold M, we let $\dot{T}^*M := T^*M - 0_M$.

Because the set theoretic product of two contact manifolds is not contact, when writing $V \times V'$ for two contact manifolds V, V', we will mean the contact manifold

$$V \times V' \coloneqq (\hat{V} \times \hat{V'}) / \mathbb{R}_+$$

obtained by quotienting the product of their symplectizations along the diagonal \mathbb{R}_+ -action. A similar notation will be used for morphisms in contact geometry.

2. TWISTED SHEAVES AND MICROLOCALIZATION

Here we recall (in somewhat updated language) the notion of 'twisted sheaves' from [22], and the direct generalization to this context of the microlocal sheaf theory of [28].

Given a topological space X, we denote the category of sheaves on X by sh(X; C), or just sh(X). The assignment $X \supseteq U \mapsto sh(U)$ is itself a sheaf of categories on X which we denote by sh_X .

2.1. Linearity. Let $(\mathcal{C}, \otimes, 1_{\mathcal{C}})$ be a symmetric monoidal category and X a topological space admitting good covers (all opens and all non-empty intersections of finitely-many opens are contractible). Consider the constant presheaf of categories which assigns to $U \subseteq X$ the category C. The sheafification of this presheaf is the sheaf Loc_X , which assigns to $U \subseteq X$ the category Loc(U) of C-valued local systems on U. Explicitly, this is because the presheaf map which is defined by

$$\mathcal{C} \to \operatorname{Loc}(U)$$

 $c \mapsto c_U,$

where c_U is the constant local system on U with stalk c, is an isomorphism on an open cover by the locally contractible assumption.

Assume further that C is presentable. One can form the category \Pr_{st}^{L} of C-linear presentable categories with C-linear colimit-preserving functors (or equivalently functors which are left adjoints). The category \Pr_{st}^{L} admits a symmetric monoidal structure \otimes , sometimes referred as the stable presentable product, and the unit object is given by C. We will suppress this notation and stop emphasizing C-linearity when it is clear from the context.

We shall be interested in the category $\operatorname{sh}(X; \operatorname{Pr}_{st}^{L})$ of $\operatorname{Pr}_{st}^{L}$ -valued sheaves. An important object in this category is sh_X , the sheaf of \mathcal{C} -valued sheaves on X. The category $\operatorname{sh}(X; \operatorname{Pr}_{st}^{L})$ inherits the tensor product structure from $\operatorname{Pr}_{st}^{L}$. That is, for $\mathcal{A}, \mathcal{B} \in \operatorname{sh}(X; \operatorname{Pr}_{st}^{L})$, there is a $\operatorname{Pr}_{st}^{L}$ -valued sheaf $\mathcal{A} \otimes \mathcal{B}$ which is given by the sheafification of the presheaf

$$U \mapsto \mathcal{A}(U) \otimes \mathcal{B}(U).$$

Example 2.1. The discussion from the above paragraph implies that there is an action, for any open $U \subseteq X$,

$$\operatorname{Loc}(U) \otimes \mathcal{A}(U) \to \mathcal{A}(U)$$

induced by the composition

$$\operatorname{Loc}(U) \otimes \mathcal{A}(U) \to (\operatorname{Loc}_X \otimes \mathcal{A})(U) = \mathcal{A}(A).$$

Here the later identification is induced by the sheafifying the presheaf morphism which, on U, is given by the C-linear structure $\mathcal{C} \otimes \mathcal{A}(U) \xrightarrow{\sim} \mathcal{A}(U)$ and the fact that sheafification commutes with tensor product.

Example 2.2. When $\mathcal{A} = \operatorname{sh}_X$, this recovers the usual action of local system on sheaves. That is, for a local system $\mathcal{L} \in \operatorname{Loc}_X(U)$ and section $\mathcal{F} \in \mathcal{A}(U)$, we may form the sheaf $\mathcal{L} \otimes \mathcal{F} \in \mathcal{A}(U)$.

2.2. Twisting.

Definition 2.3. A twisting \mathcal{P}_{η} on X is a local system whose stalks are given by $Pic(\mathcal{C})$, the subcategory of invertible objects in \mathcal{C} , which is particular is a(n) (∞ -)group. Such local systems correspond to maps of the form

(1)
$$\eta: X \to BPic(\mathcal{C})$$

and pulling back along $(\{*\} \rightarrow BPic(\mathcal{C}))$ recovers \mathcal{P}_{η} .

An alternative way to view \mathcal{P}_{η} is that it is a (∞) -principal bundle. Recall the situation one category level down: one can twist a rank k vector bundle E with a principal G-bundle after fixing a representation $\rho : G \to GL(k; \mathbb{R})$ by forming $E \times^{\rho} G$. In our case, the most natural category acted by $Pic(\mathcal{C})$ is \mathcal{C} itself where the action is given by

$$i: Pic(\mathcal{C}) \to Aut(\mathcal{C})$$

 $c \mapsto c \otimes (-)$

as invertible objects.

Definition 2.4. The sheaf $\operatorname{Loc}_X^{\eta}$ of η -twisted local systems is the sheafification of the presheaf $(U \mapsto \mathcal{C} \times^i \mathcal{P}_{\eta}(U))$. Here the category $\mathcal{C} \times^i \mathcal{P}_{\eta}(U)$ is given by the colimit

$$\operatorname{colim}\left(\ldots \rightrightarrows \mathcal{C} \times \operatorname{Pic}(\mathcal{C}) \times \operatorname{Pic}(\mathcal{C}) \times \mathcal{P}_{\eta}(U) \rightrightarrows \mathcal{C} \times \operatorname{Pic}(\mathcal{C}) \times \mathcal{P}_{\eta}(U) \rightrightarrows \mathcal{C} \times \mathcal{P}_{\eta}(U)\right)$$

Note that the assumption of local contractibility implies that for small open sets U, $\mathcal{P}_{\eta}(U) = Pic(\mathcal{C})$, so, on such open sets, sheafification is not needed. As a result, one can compute Loc_X^{η} by choosing a cover $\{U_{\alpha}\}$ with trivializations $f_{\alpha} : \mathcal{P}_{\eta}|_{U_{\alpha}} = Pic(\mathcal{C})_{U_{\alpha}}$, similarly to the computation in differential geometry for principal G-bundles and associated vector bundles. Then Loc_X^{η} can be computed by the limit

(2)
$$\lim \left(\prod_{\alpha} \operatorname{Loc}_{U_{\alpha}} \rightrightarrows \prod_{\alpha\beta} \operatorname{Loc}_{U_{\beta\alpha}} \rightrightarrows \cdots \right)$$

where the restriction maps are appropriately twisted by the chosen trivialization f_{α} 's. This justifies the name 'twisted local systems'.

Analogously to how we can tensor a vector bundle by a line bundle, twisted local systems act on sheaves of C-linear presentable categories.

Definition 2.5. For $\mathcal{A} \in \operatorname{sh}(X; \operatorname{Pr}_{st}^{\operatorname{L}})$, we set $\mathcal{A}^{\eta} \coloneqq \mathcal{A} \otimes \operatorname{Loc}_{X}^{\eta}$.

The following lemma will be used when studying twisted sheaves in the next Subsection 2.3

Lemma 2.6. Assume, for any open set $U \subseteq X$, the category $\mathcal{A}(U)$ is dualizable and, for any inclusion $U \subseteq V$, the restriction map $i_{U,V}^* : \mathcal{A}(V) \to \mathcal{A}(U)$ is limit-preserving in addition to colimit-preversing. Then, for any $U \subseteq X$, the category $\mathcal{A}^{\tau}(U)$ is dualizable for all $U \subseteq X$.

Proof. For any cover $\{U_{\alpha}\}$ of U, the category $\mathcal{A}^{\tau}(U)$ is computed as the limit $\mathcal{A}^{\tau}(U) = \lim \mathcal{A}^{\tau}(U_{\alpha})$. The limit-preserving assumption implies that the restriction map $i_{U,V}^*$ has a left adjoint i_{U,V_1} and $\mathcal{A}^{\tau}(U)$ can be computed as a colimit

$$\mathcal{A}^{\tau}(U) = \operatorname{colim} \mathcal{A}^{\tau}(U_{\alpha})$$

where the transition maps are now given by the left adjoints i_1 's. Here we use the equivalence $\Pr_{st}^{R} = \Pr_{st}^{L^{op}}$. Because of local contractibility, we can assume the U_{α} 's are contractible and thus $\mathcal{A}^{\tau}(U_{\alpha}) = \mathcal{A}(U_{\alpha})$. The proposition then follows from [11, Proposition 6.3.4].

Example 2.7. To recover the notion of twists in [22, 3.14], recall that if R is a (discrete) ring and C = R - mod, then $Pic(C) = \mathbb{Z} \times B(R^{\times})$. Fix a cover $\{U_{\alpha}\}$ and a trivialization

$$f_{\alpha}: \mathcal{P}_{\eta}|_{U_{\alpha}} \xrightarrow{\sim} U_{\alpha} \times \mathbb{Z} \times B(R^{\times})$$

for each α . On the double overlaps $U_{\beta\alpha}$, the composition $g_{\beta\alpha} \coloneqq f_{\beta}f_{\alpha}^{-1}$ corresponds to a map which (by abuse of notation) we still denote by

$$g_{\beta\alpha} \in \operatorname{Map}(U_{\beta\alpha}, Pic(\mathcal{C})) = H^0(U_{\beta\alpha}; \mathbb{Z}) \times \operatorname{Map}(U_{\beta\alpha}, B(R^{\times}))$$

Thus, upon linearization by $\operatorname{Loc}(X)^{\eta} := \operatorname{Loc}(X) \times_{\mathcal{C}} \mathcal{P}_{\eta}$, the $g_{\beta\alpha}$ twists the local systems by the identification $L_{\beta\alpha}[n_{\beta\alpha}] \otimes (-)$ on the double overlap where $n_{\beta\alpha} \in \mathbb{Z}$ and $L_{\beta\alpha}$ is a local system with stalk R. The fact that $g_{\gamma\beta}g_{\beta\alpha} = g_{\gamma\alpha}$ gives the identification

$$L_{\gamma\beta}[n_{\gamma\beta}] \otimes L_{\beta\alpha}[n_{\beta\alpha}] \xrightarrow{\sim} L_{\gamma\alpha}[n_{\gamma\alpha}]$$

and the required conditions in [22, 3.13] can be deduced similarly from the fact that $\Omega_*B(R^{\times}) = R^{\times}$ is discrete.



FIGURE 1. Čech 3-cocycle condition

Even more concretely, the space $Map(X, B(R^{\times}))$ has its connected components given by $H^1(X, R)$, and, when X = M is a manifold, by assuming $\{U_{\alpha}\}$ to be a good cover, one can canonically trivialize the $L_{\beta\alpha}$'s and the identifications on triple overlaps will be simply given by some usual Čech 3-cocycles $\{c_{\gamma\beta\alpha}\}$, satisfying conditions $c_{\delta\gamma\alpha}c_{\gamma\beta\alpha} = c_{\delta\gamma\beta}c_{\delta\beta\alpha}$ as illustrated in Figure 1.

In other words, over R, if we decompose \mathcal{A} to the gluing data $(\mathcal{A}|_{U_{\alpha}}, \mathrm{id}_{\mathcal{A}}, \mathrm{id}_{\mathrm{id}_{\mathcal{A}}})$, the sheaf \mathcal{A}^{τ} can be obtained simply by modifying the 2-morphisms to $(C|_{U_{\alpha}}, id, c_{\delta\gamma\alpha})$. Thus an object in $\mathcal{A}^{\tau}(X)$ is a family $\{A_{\alpha}\}$, for $A_{\alpha} \in \mathcal{A}(U_{\alpha})$, with identifications $g_{\beta\alpha} : A_{\alpha}[n_{\beta\alpha}] = A_{\beta}$, on double overlaps, so that $g_{\gamma\beta} \circ g_{\beta\alpha}$ and $g_{\gamma\alpha}$ differ by $c_{\gamma\beta\alpha}$, on triple overlaps.

Remark 2.8. In fact, the description in the above Example 2.7 holds more generally. For two twisting $\eta_1, \eta_2 : X \to BPic(\mathcal{C})$, a homotopy $h : j^*\eta_1 = j^*\eta_2$ on some open set $j : U \hookrightarrow X$, i.e., a path in $Map(U, BPic(\mathcal{C}))$ induces an identification of the bundle $h_1 : \mathcal{P}_{\eta_1}|_U = \mathcal{P}_{\eta_2}|_U$. Two such homotopies h_1, h_2 differs necessarily by a loop $g := h_2 \# h_1^{rev} \in \Omega_{\eta_1} Map(U, BPic(\mathcal{C})) = Map(U, Pic(\mathcal{C}))$. The discussion in Remark B.6 implies that the auto-equivalence

$$h_2 \# h_1^{\text{rev}} : \mathcal{P}_{\eta_1} = \mathcal{P}_{\eta_2} = \mathcal{P}_{\eta_1}$$

is exactly given by tensoring with g. The above example is the case when $\eta_2 = *$ is the constant map.

2.3. Twisted sheaves. We specialize to the case $\mathcal{A} = \operatorname{sh}_X$, the sheaf of sheaves on a locally compact Hausdorff space X, and keep the locally contractible assumption. For such spaces X and Y, colimit-preserving functors from $\operatorname{sh}(Y)$ to $\operatorname{sh}(X)$ are given by sheaf kernels. More precisely, there is an equivalence of categories

$$sh(X \times Y) \xrightarrow{\sim} Fun^{L}(sh(X), sh(Y))$$
$$K \mapsto (F \mapsto K \circ F),$$

through convolution $K \circ F := \pi_{Y!}(K \otimes \pi_X^* F)$, which follows tracing the computation

$$\operatorname{Fun}^{L}(\operatorname{sh}(X), \operatorname{sh}(Y)) = \operatorname{sh}(X)^{\vee} \otimes \operatorname{sh}(Y) = \operatorname{sh}(X) \otimes \operatorname{sh}(Y) = \operatorname{sh}(X \times Y).$$

Here the first equality follows from abstract properties of dualizable objects, the second equality follows from the fact that sh(X) is self-dual, and the last equality is the Künneth formula. (See for example [31, Corollary 1.3.1.8])

Lemma 2.9 (Twisted Künneth formula). Let X and Y be locally contractible and locally compact Hausdorff spaces, and $\eta \in Map(X, BPic(\mathcal{C}))$ and $\xi \in Map(Y, BPic(\mathcal{C}))$ be twistings. Then there is an equivalence $\operatorname{sh}^{\eta}(X) \otimes \operatorname{sh}^{\xi}(Y) = \operatorname{sh}^{\eta \Box \xi}(X \times Y)$ where $\eta \boxdot \xi$ is given by the composition

$$X \times Y \xrightarrow{\eta \times \xi} BPic(\mathcal{C}) \times BPic(\mathcal{C}) \to BPic(\mathcal{C})$$

with the last map being the multiplication.

Proof. The proof is a very standard argument in the set up of $\operatorname{Pr}_{st}^{L}$ and can be found in [11, III.3.1.5]. That is, for any choose of over cover $\{U_{\alpha}\}$, the category $\operatorname{sh}^{\tau}(X) = \limsup \operatorname{sh}^{\tau}(U_{\alpha}) = \operatorname{colim} \operatorname{sh}^{\tau}(U_{\alpha})$ is both a limit and a colimit, and thus commutes with \otimes . As a result,

$$\operatorname{sh}^{\eta}(X) \otimes \operatorname{sh}^{\xi}(Y) = \lim_{\alpha,\alpha'} \left(\operatorname{sh}^{\eta}(U_{\alpha}) \otimes \operatorname{sh}^{\xi}(V_{\alpha'}) \right)$$

where $\{V_{\alpha'}\}$ is a open cover of Y. Assume the covers are given by contractible open sets and the Künneth formula follows from the usual one.

Corollary 2.10. For locally Hausdorff topological spaces X and Y, twists $\eta : X \to BPic(\mathcal{C})$ and $\xi : Y \to BPic(\mathcal{C})$, there is an equivalence

$$\operatorname{Fun}^{L}(\operatorname{sh}(X)^{\eta}, \operatorname{sh}(Y)^{\xi}) = \operatorname{sh}(X \times Y)^{\eta^{-1} \boxdot \xi}$$

Proof. By Lemma 2.6 and Lemma 2.9, it remains to show $(\operatorname{sh}^{\eta}(X))^{\vee} = \operatorname{sh}^{\eta^{-1}}(X)$. To see this one chase through the proof of [11, Proposition 6.3.4] and see that the unit and counit are both given by the diagonal $\Delta : X \hookrightarrow X \times X$. More precisely, one can twist the inclusion $\operatorname{sh}_{\Delta} \hookrightarrow \operatorname{sh}_{X \times X}$ of sheaves supported on the diagonal to sheaves on the product by $\eta \boxdot \eta^{-1}$. Because $\Delta^*(\eta \boxdot \eta^{-1}) = *$ and thus $(\operatorname{sh}_{\Delta})^{\eta \boxdot \eta^{-1}} = \operatorname{sh}_{\Delta}$, we obtain an adjunction

$$\Delta^* : \operatorname{sh}^{\eta \Box \eta^{-1}}(X \times X) \rightleftharpoons \operatorname{sh}(X) : \Delta_*,$$

and the unit and counit exhibiting $(\operatorname{sh}^{\eta}(X))^{\vee} = \operatorname{sh}^{\eta^{-1}}(X)$ is given by the two compositions

$$\mathcal{C} \xrightarrow{a^*} \operatorname{sh}(X) \xrightarrow{\Delta_*} \operatorname{sh}^{\eta \Box \eta^{-1}}(X \times X), \text{ and } \operatorname{sh}^{\eta \Box \eta^{-1}}(X \times X) \xrightarrow{\Delta^*} \operatorname{sh}(X) \xrightarrow{a_!} \mathcal{C}$$

where $a: X \to \{*\}$ is the projection to the point.

Example 2.11. Following the notation of Example 2.7, the above identification in the case when C = R - mod for a (discrete) ring R is simply given by convolution of the gluing data. That is, for $\{F_{\alpha}\} \in \operatorname{sh}(X)^{\eta}$ and $\{K_{\alpha'\alpha}\} \in \operatorname{sh}(X \times Y)^{\eta^{-1} \Box \xi}$, one can check directly that the family $\{K_{\alpha'\alpha} \circ F_{\alpha}\}_{\alpha'}$ forms an object in $\operatorname{sh}(Y)^{\xi}$.

Example 2.12. Consider a map $f : X \to Y$ and a twisting $\xi : Y \to BPic(\mathcal{C})$. The kernel

$$1_{\Gamma_f} \in \operatorname{sh}(X \times Y) = \operatorname{sh}^{f^* \xi \boxdot \xi^{-1}}(X \times Y)$$

gives the adjunction $f^* : \operatorname{sh}^{\xi}(Y) \rightleftharpoons \operatorname{sh}^{f^*\xi}(X) : f_*$.

Note that the same argument as in the proof of corollary 2.10 provides us the notion of tensor product and internal-Hom.

Definition 2.13. Let $\eta_1, \eta_2 : X \to BPic(\mathcal{C})$ be twistings. There is a tensor product

$$(-)\otimes(-):\operatorname{sh}^{\eta_1}(X)\otimes\operatorname{sh}^{\eta_2}(X)\to\operatorname{sh}^{\eta_1\cdot\eta_2}(X)$$

which is given by the composition $\operatorname{sh}^{\eta_1}(X) \otimes \operatorname{sh}^{\eta_2}(X) \xrightarrow{\sim} \operatorname{sh}^{\eta_1 \boxdot \eta_2}(X \times X) \xrightarrow{\Delta^*} \operatorname{sh}^{\eta_1 \cdot \eta_2}(X)$. By passing to the right adjoint, there is an internal-Hom

$$\mathcal{H}om(-,-): \operatorname{sh}^{\eta_1}(X)^{op} \otimes \operatorname{sh}^{\eta_2}(X) \to \operatorname{sh}^{\eta_1^{-1} \cdot \eta_2}(X).$$

In particular, there is a Verdier dual $D_X : \operatorname{sh}^{\eta}(X) \to \operatorname{sh}^{\eta^{-1}}(X)$ given by Hom-ing into the dualizing sheaf $\mathcal{H}om(-,\omega_X)$.

2.4. Microlocalization for twisted sheaves. Let M be a manifold. Since local representatives of a twisted sheaf $F \in \operatorname{sh}(M)^{\eta}$ differ only by tensoring with a local system, they always have the same microsupport.

Definition 2.14. For $F \in sh(M)^{\eta}$, the microsupport of F is defined as the union

$$ss(F) := \bigcup_{U,F_U} ss(F_U),$$

where U runs over open sets where \mathcal{P}_{η} is trivializable and $F_U \in \operatorname{sh}(U)$ runs over local representatives of F.

The notion of microsupport for twisted sheaf locally matches the ordinary notion of microsupport of untwisted sheaves. Therefore, all properties of the microsupport of untwisted sheaves which can be checked locally also hold for twisted sheaves. For example, ss(F) is a closed conic coisotropic subset of T^*M .

For a closed conic subset $X \subseteq T^*M$, we use the notation

$$(\operatorname{sh}^{\eta})_X(M) \coloneqq \{F \in \operatorname{sh}^{\eta}(M) | ss(F) \subseteq X\}$$

to denote the subcategory of twisted sheaves microsupported in X.

Definition-Lemma 2.15. There is an equivalence $(sh^{\eta})_X(M) = (sh_X)^{\eta}(M)$. In other words, twisting respects the notion of microsupport and there is an unambiguous meaning of the notation $sh_X^{\eta}(M)$.

Proof. As usual, choose a cover by open balls $\{U_{\alpha}\}$ and write $(\operatorname{sh}_X)^{\eta}(M)$ by the limit $\limsup \operatorname{sh}_X(U_{\alpha})$.

Example 2.16. By the above definition-lemma, twisted sheaves with microsupport contained in the zero section $\operatorname{sh}_{0_M}^{\eta}(M) = \operatorname{Loc}^{\tau}(M)$ are given by twisted local systems.

For a manifold M and a twist $\eta \in Map(M, BPic(\mathcal{C}))$, we can consider a presheaf of categories (3) $\mu \operatorname{sh}^{\eta, pre}(-) : \Omega \mapsto \operatorname{sh}^{\eta}(M) / \operatorname{sh}^{\eta}_{\Omega^{c}}(M).$

Definition 2.17. We let $\mu \operatorname{sh}_{T^*M}^{\eta}$ be the sheafification of the presheaf (3). This is a sheaf of categories on T^*M ; the objects of $\mu \operatorname{sh}_{T^*M}^{\eta}(\Omega)$, for an open set $\Omega \subseteq M$, are called $(\eta$ -)twisted microsheaves on Ω . The notion of microsupport descends to $\mu \operatorname{sh}_{T^*M}^{\eta}$ so, for a conic closed subset $X \subseteq T^*M$, we denote by $\mu \operatorname{sh}_X^{\tau}$ the subsheaf consisting of objects microsupported on X.

Since the microsupport ss(F) of a sheaf $F \in sh(M)$ is conic, the restriction of $\mu sh_{T^*M}^{\eta}$ to \dot{T}^*M is a the pullback of a sheaf of categories on the cosphere bundle S^*M . We shall denote this later sheaf by $\mu sh_{S^*M}^{\eta}$. We use a similar notation μsh_X^{η} for objects microsupported on a closed set $X \subseteq S^*M$. Because T^*M retracts to M, η extends to T^*M uniquely so we abuse the notation to denote the extension by $\eta: T^*M \to Pic(\mathcal{C})$

Lemma 2.18. There is an equivalence $\mu \operatorname{sh}_{T^*M}^{\eta} = (\mu \operatorname{sh}_{T^*M})^{\eta}$. That is, microlocalizing $(\eta -)$ twisted sheaves is the same as twisting (by η) the usual microsheaves.

Proof. Follows from Definition-Lemma 2.15.

We note that by [28, Proposition 5.4.14], the Verdier dual $D_M : \operatorname{sh}^{\eta}(M) \to \operatorname{sh}^{\eta^{-1}}(M)^{op}$ microlocalizes to

(4)
$$\mu \mathbb{D}_M : \mu \mathrm{sh}_{T^*M}^{\eta} \to a^* \,\mu \mathrm{sh}_{T^*M}^{\eta^{-1}}$$

where $a: T^*M \to T^*M$ is the antipodal map $a(x,\xi) = (x, -\xi)$. Since Verdier duality restricts to an equivalence $D_M : \operatorname{sh}_{\mathbb{R}-c}(M)^b = \operatorname{sh}_{\mathbb{R}-c}(M)^{b,op}$ on constructible sheaves with perfect stalks, by [28, Remark 7.5.8], its microlocalization $\mu \mathbb{D}_M$ restricts to an equivalence

$$\mu \mathbb{D}_M : (\mu \mathrm{sh}_{T^*M;\mathbb{R}-c}^{\eta})^b = a^* (\mu \mathrm{sh}_{T^*M;\mathbb{R}-c}^{\eta^{-1}})^{b,op}$$

on microsheaves with Lagrangian microsupport and perfect microstalks.

2.5. Microlocalizing integral transforms. We collect and adapt material from [28, Section 7.1].

Definition 2.19 ([28, Definition 7.1.1]). For manifolds M and N and conic open subsets $\mathcal{U} \subseteq T^*M$ and $\mathcal{V} \subseteq T^*N$ one denotes by $N(\mathcal{U}, \mathcal{V})$ the subcategory of $\mu sh^{pre}(T^*M \times \mathcal{V})$ consisting of objects K satisfying:

- (1) $ss(K) \cap (T^*M \times \mathcal{V}) \subseteq \mathcal{U}^a \times \mathcal{V}$ where $(-)^a$ means the image under the antipodal map $a: (x,\xi) \mapsto (x,-\xi)$.
- (2) The projection $ss(K) \cap (T^*M \times \mathcal{V}) \to \mathcal{V}$ is proper.

Remark 2.20. When $M = \{*\}$, the category $N(\{*\}, \mathcal{V}) = \mu \operatorname{sh}^{\operatorname{pre}}(\mathcal{V})$.

Proposition 2.21 ([28, Proposition 7.1.2, Definition 7.1.3]). For i = 1, 2, 3, let M_i be a manifold and $U_i \subseteq T^*M_i$ be a conic open set. Let $K \in N(\mathcal{U}_2, \mathcal{U}_3)$. Then, $K \circ (-) : sh(M_1 \times M_2) \rightarrow sh(M_1 \times M_3)$ which is defined by

$$K \circ L \coloneqq p_{13!}(p_{23}^* K \otimes p_{12}^* L)$$

descends to a functor $K \circ (-)$: $\mu sh^{pre}(T * M_1 \times U_2) \rightarrow \mu sh^{pre}(T^*M_1 \times U_3)$ and the canonical map $K \circ L \rightarrow p_{13*}(p_{23}^*K \otimes p_{12}^*L)$ is an equivalence. Furthermore, this functor restricts to

 $K \circ (-) : N(\mathcal{U}_1, \mathcal{U}_2) \to N(\mathcal{U}_1, \mathcal{U}_3).$

In particular, for $K \in N(\mathcal{U}, \mathcal{V})$, there is a functor

$$K \circ (-) : \mu \mathrm{sh}^{\mathrm{pre}}(\mathcal{U}) \to \mu \mathrm{sh}^{\mathrm{pre}}(\mathcal{V}),$$

and the canonical morphism $K \circ F \to p_{2*}(K \circ \pi_1^* F)$ is an isomorphism for $F \in \mu sh^{pre}(\mathcal{U})$.

One might want to use the above proposition to define an action of general microsheaf kernels on microsheaves. However, notice that for $\mathcal{V}_1 \subseteq \mathcal{V}_2$, the image of $N(\mathcal{U}, \mathcal{V}_2)$ under the restriction $\mu sh^{pre}(T^*M \times \mathcal{V}_1) \rightarrow \mu sh^{pre}(T^*M \times \mathcal{V}_2)$ is not necessarily contained in $N(\mathcal{U}, \mathcal{V}_1)$, since the intersection of a compact set with an open set is, in general, only closed and condition (2) of Definition 2.19 cannot be guaranteed.

Luckily for us, we will be interested in actions given by microsheaf kernels microsupported on graphs of homogeneous symplectomorphisms and Proposition 2.21 can be applied. Fix conic open subsets $\mathcal{U} \hookrightarrow \dot{T}^*M$, $\mathcal{V} \hookrightarrow \dot{T}^*N$, and a homogeneous symplectomorphism $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{V}$. Denote by $\Gamma_{\chi}^a \coloneqq \{(x, -\xi, \chi(x, \xi)) | (x, \xi) \in \mathcal{U}\}$ the image of the graph of χ under $(a \times id)$. Consider the subpresheaf $\mu \operatorname{sh}_{\Gamma_{\chi}^a}^{\operatorname{pre}}$ of $\mu \operatorname{sh}^{\operatorname{pre}}$ on $\mathcal{U}^a \times \mathcal{V}$ which is given by the assignment

$$(\Omega \subseteq \mathcal{U}^a \times \mathcal{V}) \mapsto \{F \in \mu \mathrm{sh}^{\mathrm{pre}}(\Omega) | ss_{\Omega}(F) \subseteq \Gamma^a_{\chi} \}$$

Here, we use the notation $ss_{\Omega}(F) \coloneqq ss(F) \cap \Omega$ to denote the microsupport on Ω .

Denote by q_i the projection of $T^*M \times T^*N$ to the corresponding component. For an open set $\Omega \subseteq \mathcal{U}^a \times \mathcal{V}$, set $\mathcal{U}(\Omega) \coloneqq q_1(\Omega \cap \Gamma_{\chi}^a)^a$ and $\mathcal{V}(\Omega) \coloneqq \chi(\mathcal{U}(\Omega))$. Then observe that, because of the

microsupport condition Γ_{χ}^{a} , $\mu \operatorname{sh}_{\Gamma_{\chi}^{a}}^{\operatorname{pre}}(\Omega) = \mu \operatorname{sh}_{\Gamma_{\chi}^{a}}^{\operatorname{pre}}(\mathcal{U}(\Omega) \times \mathcal{V}(\Omega))$ and so, combined with the fact that $q_{2}: \Gamma_{\chi}^{a} \to \mathcal{V}$ is a diffeomorphism, Proposition 2.21 can be applied. Thus, we obtain an action $\mu \operatorname{sh}_{\Gamma_{\chi}^{a}}^{\operatorname{pre}}(\Omega) \otimes \mu \operatorname{sh}^{\operatorname{pre}}(\mathcal{U}(\Omega)) \to \mu \operatorname{sh}^{\operatorname{pre}}(\mathcal{V}(\Omega))$.

(5)
$$\mu \mathrm{sh}_{\Gamma^a_{\mathcal{V}}} \otimes q_1^* \,\mu \mathrm{sh}_{\mathcal{U}} \to q_2^* \,\mu \mathrm{sh}_{\mathcal{V}}$$

Proposition 2.22. The morphism $\mu \operatorname{sh}_{\Gamma^a_{\chi}} \otimes q_1^* \mu \operatorname{sh}_{\mathcal{U}} \to q_2^* \mu \operatorname{sh}_{\mathcal{V}}$ from (5) is an equivalence.

Proof. We check at stalks in the sense of sheaves valued in stable categories. That is, for (x, ξ, y, η) , we need to check that $\mu \operatorname{sh}_{\Gamma_{\chi}^{a},(x,-\xi,y,\eta)} \otimes \mu \operatorname{sh}_{(x,\xi)} \to \mu \operatorname{sh}_{(y,\eta)}$ is an equivalence. We recall that, over stable categories, an objects of $\mu \operatorname{sh}_{(x,\xi)}$ is given by germs of objects in $\mu \operatorname{sh}^{\operatorname{pre}}(U_0)$ for $U_0 \ni (x,\xi)$. See for example [36]. By [28, Corollary 7.2.2], on small open sets near $(x, -\xi, y, \eta)$, there exists \mathcal{K} of microlocal rank 1, which induces an equivalence $\mathcal{K} \circ (-) : \mu \operatorname{sh}_{(x,\xi)} \xrightarrow{\sim} \mu \operatorname{sh}_{(y,\eta)}$ by [28, Proposition 7.1.10]. Thus, the identification of $\mathcal{C} = \mu \operatorname{sh}_{\Gamma_{\chi}^{a},(x,-\xi,y,\eta)}$ through $1_{\mathcal{C}} \mapsto \mathcal{K}$ identify the desired equivalence as

$$\mu \mathrm{sh}_{\Gamma^a_{\chi},(x,-\xi,y,\eta)} \otimes \mu \mathrm{sh}_{(x,\xi)} = 1_{\mathcal{C}} \otimes \mu \mathrm{sh}_{(x,\xi)} = \mu \mathrm{sh}_{(y,\eta)}.$$

Note that convolution of twisted sheaves from Corollary 2.10 can be microlocalized directly, since microsupport is local in nature as discussed in Section 2.4. In particular, we have the following proposition.

Proposition 2.23. Let $\eta \in Map(M, Pic(\mathcal{C}))$ and $\xi \in Map(N, Pic(\mathcal{C}))$ be twistings. Following the same notation as above, there is an action of $(\eta^{-1} \boxdot \xi)$ -twisted microsheaf kernel, sending η -twisted microsheaves on \mathcal{U} to ξ -twisted microsheaves on \mathcal{V} ,

$$\mu \mathrm{sh}_{\Gamma^a_{\chi}}^{\eta^{-1} \boxdot \xi} \otimes q_1^* \, \mu \mathrm{sh}_{\mathcal{U}}^{\eta} \to q_2^* \, \mu \mathrm{sh}_{\mathcal{V}}^{\xi}$$

and it is an equivalence.

Lemma 2.24. Let $Z \subseteq M$ be a closed submanifolds and denote by $N^*(Z)$ the conormal bundle of Z in M. Then $\mu sh_{N^*(Z)} = Loc_{N^*(Z)}$.

Proof. Denote by $\mathcal{C}_{N^*(Z)}^{\text{pre}}$ the constant presheaf, i.e., $\mathcal{C}_{N^*(Z)}^{\text{pre}}(\Omega) = \mathcal{C}$ for all $\Omega \subseteq N^*(Z)$. There is a morphism between presheaves whose effect over Ω is given by

$$\mathcal{C}_{N^*(Z)}^{\text{pre}}(\Omega) = \mathcal{C} \to \mu \text{sh}_{N^*(\Delta)}^{\text{pre}}(\Omega)$$
$$C \mapsto C_Z$$

where C_Z is the constant sheaf supported on the submanifold Z with stalk C. By [28, Proposition 6.6.1], this map induces equivalences on stalks, and thus becomes an equivalence $\text{Loc}_{N^*(Z)} = \mu \text{sh}_{N^*(Z)}$ upon sheafification.

Example 2.25. We note that $\Gamma^a_{\mathrm{id}_{T^*M}} = N^*(\Delta) \cong T^*M$ where $\Delta \subseteq M \times M$ is the diagonal and the later diffeomorphism can be obtained by dualizing the short exact sequence

$$1 \to TM \xrightarrow{d\Delta} \Delta^* T(M \times M) \to N(\Delta) \to 1.$$

By base change, for any sheaf $F \in sh(M)$,

$$C_{\Delta} \circ F = p_{2!}(\Delta_! C_M \otimes p_1^* F) = p_{2!} \Delta_! (C_M \otimes \Delta^* p_1^* F) = C_M \otimes F.$$

That is, the action of $\mu sh_{N^*(\Delta)}$ on μsh_{T^*M} under the equivalence $Loc_{T^*M} = \mu sh_{T^*M}$ is canonically the by local system discussed in Example 2.1.

Lemma 2.26. Assume \mathcal{U} is contractible. Then the category $\mu \operatorname{sh}_{\Gamma^a_{\chi}}(-\mathcal{U} \times \mathcal{V})$ is non-empty and the functor

$$\mu \mathrm{sh}_{\Gamma^a_{\chi}}^{\eta^{-1} \boxdot \xi} (-\mathcal{U} \times \mathcal{V}) \to Fun^{ex} \left(\mu \mathrm{sh}^{\eta}(\mathcal{U}), \mu \mathrm{sh}^{\xi}(\mathcal{V})\right)$$
$$\mathcal{K} \mapsto (\mathcal{F} \mapsto \mathcal{K} \circ \mathcal{F})$$

is conservative.

Proof. We refer to Theorem 3.8 for the non-emptiness of $\mu \operatorname{sh}_{\Gamma^a_{\chi}}(-\mathcal{U} \times \mathcal{V})$. We remark that we will need the weaker version discussed in [28, Theorem 7.2.1] and only include this version for completeness.

By symmetry, the category $\mu \operatorname{sh}_{\Gamma^a_{\chi^{-1}}}(-\mathcal{V} \times \mathcal{U})$ is non-empty. In fact, the proof of [28, Theorem 7.2.1] implies that the convolution inverse of \mathcal{K} is given by

$$\mathcal{K}^* \coloneqq (dv^{\vee})^* \, \mu \mathbb{D}_{M \times N}(\mathcal{K} \otimes p_1^* \omega_M)$$

where $v: N \times M = M \times N$ is the coordinates swapping map is the coordinate swapping map. If two microkernel $\mathcal{K}_1, \mathcal{K}_2$, induces the same functor, then $\mathcal{K}_2^* \circ \mathcal{K}_1$ will acts as the identity. But we know from Example 2.25 that $\mathcal{K}_2^* \circ \mathcal{K}_1 = \mathcal{I}_{\mathcal{U}}$, since \mathcal{U} is contractible, and so $\mathcal{K}_2 = \mathcal{K}_1$. \Box

Corollary 2.27. Let $\mathcal{K} \in \mu \operatorname{sh}_{\Gamma^a_{\mathcal{V}}}(-\mathcal{U} \times \mathcal{V})$ be an invertible object. Then the morphism

$$\operatorname{Loc}_{\mathcal{U}} \to (a_M \times \chi)^* \mu \operatorname{sh}_{\Gamma^a_{\chi}}(-\mathcal{U} \times \mathcal{V})$$
$$L \mapsto L \otimes \mathcal{K}$$

is an equivalence.

Proof. Since this is morphism between sheaves, it is sufficient to check on an open basis. That is, we only need to evaluate at small enough open balls $\Omega \subseteq \mathcal{U}$. In this case, fully-faithfulness is [28, (7.2.4) in Theorem 7.2.1.]. To see it's an surjection, consider any other microkernel \mathcal{K}' . The above Lemma 2.26 implies that $\mathcal{K}^* \circ \mathcal{K}' = c_{\mathcal{U}}$ for some $c \in Pic(\mathcal{C})$ and thus $\mathcal{K}' = c \otimes \mathcal{K}$.

3. INTEGRAL TRANSFORMS AND MASLOV DATA

We have seen in Section 2.4, that to a manifold M and a twisting $\eta : M \to Pic(\mathcal{C})$, we may associate a sheaf of categories $\mu sh_{S^*M}^{\eta}$. More generally, we could have allowed twistings $\eta : S^*M \to Pic(\mathcal{C})$.

In fact, for any contact manifold \mathcal{U} , subject to the vanishing of a certain topological obstruction, the space of twistings has a natural torsor called the space of 'Maslov data', and for each Maslov datum τ , there is a corresponding sheaf-of-categories $\mu \operatorname{sh}_{\mathcal{U},\tau}$ on \mathcal{U} [33]. When $\mathcal{U} = S^*M$, there is a distinguished Maslov datum ϕ arising ultimately from the natural polarization of the cotangent bundle, and an identification of $\mu \operatorname{sh}_{S^*M,\phi}$ with $\mu \operatorname{sh}_{S^*M}$ in the previous sense [8].

In particular, given a contactomorphism $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{U}'$ along with some Maslov datum τ' on \mathcal{U}' and a homotopy $h : \tau \sim \chi^* \tau'$, one obtains an equivalence $h^* : \mu \operatorname{sh}_{\mathcal{U},\tau} \xrightarrow{\sim} \chi^* \mu \operatorname{sh}_{\mathcal{U}',\tau'}$. These functors were originally constructed by an abstract argument. The purpose of this Section is explain how to more explicitly describe such functors h^* in local charts via integral transforms.

3.1. **Maslov data.** We recall results from [33, Section 11]. Some homotopy theory background can be found in Appendix B. The central construction is a group homomorphism [33, (28), Proposition 11.11]

(6)
$$\mathfrak{M}: U/O \to BPic(\mathcal{C}),$$

For any co-oriented contact manifold (V, ξ) , we denote by $BPic(\mathcal{C})(\xi)$ the principal $BPic(\mathcal{C})$ bundle classified by the composition

(7)
$$V \xrightarrow{\xi} BU \to B(U/O) \xrightarrow{B\mathfrak{M}} B^2 Pic(\mathcal{C}).$$

Definition 3.1.

(1) *Maslov data* is a null-homotopy τ of the map $V \to B(U/O) \to B^2 Pic(\mathcal{C})$ from (7).

(2) A *polarization* is a null-homotopy ϕ of the classifying map $V \to B(U/O)$.

Evidently a polarization induces Maslov data. If ϕ is a polarization, we will routinely abuse notation by also denoting the induced Maslov datum by ϕ .

Let (V,ξ) be a contact manifold. Then we can consider the (stable) Lagrangian Grassmannian $\mathfrak{f}: U/O(\xi) \to V$. Its relative cotangent bundle³ $T^*\mathfrak{f} \to U/O(\xi)$ is naturally a contact manifold and carries a canonical polarization. Using the large codimensional embedding trick (reviewed in Section 3.2.1 below), one may define microlocal sheaves on $T^*\mathfrak{f}$. The full subcategory of microlocal sheaves supported on the zero section is then a sheaf of categories on $U/O(\xi)$ which we denote by $\mu \mathrm{sh}_{U/O(\xi)}$. As reviewed in Proposition 3.14, this descends to a sheaf of categories on $BPic(\mathcal{C})(\xi)$ which is denoted by $\mu \mathrm{sh}_{BPic(\mathcal{C})(\xi)}$.

Definition 3.2. Given Maslov data τ on V, we define $\mu \operatorname{sh}_{V;\tau} \coloneqq \tau^* \mu \operatorname{sh}_{BPic(\mathcal{C})(\xi)}$.

The main feature of the sheaf $\mu sh_{BPic(\mathcal{C})(\xi)}$ is that it is locally constant in the fiber direction. In fact, locally on a Darboux chart \mathcal{U} and along a section, the sheaf $\mu sh_{BPic(\mathcal{C})(\xi)}$ is non-canonically equivalent to μsh_{S^*M} for some M of the correct dimension. Such a local identification respects the notion of microsupport and we can thus consider, for a closed subset $X \subseteq V$, the subsheaf $\mu sh_{X;\tau}$ of objects microsupported on X.

Proposition 3.3 ([8, Corollary 4.13]). Denote by $\phi_M : S^*M \to BO$ the (stable) fiber polarization. *There is a canonical equivalence*

(8)
$$\mu \mathrm{sh}_{S^*M;\phi_M} = \mu \mathrm{sh}_{S^*M}$$

where the right hand side is defined as in Definition 2.17. Moreover, the notion of microsupport defined in Definition 3.4 coincides with the usual notion.

We shall revisit the proof of Proposition 3.3 in the next subsection, after introducing some more notation. For now we continue our discussion of Maslov data.

Let τ_0, τ_1 be two choices of Maslov data. A homotopy $h : \tau_0 = \tau_1$ between them is equivalent to a homotopy between the corresponding sections, which we denote it by the same thing. Thus Definition 3.2 implies that there exists an equivalence $\mu sh_{V;\tau_0} = \mu sh_{V;\tau_1}$ induced by h.

We can now give a precise formulation of the problem we will solve in this section. Consider open sets $\mathcal{U} \subset S^*M$, $\mathcal{V} \subset S^*N$, and a contactomorphism $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{V}$. Proposition 3.3 implies that there is a canonical equivalence $\mu \mathrm{sh}_{\mathcal{U};\chi^*\phi_N} = \chi^* \mu \mathrm{sh}_{\mathcal{V}}$. In particular, a homotopy of Maslov data $h : \phi_M \simeq \chi^*\phi_N$ induces an equivalence $\mu \mathrm{sh}_{\mathcal{U}} = \chi^* \mu \mathrm{sh}_{\mathcal{V}}$.

³If $\pi : E \to B$ is a smooth fiber bundle, then its relative cotangent bundle is the vector bundle $T^*\pi \to E$ defined by the exact sequence $0 \to \pi^*T^*B \to T^*E \to T^*\pi \to 0$.

On the other hand, we may use the same homotopy of Maslov data to compare the fiber polarization along the graph of χ to the fiber polarization along the graph of the diagonal. Again using Proposition 3.3 along with Lemma 2.24, we find:

(9)
$$\mu \mathrm{sh}_{\Gamma^a_{\chi}} = \mu \mathrm{sh}_{\Gamma^a_{\mathrm{id}_{\mathcal{U}}}; \phi_M \boxdot \chi^* \phi_N} \stackrel{h}{=} \mu \mathrm{sh}_{N^*(\Delta) \cap (\mathcal{U}^a \times \mathcal{U})} = \mathrm{Loc}_{\mathcal{U}}.$$

Let $\mathcal{K} \in \mu \mathrm{sh}_{\Gamma^a_{\chi}}(\mathcal{U}^a \times \mathcal{V})$ be the image of $1_{\mathcal{U}}$ under the above equivalence. Then convolution of microsheaf kernels from equation (5) induces another equivalence $\mu \mathrm{sh}_{\mathcal{U}} = \chi^* \mu \mathrm{sh}_{\mathcal{V}}$ by [28, Proposition 7.1.10].

The purpose of the present section is to show that these two equivalences agree. The result is ultimately formulated as Theorem 3.17 below.

3.2. **Polarization-to-kernel.** In the present subsection, we shall consider open subsets $\mathcal{U} \hookrightarrow S^*M$ and $\mathcal{V} \hookrightarrow S^*N$ and a contactomorphism $\chi : \mathcal{U} \to \mathcal{V}$. Let $h : \phi_M \simeq \chi^* \phi_N$ be a homotopy of polarizations. Our goal is to prove that (under the identification (8)), the induced equivalence $\mu \operatorname{sh}_{\mathcal{U};\phi_M} = \mu \operatorname{sh}_{\mathcal{V};\chi_N}$ is induced by convolution with a (twisted) microsheaf kernel, and to characterize this kernel. In the next subsection, we will explain how to generalize this statement to the case where h is merely a homotopy between the Maslov data induced by $\phi_M, \chi^* \phi_N$.

3.2.1. Large codimensional embeddings. The sheaf $\mu sh_{U/O(\xi)}$ is constructed using the large codimensional embedding method which first appeared in [41] and which we now briefly review. For any co-oriented contact manifold (V^{2n-1}, ξ) , Gromov's *h*-principle provides contact embeddings

$$i:V \hookrightarrow \mathbb{R}^{2k+1}$$

for $k \gg 0$ and guarantees that the space of such embeddings can be made as connected as desired by increasing k. Such embeddings realize the inverse of the image of ξ in Map(V, BU) as its stable symplectic normal bundle ν_V .

Recall also that, because of the fiber sequence $O \to U \to U/O$, a null-homotopy of $\xi : V \to BU \to B(U/O)$ is equivalent to a map $\rho : V \to BO$ which factorizes $\xi = (BO \to BU) \circ \rho$. Since $(BO \to BU)$ is a group, the inverse $\sigma := \rho^{-1}$ composes with $BO \to BU$ to ν_V . Geometrically, this is a Lagrangian sub-bundle $\sigma \subseteq \nu_V$ of the stable symplectic normal bundle (a "stable normal polarization"). Given such a Lagrangian subbundle, one then *defines*

(10)
$$\mu \mathrm{sh}_{V;\rho} := \mu \mathrm{sh}_{V^{\sigma}} |_{V}.^{4}$$

A priori, $\mu \operatorname{sh}_{V;\rho}$ depends on the choice of embedding and thickening. To verify that it is well-defined [33, Lemma 6.3] argues by local constancy. Namely, let $h : \rho_0 = \rho_1$ be a homotopy between two polarizations. Up to stabilization, this consists of a family of Lagrangian sub-bundles $\sigma_t \subseteq \nu_V$, extending the isotopy of contact embedding $i_t : V \hookrightarrow \mathbb{R}^{2k+1}$. The total family $\mu \operatorname{sh}_{V^{\sigma_t} \times [0,1]}|_{V \times [0,1]}$ is constant along the *t*-direction and one thus obtains an equivalence $\mu \operatorname{sh}_{V;\rho_0} = \mu \operatorname{sh}_{V;\rho_1}$.

Objects of $\mu sh_{V;\sigma}$ have a well-defined (micro)support which we typically denote by ss(-).

Definition 3.4. Choose any contact embedding $i: V \hookrightarrow S^* \mathbb{R}^k$, for an object

(11)
$$\mathcal{F} \in \mu \mathrm{sh}_{V;\rho}(V) = \mu \mathrm{sh}_{V^{\sigma}}(i(V)) \coloneqq \mathrm{colim}_{\mathcal{U} \supset i(\mathcal{V})} \, \mu \mathrm{sh}_{V^{\sigma}}(\mathcal{U}),$$

its (micro)support is the subset $ss(\mathcal{F}) := i(V) \cap ss_{\mathbb{R}^k}(\mathcal{F}) \subset V$, where $ss_{\mathbb{R}^k}(-)$ refers to the usual notion of microsupport for a microsheaf on $S^*\mathbb{R}^k$.

⁴Here we once and for all fix an embedding $\mathbb{R}^{2N+1} \hookrightarrow S^*(\mathbb{R}^{N+1})$ to view $\mu \operatorname{sh}_{V^{\sigma}}|_V$ as the subsheaf of $\mu \operatorname{sh}_{S^*(\mathbb{R}^{N+1})}$ on objects microsupported in V^{σ} .

Proof of Proposition 3.3. Pick any embedding $f : M \hookrightarrow \mathbb{R}^k$ for some $k \gg 0$. By Lemma C.3 the choice of a section of $f^* : T^*\mathbb{R}^n|_M \to T^*M$ is contractible and any such section induces an embedding of Liouville manifolds $T^*M \hookrightarrow T^*\mathbb{R}^k$.

Observe also that the complexification of the normal bundle of $M \subset \mathbb{R}^k$ is naturally isomorphic, as a symplectic vector bundle, to the normal bundle of of $T^*M \hookrightarrow T^*\mathbb{R}^k$; hence $-\phi_M = \nu^*$. We let $S^*M(\nu^*)$ be the thickening of S^*M in the conormal direction. Then the desired map $\mu \operatorname{sh}_{S^*M;\phi_M} := \mu \operatorname{sh}_{S^*\mathbb{R}^k;S^*M(\nu^*)}|_{S^*M} \xrightarrow{f^*} \mu \operatorname{sh}_{S^*M}$ is obtained from microlocalizing the equivalence $i^* : sh(\mathbb{R}^k)_{supp(M)} = sh(M)$.

Now to see the statement on microsupport, we choose a local coordinate (x, v) for \mathbb{R}^k such that M is given by $\{v = 0\}$. Then $S^*\mathbb{R}^k$ will be given by $(x, v, [\xi, \lambda])$ for $\xi \in T^*_x M$ and $\lambda \in \nu^*_x$. The subset $S^*M(\nu^*)|_{S^*M}$ is then given by v = 0 and, which is exactly the image of $\mu \operatorname{sh}_{S^*M;\phi_M} \xrightarrow{\sim} \mu \operatorname{sh}_{S^*\mathbb{R}^k;S^*M(\nu^*)}|_{S^*M}$.

 $U/O(\xi)$ admits a canonical stable normal polarization and $\mu sh_{U/O(\xi)}$ is thus defined unambiguously using (10). For a general contact manifold V equipped with a stable normal polarization ρ , we may define $\mu sh_{V;\rho}$ equivalently either via (10) or by pullback $\rho^* \mu sh_{U/O(\xi)}$ [33].

3.2.2. *GKS quantization.* We begin by reviewing the GKS quantization method [18, Theorem 3.7]. For a contact isotopy $\Phi : S^*M \times I \to S^*M$, there exists a unique sheaf kernel $K(\Phi) \in \operatorname{sh}(M \times M \times I)$ so that its restriction to the time 0-slice is $K(\Phi)|_0 = 1_\Delta$ and its microsupport at infinity $ss^{\infty}(K(\Phi)) \subseteq \Lambda_{\Phi}$ is contained in the contact movie. A standard microsupport estimate implies that, for any $F \in \operatorname{sh}(M)$, the kernel $K(\Phi)$ moves the microsupport of F by the isotopy, or, when setting $F_t = K(\Phi)|_t \circ F$, we have $ss^{\infty}(F_t) = \varphi_t(ss^{\infty}(F))$. A consequence is that, for $t \in I$, convolving with $K(\Phi)|_t$ induces an equivalence $\operatorname{sh}_X(M) = \operatorname{sh}_{\varphi_t(X)}(M)$ for any closed subset $X \subseteq S^*M$.

In particular, suppose that $(\phi_t)_{t \in [0,1]}$ is a family of polarizations for the contact manifolds V. Then [33, Lemma 6.3] furnishes a map

(12)
$$\mu \operatorname{sh}_{V;\phi_0} \xrightarrow{\sim} \mu \operatorname{sh}_{V;\phi_1}$$

This existence of this map is deduced in loc. cit. by an abstract argument establishing that $\mu sh_{V;\phi_t}$ is a locally constant family of categories. In fact, this locally constant family can be described concretely via GKS quantization.

To this end, pick a high-codimensional embedding $V \hookrightarrow \mathbb{R}^{2k+1}$; then the homotopy $\phi_0 \rightsquigarrow \phi_1$ induces a family of thickenings $V_0 \rightsquigarrow V_1$, and we may assume that $V_t = \Phi_t(V_0)$ for a global contact isotopy $\Phi : [0,1] \times S^* \mathbb{R}^{2k+1} \to S^* \mathbb{R}^{2k+1}$; see Corollary C.13. We now apply the GKS sheaf quantization, which furnishes a ([0,1]-family) sheaf kernel $K(\Phi) \in \operatorname{sh}(\mathbb{R}^k \times \mathbb{R}^k \times [0,1])$. Because the equivalences induced by $K(\Phi)$ on $\operatorname{sh}(M)$ respect microsupport, we obtain an equivalence of sheaves of categories:

(13)
$$K(\Phi) \circ (-) : \mu \operatorname{sh}_{S^* \mathbb{R}^{N+1}} = \Phi^* \mu \operatorname{sh}_{\Lambda_\Phi \circ S^* M}$$

where $\Lambda_{\Phi} \circ S^* \mathbb{R}^k$ is the movie of $S^* M$ under Φ in $S^* (\mathbb{R}^k \times [0, 1])$, and it restricts to

(14)
$$K(\Phi)_1 \circ (-) : \mu \operatorname{sh}_{V_0} = \varphi_1^* \, \mu \operatorname{sh}_{V_1}$$

where we use the notation $\varphi_t \coloneqq \Phi(-, t), t \in [0, 1]$.

Lemma 3.5. *The maps* (12) *and* (14) *agree.*

Proof. As discussed in [18, 3.4], for any $t \in [0, 1]$, the restriction $i_t^* : \operatorname{sh}_{\Lambda_{\Phi} \circ T^*M}(M \times [0, 1]) \to \operatorname{sh}(M)$ is an equivalence with the inverse of i_0^* given by convolving with $K(\Phi) \circ (-)$. In fact, [18, Proposition 3.12] implies that this equivalence microlocalizes to an equivalence between sheaves of categories

$$K(\Phi) \circ (-) : \mu \mathrm{sh}_{S^* \mathbb{R}^k} = \Phi^* \mu \mathrm{sh}_{\Lambda_{\Phi} \circ S^* \mathbb{R}^k}$$

and the same proposition shows that it restricts to equivalences on subsheaves

$$K(\Phi) \circ (-) : \mu \operatorname{sh}_{V_0} = \Phi^* \mu \operatorname{sh}_{\Lambda_{\Phi} \circ V_0}$$
.

Here, the sheaf $\mu sh_{\Lambda_{\Phi} \circ V_0}$ is exactly the constant family in which gives rise to the family of sheaves $\{\mu sh_{V;\phi_t}\}_{t\in[0,1]}$ by restricting to the *t*-slice. Similarly, one can microlocalize i_t^* and obtain

$$i_t^*: \Phi^* \mu \operatorname{sh}_{\Lambda_\Phi \circ V_0} = \mu \operatorname{sh}_{V_t}.$$

By composing the microlocalization of $K(\Phi)$ with that of i_1^* , we see that the equivalence (12) is realized by the equivalence (14).

Remark 3.6. The proof of Lemma 3.5 shows that the action of $\mathcal{K}(\Phi)|_t$ on the sheaf of categories μsh_{V_0} is independent of the extension Φ , and only depends on the family of thickenings $V_0 \rightsquigarrow V_1$, and not on the extension to a global contact isotopy. (Indeed, a GKS kernel associated to a contact isotopy which is the identity on $V \subset \mathbb{R}^{2n+1}$ acts by the identity on μsh_V).

It will also be important to us to consider the case where $V = -\mathcal{U} \times \mathcal{U}$ and we have an isotopy of polariations $\phi_0 \boxdot \phi_0 \rightsquigarrow \phi_0 \boxdot \phi_1$. In this setting, we again have abstractly by [33] an induced equivalence

(15)
$$\mu \mathrm{sh}_{-\mathcal{U} \times \mathcal{U}; \phi_0 \boxdot \phi_0} \xrightarrow{\sim} \mu \mathrm{sh}_{-\mathcal{U} \times \mathcal{U}; \phi_0 \boxdot \phi_1}$$

Meanwhile, there is another way to produce such an equivalence. Namely, consider the family of equivalences

(16)
$$K(\Phi)|_{t} \circ (-) : \mu \mathrm{sh}_{-\mathcal{U} \times \mathcal{U}; \phi_{0} \Box \phi_{0}} \to \mu \mathrm{sh}_{-\mathcal{U} \times \mathcal{U}; \phi_{0} \Box \phi_{t}}.$$

Lemma 3.7. (15) and (16) agree

Proof. Same argument as the proof of Lemma 3.5, replacing (the microlocalization of) convolution on sheaves

$$K \circ F \coloneqq p_{2!}(K \otimes \pi_1^* F)$$

by (the microlocalization of) convolution between sheaf kernels

$$K \circ H \coloneqq p_{13!}(\pi_{23}^* K \otimes p_{12}^* H).$$

We now come to the main result of this section. To set the stage, let $\mathcal{U} \hookrightarrow S^*M$, $\mathcal{V} \hookrightarrow S^*N$ be opens, and $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{V}$ be a contactomorphism. If $h : \phi_M|_{\mathcal{U}} = \chi^*(\phi_N|_{\mathcal{V}})$ is a homotopy between polarizations, then we have

$$\operatorname{Loc}_{S^*M} = \mu \operatorname{sh}_{N^*(\Delta_M)} = \mu \operatorname{sh}_{N^*(\Delta_M);\phi_M \boxdot \phi_M} = \mu \operatorname{sh}_{N^*(\Delta_M);\phi_M \boxdot \chi^* \phi_N} = \mu \operatorname{sh}_{\Gamma^a_{\chi};\phi_M \boxtimes \phi_N} = \mu \operatorname{sh}_{\Gamma^a_{\chi}}$$

Theorem 3.8. Under the identification (8), the induced equivalence $\mu sh_{\mathcal{U};\phi_M} = \mu sh_{\mathcal{V}:\chi^*\phi_N}$ is given by microkernel convolution defined by (5)

$$\mathcal{K}(h) \circ (-) : \mu \mathrm{sh}_{\mathcal{U}} = \chi^* \mu \mathrm{sh}_{\mathcal{V}}$$

where $\mathcal{K}(h)$ is the image of $1_{\mathcal{U}}$ under (17).

Proof. By Lemma 3.7, the third equality of (17) can be described via (16). So the image of $1_{\mathcal{U}}$ under the first three equalities is $K(\Phi)|_1 \circ 1_{\Delta_M}$. Associativity of convolution implies that, for $\mathcal{F} \in \mu \operatorname{sh}(\mathcal{U})$,

$$\mathcal{K}(h) \circ \mathcal{F} = (K(\Phi)_1 \circ 1_{\Delta_M}) \circ \mathcal{F} = K(\Phi)_1 \circ (1_{\Delta_M} \circ \mathcal{F}) = K(\Phi)_1 \circ \mathcal{F},$$

and we thus see that the equivalence induced by homotopy of polarizations, $\mu sh_{\mathcal{U};\phi_M} = \mu sh_{\mathcal{V}:\chi^*\phi_N}$, is realized by convolving with $\mathcal{K}(h)$.

Corollary 3.9. In the same setting as Theorem 3.8, denote by $h^{\text{rev}} : \chi^*(\phi_N|_{\mathcal{V}}) = \phi_M|_{\mathcal{U}}$, then

$$\mathcal{K}(h^{\mathrm{rev}}) = v^* \, \mu \mathbb{D}_{M \times N}(\mathcal{K} \otimes p_1^* \omega_M)$$

where $v: N \times M = M \times N$ is the coordinates swapping map.

Proof. In the same setting of the proof above. We further notice that, if we write Φ^{-1} for the isotopy such that $\Phi^{-1}(-,t) = \varphi_t^{-1}$ where $\varphi_t := \Phi(-,t)$, then

$$K(\Phi^{-1}) = (v_{\mathbb{R}^k} \times \mathrm{id}_I)^* \mathcal{H}\mathrm{om}(K(\Phi), \mathbb{1}_M \boxtimes \omega_M \otimes \mathbb{1}_I),$$

where I = [0, 1], by [18, Proposition 3.2].

We end this section with a statement concerning the compatibility between Theorem 3.8 and composition.

Lemma 3.10. In the same setting as Theorem 3.8, let $h_0, h_1 : \phi_M|_{\mathcal{U}} = \chi^*(\phi_N|_{\mathcal{V}})$ be two homotopies between the polarizations. Then, a 2-homotopy $k : h_0 = h_1$ induces an equivalence $\mathcal{K}(h_0) = \mathcal{K}(h_1)$ between microkernels.

Proof. Consider the same high codimensional embedding as in the proof of Theorem 3.8. The 2-homotopy k can be realized as a family of Lagrangian thickenings $\sigma_{t,s}$, $t, s \in [0, 1]$, such that $\sigma_{t,i}$ is the homotopy between the thickening $S^*M(\nu_M^*)|_{\mathcal{U}}$ and $S^*N(\nu_N^*)|_{\mathcal{V}}$ given by h_i , i = 0, 1, and $\sigma_{0,s} = S^*M(\nu_M^*)|_{\mathcal{U}}$ and $\sigma_{1,s} = S^*N(\nu_N^*)|_{\mathcal{V}}$ for all $s \in [0, 1]$. Extend $\sigma_{t,s}$ to an ambient homotopy between isotopies $\Phi : S^*\mathbb{R}^k \times [0, 1]_t \times [0, 1]_s \to S^*\mathbb{R}^k$ such that $\Phi_{0,s} = \mathrm{id}_{S^*\mathbb{R}^k}$ and $\Phi_{1,s}$ is a constant contactomorphism for all $s \in [0, 1]$. As explained in [18, Remark 3.9], the GKS sheaf quantization applies for any contractible parameter space and we thus have a sheaf kernel $K(\Phi) \in sh(M \times M \times [0, 1] \times [0, 1])$. By construction and the uniqueness, one recovers $\mathcal{K}(h_i)$ by setting

$$\mathcal{K}(h_i) \coloneqq K(\Phi)|_{t=1,s=i} \circ 1_{\Delta_M}.$$

However, the constancy condition at t = 1 implies that $K(\Phi)|_{t=1} \in \operatorname{sh}(M \times M \times \{1\} \times [0, 1]_s)$ is constant on the *s*-direction. Thus $K(\Phi)|_{t=1,s=0} = K(\Phi)|_{t=1,s=1}$ canonically and it induces the identification $\mathcal{K}(h_0) = \mathcal{K}(h_1)$.

Corollary 3.11. Let $\mathcal{U}_i \hookrightarrow S^*M_i$, i = 0, 1, 2 be opens, $\chi_i : \mathcal{U}_i \xrightarrow{\sim} \mathcal{U}_{i+1}$, i = 0, 1 be contactomorphisms, $h_i : \phi_{M_i} = \chi_i^* \phi_{M_{i+1}}$, i = 0, 1 and $h_2 : \phi_{M_0} = (\chi_1 \circ \chi_0)^* \phi_{M_2}$ be homotopy between polarizations. Then a 2-homotopy $k : h_2 = h_1 \# h_0$ induces an identification

$$\mathcal{K}(h_2) = \mathcal{K}(h_1) \circ \mathcal{K}(h_0)$$

between microkernels and the is identification is compatible with the identification of the corresponding identifications between microsheaf categories induced from identifications between polarizations

$$k: \left(\mu \mathrm{sh}_{\mathcal{U}_0} \stackrel{h_0}{=} \mu \mathrm{sh}_{\mathcal{U}_1} \stackrel{h_1}{=} \mu \mathrm{sh}_{\mathcal{U}_2}\right) = \left(\mu \mathrm{sh}_{\mathcal{U}_0} \stackrel{h_2}{=} \mu \mathrm{sh}_{\mathcal{U}_2}\right).$$

$$\square$$

Proof. Consider a similar high codimensional embedding as in the proof of the above Lemma 3.10 and one concludes that $\mathcal{K}(h_2) = \mathcal{K}(h_1 \# h_0)$. Call the ambient isotopies which give h_0 and h_1 by Φ and Ψ . We note that up to a scaling, $\Psi \circ \Phi$ and $\Psi \# \Phi$ are homotopic to each other. Thus, a similar constancy argument at end point implies that the GKS sheaf kernel $K(\Psi \# \Phi) = K(\Psi \circ \Phi) =$ $K(\Psi) \circ K(\Phi)$ where we use the compatibility of GKS sheaf quantization between composition of isotopies for the second quality. The equivalence between microkernels is then induced by applying $(-) \circ 1_{\Delta_M}$.

Remark 3.12. Higher compatibility, e.g., associativity for compositions hold as well since it follows from the same property for the GKS sheaf quantization. We leave the details to the reader.

3.3. Some remarks on principal bundles. Let G be a group object in spaces and A an abelian group object, i.e., grouplike E_1 and E_{∞} -monoids. Consider a space X and a map $f: X \to BG$, i.e., a principal G-bundle $G(f) \to X$. Assume \mathcal{D} is a sheaf of categories on G(f) such that \mathcal{D} is locally constant on the fiber direction and, when restricting to any fiber $G \hookrightarrow G(f)$, $\mathcal{D}|_G = L$ for some fixed local system L with stalk category F. Furthermore, the monodromy of L comes from some representation $\rho: A \to Aut(F)$. In other words, there is an A-bundle $A(\alpha) \to A$, classified by a map $\alpha: G \to BA$, such that $L = A(\alpha) \times^{\rho} F$.

Suppose in addition that the map α is in fact a group homomorphism. In other words, $A(\alpha)$, the fiber of α , is a group, as explained in Example B.5, and we will denote it as K. One can apply B(-) to α and obtain a map $B(\alpha) : BG \to B^2A$ in spaces and obtain the following diagram:



Corollary 3.13. If the monodromy $\alpha : G \to BA$ is a group homomorphism, then there is a principal B(A)-bundle $B(A)(f) \to X$ such that the original principal G-bundle $G(f) \to X$ is a principal K-bundle $G(f) \to B(A)(f)$ over it.

The main Theorem [33, Theorem 11.17] is that \mathcal{D} is pullback from B(A)(f). We sketch the proof since a similar discussion will be relevant.

Proposition 3.14. There is a sheaf of categories $\overline{\mathcal{D}}$ on B(A)(f) such that $\mathcal{D} = f_{\alpha}^* \overline{\mathcal{D}}$.

Proof. We take the argument from [33, Theorem 11.17], which concerned the special case (of eventual interest to us) of $\alpha = \mathfrak{M}$. Here we formulate it abstractly for clarity.

Because $G(f) \to B(A)(f)$ is a principal K-bundle, it is enough to show that \mathcal{D} is K-equivariant. That is, there is an identification $p^*\mathcal{D} = a^*\mathcal{D}$ (with higher coherence data) where $p, a : K \times G(f) \to G(f)$ are the trivial projection and the action.

To see this, we recall that the action $a: K \times G(f) \to G(f)$ is restricted from, $\tilde{a}: G \times G(f) \to G(f)$, the action given by G. The pullback of $\tilde{a}^*\mathcal{D}$ equals $L \boxtimes_{\rho} \mathcal{D}$. By the definition, $K = A(\alpha)$ is

the fiber of $G \xrightarrow{\alpha} BA$, which classifies $A(\alpha)$, and thus the pull back $a^*\mathcal{D} = A_K \boxtimes \mathcal{D}$ is trivial.



We will also consider the situation when there are two such sheaves \mathcal{D}_0 and \mathcal{D}_1 and a morphism $T : \mathcal{D}_0 \to \mathcal{D}_1$ between them so that, on any fiber, the morphism between the local systems

$$(L_1 \xrightarrow{T|_G} L_2) = (A(\alpha) \times^{\rho} F_1 \xrightarrow{A(\alpha) \times^{\rho} t} A(\alpha) \times^{\rho} F_2)$$

are induced from a functor $t: F_1 \to F_2$ between the stalk categories. The same argument as the above Proposition 3.14 implies that we can descend morphisms as well.

Proposition 3.15. There is a morphism between sheaves of categories $\overline{T} : \overline{\mathcal{D}_0} \to \overline{\mathcal{D}_1}$ on B(A)(f) such that $T = f_{\alpha}^* \overline{T}$.

Now consider two sections s_0 , s_1 of $G(f) \to X$ and a homotopy $h : \overline{s_0} = \overline{s_1}$ between their projection to $G(f) \to B(A)(f)$. Observe that there exists a unique map $g : X \to G$ such that $s_1 = g \cdot s_0$. The homotopy $h : \overline{s_0} = \overline{s_1} = \alpha(g) \cdot s_0$ thus corresponds to a null-homotopy $k : \alpha(g) = *$. Because $K = \operatorname{fib}(\alpha)$, this data k further corresponds to a lifting which we will, by abusing of notation, denote by $k : X \to K$.

Proposition 3.16. Let s_0 , s_1 be two sections of $G(f) \to X$. A homotopy $h : \overline{s_0} = \overline{s_1}$ induces an equivalence $s_0^* \mathcal{D} = s_1^* \mathcal{D}$. A similar statement for morphisms holds.

Proof. The above discussion implies that $s_1 : X \to G(f)$ can be factored as

$$X \xrightarrow{(k,s_0)} K \times G(f) \xrightarrow{a} G(f).$$

But, as mentioned in the proof of Proposition 3.14, the pullback $a^*\mathcal{D} = K_A \boxtimes_{\rho} \mathcal{D}$ is constant on the first factor and thus $s_1^*\mathcal{D} = s_0^*\mathcal{D}$.

3.4. **From polarization-to-kernel to Maslov-to-kernel.** The goal of this section is to deduce the following from Theorem 3.8.

Theorem 3.17. Let $\mathcal{U} \hookrightarrow S^*M$, $\mathcal{V} \hookrightarrow S^*N$ be opens and let $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{V}$ be a contactomorphism. Suppose that $h : \phi_M|_{\mathcal{U}} = \chi^*(\phi_N|_{\mathcal{V}})$ is a homotopy of Maslov data (i.e. of the Maslov data induced by the polarizations). Then under the identification (8), the induced equivalence $\mu \mathrm{sh}_{\mathcal{U}; \varphi_M} = \mu \mathrm{sh}_{\mathcal{V}: \chi^* \phi_N}$ is given by microkernel convolution defined by (5)

$$\mathcal{K}(h) \circ (-) : \mu \mathrm{sh}_{\mathcal{U}} = \chi^* \, \mu \mathrm{sh}_{\mathcal{V}}$$

where $\mathcal{K}(h)$ is the image of $1_{\mathcal{U}}$ under (17).

Proof. Note that the considerations of Section 3.3 apply to $\alpha = \mathfrak{M} : U/O \to BPic(\mathcal{C})$, since this is shown to be a group homomorphism by [33, Theorem 11.10, Proposition 11.11].

The discussion before Proposition 3.16 implies that there is an equivalence $h : \phi_M|_{\mathcal{U}} = k \cdot \chi^*(\phi_N|_{\mathcal{V}})$ for some $k \in \operatorname{Map}(\mathcal{U}, K)$ where K is the fiber of the classifying map $\mathfrak{M} : U/O \to BPic(\mathcal{C})$. We cannot apply Theorem 3.8 directly but, since the action of k depends only on its composition to $K \to U/O$, and the thickening $k \cdot \chi^*(\phi_N|_{\mathcal{V}})$ differs from $\chi^*(\phi_N|_{\mathcal{V}})$ by a rotation by some element in U(l), for some $l \gg 0$, so the same proof still applies. That is, the equivalence $\mu \operatorname{sh}_{\mathcal{U}} = \mu \operatorname{sh}_{\mathcal{V};k \cdot \chi^*(\phi_N|_{\mathcal{V}})}$ is induced by microsheaf convolution and, by Proposition 3.16, the latter equals $\chi^* \mu \operatorname{sh}_{\mathcal{V}}$ trivially.

Corollary 3.18. Let $\mathcal{U}_i \hookrightarrow S^*M_i$, i = 0, 1, 2 be opens, $\chi_i : \mathcal{U}_i \xrightarrow{\sim} \mathcal{U}_{i+1}$, i = 0, 1 be contactomorphisms, $h_i : \phi_{M_i} = \chi_i^* \phi_{M_{i+1}}$, i = 0, 1 and $h_2 : \phi_{M_0} = (\chi_1 \circ \chi_0)^* \phi_{M_2}$ be homotopy between Maslov data. Then a 2-homotopy $k : h_2 = h_1 \# h_0$ induces an identification

$$\mathcal{K}(h_2) = \mathcal{K}(h_1) \circ \mathcal{K}(h_0)$$

between microkernels and the is identification is compatible with the identification of the corresponding identifications between microsheaf categories induced from identifications between Maslov data

$$k: \left(\mu \mathrm{sh}_{\mathcal{U}_0} \stackrel{h_0}{=} \mu \mathrm{sh}_{\mathcal{U}_1} \stackrel{h_1}{=} \mu \mathrm{sh}_{\mathcal{U}_2}\right) = \left(\mu \mathrm{sh}_{\mathcal{U}_0} \stackrel{h_2}{=} \mu \mathrm{sh}_{\mathcal{U}_2}\right).$$

Proof. Similar to the discussion before 3.16, if there are sections s_i , i = 0, 1, 2 and homotopies $h_i : s_i = s_{i+1}, i = 0, 1$ and $h_2 : s_0 = s_2$. Then a 2-homotopy $h_2 = h_1 \# h_0$ induces an identification between the liftings. The result then follows from Corollary 3.11 and the proof of Theorem 3.17.

Corollary 3.19. Let $\eta \in Map(M, Pic(\mathcal{C}))$ and $\xi \in Map(N, Pic(\mathcal{C}))$ be twistings. In the same setting as Theorem 3.17, for a homotopy $h : (\eta \cdot \phi_M)|_{\mathcal{U}} = \chi^*((\xi \cdot \phi_N)|_{\mathcal{V}})$, the induced equivalence $\mu sh_{\mathcal{U};\eta\cdot\phi_M} = \mu sh_{\mathcal{V}:\chi^*(\xi\cdot\phi_N)}$ is realized by convolution

$$\mathcal{K}(h) \circ (-) : \mu \mathrm{sh}_{\mathcal{U}}^{\eta} = \chi^* \mu \mathrm{sh}_{\mathcal{V}}^{\xi}$$

using Formula 5, where $\mathcal{K}(h)$ is given by an equivalence similar to (17). Furthermore, denoting by h^{rev} the reverse isotopy, then

$$\mathcal{K}(h^{\mathrm{rev}}) = v^* \, \mu \mathbb{D}_{M \times N}(\mathcal{K} \otimes p_1^* \omega_M)$$

where $v: N \times M = M \times N$ is the coordinates swapping map.

Proof. The identification $\mu \operatorname{sh}_{\mathcal{U};\eta\cdot\phi_M} = \mu \operatorname{sh}_{\mathcal{U}}^{\eta}$ follows from [8, Corollary 4.13] and Lemma 2.18. To apply Theorem 3.17, one choose a cover $\{U_{\alpha}\}$ of M and $\{V_{\beta}\}$ of N with trivializations of η and ξ . For each $S^*U_{\alpha} \cap \mathcal{U}$ and $S^*V_{\beta} \cap \mathcal{V}$, over their intersection through χ , there is an identification

$$\phi_M = \eta \cdot \phi_M = \chi^*(\xi \cdot \phi_N) = \chi^* \phi_N$$

and Theorem 3.17 provides a microsheaf kernel $\mathcal{K}(h)_{\alpha\beta}$ which realizes the equivalence by convolution. But the same gluing description is used to for Corollary 2.10, which microlocalizes to Formula 5. The statement for the reverse homotopy follows from Corollary 3.9 by a similar argument.

Corollary 3.20. Let $\eta_i \in \text{Map}(M_1, Pic(\mathcal{C}))$, for i = 0, 1, 2 be twistings. In the same setting as Corollary 3.20. If $h_i : \eta_i \cdot \phi_{M_i} = \chi_i^*(\eta_{i+1} \cdot \phi_{M_{i+1}})$, i = 0, 1 and $h_2 : \phi_{M_0} = (\chi_1 \circ \chi_0)^*(\eta_2 \phi_{M_2})$ are homotopy between Malsov data. Then a 2-homotopy $k : h_2 = h_1 \# h_0$ induces an identification

$$\mathcal{K}(h_2) = \mathcal{K}(h_1) \circ \mathcal{K}(h_0)$$

between microkernels and the is identification is compatible with the identification of the corresponding identifications between microsheaf categories induced from identifications between Maslov data

$$k: \left(\mu \mathrm{sh}_{\mathcal{U}_0}^{\eta_0} \stackrel{h_0}{=} \mu \mathrm{sh}_{\mathcal{U}_1}^{\eta_1} \stackrel{h_1}{=} \mu \mathrm{sh}_{\mathcal{U}_2}^{\eta_2}\right) = \left(\mu \mathrm{sh}_{\mathcal{U}_0}^{\eta_0} \stackrel{h_2}{=} \mu \mathrm{sh}_{\mathcal{U}_2}^{\eta_2}\right).$$

Proof. By trivializing the twistings over the corresponding covers as in the proof of Corollary 3.19, we obtain the identification $\mathcal{K}(h_2) = \mathcal{K}(h_1) \circ \mathcal{K}(h_0)$ as (untwisted) microkernels locally by Corollary 3.18. Now, the transition maps needed for gluing them back as twisted microkernels are given by tensoring with local systems, which are compatible with convolutions and thus the identifications glue as well.

Remark 3.21. As mentioned in Remark 3.12 that polarization-to-kernel satisfied higher compatibility. A similar argument as the above Corollary 3.20 implies that it descends to the level of Maslov-to-kernel as well.

4. VERDIER DUALITY ON CONTACT MANIFOLDS

In (4) we observed that Verdier duality microlocalizes to cotangent/cosphere bundles. We shall now explain how to globalize Verdier duality to arbitrary (co-oriented) contact manifolds.

Let V be a co-oriented contact manifold and let \overline{V} denote the same contact manifold with the coorientation reversed. For a Maslov datum τ , we will explain the existence of a Verdier dualization functor

(18)
$$\mu \mathbb{D}_{V;\tau} : \mu \mathrm{sh}_{V;\tau} \to \mu \mathrm{sh}_{\overline{V};\tau}^{op}$$

which specializes to (4) on Darboux charts. In particular, (18) restricts to an equivalence on constructible microsheaves with perfect microstalks.

To set the stage, we recall the standard embedding $(\mathbb{R}^{2k+1}, \sum_{i=0}^{k-1} p_i dq_i + dq_k) \stackrel{\iota}{\hookrightarrow} S^* \mathbb{R}^{k+1}$ taking $(q_0, \ldots, q_k, p_0, \ldots, p_{k-1}) \mapsto (q_0, \ldots, q_k, p_0, \ldots, p_{k-1}, -1)$. We will also consider the opposite embedding $\mathbb{R}^{2k+1} = (\mathbb{R}^{2k+1}, -(\sum_{0}^{k-1} p_i dq_i + dq_k)) \stackrel{\overline{\iota}:=a\circ\iota}{\longrightarrow} S^* \mathbb{R}^{k+1}$, where *a* is the antipodal map. In the forthcoming discussion, it will be important to remember that Gromov's h-principle applies to *co-oriented* contact manifolds; in particular, a high codimensional embedding $V \hookrightarrow \mathbb{R}^{2k+1}$.

We begin by assuming that our Maslov datum is induced by a polarization. So let ρ be a polarization of $\xi : V \to B(U/O)$ and $\sigma \coloneqq \rho^{-1}$ a polarization of the stable normal bundle. Fix a large codimensional (co-oriented) contact embedding $i : V \hookrightarrow \mathbb{R}^{2k+1} \stackrel{\iota}{\hookrightarrow} S^*(\mathbb{R}^{k+1})$. So σ is realized as a thickening V^{σ} of V, which is Lagrangian in the normal direction, and the sheaf $\mu \mathrm{sh}_{V;\rho}$ is given by $\mu \mathrm{sh}_{S^*(\mathbb{R}^{k+1});V^{\sigma}}|_V$. The microlocal Verdier dual $\mu \mathbb{D}_{\mathbb{R}^{k+1}} : \mu \mathrm{sh}_{S^*(\mathbb{R}^{k+1})} \to \mu \mathrm{sh}_{S^*(\mathbb{R}^{k+1})}^{op}$ then restricts to

$$\mu \mathbb{D}_{\mathbb{R}^{k+1}} : \mu \mathrm{sh}_{S^*(\mathbb{R}^{k+1});V^{\sigma}} |_V \to a^*_{\mathbb{R}^{k+1}} \, \mu \mathrm{sh}_{S^*(\mathbb{R}^{k+1});(V^{\sigma})^a} |_{V^a}.$$

But the target is exactly \overline{V} with the same polarization ρ , so we obtain a Verdier dual

(19)
$$\mu \mathbb{D}_{V;\rho} : \mu \mathrm{sh}_{V;\rho} \to \mu \mathrm{sh}_{\overline{V};\rho}^{op}$$

On a cosphere bundle equipped with its fiber polarization, we now have two Verdier duality functors. They agree:

Proposition 4.1. The identification $\mu sh_{S^*M;\phi_M} = \mu sh_{S^*M}$ in (8) intertwines the Verdier duality functors (19) and (4).

Proof. As usual, choose a large codimensional embedding of the base manifold $M \hookrightarrow \mathbb{R}^k$ and choose, up to a contractible space of choices, a splitting of the projection map $T^*\mathbb{R}^k|_M \to T^*M$ to obtain an inclusion $i: S^*M \hookrightarrow S^*\mathbb{R}^k$. One first sees that the antipodal map $a_{\mathbb{R}^k}$ on $S^*\mathbb{R}^k$ restricts to that on M so $\mu \operatorname{sh}_{\overline{S^*M};\phi_M} = a^* \mu \operatorname{sh}_{S^*M}$. The statement, up to microlocalization, then follows from the fact that, if $\iota: Z \hookrightarrow X$ is a closed inclusion of manifolds, then $\iota^*D_X(F) = D_Z(\iota^*F)$ for $F \in \operatorname{sh}(X)$.

We now extend the construction of a Verdier duality functor to contact manifolds equipped with a Maslov datum which does not necessarily come from a polarization. Tracing through the stabilization argument in [33, Section 11.1], we see that the above construction defines a morphism $\mu \mathbb{D}_{U/O(\xi)} : \mu \mathrm{sh}_{U/O(\xi)} \to \mu \mathrm{sh}_{\overline{U/O(\xi)}}^{op}$. We elaborate the notation $\overline{U/O(\xi)}$. A Lagrangian submanifold will remain Lagrangian when changing the symplectic form to its minus. Thus both V and \overline{V} shares the same Lagrangian Grassmannian bundle $\mathfrak{f} : LGr(\xi) \to V = \overline{V}$, viewed as fiber bundles. However, the relative cotangent bundle $T^*\mathfrak{f}$, for V, comes with a co-oriented contact structure λ_f described in [33, Lemma 10.5]. For the co-orientation reversing contact manifold \overline{V} , we will take the reversed co-orientation $-\lambda_f$ on $T^*\mathfrak{f}$ Now, the antipodal map $a : T^*\mathfrak{f} \to T^*\mathfrak{f}$ turn λ_f to $-\lambda_f$ and fixes the zero section $LGr(\xi) \hookrightarrow T^*\mathfrak{f}$. The discussion from the above paragraph thus produces a morphism $\mu \mathbb{D}_{LGr(\xi)} : \mu \mathrm{sh}_{LGr(\xi)} \to \mu \mathrm{sh}_{\overline{C}r(\xi)}^{op}$ where we use $\overline{LGr(\xi)}$ to emphasize that it is a set inside the co-orientation reserving contact manifold. Stabilizing the construction as in [33, Section 11.1] and we obtain the morphism $\mu \mathbb{D}_{U/O(\xi)} : \mu \mathrm{sh}_{U/O(\xi)} \to \mu \mathrm{sh}_{\overline{U}/O(\xi)}^{op}$.

Recall for a manifold M, we have $\mu \mathbb{D}_M(\mathcal{F} \otimes L) = \mu \mathbb{D}_M(F) \otimes L^{\vee}$, for any microsheaf \mathcal{F} and local system L on S^*M . Similar formulas holds in various situations, e.g. for the high codimensional embedding construction or when considering the relative cotangent bundle $T^*\mathfrak{f}$, we see that we are in the situation to apply Proposition 3.15. That is, we will apply it to the case when the functor between the stalk categories $t: F_1 \to F_2$ is given by talking the naive dual $(-)^{\vee}: \mathcal{C} \to \mathcal{C}$ which is given by $x^{\vee} := \underline{\mathrm{Hom}}(x, 1_{\mathcal{C}})$, and this allows us to descend the morphism $\mu \mathbb{D}_{U/O(\mathcal{E})}$

(20)
$$\mu \mathbb{D}_{BPic(\mathcal{C})} : \mu \mathrm{sh}_{BPic(\mathcal{C})(\xi)} \to \mu \mathrm{sh}_{\overline{BPic(\mathcal{C})(\xi)}}^{op}$$

and thus, for any Maslov data τ on V, we obtain the morphism $\mu \mathbb{D}_{V;\tau}$ in (18).

We end this section by discussing the compatibility of this global Verdier dual with the sheaf kernels obtained from the process of Maslov-to-kernel. We first recall a lemma regarding the compatibility of the usual Verdier dual with sheaf kernel convolution.

Lemma 4.2. Let M and N be manifolds and $K \in sh(M \times N)$ be a sheaf kernel. Assume K is constructible with perfect stalks, $ss(K) \cap T^*M \times 0_N \subseteq 0_{M \times N}$, and $supp(K) \hookrightarrow M \times N \to N$ is proper. Then for any $F \in sh(M)$, we have

$$D_N(K \circ (F)) = D_{M \times N}(K \otimes p_1^* \omega_M) \circ D_M(F)$$

Proof. Standard exercise using base change.

Proposition 4.3. In the setting of Theorem 3.8, there is an equivalence of morphisms

$$\mu \mathbb{D}_N(\mathcal{K}(h) \circ (-)) = \mu \mathbb{D}_{M \times N}(\mathcal{K}(h) \otimes p_1^* \omega_M) \circ \mu \mathbb{D}_M(-) : \mu \mathrm{sh}_{\mathcal{U}} \to \chi^* a_N^* \, \mu \mathrm{sh}_{\mathcal{V}}^{op}$$

Proof. Choose a large codimensional contact embedding $-\mathcal{U} \times \mathcal{V} \hookrightarrow S^*(\mathbb{R}^k \times \mathbb{R}^k)$ obtained from a large codimensional embedding of the form $M \times N \hookrightarrow \mathbb{R}^k \times \mathbb{R}^k$ as in the proof of Theorem 3.8. Now apply Lemma 4.2 to the GKS sheaf kernel used in Theorem 3.8. Note that we implicitly use the properness assumption when invoking [28, Proposition 6.3.3].

Proposition 4.4. In the setting of Corollary 3.19, there is an equivalence of morphisms

$$\mu \mathbb{D}_N(\mathcal{K}(h) \circ (-)) = \mu \mathbb{D}_{M \times N}(\mathcal{K}(h) \otimes p_1^* \omega_M) \circ \mu \mathbb{D}_M(-) : \mu \mathrm{sh}_{\mathcal{U}}^\eta \to \chi^* a_N^* \, \mu \mathrm{sh}_{\mathcal{V}}^{\xi, op} \, .$$

5. MICROSHEAVES ON COMPLEX CONTACT MANIFOLDS

From now on, we assume the coefficients C to be R - mod for some (discrete) ring R. We write $det^2 : U \to S^1 = B\mathbb{Z}$ for the colimit of $det_n^2 : U(n) \to S^1$. We also consider the second Stiefel-Whitney class $w_2 : BO \to B(\mathbb{Z}/2)$, induced from the exact sequence

$$1 \to \mathbb{Z}/2 \to \text{Pin} \to O \to 1.$$

Consider the ring homomorphism $\mathbb{Z} \to R$, which restricts to a group homomorphism $\mathbb{Z}/2 = \mathbb{Z}^{\times} \to R^{\times}$. We shall abuse notation and use w_2 to also denote the composition

$$w_2: BO \to B(\mathbb{Z}/2) \to BR^{\times}$$

As recalled in Section 3.1, for a real contact manifold V, there is a universal sheaf $\mu sh_{BPic(R)(\xi)}$ on $BPic(R)(\xi)$, in the case when $\mathcal{C} = R - mod$, the obstruction for $\mu sh_{BPic(R)(\xi)}$ to descend to V (here termed the *Maslov obstruction*) is given by

$$V \xrightarrow{\xi} BU \to B(U/O) \to B^2 Pic(R) = B^2 \mathbb{Z} \times B^3(R^{\times})$$

where the last map is given by the map $B \det^2 : B(U/O) \to B^2 \mathbb{Z}$ and the composition $B(U/O) \to B^2 O \xrightarrow{Bw_2} B^3 R^{\times}$, and a null-homotopy of the first is referred as a *grading* and that of the second an *orientation*. Here we notice that \det^2 vanishes on O and the map thus descends. Note that there is always a canonical orientation o_{can} which is given by composing with the canonical null-homotopy from the fiber sequence $BU \to B(U/O) \to B^2 O$.

Example 5.1. Let M be a (real) manifold. The fiber polarization ϕ_M provides both a grading gr_M and an orientation o_M . We've mentioned in Proposition 3.3 that $\mu \text{sh}_{S^*M;gr_M \times o_M} = \mu \text{sh}_{S^*M}$. By comparing the difference between o_M with the canonical orientation o_{can} , as detailed in Proposition B.7, one can conclude that, when replacing the fiber orientation by the canonical one, the associated microsheaves $\mu \text{sh}_{S^*M;gr_M \times o_{can}} = \mu \text{sh}_{S^*M}^{w_2(M)}$ is given by the usual microsheaves twisted by the second Stiefel-Whitney class.

Thus, in the situation when there are open subsets $\mathcal{U} \hookrightarrow S^*M$ and $\mathcal{V} \hookrightarrow S^*N$ and a contactomorphism $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{V}$, one needs only a homotopy $h : o_M|_{\mathcal{U}} = \chi^*(o_N|_{\mathcal{V}})$ in order to obtain an identification $\mu \operatorname{sh}_{\mathcal{U}}^{w_2(M)} = \mu \operatorname{sh}_{\mathcal{V}}^{w_2(N)}$.

Let (V, ξ) be a real contact manifold. The main observation in [8, Section 3] is that, if, up to stabilization, ξ is the underlying real symplectic vector bundle of a complex symplectic bundle Eon V, then an identification $E_{\mathbb{R}} = \xi$ provides ξ a grading. Now consider the setting when (V, ξ) is a complex contact manifold and we use $\pi : \tilde{V} \xrightarrow{\mathbb{C}^{\times}:1} V$ to denote its complex symplectization, which is a \mathbb{C}^{\times} -bundle over it. In this case, the real contact manifold $p : V_0 := \tilde{V}/\mathbb{R}_+$, as discussed in [8, Section 2], admits a grading coming from the complex symplectic vector bundle ξ .

Definition 5.2 ([8, Definition 5.1, Remark 5.4.]). Let o be an orientation of V_0 . We denote by $\mu \operatorname{sh}_{V_0,o}$ the microsheaves induced by the grading from the complex symplectic structure of ξ and the orientation o. When $o = o_{can}$ is the canonical orientation, we write simply $\mu \operatorname{sh}_{V_0}$. The sheaf $\mathbb{P} \, \mu \operatorname{sh}_V := p_* \, \mu \operatorname{sh}_{V_0}$ is defined to be the pushforward of $\mu \operatorname{sh}_{V_0}$.

As observed in [8, Lemma 5.6], a complex contactomorphism $\chi : V \to U$ between two complex contact manifold is, by definition, a biholomorphic map such that $\chi_*(\xi_V) = \xi_U$, so such a map automatically preserves the complex grading, even before stabilization, and induces canonically an identification $\chi^* \mu sh_U = \mu sh_V$.

In fact, as explained in [8, Remark 4.20], for a complex contact manifold V, the classifying map $V_0 \to B(U/O) \to B^2 Pic(\mathcal{C})$ always factorizes to $V_0 \to B^2 Pic(\mathcal{C})_0$ where we denote by $Pic(\mathcal{C})_0$ the connected component of $Pic(\mathcal{C})$. The sheaf $\mu \operatorname{sh}_{V_0}$ is thus a pullback of a universal sheaf $\mu \operatorname{sh}_{BPic(\mathcal{C})_0}(\xi)$ on $BPic(\mathcal{C})_0(\xi)$ whose monodromy on fibers are given by $Pic(\mathcal{C})_0$ instead of general objects in $Pic(\mathcal{C})$.

Example 5.3. Let X be a complex manifold. Its coprojective bundle \mathbb{P}^*X is a complex contact manifold, whose complex symplectization is the complex cotangent bundle T^*X . The main observation in [8, Lemma 3.2] is that its canonical grading is the same as the fiber polarization grading $gr_{X_{\mathbb{R}}}$, when viewing X as a real manifold $X_{\mathbb{R}}$. Concretely, both the complex symplectic vector bundle $T(T^*X)$ and the real vector bundle $TX_{\mathbb{R}}$ are obtained from the complex vector bundle TX, one by forming the quaternion bundle $(-) \otimes_{\mathbb{C}} \mathbb{H}$ and the other by taking the underlying real bundle $(-)_{\mathbb{R}}$, and both gives rise to the real symplectic vector bundle

$$T(T^*X_{\mathbb{R}}) = (T(T^*X))_{\mathbb{R}} = (TX_{\mathbb{R}}) \otimes_{\mathbb{R}} \mathbb{C}.$$

In summary, on the real contact manifold $S^*X_{\mathbb{R}}$, we have that the canonical microsheaves $\mu \mathrm{sh}_{S^*X_{\mathbb{R}}} = \mu \mathrm{sh}_{S^*X_{\mathbb{R}}}^{w_2(X)}$ are given by w_2 twisted microsheaves by Example 5.1, which we can push to \mathbb{P}^*X to obtain $\mathbb{P} \mu \mathrm{sh}_{P^*X} = \pi_* \mu \mathrm{sh}_{\dot{T}^*X}^{w_2(X)}$. Furthermore, if there is a complex contactomorphism $\chi : \mathcal{U} \to \mathcal{V}$ between $\mathcal{U} \subseteq \mathbb{P}^*X$ and $\mathcal{V} \subseteq \mathbb{P}^*Y$, there is a canonical identification $\mu \mathrm{sh}_{\mathcal{U}_0}^{w_2(X)} = \chi^* \mu \mathrm{sh}_{\mathcal{V}_0}^{w_2(Y)}$, which by Corollary 3.19, is induced by a twisted microsheaf kernel

(21)
$$\mathcal{K}(\chi) \in \mathbb{P} \mu \mathrm{sh}_{\Gamma^a_{\chi}}^{w_2(X \times Y)}(\mathcal{U} \times \mathcal{V}).$$

The next lemma shows that $\mathcal{K}(h)$ admits extra symmetry.

Lemma 5.4. We have a canonical identification $\mathcal{K}(\chi^{-1}) = v^* \mathcal{K}(\chi)$ where $v : X \times Y = Y \times X$ is the coordinates swapping map. In particular, $\mathcal{K}(\chi)[n]$ is Verdier self-dual (where $n = \dim_{\mathbb{C}} X$).

Remark 5.5. The case where \mathcal{U} is a ball is used implicitly in the proof of [28, Thm. 11.4.9].

Proof. The second statement is a direct consequence of the first statement and Corollary 3.19. To see the first statement, we apply [8, Lem. 5.6] to the induced map

$$(dv^{\vee}): P^*(Y \times X) \to P^*(X \times Y)$$

on the coprojective bundles. This implies that $\mathbb{P} \mu \operatorname{sh}_{P^*(X \times Y)} = (dv^{\vee})^* \mathbb{P} \mu \operatorname{sh}_{P^*(Y \times X)}$ canonically and it restricts to the subsheaves

$$\mathbb{P} \,\mu \mathrm{sh}_{P^*(X \times Y);\Gamma^a_{\chi}} = (dv^{\vee})^* \mathbb{P} \,\mu \mathrm{sh}_{P^*(Y \times X);\Gamma^a_{\chi^{-1}}},$$

and $\mathcal{K}(\chi)$ is sent to $\mathcal{K}(\chi^{-1})$ under this identification. However, realizing the equivalence under the identification $\mathbb{P} \mu \mathrm{sh}_{P^*X} = \pi_* \mu \mathrm{sh}_{T^*X}^{w_2(X)}$ implies that the equivalence is given by

(22)
$$v^* : \mathbb{P} \mu \mathrm{sh}_{P^*(X \times Y)}^{w_2(X \times Y)} = \mathbb{P} \mu \mathrm{sh}_{P^*(Y \times X)}^{w_2(Y \times X)},$$

microlocalized from $v^* : sh(X \times Y) = sh(Y \times X)$.

When there is another $\chi' : \mathcal{V} \to \mathcal{W}$ for some $\mathcal{W} \hookrightarrow \mathbb{P}^*Z$, the strict identification $(\chi' \circ \chi)_*\xi_{\mathcal{U}} = \chi'_*(\chi_*\xi_{\mathcal{U}}) = \chi'_*\xi_{\mathcal{V}}$ implies that there is a canonical identification

(23)
$$c_{\chi',\chi}: \mathcal{K}(\chi') \circ \mathcal{K}(\chi) = \mathcal{K}(\chi' \circ \chi) \in \mathbb{P} \mu \mathrm{sh}_{\Gamma^a_{\chi' \circ \chi}}^{w_2(X \times Z)}(\mathcal{U} \times \mathcal{W}).$$

A similar statement holds for 2-morphisms between such identification, in that the Čech 3-cocycle condition exhibited in Figure 1 holds automatically. That is, assume there is a further contactomorphism $\chi'' : W \to Q$, we will need to show that the two different compositions of identifications between microkernels,

$$\mathcal{K}(\chi'') \circ \mathcal{K}(\chi') \circ \mathcal{K}(\chi) \xrightarrow{\mathcal{K}(\chi'') \bigcirc c_{\chi',\chi}} \mathcal{K}(\chi'') \circ \mathcal{K}(\chi' \circ \chi) \xrightarrow{c_{\chi'',\chi' \circ \chi}} \mathcal{K}(\chi'' \circ \chi' \circ \chi),$$

and

$$\mathcal{K}(\chi'') \circ \mathcal{K}(\chi') \circ \mathcal{K}(\chi) \xrightarrow{c_{\chi'',\chi'} \bigcirc \mathcal{K}(\chi)} \mathcal{K}(\chi'' \circ \chi') \circ \mathcal{K}(\chi) \xrightarrow{c_{\chi''\circ\chi',\chi}} \mathcal{K}(\chi'' \circ \chi' \circ \chi)$$

are the same. Per Remark 3.21, it will be implied by the same equality between Maslov data. But, in this case, it is just the associativity for the identification $\chi''_*\chi'_*\chi_*\xi_{\mathcal{U}} = \xi_{\mathcal{Q}}$.

We end this section with a gluing description for $\mathbb{P} \mu \mathrm{sh}_V$. First, take a cover of V by complex Darboux charts $\{\mathcal{U}_{\alpha}\}$ with complex contact embeddings $f_{\alpha} : \mathcal{U}_{\alpha} \hookrightarrow \mathbb{P}^* X_{\alpha}$ for some complex manifold X_{α} . The above Example 5.3 implies that $\mathbb{P} \mu \mathrm{sh}_V |_{\mathcal{U}_{\alpha}} = f_{\alpha}^* \mu \mathrm{sh}_{P^* X_{\alpha}}^{w_2(\alpha)}$ where we use the notation $w_2(\alpha) := w_2(X_{\alpha})$, and we will further simplify the notation by $\mu \mathrm{sh}_{\alpha}^{w_2} := f_{\alpha}^* \mu \mathrm{sh}_{P^* X_{\alpha}}^{w_2(\alpha)}$. The complex contactomorphism, $\chi_{\beta\alpha} : f_{\alpha}(\mathcal{U}_{\alpha\beta}) \xrightarrow{\sim} f_{\beta}(\mathcal{U}_{\alpha\beta})$ produces a w_2 -microkernel and we denote its pullback on V by $\mathcal{K}_{\beta\alpha}$.

On triple overlaps, by Corollary 3.20, there is a canonical isomorphism between microkernels

$$c_{\gamma\beta\alpha}:\mathcal{K}_{\gamma\beta}\circ\mathcal{K}_{\beta\alpha}=\mathcal{K}_{\gamma\alpha}$$

Theorem 5.6. Once fixed a Darboux chart $\{(\mathcal{U}_{\alpha}, f_{\alpha})\}$, the sheaf $\mathbb{P} \mu sh_V$ can be obtained by the gluing data

(24) $(\{\mu sh_{\alpha}\}, \{\mathcal{K}_{\beta\alpha}\}, \{c_{\gamma\beta\alpha}\})$

where $\mathcal{K}_{\beta\alpha}$ are w_2 -microsheaf kernel satisfying $\mu \mathbb{D}_{\beta\alpha}(\mathcal{K}_{\beta\alpha}[2n]) = \mathcal{K}_{\beta\alpha}$ compatible with the identifications $c_{\gamma\beta\alpha}$. where $\dim_{\mathbb{C}} V = 2n - 1$ and $\mu \mathbb{D}_{\beta\alpha}$ is the corresponding Verdier dual defined in (18).

Remark 5.7. A priori, there is an infinite layer of compatible identifications when gluing a sheaf of categories. However, our identifications comes from the homotopical structure of Pic(R), which has $\pi_k(Pic(R)) = 0$ for k > 1, so all k-morphisms for k > 2 are automatically trivial.

Proof. By Proposition 4.4, the sheaf $\mu \operatorname{sh}_{\overline{V_0}}$ is glued by $\mu \mathbb{D}_{\beta\alpha}(\mathcal{K}_{\beta\alpha}[2n])$. However, reversing the co-orientation is invisible on the complex contact level and so $p_* \mu \operatorname{sh}_{\overline{V_0}} = \mathbb{P} \mu \operatorname{sh}_V$ as well.

Remark 5.8. Because the null-homotopy on the R^{\times} -component already happen before composing with $\mathbb{Z}^{\times} \to R^{\times}$. The above data can in fact be obtained from tensoring the same data over \mathbb{Z} with $(-) \otimes_{\mathbb{Z}} R$.

6. MODULES OVER TWISTED DIFFERENTIAL OPERATORS

In Section 2, we discussed twisting sheaves of categories. In the special case when the sheaf is formed by taking modules of an algebroid, we can instead twist the algebroid and consider ordinary modules. We will not try to pursue the general theory in this section but restrict ourselves to the case of the ring of differential operators \mathcal{D}_X .

6.1. Algebroids. Given a sheaf of categories \mathcal{A} on a space X, we denote by $\pi_0(\mathcal{A})$ the sheafification of the presheaf of sets $X \supseteq U \mapsto \pi_0(\mathcal{A}(U))$. (Recall that $\pi_0(-)$ of a category is the set of isomorphism classes of objects).

Recall that a \mathbb{C} -algebroid \mathcal{A} is a sheaf of \mathbb{C} -linear 1-categories such that locally $\pi_0(\mathcal{A}) = \{*\}$, i.e., \mathcal{A} is locally non-empty and all sections are isomorphic to each other. If A is a sheaf of \mathbb{C} -algebras, then one can consider the presheaf of categories $A^{+,pre}$ which assigns an open set U to the category with one object $\{*_U\}$ whose endomorphism is A(U). Its sheafification A^+ is a \mathbb{C} -algebroid. Although A^+ does not contain more data than A, there can be more algebroid morphisms than algebra morphisms.

A leisurely account of algebroids and their use in D-module theory can be found in [29, Sec. 2.1].

6.2. Line bundles. Let X be a complex manifold. We let $Pic(\mathcal{O}_X)$ be the algebroid whose objects are the (holomorphic) line bundles on X, whose morphisms are isomorphisms of line bundles, and whose multiplication is tensor product. More generally, we consider the "Picard stack" of \mathbb{C} -algebroids $Pic_X(-)$, which assigns to $U \subseteq X$ the algebroid $Pic(\mathcal{O}_U)$.

Lemma 6.1. $Pic_X(-) = \mathcal{O}_X^+$. Furthermore, under this identification, the canonical ring antiisomorphism $\mathcal{O}_X \xrightarrow{\sim} \mathcal{O}_X^{op}$ is identified with taking the dual line bundle

$$Pic_X \xrightarrow{(-)^{\vee}} Pic_X$$
$$L \mapsto L^{\vee} (= L^{-1})$$

Proof. For $U \subseteq X$, the category $\mathcal{O}_X^+(U)$ is given by $Pic(\mathcal{O}_U)$, the category of (holomorphic) line bundles on U. Indeed there is a morphism $\mathcal{O}_X^{+,pre}(U) \to Pic(\mathcal{O}_U)$ which send the point $\{*_U\}$ to the trivial line bundle \mathcal{O}_U with its effect on morphisms is the identity on $\mathcal{O}(U)$. This morphism is an equivalence by $\bar{\partial}$ -Poincaré lemma. For the second statement, we notice that when reversing the order of multiplication, a transition map $g_{\beta\alpha} \in \mathcal{O}^{\times}$ of L will be seen as going the reverse direction. To have the correct direction, one turns $g_{\beta\alpha}$ to $g_{\beta\alpha}^{-1}$ but the later is a cocycle of L^{-1} .

Remark 6.2. Let us recall the Čech description of line bundles, i.e. elements of $\pi_0(Pic(\mathcal{O}_X)) \simeq H^1(X, \mathcal{O}_X^{\times})$. It will be useful later to spell this out explicitly. Let U_{α} be a cover of X such that $L|_{U_{\alpha}}$ admits a section s_{α} and, on the overlap $U_{\beta\alpha}$, there exists a holomorphic function $g_{\beta\alpha} : U_{\beta\alpha} \to \mathbb{C}^{\times}$ such that $s_{\beta} = g_{\beta\alpha}s_{\alpha}$. Note that L is a (holomorphic) line bundle implies that $g_{\gamma\beta}g_{\beta\alpha} = g_{\gamma\alpha}$. In addition, the natural equivalence $L \otimes_{\mathcal{O}} L^{-1} = \mathcal{O}_X$ picks out sections s_{α}^* such that $\langle s_{\alpha}^*, s_{\alpha} \rangle = 1$. This later equality implies that $s_{\beta}^* = g_{\beta\alpha}^{-1}s_{\alpha}$. In more concise term, the there is an isomorphism of groups

$$\pi_0(Pic(X)) \xrightarrow{\sim} H^1(X; \mathcal{O}_X^{\times})$$
$$L \mapsto [(U_{\alpha}, g_{\beta\alpha})]$$

between holomorphic line bundles and equivalence classes of Čech cocycles.

We now discuss "fractional" line bundles. Let G be a closed subgroup of \mathbb{C}^* and fix $\eta : X \to B^2G$. Then we can consider the stack

$$Pic_X^{\eta}(-) := Pic_X(-) \otimes_{BG} P_{\eta}.$$

Here BG acts on Pic(U) for any $U \subset X$ by $G \to \Omega^*BG \to \Omega^*Aut(Pic(U)) = End_{\mathcal{O}_U-mod}(\mathcal{O}_U) = \mathcal{O}_X(U)$, which is the inclusion of G as constant functions into the ring of holomorphic function on U.

A $(\eta$ -)twisted line bundle is a global section of $Pic_X^{\eta}(-)$. Note that we have maps $Pic_X^{\eta}(-) \otimes Pic_X^{\mu+\mu}(-)$. In particular, if $G = \mathbb{Z}/k$, we have $Pic_X^{\eta}(X)^{\otimes k} \to Pic_X(X)$.

Example 6.3. If L is a line bundle on X, a k-th root of L is a line bundle $L^{1/k}$ along with an isomorphism $(L^{1/k})^{\otimes k} \simeq L$. Such a k-th root need not exist in general as a line bundle. However, we can always construct a k-th root as a *twisted* line bundle.

To do this, consider the exact sequence $0 \to \mathbb{Z}/k \to \mathcal{O}_X^* \xrightarrow{z \mapsto z^k} \mathcal{O}_X^* \to 1$. Let $\eta : X \to B^2(\mathbb{Z}/k)$ be image of $L \in H^1(X, \mathcal{O}_X^*) \to H^2(X, \overline{\mathbb{Z}/k})$ under the connecting map.

Then we can define a twisted sheaf $L^{1/k} \in Pic^{\eta}(X)$ concretely as follows: choose a cover $\{U_{\alpha}\}$ and choose a k-th root of the transition functions $f_{\alpha\beta}^{1/k}$. On triple overlaps, the cocycle condition is satisfied up to a \mathbb{Z}/k ambiguity, which (tracing through the definitions) is exactly what is needed to define an object $L^{1/k} \in Pic^{\eta}(X)$. By construction, $(L^{1/k})^{\otimes k} \simeq L$.

Note that a sufficient condition for L to admit a k-th root as a genuine line bundle is for the image of $L \in H^1(X, \mathcal{O}_X^*) \to H^2(X, \mathbb{Z}/k)$ to vanish. This condition is evidently also necessary since $H^2(X, \mathbb{Z}/k)$ is k-torsion.

Example 6.4. The case k = 2 shows that a line bundle L admits a square root iff $w_2(L) = 0$. In particular, the canonical bundle Ω_X admits a square root iff $w_2(\Omega_X) = 0$ iff $w_2(TX) = 0$ iff X is spin.

6.3. Twisted differential operators. Let X be a complex manifold and let \mathcal{D}_X be the sheaf of differential operators of X. This is a sheaf of non-commutative unital rings, and we have inclusions

$$\mathcal{O}_X \subset \mathcal{D}_X \subset \mathcal{E}nd_{\mathbb{C}}(\mathcal{O}_X).$$

The sheaf \mathcal{D}_X can be defined in multiple equivalent ways. Most concretely a section $D \in \mathcal{D}_X(U)$ is defined to be a section $D \in Hom_{\mathbb{C}}(\mathcal{O}_X(U), \mathcal{O}_X(U))$ which is locally of the form

$$D(x) = \sum_{\alpha} a_{\alpha}(x) \partial^{\alpha}.$$

Here $\alpha = (\alpha_1, \ldots, \alpha_k) \in \mathbb{N}^k$ ranges over all multi-indices of length $k \geq 0$, and we write $\partial^{\alpha} = \partial^{\alpha_1} \ldots \partial^{\alpha_n}$. We also mention the coordinate-free description, which goes back to Sato, as $\Gamma_{[\Delta]}(\mathcal{O}_X \boxtimes_{\mathcal{O}} \Omega_X)[-n]$ where for complex variety $V \subseteq X$, the functor $\Gamma_{[V]}$ is the temperate support in V [5, II.5].

Let L be a holomorphic line bundle on X. We denote by $\mathcal{D}_X^L \coloneqq L \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} \otimes L^{-1}$ the \mathcal{O}_X -algebra of differential operators twisted by L, whose multiplication is given by,

$$(L \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} \otimes L^{-1}) \otimes_{\mathcal{O}} (L \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} \otimes L^{-1}) = L \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} \otimes L^{-1} \to L \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} \otimes L^{-1}$$

where we use $L \otimes_{\mathcal{O}} L^{-1} = \mathcal{O}_X$ for the first equality.

Equivalently and more concretely, \mathcal{D}_X^L can be described by gluing data. Following the notation of Remark 6.2, let U_{α} be a cover of X such that $L|_{U_{\alpha}}$ admits a section s_{α} and, on the overlap $U_{\beta\alpha}$, there exists a holomorphic function $g_{\beta\alpha}: U_{\beta\alpha} \to \mathbb{C}^{\times}$ such that $s_{\beta} = g_{\beta\alpha}s_{\alpha}$. Then on U_{β} , a section of \mathcal{D}_X^L has the form $s_{\beta} \otimes P \otimes s_{\beta}^*$, which when restricting to $U_{\beta\alpha}$, becomes the same as

$$g_{\beta\alpha}s_{\alpha}\otimes P\otimes g_{\beta\alpha}^{-1}s_{\alpha}^{*}=s_{\alpha}\otimes g_{\beta\alpha}Pg_{\beta\alpha}^{-1}\otimes s_{\alpha}^{*}=s_{\alpha}\otimes \operatorname{Ad}(g_{\beta\alpha})(P)\otimes s_{\alpha}^{*}.$$

In short, \mathcal{D}_X^L is glued from $(\mathcal{D}_{U_{\alpha}}, \operatorname{Ad}(g_{\beta \alpha}))$ as an algebra.

We can also twist the sheaf of rings \mathcal{D}_X by a *fractional* line bundle.

Definition 6.5. We denote by $\mathcal{D}_X^{\sqrt{L}}$ the sheaf of \mathcal{O} -algebras $L^{1/2} \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} L^{-1/2}$.

The expression $L^{1/2} \otimes_{\mathcal{O}} \mathcal{D}_X \otimes_{\mathcal{O}} L^{-1/2}$ is a sheaf since $L^{1/2}$ and $L^{-1/2}$ have twistings inverse to each other. More concretely, by shrinking the open cover, we may assume that $g_{\beta\alpha}$ admits a square root $\sqrt{g_{\beta\alpha}} : U_{\alpha} \to \mathbb{C}^{\times}$. Then the sheaf $\mathcal{D}_X^{\sqrt{L}}$ is glued by the data $(\mathcal{D}_{U_{\alpha}}, \operatorname{Ad}(\sqrt{g_{\beta\alpha}}))$. We note that since $\operatorname{Ad}(\pm 1) = 1$, the expression does not depend on the choice of the square root $\sqrt{g_{\beta\alpha}}$.

6.4. **Twisted** \mathcal{D} -modules. A left/right \mathcal{D}_X -module is simply a left/right module over the sheaf of non-commutative rings \mathcal{D}_X . Similarly one defines left/right \mathcal{D}_X^L -module for any possibly twisted line bundle L. When we refer to \mathcal{D}_X -modules without any further adjective, we always mean left \mathcal{D}_X -modules.

Example 6.6 ([5, 2.5.17]). Tautologically, the structure ring \mathcal{O}_X is a left \mathcal{D}_X -module. Let $T \subseteq X$ be an analytic hypersurface. Then $\mathcal{O}(*T)$, the sheaf of meromorphic functions with poles contained in T, is a left \mathcal{D}_X -module which contains \mathcal{O}_X as a submodule.

Example 6.7. Recall that a vector field v acts on top forms Ω_X by the Lie derivative L_v . By Cartan's formula, for $v \in \Theta$ and $\omega \in \Omega_X$, it is simply given by

$$L_v(\omega) = (d\iota_v + \iota_v d)\omega$$

where ι_v is the natural contraction of forms by vector fields (at the first component). One can check that

$$\omega v \coloneqq -L_v \omega$$

equips Ω_X with the structure of a right \mathcal{D}_X -module. See for example [5, Thm. 1.2.14] or [25, Lem. 1.8].

In general, the ring \mathcal{D}_X is different from \mathcal{D}_X^L . But one categorical level up, the twisting is trivial.

Lemma 6.8 ([25, Prop. 1.9]). There is an equivalence of sheaves $\mathcal{D}_X^L - mod = \mathcal{D}_X - mod$.

Proof. The equivalence is simply given by

$$\mathcal{D}_X - mod \xrightarrow{\sim} \mathcal{D}_X^L - mod$$

 $\mathcal{M} \mapsto L \otimes_\mathcal{O} \mathcal{M}$

However, for the purpose of Proposition 6.9, we give a more involved proof from the point of view of gluing. We recall that, for sheaves of sets G, F, the limit

$$\operatorname{Hom}(G, F) = \lim \left(\prod_{\alpha \in I} \operatorname{Hom}(G_{\alpha}, F_{\alpha}) \rightrightarrows \prod_{\alpha, \beta \in I} \operatorname{Hom}(G_{\beta\alpha}, F_{\beta\alpha}) \right),$$

where the subscript indicates restrictions, states that, in order to define a morphism from G to F, it suffices to define them on each U_{α} and check that they agree on the overlap $U_{\beta\alpha}$. When C and D are sheaves of abelian categories, the existence of non-trivial 2-morphisms increase the length of the limit by one to

$$\operatorname{Hom}(C,D) = \lim \left(\prod_{\alpha \in I} \operatorname{Hom}(C_{\alpha}, D_{\alpha}) \rightrightarrows \prod_{\alpha, \beta \in I} \operatorname{Hom}(C_{\beta\alpha}, D_{\beta\alpha}) \rightrightarrows \prod_{\alpha, \beta, \gamma \in I} \operatorname{Hom}(C_{\gamma\beta\alpha}, D_{\gamma\beta\alpha}) \right).$$

That is, equality between two functors are now a structure, a natural equivalence, so we have to check whether they are compatible on triple overlap. (Since there is no nontrivial 3-morphism, equality on that level is again a property.)

By Corollary A.2, we have $L_{g_{\beta\alpha}}$: $\mathrm{id}_{\mathcal{D}-mod} \xrightarrow{\sim} _{\mathrm{Ad}(g_{\beta\alpha})}(-)$, and we can define a isomorphism from $\mathcal{D}^L - mod$ to $\mathcal{D} - mod$ by the following diagram:



To check that this functor is well-defined, we need to show that $L_{g_{\gamma\beta}} \circ L_{g_{\beta\alpha}} = L_{g_{\gamma\alpha}}$, which follows from the fact that $\{g_{\beta\alpha}\}$ is a Čech cocycle.

Proposition 6.9. There is an equivalence of sheaves $\mathcal{D}_X^{\sqrt{L}} - mod = (\mathcal{D}_X - mod)^{w_2(L)}$

Proof. Similarly to the proof of Lemma 6.8, we would like to untwist $\mathcal{D}^{\sqrt{L}} - mod$ by the following diagram:



On the left side, the natural transformations over the triple overlaps are given by id, since $Ad(\pm 1) = 1$. To absorb the twisting coming from the $L_{\sqrt{g_{\beta\alpha}}}$'s, those for the right hand side have to be the $c_{\gamma\beta\alpha}$'s.

6.5. The star anti-involution. The right module structure of Ω_X discussed in Example 6.7 provides an equivalence

$$\Omega_X \otimes_{\mathcal{O}_{\dot{T}^*X}} (-) : \mathcal{D}_X - \mathfrak{Mod} \xrightarrow{\sim} \mathcal{D}_X^{op} - \mathfrak{Mod}$$
$$\mathcal{M} \mapsto \Omega_X \otimes_{\mathcal{O}_X} \mathcal{M}.$$

This follows from the general fact that the tensor product of a left and a right *D*-module over \mathcal{O} has a right module structure as explained in [5, (4), 1.3.1 Theorem]. In our case, for $\mathcal{M} \in \mathcal{D}_X - mod$, the right module structure on $\Omega_X \otimes_{\mathcal{O}_X} \mathcal{M}$ is given by

$$(\omega \otimes m)v \coloneqq (-L_v \omega) \otimes m + \omega \otimes (-vm)$$

for $v \in \Theta$, $\omega \in \Omega$, and $m \in \mathcal{M}$. One can also view the equivalence as giving by the tautological left \mathcal{D}_X^{op} and right \mathcal{D}_X -module $\Omega_X \otimes_{\mathcal{O}_X} \mathcal{D}_X$ where the 'interesting' left \mathcal{D}_X^{op} -module structure is the one described above and the right \mathcal{D}_X -module structure coming from simply multiplying on the right. General Morita theory discussed in Lemma A.3, or rather its sheaf version studied in [9, Section 3], thus produces an \mathcal{O}_X -algebra isomorphism

(25)
$$*: \mathcal{D}_X \xrightarrow{\sim} \mathcal{D}_X^{\Omega_X, op}.$$

Proposition 6.10. The anti-involution $*: \mathcal{D}_X \xrightarrow{\sim} \mathcal{D}_X^{\Omega_X, op}$ is the usual star anti-involution (or formal adjoint). That is, in local coordinates (x_1, \ldots, x_n) , we have a natural section $dx_1 \wedge \cdots \wedge dx_n$ of Ω_X , and (25) is then just the map $P = \sum_{\alpha} a^{\alpha} \partial_{\alpha} \mapsto *(P) := P^* := \sum_{\alpha} (-1)^{|\alpha|} \partial_{\alpha} a^{\alpha}$.

Proof. Set $dx = dx_1 \wedge \cdots \wedge dx_n$. Then $dx \otimes 1 \in \Omega_X \otimes_{\mathcal{O}_X} \mathcal{D}_X$ forms a common generator of the two module structures. To see its effect, according to Lemma A.3, we have to apply P on $dx \otimes 1$ by the interesting \mathcal{D}_X^{op} action. But for any k, $\partial_k(dx \otimes 1) \coloneqq (-L_{\partial_k}dx \otimes 1) - (dx \otimes 1 \cdot \partial_k) = -dx \otimes \partial_k$. Thus, for any monomial $a(x)\partial^{\alpha}$, we have

$$(a(x)\partial^{\alpha}) \cdot (dx \otimes 1) \coloneqq (a(x) \cdot (\partial^{\alpha} \cdot (dx \otimes 1)) = a(x) \cdot (dx \otimes ((-1)^{|\alpha|}\partial^{\alpha}))$$
$$= dx \otimes ((-1)^{|\alpha|}\partial^{\alpha}) \cdot a(x)).$$

Lemma 6.11. There is an identification $(\mathcal{D}_X - \mathfrak{M}od)^{w_2(X)} = \mathcal{D}_X^{\sqrt{\Omega_X}} - \mathfrak{M}od$ where $\mathcal{D}_X^{\sqrt{\Omega_X}}$ is the ring of differential operators twisted by any square root of Ω_X . Under this identification, the equivalence $\Omega_X \otimes_{\mathcal{O}_X} (-) : (\mathcal{D}_X - \mathfrak{M}od)^{w_2(X)} = \mathcal{D}_X^{op} - \mathfrak{M}od^{w_2(X)}$ is identified with

$$\mathcal{D}_X^{\sqrt{\Omega_X}} - \mathfrak{Mod} = \mathcal{D}_X^{\sqrt{\Omega_X}, op} - \mathfrak{Mod}$$

 $\mathcal{M} \mapsto \mathcal{M}.$

Here we use the fact that any left $\mathcal{D}_X^{\sqrt{\Omega_X}}$ -module automatically admits right $\mathcal{D}_X^{\sqrt{\Omega_X}}$ -module structure to define the map.

Proof. By Proposition 6.9, the equivalence $\mathcal{D}_X^{\sqrt{\Omega_X}} - mod = (\mathcal{D}_X - mod)^{w_2(L)}$ is given by $\Omega_X^{-1/2} \otimes_{\mathcal{O}} (-)$. Since $(\mathcal{D}_X^L)^{op} = (\mathcal{D}_X^{op})^{L^{-1}}$, the same equivalence for the 'op' version is given by $\Omega_X^{1/2} \otimes_{\mathcal{O}} (-)$. But then the composition

$$\mathcal{D}_X^{\sqrt{\Omega_X}} - mod = (\mathcal{D}_X - mod)^{w_2(L)} = (\mathcal{D}_X^{op} - mod)^{w_2(L)} = \mathcal{D}_X^{\sqrt{\Omega_X, op}} - mod$$

is given by $\Omega_X^{-1/2} \otimes_{\mathcal{O}_X} \Omega_X \otimes_{\mathcal{O}_X} \Omega^{-1/2} \otimes_{\mathcal{O}_X} (-) = \mathcal{O}_X \otimes_{\mathcal{O}_X} (-)$, which does nothing on the underlying \mathcal{O}_X -module. Said differently, the bimodule which induces the equivalence $\mathcal{D}_X^{\sqrt{\Omega_X}} - mod = \mathcal{D}_X^{\sqrt{\Omega_X,op}} - mod$ is obtained from the bimodule $\Omega_X \otimes_{\mathcal{O}_X} \mathcal{D}_X$ by averaging the twisting

$$\Omega_X^{-1/2} \otimes_{\mathcal{O}_X} \Omega_X \otimes_{\mathcal{O}_X} \mathcal{D}_X \otimes_{\mathcal{O}_X} \Omega^{1/2} = \mathcal{D}_X^{\sqrt{\Omega_X}}$$

and thus is invisible on the O-module level.

7. THE CANONICAL SHEAF OF MICRODIFFERENTIAL OPERATORS

7.1. **Microlocal operators.** Let X be a complex manifold. There is a sheaf of algebras $\mathcal{E}_X^{\mathbb{R}}$ on T^*X whose sections are sometimes called *microlocal operators*. It was originally introduced in the foundational paper of Sato, Kawai, and Kashiwara [38, II.1.1, II.1.2].

In this subsection, we summarize the construction and some key properties of $\mathcal{E}_X^{\mathbb{R}}$ following [28, Sec. 11.4]. We will ultimately be interested in a "tempered" variant $\mathcal{E}_X^{\mathbb{R},f}$ which will be introduced in the next subsection. However, since both versions enjoy the same formal properties, it seems pedagogically preferable to begin our discussion with $\mathcal{E}_X^{\mathbb{R}}$.

Let X_1, X_2, X_3 be manifolds. Following the notation of [28, Section 7.1], we let $q_{ij} : X_1 \times X_2 \times X_j \to X_i \times X_j$ be the projection. Similarly we let $p_{ij} := T^*X_1 \times T^*X_2 \times T^*X_j \to T^*X_i \times T^*X_j$ be the projection on cotangent bundles, and write p_{ij}^a for the composition of p_{ij} with the antipodal map on the *j*-th component. For sheaf kernels $K_1 \in sh(X_1 \times X_2)$, and $K_2 \in sh(X_2 \times X_3)$ we have the convolution

(26)
$$K_1 \circ K_2 := q_{13!}(q_{23}^* K_1 \otimes q_{12}^* K_2)$$

The functor μ hom(-, -) intertwines convolution with tensor product [28, Proposition 4.4.11]. More precisely, given sheaf kernels $K_1, F_1 \in sh(X_1 \times X_2)$ and $K_2, F_2 \in sh(X_X X_3)$, we have

(27)
$$p_{13!}^a(p_{23}^{a*}\mu \hom(K_1, F_1) \otimes p_{12}^{a*}\mu \hom(K_2, F_2)) \to \mu \hom(K_1 \circ K_2, F_1 \circ F_2).$$

Let us now identify $N^*_{\Delta_X}(X \times X) = T^*X$ by projecting onto the first component. We shall also write $\Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_Y := q_1^* \Omega_X \otimes_{q_1^* \mathcal{O}_X} \mathcal{O}_{X \times Y}$

We now consider the sheaf of abelian groups

(28)
$$\mathcal{E}_X^{\mathbb{R}} \coloneqq \mu \hom(\mathbb{C}_{\Delta_X}[-n], \Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_X) \in sh(T^*X)^{\heartsuit}$$

It can be shown [28, Prop. 11.4.1, 11.4.4] that $\mathcal{E}_X^{\mathbb{R}}$ has a ring structure. This is ultimately because of the fact that $\mathbb{C}_{\Delta_X} \circ \mathbb{C}_{\Delta_X} = \mathbb{C}_{\Delta_X}$. The 'multiplication' comes from the 'integration' morphism [28, Theorem 11.1.4]

(29)
$$f_!\Omega_Y[\dim_{\mathbb{C}} Y] \to \Omega_Z[\dim_{\mathbb{C}} Z],$$

which exists for any holomorphic map $f: Y \to Z$, specializing to the case of $q_2: X \times X \to X$. Similarly, for any microsheaf F, the sheaf $H^k \mu \text{hom}(F, \mathcal{O}_X)$ (resp. $H^k \mu \text{hom}(F, \Omega_X)$) has left (resp. right) $\mathcal{E}_X^{\mathbb{R}}$ -module structure. In particular, when taking $F = 1_X$, we see that \mathcal{O}_X (resp. Ω_X) is a left (resp.) $\mathcal{E}_X^{\mathbb{R}}$ -module. See also [27, Proposition 9.3.1].

7.2. From microsheaf kernels to \mathcal{E} kernels. Let $\mathcal{U} \subseteq \dot{T}^*X$, $\mathcal{V} \subseteq \dot{T}^*Y$ and let $\chi : \mathcal{U} \xrightarrow{\sim} \mathcal{V}$ be a complex homogeneous symplectomorphism. As discussed in Example 5.3, there is a canonical (twisted)-microkernel

$$\mathcal{K}(\chi) \in \mu \mathrm{sh}^{w_2(X \times Y)}(-\mathcal{U} \times \mathcal{V})$$

such that convolution with $\mathcal{K}(\chi)$ induces an equivalence

$$\mathcal{K}(\chi): \mu \mathrm{sh}^{w_2(X)}(\mathcal{U}) \xrightarrow{\sim} \mu \mathrm{sh}^{w_2(Y)}(\mathcal{V})$$

between microsheaves. Define

(30)
$$\mathcal{H}(\chi) \coloneqq \mu \hom(\mathcal{K}(\chi)[-n], \Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_Y) \in sh^{w_2(X \times Y)}(-\mathcal{U} \times \mathcal{V})^{\heartsuit}$$

For the next proposition, we will use the notation $\mathcal{E}_X^{\mathbb{R}} - \mathfrak{M}$ od to denote the sheaf of categories whose sections on \mathcal{U} is $\mathcal{E}_X^{\mathbb{R}} - \mathfrak{M}$ od $(\mathcal{U}) := \mathcal{E}_X^{\mathbb{R}}|_{\mathcal{U}} - mod$.

Proposition 7.1. $\mathcal{H}(\chi)$ is a right $r_1^* \mathcal{E}_X^{\mathbb{R}}$ and a left $r_2^* \mathcal{E}_Y^{\mathbb{R}}$ -module where $r_1 : \Gamma_{\chi}^a \xrightarrow{\sim} \mathcal{U}$ is the projection and similarly for r_2 . Furthermore, tensoring with the bimodule $\mathcal{H}(h)$ induces an equivalence

$$\mathcal{H}(\chi) \otimes_{\mathcal{E}_X^{\mathbb{R}}} (-) : (\mathcal{E}_X^{\mathbb{R}} - \mathfrak{Mod})^{w_2(X)}|_{\mathcal{U}} \xrightarrow{\sim} (\mathcal{E}_Y^{\mathbb{R}} - \mathfrak{Mod})^{w_2(Y)}|_{\mathcal{V}},$$

where $\mathcal{H}(\chi) \otimes_{\mathcal{E}_X^{\mathbb{R}}} \mathcal{M} \coloneqq r_{2*}(\mathcal{H}(\chi) \otimes_{r_1^* \mathcal{E}_X^{\mathbb{R}}|_{\mathcal{U}}} r_1^* \mathcal{M})$, between the (twisted) module categories.

Proof. The bimodule structure is explained in [28, Lemma 11.4.3], and is the same structure used to define the ring structure of $\mathcal{E}_X^{\mathbb{R}}$. To show the equivalence, it is enough by Morita theory to find an inverse $(\mathcal{E}_X^{\mathbb{R}}, \mathcal{E}_Y^{\mathbb{R}})$ -bimodule. We claim that this bimodule is

$$\mathcal{H}(\chi^{-1}) \coloneqq \mu \hom(\mathcal{K}(\chi^{-1})[n], \Omega_Y \boxtimes_{\mathcal{O}} \mathcal{O}_X).$$

Indeed, the tensor product

$$\mathcal{H}(\chi^{-1}) \otimes_{\mathcal{E}_Y^{\mathbb{R}}} \mathcal{H}(\chi) \to \mu \mathrm{hom} \left(\mathcal{K}(\chi^{-1})[-n] \circ \mathcal{K}(\chi)[-n], (\Omega_Y \boxtimes_{\mathcal{O}} \mathcal{O}_X) \circ (\Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_Y) \right) \\ \to \mu \mathrm{hom} \left(1_{\Delta_X}[-n], \Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_X \right) = \mathcal{E}_X^{\mathbb{R}}.$$

admits a canonical map to $\mathcal{E}_X^{\mathbb{R}}$ and similarly to $\mathcal{E}_Y^{\mathbb{R}}$. To define the second arrow, we are using (i) the fact that $\mathcal{K}(\chi^{-1}) \circ \mathcal{K}(\chi) = 1_{\Delta_X}$ canonically and (ii) the integration map $(\Omega_Y \boxtimes_{\mathcal{O}} \mathcal{O}_X) \circ (\Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_X)$

 \mathcal{O}_Y) $\rightarrow \Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_X[-n]$; see [28, Lem. 11.4.3]. Thus it is sufficient to check on stalks that this map is an equivalence. But this is [28, Theorem 11.4.9] once we restrict to smaller open sets. \Box

Similarly to the discussion in Section 6.5, there is an anti-involution on $\mathcal{E}_X^{\mathbb{R}}$. The slight complication, as explained in [39, 5.2] or [25, Theorem 7.7], is that the map $*: \mathcal{E}_X^{\mathbb{R}} \to a^* \mathcal{E}_X^{\mathbb{R},\Omega_X,op}$ now reverses the cotangent direction, due to the fact that ξ is sent to $-\xi$. Similarly to Lemma 6.8, the twisting is invisible at the module level and is given by tensoring with the line bundle Ω_X .

Lemma 7.2. There is an equivalence of sheaves of categories

$$\Omega_X \otimes_{\mathcal{O}_{T^*X}} (-) : \mathcal{E}_X^{\mathbb{R}} - \mathfrak{M} \mathrm{od} \xrightarrow{\sim} a^* \mathcal{E}_X^{\mathbb{R}, op} - \mathfrak{M} \mathrm{od}$$

whose inverse is given by $\Omega_X^{-1} \otimes_{\mathcal{O}_{T^*X}} (-)$.

(

Proposition 7.3. We have the following commuting diagram:

Here, for the bottom row, we identify $(a_X^* \mathcal{E}_X^{\mathbb{R},op} - \mathfrak{Mod})^{w_2(X)}|_{\mathcal{U}}$ with $(\mathcal{E}_X^{\mathbb{R},op} - \mathfrak{Mod})^{w_2(X)}|_{\mathcal{U}^a}$ and similarly for Y.

Proof. We see in the proof of Proposition 7.1 that $\mathcal{H}(\chi^{-1}) \in \operatorname{sh}(\mathcal{V} \times \mathcal{U})^{\heartsuit}$ is a sheaf viewed as an $(\mathcal{E}_X^{\mathbb{R}}, \mathcal{E}_Y^{\mathbb{R}})$ -bimodule using χ . Since the bimodule structure is given by [28, Lemma 11.4.3], a microlocal version of convolution, in order to consider its opposite bimodule, we have to swap its coordinate in a way similar to Lemma A.4. That is,

$$\mathcal{H}(\chi^{-1})^{op} = v^* \,\mu \mathrm{hom}(\mathcal{K}(\chi^{-1})[-n], \Omega_Y \boxtimes_{\mathcal{O}} \mathcal{O}_X) \in \mathrm{sh}(\mathcal{U} \times \mathcal{V})^{\heartsuit}.$$

We thus begin in the bottom left corner of the diagram; we shall argue that traveling around the left, top, and right arrows is the same as traveling along the bottom arrow and we would like to see $\mathcal{H}(\chi^{-1})^{op}$.

The first itinerary is realized by the composition of bimodules

(31)

$$\Omega_{Y} \otimes \mathcal{H}(\chi) \otimes_{\mathcal{E}_{X}^{\mathbb{R}}} \Omega_{X}^{-1} \simeq \mu \hom(1_{Y}, \Omega_{Y}) \otimes \mathcal{H}(\chi) \otimes_{\mathcal{E}_{X}^{\mathbb{R}}} \mu \hom(1_{X}, \Omega_{X}^{-1}) \simeq \mu \hom(\mathcal{K}(\chi)[-n], \mathcal{O}_{X} \boxtimes_{\mathcal{O}} \Omega_{Y}) \in sh(\mathcal{U} \times \mathcal{V})^{\heartsuit}$$

Applying v^* (see 22) transforms (31) into

(32)
$$\mu \hom(v^* \mathcal{K}(\chi)[-n], \Omega_Y \boxtimes_{\mathcal{O}} \mathcal{O}_X) \in \operatorname{sh}(\mathcal{V} \times \mathcal{U})^{\heartsuit}.$$

But by Lemma 5.4, (32) may be rewritten as

(33)
$$\mu \hom(\mathcal{K}(\chi^{-1})[-n], \Omega_Y \boxtimes_{\mathcal{O}} \mathcal{O}_X) \in \operatorname{sh}(\mathcal{V} \times \mathcal{U})^{\heartsuit}.$$

As explained in the proof of Proposition 7.1, (33) is the bimodule

$$\mathcal{H}(\chi^{-1}) \otimes_{\mathcal{E}_Y^{\mathbb{R}}} (-) : (\mathcal{E}_Y^{\mathbb{R}} - \mathfrak{M}od)^{w_2(Y)}|_{\mathcal{V}} \xrightarrow{\sim} (\mathcal{E}_X^{\mathbb{R}} - \mathfrak{M}od)^{w_2(X)}|_{\mathcal{U}}.$$

Finally, we apply v^* (undoing our previous application of v^*) to obtain

(34)
$$\mathcal{H}(\chi^{-1})^{op} \otimes_{\mathcal{E}_X^{\mathbb{R},op}} (-) : (\mathcal{E}_X^{\mathbb{R},op} - \mathfrak{M}od)^{w_2(X)}|_{\mathcal{U}^a} \to (\mathcal{E}_Y^{\mathbb{R},op} - \mathfrak{M}od)^{w_2(Y)}|_{\mathcal{V}^a}$$
as desired.

Remark 7.4. The only non-formal step in the proof of Proposition 7.3 is the appeal to Lemma 5.4. This relies on our assumption that χ is a *complex* homogeneous symplectomorphism, and can be understood as a manifestation of Verdier duality. Proposition 7.3 would be false if we only assumed that χ was a real homogeneous symplectomorphism.

Corollary 7.5. In the situation of Proposition 7.3, if there is another $\chi' : \mathcal{V} \to \mathcal{W}$ for some $\mathcal{W} \hookrightarrow \mathbb{P}^*Z$, then the commuting diagrams for χ and χ' composes to the same diagram for $\chi' \circ \chi$.

Proof. Because $\Omega_Y^{-1} \otimes_{\Omega_{T^*Y}} \Omega_Y = \mathcal{O}_{T^*Y}$ canonically, the composition of the diagram of χ and χ' can thus be identified as the diagram

$$(\mathcal{E}_{X}^{\mathbb{R}} - \mathfrak{Mod})^{w_{2}(X)}|_{\mathcal{U}} \xrightarrow{\mathcal{H}(\chi) \otimes_{\mathcal{E}_{X}^{\mathbb{R}}} (-)} (\mathcal{E}_{Y}^{\mathbb{R}} - \mathfrak{Mod})^{w_{2}(Y)}|_{\mathcal{V}} \xrightarrow{\mathcal{H}(\chi') \otimes_{\mathcal{E}_{Y}^{\mathbb{R}}} (-)} (\mathcal{E}_{Z}^{\mathbb{R}} - \mathfrak{Mod})^{w_{2}(Z)}|_{\mathcal{W}}} \int_{\Omega_{Z} \otimes_{\mathcal{O}_{T^{*}Y}} (-)} (\mathcal{O}_{X} \otimes_{\mathcal{O}_{T^{*}X}} (-)) \qquad (\mathcal{O}_{X} \otimes_{\mathcal{O}_{T^{*}X}} (-)) \qquad (\mathcal{O}_{X} \otimes_{\mathcal{O}_{T^{*}Y}} (-)) = (a_{X}^{*} \mathcal{E}_{X}^{\mathbb{R},op} - \mathfrak{Mod})^{w_{2}(Y)}|_{\mathcal{V}} \xrightarrow{\mathcal{O}_{Z} \otimes_{\mathcal{O}_{T^{*}Y}} (-)} (a_{Z}^{*} \mathcal{E}_{Z}^{\mathbb{R},op} - \mathfrak{Mod})^{w_{2}(Z)}|_{\mathcal{W}}$$

where the bottom arrows are similarly given by the 'op' version of the top arrows. But similar to the proof of Proposition 7.1, there is a morphism $\mathcal{H}(\chi') \otimes_{\mathcal{E}_{V}^{\mathbb{R}}} \mathcal{H}(\chi) \to \mathcal{H}(\chi' \circ \chi)$ given by the morphism

$$p_{13!}^{a}(p_{23}^{a*} \mu \hom(\mathcal{K}(\chi'), \Omega_{Y} \boxtimes_{\mathcal{O}} \mathcal{O}_{Z}) \otimes p_{12}^{a*} \mu \hom(\mathcal{K}(h), \Omega_{X} \boxtimes_{\mathcal{O}} \mathcal{O}_{Y})) \rightarrow \mu \hom(\mathcal{K}(\chi') \circ \mathcal{K}(\chi), (\Omega_{Y} \boxtimes_{\mathcal{O}} \mathcal{O}_{Z}) \circ (\Omega_{X} \boxtimes_{\mathcal{O}} \mathcal{O}_{Y})) \rightarrow \mu \hom(\mathcal{K}(\chi') \circ \mathcal{K}(\chi), \Omega_{X} \boxtimes_{\mathcal{O}} \mathcal{O}_{Z}) = \mu \hom(\mathcal{K}(\chi' \circ \chi), \Omega_{X} \boxtimes_{\mathcal{O}} \mathcal{O}_{Z})$$

where we use the (23) for the last equality. But [28, Proposition 11.4.7] then implies that this map is an equivalences. A similar computation holds for the 'op' version as well.

7.3. E-modules and microlocal Riemann-Hilbert. For the purpose of microlocal Riemann-Hilbert, the sheaf $\mathcal{E}_X^{\mathbb{R}}$ is too large. In particular, this sheaf is only real conic, i.e., conic with respect to \mathbb{R}_+ , but not complex conic, i.e., conic with respect to \mathbb{C}^{\times} . This is reflected in the fact that $\mathcal{E}_X^{\mathbb{R}}$ can contain functions of the form $\log(\xi)$.

A naive way to enforce respect for \mathbb{C}^{\times} is the following. Let $\gamma: \dot{T}^*X \to \mathbb{P}^*X$ be the quotient map by \mathbb{C}^{\times} , and consider

$$\mathcal{E}_X^\infty := \gamma^* \gamma_* \mathcal{E}_X^\mathbb{R}.$$

Sections of this were called "microdifferential operators of infinite order" in [5, 8.2.15], and "pseudo-differential operators" in [38].

Remark 7.6. As mentioned at the beginning of [38, 2.1] or [28, Remark 11.4.5], the sheaf of (infinite order) differential operator can be obtained by

$$\mathcal{D}_X^{\infty} = \Gamma_{\Delta_X}(\Omega_X \boxtimes_\mathcal{O} \mathcal{O}_X)[-n].$$

In fact, certain important 'finite degree' properties are lost in \mathcal{E}_X^{∞} . This is a microlocal version of the following issue in the theory of \mathcal{D} -modules: for a closed analytic set $Z \subseteq X$, there is an important difference between sections with support in Z and sections with 'temperate' support in Z (see e.g. [5, II.5]).

There is a better version, denoted by \mathcal{E}_X ⁵, whose sections are called (finite-order) microdifferential operators. In [25, 5, 39], this sheaf of rings is introduced by taking formal series of differential operators, adjoining ' ∂_x^{-1} ', to obtain what is commonly denoted as $\hat{\mathcal{E}}_X$ ⁶, and imposing convergence conditions. More convenient to us is a variant of the original approach to \mathcal{E}_X in [38], developed further by Andronikof [2], where the ring structure is functorially extracted from μ hom. Some discussions from a modern point of view can be found in [16, 35].

The key ingredient is the tempered microlocal hom [2, Def. 2.3.1]:

$$t$$
- μ hom $(-, \mathcal{O}_X)$: sh _{\mathbb{R},c} $(X) \to$ sh (T^*X) ,

where $sh_{\mathbb{R},c}(X)$ is the category of \mathbb{R} -constructible sheaves.

The construction of \mathcal{E}_X is completely parallel to that of $\mathcal{E}^{\mathbb{R}}$ in the previous subsection, except that all occurrences of μ hom are replaced with t- μ hom. Namely, one first defines

$$\mathcal{E}_X^{\mathbb{R},f} \coloneqq \operatorname{t-}\mu\operatorname{hom}(\mathbb{C}_{\Delta_X}[-1], \Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_X) \in \operatorname{sh}(T^*X)^{\heartsuit}.$$

Then one sets

$$\mathcal{E}_X := \gamma^* \gamma_* \mathcal{E}_X^{\mathbb{R}, f} \hookrightarrow \mathcal{E}_X^{\mathbb{R}, f},$$

where we recall that $\gamma : \dot{T}^*X \to \mathbb{P}^*X$ is the quotient map. Andronikof checks that this agrees with other definitions [2, Cor. 5.5.2, Cor. 5.6.1].

Since t- μ hom and μ hom enjoy the same formal properties, the algebras $\mathcal{E}_X^{\mathbb{R},f}$ and \mathcal{E}_X correspondingly satisfy all the properties enjoyed by $\mathcal{E}^{\mathbb{R}}$ which were discussed in the previous subsection ([2, Cor. 5.5.2, Cor. 5.6.1]). For example, H^j t- μ hom (F, \mathcal{O}_X) (resp. $H^j\gamma^*\gamma_*$ t- μ hom (F, \mathcal{O}_X)) is a left $\mathcal{E}_X^{\mathbb{R},f}$ (resp. \mathcal{E}_X) module. Similarly, quantized contact transforms exist both for $\mathcal{E}^{\mathbb{R},f}$ -modules and \mathcal{E} -modules.

The works of Andronikov and Waschkies [1, 43] microlocalized the usual Riemann-Hilbert correspondence to projectivized cotangent bundles:

Theorem 7.7 (Local microlocal Riemann–Hilbert; Andronikov [1] Waschkies [43]). Let X be a complex manifold. There is an equivalence

$$\mu \operatorname{RH}_X : \mathbb{P}erv_{\mathbb{P}^*X} \xrightarrow{\sim} \mathcal{E}_X - \mathfrak{M}od_{rh}$$

between perverse microsheaves on \mathbb{P}^*X and regular holonomic \mathcal{E}_X -modules. Furthermore, the equivalence respects contact transform [43, Corollary 3.4.3]. That is, in the setting of Section 7.2, we have the following commuting diagram:

$$\begin{aligned} & \mathbb{P}erv_{\mathcal{U}}^{w_{2}(X)} \xrightarrow{\mathcal{K}(\chi) \circ (-)} & \mathbb{P}erv_{\mathcal{V}}^{w_{2}(Y)} \\ & \downarrow & \mu \mathrm{RH}_{X} \mid_{\mathcal{U}} & (\overset{\circlearrowright}{\longrightarrow} & \downarrow & \mu \mathrm{RH}_{Y} \mid_{\mathcal{V}} \\ & \downarrow & \mathcal{H}(\chi) \otimes_{\mathcal{E}_{X}^{\mathbb{R}}} (-) & (\mathcal{E}_{Y}^{\mathbb{R}} - \mathfrak{M}\mathrm{od}_{rh})^{w_{2}(Y)} \mid_{\mathcal{V}} \end{aligned}$$

⁵Originally denoted by \mathcal{P}_X^f in [38, Definition1.5.6]

⁶This is the ring of 'formal' microdifferential operators, which [25] mostly works with.

Lastly, we mention that the algebra \mathcal{E}_X has the nice Morita-theoretical property of being Picard good. See Definition A.5 for the definition.

Theorem 7.8 ([9, Theorem 4.3.6]). The \mathbb{C} -algebra \mathcal{E}_X is Picard good.

7.4. The microlocal Riemann–Hilbert correspondence for complex contact manifolds. We recall Kashiwara's quantization of contact manifolds and Polsello's uniqueness criterion.

Theorem 7.9 ([24, Theorem 2], [34, Theorem 3.3]). Let (V, \mathcal{H}) be a complex contact manifold. There exists a unique \mathbb{C} -algebroid \mathcal{E}_V admitting the following structures

- (i) \mathcal{E}_V is filtered. (This means that for any $U \subset V$ and an objects $K, L \in \mathcal{E}_V(U)$, the \mathbb{C} -module $Hom_{\mathcal{E}_V(U)}(K, L)$ is filtered, the filtrations are compatible with composition, etc. see [34, Sec. 2]).
- (ii) there is an isomorphism of commutative algebroid stacks

$$\sigma: Gr(\mathcal{E}_V) \simeq \left(\bigoplus_{m \in \mathbb{Z}} \mathcal{L}^m\right)^{-1}$$

where \mathcal{L} is the line bundle $(TV/\mathcal{H})^{\vee}$, and its zeroth degree associated graded is given by taking dual line bundle $(-)^{\vee}$, and on each local Darboux chart $(\mathcal{E}_V, *)$ is equivalent to the canonical filtered algebroid \mathcal{E}_Y on $(\mathbb{P}^*Y, *)$.

(iii) \mathcal{E}_V is equipped with an anti-involution * (by definition: an equivalence of stacks $*: \mathcal{E}_V \to \mathcal{E}_V^{op}$ along with an invertible natural transformation $\epsilon: *^2 \Rightarrow \mathrm{id}$ such that $\epsilon \circ id_*: *^3 \Rightarrow *$ and $id_* \circ \epsilon: * \Rightarrow ^3$ are inverse).

These structures must be compatible; this means that * is a filtered functor, ϵ is a filtered natural transformation, and there exists an invertible natural transformation

(35)
$$\delta_0: \sigma_0 \circ Gr_0(*) \Rightarrow D \circ \sigma_0$$

such that the following diagram commutes:

(36)

$$\sigma_{0} \circ Gr_{0}(*^{2}) \xrightarrow{\delta_{0} \circ id_{Gr_{0}(*)}} D \circ \sigma_{0} \circ Gr_{0}(*)$$

$$\downarrow id_{\sigma_{0}} \circ Gr_{0}(\epsilon) \qquad id_{D} \circ \delta_{0} \uparrow \uparrow$$

$$\sigma_{0} \xrightarrow{\simeq} D^{2} \circ \sigma_{0}^{2}$$

Here $D: Pic_V \to Pic_V^{op}$ is the functor sending a line bundle to its dual.

We note that the notion of holonomic and regular holonomic is well-defined microlocally ([25, Chapter 8], [5, Chapter VIII]) and the work of Andronikov [2, Chapter 5] and Waschkies [43, Section 3] shows that they are invariant under quantized contact transform. Similarly on the microsheaf side, we show in our previous paper [8, Section 6], the notion of perverse *t*-structure exists on the canonical microsheaf category μsh_V and, on Darboux charts, it coincides with the usual notion.

Definition 7.10. We denote by $\mathbb{P} \mu \text{sh}_{V;\mathbb{C}-c}^b$ the subsheaf of $\mathbb{P} \mu \text{sh}_V$ consisting of complex constructible microsheaves with perfect microstalks, $\mathbb{P}erv_V \subseteq \mathbb{P} \mu \text{sh}_{V;\mathbb{C}-c}^b$ the subsheaf (of abelian categories) given by the heart of the perverse *t*-structure. Similarly, we denote by $\mathcal{E}_V - \mathfrak{M}od_h$ (resp. $\mathcal{E}_V - \mathfrak{M}od_{rh}$) the subsheaf of $\mathcal{E}_V - \mathfrak{M}od$ consisting of holonomic (resp. regular holonomic) \mathcal{E}_V -modules. We turn now to our main task, of globalizing the Andronikov-Waschkies results to a comparison of $\mathbb{P}erv_V$ and $\mathcal{E}_V - \mathfrak{M}od_{rh}$ on an arbitrary complex contact manifold V.

Let $\mathcal{U} \hookrightarrow \mathbb{P}^*X$, $\mathcal{V} \hookrightarrow \mathbb{P}^*Y$ be open subsets and $\chi : \mathcal{U} \to \mathcal{V}$ be a complex contactomorphism. Then we define the following "tempered" analog of (30):

(37)
$$\mathcal{H}(\chi) \coloneqq \gamma^* \gamma_* \operatorname{t-}\mu \operatorname{hom}(\mathcal{K}(\tilde{\chi})[-n], \Omega_X \boxtimes_{\mathcal{O}} \mathcal{O}_Y) \in \operatorname{sh}^{w_2(X \times Y)}(\mathcal{U} \times \mathcal{V})^{\heartsuit}$$

Proposition 7.11. Tensoring with the bimodule $\mathcal{H}(\chi)$ (resp. $\mathcal{H}(\chi^{-1})^{op}$) induces an equivalence

$$\mathcal{H}(\chi) \otimes_{\mathcal{E}_X} (-) : (\mathcal{E}_X - \mathfrak{M}\mathrm{od})^{w_2(X)}|_{\mathcal{U}} \xrightarrow{\sim} (\mathcal{E}_Y - \mathfrak{M}\mathrm{od})^{w_2(Y)}|_{\mathcal{V}}$$
$$(resp. \,\mathcal{H}(\chi^{-1})^{op} \otimes_{\mathcal{E}_X^{op}} (-) : (\mathcal{E}_X^{op} - \mathfrak{M}\mathrm{od})^{w_2(X)}|_{\mathcal{U}} \xrightarrow{\sim} (\mathcal{E}_Y^{op} - \mathfrak{M}\mathrm{od})^{w_2(Y)}|_{\mathcal{V}}.)$$

Furthermore, the obvious tempered analog of Proposition 7.3 and Corollary 7.5 hold.

Proof. As in Proposition 7.1 and Proposition 7.3, replacing everywhere when appropriate μ hom by t- μ hom.

This proposition gives us (what we will eventually show to be) an alternative construction of Kashiwara's algebroid \mathcal{E}_V : Choose a Darboux cover $\{(\mathcal{U}_\alpha, f_\alpha)\}$ as in Theorem 5.6; this provise gluing data for $\mathbb{P}erv_V$. Apply the construction of (37) to said gluing data to obtain the \mathcal{H} and c of

$$(\{(\mathcal{E}_{\alpha} - \mathfrak{M} \mathrm{od})^{w_2(\alpha)}\}, \{\mathcal{H}_{\beta\alpha}\}, \{c_{\gamma\beta\alpha}\})$$

We have applied a functor to gluing data, hence obtain gluing data for some sheaf of categories which is locally isomorphic to $\mathcal{E}_V - \mathfrak{M}od_{rh}$. We denote the resulting sheaf of categories by \mathcal{C}_V .

Definition 7.12 ([9, Definition 5.2.1]). Let V be a contact manifold.

- (1) An *E*-algebroid \mathcal{A} on V is a \mathbb{C} -algebroid with the property that for any Darboux ball $V \supset \mathcal{U} \subseteq f \to \mathbb{D}^* V$ there is an equivalence $\mathcal{A} \models \cong (f^* \mathcal{E}_{-})^+$
- U → P*X, there is an equivalence A|_U ≃ (f*E_X)⁺.
 (2) A sheaf of C-linear categories C on V is called a *sheaf of twisted E-modules* if for any Darboux ball V ⊃ U → P*X, there is an equivalence C|_U ≃ f*(E_X mod).

The Picard good property of \mathcal{E}_X , recalled in Theorem 7.8, implies that there is a simple description of sheaves of twisted \mathcal{E} -modules.

Theorem 7.13 ([9, Theorem 5.2.3]).

- (1) Any sheaf of twisted \mathcal{E} -modules \mathcal{C} is equivalent to $\mathcal{A} mod$ for some \mathcal{E} -algebra.
- (2) Two *E*-algebroids *A* and *B* are equivalent if and only if the associated sheaves of twisted *E*-modules *A* mod and *B* mod are equivalent.

Proof. This is a direct consequence of Theorem 7.8 and Proposition A.6.

The above Theorem 7.13 implies that C_V is of the form $\mathcal{E}'_V - mod$ for some \mathcal{E} -algebroid \mathcal{E}'_V .

Theorem 7.14. The algebroid \mathcal{E}'_V is equivalent to the canonical algebroid \mathcal{E}_V .

Proof. We will eventually use Polsello's criterion, for which we will need a $*: \mathcal{E}'_V \xrightarrow{\sim} \mathcal{E}'_V^{,op}$. Let us construct it.

First note that appealing to the statement about right modules in Proposition 7.11, the same choice of Darboux charts $\{(\mathcal{U}_{\alpha}, f_{\alpha})\}$, gives us the gluing data

$$\left(\{(a_{\alpha}^{*}\mathcal{E}_{\alpha}^{op}-\mathfrak{Mod})^{w_{2}(\alpha)}\},\{\mathcal{H}_{\alpha\beta}^{op}\},\{c_{\gamma\beta\alpha}\}\right)$$

where $\mathcal{H}_{\alpha\beta}^{op} = \mathcal{H}(f_{\alpha} \circ f_{\beta}^{-1})^{op}$ and, by the proof of Theorem 7.13, it glues to $\mathcal{E}_{V}^{\prime,op} - mod$. Furthermore, the tempered version of Proposition 7.3 and Corollary 7.5 implies that there exists an equivalence

$$*: \mathcal{E}'_V \xrightarrow{\sim} \mathcal{E}'^{,op}_V$$

which can be recovered from its corresponding twisted \mathcal{E} -module equivalence, glued from

$$\Omega_X \otimes_{\mathcal{O}_{\dot{T}^*X}} (-) : \mathcal{E}_X - \mathfrak{M}d^{w_2(X)} \xrightarrow{\sim} a^* \mathcal{E}_X^{op} - \mathfrak{M}d^{w_2(X)}$$

on Darboux charts. Indeed, in order to glue, we have to check that the diagram of Proposition 7.3 is functorial with respect to horizontal compositions; since we are gluing morphisms between (2, 1)-sheaves, we need to check only that, for triple intersections indexed by α , β , and γ , the commuting diagram compose in the way illustrated below:

$$\begin{array}{c} \alpha & \xrightarrow{\sim} & \beta & \beta & \xrightarrow{\sim} & \gamma & \alpha & \xrightarrow{\sim} & \gamma \\ \downarrow & \bigcirc & \downarrow & \bigcirc & \downarrow & \bigcirc & \downarrow & \bigcirc & \downarrow & \uparrow \\ \alpha & \xrightarrow{\sim} & \beta & \beta & \xrightarrow{\sim} & \gamma & \alpha & \xrightarrow{\sim} & \gamma \end{array}$$

But this is Corollary 7.5.

Up until now, our strategy is to define the relevant functors by first defining them as functors between module categories using bimodules, and turn them into algebroid morphisms by appealing to (the proof of) Theorem 7.13. However, as explained in Proposition A.6, on small open sets where the algebroid admits a section so there exists a common generator for the invertible bimodules, the passage from bimodules to ring homomorphisms is given by Lemma A.3. In fact, this process is exactly how the quantized contact transform, as ring isomorphisms, are classically obtained in for example [28, Theorem 11.4.9].

Thus, refining the Darboux cover if needed, we can assume the transition maps comes from genuine ring isomorphisms $\Phi(\chi) : \mathcal{E}_X|_{\mathcal{U}} \xrightarrow{\sim} \mathcal{E}_Y^{op}|_{\mathcal{V}}$. By [21, Theorem 7.2.2], this isomorphism is filtered and compatible with the symbols, i.e., there is a commuting diagram



Now (i) is automatic since the filtration is defined locally and is preserved by $\Phi(\chi)$.

For (ii), the above commuting diagram implies that the identification $\sigma_X : Gr(\mathcal{E}_X) \xrightarrow{\sim} \mathcal{S}_{\mathcal{O}_X}(\Theta_X)$ can be glued to a global identification $\sigma : Gr(\mathcal{E}'_V) \simeq \left(\bigoplus_{m \in \mathbb{Z}} \mathcal{L}^m\right)^+$. In fact, the object $Gr(\mathcal{E}'_V)$ is well-defined as an algebra (not merely as an algebroid) and σ comes from an algebra isomorphism $\sigma : Gr(\mathcal{E}'_V) \simeq \bigoplus_{m \in \mathbb{Z}} \mathcal{L}^m$. This is because, on any triple overlap indexed by α, β, γ , the composition $\Phi_{\alpha\gamma} \Phi_{\gamma\beta} \Phi_{\beta\alpha}$ is a filtered and symbol-preserving ring isomorphism on \mathcal{E}_{α} and so it becomes the identity when taking Gr. (In particular, the ambiguity P_{ijk} in [24, (0.1)], which leads to the necessity of working with algebroids, disappear when passing to graded algebras.)

Passing now to (iii): on local charts, we have natural transformation of bimodules

$$\Omega_X^{-1} \otimes_{\mathcal{O}} \Omega_X \to \mathcal{O}_X$$

which clearly glue to a natural transformation of bimodules on V. This in turn induces the natural transformation $*^2 \rightarrow id$ by Proposition A.6. The fact that this natural transformation is a natural isomorphism can be checked on Darboux charts, where it is obvious.

Finally, we exhibit the natural transformation δ_0 : in a Darboux chart $V \supset \mathcal{U} \hookrightarrow P^*X$, δ_0 is just given by the diagram

That this diagram commutes is a restatement of Lemma 6.1. It follows from (ii) that these locally defined natural transformations induce a globally defined natural transformation. (We remark that a more intrinsic way to see this is to use Lemma 6.11 and trade $(\mathcal{E}_X - mod)^{w_2(X)}$ for $\mathcal{E}_X^{\sqrt{\Omega_X}} - mod$; under this identification, $Gr_0(*)$ is exactly D.) This completes the proof.

Theorem 7.15 (Global microlocal Riemann–Hilbert correspondence). Let V be a complex manifold. There is an equivalence

$$\mu \operatorname{RH}_V : \mathbb{P}erv_V \xrightarrow{\sim} \mathcal{E}_V - \mathfrak{M}od_{rh}$$

between perverse microsheaves on V and regular holonomic \mathcal{E}_V -modules.

Proof. The sheaves $\operatorname{Perv}_V \subseteq \mu \operatorname{sh}_V$ and $\mathcal{E}_V - \mathfrak{M} \operatorname{od}_{rh} \subseteq \mathcal{E}_V - \mathfrak{M} \operatorname{od}$ can be reconstructed by the gluing data

 $(\{\mathbb{P}erv_{\alpha}^{w_2}\}, \{\mathcal{K}_{\beta\alpha}\}, \{c_{\gamma\beta\alpha}\}) \text{ and } (\{(\mathcal{E}_{\alpha} - \mathfrak{M}od_{rh})^{w_2(\alpha)}\}, \{\mathcal{H}_{\beta\alpha}\}, \{c_{\gamma\beta\alpha}\})$

where we use Theorem 7.14 to conclude for the latter. But the compatibility statement in Theorem 7.7 implies that the local equivalence μRH_{α} glues to an equivalence on V.

Remark 7.16. To get a feel for Polesello's criterion Theorem 7.9, it is instructive to consider the example of a complex manifold X. Let $L = \Omega_X$ be the canonical bundle. Then Kashiwara's stack is $(\mathcal{D}_X^{L/2})^+$ which is generally different from \mathcal{D}_X^+ . So let us understand where Polesello's criterion fails for \mathcal{D}_X^+ .

Although $\mathcal{D}_X \neq \mathcal{D}_X^L$ in general, combining Lemma 6.8 and Morita theory in the sheaf setting, e.g. [9, Corollary 3.3.8], we see that the *algebroids* \mathcal{D}_X^+ and $\mathcal{D}_X^{L,+}$ are equivalent. (The bimodule is $L \otimes_{\mathcal{O}} D_X$). As explained in Section 6.5, * gives an algebroid isomorphism $D_X^+ - \mathfrak{M}od \rightarrow D_X^{L,op} - \mathfrak{M}od$. And by Morita theory (applied to the sheaf of categories $D_X^+ - \mathfrak{m}od$), we know that $D_X^+ - \mathfrak{m}od$ and $(D_X^L)^+ - \mathfrak{m}od$ are isomorphic. This allows us to construct an anti-involution on $D_X^+ - \mathfrak{M}od$, which verifies (iii) in Theorem 7.9.

Consider further the fact that the associated graded $Gr(\mathcal{D}_X) = S_{\mathcal{O}}(\Theta)$ is given by symmetric products of holomorphic vector fields, i.e., functions of the cotangent bundle which are polynomials in the fiber direction. In particular, Gr_0 is simply \mathcal{O}_X . Because $S_{\mathcal{O}}(\Theta)$ is commutative, $Gr(\mathcal{D}_X) = Gr(\mathcal{D}_X^L)$ as algebras. This verifies (i) and (ii) in Theorem 7.9.

What breaks is the compatibility condition (35). Namely, the equivalence $\mathcal{D}_X^+ = \mathcal{D}_X^{L,+}$ after applying Gr_0 is not the identity, since on the level of modules, it is given by

$$\mathcal{O}_X - mod \xrightarrow{\sim} \mathcal{O}_X - mod$$

 $\mathcal{M} \mapsto L \otimes_{\mathcal{O}} \mathcal{M}.$

APPENDIX A. RECOLLECTIONS FROM MORITA THEORY

To study the untwisting procedure in Lemma 6.8 and Proposition 6.9, we consider the following situation: Let R and S be rings and $f : R \to S$ be a ring homomorphism. Viewing S as a right R-module and a left S-module, we obtain a tensor-forgetful adjunction

$$S \otimes_R (-) : R - mod \rightleftharpoons S - mod : {}_f(-)$$

where for a S-module N, we use ${}_{f}N$ to denote the R-module whose structure is given by $r \cdot {}_{f}n := f(r)n$ for $r \in R$ and $n \in N$. We note that, if $\alpha : N_1 \to N_2$ is a S-module homomorphism, the functions ${}_{f}\alpha$ and ${}_{a}\alpha$ are both set-theoretically the same as α .

Lemma A.1. Let $f, g : R \to S$ be two ring homomorphisms. Any natural transformation $T : {}_{f}(-) \to {}_{g}(-)$ has the form L_{s} , for $s \in S$ such that sf(r) = g(r)s for all $r \in R$, where L_{s} is the family of function which is defined by left multiplication

$$L_s(N): N \to N$$
$$n \mapsto sn.$$

In particular, the functors f(-) and g(-) are equivalent if and if only there is $s \in S^{\times}$ such that $g = \operatorname{Ad}(s) \circ f$.

Proof. The natural transformation $T : {}_{f}(-) \to {}_{g}(-)$ is determined by T(N)(n) for any $N \in S - mod$ and $n \in N$. Fix such an $n \in N$, the function

$$\phi_n: S \to N$$
$$s \mapsto sn$$

is an S-module homomorphism. Thus, we have he equality $T(N) \circ_f \phi_n = {}_g \phi_n \circ T(S)$. In other words, for any $s \in S$, we have T(N)(sn) = (T(S)(s)) n. Taking s = 1, we see that T(N)(n) = (T(S)(1)) n is determined by the element $s_T \coloneqq T(S)(1) \in S$, or $T = L_{s_T}$. To see that s_T satisfies the desired property, we take N = S and n = f(r) for $r \in R$. Then we see that

$$s_T f(r) = T(S)(f(r)) = T(S)(r \cdot_f 1) = r \cdot_g T(S)(1) = g(r)s_T.$$

One can similarly show that T is a natural equivalence if $s_T \in S^{\times}$ and in this case $g(r) = s_T^{-1} f(r) s_T$, or $g = \operatorname{Ad}(s_T) \circ f$.

Corollary A.2. For a ring R, a ring automorphism $f : R \to R$ induces a functor $_f(-)$ equivalent to the identity id_{R-mod} if and only if f = Ad(r) for some $r \in R^{\times}$. In this case, we have

$$L_r: \operatorname{id}_{R-mod} \xrightarrow{\sim} \operatorname{Ad}(r)(-).$$

Another fact which we will use is that a (R, S)-bimodule with a common left R-module and right S-module generator t gives rise to a anti-ring homomorphism. The fact is well-known but we spell out the details since we will need it in Section 6.5. We also remark that this is the same procedure in [28, Theorem 11.4.9] to obtain the quantized contact transform.

Lemma A.3. Let M be a (R, S)-bimodule with a common generator t. That is the left R-module morphism $R \to M, r \mapsto r \cdot t$ and the right S-module homomorphism $S \to M, s \mapsto t \cdot s$ are both an isomorphism. Then, M defines an ring homomorphism $f_M : R \to S$.

Proof. For any $r \in R$, by the assumption on t, there exists a unique $f_M(r) \in S$ such that $r \cdot t = t \cdot f_M(r)$, which one can check satisfies $f_M(1_R) = 1_S$ and $f_M(r_1r_2) = f_M(r_1) \cdot f_M(r_2)$.

We also need to consider passing to 'op' modules. That is, a (R, S)-bimodule M can be viewed tautologically as an (S^{op}, R^{op}) -bimodule, which we denote it by M^{op} . For the sheaf-theoretical situation, since convolution has a prefer direction, the coordinates have to be swapped. That is, we let $f: X \to Y$ be a homeomorphism between topological spaces and we denote by $v: X \times Y \xrightarrow{\sim} Y \times X$ the coordinate swapping map v(x, y) = (y, x). Let $S \in sh(X), R \in sh(Y)$ be ring-valued sheaves and $M \in sh(X \times Y)$ be a (p_2^*R, p_1^*S) -bimodule supported on the graph Γ_f so v^*M is supported in $\Gamma_{f^{-1}} \subset Y \times X$.

Lemma A.4. Convolution defines functors

$$M \circ_S (-) : S - mod \to R - mod$$

and

$$v^*M^{op} \circ_{R^{op}} (-) : R^{op} - mod \to S^{op} - mod.$$

Finally, we mention that Morita theory generalizes mostly naturally in the algebroid set up. Our main reference for this discussion will be [9]. For an algebroid \mathcal{A} on X (over \mathbb{C}), the category of modules is defined by

(39)
$$\mathcal{A} - mod \coloneqq \operatorname{Fun}_{\mathbb{C}}(\mathcal{A}, \mathbb{C}_X - mod).$$

Note that such an assignment naturally organizes to a sheaf $\mathcal{A} - \mathfrak{M}$ od. Similarly to the case of algebras, for two algebroids \mathcal{A} , \mathcal{B} , one can form their tensor algebroid $\mathcal{A} \otimes_{\mathbb{C}} \mathcal{B}$ and a $(\mathcal{A}, \mathcal{B})$ -bimodule is an object $\mathcal{P} \in \mathcal{A} \otimes_{\mathbb{C}} \mathcal{B}^{op}$. A $(\mathcal{A}, \mathcal{B})$ -bimodule is said to be invertible if there exists a $(\mathcal{B}, \mathcal{A})$ -bimodule \mathcal{Q} such that

$$\mathcal{P} \otimes_{\mathcal{B}} \mathcal{Q} \cong \Delta_{\mathcal{A}} \text{ and } \mathcal{Q} \otimes_{\mathcal{A}} \mathcal{P} \cong \Delta_{\mathcal{B}}$$

where $\Delta_{\mathcal{A}}$ is the diagonal bimodule which is given by $((x, y) \mapsto \mathcal{A}(x, y))$. An equivalence of bimodules $f : \mathcal{A} \xrightarrow{\sim} \mathcal{B}$ naturally induces an equivalence $f \circ (-) : \mathcal{B} - mod \xrightarrow{\sim} \mathcal{A} - mod$. In fact, this assignment organizes to a 2-functor

(40)
$$\operatorname{Equiv}_{\mathbb{C}}(\mathcal{A}, \mathcal{B}) \to \operatorname{Equiv}_{\mathbb{C}}(\mathcal{B} - mod, \mathcal{A} - mod).$$

Definition A.5 ([9, Definition 3.4.1]). An algebroid \mathcal{A} is said to be Picard good if all invertible $\mathcal{A} \otimes_{\mathbb{C}} \mathcal{A}^{op}$ bimodules are locally free of rank one over \mathcal{A} (or, equivalently, over \mathcal{A}^{op}).

Proposition A.6 ([9, Proposition 3.4.3]). When A is Picard good and B is locally equivalent to A, the (40) is an equivalence.

Proof. D'Agnolo and Polesello prove the more general statement, in [9, Proposition 3.3.7], that for linear stacks the image of (40) consists of bimodules locally free of rank one over A. In our case, the classification of the image is implied by the discussion at the beginning of this Appendix section, and Lemma A.3 gives the explicit way to reconstruct the algebroid map from the bimodule on small open set when there is a section.

APPENDIX B. RECOLLECTIONS FROM ALGEBRAIC TOPOLOGY

Here we trace through some relationships classifying certain null-homotopies and homotopies of null-homotopies etc. which we have needed in the text. Let us first recall the basic result on (de-)looping and commutativity. It is due to Boardman-Vogt and May; we give references to the treatment in Lurie's books.

Theorem B.1 ([30, Theorem 5.2.6.10]). For $\infty > k \ge 0$, the functor of taking k-fold loop space

$$\operatorname{Spc}_{*}^{k \geq} \xrightarrow{\sim} Mon_{\mathbb{E}_{k}}^{gp}(\operatorname{Spc})$$

 $(X, *) \mapsto \Omega_{*}^{k}(X)$

is an equivalence between k-connective space and group-like \mathbb{E}_k -monoid in spaces.

As a corollary, one obtain a similar statement for group-like \mathbb{E}_{∞} -monoid in spaces; they are the abelian group objects in this setting:

Corollary B.2 ([30, Corollary 5.2.6.27]). There exists an equivalence

$$\Omega^{\infty}: \operatorname{Sp}^{cn} \xrightarrow{\sim} Mon^{gp}_{\mathbb{E}_{\infty}}(\operatorname{Spc})$$

between connective spectra, spectra with no non-trivial negative homotopy group, and group-like \mathbb{E}_{∞} -monoid in spaces.

Example B.3. Any abelian group A can be seen as a discrete topological group and hence an abelian group object in spaces. The theorem above thus implies that there exists a k-connective space $B^k(A)$ such that $\Omega^l_*B^k(A) = B^{k-l}(A)$, for $l \leq k$ where we use the notation $B^0A = A$. In particular, $\Omega_*B(A) = A$ as a group in spaces.

Example B.4. By Bott periodicity, O, U, and U/O can be regarded as an object in $Mon_{\mathbb{E}_{\infty}}^{gp}(\text{Spc})$ and thus in Sp^{cn} . However, taking $\text{Sp}^{cn} \hookrightarrow \text{Sp}$ is invariant undertaking cofibers and we thus have fiber sequences $U \to U/O \to BO, U/O \to BO \to B^2U$, etc..

Example B.5. Let $G \in \mathbb{E}_1^{gp}(\operatorname{Spc})$ and $A \in \mathbb{E}_2^{gp}(\operatorname{Spc})$. We mention in the last example that $Mon_{\mathbb{E}_{\infty}}^{gp}(\operatorname{Spc})$ in general is only closed under taking cofibers. However, in case when there is a group homomorphism $\alpha : G \to BA$, its fiber $K := \operatorname{fib}(\alpha)$ in fact admits a group structure. Indeed, passing to connective spaces by Theorem B.1, we need to argue that the fiber $F := \operatorname{fib}(BG \to B^2A)$ has a vanishing π_0 . But this follows from the homotopy long exact sequence

$$\cdots \to \pi_1(B^2A) \to \pi_0(F) \to \pi_0(BG) \to \cdots$$

and the fact that $\pi_1(B^2A) = \pi_0(BA) = \{*\}.$

Remark B.6. Let $G \in Mon_{\mathbb{E}_1}^{gp}(\operatorname{Spc})$ be a group in spaces. Consider a nice topological space X, e.g., a CW complex, and a classifying map $\eta : X \to BG$, classifying the principal G-bundle \mathcal{P}_{η} . We know that a null-homotopy τ of $\eta : X \to BG$, i.e., an equivalence $\tau : \eta \xrightarrow{\sim} *$ to the constant map, corresponds to a section, which we will abuse the notation and denote it by $\tau : X \to \mathcal{P}_{\tau}$. For two of such sections τ_1 and τ_2 , if we denote by $\tau_1^{\text{rev}} : * \xrightarrow{\sim} \eta$ the reverse equivalence, then the concatenation $\tau_2 \# \tau_1^{\text{rev}}$ will be a loop at the constant map *. Thus,

$$\tau_2 \# \tau_1^{\text{rev}} \in \Omega_* Map(X, BG) = Map(X, \Omega_* BG) = Map(X, G)$$

is given by a "G-valued function" $g: X \to G$. Now, the k = 1 case of Theorem B.1 implies that $\Omega_*BG = G$ as groups, i.e., concatenation of loops in BG is given the group multiplication of G. This implies that, as sections, $\tau_2 = g \cdot \tau_1$ where \cdot comes from the G-action on \mathcal{P}_{τ} .

Now let (N, ξ) be a manifold N with a symplectic/complex vector bundle ξ , or equivalently, a map $\xi : M \to BU(n)$. Abuse the notation and denote again by ξ the composition

$$M \xrightarrow{\xi} BU(n) \to BU \to BU/O \to BO^2 \xrightarrow{Bw_2} B^3(\mathbb{Z}/2).$$

Here, we note that the universal second Stiefel-Whitney, $w_2 : BO \to B^2 \mathbb{Z}/2$, is induced by the exact sequence $1 \to \mathbb{Z}/2 \to \text{Pin} \to O \to 1$. We note that since $BU \to BU/O \to BO^2$ is a fiber sequence, it is canonically null-homotopic, and we will use the notation γ_{can} to denote the induced null-homotopy for ξ .

Assume now that there exists an open cover $\{U_{\alpha}\}$ with an open embedding $f_{\alpha}: U_{\alpha} \hookrightarrow T^*M_{\alpha}$, for some manifold M_{α} of dimension n, so that the vector bundle $\phi_{\alpha}: T^*M_{\alpha} \to BO(n)$ given by the fiber $\phi_{\alpha}(x_{\alpha}, \xi_{\alpha}) = T^*_{x_{\alpha}}M_{\alpha}$ is a polarization of ξ on U_{α} , i.e., the following diagram commutes:



Passing to stabilization, this fiber polarization provides another null-homotopy for $\xi|_{U_{\alpha}}$ since $BO \to BU \to BU/O$ is a fiber sequence and we abuse the notation and still use ϕ_{α} to denote it. Denote by τ_{can}^{rev} the reverse homotopy from the constant map to $\xi_{U_{\alpha}}$.

Proposition B.7. The concatenation $\phi_{\alpha} \# \tau_{can}^{rev} \in \Omega_* \operatorname{Map} (U_{\alpha}, B^3(\mathbb{Z}/2)) = \operatorname{Map} (U_{\alpha}, B^2(\mathbb{Z}/2))$ is given by $\phi_{\alpha} \# \tau_{can}^{rev} = f_{\alpha}^* w_2(M_{\alpha})$.

Proof. We note that both null-homotopies are from composing with null-homotopies which happens in the sequence

$$BO \to BU \to BU/O \to B^2O$$

and hence the following lemma implies that the corresponding point in Map $(U_{\alpha}, B^2(\mathbb{Z}/2))$ is given by the composition

$$U_{\alpha} \xrightarrow{f_{\alpha}} T^* M_{\alpha} \xrightarrow{\phi_{\alpha}} BO \xrightarrow{w_2} B^2(\mathbb{Z}/2),$$

i.e., $w_2(M_{\alpha})$.

Lemma B.8. Let *C* be a stable category, $A \xrightarrow{\alpha} B \xrightarrow{\beta} C$ be a fiber sequence and choose, up to a contractible ambiguity, a homotopy $h : \beta \circ \alpha = 0$. Extend the sequence and get $B \xrightarrow{\beta} C \xrightarrow{\gamma} A[1]$ with a similar choice of homotopy g. Then under the identification $\Omega_0 \operatorname{Map}(A, A[1]) = \operatorname{Map}(A, A)$, the point $(g^{\operatorname{rev}} \bigcirc \alpha) \# (\gamma \bigcirc h)$ corresponds to id_A . Here ' \bigcirc ' denotes the horizontal composition between a 1-morphism and a 2-morphism.

Proof. The follows from tracing through universal properties and the fact that composition of pullbacks is a pullback, i.e., $\Omega_0 A[1] = A$ is given by both of the following diagrams:



Denote by $B^2(\mathbb{Z}/2)(\xi)$ the (∞ -)principal $B^2(\mathbb{Z}/2)$ -bundle over M classify by $\xi : M \to B^3(\mathbb{Z}/2)$. Recall that null-homotopies of $\xi : M \to B^3(\mathbb{Z}/2)$ corresponds to sections of $B^2(\mathbb{Z}/2)(\xi) \to M$. More precisely, since we have the pullback diagram



where we use $\Omega_*B^3(\mathbb{Z}/2) = B^2(\mathbb{Z}/2)$ for the right square, a null-homotopy identifies $B^2(\mathbb{Z}/2)(\xi)$ with $M \times B^2(\mathbb{Z}/2)$, which admits an obvious section given by $M = M \times \{*\} \hookrightarrow M \times B^2(\mathbb{Z}/2)$. In our case, denote by τ_{can} and ϕ_{α} , the sections which corresponds to the homotopies h_{can} and h_{α} . Since $\Omega_*B^3(\mathbb{Z}/2) = B^2(\mathbb{Z}/2)$ as groups by Theorem B.1, the previous Proposition B.7 implies that

(41)
$$\phi_{\alpha} = (f_{\alpha}^* w_2(M_{\alpha})) \cdot \tau_{can}$$

where we view $f_{\alpha}^* w_2(M_{\alpha}) \in \text{Map}(U_{\alpha}, B^2(\mathbb{Z}/2))$ and the multiplication \cdot is induced from the principal bundle structure $a: B^2(\mathbb{Z}/2) \times B^2(\mathbb{Z}/2)(\xi) \to B^2(\mathbb{Z}/2)(\xi)$.

APPENDIX C. RECOLLECTIONS FROM SYMPLECTIC AND CONTACT GEOMETRY

Definition C.1. A contact manifold $V = (V, \xi)$ is the data of a manifold V of dimension 2n + 1 along with a codimension-1 sub-bundle $\xi \subset TV$ which is maximally non-integrable. Concretely, this means that we can locally write $\xi = \ker \alpha$, for some 1-form α having the property that $\alpha \land (d\alpha)^n \neq 0$.

If (V,ξ) is a contact manifold, then $\xi \to V$ has the structure of a conformally symplectic vector bundle. (Indeed, if $f\alpha = \alpha'$, then $d\alpha' = df \wedge \alpha + fd\alpha$, so $d\alpha'|_{\xi} = fd\alpha|_{\xi}$.)

Definition C.2. Let (V, ξ) be a contact manifold. A submanifold $W \subset V$ is said to be:

- a *contact submanifold* if $\xi \cap TW \subset TV$ is a contact structure on W
- *isotropic* if $TW \subset \xi \subset TV$
- coisotropic if (TW ∩ ξ) is an coisotropic subspace of (ξ, dα) at all points in W, for some (equivalently any) locally defined 1-form ξ = ker α.

There are obvious notions of contact/isotropic/coisotropic embeddings.

The above definitions are equally valid in the category of real manifold with smooth maps and the category of complex manifolds with holomorphic maps. *However, for the remainder of this appendix, we will exclusively work in the category of real contact manifolds*. We refer to [8, Section 2] for background on the complex case.

Given a contact manifold (Y,ξ) , its symplectization is the manifold $SY := \{\alpha \in T^*Y \mid \ker \alpha = \xi\}$ which is symplectic with respect to the restriction of the canonical 1-form on T^*Y . We say that Y is co-orientable iff SY is disconnected; a co-orientation amounts to labeling one of the two components as positive.

An exact symplectic manifold (M, λ) is said to be *Liouville* if it admits a proper embedding $S^+Y \hookrightarrow (M, \lambda)$ of the positive symplectization of a closed contact manifold. If (M, λ) is Liouville, then $[0, 1] \times M, dt + \lambda$ is a contact manifold which is often called the *contactization*.

Lemma C.3. Let $\iota : N \hookrightarrow M$ be an embedding and consider the exact sequence of bundle maps $0 \to \ker \iota^* \to T^*M|_N \xrightarrow{\iota^*} T^*N \to 0$. The space of sections of ι^* is contractible. Any section σ induces an inclusion of Liouville manifolds $T^*N \hookrightarrow T^*M$.

Definition C.4. Let (W, ξ) be a co-orientable contact manifold. A (possibly time-dependent) vector field Z is called a *contact vector field* it its flow preserves the contact structure.

Lemma C.5. [15, Thm. 2.3.1] Let (W, ξ) be a co-orientable contact manifold. A choice of contact form $\xi = \ker \alpha$ induces a bijection of sets

(42) $\{ contact vector fields on W \} \leftrightarrow C^{\infty}(W)$

This bijection takes $Z \mapsto \alpha(Z)$. The inverse sends a Hamiltonian H to a contact vector field Z uniquely defined by the equations $\alpha(Z_H) = H$ and $i_{Z_H} d\alpha = dH(R_\alpha)\alpha - dH$, where R_α the Reeb vector field.

Corollary C.6. Let (W, ξ) be a co-orientable contact manifold and let $\phi_t : (V, \zeta) \hookrightarrow (W, \xi), t \in [0, 1]$ be a 1-parameter family of codimension zero contact embeddings. Then, after possibly shrinking W, ϕ_t extends to a family of contactomorphisms of (V, ξ) . More precisely, there exists a family of contactomorphisms $\tilde{\phi}_t : (W, \xi) \to (W, \xi), t \in [0, 1]$ such that $\tilde{\phi}_t \circ \phi_0 = \phi_t$ holds on any compact subset of W.

Proof. Fix a contact form $\xi = \ker \alpha$ and consider the contact vector field $Z_t := \frac{d}{dt}\phi_t$ defined on $\phi_t(W)$. Let $H_t := \alpha(Z_t)$ be the corresponding family of Hamiltonians and let \tilde{H}_t be an extension of H_t to all of W. Now define $\tilde{\phi}_t$ to be the flow of \tilde{H}_t .

Proposition C.7. Let (V,ξ) be a co-orientable contact manifold and let $(E,\omega) \to V$ be a symplectic vector bundle.

- (i) a neighborhood of the zero section admits a contact structure $\hat{\xi}$ whose restriction to the tangent space of the zero section agrees with ξ , and such that the conformal symplectic normal bundle of the zero section is precisely E.
- (ii) the space of (germs of) such structures is weakly contractible.

Proof. Fix a contact form $\xi = \ker \alpha$. It is explained by Avdek [4, Thm. 5.4] how to construct a contact form $\hat{\alpha}$ on $Op(0_E) \subset E$ such that (a) the restriction of $\hat{\alpha}$ to the tangent space of the zero section agrees with α and (b) $d\alpha = \omega$ on $\xi|_{0_V} \subset TE|_{0_V}$. For the sake of checking Lemma C.9, here is a summary of the construction: Fix a complex structure on E compatible with ω , thus making E into a unitary vector bundle. Fix unitary bundle trivializations $\pi^{-1}(U_i) \simeq U_i \times \mathbb{R}^{2n}$ corresponding to a covering $\{U_i\}$ of V, and where we denote the fiber coordinates by (q, p). Let $\{h_i\}$ be a partition of unity subordinate to the covering $\{U_i\}$ and define $\lambda_i = \frac{h_i}{2} (\sum_j p^j dq_j - q^j dp_j)$. Finally, set $\lambda = \sum_i \lambda_i$ and set $\hat{\alpha} = \pi^* \alpha + \lambda$.

To prove (ii), suppose that $\hat{\xi}_0$ and $\hat{\xi}_1$ both satisfy (i). Choose corresponding contact forms $\hat{\alpha}_0$ and $\hat{\alpha}_1$ and observe that the linear interpolation ker $((1 - t)\alpha_0 + t\alpha_1)$ is contact in some possibly smaller neighborhood of 0_V . A similar argument works in higher dimensions.

Lemma C.8. Let $(V, \xi = TV \cap \eta) \hookrightarrow (W, \eta)$ be a contact embedding. Let $(O_P(0_V), \hat{\xi}) \to V$ be a neighborhood of the zero section in the symplectic normal bundle with the contact structure

furnished by Proposition C.7. The space of extensions to a codimension zero contact embedding



is (weakly-)contractible.

Proof. Follows from the parametric relative neighborhood theorem in [37, Thm. 2.10]. \Box

Lemma C.9. Fix a polarization $\tau : V \to LGr(E)$ and let $L_{\tau} \subset E$ be the associated bundle of Lagrangian subspaces. Then $L_{\tau} \subset (E, \hat{\alpha})$ is coisotropic; its characteristic foliation is the natural foliation by the fibers of $L_{\tau} \to V$.

Proof. It is enough to check, for arbitrary $x \in V$, that any element in the tangent space of $(L_{\tau})_x \subset E_x$ pairs to zero with any vector in $TL_{\tau} \cap \xi$ with respect to $d\hat{\alpha}$ (indeed, for dimension reasons, the tangent space of the fibers $(L_{\tau})_x$ must then be the whole orthogonal complement of $(TL_{\tau} \cap \xi, d\hat{\alpha})$).

We follow the notation of Proposition C.7. Suppose v is tangent to $(L_{\tau})_x$; then $d\hat{\alpha}(v, -) = d\pi^* \alpha(v, -) + d\lambda(v, -) = d\lambda(v, -)$. Now $d\lambda = \sum d(h_i\lambda_i) = \sum dh_i \wedge \lambda_i + \sum h_i d\lambda_i$. To verify that $d\lambda(v, -) = 0$ on L_{τ} , it is enough to check for each i that $dh_i \wedge \lambda_i(v, -) = 0$ and $h_i d\lambda_i(v, -) = 0$. We may check this in the local chart $U_i \times \mathbb{R}^{2n} \simeq \pi^{-1}(U_i)$.

We first argue that $\lambda_i(v) = 0$. Indeed, $(L_{\tau})_x \subset E_x$ is a linear Lagrangian subspace. $(L_{\tau})_x$ is therefore preserved by the radial Liouville vector field $\sum_j (q_j \partial_{q_j} + p_j \partial_{p_j})$; it follows that $\sum_j p^j dq_j - q^j dp_j$ vanishes identically on $(L_{\tau})_x$. (Note that this assertion holds at all points in the vector space $(L_{\tau})_x -$ not merely at the origin.)

We also have $\sum h_i d\lambda_i(v, -) = 0$: indeed, for any vector w tangent to $(L_\tau)_x$, $d\lambda_i(v, w) = 0$ because $(L_\tau)_x$ is Lagrangian, while for any w tangent to the base, $d\lambda_i(v, w) = 0$ trivially (because λ_i only involves fiber coordinates).

Lemma C.10. Let (E,ξ) be a contact manifold and let $(V, TV \cap \xi) \subset E$ be a contact submanifold. Suppose that V is contained in some coisotropic submanifold $C \subset E$.

Let $(\phi_t)_{t \in [0,1]}$ be a diffeotopy enjoying the following properties:

- ϕ_t fixes C pointwise
- $C_t = \phi_t(C) \subset E$ is coisotropic for all t (where $C := C_0$)
- $\phi_t : C \to C_t$ carries the characteristic foliation of C to that of C_t

Then there is a contact diffeotopy $\Phi_t : Op(V) \to Op(V)$ such that (after possibly replacing C with $C \cap Op(V)$), we have $\Phi_t(C) = \psi_t(C)$.

Proof. Fix any contact form α on E. As explained in [37, Sec. 2.4], the contact structure in a neighborhood of a coisotropic is entirely determined by the restriction of the contact form to the leaf space of the characteristic foliation. By assumption, ϕ_t preserves the characteristic foliation, and the leaf space is canonically identified with $(V, TV \cap \xi)$. But since ϕ_t preserves V pointwise, it also preserves the contact form $\alpha|_V$.

Definition C.11. Let $(V^{2n+1}, \xi \cap TV) \hookrightarrow (W^{2N+1}, \xi)$ be a contact embedding. A *thickening* is a (germ of a) co-isotropic submanifold of dimension $(\dim W - \dim V)/2 + \dim V$ which contains V and whose characteristic foliation is non-singular and transverse to V.

Lemma C.10 and Corollary C.6 together imply

Corollary C.12. Any isotopy of thickenings is induced by a global contact isotopy.

Given a contact embedding $\iota : (V^{2n+1}, \xi \cap TV) \hookrightarrow (W^{2N+1}, \xi)$ and a normal polarization $\tau \in LGr(NV)$, its image under the (contractible choice of) extension furnished by Lemma C.8 is a thickening. In other words, there is a natural map from the space of pairs (ι, τ) to the space of pairs (ι, C) where C is a thickening. Since all choices involved in the construction of this map are contractible:

Corollary C.13. *This map is a homotopy equivalence.*

REFERENCES

- [1] Emmanuel Andronikof. A microlocal version of the Riemann-Hilbert correspondence. *Topological methods in nonlinear analysis*, 4(2):417–425, 1994. 2, 4, 34
- [2] Emmanuel Andronikof. Microlocalisation tempérée. Number 57. Société mathématique de France, 1994. 34, 35
- [3] Emmanuel Andronikof. Microlocalization of perverse sheaves. *Journal of Mathematical Sciences*, 82(6):3754–3758, 1996. 3
- [4] Russell Avdek. Liouville hypersurfaces and connect sum cobordisms. *Journal of Symplectic Geometry*, 19(4):865–957, 2021. 44
- [5] Jan-Erik Björk. Analytic D-modules and applications, volume 247. Springer Science & Business Media, 2013.
 27, 28, 29, 33, 34, 35
- [6] Tom Braden, Anthony Licata, Nicholas Proudfoot, and Ben Webster. Quantizations of conical symplectic resolutions II: category O and symplectic duality. arXiv preprint arXiv:1407.0964, 2014. 3
- [7] Tom Braden, Nicholas Proudfoot, and Ben Webster. Quantizations of conical symplectic resolutions I: local and global structure. *arXiv preprint arXiv:1208.3863*, 2012. 3
- [8] Laurent Côte, Christopher Kuo, David Nadler, and Vivek Shende. Perverse microsheaves. *arXiv Preprint*, arXiv:2209.12998:1–25, 2022. 2, 3, 4, 12, 13, 20, 23, 24, 35, 43
- [9] Andrea D'Agnolo and Pietro Polesello. Morita classes of microdifferential algebroids. *Publications of the Research Institute for Mathematical Sciences*, 51(2):373–416, 2015. 29, 35, 36, 38, 40
- [10] Andrea D'Agnolo and Pierre Schapira. Quantization of complex Lagrangian submanifolds. Advances in Mathematics, 213(1):358–379, 2007. 3
- [11] Dennis Gaitsgory and Nick Rozenblyum. A study in derived algebraic geometry: Volume I: correspondences and duality, volume 221. American Mathematical Society, 2019. 6, 8
- [12] Sheel Ganatra, John Pardon, and Vivek Shende. Covariantly functorial wrapped Floer theory on Liouville sectors. *Publications mathématiques de l'IHÉS*, 131(1):73–200, 2020. 3
- [13] Sheel Ganatra, John Pardon, and Vivek Shende. Microlocal Morse theory of wrapped Fukaya categories. *Annals of Mathematics*, 199(3):943–1042, 2024. 3
- [14] Sheel Ganatra, John Pardon, and Vivek Shende. Sectorial descent for wrapped Fukaya categories. *Journal of the American Mathematical Society*, 37(2):499–635, 2024. 3
- [15] Hansjörg Geiges. An introduction to contact topology, volume 109. Cambridge University Press, 2008. 44
- [16] Stephane Guillermou. dg-methods for microlocalization. *Publications of the Research Institute for Mathematical Sciences*, 47(1):99–140, 2011. 34
- [17] Stéphane Guillermou. Quantization of conic Lagrangian submanifolds of cotangent bundles. *arXiv preprint arXiv:1212.5818*, 2012. 3
- [18] Stéphane Guillermou, Masaki Kashiwara, and Pierre Schapira. Sheaf quantization of Hamiltonian isotopies and applications to nondisplaceability problems. *Duke Mathematical Journal*, 161(2):201–245, 2012. 15, 16, 17
- [19] Masaki Kashiwara. On the maximally overdetermined system of linear differential equations, i. *Publications of the Research Institute for Mathematical Sciences*, 10(2):563–579, 1975. 2
- [20] Masaki Kashiwara. The Riemann-Hilbert problem for holonomic systems. *Publications of the Research Institute for Mathematical Sciences*, 20(2):319–365, 1984. 2
- [21] Masaki Kashiwara. Introduction to Microlocal Analysis: Lectures Given at the University of Berne in June 1984, Under the Sponsorship of the International Mathematical Union. L'Enseignement Mathématique, Université de Genève, 1986. 37
- [22] Masaki Kashiwara. *Representation theory and D-modules on flag varieties*. Kyoto University. Research Institute for Mathematical Sciences [RIMS], 1988. 4, 6

- [23] Masaki Kashiwara. Algebraic study of systems of partial differential equations, volume 63. Société mathématique de France, 1995. 2
- [24] Masaki Kashiwara. Quantization of contact manifolds. Publications of the Research Institute for Mathematical Sciences, 32(1):1–7, 1996. 2, 3, 4, 35, 37
- [25] Masaki Kashiwara. *D-modules and microlocal calculus*, volume 217. American Mathematical Soc., 2003. 2, 28, 32, 34, 35
- [26] Masaki Kashiwara and Raphaël Rouquier. Microlocalization of rational Cherednik algebras. Duke Mathematical Journal, 144(3):525–573, 2008. 3
- [27] Masaki Kashiwara and Pierre Schapira. *Microlocal study of sheaves*, volume 469. Société mathématique de France, 1985. 31
- [28] Masaki Kashiwara and Pierre Schapira. *Sheaves on manifolds*, volume 292 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1990. 2, 4, 9, 10, 11, 12, 14, 22, 24, 30, 31, 32, 33, 37, 39
- [29] Masaki Kashiwara and Pierre Schapira. Deformation quantization modules. Astérisque, (1), 2012. 26
- [30] Jacob Lurie. Higher algebra. 2014. Preprint, available at http://www.math.harvard.edu/~lurie, 2016. 41
- [31] Jacob Lurie. Spectral algebraic geometry. Preprint, available at https://www.math.ias.edu/ lurie, 2018. 7
- [32] Zoghman Mebkhout. Une équivalence de catégories. Compositio mathematica, 51(1):51–62, 1984. 2
- [33] David Nadler and Vivek Shende. Sheaf quantization in Weinstein symplectic manifolds. *arXiv preprint* arXiv:2007.10154, 2020. 2, 3, 12, 13, 14, 15, 16, 18, 19, 22
- [34] Pietro Polesello. Uniqueness of quantization of complex contact manifolds. arXiv preprint math/0512501, 2005.
 4, 35
- [35] Luca Prelli. Specialization and microlocalization of subanalytic sheaves. arXiv preprint math/0702459. 34
- [36] Nick Rozenblyum. Filtered colimits of ∞-categories. *Dennis Gaitsgory's webpage, Notes on Geometric Langlands*, 2012. 11
- [37] Kevin Sackel. Getting a handle on contact manifolds. arXiv preprint arXiv:1905.11965, 2019. 45
- [38] Mikio Sato, Masaki Kashiwara, and Takahiro Kawai. Micro functions and pseudo-differential equations. *Springer Lecture Notes*, 287:265–529, 1973. 2, 30, 33, 34
- [39] Pierre Schapira. *Microdifferential systems in the complex domain*, volume 269. Springer Science & Business Media, 2012. 32, 34
- [40] Paul Seidel and Ivan Smith. A link invariant from the symplectic geometry of nilpotent slices. Duke Mathematical Journal, 134(3):453–514, 2006. 3
- [41] Vivek Shende. Microlocal category for Weinstein manifolds via the h-principle. *Publications of the Research Institute for Mathematical Sciences*, 57(3):1041–1048, 2021. 2, 14
- [42] Ingo Waschkies. The stack of microlocal perverse sheaves. Bulletin de la société mathématique de France, 132(3):397–462, 2004. 3
- [43] Ingo Waschkies. Microlocal Riemann-Hilbert correspondence. Publications of the Research Institute for Mathematical Sciences, 41(1):37–72, 2005. 2, 4, 34, 35