

THE GENERAL KASTLER-KALAU-WALZE TYPE THEOREM FOR THE J -TWIST D_J OF THE DIRAC OPERATOR

SIYAO LIU YONG WANG*

ABSTRACT. In [21] and [22], we proved the Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator on 3-dimensional, 4-dimensional and 6-dimensional almost product Riemannian spin manifold with boundary. In this paper, we generalize our previous conclusions and establish the proof of the general Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator on even-dimensional almost product Riemannian spin manifold with boundary.

1. INTRODUCTION

The noncommutative residue was found in [1, 2]. Since noncommutative residues are of great importance to the study of noncommutative geometry, more and more attention has been attached to the study of noncommutative residues. Connes put forward that the noncommutative residue of the square of the inverse of the Dirac operator was proportioned to the Einstein-Hilbert action, which is called the Kastler-Kalau-Walze theorem now [3, 4]. Kastler gave a brute-force proof of this theorem [5]. In the same time, Kalau-Walze and Ackermann proved this theorem by using the normal coordinates system and the heat kernel expansion, respectively [6, 7]. The result of Connes was extended to the higher dimensional case [8]. Wang generalized the Connes' results to the case of manifolds with boundary in [9, 10], and proved the Kastler-Kalau-Walze type theorem for the Dirac operator and the signature operator on lower-dimensional manifolds with boundary. In [11], for the Dirac operator, Wang computed $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-1}]$, in these cases the boundary term vanished. Most of the operators studied in the literature have the leading symbol $\sqrt{-1}c(\xi)$ [12, 13, 14, 15, 16]. However, Wu and Wang studied operators with the leading symbol $-\widehat{c}(V)c(\xi)$. In [17], Wu and Wang gave the proof of Kastler-Kalau-Walze type theorems of the operators $\sqrt{-1}\widehat{c}(V)(d+\delta)$ and $-\sqrt{-1}\widehat{c}(V)(d+\delta)$ on 3,4-dimensional oriented compact manifolds with or without boundary. On the other hand, some preliminaries and lemmas about the Dirac operator D and the J -twist are given in [18]. In [19, 20], the author got estimates on the higher eigenvalues of the Dirac operator on locally reducible Riemannian manifolds by the J -twist of the Dirac operator. It can be obtained by simple calculations that the leading symbol of the J -twist D_J of the Dirac

2010 Mathematics Subject Classification: 53C40; 53C42.

Key words and phrases: Dirac operator; the J -twist of the Dirac operator; Kastler-Kalau-Walze type theorems.

* Corresponding author.

operator is not $\sqrt{-1}c(\xi)$. Therefore, Liu and Wang proved the Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator on 3-dimensional, 4-dimensional and 6-dimensional almost product Riemannian spin manifold with boundary [21, 22].

In [23] and [24], the authors established the general Kastler-Kalau-Walze type theorems for any dimensional manifolds with boundary which generalizes the results in [12, 13]. **The motivation of this paper** is to prove the general Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator on even-dimensional almost product Riemannian spin manifold with boundary. The novelty of our paper is to the computations related to the negative higher power of D_J . Motivated by [21] and [22], we compute the generalized noncommutative residue $\widetilde{\text{Wres}}[\pi^+(D_J^{-1}) \circ \pi^+(D_J^{-n+3})]$ on even-dimensional manifold with boundary. Our main theorem is as follows.

Theorem 1.1. *Let M be an n -dimensional almost product Riemannian spin manifold with the boundary ∂M , then we get the following equality:*

(1.1)

$$\begin{aligned}
\widetilde{\text{Wres}}[\pi^+(D_J^{-1}) \circ \pi^+(D_J^{-n+3})] &= \frac{(n-2)\pi^{\frac{n}{2}}}{(\frac{n}{2}-1)!} \int_M 2^{\frac{n}{2}} \left(\frac{1}{4} \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i) \right. \\
&\quad - \frac{1}{2} \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) - \frac{1}{2} \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j \\
&\quad - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j) \\
&\quad \cdot g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) + \frac{1}{4} \sum_{\nu,j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{12}s \Big) d\text{Vol}_M + \int_{\partial M} \left[2^{\frac{4-n}{2}} K \right. \\
&\quad \cdot \left(\frac{\pi(n-2)^2}{n^2+n-2} \cdot \frac{(n-3)!}{(\frac{n}{2})!(\frac{n}{2}-2)!} + \sum_{i=1}^n \langle J(e_i), e_n \rangle^2 \cdot \frac{\pi n(n^2-8n+12)}{8(n-1)} \cdot \frac{(n-3)!}{(\frac{n}{2}+1)!(\frac{n}{2})!} \right. \\
&\quad \left. - \langle J(e_n), e_n \rangle^2 \cdot \frac{\pi(n^2-3n-2)}{n^2+n-2} \cdot \frac{(n-3)!}{(\frac{n}{2})!(\frac{n}{2}-2)!} \right) - 2^{-\frac{2+n}{2}} \sum_{i=1}^n g^M(J(e_i), (\nabla_{e_i}^L J)e_n) \\
&\quad \left. \cdot \pi(n^2-8n+12) \cdot \frac{(n-2)!}{(\frac{n}{2}+1)!(\frac{n}{2}-1)!} \right] \text{Vol}(S^{n-2}) d\text{Vol}_{\partial M},
\end{aligned}$$

where n is the even number, s is the scalar curvature, $\text{Vol}(S^{n-2})$ is the canonical volume of S^{n-2} and K are defined in section 3.

A brief description of the organization of this paper is as follows. In Section 2, this paper will first introduce the basic notions of Boutet de Monvel's calculus and the definition of the noncommutative residue for manifolds with boundary. In the next section, we recall some basic facts and formulas about the J -twist D_J of the Dirac operator and prove the proof of the general Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator on even-dimensional almost product Riemannian spin manifold with boundary.

2. BOUTET DE MONVEL'S CALCULUS AND THE DEFINITION OF THE NONCOMMUTATIVE RESIDUE

In this section, we explain the basic notions of Boutet de Monvel's calculus and the definition of the noncommutative residue for manifolds with boundary that will be used throughout the paper. For the details, see Ref.[11].

Let M be an n -dimensional compact oriented manifold with the boundary ∂M . We assume that the metric g^{TM} on M has the following form near the boundary,

$$(2.1) \quad g^M = \frac{1}{h(x_n)} g^{\partial M} + dx_n^2,$$

where $g^{\partial M}$ is the metric on ∂M and $h(x_n) \in C^\infty([0, 1]) := \{\widehat{h}|_{[0,1]} | \widehat{h} \in C^\infty((-\varepsilon, 1))\}$ for some $\varepsilon > 0$ and $h(x_n)$ satisfies $h(x_n) > 0$, $h(0) = 1$, where x_n denotes the normal directional coordinate. Let $U \subset M$ be a collar neighborhood of ∂M which is diffeomorphic with $\partial M \times [0, 1]$. By the definition of $h(x_n) \in C^\infty([0, 1])$ and $h(x_n) > 0$, there exists $\widehat{h} \in C^\infty((-\varepsilon, 1))$ such that $\widehat{h}|_{[0,1]} = h$ and $\widehat{h} > 0$ for some sufficiently small $\varepsilon > 0$. Then there exists a metric g' on $\widetilde{M} = M \cup_{\partial M} \partial M \times (-\varepsilon, 0]$ which has the form on $U \cup_{\partial M} \partial M \times (-\varepsilon, 0]$

$$(2.2) \quad g' = \frac{1}{\widehat{h}(x_n)} g^{\partial M} + dx_n^2,$$

such that $g'|_M = g$. We fix a metric g' on the \widetilde{M} such that $g'|_M = g$.

Let the Fourier transformation F' be

$$F' : L^2(\mathbf{R}_t) \rightarrow L^2(\mathbf{R}_v); \quad F'(u)(v) = \int_{\mathbf{R}} e^{-ivt} u(t) dt$$

and let

$$r^+ : C^\infty(\mathbf{R}) \rightarrow C^\infty(\widetilde{\mathbf{R}}^+); \quad f \rightarrow f|_{\widetilde{\mathbf{R}}^+}; \quad \widetilde{\mathbf{R}}^+ = \{x \geq 0; x \in \mathbf{R}\}.$$

We define $H^+ = F'(\Phi(\widetilde{\mathbf{R}}^+))$; $H_0^- = F'(\Phi(\widetilde{\mathbf{R}}^-))$ which satisfies $H^+ \perp H_0^-$, where $\Phi(\widetilde{\mathbf{R}}^+) = r^+ \Phi(\mathbf{R})$, $\Phi(\widetilde{\mathbf{R}}^-) = r^- \Phi(\mathbf{R})$ and $\Phi(\mathbf{R})$ denotes the Schwartz space. We have the following property: $h \in H^+$ (resp. H_0^-) if and only if $h \in C^\infty(\mathbf{R})$ which has an analytic extension to the lower (resp. upper) complex half-plane $\{\text{Im} \xi < 0\}$ (resp. $\{\text{Im} \xi > 0\}$) such that for all nonnegative integer l ,

$$\frac{d^l h}{d\xi^l}(\xi) \sim \sum_{k=1}^{\infty} \frac{d^k}{d\xi^k} \left(\frac{c_k}{\xi^k} \right),$$

as $|\xi| \rightarrow +\infty$, $\text{Im} \xi \leq 0$ (resp. $\text{Im} \xi \geq 0$) and where $c_k \in \mathbf{C}$ are some constants.

Let H' be the space of all polynomials and $H^- = H_0^- \oplus H'$; $H = H^+ \oplus H^-$. Denote by π^+ (resp. π^-) the projection on H^+ (resp. H^-). Let $\widetilde{H} = \{\text{rational functions having no poles on the real axis}\}$. Then on \widetilde{H} ,

$$(2.3) \quad \pi^+ h(\xi_0) = \frac{1}{2\pi i} \lim_{u \rightarrow 0^-} \int_{\Gamma^+} \frac{h(\xi)}{\xi_0 + iu - \xi} d\xi,$$

where Γ^+ is a Jordan closed curve included $\text{Im}(\xi) > 0$ surrounding all the singularities of h in the upper half-plane and $\xi_0 \in \mathbf{R}$. In our computations, we only compute π^+h for h in \tilde{H} . Similarly, define π' on \tilde{H} ,

$$(2.4) \quad \pi'h = \frac{1}{2\pi} \int_{\Gamma^+} h(\xi) d\xi.$$

So $\pi'(H^-) = 0$. For $h \in H \cap L^1(\mathbf{R})$, $\pi'h = \frac{1}{2\pi} \int_{\mathbf{R}} h(v) dv$ and for $h \in H^+ \cap L^1(\mathbf{R})$, $\pi'h = 0$.

An operator of order $m \in \mathbf{Z}$ and type d is a matrix

$$\tilde{A} = \begin{pmatrix} \pi^+P + G & K \\ T & \tilde{S} \end{pmatrix} : \begin{array}{c} C^\infty(M, E_1) \\ \oplus \\ C^\infty(\partial M, F_1) \end{array} \longrightarrow \begin{array}{c} C^\infty(M, E_2) \\ \oplus \\ C^\infty(\partial M, F_2) \end{array},$$

where M is a manifold with boundary ∂M and E_1, E_2 (resp. F_1, F_2) are vector bundles over M (resp. ∂M). Here, $P : C_0^\infty(\Omega, \overline{E_1}) \rightarrow C^\infty(\Omega, \overline{E_2})$ is a classical pseudo-differential operator of order m on Ω , where Ω is a collar neighborhood of M and $\overline{E_i}|_M = E_i$ ($i = 1, 2$). P has an extension: $\mathcal{E}'(\Omega, \overline{E_1}) \rightarrow \mathcal{D}'(\Omega, \overline{E_2})$, where $\mathcal{E}'(\Omega, \overline{E_1})$ ($\mathcal{D}'(\Omega, \overline{E_2})$) is the dual space of $C^\infty(\Omega, \overline{E_1})$ ($C_0^\infty(\Omega, \overline{E_2})$). Let $e^+ : C^\infty(M, E_1) \rightarrow \mathcal{E}'(\Omega, \overline{E_1})$ denote extension by zero from M to Ω and $r^+ : \mathcal{D}'(\Omega, \overline{E_2}) \rightarrow \mathcal{D}'(\Omega, E_2)$ denote the restriction from Ω to X , then define

$$\pi^+P = r^+Pe^+ : C^\infty(M, E_1) \rightarrow \mathcal{D}'(\Omega, E_2).$$

In addition, P is supposed to have the transmission property; this means that, for all j, k, α , the homogeneous component p_j of order j in the asymptotic expansion of the symbol p of P in local coordinates near the boundary satisfies:

$$\partial_{x_n}^k \partial_\xi^\alpha p_j(x', 0, 0, +1) = (-1)^{j-|\alpha|} \partial_{x_n}^k \partial_\xi^\alpha p_j(x', 0, 0, -1),$$

then $\pi^+P : C^\infty(M, E_1) \rightarrow C^\infty(M, E_2)$. Let G, T be respectively the singular Green operator and the trace operator of order m and type d . Let K be a potential operator and S be a classical pseudo-differential operator of order m along the boundary. Denote by $B^{m,d}$ the collection of all operators of order m and type d , and \mathcal{B} is the union over all m and d .

Recall that $B^{m,d}$ is a Fréchet space. The composition of the above operator matrices yields a continuous map: $B^{m,d} \times B^{m',d'} \rightarrow B^{m+m', \max\{m'+d, d'\}}$. Write

$$\tilde{A} = \begin{pmatrix} \pi^+P + G & K \\ T & \tilde{S} \end{pmatrix} \in B^{m,d}, \tilde{A}' = \begin{pmatrix} \pi^+P' + G' & K' \\ T' & \tilde{S}' \end{pmatrix} \in B^{m',d'}.$$

The composition $\tilde{A}\tilde{A}'$ is obtained by multiplication of the matrices (For more details see [25]). For example $\pi^+P \circ G'$ and $G \circ G'$ are singular Green operators of type d' and

$$\pi^+P \circ \pi^+P' = \pi^+(PP') + L(P, P').$$

Here PP' is the usual composition of pseudo-differential operators and $L(P, P')$ called leftover term is a singular Green operator of type $m' + d$. For our case, P, P' are classical pseudo-differential operators, in other words $\pi^+P \in \mathcal{B}^\infty$ and $\pi^+P' \in \mathcal{B}^\infty$.

Let M be an n -dimensional compact oriented manifold with the boundary ∂M . Denote by \mathcal{B} the Boutet de Monvel's algebra. We recall that the main theorem in [26].

Theorem 2.1. [26] (**Fedosov-Golse-Leichtnam-Schrohe**) *Let M and ∂M be connected, $\dim M = n \geq 3$, and let \tilde{S} (resp. \tilde{S}') be the unit sphere about ξ (resp. ξ') and $\sigma(\xi)$ (resp. $\sigma(\xi')$) be the corresponding canonical $n - 1$ (resp. $(n - 2)$) volume form. Set $\tilde{A} = \begin{pmatrix} \pi^+P + G & K \\ T & \tilde{S} \end{pmatrix} \in \mathcal{B}$, and denote by p, b and s the local symbols of P, G and \tilde{S} respectively. Define:*

$$(2.5) \quad \begin{aligned} \widetilde{\text{Wres}}(\tilde{A}) &= \int_X \int_{\tilde{S}} \text{tr}_E [p_{-n}(x, \xi)] \sigma(\xi) dx \\ &+ 2\pi \int_{\partial X} \int_{\tilde{S}'} \{ \text{tr}_E [(\text{tr} b_{-n})(x', \xi')] + \text{tr}_F [s_{1-n}(x', \xi')] \} \sigma(\xi') dx', \end{aligned}$$

where $\widetilde{\text{Wres}}$ denotes the noncommutative residue of an operator in the Boutet de Monvel's algebra. Then a) $\widetilde{\text{Wres}}([\tilde{A}, B]) = 0$, for any $\tilde{A}, B \in \mathcal{B}$; b) It is the unique continuous trace on $\mathcal{B}/\mathcal{B}^{-\infty}$.

Let p_1, p_2 be non-negative integers and $p_1 + p_2 \leq n$.

Definition 2.2. [11] *Lower dimensional volumes of spin manifolds with boundary are defined by*

$$(2.6) \quad \text{Vol}_n^{(p_1, p_2)} M := \widetilde{\text{Wres}}[\pi^+ D^{-p_1} \circ \pi^+ D^{-p_2}].$$

Denote by $\sigma_l(D)$ the l -order symbol of the Dirac operator D . By [11], we get

$$(2.7) \quad \widetilde{\text{Wres}}[\pi^+ D^{-p_1} \circ \pi^+ D^{-p_2}] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)} [\sigma_{-n}(D^{-p_1-p_2})] \sigma(\xi) dx + \int_{\partial M} \Phi$$

and

$$(2.8) \quad \begin{aligned} \Phi &= \int_{|\xi|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{trace}_{S(TM)} [\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D^{-p_1})(x', 0, \xi', \xi_n)] \\ &\times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D^{-p_2})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned}$$

where the sum is taken over $r + l - k - |\alpha| - j - 1 = -n$, $r \leq -p_1, l \leq -p_2$.

Since $[\sigma_{-n}(D^{-p_1-p_2})]|_M$ has the same expression as $\sigma_{-n}(D^{-p_1-p_2})$ in the case of manifolds without boundary, so locally we can compute the first term by [5, 6, 11, 27].

3. THE NONCOMMUTATIVE RESIDUE $\widetilde{\text{Wres}}[\pi^+(D_J^{-1}) \circ \pi^+(D_J^{-n+3})]$ ON
EVEN-DIMENSIONAL MANIFOLDS WITH BOUNDARY

We give some definitions and basic notions which we will use in this paper. Let M be an n -dimensional ($n \geq 6$) oriented compact Riemannian manifold with a Riemannian metric g^M . We recall that the Dirac operator D is locally given as follows:

$$(3.1) \quad D = \sum_{i,j=1}^n g^{ij} c(\partial_i) \nabla_{\partial_j}^S = \sum_{i=1}^n c(e_i) \nabla_{e_i}^S,$$

where ∂_i is a natural local frame on TM , $(g^{ij})_{1 \leq i,j \leq n}$ is the inverse matrix associated to the metric matrix $(g_{ij})_{1 \leq i,j \leq n}$ on M and $c(e_i)$ be the Clifford action which satisfies the relation

$$(3.2) \quad c(e_i)c(e_j) + c(e_j)c(e_i) = -2g^M(e_i, e_j) = -2\delta_{ij}^j,$$

$$(3.3) \quad \nabla_{\partial_j}^S = \partial_j + \sigma_j$$

and

$$(3.4) \quad \sigma_i = \frac{1}{4} \sum_{j,k=1}^n \langle \nabla_{\partial_i}^L e_j, e_k \rangle c(e_j)c(e_k).$$

Let J be a $(1, 1)$ -tensor field on (M, g^M) such that $J^2 = \text{id}$,

$$(3.5) \quad g^M(J(X), J(Y)) = g^M(X, Y),$$

for all vector fields $X, Y \in \Gamma(TM)$. Here id stands for the identity map. (M, g^M, J) is an almost product Riemannian manifold. We can also define on almost product Riemannian spin manifold the following J -twist D_J of the Dirac operator D by

$$(3.6) \quad D_J := \sum_{i=1}^n c(e_i) \nabla_{J(e_i)}^S = \sum_{i=1}^n c[J(e_i)] \nabla_{e_i}^S.$$

Let $\xi = \sum_k \xi_j dx_j$, $\nabla_{\partial_i}^L \partial_j = \sum_k \Gamma_{ij}^k \partial_k$, we denote that

$$\Gamma^k = g^{ij} \Gamma_{ij}^k; \quad \sigma^j = g^{ij} \sigma_i; \quad \partial^j = g^{ij} \partial_i,$$

where Γ_{ij}^k is the Christoffel coefficient of Levi-Civita connection ∇^L .

In this section, we want to compute the noncommutative residue $\widetilde{\text{Wres}}[\pi^+(D_J^{-1}) \circ \pi^+(D_J^{-n+3})]$ on n -dimensional almost product Riemannian spin manifold with the boundary. By (2.7) and (2.8), we can rewrite the noncommutative residue as

$$(3.7) \quad \widetilde{\text{Wres}}[\pi^+(D_J^{-1}) \circ \pi^+(D_J^{-n+3})] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)}[\sigma_{-n}(D_J^{-n+2})] \sigma(\xi) dx + \int_{\partial M} \Phi$$

and

$$(3.8) \quad \begin{aligned} \Phi = & \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{trace}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D_J^{-1})(x', 0, \xi', \xi_n) \\ & \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D_J^{-n+3})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned}$$

where the sum is taken over $r+l-k-|\alpha|-j-1=-n$, $r \leq -1, l \leq -n+3$.

By [21] and [22], we have

Theorem 3.1. [21, 22] *If M is an n -dimensional almost product Riemannian spin manifold without boundary, we have the following:*

$$(3.9) \quad \begin{aligned} \text{Wres}(D_J^{-n+2}) = & \frac{(n-2)\pi^{\frac{n}{2}}}{(\frac{n}{2}-1)!} \int_M 2^{\frac{n}{2}} \left(\frac{1}{4} \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i) - \frac{1}{2} \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) \right. \\ & - \frac{1}{2} \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j) \\ & - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) \\ & - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j) g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) \\ & \left. + \frac{1}{4} \sum_{\nu,j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{12}s \right) d\text{Vol}_M. \end{aligned}$$

Locally we can use Theorem 3.1 to compute the interior term of (3.7), then

$$(3.10) \quad \begin{aligned} & \int_M \int_{|\xi|=1} \text{trace}_{S(TM)}[\sigma_{-n}(D_J^{-n+2})]\sigma(\xi) dx \\ & = \frac{(n-2)\pi^{\frac{n}{2}}}{(\frac{n}{2}-1)!} \int_M 2^{\frac{n}{2}} \left(\frac{1}{4} \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i) - \frac{1}{2} \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) \right. \\ & - \frac{1}{2} \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j) \\ & \left. - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) \right) \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{4} \sum_{\alpha, \nu, j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j) g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) \\
& + \frac{1}{4} \sum_{\nu, j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{12} s) d\text{Vol}_M.
\end{aligned}$$

In the following, we will compute $\int_{\partial M} \Phi$. As shown in [21] and [22], we see that

Lemma 3.2. [21] *The following identities hold:*

$$(3.11) \quad \sigma_1(D_J) = ic[J(\xi)];$$

$$(3.12) \quad \sigma_0(D_J) = -\frac{1}{4} \sum_{i, j, k=1}^n \omega_{j, k}(e_i) c[J(e_i)] c(e_j) c(e_k),$$

where $(\omega_{j, k})$ denotes the connection matrix of Levi-Civita connection ∇^L .

Lemma 3.3. [21] *The following identities hold:*

$$(3.13)$$

$$\sigma_{-1}(D_J^{-1}) = \frac{ic[J(\xi)]}{|\xi|^2};$$

$$(3.14)$$

$$\sigma_{-2}(D_J^{-1}) = \frac{c[J(\xi)]\sigma_0(D_J)c[J(\xi)]}{|\xi|^4} + \frac{c[J(\xi)]}{|\xi|^6} \sum_{j=1}^n c[J(dx_j)] \left[\partial_{x_j}(c[J(\xi)])|\xi|^2 - c[J(\xi)]\partial_{x_j}(|\xi|^2) \right].$$

Lemma 3.4. [22] *The following identities hold:*

$$(3.15) \quad \sigma_3(D_J^3) = ic[J(\xi)]|\xi|^2;$$

$$\begin{aligned}
(3.16) \quad \sigma_2(D_J^3) &= \sum_{i, j, l=1}^n c[J(dx_l)] \partial_l(g^{ij}) \xi_i \xi_j + c[J(\xi)](2\sigma^k - \Gamma^k) \xi_k \\
&\quad - \sum_{\alpha=1}^n c[J(\xi)] c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J)(\xi^*)] - \frac{1}{4} |\xi|^2 \sum_{s, t, l=1}^n \omega_{s, t}(e_l) c[J(e_l)] c(e_s) c(e_t),
\end{aligned}$$

where $\xi^* = \sum_{\beta=1}^n \langle e_\beta, \xi \rangle e_\beta$.

According to (2.13) in [21], we have

Lemma 3.5. *The following identities hold:*

$$(3.17) \quad \sigma_2(D_J^2) = |\xi|^2;$$

$$(3.18) \quad \sigma_1(D_J^2) = i(-2\sigma^k + \Gamma^k) \xi_k + i \sum_{\alpha=1}^n c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J)(\xi^*)].$$

Write

$$(3.19) \quad D_x^\alpha = (-i)^{|\alpha|} \partial_x^\alpha; \quad \sigma(D_J^2) = p_2 + p_1 + p_0; \quad \sigma(D_J^{-2}) = \sum_{j=2}^{\infty} q_{-j}.$$

By the composition formula of pseudodifferential operators, we have

$$(3.20) \quad \begin{aligned} 1 = \sigma(D_J^2 \circ D_J^{-2}) &= \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma(D_J^2)] D_x^{\alpha} [\sigma(D_J^{-2})] \\ &= (p_2 + p_1 + p_0)(q_{-2} + q_{-3} + \cdots) \\ &\quad + \sum_j (\partial_{\xi_j} p_2 + \partial_{\xi_j} p_1 + \partial_{\xi_j} p_0)(D_{x_j} q_{-2} + D_{x_j} q_{-3} + \cdots) \\ &= p_2 q_{-2} + (p_2 q_{-3} + p_1 q_{-2} + \sum_j \partial_{\xi_j} p_2 D_{x_j} q_{-2}) + \cdots, \end{aligned}$$

so

$$(3.21) \quad q_{-2} = p_2^{-1}; \quad q_{-3} = -p_2^{-1} [p_1 p_2^{-1} + \sum_j \partial_{\xi_j} p_2 D_{x_j} (p_2^{-1})].$$

Lemma 3.6. *The following identities hold:*

$$(3.22) \quad \sigma_{-2}(D_J^{-2}) = |\xi|^{-2};$$

$$(3.23) \quad \begin{aligned} \sigma_{-3}(D_J^{-2}) &= -i|\xi|^{-4}(-2\sigma^k + \Gamma^k)\xi_k - i|\xi|^{-4} \sum_{\alpha=1}^n c[J(e_{\alpha})]c[(\nabla_{e_{\alpha}}^L J)(\xi^*)] \\ &\quad - i|\xi|^{-6} \sum_{j=1}^n 2\xi^j \xi_{\alpha} \xi_{\beta} \partial_{x_j} (g^{\alpha\beta}). \end{aligned}$$

Since $\sigma_{-2}(D_J^{-2}) = |\xi|^{-2}$, it follows that $\sigma_{-n}(D_J^{-n}) = |\xi|^{-n}$ and

$$(3.24) \quad \begin{aligned} \sigma_{-n+3}(D_J^{-n+3}) &= \sigma_{-n+3}(D_J^{-n} \cdot D_J^3) \\ &= |\xi|^{-n} \cdot ic[J(\xi)]|\xi|^2 = ic[J(\xi)]|\xi|^{-n+2}. \end{aligned}$$

From [24], we similarly can obtain

$$(3.25) \quad \begin{aligned} \sigma_{-n+2}(D_J^{-n+3}) &= \sigma_{-n+2}(D_J^{-n+2} \cdot D_J) \\ &= \left\{ \sum_{|\alpha|=0}^{+\infty} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma(D_J^{-n+2})] D_x^{\alpha} [\sigma(D_J)] \right\}_{-n+2} \\ &= \sigma_{-n+2}(D_J^{-n+2})\sigma_0(D_J) - i \sum_{j=1}^n \partial_{\xi_j} [\sigma_{-n+2}(D_J^{-n+2})] \partial_{x_j} [\sigma_1(D_J)] \\ &\quad + \left\{ \frac{n-2}{2} \sigma_2^{(-\frac{n}{2}+2)}(D_J^2) \sigma_{-3}(D_J^{-2}) - i \sum_{k=0}^{\frac{n}{2}-3} \sum_{\mu=1}^n \partial_{\xi_{\mu}} \sigma_2^{(-\frac{n}{2}+k+2)}(D_J^2) \right\} \end{aligned}$$

$$\times \partial_{x_\mu} \sigma_2^{-1}(D_J^2)(\sigma_2(D_J^2))^{(-k)} \Big\} \sigma_1(D_J).$$

Since Φ is a global form on ∂M , so for any fixed point $x_0 \in \partial M$, we can choose the normal coordinates U of x_0 in ∂M (not in M) and compute $\Phi(x_0)$ in the coordinates $\tilde{U} = U \times [0, 1) \subset M$ and the metric $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$. The dual metric of g^M on \tilde{U} is $h(x_n)g^{\partial M} + dx_n^2$. Write $g_{ij}^M = g^M(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j})$; $g_M^{ij} = g^M(dx_i, dx_j)$, then

$$(3.26) \quad [g_{ij}^M] = \begin{bmatrix} \frac{1}{h(x_n)}[g_{ij}^{\partial M}] & 0 \\ 0 & 1 \end{bmatrix}; \quad [g_M^{ij}] = \begin{bmatrix} h(x_n)[g_{\partial M}^{ij}] & 0 \\ 0 & 1 \end{bmatrix}$$

and

$$(3.27) \quad \partial_{x_s} g_{ij}^{\partial M}(x_0) = 0, 1 \leq i, j \leq n-1; \quad g_{ij}^M(x_0) = \delta_i^j.$$

$\{e_1, \dots, e_n\}$ be an orthonormal frame field in U about $g^{\partial M}$ which is parallel along geodesics and $e_i(x_0) = \frac{\partial}{\partial x_i}(x_0)$. We review the following three lemmas.

Lemma 3.7. [11] *With the metric g^M on M near the boundary*

$$(3.28) \quad \partial_{x_j}(|\xi|_{g^M}^2)(x_0) = \begin{cases} 0, & \text{if } j < n, \\ h'(0)|\xi'|_{g^{\partial M}}^2, & \text{if } j = n; \end{cases}$$

$$(3.29) \quad \partial_{x_j}[c(\xi)](x_0) = \begin{cases} 0, & \text{if } j < n, \\ \partial_{x_n}(c(\xi'))(x_0), & \text{if } j = n, \end{cases}$$

where $\xi = \xi' + \xi_n dx_n$.

Lemma 3.8. [11] *With the metric g^M on M near the boundary*

$$(3.30) \quad \omega_{s,t}(e_i)(x_0) = \begin{cases} \omega_{n,i}(e_i)(x_0) = \frac{1}{2}h'(0), & \text{if } s = n, t = i, i < n; \\ \omega_{i,n}(e_i)(x_0) = -\frac{1}{2}h'(0), & \text{if } s = i, t = n, i < n; \\ \omega_{s,t}(e_i)(x_0) = 0, & \text{other cases.} \end{cases}$$

Lemma 3.9. [11]

$$(3.31) \quad \Gamma_{st}^k(x_0) = \begin{cases} \Gamma_{ii}^n(x_0) = \frac{1}{2}h'(0), & \text{if } s = t = i, k = n, i < n; \\ \Gamma_{ni}^i(x_0) = -\frac{1}{2}h'(0), & \text{if } s = n, t = i, k = i, i < n; \\ \Gamma_{in}^i(x_0) = -\frac{1}{2}h'(0), & \text{if } s = i, t = n, k = i, i < n, \\ \Gamma_{st}^i(x_0) = 0, & \text{other cases.} \end{cases}$$

When n is even, then $\text{tr}[\text{id}] = 2^{\frac{n}{2}}$, where tr as shorthand of $\text{traces}_{S(TM)}$, since the sum is taken over $r+l-k-j-|\alpha|-1 = -n$, $r \leq -1, l \leq -n+3$, then we have the following five cases:

case a) I) $r = -1, l = -n+3, k = j = 0, |\alpha| = 1$.

By applying the formula shown in (3.8), we can calculate

$$(3.32) \quad \Phi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{x'}^{\alpha} \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx'.$$

For $i < n$, (3.39) in [21] makes it obvious that

(3.33)

$$\begin{aligned} \pi_{\xi_n}^+ \partial_{\xi_i} \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} &= \frac{1}{2(\xi_n - i)} c[J(dx_i)] + \frac{2i - \xi_n}{2(\xi_n - i)^2} \sum_{q=1}^{n-1} \xi_i \xi_q c[J(dx_q)] \\ &\quad - \frac{1}{2(\xi_n - i)^2} \xi_i c[J(dx_n)] \\ &= \frac{1}{2(\xi_n - i)} \sum_{\alpha=1}^n a_{\alpha}^i c(dx_{\alpha}) + \frac{2i - \xi_n}{2(\xi_n - i)^2} \sum_{\beta=1}^n \sum_{q=1}^{n-1} \xi_i \xi_q a_{\beta}^q c(dx_{\beta}) \\ &\quad - \frac{1}{2(\xi_n - i)^2} \sum_{\gamma=1}^n \xi_i a_{\gamma}^n c(dx_{\gamma}), \end{aligned}$$

where $c[J(dx_p)] = \sum_{h=1}^n a_h^p c(dx_h)$.

We get

(3.34)

$$\partial_{x_i} (ic[J(\xi)]|\xi|^{-n+2}) (x_0) = \frac{i\partial_{x_i}(c[J(\xi)])(x_0)}{|\xi|^{n-2}} - \frac{ic[J(\xi)]\partial_{x_i}(|\xi|^{n-2})(x_0)}{|\xi|^{2n-4}} = \frac{i\partial_{x_i}(c[J(\xi)])(x_0)}{|\xi|^{n-2}},$$

of course,

(3.35)

$$\begin{aligned} \partial_{x_i} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0) &= \frac{i\partial_{x_i}(c[J(\sum_{p=1}^n \xi_p dx_p)])(x_0)}{|\xi|^{n-2}} = \frac{i \sum_{p=1}^n \xi_p \partial_{x_i}(c(\sum_{h=1}^n a_h^p dx_h))(x_0)}{|\xi|^{n-2}} \\ &= \frac{i \sum_{p,h=1}^n \xi_p \partial_{x_i}(a_h^p) c(dx_h)(x_0)}{|\xi|^{n-2}} + \frac{i \sum_{p,h=1}^n \xi_p a_h^p \partial_{x_i}(c(dx_h))(x_0)}{|\xi|^{n-2}} \\ &= \frac{i \sum_{p,h=1}^n \xi_p \partial_{x_i}(a_h^p) c(dx_h)(x_0)}{|\xi|^{n-2}}. \end{aligned}$$

When $|\xi'| = 1$, we see that

(3.36)

$$\partial_{x_i} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0)|_{|\xi'|=1} = \frac{i \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_i}(a_h^p) c(dx_h)(x_0)}{(1 + \xi_n^2)^{\frac{n}{2}-1}} + \frac{i \sum_{h=1}^n \xi_n \partial_{x_i}(a_h^n) c(dx_h)(x_0)}{(1 + \xi_n^2)^{\frac{n}{2}-1}}.$$

An easy computation shows that

$$(3.37) \quad \partial_{\xi_n} \partial_{x_i} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0)|_{|\xi'|=1} = i \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_i}(a_h^p) c(dx_h) \partial_{\xi_n} \left(\frac{1}{(1 + \xi_n^2)^{\frac{n}{2}-1}} \right)$$

$$\begin{aligned}
& + i \sum_{h=1}^n \partial_{x_i}(a_h^n) c(dx_h) \partial_{\xi_n} \left(\frac{\xi_n}{(1 + \xi_n^2)^{\frac{n-1}{2}}} \right) \\
& = - \frac{i(n-2)\xi_n}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_i}(a_h^p) c(dx_h) \\
& + \frac{i[1 - (n-3)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \partial_{x_i}(a_h^n) c(dx_h).
\end{aligned}$$

Hence, we have

$$\begin{aligned}
(3.38) \quad & \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) |_{|\xi'|=1} \\
& = - \frac{i(n-2)\xi_n}{2(\xi_n - i)^{\frac{n}{2}+1}(\xi_n + i)^{\frac{n}{2}}} \sum_{\alpha, h=1}^n \sum_{i, p=1}^{n-1} \text{tr}[\xi_p a_\alpha^i \partial_{x_i}(a_h^p) c(dx_\alpha) c(dx_h)] \\
& + \frac{i[1 - (n-3)\xi_n^2]}{2(\xi_n - i)^{\frac{n}{2}+1}(\xi_n + i)^{\frac{n}{2}}} \sum_{\alpha, h=1}^n \sum_{i=1}^{n-1} \text{tr}[a_\alpha^i \partial_{x_i}(a_h^n) c(dx_\alpha) c(dx_h)] \\
& - \frac{i(2i - \xi_n)(n-2)\xi_n}{2(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} \sum_{\beta, h=1}^n \sum_{i, q, p=1}^{n-1} \text{tr}[\xi_i \xi_q \xi_p a_\beta^q \partial_{x_i}(a_h^p) c(dx_\beta) c(dx_h)] \\
& + \frac{i(2i - \xi_n)[1 - (n-3)\xi_n^2]}{2(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} \sum_{\beta, h=1}^n \sum_{i, q=1}^{n-1} \text{tr}[\xi_i \xi_q a_\beta^q \partial_{x_i}(a_h^n) c(dx_\beta) c(dx_h)] \\
& + \frac{i(n-2)\xi_n}{2(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} \sum_{\gamma, h=1}^n \sum_{i, p=1}^{n-1} \text{tr}[\xi_i \xi_p a_\gamma^n \partial_{x_i}(a_h^p) c(dx_\gamma) c(dx_h)] \\
& - \frac{i[1 - (n-3)\xi_n^2]}{2(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} \sum_{\gamma, h=1}^n \sum_{i=1}^{n-1} \text{tr}[\xi_i a_\gamma^n \partial_{x_i}(a_h^n) c(dx_\gamma) c(dx_h)].
\end{aligned}$$

We note that $\int_{|\xi'|=1} \{\xi_{i_1} \cdots \xi_{i_{2d+1}}\} \sigma(\xi') = 0$, this gives

$$\begin{aligned}
(3.39) \quad & \Phi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
& = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i[1 - (n-3)\xi_n^2]}{2(\xi_n - i)^{\frac{n}{2}+1}(\xi_n + i)^{\frac{n}{2}}} \sum_{\alpha, h=1}^n \sum_{i=1}^{n-1} \text{tr}[a_\alpha^i \partial_{x_i}(a_h^n) c(dx_\alpha) c(dx_h)] d\xi_n \sigma(\xi') dx' \\
& - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(2i - \xi_n)[1 - (n-3)\xi_n^2]}{2(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} \sum_{\beta, h=1}^n \sum_{i, q=1}^{n-1} \text{tr}[\xi_i \xi_q a_\beta^q \partial_{x_i}(a_h^n) c(dx_\beta) c(dx_h)] d\xi_n \sigma(\xi') dx'
\end{aligned}$$

$$- \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(n-2)\xi_n}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{\gamma,h=1}^n \sum_{i,p=1}^{n-1} \text{tr}[\xi_i \xi_p a_\gamma^n \partial_{x_i}(a_h^p) c(dx_\gamma) c(dx_h)] d\xi_n \sigma(\xi') dx'.$$

Because $c(e_i)c(e_j) + c(e_j)c(e_i) = -2\delta_i^j$ then by the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, we have the following equalities:

$$(3.40) \quad \sum_{\alpha,h=1}^n \sum_{i=1}^{n-1} \text{tr}[a_\alpha^i \partial_{x_i}(a_h^n) c(dx_\alpha) c(dx_h)] = - \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \text{tr}[\text{id}];$$

$$(3.41) \quad \sum_{\beta,h=1}^n \sum_{i,q=1}^{n-1} \text{tr}[\xi_i \xi_q a_\beta^q \partial_{x_i}(a_h^n) c(dx_\beta) c(dx_h)] = - \sum_{h=1}^n \sum_{i,q=1}^{n-1} \xi_i \xi_q a_h^q \partial_{x_i}(a_h^n) \text{tr}[\text{id}];$$

$$(3.42) \quad \sum_{\gamma,h=1}^n \sum_{i,p=1}^{n-1} \text{tr}[\xi_i \xi_p a_\gamma^n \partial_{x_i}(a_h^p) c(dx_\gamma) c(dx_h)] = - \sum_{h=1}^n \sum_{i,p=1}^{n-1} \xi_i \xi_p a_h^n \partial_{x_i}(a_h^p) \text{tr}[\text{id}].$$

And by $\int_{|\xi'|=1} \sum_{i,j=1}^{n-1} \xi_i \xi_j \sigma(\xi') = \frac{1}{n-1} \sum_{i,j=1}^{n-1} \delta_j^i \text{Vol}(S^{n-2})$ in [28], we have

$$(3.43) \quad \begin{aligned} \Phi_1 &= \int_{-\infty}^{+\infty} \frac{i[1-(n-3)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+1}(\xi_n+i)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\ &+ \int_{-\infty}^{+\infty} \frac{i(2i-\xi_n)[1-(n-3)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\ &+ \int_{-\infty}^{+\infty} \frac{i(n-2)\xi_n}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^n \partial_{x_i}(a_h^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'. \end{aligned}$$

It is easily seen that

$$(3.44) \quad \begin{aligned} \Phi_1 &= \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2})!} \mathcal{A}_0 dx' \\ &+ \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_1 dx' \\ &+ \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^n \partial_{x_i}(a_h^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_2 dx', \end{aligned}$$

where

$$(3.45) \quad \mathcal{A}_0 = \left[\frac{-i(n-3)\xi_n^2 + i}{2(\xi_n+i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}\right)};$$

$$(3.46) \quad \mathcal{A}_1 = \left[\frac{i(n-3)\xi_n^3 + 2(n-3)\xi_n^2 - i\xi_n - 2}{2(\xi_n+i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}+1\right)};$$

$$(3.47) \quad \mathcal{A}_2 = \left[\frac{i(n-2)\xi_n}{2(\xi_n+i)^{\frac{n}{2}}} \right]^{\left(\frac{n}{2}+1\right)} \Big|_{\xi_n=i}.$$

case (a) (II) $r = -1, l = -n + 3, |\alpha| = k = 0, j = 1$.

It is easy to check that

$$(3.48) \quad \Phi_2 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{\xi_n}^2 \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx'.$$

We can assert that

$$(3.49) \quad \begin{aligned} & \pi_{\xi_n}^+ \partial_{x_n} \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0) \Big|_{|\xi'|=1} \\ &= \frac{1}{2(\xi_n - i)} \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_n} (a_h^p) c(dx_h) + \frac{i}{2(\xi_n - i)} \sum_{h=1}^n \partial_{x_n} (a_h^n) c(dx_h) \\ &+ \frac{1}{2(\xi_n - i)} \sum_{h,p=1}^{n-1} \xi_p a_h^p \partial_{x_n} (c(dx_h)) + \frac{i}{2(\xi_n - i)} \sum_{h=1}^{n-1} a_h^n \partial_{x_n} (c(dx_h)) \\ &+ \frac{2i - \xi_n}{4(\xi_n - i)^2} h'(0) \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p a_h^p c(dx_h) - \frac{1}{4(\xi_n - i)^2} h'(0) \sum_{h=1}^n a_h^n c(dx_h), \end{aligned}$$

where $\sum_{h=1}^{n-1} \partial_{x_n} (c(dx_h)) = \sum_{h=1}^{n-1} \frac{1}{2} h'(0) c(dx_h)$.

By calculation, we have

$$(3.50) \quad \begin{aligned} \partial_{\xi_n}^2 \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0) &= \partial_{\xi_n}^2 \left(\frac{i \sum_{i=1}^n \xi_i c[J(dx_i)]}{|\xi|^{n-2}} \right) (x_0) \\ &= \partial_{\xi_n}^2 \left(\frac{i \sum_{i,\beta=1}^n \xi_i a_{\beta}^i c(dx_{\beta})}{|\xi|^{n-2}} \right) (x_0) \end{aligned}$$

and

$$(3.51) \quad \begin{aligned} \partial_{\xi_n}^2 \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0) \Big|_{|\xi'|=1} &= i \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_{\beta}^i c(dx_{\beta}) \partial_{\xi_n}^2 \left(\frac{1}{(1 + \xi_n^2)^{\frac{n}{2}-1}} \right) \\ &+ i \sum_{\beta=1}^n a_{\beta}^n c(dx_{\beta}) \partial_{\xi_n}^2 \left(\frac{\xi_n}{(1 + \xi_n^2)^{\frac{n}{2}-1}} \right) \\ &= \frac{i(n-2)[-1 + (n-1)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}+1}} \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_{\beta}^i c(dx_{\beta}) \\ &+ \frac{i(n-2)\xi_n[-3 + (n-3)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}+1}} \sum_{\beta=1}^n a_{\beta}^n c(dx_{\beta}). \end{aligned}$$

A trivial verification shows that

(3.52)

$$\begin{aligned}
\Phi_2 &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{\xi_n}^2 \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(n-2)[-1+(n-1)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} \sum_{h,\beta=1}^n \sum_{p,i=1}^{n-1} \text{tr}[\xi_p \xi_i a_\beta^i \partial_{x_n}(a_h^p) c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} -\frac{(n-2)\xi_n[-3+(n-3)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} \sum_{h,\beta=1}^n \text{tr}[a_\beta^n \partial_{x_n}(a_h^n) c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(n-2)[-1+(n-1)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} \sum_{\beta=1}^n \sum_{h,p,i=1}^{n-1} \text{tr}[\xi_p \xi_i a_h^p a_\beta^i \partial_{x_n}(c(dx_h)) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} -\frac{(n-2)\xi_n[-3+(n-3)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} \sum_{\beta=1}^n \sum_{h=1}^{n-1} \text{tr}[a_h^n a_\beta^n \partial_{x_n}(c(dx_h)) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(n-2)(2i-\xi_n)[-1+(n-1)\xi_n^2]}{4(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h,\beta=1}^n \sum_{p,i=1}^{n-1} \text{tr}[\xi_p \xi_i a_h^p a_\beta^i c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} -\frac{i(n-2)\xi_n[-3+(n-3)\xi_n^2]}{4(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h,\beta=1}^n \text{tr}[a_h^n a_\beta^n c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx'.
\end{aligned}$$

As in the proof of case (a) (I), equation (3.52) gives

(3.53)

$$\begin{aligned}
\Phi_2 &= -\frac{1}{2} \int_{-\infty}^{+\infty} -\frac{i(n-2)[-1+(n-1)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_n}(a_h^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n[-3+(n-3)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} \sum_{h=1}^n a_h^n \partial_{x_n}(a_h^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} -\frac{i(n-2)[-1+(n-1)\xi_n^2]}{4(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h,i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n[-3+(n-3)\xi_n^2]}{4(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^{n-1} (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} -\frac{i(n-2)(2i-\xi_n)[-1+(n-1)\xi_n^2]}{4(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'
\end{aligned}$$

$$-\frac{1}{2} \int_{-\infty}^{+\infty} \frac{i(n-2)\xi_n[-3+(n-3)\xi_n^2]}{4(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^n (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'.$$

We can certainly assume that $J(\frac{\partial}{\partial x_l}) = \sum_{q=1}^n \tilde{a}_l^q \frac{\partial}{\partial x_q}$, since

$$(3.54) \quad \left\langle \frac{\partial}{\partial x_l}, J(dx_p) \right\rangle = \left\langle J\left(\frac{\partial}{\partial x_l}\right), dx_p \right\rangle,$$

then we have $a_h^q = \tilde{a}_h^q$. As

$$(3.55) \quad J^2(x_0) = \text{id}, J^T J(x_0) = \text{id},$$

we have $a_l^p a_p^j(x_0) = \delta_l^j$, $a_l^p(x_0) = a_p^l(x_0)$. Let $J(e_l) = \sum_{q=1}^n b_l^q e_q$, then $\partial x_j(b_l^q)(x_0) = \partial x_j(a_l^q)(x_0)$ and $b_l^q(x_0) = a_l^q(x_0)$. We next show that

$$(3.56) \quad 0 = \sum_{i=1}^{n-1} \partial_{x_n}(\delta_i^i) = \sum_{h=1}^n \sum_{i=1}^{n-1} \partial_{x_n}(a_h^i a_h^i) = 2 \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_n}(a_h^i);$$

$$(3.57) \quad 0 = \partial_{x_n}(\delta_n^n) = \sum_{h=1}^n \partial_{x_n}(a_h^n a_h^n) = 2 \sum_{h=1}^n a_h^n \partial_{x_n}(a_h^n).$$

Therefore

$$(3.58) \quad \begin{aligned} \Phi_2 &= h'(0) \sum_{h,i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_3 dx' \\ &+ h'(0) \sum_{i=1}^{n-1} (a_i^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_4 dx' \\ &+ h'(0) \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{A}_5 dx' \\ &+ h'(0) \sum_{h=1}^n (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{A}_6 dx', \end{aligned}$$

where

$$(3.59) \quad \mathcal{A}_3 = \left[\frac{i(n^2 - 3n + 2)\xi_n^2 - i(n-2)}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i};$$

$$(3.60) \quad \mathcal{A}_4 = \left[\frac{-(n^2 - 5n + 6)\xi_n^3 + 3(n-2)\xi_n}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i};$$

$$(3.61) \quad \mathcal{A}_5 = \left[\frac{-i(n^2 - 3n + 2)\xi_n^3 - 2(n^2 - 3n + 2)\xi_n^2 + i(n-2)\xi_n + 2(n-2)}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i};$$

$$(3.62) \quad \mathcal{A}_6 = \left[\frac{-i(n^2 - 5n + 6)\xi_n^3 + 3i(n-2)\xi_n}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}.$$

case (a) (III) $r = -1, l = -n + 3, |\alpha| = j = 0, k = 1$.

Using (3.8), we get

$$(3.63) \quad \Phi_3 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx'.$$

We check at once that

$$(3.64) \quad \pi_{\xi_n}^+ \partial_{\xi_n} \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} = -\frac{1}{2(\xi_n - i)^2} \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_{\beta}^i c(dx_{\beta}) - \frac{i}{2(\xi_n - i)^2} \sum_{\beta=1}^n a_{\beta}^n c(dx_{\beta}).$$

Likewise,

$$(3.65) \quad \begin{aligned} \partial_{x_n} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0) &= \frac{i \partial_{x_n}(c[J(\xi)])(x_0)}{|\xi|^{n-2}} - \frac{ic[J(\xi)] \partial_{x_n}(|\xi|^{n-2})(x_0)}{|\xi|^{2n-4}} \\ &= \frac{i \sum_{p,h=1}^n \xi_p \partial_{x_n}(a_h^p) c(dx_h)(x_0)}{|\xi|^{n-2}} + \frac{i \sum_{p,h=1}^n \xi_p a_h^p \partial_{x_n}(c(dx_h))(x_0)}{|\xi|^{n-2}} \\ &\quad - \frac{i(\frac{n}{2} - 1) h'(0) |\xi'|^2 \sum_{p,h=1}^n \xi_p a_h^p c(dx_h)(x_0)}{|\xi|^n}. \end{aligned}$$

Then we have

$$(3.66) \quad \begin{aligned} \partial_{x_n} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0)|_{|\xi'|=1} &= \frac{i \sum_{p,h=1}^n \xi_p \partial_{x_n}(a_h^p) c(dx_h)(x_0)}{(1 + \xi_n^2)^{\frac{n}{2}-1}} + \frac{i \sum_{p,h=1}^n \xi_p a_h^p \partial_{x_n}(c(dx_h))(x_0)}{(1 + \xi_n^2)^{\frac{n}{2}-1}} \\ &\quad - \frac{i(\frac{n}{2} - 1) h'(0) \sum_{p,h=1}^n \xi_p a_h^p c(dx_h)(x_0)}{(1 + \xi_n^2)^{\frac{n}{2}}}. \end{aligned}$$

Thus

$$(3.67) \quad \begin{aligned} \partial_{\xi_n} \partial_{x_n} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0)|_{|\xi'|=1} &= -\frac{i(n-2)\xi_n}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_n}(a_h^p) c(dx_h) \\ &\quad + \frac{i[1 - (n-3)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \partial_{x_n}(a_h^n) c(dx_h) - \frac{i(n-2)\xi_n}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{h,p=1}^{n-1} \xi_p a_h^p \partial_{x_n}(c(dx_h)) \\ &\quad + \frac{i[1 - (n-3)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^{n-1} a_h^n \partial_{x_n}(c(dx_h)) + \frac{i(\frac{n}{2} - 1)n\xi_n}{(1 + \xi_n^2)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p a_h^p c(dx_h) \\ &\quad - \frac{i(\frac{n}{2} - 1)[1 - (n-1)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^n a_h^n c(dx_h). \end{aligned}$$

We see at once that

(3.68)

$$\begin{aligned}
\Phi_3 &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(n-2)\xi_n}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{h,\beta=1}^n \sum_{p,i=1}^{n-1} \text{tr}[\xi_p \xi_i a_\beta^i \partial_{x_n}(a_h^p) c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(n-2)\xi_n}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{\beta=1}^n \sum_{h,p,i=1}^{n-1} \text{tr}[\xi_p \xi_i a_h^p a_\beta^i \partial_{x_n}(c(dx_h)) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} -\frac{i(\frac{n}{2}-1)n\xi_n}{2(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h,\beta=1}^n \sum_{p,i=1}^{n-1} \text{tr}[\xi_p \xi_i a_h^p a_\beta^i c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{1-(n-3)\xi_n^2}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{h,\beta=1}^n \text{tr}[a_\beta^n \partial_{x_n}(a_h^n) c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{1-(n-3)\xi_n^2}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{\beta=1}^n \sum_{h=1}^{n-1} \text{tr}[a_h^n a_\beta^n \partial_{x_n}(c(dx_h)) c(dx_\beta)] d\xi_n \sigma(\xi') dx' \\
&\quad - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} -\frac{(\frac{n}{2}-1)[1-(n-1)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h,\beta=1}^n \text{tr}[a_h^n a_\beta^n c(dx_h) c(dx_\beta)] d\xi_n \sigma(\xi') dx'.
\end{aligned}$$

From this, we have

(3.69)

$$\begin{aligned}
\Phi_3 &= -\frac{1}{2} \int_{-\infty}^{+\infty} -\frac{i(n-2)\xi_n}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_n}(a_h^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} -\frac{i(n-2)\xi_n}{4(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{h,i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} \frac{i(\frac{n}{2}-1)n\xi_n}{2(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} -\frac{1-(n-3)\xi_n^2}{2(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} \sum_{h=1}^n a_h^n \partial_{x_n}(a_h^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} -\frac{1-(n-3)\xi_n^2}{4(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{h=1}^{n-1} (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&\quad - \frac{1}{2} \int_{-\infty}^{+\infty} \frac{(\frac{n}{2}-1)[1-(n-1)\xi_n^2]}{2(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}+1}} h'(0) \sum_{h=1}^n (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'.
\end{aligned}$$

It is immediate that

$$\begin{aligned}
(3.70) \quad \Phi_3 &= h'(0) \sum_{h,i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_7 dx' \\
&+ h'(0) \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{A}_8 dx' \\
&+ h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_9 dx' \\
&+ h'(0) \sum_{h=1}^n (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{A}_{10} dx',
\end{aligned}$$

where

$$(3.71) \quad \mathcal{A}_7 = \left[\frac{i(n-2)\xi_n}{8(\xi_n+i)^{\frac{n}{2}}} \right]^{\left(\frac{n}{2}+1\right)} \Big|_{\xi_n=i};$$

$$(3.72) \quad \mathcal{A}_8 = \left[\frac{-in(n-2)\xi_n}{8(\xi_n+i)^{\frac{n}{2}+1}} \right]^{\left(\frac{n}{2}+2\right)} \Big|_{\xi_n=i};$$

$$(3.73) \quad \mathcal{A}_9 = \left[\frac{-(n-3)\xi_n^2+1}{8(\xi_n+i)^{\frac{n}{2}}} \right]^{\left(\frac{n}{2}+1\right)} \Big|_{\xi_n=i};$$

$$(3.74) \quad \mathcal{A}_{10} = \left[\frac{(n^2-3n+2)\xi_n^2-(n-2)}{8(\xi_n+i)^{\frac{n}{2}+1}} \right]^{\left(\frac{n}{2}+2\right)} \Big|_{\xi_n=i}.$$

In combination with the calculation,

$$\begin{aligned}
(3.75) \quad \Phi_1 + \Phi_2 + \Phi_3 &= \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i} (a_h^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2})!} \mathcal{A}_0 dx' \\
&+ \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i} (a_h^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_1 dx' \\
&+ \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^n \partial_{x_i} (a_h^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_2 dx' \\
&+ h'(0) \sum_{h,i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_{11} dx' \\
&+ h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{A}_{12} dx'
\end{aligned}$$

$$\begin{aligned}
& + h'(0) \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{A}_{13} dx' \\
& + h'(0) \sum_{h=1}^n (a_h^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{A}_{14} dx',
\end{aligned}$$

where

(3.76)

$$\mathcal{A}_{11} = \left[\frac{in(n-2)\xi_n^2 - (n-2)\xi_n - i(n-2)}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}^{\frac{n}{2}+1};$$

(3.77)

$$\mathcal{A}_{12} = \left[\frac{-(n^2-4n+3)\xi_n^3 - i(n-3)\xi_n^2 + (3n-5)\xi_n + i}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}^{\frac{n}{2}+1};$$

(3.78)

$$\mathcal{A}_{13} = \left[\frac{-i(n^2-3n+2)\xi_n^3 - 2(n^2-3n+2)\xi_n^2 - i(n^2-3n+2)\xi_n + 2(n-2)}{8(\xi_n+i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}^{\frac{n}{2}+2};$$

(3.79)

$$\mathcal{A}_{14} = \left[\frac{-i(n^2-5n+6)\xi_n^2 + 2(n-2)\xi_n + i(n-2)}{8(\xi_n+i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\frac{n}{2}+2}.$$

case (b) $r = -1, l = -n + 2, |\alpha| = j = k = 0$.

It is easily seen that

$$\begin{aligned}
(3.80) \quad \Phi_4 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \partial_{\xi_n} \sigma_{-n+2}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_J^{-1}) \times \sigma_{-n+2}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx'.
\end{aligned}$$

In the normal coordinate, $g^{ij}(x_0) = \delta_i^j$ and $\partial_{x_j}(g^{\alpha\beta})(x_0) = 0$, if $j < n$; $\partial_{x_j}(g^{\alpha\beta})(x_0) = h'(0)\delta_\beta^\alpha$, if $j = n$. So by [11], when $k < n$, we have $\Gamma^n(x_0) = \frac{n-1}{2}h'(0)$ and $\Gamma^k(x_0) = 0$. We thus get

(3.81)

$$\begin{aligned}
& \sigma_{-n+2}(D_J^{-n+3})(x_0)|_{|\xi'|=1} \\
&= -\frac{1}{4(1+\xi_n^2)^{\frac{n}{2}-1}} h'(0) \sum_{\mu=1}^n \sum_{\nu=1}^{n-1} a_\nu^\mu c(dx_\mu) c(dx_n) c(dx_\nu) \\
&\quad - \frac{n-2}{4(1+\xi_n^2)^{\frac{n}{2}}} h'(0) \sum_{\omega=1}^n \sum_{\lambda,k=1}^{n-1} \xi_k \xi_\lambda a_\omega^\lambda c(dx_k) c(dx_n) c(dx_\omega)
\end{aligned}$$

$$\begin{aligned}
& - \frac{(n-2)\xi_n}{4(1+\xi_n^2)^{\frac{n}{2}}} h'(0) \sum_{\omega=1}^n \sum_{k=1}^{n-1} \xi_k a_\omega^n c(dx_k) c(dx_n) c(dx_\omega) \\
& + \frac{(n^2-3n+2)\xi_n^3 + (2n^2-5n+2)\xi_n}{4(1+\xi_n^2)^{\frac{n}{2}+1}} h'(0) \sum_{\omega=1}^n \sum_{\lambda=1}^{n-1} \xi_\lambda a_\omega^\lambda c(dx_\omega) \\
& + \frac{(n^2-3n+2)\xi_n^4 + (2n^2-5n+2)\xi_n^2}{4(1+\xi_n^2)^{\frac{n}{2}+1}} h'(0) \sum_{\omega=1}^n a_\omega^n c(dx_\omega) \\
& + \frac{n-2}{2(1+\xi_n^2)^{\frac{n}{2}}} \sum_{\alpha,\beta,\omega=1}^n \sum_{\lambda=1}^{n-1} \xi_\lambda a_\alpha^\beta a_\omega^\lambda c(dx_\beta) c[(\nabla_{e_\alpha}^L J)(\xi^*)] c(dx_\omega) \\
& + \frac{(n-2)\xi_n}{2(1+\xi_n^2)^{\frac{n}{2}}} \sum_{\alpha,\beta,\omega=1}^n a_\alpha^\beta a_\omega^n c(dx_\beta) c[(\nabla_{e_\alpha}^L J)(\xi^*)] c(dx_\omega) \\
& - \frac{n-2}{(1+\xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{j,p=1}^{n-1} \xi_j \xi_p \partial_{x_j}(a_h^p) c(dx_h) \\
& - \frac{(n-2)\xi_n}{(1+\xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_n}(a_h^p) c(dx_h) \\
& - \frac{(n-2)\xi_n}{(1+\xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \sum_{j=1}^{n-1} \xi_j \partial_{x_j}(a_h^n) c(dx_h) \\
& - \frac{(n-2)\xi_n^2}{(1+\xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^n \partial_{x_n}(a_h^n) c(dx_h) \\
& - \frac{(n-2)\xi_n}{(1+\xi_n^2)^{\frac{n}{2}}} \sum_{p,h=1}^{n-1} \xi_p a_h^p \partial_{x_n}(c(dx_h)) \\
& - \frac{(n-2)\xi_n^2}{(1+\xi_n^2)^{\frac{n}{2}}} \sum_{h=1}^{n-1} a_h^n \partial_{x_n}(c(dx_h)).
\end{aligned}$$

Summarizing, we have

$$\begin{aligned}
(3.82) \quad \Phi_4 & = h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{B}_0 dx' \\
& + h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2})!} \mathcal{B}_1 dx' \\
& + h'(0) \sum_{i=1}^{n-1} a_i^i a_n^n \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \cdot -\mathcal{B}_0 dx'
\end{aligned}$$

$$\begin{aligned}
& + h'(0) \sum_{i=1}^{n-1} a_i^i a_n^n \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2}\right)!} \cdot -\mathcal{B}_1 dx' \\
& + h'(0) \sum_{l=1}^n (a_n^l)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 2\right)!} \mathcal{B}_2 dx' \\
& + h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \mathcal{B}_3 dx' \\
& + h'(0) \sum_{l=1}^n \sum_{i=1}^{n-1} (a_i^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 2\right)!} \mathcal{B}_4 dx' \\
& + h'(0) \sum_{i,j=1}^{n-1} (a_j^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \mathcal{B}_5 dx' \\
& + \sum_{l,\alpha=1}^n \sum_{i=1}^{n-1} (a_i^i)^2 g^M(J(dx_\alpha), (\nabla_{e_\alpha}^L J)e_n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \mathcal{B}_5 dx' \\
& + \sum_{i=1}^{n-1} g^M(J(dx_i), (\nabla_{e_n}^L J)e_i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \cdot -2\mathcal{B}_0 dx' \\
& + \sum_{i=1}^{n-1} g^M(J(dx_n), (\nabla_{e_i}^L J)e_i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \cdot 2\mathcal{B}_0 dx' \\
& + \sum_{l,\alpha=1}^n (a_l^n)^2 g^M(J(dx_\alpha), (\nabla_{e_\alpha}^L J)e_n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \mathcal{B}_3 dx' \\
& + \sum_{l=1}^n \sum_{i=1}^{n-1} a_i^i \partial_{x_i} (a_l^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{\left(\frac{n}{2} + 1\right)!} \cdot 4\mathcal{B}_0 dx',
\end{aligned}$$

where

$$(3.83) \quad \mathcal{B}_0 = \left[-\frac{i(n-2)\xi_n + n-2}{8(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}+1\right)} ;$$

$$(3.84) \quad \mathcal{B}_1 = \left[-\frac{1}{8(\xi_n + i)^{\frac{n}{2}-1}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}\right)} ;$$

$$(3.85) \quad \mathcal{B}_2 = \left[-\frac{(n^2 - 3n + 2)\xi_n^4 + (2n^2 - 5n + 2)\xi_n^2}{8(\xi_n + i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}+2\right)} ;$$

$$(3.86) \quad \mathcal{B}_3 = \left[\frac{(n-2)\xi_n^2}{4(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}+1\right)} ;$$

$$(3.87) \quad \mathcal{B}_4 = \left[\frac{i(n^2 - 3n + 2)\xi_n^3 + i(2n^2 - 5n + 2)\xi_n}{8(\xi_n + i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}+2\right)} ;$$

$$(3.88) \quad \mathcal{B}_5 = \left[-\frac{i(n-2)\xi_n}{4(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\left(\frac{n}{2}+1\right)} .$$

case (c) $r = -2, l = -n + 3, |\alpha| = j = k = 0$.

We calculate

$$(3.89) \quad \Phi_5 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-2}(D_J^{-1}) \times \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx'.$$

We follow the notation of [21].

$$(3.90)$$

$$A_1(x_0) = \frac{c[J(\xi)]\sigma_0(D_J)(x_0)c[J(\xi)]}{(1 + \xi_n^2)^2};$$

$$(3.91)$$

$$A_2(x_0) = \frac{c[J(\xi)]}{(1 + \xi_n^2)^2} \left[\sum_{j,p,h=1}^n \xi_p \partial_{x_j} (a_h^p) c[J(dx_j)] c(dx_h) + \sum_{p=1}^n \sum_{h=1}^{n-1} \xi_p a_h^p c[J(dx_n)] \partial_{x_n} (c(dx_h)) \right];$$

$$(3.92)$$

$$A_3(x_0) = \frac{c[J(\xi)]}{(1 + \xi_n^2)^3} c[J(dx_n)] c[J(\xi)],$$

means that

$$(3.93) \quad \pi_{\xi_n}^+ \sigma_{-2}(D_J^{-1})(x_0)|_{|\xi'|=1} = \pi_{\xi_n}^+(A_1(x_0)) + \pi_{\xi_n}^+(A_2(x_0)) - h'(0)\pi_{\xi_n}^+(A_3(x_0)).$$

Computations show that

$$(3.94)$$

$$\begin{aligned} \pi_{\xi_n}^+(A_1(x_0)) &= \frac{i\xi_n}{16(\xi_n - i)^2} h'(0) \sum_{l,\gamma,\mu=1}^n \sum_{\nu=1}^{n-1} a_l^n a_\gamma^n a_\nu^\mu c(dx_l) c(dx_\mu) c(dx_n) c(dx_\nu) c(dx_\gamma) \\ &+ \frac{i}{16(\xi_n - i)^2} h'(0) \sum_{l,\gamma,\mu=1}^n \sum_{q,\nu=1}^{n-1} \xi_q a_l^q a_\gamma^n a_\nu^\mu c(dx_l) c(dx_\mu) c(dx_n) c(dx_\nu) c(dx_\gamma) \\ &+ \frac{i}{16(\xi_n - i)^2} h'(0) \sum_{l,\gamma,\mu=1}^n \sum_{\alpha,\nu=1}^{n-1} \xi_\alpha a_l^n a_\gamma^\alpha a_\nu^\mu c(dx_l) c(dx_\mu) c(dx_n) c(dx_\nu) c(dx_\gamma) \\ &+ \frac{i\xi_n + 2}{16(\xi_n - i)^2} h'(0) \sum_{l,\gamma,\mu=1}^n \sum_{q,\alpha,\nu=1}^{n-1} \xi_q \xi_\alpha a_l^q a_\gamma^\alpha a_\nu^\mu c(dx_l) c(dx_\mu) c(dx_n) c(dx_\nu) c(dx_\gamma). \end{aligned}$$

By computation, we have

$$\begin{aligned}
(3.95) \quad \partial_{\xi_n} \left(\frac{ic[J(\xi)]}{|\xi|^{n-2}} \right) (x_0)|_{|\xi'|=1} &= i \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_\beta^i c(dx_\beta) \partial_{\xi_n} \left(\frac{1}{(1 + \xi_n^2)^{\frac{n-1}{2}}} \right) \\
&+ i \sum_{\beta=1}^n a_\beta^n c(dx_\beta) \partial_{\xi_n} \left(\frac{\xi_n}{(1 + \xi_n^2)^{\frac{n-1}{2}}} \right) \\
&= -\frac{i(n-2)\xi_n}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_\beta^i c(dx_\beta) \\
&+ \frac{i[1 - (n-3)\xi_n^2]}{(1 + \xi_n^2)^{\frac{n}{2}}} \sum_{\beta=1}^n a_\beta^n c(dx_\beta).
\end{aligned}$$

By $\int_{|\xi'|=1} \{\xi_{i_1} \cdots \xi_{i_{2d+1}}\} \sigma(\xi') = 0$ and $\int_{|\xi'|=1} \sum_{i,j=1}^{n-1} \xi_i \xi_j \sigma(\xi') = \frac{1}{n-1} \sum_{i,j=1}^{n-1} \delta_j^i \text{Vol}(S^{n-2})$, then we have

$$\begin{aligned}
(3.96) \quad &-i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+(A_1(x_0)) \times \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= -i \int_{-\infty}^{+\infty} \frac{(n-3)\xi_n^3 - \xi_n}{16(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} h'(0) \\
&\cdot \sum_{l=1}^n \sum_{\nu=1}^{n-1} (-(a_\nu^n)^2 (a_l^n)^2 + (a_l^n)^2 a_\nu^\nu a_n^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&-i \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n}{16(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} h'(0) \\
&\cdot \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (-(a_\nu^n)^2 (a_i^i)^2 + (a_i^i)^2 a_\nu^\nu a_n^n) \cdot \frac{2}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
&-i \int_{-\infty}^{+\infty} \frac{(n-3)\xi_n^3 - 2i(n-3)\xi_n^2 - \xi_n + 2i}{16(\xi_n - i)^{\frac{n}{2}+2}(\xi_n + i)^{\frac{n}{2}}} h'(0) \\
&\cdot \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} ((a_\nu^n)^2 (a_i^i)^2 - (a_i^i)^2 a_\nu^\nu a_n^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'.
\end{aligned}$$

[21] also shown that

$$\begin{aligned}
(3.97) \quad \pi_{\xi_n}^+(A_2(x_0)) &= -\frac{i\xi_n}{4(\xi_n - i)^2} \sum_{l,j,h,y=1}^n a_l^n a_y^j \partial_{x_j} (a_h^n) c(dx_l) c(dx_y) c(dx_h) \\
&- \frac{i}{4(\xi_n - i)^2} \sum_{l,j,h,y=1}^n \sum_{q=1}^{n-1} \xi_q a_l^q a_y^j \partial_{x_j} (a_h^n) c(dx_l) c(dx_y) c(dx_h)
\end{aligned}$$

$$\begin{aligned}
& - \frac{i}{4(\xi_n - i)^2} \sum_{l,j,h,y=1}^n \sum_{p=1}^{n-1} \xi_p a_l^n a_y^j \partial_{x_j} (a_h^p) c(dx_l) c(dx_y) c(dx_h) \\
& - \frac{i\xi_n + 2}{4(\xi_n - i)^2} \sum_{l,j,h,y=1}^n \sum_{q,p=1}^{n-1} \xi_q \xi_p a_l^q a_y^j \partial_{x_j} (a_h^p) c(dx_l) c(dx_y) c(dx_h) \\
& - \frac{i\xi_n}{8(\xi_n - i)^2} h'(0) \sum_{l,z=1}^n \sum_{h=1}^{n-1} a_l^n a_h^n a_z^n c(dx_l) c(dx_z) c(dx_h) \\
& - \frac{i}{8(\xi_n - i)^2} h'(0) \sum_{l,z=1}^n \sum_{q,h=1}^{n-1} \xi_q a_l^q a_h^n a_z^n c(dx_l) c(dx_z) c(dx_h) \\
& - \frac{i}{8(\xi_n - i)^2} h'(0) \sum_{l,z=1}^n \sum_{p,h=1}^{n-1} \xi_p a_l^n a_h^p a_z^n c(dx_l) c(dx_z) c(dx_h) \\
& - \frac{i\xi_n + 2}{8(\xi_n - i)^2} h'(0) \sum_{l,z=1}^n \sum_{q,p,h=1}^{n-1} \xi_q \xi_p a_l^q a_h^p a_z^n c(dx_l) c(dx_z) c(dx_h).
\end{aligned}$$

Accordingly, we have

(3.98)

$$\begin{aligned}
& - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+(A_2(x_0)) \times \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
& = - i \int_{-\infty}^{+\infty} \frac{-(n-3)\xi_n^3 + \xi_n}{4(\xi_n - i)^{\frac{n}{2}+2} (\xi_n + i)^{\frac{n}{2}}} \sum_{j,\beta,\eta=1}^n (a_\beta^n)^2 a_\eta^j \partial_{x_j} (a_\eta^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& - i \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n}{4(\xi_n - i)^{\frac{n}{2}+2} (\xi_n + i)^{\frac{n}{2}}} \sum_{j,\beta,\eta=1}^n \sum_{i=1}^{n-1} (a_\beta^n)^2 a_\eta^j \partial_{x_j} (a_\eta^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& - i \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n}{4(\xi_n - i)^{\frac{n}{2}+2} (\xi_n + i)^{\frac{n}{2}}} \sum_{l=1}^n \sum_{i=1}^{n-1} a_l^n \partial_{x_i} (a_l^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& - i \int_{-\infty}^{+\infty} \frac{(n-3)\xi_n^3 - 2i(n-3)\xi_n^2 - \xi_n + 2i}{4(\xi_n - i)^{\frac{n}{2}+2} (\xi_n + i)^{\frac{n}{2}}} \sum_{l=1}^n \sum_{i=1}^{n-1} a_l^n \partial_{x_i} (a_l^i) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& - i \int_{-\infty}^{+\infty} \frac{-(n-3)\xi_n^3 + \xi_n}{8(\xi_n - i)^{\frac{n}{2}+2} (\xi_n + i)^{\frac{n}{2}}} h'(0) \sum_{l=1}^n \sum_{\nu=1}^{n-1} (a_\nu^n)^2 (a_l^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& - i \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n}{8(\xi_n - i)^{\frac{n}{2}+2} (\xi_n + i)^{\frac{n}{2}}} h'(0) \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'
\end{aligned}$$

$$\begin{aligned}
& -i \int_{-\infty}^{+\infty} -\frac{(n-2)\xi_n}{8(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& -i \int_{-\infty}^{+\infty} \frac{(n-3)\xi_n^3 - 2i(n-3)\xi_n^2 - \xi_n + 2i}{8(\xi_n-i)^{\frac{n}{2}+2}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'.
\end{aligned}$$

Since

(3.99)

$$\begin{aligned}
-h'(0)\pi_{\xi_n}^+(A_3(x_0)) &= \frac{3\xi_n + i\xi_n^2}{16(\xi_n - i)^3} h'(0) \sum_{l,w,\gamma=1}^n a_l^n a_w^n a_\gamma^n c(dx_l) c(dx_w) c(dx_\gamma) \\
&+ \frac{i\xi_n + 3}{16(\xi_n - i)^3} h'(0) \sum_{l,w,\gamma=1}^n \sum_{q=1}^{n-1} \xi_q a_l^q a_w^n a_\gamma^n c(dx_l) c(dx_w) c(dx_\gamma) \\
&+ \frac{i\xi_n + 3}{16(\xi_n - i)^3} h'(0) \sum_{l,w,\gamma=1}^n \sum_{\alpha=1}^{n-1} \xi_\alpha a_l^n a_w^n a_\gamma^\alpha c(dx_l) c(dx_w) c(dx_\gamma) \\
&+ \frac{-8i + 9\xi_n + 3i\xi_n^2}{16(\xi_n - i)^3} h'(0) \sum_{l,w,\gamma=1}^n \sum_{q,\alpha=1}^{n-1} \xi_q \xi_\alpha a_l^q a_w^n a_\gamma^\alpha c(dx_l) c(dx_w) c(dx_\gamma),
\end{aligned}$$

it is sufficient to show that

(3.100)

$$\begin{aligned}
& -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[-h'(0)\pi_{\xi_n}^+(A_3(x_0)) \times \partial_{\xi_n} \sigma_{-n+3}(D_J^{-n+3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= -i \int_{-\infty}^{+\infty} \frac{(n-3)\xi_n^4 - 3i(n-3)\xi_n^3 - \xi_n^2 + 3i\xi_n}{16(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{\beta,l=1}^n (a_\beta^n)^2 (a_l^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& -i \int_{-\infty}^{+\infty} \frac{(n-2)\xi_n^2 - 3i(n-2)\xi_n}{16(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{\beta,\omega=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 (a_\omega^n)^2 \cdot \frac{2}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx' \\
& -i \int_{-\infty}^{+\infty} -\frac{3(n-3)\xi_n^4 - 9i(n-3)\xi_n^3 - (8n-21)\xi_n^2 + 9i\xi_n + 8}{16(\xi_n-i)^{\frac{n}{2}+3}(\xi_n+i)^{\frac{n}{2}}} h'(0) \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 (a_\beta^n)^2 \\
& \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] d\xi_n dx'.
\end{aligned}$$

It follows immediately that

$$(3.101) \quad \sum_{l=1}^n \sum_{i=1}^{n-1} a_l^n \partial_{x_i} (a_l^i) = \sum_{l,i=1}^n a_l^n \partial_{x_i} (a_l^i) - \sum_{l=1}^n a_l^n \partial_{x_n} (a_l^n)$$

$$= \sum_{l,i=1}^n a_l^n \partial_{x_i}(a_l^i) - \partial_{x_n}(\delta_n^n) = \sum_{l,i=1}^n a_l^n \partial_{x_i}(a_l^i).$$

On account of the above result,

$$\begin{aligned}
(3.102) \quad \Phi_5 &= \sum_{j,\beta,\eta=1}^n (a_\beta^n)^2 a_\eta^j \partial_{x_j}(a_\eta^n) \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{C}_0 dx' \\
&+ \sum_{j,\beta,\eta=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 a_\eta^j \partial_{x_j}(a_\eta^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{C}_1 dx' \\
&+ h'(0) \sum_{l=1}^n \sum_{\nu=1}^{n-1} (a_l^n)^2 a_\nu^n \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \cdot -\frac{1}{4} \mathcal{C}_0 dx' \\
&+ h'(0) \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_l^i)^2 a_\nu^n \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{C}_2 dx' \\
&+ \sum_{l,j=1}^n a_l^j \partial_{x_j}(a_l^n) \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{C}_3 dx' \\
&+ h'(0) \sum_{\beta,l=1}^n (a_\beta^n)^2 (a_l^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{C}_4 dx' \\
&+ h'(0) \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_\beta^n)^2 (a_l^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+2)!} \mathcal{C}_5 dx' \\
&+ h'(0) \sum_{l=1}^n \sum_{\nu=1}^{n-1} (a_\nu^n)^2 (a_l^n)^2 \cdot \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \cdot \frac{3}{4} \mathcal{C}_0 dx' \\
&+ h'(0) \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \mathcal{C}_6 dx' \\
&+ h'(0) \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \cdot \frac{1}{n-1} \text{Vol}(S^{n-2}) \text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2}+1)!} \cdot \frac{1}{2} \mathcal{C}_3 dx',
\end{aligned}$$

where

$$(3.103) \quad \mathcal{C}_0 = \left[\frac{i(n-3)\xi_n^3 - i\xi_n}{4(\xi_n+i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\frac{(n+1)}{2}};$$

$$(3.104) \quad \mathcal{C}_1 = \left[\frac{i(n-2)\xi_n}{4(\xi_n+i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\frac{(n+1)}{2}};$$

$$(3.105) \quad \mathcal{C}_2 = \left[\frac{i(n-3)\xi_n^3 + 2(n-3)\xi_n^2 - i(2n-3)\xi_n - 2}{16(\xi_n+i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{\frac{(n+1)}{2}};$$

$$(3.106) \quad \mathcal{C}_3 = \left[\frac{-i(n-3)\xi_n^3 - 2(n-3)\xi_n^2 + i(n-1)\xi_n + 2}{4(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{(\frac{n}{2}+1)} ;$$

$$(3.107) \quad \mathcal{C}_4 = \left[\frac{-i(n-3)\xi_n^4 - 3(n-3)\xi_n^3 + i\xi_n^2 + 3\xi_n}{16(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{(\frac{n}{2}+2)} ;$$

$$(3.108) \quad \mathcal{C}_5 = \left[\frac{3i(n-3)\xi_n^4 + 9(n-3)\xi_n^3 - 5i(2n-5)\xi_n^2 - 3(2n-1)\xi_n + 8i}{16(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{(\frac{n}{2}+2)} ;$$

$$(3.109) \quad \mathcal{C}_6 = \left[\frac{-i(n-3)\xi_n^3 - 2(n-3)\xi_n^2 - i(-4n+7)\xi_n + 2}{16(\xi_n + i)^{\frac{n}{2}}} \right] \Big|_{\xi_n=i}^{(\frac{n}{2}+1)} .$$

A simple calculation shows that

$$(3.110) \quad \begin{aligned} \sum_{i=1}^{n-1} g^M(J(dx_i), (\nabla_{e_n}^L J)e_i)(x_0) &= \sum_{i=1}^{n-1} g^M(J(e_i), (\nabla_{e_n}^L J)e_i)(x_0) \\ &= \sum_{i=1}^{n-1} g^M(J(e_i), \nabla_{e_n}^L J(e_i) - J(\nabla_{e_n}^L e_i))(x_0) \\ &= \sum_{i=1}^{n-1} g^M(J(e_i), \nabla_{e_n}^L J(e_i))(x_0) - \sum_{i=1}^{n-1} g^M(J(e_i), J(\nabla_{e_n}^L e_i))(x_0) \\ &= \sum_{i=1}^{n-1} g^M(J(e_i), \nabla_{e_n}^L J(e_i))(x_0) - \sum_{i=1}^{n-1} g^M(e_i, \nabla_{e_n}^L e_i)(x_0) \\ &= \frac{1}{2} \sum_{i=1}^{n-1} e_n(g^M(J(e_i), J(e_i)))(x_0) - \frac{1}{2} \sum_{i=1}^{n-1} e_n(g^M(e_i, e_i))(x_0) = 0. \end{aligned}$$

Since

$$(3.111) \quad \begin{aligned} &\sum_{i=1}^{n-1} g^M(J(dx_n), (\nabla_{e_i}^L J)e_i)(x_0) + \sum_{i=1}^{n-1} g^M(J(dx_i), (\nabla_{e_i}^L J)e_n)(x_0) \\ &= \sum_{i=1}^{n-1} g^M(J(e_n), \nabla_{e_i}^L J(e_i))(x_0) - \sum_{i=1}^{n-1} g^M(J(e_n), J(\nabla_{e_i}^L e_i))(x_0) \\ &+ \sum_{i=1}^{n-1} g^M(J(e_i), \nabla_{e_i}^L J(e_n))(x_0) - \sum_{i=1}^{n-1} g^M(J(e_i), J(\nabla_{e_i}^L e_n))(x_0) = 0, \end{aligned}$$

we get

$$(3.112) \quad \sum_{i=1}^{n-1} g^M(J(dx_n), (\nabla_{e_i}^L J)e_i)(x_0) = - \sum_{i=1}^{n-1} g^M(J(e_i), (\nabla_{e_i}^L J)e_n)(x_0).$$

Similarly, we have

$$(3.113) \quad \sum_{\alpha=1}^n g^M(J(dx_\alpha), (\nabla_{e_\alpha}^L J)e_n)(x_0) = \sum_{\alpha=1}^{n-1} g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_n)(x_0).$$

In summary,

$$(3.114) \quad \begin{aligned} \Phi &= \Phi_1 + \Phi_2 + \Phi_3 + \Phi_4 + \Phi_5 \\ &= h'(0) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 2)!} \mathcal{D}_0 dx' \\ &\quad + \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2})!} \mathcal{D}_1 dx' \\ &\quad + \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_i}(a_h^n) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 1)!} \mathcal{D}_2 dx' \\ &\quad + h'(0) \sum_{\nu, i=1}^{n-1} (a_\nu^i)^2 \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 1)!} \mathcal{D}_3 dx' \\ &\quad + h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2})!} \mathcal{D}_4 dx' \\ &\quad + h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 1)!} \mathcal{D}_5 dx' \\ &\quad + h'(0) \sum_{i=1}^{n-1} a_i^i a_n^n \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2})!} \cdot -\mathcal{D}_4 dx' \\ &\quad + h'(0) \sum_{i=1}^{n-1} a_i^i a_n^n \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 1)!} \mathcal{D}_6 dx' \\ &\quad - \sum_{i=1}^{n-1} g^M(J(e_i), (\nabla_{e_i}^L J)e_n) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 1)!} \cdot \mathcal{D}_7 dx' \\ &\quad + \sum_{i=1}^{n-1} g^M(J(e_i), (\nabla_{e_i}^L J)e_n) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{2\pi i}{(\frac{n}{2} + 1)!} \cdot \mathcal{D}_8 dx', \end{aligned}$$

where

$$(3.115) \quad \mathcal{D}_0 = \left[\frac{i(n-3)\xi_n^5 - (n^2 - 5n + 8)\xi_n^4 - i(n^2 - 3n + 2)\xi_n^3 - (3n^2 - 10n + 14)\xi_n^2}{8(\xi_n + i)^{\frac{n}{2}+1}} + \frac{i(n^2 - 2n + 1)\xi_n + n - 6}{8(\xi_n + i)^{\frac{n}{2}+1}} \right] \Big|_{\xi_n=i}^{\frac{n}{2}+2};$$

(3.116)

$$\mathcal{D}_1 = \left[\frac{-i(n-3)\xi_n^2 + i}{2(\xi_n + i)^{\frac{n}{2}}} \right]^{\binom{n}{2}} \Big|_{\xi_n=i};$$

(3.117)

$$\mathcal{D}_2 = \left[\frac{in(n-3)\xi_n^3 + 2(n-3)\xi_n^2 + i(n^2 - 7n + 8)\xi_n - 2n + 2}{4(n-1)(\xi_n + i)^{\frac{n}{2}}} \right]^{\binom{n}{2}+1} \Big|_{\xi_n=i};$$

(3.118)

$$\mathcal{D}_3 = \left[\frac{-i(n-3)\xi_n^4 - (n-3)\xi_n^3 + i(n^2 - 5n + 9)\xi_n^2 + \xi_n - in + 4i}{8(n-1)(\xi_n + i)^{\frac{n}{2}+1}} \right]^{\binom{n}{2}+1} \Big|_{\xi_n=i};$$

(3.119)

$$\mathcal{D}_4 = \left[-\frac{1}{8(\xi_n + i)^{\frac{n}{2}-1}} \right]^{\binom{n}{2}} \Big|_{\xi_n=i};$$

(3.120)

$$\mathcal{D}_5 = \left[\frac{i(n^2 - 4n + 3)\xi_n^4 - (n^3 - 5n^2 + 5n - 1)\xi_n^3 + i(2n^2 - 6n + 5)\xi_n^2 + (n^2 - 1)\xi_n + in}{8(n-1)(\xi_n + i)^{\frac{n}{2}+1}} \right]^{\binom{n}{2}+1} \Big|_{\xi_n=i};$$

(3.121)

$$\mathcal{D}_6 = \left[\frac{(n^2 - 4n + 3)\xi_n^2 - i(n^2 - 4n + 4)\xi_n - 1}{8(n-1)(\xi_n + i)^{\frac{n}{2}}} \right]^{\binom{n}{2}+1} \Big|_{\xi_n=i};$$

(3.122)

$$\mathcal{D}_7 = \left[-\frac{i(n-2)\xi_n + n - 2}{4(n-1)(\xi_n + i)^{\frac{n}{2}}} \right]^{\binom{n}{2}+1} \Big|_{\xi_n=i};$$

(3.123)

$$\mathcal{D}_8 = \left[\frac{(n^2 - 3n + 2)\xi_n^2 - 5i(n-2)\xi_n}{4(n-1)(\xi_n + i)^{\frac{n}{2}}} \right]^{\binom{n}{2}+1} \Big|_{\xi_n=i}.$$

Through a complex series of calculations, we obtain

(3.124)

$$\begin{aligned} \Phi &= -h'(0) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \pi n(n^4 - 5n^3 - 16n^2 + 68n - 48)2^{-4-n} \cdot \frac{(n-3)!}{\left(\frac{n}{2}+2\right)!\left(\frac{n}{2}\right)!} dx' \\ &+ h'(0) \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \pi n(n^2 - 8n + 12)2^{-3-n} \cdot \frac{(n-3)!}{\left(\frac{n}{2}+1\right)!\left(\frac{n}{2}\right)!} dx' \\ &+ h'(0) \sum_{i=1}^{n-1} (a_n^i)^2 \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \frac{\pi(-2n^2 + 7n - 2)}{n+2} 2^{-n} \cdot \frac{(n-3)!}{\left(\frac{n}{2}\right)!\left(\frac{n}{2}-2\right)!} dx' \\ &- \sum_{i=1}^{n-1} g^M(J(e_i), (\nabla_{e_i}^L J)e_n) \cdot Vol(S^{n-2})\text{tr}[\text{id}] \cdot \pi(n^2 - 8n + 12)2^{-1-n} \cdot \frac{(n-2)!}{\left(\frac{n}{2}+1\right)!\left(\frac{n}{2}-1\right)!} dx'. \end{aligned}$$

Recall the Einstein-Hilbert action for manifolds with boundary (see [12] for more details)

$$(3.125) \quad I_{G_r} = \frac{1}{16\pi} \int_M sd\text{Vol}_M + 2 \int_{\partial M} K d\text{Vol}_{\partial M} := I_{G_{r,i}} + I_{G_{r,b}},$$

where

$$(3.126) \quad K = \sum_{1 \leq i, j \leq n-1} K_{ij} g_{\partial M}^{ij}; \quad K_{ij} = -\Gamma_{ij}^n,$$

and K_{ij} is the second fundamental form, or extrinsic curvature. Take the metric of (2.1), then by [11], $K_{ij}(x_0) = -\Gamma_{ij}^n(x_0)$, when $i = j < n$, otherwise it is zero. Then

$$(3.127) \quad \begin{aligned} K(x_0) &= \sum_{i, j < n-1} K_{ij}(x_0) g_{\partial M}^{ij}(x_0) \\ &= \sum_{i=1}^{n-1} K_{ii}(x_0) = -\frac{n-1}{2} h'(0). \end{aligned}$$

A trivial verification shows that

$$(3.128) \quad \begin{aligned} \sum_{\nu, i=1}^{n-1} (a_\nu^i)^2 &= \sum_{\nu, i=1}^n (a_\nu^i)^2 - \sum_{\nu=1}^{n-1} (a_\nu^n)^2 - \sum_{i=1}^{n-1} (a_n^i)^2 - (a_n^n)^2 \\ &= \sum_{i=1}^n \delta_i^i - 2 \sum_{i=1}^{n-1} \langle J(e_i), e_n \rangle^2 - \langle J(e_n), e_n \rangle^2 \\ &= n - 2 \sum_{i=1}^n \langle J(e_i), e_n \rangle^2 + \langle J(e_n), e_n \rangle^2 \end{aligned}$$

and

$$\sum_{i=1}^{n-1} (a_n^i)^2 = \sum_{i=1}^n (a_n^i)^2 - (a_n^n)^2 = 1 - \langle J(e_n), e_n \rangle^2.$$

Hence we have proved the Theorem 1.1.

4. DECLARATIONS

Ethics approval and consent to participate No applicable.

Consent for publication No applicable.

Availability of data and material The authors confirm that the data supporting the findings of this study are available within the article.

Competing interests The authors declare no conflict of interest.

Funding This research was funded by National Natural Science Foundation of China: No.11771070.

Authors' contributions All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Siyao Liu and Yong Wang. The first draft of the manuscript was written by Siyao Liu and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

REFERENCES

- [1] Guillemin V W. A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. *Adv. Math.* 1985, 55(2): 131-160.
- [2] Wodzicki M. Local invariants of spectral asymmetry. *Invent. Math.* 1995, 75(1): 143-178.
- [3] Connes A. Quantized calculus and applications. 11th International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 1995, 15-36.
- [4] Connes A. The action functinal in noncommutative geometry. *Comm. Math. Phys.* 1998, 117: 673-683.
- [5] Kastler D. The Dirac operator and gravitation. *Comm. Math. Phys.* 1995, 166: 633-643.
- [6] Kalau W, Walze M. Gravity, noncommutative geometry and the Wodzicki residue. *J. Geom. Physics.* 1995, 16: 327-344.
- [7] Ackermann T. A note on the Wodzicki residue. *J. Geom.Phys.* 1996, 20: 404-406.
- [8] Ugalde W J. Differential forms and the Wodzicki residue. *J. Geom. Phys.* 2008, 58: 1739-1751.
- [9] Wang Y. Differential forms and the Wodzicki residue for manifolds with boundary. *J. Geom. Physics.* 2006, 56: 731-753.
- [10] Wang Y. Differential forms and the noncommutative residue for manifolds with boundary in the non-product case. *Lett. Math. Phys.* 2006, 77: 41-51.
- [11] Wang Y. Gravity and the noncommutative residue for manifolds with boundary. *Lett. Math. Phys.* 2007, 80: 37-56.
- [12] Wang Y. Lower-dimensional volumes and Kastler-Kalau-Walze type theorem for manifolds with boundary. *Commun. Theor. Phys.* 2010, 54: 38-42.
- [13] Wang J, Wang Y. The Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary. *J. Math. Phys.* 2015, 56: 052501.
- [14] Wei S, Wang Y. Modified Novikov operators and the Kastler-Kalau-Walze type theorem for manifolds with boundary. *Adv. Math. Phys.* 2020, 9090656: 1-28.
- [15] Wu T, Wang J, Wang Y. Dirac-witten operators and the Kastler-Kalau-Walze type theorem for manifolds with boundary. *J. Nonlinear Math. Phys.* 2022, 29: 1-40.
- [16] Wang Y. A Kastler-Kalau-Walze type theorem and the spectral action for perturbations of Dirac operators on manifolds with boundary. *Abstr. Appl. Anal.* 2014, 17: 286-299.
- [17] Wu T, Wang Y. The operator $\sqrt{-1}c(V)(d + \delta)$ and the Kastler-Kalau-Walze type theorems. *Adv. Oper. Theory.* 2022, 7(4): 47-53.
- [18] Kim E C. Lower bounds of the Dirac eigenvalues on compact Riemannian spin manifolds with locally product structure. *arXiv:math/0402427*.
- [19] Chen Y. The Dirac operator on locally reducible Riemannian manifolds. *J. Geom. Phys.* 2019, 139: 17-24.

- [20] Chen Y. The Dirac operator on manifold admitting parallel one-form. *J. Geom. Phys.* 2017, 117: 214-221.
- [21] Liu S, Wang Y. A Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator. *J. Nonlinear Math. Phys.* 2023, 30(2): 743-749.
- [22] Liu S, Wang Y. The J -twist D_J of the Dirac operator and the Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary. *Int. J. Geom. Methods M. Phys.* 2023, 20(11): 2350188.
- [23] Wang J, Wang Y. A general Kastler-Kalau-Walze type theorem for manifolds with boundary. *Int. J. Geom. Methods M. Phys.* 2016, 13(1): 1650003.
- [24] Wang Y. General Kastler-Kalau-Walze type theorems for manifolds with boundary II. *Int. J. Geom. Methods M. Phys.* 2019, 16(2): 1950028.
- [25] Schrohe E. Noncommutative residue, Dixmier's trace, and heat trace expansions on manifolds with boundary. *Contemp. Math.* 1999, 242: 161-186.
- [26] Fedosov B V, Golse F, Leichtnam E, Schrohe E. The noncommutative residue for manifolds with boundary. *J. Funct. Anal.* 1996, 142: 1-31.
- [27] Ponge R. Noncommutative geometry and lower dimensional volumes in Riemannian geometry. *Lett. Math. Phys.* 2008, 83(1): 19-32.
- [28] Li H, Wang Y. The general Dabrowski-Sitarz-Zalecki type theorem for odd dimensional manifolds with boundary II. arXiv:2312.00154.

S. Liu

School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China

Email: liusy719@nenu.edu.cn

Y. Wang

School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China

Email: wangy581@nenu.edu.cn