

Dyck Paths Enumerated by the \mathbb{Q} -bonacci Numbers

Elena Barucci Antonio Bernini Stefano Bilotta Renzo Pinzani

Dipartimento di Matematica e Informatica “U. Dini”

Università degli Studi di Firenze
Viale G.B. Morgagni 65, 50134 Firenze, Italy.

`elena.barucci@unifi.it`
`antonio.bernini@unifi.it`
`stefano.bilotta@unifi.it`
`renzo.pinzani@unifi.it`

1 Introduction

The q -generalized Fibonacci numbers [5, 7] can be combinatorially interpreted in many different ways, for $q \in \mathbb{N}^+$. One of them involves the Dyck paths having height at most two. More precisely, the subsets of these bounded Dyck paths avoiding q consecutive valleys at height 1 are enumerated according to their semilength (i.e. the number of steps of the path divided by 2) by the mentioned sequence [1, 2]. The same numbers count the binary strings avoiding $q + 1$ consecutive 1’s, with $q \geq 1$, according to their length [4, 8]. Note that to be exact we should write that the enumerating sequences are the $(q + 1)$ -generalized Fibonacci numbers, for $q \geq 1$.

Not long ago Baril, Kirgizov, and Vajnovszki [3] introduced the set \mathcal{W}_n^q of the q -decreasing strings which are binary strings of length n where each maximal factor $0^a 1^b$, with $a > 0$, satisfies $q \cdot a > b$, for $q \geq 1$. Moreover, among other results, the authors give a bijection between \mathcal{W}_n^q and the set $\mathcal{B}_n(1^{q+1})$ of the binary strings of length n avoiding $q + 1$ consecutive 1’s, for $q \geq 1$.

Recently [6], Kirgizov generalized the q -decreasing strings to the case where q is a positive rational number, $q \in \mathbb{Q}^+$, and they are enumerated by the numbers called \mathbb{Q} -bonacci by the author. In this paper we provide a class of restricted Dyck paths that result in having the same enumeration (according to their semilength). More precisely, we consider the Dyck paths having height at most two and introduce some constraints on the numbers of consecutive valleys at height one which must be followed by a suitable number of valleys at height zero, depending on the value of $q \in \mathbb{Q}^+$.

2 Preliminaries

In the paper we indicate a Dyck path in linear notation as a string over the alphabet $\{U, D\}$, where U and D replace the up and the down steps of the path, respectively. We figure a Dyck path in a Cartesian coordinate system, starting from the origin and ending in a point of the x -axis. A valley is a substring DU , while a peak is a substring UD . The height of a valley is the ordinate reached by the D step. We refer to a valley at height 1 or at height 0 with 1 -valley or 0 -valley, respectively. For the peaks, with 1 -peak (or 0 -peak) we mean a peak UD whose D step reaches the ordinate 1 (or 0).

Since we are going to deal only with Dyck paths having height at most 2, we do not record this restriction in our notation.

Let \mathcal{D}_n^q , with $q \geq 1$, denote the set of the Dyck paths having height at most 2, and avoiding $q + 1$ consecutive 1 -peaks (which is the same as avoiding q consecutive 1 -valleys), and having semilength n ,

where $n \geq 0$.

Clearly, a Dyck path $P \in \mathcal{D}_n^q$ starts with one of the factors $UD, UUDD, UUDUDD, \dots, U(UD)^q D$ (where $(UD)^q$ is the string obtained by concatenating UD to itself q times). Then the path P is obtained by concatenating one of these factors to a path of suitable length, and the set \mathcal{D}_n^q can be generated by

$$\mathcal{D}_n^q = \begin{cases} \varepsilon, & \text{if } n = 0; \\ \bigcup_{j=0}^q U(UD)^j D \cdot \mathcal{D}_{n-1-j}^q, & \text{if } n \geq 1. \end{cases}$$

It is thus enumerated by the sequence of the $(q+1)$ -generalized Fibonacci numbers

$$f_n^{(k)} = \begin{cases} 1, & \text{if } n = 0; \\ \sum_{i=1}^k f_{n-i}^{(k)}, & \text{if } n \geq 1 \quad (f_\ell^{(k)} = 0 \text{ if } \ell < 0). \end{cases}$$

Note that for $q = 1$ we get the classical Fibonacci numbers.

3 Construction in the case $q \in \mathbb{Q}^+$

3.1 The particular case $q = 1/s$

If $P \in \mathcal{D}_n^q$ with $q \in \mathbb{N}^+$, then q consecutive 1-peaks in P are necessarily followed by at least one 0-valley. In the case where $q \in \mathbb{Q}^+$, we require that the number of 0-valleys have some constraints.

We start with the particular value $q = 1/s$. In this case we impose that each 1-peak of a Dyck path P must be followed by at least s consecutive 0-valleys if after the 1-peak there is enough space to contain s consecutive 0-valleys. If a 1-peak occurs near the end of P and there is no space to contain s consecutive 0-valleys, then no other 1-peaks can occur up to the end of P . Summarizing, we give the following definition.

Definition 3.1 Let $\mathcal{D}_n^{1/s}$ denote the set of Dyck paths P of semilength n having height at most 2, where either P has no 1-peaks ($P = (UD)^n$) or each 1-peak in P is followed by at least s consecutive 0-valleys, except the last 1-peak which can be followed by less than s consecutive 0-valleys.

The construction is straightforward: a path $P \in \mathcal{D}_n^{1/s}$ starts with one of the UD or $UUD(DU)^{s-1}D$ factors (or P is a suitable prefix of this last one factor, if $n \leq s$), then P is obtained by concatenating one of these factor to a Dyck path $Q \in \mathcal{D}_{n-1}^{1/s}$ or $Q \in \mathcal{D}_{n-s-1}^{1/s}$. In the case P begins with the longer factor, since Q starts, of course, with an up step U , then really the first 1-peak of P is followed by s consecutive 0-valleys (so that $P \in \mathcal{D}_n^{1/s}$): the first $s-1$ ones are the $s-1$ consecutive 0-valleys of the factor, and the last one is given by the last step D of the factor and the first U step of Q .

Not considering, for the moment, paths with a semilength less or equal to s , we can write

$$\mathcal{D}_n^{1/s} = UD \cdot \mathcal{D}_{n-1}^{1/s} \cup UUD(DU)^{s-1}D \cdot \mathcal{D}_{n-s-1}^{1/s} \quad (\text{for } n > s). \quad (1)$$

Thus, the set $\mathcal{D}_n^{1/s}$ is enumerated by

$$w_n = w_{n-1} + w_{n-s-1}$$

omitting at this stage the initial conditions.

We note that this recurrence relation matches the one enumerating the q -decreasing strings in the case $q = 1/s$ stated in [6].

As far as the initial conditions are concerned, we observe that prepending the factor UD generates Dyck paths having semilength starting from 1, so that the empty Dyck path ε must be considered as a legal path of $\mathcal{D}_0^{1/s}$ (actually, the only one!).

Prepending the factor $UUD(DU)^{s-1}D$ generates Dyck paths having semilength starting from $s+1$. For semilengths less or equal to s , we note that the construction described in equation (1) does not generate the paths $UUD, UUDUD, \dots, UUD(DU)^tD$ with $t = 0, 1, \dots, s-2$. These paths are suitable prefixes of $UUD(DU)^{s-1}D$ which however satisfy Definition 3.1, so that they must be considered among the initial conditions.

Therefore, the generation of the set $\mathcal{D}_n^{1/s}$ can be completely described as follows:

$$\mathcal{D}_n^{1/s} = \begin{cases} \varepsilon, & \text{if } n = 0; \\ UD, & \text{if } n = 1; \\ UD \cdot \mathcal{D}_{n-1}^{1/s} \cup U \cdot p_{n-1}(UD(DU)^{s-1}) \cdot D, & \text{if } 2 \leq n \leq s+1; \\ UD \cdot \mathcal{D}_{n-1}^{1/s} \cup UUD(DU)^{s-1}D \cdot \mathcal{D}_{n-s-1}^{1/s}, & \text{if } n > s+1. \end{cases}$$

In the above formula $p_{n-1}(UD(DU)^{s-1})$ is the prefix of semilength $n-1$ of $UD(DU)^{s-1}$.

It is not difficult to see that $\mathcal{D}_n^{1/s}$ is enumerated by

$$w_n = \begin{cases} 1, & \text{if } n = 0; \\ n, & \text{if } 1 \leq n \leq s+1; \\ w_{n-1} + w_{n-s-1}, & \text{if } n > s+1. \end{cases}$$

This sequence matches the one enumerating the q -decreasing strings for $q = 1/s$ that can be deduced from [6].

3.2 The general case $q = r/s$

Following the outline of the constructions in the cases where q is an integer, and where $q = 1/s$, in the general case $q = r/s$ (we suppose r and s to be coprime) we require that a path P avoid $r+1$ consecutive 1-peaks, and if r consecutive 1-peaks occur in P , then they must be followed by at least s consecutive 0-valleys. Clearly, we have to deal with the case where p consecutive 1-peaks, with $p = 1, 2, \dots, r-1$, occur in P . When this happens, we impose that the p consecutive 1-peaks must be followed by a number v of consecutive 0-valleys such that

$$\frac{p}{v} \leq \frac{r}{s} \quad \text{for } p = 1, 2, \dots, r-1. \quad (2)$$

Moreover, we have to deal with the case where the rightmost block of r consecutive 1-peaks occurs near the end of P (more precisely, when after this block there is no space to contain s consecutive 0-valleys). In this case, no other 1-peaks can occur up to the end of P . Finally, we allow the paths end with a consecutive block of 1-peaks (clearly, less than $r+1$). Summarizing, we give the following definition.

Definition 3.2 Let $\mathcal{D}_n^{r/s}$ denote the set of Dyck paths P of semilength n having height at most 2, where

- each block B of r consecutive 1-peaks in P is followed by at least s consecutive 0-valleys, except the rightmost block B which can be followed by less than s consecutive 0-valleys, and

- each block C of p consecutive 1-peaks in P , with $p = 1, 2, \dots, r-1$, is followed by at least ν consecutive 0-valleys such that $p/\nu \leq r/s$, and
- the path P can end with $(UD)^t D$, with $t = 1, 2, \dots, r$ (in other words, P ends with t consecutive 1-peaks ($t \leq r$) followed by a down step).

Clearly, the number ν of consecutive 0-valleys, since ν is an integer, satisfies our request (2) when $\nu \geq \lceil p \cdot \frac{s}{r} \rceil$. Moreover, we note that in the case where p consecutive 1-peaks, with $p = 1, 2, \dots, r-1$, occur near the end of P and there is not enough space to contain ν consecutive 0-valleys, then the path P does not belong to $\mathcal{D}_n^{r/s}$, according to the second bullet in Definition (3.2). For example, if $q = 4/5$, the path $P = UUDUDDUDUD$ is not allowed, since after two consecutive 1-peaks ($p = 2$) at least three consecutive 0-valleys ($\nu = 3$, according to request (2)) must occur. On the other hand, the path $P = UUDUDUDUDDUDUDUD$ is allowed.

Also in this (general) case, the construction is straightforward. A path $P \in \mathcal{D}_n^{r/s}$ starts with one of the factors UD or $U(UD)^p(DU)^{\lceil ps/r \rceil - 1}D$, with $p = 1, 2, \dots, r$ (or P is a suitable prefix of $U(UD)^r(DU)^{s-1}D$). With an argument similar to the one used in the case where $q = 1/s$, it is not difficult to get the following construction:

$$\mathcal{D}_n^{r/s} = \begin{cases} \varepsilon, & \text{if } n = 0; \\ UD, & \text{if } n = 1; \\ UD \cdot \mathcal{D}_{n-1}^{r/s} \cup U \cdot p_{n-1}((UD)^r(DU)^{s-1}) \cdot D \\ \cup U(UD)^p(DU)^{\lceil ps/r \rceil - 1}D \cdot \mathcal{D}_{n-p-\lceil ps/r \rceil}^{r/s}, & \text{if } 2 \leq n \leq r+s \\ & \text{and } n-p-\lceil ps/r \rceil \geq 1 \\ & \text{with } 1 \leq p \leq r-1; \\ UD \cdot \mathcal{D}_{n-1}^{r/s} \cup U(UD)^p(DU)^{\lceil ps/r \rceil - 1}D \cdot \mathcal{D}_{n-p-\lceil ps/r \rceil}^{r/s}, & \text{if } n > r+s \\ & \text{and } 1 \leq p \leq r; \end{cases}$$

Then we have (in the following $\chi(f) = 1$ if f is true, and $\chi(f) = 0$ otherwise)

$$w_n = \begin{cases} 1, & \text{if } n = 0; \\ 1, & \text{if } n = 1; \\ w_{n-1} + 1 + \sum_{p=1}^{r-1} \chi(n-p-\lceil ps/r \rceil \geq 1) w_{n-p-\lceil ps/r \rceil}, & \text{if } 2 \leq n \leq r+s; \\ w_{n-1} + \sum_{p=1}^r w_{n-p-\lceil ps/r \rceil}, & \text{if } n > r+s. \end{cases}$$

In order to respect the second bullet in Definition 3.2 we added the factor $\chi(n-p-\lceil ps/r \rceil \geq 1)$ in the case $2 \leq n \leq r+s$ of the definition of w_n , and the statement $n-p-\lceil ps/r \rceil \geq 1$ since at least $\lceil ps/r \rceil$ consecutive 0-valleys must occur after p consecutive 1-peaks, with $1 \leq p \leq r-1$ (in order to have $\mathcal{D}_{n-p-\lceil ps/r \rceil}^{r/s} \neq \emptyset$).

Also, in this case, the recurrence relations match the ones enumerating the q -decreasing strings in the case $q = r/s$ founded in [6].

4 Acknowledgements

This work is partially supported by the INdAM – GNCS Project 2023 “Aspetti combinatori ed enumerativi di strutture discrete: stringhe, ipergrafi e permutazioni”, code CUP_E53C22001930001.

References

- [1] E. Barcucci, A. Bernini & R. Pinzani (2022): *From the Fibonacci to Pell numbers and beyond via Dyck paths*. *Pure Math. Appl. (P.U.M.A.)* 30, pp. 17–22, doi:10.2478/puma-2022-0004.
- [2] E. Barcucci, A. Bernini & R. Pinzani (2024): *Sequences from Fibonacci to Catalan: A combinatorial interpretation via Dyck paths*. *RAIRO Theor. Inform. Appl.* 58:8, doi:10.1051/ita/2024007.
- [3] J.-L. Baril, S. Kirgizov & V. Vajnovszki (2022): *Gray codes for Fibonacci q -decreasing words*. *Theoret. Comput. Sci.* 927, pp. 120–132, doi:10.1016/j.tcs.2022.06.003.
- [4] A. Bernini (2017): *Restricted binary strings and generalized Fibonacci numbers*. In A. Dennunzio, E. Formenti, L. Manzoni & A. Porreca, editors: *Cellular Automata and Discrete Complex Systems. AUTOMATA 2017. Lecture Notes in Computer Science*, 10248, Springer International Publishing, Cham, pp. 32–43, doi:10.1007/978-3-319-58631-1_3.
- [5] M. Feinberg (1963): *Fibonacci-Tribonacci*. *Fibonacci Quart.* 1, pp. 71–74.
- [6] S. Kirgizov (2022): *\mathbb{Q} -bonacci numbers and words*. *Fibonacci Quart.* 60(5), pp. 187–195.
- [7] E. P. Miles Jr. (1960): *Generalized Fibonacci numbers and associated matrices*. *Amer. Math. Monthly* 67, pp. 745–752, doi:10.1080/00029890.1960.11989593.
- [8] V Vajnovszki (2001): *A loopless generation of bitstrings without p consecutive ones*. In C. S. Calude, M. J. Dinneen & S. Sburlan, editors: *Combinatorics, Computability and Logic. Discrete Mathematics and Theoretical Computer Science*, Springer, London, pp. 227–240, doi:10.1007/978-1-4471-0717-0_19.