

Bijjective Enumeration and Sign-Imbalance for Permutation Depth and Excedances

Sen-Peng Eu

Department of Mathematics
National Taiwan Normal University
Taiwan, ROC

Chinese Air Force Academy
Taiwan, ROC

speu@math.ntnu.edu.tw

Tung-Shan Fu

Department of Applied Mathematics
National Pingtung University
Taiwan, ROC

tsfu@mail.nptu.edu.tw

Yuan-Hsun Lo

yhlo@mail.nptu.edu.tw

We present a simplified variant of Biane's bijection between permutations and 3-colored Motzkin paths with weight that keeps track of the inversion number, excedance number and a statistic so-called depth of a permutation. This generalizes a result by Guay-Paquet and Petersen about a continued fraction of the generating function for depth on the symmetric group \mathfrak{S}_n of permutations. In terms of weighted Motzkin path, we establish an involution on \mathfrak{S}_n that reverses the parities of depth and excedance numbers simultaneously, which proves that the numbers of permutations with even and odd depth (excedance numbers, respectively) are equal if n is even and differ by the tangent number if n is odd. Moreover, we present some interesting sign-imbalance results on permutations and derangements, refined with respect to depth and excedance numbers.

1 Introduction

1.1 Preliminaries and background

Let \mathfrak{S}_n be the symmetric group of permutations on $[n] := \{1, 2, \dots, n\}$. Given a permutation $\sigma = \sigma(1) \cdots \sigma(n) \in \mathfrak{S}_n$, an *inversion* of σ is a pair (i, j) of indices such that $1 \leq i < j \leq n$ and $\sigma(i) > \sigma(j)$. The *inversion number* of σ , denoted by $\text{inv}(\sigma)$, is defined to be the number of inversions of σ . Petersen and Tenner [11] presented a statistic called *depth* for a Coxeter group, which is defined in terms of factorizations of the elements into reflections. In the case of symmetric group, the reflections of \mathfrak{S}_n are the transpositions (ij) , $1 \leq i < j \leq n$, and the depth of a permutation $\sigma \in \mathfrak{S}_n$, denoted by $\text{depth}(\sigma)$, is defined as

$$\text{depth}(\sigma) = \min \left\{ \sum_{r=1}^k (j_r - i_r) : \sigma = (i_1 j_1)(i_2 j_2) \cdots (i_k j_k) \right\}. \quad (1.1)$$

They obtained a simple formula for the calculation of depth

$$\text{depth}(\sigma) = \sum_{\sigma(i) > i} (\sigma(i) - i), \quad (1.2)$$

which turns out to be one half of the *total displacement* of σ [9, Problem 5.1.1.28] (also called *Spearman's disarray* for \mathfrak{S}_n in [3]).

For any two sequences $\{\gamma_h\}_{h \geq 0}$ and $\{\lambda_h\}_{h \geq 1}$, let $\mathfrak{F}(\gamma_h, \lambda_h)$ denote the series in z defined by

$$\mathfrak{F}(\gamma_h, \lambda_h) = \frac{1}{1 - \gamma_0 z - \frac{\lambda_1 z^2}{1 - \gamma_1 z - \frac{\lambda_2 z^2}{1 - \gamma_2 z - \frac{\lambda_3 z^2}{\ddots}}}} \quad (1.3)$$

These are Jacobi-type continued fractions (also known as *J-fractions*). Flajolet [6] gave a combinatorial interpretation of (1.3) as the generating function for weighted Motzkin paths.

A *Motzkin path* of length n is a lattice path from the origin to the point $(n, 0)$ staying weakly above the x -axis, using the *up step* $(1, 1)$, *down step* $(1, -1)$ and *horizontal step* $(1, 0)$. Let U , D and H denote an up step, a down step and a horizontal step, respectively. Let \mathcal{M}_n denote the set of Motzkin paths of length n .

For a Motzkin path $\mu = x_1 \cdots x_n \in \mathcal{M}_n$, the *height* of each step x_j is the maximum y -coordinate achieved by x_j . For $x \in \{U, D, H\}$ and a nonnegative integer h , let $x^{(h)}$ denote a step x at height h . The *weight* $\omega(\mu)$ of μ is defined to be the product of the weight $\omega(x_j)$ of each step x_j for all $j \in [n]$. By the *area* under μ , denoted by $\text{area}(\mu)$, we mean the area of the region enclosed by μ and the x -axis. Guay-Paquet and Petersen [8] defined a weight of μ by setting

$$\omega(x_i) = \begin{cases} ht^{(2h-1)/2} & \text{if } x_i = U^{(h)} \text{ or } D^{(h)}; \\ (2h+1)t^h & \text{if } x_i = H^{(h)} \end{cases}$$

and established a surjective map $\Phi: \sigma \mapsto \mu$ of \mathfrak{S}_n onto \mathcal{M}_n with $\text{depth}(\sigma) = \text{area}(\mu)$, which yields the following generating function for permutations with respect to depth:

$$\sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} t^{\text{depth}(\sigma)} z^n = \mathfrak{F}(\gamma_h, \lambda_h) \quad (1.4)$$

with $\lambda_h = h^2 t^{2h-1}$ and $\gamma_h = (2h+1)t^h$, i.e.,

$$\sum_{n \geq 0} \sum_{\sigma \in \mathfrak{S}_n} t^{\text{depth}(\sigma)} z^n = \frac{1}{1 - z - \frac{tz^2}{1 - 3tz - \frac{4t^3 z^2}{1 - 5t^2 z - \frac{9t^5 z^2}{1 - 7t^3 z - \frac{16t^7 z^2}{1 - \dots}}}}}$$

Guay-Paquet and Petersen [8] commented that map $\Phi: \mathfrak{S}_n \rightarrow \mathcal{M}_n$ is due to Foata and Zeilberger [7] and that the statistic depth coincides with a statistic, Edif, studied by Clarke, Steingrímsson and Zeng [2].

Following [13], a *3-colored Motzkin path* is a Motzkin path with three kinds of horizontal steps, denoted by H_1 , H_2 and H_3 . Using a double labeling scheme, Biane [1] established a bijection between permutations and 3-colored Motzkin paths with weight, which keeps track of the number of inversions. We refer to an alternative version of this bijection given by Sokal and Zeng in [13, Section 6.2]. Elizalde [5] used this bijection to study the joint distribution of multiple statistics on \mathfrak{S}_n . Using Biane's method, we propose a multivariate weight function on the steps of a 3-colored Motzkin path and turn the surjective map $\Phi: \mathfrak{S}_n \rightarrow \mathcal{M}_n$ of Guay-Paquet and Petersen into a bijection that gives a generalization of (1.4). (This resolves a conjecture raised by Petersen in an unpublished note [10].)

1.2 Main results

Let \mathscr{W}_n denote the set of 3-colored Motzkin paths of length n with a weight function ω defined by setting $\omega(H_3^{(0)}) = p$ and

$$\begin{cases} \omega(U^{(h)}) \in \{st^{2h-1}, st^{2h-1}q, \dots, st^{2h-1}q^{h-1}\}; \\ \omega(D^{(h)}) \in \{q^{2h-1}, q^{2h}, \dots, q^{3h-2}\}; \\ \omega(H_1^{(h)}) \in \{st^h q^h, st^h q^{h+1}, \dots, st^h q^{2h-1}\}; \\ \omega(H_2^{(h)}) \in \{t^h q^h, t^h q^{h+1}, \dots, t^h q^{2h-1}\}; \\ \omega(H_3^{(h)}) = pt^h q^{2h} \end{cases} \quad (1.5)$$

for all $h \geq 1$.

Let $\sigma = \sigma(1) \cdots \sigma(n) \in \mathfrak{S}_n$. We say that an index $i \in [n]$ is an *excedance* if $\sigma(i) > i$ and a *fixed point* if $\sigma(i) = i$. Let $\text{exc}(\sigma)$ and $\text{fix}(\sigma)$ denote the number of excedances and fixed points of σ , respectively. For all positive integers k , we use the notations for q -integer $[k]_q = 1 + q + \cdots + q^{k-1}$ and $[0]_q = 0$. One of our main results is the following bijection.

Theorem 1.1. *There is a bijection $\Psi : \mathfrak{S}_n \rightarrow \mathscr{W}_n$ such that a permutation $\sigma \in \mathfrak{S}_n$ with i inversions, j fixed points, k excedances and a depth of ℓ is carried to a 3-colored Motzkin path $\mu \in \mathscr{W}_n$ with weight $\omega(\mu) = q^i p^j s^k t^\ell$. Therefore, we have*

$$\sum_{n \geq 0} \left(\sum_{\sigma \in \mathfrak{S}_n} q^{\text{inv}(\sigma)} p^{\text{fix}(\sigma)} s^{\text{exc}(\sigma)} t^{\text{depth}(\sigma)} \right) z^n = \mathfrak{F}(\gamma_h, \lambda_h) \quad (1.6)$$

with $\lambda_h = s[h]_q^2 (qt)^{2h-1}$ and $\gamma_h = ((1+s)[h]_q + pq^h)(qt)^h$.

Remarks. By the weight given in (1.5), notice that a horizontal step $x \in \{H_2^{(h)}, H_3^{(h)}\}$ can be distinguished by the q -factor of $\omega(x)$ for each $h \geq 1$. That is, if $\omega(x)$ contains q^{h+d} then $x = H_2^{(h)}$ ($H_3^{(h)}$, respectively) if $0 \leq d \leq h-1$ ($d = h$, respectively). Thus, the bijection $\Psi : \mathfrak{S}_n \rightarrow \mathscr{W}_n$ can still be established with $H_2^{(h)}, H_3^{(h)}$ combined and the parameter p omitted.

The classical *Euler numbers* E_n , defined by

$$\begin{aligned} \sum_{n \geq 0} E_n \frac{z^n}{n!} &= \tan z + \sec z \\ &= 1 + z + \frac{z^2}{2!} + 2\frac{z^3}{3!} + 5\frac{z^4}{4!} + 16\frac{z^5}{5!} + 61\frac{z^6}{6!} + 272\frac{z^7}{7!} + 1385\frac{z^8}{8!} + \cdots, \end{aligned}$$

count the the number of *alternating permutations* in \mathfrak{S}_n , i.e., $\sigma \in \mathfrak{S}_n$ such that $\sigma_1 > \sigma_2 < \sigma_3 > \cdots \sigma_n$. The numbers E_{2n} are called the *secant numbers* and the numbers E_{2n+1} are called the *tangent numbers*.

Based on the bijection $\Psi : \mathfrak{S}_n \rightarrow \mathscr{W}_n$, we shall establish an involution on \mathfrak{S}_n in terms of weighted Motzkin paths, which reverses the parities of depth and the number of excedances simultaneously. This proves that the numbers of permutations with even and odd depth (excedance numbers, respectively) are equal if n is even and differ by E_n (E_n up to sign, respectively) if n is odd.

Theorem 1.2. *There is an involution $\sigma \mapsto \sigma'$ on \mathfrak{S}_n satisfying*

$$\text{depth}(\sigma) - \text{depth}(\sigma') = \text{exc}(\sigma) - \text{exc}(\sigma') = \text{inv}(\sigma) - \text{inv}(\sigma') \in \{1, 0, -1\}$$

and resulting in the following identities

$$(i) \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{depth}(\sigma)} = \begin{cases} E_n & \text{for } n \text{ odd} \\ 0 & \text{for } n \text{ even.} \end{cases}$$

$$(ii) \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{exc}(\sigma)} = \begin{cases} (-1)^{\frac{n-1}{2}} E_n & \text{for } n \text{ odd} \\ 0 & \text{for } n \text{ even.} \end{cases}$$

Note that the identity in Theorem 1.2(ii) is a classical result of Euler. When the fixed points of σ are ignored, we present some interesting results on the sign imbalances of permutations and derangements, refined with respect to depth and excedance numbers.

Theorem 1.3. For $n \geq 1$, we have

$$\sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{inv}(\sigma)} s^{\text{exc}(\sigma)} t^{\text{depth}(\sigma)} = (1-st)^{n-1}. \quad (1.7)$$

A *derangement* of size n is a permutation in \mathfrak{S}_n that contains no fixed point. Let $\mathcal{D}_n \subset \mathfrak{S}_n$ be the set of derangements of size n . Define

$$F_n = F_n(s, t) = \sum_{\sigma \in \mathcal{D}_n} (-1)^{\text{inv}(\sigma)} s^{\text{exc}(\sigma)} t^{\text{depth}(\sigma)}. \quad (1.8)$$

Several of the initial polynomials $F_n(s, t)$ are listed below:

$$\begin{aligned} F_1(s, t) &= 0, \\ F_2(s, t) &= -st, \\ F_3(s, t) &= s(1+s)t^2, \\ F_4(s, t) &= s^2t^2 - s(1+s)^2t^3, \\ F_5(s, t) &= -2s^2(1+s)t^3 + s(1+s)^3t^4. \end{aligned}$$

Collected in powers of t , the coefficient s -polynomials of $F_n(s, t)$ for $2 \leq n \leq 9$ are listed in Table 1. We obtain a neat expression for the sign imbalance of the joint distribution of depth and excedance numbers over \mathcal{D}_n .

Table 1: The s -polynomial coefficients of $F_n(s, t)$ in powers of t for $2 \leq n \leq 9$.

	t	t^2	t^3	t^4	t^5	t^6	t^7	t^8
F_2	$-s$							
F_3		$s(1+s)$						
F_4		s^2	$-s(1+s)^2$					
F_5			$-2s^2(1+s)$	$s(1+s)^3$				
F_6			$-s^3$	$3s^2(1+s)^2$	$-s(1+s)^4$			
F_7				$3s^3(1+s)$	$-4s^2(1+s)^3$	$s(1+s)^5$		
F_8				s^4	$-6s^3(1+s)^2$	$5s^2(1+s)^4$	$-s(1+s)^6$	
F_9					$-4s^4(1+s)$	$10s^3(1+s)^3$	$-6s^2(1+s)^5$	$s(1+s)^7$

Theorem 1.4. We have

$$\sum_{n \geq 1} \left(\sum_{\sigma \in \mathcal{D}_n} (-1)^{\text{inv}(\sigma)} s^{\text{exc}(\sigma)} t^{\text{depth}(\sigma)} \right) z^n = \sum_{k \geq 1} (-1)^k \left(\sum_{i=0}^{k-1} \binom{k-1}{i} s^{1+i} (1+s)^{k-1-i} z^{k+1+i} \right) t^k.$$

References

- [1] P. Biane (1993): *Permutations suivant le type d'excédance et le nombre d'inversions et interprétation combinatoire d'une fraction continue de Heine*. *European J. Combin.* 14(4), pp. 277–284, doi:10.1006/eujc.1993.1031.
- [2] R.J. Clarke, E. Steingrímsson & J. Zeng (1997): *New Euler–Mahonian statistics on permutations and words*. *Adv. Appl. Math.* 18(3), pp. 237–270, doi:10.1006/aama.1996.0506.
- [3] P. Diaconis & R.L. Graham (1977): *Spearman's footrule as a measure of disarray*. *J. Roy. Statist. Soc. Ser. B* 39(2), pp. 262–268, doi:10.1111/j.2517-6161.1977.tb01624.x.
- [4] D. Dumont (1995): *Further triangles of Seidel–Arnold type and continued fractions related to Euler and Springer numbers*. *Adv. Appl. Math.* 16(3), pp. 275–296, doi:10.1006/aama.1995.1014.
- [5] S. Elizalde (2018): *Continued fractions for permutation statistics*. *Discrete Math. Theor. Comput. Sci.* 19(2), doi:10.23638/DMTCS-19-2-11. arXiv:1703.08742.
- [6] P. Flajolet (1980): *Combinatorial aspects of continued fractions*. *Discrete Math.* 32(2), pp. 125–161, doi:10.1016/0012-365X(80)90050-3.
- [7] D. Foata & D. Zeilberger (1990): *Denert's permutation statistic is indeed Euler–Mahonian*. *Stud. Appl. Math.* 83(1), pp. 31–59, doi:10.1002/sapm199083131.
- [8] M. Guay-Paquet & T.K. Petersen (2014): *The generating function for total displacement*. *Electron. J. Combin.* 21(3):P3.37, doi:10.37236/4329.
- [9] D.E. Knuth (1998): *The Art of Computer Programming, vol. 3*, second edition. Addison Wesley Longman Publishing Co., Inc., USA.
- [10] T.K. Petersen: *A note for Doron: the generation function for total displacement (Spearman's footrule) and inversion*. Available at <https://sites.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/noga12yleFeedback.pdf>.
- [11] T.K. Petersen & B.E. Tenner (2015): *The depth of a permutation*. *J. Combin.* 6(1–2), pp. 145–178, doi:10.4310/JOC.2015.v6.n1.a9.
- [12] N.J.A. Sloane: *The On-Line Encyclopedia of Integer Sequences*. Available at <http://oeis.org>.
- [13] A.D. Sokal & J. Zeng (2022): *Some multivariate master polynomials for permutations, set partitions, and perfect matchings, and their continued fractions*. *Adv. Appl. Math.* 138:102341, doi:10.1016/j.aam.2022.102341.