

# Counting Colored Tilings on Grids and Graphs

José L. Ramírez

Departamento de Matemáticas  
Universidad Nacional de Colombia  
Bogotá, Colombia  
jlamirezr@unal.edu.co

Diego Villamizar

Escuela de Ciencias Exactas e Ingeniería  
Universidad Sergio Arboleda  
Bogotá, Colombia  
diego.villamizarr@usa.edu.co

In this paper, we explore some generalizations of a counting problem related to tilings in grids of size  $2 \times n$ , which was originally posed as a question on Mathematics Stack Exchange (Question 3972905). In particular, we consider this problem for the product of two graphs  $G$  and  $P_n$ , where  $P_n$  is the path graph of  $n$  vertices. We give explicit bivariate generating functions for some specific cases.

## 1 Introduction

Question 3972905 in Mathematics Stack Exchange asks for the number of ways to partition a tile  $2 \times n$  into  $s$  parts. That is the number of different configurations (tilings) in a grid of size  $2 \times n$  with exactly  $s$  polyominoes using 2 colors. For example, if  $n = 4$  we have 12 configurations with exactly 4 polyominoes, see Figure 1.




Figure 1: Configurations of a grid  $2 \times 3$  with exactly 4 polyominoes.

In [4], we study this problem for a general grid of size  $m \times n$  and  $k$  colors. We employ generating functions to provide a partial solution to this problem for the cases  $m = 1, 2, 3$ . Specifically, if  $c(n, i)$  represents the number of different tilings of a  $2 \times n$  grid with exactly  $i$  polyominoes and using two colors, then

$$\begin{aligned} \sum_{n, i \geq 1} c(n, i) x^n y^i &= \frac{2xy(1+y-x(1-y)(1-2y))}{1-x(2+y+y^2)+x^2(1-y)(1-5y^2-2y(1-2y))} \\ &= (2y+2y^2)x + (2y+12y^2+2y^4)x^2 + (2y+30y^2+18y^3+12y^4+2y^6)x^3 \\ &\quad + (2y+56y^2+102y^3+56y^4+24y^5+14y^6+2y^8)x^4 + O(x^5). \end{aligned}$$

Figure 1 shows the colored tilings corresponding to the bold coefficient in the above series.

This counting problem was explored by Richey [5] in 2014. Specifically, he showed that  $\lim_{n, m \rightarrow \infty} e(m, n)/mn$  exists and is finite, where  $e(m, n)$  is the expected number of polyominoes on the  $m \times n$  grid. Mansour [3] considers this problem for bicolored tilings ( $k = 2$ ) for  $m = 1, 2, 3$  using automata. A related problem was addressed by Bodini during GASCOM 2022, referred to as *rectangular shape partitions* [1].

## 2 Colored Tilings of Grids

Let  $\mathcal{T}_{m,n}^{(k)}$  denote the set of tilings of an  $m \times n$  grid with polyominoes colored with one of  $k$  colors, such that adjacent polyominoes are colored with different colors. An element of  $\mathcal{T}_{m,n}^{(k)}$  is called a  $k$ -colored tiling. Given a  $k$ -colored tiling  $T$  in  $\mathcal{T}_{m,n}^{(k)}$ , we use  $\rho(T)$  to denote the number of polyominoes in  $T$ . For fixed positive integers  $m$  and  $k$ , we define the bivariate generating function

$$C_m^{(k)}(x, y) := \sum_{n \geq 1} x^n \sum_{T \in \mathcal{T}_{m,n}^{(k)}} y^{\rho(T)}. \tag{1}$$

Note that the coefficient of  $x^n y^i$  in  $C_m^{(k)}(x, y)$  is the number of  $k$ -colored tilings of an  $m \times n$  grid with exactly  $i$  polyominoes. Let  $c_{m,k}(n, i)$  denote the coefficient of  $x^n y^i$  in the generating function  $C_m^{(k)}(x, y)$ . In [4], we derive explicit generating functions for the cases  $m = 1, 2, 3$ . Additionally, we introduce a variation of this problem for hexagonal grids.

The combinatorial problem can be described in terms of graphs. Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two undirected graphs. The product of  $G_1$  and  $G_2$  is defined as  $G_1 \times G_2 = (V_1 \times V_2, E_{G_1 \times G_2})$ , where

$$E_{G_1 \times G_2} = \{ \{(v_1, v_2), (w_1, w_2)\} : (v_1 = w_1 \text{ and } \{v_2, w_2\} \in E_2) \text{ or } (v_2 = w_2 \text{ and } \{v_1, w_1\} \in E_1) \}.$$

Let  $P_n$  be a *path graph*, that is a simple graph with  $n$  vertices arranged in a linear sequence in such a way that two vertices are adjacent if they are consecutive in the sequence, and are non-adjacent otherwise. A *grid graph* of size  $m \times n$  is defined as the product  $P_m \times P_n$ , and it is denoted by  $L_{m,n}$ .

Let  $G = (V, E)$  be an undirected graph. Two non-empty disjoint subsets  $V_1, V_2 \subseteq V$  are *neighbors* if there is an edge  $(v_1, v_2) \in E$  such that  $v_1 \in V_1$  and  $v_2 \in V_2$ . A  $k$ -colored partition of size  $s$  of the vertices  $V$  of  $G$  is a partition of the set  $V = \bigcup_{i=1}^s V_i$  such that for each  $V_i$ , the induced graph is connected, all vertices in  $V_i$  are colored with exactly one of  $k$  colors, and any pair  $V_i$  and  $V_j$  of neighbors are colored with different colors.

For example, Figure 2 (left) shows a 3-colored partition of size 7 of the grid graph  $L_{3,8}$ , and the Figure 2 (right) shows the corresponding tiling in  $\mathcal{T}_{3,8}^{(3)}$ .




Figure 2: A 3-colored partition of size 7 of  $L_{3,8}$ .

Let  $G$  be an undirected graph. We denote by  $\mathcal{T}_n^{(k)}(G)$  the set of  $k$ -colored partitions of  $G \times P_n$ . Given a  $k$ -colored partition  $T$  in  $\mathcal{T}_n^{(k)}(G)$ , we use  $\rho(T)$  to denote the size of the partition. For fixed positive integers  $m$  and  $k$ , we define the bivariate generating function

$$C_G^{(k)}(x, y) := \sum_{n \geq 1} x^n \sum_{T \in \mathcal{T}_n^{(k)}(G)} y^{\rho(T)}.$$

It is clear that  $C_m^{(k)}(x, y) = C_{P_m}^{(k)}(x, y)$ .

### 3 The complete graph case.

In this section we analyze the case when  $G = K_m$ , where  $K_m$  is the complete graph of size  $m$ . For example, Figure 3 shows a 2-colored partition of size 4 of the graph  $K_5 \times P_4$ .




Figure 3: A 2-colored partition of size 4 of  $K_5 \times P_4$ .

#### 3.1 The case $m = 3$ .

In this section we give the explicit bivariate generating function for the 2-colored partitions of  $K_3 \times P_n$  for all  $n \geq 1$ .

**Theorem 3.1.** *The bivariate generating function  $T(x, y) = C_{K_3}^{(2)}(x, y)$  is given by*

$$T(x, y) = \frac{2xy(1 + 3y - x(3 - 7y + 4y^2))}{1 - x(4 + 3y + y^2) + x^2(3 - 7y + 3y^2 + y^3)}.$$

Moreover,  $[x^n]T(x, 1) = 8^n$ .

*Proof.* Let  $\mathcal{A}_n$  and  $\mathcal{B}_n$  denote the sets of colored tilings in  $\mathcal{T}_n^{(2)}(K_3)$ , such that in the first case the last triangle is colored with only one color, while in  $\mathcal{B}_n$ , the last triangle is colored with the two colors.

Now, we define the bivariate generating functions:

$$T_1(x, y) := \sum_{n \geq 1} x^n \sum_{T \in \mathcal{A}_n} y^{\rho(T)} \quad \text{and} \quad T_2(x, y) := \sum_{n \geq 1} x^n \sum_{T \in \mathcal{B}_n} y^{\rho(T)}.$$

It is clear that  $T(x, y) = T_1(x, y) + T_2(x, y)$ .

Let  $T$  be a 2-colored partition in  $\mathcal{A}_n$ . If  $n = 1$ , then  $T = K_3$ , and its contribution to the generating function is the term  $2xy$  because it has to be monochromatic. If  $n > 1$ , then  $T$  may be decomposed as either  $T_1K_3$  or  $T_2K_3$ , where  $T_1 \in \mathcal{A}_{n-1}$ , and  $T_2 \in \mathcal{B}_{n-1}$ . Depending on whether the colors of the last two triangles coincide or not, we obtain the cases given in Table 1.

From this decomposition, we obtain the functional equation

$$T_1(x, y) = 2xy + xT_1(x, y) + xyT_1(x, y) + xT_2(x, y) + xT_2(x, y).$$

For the colored tilings in  $\mathcal{B}_n$  we obtain the different decompositions given in Table 2. From this decomposition we obtain the functional equation:

$$T_2(x, y) = 6xy^2 + 3xyT_1(x, y) + 3xyT_1(x, y) + 3xT_2(x, y) + xy^2T_2(x, y) + 2xyT_2(x, y).$$




Table 1: Cases for the generating function  $T_1(x,y)$ .




Table 2: Cases for the generating function  $T_2(x,y)$ .

Since  $T(x,y) = T_1(x,y) + T_2(x,y)$ , we have a system of three linear equations with three unknowns  $T(x,y)$ ,  $T_1(x,y)$ , and  $T_2(x,y)$ . Solving the system for  $T(x,y)$  we obtain the desired result.  $\square$

As a series expansion, the generating function  $T(x,y)$  begins with

$$T(x,y) = (2y + 6y^2)x + (2y + 44y^2 + 12y^3 + 6y^4)x^2 + (2y + 178y^2 + 218y^3 + 84y^4 + 24y^5 + \mathbf{6y^6})x^3 + (2y + 600y^2 + 1674y^3 + 1100y^4 + 528y^5 + 150y^6 + 36y^7 + 6y^8)x^4 + O(x^5).$$

Figure 4 shows the 2-colored partitions corresponding to the bold coefficient in the above series.




Figure 4: All 2-colored partitions in  $\mathcal{T}_3^{(2)}(K_3)$ .

**Corollary 3.2.** *The expected number for the size of the partition when the colors assigned to each vertex are selected uniformly in  $\mathcal{T}_3^{(2)}(K_3)$  is given by*

$$\frac{2^{3n-5}(37 + 19n)}{2^{3n}}.$$

## References

- [1] O. Bodini (2022): *On the strange kinetic aesthetic of rectangular shape partitions*. *Pure Math. Appl. (P.U.M.A.)* 30, doi:10.2478/puma-2022-0007. Available at <https://sciendo.com/article/10.2478/puma-2022-0007>.
- [2] Mathematics Stack Exchange (2021): *Question 3972905 in Mathematics Stack Exchange*. *Number of ways to partition  $2 \times N$ . Tile into  $m$  parts*. Available at <https://math.stackexchange.com/q/3972905>.
- [3] R. Mansour (2021): *Counting clusters in a coloring grid*. *Discrete Math. Lett.* 5. Available at [https://www.dmllett.com/archive/v5/DML21\\_v5\\_pp20-23.pdf](https://www.dmllett.com/archive/v5/DML21_v5_pp20-23.pdf).
- [4] J. L. Ramírez & D. Villamizar (2024): *Colored random tilings on grids*. To appear in *Journal of Automata, Languages and Combinatorics*.
- [5] J. Richey (2014): *Counting clusters on a grid*. Undergraduate Honors Thesis. Dartmouth College.