

ENUMERATION OF TERRACINI SCHEMES

CIRO CILIBERTO

ABSTRACT. In this note we outline a way of computing the expected length of the Terracini scheme of a curve, when this scheme is expected to be finite and we give a closed formula for curves in \mathbb{P}^4 . We also discuss the widely open case of varieties of higher dimension.

INTRODUCTION

Let $X \subset \mathbb{P}^r$ be an irreducible, projective variety, that we assume to be smooth and non-degenerate. Let p_1, \dots, p_h be $h \geq 2$ general points of X . The famous *Terracini's lemma* says that the span of the union of the tangent spaces to X at p_1, \dots, p_h is the tangent space to the $(h-1)$ -th secant variety $\text{Sec}_{h-1}(X)$ to X at a general point x of the linear space $\langle p_1, \dots, p_h \rangle$, that is also a general point of $\text{Sec}_{h-1}(X)$. The expected dimension of $\text{Sec}_{h-1}(X)$, and therefore of $T_{\text{Sec}_r(X), x}$, is $\min\{r, h(n+1) - 1\}$, and the dimension $\sigma_{h-1}(X)$ of $\text{Sec}_{h-1}(X)$ is bounded above by the expected dimension. If $\sigma_{h-1}(X) < \min\{r, h(n+1) - 1\}$, X is said to be $(h-1)$ -defective. Curves are never defective, whereas in higher dimension there are a lot of defective varieties.

In [2] the Authors introduced the concept of *Terracini loci* of X . According to [2], the h -th *Terracini locus* of X is the subset of X_h , the Hilbert scheme of 0-dimensional schemes of length h in X , of h -tuples of distinct points p_1, \dots, p_h of X such that the span of the union of the tangent spaces to X at p_1, \dots, p_h is strictly smaller than $\sigma_{h-1}(X)$. In particular, if X is not $(h-1)$ -defective, the h -th Terracini locus of X is the set of distinct h -tuples of points p_1, \dots, p_h of X such that

$$\dim \left(\left\langle \bigcup_{i=1}^h T_{X, p_i} \right\rangle \right) < \min\{r, h(n+1) - 1\}.$$

One can actually give a more formal definition of *Terracini scheme* in X_h (see §1 for the case of curves).

In this paper we look at the case of non- $(h-1)$ -defective varieties X for which the h -th Terracini scheme is expected to be finite and we consider the problem of computing the expected length this scheme. In §1 we outline a way of solving this problem in the case of curves, and we provide a closed formula for the aforementioned length in the case of curves in \mathbb{P}^4 . In §2 we briefly discuss the case of higher dimensional varieties, which is widely open.

Acknowledgement: This problem arose in the course of a problem session during the Workshop “Geometry of secants” at the IMPAM of Warsaw in October

2020 *Mathematics Subject Classification.* Primary 14N10; Secondary 14N07.
Key words and phrases. Terracini schemes, secant varieties.

2021. The author thanks the organizers and the participants for the very stimulating and nice atmosphere. The author also thanks Paolo Aluffi for interesting discussions about the topic of this note.

The author is a member of GNSAGA of the Istituto Nazionale di Alta Matematica “F. Severi”.

1. THE CURVE CASE

Let $n \geq 2$ be an integer and let $C \subset \mathbb{P}^{3n-2}$ be a smooth, irreducible, projective, non-degenerate curve of degree d and genus g . We will denote by L the hyperplane class bundle on C and by $V \subset H^0(C, L)$ the $(3n-1)$ -dimensional vector space corresponding to the embedding of C in \mathbb{P}^4 .

Let C_n be the n -fold symmetric product of C . Consider $C \times C_n$ with the two projections π_1, π_2 to the first and second factor. Let $D \subset C \times C_n$ be the universal divisor. We define a rank $2n$ vector bundle \mathcal{E}_C on C_n as follows

$$\mathcal{E}_C = (\pi_2)_*(\mathcal{O}_{2D} \otimes \pi_1^*(L))$$

and we can look at the evaluation map

$$\varphi : V \otimes \mathcal{O}_{C_n} \longrightarrow \mathcal{E}_C.$$

We can consider the subscheme $\text{Terr}(C)$ of C_n of points $E \in C_n$ where $\text{rk}(\varphi) < 2n$. The scheme $\text{Terr}(C)$ is called the *Terracini scheme* of C . A divisor $E \in C_n$ belongs to $\text{Terr}(C)$ if and only if the subspace $V(-2E) \subset V$ of sections $s \in V$ such that $\text{div}(s) - 2E$ is effective, has dimension at most $2n-1$.

In this setting one may expect that $\text{Terr}(C)$ is finite. If so its length will be denoted by $t(C)$, and, by Porteous formula (see [1, p. 86]), one has

$$(1) \quad t(C) = \begin{vmatrix} c_1 & \dots & c_{n-1} & c_n \\ c_0 & \dots & c_{n-2} & c_{n-1} \\ \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & c_1 \end{vmatrix}$$

where $c_i = c_i(\mathcal{E}_C)$ for $0 \leq i \leq n$. To compute $t(C)$ one has to compute the Chern classes $c_i = c_i(\mathcal{E}_C)$. To do this, one can proceed as follows.

By Grothendieck–Riemann–Roch theorem, one has

$$\text{td}(C_n) \cdot \text{ch}(\mathcal{E}_C) = (\pi_2)_*(\text{td}(C \times C_n) \cdot \text{ch}(\mathcal{O}_{2D} \otimes \pi_1^*(L))).$$

Eliminating the factor $\text{td}(C_n)$, we find

$$(2) \quad \text{ch}(\mathcal{E}_C) = (\pi_2)_*(\text{td}(C) \cdot \text{ch}(\mathcal{O}_{2D} \otimes \pi_1^*(L))).$$

From the exact sequence

$$0 \longrightarrow \pi_1^*(L) \otimes \mathcal{O}_{C \times C_2}(-2D) \longrightarrow \pi_1^*(L) \longrightarrow \mathcal{O}_{2D} \otimes \pi_1^*(L) \longrightarrow 0$$

we deduce

$$(3) \quad \text{ch}(\mathcal{O}_{2D} \otimes \pi_1^*(L)) = \text{ch}(\pi_1^*(L)) - \text{ch}(\pi_1^*(L) \otimes \mathcal{O}_{C \times C_2}(-2D)).$$

Let us denote by η the class of the pull back by π_1 of a point p of C , i.e., the class of $\{p\} \times C_n \cong C_n$. Of course, one has

$$\eta^2 = 0.$$

We denote by δ the class of D and by θ the pull back of the theta divisor in the jacobian $J(C)$ via the Abel–Jacobi map $a : C_n \rightarrow J(C)$. By abuse of notation, θ will also denote the pull back on $C \times C_n$ via π_2 of this class. One has

$$\delta = n\eta + \gamma + x$$

where x denotes the class of a divisor of the form $p + C_{n-1} \subset C_n$, with $p \in C$ (by abuse of notation, x will also denote the pull back of this class on $C \times C_n$ via π_2), and γ is a suitable class on $C \times C_n$ (see [1, p. 338]), that is such that

$$\gamma^2 = -2\eta\theta, \quad \gamma^3 = \eta\gamma = 0.$$

One has

$$\text{ch}(\pi_1^*(L)) = e^{d\eta}, \quad \text{ch}(\pi_1^*(L) \otimes \mathcal{O}_{C \times C_2}(-2D)) = e^{d\eta - 2\delta} = e^{(d-2n)\eta - 2\gamma} e^{-2x}$$

and

$$\text{td}(C) = 1 + (1 - g)\eta.$$

Hence, from (3) we get

$$\text{td}(C) \cdot \text{ch}(\mathcal{O}_{2D} \otimes \pi_1^*(L)) = 1 + (d - g + 1)\eta - (1 + (d - g - 2n + 1)\eta - 2\gamma - 4\eta\theta)e^{-2x}.$$

By (2), and taking into account that $\eta\gamma = 0$ implies $(\pi_2)_*(\gamma) = 0$, we find

$$(4) \quad \text{ch}(\mathcal{E}_C) = (d - g + 1) - (d - g - 2n + 1 - 4\theta)e^{-2x}$$

Expanding e^{-2x} , taking into account that $x^n = 1$ and using Poincaré's formula to compute $x^i \theta^{n-i}$ for $0 \leq i \leq n$, one computes $\text{ch}(\mathcal{E}_C)$. Now

$$\text{ch}(\mathcal{E}_C) = 2n + \sum_{i=1}^n \frac{p_i}{n!}$$

where $p_1 = c_1$ and p_i is determined inductively by *Newton's formula*

$$p_i - c_1 p_{i-1} + c_2 p_{i-2} + \cdots + (-1)^{i-1} c_{i-1} p_1 + (-1)^i i c_i = 0$$

(see [3, p. 56]). From this, in principle one can recursively compute the Chern classes of \mathcal{E}_C and plugging into (1) one finds the expression for $t(C)$.

The general computation is cumbersome and it is not easy to come up with a closed formula for $t(C)$ for all $n \geq 2$. This is however possible for low values of n . For example, one has:

Proposition 1.1. *For $n = 2$, i.e., for curves in \mathbb{P}^4 , one has*

$$t(C) = 2(d - g - 3)(d - g - 4) + 8g(d - 5).$$

Proof. Formula (1) now reads

$$(5) \quad t(C) = c_1^2(\mathcal{E}_C) - c_2(\mathcal{E}_C)$$

so we have to compute only $c_1^2(\mathcal{E}_C)$ and $c_2(\mathcal{E}_C)$.

Formula (4) in this case reads

$$\text{ch}(\mathcal{E}_C) = 4 + 2(d - g - 3)x + 4\theta - 2(d - g - 3)x^2 - 8\theta x.$$

From this, taking into account that $x^2 = 1, \theta x = g$, we get that

$$c_1(\mathcal{E}_C) = 2(d - g - 3)x + 4\theta$$

$$\frac{c_1(\mathcal{E}_C)^2 - 2c_2(\mathcal{E}_C)}{2} = -2(d - g - 3)x^2 - 8\theta x = -2(d - g - 3) - 8g.$$

Since $\theta^2 = g(g-1)$ we deduce that

$$\begin{aligned} c_1(\mathcal{E}_C)^2 &= 4(d-g-3)^2 + 16(d-g-3)g + 16g(g-1) \\ c_2(\mathcal{E}_C) &= 2(d-g-3)(d-g-2) + 8(d-3-g)g + 8g^2. \end{aligned}$$

and, plugging into (5), we see that the assertion holds. \square

2. OPEN PROBLEMS

One can consider Terracini schemes also for higher dimensional varieties and one can consider enumerative problems as in the curve case we treated above. Indeed, let $n \geq 2$ be an integer, and let $X \subset \mathbb{P}^{n(2m+1)-2}$ be a smooth, projective, irreducible, non-degenerate, not $(n-1)$ -defective variety of dimension $m \geq 2$. Let L be the hyperplane class bundle on X and let $V \subset H^0(C, L)$ be the $(n(2m+1)-1)$ -dimensional vector space corresponding to the embedding of X in $\mathbb{P}^{n(2m+1)-2}$.

Let X_n be the Hilbert scheme of 0-dimensional subschemes of length n of X . Consider $X \times X_n$ with the two projections π_1, π_2 to the first and second factor. Let $D \subset X \times X_n$ be the universal scheme. We define a rank $n(m+1)$ vector bundle \mathcal{E}_X on X_n as follows

$$\mathcal{E}_X = (\pi_2)_*(\mathcal{O}_{X \times X_n} / \mathcal{I}_D^2 \otimes \pi_1^*(L)),$$

where \mathcal{I}_D is the ideal sheaf of D in $X \times X_n$. One can look at the evaluation map

$$\varphi : V \otimes \mathcal{O}_{X_n} \longrightarrow \mathcal{E}_X$$

and consider the subscheme $\text{Terr}(X)$ of X_n of points $Z \in X_n$ where $\text{rk}(\varphi) < n(m+1)$. The scheme $\text{Terr}(X)$ is called the *Terracini scheme* of X . A scheme $Z \in X_n$ belongs to $\text{Terr}(X)$ if and only if the subspace $V(-2Z) \subset V$, with $V(-2Z) = V \cap H^0(X, L \otimes \mathcal{I}_{Z, X}^2)$, has dimension at most $n(m+1)-1$.

In this setting, one may expect that $\text{Terr}(X)$ is finite and one may want to compute its expected length. In principle, one can follow the same argument outlined in the curve case, but, since the universal scheme $D \subset X \times X_n$ is no longer a divisor, the computations become soon very difficult, even in the simplest case $m = n = 2$ of surfaces in \mathbb{P}^8 . It is therefore an open problem to find formulas for the length of $\text{Terr}(X)$ in the above cases.

REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths, J. Harris, *Geometry of Algebraic Curves, Vol. I*, Springer Verlag, Grundlehren der mathematischen Wissenschaften, **267**, 1985.
- [2] E. Ballico, L. Chiantini, *On the Terracini locus of projective varieties*, Milan Journal of Mathematics, **89** (2021), 1–17.
- [3] W. Fulton, *Intersection Theory*, Springer Verlag, Ergebnisse der Math. und ihre Grenzgeb., **2**, 1984.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA TOR VERGATA, VIA O. RAIMONDO 00173, ROMA, ITALIA

Email address: `cilibert@mat.uniroma2.it`