SELF-SIMILAR BLOWUP FOR THE CUBIC SCHRÖDINGER EQUATION

ROLAND DONNINGER AND BIRGIT SCHÖRKHUBER

ABSTRACT. We give a rigorous proof for the existence of a finite-energy, self-similar solution to the focusing cubic Schrödinger equation in three spatial dimensions.

1. INTRODUCTION

This paper is concerned with the cubic, focusing Schrödinger equation

$$i\partial_t \psi(t,x) + \Delta_x \psi(t,x) + \psi(t,x)|\psi(t,x)|^2 = 0$$
(1.1)

for an unknown $\psi : I \times \mathbb{R}^3 \to \mathbb{C}$, where $I \subset \mathbb{R}$ is an interval. Eq. (1.1) is the basic model for describing the competition of dispersion and nonlinear, focusing effects. As a consequence, the cubic Schrödinger equation appears in virtually every problem where nonlinear wave interaction is crucial, e.g. in nonlinear optics or in the study of propagation of waves in plasmas. We refer the interested reader to [25] and references therein for the background. In addition, nonlinear Schrödinger equations are fundamental for many-particle quantum mechanics where they occur as effective equations in suitable limits of large particle numbers. From a pure mathematical perspective, the desire to understand nonlinear Schrödinger equations is one of the main driving forces in many branches of contemporary analysis.

The natural setup for studying Eq. (1.1) is the *Cauchy problem*, that is to say, one prescribes a function $\psi_0 : \mathbb{R}^3 \to \mathbb{C}$ (the *data*) and attempts to understand the behavior of the solution ψ of Eq. (1.1) that satisfies $\psi(0, \cdot) = \psi_0$. It is well-known that at least for short times t, this is a wellposed problem in the sense that a unique solution exists that depends continuously on the data, provided ψ_0 is sufficiently regular, see e.g. [25] and references therein. For large times, however, the situation is much more delicate. In fact, it is known that solutions may cease to exist after a finite time. This can be shown by a qualitative argument [7] which, however, gives little information on what actually happens to the solution. Consequently, there have been many attempts to quantify the breakdown and numerical studies indicate the existence of a nontrivial (meaning not identically zero) *self-similar solution* ψ_* of the form

$$\psi_*(t,x) = \frac{1}{\sqrt{2\alpha}} (1-t)^{-\frac{1}{2} - \frac{i}{2\alpha}} Q\left(\frac{x}{\sqrt{2\alpha(1-t)}}\right),\tag{1.2}$$

defined on $(-\infty, 1) \times \mathbb{R}^3$, where $Q : \mathbb{R}^3 \to \mathbb{C}$ is a smooth, radial function in $L^4(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3)$ and $\alpha \approx 0.917$, see [25], p. 121. Such a solution provides a quantitative example of the breakdown but what is even more exciting is the conjectured *universality* of ψ_* . Indeed, in numerical simulations one observes that for sufficiently large but otherwise arbitrary data, the solution develops a singularity and close to the breakdown, the shape of the solution is independent of the particular form of the data and approaches ψ_* (modulo symmetries of the equation). In other words, shortly before breaking down, the solution "forgets" all of its features and takes on a universal shape given by ψ_* . This means that the self-similar solution ψ_* appears to describe the *generic* breakdown behavior

This work was supported by the Austrian Science Fund FWF, Project P34560. Furthermore, the authors would like to thank the Erwin Schrödinger International Institute for Mathematics and Physics (ESI) for hospitality. This work was finalized during the thematic program "Nonlinear Waves and Relativity" at ESI in 2024.

of Eq. (1.1) and is thus a key feature of the model. As a consequence, in order to rigorously understand the dynamics of Eq. (1.1), it is of tantamount importance to prove the existence of ψ_* . Unfortunately, this seems to be a hard problem that resisted all attempts so far. In this paper, we give the first rigorous proof for the existence of the self-similar solution ψ_* .

Theorem 1.1 (Main theorem, qualitative version). There exists a nontrivial, radial function $Q \in L^4(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3) \cap C^{\infty}(\mathbb{R}^3)$ and an $\alpha > 0$ such that $\psi_* : (-\infty, 1) \times \mathbb{R}^3 \to \mathbb{C}$, defined by

$$\psi_*(t,x) := \frac{1}{\sqrt{2\alpha}} (1-t)^{-\frac{1}{2} - \frac{i}{2\alpha}} Q\left(\frac{x}{\sqrt{2\alpha(1-t)}}\right),$$

satisfies

$$i\partial_t \psi_*(t,x) + \Delta_x \psi_*(t,x) + \psi_*(t,x) |\psi_*(t,x)|^2 = 0$$

for all $(t,x) \in (-\infty,1) \times \mathbb{R}^3$. In particular, Q satisfies the profile equation

$$\Delta Q(\xi) - Q(\xi) + i\alpha \left[\xi \cdot \nabla Q(\xi) + Q(\xi)\right] + Q(\xi)|Q(\xi)|^2 = 0$$
(1.3)

for all $\xi \in \mathbb{R}^3$.

In addition to the pure existence statement in Theorem 1.1, we obtain an extremely precise approximation to Q with rigorously proven error bounds.

Definition 1.2 (Chebyshev polynomials). Let $n \in \mathbb{N}_0$. Then $T_n : \mathbb{C} \to \mathbb{C}$ is defined to be the unique polynomial that satisfies

$$T_n(\cos(z)) = \cos(nz)$$

for all $z \in \mathbb{C}$.

Definition 1.3. Let

$$P_*(y) := \sum_{n=0}^{50} c_n(P_*)T_n(y)$$

with $(c_n(P_*))_{n=0}^{50} \subset \mathbb{C}$ given in Appendix A.1. We set

$$a_* := \frac{772201763088846}{841768781900003}$$

and define $g_*: [0,\infty) \times \mathbb{R} \to \mathbb{C}$ by

$$g_*(r,a) := P_*\left(\frac{r-1}{r+1}\right) + \frac{2i(a_*+a)}{1-i(a_*+a)}P'_*(1) - P_*(1).$$

Theorem 1.4 (Main theorem, quantitative version). There exists an $a \in [-10^{-10}, 10^{-10}]$ and a $g: [0, \infty) \to \mathbb{C}$ satisfying

$$2\sup_{r>0} r|g'(r)| + \sup_{r>0} |g(r)| \le 1.2 \cdot 10^{-6}$$

such that $Q: \mathbb{R}^3 \to \mathbb{C}$, defined by

$$Q(x) := (1+|x|)^{-1-\frac{i}{a_*+a}} \left[g_*(|x|,a) + g(|x|) \right],$$

belongs to $L^4(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3) \cap C^{\infty}(\mathbb{R}^3)$ and

$$\psi_*(t,x) := \frac{1}{\sqrt{2(a_*+a)}} (1-t)^{-\frac{1}{2} - \frac{i}{2(a_*+a)}} Q\left(\frac{x}{\sqrt{2(a_*+a)(1-t)}}\right)$$

satisfies

$$i\partial_t \psi_*(t,x) + \Delta_x \psi_*(t,x) + \psi_*(t,x) |\psi_*(t,x)|^2 = 0$$

for all $(t, x) \in (-\infty, 1) \times \mathbb{R}^3$.

Remark 1.5. Note that

$$a_* = \frac{772201763088846}{841768781900003} = 0.9173561430..$$

and

 $g_*(0,0) = -1.88566\ldots$

These particular values appear in the numerical literature [28] and strongly suggest that the solution we construct coincides with the one that has been studied numerically for a long time.

Evidently, Theorem 1.4 implies Theorem 1.1.

1.1. History of the problem. The formation of singularities in the three-dimensional focusing cubic Schrödinger equation has received considerable attention ever since the model was derived by Zakharov [29], [30] in the context of plasma physics. There, Eq. (1.1) arises as a limit of the so-called Zakharov system which provides a simplified description of the propagation of Langmuir waves in an ionized plasma. With ψ modelling the envelope of the electric field, finite-time blowup (referred to as *wave collapse*) via radially symmetric self-similar solutions of the form (1.2) was proposed as a mechanism for energy dissipation.

From a rigorous point of view, a first "obstructive" blowup result for Eq. (1.1) was obtained by Glassey [7] for initial data with finite variance and negative energy, where the latter is a conserved quantity of the flow,

$$E(\psi(t,\cdot)) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla_x \psi(t,x)|^2 dx - \frac{1}{4} \int_{\mathbb{R}^3} |\psi(t,x)|^4 dx = E(\psi(0,\cdot)),$$

see also [20] for an analogous statement in the radial case without requiring finite variance. However, it is in the nature of these types of results that they do not provide much information on the details of singularity formation.

McLaughlin, Papanicolaou, Sulem, and Sulem [15] investigated the problem numerically and confirmed Zakharov's predictions on the asymptotically self-similar nature of the blowup for Eq. (1.1) by looking at different types of radially symmetric initial data. More importantly, they observed convergence to a universal limiting blowup profile corresponding to a solution of Eq. (1.3) for $\alpha \approx 0.917$. Later, Landman, Papanicolaou, Sulem, Sulem, and Wang [12] came to the same conclusion even for nonradial data. Recent studies by Yang, Roudenko, and Zhao [28] confirmed the earlier findings in the radial setting. Due to these results it is widely believed that the asymptotic behavior of solutions to Eq. (1.1) corresponding to "generic" large initial data is well described (up to symmetries) by a self-similar solution of the above form with a radially symmetric blowup profile Q corresponding to $\alpha \approx 0.917$, where |Q| decreases monotonically.

Rigorous results on the profile equation (1.3) were first obtained by Wang [27], who proved the existence of radially symmetric solutions for any $\alpha \in \mathbb{R} \setminus \{0\}$ and prescribed $Q(0) \in \mathbb{R}$. However, the solutions display highly oscillatory behavior at infinity in general. More precisely, for large values of $|\xi|$, $Q = c_1Q_1 + c_2Q_2$, for $c_1, c_2 \in \mathbb{C}$, where

$$Q_1(\xi) \sim |\xi|^{-1-\frac{i}{\alpha}}, \qquad Q_2(\xi) \sim |\xi|^{-2+\frac{i}{\alpha}} e^{-i\alpha \frac{\xi^2}{2}},$$

see e.g. [13]. In particular, solutions have finite energy¹ only if $c_2 = 0$. The latter property is conjectured to hold only for specific values of α . In this respect we would like to mention the recent preprint [26] that erroneously² claims that this is the case for every $\alpha > 0$ and Q(0) > 0. In fact, the solution provided by Theorem 1.4 is the first rigorous example of this kind and its properties strongly suggest that it corresponds to the numerically observed limiting blowup profile.

¹In fact, in this case the energy must be 0, see [25].

²The flaw in the argument of [26] seems to be in the step from Eq. (2.35) to (2.36).

1.2. Related results. The dynamics for the cubic NLS crucially depend on the underlying space dimension. In the one-dimensional case, the equation is completely integrable and solutions exist globally in time. For d = 2, the problem is mass critical, i.e., the L^2 -norm is invariant under the natural scaling of the equation. In this case, singularities may form in finite time and the generic blowup behavior is conjectured to be described by a dynamically rescaling soliton with a nonself-similar blowup rate. The literature in this field is vast and since the present paper is solely concerned with the mass supercritical case, we restrict our discussion to this situation. We remark that the critical space for Eq. (1.1) is $\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)$, which renders the problem mass supercritical and energy subcritical.

For the cubic NLS in dimensions d > 2 sufficiently close to the mass critical case, Kopell and Landman [11] as well as Rottschäfer and Kaper [23] proved the existence of radial self-similar profiles, which are monotonically decreasing in absolute value and display the "good" asymptotic behavior at infinity leading to finite energy. The condition on the dimension translates into smallness of the parameter $\alpha = \alpha(d)$ in the corresponding profile equation. In a similar spirit, but using different methods, radially symmetric self-similar solutions have been constructed recently by Bahri, Martel, and Raphaël [1] in the more general framework of slightly L^2 -supercritical nonlinear Schrödinger equations,

$$i\partial_t \psi(t,x) + \Delta_x \psi(t,x) + \psi(t,x) |\psi(t,x)|^{p-1} = 0,$$
(1.4)

for $d \ge 1$ and $0 , where <math>p_c = 1 + \frac{4}{d}$. More precisely, they proved that for any p sufficiently close to p_c , there exists an $\alpha = \alpha(p)$ and a corresponding nontrivial finite-energy solution of the corresponding profile equation. We note that in this slightly supercritical case also non-monotonic self-similar profiles are known to exist, see the work of Rottschäfer and Kaper [24] as well as Budd, Chen, and Russel [3], and Budd [2], see also [28].

Merle, Raphaël, and Szeftel [17] investigated more general dynamics in this perturbative regime and proved for $1 \leq d \leq 5$ the existence of an open set of initial data in $H^1(\mathbb{R}^d)$ which lead to solutions blowing up at a self-similar rate. Their approach does not rely on the knowledge of exact self-similar profiles, instead they use rough approximations and extend techniques from the mass critical case.

A completely different type of blowup solutions, so-called "standing ring solutions", have been constructed for Eq. (1.4) with p = 5 by Raphaël [21] (d = 2), and Raphaël and Szeftel [22] $(d \ge 3)$ in the radial setting, see also [10]. The considered dimensions correspond to mass supercritical (but energy subcritical), energy critical, and energy supercritical regimes. Moreover, these solutions are stable under small, smooth radial perturbations. Adopting this approach, Holmer and Roudenko [9] and Zwiers [31] studied Eq. (1.1) in axial symmetry and constructed solutions blowing up in finite time on a circle with a nonself-similar rate. Stability of these solutions holds within this particular symmetry class. Fibich, Gavish and Wang [5] revealed yet another blowup mechanism for Eq. (1.1) by numerically observing solutions which focus on a sphere that collapses towards the origin, see also [6]. The existence of such "collapsing ring solutions" was then proved by Merle, Raphaël, and Szeftel [18] for Eq. (1.4) for $d \ge 2$, $p_c and independently by Holmer, Perelman, Roudenko [8] for <math>d = 3$ and p = 3. Both standing and collapsing ring solutions are conjectured to be unstable under nonradial perturbations.

Finally, we note that for Eq. (1.4) in the energy supercritical regime $d \ge 11$ and p sufficiently large, blowup via concentration of a soliton profile has been proved by Merle, Raphaël and Rodnianski [16].

Summarizing the situation for the three-dimensional cubic NLS (1.1), the ring-type solutions of [9, 31, 18, 8] and the self-similar solution ψ_* provided by Theorem 1.4 are, to the best of our knowledge, the only explicit examples for finite-time blowup. Recently, a general framework for analyzing the stability of self-similar blowup solutions in the mass supercritical (but energy

subcritical) regime has been presented by Li [14]. His results imply finite-codimension asymptotic stability of ψ_* in $\dot{H}^s(\mathbb{R}^3)$ for $0 < s - \frac{1}{2} \ll 1$. However, in view of the numerical observations, a much stronger stability result is expected to hold.

Lastly, we would like to mention that Eq. (1.1) admits solitary waves. They have been studied extensively in the past and play a key role in the description of large data dynamics in $H^1(\mathbb{R}^3)$, see in particular the works by Nakanishi and Schlag [19] and Duyckaerts and Roudenko [4] as well as the references therein. The blowup solution ψ_* (as any self-similar solution) does not have finite mass and hence does not fit into this framework.

2. Idea of proof

We give an outline of the proof describing the main difficulties and how we overcome them.

2.1. Setup of the problem. To begin with, we rewrite Eq. (1.1) in self-similar variables, i.e., we define a new unknown u by

$$\psi(t,x) = \frac{1}{\sqrt{2\alpha(1-t)}} u\left(-\frac{1}{2\alpha}\log(1-t), \frac{x}{\sqrt{2\alpha(1-t)}}\right),$$

where $\alpha > 0, t < 1$, and $x \in \mathbb{R}^3$. Then ψ satisfies Eq. (1.1) if and only if u satisfies

$$i\partial_{\tau}u(\tau,\xi) + \Delta_{\xi}u(\tau,\xi) + i\alpha \left[\xi \cdot \nabla_{\xi}u(\tau,\xi) + u(\tau,\xi)\right] + u(\tau,\xi)|u(\tau,\xi)|^2 = 0$$
(2.1)

for $\tau \in \mathbb{R}$ and $\xi \in \mathbb{R}^3$. We are looking for self-similar profiles, that is to say, we insert the ansatz $u(\tau,\xi) = e^{i\tau}Q(\xi)$. This yields the profile equation Eq. (1.3) for the function $Q : \mathbb{R}^3 \to \mathbb{C}$. By writing $Q(\xi) = q(|\xi|)$, Eq. (1.3) transforms into

$$\mathcal{E}(\alpha, q)(r) := q''(r) + \left(\frac{2}{r} + i\alpha r\right)q'(r) + (i\alpha - 1)q(r) + q(r)|q(r)|^2 = 0$$
(2.2)

for r > 0. Consequently, the proof boils down to showing the existence of a suitable solution of Eq. (2.2).

2.2. **Difficulties.** The following observations indicate that solving Eq. (2.2) might be a challenging problem.

- The domain on which Eq. (2.2) is posed is unbounded.
- The equation is nonlinear and depends on a parameter α that is unknown itself.
- The equation is singular at 0 and at ∞ . Moreover, the singularity at ∞ is irregular in the Fuchsian sense and the value of α determines the leading order asymptotics near ∞ .
- There is a degeneracy due to phase invariance: If q is a solution then so is $e^{i\theta}q$ for any $\theta \in \mathbb{R}$.
- There is no hope for a perturbative treatment as there is no known solution that is in any sense "close" to the desired solution.

Since there is no object to "perturb off", one obvious strategy is to first construct an approximate solution (α_*, q_*) with the desired properties by some numerical means. In a second step, one may try to find a correction $(\tilde{\alpha}, \tilde{q})$ so that $(\alpha_* + \tilde{\alpha}, q_* + \tilde{q})$ becomes a true solution. To this end, it is natural to linearize at (α_*, q_*) and write

$$0 = \mathcal{E}(\alpha_* + \widetilde{\alpha}, q_* + \widetilde{q}) = \mathcal{E}(\alpha_*, q_*) + \mathcal{E}'(\alpha_*, q_*)(\widetilde{\alpha}, \widetilde{q}) + \text{nonlinear terms.}$$

If the approximation (α_*, q_*) is accurate enough, i.e., if $\mathcal{E}(\alpha_*, q_*)$ is sufficiently small, one might hope to solve this equation by a fixed point argument. To this end, it is necessary to invert the linear operator $\mathcal{E}'(\alpha_*, q_*)$ and the operator norm of its inverse will determine the required quality of the approximation (α_*, q_*) . This naive approach faces some serious difficulties and is ultimately bound to fail for a number of reasons:

- Already from a pure numerical perspective, constructing highly accurate approximate solutions to Eq. (2.2) is challenging because of the unbounded domain and the singularity at infinity. In fact, a substantial number of numerics papers is concerned with just that.
- The linear operator is not C-linear but merely R-linear due to the form of the nonlinearity. This is more of a technical complication but nevertheless increases the algebraic complexity of the problem considerably.
- The linearized equation

$$\mathcal{E}'(\alpha_*, q_*)(\widetilde{\alpha}, \widetilde{q}) = 0$$

cannot be solved explicitly. Thus, it is unclear how to obtain a fundamental system in order to invert $\mathcal{E}'(\alpha_*, q_*)$.

- In fact, the linear operator $\mathcal{E}'(\alpha_*, q_*)$ has a nontrivial kernel due to the phase invariance and is hence not invertible.
- The correction $\tilde{\alpha}$ cannot be obtained by a purely perturbative argument because it determines the leading order asymptotics of the solution.

2.3. Strategy of the proof. In order to overcome all of these difficulties, we choose a more refined strategy.

• First, we compactify the problem by setting

$$q(r) = (1+r)^{-1-\frac{i}{\alpha}} f\left(\frac{r-1}{r+1}\right).$$

In particular, we factor out the expected asymptotic oscillations of the solution. Then, q satisfies Eq. (2.2) if and only if $\mathcal{R}(\alpha, f) = 0$ on (-1, 1), where

$$\mathcal{R}(\alpha, f)(y) := f''(y) + p_0(y, \alpha)f'(y) + q_0(y, \alpha)f(y) + \frac{f(y)|f(y)|^2}{(1-y)^2},$$

and

$$p_0(y,\alpha) := \frac{4i\alpha}{(1-y)^3} - \frac{2i\alpha}{(1-y)^2} - \frac{2 + \frac{2i}{\alpha}}{1-y} + \frac{2}{1+y}$$
$$q_0(y,\alpha) := -\frac{2 - 2i\alpha}{(1-y)^3} - \frac{\frac{1}{\alpha^2} - \frac{i}{\alpha}}{(1-y)^2} - \frac{1 + \frac{i}{\alpha}}{1-y} - \frac{1 + \frac{i}{\alpha}}{1+y}$$

This removes the problems connected to the unboundedness of the original domain and the oscillatory behavior of the solution near ∞ .

• Next, we employ a Chebyshev pseudo-spectral method in order to obtain an approximate solution (a_*, f_*) that is defined by

$$f_*(y,a) := P_*(y) + \frac{2i(a_* + a)}{1 - i(a_* + a)} P'_*(1) - P_*(1),$$

where the polynomial $P_* : \mathbb{Q} \to \mathbb{Q}$ and $a_* \in \mathbb{Q}$ are given in Definition 1.3. Observe the *a*-dependent correction that accounts for the fact that the parameter $\alpha = a_* + a$ determines the leading-order asymptotics. Furthermore, it is important to note that the origin of the coefficients in Appendix A.1 that define P_* is completely irrelevant. Once they are given, our whole argument is rigorous and independent of numerics.

• Now we would like to correct (a_*, f_*) to an actual solution, i.e., we need to find (a, f) such that $\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = 0$. As usual, the idea is to apply a fixed point argument and hence we linearize at (a_*, f_*) and write

$$\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = \mathcal{R}(a_* + a, f_*(\cdot, a)) + \mathcal{L}_a(f) + \mathcal{N}_a(f),$$

where \mathcal{L}_a is \mathbb{R} -linear and $\mathcal{N}_a(f)$ contains all the terms that are quadratic or cubic in f. Note that we only linearize in f. Thus, our goal is to solve

$$\mathcal{L}_a(f) = -\mathcal{R}(a_* + a, f_*(\cdot, a)) - \mathcal{N}_a(f).$$

- Due to the phase invariance, the operator \mathcal{L}_a has a one-dimensional kernel. We construct an approximate fundamental system for \mathcal{L}_a that is defined by the coefficients in Appendices A.2 and A.3. This fundamental system defines an approximate linear operator $\widetilde{\mathcal{L}}_a$ that is shown to be close to \mathcal{L}_a in a suitable sense. While $\widetilde{\mathcal{L}}_a$ may not have a nontrivial kernel, it certainly has an eigenvalue very close to 0 for perturbative reasons and it is not feasible to invert it directly.
- We employ the variation of constants formula and the explicit fundamental system for $\tilde{\mathcal{L}}_a$ to construct a formal inverse \mathcal{H}_a of $\tilde{\mathcal{L}}_a$ in the ODE sense. That is to say, $\tilde{\mathcal{L}}_a\mathcal{H}_a g = g$ on (-1,1) but \mathcal{H}_a is not an inverse of $\tilde{\mathcal{L}}_a$ in the functional-analytic sense, i.e., \mathcal{H}_a does not map into the right space. Concretely, $\mathcal{H}_a g$ is singular at the left endpoint -1, unless $\psi_a(g) = 0$ for a suitable \mathbb{R} -linear functional ψ_a that is easily read off from the variation of constants formula. From the spectral theory point of view, the kernel of ψ_a is precisely the complement of the eigenspace of the eigenvalue of $\tilde{\mathcal{L}}_a$ close to 0.
- Next, we add a regularization term of the form $\psi_a(g)f_0$, where f_0 is a suitable fixed function that is supported near the left endpoint -1 and singular at -1. With this, we define a new operator

$$\mathcal{J}_a(g) := \mathcal{H}_a(g) + \psi_a(g) f_0$$

which now has nice mapping properties but is of course no longer a formal inverse of \mathcal{L}_a , unless $\psi_a(g) = 0$.

• We then solve the equation

$$f = \mathcal{J}_a\left(\widetilde{\mathcal{L}}_a f - \mathcal{L}_a f - \mathcal{N}_a(f) - \mathcal{R}(a_* + a, f_*(\cdot, a))\right)$$

by applying the contraction mapping principle. This yields a solution $f = f_a$ for any sufficiently small a. Note that the implementation of this fixed point argument requires quantitative bounds for all of the terms involved.

• Finally, we show that

$$a \mapsto \psi_a \left(\widetilde{\mathcal{L}}_a f_a - \mathcal{L}_a f_a - \mathcal{N}_a(f_a) - \mathcal{R}(a_* + a, f_*(\cdot, a)) \right)$$

changes sign and hence, by continuity, vanishes at some value of a. For such a value of a, we have

$$\widetilde{\mathcal{L}}_a f = \widetilde{\mathcal{L}}_a \mathcal{J}_a \left(\widetilde{\mathcal{L}}_a f_a - \mathcal{L}_a f_a - \mathcal{N}_a (f_a) - \mathcal{R} (a_* + a, f_* (\cdot, a)) \right)$$
$$= \widetilde{\mathcal{L}}_a f_a - \mathcal{L}_a f_a - \mathcal{N}_a (f_a) - \mathcal{R} (a_* + a, f_* (\cdot, a))$$

or, equivalently, $\mathcal{R}(a_* + a, f_*(\cdot, a) + f_a) = 0$ and the existence of the desired solution follows. Technically, this is one of the most challenging parts of the proof because extracting the sign change requires very precise quantitative estimates for highly oscillatory integrals.

2.4. Function spaces. As a motivation for our choice of function spaces, it is instructive to consider the linearization around 0, i.e., the \mathbb{C} -linear equation

$$f''(y) + p_0(y,\alpha)f'(y) + q_0(y,\alpha)f(y) = 0.$$

This is not the linear equation we have to deal with but to leading order the behavior near the singular endpoints ± 1 is the same. In order to determine the asymptotics of solutions, we make

the rough ansatz $f(y) \sim e^{(1-y)^{-\sigma}}$. Since $p_0(y, \alpha) \sim (1-y)^{-3}$, we must have $\sigma = 2$. This motivates the refined ansatz

$$f(y) \sim e^{a_2(1-y)^{-2} + a_1(1-y)^{-1} + a_0 \log(1-y)}$$

and by comparing orders we determine the coefficients $a_2 = -2i\alpha$, $a_1 = 2i\alpha$, and $a_0 = 1 - \frac{2i}{\alpha}$. Consequently, near 1, we expect a singular oscillatory solution $\hat{f}_1(y,\alpha) \sim (1-y)e^{i\varphi(y,\alpha)}$ with

$$\varphi(y, \alpha) = -\frac{2\alpha}{(1-y)^2} + \frac{2\alpha}{1-y} - \frac{2}{\alpha}\log(1-y).$$

Furthermore, by a standard power series ansatz we expect a smooth solution $f_1(y,\alpha) \sim 1$. Near -1 the situation is more classical because the singularity at y = -1 is regular. By employing the Frobenius method, we find two solutions $f_{-1}(y,\alpha) \sim 1$ and $\hat{f}_{-1}(y,\alpha) \sim (1+y)^{-1}$. For the Wronskian we obtain the expression

$$W(f_{-1}(\cdot,\alpha), f_1(\cdot,\alpha))(y) = W(f_{-1}(\cdot,\alpha), f_1(\cdot,\alpha))(0)e^{-\int_0^y p_0(x,\alpha)dx}$$

= $W(f_{-1}(\cdot,\alpha), f_1(\cdot,\alpha))(0)(1-y^2)^{-2}e^{i\varphi(y,\alpha)} =: W_*(y,\alpha).$

Assuming that $W(f_{-1}(\cdot, \alpha), f_1(\cdot, \alpha))(0) \neq 0$, the variation of constants formula yields the inverse operator

$$\mathcal{L}^{-1}(\alpha, g)(y) = f_{-1}(y, \alpha) \int_{y}^{1} \frac{f_{1}(x, \alpha)}{W_{*}(x, \alpha)} g(x) dx + f_{1}(y, \alpha) \int_{-1}^{y} \frac{f_{-1}(x, \alpha)}{W_{*}(x, \alpha)} g(x) dx.$$

By the above, we have

$$|f_{-1}(y,\alpha)| \lesssim 1, \qquad |f'_{-1}(y,\alpha)| \lesssim (1-y)^{-2}$$

and

$$|f_1(y,\alpha)| \lesssim (1+y)^{-1}, \qquad |f_1'(y,\alpha)| \lesssim (1+y)^{-2}.$$

Consequently, we infer the bounds

$$\begin{aligned} |\mathcal{L}^{-1}(\alpha,g)(y)| &\lesssim \int_{y}^{1} (1+x)(1-x)^{2} |g(x)| dx + (1+y)^{-1} \int_{-1}^{y} (1-x^{2})^{2} |g(x)| dx \\ &\lesssim \sup_{x \in (-1,1)} (1+x)(1-x)^{2} |g(x)| \end{aligned}$$

and

$$\begin{aligned} |(1-y^2)\mathcal{L}^{-1}(\alpha,g)'(y)| &\lesssim (1-y)^{-1} \int_y^1 (1+x)(1-x)^2 |g(x)| dx + (1+y)^{-1} \int_{-1}^y (1-x^2)^2 |g(x)| dx \\ &\lesssim \sup_{x \in (-1,1)} (1+x)(1-x)^2 |g(x)|. \end{aligned}$$

Motivated by this, we define the norms

$$||f||_X := \sup_{y \in (-1,1)} (1 - y^2) |f'(y)| + ||f||_{L^{\infty}(-1,1)}$$
$$||f||_Y := \sup_{y \in (-1,1)} (1 + y)(1 - y)^2 |f(y)|$$

and obtain the bound

$$\|\mathcal{L}^{-1}(\alpha,g)\|_X \lesssim \|g\|_Y.$$

This means that all terms that are small with respect to $\|\cdot\|_{Y}$ are expected to be perturbative.

These heuristic considerations suggest to base the functional-analytic treatment on the spaces X and Y as given in Definition 3.1 below. Note, however, that $\mathcal{R}(\alpha, f)$ is cubically singular at 1 and hence does not belong to Y in general. As a consequence, we are forced to include a

suitable correction of the approximate solution at the right endpoint. Luckily, this correction can be determined explicitly because the condition

$$\lim_{y \to 1-} (1-y)^3 \mathcal{R}(\alpha, f)(y) = 0$$

involves only terms that are *linear* in f. Finally, the spaces X and Y are also compatible with the nonlinear structure. Indeed, the required local Lipschitz bound

$$\|\mathcal{N}_a(f) - \mathcal{N}_a(g)\|_Y \lesssim (\|f\|_X + \|g\|_X)\|f - g\|_X,$$

is easy to establish.

In the true implementation we are facing the additional technical difficulty that the linearized operator is not \mathbb{C} -linear but only \mathbb{R} -linear. As a consequence, we need to separate real and imaginary parts and treat the linearized equation as a coupled system. Fortunately, the terms that spoil \mathbb{C} -linearity are of higher order and the above asymptotic analysis remains essentially valid. A more substantial difference to the simple heuristics is the fact that we will be forced to choose a fundamental system that involves a function that is bad at both endpoints. This is due to the nontrivial kernel of the linearized operator induced by the phase invariance and eventually leads to the vanishing condition on the functional ψ_a as described before.

2.5. Computer assistance. Our approach splits into a theoretical part and a concrete implementation. The latter is mildly computer-assisted and based on polynomial approximations. More precisely, we use polynomials with *rational coefficients* that are provided explicitly in tables, see Appendix A. At rational points these polynomials can be evaluated *exactly* using elementary fraction arithmetic and this is one of the key features of our approach. In fact, evaluating polynomials with rational coefficients at rational points is the only component of our proof that requires computer assistance. We do not use any kind of symbolic computation, interval arithmetic or more sophisticated computer-assisted tools. In particular, all of our computations are exact and we never have to deal with round-off errors in floating-point arithmetic. Of course, the polynomials we use are initially obtained by some suitable numerical method (usually Chebyshev pseudo-spectral techniques) but their origin is completely irrelevant to the argument and logically independent of the proof.

In the technical implementation, we work with three different polynomial representations in parallel. First, we use the Lagrange representation that is defined by evaluating the polynomial at a prescribed set of (rational) points. This representation is the easiest and most straightforward one to obtain, regardless in what form the polynomial is given. Second, we use the monomial representation in order to compute derivatives of polynomials without symbolic computation. Lastly, we use the Chebyshev representation, where the polynomial is expanded in a Chebyshev basis. The latter is crucial in order to compute what we call the *T*-norm, essentially the ℓ^1 -norm of the Chebyshev expansion coefficients. This norm can be computed exactly and acts as a substitute for the L^{∞} -norm. The *T*-norm together with explicit evaluation turns out to be sufficient to rigorously prove all the estimates we need. The linear maps that are used to switch between the three representations are provided explicitly and all of our computations can be verified straight away by using any programming language that supports fraction arithmetic.

2.6. Notation. To conclude the introduction, we finally explain some less standard notation that we prefer to use in the context of matrices. Let $A \in \mathbb{R}^{n \times n}$. We denote by A_k the k-th column vector and by A^j_k the element in the j-th row and the k-th column, e.g.

$$A = (A_1, A_2, A_3, A_4) = \begin{pmatrix} A^1_1 & A^1_2 & A^1_3 & A^1_4 \\ A^2_1 & A^2_2 & A^2_3 & A^2_4 \\ A^3_1 & A^3_2 & A^3_3 & A^3_4 \\ A^4_1 & A^4_2 & A^4_3 & A^4_4 \end{pmatrix}$$

for n = 4. Accordingly, δ is the unit matrix with elements δ^{j}_{k} given by the Kronecker symbol and δ_{k} is the k-th unit vector. The determinant det is the unique n-form on \mathbb{R}^{n} that satisfies $\det(\delta_{1}, \delta_{2}, \ldots, \delta_{n}) = 1$ and as usual, we write

$$\det(A) := \det(A_1, A_2, \dots, A_n)$$

The adjugate matrix adj(A) is defined by

$$\operatorname{adj}(A)^{j}_{k} := \operatorname{det}(A_{1}, \dots, A_{j-1}, \delta_{k}, A_{j+1}, \dots, A_{n}).$$

We have

$$\sum_{\ell=1}^{n} \operatorname{adj}(A)^{j}{}_{\ell}A^{\ell}{}_{k} = \sum_{\ell=1}^{n} \det(A_{1}, \dots, A_{j-1}, A^{\ell}{}_{k}\delta_{\ell}, A_{j+1}, \dots, A_{n})$$

= $\det(A_{1}, \dots, A_{j-1}, A_{k}, A_{j+1}, \dots, A_{n})$
= $\delta^{j}{}_{k} \det(A),$

which, if $det(A) \neq 0$, yields Cramer's rule

$$A^{-1} = \frac{\operatorname{adj}(A)}{\det(A)}.$$

Finally, we remark that for the sake of readability we have occasionally simplified the notation in this introductory section. In the technical part we often use more complicated symbols that carry more information. This is just as a warning in case the reader wants to quickly browse through the technical part, looking for a specific object that is mentioned in the introduction.

2.7. Structure of the paper. In Section 3, we develop the theoretical framework to construct a solution near a given perturbative solution. This is a purely theoretical part that makes no use of a specific polynomial approximation. In Section 4, we provide quantitative estimates for the operator norms in play, again on a purely theoretical basis without making reference to a particular approximation. In Section 5, we introduce our methods for estimating polynomials and rational functions. In Section 6, we define the concrete approximations and prove basic properties. In Section 7, we use the bounds from Section 4 to obtain explicit bounds on the operator norms. In Section 8, we prove bounds on the approximate linear operator. In Section 9, we prove the oscillatory estimates on the functional ψ_a . Finally, in Section 10, we complete the proof of Theorem 1.4. Appendix A contains the tables with the explicit coefficients of the polynomials we use throughout.

3. Theoretical framework

In this section we develop the theory for the construction of a solution to $\mathcal{R}(\alpha, f) = 0$ close to a given "approximate solution".

3.1. Function spaces and basic properties.

Definition 3.1. We define the norms

$$\|f\|_X := \sup_{y \in (-1,1)} (1-y^2) |f'(y)| + \|f\|_{L^{\infty}(-1,1)}$$
$$\|f\|_Y := \sup_{y \in (-1,1)} (1+y)(1-y)^2 |f(y)|.$$

Furthermore, we let X be the vector space of all $f \in C([-1,1]) \cap C^1(-1,1)$ such that $y \mapsto (1-y^2)f'(y)$ extends to a continuous function on [-1,1]. Similarly, we let Y be the vector space of all functions $f \in C(-1,1)$ such that $y \mapsto (1+y)(1-y)^2 f(y)$ extends to a continuous function on [-1,1].

Remark 3.2. Note that $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are Banach spaces. **Definition 3.3.** For $(\alpha, f) \in \mathbb{R} \setminus \{0\} \times C^2(-1, 1)$ and $y \in (-1, 1)$, we set

$$\mathcal{R}(\alpha, f)(y) := f''(y) + p_0(y, \alpha)f'(y) + q_0(y, \alpha)f(y) + \frac{f(y)|f(y)|^2}{(1-y)^2},$$

where

$$p_0(y,\alpha) := \frac{4i\alpha}{(1-y)^3} - \frac{2i\alpha}{(1-y)^2} - \frac{2 + \frac{2i}{\alpha}}{1-y} + \frac{2}{1+y}$$
$$q_0(y,\alpha) := -\frac{2 - 2i\alpha}{(1-y)^3} - \frac{\frac{1}{\alpha^2} - \frac{i}{\alpha}}{(1-y)^2} - \frac{1 + \frac{i}{\alpha}}{1-y} - \frac{1 + \frac{i}{\alpha}}{1+y}$$

Definition 3.4. Let $a_* \in \mathbb{R} \setminus \{0\}$ and $f_* \in C^2([-1,1] \times \mathbb{R} \setminus \{-a_*\})$. Then the pair (a_*, f_*) is called an *admissible approximation* if³

$$f'_*(1,a) = \frac{1 - i(a_* + a)}{2i(a_* + a)} f_*(1,a)$$

for all $a \in \mathbb{R} \setminus \{-a_*\}$. An admissible approximation (a_*, f_*) is called *trivial* if $f_* = 0$.

Lemma 3.5. Let (a_*, f_*) be an admissible approximation. Then we have $\mathcal{R}(a_* + a, f_*(\cdot, a)) \in Y$ for all $a \in \mathbb{R} \setminus \{-a_*\}$. Furthermore, the map

$$a \mapsto \mathcal{R}(a_* + a, f_*(\cdot, a)) : \mathbb{R} \setminus \{-a_*\} \to Y$$

is continuous.

Proof. Fix $a \in \mathbb{R} \setminus \{-a_*\}$. To begin with, we only have $\mathcal{R}(a_*+a, f_*(\cdot, a))(y) = O((1+y)^{-1}(1-y)^{-3})$ but

$$\lim_{y \to 1} (1-y)^3 \mathcal{R}(a_* + a, f_*(\cdot, a))(y) = 4i(a_* + a)f'_*(1, a) - [2 - 2i(a_* + a)]f_*(1, a) = 0$$

and $f_*(\cdot, a) \in C^2([-1, 1])$ show that in fact $\mathcal{R}(a_* + a, f_*(\cdot, a))(y) = O((1+y)^{-1}(1-y)^{-2})$. Furthermore, the map

$$(y,a) \mapsto (1+y)(1-y)^2 \mathcal{R}(a_*+a, f_*(\cdot, a))(y) : (-1,1) \times \mathbb{R} \setminus \{-a_*\} \to \mathbb{C}$$

extends continuously to $[-1,1] \times \mathbb{R} \setminus \{-a_*\}$ and is thus uniformly continuous with respect to y. \Box

3.2. Perturbative decomposition.

Definition 3.6. Let (a_*, f_*) be an admissible approximation. For $(a, f) \in \mathbb{R} \setminus \{-a_*\} \times C^2(-1, 1)$ and $y \in (-1, 1)$, we set

$$\mathcal{L}(a_*, f_*, a, f)(y) := f''(y) + p_0(y, a_* + a)f'(y) + q_0(y, a_* + a)f(y) + \frac{2|f_*(y, a)|^2}{(1-y)^2}f(y) + \frac{f_*(y, a)^2}{(1-y)^2}\overline{f(y)}$$

and

$$\mathcal{N}(a_*, f_*, a, f)(y) := \frac{\overline{f_*(y, a)}f(y)^2 + 2f_*(y, a)|f(y)|^2 + f(y)|f(y)|^2}{(1-y)^2}$$

Lemma 3.7 (Perturbative decomposition). Let (a_*, f_*) be an admissible approximation and $(a, f) \in \mathbb{R} \setminus \{-a_*\} \times C^2(-1, 1)$. Then we have

$$\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = \mathcal{L}(a_*, f_*, a, f) + \mathcal{N}(a_*, f_*, a, f) + \mathcal{R}(a_* + a, f_*(\cdot, a)).$$

Proof. This follows immediately by inserting the definitions.

³A prime denotes the derivative with respect to the first variable, i.e., $f'_*(y,a) := \partial_y f_*(y,a)$.

3.3. The admissible right fundamental matrix. We construct an approximate inverse of the operator $f \mapsto \mathcal{L}(a_*, f_*, a, f)$.

Definition 3.8. We define $\phi : \mathbb{C} \to \mathbb{R}^{2 \times 2}$ by

$$\phi(z) := \begin{pmatrix} \operatorname{Re} z & -\operatorname{Im} z \\ \operatorname{Im} z & \operatorname{Re} z \end{pmatrix}$$

Definition 3.9. Let $f : \mathbb{C} \to \mathbb{C}$. We write $f(y) = \mathcal{P}(y)$ if f is a polynomial.

Definition 3.10. For $\alpha \in \mathbb{R} \setminus \{0\}$ and $y \in (-1, 1)$, we set

$$\varphi(y, \alpha) := -\frac{2\alpha}{(1-y)^2} + \frac{2\alpha}{1-y} - \frac{2}{\alpha}\log(1-y).$$

Definition 3.11. For $a_* \in \mathbb{R} \setminus \{0\}$ let $J \subset \mathbb{R} \setminus \{-a_*\}$ be compact. Furthermore, let $f_1, f_2, f_3, f_4 : [0,1) \times J \to \mathbb{C}$ and set

$$F(y,a) := \begin{pmatrix} \operatorname{Re} f_1(y,a) & \operatorname{Re} f_2(y,a) & \operatorname{Re} f_3(y,a) & \operatorname{Re} f_4(y,a) \\ \operatorname{Im} f_1(y,a) & \operatorname{Im} f_2(y,a) & \operatorname{Im} f_3(y,a) & \operatorname{Im} f_4(y,a) \\ \operatorname{Re} f_1'(y,a) & \operatorname{Re} f_2'(y,a) & \operatorname{Re} f_3'(y,a) & \operatorname{Re} f_4'(y,a) \\ \operatorname{Im} f_1'(y,a) & \operatorname{Im} f_2'(y,a) & \operatorname{Im} f_3'(y,a) & \operatorname{Im} f_4'(y,a) \end{pmatrix}.$$

Then F is called an *admissible right fundamental matrix on* J if $det(F(y, a)) \neq 0$ for all $(y, a) \in [0, 1) \times J$ and

$$\begin{split} f_1(y,a) &= 1 + \frac{a_* + a + i}{2(a_* + a)}(1 - y) + (1 - y)^2 \mathcal{P}(y) \\ f_2(y,a) &= i + i\frac{a_* + a + i}{2(a_* + a)}(1 - y) + (1 - y)^2 \mathcal{P}(y) \\ f_3(y,a) &= e^{i\varphi(y,a_* + a)}(1 - y) \left[1 + \frac{2(a_* + a) - i}{2(a_* + a)}(1 - y) + (1 - y)^2 \mathcal{P}(y) \right] \\ &+ e^{-i\varphi(y,a_* + a)}(1 - y)^5 \mathcal{P}(y) \\ f_4(y,a) &= ie^{i\varphi(y,a_* + a)}(1 - y) \left[1 + \frac{2(a_* + a) - i}{2(a_* + a)}(1 - y) + (1 - y)^2 \mathcal{P}(y) \right] \\ &+ ie^{-i\varphi(y,a_* + a)}(1 - y)^5 \mathcal{P}(y). \end{split}$$

The number a_* is called the *parameter of* F.

Remark 3.12. Observe that $f_j : [0,1) \times J \to \mathbb{C}$ for $j \in \{1,2,3,4\}$ extends continuously to the compact domain $[0,1] \times J$ and thus, f_j is uniformly continuous.

Lemma 3.13. Let F be an admissible right fundamental matrix on J with parameter a_* and for $j \in \{1, 2, 3, 4\}$ set $f_j := F_j^1 + iF_j^2$. Then there exists a $\delta \in (0, 1]$ such that

$$F(y,a)^{-1} = \begin{pmatrix} \phi(\frac{f'_3(y,a)}{W(y,a)}) & -\phi(\frac{f_3(y,a)}{W(y,a)}) \\ -\phi(\frac{f'_1(y,a)}{W(y,a)}) & \phi(\frac{f_1(y,a)}{W(y,a)}) \end{pmatrix} [1 + O((1-y)^2 a^0)]$$

for all $(y, a) \in [1 - \delta, 1) \times J$, where

$$W(y,a) := f_1(y,a)f'_3(y,a) - f'_1(y,a)f_3(y,a)$$

= $-\frac{4i(a_*+a)}{(1-y)^2}e^{i\varphi(y,a_*+a)}\left[1 + (1-y) + O((1-y)^2a^0)\right].$

Remark 3.14. More precisely, we mean that

$$F(y,a)^{-1} = \begin{pmatrix} \phi(\frac{f'_3(y,a)}{W(y,a)})[1+O((1-y)^2a^0)] & -\phi(\frac{f_3(y,a)}{W(y,a)})[1+O((1-y)^2a^0)] \\ -\phi(\frac{f'_1(y,a)}{W(y,a)})[1+O((1-y)^2a^0)] & \phi(\frac{f_1(y,a)}{W(y,a)})[1+O((1-y)^2a^0)] \end{pmatrix}$$

but here and in the following, we will use the sloppier but more compact notation of Lemma 3.13. *Proof of Lemma 3.13.* We commence with the statement on the Wronskian W. Note that $f'_1(y, a) = O(y^0 a^0)$ and

$$f'_{3}(y,a) = i\varphi'(y,a_{*}+a)e^{i\varphi(y,a_{*}+a)}(1-y)\left[1 + \frac{2(a_{*}+a)-i}{2(a_{*}+a)}(1-y) + O((1-y)^{2}a^{0})\right] + O(y^{0}a^{0}).$$

Thus, since

$$\varphi'(y, a_* + a) = -\frac{4(a_* + a)}{(1 - y)^3} + \frac{2(a_* + a)}{(1 - y)^2} + \frac{2}{(a_* + a)(1 - y)}$$
$$= -\frac{4(a_* + a)}{(1 - y)^3} \left[1 - \frac{1}{2}(1 - y) + O((1 - y)^2 a^0)\right],$$

we obtain

$$\begin{split} f_3'(y,a) &= -\frac{4i(a_*+a)}{(1-y)^2} e^{i\varphi(y,a_*+a)} \left[1 - \frac{1}{2}(1-y) + O((1-y)^2 a^0) \right] \\ &\times \left[1 + \frac{2(a_*+a) - i}{2(a_*+a)} (1-y) + O((1-y)^2 a^0) \right] \\ &= -\frac{4i(a_*+a)}{(1-y)^2} e^{i\varphi(y,a_*+a)} \left[1 + \frac{a_*+a-i}{2(a_*+a)} (1-y) + O((1-y)^2 a^0) \right]. \end{split}$$

Consequently,

$$\begin{split} W(y,a) &= f_1(y,a) f'_3(y,a) + O((1-y)a^0) \\ &= -\frac{4i(a_*+a)}{(1-y)^2} e^{i\varphi(y,a_*+a)} \left[1 + \frac{a_*+a+i}{2(a_*+a)}(1-y) + O((1-y)^2a^0) \right] \\ &\times \left[1 + \frac{a_*+a-i}{2(a_*+a)}(1-y) + O((1-y)^2a^0) \right] \\ &= -\frac{4i(a_*+a)}{(1-y)^2} e^{i\varphi(y,a_*+a)} \left[1 + (1-y) + O((1-y)^2a^0) \right], \end{split}$$

as claimed. Furthermore, it follows that we can find a $\delta \in (0,1]$ such that $W(y,a) \neq 0$ for all $(y,a) \in [1-\delta,1) \times J$.

Now observe that

$$f_2(y,a) = if_1(y,a) + O((1-y)^2)$$

$$f'_2(y,a) = if'_1(y,a) + O(1-y)$$

$$f_4(y,a) = if_3(y,a) + O((1-y)^3a^0)$$

$$f'_4(y,a) = if'_3(y,a) + O(y^0a^0)$$

and thus,

$$F(y,a) = \begin{pmatrix} \phi(f_1(y,a)) & \phi(f_3(y,a)) \\ \phi(f'_1(y,a)) & \phi(f'_3(y,a)) \end{pmatrix} + \begin{pmatrix} 0 & O((1-y)^2) & 0 & O((1-y)^3 a^0) \\ 0 & O((1-y)^2) & 0 & O((1-y)^3 a^0) \\ 0 & O(1-y) & 0 & O(y^0 a^0) \\ 0 & O(1-y) & 0 & O(y^0 a^0) \end{pmatrix}.$$

Since $\phi(z_1z_2) = \phi(z_1)\phi(z_2)$ for all $z_1, z_2 \in \mathbb{C}$, the first matrix can be formally inverted like a 2×2 -matrix and we obtain

$$\begin{split} F(y,a) &= \begin{pmatrix} \phi(f_1(y,a)) & \phi(f_3(y,a)) \\ \phi(f_1'(y,a)) & \phi(f_3'(y,a)) \end{pmatrix} \\ &\times \left[1 + \begin{pmatrix} \phi(\frac{f_3'(y,a)}{W(y,a)}) & -\phi(\frac{f_3(y,a)}{W(y,a)}) \\ -\phi(\frac{f_1'(y,a)}{W(y,a)}) & \phi(\frac{f_1(y,a)}{W(y,a)}) \end{pmatrix} \begin{pmatrix} 0 & O((1-y)^2) & 0 & O((1-y)^3 a^0) \\ 0 & O((1-y)^2) & 0 & O((1-y)^3 a^0) \\ 0 & O(1-y) & 0 & O(y^0 a^0) \\ 0 & O(1-y) & 0 & O(y^0 a^0) \end{pmatrix} \right] \\ &= \begin{pmatrix} \phi(f_1(y,a)) & \phi(f_3(y,a)) \\ \phi(f_1'(y,a)) & \phi(f_3'(y,a)) \end{pmatrix} \left[1 + O((1-y)^2 a^0) \right] \end{split}$$

for all $(y,a) \in [1-\delta,1) \times J$. This yields the stated form of F^{-1} .

3.4. The admissible left fundamental matrix.

Definition 3.15. Let $f_1, f_2, f_3, f_4 : (-1, 0] \to \mathbb{C}$ and set

$$F := \begin{pmatrix} \operatorname{Re} f_1 & \operatorname{Re} f_2 & \operatorname{Re} f_3 & \operatorname{Re} f_4 \\ \operatorname{Im} f_1 & \operatorname{Im} f_2 & \operatorname{Im} f_3 & \operatorname{Im} f_4 \\ \operatorname{Re} f'_1 & \operatorname{Re} f'_2 & \operatorname{Re} f'_3 & \operatorname{Re} f'_4 \\ \operatorname{Im} f'_1 & \operatorname{Im} f'_2 & \operatorname{Im} f'_3 & \operatorname{Im} f'_4 \end{pmatrix}$$

Then F is called an *admissible left fundamental matrix* if $det(F(y)) \neq 0$ for all $y \in (-1, 0]$ and

$$f_1(y) = 1 + (1+y)\mathcal{P}(y) \qquad f_2(y) = i + (1+y)\mathcal{P}(y)$$

$$f_3(y) = \frac{1}{1+y}[1 + (1+y)\mathcal{P}(y)] \qquad f_4(y) = \frac{i}{1+y}[1 + (1+y)\mathcal{P}(y)].$$

Lemma 3.16. Let F be an admissible left fundamental matrix. Then we have

$$F(y)^{-1} = \begin{pmatrix} 1 & O(1+y) & 1+y & O((1+y)^2) \\ O(1+y) & 1 & O((1+y)^2) & 1+y \\ O((1+y)^2) & O((1+y)^2) & -(1+y)^2 & O((1+y)^3) \\ O((1+y)^2) & O((1+y)^2) & O((1+y)^3) & -(1+y)^2 \end{pmatrix} [1+O(1+y)].$$

Proof. For $j \in \{1, 2, 3, 4\}$, set $f_j := F^1{}_j + iF^2{}_j$. By definition, we have

$$f_1(y) = 1 + O(1+y) \qquad f_3(y) = \frac{1}{1+y} [1 + O(1+y)]$$

$$f'_1(y) = O(y^0) \qquad f'_3(y) = -\frac{1}{(1+y)^2} [1 + O(1+y)]$$

and thus,

$$W(y) := f_1(y)f'_3(y) - f'_1(y)f_3(y) = -\frac{1}{(1+y)^2}[1+O(1+y)].$$

As in the proof of Lemma 3.13, we have

$$\begin{split} F(y) &= \begin{pmatrix} \phi(f_1(y)) & \phi(f_3(y)) \\ \phi(f'_1(y)) & \phi(f'_3(y)) \end{pmatrix} \\ &\times \left[1 + \begin{pmatrix} \phi(\frac{f'_3(y)}{W(y)}) & -\phi(\frac{f_3(y)}{W(y)}) \\ -\phi(\frac{f'_1(y)}{W(y)}) & \phi(\frac{f_1(y)}{W(y)}) \end{pmatrix} \begin{pmatrix} 0 & O(1+y) & 0 & O(1+y) \\ 0 & O(1+y) & 0 & O(1+y) \\ 0 & O(y^0) & 0 & O(y^0) \\ 0 & O(y^0) & 0 & O(y^0) \end{pmatrix} \right], \end{split}$$

provided that y is sufficiently close to -1 and thus,

$$F(y)^{-1} = \begin{pmatrix} \phi(\frac{f'_3(y)}{W(y)}) & -\phi(\frac{f_3(y)}{W(y)}) \\ -\phi(\frac{f_1(y)}{W(y)}) & \phi(\frac{f_1(y)}{W(y)}) \end{pmatrix} [1 + O(1+y)]$$

This implies the claim.

3.5. The global fundamental matrix.

Definition 3.17. Let F_R be an admissible right fundamental matrix on J and let F_L be an admissible left fundamental matrix. Then

$$F(y,a) := 1_{(-1,0]}(y)F_L(y)F_L(0)^{-1}F_R(0,a) + 1_{(0,1)}(y)F_R(y,a)$$

is called an *admissible fundamental matrix on* J if

$$[F_L(0)^{-1}F_R(0,a)]^3_1 \neq 0$$

for all $a \in J$. Furthermore, F_L and F_R are called the *left part of* F and the *right part of* F, respectively. Finally, the *parameter of* F is defined as the parameter of F_R .

Remark 3.18. The condition $[F_L(0)^{-1}F_R(0,a)]^3_1 \neq 0$ ensures that the function $F^1_1 + iF^2_1$ is singular at the left endpoint -1. This will be crucial.

Definition 3.19. Let F be an admissible fundamental matrix on J. Then we define $M_F: J \to \mathbb{R}^{4 \times 4}$ by

$$M_F(a) := F_L(0)^{-1} F_R(0, a)$$

where F_L is the left part of F and F_R is the right part of F.

Definition 3.20. Let *F* be an admissible fundamental matrix on *J*. Then the functions (p_1, p_2, q_1, q_2) , defined on $(-1, 1) \setminus \{0\} \times J$ by

$$p_1 := \frac{1}{2}(A^3_3 + A^4_4) + \frac{i}{2}(A^4_3 - A^3_4), \qquad p_2 := \frac{1}{2}(A^3_3 - A^4_4) + \frac{i}{2}(A^4_3 + A^3_4), q_1 := \frac{1}{2}(A^3_1 + A^4_2) + \frac{i}{2}(A^4_1 - A^3_2), \qquad q_2 := \frac{1}{2}(A^3_1 - A^4_2) + \frac{i}{2}(A^4_1 + A^3_2),$$

where we abbreviate $A(y,a) = -\partial_y F(y,a)F(y,a)^{-1}$, are called the *coefficients associated to F*.

Proposition 3.21. Let F be an admissible fundamental matrix on J with parameter a_* and let

$$f_j(y,a) := F(y,a)^1{}_j + iF(y,a)^2{}_j$$

for $j \in \{1, 2, 3, 4\}$, $y \in (-1, 1)$, and $a \in J$. Then $f_j(\cdot, a) \in C^2((-1, 1) \setminus \{0\}) \cap C^1(-1, 1)$ for any $a \in J$ and

$$f_j''(y,a) + p_1(y,a)f_j'(y,a) + p_2(y,a)\overline{f_j'(y,a)} + q_1(y,a)f_j(y,a) + q_2(y,a)\overline{f_j(y,a)} = 0$$
15

for all $(y, a) \in (-1, 1) \setminus \{0\} \times J$ and $j \in \{1, 2, 3, 4\}$, where (p_1, p_2, q_1, q_2) are the coefficients associated to F. Furthermore,

$$p_1(y,a) = \frac{4i(a_*+a)}{(1-y)^3} - \frac{2i(a_*+a)}{(1-y)^2} + \frac{2}{1+y} + O((1-y)^{-1}a^0)$$

$$p_2(y,a) = O((1-y)^{-1}a^0)$$

$$q_1(y,a) = -\frac{2-2i(a_*+a)}{(1-y)^3} + O((1-y)^{-2}a^0) + O((1+y)^{-1})$$

$$q_2(y,a) = O((1-y)^{-2}a^0) + O((1+y)^{-1}).$$

Proof. Let $a \in J$ and denote by F_L and F_R the left part and the right part of F, respectively. By definition, we have

$$F(y,a) = \begin{pmatrix} \operatorname{Re} f_1(y,a) & \operatorname{Re} f_2(y,a) & \operatorname{Re} f_3(y,a) & \operatorname{Re} f_4(y,a) \\ \operatorname{Im} f_1(y,a) & \operatorname{Im} f_2(y,a) & \operatorname{Im} f_3(y,a) & \operatorname{Im} f_4(y,a) \\ \operatorname{Re} f_1'(y,a) & \operatorname{Re} f_2'(y,a) & \operatorname{Re} f_3'(y,a) & \operatorname{Re} f_4'(y,a) \\ \operatorname{Im} f_1'(y,a) & \operatorname{Im} f_2'(y,a) & \operatorname{Im} f_3'(y,a) & \operatorname{Im} f_4'(y,a) \end{pmatrix}$$

for all $y \in (-1,1) \setminus \{0\}$ and $F(\cdot,a) \in C((-1,1), \mathbb{R}^{4\times 4})$. Moreover, $F_L \in C^2((-1,0), \mathbb{R}^{4\times 4})$ and $F(\cdot,a) \in C^1((0,1), \mathbb{R}^{4\times 4})$. Consequently, $f_j(\cdot,a) \in C^2((-1,1) \setminus \{0\}) \cap C^1(-1,1)$ for all $j \in \{1,2,3,4\}$. Let $A(y,a) := -F'(y,a)F(y,a)^{-1}$. Then we have

$$F'(y,a) + A(y,a)F(y,a) = 0$$

Since $F'(y,a)^j = F(y,a)^{j+2}$ for $j \in \{1,2\}$, it follows that

$$A(y,a)^{j}{}_{k} = -\sum_{\ell=1}^{4} F'(y,a)^{j}{}_{\ell}F^{-1}(y,a)^{\ell}{}_{k} = -\sum_{\ell=1}^{4} F(y,a)^{j+2}{}_{\ell}F^{-1}(y,a)^{\ell}{}_{k} = -\delta^{j+2}{}_{k}F^{-1}(y,a)^{\ell}{}_{k} = -\delta^{j+2}{}_{k}F^$$

and thus, A is of the form

$$A = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ A_{1}^{3} & A_{2}^{3} & A_{3}^{3} & A_{4}^{3} \\ A_{1}^{4} & A_{2}^{4} & A_{3}^{4} & A_{4}^{4} \end{pmatrix}$$

Consequently, F'(y, a) + A(y, a)F(y, a) = 0 is equivalent to

$$f_j''(y,a) + p_1(y,a)f_j'(y,a) + p_2(y,a)\overline{f_j'(y,a)} + q_1(y,a)f_j(y,a) + q_2(y,a)\overline{f_j(y,a)} = 0$$

for $j \in \{1, 2, 3, 4\}$, where

$$p_{1} = \frac{1}{2}(A^{3}_{3} + A^{4}_{4}) + \frac{i}{2}(A^{4}_{3} - A^{3}_{4}), \qquad p_{2} = \frac{1}{2}(A^{3}_{3} - A^{4}_{4}) + \frac{i}{2}(A^{4}_{3} + A^{3}_{4}),$$

$$q_{1} = \frac{1}{2}(A^{3}_{1} + A^{4}_{2}) + \frac{i}{2}(A^{4}_{1} - A^{3}_{2}), \qquad q_{2} = \frac{1}{2}(A^{3}_{1} - A^{4}_{2}) + \frac{i}{2}(A^{4}_{1} + A^{3}_{2}).$$

If $y \in (-1, 0)$, we have

$$A(y,a) = -F'(y,a)F(y,a)^{-1} = -F'_L(y)F_L(0)^{-1}F_R(0,a)F_R(0,a)^{-1}F_L(0)F_L(y)^{-1}$$

= $-F'_L(y)F_L(y)^{-1}$

and thus, by definition and Lemma 3.16,

$$\begin{split} A(y,a) &= - \begin{pmatrix} O(y^0) & O(y^0) & -(1+y)^{-2}[1+O(1+y)] & O((1+y)^{-1}) \\ O(y^0) & O(y^0) & O((1+y)^{-1}) & -(1+y)^{-2}[1+O(1+y)] \\ O(y^0) & O(y^0) & 2(1+y)^{-3}[1+O(1+y)] & O((1+y)^{-2}) \\ O(y^0) & O(y^0) & O((1+y)^{-2}) & 2(1+y)^{-3}[1+O(1+y)] \end{pmatrix} \\ &\times \begin{pmatrix} 1 & O(1+y) & 1+y & O((1+y)^2) \\ O(1+y) & 1 & O((1+y)^2) & 1+y \\ O((1+y)^2) & O((1+y)^2) & -(1+y)^2 & O((1+y)^3) \\ O((1+y)^2) & O((1+y)^2) & O((1+y)^3) & -(1+y)^2 \end{pmatrix} \begin{bmatrix} 1+O(1+y) \end{bmatrix} \\ &= \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ O((1+y)^{-1}) & O((1+y)^{-1}) & 2(1+y)^{-1}[1+O(1+y)] \\ O((1+y)^{-1}) & O((1+y)^{-1}) & O(y^0) \\ O((1+y)^{-1}[1+O(1+y)] \end{pmatrix} , \end{split}$$

which yields the claimed asymptotics of the coefficients on the left-hand side. In the case $y \in (0, 1)$, we have $A(y, a) = -F'_R(y, a)F_R(y, a)^{-1}$ and by definition,

$$f_2(y,a) = if_1(y,a) + O((1-y)^2)$$

$$f'_2(y,a) = if'_1(y,a) + O(1-y)$$

$$f''_2(y,a) = if''_1(y,a) + O(y^0),$$

as well as

$$f_4(y,a) = if_3(y,a) + O((1-y)^3 a^0)$$

$$f'_4(y,a) = if'_3(y,a) + O(y^0 a^0)$$

$$f''_4(y,a) = if''_3(y,a) + O((1-y)^{-3} a^0).$$

Consequently,

$$F_{R}'(y,a) = \begin{pmatrix} \phi(f_{1}'(y,a)) & \phi(f_{3}'(y,a)) \\ \phi(f_{1}''(y,a)) & \phi(f_{3}''(y,a)) \end{pmatrix} + \begin{pmatrix} 0 & O(1-y) & 0 & O(y^{0}a^{0}) \\ 0 & O(1-y) & 0 & O(y^{0}a^{0}) \\ 0 & O(y^{0}) & 0 & O((1-y)^{-3}a^{0}) \\ 0 & O(y^{0}) & 0 & O((1-y)^{-3}a^{0}) \end{pmatrix}$$

and, in view of Lemma 3.13,

$$\begin{split} A(y,a) &= -F'(y,a)F(y,a)^{-1} \\ &= F'_R(y,a) \begin{pmatrix} -\phi(\frac{f'_3(y,a)}{W(y,a)}) & \phi(\frac{f_3(y,a)}{W(y,a)}) \\ \phi(\frac{f'_1(y,a)}{W(y,a)}) & -\phi(\frac{f_1(y,a)}{W(y,a)}) \end{pmatrix} [1 + O((1-y)^2 a^0)] \\ &= \begin{pmatrix} \phi(0) & -\phi(1) \\ \phi(\frac{f'_1(y,a)f''_3(y,a) - f''_1(y,a)f'_3(y,a)}{W(y,a)}) & -\phi(\frac{f_1(y,a)f''_3(y,a) - f''_1(y,a)f_3(y,a)}{W(y,a)}) \end{pmatrix} [1 + O((1-y)^2 a^0)] \\ &+ \begin{pmatrix} O((1-y)a^0) & O((1-y)a^0) & O((1-y)^2 a^0) & O((1-y)^2 a^0) \\ O((1-y)a^0) & O((1-y)a^0) & O((1-y)^2 a^0) & O((1-y)^2 a^0) \\ O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) \end{pmatrix} . \end{split}$$

In other words,

$$\begin{split} A(y,a) &= \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ \operatorname{Re}h_1(y,a) &-\operatorname{Im}h_1(y,a) & \operatorname{Re}h_2(y,a) &-\operatorname{Im}h_2(y,a) \\ \operatorname{Im}h_1(y,a) & \operatorname{Re}h_1(y,a) & \operatorname{Im}h_2(y,a) & \operatorname{Re}h_2(y,a) \end{pmatrix} [1 + O((1-y)^2 a^0)] \\ &+ \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) \\ O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) & O((1-y)^{-1}a^0) \end{pmatrix} \\ \end{split}$$

with

$$h_1(y,a) := \frac{f'_1(y,a)f''_3(y,a) - f''_1(y,a)f'_3(y,a)}{W(y,a)}$$
$$h_2(y,a) := -\frac{f_1(y,a)f''_3(y,a) - f''_1(y,a)f_3(y,a)}{W(y,a)}.$$

From Lemma 3.13 we recall that

$$W(y,a) = -\frac{4i(a_*+a)}{(1-y)^2} e^{i\varphi(y,a_*+a)} \left[1 + (1-y) + O((1-y)^2 a^0)\right]$$

and by definition, we have $f'_3(y,a) = O((1-y)^{-2}a^0)$ as well as

$$f_3''(y,a) = -\varphi'(y,a)^2 e^{i\varphi(y,a_*+a)}(1-y) \left[1 + \frac{2(a_*+a)-i}{2(a_*+a)}(1-y) + O((1-y)^2 a^0) \right] + O((1-y)^{-3}a^0).$$

Note that

$$\varphi'(y,a)^2 = \left[-\frac{4(a_*+a)}{(1-y)^3} + \frac{2(a_*+a)}{(1-y)^2} + \frac{2}{(a_*+a)(1-y)} \right]^2$$
$$= \frac{16(a_*+a)^2}{(1-y)^6} - \frac{16(a_*+a)^2}{(1-y)^5} + O((1-y)^{-4}a^0)$$
$$= \frac{16(a_*+a)^2}{(1-y)^6} \left[1 - (1-y) + O((1-y)^2a^0) \right]$$

and thus,

$$\begin{split} f_3''(y,a) &= -\frac{16(a_*+a)^2}{(1-y)^5} e^{i\varphi(y,a_*+a)} \left[1 - (1-y) + O((1-y)^2 a^0) \right] \\ &\times \left[1 + \frac{2(a_*+a) - i}{2(a_*+a)} (1-y) + O((1-y)^2 a^0) \right] \\ &= -\frac{16(a_*+a)^2}{(1-y)^5} e^{i\varphi(y,a_*+a)} \left[1 - \frac{i}{2(a_*+a)} (1-y) + O((1-y)^2 a^0) \right]. \end{split}$$

This yields

$$\frac{f_3''(y,a)}{W(y,a)} = -\frac{4i(a_*+a)}{(1-y)^3} \left[1 - \frac{i}{2(a_*+a)}(1-y) + O((1-y)^2 a^0) \right] \left[1 - (1-y) + O((1-y)^2 a^0) \right]$$
$$= -\frac{4i(a_*+a)}{(1-y)^3} \left[1 - \frac{2(a_*+a)+i}{2(a_*+a)}(1-y) + O((1-y)^2 a^0) \right]$$

and by recalling that $f_1(y,a) = 1 + \frac{a_* + a + i}{2(a_* + a)}(1-y) + (1-y)^2 \mathcal{P}(y)$, we obtain

$$h_1(y,a) = f_1'(y,a)\frac{f_3''(y,a)}{W(y,a)} + O((1-y)^0 a^0) = \left[-\frac{a_* + a + i}{2(a_* + a)} + O(1-y)\right]\frac{f_3''(y,a)}{W(y,a)} + O((1-y)^0 a^0)$$
$$= -\frac{2 - 2i(a_* + a)}{(1-y)^3}[1 + O((1-y)a^0)]$$

as well as

$$h_{2}(y,a) = -f_{1}(y,a)\frac{f_{3}''(y,a)}{W(y,a)} + O((1-y)^{3}a^{0})$$

$$= \frac{4i(a_{*}+a)}{(1-y)^{3}} \left[1 + \frac{a_{*}+a+i}{2(a_{*}+a)}(1-y) + O((1-y)^{2}a^{0}) \right]$$

$$\times \left[1 - \frac{2(a_{*}+a)+i}{2(a_{*}+a)}(1-y) + O((1-y)^{2}a^{0}) \right]$$

$$= \frac{4i(a_{*}+a)}{(1-y)^{3}} \left[1 - \frac{1}{2}(1-y) + O((1-y)^{2}a^{0}) \right].$$

Consequently,

$$p_1(y,a) = \frac{1}{2}(A(y,a)^3 + A(y,a)^4) + \frac{i}{2}(A(y,a)^4 - A(y,a)^3)$$
$$= h_2(y,a)[1 + O((1-y)^2a^0)] + O((1-y)^{-1}a^0)$$
$$= \frac{4i(a_*+a)}{(1-y)^3} - \frac{2i(a_*+a)}{(1-y)^2} + O((1-y)^{-1}a^0)$$

and

$$p_2(y,a) = \frac{1}{2}(A(y,a)^3 - A(y,a)^4) + \frac{i}{2}(A(y,a)^4 + A(y,a)^3)$$

= $h_2(y,a)O((1-y)^2a^0) + O((1-y)^{-1}a^0)$
= $O((1-y)^{-1}a^0),$

as well as

$$q_1(y,a) = \frac{1}{2}(A(y,a)^3 + A(y,a)^4) + \frac{i}{2}(A(y,a)^4 - A(y,a)^3)$$

= $h_1(y,a)[1 + O((1-y)^2a^0)] + O((1-y)^{-1}a^0)$
= $-\frac{2 - 2i(a_* + a)}{(1-y)^3} + O((1-y)^{-2}a^0)$

and

$$q_2(y,a) = \frac{1}{2}(A(y,a)^3_1 - A(y,a)^4_2) + \frac{i}{2}(A(y,a)^4_1 + A(y,a)^3_2)$$

= $h_1(y,a)O((1-y)^2a^0) + O((1-y)^{-1}a^0)$
= $O((1-y)^{-1}a^0).$

Definition 3.22. Let F be an admissible fundamental matrix on J and let (p_1, p_2, q_1, q_2) be the associated coefficients. Furthermore, let $I \subset (-1, 1)$ be open. Then, for $f \in C^2(I)$ and $(y, a) \in I \times J$, we set

$$\mathcal{L}_F(a,f)(y) := f''(y) + p_1(y,a)f'(y) + p_2(y,a)\overline{f'(y)} + q_1(y,a)f(y) + q_2(y,a)\overline{f(y)} + q_2(y$$

3.6. The inhomogeneous equation.

Definition 3.23. Let F be an admissible fundamental matrix on J. Then, for $(y, a) \in (-1, 1) \times J$, $k \in \{1, 2, 3, 4\}$, and $g \in Y$, we set

$$\alpha_F^k(a,g)(y) := F^{-1}(y,a)^k{}_3\operatorname{Re} g(y) + F^{-1}(y,a)^k{}_4\operatorname{Im} g(y)$$

Lemma 3.24. Let F be an admissible fundamental matrix on J. Then we have the bound

$$\left\|\alpha_F^k(a,g)\right\|_{L^{\infty}(-1,1)} \lesssim \|g\|_{Y}$$

for all $(a, g) \in J \times Y$ and $k \in \{1, 2, 3, 4\}$.

Proof. From Lemma 3.13 we have

$$\left|F^{-1}(y,a)^{k}_{3}\right| + \left|F^{-1}(y,a)^{k}_{4}\right| \lesssim (1-y)^{2}$$

for all $(y, a) \in [0, 1) \times J$. Furthermore, if $y \in (-1, 0]$,

$$F^{-1}(y,a)^{k}{}_{\ell} = \sum_{m=1}^{4} M_{F}^{-1}(a)^{k}{}_{m}F_{L}^{-1}(y)^{m}{}_{\ell}$$

and Lemma 3.16 yields

$$\left|F^{-1}(y,a)^{k}{}_{3}\right| + \left|F^{-1}(y,a)^{k}{}_{4}\right| \lesssim 1 + y$$

for all $(y,a) \in (-1,0] \times J$.

Definition 3.25. Let F be an admissible fundamental matrix on J. Then, for $(y, a) \in (-1, 1) \times J$ and $g \in Y$, we set

$$\begin{split} \mathcal{L}_{F}^{-1}(a,g)(y) &:= \sum_{k=1}^{2} \left[F(y,a)^{1}{}_{k} + iF(y,a)^{2}{}_{k} \right] \int_{-1}^{y} \alpha_{F}^{k}(a,g)(x) dx \\ &- \sum_{k=3}^{4} \left[F(y,a)^{1}{}_{k} + iF(y,a)^{2}{}_{k} \right] \int_{y}^{1} \alpha_{F}^{k}(a,g)(x) dx \\ &+ \left[F(y,a)^{1}{}_{1} + iF(y,a)^{2}{}_{1} \right] \sum_{k=3}^{4} \frac{M_{F}(a)^{3}{}_{k}}{M_{F}(a)^{3}{}_{1}} \int_{-1}^{1} \alpha_{F}^{k}(a,g)(x) dx \end{split}$$

Remark 3.26. The first two terms in the definition of \mathcal{L}_F^{-1} are straightforward from the variation of constants formula. However, the last term requires some explanation. Without this term we would have to satisfy two real vanishing conditions in order to ensure that $\mathcal{L}_F^{-1}(a,g)$ is regular at y = -1. However, we only have the single real parameter a to vary. Thus, we need to cancel one of the singular terms near y = -1 by some other means. This is provided by the last term in the definition of \mathcal{L}_F^{-1} , see below. Indeed, the condition $M_F(a)^{3}_1 \neq 0$ ensures that the function $F^{1}_{1} + iF^{2}_{1}$ (which is regular at y = 1) is singular at y = -1 and can therefore be used to cancel one of the singular terms. The remaining singular term will then give rise to a single real vanishing condition that will be satisfied by varying the parameter a. This will determine the value of a.

At this point the reader might wonder why it is not possible to cancel both singular contributions by such a construction, without getting a condition on a. This would require the function $F_2^1+iF_2^2$ to be singular at y = -1. However, there is no condition in Definition 3.17 that would ensure this. In other words, \mathcal{L}_F is allowed to have a one-dimensional kernel and this is necessary because of the phase invariance of the problem. **Lemma 3.27.** Let F be an admissible fundamental matrix on J and $g \in Y$. Then $\mathcal{L}_F^{-1}(a,g) \in C^2((-1,1) \setminus \{0\}) \cap C^1(-1,1)$ and

$$\mathcal{L}_F\left(a, \mathcal{L}_F^{-1}(a, g)\right)(y) = g(y)$$

for all $(y, a) \in (-1, 1) \setminus \{0\} \times J$.

Proof. For $j \in \{1, 2, 3, 4\}$, we set

$$f^{j}(y,a) := \sum_{k=1}^{4} F(y,a)^{j}{}_{k} \int_{y_{k}}^{y} \left[F^{-1}(x,a)^{k}{}_{3}\operatorname{Re} g(x) + F^{-1}(x,a)^{k}{}_{4}\operatorname{Im} g(x) \right] dx + c(a,g)F(y,a)^{j}{}_{1},$$

where $(y_1, y_2, y_3, y_4) = (-1, -1, 1, 1)$ and

$$c(a,g) := \sum_{k=3}^{4} \frac{M_F(a)^3{}_k}{M_F(a)^3{}_1} \int_{-1}^{1} \alpha_F^k(a,g)(x) dx.$$

Then we have

$$\partial_{y}f^{j}(y,a) = \sum_{k=1}^{4} \partial_{y}F(y,a)^{j}{}_{k} \int_{y_{k}}^{y} \left[F^{-1}(x,a)^{k}{}_{3}\operatorname{Re}g(x) + F^{-1}(x,a)^{k}{}_{4}\operatorname{Im}g(x) \right] dx + \delta^{j}{}_{3}\operatorname{Re}g(y) + \delta^{j}{}_{4}\operatorname{Im}g(y) + c(a,g)\partial_{y}F(y,a)^{j}{}_{1}$$
(3.1)
$$= -\sum_{k=1}^{4} A(y,a)^{j}{}_{k}f^{k}(y,a) + \delta^{j}{}_{3}\operatorname{Re}g(y) + \delta^{j}{}_{4}\operatorname{Im}g(y)$$

with $A(y,a) = -\partial_y F(y,a)F(y,a)^{-1}$. Note that $\partial_y f^j(y,a) = f^{j+2}(y,a)$ for $j \in \{1,2\}$ and thus, in vector notation, Eq. (3.1) reads

$$\partial_y \begin{pmatrix} f^1(y,a) \\ f^2(y,a) \\ \partial_y f^1(y,a) \\ \partial_y f^2(y,a) \end{pmatrix} + A(y,a) \begin{pmatrix} f^1(y,a) \\ f^2(y,a) \\ \partial_y f^1(y,a) \\ \partial_y f^2(y,a) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \operatorname{Re} g(y) \\ \operatorname{Im} g(y) \end{pmatrix}.$$

Consequently, in view of the definition of the coefficients (p_1, p_2, q_1, q_2) associated to F,

$$\mathcal{L}_{F}^{-1}(a,g)(y) = f^{1}(y,a) + if^{2}(y,a)$$

satisfies the stated equation.

3.7. Regularization of \mathcal{L}_F^{-1} . The problem with the inverse \mathcal{L}_F^{-1} is that it does not map from Y to X because of a singularity at the endpoint -1. We now correct \mathcal{L}_F^{-1} with an additional term that cancels this singularity.

Definition 3.28. Let F be an admissible fundamental matrix on J. Then we define $\psi_F : J \times Y \to \mathbb{R}$ by

$$\psi_F(a,g) := \sum_{k=3}^4 \left[M_F(a)^4 \frac{M_F(a)^3}{M_F(a)^3} - M_F(a)^4 \right] \int_{-1}^1 \alpha_F^k(a,g)(x) dx.$$

Lemma 3.29. Let F be an admissible fundamental matrix on J. Then we have the bound

 $|\psi_F(a,g)| \lesssim ||g||_Y$

for all $g \in Y$ and $a \in J$. Furthermore, for every $g \in Y$, the map

$$a \mapsto \psi_F(a,g) : J \to \mathbb{R}$$

is continuous.

Proof. The bound follows directly from Lemma 3.24. For the continuity statement it suffices to note that, given $\epsilon > 0$, there exist $\eta, \delta > 0$ such that

$$\int_{1-\eta}^1 \left| \alpha_F^k(g,a) - \alpha_F^k(g,b) \right| < \frac{\epsilon}{2}$$

for all $a, b \in J$, as well as

$$\int_{-1}^{1-\eta} \left| \alpha_F^k(a,g) - \alpha_F^k(b,g) \right| < \frac{\epsilon}{2},$$

provided that $|a - b| < \delta$.

Definition 3.30. Let F be an admissible fundamental matrix on J and $(a, g) \in J \times Y$. Furthermore, let $\chi : \mathbb{R} \to [0, 1]$ be a smooth cut-off that satisfies $\chi(y) = 1$ for $y \leq -\frac{1}{2}$, $\chi(y) = 0$ for $y \geq 0$, and $|\chi'(y)| \leq 3$ for all $y \in \mathbb{R}$. Then we set

$$\mathcal{J}_F(a,g)(y) := \mathcal{L}_F^{-1}(a,g)(y) - \chi(y) \left[F_L(y)^1_4 + iF_L(y)^2_4 \right] \psi_F(a,g)$$

for $y \in (-1, 1)$, where F_L denotes the left part of F.

Remark 3.31. Note that \mathcal{J}_F is not an inverse of \mathcal{L}_F , unless $\psi_F(a,g) = 0$.

Proposition 3.32. Let F be an admissible fundamental matrix on J. Then we have the bound

$$\|\mathcal{J}_F(a,g)\|_X \lesssim \|g\|_Y$$

for all $(a,g) \in J \times Y$.

Proof. From Lemma 3.24 we obtain

$$\begin{aligned} |\mathcal{J}_{F}(a,g)(y)| &\lesssim \sum_{k=1}^{2} \left(|F(y,a)^{1}{}_{k}| + |F(y,a)^{2}{}_{k}| \right) \int_{-1}^{y} \|g\|_{Y} dx \\ &+ \sum_{k=3}^{4} \left(|F(y,a)^{1}{}_{k}| + |F(y,a)^{2}{}_{k}| \right) \int_{y}^{1} \|g\|_{Y} dx + \left(|F(y,a)^{1}{}_{1}| + |F(y,a)^{2}{}_{1}| \right) \|g\|_{Y} \\ &\lesssim \|g\|_{Y} \end{aligned}$$

and

$$\begin{aligned} |\mathcal{J}_{F}(a,g)'(y)| &\lesssim \sum_{k=1}^{2} \left(|F(y,a)^{3}_{k}| + |F(y,a)^{4}_{k}| \right) \int_{-1}^{y} ||g||_{Y} dx \\ &+ \sum_{k=3}^{4} \left(|F(y,a)^{3}_{k}| + |F(y,a)^{4}_{k}| \right) \int_{y}^{1} ||g||_{Y} dx + \left(|F(y,a)^{3}_{1}| + |F(y,a)^{4}_{1}| \right) ||g||_{Y} \\ &\lesssim \frac{||g||_{Y}}{1-y} \end{aligned}$$

for all $(y, a) \in [0, 1) \times J$.

In the case $y \in (-1,0]$, we have $F(y,a) = F_L(y)M_F(a)$, where F_L denotes the left part of F. Consequently, if $y \in (-1, -\frac{1}{2}]$,

$$\begin{aligned} \mathcal{J}_{F}(a,g)(y) &= \mathcal{L}_{F}^{-1}(a,g)(y) - \left[F_{L}(y)^{1}_{4} + iF_{L}(y)^{2}_{4}\right]\psi_{F}(a,g) \\ &= \sum_{k=1}^{2}\sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell}\right]M_{F}(a)^{\ell}_{k}\int_{-1}^{y}\alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4}\sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell}\right]M_{F}(a)^{\ell}_{k}\int_{y}^{1}\alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4}\sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell}\right]M_{F}(a)^{\ell}_{1}\frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}}\int_{-1}^{1}\alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \left[M_{F}(a)^{4}_{1}\frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} - M_{F}(a)^{4}_{k}\right]\left[F_{L}(y)^{1}_{4} + iF_{L}(y)^{2}_{4}\right]\int_{-1}^{1}\alpha_{F}^{k}(a,g). \end{aligned}$$

Since $|F_L(y)^j{}_k| \lesssim (1+y)^{-1}$ for $j \in \{1,2\}$ and $|F_L(y)^j{}_k| \lesssim 1$ for $k \in \{1,2\}$, we obtain

$$\mathcal{L}_{F}^{-1}(a,g)(y) = O((1+y)^{-1}a^{0}) \sum_{k=1}^{2} \int_{-1}^{y} \alpha_{F}^{k}(a,g) + O(y^{0}a^{0}) \sum_{k=3}^{4} \int_{y}^{1} \alpha_{F}^{k}(a,g) + \sum_{k=3}^{4} \left[F_{L}(y)^{1}_{3} + iF_{L}(y)^{2}_{3} \right] M_{F}(a)^{3}_{k} \left[\int_{1}^{y} \alpha_{F}^{k}(a,g) + \int_{-1}^{1} \alpha_{F}^{k}(a,g) \right] + \sum_{k=3}^{4} \left[F_{L}(y)^{1}_{4} + iF_{L}(y)^{2}_{4} \right] M_{F}(a)^{4}_{k} \left[\int_{1}^{y} \alpha_{F}^{k}(a,g) + \int_{-1}^{1} \alpha_{F}^{k}(a,g) \right]$$

and $\|\alpha_F^k(a,g)\|_{L^{\infty}(-1,1)} \lesssim \|g\|_Y$ yields

$$\mathcal{J}_F(a,g)(y)| \lesssim \|g\|_Y$$

for all $(y,a) \in (-1, -\frac{1}{2}] \times J$. Analogously, we find

$$\left|\mathcal{J}_F(a,g)'(y)\right| \lesssim \frac{\|g\|_Y}{1+y}$$

for all $(y,a) \in (-1, -\frac{1}{2}] \times J$. Finally, in the domain $y \in [-\frac{1}{2}, 0]$, the desired bound is obvious.

Next, we turn to the continuity of $\mathcal{J}_F(a,g)$ which requires a bit of thought because of the oscillatory terms near y = 1. We start with a simple lemma on a particular type of oscillatory integral that provides the basis for the treatment.

Lemma 3.33. Let $J \subset \mathbb{R} \setminus \{0\}$ be compact. Then we have the bound

$$\left|\int_{y}^{1} e^{-i\varphi(x,\alpha)} dx\right| \lesssim (1-y)^{3}$$

for all $y \in [0, 1)$ and all $\alpha \in J$.

Proof. An integration by parts yields

$$\int_{y}^{1} e^{-i\varphi(x,\alpha)} dx = \int_{y}^{1} \frac{i}{\varphi'(x,\alpha)} \partial_{x} e^{-i\varphi(x,\alpha)} dx = -\frac{i}{\varphi'(y,\alpha)} e^{-i\varphi(y,\alpha)} - \int_{y}^{1} e^{-i\varphi(x,\alpha)} \partial_{x} \frac{i}{\varphi'(x,\alpha)} dx$$

d the stated bound follows.

and the stated bound follows.

Lemma 3.34. Let $J \subset \mathbb{R} \setminus \{0\}$ be compact and $f \in C([0,1])$. Define $\psi : [0,1) \times J \to \mathbb{C}$ by

$$\psi(y,\alpha) := \frac{e^{i\varphi(y,\alpha)}}{1-y} \int_y^1 e^{-i\varphi(x,\alpha)} f(x) dx$$

Then ψ extends to a continuous function on $[0,1] \times J$.

Proof. Let $\epsilon > 0$ be arbitrary. Then there exists a $\delta > 0$ such that $|f(x) - f(1)| < \epsilon$ for all $x \in [1 - \delta, 1)$. In view of Lemma 3.33,

$$\begin{aligned} \left|\frac{1}{1-y}\int_{y}^{1}e^{-i\varphi(x,\alpha)}f(x)dx\right| &\leq \left|\frac{1}{1-y}\int_{y}^{1}e^{-i\varphi(x,\alpha)}f(1)dx\right| + \left|\frac{1}{1-y}\int_{y}^{1}e^{-i\varphi(x,\alpha)}[f(x)-f(1)]dx\right| \\ &\lesssim (1-y)^{2} + \epsilon \end{aligned}$$

for all $y \in [1 - \delta, 1)$ and all $\alpha \in J$. As a consequence, $\lim_{y \to 1^-} \psi(y, \alpha) = 0$ and we extend ψ to $[0, 1] \times J$ by setting $\psi(1, \alpha) := 0$. By construction, $\psi(\cdot, \alpha) : [0, 1] \to \mathbb{C}$ is continuous and the continuity is uniform with respect to $\alpha \in J$. Furthermore, $\psi(y, \cdot) : J \to \mathbb{C}$ is continuous for every $y \in [0, 1]$. By putting all of this together, we arrive at the claim.

Lemma 3.35. Let F be an admissible fundamental system on J and let $(a,g) \in J \times Y$. Then $\mathcal{J}_F(a,g) \in X$ and the map

$$a \mapsto \mathcal{J}_F(a,g) : J \to X$$

is continuous.

Proof. In view of the proof of Proposition 3.32 it is evident that $(y, a) \mapsto \mathcal{J}_F(a, g)(y)$ and $(y, a) \mapsto (1-y^2)\mathcal{J}_F(a, g)'(y)$ extend continuously to $[-1, 1-\delta] \times J$ for any $\delta \in (0, 1]$. Consequently, the only terms that require an argument are the oscillatory ones near y = 1. Inspection of Definitions 3.23 and 3.25 in conjunction with Lemma 3.13 shows that the most delicate term is

$$\sum_{k=3}^{4} \left[F(y,a)^{3}{}_{k} + iF(y,a)^{4}{}_{k} \right] \int_{y}^{1} \alpha_{F}^{k}(a,g)(x) dx,$$

which occurs in the expression for $\mathcal{J}_F(a,g)'$. However, this term is exactly of the type that is handled in Lemma 3.34 and we conclude that the map $(y,a) \mapsto (1-y^2)\mathcal{J}_F(a,g)'(y)$ extends continuously to $[-1,1] \times J$.

3.8. The solution system. Now we have all components in place in order to solve the equation $\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = 0.$

Definition 3.36. Let (a_*, f_*) be an admissible approximation and let F be an admissible fundamental matrix on J with parameter a_* . Furthermore, let (p_1, p_2, q_1, q_2) be the associated coefficients. Then, for $(a, f) \in J \times X$ and $y \in (-1, 1)$, we set

$$\begin{aligned} \mathcal{G}_F(a_*, f_*, a, f)(y) &:= [p_1(y, a) - p_0(y, a_* + a)]f'(y) + p_2(y, a)\overline{f'(y)} \\ &+ \left[q_1(y, a) - q_0(y, a_* + a) - \frac{2|f_*(y, a)|^2}{(1 - y)^2}\right]f(y) \\ &+ \left[q_2(y, a) - \frac{f_*(y, a)^2}{(1 - y)^2}\right]\overline{f(y)} - \mathcal{N}(a_*, f_*, a, f)(y) - \mathcal{R}(a_* + a, f_*(\cdot, a))(y). \end{aligned}$$

The operator \mathcal{G}_F represents the full "right-hand side" of the problem, taking into account also the error we make by replacing the linear operator \mathcal{L} by \mathcal{L}_F . The point is that \mathcal{G}_F maps into Y and by choosing "good" approximations, the right-hand side can be made arbitrarily small and hence treated perturbatively. **Lemma 3.37.** Let (a_*, f_*) be an admissible approximation and let F be an admissible fundamental matrix on J with parameter a_* . Then $\mathcal{G}_F(a_*, f_*, a, f) \in Y$ for all $(a, f) \in J \times X$.

Proof. Let $(a, f) \in J \times X$. We show that each of the terms that constitute $\mathcal{G}_F(a_*, f_*, a, f)$ belongs to Y. From Lemma 3.5 we already know that $\mathcal{R}(a_* + a, f_*(\cdot, a)) \in Y$. For the nonlinearity note that $f \in X$ implies $f \in L^{\infty}(-1,1)$ and thus, $\mathcal{N}(a_*, f_*, a, f)(y) = O((1-y)^{-2})$. This shows that $\mathcal{N}(a_*, f_*, a, f) \in Y$. Next, recall from Definition 3.3 that

$$p_0(y, a_* + a) = \frac{4i(a_* + a)}{(1 - y)^3} - \frac{2i(a_* + a)}{(1 - y)^2} - \frac{2 + \frac{2i}{(a_* + a)}}{1 - y} + \frac{2}{1 + y}$$
$$q_0(y, a_* + a) = -\frac{2 - 2i(a_* + a)}{(1 - y)^3} - \frac{\frac{1}{(a_* + a)^2} - \frac{i}{(a_* + a)}}{(1 - y)^2} - \frac{1 + \frac{i}{a_* + a}}{1 - y} - \frac{1 + \frac{i}{a_* + a}}{1 + y}$$

and from Proposition 3.21 we have

$$p_1(y,a) = \frac{4i(a_*+a)}{(1-y)^3} - \frac{2i(a_*+a)}{(1-y)^2} + \frac{2}{1+y} + O((1-y)^{-1})$$
$$q_1(y,a) = -\frac{2-2i(a_*+a)}{(1-y)^3} + O((1-y)^{-2}) + O((1+y)^{-1})$$

as well as

$$p_2(y,a) = O((1-y)^{-1})$$

$$q_2(y,a) = O((1-y)^{-2}) + O((1+y)^{-1}).$$

Consequently,

$$p_1(y,a) - p_0(y,a_* + a) = O((1-y)^{-1})$$

$$q_1(y,a) - q_0(y,a_* + a) = O((1-y)^{-2}) + O((1+y)^{-1})$$

and since $f \in X$ implies that $f'(y) = O((1-y^2)^{-1})$, we obtain

$$[p_1(y,a) - p_0(y,a_*+a)]f'(y) = O((1+y)^{-1}(1-y)^{-2})$$
$$p_2(y,a)\overline{f'(y)} = O((1+y)^{-1}(1-y)^{-2})$$

as well as

$$\left[q_1(y,a) - q_0(y,a_*+a) - \frac{2|f_*(y,a)|^2}{(1-y)^2} \right] f(y) = O((1+y)^{-1}(1-y)^{-2})$$
$$\left[q_2(y,a) - \frac{f_*(y,a)^2}{(1-y)^2} \right] \overline{f(y)} = O((1+y)^{-1}(1-y)^{-2}),$$

which completes the proof.

Lemma 3.38. Let (a_*, f_*) be an admissible approximation and let F be an admissible fundamental matrix on J with parameter a_* . Then, for every $f \in X$, the map

$$a \mapsto \mathcal{G}_F(a_*, f_*, a, f) : J \to Y$$

is continuous.

Proof. This follows immediately from the explicit form of \mathcal{G}_F and Lemma 3.5.

Lemma 3.39. Let (a_*, f_*) be an admissible approximation and let F be an admissible fundamental matrix on J with parameter a_* . Suppose that $(a, f) \in J \times X$ satisfies

$$\begin{cases} f(y) = \mathcal{J}_F(a, \mathcal{G}_F(a_*, f_*, a, f))(y) \\ \psi_F(a, \mathcal{G}_F(a_*, f_*, a, f)) = 0 \\ 25 \end{cases}$$
(3.2)

for all $y \in (-1, 1)$. Then $f \in C^2(-1, 1)$ and

$$\mathcal{R}(a_* + a, f_*(\cdot, a) + f)(y) = 0$$

for all $y \in (-1, 1)$.

Proof. Since $\psi_F(a, \mathcal{G}_F(a_*, f_*, a, f)) = 0$, we have $\mathcal{J}_F(a, \mathcal{G}_F(a_*, f_*, a, f)) = \mathcal{L}_F^{-1}(a, \mathcal{G}_F(a_*, f_*, a, f))$. By Lemma 3.37 and Lemma 3.27, $f \in C^2((-1, 1) \setminus \{0\}) \cap C^1(-1, 1)$ and by Eq. (3.2) together with Lemma 3.27,

$$\mathcal{L}_F(a,f)(y) = \mathcal{L}_F\left(a, \mathcal{L}_F^{-1}\left(a, \mathcal{G}_F(a_*, f_*, a, f)\right)\right)(y) = \mathcal{G}_F(a_*, f_*, a, f)(y)$$

for all
$$y \in (-1,1) \setminus \{0\}$$
. Furthermore, by the perturbative decomposition (Lemma 3.7), we obtain
 $\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = \mathcal{L}(a_*, f_*, a, f) + \mathcal{N}(a_*, f_*, a, f) + \mathcal{R}(a_* + a, f_*(\cdot, a))$
 $= \mathcal{L}_F(a, f) + \mathcal{L}(a_*, f_*, a, f) - \mathcal{L}_F(a, f) + \mathcal{N}(a_*, f_*, a, f) + \mathcal{R}(a_* + a, f_*(\cdot, a))$
 $= \mathcal{G}_F(a_*, f_*, a, f) + \mathcal{L}(a_*, f_*, a, f) - \mathcal{L}_F(a, f) + \mathcal{N}(a_*, f_*, a, f)$
 $+ \mathcal{R}(a_* + a, f_*(\cdot, a))$
 $= 0$

on $(-1,1) \setminus \{0\}$. Since $f \in C^1(-1,1)$ and $\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = 0$ is a second-order ODE with coefficients and forcing term in C(-1,1), it follows that $f \in C^2(-1,1)$ and $\mathcal{R}(a_*+a, f_*(\cdot, a)+f) = 0$ on all of (-1, 1).

Remark 3.40. Lemma 3.39 provides us with the means to practically solve the equation $\mathcal{R}(a_* +$ $a, f_*(\cdot, a) + f = 0$ since Eq. (3.2) only involves quantities that can be constructed explicitly.

4. Quantitative bounds

In order to really solve Eq. (3.2), it will be necessary to quantify all the bounds involved. In this section we develop the corresponding theoretical foundation.

Definition 4.1. Let F be an admissible fundamental matrix on J. In view of Lemma 3.24, we define

$$C_{\alpha}^{k}(F) := \sup\left\{ \left\| \alpha_{F}^{k}(a,g) \right\|_{L^{\infty}(-1,1)} : g \in Y, \|g\|_{Y} \le 1, a \in J \right\}$$

for $k \in \{1, 2, 3, 4\}$.

Lemma 4.2. Let F be an admissible fundamental matrix on J. Then we have the bound

$$C_{\alpha}^{k}(F) \leq \sup_{(y,a)\in(-1,1)\times J} \frac{|F^{-1}(y,a)^{k}| + |F^{-1}(y,a)^{k}|}{(1+y)(1-y)^{2}}$$

for $k \in \{1, 2, 3, 4\}$.

Proof. Evidently,

$$\left| \alpha_{F}^{k}(a,g)(y) \right| \leq \left(\left| F^{-1}(y,a)^{k}{}_{3} \right| + \left| F^{-1}(y,a)^{k}{}_{4} \right| \right) |g(y)|$$
 follows

and the stated bound follows.

Definition 4.3. Let F be an admissible fundamental matrix on J. Then we define

$$C_{\psi}(F) := \sup \{ |\psi_F(a,g)| : g \in Y, ||g||_Y \le 1, a \in J \}$$

Lemma 4.4. Let F be an admissible fundamental matrix on J. Then we have the bound

$$C_{\psi}(F) \le 2 \max_{a \in J} \sum_{k=3}^{4} \left| M_F(a)^4 \frac{M_F(a)^3}{M_F(a)^3} - M_F(a)^4 \right| C_{\alpha}^k(F).$$

Proof. By definition, we have

$$|\psi_F(a,g)| \le \sum_{k=3}^4 \left| M_F(a)^4 \frac{M_F(a)^3}{M_F(a)^3} - M_F(a)^4 \right| \int_{-1}^1 \left| \alpha_F^k(a,g)(x) \right| dx$$

ollows.

and the claim follows.

Definition 4.5. Let *F* be an admissible fundamental matrix on *J* with left part F_L . For $n \in \{0, 1\}$, we define $\beta_{F,L}^n : (-1, -\frac{1}{2}) \times J \to \mathbb{R}, \ \beta_{F,M}^n : [-\frac{1}{2}, 0] \times J \to \mathbb{R}, \ \text{and} \ \beta_{R,M}^n : (0, 1) \times J \to \mathbb{R}$ by

$$\begin{split} \beta_{F,L}^{n}(y,a) &:= (1+y)(1-y^{2})^{n} \sum_{k=1}^{2} \sum_{m=1}^{2} \left| \sum_{\ell=1}^{4} F_{L}(y)^{m+2n} {}_{\ell} M_{F}(a)^{\ell}{}_{k} \right| C_{\alpha}^{k}(F) \\ &+ (1+y)(1-y^{2})^{n} \\ &\times \sum_{k=3}^{4} \sum_{m=1}^{2} \left| \sum_{\ell=1}^{3} F_{L}(y)^{m+2n} {}_{\ell} M_{F}(a)^{\ell}{}_{1} \frac{M_{F}(a)^{3}{}_{k}}{M_{F}(a)^{3}{}_{1}} + F_{L}(y)^{m+2n} {}_{4} M_{F}(a)^{4}{}_{k} \right| C_{\alpha}^{k}(F) \\ &+ (1-y)(1-y^{2})^{n} \\ &\times \sum_{k=3}^{4} \sum_{m=1}^{2} \left| \sum_{\ell=1}^{2} F_{L}(y)^{m+2n} {}_{\ell} \left(M_{F}(a)^{\ell}{}_{1} \frac{M_{F}(a)^{3}{}_{k}}{M_{F}(a)^{3}{}_{1}} - M_{F}(a)^{\ell}{}_{k} \right) \right| C_{\alpha}^{k}(F), \end{split}$$

$$\begin{split} \beta_{F,M}^{n}(y,a) &:= (1+y)(1-y^{2})^{n} \sum_{k=1}^{2} \sum_{m=1}^{2} \left| \sum_{\ell=1}^{4} F_{L}(y)^{m+2n} {}_{\ell} M_{F}(a)^{\ell} {}_{k} \right| C_{\alpha}^{k}(F) \\ &+ (1+y)(1-y^{2})^{n} \sum_{k=3}^{4} \sum_{m=1}^{2} \left| \sum_{\ell=1}^{4} F_{L}(y)^{m+2n} {}_{\ell} M_{F}(a)^{\ell} {}_{1} \frac{M_{F}(a)^{3} {}_{k}}{M_{F}(a)^{3} {}_{1}} \right| C_{\alpha}^{k}(F) \\ &+ (1-y)(1-y^{2})^{n} \\ &\times \sum_{k=3}^{4} \sum_{m=1}^{2} \left| \sum_{\ell \in \{1,2,4\}} F_{L}(y)^{m+2n} {}_{\ell} \left(M_{F}(a)^{\ell} {}_{1} \frac{M_{F}(a)^{3} {}_{k}}{M_{F}(a)^{3} {}_{1}} - M_{F}(a)^{\ell} {}_{k} \right) \right| C_{\alpha}^{k}(F) \\ &+ (1-y^{2})^{n} \sum_{m=1}^{2} \left| F_{L}(y)^{m+2n} {}_{4} \right| C_{\psi}(F) + 3n(1-y^{2})^{n} \sum_{m=1}^{2} |F_{L}(y)^{m} {}_{4}| C_{\psi}(F), \end{split}$$

and

$$\begin{split} \beta_{F,R}^n(y,a) &:= (1+y)(1-y^2)^n \sum_{k=1}^2 \sum_{m=1}^2 \left| F(y,a)^{m+2n}{}_k \right| C_\alpha^k(F) \\ &+ (1-y)(1-y^2)^n \sum_{k=3}^4 \sum_{m=1}^2 \left| \frac{M_F(a)^3{}_k}{M_F(a)^3{}_1} F(y,a)^{m+2n}{}_1 - F(y,a)^{m+2n}{}_k \right| C_\alpha^k(F) \\ &+ (1+y)(1-y^2)^n \sum_{k=3}^4 \sum_{m=1}^2 \left| F(y,a)^{m+2n}{}_1 \frac{M_F(a)^3{}_k}{M_F(a)^3{}_1} \right| C_\alpha^k(F). \end{split}$$

Definition 4.6. Let F be an admissible fundamental matrix on J. We define

$$C_{\mathcal{J}}(F) := \sup \{ \|\mathcal{J}_F(a,g)\|_X : g \in Y, \|g\|_Y \le 1, a \in J \}.$$

Proposition 4.7. Let F be an admissible fundamental matrix on J. Then we have the bound

$$C_{\mathcal{J}}(F) \le \max\left\{\sum_{n=0}^{1} \sup_{(-1,-\frac{1}{2})\times J} \beta_{F,L}^{n}, \sum_{n=0}^{1} \sup_{(-\frac{1}{2},0)\times J} \beta_{F,M}^{n}, \sum_{n=0}^{1} \sup_{(0,1)\times J} \beta_{F,R}^{n}\right\}.$$

Proof. Let $g \in Y$ with $||g||_Y \leq 1$ and $a \in J$. By definition, we have

$$\begin{aligned} \mathcal{L}_{F}^{-1}(a,g)(y) &= \sum_{k=1}^{2} \left[F(y,a)^{1}{}_{k} + iF(y,a)^{2}{}_{k} \right] \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \left[F(y,a)^{1}{}_{k} + iF(y,a)^{2}{}_{k} \right] \int_{y}^{1} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \left[F(y,a)^{1}{}_{1} + iF(y,a)^{2}{}_{1} \right] \frac{M_{F}(a)^{3}{}_{k}}{M_{F}(a)^{3}{}_{1}} \left[\int_{-1}^{y} \alpha_{F}^{k}(a,g) + \int_{y}^{1} \alpha_{F}^{k}(a,g) \right] \end{aligned}$$

and

$$\mathcal{J}_F(a,g)(y) = \mathcal{L}_F^{-1}(a,g)(y) - \chi(y) \left[F_L(y)^1_4 + iF_L(y)^2_4 \right] \psi_F(a,g),$$

with

$$\psi_F(a,g) = \sum_{k=3}^4 \left[M_F(a)^4 \frac{M_F(a)^3}{M_F(a)^3} - M_F(a)^4 \right] \int_{-1}^1 \alpha_F^k(a,g)(x) dx.$$

Consequently, $|\mathcal{J}_F(a,g)(y)| \leq \beta_{F,R}^0(y,a)$ for all $(y,a) \in (0,1) \times J$. For the derivative, we note that

$$\begin{split} \mathcal{L}_{F}^{-1}(a,g)'(y) &= \sum_{k=1}^{2} \left[F(y,a)^{3}{}_{k} + iF(y,a)^{4}{}_{k} \right] \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \left[F(y,a)^{3}{}_{k} + iF(y,a)^{4}{}_{k} \right] \int_{y}^{1} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \left[F(y,a)^{3}{}_{1} + iF(y,a)^{4}{}_{1} \right] \frac{M_{F}(a)^{3}{}_{k}}{M_{F}(a)^{3}{}_{1}} \left[\int_{-1}^{y} \alpha_{F}^{k}(a,g) + \int_{y}^{1} \alpha_{F}^{k}(a,g) \right] \end{split}$$

since $\partial_y F(y,a)^j{}_k = F(y,a)^{j+2}{}_k$ for all $(y,a) \in (-1,1) \times J, \ j \in \{1,2\}, \ k \in \{1,2,3,4\}$, and the evaluation terms vanish because

$$\sum_{k=1}^{4} \left[F(y,a)^{1}{}_{k} + iF(y,a)^{2}{}_{k} \right] \alpha_{F}^{k}(a,g)(y) = \sum_{k=1}^{4} \left[F(y,a)^{1}{}_{k} + iF(y,a)^{2}{}_{k} \right] \left[F^{-1}(y,a)^{k}{}_{3}\operatorname{Re} g(y) + F^{-1}(y,a)^{k}{}_{4}\operatorname{Im} g(y) \right] = 0.$$

Accordingly, we obtain $(1-y^2)|\mathcal{J}_F(a,g)'(y)| \leq \beta_{F,R}^1(y,a)$ for all $(y,a) \in (0,1) \times J$.

In the case $y \in (-1,0]$, we have $F(y,a)^j{}_k = \sum_{\ell=1}^4 F_L(y)^j{}_\ell M_F(a)^\ell{}_k$ and by observing that

$$\begin{split} &\sum_{k=3}^{4}\sum_{\ell=1}^{4}\left[F_{L}(y)^{1}_{\ell}+iF_{L}(y)^{2}_{\ell}\right]M_{F}(a)^{\ell}_{1}\frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}}\int_{y}^{1}\alpha_{F}^{k}(a,g)\\ &=\sum_{k=3}^{4}\sum_{\ell\in\{1,2,4\}}\left[F_{L}(y)^{1}_{\ell}+iF_{L}(y)^{2}_{\ell}\right]M_{F}(a)^{\ell}_{1}\frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}}\int_{y}^{1}\alpha_{F}^{k}(a,g)\\ &+\sum_{k=3}^{4}\left[F_{L}(y)^{1}_{3}+iF_{L}(y)^{2}_{3}\right]M_{F}(a)^{3}_{k}\int_{y}^{1}\alpha_{F}^{k}(a,g), \end{split}$$

we find

$$\begin{aligned} \mathcal{L}_{F}^{-1}(a,g)(y) &= \sum_{k=1}^{2} \sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{k} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \sum_{\ell \in \{1,2,4\}} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{k} \int_{y}^{1} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \sum_{\ell \in \{1,2,4\}} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} \int_{y}^{1} \alpha_{F}^{k}(a,g). \end{aligned}$$

By recalling that $|\chi'| \leq 3$, we therefore obtain $|\mathcal{J}_F(a,g)(y)| \leq \beta_{F,M}^0(y,a)$ and $|(1-y^2)\mathcal{J}_F(a,g)'(y)| \leq \beta_{F,M}^1(y,a)$ for all $(y,a) \in [-\frac{1}{2},0] \times J$. In the case $y \in (-1,-\frac{1}{2}]$, we further compute

$$\begin{split} \mathcal{J}_{F}(a,g)(y) &= \mathcal{L}_{F}^{-1}(a,g)(y) - \sum_{k=3}^{4} \left[F_{L}(y)^{1}_{4} + iF_{L}(y)^{2}_{4} \right] \left[M_{F}(a)^{4}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} - M_{F}(a)^{4}_{k} \right] \\ &\times \left[\int_{-1}^{y} \alpha_{F}^{k}(a,g) + \int_{y}^{1} \alpha_{F}^{k}(a,g) \right] \\ &= \sum_{k=1}^{2} \sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{k} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \left[F_{L}(y)^{1}_{4} + iF_{L}(y)^{2}_{4} \right] \left[M_{F}(a)^{4}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} - M_{F}(a)^{4}_{k} \right] \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \sum_{\ell=1}^{2} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{k} \int_{y}^{1} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \sum_{\ell=1}^{2} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} \int_{y}^{1} \alpha_{F}^{k}(a,g), \end{split}$$

which simplifies to

$$\begin{aligned} \mathcal{J}_{F}(a,g)(y) &= \sum_{k=1}^{2} \sum_{\ell=1}^{4} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{k} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \sum_{\ell=1}^{3} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \left[F_{L}(y)^{1}_{4} + iF_{L}(y)^{2}_{4} \right] M_{F}(a)^{4}_{k} \int_{-1}^{y} \alpha_{F}^{k}(a,g) \\ &- \sum_{k=3}^{4} \sum_{\ell=1}^{2} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{k} \int_{y}^{1} \alpha_{F}^{k}(a,g) \\ &+ \sum_{k=3}^{4} \sum_{\ell=1}^{2} \left[F_{L}(y)^{1}_{\ell} + iF_{L}(y)^{2}_{\ell} \right] M_{F}(a)^{\ell}_{1} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} \int_{y}^{1} \alpha_{F}^{k}(a,g) \end{aligned}$$

and this yields the bounds $|\mathcal{J}_F(a,g)(y)| \leq \beta_{F,L}^0(y,a)$ and $|(1-y^2)\mathcal{J}_F(a,g)'(y)| \leq \beta_{F,L}^1(y,a)$ for all $(y,a) \in (-1,-\frac{1}{2}] \times J.$

5. Tools for polynomials and rational functions

In order to apply the theory developed so far to the concrete problem at hand, we need good approximations to the desired solution and the fundamental matrix of the linear operator. These will be provided by explicit polynomials with rational coefficients. For estimating these polynomials, we employ some elementary methods that we describe now.

5.1. The *T*-norm. One of the most frequent problems we will face is to rigorously estimate the L^{∞} -norm of explicitly given but complicated polynomials of high degree. The key tool for this is an auxiliary norm that is based on the Chebyshev expansion.

Definition 5.1. Let $p : \mathbb{C}^n \to \mathbb{C}$ be a polynomial. For $j \in [1, n] \cap \mathbb{Z}$, we denote by $\deg_j p$ the degree of p in the j-th variable and set $\deg p := (\deg_1 p, \deg_2 p, \ldots, \deg_n p)$. Furthermore, we say that $\deg p \leq (d_1, d_2, \ldots, d_n) \in \mathbb{N}_0^n$ if $\deg_j p \leq d_j$ for every $j \in [1, n] \cap \mathbb{Z}$ and analogously for \leq replaced by \geq .

Definition 5.2 (*T*-norm on the standard cube). Let $p : \mathbb{C}^n \to \mathbb{C}$ be a polynomial with deg $p \ge 0$ and set $d_j := \deg_j p$ for $j \in [1, n] \cap \mathbb{Z}$. Furthermore, let $c_{j_1 j_2 \dots j_n} \in \mathbb{C}$ be the unique coefficients that satisfy

$$p(x^1, \dots, x^n) = \sum_{j_1=0}^{d_1} \cdots \sum_{j_n=0}^{d_n} c_{j_1\dots j_n} T_{j_1}(x^1) \dots T_{j_n}(x^n)$$

for all $(x^1, \ldots, x^n) \in \mathbb{C}^n$. Then we define the *T*-norm of *p* on $[-1, 1]^n$ by

$$\|p\|_{T([-1,1]^n)} := \sum_{j_1=0}^{d_1} \cdots \sum_{j_n=0}^{d_n} |c_{j_1\dots j_n}|$$

and $||0||_{T([-1,1]^n)} := 0.$

The T-norm on an arbitrary rectangular domain is now simply defined by composition with suitable affine-linear functions.

Definition 5.3. For $a, b \in \mathbb{R}$ with a < b, we define $\phi_{a,b} : [-1,1] \to [a,b]$ by

$$\phi_{a,b}(x) := \frac{b-a}{2}x + \frac{a+b}{2}.$$

Definition 5.4 (*T*-norm on a rectangular domain). Let $n \in \mathbb{N}$ and for $j \in [1, n] \cap \mathbb{Z}$, let $a_j, b_j \in \mathbb{R}$ with $a_j < b_j$. Let $p : \mathbb{C}^n \to \mathbb{C}$ be a polynomial. Then we define the *T*-norm of p on $[a_1, b_1] \times \cdots \times [a_n, b_n]$ by

$$\|p\|_{T([a_1,b_1]\times\cdots\times[a_n,b_n])} := \|(x^1,\ldots,x^n)\mapsto p(\phi_{a_1,b_1}(x^1),\ldots,\phi_{a_n,b_n}(x^n))\|_{T([-1,1]^n)}$$

One crucial feature is that the T-norm controls the L^{∞} -norm.

Lemma 5.5. Let $p : \mathbb{C}^n \to \mathbb{C}$ be a polynomial and for $j \in [1, n] \cap \mathbb{Z}$, let $a_j, b_j \in \mathbb{R}$ with $a_j < b_j$. Then we have

$$\|p\|_{L^{\infty}([a_1,b_1]\times\cdots\times[a_n,b_n])} \le \|p\|_{T([a_1,b_1]\times\cdots\times[a_n,b_n])}.$$

Proof. It suffices to prove the statement for n = 1 and we write $a = a_1$ and $b = b_1$. The case p = 0 is trivial and thus, we assume deg $p \ge 0$. Let $(c_j)_{j=0}^{\deg p} \subset \mathbb{C}$ be the unique coefficients that satisfy

$$(p \circ \phi_{a,b})(x) = \sum_{j=0}^{\deg p} c_j T_j(x)$$

Then we have

$$\|p\|_{L^{\infty}([a,b])} = \|p \circ \phi_{a,b}\|_{L^{\infty}([-1,1])} \le \sum_{j=0}^{\deg p} |c_j| = \|p \circ \phi_{a,b}\|_{T([-1,1])} = \|p\|_{T([a,b])},$$

since $||T_n||_{L^{\infty}([-1,1])} \leq 1$ for all $n \in \mathbb{N}_0$,

Remark 5.6. One of the key observations in the following is that the *T*-norm not only controls the L^{∞} -norm but is in fact very close to it. As a consequence, in many cases one can obtain a sufficiently precise estimate on the L^{∞} -norm of a polynomial by simply computing its *T*-norm.

5.2. Computation of the T-norm. Next, we derive an algorithm to explicitly compute the T-norm of a given polynomial. The first ingredient is the representation of monomials in terms of Chebyshev polynomials.

Lemma 5.7. Let $n \in \mathbb{N}_0$ and $x \in \mathbb{C}$. Then we have

$$x^{n} = \frac{1}{2^{n-1}} \sum_{k=0}^{\frac{n-1}{2}} \binom{n}{\frac{n-1}{2}-k} T_{2k+1}(x)$$

if n is odd and

$$x^{n} = \frac{1}{2^{n}} \binom{n}{\frac{n}{2}} + \frac{1}{2^{n-1}} \sum_{k=1}^{\frac{n}{2}} \binom{n}{\frac{n}{2}-k} T_{2k}(x)$$

if n is even.

Proof. These formulas are of course well-known and can easily be derived by combining the defining property $T_n(\cos(z)) = \cos(nz), z \in \mathbb{C}$, with the binomial theorem applied to

$$\cos(z)^n = \frac{1}{2^n} \left(e^{iz} + e^{-iz} \right)^n.$$

Lemma 5.7 motivates the following definition.

Definition 5.8. Let $j, k \in \mathbb{N}_0$. If j > k, we set $C^j{}_k = 0$. If $j \le k$, we define

$$\begin{split} C^{j}{}_{k} &:= \frac{1}{2^{k-1}} \binom{k}{\frac{k-1}{2} - \frac{j-1}{2}} & \text{if } k \text{ odd and } j \text{ odd} \\ C^{0}{}_{k} &:= \frac{1}{2^{k}} \binom{k}{\frac{k}{2}} & \text{if } k \text{ even} \\ C^{j}{}_{k} &:= \frac{1}{2^{k-1}} \binom{k}{\frac{k}{2} - \frac{j}{2}} & \text{if } k \text{ even, } j \text{ even, and } j \ge 2 \\ C^{j}{}_{k} &:= 0 & \text{otherwise.} \end{split}$$

According to Lemma 5.7, we have

$$x^n = \sum_{k=0}^n C^k{}_n T_k(x)$$

and this allows one to go from the monomial representation to the Chebyshev representation. In practice, however, a polynomial p is typically given in *Lagrange representation*, i.e., by evaluation at particular points. In order to capture this, one uses the *Lagrange basis*.

Definition 5.9. Let $d \in \mathbb{N}$ and $\xi = (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{C}^{d+1}$ with $\xi^j \neq \xi^k$ if $j \neq k$. Then, for $j \in [0, d] \cap \mathbb{Z}$, we define $\ell_j^{\xi} : \mathbb{C} \to \mathbb{C}$ by

$$\ell_j^{\xi}(x) := \prod_{k \in [0,d] \cap \mathbb{Z} \setminus \{j\}} \frac{x - \xi^k}{\xi^j - \xi^k}.$$

Note that $\ell_j^{\xi}(\xi^k) = \delta_j{}^k$ and thus, the corresponding Lagrange representation of a polynomial p of degree d reads

$$p(x) = \sum_{j=0}^{d} p(\xi^{j}) \ell_{j}^{\xi}(x).$$

In order to obtain the monomial representation, we need to express the polynomials ℓ_j^{ξ} in terms of monomials. This is a classical problem in polynomial approximation that amounts to inverting the Vandermonde matrix. To keep the technical prerequisites at a bare minimum, we prefer to state an explicit recursion formula that can be implemented straight away without the need for solving a linear system.

Definition 5.10. Let $n \in \mathbb{N}$ and $\xi = (\xi^1, \xi^2, \dots, \xi^n) \in \mathbb{C}^n$. For $j \in [0, n] \cap \mathbb{Z}$ and $k \in [0, j] \cap \mathbb{Z}$, we define $a_{j,k}(\xi) \in \mathbb{C}$ recursively by

$$a_{j,0}(\xi) := \prod_{\ell=1}^{j} \xi^{\ell}$$

$$a_{j,k}(\xi) := a_{j-1,k-1}(\xi) + \xi^{j} a_{j-1,k}(\xi), \qquad k \in [1, j-1] \cap \mathbb{Z}$$

$$a_{j,j}(\xi) := 1.$$

Lemma 5.11. Let $n \in \mathbb{N}$, $\xi = (\xi^1, \xi^2, \dots, \xi^n) \in \mathbb{C}^n$, and $x \in \mathbb{C}$. Then we have

$$\prod_{j=1}^{n} (x+\xi^j) = \sum_{k=0}^{n} a_{n,k}(\xi) x^k.$$

Proof. This follows immediately by a simple induction.

Definition 5.12. Let $d \in \mathbb{N}$ and $\xi = (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{C}^{d+1}$. For $j \in [0, d] \cap \mathbb{Z}$, we define $\xi_{(j)} \in \mathbb{C}^d$ by

$$\xi_{(j)} := (\xi^1, \dots, \xi^{j-1}, \xi^{j+1}, \dots, \xi^d)$$

i.e., $\xi_{(j)}$ is obtained from ξ by removing the *j*-th element.

Definition 5.13. Let $d \in \mathbb{N}$ and $\xi = (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{C}^{d+1}$ with $\xi^j \neq \xi^k$ if $j \neq k$. For $j, k \in [0, d] \cap \mathbb{Z}$, we define $L(\xi)^j_k \in \mathbb{C}$ by

$$L(\xi)^{j}_{k} := \frac{a_{d,j}(-\xi_{(k)})}{\prod_{\ell \in [0,d] \cap \mathbb{Z} \setminus \{k\}} (\xi^{k} - \xi^{\ell})}.$$

Lemma 5.14. Let $d \in \mathbb{N}$, $\xi = (\xi_0, \xi_1, \dots, \xi_d) \in \mathbb{C}^{d+1}$ with $\xi^j \neq \xi^k$ if $j \neq k, x \in \mathbb{C}$, and $j \in [0, d] \cap \mathbb{Z}$. Then we have

$$\ell_j^{\xi}(x) = \sum_{k=0}^d L(\xi)^k{}_j x^k.$$

Proof. This follows immediately by inserting the definitions and using Lemma 5.11.

Definition 5.15. Let $d \in \mathbb{N}$ and $\xi = (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{C}^{d+1}$ with $\xi^j \neq \xi^k$ if $j \neq k$. Then, for $j, k \in [0, d] \cap \mathbb{Z}$, we define $T(\xi)^j{}_k \in \mathbb{C}$ by

$$T(\xi)^{j}{}_{k} := \sum_{\ell=0}^{d} C^{j}{}_{\ell}L(\xi)^{\ell}{}_{k}.$$

With these preparations we can now calculate the T-norm directly from the Lagrange representation.

Lemma 5.16. Let $d \in \mathbb{N}$, $\xi \in (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{C}^{d+1}$ with $\xi^j \neq \xi^k$ if $j \neq k$. Let $p : \mathbb{C} \to \mathbb{C}$ be a polynomial of degree d. Then we have

$$\|p\|_{T([-1,1])} = \sum_{j=0}^{d} \left| \sum_{k=0}^{d} T(\xi)^{j}{}_{k} p(\xi^{k}) \right|$$

Proof. By Lemmas 5.14 and 5.7, we have

$$\sum_{j=0}^{d} p(\xi^{j})\ell_{j}^{\xi}(x) = \sum_{j=0}^{d} p(\xi^{j}) \sum_{k=0}^{d} L(\xi)^{k}{}_{j}x^{k} = \sum_{j=0}^{d} p(\xi^{j}) \sum_{k=0}^{d} L(\xi)^{k}{}_{j}\sum_{\ell=0}^{d} C^{\ell}{}_{k}T_{\ell}(x)$$
$$= \sum_{j=0}^{d} \sum_{k=0}^{d} T(\xi)^{j}{}_{k}p(\xi^{k})T_{j}(x)$$

and the claim follows.

Remark 5.17. One crucial point is that if the polynomial p has rational coefficients and $a, b \in \mathbb{Q}$ then $\|p\|_{T([a,b])}$ can be computed by performing a finite number of elementary operations in \mathbb{Q} . Consequently, $\|p\|_{T([a,b])}$ can be computed *exactly*.

The generalization to polynomials of more than one variable is immediate and at most notationally challenging.

Lemma 5.18. Let $n \in \mathbb{N}$ and for $j \in [1, n] \cap \mathbb{Z}$, let $d_j \in \mathbb{N}$ and $\xi_j = (\xi_j^0, \xi_j^1, \dots, \xi_j^{d_j}) \in \mathbb{C}^{d_j+1}$ with $\xi_j^k \neq \xi_j^\ell$ if $k \neq \ell$. Suppose that $p : \mathbb{C}^n \to \mathbb{C}$ is a polynomial of degree d_j in the *j*-th variable. Then

 \square

we have

$$\|p\|_{T([-1,1]^n)} = \sum_{j_1=0}^{d_1} \cdots \sum_{j_n=0}^{d_n} \left| \sum_{k_1=0}^{d_1} \cdots \sum_{k_n=0}^{d_n} T(\xi_1)^{j_1}{}_{k_1} \dots T(\xi_n)^{j_n}{}_{k_n} p(\xi_1^{k_1}, \dots, \xi_n^{k_n}) \right|.$$

5.3. **Derivatives of polynomials.** With the tools developed so far it is also straightforward to compute derivatives of polynomials.

Lemma 5.19. Let $d \in \mathbb{N}_0$ and $\xi = (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{C}^{d+1}$ with $\xi^j \neq \xi^k$ if $j \neq k$. Furthermore, let $p : \mathbb{C} \to \mathbb{C}$ be a polynomial of degree d and $m \in [0, d] \cap \mathbb{Z}$. Then we have

$$p^{(m)}(x) = \sum_{j=0}^{d-m} \frac{(j+m)!}{j!} \sum_{k=0}^{d} L(\xi)^{j+m}{}_k p(\xi^k) x^j.$$

Proof. This is straightforward from the representation

$$p(x) = \sum_{j=0}^{d} \sum_{k=0}^{d} L(\xi)^{j}{}_{k} p(\xi^{k}) x^{j}.$$

5.4. The approximate maximum. Finally, we provide a simple method to rigorously estimate the minima and maxima of rational functions by evaluating them on a discrete grid.

Lemma 5.20. Let $n \in \mathbb{N}$ and for $j \in \{1, 2..., n\}$, let $M_j \in \mathbb{N}$. Furthermore, let $a_j, b_j \in \mathbb{R}$ with $a_j < b_j$ and set

$$\Lambda := \left\{ \left(a_1 + \frac{b_1 - a_1}{M_1} k_1, a_2 + \frac{b_2 - a_2}{M_2} k_2, \dots, a_n + \frac{b_n - a_n}{M_n} k_n \right) : k_j \in [0, M_j] \cap \mathbb{Z}, j \in [1, n] \cap \mathbb{Z} \right\}$$

as well as $\Omega := [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$. Let $f : \Omega \to \mathbb{R}$ be continuously differentiable. Then we have the bound

$$\max_{\Omega} f \le \max_{\Lambda} f + \sum_{j=1}^{n} \frac{b_j - a_j}{M_j} \|\partial_j f\|_{L^{\infty}(\Omega)}.$$

Proof. The statement is a consequence of the mean value theorem.

Definition 5.21. Let $n \in \mathbb{N}$. For $j \in \{1, 2, ..., n\}$ let $a_j, b_j \in \mathbb{R}$ with $a_j < b_j$ and set $\Omega := [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$. Furthermore, let $p, q : \mathbb{R}^n \to \mathbb{R}$ be polynomials with $\min_{\Omega} |q| > 0$, $\epsilon > 0$, and set

$$M_j := \min\left\{m \in \mathbb{N} : m \ge \frac{n}{\epsilon}(b_j - a_j) \|\partial_j pq - p\partial_j q\|_{T(\Omega)}\right\}$$

for $j \in \{1, 2, \ldots, n\}$, as well as

$$\Lambda := \left\{ \left(a_1 + \frac{b_1 - a_1}{M_1} k_1, a_2 + \frac{b_2 - a_2}{M_2} k_2, \dots, a_n + \frac{b_n - a_n}{M_n} k_n \right) : k_j \in [0, M_j] \cap \mathbb{Z}, j \in [0, n] \cap \mathbb{Z} \right\}.$$

Then we define

$$\max_{\Omega,\epsilon} \frac{p}{q} := \max_{\Lambda} \frac{p}{q}, \qquad \min_{\Omega,\epsilon} \frac{p}{q} := \min_{\Lambda} \frac{p}{q}, \qquad \left\| \frac{p}{q} \right\|_{L^{\infty}(\Omega,\epsilon)} := \max_{\Lambda} \left| \frac{p}{q} \right|.$$

Lemma 5.22. Let $p, q : \mathbb{R}^n \to \mathbb{R}$ be polynomials, $\epsilon > 0$, and $a_j, b_j \in \mathbb{R}$ with $a_j < b_j$ for $j \in [1, n] \cap \mathbb{Z}$. Furthermore, set $\Omega := [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$ and assume that $\min_{\Omega} |q| > 0$. Then we have

$$\left|\min_{\Omega,\epsilon} \frac{p}{q} - \min_{\Omega} \frac{p}{q}\right| \le \frac{\epsilon}{\min_{\Omega} q^2}, \qquad \left|\max_{\Omega,\epsilon} \frac{p}{q} - \max_{\Omega} \frac{p}{q}\right| \le \frac{\epsilon}{\min_{\Omega} q^2}$$

and

$$\left\|\frac{p}{q}\right\|_{L^{\infty}(\Omega,\epsilon)} - \left\|\frac{p}{q}\right\|_{L^{\infty}(\Omega)}\right| \leq \frac{\epsilon}{\min_{\Omega} q^2}$$

Proof. Let M_j and Λ be as in Definition 5.21. Obviously, we have $\max_{\Omega} \frac{p}{q} \ge \max_{\Omega, \epsilon} \frac{p}{q}$ and thus, by Lemmas 5.20 and 5.5, we obtain

$$\begin{split} \left| \max_{\Omega,\epsilon} \frac{p}{q} - \max_{\Omega} \frac{p}{q} \right| &= \max_{\Omega} \frac{p}{q} - \max_{\Lambda} \frac{p}{q} \le \sum_{j=1}^{n} \frac{b_j - a_j}{M_j} \left\| \partial_j \frac{p}{q} \right\|_{L^{\infty}(\Omega)} = \sum_{j=1}^{n} \frac{b_j - a_j}{M_j} \left\| \frac{\partial_j pq - p \partial_j q}{q^2} \right\|_{L^{\infty}(\Omega)} \\ &\le \frac{1}{\min_{\Omega} q^2} \sum_{j=1}^{n} \frac{b_j - a_j}{M_j} \| \partial_j pq - p \partial_j q \|_{T(\Omega)} \\ &\le \frac{\epsilon}{\min_{\Omega} q^2}. \end{split}$$

The remaining estimates are a direct consequence of this because $\min \frac{p}{q} = -\max(-\frac{p}{q})$ and $\|\frac{p}{q}\|_{L^{\infty}} = \max\{|\min \frac{p}{q}|, |\max \frac{p}{q}|\}$.

Remark 5.23. Again, the crucial point is that for polynomials p, q with rational coefficients and rational endpoints $a_j, b_j, \max_{\Omega, \epsilon} \frac{p}{q}$ can be computed *exactly*.

6. Approximations

6.1. The approximate solution. We define the approximate solution and rigorously estimate the remainder $\mathcal{R}(a_* + a, f_*(\cdot, a))$.

Definition 6.1. Let

$$P_*(y) := \sum_{n=0}^{50} c_n(P_*)T_n(y)$$

with $(c_n(P_*))_{n=0}^{50} \subset \mathbb{C}$ given in Appendix A.1. We set $a_* := \frac{772201763088846}{841768781900003}$ and define $f_* : [-1, 1] \times \mathbb{R} \setminus \{-a_*\}$ by

$$f_*(y,a) := P_*(y) + \frac{2i(a_* + a)}{1 - i(a_* + a)} P'_*(1) - P_*(1).$$

Lemma 6.2. The pair (a_*, f_*) is a nontrivial admissible approximation.

Proof. Evidently, $(a_*, f_*) \in \mathbb{R} \setminus \{0\} \times C^2([-1, 1] \times \mathbb{R} \setminus \{-a_*\})$ and

$$f'_*(1,a) = \frac{1 - i(a_* + a)}{2i(a_* + a)} f_*(1,a)$$

for all $a \in \mathbb{R} \setminus \{-a_*\}$.

Definition 6.3. For brevity, we set

$$J_* := [-10^{-10}, 10^{-10}]$$

Proposition 6.4. We have

 $\|\mathcal{R}(a_* + a, f_*(\cdot, a))\|_Y \le 4 \cdot 10^{-9}$

for all $a \in J_*$.

Proof. By definition, we have

$$\mathcal{R}(a_* + a, f_*(\cdot, a))(y) = f_*''(y, a) + p_0(y, a_* + a)f_*'(y, a) + q_0(y, a_* + a)f_*(y, a) + \frac{f_*(y, a)^2 f_*(y, a)}{(1 - y)^2}$$

with

$$p_0(y, a_* + a) = \frac{4i(a_* + a)}{(1 - y)^3} - \frac{2i(a_* + a)}{(1 - y)^2} - \frac{2 + \frac{2i}{a_* + a}}{1 - y} + \frac{2}{1 + y}$$
$$q_0(y, a_* + a) = -\frac{2 - 2i(a_* + a)}{(1 - y)^3} - \frac{\frac{1}{(a_* + a)^2} - \frac{i}{a_* + a}}{(1 - y)^2} - \frac{1 + \frac{i}{a_* + a}}{1 - y} - \frac{1 + \frac{i}{a_* + a}}{1 + y}$$

We set

 $R(y,a) := (a_* + a)^2 [1 - i(a_* + a)]^2 [1 + i(a_* + a)] (1 + y)(1 - y)^2 \mathcal{R}(a_* + a, f_*(\cdot, a))(y).$

By construction and Lemma 3.5, R is a polynomial of degree at most (151, 6). Furthermore, we have

$$|(a_*+a)^2[1-i(a_*+a)]^2[1+i(a_*+a)]| \ge 2$$

for all $a \in J_*$ and this yields

$$\begin{aligned} \|\mathcal{R}(a_* + a, f_*(\cdot, a))\|_Y &= \sup_{y \in (-1, 1)} (1 + y)(1 - y)^2 \, |\mathcal{R}(a_* + a, f_*(\cdot, a))(y) \\ &\leq \frac{1}{2} \sqrt{\|\operatorname{Re} R\|_{T([-1, 1] \times J_*)}^2 + \|\operatorname{Im} R\|_{T([-1, 1] \times J_*)}^2}. \end{aligned}$$

We compute

$$\|\operatorname{Re} R\|_{T([-1,1]\times J_*)}^2 + \|\operatorname{Im} R\|_{T([-1,1]\times J_*)}^2 \le 4\left(4\cdot 10^{-9}\right)^2$$

which implies the claimed bound.

6.2. The approximate left fundamental matrix.

Definition 6.5. Let

$$P_{L,j}(y) := \sum_{n=0}^{16} c_n(P_{L,j}) T_n(2y+1), \qquad j \in \{1, 2, 3, 4\}$$

with $(c_n(P_{L,j}))_{n=0}^{16} \subset \mathbb{C}$ given in Appendix A.2. For $y \in (-1,0]$ we set

$$f_{L,1}(y) := 1 + (1+y)P_{L,1}(y) \qquad f_{L,2}(y) := i + (1+y)P_{L,2}(y)$$

$$f_{L,3}(y) := \frac{1}{1+y}[1 + (1+y)^2P_{L,3}(y)] \qquad f_{L,4}(y) := \frac{i}{1+y}[1 + (1+y)^2P_{L,4}(y)]$$

and

$$F_L := \begin{pmatrix} \operatorname{Re} f_{L,1} & \operatorname{Re} f_{L,2} & \operatorname{Re} f_{L,3} & \operatorname{Re} f_{L,4} \\ \operatorname{Im} f_{L,1} & \operatorname{Im} f_{L,2} & \operatorname{Im} f_{L,3} & \operatorname{Im} f_{L,4} \\ \operatorname{Re} f'_{L,1} & \operatorname{Re} f'_{L,2} & \operatorname{Re} f'_{L,3} & \operatorname{Re} f'_{L,4} \\ \operatorname{Im} f'_{L,1} & \operatorname{Im} f'_{L,2} & \operatorname{Im} f'_{L,3} & \operatorname{Im} f'_{L,4} \end{pmatrix}$$

Definition 6.6. We set $P_{d,L}(y) := (1+y)^4 \det(F_L(y)).$

Remark 6.7. Note that deg $P_{d,L} \leq 72$.

Lemma 6.8. We have

$$\min_{[-1,0]} P_{d,L} \ge \frac{99}{100}.$$

In particular, F_L is an admissible left fundamental matrix.
Proof. It suffices to compute

$$\min_{[-1,0],\frac{1}{100}} P_{d,L} = 1.$$

6.3. The approximate right fundamental matrix.

Definition 6.9 (Polynomials for fundamental system). Let

$$P_{R,j}(y) := \sum_{n=0}^{30} c_n(P_{R,j}) T_n(2y-1), \qquad j \in \{1, 2, 3, 4\}$$

and

$$Q_{R,j}(y) := \sum_{n=0}^{26} c_n(Q_{R,j})T_n(2y-1), \qquad j \in \{3,4\},$$

where the coefficients $(c_n(P_{R,j}))_{n=0}^{30} \subset \mathbb{C}$ and $(c_n(Q_{R,j}))_{n=0}^{26} \subset \mathbb{C}$ are given in Appendix A.3. For $(y,a) \in [0,1) \times \mathbb{R} \setminus \{-a_*\}$ we set

$$\begin{split} f_{R,1}(y,a) &:= 1 + \frac{a_* + a + i}{2(a_* + a)}(1 - y) + (1 - y)^2 P_{R,1}(y) \\ f_{R,2}(y,a) &:= i + i \frac{a_* + a + i}{2(a_* + a)}(1 - y) + (1 - y)^2 P_{R,2}(y) \\ f_{R,3}(y,a) &:= e^{i\varphi(y,a_* + a)}(1 - y) \left[1 + \frac{2(a_* + a) - i}{2(a_* + a)}(1 - y) + (1 - y)^2 P_{R,3}(y) \right] \\ &+ e^{-i\varphi(y,a_* + a)}(1 - y)^5 Q_{R,3}(y) \\ f_{R,4}(y,a) &:= i e^{i\varphi(y,a_* + a)}(1 - y) \left[1 + \frac{2(a_* + a) - i}{2(a_* + a)}(1 - y) + (1 - y)^2 P_{R,4}(y) \right] \\ &+ i e^{-i\varphi(y,a_* + a)}(1 - y)^5 Q_{R,4}(y) \end{split}$$

and define

$$F_{R}(y,a) := \begin{pmatrix} \operatorname{Re} f_{R,1}(y,a) & \operatorname{Re} f_{R,2}(y,a) & \operatorname{Re} f_{R,3}(y,a) & \operatorname{Re} f_{R,4}(y,a) \\ \operatorname{Im} f_{R,1}(y,a) & \operatorname{Im} f_{R,2}(y,a) & \operatorname{Im} f_{R,3}(y,a) & \operatorname{Im} f_{R,4}(y,a) \\ \operatorname{Re} f'_{R,1}(y,a) & \operatorname{Re} f'_{R,2}(y,a) & \operatorname{Re} f'_{R,3}(y,a) & \operatorname{Re} f'_{R,4}(y,a) \\ \operatorname{Im} f'_{R,1}(y,a) & \operatorname{Im} f'_{R,2}(y,a) & \operatorname{Im} f'_{R,3}(y,a) & \operatorname{Im} f'_{R,4}(y,a) \end{pmatrix}$$

Definition 6.10 (Polynomials for fundamental matrix). For $(y, a) \in [0, 1] \times \mathbb{R}$ and $j \in \{3, 4\}$, we set

$$P_{C,j}(y,a) := (a_* + a)(1-y) \left[2 - y + (1-y)^2 \operatorname{Re} P_{R,j}(y) + (1-y)^4 \operatorname{Re} Q_{R,j}(y) \right]$$
$$P_{S,j}(y,a) := (1-y)^2 \left[\frac{1}{2} - (a_* + a)(1-y) \operatorname{Im} P_{R,j}(y) + (a_* + a)(1-y)^3 \operatorname{Im} Q_{R,j}(y) \right]$$

as well as

$$Q_{C,j}(y,a) := -(1-y)^2 \left[\frac{1}{2} - (a_* + a)(1-y) \operatorname{Im} P_{R,j}(y) - (a_* + a)(1-y)^3 \operatorname{Im} Q_{R,j}(y) \right]$$
$$Q_{S,j}(y,a) := (a_* + a)(1-y) \left[2 - y + (1-y)^2 \operatorname{Re} P_{R,j}(y) - (1-y)^4 \operatorname{Re} Q_{R,j}(y) \right].$$

Remark 6.11. Note that deg $P_{X,j} \leq (33,1)$ and deg $Q_{X,j} \leq (33,1)$ for $X \in \{C,S\}$ and $j \in \{3,4\}$.

Definition 6.12. (Polynomials for derivatives of fundamental system) For $(y, a) \in [0, 1) \times \mathbb{R}$ and $j \in \{3, 4\}$, we set

$$P_{C,j}^{(1)}(y,a) := (a_* + a)(1 - y)^2 \left[\varphi'(y, a_* + a) P_{S,j}(y,a) + P_{C,j}'(y,a) \right]$$
$$P_{S,j}^{(1)}(y,a) := (a_* + a)(1 - y)^2 \left[-\varphi'(y, a_* + a) P_{C,j}(y,a) + P_{S,j}'(y,a) \right]$$

as well as

$$Q_{C,j}^{(1)}(y,a) := (a_* + a)(1 - y)^2 \left[\varphi'(y, a_* + a)Q_{S,j}(y,a) + Q'_{C,j}(y,a) \right]$$
$$Q_{S,j}^{(1)}(y,a) := (a_* + a)(1 - y)^2 \left[-\varphi'(y, a_* + a)Q_{C,j}(y,a) + Q'_{S,j}(y,a) \right].$$

Remark 6.13. By explicit evaluation it follows that

$$P_{C,j}(1,a) = P_{S,j}(1,a) = Q_{C,j}(1,a) = Q_{S,j}(1,a) = 0$$

for $j \in \{3,4\}$ and $a \in \{0,1\}$ and thus, the functions defined in Definition 6.14 are polynomials of degree at most (34,3).

Definition 6.14 (Polynomials for decomposition of fundamental matrix). For $(y, a) \in [0, 1) \times J_*$, we set

$$P_{0}(y,a) := \begin{pmatrix} (a_{*}+a)\operatorname{Re} f_{R,1}(y,a) & (a_{*}+a)\operatorname{Re} f_{R,2}(y,a) & 0 & 0\\ (a_{*}+a)\operatorname{Im} f_{R,1}(y,a) & (a_{*}+a)\operatorname{Im} f_{R,2}(y,a) & 0 & 0\\ (a_{*}+a)\operatorname{Re} f'_{R,1}(y,a) & (a_{*}+a)\operatorname{Re} f'_{R,2}(y,a) & 0 & 0\\ (a_{*}+a)\operatorname{Im} f'_{R,1}(y,a) & (a_{*}+a)\operatorname{Im} f'_{R,2}(y,a) & 0 & 0 \end{pmatrix},$$

$$P_{C}(y,a) := \begin{pmatrix} 0 & 0 & (a_{*}+a)(1-y)^{2}P_{C,3}(y,a) & -(a_{*}+a)(1-y)^{2}Q_{C,4}(y,a)\\ 0 & 0 & (a_{*}+a)(1-y)^{2}Q_{C,3}(y,a) & (a_{*}+a)(1-y)^{2}P_{C,4}(y,a)\\ 0 & 0 & P_{C,3}^{(1)}(y,a) & -Q_{C,4}^{(1)}(y,a)\\ 0 & 0 & Q_{C,3}^{(1)}(y,a) & P_{C,4}^{(1)}(y,a) \end{pmatrix},$$

and

$$P_{S}(y,a) := \begin{pmatrix} 0 & 0 & (a_{*}+a)(1-y)^{2}P_{S,3}(y,a) & -(a_{*}+a)(1-y)^{2}Q_{S,4}(y,a) \\ 0 & 0 & (a_{*}+a)(1-y)^{2}Q_{S,3}(y,a) & (a_{*}+a)(1-y)^{2}P_{S,4}(y,a) \\ 0 & 0 & P_{S,3}^{(1)}(y,a) & -Q_{S,4}^{(1)}(y,a) \\ 0 & 0 & Q_{S,3}^{(1)}(y,a) & P_{S,4}^{(1)}(y,a) \end{pmatrix}$$

Remark 6.15. Note that $\deg(P_X)^j{}_k \leq (35,3)$ for $X \in \{0, C, S\}$ and $j, k \in \{1, 2, 3, 4\}$. Lemma 6.16. We have the decomposition

$$F_R(y,a) = \frac{1}{a_* + a} P_0(y,a) + \frac{\cos(\varphi(y,a_* + a))}{(a_* + a)^2(1 - y)^2} P_C(y,a) + \frac{\sin(\varphi(y,a_* + a))}{(a_* + a)^2(1 - y)^2} P_S(y,a).$$

Proof. By definition, we have

$$\operatorname{Re} f_{R,3}(y,a) = \frac{\cos(\varphi(y,a_*+a))}{a_*+a} P_{C,3}(y,a) + \frac{\sin(\varphi(y,a_*+a))}{a_*+a} P_{S,3}(y,a)$$
$$\operatorname{Im} f_{R,3}(y,a) = \frac{\cos(\varphi(y,a_*+a))}{a_*+a} Q_{C,3}(y,a) + \frac{\sin(\varphi(y,a_*+a))}{a_*+a} Q_{S,3}(y,a)$$

and

$$\operatorname{Re} f_{R,4}(y,a) = -\frac{\cos(\varphi(y,a_*+a))}{a_*+a} Q_{C,4}(y,a) - \frac{\sin(\varphi(y,a_*+a))}{a_*+a} Q_{S,4}(y,a)$$
$$\operatorname{Im} f_{R,4}(y,a) = \frac{\cos(\varphi(y,a_*+a))}{a_*+a} P_{C,4}(y,a) + \frac{\sin(\varphi(y,a_*+a))}{a_*+a} P_{S,4}(y,a).$$

Furthermore,

$$\operatorname{Re} f_{R,3}'(y,a) = \frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} P_{C,3}^{(1)}(y,a) + \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} P_{S,3}^{(1)}(y,a)$$
$$\operatorname{Im} f_{R,3}'(y,a) = \frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} Q_{C,3}^{(1)}(y,a) + \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} Q_{S,3}^{(1)}(y,a)$$

as well as

$$\operatorname{Re} f_{R,4}'(y,a) = -\frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} Q_{C,4}^{(1)}(y,a) - \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} Q_{S,4}^{(1)}(y,a)$$
$$\operatorname{Im} f_{R,4}'(y,a) = \frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} P_{C,4}^{(1)}(y,a) + \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2} P_{S,4}^{(1)}(y,a)$$

and the stated decomposition follows.

Definition 6.17. We set

$$P_{d,R} := \det\left(P_0 + P_C\right).$$

Remark 6.18. Note that deg $P_{d,R} \leq (140, 12)$.

Lemma 6.19. We have

$$(a_* + a)^6 (1 - y)^4 \det(F_R(y, a)) = P_{d,R}(y, a) + \varepsilon_{d,R}(y, a)$$

for all $(y, a) \in [0, 1) \times J_*$, where

$$\sup_{(y,a)\in[0,1)\times J_*} \left| \frac{\varepsilon_{d,R}(y,a)}{(1-y)^2} \right| \le 10^{-17}$$

and

$$\min_{[0,1]\times J_*} P_{d,R} \ge 8$$

In particular, the matrix F_R is an admissible right fundamental matrix on J_* with parameter a_* . Proof. By Lemma 6.16 and multilinearity, we obtain the decomposition

$$(a_* + a)^6 (1 - y)^4 \det(F_R(y, a)) = \cos(\varphi(y, a_* + a))^2 P_{d,R}(y, a) + \cos(\varphi(y, a_* + a)) \sin(\varphi(y, a_* + a)) Q_2(y, a) + \sin(\varphi(y, a_* + a))^2 Q_3(y, a) = P_{d,R}(y, a) + \cos(\varphi(y, a_* + a)) \sin(\varphi(y, a_* + a)) Q_2(y, a) + \sin(\varphi(y, a_* + a))^2 [Q_3(y, a) - P_{d,R}(y, a)],$$

where we abbreviate

$$Q_2 := \frac{1}{2} \det \left(P_0 + P_C + P_S \right) - \frac{1}{2} \det \left(P_0 + P_C - P_S \right)$$

and $Q_3 := \det(P_0 + P_S)$. Note that $\deg P_{d,R} \leq (140, 12)$ and $\deg Q_j \leq (140, 12)$ for $j \in \{2, 3\}$. By explicit evaluation, we find

$$\partial_1^j Q_3(1,a) - \partial_1^j P_{d,R}(1,a) = \partial_1^j Q_2(1,a) = 0$$

for $j \in \{0, 1\}$ and $a \in \{0, 1, 2, \dots, 12\}$. Consequently,

$$(y,a) \mapsto \frac{Q_3(y,a) - P_{d,R}(y,a)}{(1-y)^2}, \qquad (y,a) \mapsto \frac{Q_2(y,a)}{(1-y)^2}$$

| _ | | |
|-----|--|--|
| - 1 | | |
| | | |
| - 1 | | |

are polynomials and computing their T-norms on $[0,1] \times J_*$ yields the claimed bound on $\varepsilon_{d,R}$. Furthermore, we compute

$$\min_{[0,1]\times J_*,\frac{1}{100}} P_{d,R} \ge 8.02,$$

which finishes the proof.

6.4. The global fundamental matrix.

Definition 6.20. For $(y, a) \in (-1, 1) \times \mathbb{R} \setminus \{-a_*\}$ we set

$$F(y,a) := 1_{(-1,0]}(y)F_L(y)M_F(a) + 1_{(0,1)}(y)F_R(y,a)$$

with $M_F(a) = F_L(0)^{-1} F_R(0, a)$.

Remark 6.21. Note that by Remark 6.15, Lemma 6.16, and $\varphi(0, a_* + a) = 0$ for all $a \in J_*$, it follows that $(a_* + a)^2 M_F(a)$ is a matrix with polynomial entries of degree at most 4 in a.

Lemma 6.22. The matrix F is an admissible fundamental matrix on J_* with parameter a_* .

Proof. Let F_L and F_R denote the left and right part of F, respectively. By Lemmas 6.8 and 6.19, we only have to show that

$$0 \neq [F_L(0)^{-1}F_R(0,a)]^3{}_1 = \sum_{\ell=1}^4 [F_L(0)^{-1}]^3{}_\ell F_R(0,a)^\ell{}_1$$

for all $a \in J_*$. Note that $\varphi(0, a_* + a) = 0$ for all $a \in \mathbb{R} \setminus \{-a_*\}$ and thus, by Lemma 6.16,

$$F_R(0,a) = \frac{P_0(0,a)}{a_* + a} + \frac{P_C(0,a)}{(a_* + a)^2}.$$

This shows that $a \mapsto (a_* + a)^2 F_R(0, a)^{\ell_1}$ is a polynomial of degree at most 4 and so is $P(a) := (a_* + a)^2 [F_L(0)^{-1} F_R(0, a)]^3_1$. We compute

$$\max_{J_*,\frac{1}{10}} P \le -\frac{1}{5}.$$

7. Explicit bounds on the operator norms

7.1. The operator norm of α_F^k .

Definition 7.1 (Polynomials for $\operatorname{adj}(F_R(y, a))$). For $j \in \{1, 2\}$ and $k \in \{1, 2, 3, 4\}$, we set

$$(Q_C)^{j_k} := 0 \qquad (Q_C)^{j+2}{}_k := \operatorname{adj} (P_0 + P_C)^{j+2}{}_k (Q_S)^{j_k} := 0 \qquad (Q_S)^{j+2}{}_k := \operatorname{adj} (P_0 + P_S)^{j+2}{}_k$$

as well as

$$\begin{aligned} (Q_{CC})^{j}{}_{k} &:= \operatorname{adj} (P_{0} + P_{C})^{j}{}_{k} & (Q_{CC})^{j+2}{}_{k} &:= 0 \\ (Q_{CS})^{j}{}_{k} &:= \frac{1}{2} \operatorname{adj} (P_{0} + P_{C} + P_{S})^{j}{}_{k} - \frac{1}{2} \operatorname{adj} (P_{0} + P_{C} - P_{S})^{j}{}_{k} & (Q_{CS})^{j+2}{}_{k} &:= 0 \\ (Q_{SS})^{j}{}_{k} &:= \operatorname{adj} (P_{0} + P_{S})^{j}{}_{k} & (Q_{SS})^{j+2}{}_{k} &:= 0. \end{aligned}$$

Remark 7.2. Note that $\deg(Q_X)^{j_k} \leq (105,9)$ and $\deg(Q_{XY})^{j_k} \leq (105,9)$ for $X \in \{C, S\}$, $XY \in \{CC, CS, SS\}$, and $j, k \in \{1, 2, 3, 4\}$.

Lemma 7.3. We have the decomposition

$$\begin{aligned} \operatorname{adj}(F_R(y,a)) &= \frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^4(1-y)^2} Q_C(y,a) + \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^4(1-y)^2} Q_S(y,a) \\ &+ \frac{\cos(\varphi(y,a_*+a))^2}{(a_*+a)^5(1-y)^4} Q_{CC}(y,a) + \frac{\sin(\varphi(y,a_*+a))^2}{(a_*+a)^5(1-y)^4} Q_{SS}(y,a) \\ &+ \frac{\cos(\varphi(y,a_*+a))\sin(\varphi(y,a_*+a))}{(a_*+a)^5(1-y)^4} Q_{CS}(y,a). \end{aligned}$$

Proof. By definition, Lemma 6.16, and multilinearity, we have

$$\begin{aligned} \operatorname{adj}(F_{R}(y,a))^{1}_{k} &= \det\left(\delta_{k}, \frac{P_{0}(y,a)_{2}}{a_{*}+a}, \frac{\cos(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{C}(y,a)_{3}, \frac{\cos(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{C}(y,a)_{4}\right) \\ &+ \det\left(\delta_{k}, \frac{P_{0}(y,a)_{2}}{a_{*}+a}, \frac{\cos(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{C}(y,a)_{3}, \frac{\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{S}(y,a)_{4}\right) \\ &+ \det\left(\delta_{k}, \frac{P_{0}(y,a)_{2}}{a_{*}+a}, \frac{\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{S}(y,a)_{3}, \frac{\cos(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{C}(y,a)_{4}\right) \\ &+ \det\left(\delta_{k}, \frac{P_{0}(y,a)_{2}}{a_{*}+a}, \frac{\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{S}(y,a)_{3}, \frac{\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)^{2}(1-y)^{2}}P_{S}(y,a)_{4}\right) \end{aligned}$$

and accordingly for $\operatorname{adj}(F_R(y, a))^2_k$. Furthermore,

$$\operatorname{adj}(F_R(y,a))^3_k = \det\left(\frac{P_0(y,a)_1}{a_*+a}, \frac{P_0(y,a)_2}{a_*+a}, \delta_k, \frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2}P_C(y,a)_4\right) \\ + \det\left(\frac{P_0(y,a)_1}{a_*+a}, \frac{P_0(y,a)_2}{a_*+a}, \delta_k, \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^2(1-y)^2}P_S(y,a)_4\right)$$

and accordingly for $\operatorname{adj}(F_R(y, a))^4_k$. Thus, the claim follows straightforwardly by multilinearity and the observation that

$$\frac{1}{2} \operatorname{adj} (P_0 + P_C + P_S)^1_k - \frac{1}{2} \operatorname{adj} (P_0 + P_C - P_S)^1_k$$

= $\frac{1}{2} \operatorname{det} (\delta_k, (P_0)_2, (P_C)_3 + (P_S)_3, (P_C)_4 + (P_S)_4)$
- $\frac{1}{2} \operatorname{det} (\delta_k, (P_0)_2, (P_C)_3 - (P_S)_3, (P_C)_4 - (P_S)_4)$
= $\operatorname{det} (\delta_k, (P_0)_2, (P_C)_3, (P_S)_4) + \operatorname{det} (\delta_k, (P_0)_2, (P_S)_3, (P_C)_4).$

Proposition 7.4. We have the bounds

 $C^1_{\alpha}(F) \le 0.2, \qquad C^2_{\alpha}(F) \le 0.51, \qquad C^3_{\alpha}(F) \le 0.47, \qquad C^4_{\alpha}(F) \le 0.47.$

Proof. From Lemma 4.2, we have the bound

$$C_{\alpha}^{k}(F) \leq \sup_{(y,a)\in(-1,1)\times J_{*}} \frac{|F^{-1}(y,a)^{k}_{3}| + |F^{-1}(y,a)^{k}_{4}|}{(1+y)(1-y)^{2}}.$$

As always, we denote by F_L and F_R the left and right part of F, respectively. We distinguish between $y \in (-1, 0]$ and $y \in (0, 1)$ and start with the latter case. By Lemma 6.19, we have

$$\frac{F^{-1}(y,a)}{(1+y)(1-y)^2} = \frac{\operatorname{adj}(F_R(y,a))}{(1+y)(1-y)^2 \operatorname{det}(F_R(y,a))} = \frac{(a_*+a)^6(1-y)^2 \operatorname{adj}(F_R(y,a))}{(1+y)[P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)]}$$

and Lemma 7.3 yields the bound

$$(a_{*}+a)^{6}(1-y)^{2} |\operatorname{adj}(F_{R}(y,a))^{k}_{\ell}| \leq (a_{*}+a)^{2} |Q_{C}(y,a)^{k}_{\ell}| + (a_{*}+a)^{2} |Q_{S}(y,a)^{k}_{\ell}| + \left| \frac{(a_{*}+a)Q_{CC}(y,a)^{k}_{\ell}}{(1-y)^{2}} \right| + \left| \frac{(a_{*}+a)Q_{CS}(y,a)^{k}_{\ell}}{(1-y)^{2}} \right| + \left| \frac{(a_{*}+a)[Q_{SS}(y,a)^{k}_{\ell} - Q_{CC}(y,a)^{k}_{\ell}]}{(1-y)^{2}} \right|$$
(7.1)

and by explicit evaluation, we find

$$\partial_1^j Q_{XY}(1,a)^k{}_\ell = 0$$

for $k \in \{1, 2\}, \ell \in \{3, 4\}, j \in \{0, 1\}, a \in \{0, 1, 2, \dots, 9\}$, and $XY \in \{CC, CS, SS\}$. Consequently, since $\deg(Q_{XY})^k_{\ell} \leq (105, 9)$ by Remark 7.2, we see that

$$(y,a) \mapsto \frac{(a_*+a)Q_{XY}(y,a)^k_{\ell}}{(1-y)^2}$$

are polynomials of degree at most (105, 10). We compute

$$\left\| (y,a) \mapsto \frac{(a_* + a)Q_{CS}(y,a)^k{}_{\ell}}{(1-y)^2} \right\|_{T([0,1] \times J_*)} \le 10^{-18}$$
$$\left\| (y,a) \mapsto \frac{(a_* + a)[Q_{SS}(y,a)^k{}_{\ell} - Q_{CC}(y,a)^k{}_{\ell}]}{(1-y)^2} \right\|_{T([0,1] \times J_*)} \le 10^{-18}$$

and thus, since $(Q_{XY})^k_{\ell} = 0$ for $XY \in \{CC, CS, SS\}$ and $k \in \{3, 4\}$, the last two terms in Eq. (7.1) are negligible. Next, we compute

$$\max_{\substack{[0,1]\times J_{*,1}\\[0,1]\times J_{*,1}}} \left| (y,a) \mapsto \frac{(a_{*}+a)\frac{Q_{CC}(y,a)^{1}_{3}}{(1-y)^{2}}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.045, \quad \max_{\substack{[0,1]\times J_{*,1}\\[0,1]\times J_{*,1}}} \left| (y,a) \mapsto \frac{(a_{*}+a)\frac{Q_{CC}(y,a)^{2}_{3}}{(1-y)^{2}}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.19, \quad \max_{\substack{[0,1]\times J_{*,1}\\[0,1]\times J_{*,1}}} \left| (y,a) \mapsto \frac{(a_{*}+a)\frac{Q_{CC}(y,a)^{2}_{4}}{(1-y)^{2}}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.28.$$

Furthermore, $(Q_C)^k_{\ell} = (Q_S)^k_{\ell} = 0$ for $k \in \{1, 2\}$ and

$$\max_{\substack{[0,1]\times J_{*},1\\[0,1]\times J_{*},1}} \left| (y,a) \mapsto \frac{(a_{*}+a)^{2}Q_{C}(y,a)^{3}{}_{3}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.08, \quad \max_{\substack{[0,1]\times J_{*},1\\[0,1]\times J_{*},1}} \left| (y,a) \mapsto \frac{(a_{*}+a)^{2}Q_{C}(y,a)^{4}{}_{3}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.15, \quad \max_{\substack{[0,1]\times J_{*},1\\[0,1]\times J_{*},1\\[0,1$$

as well as

$$\max_{\substack{[0,1]\times J_{*},1\\[0,1]\times J_{*},1}} \left| (y,a) \mapsto \frac{(a_{*}+a)^{2}Q_{S}(y,a)^{3}{}_{3}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.15, \quad \max_{\substack{[0,1]\times J_{*},1\\[0,1]\times J_{*},1}} \left| (y,a) \mapsto \frac{(a_{*}+a)^{2}Q_{S}(y,a)^{4}{}_{3}}{(1+y)P_{d,R}(y,a)} \right| \leq 0.08, \quad \max_{\substack{[0,1]\times J_{*},1\\[0,1]\times J_{*},1\\[0,1$$

With $\|\varepsilon_{d,R}\|_{L^{\infty}([0,1)\times J_*)} \leq 10^{-17}$ and $\min_{[0,1]\times J_*} P_{d,R} \geq 8$ from Lemma 6.19, this yields the bounds

$$\sup_{\substack{(y,a)\in(0,1)\times J_*\\(y,a)\in(0,1)\times J_*}} \frac{|F^{-1}(y,a)^1_3| + |F^{-1}(y,a)^1_4|}{(1+y)(1-y)^2} \le 0.2$$

$$\sup_{\substack{(y,a)\in(0,1)\times J_*\\(1+y)(1-y)^2}} \frac{|F^{-1}(y,a)^2_3| + |F^{-1}(y,a)^2_4|}{(1+y)(1-y)^2} \le 0.51$$

and

$$\sup_{\substack{(y,a)\in(0,1)\times J_{*}}} \frac{|F^{-1}(y,a)^{3}_{3}| + |F^{-1}(y,a)^{3}_{4}|}{(1+y)(1-y)^{2}} \leq 0.47$$
$$\sup_{\substack{(y,a)\in(0,1)\times J_{*}}} \frac{|F^{-1}(y,a)^{4}_{3}| + |F^{-1}(y,a)^{4}_{4}|}{(1+y)(1-y)^{2}} \leq 0.47$$

Next, we turn to the left-hand side $y \in (-1, 0]$. In this case, we have

$$\frac{F^{-1}(y,a)}{(1+y)(1-y)^2} = \frac{M_F(a)^{-1}F_L(y)^{-1}}{(1+y)(1-y)^2} = M_F(a)^{-1}\frac{(1+y)^3\operatorname{adj}(F_L(y))}{(1-y)^2P_{d,L}(y)}$$

and thus,

$$\frac{F^{-1}(y,a)^k_{\ell}}{(1+y)(1-y)^2} = \sum_{j=1}^4 \frac{\operatorname{adj}(M_F(a))^k_j (1+y)^3 \operatorname{adj}(F_L(y))^j_{\ell}}{\det(M_F(a))(1-y)^2 P_{d,L}(y)}.$$

Note that for $k \in \{1, 2, 3, 4\}$ and $\ell \in \{3, 4\}$, $y \mapsto (1+y)^3 \operatorname{adj}(F_L(y))^k_\ell$ are polynomials of degree at most 54 and $\operatorname{deg}(y \mapsto (1-y)^2 P_{d,L}(y)) \leq 74$. Furthermore, by Remark 6.21, $(a_* + a)^2 M_F(a)$ is a matrix with polynomial entries of degree at most 4. Consequently, $a \mapsto \operatorname{det}((a_* + a)^2 M_F(a))$ is a polynomial of degree at most 16 and with this information, we compute

$$\max_{\substack{[-1,0]\times J_{*},1}} \left| (y,a) \mapsto \frac{F^{-1}(y,a)^{1}_{3}}{(1+y)(1-y)^{2}} \right| \leq 0.07, \qquad \max_{\substack{[-1,0]\times J_{*},1}} \left| (y,a) \mapsto \frac{F^{-1}(y,a)^{1}_{4}}{(1+y)(1-y)^{2}} \right| \leq 0.1, \\
\max_{\substack{[-1,0]\times J_{*},1}} \left| (y,a) \mapsto \frac{F^{-1}(y,a)^{2}_{3}}{(1+y)(1-y)^{2}} \right| \leq 0.16, \qquad \max_{\substack{[-1,0]\times J_{*},1}} \left| (y,a) \mapsto \frac{F^{-1}(y,a)^{2}_{4}}{(1+y)(1-y)^{2}} \right| \leq 0.34,$$

as well as

$$\max_{\substack{[-1,0]\times J_{*},1\\[-1,0]\times J_{*},1}} \left| (y,a) \mapsto \frac{F^{-1}(y,a)^{3}_{3}}{(1+y)(1-y)^{2}} \right| \leq 0.09, \qquad \max_{\substack{[-1,0]\times J_{*},1\\[-1,0]\times J_{*},1\\[-1,0]\times J_{*},1}} \left| (y,a) \mapsto \frac{F^{-1}(y,a)^{4}_{3}}{(1+y)(1-y)^{2}} \right| \leq 0.09, \qquad \max_{\substack{[-1,0]\times J_{*},1\\[-1,0]\times J_$$

Furthermore, we compute

$$\min_{[-1,0]\times J_*,1} \left((y,a) \mapsto \det(M_F(a))(1-y)^2 P_{d,L}(y) \right) \ge 50$$

and the claimed bounds follow.

7.2. The operator norm of ψ_F .

Lemma 7.5. We have $C_{\psi}(F) \leq 13$.

Proof. By Lemma 4.4, we have

$$C_{\psi}(F) \leq 2 \max_{a \in J} \sum_{k=3}^{4} \left| M_{F}(a)^{4} \frac{M_{F}(a)^{3}_{k}}{M_{F}(a)^{3}_{1}} - M_{F}(a)^{4}_{k} \right| C_{\alpha}^{k}(F)$$

= $2 \max_{a \in J} \sum_{k=3}^{4} \left| \frac{(a_{*} + a)^{4} M_{F}(a)^{4} M_{F}(a)^{3}_{k} - (a_{*} + a)^{4} M_{F}(a)^{3}_{1} M_{F}(a)^{4}_{k}}{(a_{*} + a)^{4} M_{F}(a)^{3}_{1}} \right| C_{\alpha}^{k}(F)$

and from Remark 6.21 we recall that $(a_* + a)^2 M_F(a)$ has polynomial entries of degree at most 4. We compute

$$\max_{J_*,\frac{1}{1000}} \left| a \mapsto \frac{(a_*+a)^4 M_F(a)^4 M_F(a)^3 - (a_*+a)^4 M_F(a)^3 M_F(a)^4}{(a_*+a)^4 M_F(a)^3} \right| \le 9.9$$

$$\max_{J_*,\frac{1}{1000}} \left| a \mapsto \frac{(a_*+a)^4 M_F(a)^4 M_F(a)^3 - (a_*+a)^4 M_F(a)^3 M_F(a)^4}{(a_*+a)^4 M_F(a)^3} \right| \le 3.5$$

as well as

$$\min_{J_*, 10^{-4}} \left(a \mapsto \left| (a_* + a)^4 M_F(a)^3{}_1 \right| \right) \ge 0.2008$$

and the claimed bound follows.

7.3. The operator norm of \mathcal{J}_F .

Proposition 7.6. We have $C_{\mathcal{J}}(F) \leq 232$.

Proof. By Definition 4.5 and Proposition 7.4, we have

$$\beta_{F,L}^n(y,a) \le \sum_{k=1}^2 \sum_{m=1}^2 \frac{|P_k^{nm}(y,a)|}{(a_*+a)^2} + \sum_{k=3}^4 \sum_{m=1}^2 \frac{|Q_k^{nm}(y,a)|}{|(a_*+a)^4 M_F(a)^3_1|} + \sum_{k=3}^4 \sum_{m=1}^2 \frac{|R_k^{nm}(y,a)|}{|(a_*+a)^4 M_F(a)^3_1|} + \sum_{k=3}^4 \sum_{m=1}^4 \frac{|R_k^{nm}(y,a)|}{|(a_*+a)^4 M_F(a)^4} + \sum_{m=1}^4 \sum_{m=1}^4 \sum_{m=1}^4 \frac{|R_k^{nm}(y,a)|}{|(a_*+a)^4 M_F(a)^4} + \sum_{m=1}^4 \sum$$

with

$$\begin{split} P_k^{nm}(y,a) &\coloneqq c_k(1+y)(1-y^2)^n(a_*+a)^2[F_L(y)M_F(a)]^{m+2n}{}_k \\ Q_k^{nm}(y,a) &\coloneqq c_k(1+y)(1-y^2)^n(a_*+a)^4 \\ &\times \left(\sum_{\ell=1}^3 F_L(y)^{m+2n}{}_\ell M_F(a)^\ell{}_1 M_F(a)^3{}_k + F_L(y)^{m+2n}{}_4 M_F(a)^4{}_k M_F(a)^3{}_1\right) \\ R_k^{nm}(y,a) &\coloneqq c_k(1-y)(1-y^2)^n(a_*+a)^4 \\ &\times \sum_{\ell=1}^2 F_L(y)^{m+2n}{}_\ell \left(M_F(a)^\ell{}_1 M_F(a)^3{}_k - M_F(a)^\ell{}_k M_F(a)^3{}_1\right) \end{split}$$

and $(c_k) = (\frac{1}{5}, \frac{51}{100}, \frac{47}{100}, \frac{47}{100})$. By Remark 6.21, X_k^{nm} for $X \in \{P, Q, R\}$ are polynomials with deg $X_k^{nm} \leq (20, 12)$. Furthermore, from the proof of Lemma 7.5 we recall the estimate $|(a_* + a)^4 M_F(a)^3_1| \geq \frac{1}{5}$ and with $(a_* + a)^2 \geq \frac{84}{100}$, we obtain

$$\sum_{n=0}^{1} \sup_{(-1,-\frac{1}{2})\times J_{*})} \beta_{F,L}^{n} \leq \frac{100}{84} \sum_{n=0}^{1} \sum_{k=1}^{2} \sum_{m=1}^{2} \|P_{k}^{nm}\|_{T([-1,-\frac{1}{2}]\times J_{*})} + 5 \sum_{n=0}^{1} \sum_{k=3}^{4} \sum_{m=1}^{2} \|Q_{k}^{nm}\|_{T([-1,-\frac{1}{2}]\times J_{*})} + 5 \sum_{n=0}^{1} \sum_{k=3}^{4} \sum_{m=1}^{2} \|R_{k}^{nm}\|_{T([-1,-\frac{1}{2}]\times J_{*})} \leq 54.$$

Next, we have

$$\begin{split} \beta_{F,M}^n(y,a) &\leq \sum_{k=1}^2 \sum_{m=1}^2 \frac{|P_k^{nm}(y,a)|}{(a_*+a)^2} + \sum_{k=3}^4 \sum_{m=1}^2 \frac{|Q_k^{nm}(y,a)|}{|(a_*+a)^4 M_F(a)^3_1|} \\ &+ \sum_{k=3}^4 \sum_{m=1}^2 \frac{|R_k^{nm}(y,a)|}{|(a_*+a)^4 M_F(a)^3_1|(1+y)} + \sum_{m=1}^2 \frac{|p^{nm}(y)|}{1+y} + \sum_{m=1}^2 |q^{nm}(y)| \end{split}$$

with

$$P_k^{nm}(y,a) := c_k(1+y)(1-y^2)^n (a_*+a)^2 [F_L(y)M_F(a)]^{m+2n} k$$

$$Q_k^{nm}(y,a) := c_k(1+y)(1-y^2)^n (a_*+a)^4 [F_L(y)M_F(a)]^{m+2n} M_F(a)^3 k$$

$$R_k^{nm}(y,a) := c_k(1-y^2)^{1+n} (a_*+a)^4 \sum_{\ell \in \{1,2,4\}} F_L(y)^{m+2n} \ell \left(M_F(a)^\ell M_F(a)^3 - M_F(a)^3 M_F(a)^\ell k \right)$$

as well as

$$p^{nm}(y) := 13(1+y)(1-y^2)^n F_L(y)^{m+2n} q^{nm}(y) := 39n(1-y^2)^n F_L(y)^m q_4.$$

We have $\deg(X_k^{nm}) \leq (21, 12)$ for $X \in \{P, Q, R\}$ and compute

$$\begin{split} \|P_k^{nm} - P_k^{nm}(\cdot, 0)\|_{T([-\frac{1}{2}, 0] \times J_*)} &\leq 3 \cdot 10^{-10} & (n, m, k) \in \{0, 1\} \times \{1, 2\} \times \{1, 2\} \\ \|Q_k^{nm} - Q_k^{nm}(\cdot, 0)\|_{T([-\frac{1}{2}, 0] \times J_*)} &\leq 2 \cdot 10^{-9} & (n, m, k) \in \{0, 1\} \times \{1, 2\} \times \{3, 4\} \\ \|R_k^{nm} - R_k^{nm}(\cdot, 0)\|_{T([-\frac{1}{2}, 0] \times J_*)} &\leq 2 \cdot 10^{-9} & (n, m, k) \in \{0, 1\} \times \{1, 2\} \times \{3, 4\}, \end{split}$$

as well as

$$\|\partial_1 X_k^{nm}(\cdot, 0)\|_{T([-\frac{1}{2}, 0])} \le 9, \qquad \|(q^{nm})'\|_{T([-\frac{1}{2}, 0])} \le 134.$$

Furthermore,

$$||y \mapsto (1+y)\partial_1 R_k^{nm}(y,0) - R_k^{nm}(y,0)||_{T([-\frac{1}{2},0]]} \le 6$$

and

$$||y \mapsto (1+y)(p^{nm})'(y) - p^{nm}(y)||_{T([-\frac{1}{2},0])} \le 43.$$

With this information and the bound

$$\begin{split} \beta_{F,M}^n(y,a) &\leq \frac{100}{84} \sum_{k=1}^2 \sum_{m=1}^2 |P_k^{nm}(y,a)| + 5 \sum_{k=3}^4 \sum_{m=1}^2 |Q_k^{nm}(y,a)| \\ &+ 5 \sum_{k=3}^4 \sum_{m=1}^2 \frac{|R_k^{nm}(y,a)|}{1+y} + \sum_{m=1}^2 \frac{|p^{nm}(y)|}{1+y} + \sum_{m=1}^2 |q^{nm}(y)|, \end{split}$$

we compute by explicit evaluation the estimates

$$\sup_{(-\frac{1}{2},0)\times J_*} \beta_{F,M}^0 \le 74, \qquad \qquad \sup_{(-\frac{1}{2},0)\times J_*} \beta_{F,M}^1 \le 158.$$

Finally, for $y \in (0,1)$, we have $F(y,a) = F_R(y,a)$ and by Lemma 6.16, we obtain the bound

$$\begin{split} \beta_{F,R}^n(y,a) &\leq \sum_{k=1}^2 \sum_{m=1}^2 \sum_{X \in \{0,C,S\}} \frac{|P_{X,k}^{nm}(y,a)|}{(a_*+a)^2} + \sum_{k=3}^4 \sum_{m=1}^2 \sum_{X \in \{0,C,S\}} \frac{|Q_{X,k}^{nm}(y,a)|}{(a_*+a)^4 M_F(a)^{3}_1} \\ &+ \sum_{k=3}^4 \sum_{m=1}^2 \sum_{X \in \{0,C,S\}} \frac{|R_{X,k}^{nm}(y,a)|}{(a_*+a)^4 M_F(a)^{3}_1} \end{split}$$

with

$$P_{0,k}^{nm}(y,a) := c_k(a_* + a)(1+y)(1-y^2)^n P_0(y,a)^{m+2n} k$$
$$P_{X,k}^{nm}(y,a) := c_k \frac{1+y}{(1-y)^2} (1-y^2)^n P_X(y,a)^{m+2n} k,$$

$$\begin{aligned} Q_{0,k}^{nm}(y,a) &:= c_k (1-y)(1-y^2)^n (a_*+a)^3 \left[M_F(a)^3{}_k P_0(y,a)^{m+2n}{}_1 - M_F(a)^3{}_1 P_0(y,a)^{m+2n}{}_k \right] \\ Q_{X,k}^{nm}(y,a) &:= c_k \frac{(1-y^2)^n}{1-y} (a_*+a)^2 \left[M_F(a)^3{}_k P_X(y,a)^{m+2n}{}_1 - M_F(a)^3{}_1 P_X(y,a)^{m+2n}{}_k \right], \end{aligned}$$

and

$$R_{0,k}^{nm}(y,a) := c_k (1+y)(1-y^2)^n (a_*+a)^3 M_F(a)^3{}_k P_0(y,a)^{m+2n} {}_1$$
$$R_{X,k}^{nm}(y,a) := c_k \frac{1+y}{(1-y)^2} (1-y^2)^n (a_*+a)^2 M_F(a)^3{}_k P_X(y,a)^{m+2n} {}_1$$

for $X \in \{C, S\}$. By construction and Remarks 6.15 and 6.21, these are all polynomials of degree at most (38, 8). We compute

$$\begin{split} \sum_{n=0}^{1} \sup_{(0,1)\times J_{*}} \beta_{F,R}^{n} &\leq \frac{100}{84} \sum_{n=0}^{1} \sum_{k=1}^{2} \sum_{m=1}^{2} \sum_{X \in \{0,C,S\}} \|P_{X,k}^{nm}\|_{T([0,1]\times J_{*})} \\ &+ 5 \sum_{n=0}^{1} \sum_{k=3}^{4} \sum_{m=1}^{2} \sum_{X \in \{0,C,S\}} \|Q_{X,k}^{nm}\|_{T([0,1]\times J_{*})} \\ &+ 5 \sum_{n=0}^{1} \sum_{k=3}^{4} \sum_{m=1}^{2} \sum_{X \in \{0,C,S\}} \|R_{X,k}^{nm}\|_{T([0,1]\times J_{*})} \\ &\leq 171 \end{split}$$

and Proposition 4.7 yields the claimed estimate.

8. ESTIMATES ON THE COEFFICIENTS

8.1. The left-hand side.

Lemma 8.1. For the coefficients (p_1, p_2, q_1, q_2) associated to F, we have the bounds

$$\sup_{\substack{(y,a)\in(-1,0)\times J_*}} (1-y) |p_1(y,a) - p_0(y,a_*+a)| \le 3 \cdot 10^{-7}$$
$$\sup_{\substack{(y,a)\in(-1,0)\times J_*}} (1-y) |p_2(y,a)| \le 2 \cdot 10^{-7}$$

as well as

$$\sup_{\substack{(y,a)\in(-1,0)\times J_{*}}} (1+y)(1-y)^{2} \left| q_{1}(y,a) - q_{0}(y,a_{*}+a) - \frac{2|f_{*}(y,a)|^{2}}{(1-y)^{2}} \right| \leq 7 \cdot 10^{-7}$$
$$\sup_{\substack{(y,a)\in(-1,0)\times J_{*}}} (1+y)(1-y)^{2} \left| q_{2}(y,a) - \frac{f_{*}(y,a)^{2}}{(1-y)^{2}} \right| \leq 2 \cdot 10^{-6}.$$

Proof. Let $y \in (-1, 0)$. By definition, we have

$$F'(y,a)F(y,a)^{-1} = F'_L(y)M_F(a)M_F(a)^{-1}F_L(y)^{-1} = F'_L(y)F_L(y)^{-1} = F'_L(y)\frac{\operatorname{adj}(F_L(y))}{\operatorname{det}(F_L(y))}$$
$$= \frac{(1+y)^4F'_L(y)\operatorname{adj}(F_L(y))}{P_{d,L}(y)} = \frac{1}{(1+y)^3}\frac{P_L(y)}{P_{d,L}(y)}$$

with $P_L(y) := (1+y)^7 F'_L(y) \operatorname{adj}(F_L(y))$. Note that P_L is a matrix with polynomial entries of degree at most 74. Consequently, for $f \in \{p_1, p_2, q_1, q_2\}$, we obtain

$$f(y,a) = -\frac{1}{2P_{d,L}(y)} \frac{P_L^f(y)}{(1+y)^3}$$

with

$$\begin{split} P_L^{p_1}(y) &:= P_L(y)^3{}_3 + P_L(y)^4{}_4 + iP_L(y)^4{}_3 - iP_L(y)^3{}_4 \\ P_L^{p_2}(y) &:= P_L(y)^3{}_3 - P_L(y)^4{}_4 + iP_L(y)^4{}_3 + iP_L(y)^3{}_4 \\ P_L^{q_1}(y) &:= P_L(y)^3{}_1 + P_L(y)^4{}_2 + iP_L(y)^4{}_1 - iP_L(y)^3{}_2 \\ P_L^{q_2}(y) &:= P_L(y)^3{}_1 - P_L(y)^4{}_2 + iP_L(y)^4{}_1 + iP_L(y)^3{}_2, \end{split}$$

see Definition 3.20. Next, we recall that

$$p_0(y, a_* + a) = \frac{4i(a_* + a)}{(1 - y)^3} - \frac{2i(a_* + a)}{(1 - y)^2} - \frac{2 + \frac{2i}{a_* + a}}{1 - y} + \frac{2}{1 + y}$$
$$q_0(y, a_* + a) = -\frac{2 - 2i(a_* + a)}{(1 - y)^3} - \frac{\frac{1}{(a_* + a)^2} - \frac{i}{a_* + a}}{(1 - y)^2} - \frac{1 + \frac{i}{a_* + a}}{1 - y} - \frac{1 + \frac{i}{a_* + a}}{1 + y}$$

as well as

$$f_*(y,a) = P_*(y) + \frac{2i(a_*+a)}{1-i(a_*+a)}P'_*(1) - P_*(1),$$

where P_* is a polynomial of degree 50. Based on this, we obtain

$$\begin{aligned} f_1(y,a) &:= (1-y)[p_1(y,a) - p_0(y,a_*+a)] = \frac{P_L^{f_1}(y,a)}{2(a_*+a)(1-y)^2 P_{d,L}(y)} \\ f_2(y,a) &:= (1-y)p_2(y,a) = \frac{P_L^{f_2}(y)}{2P_{d,L}(y)} \end{aligned}$$

and

$$g_{1}(y,a) := (1+y)(1-y)^{2} \left[q_{1}(y,a) - q_{0}(y,a_{*}+a) - \frac{2|f_{*}(y,a)|^{2}}{(1-y)^{2}} \right]$$

$$= \frac{P_{L}^{g_{1}}(y,a)}{2(a_{*}+a)^{2}|1-i(a_{*}+a)|^{2}(1-y)P_{d,L}(y)}$$

$$g_{2}(y,a) := (1+y)(1-y)^{2} \left[q_{2}(y,a) - \frac{f_{*}(y,a)^{2}}{(1-y)^{2}} \right] = \frac{P_{L}^{g_{2}}(y,a)}{2[1-i(a_{*}+a)]^{2}P_{d,L}(y)},$$

where

$$P_L^{f_1}(y,a) := -(a_* + a) \frac{(1-y)^3 P_L^{p_1}(y)}{(1+y)^3} - 2(a_* + a)(1-y)^3 P_{d,L}(y) p_0(y,a_* + a)$$
$$P_L^{f_2}(y) := -\frac{(1-y) P_L^{p_2}(y)}{(1+y)^3}$$

and

$$P_L^{g_1}(y,a) := -(a_* + a)^2 |1 - i(a_* + a)|^2 \Big[\frac{(1 - y)^3 P_L^{q_1}(y)}{(1 + y)^2} + 2(1 + y)(1 - y)^3 P_{d,L}(y) q_0(y, a_* + a) + 4(1 + y)(1 - y) P_{d,L}(y) |f_*(y, a)|^2 \Big]$$
$$P_L^{g_2}(y,a) := -[1 - i(a_* + a)]^2 \left[\frac{(1 - y)^2 P_L^{q_2}(y)}{(1 + y)^2} + 2(1 + y) P_{d,L}(y) f_*(y, a)^2 \right].$$

By explicit evaluation, we find

$$P_L^{p_1}(-1) = (P_L^{p_1})'(-1) = P_L^{p_2}(-1) = (P_L^{p_2})'(-1) = (P_L^{p_2})''(-1) = 0$$

as well as

$$P_L^{q_1}(-1) = (P_L^{q_1})'(-1) = P_L^{q_2}(-1) = (P_L^{q_2})'(-1) = 0$$

and thus, by Proposition 3.21, P_L^f for $f \in \{f_1, f_2, g_1, g_2\}$ are polynomials with deg $P_L^f \leq (174, 6)$. We compute

$$\begin{split} \left\| P_L^{f_1} \right\|_{T([-1,0] \times J_*)} &\leq 8.8 \cdot 10^{-7}, \\ \left\| P_L^{g_1} \right\|_{T([-1,0] \times J_*)} &\leq 4 \cdot 10^{-6}, \\ \\ \left\| P_L^{g_2} \right\|_{T([-1,0] \times J_*)} &\leq 4.7 \cdot 10^{-6}, \\ \end{split}$$

and recall that $\min_{[-1,0]} P_{d,L} \geq \frac{99}{100}$ (Lemma 6.8). Furthermore,

$$\min_{[-1,0],\frac{1}{100}} \left[y \mapsto (1-y) P_{d,L}(y) \right] \ge 2$$

and from these bounds, the claimed estimates follow.

8.2. The right-hand side.

Definition 8.2 (Polynomials for $F'_R(y, a)$). For $(y, a) \in (0, 1) \times J_*$, we set

$$P_C^{(1)}(y,a) := (a_* + a)(1 - y)^3 \varphi'(y, a_* + a) P_S(y,a) + 2(a_* + a)(1 - y)^2 P_C(y,a) + (a_* + a)(1 - y)^3 P_C'(y,a)$$

and

$$P_S^{(1)}(y,a) := -(a_* + a)(1-y)^3 \varphi'(y,a_* + a) P_C(y,a) + 2(a_* + a)(1-y)^2 P_S(y,a) + (a_* + a)(1-y)^3 P'_S(y,a),$$

see Definition 6.14.

Remark 8.3. Note that $\deg(P_X^{(1)})^j{}_k \leq (37,5)$ for $X \in \{C,S\}$ and $j,k \in \{1,2,3,4\}$. **Definition 8.4** (Polynomials for $F'_R(y,a)F_R(y,a)^{-1}$). For $(y,a) \in (0,1) \times J_*$, we set

$$P_{CC}(y,a) := P_C^{(1)}(y,a)Q_C(y,a) + (a_* + a)(1-y)^3 P_0'(y,a)Q_{CC}(y,a)$$

$$P_{CS}(y,a) := P_C^{(1)}(y,a)Q_S(y,a) + P_S^{(1)}(y,a)Q_C(y,a) + (a_* + a)(1-y)^3 P_0'(y,a)Q_{CS}(y,a)$$

$$P_{SS}(y,a) := P_S^{(1)}(y,a)Q_S(y,a) + (a_* + a)(1-y)^3 P_0'(y,a)Q_{SS}(y,a),$$

see Definition 7.1.

Remark 8.5. Note that $\deg(P_{XY})^j{}_k \leq (142, 14)$ for $XY \in \{CC, CS, SS\}$ and $j, k \in \{1, 2, 3, 4\}$. **Definition 8.6** (Polynomials for coefficients). For $XY \in \{CC, CS, SS\}$, we define

$$P_{XY}^{p_1} := -\frac{1}{2} \left[(P_{XY})^3{}_3 + (P_{XY})^4{}_4 \right] - \frac{i}{2} \left[(P_{XY})^4{}_3 - (P_{XY})^3{}_4 \right] P_{XY}^{p_2} := -\frac{1}{2} \left[(P_{XY})^3{}_3 - (P_{XY})^4{}_4 \right] - \frac{i}{2} \left[(P_{XY})^4{}_3 + (P_{XY})^3{}_4 \right]$$

as well as

$$P_{XY}^{q_1} := -\frac{1}{2} \left[(P_{XY})^3{}_1 + (P_{XY})^4{}_2 \right] - \frac{i}{2} \left[(P_{XY})^4{}_1 - (P_{XY})^3{}_2 \right] P_{XY}^{q_2} := -\frac{1}{2} \left[(P_{XY})^3{}_1 - (P_{XY})^4{}_2 \right] - \frac{i}{2} \left[(P_{XY})^4{}_1 + (P_{XY})^3{}_2 \right].$$

Lemma 8.7. For $j \in \{1, 2\}$ we have the bounds

$$\sup_{(y,a)\in[0,1)\times J_*} \left| \frac{P_{CC}^{p_j}(y,a) - P_{SS}^{p_j}(y,a)}{(1-y)^2} \right| \le 10^{-16}, \qquad \sup_{(y,a)\in[0,1)\times J_*} \left| \frac{P_{CS}^{p_j}(y,a)}{(1-y)^2} \right| \le 10^{-16},$$

and

$$\sup_{(y,a)\in[0,1)\times J_*} \left| \frac{P_{CC}^{q_j}(y,a) - P_{SS}^{q_j}(y,a)}{1-y} \right| \le 10^{-15}, \qquad \sup_{(y,a)\in[0,1)\times J_*} \left| \frac{P_{CS}^{q_j}(y,a)}{1-y} \right| \le 10^{-15}.$$

Proof. By explicit evaluation we check that

$$P_{CC}^{f}(1,a) - P_{SS}^{f}(1,a) = (P_{CC}^{f})'(1,a) - (P_{SS}^{f})'(1,a) = P_{CS}^{f}(1,a) = (P_{CS}^{f})'(1,a) = 0$$

for $f \in \{p_1, p_2, q_1, q_2\}$, $j \in \{1, 2\}$, and $a \in \{0, 1, 2, \dots, 14\}$. Thus, since $\deg_2 P_{XY}^f \leq 14$ for $XY \in \{CC, SS, CS\}$, we see that

$$(y,a) \mapsto \frac{P_{CC}^{p_j}(y,a) - P_{SS}^{p_j}(y,a)}{(1-y)^2}, \qquad (y,a) \mapsto \frac{P_{CS}^{p_j}(y,a)}{(1-y)^2}$$

and

$$(y,a) \mapsto \frac{P_{CC}^{q_j}(y,a) - P_{SS}^{q_j}(y,a)}{1 - y}, \qquad (y,a) \mapsto \frac{P_{CS}^{q_j}(y,a)}{1 - y}$$

are polynomials and by computing their T-norms on $[0,1] \times J_*$, the claim follows.

Lemma 8.8. Let (p_1, p_2, q_1, q_2) be the coefficients associated to F. Then, for $(y, a) \in (0, 1) \times J_*$ and $f \in \{p_1, p_2, q_1, q_2\}$, we have

$$f(y,a) = \frac{\cos(\varphi(y,a_*+a))^2}{(a_*+a)(1-y)^3} \frac{P_{CC}^f(y,a)}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)} + \frac{\sin(\varphi(y,a_*+a))^2}{(a_*+a)(1-y)^3} \frac{P_{SS}^f(y,a)}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)} + \frac{\cos(\varphi(y,a_*+a))\sin(\varphi(y,a_*+a))}{(a_*+a)(1-y)^3} \frac{P_{CS}^f(y,a)}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)}.$$

Proof. Recall the decomposition

$$F_R(y,a) = \frac{1}{a_* + a} P_0(y,a) + \frac{\cos(\varphi(y,a_* + a))}{(a_* + a)^2(1 - y)^2} P_C(y,a) + \frac{\sin(\varphi(y,a_* + a))}{(a_* + a)^2(1 - y)^2} P_S(y,a)$$

from Lemma 6.16, which immediately yields

$$F_{R}'(y,a) = \frac{1}{a_{*}+a}P_{0}'(y,a) + \frac{\cos(\varphi(y,a_{*}+a))}{(a_{*}+a)^{3}(1-y)^{5}}P_{C}^{(1)}(y,a) + \frac{\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)^{3}(1-y)^{5}}P_{S}^{(1)}(y,a).$$

Next, recall that

 $(Q_{CC})^{j+2}{}_k = (Q_{CS})^{j+2}{}_k = (Q_{SS})^{j+2}{}_k = (Q_C)^j{}_k = (Q_S)^j{}_k = (P_0)^k{}_{j+2} = (P_C)^k{}_j = (P_S)^k{}_j = 0$ for $j \in \{1, 2\}$ and $k \in \{1, 2, 3, 4\}$, which implies that

$$P_0'Q_C = P_0'Q_S = P_C^{(1)}Q_{CC} = P_C^{(1)}Q_{CS} = P_C^{(1)}Q_{SS} = P_S^{(1)}Q_{CC} = P_S^{(1)}Q_{CS} = P_S^{(1)}Q_{SS} = 0.$$

Consequently, in view of the decomposition from Lemma 7.3, we obtain

$$F_{R}'(y,a) \operatorname{adj}(F_{R}(y,a)) = \frac{\cos(\varphi(y,a_{*}+a))^{2}}{(a_{*}+a)^{7}(1-y)^{7}} P_{CC}(y,a) + \frac{\sin(\varphi(y,a_{*}+a))^{2}}{(a_{*}+a)^{7}(1-y)^{7}} P_{SS}(y,a) + \frac{\cos(\varphi(y,a_{*}+a))\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)^{7}(1-y)^{7}} P_{CS}(y,a),$$

which, by Lemma 6.19, yields

$$\begin{split} F_{R}'(y,a)F_{R}(y,a)^{-1} &= F_{R}'(y,a)\frac{\operatorname{adj}(F_{R}(y,a))}{\det(F_{R}(y,a))} = F_{R}'(y,a)\frac{(a_{*}+a)^{6}(1-y)^{4}\operatorname{adj}(F_{R}(y,a))}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)} \\ &= \frac{\cos(\varphi(y,a_{*}+a))^{2}}{(a_{*}+a)(1-y)^{3}}\frac{P_{CC}(y,a)}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)} \\ &\quad + \frac{\sin(\varphi(y,a_{*}+a))^{2}}{(a_{*}+a)(1-y)^{3}}\frac{P_{SS}(y,a)}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)} \\ &\quad + \frac{\cos(\varphi(y,a_{*}+a))\sin(\varphi(y,a_{*}+a))}{(a_{*}+a)(1-y)^{3}}\frac{P_{CS}(y,a)}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)}. \end{split}$$

Thus, the claim follows by recalling Definition 3.20.

Lemma 8.9. Let (p_1, p_2, q_1, q_2) be the coefficients associated to F. Then we have the bounds

$$\sup_{\substack{(y,a)\in(0,1)\times J_*}} (1-y) |p_1(y,a) - p_0(y,a_*+a)| \le 3 \cdot 10^{-7}$$
$$\sup_{\substack{(y,a)\in(0,1)\times J_*}} (1-y) |p_2(y,a)| \le 6 \cdot 10^{-7}$$

as well as

$$\sup_{\substack{(y,a)\in(0,1)\times J_{*}}} (1+y)(1-y)^{2} \left| q_{1}(y,a) - q_{0}(y,a_{*}+a) - \frac{2|f_{*}(y,a)|^{2}}{(1-y)^{2}} \right| \leq 1.5 \cdot 10^{-6}$$
$$\sup_{\substack{(y,a)\in(0,1)\times J_{*}}} (1+y)(1-y)^{2} \left| q_{2}(y,a) - \frac{f_{*}(y,a)^{2}}{(1-y)^{2}} \right| \leq 1.3 \cdot 10^{-6}.$$

Proof. We define

$$\widetilde{f}(y,a) := \frac{P^f_{CC}(y,a)}{(a_*+a)(1-y)^3 P_{d,R}(y,a)}$$

for $f \in \{p_1, p_2, q_1, q_2\}$. By Lemmas 6.19, 8.7, 8.8, and

$$||P_{CC}^{p_j}||_{T([0,1]\times J_*)} \le 10^3, \qquad ||P_{CC}^{q_j}||_{T([0,1]\times J_*)} \le 10^4$$

for $j \in \{1, 2\}$, we infer that

$$\sup_{(y,a)\in(0,1)\times J_*} (1-y) |p_j(y,a) - \widetilde{p}_j(y,a)| \le 10^{-15}$$

and

$$\sup_{(y,a)\in(0,1)\times J_*} (1+y)(1-y)^2 |q_j(y,a) - \widetilde{q}_j(y,a)| \le 10^{-14}.$$

We set

$$f_1(y,a) := (1-y) \left[\widetilde{p}_1(y,a) - p_0(y,a_*+a) \right] =: \frac{P_R^{f_1}(y,a)}{(a_*+a)(1+y)P_{d,R}(y,a)}$$
$$f_2(y,a) := (1-y)\widetilde{p}_2(y,a) =: \frac{P_R^{f_2}(y,a)}{(a_*+a)P_{d,R}(y,a)}$$

as well as

$$g_{1}(y,a) := (1+y)(1-y)^{2} \left[\tilde{q}_{1}(y,a) - q_{0}(y,a_{*}+a) - \frac{2|f_{*}(y,a)|^{2}}{(1-y)^{2}} \right]$$
$$=: \frac{P_{R}^{g_{1}}(y,a)}{(a_{*}+a)^{2}|1-i(a_{*}+a)|^{2}P_{d,R}(y,a)}$$
$$g_{2}(y,a) := (1+y)(1-y)^{2} \left[\tilde{q}_{2}(y,a) - \frac{f_{*}(y,a)^{2}}{(1-y)^{2}} \right] =: \frac{P_{R}^{g_{2}}(y,a)}{(a_{*}+a)|1-i(a_{*}+a)|^{4}P_{d,R}(y,a)}$$

and Proposition 3.21 implies that P_R^f for $f \in \{f_1, f_2, g_1, g_2\}$ are polynomials of degree at most (245, 18). Recall from Lemma 6.19 that $\min_{[0,1] \times J_*} P_{d,R} \ge 8$. Consequently, with $\Omega_j := [\frac{j}{10}, \frac{j+1}{10}] \times J_*$, we compute

$$\|\operatorname{Re} f_1\|_{L^{\infty}((0,1)\times J_*)} \le \max\left\{\frac{\|\operatorname{Re} P_R^{f_1}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)(1+y)P_{d,R}(y,a)]-1} : j\in[0,9]\cap\mathbb{Z}\right\}$$
$$\le 2\cdot 10^{-7}$$

and

$$\|\operatorname{Im} f_1\|_{L^{\infty}((0,1)\times J_*)} \le \max\left\{\frac{\|\operatorname{Im} P_R^{f_1}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)(1+y)P_{d,R}(y,a)]-1} : j\in[0,9]\cap\mathbb{Z}\right\}$$
$$\le 2\cdot 10^{-7}$$

and

$$\|\operatorname{Re} f_2\|_{L^{\infty}((0,1)\times J_*)} \le \max\left\{\frac{\|\operatorname{Re} P_R^{f_2}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)P_{d,R}(y,a)]-1} : j\in[0,9]\cap\mathbb{Z}\right\}$$
$$\le 2\cdot 10^{-7}$$

and

$$\|\operatorname{Im} f_2\|_{L^{\infty}((0,1)\times J_*)} \le \max\left\{\frac{\|\operatorname{Im} P_R^{f_2}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)P_{d,R}(y,a)]-1} : j\in[0,9]\cap\mathbb{Z}\right\}$$
$$\le 5\cdot 10^{-7}$$

and

$$\|\operatorname{Re} g_1\|_{L^{\infty}((0,1)\times J_*)} \leq \max\left\{ \frac{\|\operatorname{Re} P_R^{g_1}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)^2|1-i(a_*+a)|^2P_{d,R}(y,a)]-1} : j \in [0,9] \cap \mathbb{Z} \right\} \leq 10^{-6}$$

and

 $\|\operatorname{Im} g_1\|_{L^{\infty}((0,1)\times J_*)}$

$$\leq \max\left\{\frac{\|\operatorname{Im} P_R^{g_1}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)^2|1-i(a_*+a)|^2P_{d,R}(y,a)]-1}: j\in[0,9]\cap\mathbb{Z}\right\}$$

$$\leq 5\cdot 10^{-7}$$

and

$$\|\operatorname{Re} g_2\|_{L^{\infty}((0,1)\times J_*)} \leq \max\left\{ \frac{\|\operatorname{Re} P_R^{g_2}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)|1-i(a_*+a)|^4 P_{d,R}(y,a)]-1} : j \in [0,9] \cap \mathbb{Z} \right\} \leq 8 \cdot 10^{-7}$$

and

$$\|\operatorname{Im} g_2\|_{L^{\infty}((0,1)\times J_*)} \leq \max \left\{ \frac{\|\operatorname{Im} P_R^{g_2}\|_{T(\Omega_j)}}{\min_{\Omega_j,1}[(y,a)\mapsto (a_*+a)|1-i(a_*+a)|^4 P_{d,R}(y,a)]-1} : j \in [0,9] \cap \mathbb{Z} \right\} \leq 10^{-6}.$$

These estimates and the triangle inequality imply the claim.

9. Estimates on the functional ψ_F

Unfortunately, the functional ψ_F involves highly oscillatory integrals which require special care. As a consequence, we need to split the domain of integration and to this end, it is convenient to introduce a more efficient notation.

Definition 9.1. For $y_0, y_1 \in [-1, 1]$ and $a \in J_*$, we define

$$I_F(a, y_0, y_1) := \sum_{k=3}^4 \int_{y_0}^{y_1} \gamma_{F,k}(a) \alpha_F^k(a, \mathcal{R}(a_* + a, f_*(\cdot, a))),$$

where

$$\gamma_{F,k}(a) := \frac{M_F(a)^4}{M_F(a)^3} M_F(a)^3_k - M_F(a)^4_k.$$

Note that

$$\psi_F(a, \mathcal{R}(a_* + a, f_*(\cdot, a))) = I_F(a, -1, 1).$$

Definition 9.2 (Polynomials for ψ_F). For $y \in [0, 1) \times J_*$, we set

$$P_{C}^{\psi}(y,a) := (a_{*}+a)^{2}(1+y)(1-y)^{2} \sum_{k=3}^{4} \gamma_{F,k}(a) \left[Q_{C}(y,a)^{k}{}_{3}\operatorname{Re}r(y,a) + Q_{C}(y,a)^{k}{}_{4}\operatorname{Im}r(y,a) \right]$$
$$P_{S}^{\psi}(y,a) := (a_{*}+a)^{2}(1+y)(1-y)^{2} \sum_{k=3}^{4} \gamma_{F,k}(a) \left[Q_{S}(y,a)^{k}{}_{3}\operatorname{Re}r(y,a) + Q_{S}(y,a)^{k}{}_{4}\operatorname{Im}r(y,a) \right],$$
here $r(y,a) := \mathcal{R}(a_{*}+a,f_{*}(\cdot,a))(y)$

where $r(y,a) := \mathcal{R}(a_* + a, f_*(\cdot, a))(y).$

Remark 9.3. Note that by definition, P_X^{ψ} for $X \in \{C, S\}$ is a polynomial in the first variable and by Remark 7.2, we have $\deg_1 P_X^{\psi} \leq 256$.

Definition 9.4. For $y_0, y_1 \in [0, 1]$ and $a \in J_*$, we set

$$\begin{split} \widetilde{I}_{F}(a, y_{0}, y_{1}) &:= \int_{y_{0}}^{y_{1}} \cos(\varphi(y, a_{*} + a)) \frac{P_{C}^{\psi}(y, a)}{(1+y)P_{d,R}(y, a)} dy \\ &+ \int_{y_{0}}^{y_{1}} \sin(\varphi(y, a_{*} + a)) \frac{P_{S}^{\psi}(y, a)}{(1+y)P_{d,R}(y, a)} dy. \end{split}$$

Furthermore, for brevity we set $a_{\pm} := \pm 10^{-10}$.

Lemma 9.5. Let $0 \le y_0 \le y_1 \le 1$. Then we have the bounds

$$\left|I_F(a_{\pm}, y_0, y_1) - \widetilde{I}_F(a_{\pm}, y_0, y_1)\right| \le 10^{-18}.$$

Proof. From Lemma 7.3 and Definition 7.1, we obtain

$$\operatorname{adj}(F_R(y,a))^j{}_k = \frac{\cos(\varphi(y,a_*+a))}{(a_*+a)^4(1-y)^2} Q_C(y,a)^j{}_k + \frac{\sin(\varphi(y,a_*+a))}{(a_*+a)^4(1-y)^2} Q_S(y,a)^j{}_k$$

for $j \in \{3, 4\}$ and $k \in \{1, 2, 3, 4\}$. Consequently, by Lemma 6.19,

$$[F_{R}(y,a)^{-1}]^{j}_{k} = \frac{\operatorname{adj}(F_{R}(y,a))^{j}_{k}}{\operatorname{det}(F_{R}(y,a))}$$
$$= \cos(\varphi(y,a_{*}+a))\frac{(a_{*}+a)^{2}(1-y)^{2}Q_{C}(y,a)^{j}_{k}}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)}$$
$$+ \sin(\varphi(y,a_{*}+a))\frac{(a_{*}+a)^{2}(1-y)^{2}Q_{S}(y,a)^{j}_{k}}{P_{d,R}(y,a) + \varepsilon_{d,R}(y,a)}$$

and by recalling that

$$\alpha_F^k(a,g)(y) = [F(y,a)^{-1}]^k{}_3\operatorname{Re} g(y) + [F(y,a)^{-1}]^k{}_4\operatorname{Im} g(y),$$

we find

$$I_F(a, y_0, y_1) = \int_{y_0}^{y_1} \cos(\varphi(y, a_* + a)) \frac{P_C^{\psi}(y, a)}{(1+y)[P_{d,R}(y, a) + \varepsilon_{d,R}(y, a)]} dy + \int_{y_0}^{y_1} \sin(\varphi(y, a_* + a)) \frac{P_S^{\psi}(y, a)}{(1+y)[P_{d,R}(y, a) + \varepsilon_{d,R}(y, a)]} dy.$$

We compute

$$\|P_C^{\psi}(\cdot, a_{\pm})\|_{T([0,1])} + \|P_S^{\psi}(\cdot, a_{\pm})\|_{T([0,1])} \le 10^{-6}.$$

Thanks to the bounds $\|\varepsilon_{d,R}\|_{L^{\infty}([0,1]\times J_*)} \leq 10^{-17}$ and $\min_{[0,1]\times J_*} P_{d,R} \geq 8$ from Lemma 6.19, we see that

$$\left| I_F(a_{\pm}, y_0, y_1) - \widetilde{I}_F(a_{\pm}, y_0, y_1) \right| \le 10^{-18},$$

as claimed.

9.1. Close to the right endpoint. Next, we show that the contribution near the irregular singularity at y = 1 is negligible.

Lemma 9.6. We have the bound

$$\left|I_F\left(a_{\pm}, \frac{9}{10}, 1\right)\right| \le 10^{-11}.$$

Proof. By Lemma 9.5, it suffices to consider \widetilde{I}_F . In order to exploit the oscillations, we write

$$\begin{split} \widetilde{I}_{F}(a, \frac{9}{10}, 1) &= \int_{\frac{9}{10}}^{1} \partial_{y} \sin(\varphi(y, a_{*} + a)) \frac{P_{C}^{\psi}(y, a)}{(1+y)P_{d,R}(y, a)\varphi'(y, a_{*} + a)} dy \\ &- \int_{\frac{9}{10}}^{1} \partial_{y} \cos(\varphi(y, a_{*} + a)) \frac{P_{S}^{\psi}(y, a)}{(1+y)P_{d,R}(y, a)\varphi'(y, a_{*} + a)} dy \end{split}$$

and integrate by parts, which yields

$$\begin{split} \widetilde{I}_{F}(a, \frac{9}{10}, 1) &= -\frac{\sin(\varphi(\frac{9}{10}, a_{*} + a))P_{C}^{\psi}(\frac{9}{10}, a)}{\frac{19}{10}P_{d,R}(\frac{9}{10}, a)\varphi'(\frac{9}{10}, a_{*} + a)} + \frac{\cos(\varphi(\frac{9}{10}, a_{*} + a))P_{S}^{\psi}(\frac{9}{10}, a)}{\frac{19}{10}P_{d,R}(\frac{9}{10}, a)\varphi'(\frac{9}{10}, a_{*} + a)} \\ &- \int_{\frac{9}{10}}^{1}\sin(\varphi(y, a_{*} + a))\partial_{y}\left[\frac{P_{C}^{\psi}(y, a)}{(1 + y)P_{d,R}(y, a)\varphi'(y, a_{*} + a)}\right]dy \\ &+ \int_{\frac{9}{10}}^{1}\cos(\varphi(y, a_{*} + a))\partial_{y}\left[\frac{P_{S}^{\psi}(y, a)}{(1 + y)P_{d,R}(y, a)\varphi'(y, a_{*} + a)}\right]dy \end{split}$$

since |

$$\frac{1}{\varphi'(y,a_*+a)} = -\frac{(a_*+a)(1-y)^3}{2(a_*+a)^2(1+y) - 2(1-y)^2}$$

As a consequence, we obtain the bound

$$\begin{aligned} |\widetilde{I}_{F}(a,\frac{9}{10},1)| &\leq \frac{10}{19} \frac{|P_{C}^{\psi}(\frac{9}{10},a)| + |P_{S}^{\psi}(\frac{9}{10},a)|}{|P_{d,R}(\frac{9}{10},a)\varphi'(\frac{9}{10},a_{*}+a)|} + \frac{1}{10} \max_{y \in [\frac{9}{10},1]} \left| \partial_{y} \frac{P_{C}^{\psi}(y,a)}{(1+y)P_{d,R}(y,a)\varphi'(y,a_{*}+a)} \right| \\ &+ \frac{1}{10} \max_{y \in [\frac{9}{10},1]} \left| \partial_{y} \frac{P_{S}^{\psi}(y,a)}{(1+y)P_{d,R}(y,a)\varphi'(y,a_{*}+a)} \right|. \end{aligned}$$
(9.1)

We have

$$\partial_y \frac{P_C^{\psi}(y,a)}{(1+y)P_{d,R}(y,a)\varphi'(y,a_*+a)} = \partial_y \frac{\widetilde{P}_C^{\psi}(y,a)}{\widetilde{P}_{d,R}(y,a)} = \frac{\partial_y \widetilde{P}_C^{\psi}(y,a)\widetilde{P}_{d,R}(y,a) - \widetilde{P}_C^{\psi}(y,a)\partial_y \widetilde{P}_{d,R}(y,a)}{\widetilde{P}_{d,R}(y,a)^2}$$

with $\tilde{P}_{C}^{\psi}(y,a) := -(a_{*}+a)(1-y)^{3}P_{C}^{\psi}(y,a)$ and

$$\widetilde{P}_{d,R}(y,a) := \left[2(a_*+a)^2(1+y) - 2(1-y)^2\right](1+y)P_{d,R}(y,a)$$

Analogously, we set $\widetilde{P}_{S}^{\psi}(y,a) := -(a_* + a)(1 - y)^3 P_{S}^{\psi}(y,a)$. By Remarks 6.18 and 9.3, we have $\deg_1 \widetilde{P}_{d,R} \leq 143$ and $\deg_1 \widetilde{P}_{X}^{\psi} \leq 259$ for $X \in \{C, S\}$. Thus, we compute

$$\min_{\left[\frac{9}{10},1\right],1}\widetilde{P}_{d,R}(\cdot,a_{\pm}) \ge 54$$

and

$$\left\| \widetilde{P}_{C}^{\psi}(\cdot, a_{\pm})' \widetilde{P}_{d,R}(\cdot, a_{\pm}) - \widetilde{P}_{C}^{\psi}(\cdot, a_{\pm}) \widetilde{P}_{d,R}(\cdot, a_{\pm})' \right\|_{T([\frac{9}{10}, 1])} \leq 7 \cdot 10^{-8} \\ \left\| \widetilde{P}_{S}^{\psi}(\cdot, a_{\pm})' \widetilde{P}_{d,R}(\cdot, a_{\pm}) - \widetilde{P}_{S}^{\psi}(\cdot, a_{\pm}) \widetilde{P}_{d,R}(\cdot, a_{\pm})' \right\|_{T([\frac{9}{10}, 1])} \leq 8 \cdot 10^{-8}.$$

These bounds imply

$$\begin{split} \max_{y \in [\frac{9}{10},1]} \left| \partial_y \frac{P_C^{\psi}(y, a_{\pm})}{(1+y) P_{d,R}(y, a_{\pm}) \varphi'(y, a_{\ast} + a_{\pm})} \right| + \max_{y \in [\frac{9}{10},1]} \left| \partial_y \frac{P_S^{\psi}(y, a_{\pm})}{(1+y) P_{d,R}(y, a) \varphi'(y, a_{\ast} + a)} \right| \\ & \leq \frac{1.5 \cdot 10^{-7}}{53^2} \leq 6 \cdot 10^{-11}. \end{split}$$

Finally, by explicit evaluation, we find

$$\frac{10}{19} \frac{|P_C^{\psi}(\frac{9}{10}, a_{\pm})| + |P_S^{\psi}(\frac{9}{10}, a_{\pm})|}{|P_{d,R}(\frac{9}{10}, a_{\pm})\varphi'(\frac{9}{10}, a_{\ast} + a_{\pm})|} \le 10^{-12}$$

and the claim follows from Eq. (9.1).

9.2. The highly oscillatory regime. Next, we consider $I_F(a_{\pm}, \frac{7}{10}, \frac{9}{10})$. This is the most delicate regime with a high number of oscillations that need to be resolved. The basis for the treatment is provided by the following general observation.

Lemma 9.7. Let a < b and $f \in C([a,b])$. Furthermore, let $g \in C^1([a,b])$ be real-valued with |g'| > 0 on [a,b] and let $p : \mathbb{C} \to \mathbb{C}$ be a polynomial. Then we have the bound

$$\left| \int_{a}^{b} e^{ig(x)} f(x) dx - e^{ig(a)} \sum_{k=0}^{\deg p} i^{k+1} p^{(k)}(g(a)) + e^{ig(b)} \sum_{k=0}^{\deg p} i^{k+1} p^{(k)}(g(b)) \right| \le \left\| f - (p \circ g)g' \right\|_{L^{1}(a,b)}$$

Proof. Since |g'| > 0 on [a, b], g is invertible and by a change of variables, we find

$$\int_{a}^{b} e^{ig(x)} f(x) dx = \int_{g(a)}^{g(b)} e^{iy} \frac{f(g^{-1}(y))}{g'(g^{-1}(y))} dy = \int_{g(a)}^{g(b)} e^{iy} p(y) dy + \int_{g(a)}^{g(b)} e^{iy} \left[\frac{f(g^{-1}(y))}{g'(g^{-1}(y))} - p(y) \right] dy.$$

By repeated integration by parts, we obtain

$$\int_{g(a)}^{g(b)} e^{iy} p(y) dy = \sum_{k=0}^{\deg p} i^{k+1} \left[e^{ig(a)} p^{(k)}(g(a)) - e^{ig(b)} p^{(k)}(g(b)) \right]$$

and the claim follows from the estimate

$$\left| \int_{g(a)}^{g(b)} e^{iy} \left[\frac{f(g^{-1}(y))}{g'(g^{-1}(y))} - p(y) \right] dy \right| = \left| \int_{a}^{b} e^{ig(x)} \left[f(x) - p(g(x))g'(x) \right] dx \right| \le \|f - (p \circ g)g'\|_{L^{1}(a,b)}.$$

Remark 9.8. Lemma 9.7 provides a means to compute oscillatory integrals to a high degree of precision provided one is able to find a polynomial p such that $||f - (p \circ g)g'||_{L^1(a,b)}$ is sufficiently small. The point is that the function g is not oscillatory and thus, if f is well-behaved, there is no need to approximate oscillatory functions as would be the case if one applied a standard quadrature method.

In order to implement this idea for our problem, we first need a polynomial approximation to the phase φ .

Definition 9.9. For $y \in [0, \frac{9}{10}]$, we set

$$\widetilde{\varphi}(y) := \sum_{n=0}^{40} c_n(\widetilde{\varphi}) T_n\left(\frac{10}{9} \left(2y - \frac{9}{10}\right)\right),$$

where the coefficients $(c_n(\widetilde{\varphi}))_{n=0}^{40} \subset \mathbb{Q}$ are given in Appendix A.4.

Lemma 9.10. We have the bounds

$$\|\varphi(\cdot, a_* + a_{\pm}) - \widetilde{\varphi}\|_{L^{\infty}(0, \frac{9}{10})} \le 10^{-7}.$$

Proof. From the summation formula of the geometric series we have the identity

$$\log(1-y) + \sum_{k=1}^{n} \frac{y^k}{k} = -\int_0^y \frac{x^n}{1-x} dx,$$

which yields the bound

$$\left|\log(1-y) + \sum_{k=1}^{n} \frac{y^k}{k}\right| \le 10 \int_0^{\frac{9}{10}} x^n dx = \frac{10}{n+1} \left(\frac{9}{10}\right)^{n+1}$$

for all $y \in [0, \frac{9}{10}]$ and $n \in \mathbb{N}$. Consequently, by setting

$$\widehat{\varphi}_{\pm}(y) := -\frac{2(a_* + a_{\pm})}{(1 - y)^2} + \frac{2(a_* + a_{\pm})}{1 - y} + \frac{2}{a_* + a_{\pm}} \sum_{k=1}^{200} \frac{y^k}{k},$$

we obtain

$$\|\varphi(\cdot, a_* + a_{\pm}) - \widehat{\varphi}_{\pm}\|_{L^{\infty}(0, \frac{9}{10})} \le 7 \cdot 10^{-11}.$$

Furthermore,

$$\widehat{\varphi}_{\pm}(y) - \widetilde{\varphi}(y) = \frac{P_{\pm}(y)}{(a_* + a_{\pm})(1 - y)^2},$$

with

$$P_{\pm}(y) := -2(a_* + a_{\pm})^2 + 2(a_* + a_{\pm})^2(1 - y) + 2(1 - y)^2 \sum_{k=1}^{200} \frac{y^k}{k} - (a_* + a_{\pm})(1 - y)^2 \widetilde{\varphi}(y)$$

000

and we compute

$$\max_{[0,\frac{9}{10}], 5 \cdot 10^{-12}} \left| y \mapsto \frac{P_{\pm}(y)}{(a_* + a_{\pm})(1 - y)^2} \right| \le 3 \cdot 10^{-8},$$

which yields the claim.

Next, we define the polynomials that will be used to approximate the oscillatory integrals. Definition 9.11. For $x \in [-161, -10]$, we set

$$P_{\pm}^{\sharp}(x) := 10^{-10} \sum_{n=0}^{40} c_n (P_{\pm}^{\sharp}) T_n \left(\frac{2x+171}{151}\right)$$
$$Q_{\pm}^{\sharp}(x) := 10^{-10} \sum_{n=0}^{40} c_n (Q_{\pm}^{\sharp}) T_n \left(\frac{2x+171}{151}\right),$$

where the coefficients $(c_n(P_{\pm}^{\sharp}))_{n=0}^{40} \subset \mathbb{Q}$ and $(c_n(Q_{\pm}^{\sharp}))_{n=0}^{40} \subset \mathbb{Q}$ are given in Appendix A.5. Lemma 9.12. We have the bounds

$$\sup_{\substack{y \in (\frac{7}{10}, \frac{9}{10})\\ y \in (\frac{7}{10}, \frac{9}{10})}} \left| \frac{P_C^{\psi}(y, a_{\pm})}{(1+y) P_{d,R}(y, a_{\pm})} - P_{\pm}^{\sharp}\left(\widetilde{\varphi}(y)\right) \widetilde{\varphi}'(y) \right| \le 10^{-12}$$

$$\sup_{\substack{y \in (\frac{7}{10}, \frac{9}{10})\\ (1+y) P_{d,R}(y, a_{\pm})}} - Q_{\pm}^{\sharp}\left(\widetilde{\varphi}(y)\right) \widetilde{\varphi}'(y) \right| \le 10^{-12}.$$

Proof. We define

$$P_{\pm}(y) := P_C^{\psi}(y, a_{\pm}) - (1+y)P_{d,R}(y, a_{\pm})P_{\pm}^{\sharp}\left(\widetilde{\varphi}(y)\right)\widetilde{\varphi}'(y)$$
$$Q_{\pm}(y) := P_S^{\psi}(y, a_{\pm}) - (1+y)P_{d,R}(y, a_{\pm})Q_{\pm}^{\sharp}\left(\widetilde{\varphi}(y)\right)\widetilde{\varphi}'(y)$$

and note that by Remark 6.18, deg $X_{\pm} \leq 1780$ for $X \in \{P, Q\}$. We compute

$$||X_{\pm}||_{T([\frac{7}{10},\frac{9}{10}])} \le 3 \cdot 10^{-12}$$

and together with $\min_{[0,1]\times J_*} P_{d,R} \ge 8$ from Lemma 6.19, the claimed bounds follow. Lemma 9.13. We have the bounds

$$I_F(a_-, \frac{7}{10}, \frac{9}{10}) \ge 2 \cdot 10^{-11}$$

$$I_F(a_+, \frac{7}{10}, \frac{9}{10}) \le -2 \cdot 10^{-11}.$$

Proof. We explicitly compute

$$\operatorname{Re}\left[e^{i\widetilde{\varphi}(\frac{7}{10})}\sum_{k=0}^{40}i^{k+1}(P_{-}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{7}{10})) - e^{i\widetilde{\varphi}(\frac{9}{10})}\sum_{k=0}^{40}i^{k+1}(P_{-}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{9}{10}))\right] \ge 1.5 \cdot 10^{-11}$$
$$\operatorname{Im}\left[e^{i\widetilde{\varphi}(\frac{7}{10})}\sum_{k=0}^{40}i^{k+1}(Q_{-}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{7}{10})) - e^{i\widetilde{\varphi}(\frac{9}{10})}\sum_{k=0}^{40}i^{k+1}(Q_{-}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{9}{10}))\right] \ge 8 \cdot 10^{-12}$$

as well as

$$\operatorname{Re}\left[e^{i\widetilde{\varphi}(\frac{7}{10})}\sum_{k=0}^{40}i^{k+1}(P_{+}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{7}{10})) - e^{i\widetilde{\varphi}(\frac{9}{10})}\sum_{k=0}^{40}i^{k+1}(P_{+}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{9}{10}))\right] \leq -1.9 \cdot 10^{-11} \operatorname{Im}\left[e^{i\widetilde{\varphi}(\frac{7}{10})}\sum_{k=0}^{40}i^{k+1}(Q_{+}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{7}{10})) - e^{i\widetilde{\varphi}(\frac{9}{10})}\sum_{k=0}^{40}i^{k+1}(Q_{+}^{\sharp})^{(k)}(\widetilde{\varphi}(\frac{9}{10}))\right] \leq -7 \cdot 10^{-12}.$$

Consequently, Lemmas 9.7 and 9.12 yield

$$\int_{\frac{7}{10}}^{\frac{9}{10}} \cos(\widetilde{\varphi}(y)) \frac{P_C^{\psi}(y, a_-)}{(1+y)P_{d,R}(y, a_-)} dy = \operatorname{Re} \int_{\frac{7}{10}}^{\frac{9}{10}} e^{i\widetilde{\varphi}(y)} \frac{P_C^{\psi}(y, a_-)}{(1+y)P_{d,R}(y, a_-)} dy \ge 1.4 \cdot 10^{-11}$$
$$\int_{\frac{7}{10}}^{\frac{9}{10}} \sin(\widetilde{\varphi}(y)) \frac{P_S^{\psi}(y, a_-)}{(1+y)P_{d,R}(y, a_-)} dy = \operatorname{Im} \int_{\frac{7}{10}}^{\frac{9}{10}} e^{i\widetilde{\varphi}(y)} \frac{P_S^{\psi}(y, a_-)}{(1+y)P_{d,R}(y, a_-)} dy \ge 7 \cdot 10^{-12}$$

as well as

$$\int_{\frac{7}{10}}^{\frac{9}{10}} \cos(\widetilde{\varphi}(y)) \frac{P_C^{\psi}(y, a_+)}{(1+y)P_{d,R}(y, a_+)} dy = \operatorname{Re} \int_{\frac{7}{10}}^{\frac{9}{10}} e^{i\widetilde{\varphi}(y)} \frac{P_C^{\psi}(y, a_+)}{(1+y)P_{d,R}(y, a_+)} dy \le -1.8 \cdot 10^{-11}$$
$$\int_{\frac{7}{10}}^{\frac{9}{10}} \sin(\widetilde{\varphi}(y)) \frac{P_S^{\psi}(y, a_+)}{(1+y)P_{d,R}(y, a_+)} dy = \operatorname{Im} \int_{\frac{7}{10}}^{\frac{9}{10}} e^{i\widetilde{\varphi}(y)} \frac{P_S^{\psi}(y, a_+)}{(1+y)P_{d,R}(y, a_+)} dy \le -6 \cdot 10^{-12}.$$

The claim now follows by combining these bounds with Lemmas 9.10, 9.5,

$$\|P_C^{\psi}(\cdot, a_{\pm})\|_{T([0,1])} + \|P_S^{\psi}(\cdot, a_{\pm})\|_{T([0,1])} \le 10^{-6},$$

and $\min_{[0,1] \times J_*} P_{d,R} \ge 8$ (Lemma 6.19).

9.3. The remaining contribution from the right-hand side. Next, we turn to the final contribution from the right-hand side and estimate $I_F(a_{\pm}, 0, \frac{7}{10})$. Again, we need some auxiliary polynomials.

Definition 9.14. For $y \in [0, \frac{7}{10}]$, we set

$$P_{\pm}^{\flat}(y) := 10^{-8} \sum_{n=0}^{40} c_n (P_{\pm}^{\flat}) T_n \left(\frac{10}{7} (2y - \frac{7}{10})\right)$$
$$Q_{\pm}^{\flat}(y) := 10^{-8} \sum_{n=0}^{40} c_n (Q_{\pm}^{\flat}) T_n \left(\frac{10}{7} (2y - \frac{7}{10})\right),$$

where the coefficients $(c_n(P_{\pm}^{\flat}))_{n=0}^{40} \subset \mathbb{Q}$ and $(c_n(Q_{\pm}^{\flat}))_{n=0}^{40} \subset \mathbb{Q}$ are given in Appendix A.6.

Lemma 9.15. We have the bounds

$$\sup_{y \in (0, \frac{7}{10})} \left| \cos(\varphi(y, a_* + a_{\pm})) \frac{P_C^{\psi}(y, a_{\pm})}{(1+y)P_{d,R}(y, a_{\pm})} - P_{\pm}^{\flat}(y) \right| \le 1.5 \cdot 10^{-12}$$
$$\sup_{y \in (0, \frac{7}{10})} \left| \sin(\varphi(y, a_* + a_{\pm})) \frac{P_S^{\psi}(y, a_{\pm})}{(1+y)P_{d,R}(y, a_{\pm})} - Q_{\pm}^{\flat}(y) \right| \le 3.2 \cdot 10^{-12}.$$

Proof. Let $y \in [0, \frac{7}{10}]$. From Lemma 9.10, we have the bound

$$\left|\cos(\varphi(y, a_* + a_{\pm})) - \cos(\widetilde{\varphi}(y))\right| \le \left\|\varphi(\cdot, a_* + a_{\pm}) - \widetilde{\varphi}\right\|_{L^{\infty}(0, \frac{7}{10})} \le 10^{-7}.$$

Furthermore, $\widetilde{\varphi}(y) \in [-12, 0]$ and from the bound

$$\left|\cos(x) - \sum_{k=0}^{n-1} (-1)^k \frac{(x+2\pi)^{2k}}{(2k)!}\right| \le \frac{|x+2\pi|^{2n}}{(2n)!},$$

valid for all $x \in \mathbb{R}$ and $n \in \mathbb{N}$, we obtain

$$\left|\cos(\tilde{\varphi}(y)) - \sum_{k=0}^{18} (-1)^k \frac{[\tilde{\varphi}(y) + 2\tilde{\pi}]^{2k}}{(2k)!}\right| \le 10^{-12},$$

where $\widetilde{\pi} := 3.1415926535897932 \in \mathbb{Q}$. We set

$$P_{\pm}(y) := P_C^{\psi}(y, a_{\pm}) \sum_{k=0}^{18} (-1)^k \frac{[\widetilde{\varphi}(y) + 2\widetilde{\pi}]^{2k}}{(2k)!} - (1+y) P_{d,R}(y, a_{\pm}) P_{\pm}^{\flat}(y)$$

and note that by Remarks 6.18 and 9.3, deg $P_{\pm} \leq 1696$. Thus, we compute

$$||P_{\pm}||_{T([0, \frac{7}{10}])} \le 1.1 \cdot 10^{-11}$$

and by combining this with the bounds

$$\|P_C^{\psi}(\cdot, a_{\pm})\|_{T([0,1])} + \|P_S^{\psi}(\cdot, a_{\pm})\|_{T([0,1])} \le 10^{-6}$$

from the proof of Lemma 9.13 and $\min_{[0,1]} P_{d,R} \ge 8$ (Lemma 6.19), the claimed estimate for P_{\pm}^{\flat} follows from the triangle inequality. The bound for Q_{\pm}^{\flat} is proved in an analogous fashion by considering

$$Q_{\pm}(y) := P_{S}^{\psi}(y, a_{\pm}) \sum_{k=0}^{18} (-1)^{k} \frac{[\widetilde{\varphi}(y) + 2\widetilde{\pi}]^{2k+1}}{(2k+1)!} - (1+y) P_{d,R}(y, a_{\pm}) Q_{\pm}^{\flat}(y),$$
58

noting that deg $Q_{\pm} \leq 1736$ (Remarks 6.18 and 9.3), and computing

$$|Q_{\pm}||_{T([0, \frac{7}{10}])} \le 2.4 \cdot 10^{-11}.$$

Lemma 9.16. We have the bounds

$$I_F\left(a_{-}, 0, \frac{7}{10}\right) \ge 6 \cdot 10^{-10}$$
$$I_F\left(a_{+}, 0, \frac{7}{10}\right) \le -6 \cdot 10^{-10}$$

Proof. By explicit integration, we find

$$\int_{0}^{\frac{1}{10}} \left[P_{-}^{\flat}(y) + Q_{-}^{\flat}(y) \right] dy \ge 6.04 \cdot 10^{-10}$$
$$\int_{0}^{\frac{7}{10}} \left[P_{+}^{\flat}(y) + Q_{+}^{\flat}(y) \right] dy \le -6.47 \cdot 10^{-10}$$

and the claim follows by combining these estimates with Lemmas 9.15 and 9.5.

In summary, we obtain the following estimates for the contribution of the right-hand side. Corollary 9.17. We have the bounds

$$I_F(a_-, 0, 1) \ge 6 \cdot 10^{-10}$$

 $I_F(a_+, 0, 1) \le -6 \cdot 10^{-10}$

Proof. The claim follows by combining Lemmas 9.6, 9.13, and 9.16.

9.4. The contribution from the left-hand side.

Definition 9.18. For $(y, a) \in (-1, 0] \times J_*$, we set

$$P_L^{\psi}(y,a) := (1+y)^4 (1-y)^2 \sum_{k=3}^4 \gamma_{F,k}(a) \\ \times \sum_{\ell=1}^4 M_F^{-1}(a)^k{}_\ell \left[\operatorname{adj}(F_L(y))^\ell{}_3 \operatorname{Re} r(y,a) + \operatorname{adj}(F_L(y))^\ell{}_4 \operatorname{Im} r(y,a) \right].$$

Remark 9.19. Note that $P_L^{\psi}(\cdot, a_{\pm})$ is a polynomial of degree at most 205. Furthermore, since

$$F_L(y)^{-1} = \frac{(1+y)^4 \operatorname{adj}(F_L(y))}{(1+y)^4 \operatorname{det}(F_L(y))} = \frac{(1+y)^4 \operatorname{adj}(F_L(y))}{P_{d,L}(y)}$$

and $F(y,a)^{-1} = M_F(a)^{-1} F_L(y)^{-1}$ for $y \in [-1,0]$, we see that

$$I_F(a_{\pm}, -1, 0) = \int_{-1}^0 \frac{P_L^{\psi}(y, a_{\pm})}{(1-y)^2 P_{d,L}(y)} dy.$$
(9.2)

Definition 9.20. For $y \in [-1, 0]$, we set

$$P_{\pm}^{L}(y) := 10^{-10} \sum_{n=0}^{50} c_n (P_{\pm}^{L}) T_n (2y+1),$$

where the coefficients $(c_n(P_{\pm}^L))_{n=0}^{50} \subset \mathbb{Q}$ are given in Appendix A.7.

Lemma 9.21. We have the bounds

$$\sup_{y \in (-1,0)} \left| \frac{P_L^{\psi}(y, a_{\pm})}{(1-y)^2 P_{d,L}(y)} - P_{\pm}^L(y) \right| \le 3 \cdot 10^{-13}.$$

Proof. We define

$$P_{\pm}(y) := P_L^{\psi}(y, a_{\pm}) - (1-y)^2 P_{d,L}(y) P_{\pm}^L(y)$$

and note that by Remarks 6.7 and 9.19, we infer that deg $P_{\pm} \leq 205$. We compute

$$||P_{\pm}||_{T([-1,0])} \le 2.6 \cdot 10^{-13}$$

and with $\min_{[-1,0]} P_{d,L} \geq \frac{99}{100}$ from Lemma 6.8, the claimed bounds follow. Lemma 9.22. We have the bounds

$$I_F(a_-, -1, 0) \ge -2 \cdot 10^{-10}$$

 $I_F(a_+, -1, 0) \le 2 \cdot 10^{-10}$.

Proof. By explicit integration, we find

$$\begin{split} &\int_{-1}^{0}P_{-}^{L}(y)dy\geq -1.9\cdot 10^{-10}\\ &\int_{-1}^{0}P_{+}^{L}(y)dy\leq 1.9\cdot 10^{-10}, \end{split}$$

and the claim follows by combining these estimates with Lemma 9.21 and Eq. (9.2).

Corollary 9.23. We have the bounds

$$\psi_F(a_-, \mathcal{R}(a_* + a_-, f_*(\cdot, a_-))) \ge 4 \cdot 10^{-10}$$

$$\psi_F(a_+, \mathcal{R}(a_* + a_+, f_*(\cdot, a_+))) \le -4 \cdot 10^{-10}.$$

Proof. Recall that

$$\psi_F(a_{\pm}, \mathcal{R}(a_* + a_{\pm}, f_*(\cdot, a_{\pm}))) = I_F(a_{\pm}, -1, 1) = I_F(a_{\pm}, -1, 0) + I_F(a_{\pm}, 0, 1)$$

and thus, the claim follows by combining Corollary 9.17 and Lemma 9.22.

10. Proof of the main result

Finally, we are in the position to conclude the proof of the main result of this paper.

10.1. Solution of Eq. (3.2).

Lemma 10.1. We have the bound

$$\begin{aligned} \|\mathcal{N}(a_*, f_*, a, f) - \mathcal{N}(a_*, f_*, a, g)\|_Y &\leq 10 \left(\|f\|_X + \|g\|_X \right) \|f - g\|_X \\ &+ 3 \left(\|f\|_X^2 + \|g\|_X^2 \right) \|f - g\|_X \end{aligned}$$

for all $f, g \in X$ and $a \in J_*$.

Proof. Recall that

$$\mathcal{N}(a_*, f_*, a, f)(y) = \frac{\overline{f_*(y, a)}f(y)^2 + 2f_*(y, a)|f(y)|^2 + f(y)|f(y)|^2}{(1-y)^2},$$

which yields

$$\begin{aligned} (1+y)(1-y)^2 \left| \mathcal{N}(a_*, f_*, a, f)(y) - \mathcal{N}(a_*, f_*, a, g)(y) \right| \\ &\leq (1+y) |f_*(y, a)| \left| f(y)^2 - g(y)^2 \right| + 2(1+y) |f_*(y, a)| \left| |f(y)|^2 - |g(y)|^2 \right| \\ &+ (1+y) \left| f(y) |f(y)|^2 - g(y) |g(y)|^2 \right| \\ &\leq 3(1+y) |f_*(y, a)| \left[|f(y)| + |g(y)| \right] |f(y) - g(y)| + \frac{3}{2}(1+y) \left[|f(y)|^2 + |g(y)|^2 \right] |f(y) - g(y)| \end{aligned}$$

for all $(y, a) \in (-1, 1) \times J_*$ and by checking that

$$\max_{[-1,1] \times J_*, \frac{1}{10}} |(y,a) \mapsto (1+y)f_*(y,a)| \le 3.2,$$

the claim follows.

Proposition 10.2. For every $a \in J_*$ there exists a unique $f_a \in X$ with $||f_a||_X \leq 1.2 \cdot 10^{-6}$ such that

$$f_a(y) = \mathcal{J}_F\left(a, \mathcal{G}_F(a_*, f_*, a, f_a)\right)(y)$$

for all $y \in (-1,1)$. Furthermore, the map $a \mapsto f_a : J_* \to X$ is continuous.

Proof. Let $a \in J_*$ and recall that

$$\begin{aligned} \mathcal{G}_F(a_*, f_*, a, f)(y) = & [p_1(y, a) - p_0(y, a_* + a)]f'(y) + p_2(y, a)f'(y) \\ & + \left[q_1(y, a) - q_0(y, a_* + a) - \frac{2|f_*(y, a)|^2}{(1 - y)^2}\right]f(y) \\ & + \left[q_2(y, a) - \frac{f_*(y, a)^2}{(1 - y)^2}\right]\overline{f(y)} - \mathcal{N}(a_*, f_*, a, f)(y) - \mathcal{R}(a_* + a, f_*(\cdot, a))(y), \end{aligned}$$

where (p_1, p_2, q_1, q_2) are the coefficients associated to F. From Lemmas 8.1, 8.9, 10.1, and Propositions 6.4, 7.6, we obtain the bounds

$$\begin{aligned} \|\mathcal{J}_F(a, \mathcal{G}_F(a_*, f_*, a, f))\|_X &\leq C_{\mathcal{J}}(F) \|\mathcal{G}_F(a_*, f_*, a, f)\|_Y \\ &\leq C_{\mathcal{J}}(F) \left(5 \cdot 10^{-6} \|f\|_X + 10 \|f\|_X^2 + 3 \|f\|_X^3 + 5 \cdot 10^{-9}\right) \\ &\leq 1.2 \cdot 10^{-6} \end{aligned}$$

and

$$\begin{aligned} \|\mathcal{J}_{F}\left(a,\mathcal{G}_{F}(a_{*},f_{*},a,f)\right) - \mathcal{J}_{F}\left(a,\mathcal{G}_{F}(a_{*},f_{*},a,g)\right)\|_{X} \\ &\leq C_{\mathcal{J}}(F) \|\mathcal{G}_{F}(a_{*},f_{*},a,f) - \mathcal{G}_{F}(a_{*},f_{*},a,g)\|_{Y} \\ &\leq C_{\mathcal{J}}(F) \left[5 \cdot 10^{-6} + 10\left(\|f\|_{X} + \|g\|_{X}\right) + 3\left(\|f\|_{X}^{2} + \|g\|_{X}^{2}\right)\right] \|f - g\|_{X} \\ &\leq \frac{1}{2}\|f - g\|_{X} \end{aligned}$$

for all $f, g \in X$ satisfying $||f||_X, ||g||_X \leq 1.2 \cdot 10^{-6}$. Consequently, $f \mapsto \mathcal{J}_F(a, \mathcal{G}_F(a_*, f_*, a, f))$ is a contractive self-map on the ball $\{f \in X : ||f||_X \leq 1.2 \cdot 10^{-6}\}$ and the existence of a fixed point f_a is a consequence of the contraction mapping principle.

As for the continuity, it suffices to note that

$$\begin{aligned} \|f_a - f_b\|_X &= \|\mathcal{J}_F(a, \mathcal{G}_F(a_*, f_*, a, f_a)) - \mathcal{J}_F(b, \mathcal{G}_F(a_*, f_*, b, f_b))\|_X \\ &\leq \frac{1}{2} \|f_a - f_b\|_X + \|\mathcal{J}_F(a, \mathcal{G}_F(a_*, f_*, a, f_a)) - \mathcal{J}_F(b, \mathcal{G}_F(a_*, f_*, b, f_a))\|_X \end{aligned}$$

and the claim follows from the continuity of

$$b \mapsto \mathcal{J}_F(b, \mathcal{G}_F(a_*, f_*, b, f_a)) : J_* \to X,$$

see Lemmas 3.35 and 3.38.

Proposition 10.3. Let $a \mapsto f_a : J_* \to X$ be the continuous map constructed in Proposition 10.2. Then there exists an $a \in J_*$ such that

$$\psi_F\left(a, \mathcal{G}_F\left(a_*, f_*, a, f_a\right)\right) = 0.$$

Proof. By Lemmas 7.5, 8.1, 8.9, and 10.1, we obtain

$$\begin{aligned} &|\psi_F(a, \mathcal{G}_F(a_*, f_*, a, f_a)) + \psi_F(a, \mathcal{R}(a_* + a, f_*(\cdot, a)))| \\ &\leq 13 \left(5 \cdot 10^{-6} \|f_a\|_X + 10 \|f_a\|_X^2 + 3 \|f_a\|_X^3 \right) \\ &\leq 2.7 \cdot 10^{-10} \end{aligned}$$

for all $a \in J_*$ and Corollary 9.23 yields

$$\psi_F(a_-, \mathcal{G}_F(a_*, f_*, a_-, f_{a_-})) < 0, \qquad \psi_F(a_+, \mathcal{G}_F(a_*, f_*, a_+, f_{a_+})) > 0.$$

Consequently, the claim follows by applying the intermediate value theorem to the continuous map

 $a \mapsto \psi_F \left(a, \mathcal{G}_F \left(a_*, f_*, a, f_a \right) \right) : J_* \to \mathbb{R},$

see Lemmas 3.29 and 3.38.

Corollary 10.4. There exist
$$(a, f) \in J_* \times C^2(-1, 1)$$
 with $||f||_X \le 1.2 \cdot 10^{-6}$ such that
 $\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = 0$

on (-1, 1).

Proof. By Propositions 10.2 and 10.3, there exist $(a, f) \in J_* \times X$ with $||f||_X \leq 1.2 \cdot 10^{-6}$ that satisfy Eq. (3.2). By Lemma 3.39, the claim follows.

10.2. Proof of Theorem 1.4. It remains to prove the stated regularity properties.

Lemma 10.5. Let $f \in X$, $\alpha \in \mathbb{R} \setminus \{0\}$, and define $h : \mathbb{R}^3 \to \mathbb{C}$ by

$$h(x) := (1+|x|)^{-1-\frac{i}{\alpha}} f\left(\frac{|x|-1}{|x|+1}\right).$$

Then $h \in L^4(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3)$.

Proof. We define $\widetilde{h}: [0,\infty) \to \mathbb{C}$ by

$$\widetilde{h}(r) := (1+r)^{-1-\frac{i}{\alpha}} f\left(\frac{r-1}{r+1}\right).$$

Then $h(x) = \tilde{h}(|x|)$ and we have

$$\|h\|_{L^4(\mathbb{R}^3)}^4 \simeq \int_0^\infty |\tilde{h}(r)|^4 r^2 dr, \qquad \|h\|_{\dot{H}^1(\mathbb{R}^3)}^2 \simeq \int_0^\infty |\tilde{h}'(r)|^2 r^2 dr.$$

For the change of variable $r = \frac{1+y}{1-y}$, we obtain $dr = \frac{2dy}{(1-y)^2}$ and thus,

$$\|h\|_{L^4(\mathbb{R}^3)}^4 \simeq \int_{-1}^1 (1-y)^4 |f(y)|^4 \left(\frac{1+y}{1-y}\right)^2 \frac{dy}{(1-y)^2} \lesssim \|f\|_X^4$$

as well as

$$\begin{aligned} \|h\|_{\dot{H}^{1}(\mathbb{R}^{3})}^{2} &\simeq \int_{-1}^{1} (1-y)^{6} |f'(y)|^{2} \left(\frac{1+y}{1-y}\right)^{2} \frac{dy}{(1-y)^{2}} + \int_{-1}^{1} (1-y)^{4} |f(y)|^{2} \left(\frac{1+y}{1-y}\right)^{2} \frac{dy}{(1-y)^{2}} \\ &\lesssim \|f\|_{X}^{2}. \end{aligned}$$

Lemma 10.6. Let $(a, f) \in J_* \times X \cap C^2(-1, 1)$ satisfy

$$\mathcal{R}(a_* + a, f_*(\cdot, a) + f) = 0$$

on (-1,1). Define $Q: \mathbb{R}^3 \to \mathbb{C}$ by

$$Q(x) := (1+|x|)^{-1-\frac{i}{a_*+a}} \left[f_*\left(\frac{|x|-1}{|x|+1}, a\right) + f\left(\frac{|x|-1}{|x|+1}\right) \right].$$

Then $Q \in L^4(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3) \cap C^{\infty}(\mathbb{R}^3)$ and Q satisfies Eq. (1.3) on \mathbb{R}^3 .

Proof. By construction and Lemmas 3.39, 10.5, $Q \in L^4(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3) \cap C^2(\mathbb{R}^3 \setminus \{0\})$ and Q satisfies Eq. (1.3) on $\mathbb{R}^3 \setminus \{0\}$. By Hölder's inequality, $Q \in H^1_{\text{loc}}(\mathbb{R}^3)$ and elliptic regularity yields $Q \in C^{\infty}(\mathbb{R}^3)$.

Finally, the stated estimate on the function g in Theorem 1.4 is a consequence of the following bound.

Lemma 10.7. Let $f \in X$ and define $g : [0, \infty) \to \mathbb{C}$ by

$$g(r) := f\left(\frac{r-1}{r+1}\right).$$

Then we have

$$2\sup_{r>0} r|g'(r)| + \sup_{r>0} |g(r)| \le ||f||_X.$$

Proof. By definition, we have

$$g'(r) = \frac{2}{(r+1)^2} f'\left(\frac{r-1}{r+1}\right)$$

and thus,

$$|2rg'(r)| = \left|\frac{4r}{(r+1)^2}f'\left(\frac{r-1}{r+1}\right)\right| \le \sup_{y \in (-1,1)}(1-y^2)|f'(y)|$$

for all r > 0. This implies the claim.

APPENDIX A. TABLES OF COEFFICIENTS

A.1. Coefficients for the approximate solution. These are the coefficients used in Definitions 1.3 and 6.1 (files Re_Pstar.dat and Im_Pstar.dat).

| n | $\operatorname{Re} c_n(P_*)$ | $\operatorname{Im} c_n(P_*)$ | n | $\operatorname{Re} c_n(P_*)$ | $\operatorname{Im} c_n(P_*)$ |
|----|---|--|----|---|--|
| 0 | $-\tfrac{2721451470460628}{1505677490833565}$ | $-rac{1078867762011625}{1320610492513519}$ | 26 | $-\tfrac{23584979}{942423027627962}$ | $\tfrac{16782531}{1527543763014338}$ |
| 1 | $\frac{591830832555107}{1325882795067685}$ | $-\tfrac{503523409161778}{1215525208542319}$ | 27 | $-\frac{10363411}{752762158039191}$ | $-\frac{9122195}{1273036310590791}$ |
| 2 | $\frac{33309682807931}{117192328161321}$ | $\frac{249431537821016}{429463288550741}$ | 28 | $-\frac{1037623}{523068233364354}$ | $-\frac{7695059}{900322851895972}$ |
| 3 | $-\frac{123831666850433}{762568250071312}$ | $\frac{58969483417430}{562787431135597}$ | 29 | $\frac{2636394}{920776893246139}$ | $-\frac{2427135}{627015566263018}$ |
| 4 | $-\frac{94362296895241}{1495365721952099}$ | $-\frac{208278440111958}{1709495256184139}$ | 30 | $\frac{1043029}{389679881702888}$ | $-\frac{137069}{885954612787875}$ |
| 5 | 27826388742075 756461404086487 | $-\frac{32027304107051}{811755765327665}$ | 31 | 939092 870036758617377 | $\frac{1397610}{1301624802001117}$ |
| 6 | 28327529175143 1200014475775705 | <u>21608223588887</u> 067282652010512 | 32 | $-\frac{36787}{026268202520441}$ | 650042 774472262050485 |
| 7 | $- \frac{4593572122680}{001729855187701}$ | $\frac{11604846306461}{81405212245211224521122}$ | 33 | $-\frac{174055}{481707012301070}$ | $\frac{210261}{711166266247052}$ |
| 8 | $-\frac{8387920120151}{1245468555600007}$ | $-\frac{2389431041534}{124187611002}$ | 34 | $-\frac{281567}{1074034146600516}$ | $-\frac{39638}{1036666750088053}$ |
| 9 | $-\frac{406742805160}{722051602550002}$ | $-\frac{4264232445251}{1077516190927017}$ | 35 | $-\frac{83389}{050495005000000000000000000000000000000$ | $-\frac{157667}{1201054950202765}$ |
| 10 | $\frac{1634411901667}{100757714701000000}$ | $-\frac{771344875358}{1077316180027017}$ | 36 | 959435095886281 28205 | $-\frac{113934}{1269179294}$ |
| 11 | $\frac{1087757145912963}{437876257661}$ | $\frac{1065295895342307}{265642346741}$ | 37 | $\frac{1730718489838961}{32611}$ | 1363173358100473 - 17361 |
| 12 | $\frac{696307764680561}{89522509038}$ | 379808011765194 244271216116 | 38 | 797073541452501 71967 | $\frac{669218624902568}{4754}$ |
| 13 | $\frac{638255170736941}{117755492100}$ | $\underline{ 598168785900209} \\ \underline{ 6820473784} \\ \underline{ 682047784} \\ $ | 39 | $\frac{2654250117756974}{4933}$ | $\frac{717927786469769}{13327}$ |
| 14 | $589974697761101 \\ \underline{31618874999}$ | $\frac{682731068968999}{76969905434}$ | 40 | $ 600713092825087 \underline{2587} $ | $\frac{968274850704167}{6388}$ |
| 15 | $\begin{array}{c} 687200813024600 \\ 74641597491 \end{array}$ | $814968793309763 \\18025759659$ | 41 | $\frac{1091724494207482}{2187}$ | $\begin{array}{c} 719277775331407 \\ 3259 \end{array}$ |
| 10 | $\frac{2908279600103189}{25500972924}$ | $-\frac{498226704994540}{1173876505}$ | 41 | $-\frac{1}{468954843525869}$ 5147 | 1225269357285833 782 |
| 10 | $\frac{1200692005935517}{1407090135}$ | 270355500536671 2453687289 | 42 | $-\frac{1711259362214795}{1793}$ | $-\frac{1032843841522687}{1186}$ |
| 17 | 324017497006981 | 261078590300893 6222314030 | 43 | 2031167694756949 | 793758766058135 |
| 18 | $-\frac{1453030732}{626828763236497}$ | 1665707814672911 | 44 | 747735360938753 | $-\frac{513}{516229281713762}$ |
| 19 | $-\frac{3196827854}{1480462540436741}$ | $\frac{88697433}{876534657690958}$ | 45 | $\frac{493}{866534589214144}$ | $-\frac{659}{1846327333506367}$ |
| 20 | $-\frac{784093409}{1033902289669424}$ | $-\frac{694824828}{951854540026661}$ | 46 | $\frac{341}{940674986436191}$ | $\frac{12}{1534685279850239}$ |
| 21 | $\frac{16311127}{845829326226001}$ | $-rac{574584500}{1127166940199931}$ | 47 | $\frac{100}{1139858083359883}$ | $\frac{149}{1160416855750006}$ |
| 22 | $\frac{61819697}{291114080642220}$ | $-\tfrac{174441984}{991147170856703}$ | 48 | $-\frac{167}{1430503381545037}$ | $\frac{237}{925315921317341}$ |
| 23 | $\frac{66841451}{441859334922159}$ | $\frac{20324816}{1031487384778575}$ | 49 | $-\frac{1299}{18049381579847260}$ | $\frac{69}{1181044907862509}$ |
| 24 | $\frac{24335317}{514920381859084}$ | $\frac{38004608}{529759164589729}$ | 50 | $\frac{31}{1479728935331543}$ | $-\frac{59}{575506931225862}$ |
| 25 | $-\frac{40105997}{2987332128606330}$ | $\frac{20671763}{439071227878684}$ | | | |

A.2. Coefficients for the left fundamental matrix. These are the coefficients used in Definition 6.5 (files Re_PL1.dat, Im_PL1.dat, Re_PL2.dat, Im_PL2.dat, Re_PL3.dat, Im_PL3.dat, Re_PL4.dat, Im_PL4.dat).

| n | $\operatorname{Re} c_n(P_{L,1})$ | $\operatorname{Im} c_n(P_{L,1})$ | $\operatorname{Re} c_n(P_{L,2})$ | $\operatorname{Im} c_n(P_{L,2})$ |
|----|---|---|---|---|
| 0 | $-\tfrac{25677447589672}{1163963890016041}$ | $\frac{530349753284166}{1047712021892891}$ | $-\tfrac{517945004884255}{724226719805958}$ | $rac{251627013871075}{660395344710337}$ |
| 1 | $-\tfrac{1082442536473116}{1635220915357765}$ | $-rac{110367753663726}{708047493550373}$ | $-\tfrac{114302268418242}{761576205136705}$ | $-\tfrac{123913358194209}{741695303878733}$ |
| 2 | $-\tfrac{185936764340935}{1296137430677681}$ | $-rac{111920181288506}{778797133744727}$ | $\tfrac{17151015646089}{541448058831086}$ | $-\tfrac{56801890756933}{1070300059045458}$ |
| 3 | $\tfrac{2345025580829}{870128025908873}$ | $-\tfrac{34365822575423}{1337355174852400}$ | $\frac{12280916504473}{899971311827585}$ | $-\tfrac{4036122681220}{978714325901481}$ |
| 4 | $\tfrac{144189079060498}{18283075520062425}$ | $\tfrac{1118439732896}{361756199062829}$ | $\frac{2367038770634}{1164366581657513}$ | $\tfrac{696494230513}{544434361047937}$ |
| 5 | $\frac{4674586796888}{2744490668379641}$ | $\frac{2557817734831}{941852677758271}$ | $-\tfrac{68218152895}{1289289733871468}$ | $\frac{628708864225}{1260394898925096}$ |
| 6 | $\frac{7687896621}{2004608749423625}$ | $\frac{799499324755}{1174208276879177}$ | $-\tfrac{69582891023}{702483920307261}$ | $\tfrac{42725908350}{563156598813481}$ |
| 7 | $-\frac{93374633783}{1038722783958176}$ | $\tfrac{18545361642}{275567875203109}$ | $-\frac{74265152988}{3065120819772145}$ | $-\tfrac{355734767}{460001483151423}$ |
| 8 | $-\tfrac{74972967052}{3069745379188931}$ | $-\frac{8395948274}{694862258440025}$ | $-\tfrac{1425956526}{580260321528581}$ | $-\tfrac{3054521565}{911188844028754}$ |
| 9 | $-\tfrac{1753498904}{892264564109643}$ | $-\tfrac{14325340118}{2370412550810421}$ | $\tfrac{350564155}{1114357638294561}$ | $-\tfrac{1090756380}{1249267251208579}$ |
| 10 | $\frac{252981210}{329942940569983}$ | $-\tfrac{1427767523}{1455622877913618}$ | $\frac{814673447}{4479667553603919}$ | $-\frac{42698153}{422756615139272}$ |
| 11 | $\frac{233811361}{675089370308832}$ | $\frac{65076613}{1748449524823006}$ | $\frac{31150042}{858129033616623}$ | $\frac{8792583}{1201122954408068}$ |
| 12 | $\tfrac{88693487}{1239673993058033}$ | $\tfrac{65452033}{1077676849558065}$ | $\tfrac{1621271}{615294864255555}$ | $\frac{7412633}{1261169303904716}$ |
| 13 | $\tfrac{16152549}{2378642812142881}$ | $\tfrac{10309096}{646555332622629}$ | $-rac{409762}{596152022550001}$ | $\tfrac{4166754}{3260798591775841}$ |
| 14 | $-rac{1165135}{1521116212026049}$ | $\frac{687636}{361267137880831}$ | $-\frac{328995}{1149047074605608}$ | $\frac{70602}{589692303242011}$ |
| 15 | $-rac{324441}{1082461988954614}$ | $-\frac{102346}{336321508849921}$ | $-rac{17309}{385662716915208}$ | $-rac{15957}{785266120923205}$ |
| 16 | $\frac{15487}{443684595725487}$ | $-\frac{347503}{1425037188487487}$ | $\frac{5141}{2306675406400974}$ | $-\frac{21625}{1828006420046424}$ |

| n | $\operatorname{Re} c_n(P_{L,3})$ | $\operatorname{Im} c_n(P_{L,3})$ | $\operatorname{Re} c_n(P_{L,4})$ | $\operatorname{Im} c_n(P_{L,4})$ |
|----|---|---|---|--|
| 0 | $-\tfrac{1880579012407712}{1264572197250759}$ | $-\tfrac{327819574106897}{494498622613644}$ | $-\tfrac{1235423862838786}{2757559154064461}$ | $-\tfrac{285132495792879}{4057283039847173}$ |
| 1 | $-\tfrac{269325222738631}{842677740476761}$ | $-rac{279059406245370}{500773457669029}$ | $-rac{278104214936677}{783316830514894}$ | $\frac{192467438451300}{2252055563919871}$ |
| 2 | $\frac{126135288121777}{786649538688151}$ | $\tfrac{13012660882672}{1325207140507713}$ | $-\tfrac{90252694047103}{1031085290502292}$ | $\tfrac{123938360444803}{7617732080365628}$ |
| 3 | $\tfrac{66966331840168}{1117986847339497}$ | $\tfrac{14126422263245}{254798882291938}$ | $-\tfrac{11814364894261}{1510073138688502}$ | $-\tfrac{725600123961}{130748046621331}$ |
| 4 | $\frac{6465480873820}{1017256720219203}$ | $\tfrac{10619128824323}{621177549874954}$ | $\frac{3295269917673}{1147301164879810}$ | $-\tfrac{1596611369498}{587329124564689}$ |
| 5 | $-\tfrac{1195359826000}{842901581462863}$ | $\tfrac{2448804123921}{1358009907248222}$ | $\frac{800208476515}{593980553933492}$ | $-\tfrac{431437902245}{1355349427602094}$ |
| 6 | $-\tfrac{767233757971}{1206025809809902}$ | $-\tfrac{295096738099}{663997884248891}$ | $\frac{320343108547}{1296813455849558}$ | $\tfrac{45943063991}{441280912817043}$ |
| 7 | $-\tfrac{56213820029}{744001059411659}$ | $-\tfrac{201950984545}{864036276330126}$ | $\frac{1678611220}{1182482488098459}$ | $\frac{77581308389}{1394322745386835}$ |
| 8 | $\frac{4008681509}{232745015528311}$ | $-\tfrac{46415387181}{975810303561101}$ | $-\tfrac{13917685903}{1132831548325475}$ | $\frac{20228340188}{1704140910599865}$ |
| 9 | $\tfrac{9101448335}{936316295545854}$ | $-rac{2666400359}{918527912661640}$ | $-\tfrac{2406729043}{668455332160438}$ | $\frac{2202762858}{2887200986290567}$ |
| 10 | $\tfrac{1140382427}{546458830753244}$ | $\frac{532397964}{433896933852091}$ | $-\frac{187103959}{451158034677887}$ | $- \frac{466366777}{1364508737553223}$ |
| 11 | $rac{136975753}{899505647020653}$ | $\tfrac{2045592387}{4885544477645626}$ | $rac{32673485}{519815835953651}$ | $-\frac{86262448}{668106598542119}$ |
| 12 | $-\tfrac{129975825}{2525855610320201}$ | $\frac{32576477}{739747340611994}$ | $\tfrac{18670649}{461399449262856}$ | $-\tfrac{25268097}{1304183977126009}$ |
| 13 | $-\tfrac{18451201}{862738719240511}$ | $-rac{6148497}{559990880016659}$ | $\frac{56345467}{5970307008258519}$ | $\frac{1277713}{1735339492096633}$ |
| 14 | $-rac{2091976}{537632796652219}$ | $-\tfrac{7839474}{1287616149628645}$ | $\frac{2514531}{2339957268256139}$ | $\frac{1294620}{1116204772221739}$ |
| 15 | $-rac{163144}{874171390981047}$ | $-\tfrac{1575517}{1122236720308525}$ | $-\frac{39037}{429277340876428}$ | $\frac{486093}{1850805774675719}$ |
| 16 | $\frac{87944}{872571465709971}$ | $-rac{172115}{1216760034973378}$ | $-rac{71911}{1010941875327823}$ | $-rac{4989}{1698292824029957}$ |

A.3. Coefficients for the right fundamental matrix. These are the coefficients used in Definition 6.9 (files Re_PR1.dat, Im_PR1.dat, Re_PR2.dat, Im_PR2.dat, Re_PR3.dat, Im_PR3.dat, Re_PR4.dat, Im_PR4.dat, Re_QR3.dat, Im_QR3.dat, Re_QR4.dat, Im_QR4.dat).

| | , <u>-</u> | , - , | , . , . | , |
|----|--|---|--|---|
| n | $\operatorname{Re} c_n(P_{R,1})$ | $\operatorname{Im} c_n(P_{R,1})$ | $\operatorname{Re} c_n(P_{R,2})$ | $\operatorname{Im} c_n(P_{R,2})$ |
| 0 | $\frac{905182892168009}{1210536500275806}$ | $-rac{964124562956389}{764588616268277}$ | $\frac{229599367281501}{2318964790029964}$ | $-rac{437213741432953}{1273601827018991}$ |
| 1 | $-\frac{29543275157879}{878678715680883}$ | $\frac{1867829427849379}{1290374648689744}$ | $-\frac{1224654218910595}{4267202327370079}$ | $\frac{1158060336659301}{2244448991962928}$ |
| 2 | $-\frac{63357277247875}{175246925958027}$ | $-\frac{313439385873153}{1248519869481776}$ | $-\frac{4615735457747}{182245197945545}$ | $-\frac{109625271108935}{721111072614097}$ |
| 3 | $\frac{171643411122718}{1439385508783457}$ | $-\frac{40427939601805}{290932896249328}$ | <u>48133140500126</u> 905133989460001 | $-\frac{51314484136660}{1873337812961177}$ |
| 4 | <u>24891665834664</u> 754600155259735 | $\frac{29414188024163}{421620319370787}$ | $-\frac{2158192286492}{888659946510293}$ | 24568723216499 813420496525666 |
| 5 | $-\frac{22895742052880}{909723752359467}$ | $\frac{2370979013401}{251620082225998}$ | $-\frac{9264298757713}{1015041263348322}$ | $-\frac{991210242019}{1074271419267915}$ |
| 6 | $-\frac{351984366023}{155503237138870}$ | $-\frac{11082072690006}{946928032602073}$ | $\frac{266663365757}{234184892421294}$ | $-\frac{3296112962521}{700243258667708}$ |
| 7 | $\frac{2684717689919}{665397340312499}$ | $-\frac{936779909321}{1917134228555035}$ | $\frac{3716211666823}{2546912601321189}$ | $\frac{561012480343}{1039110453379923}$ |
| 8 | $\frac{204495031733}{892063953585555}$ | <u>983138246457</u> 575177877165194 | $-\frac{197279420208}{965673491448683}$ | 889394790347 1283012701353518 |
| 9 | $-\frac{282079080365}{475205232780054}$ | $\frac{118544413214}{1337988439159219}$ | $-\frac{199852951835}{856374836775734}$ | $-\frac{42930913826}{544804255184983}$ |
| 10 | $-\frac{75234678489}{1146991988921768}$ | $-\frac{311455913477}{1307958831340280}$ | $\frac{12604670095}{682298764674304}$ | $-\frac{104425124649}{992700518480605}$ |
| 11 | $\frac{27446080969}{339185504352084}$ | $-rac{47454977496}{1370191231278385}$ | $\frac{39389088692}{1054442482787855}$ | $\frac{1446074809}{481257886411013}$ |
| 12 | $\frac{34471087395}{1689613159415371}$ | $\tfrac{46134027428}{1546999539500645}$ | $\frac{779106751}{405616788629073}$ | $\tfrac{25381533964}{1605587623458439}$ |
| 13 | $-rac{5693269851}{641781551348606}$ | $\frac{4691296707}{456755583552215}$ | $-\frac{5097165836}{930554422274277}$ | $\frac{2815666246}{1372958236326965}$ |
| 14 | $-\frac{1526624585}{303828308783349}$ | $-\frac{1036570712}{412578113410611}$ | $-\frac{977336355}{708469650207641}$ | $-\frac{2175607579}{1066073009498197}$ |
| 15 | $\tfrac{185663646}{629433111835183}$ | $-rac{2768069841}{1216023567070165}$ | $\frac{761796682}{1293158957171919}$ | $-rac{102953705}{129554014255937}$ |
| 16 | $\tfrac{1187495175}{1259329621322999}$ | $-rac{278541911}{1711100496646239}$ | $\frac{656489921}{1702055151458457}$ | $\frac{49547699}{348913642508819}$ |
| 17 | $\frac{432419895}{2089230096254686}$ | $\frac{328568765}{905848321145561}$ | $\frac{632938}{680946124449833}$ | $\frac{50557499}{281339303152100}$ |
| 18 | $-\tfrac{108311147}{946194004410941}$ | $\tfrac{128672807}{943052250448267}$ | $-\tfrac{15995173}{225750884329698}$ | $\tfrac{28515899}{1070558314508552}$ |
| 19 | $-\tfrac{207866887}{2812943312601145}$ | $-\tfrac{32232698}{1186108690826973}$ | $-\frac{11149469}{511236685737299}$ | $-\tfrac{31416532}{1220441323196857}$ |
| 20 | $-\frac{1157633}{1021508877076545}$ | $-\frac{24829791}{711272401151756}$ | $-\frac{1694723}{249616541638119}$ | $-rac{3511729}{263959769361999}$ |
| 21 | $\frac{10779565}{765327632541236}$ | $-\tfrac{6994135}{1142907242394279}$ | $\frac{3376427}{518049036863803}$ | $\frac{569388}{864034798879697}$ |
| 22 | $\frac{1197415}{242248789906637}$ | $\frac{3950018}{815296383956159}$ | $\frac{484957}{571359614317252}$ | $\frac{4999839}{1750261175451749}$ |
| 23 | $-rac{617141}{557185087932570}$ | $\frac{3020612}{1023745657645005}$ | $-rac{772488}{753765149157353}$ | $\frac{207458}{236124336006689}$ |
| 24 | $-rac{742697}{540112325150848}$ | $\frac{74917}{801607829597733}$ | $-\frac{551167}{1028765608432637}$ | $-rac{100702}{397829137552219}$ |
| 25 | $-rac{199319}{749381056956021}$ | $-rac{416717}{691331029767839}$ | $\frac{9719}{389320976820880}$ | $-rac{623859}{2201810291272148}$ |
| 26 | $\frac{313398}{1824536757806963}$ | $-\tfrac{345604}{1313864215913825}$ | $\frac{47930}{401620769507521}$ | $-\frac{39667}{626967832812710}$ |
| 27 | $\frac{97817}{906373931856942}$ | $\frac{111076}{2036801353243733}$ | $\frac{26736}{912224592309695}$ | $\frac{41026}{947198033341063}$ |
| 28 | $\frac{7163}{1057510687908509}$ | $\frac{66183}{835033288759151}$ | $-\frac{21319}{1526009128296655}$ | $\frac{25675}{818125449081334}$ |
| 29 | $-rac{3238}{236551642035411}$ | $\frac{1045}{751514878587959}$ | $-rac{6659}{1221774686284569}$ | $-\tfrac{1520}{644594865315677}$ |
| 30 | $-\frac{8985}{2096914333847348}$ | $-rac{7963}{719902311931441}$ | $\frac{1967}{2836559485514530}$ | $-rac{365}{71800999828723}$ |

| n | $\operatorname{Re} c_n(P_{R,3})$ | $\operatorname{Im} c_n(P_{R,3})$ | $\operatorname{Re} c_n(P_{R,4})$ | $\operatorname{Im} c_n(P_{R,4})$ |
|----|---|--|---|---|
| 0 | $rac{740650831460613}{934942792365922}$ | $\tfrac{2689410496949907}{2001499430084200}$ | $rac{761862582957073}{961718937524583}$ | $\frac{1182470673057526}{880012322737655}$ |
| 1 | $\frac{569427687308973}{601170187939750}$ | $-\tfrac{1024935540756591}{672478931558924}$ | $\frac{873060318475391}{921728689064955}$ | $-\tfrac{1624916621990511}{1066137479262242}$ |
| 2 | $-\tfrac{532979823280321}{633320510278501}$ | $-\tfrac{69957624743533}{625432248141991}$ | $-rac{599990717626501}{712947115166357}$ | $-\tfrac{399698341481122}{3573366494468815}$ |
| 3 | $\frac{28464248861417}{736126048327861}$ | $\tfrac{288594245548787}{785222675847615}$ | $\tfrac{177493227234231}{4590227854071298}$ | $\frac{737446315483133}{2006483421166275}$ |
| 4 | $\frac{73682279716033}{470643857894806}$ | $-\tfrac{135692132917951}{2621852039059466}$ | $\frac{126650470605855}{808976952394186}$ | $-\frac{34594079422525}{668429007068348}$ |
| 5 | $-\tfrac{17265510261599}{580749122199510}$ | $-\tfrac{119844136829124}{1946128179334357}$ | $-\tfrac{5291285009522}{177979629795357}$ | $-\tfrac{66233280652543}{1075550772011991}$ |
| 6 | $-\frac{3120721941593}{124168371845008}$ | $\frac{12054167583115}{841493452290073}$ | $-\frac{9755284617771}{388146663029824}$ | $\tfrac{17294129832504}{1207291745096663}$ |
| 7 | $\frac{4332350001641}{716914600094349}$ | $\tfrac{10045935101009}{1000604889763212}$ | $\frac{1801317645348}{298080930411533}$ | $\frac{11140336511600}{1109610511618277}$ |
| 8 | $\tfrac{2803101758312}{664325668187493}$ | $-\frac{1962960292922}{856141345624497}$ | $\tfrac{19349818686402}{4585841805433331}$ | $-\frac{2064861595960}{900585402398553}$ |
| 9 | $-\frac{300547693681}{389171029419735}$ | $- \frac{672359056400}{381241246995181}$ | $- \frac{217878391779}{282124800155504}$ | $-\tfrac{2445608211877}{1386709549730462}$ |
| 10 | $-\frac{690193576016}{913435044344523}$ | $\tfrac{138621865565}{667063957884079}$ | $-\frac{311859615025}{412729864748876}$ | $\tfrac{29907316969}{143917362139959}$ |
| 11 | $\frac{40653294702}{1566290108545715}$ | $\tfrac{180251714823}{569879310845225}$ | $\tfrac{16409201114}{632213688518857}$ | $\tfrac{304214814195}{961797943684436}$ |
| 12 | $\frac{177505964856}{1364189386789441}$ | $\frac{15221828077}{781193124852170}$ | $\frac{127488203139}{979787095065619}$ | $\frac{16999326924}{872415405887335}$ |
| 13 | $\frac{21592898956}{970260455485067}$ | $-\tfrac{74128523975}{1472152645404764}$ | $\tfrac{1841798254}{82759800639973}$ | $-\tfrac{44159771999}{876989338014120}$ |
| 14 | $-\frac{16993848037}{947923689335838}$ | $-\tfrac{17223588445}{1122828114044253}$ | $-\tfrac{20059202841}{1118910415155941}$ | $-\frac{7384597329}{481411496702614}$ |
| 15 | $-\tfrac{7342603046}{836112599056501}$ | $\frac{4762129268}{892540250279987}$ | $-\tfrac{8139973110}{926910256568993}$ | $\frac{4347565127}{814840725268847}$ |
| 16 | $\frac{536256341}{549789972102272}$ | $\frac{1142895923}{257839662913172}$ | $\frac{719932883}{738102003459719}$ | $\frac{3168543301}{714829863519978}$ |
| 17 | $\frac{1807540235}{896019492770726}$ | $\frac{247927811}{956783412035527}$ | $\frac{4801663923}{2380242712960663}$ | $\frac{217098274}{837808499606943}$ |
| 18 | $\frac{924598819}{2165838881659034}$ | $-\frac{1331113277}{1640260084905636}$ | $\frac{380384387}{891036499734743}$ | $-\frac{349636508}{430838470479917}$ |
| 19 | $-\frac{327603758}{1189789003082799}$ | $-\frac{232368144}{741234500984239}$ | $-\frac{70921121}{257570823873469}$ | $-\frac{376608293}{1201348236995778}$ |
| 20 | $-\frac{147579221}{820091368279374}$ | $\frac{17954603}{282176838429834}$ | $-\frac{174007412}{966951686225751}$ | $\frac{8878000}{139527784133131}$ |
| 21 | $-rac{3984606}{952669591918255}$ | $\frac{73548109}{840499395336333}$ | $-\frac{3093027}{739504174285273}$ | $\frac{68363360}{781248675518769}$ |
| 22 | $\frac{33018762}{901835766983351}$ | $\frac{24706327}{1391569899147199}$ | $\frac{4704407}{128490659190882}$ | $\frac{32585943}{1835384815157930}$ |
| 23 | $\frac{9953571}{699358207989101}$ | $-\frac{4805715}{396862377504766}$ | $\frac{28139951}{1977170374759080}$ | $-rac{2884679}{238220653176077}$ |
| 24 | $-\frac{2275043}{871158267297332}$ | $-\frac{10105489}{1194103065479818}$ | $-rac{5809473}{2224560341316904}$ | $-\frac{5127745}{605913877448069}$ |
| 25 | $-\tfrac{1508434}{370804653670483}$ | $-\tfrac{342776}{458401222363013}$ | $-\tfrac{611348}{150282135918537}$ | $-\tfrac{968837}{1295645159143331}$ |
| 26 | $-\frac{887599}{960034900081162}$ | $\tfrac{1726039}{1103539410688527}$ | $-\frac{1521889}{1646088553558104}$ | $\tfrac{2072312}{1324928326209757}$ |
| 27 | $\frac{970196}{1817456220397669}$ | $\frac{294121}{343115647286057}$ | $\frac{575969}{1078955635568715}$ | $\frac{2533931}{2956032977050961}$ |
| 28 | $\frac{152106}{468549910175669}$ | $\frac{17638}{1209308611042749}$ | $\frac{284158}{875325137573125}$ | $\frac{11638}{797932510223127}$ |
| 29 | $-\frac{5753}{513103030712072}$ | $-\frac{56845}{323718828377007}$ | $-\frac{1152}{102745470420575}$ | $-\frac{61937}{352716563860020}$ |
| 30 | $-\frac{114693}{3555716884771193}$ | $-\frac{41540}{700182517676521}$ | $-\frac{21524}{667287892267315}$ | $-\frac{70979}{1196395159416509}$ |

| n | $\operatorname{Re} c_n(Q_{R,3})$ | $\operatorname{Im} c_n(Q_{R,3})$ | $\operatorname{Re} c_n(Q_{R,4})$ | $\operatorname{Im} c_n(Q_{R,4})$ |
|----|---|--|---|--|
| 0 | $\frac{33999531320775}{130417803622346}$ | $-\tfrac{115215147451424}{803828617788823}$ | $-\tfrac{215362893343035}{826104197336093}$ | $rac{154457357338925}{1077612161277739}$ |
| 1 | $-\frac{10572359682749}{63161708216629}$ | $\frac{746784388391963}{1635700668949216}$ | $\frac{168645512739677}{1007527078849589}$ | $-\frac{317958620607164}{696432781013537}$ |
| 2 | $-\frac{175915122460926}{811520164472617}$ | $-\frac{243583922679691}{835304424568651}$ | $\frac{533699456369471}{2462027508226876}$ | $\frac{194060507809468}{665540123560885}$ |
| 3 | <u>19473410038635</u> 88507063473734 | $-\frac{21045572025109}{514527245308103}$ | $-\frac{169979945792361}{772562474762171}$ | $\frac{55499247893443}{1356859062857279}$ |
| 4 | $-\frac{27203093314906}{1270800722772467}$ | <u>107427966706222</u> <u>878712100125215</u> | $\frac{10592335313465}{527266544020241}$ | $-\frac{122037231436089}{008210515722262}$ |
| 5 | $-\frac{148538381407323}{2528542018588064}$ | $-\frac{71585036482556}{2712561127057560}$ | 50735576511685 862662548572517 | 20810520327337 788860661068021 |
| 6 | $\frac{22668386868535}{1234204302330018}$ | $-\frac{28630716086205}{1111852083268210}$ | $-\frac{10566382260292}{575207850585135}$ | $\frac{23666109621316}{010055367182821}$ |
| 7 | $\frac{16700171964815}{1539997663421869}$ | $\frac{13629148906684}{1326405874518603}$ | $-\frac{6615922888841}{610083884864192}$ | $-\frac{12226684511265}{1189916279634608}$ |
| 8 | $-\frac{9318355126389}{1825506441204187}$ | $\frac{3565203794275}{790985430829102}$ | <u>3826673106546</u> 749661963901733 | $-\frac{5739913308203}{1273472166808886}$ |
| 9 | $-\frac{567030959136}{298196167204823}$ | $- \frac{3448396151248}{1474386964872959}$ | $\tfrac{2321617326598}{1220916384464621}$ | $\frac{3061173673413}{1308827165828666}$ |
| 10 | $\frac{1008214961432}{1002184627055011}$ | $-\tfrac{1180949097475}{1434495188229149}$ | $- \frac{699863207957}{695677186797248}$ | $\frac{795907345558}{966786172173145}$ |
| 11 | $\frac{300540964851}{820355743082368}$ | $\tfrac{271331115501}{666370594916120}$ | $-\tfrac{503135583604}{1373357424879735}$ | $-\tfrac{701921307898}{1723870550801489}$ |
| 12 | $-\tfrac{71023225441}{461580005147271}$ | $\tfrac{149246212351}{901923910345033}$ | $\tfrac{131901068753}{857225162837987}$ | $-\tfrac{135042574257}{816088560763511}$ |
| 13 | $-\frac{98315576929}{1318912162058657}$ | $-\frac{24695622455}{466774332191571}$ | $\frac{102662046401}{1377220434538992}$ | $\frac{10539597265}{199209940300267}$ |
| 14 | $rac{15054662375}{970125874534439}$ | $-\frac{47489023456}{1445086203759829}$ | $-\tfrac{10157201126}{654532355474557}$ | $\frac{40965910347}{1246588528436716}$ |
| 15 | $\tfrac{13142842223}{949074650501087}$ | $\frac{5076245364}{1645116043925125}$ | $-\frac{9342805723}{674665336903952}$ | $-\tfrac{2188398995}{709219125363011}$ |
| 16 | $\frac{226298380}{767392447913163}$ | $\tfrac{6845158852}{1249648977249809}$ | $-rac{365469193}{1239329679041972}$ | $-\frac{4830796255}{881907889924153}$ |
| 17 | $- \tfrac{1005807501}{523850955588044}$ | $\tfrac{663280168}{845902234697095}$ | $\tfrac{1711840595}{891571528959154}$ | $-rac{588284646}{750258066949153}$ |
| 18 | $-\tfrac{1319447118}{2369576824364017}$ | $-\frac{88934573}{154880512839497}$ | $\frac{571497853}{1026345087399398}$ | $\frac{1399566979}{2437360906435628}$ |
| 19 | $\frac{55980467}{672098004050896}$ | $-\tfrac{249788978}{849181992024207}$ | $-\tfrac{69112865}{829764757424032}$ | $\tfrac{439436359}{1493906759582821}$ |
| 20 | $\frac{443264437}{3800279681354425}$ | $-\tfrac{21439958}{1044729047273153}$ | $- \frac{118387033}{1014978415797737}$ | $\frac{7639299}{372248750025758}$ |
| 21 | $\frac{6228350}{129225359166053}$ | $\frac{22653891}{565430884939006}$ | $- \frac{59762936}{1239957110537757}$ | $-rac{38607422}{963623811321139}$ |
| 22 | $\frac{43198}{634626999675329}$ | $\frac{27316354}{1320151184291751}$ | $-\tfrac{223990}{3290663957990577}$ | $-rac{71991424}{3479218480345129}$ |
| 23 | $-\tfrac{23340089}{1457800707355487}$ | $\frac{1851731}{1022824680125384}$ | $\frac{44561482}{2783269591663031}$ | $-\tfrac{1493525}{824965521657446}$ |
| 24 | $-rac{3303005}{481981729454043}$ | $-rac{1924111}{795104633913382}$ | $\frac{7004037}{1022044431183152}$ | $\frac{489143}{202129641141438}$ |
| 25 | $\frac{551231}{262645469615304}$ | $-\frac{2063861}{1738090169892127}$ | $-\frac{3555861}{1694263897044514}$ | $\frac{1026501}{864472606190261}$ |
| 26 | $\frac{884777}{582587972454744}$ | $-\frac{157524}{727771329356959}$ | $-\frac{2780251}{1830676874517844}$ | $\frac{157524}{727771329356959}$ |

| n | $c_n(\widetilde{\varphi})$ | n | $c_n(\widetilde{\varphi})$ | n | $c_n(\widetilde{arphi})$ | n | $c_n(\widetilde{\varphi})$ |
|----|--------------------------------|----|--------------------------------|----|--------------------------------|----|-------------------------------|
| 0 | $-\frac{355950139}{14657721}$ | 11 | $-\frac{46328383}{136777634}$ | 22 | $-\frac{1512364}{3182339229}$ | 33 | $-\frac{68819}{132457417335}$ |
| 1 | $-\frac{351922817}{8012162}$ | 12 | $-\frac{33010805}{173605428}$ | 23 | $-\frac{3688957}{14330077878}$ | 34 | $-rac{26569}{95627042824}$ |
| 2 | $-\frac{982442337}{29494607}$ | 13 | $-\frac{15683952}{147751241}$ | 24 | $-rac{554124}{3980489795}$ | 35 | $-rac{26051}{175267834537}$ |
| 3 | $-\frac{927585801}{41134127}$ | 14 | $-\frac{25169019}{426787852}$ | 25 | $-rac{295574}{3932362577}$ | 36 | $-rac{52249}{655677095601}$ |
| 4 | $-\frac{106335447}{7377385}$ | 15 | $-\frac{4349081}{133306007}$ | 26 | $-rac{574309}{14171446290}$ | 37 | $-rac{21233}{491171341160}$ |
| 5 | $-\frac{222295829}{25020823}$ | 16 | $-\frac{18067173}{1004771323}$ | 27 | $-\frac{243338}{11151617411}$ | 38 | $-rac{17458}{717858338557}$ |
| 6 | $-\frac{635353759}{118991106}$ | 17 | $-\frac{9593701}{971236636}$ | 28 | $-\frac{233428}{19892041241}$ | 39 | $-rac{9171}{594311525288}$ |
| 7 | $-\frac{99180037}{31489914}$ | 18 | $-\frac{2882805}{532844677}$ | 29 | $-rac{783613}{124316164231}$ | 40 | $-rac{5123}{802284206161}$ |
| 8 | $-\frac{90284711}{49302362}$ | 19 | $-\frac{3384291}{1145130091}$ | 30 | $-\frac{140780}{41623422413}$ | | |
| 9 | $-\frac{50679761}{48147039}$ | 20 | $-\frac{6910633}{4290928466}$ | 31 | $-\frac{181237}{99964939686}$ | | |
| 10 | $-\frac{39578500}{66027997}$ | 21 | $-\frac{3840951}{4385988911}$ | 32 | $-\frac{87979}{90614186264}$ | | |

A.4. Coefficients for approximation of phase. These are the coefficients used in Definition 9.9 (file phitw.dat).

A.5. Coefficients for functional ψ_F on the right-hand side at high frequency. These are the coefficients used in Definition 9.11 (files Pshm.dat, Pshp.dat, Qshm.dat, Qshp.dat).

| n | $c_n(P^{\sharp})$ | $c_n(P_+^\sharp)$ | n | $c_n(P^{\sharp})$ | $c_n(P_+^\sharp)$ |
|----|--------------------------------|----------------------------------|----|---------------------------------|----------------------------------|
| 0 | $-\frac{20631567}{455512634}$ | $\frac{9568160}{201881083}$ | 21 | $\frac{444379}{11176237362}$ | $\frac{464738}{9889823197}$ |
| 1 | $-\frac{69554432}{1023019957}$ | $\frac{47047045}{649018097}$ | 22 | $\frac{58611}{800400353}$ | $\frac{700381}{9033942495}$ |
| 2 | $-\frac{12329231}{269313212}$ | $\frac{15204955}{303630746}$ | 23 | $\frac{376613}{4875608921}$ | $\tfrac{1861973}{23332528565}$ |
| 3 | $-\frac{24877483}{853350175}$ | $\tfrac{13738349}{412452315}$ | 24 | $\frac{369375}{5539349437}$ | $\tfrac{856040}{12551471521}$ |
| 4 | $-\frac{7426145}{423099917}$ | $\frac{12395978}{560685879}$ | 25 | $\frac{425169}{8311530079}$ | $\frac{786717}{15112362946}$ |
| 5 | $-\frac{11757787}{1144307006}$ | $\frac{4674985}{322495783}$ | 26 | $\frac{8540}{237393733}$ | $\frac{470938}{12898482173}$ |
| 6 | $-\frac{13897943}{2368284443}$ | $\frac{7144027}{757618270}$ | 27 | $\tfrac{1192200}{50764918679}$ | $\frac{331739}{13936308798}$ |
| 7 | $-\frac{2608823}{897244087}$ | $\frac{5488408}{849193287}$ | 28 | $\frac{184101}{12908496413}$ | $\frac{280721}{19424774289}$ |
| 8 | $-rac{6500988}{7285807193}$ | $\frac{6339738}{1317306931}$ | 29 | $\tfrac{116228}{14542704637}$ | $\tfrac{260555}{32147771403}$ |
| 9 | $\frac{546214}{4782016773}$ | $\frac{6617837}{1853174972}$ | 30 | $\frac{217996}{54065032051}$ | $\frac{171382}{41809662947}$ |
| 10 | $\tfrac{829499}{5077812811}$ | $\frac{5588966}{2483699731}$ | 31 | $\frac{205018}{119920816427}$ | $\frac{20157}{11522056256}$ |
| 11 | $-rac{1461956}{3824446197}$ | $\frac{2120393}{2426452111}$ | 32 | $\frac{39536}{85699224537}$ | $\frac{55952}{115365215341}$ |
| 12 | $-\frac{1682122}{1598994471}$ | $-rac{577767}{1940668792}$ | 33 | $-\frac{14154}{106459721287}$ | $-\frac{12646}{106373436087}$ |
| 13 | $-rac{4038402}{2681337091}$ | $-\tfrac{1891895}{1794830111}$ | 34 | $-\frac{38879}{108463987388}$ | $-rac{12622}{36054520295}$ |
| 14 | $-\tfrac{2429071}{1496074848}$ | $-rac{6247835}{4617218998}$ | 35 | $-\tfrac{120350}{303451706711}$ | $-\tfrac{680501}{1737699972493}$ |
| 15 | $-\frac{3992099}{2744509183}$ | $-\frac{2951339}{2282624457}$ | 36 | $-rac{67049}{189613539928}$ | $-\frac{46268}{131964663029}$ |
| 16 | $-\frac{1677957}{1492071964}$ | $-\tfrac{1954021}{1900589114}$ | 37 | $-rac{16593}{57560035534}$ | $-rac{7513}{26229120710}$ |
| 17 | $-rac{6135114}{8103636319}$ | $-\frac{1708530}{2442318371}$ | 38 | $-\frac{142643}{621173500099}$ | $-\frac{9833}{43040520729}$ |
| 18 | $-\frac{1185727}{2724946876}$ | $-\frac{494221}{1232914518}$ | 39 | $-rac{68896}{360364723503}$ | $-rac{47271}{248344211236}$ |
| 19 | $-rac{615665}{3140688998}$ | $-\tfrac{12658987}{72084729713}$ | 40 | $-rac{7439}{83635247148}$ | $-rac{35159}{396928578929}$ |
| 20 | $-\frac{303045}{7068751433}$ | $-\frac{157896}{5140050545}$ | | | |

| n | $c_n(Q^\sharp)$ | $c_n(Q_+^\sharp)$ | n | $c_n(Q^{\sharp})$ | $c_n(Q_+^\sharp)$ |
|----|--------------------------------|--------------------------------|----|--------------------------------|--------------------------------|
| 0 | $\frac{15024113}{419334545}$ | $-\frac{84863998}{2670638095}$ | 21 | $\frac{644854}{5545449541}$ | $\frac{834944}{7476783719}$ |
| 1 | $\frac{26338241}{486449897}$ | $-\frac{9625016}{212402275}$ | 22 | $\frac{577559}{3414678615}$ | $\frac{1878683}{11290298852}$ |
| 2 | $\frac{27404960}{734705327}$ | $-\frac{33166439}{1145664248}$ | 23 | $\frac{764515}{4581725254}$ | $\frac{1108028}{6705865483}$ |
| 3 | $\frac{49269755}{1955111421}$ | $-\frac{3615029}{210454920}$ | 24 | $\tfrac{1152742}{8299295951}$ | $\frac{829552}{6014353219}$ |
| 4 | $\frac{14736779}{829173919}$ | $-\frac{20461453}{2336769147}$ | 25 | $\frac{692198}{6670546807}$ | $\frac{2114434}{20489753259}$ |
| 5 | $\frac{9864830}{799321543}$ | $-\frac{7751669}{1914492761}$ | 26 | $\frac{1104826}{15494381429}$ | $\frac{308637}{4349217023}$ |
| 6 | $\frac{5807541}{688797160}$ | $-\frac{3239765}{2016502842}$ | 27 | $\frac{768301}{16894467943}$ | $\frac{778702}{17199766551}$ |
| 7 | $\frac{2581561}{397061491}$ | $\frac{3193777}{8145377075}$ | 28 | $\frac{632143}{23507582867}$ | $\frac{376901}{14078789498}$ |
| 8 | $\frac{3477497}{601816550}$ | $\frac{20730289}{9981418642}$ | 29 | $\frac{387908}{26655806505}$ | $\frac{177583}{12263047605}$ |
| 9 | $\frac{2453911}{503078748}$ | $\frac{2919390}{1104484453}$ | 30 | $\frac{193625}{27855685003}$ | $\frac{694582}{100537691067}$ |
| 10 | $\frac{1828028}{582905913}$ | $\frac{2764644}{1543787783}$ | 31 | $\frac{102053}{38897983149}$ | $\frac{104861}{40354712764}$ |
| 11 | $\frac{1210513}{1390895317}$ | $\frac{3351387}{53798325847}$ | 32 | $\frac{54610}{139362283877}$ | $\frac{34666}{91965684667}$ |
| 12 | $-\frac{2763189}{2312900014}$ | $-\frac{3152020}{1877195641}$ | 33 | $-rac{50339}{84179707475}$ | $-\frac{66308}{109266512019}$ |
| 13 | $-\frac{3105950}{1233038341}$ | $-\frac{2501676}{890621917}$ | 34 | $-\frac{24265}{26638113301}$ | $-\frac{85546}{93373352285}$ |
| 14 | $-\frac{6430687}{2165888529}$ | $-\frac{7269183}{2313247177}$ | 35 | $-\frac{48293}{53828091949}$ | $-\frac{60407}{67096252029}$ |
| 15 | $-\frac{3392927}{1242562631}$ | $-\frac{2439474}{860760853}$ | 36 | $-\frac{141873}{187707853304}$ | $-rac{78067}{103031870398}$ |
| 16 | $-\frac{2483649}{1174604665}$ | $-\frac{3250017}{1493439299}$ | 37 | $-\frac{52688}{88842876251}$ | $-\frac{119432}{200997801417}$ |
| 17 | $-\frac{423589}{301989567}$ | $-\frac{5370137}{3730662129}$ | 38 | $-\frac{48285}{105324245683}$ | $-\frac{131581}{286559012168}$ |
| 18 | $-\frac{2486208}{3197249045}$ | $-rac{9075478}{11351091637}$ | 39 | $-\frac{25525}{68383259559}$ | $-\frac{33818}{90473887987}$ |
| 19 | $-\frac{8556765}{26851400932}$ | $-\frac{2007859}{6052955872}$ | 40 | $-\frac{39253}{228035775642}$ | $-\frac{28319}{164297023388}$ |
| 20 | $-\frac{694132}{21959417587}$ | $-\frac{976123}{24793963305}$ | | | |
A.6. Coefficients for functional ψ_F on the right-hand side at low frequency. These are the coefficients used in Definition 9.14 (files Pflm.dat, Pflp.dat, Qflm.dat, Qflp.dat).

| n | $c_n(P^{\flat})$ | $c_n(P_+^\flat)$ | n | $c_n(P^{\flat})$ | $c_n(P_+^\flat)$ |
|----|--------------------------------|--------------------------------|----|---------------------------------|--------------------------------|
| 0 | $\frac{6446693}{169820556}$ | $-\frac{14137311}{328631966}$ | 21 | $\frac{2166057}{2090984221}$ | $-\frac{8193927}{5581197824}$ |
| 1 | $-\frac{12327282}{264684349}$ | $\frac{11693343}{210302048}$ | 22 | $\frac{22085969}{2757489833}$ | $\frac{8113899}{1279324939}$ |
| 2 | $-\frac{23803678}{393848217}$ | $\frac{19064463}{376977364}$ | 23 | $\frac{11145118}{2388280363}$ | $\frac{6807137}{1806168845}$ |
| 3 | $\frac{21048531}{269406664}$ | $-\frac{14724959}{224838528}$ | 24 | $-rac{3638881}{2329474793}$ | $-\frac{1964186}{1016010475}$ |
| 4 | $\frac{28821313}{954788917}$ | $-\frac{19344345}{453315719}$ | 25 | $-\frac{5023983}{1478063533}$ | $-\frac{4227356}{1214592129}$ |
| 5 | $-\frac{20768287}{646063235}$ | $\frac{41780932}{940844507}$ | 26 | $-\frac{4084347}{2103870572}$ | $-\frac{1131210}{595573079}$ |
| 6 | $-\frac{8620650}{158968559}$ | $\frac{10086808}{280153789}$ | 27 | $-rac{3892379}{13647365033}$ | $-\frac{3456692}{16247544233}$ |
| 7 | $\frac{8822238}{1388095199}$ | $-\frac{2236863}{4220113688}$ | 28 | $\frac{939208}{2090279385}$ | $\frac{1064265}{2079124508}$ |
| 8 | $\frac{14533007}{284261596}$ | $-\frac{43264411}{685770018}$ | 29 | $\frac{6288371}{12709007896}$ | $\frac{1558501}{2903960931}$ |
| 9 | $\frac{26458656}{338746937}$ | $-\frac{13703011}{172299852}$ | 30 | $\frac{1734665}{5973228508}$ | $\frac{1896802}{6042571493}$ |
| 10 | $\frac{22678807}{320218732}$ | $-\frac{16346877}{279671722}$ | 31 | $\frac{536630}{5651461237}$ | $\frac{1307750}{12339426353}$ |
| 11 | $\frac{9797368}{378006561}$ | $-\frac{14168711}{360974638}$ | 32 | $-rac{382358}{33393758311}$ | $-\frac{474408}{63937042567}$ |
| 12 | $\frac{4288859}{436809097}$ | $\frac{9449297}{1907064162}$ | 33 | $-rac{590177}{13884609616}$ | $-\frac{411365}{9852163702}$ |
| 13 | $-\frac{70935700}{3118935447}$ | $\frac{7001722}{864204645}$ | 34 | $-\frac{213203}{5926095679}$ | $-rac{563584}{15465731401}$ |
| 14 | $-\frac{22961994}{762738821}$ | $\frac{10084709}{946783989}$ | 35 | $-\frac{9792948}{475362427189}$ | $-\frac{231121}{10909405681}$ |
| 15 | $-\frac{4520657}{1003124969}$ | $\frac{8660835}{292877546}$ | 36 | $-rac{87697}{10557331092}$ | $-\frac{196004}{22666823599}$ |
| 16 | $-\frac{6645115}{562404224}$ | $\tfrac{19207703}{2016347648}$ | 37 | $-\frac{44622}{28797895063}$ | $-\frac{106183}{64107389914}$ |
| 17 | $-\frac{15404221}{1036278934}$ | $-rac{5292167}{1012788664}$ | 38 | $\frac{56003}{96311651760}$ | $\frac{49371}{63716889130}$ |
| 18 | $\frac{31509477}{7369719805}$ | $\frac{2686367}{435921537}$ | 39 | $\frac{439154}{367625188367}$ | $\frac{97211}{63656058176}$ |
| 19 | $\frac{366127}{78792700}$ | $\frac{3349107}{1212439678}$ | 40 | $\frac{337601}{508207249578}$ | $\frac{45469}{53232547777}$ |
| 20 | $-\frac{4449739}{1029484476}$ | $-\frac{5559403}{769910962}$ | | | |

| n | $c_n(Q^\flat)$ | $c_n(Q_+^\flat)$ | n | $c_n(Q^\flat)$ | $c_n(Q_+^\flat)$ |
|----|--------------------------------|--------------------------------|----|--------------------------------|--------------------------------|
| 0 | $\tfrac{22292801}{3541672815}$ | $-\frac{7320085}{422726802}$ | 21 | $\frac{1205233}{136249056}$ | $\frac{3269741}{340039997}$ |
| 1 | $-\frac{17183123}{628486997}$ | $\frac{5370257}{763909565}$ | 22 | $\frac{7689066}{3782048833}$ | $\frac{1714269}{851396435}$ |
| 2 | $-\frac{2601881}{36861921}$ | $\frac{6497353}{117477620}$ | 23 | $-\frac{13382928}{1921692661}$ | $-\frac{6544562}{899773689}$ |
| 3 | $-\frac{12788236}{351943543}$ | $\frac{14885347}{530360674}$ | 24 | $-\frac{29677129}{3876084991}$ | $-\frac{6647810}{832327799}$ |
| 4 | $\frac{28269278}{1953066793}$ | $-\frac{16788921}{1059683705}$ | 25 | $-\frac{19841725}{8067194896}$ | $-rac{3631559}{1341549134}$ |
| 5 | $\tfrac{20469314}{2291987951}$ | $-\frac{4389249}{756215519}$ | 26 | $\frac{1962597}{1098180905}$ | $\frac{14462219}{8835686444}$ |
| 6 | $-\frac{4604735}{149513089}$ | $\frac{38096807}{1136416257}$ | 27 | $\frac{1511361}{569900654}$ | $\frac{6573893}{2551571809}$ |
| 7 | $-\frac{21852472}{393873431}$ | $\frac{10456207}{194323924}$ | 28 | $\frac{1703825}{1059871716}$ | $\frac{1508811}{956103488}$ |
| 8 | $-\frac{12134291}{257196164}$ | $\frac{12813735}{325285216}$ | 29 | $\frac{760219}{1708397708}$ | $\frac{1342975}{3059758387}$ |
| 9 | $-\frac{16365871}{971101671}$ | $\frac{3830009}{573237032}$ | 30 | $-\frac{399962}{2441922197}$ | $-rac{706790}{4402882723}$ |
| 10 | $\tfrac{6889747}{460862719}$ | $-\frac{8628562}{427825761}$ | 31 | $-\frac{1322611}{4403040654}$ | $-\frac{822089}{2786494425}$ |
| 11 | $\frac{26137905}{785469161}$ | $-\frac{18694903}{622076748}$ | 32 | $-\frac{1965647}{8887295246}$ | $-rac{914359}{4220758055}$ |
| 12 | $\frac{202329748}{5789596889}$ | $-\frac{9472768}{366945325}$ | 33 | $-\frac{848635}{8065438807}$ | $-\frac{1109517}{10858798918}$ |
| 13 | $\frac{32134415}{1418682452}$ | $-\frac{47560750}{2582279807}$ | 34 | $-rac{711346}{28065675695}$ | $-rac{244921}{10370454135}$ |
| 14 | $\frac{9236098}{1889427287}$ | $-\frac{4228843}{306292600}$ | 35 | $\tfrac{164251}{15310750346}$ | $\frac{751693}{65123424948}$ |
| 15 | $-\frac{12439359}{3554243900}$ | $-\frac{4001267}{703804442}$ | 36 | $\frac{347400}{18439634939}$ | $\frac{1362251}{71079921794}$ |
| 16 | $\frac{9589627}{5230046955}$ | $\frac{7262256}{894504715}$ | 37 | $\frac{211901}{13949644989}$ | $\frac{745769}{48570592853}$ |
| 17 | $\frac{1634541}{355455424}$ | $\frac{12443095}{956856786}$ | 38 | $\frac{176189}{18981293769}$ | $\frac{94148}{10017012737}$ |
| 18 | $-\frac{412771}{109761066}$ | $\frac{3275390}{1002114789}$ | 39 | $\frac{509239}{99105942367}$ | $\frac{258699}{49907652418}$ |
| 19 | $-\frac{24644619}{3526953038}$ | $-\frac{3518545}{1432492754}$ | 40 | $\frac{282209}{151571651735}$ | $\frac{260032}{139671794219}$ |
| 20 | $\frac{1621784}{612479555}$ | $\frac{5987552}{1215060451}$ | | | |

A.7. Coefficients for functional ψ_F on the left-hand side. These are the coefficients used in Definition 9.20 (files PLm.dat, PLp.dat).

| n | $c_n(P^L)$ | $c_n(P^L_+)$ | n | $c_n(P^L)$ | $c_n(P^L_+)$ |
|----|----------------------------------|---------------------------------|----|---------------------------------|----------------------------------|
| 0 | $-\frac{171003977}{103918830}$ | $\frac{300931817}{241723206}$ | 26 | $\frac{96976718}{349936601}$ | $\frac{92590229}{334108131}$ |
| 1 | $-\frac{166671913}{118566186}$ | $\frac{31563190}{53581289}$ | 27 | $\frac{151400197}{305788942}$ | $\frac{22436507}{45315897}$ |
| 2 | $\frac{45964957}{58483290}$ | $-\frac{99394800}{60361067}$ | 28 | $-\frac{17543371}{94545237}$ | $-\frac{46126776}{248587741}$ |
| 3 | $\frac{48279065}{56023151}$ | $-\frac{140995963}{78831741}$ | 29 | $-\frac{82657111}{170462639}$ | $-\frac{93590633}{193010693}$ |
| 4 | $\frac{8972827}{48587872}$ | $-\frac{115016911}{96883787}$ | 30 | $\frac{49807827}{180113107}$ | $\frac{9174673}{33177092}$ |
| 5 | $-\frac{41493513}{97613005}$ | $-\frac{62369530}{96861583}$ | 31 | $\frac{24408481}{84951138}$ | $\frac{28186382}{98099723}$ |
| 6 | $-\frac{124868039}{211620567}$ | $-\tfrac{85729517}{168054687}$ | 32 | $-\frac{58360311}{168209960}$ | $-rac{36952057}{106505670}$ |
| 7 | $-\frac{82444695}{147905063}$ | $-\frac{70884043}{139773967}$ | 33 | $\frac{14741103}{458100637}$ | $\frac{23630575}{734353558}$ |
| 8 | $-\frac{237703833}{506032154}$ | $-\frac{65403365}{142220181}$ | 34 | $\tfrac{23982523}{146077235}$ | $\frac{35055715}{213523903}$ |
| 9 | $-\tfrac{452690956}{1349473269}$ | $-\frac{37359657}{111070835}$ | 35 | $-\frac{23716921}{182213911}$ | $-\frac{120267449}{923998618}$ |
| 10 | $-\frac{18816773}{130013559}$ | $-\frac{60126713}{412382715}$ | 36 | $\frac{16609307}{561307166}$ | $\frac{4069334}{137522073}$ |
| 11 | $\frac{40238064}{457449389}$ | $\frac{25758991}{293806979}$ | 37 | $\frac{16141679}{720463986}$ | $\frac{10456315}{466704758}$ |
| 12 | $\frac{297981729}{933827237}$ | $\frac{169452468}{531069001}$ | 38 | $-\frac{6565187}{256652917}$ | $-\frac{19154980}{748825813}$ |
| 13 | $\frac{53894425}{111747446}$ | $\frac{50340647}{104375994}$ | 39 | $\frac{6445520}{465342699}$ | $\frac{7490204}{540765019}$ |
| 14 | $\frac{39121627}{77223319}$ | $\frac{56404525}{111337301}$ | 40 | $-\frac{4486162}{876691085}$ | $-\tfrac{8167318}{1596066945}$ |
| 15 | $\frac{25655501}{73282493}$ | $\frac{42504024}{121408339}$ | 41 | $\frac{170205}{120603977}$ | $\frac{1276468}{904480581}$ |
| 16 | $\frac{24126233}{599965344}$ | $\frac{21968416}{546305221}$ | 42 | $-\tfrac{3620268}{12159078431}$ | $-\frac{1882565}{6322806898}$ |
| 17 | $-\frac{83229183}{274756991}$ | $-\frac{62067841}{204898926}$ | 43 | $\tfrac{663057}{12981857515}$ | $\frac{602699}{11800120564}$ |
| 18 | $-\frac{19346066}{39135121}$ | $-\frac{47572821}{96234968}$ | 44 | $-rac{318302}{64657214463}$ | $-\frac{21113}{4288718797}$ |
| 19 | $-\frac{30112253}{78550984}$ | $-\frac{24747641}{64556828}$ | 45 | $\frac{157007}{87593065450}$ | $\frac{97065}{54151858808}$ |
| 20 | $\frac{8376683}{1097584769}$ | $\frac{3237094}{424151787}$ | 46 | $\frac{45784}{39375676069}$ | $\frac{144114}{123942559835}$ |
| 21 | $\frac{62855398}{152433253}$ | $\frac{31137242}{75512227}$ | 47 | $\frac{11965}{21891917444}$ | $\frac{92393}{169048051048}$ |
| 22 | $\frac{206082020}{449291361}$ | $\frac{174951822}{381422611}$ | 48 | $\frac{37259}{62899729312}$ | $\tfrac{1671737}{2822185366593}$ |
| 23 | $\frac{10197255}{358894151}$ | $\frac{26642213}{937677287}$ | 49 | $\frac{119486}{790116130157}$ | $\frac{47253}{312466375855}$ |
| 24 | $-\frac{36390813}{78742075}$ | $-\tfrac{193849193}{419448933}$ | 50 | $\frac{60673}{454939111170}$ | $\frac{10847}{81333122376}$ |
| 25 | $-\frac{40487945}{109429569}$ | $-\frac{32480152}{87786353}$ | | | |

References

- Yakine Bahri, Yvan Martel, and Pierre Raphaël. Self-similar blow-up profiles for slightly supercritical nonlinear Schrödinger equations. Ann. Henri Poincaré, 22(5):1701–1749, 2021.
- [2] C. J. Budd. Asymptotics of multibump blow-up self-similar solutions of the nonlinear Schrödinger equation. SIAM J. Appl. Math., 62(3):801–830, 2001/02.
- [3] Chris J. Budd, Shaohua Chen, and Robert D. Russell. New self-similar solutions of the nonlinear Schrödinger equation with moving mesh computations. J. Comput. Phys., 152(2):756–789, 1999.
- [4] Thomas Duyckaerts and Svetlana Roudenko. Threshold solutions for the focusing 3D cubic Schrödinger equation. Rev. Mat. Iberoam., 26(1):1–56, 2010.
- [5] Gadi Fibich, Nir Gavish, and Xiao-Ping Wang. New singular solutions of the nonlinear Schrödinger equation. Phys. D, 211(3-4):193-220, 2005.
- [6] Gadi Fibich, Nir Gavish, and Xiao-Ping Wang. Singular ring solutions of critical and supercritical nonlinear Schrödinger equations. *Phys. D*, 231(1):55–86, 2007.
- [7] R. T. Glassey. On the blowing up of solutions to the Cauchy problem for nonlinear Schrödinger equations. J. Math. Phys., 18(9):1794–1797, 1977.
- [8] Justin Holmer, Galina Perelman, and Svetlana Roudenko. A solution to the focusing 3d NLS that blows up on a contracting sphere. Trans. Amer. Math. Soc., 367(6):3847–3872, 2015.
- [9] Justin Holmer and Svetlana Roudenko. A class of solutions to the 3D cubic nonlinear Schrödinger equation that blows up on a circle. Appl. Math. Res. Express. AMRX, (1):23–94, 2011.
- [10] Justin Holmer and Svetlana Roudenko. Blow-up solutions on a sphere for the 3D quintic NLS in the energy space. Anal. PDE, 5(3):475–512, 2012.
- [11] Nancy Kopell and Michael Landman. Spatial structure of the focusing singularity of the nonlinear Schrödinger equation: a geometrical analysis. SIAM J. Appl. Math., 55(5):1297–1323, 1995.
- [12] M.J. Landman, G.C. Papanicolaou, C. Sulem, P.L. Sulem, and X.P. Wang. Stability of isotropic singularities for the nonlinear schrödinger equation. *Physica D: Nonlinear Phenomena*, 47(3):393–415, 1991.
- [13] B. J. LeMesurier, G. Papanicolaou, C. Sulem, and P.-L. Sulem. Focusing and multi-focusing solutions of the nonlinear Schrödinger equation. *Phys. D*, 31(1):78–102, 1988.
- [14] Zexing Li. On stability of self-similar blowup for mass supercritical NLS. Preprint arXiv:2304.02078, 2023.
- [15] D. W. McLaughlin, G. C. Papanicolaou, C. Sulem, and P. L. Sulem. Focusing singularity of the cubic schrödinger equation. *Phys. Rev. A*, 34:1200–1210, Aug 1986.
- [16] Frank Merle, Pierre Raphaël, and Igor Rodnianski. Type II blow up for the energy supercritical NLS. Camb. J. Math., 3(4):439–617, 2015.
- [17] Frank Merle, Pierre Raphaël, and Jeremie Szeftel. Stable self-similar blow-up dynamics for slightly L² supercritical NLS equations. Geom. Funct. Anal., 20(4):1028–1071, 2010.
- [18] Frank Merle, Pierre Raphaël, and Jeremie Szeftel. On collapsing ring blow-up solutions to the mass supercritical nonlinear Schrödinger equation. Duke Math. J., 163(2):369–431, 2014.
- [19] K. Nakanishi and W. Schlag. Global dynamics above the ground state energy for the cubic NLS equation in 3D. Calc. Var. Partial Differential Equations, 44(1-2):1–45, 2012.
- [20] Takayoshi Ogawa and Yoshio Tsutsumi. Blow-up of H^1 solution for the nonlinear Schrödinger equation. J. Differential Equations, 92(2):317–330, 1991.
- [21] Pierre Raphaël. Existence and stability of a solution blowing up on a sphere for an L^2 -supercritical nonlinear Schrödinger equation. Duke Mathematical Journal, 134(2):199 258, 2006.
- [22] Pierre Raphaël and Jérémie Szeftel. Standing ring blow up solutions to the N-dimensional quintic nonlinear Schrödinger equation. Comm. Math. Phys., 290(3):973–996, 2009.
- [23] Vivi Rottschäfer and Tasso J. Kaper. Blowup in the nonlinear Schrödinger equation near critical dimension. J. Math. Anal. Appl., 268(2):517–549, 2002.
- [24] Vivi Rottschäfer and Tasso J. Kaper. Geometric theory for multi-bump, self-similar, blowup solutions of the cubic nonlinear Schrödinger equation. Nonlinearity, 16(3):929–961, 2003.
- [25] Catherine Sulem and Pierre-Louis Sulem. The nonlinear Schrödinger equation, volume 139 of Applied Mathematical Sciences. Springer-Verlag, New York, 1999. Self-focusing and wave collapse.
- [26] William C. Troy. Zero Energy Self-Similar Solutions Describing Singularity Formation In The Nonlinear Schrodinger Equation In Dimension N=3. Preprint arXiv:2210.05645, 2022.
- [27] Xiao Ping Wang. On singular solutions of the nonlinear Schroedinger and Zakharov equations. ProQuest LLC, Ann Arbor, MI, 1990. Thesis (Ph.D.)–New York University.
- [28] Kai Yang, Svetlana Roudenko, and Yanxiang Zhao. Blow-up dynamics in the mass super-critical NLS equations. Phys. D, 396:47–69, 2019.

- [29] V. E. Zakharov. Collapse of Langmuir Waves. Soviet Journal of Experimental and Theoretical Physics, 35:908, January 1972.
- [30] V. E. Zakharov. Collapse and self-focusing of Langmuir waves. In Basic Plasma Physics: Selected Chapters, Handbook of Plasma Physics, Volume 1, page 81, January 1984.
- [31] Ian Zwiers. Standing ring blowup solutions for cubic nonlinear Schrödinger equations. Anal. PDE, 4(5):677–727, 2011.

UNIVERSITÄT WIEN, FAKULTÄT FÜR MATHEMATIK, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA *Email address*: roland.donninger@univie.ac.at

UNIVERSITÄT INNSBRUCK, INSTITUT FÜR MATHEMATIK, TECHNIKERSTRASSE 13, 6020 INNSBRUCK, AUSTRIA *Email address*: birgit.schoerkhuber@uibk.ac.at