Strictly critical snarks with girth or cyclic connectivity equal to 6

Ján Mazák, Jozef Rajník, Martin Skoviera

{mazak, rajnik, skoviera}@dcs.fmph.uniba.sk

Comenius University, Mlynská dolina, 842 48 Bratislava

June 25, 2024

Abstract

A snark – connected cubic graph with chromatic index 4 – is critical if the graph resulting from the removal of any pair of distinct adjacent vertices is 3-edgecolourable; it is bicritical if the same is true for any pair of distinct vertices. A snark is strictly critical if it is critical but not bicritical. Very little is known about strictly critical snarks. Computational evidence suggests that strictly critical snarks constitute a tiny minority of all critical snarks. Strictly critical snarks of order n exist if and only if n is even and at least 32, and for each such order there is at least one strictly critical snark with cyclic connectivity 4. A sparse infinite family of cyclically 5-connected strictly critical snarks is also known, but those with cyclic connectivity greater than 5 have not been discovered so far. In this paper we fill the gap by constructing cyclically 6-connected strictly critical snarks of each even order $n \geq 342$. In addition, we construct cyclically 5-connected strictly critical snarks of girth 6 for every even $n \geq 66$ with $n \equiv 2 \pmod{8}$.

1 Introduction

Cubic graphs that do not admit any proper edge colouring with three colours, known as *snarks*, play a crucial role in the study of a variety of problems related to flows, edge colourings, perfect matchings, or cycle covers of graphs. Over time, considerable effort has been exerted to understand the structure of snarks. Although several relevant partial results have been gathered, many deep questions remain open.

In our recent paper, *Morphology of small snarks* [15], we have analysed the structure of all snarks with at most 36 vertices. The cornerstone of our approach to structural analysis is the concept of an I-*extension*. This operation, also known as an *edge extension*, consists in specifying two edges in a cubic graph, subdividing each of them with a new vertex, and adding a new edge joining the two vertices. It is a very natural operation and in the literature it has been used many times, see for example [1, 3, 16, 21].

Edge extensions can be conveniently employed to construct new snarks from known ones: it is enough to take an existing snark, choose a *removable* pair of edges (that is, one that can be removed without breaking uncolourability), and perform an I-extension. Removable pairs of edges are present in most known snarks; actually, an overwhelming majority of known snarks can be obtained from a smaller snark by a series of I-extensions retaining uncolourability at each step. However, there exist snarks that cannot be obtained from a smaller snark by an I-extension — and these are exactly the critical snarks. A *critical* snark can be equivalently defined as one in which the removal of any two adjacent vertices produces a 3-edge-colourable graph. If the removal of any pair of vertices yields a 3-edge-colourable graph, a snark is called *bicritical*. A snark which is critical but not bicritical is called *strictly critical*.

Critical snarks have been emerging in the literature under different disguises and in different settings [6, 8, 17, 18]. In 1996, Nedela and Škoviera [17] introduced critical and bicritical snarks in the context of snark reductions and decompositions. They showed that critical snarks coincide with 6-irreducible snarks and that bicritical snarks are 7-irreducible, which means that they are the same as the irreducible ones. In 2008, da Silva et al. [5, 6] initiated the study of flow criticality of graphs and introduced flow-edge-critical and flow-vertex-critical snarks. It turned out, however, that these two approaches to snark criticality agree (Máčajová and Škoviera [13]): a snark is 4-flow-edge-critical if and only if it is critical, and is 4-flow-vertex-critical if and only if it is bicritical.

Critical snarks are known to be cyclically 4-connected with girth at least 5 [17] and therefore they are nontrivial snarks by the usual standards. As discussed in [4], every snark G with cyclic connectivity 4 can be constructed as a dot product $G_1 \cdot G_2$ of two smaller snarks. If G is bicritical, both G_1 and G_2 are bicritical. As a consequence, every bicritical snark can be decomposed into a collection of cyclically 5-connected bicritical snarks such that it can be reconstructed from them by repeated dot products. Moreover, this collection is unique up to isomorphism [4, Theorem C]. The class of bicritical snarks is thus closed under such decomposition. If G is strictly critical, then G_2 must be critical but G_1 need not; it only has to be "nearly critical" [4, Sections 4 and 6]. According to [4, Section 12], snarks arising from the decomposition of strictly critical snarks might even include snarks with cyclic connectivity 3. Overall, very little is understood about the relationship between strictly critical snarks and "nearly critical" snarks, or about the nature of graphs in either of these two classes.

Strictly critical snarks have been previously studied in [4, 9, 15, 20]. The smallest strictly critical snarks have order 32 (one is displayed in Figure 1) and such snarks have been constructed for all higher orders.

Theorem 1.1 ([4]). There exists a strictly critical snark of order n for each even $n \ge 32$.

There are exactly 64 326 024 cyclically 4-connected snarks of girth at least 5 with order not exceeding 36, of which 55 172 are critical [2]. Somewhat surprisingly, a vast majority of these snarks are bicritical, only 846 being strictly critical, a little over 1.5 percent [2, 13]. This fact does not seem to have any obvious explanation.

All strictly critical snarks discovered so far have cyclic connectivity 4 or 5. As already mentioned, the smallest strictly critical snark has order 32. Its cyclic connectivity is 4. The smallest cyclically 5-connected strictly critical snarks have order 36 (and there are 84 such snarks [15, Section 7]). It is not known how common cyclically 5-connected strictly critical snarks are. An infinite family of them was provided by Grünevald and Steffen [9], but the graphs in the family are rather large (the smallest of them has 66 vertices) and only a small proportion of orders of snarks is covered.

In Section 4, we demonstrate the existence of strictly critical snarks with cyclic connectivity 6, proving an analogue of Theorem 1.1 and solving Problem 6.3 from [4].



Figure 1: A strictly critical snark of the smallest possible order 32 with the only removable pair $\{x, y\}$ of vertices.

Theorem 1.2. There exists a cyclically 6-connected strictly critical snark of order n for each even $n \ge 342$.

We discuss the situation for orders below 342 at the end of Section 4 (see Theorem 4.8 and Problem 1).

In addition to cyclically 6-connected strictly critical snarks we present a rich family of cyclically 5-connected strictly critical snarks with girth 6. These snarks are considerably smaller than the discovered snarks with cyclic connectivity 6.

Theorem 1.3. There exists a cyclically 5-connected strictly critical snark with girth 6 and order n for each $n \ge 66$ such that $n \equiv 2 \pmod{8}$.

We prove this theorem in Section 3. The construction is inspired by specimens found among critical snarks of order 36 and analysed in Section 7 of *Morphology* [15]. Our present work can thus be considered as a supplement to Section 8 of [15] where we have generalised small examples of snarks with desirable properties into abundant families of bicritical snarks.

Snarks with cyclic connectivity or girth greater than 5 are interesting for several reasons. First, cyclic connectivity and girth are important measures of complexity of cubic graphs. For example, smallest cubic graphs of given girth, called *cages*, have been studied for decades. Second, these two parameters have strong influence on structural properties of cubic graphs, including their edge-colourability. Some 40 years ago it was conjectured [10] that snarks with cyclic connectivity or girth greater than 6 do not exist. It took 16 years to refute the girth conjecture [11], while the conjecture on cyclic connectivity remains open and is essentially intact. It is thus interesting, for every property of a snark, to ask whether there are cyclically 6-connected snarks having that property — we might discover that some nontrivial property does not hold for snarks of higher cyclic connectivity and that might shed some light on the Jaeger's connectivity conjecture. It has also been conjectured [17, Conjecture 1] that bicritical snarks have girth at most 6. We might learn more about this problem by trying to construct strictly critical and bicritical snarks with higher girth. Finally, the third reason for our interest in cyclically 6-connected strictly critical snarks is their complete absence among known snarks with at most 38 vertices, which distinguishes them from bicritical ones.

2 Preliminaries

Basics

All graphs considered in this paper are undirected; they may contain loops and parallel edges, although their presence will often be excluded by connectivity requirements. Recall that a cubic graph is cyclically k-connected if no set of fewer than k edges separates two cycles of G from each other. The cyclic connectivity of G is the largest integer k such that G is cyclically k-connected.

Multipoles

Multipoles are often used as building blocks for large graphs. Informally, a *multipole* is a graph permitted to contain dangling edges and isolated edges. Each edge has two ends, which may but need not be incident with a vertex. A *dangling edge* has only one end incident with a vertex while *isolated edge* has neither. A *link* is an edge with both ends incident with a vertex. An edge end incident with no vertex is called a *semiedge* and a multipole with k semiedges is a k-pole. The set of all semiedges of M is denoted by S(M). The semiedges of a multipole M can be grouped into pairwise disjoint *connectors* S_1, \ldots, S_t (each connector comes with a linear ordering of its semiedges). Such multipole is called a $(|S_1|, \ldots, |S_t|)$ -pole and is denoted by $M(S_1, \ldots, S_t)$. In this article, we only deal with cubic multipoles, which means that each vertex is incident with exactly three edge ends.

The width of a connector is the number of semiedges it contains. A connector of width k is called a *k*-connector. The junction of two semiedges is performed by joining the semiedges together, creating an ordinary edge from two dangling edges. We can also perform a junction of two connectors of the same width or two *k*-poles by performing individual junctions for each pair of semiedges (this is the place where the linear ordering of semiedges defined for each connector comes into play).

Formal definitions of these notions and other related information can be found in [15]. For a k-pole $M(S_1, S_2, \ldots, S_t)$ and its vertex v we denote by M - v a (k + 3)-pole $N(S_1, S_2, \ldots, S_t, S_{t+1})$ constructed by removing the vertex v from the multipole M and putting the three semiedges formerly incident with v into the connector S_{k+1} . Note that if the vertex v is incident with a dangling edge e, then e becomes an isolated edge and the ends of e are retained in the multipole M - v. To keep our notation short, we shall write $M - (v_1, v_2, \ldots, v_r)$ instead of $(((M - v_1) - v_2) - \ldots) - v_r$. Note that this operation is not commutative: the order of the connectors changes with the order of removed vertices. When u and v are adjacent vertices of a (k_1, \ldots, k_t) -pole M, we denote M - [u, v] the $(k_1, \ldots, k_t, 2, 2)$ -pole obtained from the $(k_1, \ldots, k_t, 3, 3)$ -pole M - (u, v) by deleting the isolated edge obtained from the edge uv of M.

Assume that e is a link of a multipole $M(S_1, S_2, \ldots, S_k)$. By cutting e into two dangling edges ending with semiedges f_1 and f_2 , respectively, we construct a new multipole $N(S_1, S_2, \ldots, S_k, (f_1, f_2))$ which we denote by M - e. Again, we write $M - (e_1, e_2, \ldots, e_s)$ instead of $(((M - e_1) - e_2) - \ldots) - e_s$. Finally, for a vertex v of a (k + 1)-pole M incident with exactly one dangling edge e, we denote by $M \sim v$ the k-pole obtained from M by deleting the dangling edge e and by suppressing the vertex v.

Colourings and flows

A 3-edge-colouring of a multipole M, or just a colouring for short, is an assignment of colours to the edges of M such that no two edges incident with the same vertex receive the same colour. We extend the notion of a colouring to the ends of edges in a natural fashion: an end of an edge (in particular, a semiedge) has the same colour as the edge it belongs to. Both ends of an edge thus always have the same colour.

It has become standard to take $\mathbb{K} = \mathbb{Z}_2 \times \mathbb{Z}_2 - \{(0,0)\}$ for the set of colours. With this choice, a mapping $\varphi \colon E(M) \to \mathbb{K}$ is a colouring if and only if, for any three edges e, f, and g incident with the same vertex, one has $\varphi(e) + \varphi(f) + \varphi(g) = 0$. This means that a colouring of a multipole is a nowhere-zero $(\mathbb{Z}_2 \times \mathbb{Z}_2)$ -flow and vice versa. A straightforward consequence of this fact is the following lemma.

Lemma 2.1 (Parity Lemma [7]). Let M be a k-pole and let k_1 , k_2 , k_3 be the numbers of dangling edges of colour (0, 1), (1, 0), (1, 1), respectively. Then

$$k_1 \equiv k_2 \equiv k_3 \equiv k \pmod{2}.$$

For any ordered set $A = (e_1, e_2, \ldots, e_k)$ of semiedges (in particular, a connector) we denote $(\varphi(e_1), \varphi(e_2), \ldots, \varphi(e_k))$ by $\varphi(A)$. The colouring set of a multipole M is the set

 $Col(M) = \{\varphi(S(M)) \mid \varphi \text{ is a colouring of } M\}.$

Two multipoles M_1 and M_2 are said to be *colour-equivalent* if $\operatorname{Col}(M_1) = \operatorname{Col}(M_2)$.

The flow through a connector S of M is the value

$$\varphi_*(S) = \sum_{e \in S} \varphi(e).$$

A connector S of M is called *proper* if $\varphi_*(S) \neq 0$ for each colouring φ of M. If all connectors of a multipole M are proper, we say that M itself is *proper*. Note that every uncolourable multipole is proper.

3 Cyclically 5-connected strictly critical snarks of girth 6

In this section we prove Theorem 1.3 by generalising a structure discovered in cyclically 5-connected strictly critical snarks of order 36 and described in [15, Section 7]. The structure can be built from three (2,3)-poles as follows (see Figure 2).

Definition 3.1. For three (2,3)-poles T_1 , T_2 , and T_3 , let $\text{TTT}_{sc}(T_1, T_2, T_3)$ denote the (2, 2, 2)-pole constructed from the disjoint union $T_1 \cup T_2 \cup T_3$ by adding three new vertices v_1 , v_2 , and v_3 and by attaching the dangling edges from the 3-connectors to them in such a way that each v_i becomes incident with exactly one edge from each 3-connector.

In order to produce a snark containing a (2, 2, 2)-pole TTT_{sc} we require the three (2, 3)-poles T_1, T_2 , and T_3 be proper. It is well known and easy to see that if G is a snark, e is an edge of G, and v is a vertex of G, then the (2, 3)-pole T(B, C) = (G - e) - v is proper. It means that for each colouring φ of T we have

$$\varphi_*(B) \neq 0$$
 and $\varphi_*(C) \neq 0.$ (1)

If T admits all colourings satisfying both (1) and Parity Lemma, it is called *perfect*. To fulfil the connectivity requirements of the resulting snarks it is convenient to assume that



Figure 2: A (2, 2, 2)-pole found in cyclically 5-connected strictly critical snarks of order 36

the (2,3)-pole (G-e) - v is created by using pairs e and v such that v is not adjacent to any endvertex of e. For more details on proper (2,3)-poles we refer the reader to [15, Section 5.2].

Next, we create a (2, 2, 2)-pole H_6 from a 6-cycle $(u_1u_2u_3v_4u_5u_6)$, encoded as a cyclic sequence of vertices, by attaching a dangling edge e_i to the vertex u_i for each $i \in \{1, \ldots, 6\}$. We partition the dangling edges into three connectors (e_1, e_4) , (e_2, e_5) , (e_3, e_6) and join the resulting 2, 2, 2)-pole to $\text{TTT}_{\text{sc}}(T_1, T_2, T_3)$. Strictly speaking, there are several ways of how to join H_6 to $\text{TTT}_{\text{sc}}(T_1, T_2, T_3)$, but due to the symmetry of H_6 , there is essentially only one way that preserves the connectors. All other graphs can be obtained by permuting the multipoles T_1 , T_2 , and T_3 and by switching the pairs of edges within individual 2connectors of T_1 , T_2 , and T_3 . For the purpose of our proofs, we can therefore regard the join $H_6 * \text{TTT}_{\text{sc}}(T_1, T_2, T_3)$ as uniquely determined.

Note that usually, and also in the next section, the cycles are regarded as 6-poles with their semiedges ordered in one of the natural cyclic orders. The considered partition of semiedges into three connectors in the (2, 2, 2)-pole H_6 is specifically needed for this construction and it also reflects some of the colouring properties that are specific to the 6-cycle.



Figure 3: The structure of the snark $H_6 * \text{TTT}_{sc}(T_1, T_2, T_3)$

Lemma 3.2. For any proper (2,3)-poles T_1 , T_2 , and T_3 the graph $G = H_6*TTT_{sc}(T_1, T_2, T_3)$ is a snark in which each pair of vertices v_1 , v_2 and v_3 is removable. In particular, G is not bicritical.

Proof. It is well known that H_6 is an *even* (2, 2, 2)-pole, which means that the number of proper connectors is always even (see [15, Section 5.4]). By contrast, each connector of $\text{TTT}_{sc}(T_1, T_2, T_3)$ is proper, so $H_6 * \text{TTT}_{sc}(T_1, T_2, T_3)$ is indeed a snark. Observe that

all three 2-connectors of $\text{TTT}_{sc}(T_1, T_2, T_3)$ will be proper regardless of the presence of the vertices v_1, v_2 and v_3 . Thus, any two of them form a removable pair vertices.

In general, $H_6 * \text{TTT}_{sc}(T_1, T_2, T_3)$ need not be a critical snark, and this may occur even in the case where the all three (2,3)-poles are taken from a critical snark. The reason is that a proper (2,3)-pole T = (G - e) - v constructed from a critical snark may happen to be uncolourable. Thus to ensure the criticality we need an additional rather technical property based on the following definition.

Definition 3.3 (Chladný and Škoviera [4]). A pair of distinct edges $\{e, f\}$ of a snark G is *essential* if it is non-removable and for every vertex v of the 4-pole G - (e, f) incident with some dangling edge, the 3-pole $G \sim v$ is colourable.

Definition 3.4. A proper (2,3)-pole T = (G-e) - v is called *good* if, for every endvertex w of the edge e and every pair of edges f and g such that f is incident with w and g is incident with v, the pair $\{f, g\}$ is essential in G.

The following lemma asserts that for certain multipoles, including good proper (2, 3)-poles, we can find a colouring in which two prescribed semiedges have the same colour.

Lemma 3.5. Let G be a snark and let x and y be non-adjacent vertices such that for any two edges e and f incident with x and y, respectively, the pair $\{e, f\}$ is essential in G. Then there exists a colouring φ of the 6-pole $M((e_1, e_2, e_3), (f_1, f_2, f_3)) = G - (x, y)$ such that $\varphi(e_1) = a, \varphi(e_2) = \varphi(e_3), \varphi(f_1) = c$ and $\varphi(f_2) + \varphi(f_3) = b$ for some a, b, c such that $\{a, b, c\} = \mathbb{K}.$

Proof. Since the pair of the edges $\{e_1, f_1\}$ is essential in G, the 3-pole $(G - (e_1, f_1)) \sim x$ has a colouring from which the desired colouring of G - (x, y) can be obtained in a straightforward way.

Next, we recall the following well-known fact on the colouring set of (2, 2)-poles obtained by removing a non-removable pair of adjacent vertices from a snark, also called *isochromatic* (2, 2)-poles.

Lemma 3.6 (Chladný and Škoviera [4, Section 3]). Let G be a snark and $\{u, v\}$ a pair of adjacent non-removable vertices of G. Then

$$Col(G - [u, v]) = \{(a, a, b, b) \mid a, b \in \mathbb{K}\}.$$

The following theorem is a crucial step towards the proof of Theorem 1.3. It states that three good (2,3)-poles in a TTT-multipole are sufficient to produce a critical snark.

Theorem 3.7. Let T_1 , T_2 , and T_3 be perfect good proper (2,3)-poles obtained from critical snarks. Then the snark $G = H_6 * \text{TTT}_{sc}(T_1, T_2, T_3)$ is strictly critical.

Proof. Assume $T_i = (G_i - e_i) - w_i$ for each $i \in \{1, 2, 3\}$. Since the snark G is not bicritical by Lemma 3.2, it is sufficient to show that it is critical, that is, that for an arbitrary pair of adjacent vertices x and y of G the 4-pole M = G - [x, y] is colourable. We distinguish four cases.

Case (i). Both vertices x and y belong to the 6-cycle in H_6 . The 4-pole M can be coloured as shown in Figure 4. All the (2, 3)-poles T_i admit colourings as displayed in the figure since they are perfect.

Case (ii). The vertex x is from the 6-cycle and (without loss of generality) the vertex y is from T_1 . A colouring of the 4-pole M is indicated in Figure 5. Since y has to be an end vertex of e_i , the vertices y and w_1 are non-adjacent. So, by Lemma 3.5, the 6-pole $N(S_1, S_2) = G_1 - (y, w_1)$ admits a colouring assigning the colours a, p, p to the semiedges of S_1 in the order depicted in Figure 5, and colours a, c, c to the semiedges of S_2 in some order. Due to the symmetry of the multipole $TTT_{sc}(T_1, T_2, T_3)$, it is irrelevant which semiedge of S_2 has colour a.



Figure 4: Colouring for Case (i)



Figure 5: Colouring for Case (ii)



Figure 6: Colouring for Case (iii)

Case (iii). Both x and y belong to the same proper (2,3)-pole, say, T_1 . A colouring of G - [x, y] is illustrated in Figure 6. It includes a colouring of the 9-pole $T'_1 = T_1 - [x, y]$, which can be obtained in the following way. Consider the snark G_1 which gives rise to the (2,3)-pole T_1 by removing an edge e_1 and a vertex w_1 . We start from a colouring of the 4-pole $G_1 - [x, y]$ such that all the dangling edges have the same colour p; such a colouring exists according to Lemma 3.6. Denote the colour of the edge e_1 of $G_1 - [x, y]$ by a. The links incident with the vertex w_1 have pairwise distinct colours a, b, c in some order. By symmetry, the exact order of these colours is not important. After the removal of the vertex w_1 and splitting the link e_1 , we get the required colouring of $T_1 - [x, y]$.

Case (iv). The vertex x is from T_1 and y is one of v_1 , v_2 and v_3 , say $y = v_1$. Initially, we find a colouring of the 6-pole $T_1 - x \cong (G_1 - [w_1, x]) - e_1$. Since the snark G_1 is critical,

the 4-pole $G_1 - [w_1, x]$ is colourable in such a way that all its dangling edges are coloured by the same colour c (by Lemma 3.6). Denote the colour of the link e_1 of $G_1 - [w_1, x]$ by p. After splitting the link e_1 , we obtain the 6-pole $T_1 - x \cong (G_1 - [w_1, x]) - e_1$ with its semiedges coloured by c, c, c, c, p, p. If $p \neq c$, the colouring of the 4-pole M = G - [x, y]is given in Figure 7a (with p = a). Otherwise, if p = c, we can colour the 4-pole Maccording to Figure 7b.



Figure 7: Colouring for Case (iv)

Now we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. For the proof, we take all three proper (2, 3)-poles T_1 , T_2 and T_3 from the well-known family of Isaacs snarks. It is well known that Isaacs snarks are critical excluding J_3 , which is not critical [19, Theorem 4.13]. If T_i is to be obtained from J_5 , we choose an edge and a vertex so that their removal destroys the only 5-cycle of J_5 . As a result, each T_i has girth 6. Since every pair of non-adjacent edges in any Isaacs snark except J_3 is essential [4, Example 5.5], the proper (2, 3)-poles T_1 , T_2 and T_3 are all good. Theorem 3.7 now implies that the graph $H_6 * \text{TTT}_{sc}(T_1, T_2, T_3)$ is a strictly critical snark. Moreover, it is clear that it is cyclically 5-connected and has girth 6.

The smallest member of our family has 66 vertices and is depicted in Figure 8; it is produced by taking each proper (2, 3)-pole T_i from the Isaacs snark J_5 as described above. In order to construct larger snarks, we can take the snark on 66 vertices and replace one of the (2, 3)-poles constructed from J_5 with a (2, 3)-pole constructed from J_{5+2k} where k is an arbitrary positive integer, thereby adding 8k vertices. This completes the proof of Theorem 1.3.

Remark 3.8. We are not aware of any smaller strictly critical snarks without 5-cycles.

4 Cyclically 6-connected strictly critical snarks

In this section we present a construction of cyclically 6-connected strictly critical snarks based on superposition. Superposition is a general construction method producing large cubic graphs (usually snarks) from smaller ones. Here, we restrict ourselves to basic facts about this method, which are necessary for our exposition, and refer the reader to [11, 14] for more details.



Figure 8: A cyclically 5-connected strictly critical snark of girth 6 and order 66

In order to define a superposition, we first choose a *base graph* G, which we assume to be cubic and connected. Then we replace each vertex with a *supervertex*, a multipole with tree connectors, and each edge with a *superedge*, a multipole with two connectors. Finally, we perform all the junctions between supervertices and superedges that correspond to the incidences between vertices and edges of G. The corresponding connectors are required to have the same width in order for the resulting graph \tilde{G} be cubic.

There are multiple ways how one can ensure that a superposition produces a snark. According to [11, Theorem 4], if the base graph is a snark and each of the superedges used in the superposition is proper, then the graph resulting from superposition is again a snark. Such a superposition is called a *proper superposition*.

In our construction we utilise a specific type of the junction of multipoles. Take two multipoles with three connectors each, an (i, s, r_1) -pole $M(I, S_1, R_1)$ and an (s, o, r_2) -pole $N(S_2, O, R_2)$ such that $|S_1| = |S_2|$. Following [12], we define a serial junction $M \circ N$ of M and N as the $(i, o, r_1 + r_2)$ -pole $P(I, O, R_1 \cup R_2)$ which arises by the junction of the connectors S_1 and S_2 and by merging the connectors R_1 and R_2 into a single connector $R_1 \cup R_2$. The connectors R_1 and R_2 are regarded as residual, that is, not involved in the junction. We allow r_1 or r_2 to be zero and treat a multipole $M(S_1, S_2, \emptyset)$ with an empty residual connector as $M(S_1, S_2)$. Given a multipole $M(S_1, S_2, \ldots, S_k)$ with $k \ge 2$ and $|S_1| = |S_2|$, we also define the closure \overline{M} of M to be the multipole obtained by the junction of S_1 and S_2 .



Figure 10: The superedge A_5

We now proceed to our particular superposition construction. For supervertices we will take copies of the multipole W depicted in Figure 9, which has two 3-connectors and a 1-connector in this order. Our superedges will be created from Isaacs snarks. For a detailed explanation of properties of Isaacs snarks and relevant notation we refer the

reader to [15, Section 5.5]. The *Isaacs superedge* A_5 is obtained by removing two specific vertices at distance 4 from the Isaacs snark J_5 as shown in Figure 10. For any odd $k \ge 7$, the *Isaacs superedge* A_k is constructed by substituting the 6-pole Y_{k-3} for the 6-pole Y_2 contained in A_5 . Note that A_k also arises from the Isaacs snark J_k by removing two vertices at distance 4. It is known that all the Isaacs superedges A_k are proper (3, 3)-poles [11, 15].

We will need the following two properties of multipoles constructed from Isaacs snarks.

Lemma 4.1. Let J_k be the Isaacs snark, where $k \ge 5$ is odd, and let u be a vertex and vw be an edge of J_k such that u is adjacent to neither v nor w. Then the proper (2,3)-pole $(J_k - vw) - u$ is perfect.

Proof. We proceed by induction. The statement holds for J_5 and J_7 , which we have verified by using a computer. Assume that the statement is true for some odd $k \ge 7$. Then the proper (2,3)-pole $T = (J_{k+2} - e) - v$ contains a 6-pole Y_4 which is colour-equivalent to the 6-pole Y_2 . Therefore, the proper (2,3)-pole T is colour-equivalent to some proper (2,3)-pole constructed from J_k , which is perfect by the induction hypothesis. \Box

As a straightforward consequence, we have the following lemma.

Lemma 4.2. Let $M = J_k - (u, v)$ be a (3,3)-pole constructed from an Isaacs snark J_k for some odd $k \ge 5$ by removing vertices u and v whose distance is at least 3 and let a, b, c, d, e be any colours from \mathbb{K} such that $a + b = c + d + e \ne 0$. Then

$$(a, b_2, b_3, c, d, e) \in \operatorname{Col}(M)$$

for some $b_2, b_3 \in \mathbb{K}$ with $b_2 + b_3 = b$.

Proof. Let i_1, i_2, i_3 be the links incident with u in J_k and o_1, o_2, o_3 be the links incident with v. Since the proper (2,3)-pole $N((i_1, j_2), O) = J_k - i_1 - v$ is perfect according to Lemma 4.1, it admits a colouring φ with $\varphi(S(N)) = (a, b, c, d, e)$. Then $\varphi(i_2) = b_2$ and $\varphi(i_3) = b_3$. After removing the vertex u together with the dangling edge j_2 we obtain the (3,3)-pole M with the desired colouring.

The key idea of our superposition is to choose a cycle of the base graph and replace its vertices and edges with the just introduced supervertices and superedges thereby producing a larger multipole called a supercycle. First, we recursively define a k-superpath as follows: A 1-superpath is any (3,3,1)-pole of the form $F_1 \circ W \circ F_2$, where F_1 and F_2 are arbitrary Isaacs superedges. For $k \geq 2$, a k-superpath is a (3,3,k)-pole of the form $SP_{k-1} \circ W \circ F_{k+1}$ for an arbitrary Isaacs superedge F_{k+1} and an arbitrary (k-1)-superpath SP_{k-1} . For $k \geq 2$, a k-supercycle SC_k is a k-pole of the form $\overline{SP_{k-1} \circ W}$, where SP_{k-1} is an arbitrary (k-1)-superpath.

Supercycles and superpaths have colouring properties similar to those of cycles and paths, respectively. At this point it may be convenient to regard paths and cycles as multipoles with a dangling edge attached to each vertex. To be more precise, for $k \ge 1$, the *k*-cycle is the *k*-pole $C_k(e_1, e_2, \ldots, e_k)$ consisting of a cycle $(v_1v_2 \ldots v_k)$ with a dangling edge e_i attached to v_i for each $i \in \{1, \ldots, k\}$. For $k \ge 1$, the *k*-path is the (1, 1, k)-pole $P_k((i), (o), (r_1, r_2, \ldots, r_k))$ whose underlying graph is a path $v_1 \ldots v_k$, the dangling edges *i* and *o* are incident with vertices v_1 and v_k , respectively, and the dangling edge r_i is incident with v_i for each $i \in \{1, \ldots, k\}$. Note that the *k*-path contains *k* vertices and k-1 links. **Lemma 4.3.** Let $k \ge 1$ and let SP_k be an arbitrary k-superpath. Consider an arbitrary element

$$(c_i, c_o, c_1, \ldots, c_k) \in \operatorname{Col}(P_k)$$

for a k-path P_k and arbitrary elements a_1 , a_2 , a_3 , b_1 , b_2 , $b_3 \in \mathbb{K}$ such that $a_1 + a_2 + a_3 = c_i$ and $b_1 + b_2 + b_3 = c_o$. Then

$$(a_1, a_2, a_3, b_1, b_2, b_3, c_1, \dots, c_k) \in \operatorname{Col}(SP_k).$$

Proof. We proceed by induction on k. For k = 1, we have verified that the (3, 3, 1)-pole $A_5 \circ W \circ A_5$ has the desired colouring set by using a computer. Every other superedge A_{ℓ} can be obtained from A_5 by substituting the (3, 3)-pole $Y_{\ell-3}$ for the only copy of Y_2 contained in A_5 . Since $\operatorname{Col}(Y_{\ell-3}) = \operatorname{Col}(Y_2)$, we have $\operatorname{Col}(SP_1) = \operatorname{Col}(A_x \circ W \circ A_y)$ for any odd $x, y \geq 5$.

Now, assume that the statement holds for any (k-1)-superpath. Consider a k-superpath $SP_k = F_1 \circ W_1 \circ F_2 \circ \cdots \circ W_k \circ F_{k+1}$ for some $k \ge 2$. Let $(c_i, c_o, c_1, \ldots, c_k) \in \operatorname{Col}(P_k)$. Consider arbitrary elements $a_1, a_2, a_3, b_1, b_2, b_3 \in \mathbb{K}$ such that $a_1 + a_2 + a_3 = c_i$ and $b_1 + b_2 + b_3 = c_o$. Choose $d \in \mathbb{K} - \{c_1\}$. By Lemma 4.2, we have $(a_1, a_2, a_3, d, d_1, d_2) \in \operatorname{Col}(F_1)$ for some $d_1, d_2 \in \mathbb{K}$ satisfying Parity Lemma. Then clearly, $(d, d_1, d_2, d + c_1, d_1, d_2, c_1) \in \operatorname{Col}(W_1)$. Finally, by the induction hypothesis, we have $(d + c_1, d_1, d_2, b_1, b_2, b_3, c_2, c_3, \ldots, c_k) \in \operatorname{Col}(F_2 \circ W_2 \circ F_3 \circ \cdots \circ W_k \circ F_{k+1})$ which yields the desired colouring.

Lemma 4.4. If SC_k is an arbitrary supercycle with $k \ge 2$, then $Col(SC_k) = Col(C_k)$.

Proof. Because our superposition is proper, we conclude that $\operatorname{Col}(SC_k) \subseteq \operatorname{Col}(C_k)$. To prove the converse, let $SC_k = \overline{SP_{k-1} \circ W}$, and consider a k-cycle $C_k(e_1, e_2, \ldots, e_k)$; denote by v_i the vertex incident with e_i for each $i \in \{1, 2, \ldots, k\}$. Take a colouring φ of C_k and let $c_i = \varphi(e_i)$ for each $i \in \{1, 2, \ldots, k\}$. The (3, 3, 1)-pole W is colourable in such a way that the flows through its connectors are $a = \varphi(v_{k-1}v_k)$, $b = \varphi(v_kv_1)$ and c_k , respectively. Since $(b, a, c_1, c_2, \ldots, c_{k-1}) \in \operatorname{Col}(P_{k-1})$ according to Lemma 4.3, the (k - 1)-superpath SP_{k-1} admits desired colours on its semiedges, and therefore $\varphi(SC_k) \in \operatorname{Col}(SC_k)$.

In all our superpositions, we substitute a k-supercycle SC_k for a copy C of the k-cycle C_k in a snark G. The resulting snark will be denoted by $Sup(G, C, SC_k)$.

Theorem 4.5. If G is a critical snark, then the graph $G = \text{Sup}(G, C, SC_k)$ is also a critical snark.

Proof. We start the proof by introducing some useful notation. Let G = M * C be a critical snark containing a k-cycle $C(e_1, e_2, \ldots, e_k)$; for each $i \in \{1, 2, \ldots, k\}$ let v_i denote the vertex incident with the dangling edge e_i . Let SC_k be a k-supercycle such that

$$SC_k(r_1, r_2, \dots, r_k) = \overline{F_1 \circ W_1 \circ F_2 \circ \dots \circ W_{k-1} \circ F_k \circ W_k}$$

where, for each $j \in \{1, 2, ..., k\}$, W_j is a copy of the supervertex W, w_j is the only vertex of W_j , and

$$F_j((i_j^1, i_j^2, i_j^3), (o_j^1, o_j^2, o_j^3)) = J_{\ell_j} - (u_j, t_j)$$

for some odd $\ell_j \geq 5$. The index j in all the places where we use it is taken modulo k. The notation for semiedges i_j^1, \ldots, o_j^3 is also used for their original edges (that is, the links of J_{ℓ_j} from which the semiedges arise).



Figure 11: Structure of the snark \tilde{G} used throughout the proof

Let $G = M * SC_k = \operatorname{Sup}(G, C, SC_k)$. A schematic drawing of G is given in Figure 11, where two consecutive superedges F_j and F_{j+1} from the supercycle SC_k are depicted separately. The remaining (k-3)-superpath $F_{j+2} \circ W_{j+2} \circ F_{j+3} \circ \cdots \circ W_{j-2} \circ F_{j-1}$ is denoted by SP_{k-3} (note that $F_{j-1} = F_{j+2+k-3}$).

Since \tilde{G} arises from G by a proper superposition, \tilde{G} is a snark. It remains to show that \tilde{G} is critical. Let x and y be arbitrary adjacent vertices in the snark \tilde{G} . We need to show that the 4-pole $\tilde{G} - [x, y]$ is colourable. The vertices x and y might come from the k-pole M, from a superedge contained in SC_k , or from a supervertex contained in SC_k . Accordingly, the proof splits into several cases.

Case (i). Both x and y belong to the k-pole M. Since G is critical, the 4-pole M*C-[x, y] is colourable. The colourability of $M*SC_k - [x, y]$ then follows from the fact that C and SC_k are colour-equivalent according to Lemma 4.4.

Case (ii). The vertex x belongs to M and $y = w_j$. for some j. By Lemma 3.6, the (2, 2)-pole $N((f_1, f_2), (f_3, f_4)) = G - [x, v_j]$ admits a colouring φ such that $\varphi(f_1) = \varphi(f_2) = \varphi(f_3) = \varphi(f_4) = a$. For this colouring, we have $\varphi(v_{j-1}v_{j-2}) = p$, $\varphi(e_{j-1}) = p + a$, $\varphi(v_{j+1}v_{j+2}) = q$, and $\varphi(e_{j+1}) = q + a$ for some (not necessary distinct) elements $p, q \in \mathbb{K} - \{a\}$; note that p and q must be different from a, for otherwise the colour of some edge would be zero. We colour the (k+1)-pole $SC_k - y$ as shown in Figure 12; the colour of r_i is set to $\varphi(e_i)$ for all $i \neq j$. From Lemmas 4.2 and 4.3 infer that the superedges F_j , F_{j+1} , and the superpath SP_{k-3} admit such colourings.



Figure 12: Colouring for Case (ii)

Case (iii). The vertex x is identical with w_j , for some j, and y belongs to an adjacent superedge, say F_j . Lemma 3.6 now implies that the (2, 2)-pole $N_1 = J_{\ell_j} - [y, t_j]$ can be coloured in such a way that all its dangling edges receive the same colour q. Hence, if we remove the vertex u_j from the (2, 2)-pole N_1 we get the (3, 2, 2)-pole $F'_j = (J_{\ell_j} - u_j) - [y, t_j]$ with its dangling edges coloured (a, b, c, q, q, q, q), where $\{a, b, c\} = \mathbb{K}$.

Since G is a critical snark, Lemma 3.6 implies that the (2, 2)-pole $N((f_1, f_2), (f_3, f_4)) = G - [v_{j-1}, v_j]$ admits a colouring φ such that $\varphi(f_1) = \varphi(f_2) = p \neq a$ and $\varphi(f_3) = \varphi(f_4) = q$ for some $p, q \in \mathbb{K}$, not necessary distinct. Observe that $\varphi(e_{j+1}) = r \neq q$ because e_{j+1} and the semiedge f_4 are both incident with v_{j+1} in N. The colouring of the (2, 2)-pole M - [x, y] is depicted in Figure 13; suitable colourings of F_{j+1} and SP_{k-2} exist according to Lemmas 4.2 and 4.3, respectively.



Figure 13: Colouring for Case (iii)

Case (iv). Both x and y belong to the same superedge F_j for some j. From any colouring of the (2, 2)-pole $J_{\ell_j} - [x, y]$ we can obtain a colouring of the (3, 3, 2, 2)-pole $F'_j = (J_{\ell_j} - (u_j, t_j)) - [x, y]$ assigning to its first six dangling edges colours a, b, c, p, q, r (in this order) such that a + b + c = p + q + r = 0. According to Lemma 3.6, the (2, 2)-pole $G - [v_j, v_{j+1}]$ admits a colouring (a', a', p', p') for some $a' \neq a$ and $p' \neq p$, therefore we can colour the (2, 2)-pole $\tilde{G} - [x, y]$ as shown in Figure 14. A colouring of $F_{j+1} \circ W_{j+1} \circ SP_{k-3}$, which is a (k-2)-superpath, can be obtained from Lemma 4.3.



Figure 14: Colouring for Case (iv)

Case (v). The vertices x and y belong to two different superedges, say, $x \in V(F_j)$ and $y \in V(F_{j+1})$. Set $F'_j = F_j - x$ (we discard the dangling edge adjacent to x in F'_j) and similarly $F'_{j+1} = F_{j+1} - y$ (see Figure 15). By Lemma 3.6, the exists a colouring of the (2, 2)-pole $N_1 = G - [v_j, v_{j+1}]$ that assigns the same colour a to all its semiedges. The link e_{j-1} of N_1 receives a colour $b \neq a$, because it is incident with the dangling edge incident to v_{j-1} , which is coloured a.

We proceed by finding suitable colourings for F'_j and F'_{j+1} . First, we colour F'_j . By the construction of the Isaacs superedge F_j , the vertices u_j and t_j are at distance at least 4 in J_{ℓ_j} , so the distance between u_j and x is at least 3. Thus, according to Lemma 4.2, the (3,3)-pole $N_2((i_j^1, i_j^2, i_j^3), (f_1, f_2, f_3)) = J_{\ell_j} - (u_j, x)$ admits a colouring φ such that $\varphi(i_1^j) + \varphi(i_2^j) + \varphi(i_3^j) = a, \varphi(f_1) = a = \varphi(o_1^j) + \varphi(o_2^j)$ and $\varphi(f_2) + \varphi(f_3) = 0$. If we remove the vertex t_j along with its dangling edge f_1 from N_2 , we get two more dangling edges o_j^1 and o_i^2 which receive colours b' and c', respectively, with b' + c' = a.

Next, we find a suitable colouring for F'_{j+1} . The (2, 2)-pole $N_3((i^1_{j+1}, i^2_{j+1}), (g_3, g_4)) = J_{\ell_{j+1}} - [u_{j+1}, y]$ admits a colouring ψ such that $\psi(i^1_{j+1}) = \psi(i^2_{j+1}) = c'$ and $\psi(g_3) = \psi(g_4)$. Under the colouring ψ , the links o^1_{j+1} , o^2_{j+1} and o^3_{j+1} receive pairwise distinct colours p, q, and r, respectively. We may assume that $p \neq a$ because if we had $\psi(o^1_{j+1}) = a$, then we could swap the colours a, b' in ψ and get $\psi(o^1_{j+1}) \neq a$.

By using all the described colourings and Lemma 4.3, the (2, 2)-pole G - [x, y] can be coloured as shown in Figure 15. This completes Case (v) as well as the entire proof. \Box



Figure 15: Colouring for Case (v)

We can apply Theorem 4.5 to constructing cyclically 6-connected critical snarks. For strictly critical snarks, however, we need the following lemma.

Lemma 4.6. Let G be a snark and let \hat{G} be a snark obtained from G by a proper superposition which replaces each vertex v of G with a supervertex V_v and each edge of G with a proper superedge E_e . If $\{u, v\}$ is a removable pair of vertices of G, then for each vertex \tilde{u} from V_u and each vertex \tilde{v} from V_v the pair $\{\tilde{u}, \tilde{v}\}$ forms a removable pair of vertices of \tilde{G} .

Proof. Suppose to the contrary that the 6-pole $G - (\tilde{u}, \tilde{v})$ admits a colouring $\tilde{\varphi}$. Since the superedge E_e is proper for each edge $e \in E(G)$, both connectors of E_e have the same nonzero total flow under $\tilde{\varphi}$; we denote this common value by $\varphi(e)$. The assignment $e \mapsto \varphi(e)$ defines a mapping $\varphi \colon E(G) \to \mathbb{K}$. For each vertex $w \in V - \{u, v\}$, the values of φ on the edges around w sum to zero due to Parity Lemma applied to V_w . The mapping φ thus yields a 3-edge-colouring of the 6-pole G - (u, v), contradicting the fact that $\{u, v\}$ was a removable pair of vertices of G.

Corollary 4.7. If G is a strictly critical snark with a k-cycle C, then the snark $\tilde{G} =$ Sup (G, C, SC_k) is also strictly critical.

Proof. The snark $\tilde{G} = \text{Sup}(G, C, SC_k)$ is critical due to Theorem 4.5. Since G is not bicritical, it has a pair of non-adjacent vertices whose removal leaves an uncolourable graph. According to Lemma 4.6, this pair yields a pair of removable vertices also in G', thus it cannot be bicritical.

We are now ready to prove our main result, which implies Theorem 1.2.

Theorem 4.8. There exists a cyclically 6-connected strictly critical snark of order n in each of the following cases.

- (i) $n \equiv 0 \pmod{8}$ and $n \geq 320$.
- (ii) $n \equiv 2 \pmod{8}$ and $n \ge 306$.
- (iii) $n \equiv 4 \pmod{8}$ and $n \ge 324$.

(iv) $n \equiv 6 \pmod{8}$ and $n \geq 342$.

Proof. First, we construct the smallest snark for each of the residue classes (mod 8) by taking a suitable strictly critical snark G and replacing a suitable cycle in G with a supercycle. By Corollary 4.7, the resulting snark is also strictly critical. The four base snarks used in the constructions for each even residue class (mod 8) are depicted in Figure 16 together with the cycle that we replace. In each case, all the superedges used for the corresponding supercycle are isomorphic to A_5 , which has 18 vertices.

To prove Item (i), we start with one of the smallest strictly critical snarks G_{32} (described also in [4]) which is a dot product of an I-extension of one of the two Goldblerg-Loupekine snark of order 22 and the Petersen graph (see Figure 16a). We choose the 16-cycle indicated in Figure 16a and after replacing it with the 16-supercycle containing superedges isomorphic to A_5 we obtain a strictly critical snark G_{320} of order 320. The snark G_{32} contains one pair $\{x, y\}$ of non-adjacent removable vertices and these two vertices remain removable in the snark G_{320} according to Lemma 4.6.

For the remaining three items, we take the snark G_{36} depicted in Figures 16b, 16c, and 16d which is of the form $H_6 * \text{TTT}_{sc}(T_P, T_P, T_P)$, where T_P denotes the proper (2, 3)-pole constructed from the Petersen graph (see Section 3 for the definitions of H_6 and TTT_{sc}). If we replace the 15-cycle indicated in Figure 16b, we obtain a strictly critical snark G_{306} of order 306. Replacing the 16-cycle indicated in Figure 16c yields G_{324} (of order 324) and replacing the 17-cycle in Figure 16d yields G_{342} (order 342).

Let \tilde{v} denote the only vertex contained in the supervertex that replaces a vertex v. In G_{36} , every 2-element subset of $\{x, y, z\}$ (see Figure 16) is a pair of non-adjacent removable vertices, thus according to Lemma 4.6, also every pair of vertices from the set $\{\tilde{x}, \tilde{y}, \tilde{z}\}$ is removable in G_{306} , G_{324} , and G_{342} , respectively.

For each $n \in \{320, 306, 324, 342\}$, the snark G_n is strictly critical, has order n and, as we have verified by using a computer program, it is cyclically 6-connected. If we replace one of the superedges A_5 contained in G_n with the superedge A_{5+2k} for an integer $k \ge 1$, we obtain a snark G_{n+8k} of order n + 8k. The snark G_{n+8k} is strictly critical in view of Corollary 4.7. It is cyclically 6-connected: the larger constructed snarks arise from the smallest ones by replacing Y_2 with Y_{2m} for some m, and such a substitution maintains cyclic connectivity 6 in the same way as it does for the Isaacs snarks. Hence, the snarks G_n constructed above possess all the required properties, and the proof is complete. \Box

To our best knowledge, the snark G_{306} is the smallest known cyclically 6-connected strictly critical snark. It is hard to believe that there are no smaller such snarks, but discovering them would very likely require a new method of constructing cyclically 6connected snarks, since superposition tends to produce large graphs. This suggests the following problem.

Problem 1. Does there exist a cyclically 6-connected strictly critical snark of order smaller than 306?

We believe that a statement about bicriticality analogous to Theorem 4.5 might also be true, but the proof seems too complicated if we use our current method.

Problem 2. Is it true that for every bicritical snark G any k-cycle C in G, the snark $Sup(G, C, SC_k)$ is always bicritical?

If the answer was yes (perhaps with some additional assumptions about cycles being replaced), it would provide a useful tool for constructions of cyclically 6-connected bicritical snarks.



Figure 16: The snarks G_{32} and G_{36} with cycles to replace in superposition

Acknowledgements

The authors acknowledge partial support from the research grants APVV-19-0308, VEGA 1/0743/21 and VEGA 1/0727/22.

References

- L. D. Andersen, H. Fleischner, B. Jackson, Removable edges in cyclically 4-edgeconnected cubic graphs, Graphs Combin. 4 (1988), 1–21.
- [2] G. Brinkmann, K. Coolsaet, J. Goedgebeur, H. Mélot, House of Graphs: a database of interesting graphs, Discrete Appl. Math. 161 (2013), 311-314. Available at http: //hog.grinvin.org/.
- [3] G. Brinkmann, J. Goedgebeur, J. Hägglund, K. Markström, Generation and properties of snarks, J. Combin. Theory Ser. B 103 (2013), 468–488.
- [4] M. Chladný, M. Skoviera, Factorisation of snarks, Electron. J. Combin. 71 (2010), Research Paper R32, 53 p.
- [5] C. N. da Silva, C. L. Lucchesi, Flow-critical graphs, Electron. Notes Discrete Math. 30 (2008), 165–170.
- [6] C. N. da Silva, L. Pesci, C. L. Lucchesi, Snarks and flow-critical graphs, Electron. Notes Discrete Math. 44 (2013), 299–305.

- [7] B. Descartes, Network-colourings, The Math. Gazette 32 (1948), 67–69.
- [8] M. DeVos, J. Nešetřil, A. Raspaud, On edge-maps whose inverse preserves flows or tensions, in: Graph Theory in Paris (J. A. Bondy et al., Eds.), Trends in Mathematics, Birkhäuser, Basel, 2006, pp. 109–138.
- S. Grünewald, E. Steffen, Cyclically 5-edge-connected non-bicritical critical snarks, Discuss. Math. Graph Theory 19 (1999), 5–11.
- [10] F. Jaeger, T. Swart, Problem Session, Combinatorics 1979, Part II (M. Deza and I. G. Rosenberg, eds.), Ann. Discrete Math. 9 (1980), p. 305.
- [11] M. Kochol, Snarks without small cycles, J. Combin. Theory Ser. B 67 (2006), 779– 791.
- [12] E. Máčajová, M. Škoviera, Irreducible snarks of given order and cyclic connectivity, Discrete Math. 306 (2006), 34–47.
- [13] E. Máčajová, M. Škoviera, Critical and flow-critical snarks coincide, Discuss. Math. Graph Theory 41 (2021), 503–511.
- [14] E. Máčajová, M. Škoviera, Superposition of snarks revisited, European J. Combin. 91 (2021), Article 103220.
- [15] J. Mazák, J. Rajník, M. Skoviera, Morphology of small snarks, Electron. J. Combin. 29 (2022), #P4.30.
- [16] W. McCuaig, Edge reductions in cyclically k-connected cubic graphs, J. Combin. Theory Ser. B 56 (1992), 16–44.
- [17] R. Nedela, M. Škoviera, Decompositions and reductions of snarks, J. Graph Theory 22 (1996), 253–279.
- [18] R. Šámal, Cycle-continuous mappings order structure, J. Graph Theory 85 (2017), 56–73.
- [19] E. Steffen, Classifications and characterizations of snarks, Discrete Math. 188 (1998), 183–203.
- [20] E. Steffen, Non-bicritical critical snarks, Graphs Combin. 15 (1999), 473–480.
- [21] N. C. Wormald, Classifying k-connected cubic graphs, Lecture Notes in Math. 748 (1979), 199–206.