CANONICAL FORM OF MATRIX FACTORIZATIONS FROM FUKAYA CATEGORY OF SURFACE

CHEOL-HYUN CHO AND KYUNGMIN RHO

ABSTRACT. This paper concerns homological mirror symmetry for the pair-of-pants surface (A-side) and the non-isolated surface singularity xyz = 0 (B-side). Burban-Drozd classified indecomposable maximal Cohen-Macaulay modules on the B-side. We prove that higher-multiplicity band-type modules correspond to higher-rank local systems over closed geodesics on the A-side, generalizing our previous work for the multiplicity one case. This provides a geometric interpretation of the representation tameness of the band-type maximal Cohen-Macaulay modules, as every indecomposable object is realized as a geometric object.

We also present an explicit canonical form of matrix factorizations of *xyz* corresponding to Burban-Drozd's canonical form of band-type maximal Cohen-Macaulay modules. As applications, we give a geometric interpretation of algebraic operations such as AR translation and duality of maximal Cohen-Macaulay modules as well as certain mapping cone operations.

CONTENTS

1.	Introd	uction	2
	1.1.	Proof of main theorem	6
2.	Locali	zed Mirror Functor and Its Computation	10
	2.1.	Localized mirror functor for pair-of-pants surface	10
	2.2.	Computation of localized mirror functor	12
	2.3.	Illustration with an example	14
	2.4.	Higher rank computation	16
3.	Matrix	Factorizations from Loops with a Local System	17
	3.1.	Loop words and canonical form of immersed loops	17
	3.2.	Loop data and canonical form of loops with a local system	20
	3.3.	Matrix factorizations from canonical form of loops with a local system	20
	3.4.	Periodic case	24
	3.5.	Non-cylinder-free case $w' = (2, 2, 2)^{\tau}$	26
4.	Matrix	Factorizations from Maximal Cohen-Macaulay Modules	29
	4.1.	Maximal Cohen-Macaulay modules and Macaulayfication	29
	4.2.	Band data and canonical form of maximal Cohen-Macaulay modules	30
	4.3.	Conversion formula between loop/band data	32
	4.4.	Matrix Factorizations from canonical form of maximal Cohen-Macaulay modules	33
	4.5.	Degenerate case	36
5.	Applic	ations	39
	5.1.	Flip of loops and dual of modules	39
	5.2.	Reverse of loops and shift of modules	46
	5.3.	Higher rank/multiplicity and twisted complexes	51
Ap	pendix	A. Relevant Categories	59
	A.1.	A_{∞} -category	59

A.2.	Compact Fukaya category of a surface	60
A.3.	Categories of matrix factorizations	65
A.4.	Stable categories and Eisenbud's equivalence	66
Reference	es	68

1. INTRODUCTION

A version of Homological mirror symmetry (HMS) conjecture of Kontsevich [Kon95, Kon98] says that the derived wrapped Fukaya category of a symplectic manifold Σ (A-side) and the singularity category of its mirror Landau-Ginzburg (LG) model ($X, W : X \to \mathbb{C}$) (B-side) are equivalent:

 $D^{\pi}(W\operatorname{Fuk}(\Sigma)) \simeq D_{\operatorname{sing}}(W^{-1}(0)).$

HMS between punctured Riemann spheres $\Sigma = S^2 \setminus \{n \text{ points}\}$ $(n \ge 3)$ and the corresponding LG models was established in [AAE⁺13].

In the case of the 3-punctured sphere, the corresponding LG model is given by (\mathbb{C}^3, xyz) .



This equivalence has been shown on the level of generators, which in this case consist of any two of three non-closed curves L_{xy} , L_{yz} and L_{zx} on the A-side (Figure 4), and the corresponding objects on the B-side (Remark 4.9). This work of Abouzaid-Auroux-Efimov-Katzarkov-Orlov [AAE⁺13] inspired a lot of further developments in homological mirror symmetry. However, it is hard to compare more complicated objects in both sides directly from this equivalence.

On the other hand, it is known that the following three categories are equivalent [Eis80, Buc21, Orl03]:

$$\underline{\mathrm{MF}}(xyz) \stackrel{\mathrm{Eisenbud}}{\longrightarrow} \underline{\mathrm{CM}}(A) \stackrel{\mathrm{Buchweitz}}{\longrightarrow} D_{\mathrm{sing}}(\hat{X}_0)$$

Here, $\underline{MF}(xyz)$ is the homotopy category of **matrix factorizations** of xyz, $\underline{CM}(A)$ is the stable category of **maximal Cohen-Macaulay modules** over $A := \mathbb{C}[[x, y, z]]/(xyz)$, and $D_{sing}(\hat{X}_0)$ is the singularity category of $\hat{X}_0 := \text{Spec}(A)$. In this paper, we work with power series rings instead of polynomial rings (Remark 1.5).

In a recent work [BD17a], Burban-Drozd developed a new representation-theoretic method to deal with maximal Cohen-Macaulay modules over certain non-isolated surface singularities including *A*. As a consequence, they classified all indecomposable classes of such modules, which fall into **band-type** (continuous series) and **string-type** (discrete series). This proves that those singularities have **tame** Cohen-Macaulay representation type.

Thus, a natural question is which objects of the Fukaya category correspond to the indecomposable maximal Cohen-Macaulay modules over *A* (Remark 9.8.8 in [BD17a]). Especially, it is of great interest whether their symplectic counterparts are realized as **geometric objects** in the Fukaya category. This question will be answered in the present paper by giving an explicit correspondence:

Theorem 1.1. Under homological mirror symmetry, there is a one-to-one correspondence

{closed geodesics¹ in Σ with an indecomposable local system}/~_{gauge equivalence} $\stackrel{1}{\leftarrow}$

1:1 ↔

$\{$ band-type indecomposable objects in $\underline{CM}(A)\}/\sim_{isomorphism}$,

where Σ is given a hyperbolic metric with three cusps.

In our setting of the Fukaya category of Σ , objects are oriented immersed curves in Σ with a local system. We call them **loop-type** or **arc-type** according to whether the curve is a loop (closed curve) or an arc (starting and ending at $\partial \Sigma$). Closed geodesics are representatives in certain (but not all) free homotopy classes of oriented loops in Σ . Thus, Theorem 1.1 describes a correspondence between indecomposable objects of loop-type in the Fukaya category and band-type in the category of maximal Cohen-Macaulay modules. A similar correspondence between arc-type and string-type objects can be also made, but we do not cover them in the present paper (see Remark 4.9).

In our previous work [CJKR22], we already established a correspondence between loop-type objects of rank $\rho = 1$ and band-type objects of multiplicity $\mu = 1$, and found a canonical form of matrix factorizations for that case. The main purpose of the present paper is to extend it to a correspondence between loop-type objects of arbitrary **geometric rank** ρ and band-type objects of arbitrary **algebraic multiplicity** μ .

To convert objects of Fukaya category into matrix factorizations, we use the localized mirror functor

$\mathscr{F}^{\mathbb{L}}: D^{\pi}(W\operatorname{Fuk}(\Sigma)) \longrightarrow \underline{\operatorname{MF}}(xyz)$

developed by the first author with Hong-Lau in [CHL17]. In the present work, we elaborate its computational aspect to apply it to higher-rank local systems. In particular, we give an explicit formula (2.6) for resulting matrix factorizations, and use it to deduce matrix factorizations of higher-rank local systems directly from the result on rank 1 cases (Proposition 2.5). It presents us a **canonical form of matrix factorizations** for higher-rank objects in terms of **loop data**, extending the previous version for rank 1 case in [CJKR22].

Burban-Drozd also provided a canonical form of maximal Cohen-Macaulay modules in terms of **band data**. But the corresponding matrix factorizations under Eisenbud's equivalence were not known, due to the complexity of *Macaulayfication* process. With the help of homological mirror symmetry, we now have a candidate. Indeed, we demonstrate that our canonical form of matrix factorizations fits perfectly into this framework under an explicit **conversion formula** between loop data and band data. The presence of this conversion formula suggests that it would have been hardly attainable otherwise.

For the proof, we define the notion of (λ, Λ) -**substitution pair** and use its homological property. It enables us to extend the Macaulayfication result as well as conversion formula obtained in [CJKR22] to higher-multiplicity cases (Theorem 4.14).

After all, it turns out that the geometric rank ρ and the algebraic multiplicity μ coincide in a majority of cases. But surprisingly, there are a few (countably many) exceptions called **degenerate cases**, where two parameters differ by 1 as $\rho = \mu - 1$. This can be interpreted as an inevitable phenomenon following from the elimination of the regular module *A* that occurs when we take the stable category <u>CM</u>(*A*) = CM(*A*)\{*A*}.

Analyzing the correspondence of objects in those cases is quite tricky both on geometric and algebraic sides: The geometric loop is freely homotopic to the reference loop (Seidel Lagrangian), so we perturb it to prevent an immersed cylinder (§3.5). On algebraic side, the corresponding module has one exceptional Macaulayfying element, which does not appear as a family of λ and cannot be obtained from the above (λ , Λ)-substitution process. So we perform an additional computation for this case and find that the existence of such an element causes the degeneration $\rho = \mu - 1$ (§4.5).

¹In our paper, a closed geodesic is always oriented, non-periodic, immersed (i.e., not necessarily simple) and considered up to orientation-preserving reparameterization.

Applications. The correspondence obtained in Theorem 1.1 can be used to relate natural geometric symmetries to algebraic operations. Here we present some of them, while expecting that there will be further interesting translations between two languages. The first two are from geometry to algebra, and the last two are the other way around. (These applications were not presented and have been postponed from [CJKR22] to include general relations between higher rank/multiplicity objects.)

First, taking the **duality functor** Hom_A(-, A) of modules in CM(A) corresponds to **flipping** loops in Fuk(Σ) (see figures in Example 5.7). We show the commutativity of the following diagram of functors in (A_{∞} -)categorical level (Proposition 5.9 + Proposition 5.6). Then we give a clear description of these operations in terms of loop/band data (§5.1.5).

(1.1)
$$H^{0}\operatorname{Fuk}(\Sigma) \xrightarrow{\mathscr{F}^{L}} \underline{MF}(xyz) \xrightarrow{\simeq} \underline{CM}(A)$$
$$flip \downarrow \iota \quad \text{transpose} \downarrow -\operatorname{Tr} \quad dual \downarrow \operatorname{Hom}_{A}(-,A)$$
$$H^{0}\operatorname{Fuk}(\Sigma) \xrightarrow{\mathscr{F}^{L}} \underline{MF}(xyz) \xrightarrow{\simeq} \underline{CM}(A)$$

Second, we consider the **AR translation**, which is given by the **shift functor** of the triangulated category $\underline{CM}(A)$. It is not easy to compute in terms of band data, but it is equivalent to **reversing the orientation** of underlying loops in Fuk(Σ) (Proposition 5.5 + Proposition 5.11), which we can compute in a geometric way. We will give an algorithm to compute them using conversion to the loop data (Proposition 5.12).

$$(1.2) \qquad \begin{array}{c} H^{0}\operatorname{Fuk}(\Sigma) \xrightarrow{\mathscr{F}^{\mathbb{L}}} \underline{\mathrm{MF}}(xyz) \xrightarrow{\simeq} \underline{\mathrm{CM}}(A) \\ \text{orientation} & \int_{I} \operatorname{switching}_{two \ factors} \left[1\right] & \operatorname{shift}_{I} \left[1\right] \\ H^{0}\operatorname{Fuk}(\Sigma) \xrightarrow{\mathscr{F}^{\mathbb{L}}} \underline{\mathrm{MF}}(xyz) \xrightarrow{\simeq} \underline{\mathrm{CM}}(A) \end{array}$$

We remark here that an indecomposable object in degenerate cases and its image under operations in (1.2) are invariant under operations in (1.1). Conversely, if an indecomposable object is invariant under operations in (1.1), either it or its image under operations in (1.2) is of degenerate case.

Third, categories involved in HMS typically possess natural **triangulated structures**. Along with the parameterization by band or loop data, **higher multiplicity/rank** objects are given by some iterated **mapping cones** (or **twisted complexes** in A_{∞} -categories) involving lower multiplicity/rank objects. Proposition 5.22 gives an explicit way to understand higher rank local systems in Fukaya category as twisted complexes of lower rank objects.

Finally, **periodic objects** in both sides are **decomposed** into as many pieces as the number of repetitions ². We give an explicit formula for this decomposition (Theorem 3.21). It shows that non-primitive loops in the Fukaya category are decomposable (Corollary 3.22), which is not obvious on the A-side.

Geometric interpretation of tameness. In Theorem 1.1, we showed that all band-type indecomposable maximal Cohen-Macaulay modules over *A* correspond to explicit geometric objects (rather than abstract twisted complexes) in the Fukaya category. It gives a geometric interpretation ³ of their representation-tameness, which was already proven algebraically in [BD17a]. It is geometrically intuitive that there are only countably many closed geodesics (or free homotopy classes of loops) in Σ . Then an indecomposable local system lying on a fixed loop of rank $\rho \in \mathbb{Z}_{\geq 1}$ is determined by its holonomy up to gauge equivalence, which can be represented (up to basis change) by the $\rho \times \rho$ Jordan block $J_{\rho}(\eta) \in GL_{\rho}(\mathbb{C})$ with some eigenvalue $\eta \in \mathbb{C}^{\times}$. As a result, the elements of sets in Theorem 1.1 are parameterized by closed geodesics in Σ , a rank ρ (discrete parameters), and an eigenvalue of holonomy η (continuous parameter). That is, they consist of countably many one-parameter families.

 $^{^{2}}$ We place this discussion in the middle of the main text (§3.4) since they are also needed to analyze degenerate cases.

³But we are not giving a new proof, as classification of objects in the Fukaya category here relies on the B-side result via HMS.

Relation with other mirror symmetries of surfaces. Recently, there have been many studies on homological mirror symmetry between \mathbb{Z} -graded partially wrapped Fukaya categories (or topological Fukaya categories) of graded marked surfaces (A-side), derived categories of modules over gentle algebras, and derived categories of coherent sheaves on certain **non-commutative curves** (B-side). Gentle algebras have long been an intriguing topic in representation theory since they are derived tame, closed under derived equivalences, and have well-understood indecomposable objects in their derived categories (see [SZ03], [BD17b] and references therein). Their connection with the Fukaya categories was first established in [HKK17], graded marked surfaces corresponding to them were constructed in [LP20], and independently, a closely related algebraic model was constructed in [OPS18]. Their relation with certain non-commutative curves was first found in [BD11], extended to nodal stacky curves in [LP18], and again generalized to much broader class of non-commutative nodal curves in [BD18].

It is especially remarkable that **indecomposable objects** on each side are classified, have a concrete one-to-one correspondence and therefore the derived tameness of gentle algebras can be understood in a (symplectic) geometric way. Also, many purely representation-theoretic problems concerning derived equivalence of finite-dimensional algebras have been attacked and solved using geometric insights and techniques (e.g. [PPP19, APS23, Opp19, KŠ22, CJS22, CHS23, CK24, AP24]).

The triangulated categories which are the focus of the present work are related to \mathbb{Z}_2 -graded Fukaya categories (using oriented Lagrangian curves as objects) of **surfaces** (A-side). Their mirrors are usually given by certain categories of **matrix factorizations** or equivalently, singularity categories of **Landau-Ginzburg models** (B-side). There have been many well-studied homological mirror symmetries. For example, the mirrors of genus two and higher genus closed surfaces were constructed in [Sei08] and [Efi12], respectively. In [AS21], spherical objects in \mathbb{Z}_2 -graded Fukaya categories of closed surfaces were related with simple closed curves with a rank 1 local system. Mirrors of punctured spheres and their cyclic covers were established in [AAE⁺13]. A non-commutative mirror model of punctured surfaces were also discovered in [Boc16], and a related functor was constructed in [CHL15]. Going in a different direction, [AEK21] considers Fukaya categories of singular surfaces and show the reverse direction (switching A- and B-sides) of homological mirror symmetry.

On the level of all **indecomposable objects** in \mathbb{Z}_2 -graded Fukaya categories, nevertheless, their classification and correspondence under mirror symmetry are not known in full generality. Compared to the situation of \mathbb{Z} -graded mirror symmetry, however, it is apparent that there will be much utility of establishing such a strong bridge between curves in Fukaya categories and matrix factorizations (in the global sense of [Orl12]). It will provide more applications of homological mirror symmetry, relating new tame triangulated categories arising from representation theory (other than finite-dimensional algebras) with Fukaya categories of surfaces.

Towards global \mathbb{Z}_2 -graded mirror symmetry. Our previous work [CJKR22] and the present work aim to ignite this new direction of development in the homological mirror symmetry program. There are many good reasons to start with the **pair-of-pants surface** and its mirror xyz = 0, which has been of great interest:

Most importantly, the pair serves as a **building block** to construct more complicated mirror pairs. For example, the idea of constructing the mirror of general Riemann surfaces using their pair-of-pants decompositions appeared in many places in the literature including [Lee16, Nad16, PS19, PS21, PS22]. (The last four uses a sheaf-theoretic version of Fukaya categories, which is different from (but equivalent to) the Floer-theoretic version used in this paper.) See also the well-written survey in [Boc21, §9.4] and references therein. A common approach, often referred to as a **local-to-global principle**, involves proving the compatibility of categorical gluing on both sides, based on the mirror symmetry of the local pair. Moreover, as explained in [CHL18], copies of the localized mirror functor employed in this paper (as its name implies) can be also glued together in order to obtain a mirror functor in the global setting.

Independently of the above gluing formalism, mirror symmetry of the pair-of-pants surface also played a central role in [AAE⁺13] and [HJL24]. The authors consider its cyclic and abelian covers, respectively, and construct their mirror LG model using the symmetry given by the deck transformation groups.

On the other hand, representation theory and classification of objects on B-side have been explicitly developed only for the local model $\mathbb{C}[[x, y, z]]/(xyz)$ in [BD17a]. So we will need to work out the corresponding theory for more general (non-affine) normal crossing surface singularities appearing as mirrors of other Riemann surfaces. After establishing it, we hope to generalize our present results to mirror symmetry of more general Riemann (orbi-)surfaces, which will enhance our understanding of geometric and algebraic tame categories and give many fruitful applications.

1.1. **Proof of main theorem.** In this subsection, we deduce Theorem 1.1 from several results summarized from the body of the paper. The approach to prove it will be completed through the following two steps:

- I. Compute matrix factorizations corresponding to canonical forms of local systems over loops in Σ .
- II. Convert them into Burban-Drozd's canonical form of maximal Cohen-Macaulay modules.

I. Canonical forms of loops with a local system and corresponding matrix factorizations. We take the following specific representatives of free homotopy classes of loops in Σ : Given a **loop word**

$$w' = (l'_1, m'_1, n'_1, l'_2, m'_2, n'_2, \dots, l'_{\tau}, m'_{\tau}, n'_{\tau}) \in \mathbb{Z}^{3\tau}$$

 $(\tau \in \mathbb{Z}_{\geq 1})$, consider the loop L(w') described in Figure 1. We restrict to **normal loop words** (Definition 3.5) so that they (up to shifting) produce only one loop in each hyperbolic free homotopy class. Then they are also in one-to-one correspondence with closed geodesics in Σ (Proposition 3.8).

We introduce a **loop datum** (w', η, ρ) to parameterize loops with a local system, which consists of a **normal loop word** $w' \in \mathbb{Z}^{3\tau}$ $(\tau \in \mathbb{Z}_{\geq 1})$, a **holonomy parameter** $\eta \in \mathbb{C}^{\times}$, and a **(geometric) rank** $\rho \in \mathbb{Z}_{\geq 1}$. The associated **canonical form of a loop with a local system**, denoted by $\mathscr{L}(w', \eta, \rho)$, is given by the loop L(w') together with a rank ρ local system whose holonomy is represented by $J_{\rho}(\eta)^4$ (up to basis change).

Proposition 1.2 (Corollary 3.11). There is a one-to-one correspondence

{closed geodesics in Σ with an indecomposable local system}/ ~_{gauge equivalence}

 $\stackrel{\text{l:l}}{\leftrightarrow} \quad \{\text{non-periodic loop data}\} / \sim_{\text{shifting}}.$



Figure 1. Canonical form of a loop-type object in Fuk (Σ) and <u>MF</u>(*xyz*)

⁴We denote by $J_{\rho}(\eta) \in GL_{\rho}(\mathbb{C})$ the $\rho \times \rho$ Jordan block with eigenvalue $\eta \in \mathbb{C}^{\times}$.

The **localized mirror functor** converts each loop with a local system $\mathscr{L}(w', \eta, \rho)$ into a matrix factorization $\mathscr{F}^{\mathbb{L}}(\mathscr{L}(w', \eta, \rho))$. The corresponding **canonical form of matrix factorization** of *xyz* is

(1.3)
$$\left(\varphi\left(w',\lambda,\rho\right),\psi\left(w',\lambda,\rho\right)\right)^{\sharp}$$

where λ is either η or $-\eta$ depending on w' (see Definition 4.10). Its first component is shown in Figure 1 and the second one is determined by the first (Definition 3.16).

There are some exceptions called **degenerate cases** (i.e., $w' = (2,2,2)^{\tau}$, $\eta = (-1)^{\tau}$), where the second factor $\psi(w', \lambda, \rho)$ is not defined, and we use an alternative form (Definition 3.24)

(1.4)
$$\left(\varphi_{\deg}((2,2,2)^{\tau},1,\rho),\psi_{\deg}((2,2,2)^{\tau},1,\rho)\right).$$

To integrate (1.3) and (1.4) into a unified notation, we denote them as $(\varphi_{(deg)}(w', \lambda, \rho), \psi_{(deg)}(w', \lambda, \rho))$. Namely, it defaults to (1.3) in the general case but adopts (1.4) only in the degenerate cases. Then the relation between the canonical form of loops with a local system and the canonical form of matrix factorizations is summarized as:

Theorem 1.3 (Theorem 3.17 + Theorem 3.25 + Proposition 3.27). For a loop datum (w', η, ρ) and $\lambda = \pm \eta$ determined by the conversion formula Definition 4.10, there is an isomorphism in <u>MF</u>(*xyz*)

$$\mathscr{F}^{\mathbb{L}}(\mathscr{L}(w',\eta,\rho)) \cong (\varphi_{(\mathrm{deg})}(w',\lambda,\rho),\psi_{(\mathrm{deg})}(w',\lambda,\rho)).$$

II. Canonical forms of maximal Cohen-Macaulay modules and corresponding matrix factorizations. Burban-Drozd's classification and canonical form of maximal Cohen-Macaulay modules are best handled in the **category of triples** Tri(A), which was introduced and shown to be equivalent to CM(A) in the same work. The **canonical form** $\Theta(w, \lambda, \mu)$ of a **band-type indecomposable object** in Tri(A) is described in Figure 12. It is parameterized by a **band datum** (w, λ, μ) , which consists of a **band word**

$$w = (l_1, m_1, n_1, l_2, m_2, n_2, \dots, l_{\tau}, m_{\tau}, n_{\tau}) \in \mathbb{Z}^{31}$$

 $(\tau \in \mathbb{Z}_{\geq 1})$, an eigenvalue $\lambda \in \mathbb{C}^{\times}$, and an (algebraic) multiplicity $\mu \in \mathbb{Z}_{\geq 1}$.



Figure 2. Canonical form of a band-type indecomposable object in Tri(*A*) and the corresponding loop with a local system

⁵It is also parameterized by loop data (w', λ, ρ) , but it is more convenient to distinguish two continuous parameters η (for loops with a local system) and λ (for matrix factorizations). They are related under the conversion formula in §4.3.

We denote by $M(w, \lambda, \mu)$ the corresponding object in CM (*A*), and refer to it as the **canonical form of a band-type indecomposable maximal Cohen-Macaulay module** over *A*.

When we take the stable category $\underline{CM}(A) = \underline{CM}(A) / \{A\}$, we lose exactly one isomorphism class [A] of indecomposable objects containing the regular module *A* (Definition A.14). In CM(*A*), it is written in the canonical form as A = M((0,0,0), 1, 1). This implies that for the band datum ((0,0,0), 1, 1), there is no corresponding loop datum.

In §4.3, we define a conversion formula between loop data and band data. It induces a bijection

 $\begin{array}{ll} \left\{ \text{non-periodic loop data} \right\} / \sim_{\text{shifting}} & \stackrel{1:1}{\leftrightarrow} & \left(\left\{ \text{non-periodic band data} \right\} \setminus \left\{ \left((0,0,0), 1, 1 \right) \right\} \right) / \sim_{\text{shifting}} . \\ & \left(w', \eta, \rho \right) & \leftrightarrow & \left(w, \lambda, \mu \right) \end{array}$

Note that the set on the left side already appeared in Proposition 1.2. The set on the right side is in bijection with

{band-type indecomposable objects in $\underline{CM}(A)$ }/ ~_{isomorphism}

by Burban-Drozd's classification (Theorem 4.7). In most cases, we have $\rho = \mu \in \mathbb{Z}_{\geq 1}$. However, for nonperiodic degenerate cases, the correspondence is given by

$$(w' = (2,2,2), \eta = -1, \rho) \quad \leftrightarrow \quad (w = (0,0,0), \lambda = 1, \mu)$$

with $\rho = \mu - 1 \in \mathbb{Z}_{\geq 1}$.

The conversion formula indeed relates band-type indecomposable objects in $\underline{MF}(xyz)$ and $\underline{CM}(A)$ in their canonical forms under Eisenbud's equivalence:

Theorem 1.4 (Theorem 4.14 + Theorem 4.17). For a non-periodic loop datum (w, η, ρ) and band datum (w, λ, μ) related under the conversion formula, there is an isomorphism in CM (A)

 $\operatorname{coker} \varphi_{\operatorname{(deg)}}(w',\lambda,\rho)^6 \cong M(w,\lambda,\mu).$

Mirror symmetry correspondence. Summing up, we have equivalence of categories and correspondence between loop/band-type indecomposable objects as follows (where w' and w are non-periodic):

	r	localized nirror functo	r	Eisenbud		Burban -Drozd	
(1.5)		$\varPhi^{\mathbb{L}}$		coker		\mathbb{F}_{BD}	
(1.5)	$D^{\pi}(W\operatorname{Fuk}(\Sigma))$	\longrightarrow	$\underline{\mathrm{MF}}(xyz)$	$\xrightarrow{\simeq}$	$\underline{\mathrm{CM}}(A)$	$\xrightarrow{\simeq}$	$\underline{\mathrm{Tri}}(A)$
	$\mathscr{L}(w',\eta, ho)$	≅⊥	$\varphi_{(\text{deg})}(w',\lambda,\rho)$	\Leftrightarrow	$M(w,\lambda,\mu)$	\mapsto	$\Theta(w,\lambda,\mu)$

Note that objects in each category are parameterized by loop data or band data. The conversion formula between loop data and band data realizes the one-to-one correspondence between loop/band-type in-decomposable objects (up to isomorphism) in each category as

{closed geodesics in Σ with an indecomposable local system}/ ~_{gauge equivalence}

- $\stackrel{1:1}{\leftrightarrow}$ {non-periodic loop data}/ ~_{shifting}
- $\stackrel{1:1}{\leftrightarrow} \left(\left\{ \text{non-periodic band data} \right\} \setminus \left\{ \left((0,0,0), 1, 1 \right) \right\} \right) / \sim_{\text{shifting}} \right.$
- $\stackrel{1:1}{\leftrightarrow} \quad \left\{ \text{band-type indecomposable objects in } \underline{CM}(A) \right\} / \sim_{\text{isomorphism}}.$

It proves our main Theorem 1.1.

⁶We regard a matrix factorization (φ, ψ) of xyz as a pair of homomorphisms between two free modules over $S := \mathbb{C}[[x, y, z]]$ (Definition A.11), then define $\underline{\varphi} := \varphi \otimes id_A$ as an *A*-module homomorphism (See Theorem A.16).

Periodic case. In each category, objects corresponding to **periodic** loop/band data are **decomposable**: Consider a loop datum (w', η, ρ) with a periodic normal loop word $w' = (\tilde{w}')^N \in \mathbb{Z}^{3\tau}$ $(N \in \mathbb{Z}_{\geq 2})$. Its corresponding band datum (w, λ, μ) has also a periodic band word $w = \tilde{w}^N \in \mathbb{Z}^{3\tau}$. Denote by $\lambda_0, \ldots, \lambda_{N-1} \in \mathbb{C}^{\times}$ and $\eta_0, \ldots, \eta_{N-1} \in \mathbb{C}^{\times}$ the *N*-th roots of λ and η , respectively. Then we have decompositions

$$\mathscr{L}(w',\eta,\rho) \cong \bigoplus_{k=0}^{N-1} \mathscr{L}(\tilde{w}',\eta_k,\rho), \quad \varphi_{(\text{deg})}(w',\lambda,\mu) \cong \bigoplus_{k=0}^{N-1} \varphi_{(\text{deg})}(\tilde{w}',\lambda_k,\mu)^7, \quad M(w,\lambda,\mu) \cong \bigoplus_{k=0}^{N-1} M(\tilde{w},\lambda_k,\mu)$$

in Fuk (Σ), <u>MF</u>(*xyz*), and <u>CM</u>(*A*), respectively. (See Corollary 3.22 + (3.6), Theorem 3.21 + (3.5), and (4.7).)

The loop datum $(\tilde{w}', \eta_k, \rho)$ and the band datum $(\tilde{w}, \lambda_k, \mu)$ also correspond to each other for each k under the conversion formula. Therefore, by Theorem 1.3 and Theorem 1.4, the above three decompositions are compatible with each other in non-degenerate cases. In degenerate cases, the first and the second are still compatible with each other, while the second and the third are not, due to the shifting of rank/multiplicity $\rho = \mu - 1$ (see (4.8) and (4.9)).

Remark 1.5. In this paper, we consider matrix factorizations over the power series ring $S := \mathbb{C}[[x, y, z]]$ and maximal Cohen-Macaulay modules over its quotient $A := \mathbb{C}[[x, y, z]]/(xyz)$, instead of the polynomial ring $\tilde{S} := \mathbb{C}[x, y, z]$ and its quotient $\tilde{A} := \mathbb{C}[x, y, z]/(xyz)$. To distinguish the latter, we denote by MF[xyz] the category of matrix of matrix factorizations of xyz over \tilde{S} , and by CM(\tilde{A}) the category of maximal Cohen-Macaulay modules over \tilde{A} (see [BH98]). Their stable categories and other variations are defined in the same way as in §A.3 and §A.4.

There are obvious faithful functors $\underline{MF}[xyz] \rightarrow \underline{MF}(xyz)$ and $\underline{CM}(\tilde{A}) \rightarrow \underline{CM}(A)$ (see [BH98, Theorem 2.1.3, Corollary 2.1.8]), which are essentially surjective (indeed, every indecomposable object in $\underline{CM}(A)$ is obtained as an image of the functor). Still there are some objects in $\underline{MF}[xyz]$ that are not isomorphic to each other but become isomorphic in $\underline{MF}(xyz)$. For example, the below diagram shows a family of matrix factorizations ($\lambda \in \mathbb{C}^{\times}$) that are not zero objects in $\underline{MF}[xyz]$ but becomes zero objects in $\underline{MF}(xyz)$. (Note that the vertical isomorphisms exist only in the latter category.)

$$\begin{array}{c|c} S^2 & \xrightarrow{\begin{pmatrix} z & \lambda - y \\ 0 & xy \end{pmatrix}} & S^2 & \xrightarrow{\begin{pmatrix} xy & -\lambda + y \\ 0 & z \end{pmatrix}} & S^2 \\ \begin{pmatrix} 1 & 0 \\ z(\lambda - y)^{-1} & 1 \end{pmatrix} \downarrow & \begin{pmatrix} xy & -\lambda + y \\ (\lambda - y)^{-1} & 0 \end{pmatrix} \downarrow & \begin{pmatrix} 1 & 0 \\ z(\lambda - y)^{-1} & 1 \end{pmatrix} \downarrow \\ S^2 & \xrightarrow{\begin{pmatrix} xyz & 0 \\ 0 & 1 \end{pmatrix}} & S^2 & \xrightarrow{\begin{pmatrix} 1 & 0 \\ 0 & xyz \end{pmatrix}} & S^2 \end{array}$$

They come from a loop with holonomy λ that is homotopic to one of the boundary circles in Σ , which was excluded in our correspondence in Theorem 1.1. Thus, <u>MF[xyz]</u> has more objects than <u>MF(xyz)</u>. It would be interesting to know how many objects in the gap, and whether they are also realized in a geometric way.

Acknowledgement. This work started as an attempt to understand the mirror A-side of Igor Burban and Yuriy Drozd's representation theory of maximal Cohen-Macaulay modules over non-isolated surface singularities in [BD17a], and we are grateful to Igor Burban for explanations, discussions and many great advice on this work. We thank Yong-Geun Oh, Henning Krause, Juan Omar Gomez and Sibylle Schroll for their interest on this project as well as helpful comments, Wassilij Gnedin for carefully reading and revising the draft from a representation-theoretic perspective, Severin Barmeier, Dongwook Choa, Wonbo Jeong, Sangjin Lee and Sangwook Lee for many useful discussions in symplectic geometry, and Hanwool Bae for explaining to us his thesis. The second author is also grateful to Jongil Park, Jae-Hoon Kwon, Hansol Hong, Philsang Yoo and Igor Burban for many great comments in his thesis defense [Rho23], where most parts of this work were discussed. We also thank Kyoungmo Kim for sharing his idea and helpful discussions on this topic and other joint works. The second author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(RS-2023-00248895), and the German Research Foundation SFB-TRR 358/1 2023 — 491392403.

⁷Note that if the left side is in a degenerate case, exactly one of direct summands in the right side is in a degenerate case.

2. LOCALIZED MIRROR FUNCTOR AND ITS COMPUTATION

This section is devoted to an elaboration of the computational aspect of the localized mirror functor [CHL17] applied to the *pair-of-pants surface*. Especially, we develop a formula for finding each component of the matrix factorization corresponding to a higher-rank local system (E, ∇) over a loop *L*. Every convention and notation for such an object $\mathcal{L} = (L, E, \nabla)$ is based on our geometric setting of the compact Fukaya category (with immersed loops with a local system) explained in Section A.2.

2.1. **Localized mirror functor for pair-of-pants surface.** We call a smooth surface with boundary Σ diffeomorphic to the complement in S^2 of three distinct points a **pair-of-pants surface**. Consider a marked loop $\mathbb{L} = (\mathbb{L}, e_{\mathbb{L}}, o_{\mathbb{L}})$ in Σ described in Figure 3, which is called the **Seidel Lagrangian**. (Note that its self-intersections are transversal. See also [Sei11].) Assume that the areas of two triangles bounded by \mathbb{L} in Σ are the same ⁸. We put on its domain a trivial line bundle $E_{\mathbb{L}} = S^1 \times \mathbb{C}$ equipped with a flat connection $\nabla_{\mathbb{L}}$ whose holonomy is -1 at the point \bigstar marked in Figure 3. We assume that the triple ($\mathbb{L}, E_{\mathbb{L}}, \nabla_{\mathbb{L}}$) is an object of Fuk (Σ) and still denote it as \mathbb{L} for simplicity.



Figure 3. Seidel Lagrangian \mathbb{L} in Σ



Figure 4. Generating objects of $D^{\pi}W$ Fuk (Σ)

Since L has 3 self-intersections, we have

$$\chi^{0}(\mathbb{L},\mathbb{L}) = \left\{ e_{\mathbb{L}}, \overline{X}, \overline{Y}, \overline{Z} \right\}, \quad \chi^{1}(\mathbb{L},\mathbb{L}) = \{ o_{\mathbb{L}}, X, Y, Z \}$$

We trivialize $E_{\mathbb{L}}$ over $S^1 \setminus \bigstar$ as in Proposition A.7, which yields identifications $E_{\mathbb{L}}|_{\bullet} \cong \mathbb{C}$ for each $\bullet \in \chi(\mathbb{L}, \mathbb{L})$. Then we have Hom_{\mathbb{C}} $(E_{\mathbb{L}}|_{\bullet}, E_{\mathbb{L}}|_{\bullet})^9 \cong$ Span_{\mathbb{C}} $\{\bullet\}$ as noted in Remark A.8, and hence

$$\hom^{0}(\mathbb{L},\mathbb{L}) = \operatorname{Span}_{\mathbb{C}}\left\{e_{\mathbb{L}}\overline{X},\overline{Y},\overline{Z}\right\} \quad \hom^{1}(\mathbb{L},\mathbb{L}) = \operatorname{Span}_{\mathbb{C}}\left\{o_{\mathbb{L}},X,Y,Z\right\}$$

Seidel Lagrangian \mathbb{L} is very special and useful because it has a *weak bounding cochain* (or it is *weakly unobstructed*) in the sense of [FOOO09]. It enables \mathbb{L} to serve as a reference of the localized mirror functor.

Proposition 2.1. [CHL17] A linear combination $b = xX + yY + zZ \in \text{hom}^1(\mathbb{L},\mathbb{L})$ is a weak bounding cochain for any $x, y, z \in \mathbb{C}$. That is,

(2.1)
$$\mathfrak{m}_0^b(\mathbb{L}) := \sum_{i=1}^\infty \mathfrak{m}_i(\underbrace{b,\ldots,b}_i) = xyz \cdot e_{\mathbb{L}}$$

and hence (\mathbb{L}, b) has the disk potential $W^{\mathbb{L}} = xyz$.

Proof. All \mathfrak{m}_1 -terms in the left side vanish because there are no bigons. There are several non-zero \mathfrak{m}_2 -terms such as $\mathfrak{m}_2(X, Y) = \overline{Z} = -\mathfrak{m}_2(Y, X)$ coming from the front and back triangles, but they cancle each other because the holonomy -1 contributes only to the back triangle. The only non-zero higher $\mathfrak{m}_{\geq 3}$ -term is $\mathfrak{m}_3(X, Y, Z) = xyz \cdot e_{\mathbb{L}}$, which comes from the front triangle bounded by \mathbb{L} passing through $e_{\mathbb{L}}$ (Remark A.9.(3)). This is the only surviving term in $\mathfrak{m}_0^b(\mathbb{L})$, which gives the disk potential $W^{\mathbb{L}} = xyz$.

 $^{^{8}}$ This condition is necessary when we consider the Novikov field Λ instead of our field $\mathbb{C}.$

⁹Note that although two $E_{\mathbb{L}}|_{\bullet}$'s have the same notation, they are actually fibers of *E* over different preimages (in *S*¹) of the point $\bullet \in \Sigma$ under *L*, as noted in the definition in (A.4).

Such a pair (\mathbb{L} , *b*) defines an A_{∞} -functor from Fukaya category to the A_{∞} -category MF_{A_{∞}} (*xyz*) of matrix factorizations of *xyz*, called the **localized mirror functor** [CHL17]. (Here we follow the convention in [CHL15].) It is based on the deformation theory of A_{∞} -operations, as we will see below:

For any object $\mathscr{L} = (L, E, \nabla)$ in *W* Fuk $(\Sigma)^{10}$, note that two sets $\chi^{\bullet}(L, \mathbb{L})$ ($\bullet \in \mathbb{Z}_2$) are finite sets of the same cardinality $\tau := \frac{1}{2} |L \cap \mathbb{L}|$. Therefore, two \mathbb{C} -vector spaces

(2.2)
$$\operatorname{hom}^{\bullet}(\mathscr{L},\mathbb{L}) = \bigoplus_{p \in \chi^{\bullet}(L,\mathbb{L})} \operatorname{Hom}_{\mathbb{C}}(E|_{p},\mathbb{C}) = \bigoplus_{p \in \chi^{\bullet}(L,\mathbb{L})} (E|_{p})^{*} \quad (\bullet \in \mathbb{Z}_{2})$$

have the same finite dimension $\tau \rho$, where ρ is the rank of *E*.

We define the **deformed differential**

$$\mathfrak{m}_{1}^{0,b}$$
: hom $(\mathscr{L},\mathbb{L}) \to$ hom $(\mathscr{L},\mathbb{L}), \quad f \mapsto \mathfrak{m}_{1}^{0,b}(f) := \sum_{i=0}^{\infty} \mathfrak{m}_{1+i}(f,\underbrace{b,\ldots,b}_{i}),$

which is a \mathbb{C} -linear map of degree 1 and satisfies $(\mathfrak{m}_1^{0,b})^2 = xyz \cdot \mathrm{id}_{\mathrm{hom}(\mathscr{L},\mathbb{L})}$ by the following lemma, which we recall for the reader's convenience:

Lemma 2.2. [CHL17] For a weak bounding cochain (\mathbb{L}, b) with disk potential $W^{\mathbb{L}}$, we have

$$\left(\mathfrak{m}_{1}^{0,b}\right)^{2} = W^{\mathbb{L}} \cdot \mathrm{id}_{\mathrm{hom}(\mathscr{L},\mathbb{L})}.$$

Proof. For any $f \in \hom^{\bullet}(\mathcal{L}, \mathbb{L})$ (• $\in \mathbb{Z}_2$), we can write

$$\mathfrak{m}_{1}^{0,b}\left(\mathfrak{m}_{1}^{0,b}\left(f\right)\right) = \sum_{j=0}^{\infty} \sum_{i=0}^{\infty} \mathfrak{m}_{1+j}\left(\mathfrak{m}_{1+i}\left(f,\underbrace{b,\dots,b}_{i}\right),\underbrace{b,\dots,b}_{j}\right)$$
$$= -\sum_{l_{0},l_{1}\geq0} \sum_{k=1}^{\infty} (-1)^{|f|-1} \mathfrak{m}_{2+l_{0}+l_{1}}\left(f,\underbrace{b,\dots,b}_{l_{0}},\mathfrak{m}_{k}(\underbrace{b,\dots,b}_{k}),\underbrace{b,\dots,b}_{l_{1}}\right)$$

where the second identity follows from the A_{∞} -relations (A.1). Using the identity $\sum_{k=1}^{\infty} \mathfrak{m}_k(b, \dots, b) = W^{\mathbb{L}} \cdot e_{\mathbb{L}}$ and the fact that $e_{\mathbb{L}}$ is a unit (A.2), we can rewrite it as

$$\sum_{l_0,l_1\geq 0} (-1)^{|f|} \mathfrak{m}_{2+l_0+l_1} \left(f, \underbrace{b, \dots, b}_{l_0}, W^{\mathbb{L}} \cdot e_{\mathbb{L}}, \underbrace{b, \dots, b}_{l_1} \right) = (-1)^{|f|} \mathfrak{m}_2 \left(f, W^{\mathbb{L}} \cdot e_{\mathbb{L}} \right) = W^{\mathbb{L}} \cdot f,$$

which proves the claim.

Restricting the domain of $\mathfrak{m}_1^{0,b}$ to each degree summand yields two maps

(2.3)
$$\operatorname{hom}^{0}(\mathscr{L},\mathbb{L}) = \bigoplus_{p \in \chi^{0}(L,\mathbb{L})} \left(E|_{p} \right)^{*} \xrightarrow{\Psi^{\mathbb{L}}(\mathscr{L}) := \mathfrak{m}_{1}^{0,b}} \bigoplus_{s \in \chi^{1}(L,\mathbb{L})} \left(E|_{s} \right)^{*} = \operatorname{hom}^{1}(\mathscr{L},\mathbb{L})$$

satisfying

(2.4)
$$\Psi^{\mathbb{L}}(\mathscr{L})\Phi^{\mathbb{L}}(\mathscr{L}) = xyz \cdot \mathrm{id}_{\mathrm{hom}^{0}(\mathscr{L},\mathbb{L})} \quad \text{and} \quad \Phi^{\mathbb{L}}(\mathscr{L})\Psi^{\mathbb{L}}(\mathscr{L}) = xyz \cdot \mathrm{id}_{\mathrm{hom}^{1}(\mathscr{L},\mathbb{L})}.$$

As hom[•] $(\mathcal{L}, \mathbb{L})$ (• $\in \mathbb{Z}_2$) are \mathbb{C} -vector spaces of dimension $\tau \rho$, extension of scalar to the ring $S := \mathbb{C}[[x, y, z]]$ yields two free *S*-modules $S \otimes_{\mathbb{C}}$ hom[•] $(\mathcal{L}, \mathbb{L})$ of rank $\tau \rho$. Now we view each of $\Phi^{\mathbb{L}}(\mathcal{L})$ and $\Psi^{\mathbb{L}}(\mathcal{L})$ as a map between those *S*-modules, regarding *x*, *y* and *z* as variables in the ring *S*. Then the relation (2.4) still holds just by replacing id_{hom}• $(\mathcal{L}, \mathbb{L})$ with id_{S & hom}• $(\mathcal{L}, \mathbb{L})$, so the pair

$$\mathscr{F}^{\mathbb{L}}(\mathscr{L}) := \left(\Phi^{\mathbb{L}}(\mathscr{L}), \Psi^{\mathbb{L}}(\mathscr{L}) \right)$$

produces a $\tau \rho \times \tau \rho$ matrix factorization of *xyz* in *S*.

¹⁰For definition of wrapped Fukaya category W Fuk(Σ), see [AS10] and [Abo12] (also [Aur14] for surfaces). Its objects involve not only loops but also arcs between boundaries, and it contains the compact Fukaya category Fuk(Σ) as a full subcategory.

Theorem 2.3. [CHL17] The localized mirror functor $\mathscr{F}^{\mathbb{L}}$: W Fuk $(\Sigma) \to MF_{A_{\infty}}(xyz)$ is defined as follows:

• For an object $\mathcal{L} = (L, E, \nabla)$ in W Fuk (Σ) , its mirror object in MF_{A₀}(xyz) is given by

9

$$\mathbb{F}^{\mathbb{L}}(\mathscr{L}) = \left(\Phi^{\mathbb{L}}(\mathscr{L}), \Psi^{\mathbb{L}}(\mathscr{L}) \right).$$

• Higher components $\{\mathscr{F}_k^{\mathbb{L}}\}_{k\geq 1}$ are given by

$$\mathcal{F}_{k}^{\mathbb{L}}: \hom\left(\mathcal{L}_{0}, \mathcal{L}_{1}\right) \otimes \cdots \otimes \hom\left(\mathcal{L}_{k-1}, \mathcal{L}_{k}\right) \to \hom_{\mathrm{MF}_{A_{\infty}}(xyz)}\left(\mathcal{F}^{\mathbb{L}}\left(\mathcal{L}_{0}\right), \mathcal{F}^{\mathbb{L}}\left(\mathcal{L}_{k}\right)\right)$$
$$\left(f_{1}, \dots, f_{k}\right) \mapsto \mathfrak{m}_{k+1}^{0, \dots, 0, b}\left(f_{1}, \dots, f_{k}, -\right) := \sum_{i=0}^{\infty} \mathfrak{m}_{k+1+i}(f_{1}, \dots, f_{k}, -, \underbrace{b, \dots, b}_{i}),$$

whose images are \mathbb{C} -linear maps hom $(\mathscr{L}_k, \mathbb{L}) \to \text{hom}(\mathscr{L}_0, \mathbb{L})$, also viewed as module homomorphisms $S \otimes \text{hom}(\mathscr{L}_k, \mathbb{L}) \to S \otimes \text{hom}(\mathscr{L}_0, \mathbb{L})$ over $S = \mathbb{C}[[x, y, z]]$.

Then, $\mathscr{F}^{\mathbb{L}}$ is an A_{∞} -functor. Moreover, it induces an equivalence from the derived wrapped Fukaya category $D^{\pi}W$ Fuk $(\Sigma)^{11}$ to the homotopy category $\underline{MF}[xyz]^{12}$ of matrix factorizations of xyz.

The last statement recovers the result of $[AAE^+13]$ for Σ . It follows from the fact that $\mathscr{F}^{\mathbb{L}}$ sends the three generating arcs L_{xy} , L_{yz} , and L_{zx} of $D^{\pi}W$ Fuk (Σ) in Figure 4 to the three generating matrix factorizations $z \cdot xy$, $x \cdot yz$, and $y \cdot zx$ of $\underline{MF}[xyz]$, respectively. The functor induces an isomorphism on cohomology of hom spaces between those objects and hence we can apply Theorem 4.2 in [CHL17].

2.2. **Computation of localized mirror functor.** Note in (2.2) that hom $(\mathcal{L}, \mathbb{L})$ allows a decomposition into ρ -dimensional vector spaces $(E|_p)^*$ for $p \in \chi(L, \mathbb{L})$. Correspondingly, the operation $\mathfrak{m}_1^{0,b}$ (and hence $\Phi^{\mathbb{L}}(\mathcal{L})$ and $\Psi^{\mathbb{L}}(\mathcal{L})$) is decomposed into several $((E|_s)^*, (E|_p)^*)$ -components ¹³ for $p, s \in \chi(L, \mathbb{L})$.

For $p \in \chi(L, \mathbb{L})$ and $f = f|_p \in (E|_p)^* \subseteq \text{hom}(\mathcal{L}, \mathbb{L})$, we have

$$\mathfrak{m}_{1}^{0,b}(f) = \sum_{i=0}^{\infty} \mathfrak{m}_{1+i}(f, \underbrace{b, \dots, b}_{i}) \qquad (b = xX + yY + zX \in \hom^{1}(\mathbb{L}, \mathbb{L}))$$
$$= \sum_{\substack{(x_{1}, X_{1}), \dots, (x_{i}, X_{i})\\ \in \{(x, X), (y, Y), (z, Z)\}}} x_{1} \dots x_{i} \mathfrak{m}_{1+i}(f, X_{1}, \dots, X_{i}),$$

and

$$\mathfrak{m}_{1+i}\left(f, X_1, \dots, X_i\right) = \sum_{s \in \chi(L, \mathbb{L})} \sum_{u \in \mathcal{M}\left(p, X_1, \dots, X_i, \overline{s}\right)} \operatorname{sign}(u) \operatorname{hol}_s\left(\partial u\right)\left(f, X_1, \dots, X_i\right).$$

Therefore, the $(E|_s)^*$ -component of $\mathfrak{m}_1^{0,b}(f)$ is

$$\sum_{\substack{(x_1,X_1),\dots,(x_i,X_i)\\ \varepsilon\{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p,X_1,\dots,X_i,\overline{s})} \operatorname{sign}(u) \operatorname{hol}_s(\partial u) (f, X_1,\dots,X_i).$$

So it comes from immersed polygons (also called *deformed strips*) u bounded by L and \mathbb{L} , whose angles consist of p, X_1, \ldots, X_i and \overline{s} in a counterclockwise order, for some $X_1, \ldots, X_i \in \{X, Y, Z\}$. (See Figure 5.) Such a deformed strip contributes a monomial $x_1 \cdots x_i$, where x_j is x, y or z depending on whether X_j is X, Y or Z.

¹¹It is obtained from W Fuk (Σ) by taking twisted completion (Definition 5.16), idempotent completion and then the cohomological category (Definition A.3). For more details, we refer to [Sei08].

¹²We can take the target of the functor as $MF_{A_{\infty}}[xyz]$ (see Remark 1.5) since every object in W Fuk (Σ) is quasi-isomorphic to an object whose image lies in $MF_{A_{\infty}}[xyz]$. Then MF[xyz] is the same as its cohomological category $H^0(MF_{A_{\infty}}[xyz])$.

¹³Following the standard natation in matrices, $(E|_s)^*$ and $(E|_p)^*$ are subspaces of the codomain and domain, respectively.



Figure 5. A deformed strip which contributes a monomial $x_1 \cdots x_i$ to the $(E|_s)^*$ -component of $\mathfrak{m}_1^{0,b} (f|_n)$

The sign of the deformed strip *u* is given by

sign (u) =
$$\begin{cases} 1 & \text{if orientation of } \mathbb{L} = \text{orientation of } \partial u, \\ (-1)^{i+1} & \text{otherwise} \end{cases} = (-1)^{(i+1)} \mathbb{I}_{\mathbb{Q}(\mathbb{L}) \neq \mathbf{Q}(\partial u)}.$$

It follows from formula (A.6) and the fact that the orientation of \mathbb{L} along any angle X_j is preserved (because all X_j 's have odd-degrees) and the degrees of p and s are always different.

The holonomy operation of ∂u at *s* is by definition given as

$$\operatorname{hol}_{s}(\partial u) : \operatorname{Hom}_{\mathbb{C}}\left(E_{|_{p}}, E_{\mathbb{L}}_{|_{p}}\right) \otimes \operatorname{Hom}_{\mathbb{C}}\left(E_{\mathbb{L}}_{|_{X_{1}}}, E_{\mathbb{L}}_{|_{X_{1}}}\right) \otimes \cdots \otimes \operatorname{Hom}_{\mathbb{C}}\left(E_{\mathbb{L}}_{|_{X_{i}}}, E_{\mathbb{L}}_{|_{X_{i}}}\right) \to \operatorname{Hom}_{\mathbb{C}}\left(E_{|_{s}}, E_{\mathbb{L}}_{|_{s}}\right), \\ \left(f, X_{1}, \dots, X_{i}\right) \mapsto P\left((\partial u)_{i+1}\right) \circ X_{i} \circ P\left((\partial u)_{i}\right) \circ X_{i-1} \circ \cdots \circ X_{1} \circ P\left((\partial u)_{1}\right) \circ f \circ P\left((\partial u)_{0}\right).$$

Under the identification $\operatorname{Hom}_{\mathbb{C}}\left(E_{\mathbb{L}}|_{X_{j}}, E_{\mathbb{L}}|_{X_{j}}\right) = \operatorname{Span}_{\mathbb{C}}\left\{X_{j}\right\} \cong \mathbb{C}$ $(j \in \{1, ..., i\})$, the generator X_{j} corresponds to 1. Note that $(\partial u)_{j}$ lies on \mathbb{L} for $j \in \{1, ..., i+1\}$, and under the identification $\operatorname{Hom}_{\mathbb{C}}\left(E_{\mathbb{L}}|_{X_{j-1}}, E_{\mathbb{L}}|_{X_{j}}\right) \cong \mathbb{C}$ $(j \in \{1, ..., i+1\}, X_{0} := p, X_{i+1} := \overline{s})$, in view of Proposition A.7, $P\left((\partial u)_{j}\right)$ corresponds $(-1)^{\#\left((\partial u)_{j}\cap \bigstar_{\mathbb{L}}\right)}$, where $\#\left((\partial u)_{j} \cap \bigstar_{\mathbb{L}}\right)$ is the number of times $(\partial u)_{j}$ passes through the point \bigstar on \mathbb{L} . Therefore, we can replace $\operatorname{hol}_{s}(\partial u)\left(f, X_{1}, ..., X_{i}\right)$ with

$$(-1)^{\#(\partial u \cap \bigstar_{\mathbb{L}})} f \circ P((\partial u)_0),$$

where $\#(\partial u \cap \bigstar_{\mathbb{L}})$ is the total number of times ∂u passes through the point \bigstar on \mathbb{L} , and $P((\partial u)_0) \in \text{Hom}_{\mathbb{C}}(E|_s, E|_p)$ is the parallel transport from $E|_s$ to $E|_p$ along the side of u lying in L.

We summarize the above discussion into the following formula:

Proposition 2.4. The $((E|_s)^*, (E|_p)^*)$ -component of $\mathfrak{m}_1^{0,b}$: hom $(\mathcal{L}, \mathbb{L}) \to \text{hom}(\mathcal{L}, \mathbb{L})$ (and hence of $\Phi^{\mathbb{L}}(\mathcal{L})$) or $\Psi^{\mathbb{L}}(\mathcal{L})$) is the \mathbb{C} -linear map $\mathfrak{m}_1^{0,b}: (E|_p)^* \to (E|_s)^*$ that maps $f \in (E|_p)^*$ to

(2.5)
$$\sum_{\substack{(x_1,X_1),\ldots,(x_i,X_i)\\ \in\{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p,X_1,\ldots,X_i,\overline{s})} (-1)^{(i+1)\mathbb{I}_{o(\mathbb{L})\neq o(\partial u)} + \#(\partial u \cap \bigstar_{\mathbb{L}})} f \circ P((\partial u)_0).$$

It is also an S-module map $S \otimes (E|_p)^* \to S \otimes (E|_s)^*$, by considering x, y and z as variables in $S = \mathbb{C}[[x, y, z]]$.

If we choose a point \bigstar on L so that $\operatorname{hol}_{\bigstar}(E)$ is represented by a matrix $H \in \operatorname{GL}_{\rho}(\mathbb{C})$, trivialize $E|_{S^1 \setminus \bigstar} \cong (S^1 \setminus \bigstar) \times \mathbb{C}^{\rho}$ as in Proposition A.7 and thus $(E|_p)^* \cong (E|_s)^* \cong \mathbb{C}^{\rho}$, then it is represented by the $\rho \times \rho$ matrix

(2.6)
$$\sum_{\substack{(x_1, x_1), \dots, (x_i, X_i) \\ \in \{(x, X), (y, Y), (z, Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p, X_1, \dots, X_i, \overline{s})} (-1)^{(i+1)\mathbb{I}_{o(\mathbb{L}) \neq o(\partial u)} + \#(\partial u \cap \star_{\mathbb{L}})} (H^T)^{\#(\partial u \cap \star_{\mathbb{L}})(\mathbb{I}_{o(L) = o(\partial u)} - \mathbb{I}_{o(L) \neq o(\partial u)})},$$

where $\#(\partial u \cap \bigstar_L)$ is the number of times ∂u passes through the point \bigstar on L. Note that the entries can be also viewed as elements in $\mathbb{C}[[x, y, z]]$.

2.3. **Illustration with an example.** In this section, we illustrate the computation of the localized mirror functor with a concrete example. Namely, we will find the mirror matrix factorization of the loop with a local system $\mathcal{L} := \mathcal{L}((3, -2, 2), \eta, 1)^{14}$ for any $\eta \in \mathbb{C}^{\times}$. It is an object (L, E, ∇) of Fuk (Σ) that consists of the underlying loop L := L(3, -2, 2) described in Figure 6, a trivial line bundle *E* over the domain S^1 of *L*, and a flat connection ∇ on *E* whose holonomy is η at the point \bigstar on *L* marked in Figure 6.

Note that L and L have 6 intersections, say p, q, r, and s, t, u. According to their degrees, we have

$$\chi^0(L,\mathbb{L}) = \left\{ p, q, r \right\}, \quad \chi^1(L,\mathbb{L}) = \left\{ s, t, u \right\}.$$

We trivialize *E* over $S^1 \setminus \bigstar$ as in Proposition A.7, which yields identifications $E|_{\bullet} \cong \mathbb{C}$ for each $\bullet \in \chi(L, \mathbb{L})$. Then we have Hom_{\mathbb{C}} ($E|_{\bullet}, E_{\mathbb{L}}|_{\bullet}$) \cong Span_{\mathbb{C}} { \bullet } as noted in Remark A.8, and hence

 $\hom^{0}(\mathcal{L}, \mathbb{L}) = \operatorname{Span}_{\mathbb{C}} \{ p, q, r \}, \quad \hom^{1}(\mathcal{L}, \mathbb{L}) = \operatorname{Span}_{\mathbb{C}} \{ s, t, u \}.$

Then two restricted operations $\mathfrak{m}_{1}^{0,b}$: Hom⁰ (\mathscr{L}, \mathbb{L}) \rightarrow Hom¹ (\mathscr{L}, \mathbb{L}) and $\mathfrak{m}_{1}^{0,b}$: Hom¹ (\mathscr{L}, \mathbb{L}) \rightarrow Hom⁰ (\mathscr{L}, \mathbb{L}) with respect to ordered bases {p, q, r} and {s, t, u} yield two 3 × 3 matrices $\Phi^{\mathbb{L}}(\mathscr{L})$ and $\Psi^{\mathbb{L}}(\mathscr{L})$, respectively.



Figure 6. Matrix factorization of *xyz* corresponding to $\mathcal{L} = \mathcal{L}((3, -2, 2), \eta, 1)$

Entries come from the following relations:

$$\begin{cases} \mathfrak{m}_{1}^{0,b}(p) = zs + y^{2}t - \eta x^{2}u \\ \mathfrak{m}_{1}^{0,b}(q) = x t \\ \mathfrak{m}_{1}^{0,b}(r) = -z t + y u \end{cases} \text{ and } \begin{cases} \mathfrak{m}_{1}^{0,b}(s) = xyp + (-y^{3} + \eta zx^{2})q + \eta x^{3}u \\ \mathfrak{m}_{1}^{0,b}(t) = yz q \\ \mathfrak{m}_{1}^{0,b}(u) = z^{2}q + zxu \end{cases}$$

Note that the (\blacksquare, \bullet) -entry of $\Phi^{\mathbb{L}}(\mathcal{L})$ or $\Psi^{\mathbb{L}}(\mathcal{L})$ is the coefficient of \blacksquare in $\mathfrak{m}_1^{0,b}(\bullet)$ for $\bullet, \blacksquare \in \{p, q, r, s, t, u\}$. So it can be computed from the formula (2.6), which in this case is just a single entry (in $S = \mathbb{C}[[x, y, z]]$)

(2.7)
$$\sum_{\substack{(x_1,X_1),\dots,(x_i,X_i)\\ \in \{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(\bullet,X_1,\dots,X_i,\bar{\bullet})} (-1)^{(i+1)\mathbb{I}_{o(\mathbb{L})\neq o(\partial u)} + \#(\partial u \cap \star_{\mathbb{L}})} \eta^{\#(\partial u \cap \star_{\mathbb{L}})} (\mathbb{I}_{o(\mathbb{L})=o(\partial u)} - \mathbb{I}_{o(\mathbb{L})\neq o(\partial u)})$$

That is, each deformed strip u bounded by L and \mathbb{L} , whose angles consist of •, X_1, \ldots, X_i and $\overline{\blacksquare}$ in a counterclockwise order, contributes a monomial $x_1 \cdots x_i$. Its coefficient is determined by coincidence of orientations of L, \mathbb{L} with boundary orientation of ∂u and the number of times ∂u passes through the points \bigstar_L or \bigstar_L . Figure 7 shows some deformed strips that contribute some of the entries above:

¹⁴This is the canonical form of loops with a local system corresponding to a loop datum $((3, -2, 2), \eta, 1)$. See Definition 3.9.



Figure 7. Some entries of $\Phi^{\mathbb{L}}(L)$ and $\Psi^{\mathbb{L}}(L)$

2.4. **Higher rank computation.** We can compute the matrix factorizations corresponding to a higher rank local system on a loop easily once we know the result for rank 1 local systems on the same loop.

Consider two local systems on the same loop *L* in Σ that have rank 1 and ρ , respectively:

- $\mathscr{L}_1(\eta) := (L, E_1, \nabla_1) :$ loop with a local system of rank 1 of holonomy $\eta \in \mathbb{C}^{\times}$.
- $\mathscr{L}_{\rho}(H) := (L, E_{\rho}, \nabla_{\rho})$: loop with a local system of rank ρ of holonomy $H \in GL_{\rho}(\mathbb{C})$,

Namely, there is a point \bigstar on *L* and identification $E_1|_{\bigstar} \cong \mathbb{C}$ and $E_{\rho}|_{\bigstar} \cong \mathbb{C}^{\rho}$, so that hol $_{\bigstar}(E_1)$ is represented by a scalar $\eta \in \mathbb{C}^{\times}$, while hol $_{\bigstar}(E_{\rho})$ is represented by a matrix $H \in GL_{\rho}(\mathbb{C})$.

For any $p, s \in \chi(L, \mathbb{L})$, the formula (2.7) and (2.6) imply that the (s, p)-entry of $\left(\Phi^{\mathbb{L}}\left(\mathscr{L}_{1}(\eta)\right), \Psi^{\mathbb{L}}\left(\mathscr{L}_{1}(\eta)\right)\right)$ and the $\left(\left(E_{\rho}\right|_{s}\right)^{*}, \left(E_{\rho}\right|_{p}\right)^{*}$ -component of $\left(\Phi^{\mathbb{L}}\left(\mathscr{L}_{\rho}(H)\right), \Psi^{\mathbb{L}}\left(\mathscr{L}_{\rho}(H)\right)\right)$ are given in the form

(2.8)
$$\sum_{k=-\infty}^{\infty} a_k \eta^k \in \mathbb{C}[[x, y, z]] \text{ and } \sum_{k=-\infty}^{\infty} a_k (H^T)^k \in \mathbb{C}[[x, y, z]]^{\rho \times \rho},$$

respectively, for some $a_k \in \mathbb{C}[[x, y, z]]$. (See Theorem 3.13 for the convergence issue).

Proposition 2.5 ((η, H^T) -substitution). The matrix factorizations corresponding to $\mathscr{L}_1(\eta)$ and $\mathscr{L}_{\rho}(H)$ are given in the form

$$\mathscr{F}^{\mathbb{L}}(\mathscr{L}_{1}(\eta)) = \left(\sum_{k=-\infty}^{\infty} \varphi_{k} \eta^{k}, \sum_{k=-\infty}^{\infty} \psi_{k} \eta^{k}\right) \quad and \quad \mathscr{F}^{\mathbb{L}}(\mathscr{L}_{\rho}(H)) = \left(\sum_{k=-\infty}^{\infty} \varphi_{k} \otimes \left(H^{T}\right)^{k}, \sum_{k=-\infty}^{\infty} \psi_{k} \otimes \left(H^{T}\right)^{k}\right)^{15},$$

respectively, for some $\varphi_k, \psi_k \in \mathbb{C}[[x, y, z]]^{\tau \times \tau} (\tau = \frac{1}{2} |L \cap \mathbb{L}|).$

Proof. Let φ_k be a $\tau \times \tau$ matrix over $S = \mathbb{C}[[x, y, z]]$ whose (s, p)-entry is a_k given in (2.8) (viewed as a map of free *S*-modules from $\text{Span}_S(\chi^0(L, \mathbb{L}))$ to $\text{Span}_S(\chi^1(L, \mathbb{L}))$ with respect to an appropriate order in each set $\chi^{\bullet}(L, \mathbb{L})$). Then $\sum_{k=-\infty}^{\infty} \varphi_k \eta^k$ coincides with $\Phi^{\mathbb{L}}(\mathscr{L}_1(\eta))$.

Now $\varphi_k \otimes (H^T)^k$ is a $\tau \rho \times \tau \rho$ matrix over *S*, which can be also viewed as a map of free *S*-modules

$$\bigoplus_{p \in \chi^{0}(L,\mathbb{L})} \operatorname{Span}_{S} \{p\} \otimes_{\mathbb{C}} \left(E_{\rho} \big|_{p} \right)^{*} \to \bigoplus_{s \in \chi^{1}(L,\mathbb{L})} \operatorname{Span}_{S} \{s\} \otimes_{\mathbb{C}} \left(E_{\rho} \big|_{s} \right)^{*}$$

under an order in $\chi^{\bullet}(L, \mathbb{L})$ and identification $E_{\rho}|_{p} \cong E_{\rho}|_{s} \cong \mathbb{C}^{\rho}$. Its $((E_{\rho}|_{s})^{*}, (E_{\rho}|_{p})^{*})$ -component is $a_{k}(H^{T})^{k}$, where a_{k} denotes the (s, p)-entry of φ_{k} . Therefore, the corresponding component of $\sum_{k=-\infty}^{\infty} \varphi_{k} \otimes (H^{T})^{k}$ is $\sum_{k=-\infty}^{\infty} a_{k}(H^{T})^{k}$, which is that of $\Phi^{\mathbb{L}}(\mathscr{L}_{\rho}(H))$ by (2.8). It works the same for ψ_{k} instead of φ_{k} .

The proposition says that if we know φ_k and ψ_k 's from the rank 1 cases, we immediately get the result for higher rank cases as well, just by '**substituting the matrix** H^T for the scalar η '. See the example:

Example 2.6. Consider the loop with a local system $\mathscr{L}_{\rho} := \mathscr{L}((3, -2, 2), \eta, \rho)$ of rank $\rho \in \mathbb{Z}_{\geq 1}$. It consists of the same underlying loop L := L(3, -2, 2) as in Subsection 2.3 (Figure 6), trivial vector bundle E_{ρ} of rank ρ over the domain S^1 of L, and a flat connection ∇_{ρ} on E_{ρ} whose holonomy is $J_{\rho}(\eta)$ at the point \bigstar . Then, by the result for \mathscr{L}_1 in Subsection 2.3 and Proposition 2.5, the corresponding matrix factorization is

$$\Phi^{\mathbb{L}}\left(\mathscr{L}_{\rho}\right)^{*} \begin{pmatrix} \left(E|_{q}\right)^{*} & \left(E|_{r}\right)^{*} \\ \left(E|_{q}\right)^{*} & \left(E|_{r}\right)^{*} \\ zI_{\rho} & 0 & 0 \\ y^{2}I_{\rho} & xI_{\rho} & -zI_{\rho} \\ -x^{2}J_{\rho}\left(\eta\right)^{T} & 0 & yI \end{pmatrix} and \Psi^{\mathbb{L}}\left(\mathscr{L}_{\rho}\right)^{*} \begin{pmatrix} \left(E|_{q}\right)^{*} \\ \left(E|_{q}\right)^{*} \\ \left(E|_{q}\right)^{*} \\ \left(E|_{r}\right)^{*} \end{pmatrix}^{*} \begin{pmatrix} \left(E|_{s}\right)^{*} & \left(E|_{t}\right)^{*} \\ xyI_{\rho} & 0 & 0 \\ -y^{3}I_{\rho} + zx^{2}J_{\rho}\left(\eta\right)^{T} & yzI_{\rho} & z^{2}I_{\rho} \\ x^{3}J_{\rho}\left(\eta\right)^{T} & 0 & zxI_{\rho} \end{pmatrix}$$

 $^{^{15}}$ Here \otimes for matrices refers to the *Kronecker product*.

3. MATRIX FACTORIZATIONS FROM LOOPS WITH A LOCAL SYSTEM

In this section, we first define the canonical form of immersed loops in Σ and local systems on them, parameterized by *normal loop words* (§3.1) and *loop data* (§3.2), respectively. Then we compute their mirror image under the localized mirror functor, which provides the canonical form of matrix factorizations of *xyz* (§3.3). The matrix factorizations corresponding to periodic loop words are decomposable, which implies that *non-primitive* loops (with a local system) are decomposable in the Fukaya category (§3.4). Meanwhile, an exceptional case is handled separately (§3.5).

3.1. Loop words and canonical form of immersed loops. We first recall from [CJKR22] the concept of *loop words*, which parameterize free homotopy classes of loops in Σ . They were introduced in order to pick a specific representative in each (hyperbolic) free homotopy class, motivated by the observation that two freely homotopic loops correspond to homotopically equivalent matrix factorizations under the localized mirror functor (Theorem 3.13).

Note that the fundamental group of Σ can be presented as $\pi_1(\Sigma) = \langle \alpha, \beta, \gamma | \alpha \beta \gamma = 1 \rangle$ with the based loops α , β and γ in Σ shown in Figure 8a. Also recall that there is a one-to-one correspondence between the free homotopy classes of loops in Σ and the conjugacy classes in $\pi_1(\Sigma)$.



Figure 8. Fundamental group and loop data

Definition 3.1. A loop word of length 3τ is

$$w' = (l'_1, m'_1, n'_1, l'_2, m'_2, n'_2, \dots, l'_{\tau}, m'_{\tau}, n'_{\tau}) \in \mathbb{Z}^{3\tau}$$

 $(\tau \in \mathbb{Z}_{\geq 1})$. The associated loop, denoted as

L(w'),

is illustrated in Figure 8b. It visits three holes A, B, and C in turn, winding them around the number of times specified in w'. Namely, starting from the point \bigstar marked in the figure, it winds hole $A l'_1$ -times, hole $B m'_1$ -times, hole $C n'_1$ -times, hole $A l'_2$ -times, hole $B m'_2$ -times, and so on. After finally it winds hole $C n'_{\tau}$ -times, it returns to the point \bigstar to form a closed loop. We perturb it if it is necessary to put them together into a transversal set.

Note that its free homotopy class in $[S^1, \Sigma] = \pi_1(\Sigma) / \sim_{\text{conjugation}} is$

$$[L(w')] = \left[\alpha^{l'_1}\beta^{m'_1}\gamma^{n'_1}\alpha^{l'_2}\beta^{m'_2}\gamma^{n'_2}\cdots\alpha^{l'_r}\beta^{m'_r}\gamma^{n'_r}\right].$$

Two loop words w' and \tilde{w}' are regarded as equivalent if $[L(w')] = [L(\tilde{w}')]$.

Example 3.2. The loop described in Figure 21 and Figure 6 is (a perturbation of) L(3, -2, 2).

We denote the *j*-th value of a loop word w' as w'_{j} so that

$$w' = (w'_1, w'_2, w'_3, \dots, w'_{3\tau-1}, w'_{3\tau}) \in \mathbb{Z}^{3\tau}.$$

Then any tuple $(w'_k, w'_{k+1}, ..., w'_l)$ for some distinct $k, l \in \mathbb{Z}_{3\tau}$ is called a *subword* in w'. We regard the index i of l'_i, m'_i and n'_i to be in \mathbb{Z}_{τ} (hence $3i \in \mathbb{Z}_{3\tau}$) and the index j of w'_j to be in $\mathbb{Z}_{3\tau}$. Therefore, for example, $(w'_{3\tau-1}, w'_{3\tau}, w'_1)$ is a subword. We define the 1-*shift* of a loop word w' to be

$$w'^{(1)} = (l'_2, m'_2, n'_2, \dots, l'_{\tau}, m'_{\tau}, n'_{\tau}, l'_1, m'_1, n'_1) \in \mathbb{Z}^{3n}$$

and *k*-shift to be $w'^{(k)}$ which is obtained from w' by applying the 1-shift *k*-times ($k \in \mathbb{Z}$).

For a loop word w', we define its *N*-concatenation $(w')^N$ as the *N* repetitions of w'. It is called *periodic* if it is *N*-concatenation of another loop word for some $N \in \mathbb{Z}_{\geq 2}$. For an immersed loop $L: S^1 \to \Sigma$, we define its *N*-concatenation by the immersed loop

$$L^N: S^1 \to \Sigma, \quad e^{2\pi i t} \mapsto L\left(e^{2N\pi i t}\right)$$

A loop *L* or its free homotopy class [*L*] are called *non-primitive* if *L* is freely homotopic to an *N*-concatenation of another loop for some $N \in \mathbb{Z}_{\geq 2}$. Otherwise, they are called *primitive*.

Note that if a loop word w' is periodic, then the associated loop L(w') and its free homotopy class [L(w')] are non-primitive. But the converse is not true in general as a non-periodic loop word w' can be equivalent to a periodic one. It will be fixed when we will regard only *normal* loop words (Corollary 3.7).

The following lemma is easy to check.

Lemma 3.3. The following operations on a loop word w' do not change the equivalence class of w':

- (inserting 0s) insert the subword (0,0,0) somewhere in w',
- (removing 0s) remove a subword (0,0,0) in w' if it exists,
- (adding 1s around 0) add (1,1,1) to the subword $(w'_{i-1}, 0, w'_{i+1})$ in w' where $w'_i = 0$, and
- (subtracting 1s around 1) subtract (1,1,1) from the subword $(w'_{i-1}, 1, w'_{i+1})$ in w' where $w'_i = 1$,
- (shifting) take k-shift of w' for some $k \in \mathbb{Z}$.

The converse statement is also true, but its proof involves a non-trivial word problem.

Proposition 3.4. [CJKR22] Two loop words w' and \tilde{w}' are equivalent if and only if \tilde{w}' can be obtained from w' by performing the above five operations finitely many times.

Note that several equivalent loop words can represent the same free homotopy class. To find a unique representative in each class, we introduce the following **normal form** of loop words. It will play an important role in the conversion formula between loop data and band data.

Definition 3.5. A loop word w' is said to be **normal** if it satisfies the following conditions:

- any subword of the form (a, 1, b) in w' satisfies $a, b \le 0$,
- any subword of the form (a, 0, b) in w' satisfies $a \le -1$, $b \ge 1$ or $a \ge 1$, $b \le -1$ or $a, b \ge 1$,
- w' has no subword of the form (0, -1, -1, ..., -1, 0), and
- w' does not consist only of -1, that is, $w' \neq (-1, -1, \dots, -1)$.

According to Definition 3.1, each elliptic or parabolic loop is produced by a loop word equivalent to one of (l', 0, 0), (0, m', 0) or (0, 0, n') for some $l', m', n' \in \mathbb{Z}$. Loop words in those forms are called *non-hyperbolic*, while the others are called *hyperbolic*. Therefore, hyperbolic loop words produce hyperbolic loops. Interestingly, the above normality condition automatically rules out non-hyperbolic loop words. Moreover, the normal form up to shifting gives exactly one representative among equivalent hyperbolic loop words.

Proposition 3.6. [CJKR22] Any normal loop word is hyperbolic. Conversely, any hyperbolic loop word is equivalent to a unique normal loop word up to shifting.

This also implies that two normal loop words w' and \tilde{w}' are equivalent if and only if they coincide up to shifting, that is, $\tilde{w}' = w'^{(k)}$ for some $k \in \mathbb{Z}$.

Corollary 3.7. A normal loop word w' is periodic if and only if the associated loop L(w') and its free homotopy class [L(w')] are non-primitive.

Now we give Σ a hyperbolic metric with three cusps. It can be achieved by considering the Poincaré disk as the universal cover of Σ as shown in Figure 9. In fact, such a metric is unique up to isometry (Theorem 9.8.8 in [RAR94]). It is well-known in hyperbolic geometry that there is exactly one (immersed) closed geodesic in each primitive hyperbolic free homotopy class of loops in Σ (Theorem 9.6.4 in [RAR94]). This provides another description of normal loop words.

Proposition 3.8. There is a one-to-one correspondence

 $\{\text{closed geodesics in } \Sigma\} \ \stackrel{1:1}{\leftrightarrow} \ \{\text{non-periodic normal loop words}\}/\sim_{\text{shifting}}.$



Figure 9. Fundamental domain of Σ in its universal cover (Poincaré disk)

¹⁶The terminologies come from hyperbolic geometry. In fact, if we assign a hyperbolic metric to the surface, the elliptic, parabolic, and hyperbolic loops correspond to concepts already in use in hyperbolic geometry. We refer readers to Section 9.6 in [RAR94].

3.2. Loop data and canonical form of loops with a local system.

Definition 3.9. A loop datum (w', η, ρ) consists of the following:

- (normal loop word) $w' = (l'_1, m'_1, n'_1, l'_2, m'_2, n'_2, \dots, l'_{\tau}, m'_{\tau}, n'_{\tau}) \in \mathbb{Z}^{3\tau}$ for some $\tau \in \mathbb{Z}_{\geq 1}$,
- (holonomy parameter) $\eta \in \mathbb{C}^{\times}$,
- ((geometric) rank) $\rho \in \mathbb{Z}_{\geq 1}$.

It represents an object (L, E, ∇) *of* Fuk (Σ) *, denoted as*

$$\mathscr{L}(w',\eta,\rho),$$

that consists of the loop L := L(w') defined in Definition 3.1, a trivial \mathbb{C} -vector bundle E of rank ρ over the domain S^1 of L, and a flat connection ∇ on E whose holonomy is $J_{\rho}(\eta)$ at the point \bigstar on L (marked in Figure 8b). We refer to it as the **canonical form of loops with a local system** corresponding to the loop datum (w', η, ρ) .

Proposition 3.10. Let (L, E, ∇) be a loop with a local system with L = L(w') for some normal loop word w'. Then there are finitely many pairs $(\eta_1, \rho_1), \ldots, (\eta_d, \rho_d) \in \mathbb{C}^{\times} \times \mathbb{Z}_{\geq 1}$ ($d \in \mathbb{Z}_{\geq 1}$) such that (E, ∇) is gauge equivalent to the direct sum

$$(E_1, \nabla_1) \oplus \cdots \oplus (E_d, \nabla_d)$$

where (L, E_i, ∇_i) $(i \in \{1, ..., d\})$ is the canonical form $\mathcal{L}(w', \eta_i, \rho_i)$ corresponding to the loop datum (w', η_i, ρ_i) . Moreover, the choice of the pairs is unique up to the order.

Proof. Note that every \mathbb{C} -vector bundle over S^1 is trivial. At a fixed point \bigstar on L, the holonomy hol $\bigstar(E)$ of (E, ∇) along L at \bigstar is represented by some matrix $H \in \operatorname{GL}_{\rho}(\mathbb{C})$. Then the Jordan canonical form of H has Jordan blocks $J_{\rho_1}(\eta_1), \ldots, J_{\rho_d}(\eta_d)$ $(d \in \mathbb{Z}_{\geq 1})$ for some pairs $(\eta_1, \rho_1), \ldots, (\eta_d, \rho_d) \in \mathbb{C}^{\times} \times \mathbb{Z}_{\geq 1}$, which yields the desired decomposition.

This shows that any indecomposable local system on a fixed loop L = L(w') is gauge equivalent to a unique canonical form $\mathscr{L}(w', \eta, \rho)$. Combining this with Proposition 3.8, we get the following:

Corollary 3.11. There is a one-to-one correspondence

{closed geodesics in Σ with an indecomposable local system}/ ~_{gauge equivalence}

 $\stackrel{1:1}{\leftrightarrow} \quad \{\text{non-periodic loop data}\} / \sim_{\text{shifting}} {}^{17}.$

3.3. Matrix factorizations from canonical form of loops with a local system. To get a matrix factorization defined over \mathbb{C} from the loop with a local system $\mathscr{L}(w', \eta, \rho)$ constructed in Definition 3.9, we first need to ensure that the underlying loop L(w') doesn't bound an immersed cylinder with the reference loop \mathbb{L} (Seidel Lagrangian). We briefly summarize the discussion in [CJKR22] here:

Definition 3.12. A loop L in Σ is said to be **cylinder-free** with \mathbb{L} if there is no immersion $j: S^1 \times [0,1] \to \Sigma$ that satisfies $j(e^{2\pi i t}, 0) = L(\iota(t))$ and $j(e^{2\pi i t}, 1) = \mathbb{L}(j(t))$ for some immersions $\iota, j: S^1 \to S^1$.

Theorem 3.13. [CJKR22] For an object $\mathcal{L} := (L, E, \nabla)$ in Fuk (Σ) whose underlying loop L is cylinder-free with \mathbb{L} , its mirror matrix factorization $\mathscr{F}^{\mathbb{L}}(\mathscr{L}) = (\Phi^{\mathbb{L}}(\mathscr{L}), \Psi^{\mathbb{L}}(\mathscr{L}))$ is well-defined over \mathbb{C} . More precisely, this means that the moduli spaces involved in formula 2.5 are finite.

Moreover, the homotopy class of $\mathscr{F}^{\mathbb{L}}(\mathscr{L})$ is invariant under free homotopy of the underlying curve L and gauge equivalence of the flat vector bundle $(E, \nabla)^{18}$.

Proposition 3.14. [CJKR22] For a normal loop word w' other than of the form $(2,2,2)^{\tau}$, the corresponding loop L(w') is cylinder-free with \mathbb{L} .

¹⁷We say that a loop datum is *periodic* if its normal loop word is periodic, and define a *shift* of a loop datum as the shifting of its normal loop word.

¹⁸See also [CHL19, Proposition 5.4] for its invariance under Hamiltonian isotopy of *L*.

Now we compute the matrix factorization corresponding to the canonical form $(L, E, \nabla) := \mathcal{L}(w', \eta, \rho)$ for a loop datum (w', η, ρ) with $w' \neq (2, 2, 2)^{\tau}$. For a loop word of length 3τ

$$w' = (l'_1, m'_1, n'_1, l'_2, m'_2, n'_2, \dots, l'_{\tau}, m'_{\tau}, n'_{\tau}),$$

the corresponding loop L = L(w') has 6τ intersections with \mathbb{L} . We name even-degree angles from *L* to \mathbb{L} by $p_1, q_1, r_1, \ldots, p_\tau, q_\tau, r_\tau \in \hom^0(L, \mathbb{L})$ and odd-degree angles from *L* to \mathbb{L} by $s_1, t_1, u_1, \ldots, s_\tau, t_\tau, u_\tau \in \hom^1(L, \mathbb{L})$ in the order along the orientation of *L*.



Figure 10. Canonical form $(L, E, \nabla) = \mathcal{L}(w', \eta, \rho)$ and Seidel Lagrangian \mathbb{L}

Proposition 3.15. For a loop datum (w', η, ρ) with $w' \neq (2, 2, 2)^{\tau}$ ¹⁹, the corresponding loop with a local system $(L, E, \nabla) := \mathscr{L}(w', \eta, \rho)$ is converted under the localized mirror functor into the matrix factor $\Phi^{\mathbb{L}}(\mathscr{L}(w', \eta, \rho))$ given by

where x^a , y^a , z^a are regarded as 0 if a < 0.

Proof. For $\rho = 1$ case, the computation is essentially the same as what we did for $\Phi^{\mathbb{L}}(\mathscr{L}((3, -2, 2), \eta, 1))$ in Subsection 2.3. A rigorous proof can be found in [CJKR22]. For $\rho \ge 2$, we use Proposition 2.5 to 'substitute the matrix $J_{\rho}(\eta)^{T}$ for the scalar η ', as we did in Example 2.6.

¹⁹For the case $w' = (2,2,2)^{T}$, we still get the same form even if we use the loop constructed in Definition 3.1, which is not cylinder-free with \mathbb{L} . For $\eta = -1$, however, it fails to be a matrix factor of xyz as its determinant is zero. For $\eta \neq -1$, it is a valid matrix factor of xyz but the opposite matrix factor $\Psi^{\mathbb{L}}(\mathscr{L}(w',\eta,\rho))$ cannot be directly obtained by counting polygons between L and \mathbb{L} , because it involves some moduli spaces having infinitely many elements. To justify it, we will need to develop additional explanation but we won't to do in that way here. We will rather treat them separately in Subsection 3.5.

Inspired by this observation, we propose the following definition:

Definition 3.16. For a loop datum (w', λ, ρ) with $(w', \lambda) \neq ((2, 2, 2)^{\tau}, 1)$, consider the matrix

$$\varphi(w',\lambda,\rho) \coloneqq \begin{pmatrix} zI_{\rho} & -y^{m_{1}'-1}I_{\rho} & 0 & 0 & \cdots & 0 & -x^{-l_{1}'}J_{\rho}(\lambda)^{-1} \\ -y^{-m_{1}'}I_{\rho} & xI_{\rho} & -z^{n_{1}'-1}I_{\rho} & 0 & \cdots & 0 & 0 \\ 0 & -z^{-n_{1}'}I_{\rho} & yI_{\rho} & -x^{l_{2}'-1}I_{\rho} & \cdots & 0 & 0 \\ 0 & 0 & -x^{-l_{2}'}I_{\rho} & zI_{\rho} & \ddots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & -y^{m_{\tau}'-1}I_{\rho} & 0 \\ 0 & 0 & 0 & \cdots & -y^{-m_{\tau}'}I_{\rho} & xI_{\rho} & -z^{n_{\tau}'-1}I_{\rho} \\ -x^{l_{1}'-1}J_{\rho}(\lambda) & 0 & 0 & \cdots & 0 & -z^{-n_{\tau}'}I_{\rho} & yI_{\rho} \end{pmatrix}_{3\tau\rho\times3\tau\rho}$$

where x^a , y^a , z^a are regarded as 0 if a < 0. Then it is a matrix factor of xyz with the opposite factor denoted by $\psi(w', \lambda, \rho)^{20}$. We refer to the pair ($\varphi(w', \lambda, \rho), \psi(w', \lambda, \rho)$) the **canonical form of matrix factorizations** of xyz corresponding to the loop datum (w', λ, ρ).

We can transform the matrix $\Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,\rho))$ obtained in Proposition 3.15 into $\varphi(w',\lambda,\rho)$ by some bases change and prove the following theorem:

Theorem 3.17. For a loop datum (w', η, ρ) with $w' \neq (2, 2, 2)^{\mathsf{T}}$, there is an isomorphism

$$\mathscr{F}^{\mathbb{L}}\left(\mathscr{L}\left(w',\eta,
ho
ight)
ight)\ \cong\ \left(arphi\left(w',\lambda,
ho
ight),\psi\left(w',\lambda,
ho
ight)
ight)$$

in MF(xyz), where

$$\lambda := (-1)^{l_1 + \dots + l_\tau + \tau} \eta$$

for $l_i := l'_i + 1 - \mathbb{I}_{n'_{i-1} \ge 1} - \mathbb{I}_{l'_i \ge 1} - \mathbb{I}_{m'_i \ge 1}$ $(i \in \mathbb{Z}_{\tau})^{21}$.

Proof. Note that exactly one of block components $-y^{m'_1-1}I_\rho$ and $y^{-m'_1}I_\rho$ survives in $\Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,\rho))$, for example, as y^a is regarded as 0 if a < 0. By changing the sign of some basis elements, we can replace its block components

$$y^{-m_{1}'}I_{\rho}, \ z^{-n_{1}'}I_{\rho}, \ -(-x)^{l_{2}'-1}I_{\rho}, \ -(-x)^{-l_{2}'}I_{\rho}, \ \dots \ y^{-m_{\tau}'}I_{\rho}, \ z^{-n_{\tau}'}I_{\rho}, \ -(-x)^{l_{1}'-1}J_{\rho}\left(\eta\right)^{T}, \ -(-x)^{-l_{1}'}\left(J_{\rho}\left(\eta\right)^{T}\right)^{-1}$$
with

$$-y^{-m_{1}'}I_{\rho}, \ -z^{-n_{1}'}I_{\rho}, \ -x^{l_{2}'-1}I_{\rho}, \ -x^{-l_{2}'}I_{\rho}, \ \dots \ -y^{-m_{\tau}'}I_{\rho}, \ -z^{-n_{\tau}'}I_{\rho}, \ -x^{l_{1}'-1}(-1)^{\dagger}J_{\rho}\left(\eta\right)^{T}, \ -x^{-l_{1}'}(-1)^{\dagger}\left(J_{\rho}\left(\eta\right)^{T}\right)^{-1}$$

in order, where $(-1)^{\dagger}$ is the total sign change given by

$$(-1)^{\dagger} := (-1)^{\binom{l_1'-1}{\mathbb{I}_{l_1'\geq 1}-l_1'} \mathbb{I}_{l_1'\leq 0}+\mathbb{I}_{m_1'\leq 0}+\mathbb{I}_{n_1'\leq 0}+\binom{l_2'-1}{\mathbb{I}_{2'\geq 1}-l_2'} \mathbb{I}_{l_2'\leq 0}+\mathbb{I}_{m_2'\leq 0}+\cdots+\mathbb{I}_{m_{\tau}'\leq 0}+\mathbb{I}_{n_{\tau}'\leq 0}} = (-1)^{l_1+\cdots+l_{\tau}+\tau}$$

The last two can be again replaced by $-x^{l'_1-1}J_{\rho}(\lambda)$ and $-x^{-l'_1}J_{\rho}(\lambda)^{-1}$ for $\lambda := (-1)^{\dagger}\eta$, respectively, by some bases change using the fact that the matrix $(-1)^{\dagger}J_{\rho}(\eta)^T$ is similar to $J_{\rho}(\lambda)$ from the relation

$$\begin{pmatrix} 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & 0 & (-1)^{\dagger} & 0 \\ 0 & \cdots & 1 & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ \pm 1 & \cdots & 0 & 0 & 0 \end{pmatrix} (-1)^{\dagger} \begin{pmatrix} \eta & 0 & 0 & \cdots & 0 \\ 1 & \eta & 0 & \cdots & 0 \\ 0 & 1 & \eta & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & \eta \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & \cdots & \pm 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 1 & \cdots & 0 \\ 0 & (-1)^{\dagger} & 0 & \cdots & 0 \\ 1 & 0 & 0 & \cdots & 0 \end{pmatrix} = \begin{pmatrix} \lambda & 1 & 0 & \cdots & 0 \\ 0 & \lambda & 1 & \cdots & 0 \\ 0 & 0 & \lambda & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & 1 \\ 0 & 0 & 0 & \cdots & \lambda \end{pmatrix}.$$

²⁰Note that it is a consequence of Theorem 3.13, Proposition 3.14 and Theorem 3.17. An algebraic proof is given in the proof of Theorem 4.14. Indeed, the opposite matrix $\psi(w', \lambda, \rho)$ can be explicitly written, see Corollary 9.6 in [CJKR22].

²¹This is a part of the conversion formula from loop data to band data (Definition 4.10).

To manipulate and analyze the canonical form of matrix factorizations, it is convenient to introduce notations on some special matrices.

Notation 3.18. Denote the $N \times N$ Jordan block of eigenvalue 0 and its transpose as

$$J_{N} := \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ 0 & 0 & 0 & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}_{N \times N} \quad and \quad K_{N} := J_{N}^{T} = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{pmatrix}_{N \times N}$$

Note that the $n \times n$ Jordan block of eigenvalue $\lambda \in \mathbb{C}$ is $J_n(\lambda) = \lambda I_n + J_n$. We will also frequently use

$$R_{N}(\lambda) := J_{N} + \lambda K_{N}^{N-1} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ 0 & 0 & 0 & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & 1 \\ \lambda & 0 & 0 & \cdots & 0 \end{pmatrix}_{N \times N} \quad and \quad R_{N}^{T}(\lambda) = K_{N} + \lambda J_{N}^{N-1} = \begin{pmatrix} 0 & 0 & 0 & \cdots & \lambda \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{pmatrix}_{N \times N} ^{22},$$

and their enlargement by replacing λ with $J_{\rho}(\lambda)^{\pm 1}$

$$R_{N}(J_{\rho}(\lambda)) = J_{N} \otimes I_{\rho} + K_{N}^{N-1} \otimes J_{\rho}(\lambda) = \begin{pmatrix} 0 & I_{\rho} & 0 & \cdots & 0 \\ 0 & 0 & I_{\rho} & \cdots & 0 \\ 0 & 0 & 0 & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & I_{\rho} \\ J_{\rho}(\lambda) & 0 & 0 & \cdots & 0 \end{pmatrix}_{N\rho \times N\rho},$$

$$R_{N}^{T}(J_{\rho}(\lambda)^{-1}) = K_{N} \otimes I_{\rho} + J_{N}^{N-1} \otimes J_{\rho}(\lambda)^{-1} = \begin{pmatrix} 0 & 0 & 0 & \cdots & J_{\rho}(\lambda)^{-1} \\ I_{\rho} & 0 & 0 & \cdots & 0 \\ 0 & I_{\rho} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & I_{\rho} & 0 \end{pmatrix}_{N\rho \times N\rho}$$

For square matrices A_1, \ldots, A_τ , we denote the block diagonal matrix made from them as

(3.1)
$$\bigoplus_{i=1}^{\tau} A_i := \begin{pmatrix} A_1 & 0 & 0 & \cdots & 0 \\ 0 & A_2 & 0 & \cdots & 0 \\ 0 & 0 & A_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A_{\tau} \end{pmatrix}.$$

Using these new notations, we can write the canonical form given in Definition 3.16 for $\rho = 1$ as

$$\varphi(w',\lambda,1) = \varphi(w',0,1) - \lambda x^{l'_1 - 1} K_{3\tau}^{3\tau - 1} - \lambda^{-1} x^{-l'_1} J_{3\tau}^{3\tau - 1},$$

where $\varphi(w', 0, 1)$ is obtained by putting $\lambda = \lambda^{-1} = 0$ in the expression of $\varphi(w', \lambda, 1)$. Then the general expression for arbitrary ρ is obtained from it by 'substituting the matrix $J_{\rho}(\lambda)$ for the scalar λ ', that is,

(3.2)
$$\varphi(w',\lambda,\rho) = \varphi(w',0,1) \otimes I_{\rho} - x^{l'_{1}-1} K_{3\tau}^{3\tau-1} \otimes J_{\rho}(\lambda) - x^{-l'_{1}} J_{3\tau}^{3\tau-1} \otimes J_{\rho}(\lambda)^{-1}.$$

It can be also viewed as a consequence of Proposition 2.5.

By reordering rows and columns of the canonical form $\varphi(w', \lambda, \rho)$, we obtain an **alternative canonical** form

$$(3.3) \quad \varphi_{\text{alt}}(w',\lambda,\rho) := \begin{pmatrix} zI_{\tau\rho} & -\bigoplus_{i=1}^{\tau} y^{m'_{i}-1}I_{\rho} & -\left(\bigoplus_{i=1}^{\tau} x^{-l'_{i}}I_{\rho}\right) R_{N}^{T}(J_{\rho}(\lambda)^{-1}) \\ -\bigoplus_{i=1}^{\tau} y^{-m'_{i}}I_{\rho} & xI_{\tau\rho} & -\bigoplus_{i=1}^{\tau} z^{n'_{i}-1}I_{\rho} \\ -R_{N}(J_{\rho}(\lambda)) \left(\bigoplus_{i=1}^{\tau} x^{l'_{i}-1}I_{\rho}\right) & -\bigoplus_{i=1}^{\tau} z^{-n'_{i}}I_{\rho} & yI_{\tau\rho} \end{pmatrix}_{3\tau\rho\times3\tau\rho}.$$

Sometimes it is more convenient to work with this alternative version than with the original one.

²²Note that both are just $(\lambda)_{1 \times 1}$ if N = 1.

3.4. **Periodic case.** In this subsection, we show that objects corresponding to the **periodic** normal loop words are **decomposable**. The core of the decomposition lies in the following linear algebra problem:

Lemma 3.19. The Jordan canonical form of $R_N(J_\rho(\lambda))$ and $R_N^T(J_\rho(\lambda)^{-1})$ are $\bigoplus_{k=0}^{N-1} J_\rho(\lambda_k)$ and $\bigoplus_{k=0}^{N-1} J_\rho(\lambda_k^{-1})^{23}$, respectively, where $\lambda_0, \ldots, \lambda_{N-1} \in \mathbb{C}^{\times}$ are the N-th roots of $\lambda \in \mathbb{C}^{\times}$ (i.e., distinct solutions of $x^N = \lambda$).

Proof. By the following Lemma 3.20, the matrix $R_N(J_\rho(\lambda)) = J_N \otimes I_\rho + K_N^{N-1} \otimes J_\rho(\lambda)$ is similar to

$$I_{\rho} \otimes J_{N} + J_{\rho}(\lambda) \otimes K_{N}^{N-1} = \begin{pmatrix} J_{N} + \lambda K_{N}^{N-1} & K_{N}^{N-1} & 0 & \cdots & 0 \\ 0 & J_{N} + \lambda K_{N}^{N-1} & K_{N}^{N-1} & \cdots & 0 \\ 0 & 0 & J_{N} + \lambda K_{N}^{N-1} & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & K_{N}^{N-1} \\ 0 & 0 & 0 & \cdots & J_{N} + \lambda K_{N}^{N-1} \end{pmatrix}_{\rho N \times \rho N} = \begin{pmatrix} R_{N}(\lambda) & K_{N}^{N-1} & 0 & \cdots & 0 \\ 0 & R_{N}(\lambda) & K_{N}^{N-1} & \cdots & 0 \\ 0 & 0 & R_{N}(\lambda) & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & K_{N}^{N-1} \\ 0 & 0 & 0 & \cdots & R_{N}(\lambda) \end{pmatrix}_{\rho N \times \rho N}$$

whose characteristic polynomial (in *t*) is

$$(\det(tI_N - R_N(\lambda)))^{\rho} = \begin{vmatrix} t & -1 & 0 & \cdots & 0 \\ 0 & t & -1 & \cdots & 0 \\ 0 & 0 & t & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & -1 \\ -\lambda & 0 & 0 & \cdots & t \end{vmatrix} \Big|_{N \times N}^{\rho} = (t - \lambda_0)^{\rho} \cdots (t - \lambda_{N-1})^{\rho}.$$

It is straightforward to check that the eigenspace for each eigenvalue $t = \lambda_k$ ($k \in \{0, ..., N-1\}$) has dimension only 1, which completes the proof for $R_N(J_\rho(\lambda))$. The proof for $R_N^T(J_\rho(\lambda)^{-1})$ is similar.

Lemma 3.20. For two matrices $A \in \mathbb{C}^{m_1 \times n_1}$ and $B \in \mathbb{C}^{m_2 \times n_2}$, their Kronecker products $A \otimes B \in \mathbb{C}^{m_1 m_2 \times n_1 n_2}$ and $B \otimes A \in \mathbb{C}^{m_2 m_1 \times n_2 n_1}$ are permutation equivalent. More specifically, there is a perfect shuffle matrix $S_{p,q} \in O(pq)$ for each $(p,q) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1}$ whose entries are 0 or 1, such that $S_{p,q}^T = S_{p,q}^{-1} = S_{q,p}$ and

$$S_{m_1,m_2}(A \otimes B) S_{n_2,n_1} = B \otimes A$$

for any $A \in \mathbb{C}^{m_1 \times n_1}$ and $B \in \mathbb{C}^{m_2 \times n_2}$.

In particular, if A and B are square matrices, $A \otimes B$ and $B \otimes A$ are similar.

Proof. One can get $B \otimes A$ from $A \otimes B$ (and vice versa) just by reordering rows and columns. It is straightforward to check the detail (see [Wik24]).

Theorem 3.21. Let (w', λ, ρ) be a loop datum with $(w', \lambda) \neq ((2, 2, 2)^{\tau}, 1)$. If the normal loop word w' is periodic, i.e., $w' = (\tilde{w}')^N \in \mathbb{Z}^{3\tau}$ for another normal loop word $\tilde{w}' \in \mathbb{Z}^{3\tilde{\tau}}$ $(\tau = N\tilde{\tau})$, there is an invertible matrix $V \in GL_{3\tau\rho}(\mathbb{C})$ such that

$$\varphi\left(w',\lambda,\rho\right)=V^{-1}\begin{pmatrix}\varphi(\tilde{w}',\lambda_{0},\rho) & 0 & \cdots & 0\\ 0 & \varphi(\tilde{w}',\lambda_{1},\rho) & \cdots & 0\\ \vdots & \vdots & \ddots & \vdots\\ 0 & 0 & \cdots & \varphi(\tilde{w}',\lambda_{N-1},\rho)\end{pmatrix}V,$$

where $\lambda_0, \ldots, \lambda_{N-1} \in \mathbb{C}^{\times}$ are the *N*-th roots of λ . This yields a decomposition in MF(xyz)

$$\varphi(w',\lambda,\rho) \cong \bigoplus_{k=0}^{N-1} \varphi(\tilde{w}',\lambda_k,\rho).$$

Proof. Recall the formula (3.2)

$$\varphi\left(w',\lambda,\rho\right) = \varphi\left(w',0,1\right) \otimes I_{\rho} - x^{l_{1}'-1} K_{3\tau}^{3\tau-1} \otimes J_{\rho}\left(\lambda\right) - x^{-l_{1}'} J_{3\tau}^{3\tau-1} \otimes J_{\rho}\left(\lambda\right)^{-1}.$$

Here we have

$$\varphi(w',0,1) = I_N \otimes \varphi(\tilde{w}',0,1) - x_1^{l_1'-1} J_N \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} - x_1^{-l_1'} K_N \otimes J_{3\tilde{\tau}}^{3\tilde{\tau}-1}$$

 $\overline{^{23}}$ If $\rho = 1$, they are diagonalizable via a transition matrix given by the Vandermonde matrix $\left(\lambda_j^i\right)_{0 \le i,j \le N-1}$.

for a periodic word $w' = (\tilde{w}')^N$. Denoting by $\tilde{\varphi} := \varphi(\tilde{w}', 0, 1)$, we can rewrite $\varphi(w', \lambda, \rho)$ as $\left(I_N \otimes \tilde{\varphi} - x_1^{l_1'-1} J_N \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} - x_1^{-l_1'} K_N \otimes J_{3\tilde{\tau}}^{3\tilde{\tau}-1} \right) \otimes I_\rho - x^{l_1'-1} K_N^{N-1} \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} \otimes J_\rho \left(\lambda \right) - x^{-l_1'} J_N^{N-1} \otimes J_{3\tilde{\tau}}^{3\tilde{\tau}-1} \otimes J_\rho \left(\lambda \right)^{-1},$ where we also used trivial identities $K_{3\tau}^{3\tau-1} = K_N^{N-1} \otimes K_{3\tau}^{3\tau-1}$ and $J_{3\tau}^{3\tau-1} = J_N^{N-1} \otimes J_{3\tau}^{3\tilde{\tau}-1}$.

By Lemma 3.20, we can switch the order in the Kronecker product $A \otimes B \otimes C$ above into $A \otimes C \otimes B$ to get a similar matrix. That is, $\varphi(w', \lambda, \rho)$ is similar to

$$(3.4) \qquad I_{N} \otimes I_{\rho} \otimes \tilde{\varphi} - x_{1}^{l_{1}^{\prime}-1} \left(J_{N} \otimes I_{\rho} + K_{N}^{N-1} \otimes J_{\rho} \left(\lambda \right) \right) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} - x_{1}^{-l_{1}^{\prime}} \left(K_{N} \otimes I_{\rho} + J_{N}^{N-1} \otimes J_{\rho} \left(\lambda \right)^{-1} \right) \otimes J_{3\tilde{\tau}}^{3\tilde{\tau}-1} \\ = I_{N\rho} \otimes \tilde{\varphi} - x_{1}^{l_{1}^{\prime}-1} R_{N} \left(J_{\rho} \left(\lambda \right) \right) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} - x_{1}^{-l_{1}^{\prime}} R_{N}^{T} \left(J_{\rho} \left(\lambda \right)^{-1} \right) \otimes J_{3\tilde{\tau}}^{3\tilde{\tau}-1}$$

via transition matrices given by $I_N \otimes S_{3\tilde{\tau},\rho}$ and $I_N \otimes S_{\rho,3\tilde{\tau}}$. In other words, (3.4) is another expression for

$$(I_N \otimes S_{3\tilde{\tau},\rho}) \varphi(w',\lambda,\rho) (I_N \otimes S_{\rho,3\tilde{\tau}}).$$

Now we may assume $l'_1 \ge 1$. (The other case is handled in the same way.) Then we can drop the third term in (3.4). Lemma 3.19 implies that there is an invertible matrix $P \in GL_{N\rho}(\mathbb{C})$ satisfying

$$R_N(J_\rho(\lambda))P = P\left(\bigoplus_{k=0}^{N-1} J_\rho(\lambda_k)\right).$$

Multiplying $P \otimes I_{3\tilde{\tau}}$ to (3.4) on the right yields

$$\begin{split} \left(I_N \otimes S_{3\tilde{\tau},\rho}\right) \varphi\left(w',\lambda,\rho\right) \left(I_N \otimes S_{\rho,3\tilde{\tau}}\right) (P \otimes I_{3\tilde{\tau}}) &= P \otimes \tilde{\varphi} - x_1^{l_1'-1} \left(R_N\left(J_\rho\left(\lambda\right)\right)P\right) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} \\ &= P \otimes \tilde{\varphi} - x_1^{l_1'-1} P\left(\bigoplus_{k=0}^{N-1} J_\rho\left(\lambda_k\right)\right) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1} \\ &= (P \otimes I_{3\tilde{\tau}}) \left(I_{N\rho} \otimes \tilde{\varphi} - x_1^{l_1'-1} \left(\bigoplus_{k=0}^{N-1} J_\rho\left(\lambda_k\right)\right) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1}\right) \\ &= (P \otimes I_{3\tilde{\tau}}) \bigoplus_{k=0}^{N-1} \left(I_\rho \otimes \tilde{\varphi} - x_1^{l_1'-1} J_\rho\left(\lambda_k\right) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau}-1}\right). \end{split}$$

Lemma 3.20 says that each direct summand $I_{\rho} \otimes \tilde{\varphi} - x_1^{l'_1 - 1} J_{\rho}(\lambda_k) \otimes K_{3\tilde{\tau}}^{3\tilde{\tau} - 1}$ is similar to

$$\varphi\left(\tilde{w}',\lambda_{k},\rho\right) = \tilde{\varphi} \otimes I_{\rho} - x_{1}^{l_{1}'-1} K_{3\tilde{\tau}}^{3\tilde{\tau}-1} \otimes J_{\rho}\left(\lambda_{k}\right)$$

via transition matrices $S_{\rho,3\tilde{\tau}}$ and $S_{3\tilde{\tau},\rho}$, and hence their direct sum is similar to $\bigoplus_{k=0}^{N-1} \varphi(\tilde{w}',\lambda_k,\rho)$ via transition matrices $\bigoplus_{k=0}^{N-1} S_{\rho,3\tilde{\tau}} = I_N \otimes S_{\rho,3\tilde{\tau}}$ and $\bigoplus_{k=0}^{N-1} S_{3\tilde{\tau},\rho} = I_N \otimes S_{3\tilde{\tau},\rho}$. Therefore, we can write

$$(I_N \otimes S_{\rho,3\tilde{\tau}})(P^{-1} \otimes I_{3\tilde{\tau}})(I_N \otimes S_{3\tilde{\tau},\rho})\varphi(w',\lambda,\rho)(I_N \otimes S_{\rho,3\tilde{\tau}})(P \otimes I_{3\tilde{\tau}})(I_N \otimes S_{3\tilde{\tau},\rho}) = \bigoplus_{k=0}^{N-1} \varphi(\tilde{w}',\lambda_k,\rho),$$

ch shows the claim with $V := (I_N \otimes S_{\rho,3\tilde{\tau}})(P \otimes I_{3\tilde{\tau}})(I_N \otimes S_{3\tilde{\tau},\rho}) \in \operatorname{GL}_{3\tau\rho}(\mathbb{C}).$

which shows the claim with $V := (I_N \otimes S_{\rho,3\tilde{\tau}}) (P \otimes I_{3\tilde{\tau}}) (I_N \otimes S_{3\tilde{\tau},\rho}) \in GL_{3\tau\rho}(\mathbb{C})$.

Corollary 3.22. Let (w', η, ρ) be a loop datum with $w' \neq (2, 2, 2)^{\tau}$. If the normal loop word w' is periodic, *i.e.*, $w' = (\tilde{w}')^N \in \mathbb{Z}^{3\tau}$ for another normal loop word $\tilde{w}' \in \mathbb{Z}^{3\tilde{\tau}}$ ($\tau = N\tilde{\tau}$), there is a decomposition in Fuk (Σ)

$$\mathscr{L}(w',\eta,\rho) \cong \bigoplus_{k=0}^{N-1} \mathscr{L}(\tilde{w}',\eta_k,\rho),$$

where $\eta_0, \ldots, \eta_{N-1} \in \mathbb{C}^{\times}$ are the N-th roots of η .

Remark 3.23. This shows that non-primitive loops (with a local system) are decomposable in the Fukaya category, which is not intuitively obvious. In following up works, we will see that this is indeed a general and intrinsic feature of the Fukaya category of hyperbolic Riemann surfaces.

3.5. **Non-cylinder-free case** $w' = (2,2,2)^{\tau}$. Note that a loop L in Σ is not cylinder-free with \mathbb{L} only if it is freely homotopic to \mathbb{L}^{τ} for some $\tau \in \mathbb{Z}$. Only in cases of $w' = (1,0,1,0,1,0)^{\tau}$ or $(2,2,2)^{\tau}$ for some $\tau \in \mathbb{Z}_{\geq 1}$, the corresponding loop L(w') has the same free homotopy type with \mathbb{L}^{τ} and $\mathbb{L}^{-\tau}$, respectively. Then it has a chance to bound an immersed cylinder with \mathbb{L} . In the former case, it is not the case for our canonical form given in Definition 3.1. In the latter case, however, our canonical form bounds an immersed cylinder with \mathbb{L} , which indeed produces some infinite moduli spaces involved in the formula 2.5 for the matrix factor $\Psi^{\mathbb{L}} (\mathscr{L} (w', \eta, \rho))^{24}$.

To prevent this, we take a very specific **perturbed version of the loop** for the case $w' = (2,2,2)^{\tau}$. We first define the loop L := L(2,2,2) as shown in Figure 11. Then let $L((2,2,2)^{\tau})$ be the τ -concatenation (with \bigstar as a starting point) of it. For a loop datum $((2,2,2)^{\tau},\eta,\rho)$, we define the loop with a local system $\mathscr{L}((2,2,2)^{\tau},\eta,\rho)$ as in Definition 3.9 so that it has holonomy $J_{\rho}(\eta)$ at \bigstar , using the modified underlying loop $L((2,2,2)^{\tau})$.



Figure 11. Loop with a local system $\mathcal{L} := \mathcal{L}((2,2,2),\eta,\rho)$ and the corresponding matrix factorization

Definition 3.24. The **degenerate canonical form of matrix factorizations** of xyz corresponding to a loop $datum((2,2,2)^{\tau},\lambda,\rho)$ is defined as

$$\begin{aligned} \left(\varphi_{\deg}\left((2,2,2)^{\tau},\lambda,\rho\right),\psi_{\deg}\left((2,2,2)^{\tau},\lambda,\rho\right)\right) \\ & := \left(\begin{pmatrix} -zxI_{\tau\rho} & 0 & 0 & 0 \\ zI_{\tau\rho} & -yI_{\tau\rho} & 0 & 0 \\ 0 & xI_{\tau\rho} & -zI_{\tau\rho} & 0 \\ -xR_{\tau,\rho}(\lambda) & 0 & yI_{\tau\rho} & -xyI_{\tau\rho} \end{pmatrix}_{4\tau\rho\times4\tau\rho} , \begin{pmatrix} -yI_{\tau\rho} & 0 & 0 & 0 \\ -zI_{\tau\rho} & -zxI_{\tau\rho} & 0 & 0 \\ -xI_{\tau\rho} & -x^{2}I_{\tau\rho} & -xyI_{\tau\rho} & 0 \\ -I_{\tau\rho}+R_{\tau,\rho}(\lambda) & -xI_{\tau\rho} & -zI_{\tau\rho} \end{pmatrix}_{4\tau\rho\times4\tau\rho} \right). \end{aligned}$$

Theorem 3.25. For a loop datum $((2,2,2)^{T},\eta,\rho)$, there is an isomorphism

$$\mathscr{F}^{\mathbb{L}}\left(\mathscr{L}\left((2,2,2)^{\tau},\eta,\rho\right)\right) \cong \left(\varphi_{\deg}\left((2,2,2)^{\tau},\lambda,\rho\right),\psi_{\deg}\left((2,2,2)^{\tau},\lambda,\rho\right)\right)$$

in MF (xyz), where $\lambda := (-1)^{\tau} \eta$.

²⁴Still the matrix factorization can be defined over the Novikov field. Moreover, unless $\eta = -1$, we can use the formula $1 - \eta + \eta^2 - \eta^3 + \cdots = \frac{1}{1+\eta}$ to evaluate T = 1 in some infinite series to get a matrix factorization over \mathbb{C} , which is isomorphic to the corresponding (original) canonical form. But instead of justifying that formula, we will take a detour using the perturbed loop and showing Proposition 3.27.

Proof. For $\tau = 1$ case, the matrix factorization $\mathscr{F}^{\mathbb{L}}(\mathscr{L}((2,2,2),\eta,\rho))$ is computed in Figure 11, using the same method as in Proposition 3.15. It can be transformed to $(\varphi_{deg}((2,2,2)^{\tau},\lambda,\rho),\psi_{deg}((2,2,2)^{\tau},\lambda,\rho))$ by changing the sign of some basis elements. The case of $\tau \ge 2$ is handled similarly.

For $\lambda \neq 1$, we can reduce the degenerate canonical form into the original version using the following 'matrix reduction':

Lemma 3.26. [CJKR22] Let S be the power series ring $\mathbb{C}[[x_1,...,x_n]]$ of n variables and $f \in S$ its nonzero element. Assume that the pair $\left(\begin{pmatrix} C & D \\ E^T & u \end{pmatrix}, \begin{pmatrix} F & G \\ H^T & v \end{pmatrix} \right)$ is a matrix factorization of f in S, for some matrices C, $F \in S^{k \times k}$, D, E, G, $H \in S^{k \times 1}$ and u, $v \in S$. If u or v is a unit in S, there is an isomorphism

$$\begin{pmatrix} \begin{pmatrix} C & D \\ E^T & u \end{pmatrix}, \begin{pmatrix} F & G \\ H^T & v \end{pmatrix} \cong \begin{pmatrix} C - Du^{-1}E^T, F \end{pmatrix} \oplus \begin{pmatrix} 1, f \end{pmatrix} \quad or \quad \begin{pmatrix} C, F - Gv^{-1}H^T \end{pmatrix} \oplus \begin{pmatrix} f, 1 \end{pmatrix},$$

respectively, in MF(f). Therefore, the pair is isomorphic to the reduced pair $(C - Du^{-1}E^T, F)$ or $(C, F - Gv^{-1}H^T)$, respectively, in the homotopy category <u>MF</u>(f).

Proof. Assume that *u* is a unit in *S*, and consider the following diagram:

$$S^{k} \oplus S \xrightarrow{\begin{pmatrix} C & D \\ E^{T} & u \end{pmatrix}} S^{k} \oplus S \xrightarrow{\begin{pmatrix} F & G \\ H^{T} & v \end{pmatrix}} S^{k} \oplus S$$

$$\begin{pmatrix} I_{k} & 0 \\ u^{-1}E^{T} & 1 \end{pmatrix} \underset{s}{\overset{(I_{k} - Du^{-1}E^{T} & 0)}{(0 & u^{-1})}} \underset{s}{\overset{(I_{k} - Du^{-1})}{(0 & u^{-1})}} \underset{s}{\overset{(I_{k} - Du^{-1})}{(0 & u^{-1})}} S^{k} \oplus S$$

The commutativity of the diagram is immediate from some matrix calculations using the fact that the original pair is a matrix factorization of *f*. Also, note that the vertical maps are all isomorphisms. This yields an isomorphism $\left(\begin{pmatrix} C & D \\ E^T & u \end{pmatrix}, \begin{pmatrix} F & G \\ H^T & v \end{pmatrix} \right) \cong (C - Du^{-1}E^T, F) \oplus (1, f)$ in MF(*f*).

One can construct an explicit homotopy between (1, f) or (f, 1) and the zero object $0 \rightleftharpoons 0$ in MF(*f*), which shows that it is a zero object in the homotopy category <u>MF</u>(*f*). (It is also a consequence of Proposition A.15.) This means that we can drop the direct summand (1, f) or (f, 1) in <u>MF</u>(*f*).

Proposition 3.27. If $\lambda \neq 1$, the degenerate canonical form $(\varphi_{deg}((2,2,2)^{\tau},\lambda,\rho), \psi_{deg}((2,2,2)^{\tau},\lambda,\rho))$ $(\tau,\rho \in \mathbb{Z}_{\geq 1})$ is isomorphic to the original version $(\varphi((2,2,2)^{\tau},\lambda,\rho), \psi((2,2,2)^{\tau},\lambda,\rho))$ in <u>MF</u>(xyz).

Proof. In the simplest case $\tau = \rho = 1$, the reducing process in Lemma 3.26 along a unit $-1+\lambda$ in $\psi_{deg}((2,2,2),\lambda,1)$ yields a commutative diagram, which proves the claim:

$$\begin{array}{c} S^{3} \oplus S \xrightarrow{\varphi_{\deg}((2,2,2),\lambda,1)} S \oplus S^{3} \xrightarrow{\psi_{\deg}((2,2,2),\lambda,1)} S^{3} \oplus S \\ \begin{pmatrix} I_{3} - (1-\lambda)^{-1} \begin{pmatrix} y \\ z \\ x \end{pmatrix} \end{pmatrix}_{\mathcal{U}} \xrightarrow{\psi_{\deg}((2,2,2),\lambda,1)} \downarrow_{\mathcal{U}} \xrightarrow{\psi_{\varphi}((2,2,2),\lambda,1)} \xrightarrow{\psi_{\varphi}((2,2,2),\lambda,1)} \downarrow_{\mathcal{U}} \xrightarrow{\psi_{$$

In general case, we 'substitute the matrix $R_{\tau,\rho}(\lambda)$ for the scalar λ ' to get a commutative diagram:

$$S^{3\tau\rho} \oplus S^{\tau\rho} \xrightarrow{\varphi_{deg}((2,2,2)^{\tau},\lambda,\rho)} S^{\tau\rho} \oplus S^{3\tau\rho} \xrightarrow{\psi_{deg}((2,2,2)^{\tau},\lambda,\rho)} S^{3\tau\rho} \oplus S^{\tau\rho}$$

$$\begin{pmatrix} I_{3\tau\rho} - \begin{pmatrix} yI_{\tau\rho} \\ zI_{\tau\rho} \\ xI_{\tau\rho} \end{pmatrix} (I_{\tau\rho} - R_{\tau,\rho}(\lambda))^{-1} \end{pmatrix}_{\mathcal{H}} \xrightarrow{\varphi_{deg}((2,2,2)^{\tau},\lambda,\rho)} V_{\mathcal{H}} \xrightarrow{\varphi_{$$

Therefore, the degenerate canonical form is isomorphic to the alternative canonical form (3.3) in $\underline{MF}(xyz)$, which is again isomorphic to the original canonical form in MF(xyz).

This shows that even if the normal loop word is $w' = (2,2,2)^{\tau}$ (and hence we use the perturbed loop for $\mathscr{L}((2,2,2)^{\tau},\eta,\rho)$), we can still define and use the original canonical form of matrix factorizations $(\varphi((2,2,2)^{\tau},\lambda,\rho),\psi((2,2,2)^{\tau},\lambda,\rho))$ unless $\lambda = 1$ (or equivalently $\eta = (-1)^{\tau}$). Therefore, we would not call them all degenerate cases.

Definition 3.28. A normal loop word is called **non-cylinder-free** if it is $w' = (2, 2, 2)^{\tau}$ for some $\tau \in \mathbb{Z}_{\geq 1}$, and a loop datum is called **degenerate** if it is

- (1) $(w' = (2,2,2)^{\mathsf{T}}, \eta = (-1)^{\mathsf{T}}, \rho)$ and parameterizes loops with a local system $\mathscr{L}((2,2,2)^{\mathsf{T}}, (-1)^{\mathsf{T}}, \rho)$,
- (2) $(w' = (2, 2, 2)^{\tau}, \lambda = 1, \rho)$ and parameterizes matrix factorizations $(\varphi_{deg}((2, 2, 2)^{\tau}, 1, \rho), \psi_{deg}((2, 2, 2)^{\tau}, 1, \rho))$.

Now Theorem 3.17, Theorem 3.25, and Proposition 3.27 complete the proof of Theorem 1.3 in the introduction.

For $\tau \ge 2$, the non-cylinder-free loop word $(2,2,2)^{\tau}$ is **periodic**, and the corresponding objects are **de-composable** as in the general case (Theorem 3.21):

Proposition 3.29. For a loop datum $((2,2,2)^{\tau},\lambda,\rho)$, there is a decomposition

$$\varphi_{\mathrm{deg}}\big((2,2,2)^{\tau},\lambda,\rho\big) \cong \bigoplus_{k=0}^{\tau-1} \varphi_{\mathrm{deg}}\big((2,2,2),\lambda_{\tau,k},\rho\big)$$

in MF(*xyz*), where $\lambda_{\tau,0}, \ldots, \lambda_{\tau,\tau-1} \in \mathbb{C}$ are the τ -th roots of λ .

Proof. It is proven in the same way as Theorem 3.21.

Combining Proposition 3.29 and Proposition 3.27, we get a decomposition of the matrix factorization for periodic degenerate cases ($w' = (2, 2, 2)^{T}$, $\lambda = 1$) as

(3.5)

$$\varphi_{\deg}((2,2,2)^{\tau},1,\rho) \cong \bigoplus_{k=0}^{\tau-1} \varphi_{\deg}((2,2,2),e^{2\pi i \cdot \frac{k}{\tau}},\rho) \quad \text{in MF}(xyz) \\
\cong \varphi_{\deg}((2,2,2),1,\rho) \oplus \bigoplus_{k=1}^{\tau-1} \varphi((2,2,2),e^{2\pi i \cdot \frac{k}{\tau}},\rho) \quad \text{in } \underline{\mathrm{MF}}(xyz).$$

This also implies the decomposition of the loop with a local system for periodic degenerate cases ($w' = (2,2,2)^{\tau}, \eta = (-1)^{\tau}$) as

(3.6)
$$\mathscr{L}\left((2,2,2)^{\tau},(-1)^{\tau},\rho\right) \cong \bigoplus_{k=0}^{\tau-1} \mathscr{L}\left((2,2,2),-e^{2\pi i \cdot \frac{k}{\tau}},\rho\right) \quad \text{in Fuk}(\Sigma),$$

extending the result of Corollary 3.22 to any periodic normal loop words.

4. MATRIX FACTORIZATIONS FROM MAXIMAL COHEN-MACAULAY MODULES

In this section, we first recall some general concepts on *maximal Cohen-Macaulay modules* which we will need in the rest of the section, including the *Macaulayfication* and its computation (§4.1). Then we recall Burban-Drozd's classification [BD17a] of *band-type* indecomposable maximal Cohen-Macaulay modules over $A := \mathbb{C}[[x, y, z]]/(xyz)$. Their canonical form is provided as a submodule of a free module $A^{\tau\mu}$, whose generators are given in terms of *band data* (§4.2). Those band data are in one-to-one correspondence with loop data by the *conversion formula* given in [CJKR22] (§4.3). Using this correspondence, we prove the main theorem in this section that the canonical form of matrix factorizations corresponding to loop data is related to the canonical form of band-type maximal Cohen-Macaulay modules corresponding to band data under Eisenbud's equivalence (§4.4). The *degenerate case* will be treated separately (§4.5).

4.1. Maximal Cohen-Macaulay modules and Macaulayfication. Let (A, \mathfrak{m}) be a Noetherian local ring, $\Bbbk := A/\mathfrak{m}$ its residue field, and $d := \operatorname{kr.dim}(A)$ its Krull dimension.

Definition 4.1. A Noetherian A-module M is called maximal Cohen-Macaulay if

 $\operatorname{Ext}_{A}^{i}(\mathbb{k}, M) = 0 \quad for \ i \in \{0, \dots, d-1\}.$

We denote by CM(A) the category of maximal Cohen-Macaulay modules over A, which is a full subcategory of the category A - mod of Noetherian (i.e. finitely generated) A-modules.

We refer readers, for example, to [Yos90] for general properties and representations of maximal Cohen-Macaulay modules, and [BH98] for (not necessarily maximal) Cohen-Macaulay modules. However, note that many authors, including [Yos90], refer to maximal Cohen-Macaulay modules simply as *Cohen-Macaulay modules*.

4.1.1. *Macaulayfication*. Now let us focus on our specific case where *A* is given by $\mathbb{C}[[x, y, z]]/(xyz)$. It is an example of *surface singularities* (i.e. *d* = 2), and many technics and representation-theoretic aspects for them were developed and studied in [BD08, BD17a]. We will especially use the following *Macaulayfying* process, which naturally associates any Noetherian *A*-module a maximal Cohen-Macaulay *A*-module:

Definition 4.2. [BD08] The Macaulayfication of a Noetherian A-module \tilde{M} is defined by

$$\tilde{M}^{\dagger} := \operatorname{Hom}_{A} \left(\operatorname{Hom}_{A} \left(\tilde{M}, A \right), A \right)^{25}.$$

It is indeed a maximal Cohen-Macaulay A-module ²⁶. This defines a functor \dagger : $A - \text{mod} \rightarrow \text{CM}(A)$, which is called the Macaulayfication functor.

We will use a combinatorial tool to compute the Macaulayfication of a given A-module in practice:

Definition 4.3. Let \tilde{M} be an A-submodule of a free module A^r . If there is an element $F \in A^r \setminus \tilde{M}$ such that xF, yF, $zF \in \tilde{M}$, we call it a Macaulayfying element of \tilde{M} in A^r .

Proposition 4.4. [BD17a] For an A-submodule \tilde{M} of a free module A^r , the following hold:

(1) \tilde{M} is maximal Cohen-Macaulay if and only if there is no Macaulayfying element of \tilde{M} in A^r . We have $\tilde{M}^{\dagger} = \tilde{M}$ in this case.

(2) $\tilde{M}^{\dagger} \cong \langle \tilde{M}, F \rangle_A^{\dagger}$ holds for any Macaulayfying element F of \tilde{M} in A^r .

Proof. See Proposition 4.2 in [CJKR22] for the proof of (1), and Lemma 1.5 in [BD17a] for (2).

²⁵We are using the fact that *A* is *Gorenstein in codimension one* and therefore its *canonical module* is isomorphic to *A*. See [BD08] for general definition when *A* does not have a such property.

²⁶Here the fact that the Krull dimension of A is 2 is essential. See Lemma 3.1 in [BD08].

4.2. Band data and canonical form of maximal Cohen-Macaulay modules. We recall the concept of the *band data* from [BD17a], which parameterize *band-type* indecomposable maximal Cohen Macaulay modules over $A = \mathbb{C}[[x, y, z]]/(xyz)$. Here we use a slightly modified version of band words given in [CJKR22] for our specific singularity *A*, in order to match them with the loop data in the next subsection.

Definition 4.5. A band datum (w, λ, μ) consists of the following:

- (band word) $w = (l_1, m_1, n_1, l_2, m_2, n_2, \dots, l_{\tau}, m_{\tau}, n_{\tau}) \in \mathbb{Z}^{3\tau}$ for some $\tau \in \mathbb{Z}_{\geq 1}$,
- (eigenvalue) $\lambda \in \mathbb{C}^{\times}$,
- ((algebraic) multiplicity) $\mu \in \mathbb{Z}_{\geq 1}$.

It defines an A-module, denoted by

$$\tilde{M}(w,\lambda,\mu)$$

as an A-submodule of $A^{\tau\mu}$ generated by all columns of the 6 matrices

$$\begin{aligned} x^2 y^2 I_{\tau\mu}, \quad y^2 z^2 I_{\tau\mu}, \quad z^2 x^2 I_{\tau\mu}, \quad \pi_x (w, \lambda, \mu) &:= \begin{pmatrix} x^{l_1^- + 2} y I_\mu & z x^{l_2^+ + 2} I_\mu & \cdots & 0 \\ 0 & x^{l_2^- + 2} y I_\mu & \ddots & \vdots \\ \vdots & \vdots & \ddots & z x^{l_1^+ + 2} I_\mu \\ z x^{l_1^+ + 2} J_\mu (\lambda) & 0 & \cdots & x^{l_r^- + 2} y I_\mu \end{pmatrix}_{\tau\mu \times \tau\mu} \\ \pi_y (w, \lambda, \mu) &:= \begin{pmatrix} \left(x y^{m_1^+ + 2} + y^{m_1^- + 2} z \right) I_\mu & 0 & \cdots & 0 \\ 0 & \left(x y^{m_2^+ + 2} + y^{m_2^- + 2} z \right) I_\mu & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \left(x y^{m_r^+ + 2} + y^{m_r^- + 2} z \right) I_\mu \end{pmatrix}_{\tau\mu \times \tau\mu} \\ \pi_z (w, \lambda, \mu) &:= \begin{pmatrix} \left(y z^{n_1^+ + 2} + z^{n_1^- + 2} x \right) I_\mu & 0 & \cdots & 0 \\ 0 & \left(y z^{n_2^+ + 2} + z^{n_2^- + 2} x \right) I_\mu & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \left(y z^{n_r^+ + 2} + z^{n_r^- + 2} x \right) I_\mu \end{pmatrix}_{\tau\mu \times \tau\mu} \end{aligned}$$

where $a^+ := \max\{0, a\}$ and $a^- := \max\{0, -a\}$ for $a \in \mathbb{Z}$.

The notion of *shift*, *subword*, *concatenation* and *periodicity* of a band word can be defined similarly following those of a loop word. Two band words w and \tilde{w} are considered *equivalent* if they coincide up to shifting, that is, $\tilde{w} = w^{(k)}$ for some $k \in \mathbb{Z}$.

In general, however, the *A*-module $\tilde{M}(w, \lambda, \mu)$ fails to be maximal Cohen-Macaulay. So we need to *Macaulayfy* it to get an object in CM(*A*) from a band datum.

Definition 4.6. *Given a band datum* (w, λ, μ) *, we define the corresponding maximal Cohen-Macaulay A-module*

$$M(w,\lambda,\mu) := \tilde{M}(w,\lambda,\mu)^{\dagger}$$

as the Macaulay fication of $\tilde{M}(w, \lambda, \mu)$. We refer to it as the **canonical form of band-type maximal Cohen-Macaulay modules** over A corresponding to the band datum (w, λ, μ) .

The maximal Cohen-Macaulay module $M(w, \lambda, \mu)$ constructed from any band datum is **locally free on the punctured spectrum of** A, that is, for any $\mathfrak{p} \in \text{Spec}(A) \setminus \{\mathfrak{m}\}$ the localization $M_{\mathfrak{p}}$ is a free $A_{\mathfrak{p}}$ -module, where \mathfrak{m} is the maximal ideal of A. The converse also holds, which is the following classification theorem:

Theorem 4.7. [BD17a] Any indecomposable maximal Cohen-Macaulay module over A that is locally free on the punctured spectrum is isomorphic to the canonical form $M(w, \lambda, \mu)$ for some unique non-periodic band datum (w, λ, μ) up to shifting of the band word w. The proof of the theorem actually follows from a highly non-trivial representation-theoretic study of the category CM(*A*), which is applicable to much broader class (i.e. *degenerate cusp*) of *non-isolated* surface singularities. To be more specific, they created its equivalent category Tri(*A*), called the *category of triples* (or *Burban-Drozd's triple category* in some literature), and a natural equivalence functor \mathbb{F}_{BD} : CM(*A*) $\xrightarrow{\simeq}$ Tri(*A*)²⁷. Its object consists of two modules and a 'linearized morphism' between them, and the indecomposable objects can be classified via a *matrix problem* on the linear map. As a result, they fall into the following two types: *band-type* and *string-type*. The canonical form $\Theta(w, \lambda, \mu)$ of band-type indecomposable objects in Tri(*A*) is given in Figure 12, which are parameterized by the band data (w, λ, μ) . Then they determine the corresponding objects $M(w, \lambda, \mu)$ in CM(*A*) under the equivalence, which we take as presented in Definition 4.6.



$$\Theta\left(w=(l_{i},m_{i},n_{i})_{i=1}^{\tau},\lambda,\mu\right)$$

Figure 12. Canonical form of band-type indecomposable objects in Tri (A)

Remark 4.8. One can also consider a direct functor $\underline{MF}(xyz) \rightarrow \underline{Tri}(A)$ by composing the Eisenbud's cokernel functor and Burban-Drozd's functor \mathbb{F}_{BD} . Then we have an alternative way to compute objects in $\underline{Tri}(A)$ corresponding to the canonical form of objects in $\underline{MF}(xyz)$. This gives another proof for Theorem 4.14 without going through the Macaulayfication process. However, since it involves additional algebro-geometric consideration for the category $\underline{Tri}(A)$ ([Rho23, Chapter 5]), we defer it to a separate future work.

Remark 4.9. Indecomposable maximal Cohen-Macaulay modules over A that are not locally free on the punctured spectrum were also classified in [BD17a] by using string data instead of band data. On the mirror side, they correspond to arcs starting and ending at boundaries of Σ . Most statements and proofs in the present paper apply to these cases without significant modifications, while one should take care on some technical details when proving homotopy invariance of matrix factorizations to establish a one-to-one correspondence of them with open geodesics. There we have to consider wrapped morphisms between arcs (non-compact Lagrangian submanifolds), and a discussion on the wrapped Fukaya category of immersed non-exact Lagrangians should precede this. Here we present only some correspondence between basic objects as in Figure 13, which also correspond to $z \cdot xy$, $x \cdot yz$ and $y \cdot zx$ in MF(xyz), respectively.



Figure 13. Mirror symmetry of generating objects

²⁷See also [BZ20] for another elaboration on this equivalence and its applications.

4.3. **Conversion formula between loop/band data.** We define the *conversion formula*, which underlies the correspondence between the canonical form of matrix factorizations of *xyz* (in loop data) and the canonical form of band-type indecomposable maximal Cohen-Macaulay modules over *A* (in band data). It was first introduced in [CJKR22], and now we reformulate and extend it for higher rank/multiplicity.

Definition 4.10 (Conversion from loop data to band data). A loop datum (w', η, ρ) with a normal loop word $w' = (w'_1, w'_2, w'_3, w'_4, w'_5, w'_6, \dots, w'_{3\tau-2}, w'_{3\tau-1}, w'_{3\tau}) \in \mathbb{Z}^{3\tau}$ is converted to a band datum (w, λ, μ) as follows:

• The normal loop word w' is converted to the band word $w \in \mathbb{Z}^{3\tau}$, defined as

 $w_j := w'_j + 1 - \mathbb{1}_{w'_{i-1} \ge 1} - \mathbb{1}_{w'_i \ge 1} - \mathbb{1}_{w'_{i+1} \ge 1} \quad (j \in \mathbb{Z}_{3\tau}).$

• The holonomy parameter $\eta \in \mathbb{C}^{\times}$ is converted to the eigenvalue

$$\lambda := (-1)^{l_1 + \dots + l_\tau + \tau} \eta \quad \text{where } l_i := w_{3i-2} \text{ for } i \in \mathbb{Z}_\tau.$$

• The geometric rank ρ is converted to the algebraic multiplicity ²⁸

$$\mu := \begin{cases} \rho & \text{in non-degenerate cases,} \\ \rho + 1 & \text{in the non-periodic degenerate case} \left(w' = (2, 2, 2), \eta = -1 \right). \end{cases}$$

We call a band datum **degenerate** if it is $(w = (0,0,0)^{\tau}, \lambda = 1, \mu)$ for some $\tau, \mu \in \mathbb{Z}_{\geq 1}$. Note that it corresponds to a degenerate loop datum $(w' = (2,2,2)^{\tau}, \eta = (-1)^{\tau}, \rho)$ under above the conversion formula above (while ignoring the relation between μ and ρ which is not defined if $\tau \geq 2$).

Definition 4.11 (Conversion from band data to loop data). A band datum (w, λ, μ) with a band word $w = (w_1, w_2, w_3, w_4, w_5, w_6, \dots, w_{3\tau-2}, w_{3\tau-1}, w_{3\tau}) \in \mathbb{Z}^{3\tau}$ is converted to a loop datum (w', η, ρ) as follows:

• The band word w is converted to the normal loop word $w' \in \mathbb{Z}^{3\tau}$, defined as

$$w'_j := w_j - 1 + \delta_{j-1} + \delta_j + \delta_{j+1} \quad (j \in \mathbb{Z}_{3\tau})$$

where $\delta = \delta(w) \in \{0, 1\}^{3\tau}$ is given by

 $\delta_{j} := \begin{cases} 0 & if \\ & & \\ &$

• The eigenvalue $\lambda \in \mathbb{C}^{\times}$ is converted to the holonomy parameter

$$\eta := (-1)^{l_1 + \dots + l_\tau + \tau} \lambda \quad where \ l_i := w_{3i-2} \ for \ i \in \mathbb{Z}_\tau.$$

• The algebraic multiplicity μ is converted to the geometric rank ²⁹

$$\rho := \begin{cases} \mu & \text{in non-degenerate cases,} \\ \mu - 1 & \text{in the non-periodic degenerate case } (w = (0, 0, 0), \lambda = 1). \end{cases}$$

Proposition 4.12. [CJKR22] (1) The loop word w' converted from a band word w is indeed normal.
(2) Two conversion formula above are inverse to each other.

²⁸We do not define it for periodic degenerate cases ($w' = (2, 2, 2)^{\tau}$, $\eta = (-1)^{\tau}$ with $\tau \ge 2$), because the corresponding loop with a local system and matrix factorization are decomposed into τ pieces ((3.6) and (3.5)) and they are mapped to maximal Cohen-Macaulay modules having different algebraic multiplicities (4.9).

²⁹We do not define it for periodic degenerate cases ($w = (0, 0, 0)^{\tau}$, $\lambda = 1$ with $\tau \ge 2$) in the same reason as above.

Example 4.13. The following shows the conversion between a normal loop word w' and a band word w:

$$\begin{split} w' &= (\ 8\ ,\ 2\ ,\ 3\ ,\ -1\ ,\ -1\ ,\ -1\ ,\ 0\ ,\ 5\ ,\ 0\ ,\ -2\ ,\ 1\ ,\ 0\ ,\ 4\ ,\ 6\) \\ \mathbb{1}_{w'\geq 1} = \delta(w) = (\ 1\ ,\ 1\ ,\ 1\ ,\ 0\ ,\ 0\ ,\ 0\ ,\ 0\ ,\ 0\ ,\ 0\ ,\ 0\ ,\ 1\ ,\ 0\ ,\ 1\ ,\ 0\ ,\ 1\ ,\ 1\ ,\ 1\) \\ w &= (\ 6\ ,\ 0\ ,\ 2\ ,\ -1\ ,\ 0\ ,\ -3\ ,\ 0\ ,\ 0\ ,\ 5\ ,\ 0\ ,\ -2\ ,\ 1\ ,\ -1\ ,\ 3\ ,\ 4\) \end{split}$$

The holonomy parameter η *and eigenvalue* λ *in this case are related by* $\lambda = (-1)^{6-1+0+0-1+5} \eta = -\eta$.

4.4. **Matrix Factorizations from canonical form of maximal Cohen-Macaulay modules.** Let $S := \mathbb{C}[[x, y, z]]$ be a power series ring and $A := \mathbb{C}[[x, y, z]]/(xyz)$ a hypersurface singularity. For a non-degenerate band datum (w, λ, μ) , let w' be the converted normal loop word from the band word w and $\rho := \mu$. They define objects in the canonical form as follows:

- the matrix factorization $(\varphi(w', \lambda, \rho), \psi(w', \lambda, \rho))$ corresponding to the loop datum (w', λ, ρ) ,
- the maximal Cohen-Macaulay module $M(w, \lambda, \mu)$ corresponding to the band datum (w, λ, μ) .

Now we state our main theorem in this section that they are related under Eisenbud's equivalence:

Theorem 4.14. Let (w, λ, μ) be a non-degenerate band datum, w' the converted normal loop word from the band word w and $\rho := \mu$. Then there is an isomorphism in <u>CM</u>(A)

$$\operatorname{coker} \varphi(w', \lambda, \rho) \cong M(w, \lambda, \mu),$$

or equivalently, regarding $M(w, \lambda, \mu)$ as an S-module $M_S(w, \lambda, \mu)$, it has a free resolution of the form

$$0 \longrightarrow S^n \xrightarrow{\varphi(w',\lambda,\rho)} S^n \xrightarrow{\pi} M_S(w,\lambda,\mu) \longrightarrow 0.$$

To prove the theorem, we need to compute an explicit Macaulayfication of the module in Definition 4.5 and find its resolution. The case of $\mu = 1$ was carried out in [CJKR22]. We have developed our setup so that we can extend the construction of $\mu = 1$ to the general case by '**substituting the matrix** $J_{\mu}(\lambda)$ for the scalar λ ' in all matrices. (Recall that we did the same operation in the geometric side (Proposition 2.5).) We explain the procedure in detail below:

4.4.1. *Preservation of Exactness*. Let R be a \mathbb{C} -algebra and consider a one-parameter family of matrices in the form

$$\varphi(\lambda) = \sum_{k=-N}^{N} \varphi_k \lambda^k \in \mathbb{R}^{m \times m}$$

for some matrices $\varphi_k \in \mathbb{R}^{m \times n}$. For any \mathbb{C} -valued square matrix $\Lambda \in \mathbb{C}^{\mu \times \mu}$, we associate a new matrix by

$$\varphi(\Lambda) := \sum_{k=-N}^{N} \varphi_k \otimes \Lambda^k \in R^{m\mu \times n\mu}.$$

We call the pair $(\varphi(\lambda), \varphi(\Lambda))$ a (λ, Λ) -substitution pair. We have seen many examples:

- $(\varphi(w', \lambda, 1), \varphi(w', \lambda, \rho))$ (Definition 3.16) forms a $(\lambda, J_{\rho}(\lambda))$ -substitution pair. We denote them as $\varphi(w', \lambda) := \varphi(w', \lambda, 1)$ and $\varphi(w', J_{\rho}(\lambda)) := \varphi(w', \lambda, \rho)$.
- $(\pi_{\chi}(w, \lambda, 1), \pi_{\chi}(w, \lambda, \mu))$ (Definition 4.5) forms a $(\lambda, J_{\mu}(\lambda))$ -substitution pair for each $\chi \in \{x, y, z\}$, denoted as

$$\pi_{\chi}(w,\lambda) := \pi_{\chi}(w,\lambda,1) \text{ and } \pi(w,J_{\mu}(\lambda)) := \pi(w,\lambda,\mu).$$

Such a pair enjoys a nice homological property, namely, (λ, Λ) -substitution preserves the exactness of a sequence as in the following proposition. We thank Kyoungmo Kim for providing us the idea of the proof.

Proposition 4.15. *Let*

$$\varphi(\lambda) = \sum_{k=-N}^{N} \varphi_k \lambda^k \in \mathbb{R}^{m \times n} \quad and \quad \psi(\lambda) = \sum_{k=-N}^{N} \psi_k \lambda^k \in \mathbb{R}^{l \times m}$$

be one-parameter families of matrices for some $\varphi_k \in \mathbb{R}^{m \times n}$, $\psi_k \in \mathbb{R}^{l \times n}$, which form a sequence

for each $\lambda \in \mathbb{C}$. Given a square matrix $\Lambda \in \mathbb{C}^{\mu \times \mu}$, we have new matrices

$$\varphi(\Lambda) = \sum_{k=-N}^{N} \varphi_k \otimes \Lambda^k \in \mathbb{R}^{m\mu \times n\mu} \quad and \quad \psi(\Lambda) = \sum_{k=-N}^{N} \psi_k \otimes \Lambda^k \in \mathbb{R}^{l\mu \times m\mu}$$

so that $(\varphi(\lambda), \varphi(\Lambda))$ and $(\psi(\lambda), \psi(\Lambda))$ form (λ, Λ) -substitution pairs, which also make a sequence

(4.2)
$$R^n \otimes R^\mu \xrightarrow{\varphi(\Lambda)} R^m \otimes R^\mu \xrightarrow{\psi(\Lambda)} R^l \otimes R^\mu.$$

Now if the sequence (4.1) is exact, more precisely, if we assume

- (*i*) $\psi(\lambda) \varphi(\lambda) = 0$ for any $\lambda \in \mathbb{C}$,
- (*ii*) im $\varphi(\lambda) = \ker \psi(\lambda)$ for any eigenvalue λ of Λ ,

then the sequence (4.2) is also exact, i.e., $\operatorname{im} \varphi(\Lambda) = \operatorname{ker} \psi(\Lambda)$.

Proof. The assumption (i) is equivalent to each coefficient of λ^k in the expansion of $\psi(\lambda) \varphi(\lambda)$ being zero ³⁰, that is, $\psi(\lambda) \varphi(\lambda)$ is zero as a Laurent polynomial in λ . This implies $\psi(\Lambda) \varphi(\Lambda) = 0$.

The converse is not immediate. First we show that we can replace Λ with any similar matrix J. Namely, let $J \in \mathbb{C}^{\mu \times \mu}$ be a matrix satisfying $\Lambda = P^{-1}JP$ for some invertible matrix $P \in GL_{\mu}(\mathbb{C})$. Then the computation

$$\varphi(\Lambda) = \sum_{k=-N}^{N} \varphi_k \otimes \Lambda^k = \sum_{k=-N}^{N} \varphi_k \otimes \left(P^{-1}J^kP\right) = \sum_{k=-N}^{N} \left(I_m \otimes P^{-1}\right) \left(\varphi_k \otimes J^k\right) (I_n \otimes P)$$
$$= \left(I_m \otimes P^{-1}\right) \left(\sum_{k=-N}^{N} \varphi_k \otimes J^k\right) (I_n \otimes P) = \left(I_m \otimes P^{-1}\right) \varphi(J) (I_n \otimes P)$$

and the same computation for $\psi(\Lambda)$ show the commutativity of the following diagram:

$$\begin{array}{c|c} R^{n} \otimes R^{\mu} & \xrightarrow{\varphi(\Lambda)} & R^{m} \otimes R^{\mu} & \xrightarrow{\psi(\Lambda)} & R^{l} \otimes R^{\mu} \\ I_{n} \otimes P \wr \downarrow \downarrow & & & \\ R^{n} \otimes R^{\mu} & \xrightarrow{\varphi(J)} & R^{m} \otimes R^{\mu} & \xrightarrow{\psi(J)} & R^{l} \otimes R^{\mu} \end{array}$$

Since the vertical maps are all isomorphisms, the statement is equivalent to $\operatorname{im} \varphi(J) = \ker \psi(J)$. This enables us to replace Λ with its Jordan canonical form.

Furthermore, if *J* is decomposed into block diagonals J_1, \ldots, J_k (in the sense of (3.1)), an analogous argument shows that the statement holds for *J* if and only if it holds for each block J_i . Therefore, it is enough to prove the statement for a Jordan block $J_{\mu}(\lambda)$, where λ is an eigenvalue of Λ .

We proceed by an induction on μ . For $\mu = 1$, the statement is the same as the assumption (ii). Now let $\mu \ge 2$ and assume that the statement is true for $\Lambda = J_{\mu-1}(\lambda)$. We deform $\varphi(J_{\mu}(\lambda))$ into its similar matrix $\begin{pmatrix} \varphi(J_{\mu-1}(\lambda)) & * \\ 0 & \varphi(\lambda) \end{pmatrix}$ by the following computation:

 $^{^{30}}$ This is true for any infinite (e.g. algebraically closed) field, including our field \mathbb{C} .

$$\begin{split} \varphi \left(J_{\mu} (\lambda) \right) &= \sum_{k=-N}^{N} \varphi_{k} \otimes J_{\mu} (\lambda)^{k} = \sum_{k=-N}^{N} \varphi_{k} \otimes \left(\frac{J_{\mu-1}(\lambda)}{0} \frac{*}{\lambda} \right)^{k} = \sum_{k=-N}^{N} \varphi_{k} \otimes \left(\frac{J_{\mu-1}(\lambda)^{k}}{0} \frac{*}{\lambda^{k}} \right) \\ &= \sum_{k=-N}^{N} S_{\mu,m} \left(\left(\frac{J_{\mu-1}(\lambda)^{k}}{0} \frac{*}{\lambda^{k}} \right) \otimes \varphi_{k} \right) S_{n,\mu} = S_{\mu,m} \left(\sum_{k=-N}^{N} \left(\frac{J_{\mu-1}(\lambda)^{k} \otimes \varphi_{k}}{0} \frac{*}{\lambda^{k} \varphi_{k}} \right) \right) S_{n,\mu} \\ &= S_{\mu,m} \left(\sum_{k=-N}^{N} \left(\frac{S_{m,\mu-1}(\varphi_{k} \otimes J_{\mu-1}(\lambda)^{k}) S_{\mu-1,n}}{0} \frac{*}{\lambda^{k} \varphi_{k}} \right) \right) S_{n,\mu} \\ &= S_{\mu,m} \left(S_{m,\mu-1} \oplus 1 \right) \left(\frac{\varphi (J_{\mu-1}(\lambda))}{0} \frac{*}{\varphi(\lambda)} \right) \left(S_{\mu-1,n} \oplus 1 \right) S_{n,\mu}. \end{split}$$

The same computation also works for $\psi(J_{\mu}(\lambda))$, which yields the following commutative diagram:

$$\begin{array}{c} R^{n} \otimes (R^{\mu-1} \oplus R) & \xrightarrow{\varphi(J_{\mu}(\lambda))} & R^{m} \otimes (R^{\mu-1} \oplus R) & \xrightarrow{\psi(J_{\mu}(\lambda))} & R^{l} \otimes (R^{\mu-1} \oplus R) \\ (S_{\mu-1,n} \oplus 1) S_{n,\mu} & & & \\ (S_{\mu-1,n} \oplus 1) S_{n,\mu} & & & \\ (I_{\mu-1}(\lambda)) & I_{\mu} & & \\ (I_{\mu}(I_{\mu})) & I_{\mu} & & \\ (I_{\mu-1}(\lambda)) & I_{\mu} & & \\ (I_{\mu}(I_{\mu})) & & \\ (I_{\mu}(I_{\mu})) & I_{\mu} & & \\ (I_{\mu}(I_{\mu})) & I_{\mu} & & \\ (I_{\mu}(I_{\mu})) & & \\ (I_{\mu}(I_{\mu}))$$

Note that the vertical maps are natural isomorphisms. As we know $\psi(J_{\mu}(\lambda))\varphi(J_{\mu}(\lambda)) = 0$ from our first discussion, the composition in the bottom row also vanishes. The exactness of the bottom row easily follows from the induction hypothesis and the assumption (ii), also implying that the top row is exact.

4.4.2. *Proof of the theorem.* Now recall that we have $\tilde{M}(w, \lambda, \mu) = \operatorname{im} \tilde{\pi}(w, \lambda, \mu) \subset A^{\tau\mu}$, where

is an *A*-valued matrix, or an *A*-module map $A^{6\tau\mu} \rightarrow A^{\tau\mu}$. Because it does not directly become maximal Cohen-Macaulay, we will find its Macaulayfying elements in $A^{\tau\mu}$ to Macaulayfy it. Here we briefly recall our previous discussion in [CJKR22] for $\mu = 1$ case:

For a fixed band word w, consider a one-parameter family of elements in A^{τ} of the form

(4.3)
$$F(\lambda) = F_{-}\lambda^{-1} + F_{0} + F_{+}\lambda \text{ where } F_{-}, F_{0}, F_{+} \in A^{\tau} \text{ and } \lambda \in \mathbb{C}$$

(see (9.8) in [CJKR22]) that satisfies

(4.4)
$$\chi F(\lambda) = \tilde{\pi}(w, \lambda, 1) a_{\chi}(\lambda) \text{ for each } \chi \in \{x, y, z\} \text{ and } \lambda \in \mathbb{C}^{\times}$$

for some $a_{\chi}(\lambda) = a_{\chi,-}\lambda^{-1} + a_{\chi,0} + a_{\chi,1}\lambda$ where $a_{\chi,-}, a_{\chi,0}, a_{\chi,+} \in A^{6\tau\mu}$ (see (9.10) in [CJKR22]). Then we have $\chi F(\lambda) \in \tilde{M}(w, \lambda, 1)$ for each $\chi \in \{x, y, z\}$ and $\lambda \in \mathbb{C}^{\times}$, which means that $F(\lambda)$ is a Macaulayfying element of $\tilde{M}(w, \lambda, 1)$ in A^{τ} for each $\lambda \in \mathbb{C}^{\times}$.

Theorem 4.16 (Theorem 9.1 in [CJKR22]). Let $(w, \lambda, 1)$ be a non-degenerate band datum and w' the converted normal loop word from the band word w. Then there are Macaulayfying elements $F_1(\lambda), \ldots, F_{\xi}(\lambda)$ of $\tilde{M}(w, \lambda, 1)$ in A^{τ} of the form (4.3), realizing the Macaulayfication of $\tilde{M}(w, \lambda, 1)$ as

$$M(w,\lambda,1) = \tilde{M}(w,\lambda,1)^{\dagger} = \langle \tilde{M}(w,\lambda,1), F_1(\lambda), \dots, F_{\xi}(\lambda) \rangle^{31}.$$

Moreover, denoting the right side as $\operatorname{im} \pi(w, \lambda, 1)$ for some matrix $\pi(w, \lambda, 1) \in A^{\tau \times 3\tau}$, it fits into the free resolution of $M_S(w, \lambda, 1)$, which is $M(w, \lambda, 1)$ viewed as an S-module:

(4.5)
$$0 \longrightarrow S^n \xrightarrow{\varphi(w',\lambda,1)} S^n \xrightarrow{\pi(w,\lambda,1)} M_S(w,\lambda,1) \longrightarrow 0.$$

³¹In degenerate case ($w = (0, 0, 0)^{\tau}$, $\lambda = 1$), we find only one Macaulayfying element (which doesn't belong to a one-parameter family in λ) to acheive the Macaulayfication, because there are no Macaulayfying elements for ($w = (0, 0, 0)^{\tau}$, $\lambda \neq 1$).

Now we '**substitute** $J_{\mu}(\lambda)$ for λ ' in all matrices to accomplish the same result for $\mu \ge 2$:

Proof of Theorem 4.14. For each Macaulayfying element $F(\lambda) = F_-\lambda^{-1} + F_0 + F_+\lambda \in A^{\tau}$ of $\tilde{M}(w, \lambda, 1)$ in A^{τ} given in Theorem 4.16, we associate a matrix $F(J_{\mu}(\lambda)) := F_- \otimes J_{\mu}(\lambda)^{-1} + F_0 \otimes I_{\mu} + F_+ \otimes J_{\mu}(\lambda) \in A^{\tau\mu\times\mu}$ to get a $(\lambda, J_{\mu}(\lambda))$ -substitution pair $(F(\lambda), F(J_{\mu}(\lambda)))$. Then relation (4.4) implies

 $\chi F(J_{\mu}(\lambda)) = \tilde{\pi}(w, \lambda, \mu) a_{\chi}(J_{\mu}(\lambda)) \text{ for each } \chi \in \{x, y, z\},$

which shows that each column of $F(J_{\mu}(\lambda))$ is a Macaulayfying element of $\tilde{M}(w, \lambda, \mu)$ in $A^{\tau\mu}$. Therefore,

$$M(w,\lambda,\mu) = \tilde{M}(w,\lambda,\mu)^{\dagger} = \langle \tilde{M}(w,\lambda,\mu), F_1(J_{\mu}(\lambda)), \dots, F_{\xi}(J_{\mu}(\lambda)) \rangle^{\dagger}.$$

Theorem 4.16 also implies that

$$\langle \tilde{M}(w,\lambda,\mu), F_1(J_\mu(\lambda)), \dots, F_{\xi}(J_\mu(\lambda)) \rangle = \operatorname{im} \pi(w, J_\mu(\lambda)) \subset A^{\tau\mu}.$$

On the other hand, applying Proposition 4.15 to the above resolution (4.5), we have a free resolution

$$0 \longrightarrow S^{3\tau\mu} \xrightarrow{\varphi(w', J_{\mu}(\lambda))} S^{3\tau\mu} \xrightarrow{\pi(w, J_{\mu}(\lambda))} \operatorname{im} \pi(w, J_{\mu}(\lambda))_{S} \longrightarrow 0$$

of im $\pi(w, J_{\mu}(\lambda))$ as an *S*-module. Finally, we know that $\varphi(w', J_{\mu}(\lambda))$ is a matrix factor of *xyz*, namely,

$$\rho\left(w', J_{\mu}\left(\lambda\right)\right)\psi\left(w', J_{\mu}\left(\lambda\right)\right) = x y z I_{3\tau\mu}$$

which implies that im $\pi(w, J_{\mu}(\lambda))$ is already maximal Cohen-Macaulay and hence equals $M(w, \lambda, \mu)$.

4.5. **Degenerate case.** Under the conversion formula, the degenerate band data (without multiplicity) $(w = (0,0,0)^{\tau}, \lambda = 1)$ and the degenerate loop data (without rank) $(w' = (2,2,2)^{\tau}, \lambda = 1)$ are converted to each other. Recall that we have the degenerate canonical form of matrix factorizations (Definition 3.24)

$$\left(\varphi_{\text{deg}}\left((2,2,2)^{\tau},1,\rho\right),\psi_{\text{deg}}\left((2,2,2)^{\tau},1,\rho\right)\right).$$

For $\tau \ge 2$, the normal loop word $(2,2,2)^{\tau}$ is periodic and the matrix factorization is decomposed into τ pieces, each of which corresponds to the non-periodic normal loop word (2,2,2) (Proposition 3.29). Among them, only one piece still has eigenvalue $\lambda = 1$. We compare it with the maximal Cohen-Macaulay module corresponding to the non-periodic degenerate band data ($w = (0,0,0), \lambda = 1, \mu$):

- the matrix factorization $(\varphi_{deg}((2,2,2),1,\rho), \psi_{deg}((2,2,2),1,\rho)),$
- the maximal Cohen-Macaulay module $M((0,0,0), 1, \mu)$.

It turns out that they correspond to each other under the relation $\rho = \mu - 1$:

Theorem 4.17. For a geometric rank $\rho \in \mathbb{Z}_{\geq 1}$ and algebraic multiplicity $\mu \in \mathbb{Z}_{\geq 2}$ with $\rho = \mu - 1$, we have

$$\operatorname{coker} \varphi_{\operatorname{deg}} ((2,2,2),1,\rho) \cong M((0,0,0),1,\mu)$$

in $\underline{CM}(A)$. For $\mu = 1$, the right side $M((0,0,0), 1, 1) \cong A^{32}$ is a zero object in $\underline{CM}(A)$.

Theorem 4.14 and Theorem 4.17 are combined to give Theorem 1.4 in the introduction. To prove Theorem 4.17, it is convenient to introduce a **reduced form** of our matrix factorization: We define

$$\begin{split} \left(\widetilde{\varphi_{\text{deg}}} \left((2,2,2),1,\rho \right), \widetilde{\psi_{\text{deg}}} \left((2,2,2),1,\rho \right) \right) \\ & := \left(\begin{pmatrix} -zx\mathbf{e}_{1}^{T} & 0 & 0 & 0 \\ zI_{\rho} & -yI_{\rho} & 0 & 0 \\ 0 & xI_{\rho} & -zI_{\rho} & 0 \\ -xJ_{\rho}(1) & 0 & yI_{\rho} & -xy\mathbf{e}_{\rho} \end{pmatrix}_{(3\rho+1)\times(3\rho+1)}, \begin{pmatrix} -y\mathbf{e}_{1} & -xyJ_{\rho}(0)^{T} & -yzJ_{\rho}(0)^{T} \\ -z\mathbf{e}_{1} & -zxJ_{\rho}(1)^{T} & -yzJ_{\rho}(0)^{T} & -z^{2}J_{\rho}(0)^{T} \\ -x\mathbf{e}_{1} & -x^{2}J_{\rho}(1)^{T} & -xyJ_{\rho}(0)^{T} \\ 0 & -x\mathbf{e}_{\rho}^{T} & -y\mathbf{e}_{\rho}^{T} & -z\mathbf{e}_{\rho}^{T} \end{pmatrix}_{(3\rho+1)\times(3\rho+1)} \end{split} \right)$$

³²See Remark 9.5 in [BD17a]. It also follows from our discussion below.

for $\rho \in \mathbb{Z}_{\geq 1}$, and

$$\left(\widetilde{\varphi_{\text{deg}}}((2,2,2),1,0),\widetilde{\psi_{\text{deg}}}((2,2,2),1,0)\right) := (-xyz,-1)$$

for $\rho = 0$.

Lemma 4.18. For $\rho \in \mathbb{Z}_{\geq 1}$, the degenerate canonical form $(\varphi_{deg}((2,2,2),1,\rho), \psi_{deg}((2,2,2),1,\rho))$ is isomorphic to its reduced form $(\widetilde{\varphi_{deg}}((2,2,2),1,\rho), \widetilde{\psi_{deg}}((2,2,2),1,\rho))$ in <u>MF</u>(xyz).

Proof. The submatrix $-I_{\tau\rho} + R_{\tau,\rho}(\lambda) = -I_{\rho} + J_{\rho}(1)$ of $\psi_{\text{deg}}((2,2,2),1,\rho)$ has $(\rho-1)$ units. So we can use Lemma 3.26 $(\rho-1)$ -times to reduce it to $(3\rho+1) \times (3\rho+1)$ size, the computation is straightforward.

Proof of Theorem 4.17. Let $\mu \in \mathbb{Z}_{\geq 1}$. We will find a free resolution

$$(4.6) 0 \longrightarrow S^{3\mu-2} \xrightarrow{\overline{\varphi_{\text{deg}}}((2,2,2),1,\mu-1)} S^{3\mu-2} \xrightarrow{\pi} M_S((0,0,0),1,\mu) \longrightarrow 0.$$

Then Eisenbud's equivalence and Lemma 4.18 complete the proof.

Note that

$$\tilde{M}((0,0,0),\lambda,\mu) = \operatorname{im}\tilde{\pi}((0,0,0),\lambda,\mu) = \operatorname{im}(zx^2J_{\mu}(\lambda) + x^2yI_{\mu}|(xy^2 + y^2z)I_{\mu}|(yz^2 + z^2x)I_{\mu})_A \subset A^{\mu}$$

any $\lambda \in \mathbb{C}^{\times}$ and $\mu \in \mathbb{Z}_{>1}$. It is easy to check that the sequence

for any $\lambda \in \mathbb{C}^{\times}$ and $\mu \in \mathbb{Z}_{\geq 1}$. It is easy to check that the sequence

$$S^{4} \xrightarrow{\begin{pmatrix} z & -y & 0 & 0 \\ 0 & x & -z & 0 \\ -\lambda x & 0 & y & -xy \end{pmatrix}} S^{3} \xrightarrow{(\lambda z x^{2} + x^{2} y | x y^{2} + y^{2} z | y z^{2} + z^{2} x)} \tilde{M}((0, 0, 0), \lambda, 1)_{S} \longrightarrow 0$$

is exact for any $\lambda \in \mathbb{C}^{\times}$. Then by Proposition 4.15,

$$S^{4\mu} \xrightarrow{\begin{pmatrix} zI_{\mu} & -yI_{\mu} & 0 & 0\\ 0 & xI_{\mu} & -zI_{\mu} & 0\\ -xJ_{\mu}(1) & 0 & yI_{\mu} & -xyI_{\mu} \end{pmatrix}} S^{3\mu} \xrightarrow{(zx^{2}J_{\mu}(1) + x^{2}yI_{\mu}|(xy^{2} + y^{2}z)I_{\mu}|(yz^{2} + z^{2}x)I_{\mu})} \tilde{M}((0,0,0),1,\mu)_{S} \longrightarrow 0$$

is also exact. Using

$$\begin{pmatrix} zI_{\mu} & -yI_{\mu} & 0\\ 0 & xI_{\mu} & -zI_{\mu}\\ -xJ_{\mu}(1) & 0 & yI_{\mu} \end{pmatrix} \begin{pmatrix} yI_{\mu}\\ zI_{\mu}\\ xI_{\mu} \end{pmatrix} = \begin{pmatrix} 0\\ 0\\ -xyJ_{\mu}(0) \end{pmatrix},$$

we can reduce it to the following, which is still exact:

$$S^{3\mu+1} \xrightarrow{\begin{pmatrix} zI_{\mu} & -yI_{\mu} & 0 & 0\\ 0 & xI_{\mu} & -zI_{\mu} & 0\\ -xJ_{\mu}(1) & 0 & yI_{\mu} & -xy\mathbf{e}_{\mu} \end{pmatrix}} S^{3\mu} \xrightarrow{(zx^{2}J_{\mu}(1)+x^{2}yI_{\mu}|(xy^{2}+y^{2}z)I_{\mu}|(yz^{2}+z^{2}x)I_{\mu})} \tilde{M}((0,0,0),1,\mu)_{S} \longrightarrow 0.$$

There is a Macaulayfying element $F := (xy + yz + zx)\mathbf{e}_1 := (xy + yz + zx, 0, ..., 0) \in A^{\tau}$ of $\tilde{M}((0,0,0), 1, \mu)$ in A^{τ} . So we enlarge it as $M_0((0,0,0), 1, \mu) := \langle F, \tilde{M}((0,0,0), 1, \mu) \rangle_A \subset A^{\tau}$, then

$$S^{3\mu+4} \xrightarrow{\begin{pmatrix} x & y & z & 0 & 0 & 0 & 0 \\ -\mathbf{e}_{1} & 0 & 0 & zI_{\mu} & -yI_{\mu} & 0 & 0 \\ 0 & -\mathbf{e}_{1} & 0 & 0 & xI_{\mu} & -zI_{\mu} & 0 \\ 0 & 0 & -\mathbf{e}_{1} & -xJ_{\mu}(1) & 0 & yI_{\mu} & -xy\mathbf{e}_{\mu} \end{pmatrix}}_{\underline{((xy+yz+zx)\mathbf{e}_{1}|zx^{2}J_{\mu}(1)+x^{2}yI_{\mu}|(xy^{2}+y^{2}z)I_{\mu}|(yz^{2}+z^{2}x)I_{\mu})}} M_{0}((0,0,0),1,\mu)_{S} \longrightarrow 0$$

is exact. (One can check it using Lemma 9.7 in [CJKR22].)

We can reduce matrices along three unit entries of the first matrix. (See Lemma Lemma 9.8 in [CJKR22].) As a result, for $\mu = 1$, we get a free resolution

 $0 \longrightarrow S \xrightarrow{-xyz} S \xrightarrow{xy+yz+zx} M_0((0,0,0),1,1)_S \longrightarrow 0$

and for $\mu \ge 2$, we have

$$0 \longrightarrow S^{3\mu-2} \xrightarrow{\begin{pmatrix} -zx\mathbf{e}_{1}^{T} & 0 & 0 & 0\\ zI_{\mu-1} & -yI_{\mu-1} & 0 & 0\\ 0 & xI_{\mu-1} & -zI_{\mu-1} & 0\\ -xJ_{\mu-1}(1) & 0 & yI_{\mu-1} & -xy\mathbf{e}_{\mu-1} \end{pmatrix}}_{\underbrace{\begin{pmatrix} xy+yz+zx & zx^{2}\mathbf{e}_{1}^{T} & 0 & 0\\ 0 & zx^{2}J_{\mu-1}(1)+x^{2}yI_{\mu-1} & (xy^{2}+y^{2}z)I_{\mu-1} & (yz^{2}+z^{2}x)I_{\mu-1} \end{pmatrix}}_{M_{0}((0,0,0),1,\mu)_{S}} \longrightarrow 0.$$

Note that the left matrix is $\widetilde{\varphi_{\text{deg}}}((2,2,2),1,\mu-1)$. As it is a matrix factor of xyz, the induced map $S^{3\mu-2} \rightarrow S^{3\mu-2}$ is injective and $M_0((0,0,0),1,\mu)$ is maximal Cohen-Macaulay, implying that it is the same as the Macaulayfication $M((0,0,0),1,\mu) = \tilde{M}((0,0,0),1,\mu)^{\dagger}$. So we achieved the desired free resolution (4.6) of $M((0,0,0),1,\mu)_S$ for any $\mu \ge 1$.

We finish this section with a remark on the **periodic cases**: We showed that the matrix factorizations corresponding to periodic loop data are decomposable (Theorem 3.21 for cylinder-free case and Proposition 3.29 for non-cylinder-free case). In non-degenerate cases, they are mapped to maximal Cohen-Macaulay modules corresponding to periodic band data (Theorem 4.14). It yields the decomposition

(4.7)
$$M(w,\lambda,\mu) \cong \bigoplus_{k=0}^{N-1} M(\tilde{w},\lambda_k,\mu)$$

in $\underline{CM}(A)$, where (w, λ, μ) is a non-degenerate band datum with a periodic band word $w = \tilde{w}^N$ for another band word \tilde{w} , and $\lambda_0, \ldots, \lambda_{N-1} \in \mathbb{C}^{\times}$ are the *N*-th roots of λ . In fact, an investigation in the category Tri(*A*) proves that the decomposition (4.7) is still valid in CM(*A*), even for non-degenerate band data.

Now something tricky happens in **periodic degenerate cases**: For a band datum $(w = (0, 0, 0)^{T}, \lambda = 1, \mu)$, the corresponding maximal Cohen-Macaulay module is decomposed as

$$M((0,0,0)^{\tau},1,\mu) \cong \bigoplus_{k=0}^{\tau-1} M((0,0,0),e^{2\pi i \cdot \frac{k}{\tau}},\mu).$$

Note that only the first direct summand is still degenerate, and the rank of its converted loop datum is shifted only in that piece. Namely, the corresponding matrix factorization and loop with a local system is (4.8)

$$\varphi_{\rm deg}\big((2,2,2),1,\mu-1\big) \oplus \bigoplus_{k=1}^{\tau-1} \varphi\big((2,2,2),e^{2\pi i \cdot \frac{k}{\tau}},\mu\big) \quad \text{and} \quad \mathscr{L}\big((2,2,2),-1,\mu-1\big) \oplus \bigoplus_{k=1}^{\tau-1} \mathscr{L}\big((2,2,2),-e^{2\pi i \cdot \frac{k}{\tau}},\mu\big).$$

Therefore, the object in $\underline{CM}(A)$ corresponding to a periodic degenerate band datum is mapped to objects in $\underline{MF}(xyz)$ or Fuk (Σ) that are decomposed into pieces having different geometric ranks.

Conversely, for a loop datum $(w' = (2,2,2)^{\tau}, \eta = -1, \rho)$ or $(w' = (2,2,2)^{\tau}, \lambda = 1, \rho)$, we observed in (3.6) and (3.5) the decomposition of the corresponding loop with a local system and matrix factorization. Now we know that they correspond to the decomposition of maximal Cohen-Macaulay module

(4.9)
$$M((0,0,0),1,\rho+1) \oplus \bigoplus_{k=1}^{\tau-1} M((0,0,0),e^{2\pi i \cdot \frac{k}{\tau}},\rho),$$

where only the first direct summand has a shifted multiplicity.

38

5. Applications

5.1. Flip of loops and dual of modules. Generally speaking, a symplectomorphism (diffeomorphism preserving the symplectic form) between symplectic manifolds induces an equivalence on their Fukaya categories. There are some obvious symmetries in our pair-of-pants surface Σ , each of which induces a corresponding auto-equivalence on Fuk(Σ), and hence on MF(xyz) and CM(A).

In this subsection, we take a look at the \mathbb{Z}_2 -symmetry given by flipping Σ back-and-forth, which is described in Example 5.7. It is given by an orientation-reversing diffeomorphism $\iota : \Sigma \to \Sigma$, which is an antisymplectomorphism ($\iota^* \omega = -\omega$). Such a map defines a natural *contravariant* A_{∞} -functor $\iota : Fuk(\Sigma) \to$ Fuk(Σ) (§5.1.1). We also define the *transpose functor* in MF_{A_{∞}}(f) as a contravariant A_{∞} -functor (§5.1.2), and show that, in our situation, two A_{∞} -functors are related under the localized mirror functor (§5.1.3). It is also related with the *duality functor* Hom_A(-, A) in <u>CM</u>(A) under Eisenbud's equivalence (§5.1.4). We also give a description of these operations in terms of loop/band data (§5.1.5).

5.1.1. Anti-symplectomorphism and contravariant A_{∞} -functor on Fukaya categories. We first recall some general algebraic notions following [Sei08]:

Definition 5.1. Given a \mathbb{Z}_2 -graded A_∞ -category \mathscr{A} , the **opposite** A_∞ -category \mathscr{A}^{op} consists of the same class of objects $\operatorname{Ob}(\mathscr{A}^{\operatorname{op}}) := \operatorname{Ob}(\mathscr{A})$, switched morphism spaces $\operatorname{hom}^{\bullet}_{\mathscr{A}^{\operatorname{op}}}(\mathscr{L}_0, \mathscr{L}_1) := \operatorname{hom}^{\bullet}_{\mathscr{A}}(\mathscr{L}_1, \mathscr{L}_0)$ ($\bullet \in \mathbb{Z}_2$), and A_∞ -operations $\{\mathfrak{m}_k^{\operatorname{op}}\}_{k>1}$ defined as

$$\mathfrak{m}_{k}^{\mathrm{op}}(g_{1},\ldots,g_{k}) := (-1)^{|g_{1}|+\cdots+|g_{k}|-k}\mathfrak{m}_{k}(g_{k},\ldots,g_{1})$$

for $g_i \in \hom_{\mathcal{A}^{\mathrm{op}}}^{\bullet}(\mathcal{L}_{i-1}, \mathcal{L}_i) = \hom_{\mathcal{A}}^{\bullet}(\mathcal{L}_i, \mathcal{L}_{i-1}) \ (i \in \{1, \dots, k\}, \bullet \in \mathbb{Z}_2).$

A straightforward calculation shows that \mathscr{A}^{op} is indeed an A_{∞} -category.

Definition 5.2. A contravariant A_{∞} -functor $\mathcal{G} := {\mathcal{G}_k}_{k\geq 0}$ between two A_{∞} -categories \mathcal{A} and \mathcal{B} is an A_{∞} -functor from $\mathcal{A}^{\mathrm{op}}$ to \mathcal{B} . Equivalently, it can be defined by giving a mapping

$$\mathscr{G}_0: \operatorname{Ob}(\mathscr{A}) \to \operatorname{Ob}(\mathscr{B})$$

and \Bbbk -linear maps ($k \ge 1$)

$$\mathscr{G}_k$$
: hom $_{\mathscr{A}}(\mathscr{L}_1, \mathscr{L}_0) \otimes \cdots \otimes \operatorname{hom}_{\mathscr{A}}(\mathscr{L}_k, \mathscr{L}_{k-1}) \to \operatorname{hom}_{\mathscr{B}}(\mathscr{G}_0(\mathscr{L}_0), \mathscr{G}_0(\mathscr{L}_k))$

of degree 1 - k, satisfying A_{∞} -relations

(5.1)
$$\sum_{1 \le k \le n} \sum_{1 \le i_1 < \dots < i_k = n} \mathfrak{m}_k^{\mathscr{B}} \left(\mathscr{G}_{i_1} \left(g_1, \dots, g_{i_1} \right), \dots, \mathscr{G}_{i_k} \left(g_{i_{k-1}+1}, \dots, g_n \right) \right) \\ = \sum_{0 \le i < j \le n} (-1)^{|g_1| + \dots + |g_j| - j} \mathscr{G}_{n-j+i+1} \left(g_1, \dots, g_i, \mathfrak{m}_{j-i}^{\mathscr{A}} \left(g_j, \dots, g_{i+1} \right), g_{j+1}, \dots, g_n \right)$$

for any fixed $n \in \mathbb{Z}_{\geq 1}$ and morphisms $g_i \in \hom_{\mathscr{A}}^{\bullet}(\mathscr{L}_i, \mathscr{L}_{i-1})$ $(i \in \{1, ..., n\}, \bullet \in \mathbb{Z}_2)$.

It induces an ordinary contravariant functor $H^0(\mathcal{G}): H^0(\mathcal{A}) \to H^0(\mathcal{B})$, whose mapping on objects is \mathcal{G}_0 and action on morphisms is given by $[g] \mapsto [\mathcal{G}_1(g)]$.

Now let (Σ, ω) and (Σ', ω') be 2-dimensional symplectic manifolds (possibly with boundary) of finite type (as in §A.2) and $\iota : \Sigma \to \Sigma'$ an anti-symplectomorphism $(\iota^* \omega' = -\omega)$. We will define a contravariant A_{∞} -functor $\iota := {\iota_k}_{k>0}$: Fuk $(\Sigma) \to \operatorname{Fuk}(\Sigma')$ as follows:

Any object $\mathscr{L} := (L, E, \nabla)$ of Fuk (Σ) consists of a loop $L : S^1 \to \Sigma$, a finite-rank \mathbb{C} -vector bundle E over S^1 , and a flat connection ∇ on E. We define its image under the functor ι as the triple

$$\iota_0(\mathscr{L}) := \left(\iota(L), E^*, \nabla^*\right),$$

where $\iota(L) := \iota \circ L : S^1 \to \Sigma'$ is the image of *L* under ι , E^* is the dual vector bundle of *E* over S^1 , and ∇^* is the dual connection of ∇ .

For two objects $\mathcal{L}_i := (L_i, E_i, \nabla_i)$ ($i \in \{0, 1\}$), note that there are bijections

$$\chi^{\bullet}(L_1, L_0) \stackrel{\text{1:1}}{\longleftrightarrow} \chi^{\bullet}(\iota(L_0), \iota(L_1)), \quad q \leftrightarrow \iota(q) \quad (\bullet \in \mathbb{Z}_2)$$

as shown in Figure 14 for $\bullet = 0$ case. Therefore, we have

$$\operatorname{hom}^{\bullet}(\iota(\mathscr{L}_{0}),\iota(\mathscr{L}_{1})) = \bigoplus_{q'\in\chi^{\bullet}(\iota(L_{0}),\iota(L_{1}))} \operatorname{Hom}_{\mathbb{C}}\left(E_{0}^{*}|_{q'}, E_{1}^{*}|_{q'}\right)$$
$$= \bigoplus_{q\in\chi^{\bullet}(L_{1},L_{0})} \operatorname{Hom}_{\mathbb{C}}\left(\left(E_{0}|_{q}\right)^{*}, \left(E_{1}|_{q}\right)^{*}\right) \quad (\bullet \in \mathbb{Z}_{2}).$$

We define ι_1 : hom $(\mathscr{L}_1, \mathscr{L}_0) \to$ hom $(\iota(\mathscr{L}_0), \iota(\mathscr{L}_1))$ by

(5.2)
$$g \in \operatorname{Hom}_{\mathbb{C}}\left(E_{1}|_{q}, E_{0}|_{q}\right) \mapsto (-1)^{\left|g\right|}g^{*} \in \operatorname{Hom}_{\mathbb{C}}\left(\left(E_{0}|_{q}\right)^{*}, \left(E_{1}|_{q}\right)^{*}\right)$$

for a base morphism *g* over $q \in \chi^{\bullet}(L_1, L_0)$ ($\bullet \in \mathbb{Z}_2$), and then linearly extend it to any morphisms. Higher components $\iota_{k \ge 2}$ are defined to be zero.





Figure 14. $q \in \chi^0(L_1, L_0), \iota(q) \in \chi^0(\iota(L_0), \iota(L_1))$

Figure 15. A polygon and its image under ι

Proposition 5.3. The functor ι : Fuk $(\Sigma) \rightarrow$ Fuk (Σ') defined above is indeed a contravariant A_{∞} -functor.

Proof. As $\iota_{k\geq 2} = 0$, the required A_{∞} -relations (5.1) simplify to

(5.3)
$$\mathfrak{m}_n(\iota_1(g_1),\ldots,\iota_1(g_n)) = (-1)^{|g_1|+\cdots+|g_n|-n}\iota_1(\mathfrak{m}_n(g_n,\ldots,g_1))$$

for $n \in \mathbb{Z}_{\geq 1}$ and $g_i \in \hom_{\operatorname{Fuk}(\Sigma)}^{\bullet} (\mathscr{L}_i, \mathscr{L}_{i-1})$ $(i \in \{1, \dots, n\}, \bullet \in \mathbb{Z}_2)$.

For
$$q_i \in \chi^{\bullet}(L_i, L_{i-1})$$
 and $g_i \in \operatorname{Hom}_{\mathbb{C}}(E_i|_{q_i}, E_{i-1}|_{q_i})$ $(i \in \{1, ..., n\}, \bullet \in \mathbb{Z}_2)$, the left side is
(5.4)
 $\mathfrak{m}_n\Big((-1)^{|g_1|}g_1^*, \dots, (-1)^{|g_n|}g_n^*\Big) = \sum_{r' \in \chi(\iota(L_0), \iota(L_n))} \sum_{u' \in \mathscr{M}\left[\iota(q_1), \dots, \iota(q_n), \overline{r'}\right]} (-1)^{|q_1| + \dots + |q_n|} \operatorname{sign}(u') \operatorname{hol}_{r'}(\partial u')(g_1^*, \dots, g_n^*)$

Note that there is a bijection between angles

$$\chi(L_n, L_0) \stackrel{1:1}{\longleftrightarrow} \chi(\iota(L_0), \iota(L_n)), \quad r \leftrightarrow \iota(r)$$

and between immersed polygons

$$\mathcal{M}(q_n,\ldots,q_1,\overline{r}) \stackrel{1:1}{\longleftrightarrow} \mathcal{M}(\iota(q_1),\ldots,\iota(q_n),\iota(\overline{r})), \quad u \leftrightarrow \iota(u) := \iota \circ u$$

as shown in Figure 19, for each $r \in \chi(L_n, L_0)$. Therefore, the right side of (5.4) is replaced by

(5.5)
$$\sum_{r \in \chi(L_n, L_0)} \sum_{u \in \mathcal{M}(q_n, \dots, q_1, \overline{r})} (-1)^{|q_1| + \dots + |q_n|} \operatorname{sign}(\iota(u)) \operatorname{hol}_{\iota(r)}(\partial(\iota(u)))(g_1^*, \dots, g_n^*).$$

For each pair of u and $\iota(u)$ in those sets, from the sign rule (A.6), we have

(5.6)

$$sign(u) sign(\iota(u)) = (-1)^{\wedge} \left(\sum_{i=1}^{n} |q_i| \mathbb{1}_{0(L_{i-1})\neq 0(\partial u)} + |r| \mathbb{1}_{0(L_0)\neq 0(\partial u)} + \sum_{i=1}^{n} |\iota(q_i)| \mathbb{1}_{0(\iota(L_i))\neq 0(\partial u)} + |\iota(r)| \mathbb{1}_{0(\iota(L_n))\neq 0(\partial u)} \right)$$
$$= (-1)^{\wedge} \left(\sum_{i=1}^{n} |q_i| (\mathbb{1}_{0(L_{i-1})\neq 0(\partial u)} - \mathbb{1}_{0(L_i)\neq 0(\partial u)} + 1) + |r| (\mathbb{1}_{0(L_0)\neq 0(\partial u)} - \mathbb{1}_{0(L_n)\neq 0(\partial u)} + 1) \right),$$

where we used the fact that $|q_i| = |\iota(q_i)|, |r| = |\iota(r)| \ (i \in \{1, ..., n\})$ and that the orientation of L_i coincides with that of ∂u if and only if the orientation of $\iota(L_i)$ differs from that of $\partial(\iota(u))$. Note also that

$$\left(\left|q_{i}\right|=1 \quad \Leftrightarrow \quad \mathbb{1}_{\mathsf{o}(L_{i-1})\neq\mathsf{o}(\partial u)}=\mathbb{1}_{\mathsf{o}(L_{i})\neq\mathsf{o}(\partial u)}\right) \quad \text{and} \quad \left(\left|r\right|=1 \quad \Leftrightarrow \quad \mathbb{1}_{\mathsf{o}(L_{0})\neq\mathsf{o}(\partial u)}\neq\mathbb{1}_{\mathsf{o}(L_{n})\neq\mathsf{o}(\partial u)}\right),$$

which reduce (5.6) to $(-1)^{\sum_{i=1}^{n} |q_i|}$.

On the other hand, $\operatorname{hol}_{\iota(r)}(\partial(\iota(u)))(g_1^*,\ldots,g_n^*)$ is by definition given as

$$P\left((\partial(\iota(u)))_0) \circ g_n^* \circ P\left((\partial(\iota(u)))_1\right) \circ g_{n-1}^* \circ \cdots \circ g_2^* \circ P\left((\partial(\iota(u)))_{n-1}\right) \circ g_1^* \circ P\left((\partial(\iota(u)))_n\right), g_1^* \circ P\left((\partial(\iota(u)))_n\right) \circ g_1^* \circ P\left((\partial(\iota(u)))_n\right), g_1^* \circ P\left((\partial(\iota(u)))_n\right) \circ g_1^* \circ P\left((\partial(\iota(u)))_n\right), g_1^* \circ P\left((\partial(\iota(u)))_n\right) \circ g_1^* \circ P\left($$

where each $P((\partial(\iota(u)))_i)$ is the parallel transport with respect to ∇_i^* from $E_i^*|_{\iota(q_i)}$ to $E_i^*|_{\iota(q_{i+1})}$, which is the dual $P((\partial u)_i)^*$ of the parallel transport $P((\partial u)_i)$ with respect to ∇_i from $E_i|_{q_{i+1}}$ to $E_i|_{q_i}$. Therefore, we can replace the total composition by $(\operatorname{hol}_r(\partial u)(g_n,\ldots,g_1))^*$.

Summing up, we can rewrite (5.5) as

$$\sum_{v \in \chi(L_n,L_0)} \sum_{u \in \mathcal{M}(q_n,\ldots,q_1,\overline{r})} \operatorname{sign}(u) \left(\operatorname{hol}_r(\partial u) \left(g_n,\ldots,g_1 \right) \right)^*,$$

whose degree is $|r| = 2 - n + |q_1| + \dots + |q_n|$. Therefore, it is the same as $(-1)^{|g_1| + \dots + |g_n| - n} \iota_1(\mathfrak{m}_n(g_n, \dots, g_1))$, or the right side of (5.3).

5.1.2. *Transpose functor on* $MF_{A_{\infty}}(f)$. Let *S* be the power series ring $\mathbb{C}[[x_1, ..., x_m]]$ of *m* variables, and $f \in S$ its nonzero element. Taking transpose of matrix factorizations of *f* gives rise to a contravariant A_{∞} -functor $-\operatorname{Tr}: MF_{A_{\infty}}(f) \to MF_{A_{\infty}}(f)$, called the (**minus**) **transpose functor** ³³, which we now define:

For an object $P^0 \xleftarrow{\varphi}{\psi} P^1$ with free *S*-modules P^0 , P^1 , we define its image as $(P^1)^* \xleftarrow{-\varphi^*}{-\varphi^*} (P^0)^*$, where P^* denotes the *S*-dual Hom_{*S*}(*P*,*S*) of an *S*-module *P* and φ^* , ψ^* denote the natural pull-back maps. It defines the functor $-\text{Tr} := \{-\text{Tr}_k\}_{k>0}$ on the object level as

$$-\operatorname{Tr}_{0}:\operatorname{Ob}\left(\operatorname{MF}_{A_{\infty}}(f)\right)\to\operatorname{Ob}\left(\operatorname{MF}_{A_{\infty}}(f)\right),\quad\left(\varphi,\psi\right)\to\left(-\varphi^{*},-\psi^{*}\right).$$

Given two matrix factorizations $P_0^0 \stackrel{\varphi_0}{\longleftrightarrow} P_0^1$ and $P_1^0 \stackrel{\varphi_1}{\longleftrightarrow} P_1^1$ of f and an even-degree morphism $(\alpha : P_0^0 \to P_1^0, \beta : P_0^1 \to P_1^1)$ (resp. an odd-degree morphism $(\gamma : P_0^0 \to P_1^1, \delta : P_0^1 \to P_1^0)$) (see diagrams in (A.9)), we take dual of the maps to define its image under $-\operatorname{Tr}_1$:

$$-\operatorname{Tr}_{1}: \operatorname{hom}\left(\left(\varphi_{0}, \psi_{0}\right), \left(\varphi_{1}, \psi_{1}\right)\right) \to \operatorname{hom}\left(\left(\varphi_{1}^{*}, \psi_{1}^{*}\right), \left(\varphi_{0}^{*}, \psi_{0}^{*}\right)\right), \quad \left(\alpha, \beta\right) \mapsto \left(\beta^{*}, \alpha^{*}\right) \quad \left(\operatorname{resp.}\left(\gamma, \delta\right) \mapsto \left(\delta^{*}, \gamma^{*}\right)\right).$$

The higher components $\operatorname{Tr}_{k\geq 2}$ are defined to be zero. It is straightforward to check that Tr is a contravariant A_{∞} -functor.

³³We put the minus sign here to match with the *flip functor* in the next subsection. However, the functor with a minus sign and one without a minus sign are (A_{∞}) -*quasi-isomorphic* to each other.

5.1.3. *Flip of loops and transpose of matrix factorizations.* Coming back to our specific situation, the antisymplectomorphism $\iota: \Sigma \to \Sigma$ described in Example 5.7 and the discussion so far yield the diagram:

(5.7)
$$Fuk(\Sigma) \xrightarrow{\mathscr{F}^{L}} MF_{A_{\infty}}(xyz)$$
$$flip \downarrow \iota \quad transpose \downarrow -Tr$$
$$Fuk(\Sigma) \xrightarrow{\mathscr{F}^{L}} MF_{A_{\infty}}(xyz)$$

In this subsection, we will show that the diagram commutes, in the sense that two A_{∞} -functors $\mathscr{F}^{\mathbb{L}} \circ \iota$ and $-\operatorname{Tr} \circ \mathscr{F}^{\mathbb{L}}$ are *quasi-isomorphic*³⁴ to each other.

It is based on the fact that our reference object \mathbb{L} (Seidel Lagrangian) is invariant under the flipping functor *i*. Namely, Figure 16 shows that $i(\mathbb{L})$ consists of the same underlying loop with \mathbb{L} and its local system is gauge equivalent to that of \mathbb{L} . Moreover, one can easily check (as we did in Proposition 2.1) that

$$-\iota(b) = -x\iota(X) - y\iota(Y) - z\iota(Z) \in \hom^{1}(\iota(\mathbb{L}), \iota(\mathbb{L}))$$

is a weak bounding cochain with the disk potential $W^{\iota(\mathbb{L})} = xyz$. Therefore, the pair $(\iota(\mathbb{L}), -\iota(b))$ defines a localized mirror functor $\mathscr{F}^{\iota(\mathbb{L})}$: Fuk $(\Sigma) \to MF_{A_{\infty}}(xyz)$.





Figure 16. Seidel Lagrangian and its image under *i*

Figure 17. A deformed strip and its image under ι

Now we have two localized mirror functors $\mathscr{F}^{\mathbb{L}}$ and $\mathscr{F}^{\iota(\mathbb{L})}$, but there is a trivial isomorphism between (\mathbb{L}, b) and $(\iota(\mathbb{L}), -\iota(b))$, and it has been already proven (with much greater generality) in [CHL18] that such isomorphic weak bounding cochains induce quasi-isomorphic A_{∞} -functors:

Proposition 5.4. [CHL18, Theorem 4.7.(2)] *Two localized mirror functors* $\mathscr{F}^{\mathbb{L}}$ *and* $\mathscr{F}^{\iota(\mathbb{L})}$ *are quasi-isomorphic.*

To show that the diagram (5.7) commutes, therefore, it is enough to check the following alternative:

Proposition 5.5. Two A_{∞} -functors $\mathscr{F}^{\iota(\mathbb{L})} \circ \iota$ and $-\operatorname{Tr} \circ \mathscr{F}^{\mathbb{L}}$ are the same.

Proof. Recall from §2.2 that $\mathscr{F}^{\mathbb{L}}(\mathscr{L}) = (\Phi^{\mathbb{L}}(\mathscr{L}), \Psi^{\mathbb{L}}(\mathscr{L}))$ is given as maps

$$\hom^{0}(\mathscr{L},\mathbb{L}) = \bigoplus_{p \in \chi^{0}(L,\mathbb{L})} (E|_{p})^{*} \xrightarrow{\Phi^{\mathbb{L}}(\mathscr{L}) = \mathfrak{m}_{1}^{0,b}} \bigoplus_{s \in \chi^{1}(L,\mathbb{L})} (E|_{s})^{*} = \hom^{1}(\mathscr{L},\mathbb{L}),$$

whose $((E|_s)^*, (E|_p)^*)$ -component $\mathfrak{m}_1^{0,b}: (E|_p)^* \to (E|_s)^*$ for each $p, s \in \chi(L, \mathbb{L})$ is

$$\sum_{\substack{(x_1,X_1),\ldots,(x_i,X_i)\\ \in \{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p,X_1,\ldots,X_i,\overline{s})} (-1)^{(i+1)\mathbb{I}_{0(\mathbb{L}) \neq o(\partial u)} + \#(\partial u \cap \bigstar_{\mathbb{L}})} P\left((\partial u)_0\right)^*,$$

³⁴Two A_{∞} -functors are *quasi-isomorphic* to each other if there are A_{∞} -natural transformations between them satisfying some homotopy conditions. Any A_{∞} -natural transformation induces an (ordinary) natural transformation between the induced ordinary functors. See [Sei08] for the details.

where $P((\partial u)_0) \in \operatorname{Hom}_{\mathbb{C}}(E|_s, E|_p)$ is the parallel transport from $E|_s$ to $E|_p$ along the side of u lying in L. Taking its dual yields the map $-\operatorname{Tr}(\mathscr{F}^{\mathbb{L}}(\mathscr{L})) = (-\Phi^{\mathbb{L}}(\mathscr{L})^*, -\Psi^{\mathbb{L}}(\mathscr{L})^*)$ given by

$$\left(\hom^{1}(\mathscr{L},\mathbb{L})\right)^{*} = \bigoplus_{s \in \chi^{1}(L,\mathbb{L})} E|_{s} \xrightarrow{-\Phi^{\mathbb{L}}(\mathscr{L})^{*}} \bigoplus_{p \in \chi^{0}(L,\mathbb{L})} E|_{p} = \left(\hom^{0}(\mathscr{L},\mathbb{L})\right)^{*},$$

whose $(E|_p, E|_s)$ -component $E|_s \to E|_p$ for each $p, s \in \chi(L, \mathbb{L})$ is

$$-\sum_{\substack{(x_1,X_1),\ldots,(x_i,X_i)\\\in\{(x,X),(y,Y),(z,Z)\}}} x_1\cdots x_i \sum_{u\in\mathcal{M}(p,X_1,\ldots,X_i,\overline{s})} (-1)^{(i+1)\mathbb{I}_{0(\mathbb{L})\neq 0(\partial u)}+\#(\partial u\cap \bigstar_{\mathbb{L}})} P((\partial u)_0)$$

On the other hand, the opposite side $\mathscr{F}^{\iota(\mathbb{L})}(\iota(\mathscr{L})) = (\Phi^{\iota(\mathbb{L})}(\iota(\mathscr{L})), \Psi^{\iota(\mathbb{L})}(\iota(\mathscr{L})))$ is given by (5.8)

$$\hom^{0}(\iota(\mathscr{L}),\iota(\mathbb{L})) = \bigoplus_{p' \in \chi^{0}(\iota(L),\iota(\mathbb{L}))} \left(E^{*} \big|_{p'} \right)^{*} \xrightarrow{\Phi^{\iota(\mathbb{L})}(\iota(\mathscr{L})) = \mathfrak{m}_{1}^{0,-\iota(b)}} \bigoplus_{s' \in \chi^{1}(\iota(L),\iota(\mathbb{L}))} \left(E^{*} \big|_{s'} \right)^{*} = \hom^{1}(\iota(\mathscr{L}),\iota(\mathbb{L})), \iota(\mathbb{L})$$

whose $((E^*|_{s'})^*, (E^*|_{p'})^*)$ -component $\mathfrak{m}_1^{0,-\iota(b)} : (E^*|_{p'})^* \to (E^*|_{s'})^*$ for each $p', s' \in \chi(\iota(L), \iota(\mathbb{L}))$ is (5.9) $\sum_{\substack{(x'_1, x'_1), \dots, (x'_i, x'_i) \\ \in \{(-x, \iota(X)), (-y, \iota(Y)), (-z, \iota(Z))\}}} x'_1 \cdots x'_i \sum_{u' \in \mathcal{M}(p', X'_1, \dots, X'_i, \overline{s'})} (-1)^{(i+1)\mathbb{I}_{0(\iota(\mathbb{L})) \neq 0}(\partial u') + \#(\partial u' \cap \iota(\bigstar_{\mathbb{L}}))} P((\partial u')_0)^*.$

Under the bijection between angles

$$\chi^{\bullet}(L,\mathbb{L}) \stackrel{1:1}{\longleftrightarrow} \chi^{\bullet+1}(\iota(L),\iota(\mathbb{L})), \quad p \leftrightarrow \iota(\overline{p}) \quad (\bullet \in \mathbb{Z}_2),$$

we can put $p' = \iota(\overline{s})$ and $s' = \iota(\overline{p})$ for some $p, s \in \chi(L, \mathbb{L})$. Each X'_i is $\iota(X_i)$ for some $X_i \in \{X, Y, Z\}$. There is also a bijection between deformed strips

$$\mathcal{M}\left(p, X_{i}, \dots, X_{1}, \overline{s}\right) \stackrel{1:1}{\longleftrightarrow} \mathcal{M}\left(\iota\left(\overline{s}\right), \iota\left(X_{1}\right), \dots, \iota\left(X_{i}\right), \iota\left(p\right)\right), \quad u \leftrightarrow \iota\left(u\right) := \iota \circ u$$

as shown in Figure 17. Also using the identifications $\left(E^*|_{\iota(\overline{s})}\right)^* = E|_s$ and $\left(E^*|_{\iota(\overline{p})}\right)^* = E|_p$, we can rewrite (5.16) and (5.9) as

$$\bigoplus_{s \in \chi^1(L,\mathbb{L})} E|_s \xrightarrow{\Phi^{\iota(\mathbb{L})}(\iota(\mathscr{L})) = \mathfrak{m}_1^{0,-\iota(b)}} \bigoplus_{p \in \chi^0(L,\mathbb{L})} E|_p$$

where the $(E|_p, E|_s)$ -component $E|_s \rightarrow E|_p$ is

(5.10)
$$\sum_{\substack{(x_1,X_1),\dots,(x_i,X_i)\\ \in\{(x,X),(y,Y),(z,Z)\}}} (-x_1)\cdots(-x_i) \sum_{u\in\mathcal{M}(p,X_i,\dots,X_1,\overline{s})} (-1)^{(i+1)\mathbb{I}_{0(i(\mathbb{L}))\neq 0(\partial(i(u)))} + \#(\partial(i(u))\cap i(\bigstar_{\mathbb{L}}))} P((\partial(i(u)))_0)^*$$

The obvious relations

 $\mathbb{1}_{\mathrm{o}(\mathbb{L})\neq\mathrm{o}(\partial u)} + \mathbb{1}_{\mathrm{o}(\iota(\mathbb{L}))\neq\mathrm{o}(\partial(\iota(u)))} = 1 \quad \text{and} \quad \# \big(\partial u \cap \bigstar_{\mathbb{L}}\big) = \# \big(\partial(\iota(u)) \cap \iota \big(\bigstar_{\mathbb{L}}\big)\big)$

and the fact that $P((\partial(\iota(u)))_0) \in \text{Hom}_{\mathbb{C}}(E^*|_p, E^*|_s)$ is the dual of $P((\partial u)_0) \in \text{Hom}_{\mathbb{C}}(E|_s, E|_p)$ replace (5.10) again into

$$-\sum_{\substack{(x_1,X_1),\ldots,(x_i,X_i)\\\in\{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p,X_i,\ldots,X_1,\overline{s})} (-1)^{(i+1)\mathbb{I}_{o(\mathbb{L})\neq o(\partial u)} + \#(\partial u \cap \bigstar_{\mathbb{L}})} P((\partial u)_0).$$

This shows that two functors are the same on the object level. It is also straightforward to check that they coincide on the morphism level. $\hfill \Box$

5.1.4. *Flip of loops and dual of modules.* The commutativity of diagram (5.7) induces the commutativity of the left square in the following diagram of ordinary categories and functors:

(5.11)
$$\begin{array}{ccc} H^{0}\operatorname{Fuk}(\Sigma) & \xrightarrow{\mathscr{F}^{\mathbb{L}}} & \underline{\mathrm{MF}}(xyz) & \xrightarrow{\mathrm{coker}} & \underline{\mathrm{CM}}(A) \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & \\ &$$

We check the commutativity of the right square in the general setting:

Proposition 5.6. Let S be the power series ring $\mathbb{C}[[x_1,...,x_m]]$ of m variables, $f \in S$ its nonzero element and A := S/(f) the quotient ring. Then the following diagram is commutative, that is, two compositions of functors are naturally isomorphic to each other:

$$\underbrace{\mathrm{MF}}_{\simeq}(f) \xrightarrow{\operatorname{coker}} \underline{\mathrm{CM}}(A)$$
transpose $-\operatorname{Tr}$ $\operatorname{dual}_{\operatorname{Hom}_{A}(-,A)}$

$$\underbrace{\mathrm{MF}}_{\simeq}(f) \xrightarrow{\simeq} \underline{\mathrm{CM}}(A)$$

Proof. Recall that under Eisenbud's equivalence (Theorem A.16), a matrix factorization $P^0 \stackrel{\varphi}{\underset{\psi}{\longrightarrow}} P^1$ of f corresponds to a maximal Cohen-Macaulay *A*-module $M := \operatorname{coker} \underline{\varphi}$, which admits a 2-periodic free resolution given by

$$\cdots \longrightarrow P^0 \otimes_S A \xrightarrow{\underline{\varphi}} P^1 \otimes_S A \xrightarrow{\underline{\psi}} P^0 \otimes_S A \xrightarrow{\underline{\varphi}} P^1 \otimes_S A \longrightarrow M \longrightarrow 0.$$

Taking Hom_A (-, A) yields a 2-periodic free resolution of Hom_A (M, A) as

$$0 \longrightarrow \operatorname{Hom}_{A}(M, A) \longrightarrow \operatorname{Hom}_{S}(P^{1}, A) \xrightarrow{\varphi^{*}} \operatorname{Hom}_{A}(P^{0}, A) \xrightarrow{\psi^{*}} \operatorname{Hom}_{A}(P^{1}, A) \longrightarrow \cdots,$$

where $\underline{\varphi^*} = \varphi^* \otimes \mathrm{id}_A : (P^1)^* \otimes_S A \to (P^0)^* \otimes_S A$ is induced from $\varphi^* : (P^1)^* \to (P^0)^*$. It gives natural isomorphisms

$$\operatorname{Hom}_{A}(M, A) \cong \operatorname{ker} \underline{\varphi^{*}} = \operatorname{im} \underline{\psi^{*}} \cong \operatorname{Hom}_{A}(P^{0}, A) / \operatorname{ker} \underline{\psi^{*}} = \operatorname{Hom}_{A}(P^{0}, A) / \operatorname{im} \underline{\varphi^{*}} = \operatorname{coker} \underline{\varphi^{*}} = \operatorname{coker} (\underline{-\varphi^{*}}).$$

5.1.5. *Correspondence of canonical forms*. Flipping a loop with a local system, taking transpose of a matrix factorization, and taking dual of a maximal Cohen-Macaulay modules can all be given explicitly in terms of loop/band data.

In Fuk (Σ), let (L, E, ∇) := ℒ(w', η, ρ) be the loop with a local system corresponding to a loop datum (w', η, ρ). Its flip is given by (ι(L), E*, ∇*). Recall that the free homotopy class of the loop L(w') corresponding to the given normal loop word w' = (l'₁, m'₁, n'₁,..., l'_τ, m'_τ, n'_τ) ∈ Z^{3τ} is given by

$$[L(w')] = \left[\alpha^{l'_1}\beta^{m'_1}\gamma^{n'_1}\cdots\alpha^{l'_{\tau}}\beta^{m'_{\tau}}\gamma^{n'_{\tau}}\right],$$

where α , β , γ are generators of $\pi_1(\Sigma)$ (Figure 8a). It is easy to see that the free homotopy class of the flipped loop $\iota(L(w'))$ is

$$[\iota(L(w'))] = \left[\alpha^{1-l'_1}\beta^{1-m'_1}\gamma^{1-n'_1}\cdots\alpha^{1-l'_r}\beta^{1-m'_r}\gamma^{1-n'_r}\right].$$

Defining a new normal loop word 1 - w' as the normal form of the loop word ³⁵

$$(1 - l'_1, 1 - m'_1, 1 - n'_1, \dots, 1 - l'_{\tau}, 1 - m'_{\tau}, 1 - n'_{\tau}) \in \mathbb{Z}^{3\tau},$$

two loops $\iota(L(w'))$ and L(1-w') are freely homotopic to each other.

The holonomy of (E, ∇) at some point is represented by a matrix $J_{\rho}(\eta)$ (up to conjugacy). It changes to $(J_{\rho}(\eta)^{T})^{-1}$ (which is similar to $J_{\rho}(\eta)^{-1}$ by discussion in the proof of Theorem 3.17) for the dual local system (E^*, ∇^*) . This is because the parallel transport from $E^*|_p$ to $E^*|_q$ is $(P(L_{p \to q})^*)^{-1}$, where $P(L_{p \to q})$ is the parallel transport in (E, ∇) from $E|_p$ to $E|_q$. Thus, the flipped loop with a local system $(\iota(L), E^*, \nabla^*)$ is isomorphic to the canonical form

Thus, the flipped loop with a local system $(\iota(L), E^*, \nabla^*)$ is isomorphic to the canonical form $\mathscr{L}(1-w', \eta^{-1}, \rho)$, that is, they give isomorphic matrix factorizations in <u>MF</u>(*xyz*) by Theorem 3.13.

- In <u>MF</u>(*xyz*) (or MF(*xyz*)), consider the canonical form $\varphi(w', \lambda, \rho)$ corresponding to a loop datum (w', λ, ρ) . Its transpose still remains in the canonical form up to equivalence of loop words and bases change $J_{\rho}(\lambda^{-1})^{T} \sim J_{\rho}(\lambda^{-1}) \sim J_{\rho}(\lambda)^{-1}$, which results in the canonical form $\varphi(1 w', \lambda^{-1}, \rho)$.
- In <u>CM</u>(*A*) (or CM(*A*)), the canonical form given in terms of Definition 4.6 does not directly show the relation with taking dual. However, in Tri(*A*), one can handle it in an algebraic way and see that the dual of the canonical form $M(w, \lambda, \mu)$ corresponding to a band datum (w, λ, μ) is isomorphic to $M(-w, \lambda^{-1}, \mu)$, where -w is the band word given by multiplying -1 to every entry of w.

The above discussions summarize to the following mappings (up to isomorphism) under the diagram (5.11), while two rows are consistent with our main correspondence (1.5):

$$\begin{aligned} \mathscr{L}(w',\eta,\rho) & \longmapsto^{\mathscr{F}^{\mathbb{L}}} \varphi_{(\operatorname{deg})}(w',\lambda,\rho) & \stackrel{\operatorname{coker}}{\longmapsto} M(w,\lambda,\mu) \\ & \operatorname{flip} \downarrow^{\iota} & \operatorname{transpose} \uparrow^{-\operatorname{Tr}} & \operatorname{dual} \uparrow^{\operatorname{Hom}_{A}(-,A)} \\ \mathscr{L}(1-w',\eta^{-1},\rho) & \stackrel{\mathscr{F}^{\mathbb{L}}}{\longmapsto} \varphi_{(\operatorname{deg})}(1-w',\lambda^{-1},\rho) & \stackrel{\simeq}{\longmapsto} M(-w,\lambda^{-1},\mu) \end{aligned}$$

Example 5.7. The following shows the correspondence of loops with a local system $\mathcal{L}((3, -2, 2), \eta, 1) \leftrightarrow \mathcal{L}((-2, 3, -1), \eta^{-1}, 1)$, matrix factorizations $\varphi((3, -2, 2), \lambda, 1) \leftrightarrow \varphi((-2, 3, -1), \lambda^{-1}, 1)$, and maximal Cohen-Macaulay modules $M((2, -3, 1), \lambda, 1) \leftrightarrow M((-2, 3, -1), \lambda^{-1}, 1)$ (in Tri(A)) up to isomorphism, where $\lambda = -\eta$.



³⁵It is not normal a priori in general, but Proposition 3.6 ensures that one can deform it to the unique normal loop word by performing five operations in Lemma 3.3 finitely many times.

5.2. **Reverse of loops and shift of modules.** Orientation-reversing of loops induces another auto-equivalence on Fuk (Σ) (§5.2.1). We also define the *switching functor* in MF_{A_∞}(f) (§5.2.2), and show that, in our situation, two A_∞-functors are related under the localized mirror functor (§5.2.3). They boil down to shift functors of triangulated categories, and therefore, are also related with the shift functor of <u>CM</u>(A) (§5.2.4). We give an algorithm to compute them in terms of loop/band data (§5.2.5).

5.2.1. Orientation-reversing of loops and auto-equivalence on the Fukaya category. Let (Σ, ω) be a 2-dimensional symplectic manifolds (possibly with boundary) of finite type (as in §A.2). There is an obvious symmetry of objects in Fuk (Σ) , namely, we can reverse the orientation of the underlying loop of every object. It results in an auto-equivalence on Fuk (Σ) given by a (covariant) A_{∞} -functor $J := \{J_k\}_{k\geq 0}$: Fuk $(\Sigma) \to$ Fuk (Σ) we define now:

Any object $\mathscr{L} := (L, E, \nabla)$ of Fuk (Σ) consists of a loop $L : S^1 \to \Sigma$, a finite-rank \mathbb{C} -vector bundle $E \to S^1$, and a flat connection ∇ on E. We define its image under the functor J as the triple

$$J_0(\mathscr{L}) := (J(L)) := L \circ \kappa, \kappa^* E, \kappa^* \nabla)$$

where $\kappa : S^1 \to S^1$, $e^{2\pi it} \mapsto e^{-2\pi it}$ denotes the orientation-reversing map.

Note that reversing the orientation of loops doesn't affect their (self-)intersections. Therefore, for two objects $\mathcal{L}_i := (L_i, E_i, \nabla_i)$ ($i \in \{0, 1\}$), the sets $\chi^{\bullet}(L_0, L_1)$ and $\chi^{\bullet}(J(L_0), J(L_1))$ ($\bullet \in \mathbb{Z}_2$) are identified, as shown in Figure 18. The fibers $E_i|_p$ and $\kappa^* E_i|_p$ over the preimages (in S^1) of the point $p \in \Sigma$ under L_i and $J(L_i)$, respectively, are also naturally identified. So there is also a natural identification between

$$\bigoplus_{p \in \chi^{\bullet}(L_0, L_1)} \operatorname{Hom}_{\mathbb{C}}\left(E_0|_p, E_1|_p\right) \text{ and } \bigoplus_{p \in \chi^{\bullet}\left(j(L_0), j(L_1)\right)} \operatorname{Hom}_{\mathbb{C}}\left(\kappa^* E_0|_p, \kappa^* E_1|_p\right),$$

We define j_1 : hom $(\mathscr{L}_0, \mathscr{L}_1) \to \text{hom}(j(\mathscr{L}_0), j(\mathscr{L}_1))$ by

(5.12)
$$f \in \operatorname{Hom}_{\mathbb{C}}(E_0|_p, E_1|_p) \mapsto (-1)^{|f|} f \in \operatorname{Hom}_{\mathbb{C}}(\kappa^* E_0|_p, \kappa^* E_1|_p)$$

for a base morphism f over $p \in \chi^{\bullet}(L_0, L_1)$ ($\bullet \in \mathbb{Z}_2$), and then linearly extend it to any morphisms. Higher components $J_{k\geq 2}$ are defined to be zero.



Figure 18. $p, \overline{p} \in \chi(L_0, L_1) = \chi(J(\mathcal{L}_0), J(\mathcal{L}_1))$



Proposition 5.8. The functor j: Fuk $(\Sigma) \rightarrow$ Fuk (Σ) defined above is indeed a covariant A_{∞} -functor.

Proof. As $J_{k\geq 2} = 0$, the required A_{∞} -relations (A.3) simplify to

(5.13)
$$\mathfrak{m}_n(J_1(f_1),\ldots,J_1(f_n)) = J_1(\mathfrak{m}_n(f_1,\ldots,f_n))$$

for $n \in \mathbb{Z}_{\geq 1}$ and $f_i \in \text{hom}^{\bullet}(\mathscr{L}_{i-1}, \mathscr{L}_i)$ ($i \in \{1, ..., n\}$, $\bullet \in \mathbb{Z}_2$). For $p_i \in \chi^{\bullet}(L_{i-1}, L_i)$ and $f_i \in \text{Hom}_{\mathbb{C}}(E_{i-1}|_{p_i}, E_i|_{p_i})$ ($i \in \{1, ..., n\}$, $\bullet \in \mathbb{Z}_2$), recalling the definition of \mathfrak{m}_n in (A.5), both the left and right sides are contributed by the same polygons $u \in \mathcal{M}(p_1, ..., p_k, \overline{q})$ for $q \in \chi(L_0, L_k)$. But for each polygon u, its bounding loops have different orientations in both sides. It plays a role only when we compute sign(u), whose quotient in both sides is given by $(-1)^{\sum_{i=1}^k |p_i| + |q|}$ following the sign rule (A.6). It cancels all the signs that occur when taking J_1 , which confirms (5.13). 5.2.2. Switching functor on $MF_{A_{\infty}}(f)$. Let *S* be the power series ring $\mathbb{C}[[x_1,...,x_m]]$ of *m* variables, and $f \in S$ its nonzero element. Switching two factors in matrix factorizations of *f* gives rise to a covariant A_{∞} -functor [1]: $MF_{A_{\infty}}(f) \to MF_{A_{\infty}}(f)$, called the **switching functor** ³⁶, which we now define:

For an object $P^0 \xleftarrow{\varphi}{\psi} P^1$ with free *S*-modules P^0 , P^1 , we define its image as $P^1 \xleftarrow{\psi}{\varphi} P^0$. It defines the functor $[1] := \{[1]_k\}_{k\geq 0}$ on the object level as

$$[1]_0: \operatorname{Ob}\left(\operatorname{MF}_{A_\infty}(f)\right) \to \operatorname{Ob}\left(\operatorname{MF}_{A_\infty}(f)\right), \quad \left(\varphi,\psi\right) \to \left(\psi,\varphi\right).$$

Given two matrix factorizations $P_1^0 \stackrel{\varphi_1}{\longleftrightarrow_1} P_1^1$ and $P_0^0 \stackrel{\varphi_0}{\longleftrightarrow_0} P_0^1$ of f and an even-degree morphism $(\alpha : P_1^0 \to P_0^0, \beta : P_1^1 \to P_0^1)$ (resp. an odd-degree morphism $(\gamma : P_1^0 \to P_0^1, \delta : P_1^1 \to P_0^0)$), we define its image under [1]₁ as

 $[1]_1: \hom\left((\varphi_1, \psi_1), (\varphi_0, \psi_0)\right) \to \hom\left((\psi_1, \varphi_1), (\psi_0, \varphi_0)\right), \quad (\alpha, \beta) \mapsto (\beta, \alpha) \quad (\text{resp. } (\gamma, \delta) \mapsto (\delta, \gamma)).$

The higher components $[1]_{k\geq 2}$ are defined to be zero. Then it is straightforward to check that [1] is a covariant A_{∞} -functor.

5.2.3. Orientation-reversing of loops and switching of matrix factorizations. Now in our situation, we have the diagram of A_{∞} -categories and functors:

(5.14)
$$Fuk(\Sigma) \xrightarrow{\mathscr{F}^{L}} MF_{A_{\infty}}(xyz)$$
$$\underset{\text{-reversing}}{\text{orientation}} \int_{J} \underset{\text{two factors}}{\text{switching}} \int_{[1]} [1]$$
$$Fuk(\Sigma) \xrightarrow{\mathscr{F}^{L}} MF_{A_{\infty}}(xyz)$$

In this subsection, we will show that the diagram commutes, more precisely:

Proposition 5.9. Two A_{∞} -functors $\mathscr{F}^{\mathbb{L}} \circ J$ and $[1] \circ \mathscr{F}^{\mathbb{L}}$ are the same.

Proof. Recall from §2.2 that $\mathscr{F}^{\mathbb{L}}(\mathscr{L}) = (\Phi^{\mathbb{L}}(\mathscr{L}), \Psi^{\mathbb{L}}(\mathscr{L}))$ is given as maps

(5.15)
$$\operatorname{hom}^{0}(\mathscr{L},\mathbb{L}) = \bigoplus_{p \in \chi^{0}(L,\mathbb{L})} (E|_{p})^{*} \xrightarrow{\Phi^{\mathbb{L}}(\mathscr{L}) = \mathfrak{m}_{1}^{0,b}} \bigoplus_{s \in \chi^{1}(L,\mathbb{L})} (E|_{s})^{*} = \operatorname{hom}^{1}(\mathscr{L},\mathbb{L}),$$

whose $((E|_s)^*, (E|_p)^*)$ -component $\mathfrak{m}_1^{0,b}: (E|_p)^* \to (E|_s)^*$ for each $p, s \in \chi(L, \mathbb{L})$ is

$$\sum_{\substack{(x_1,X_1),\ldots,(x_i,X_i)\\ \in \{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p,X_1,\ldots,X_i,\overline{s})} (-1)^{(i+1)\mathbb{I}_{o(\mathbb{L}) \neq o(\partial u)} + \#(\partial u \cap \bigstar_{\mathbb{L}})} P((\partial u)_0)^*,$$

where $P((\partial u)_0) \in \text{Hom}_{\mathbb{C}}(E|_s, E|_p)$ is the parallel transport from $E|_s$ to $E|_p$ along the side of u lying in L.

Switching positions of the left and right sides in (5.15), $\mathscr{F}^{\mathbb{L}}(\mathscr{L})$ [1] is given by maps

$$\hom^{1}(\mathscr{L},\mathbb{L}) = \bigoplus_{s \in \chi^{1}(L,\mathbb{L})} (E|_{s})^{*} \xrightarrow{\Psi^{\mathbb{L}}(\mathscr{L}) = \mathfrak{m}_{1}^{0,b}} \bigoplus_{p \in \chi^{0}(L,\mathbb{L})} (E|_{p})^{*} = \hom^{0}(\mathscr{L},\mathbb{L}),$$

whose $((E|_p)^*, (E|_s)^*)$ -component $\mathfrak{m}_1^{0,b}: (E|_s)^* \to (E|_p)^*$ for each $s, p \in \chi(L, \mathbb{L})$ is

$$\sum_{\substack{\{x_1,X_1\},\dots,\{x_i,X_i\}\\\{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(s,X_1,\dots,X_i,\overline{p})} (-1)^{(i+1)\mathbb{I}_{0}(\mathbb{L})\neq o(\partial u) + \#(\partial u \cap \bigstar_{\mathbb{L}})} P((\partial u)_0)^*,$$

³⁶It boils down to the shift functor of the triangulated category $\underline{MF}(f)$.

where $P((\partial u)_0) \in \text{Hom}_{\mathbb{C}}(E|_p, E|_s)$ is the parallel transport from $E|_p$ to $E|_s$ along the side of u lying in L.

On the other hand, the opposite side $\mathscr{F}^{\mathbb{L}}(J(\mathscr{L})) = (\Phi^{\mathbb{L}}(J(\mathscr{L})), \Psi^{\mathbb{L}}(J(\mathscr{L})))$ is given by (5.16)

$$\hom^{0}(J(\mathscr{L}),\mathbb{L}) = \bigoplus_{p' \in \chi^{0}(J(L),\mathbb{L})} \left(\kappa^{*}E\big|_{p'}\right)^{*} \xrightarrow{\Phi^{\mathbb{L}}(J(\mathscr{L}))=\mathfrak{m}_{1}^{0,b}} \bigoplus_{s' \in \chi^{1}(J(L),\mathbb{L})} \left(\kappa^{*}E\big|_{s'}\right)^{*} = \hom^{1}(J(\mathscr{L}),\mathbb{L}),$$

whose $((E^*|_{s'})^*, (E^*|_{p'})^*)$ -component $\mathfrak{m}_1^{0,b}: (\kappa^* E|_{p'})^* \to (\kappa^* E|_{s'})^*$ for each $p', s' \in \chi(J(L), \mathbb{L})$ is (5.17) $\sum_{\substack{(x_1, X_1), \dots, (x_i, X_i) \\ \in \{(x, X), (y, Y), (z, Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(p', X_1, \dots, X_i, \overline{s'})} (-1)^{(i+1)\mathbb{I}_{0}(\mathbb{L}) \neq 0(\partial u) + \#(\partial u \cap \bigstar_{\mathbb{L}})} P((\partial u)_0)^*,$

where $P((\partial u)_0) \in \text{Hom}_{\mathbb{C}}(\kappa^* E|_{s'}, \kappa^* E|_{p'})$ is the parallel transport from $\kappa^* E|_{s'}$ to $\kappa^* E|_{p'}$ along the side of u lying in L.

Note that reversing the orientation of *L* doesn't affect its intersection with \mathbb{L} , but their degree changes. Therefore, we can replace $p' \in \chi^0(J(L), \mathbb{L})$ and $s' \in \chi^1(J(L), \mathbb{L})$ with $s \in \chi^1(L, \mathbb{L})$ and $p \in \chi^0(L, \mathbb{L})$, respectively. The fibers $\kappa^* E|_{p'}$ and $\kappa^* E|_{s'}$ are also identified with $E|_s$ and $E|_p$, respectively. So we can rewrite (5.16) and (5.17) as

$$\bigoplus_{s \in \chi^{1}(L,\mathbb{L})} (E|_{s})^{*} \xrightarrow{\Phi^{\mathbb{L}}(J(\mathscr{L})) = \mathfrak{m}_{1}^{0,b}} \bigoplus_{p \in \chi^{0}(L,\mathbb{L})} (E|_{p})^{*}$$

whose $((E|_p)^*, (E|_s)^*)$ -component $\mathfrak{m}_1^{0,b}: (E|_s)^* \to (E|_p)^*$ is

$$\sum_{\substack{(x_1,X_1),\ldots,(x_i,X_i)\\ \in\{(x,X),(y,Y),(z,Z)\}}} x_1 \cdots x_i \sum_{u \in \mathcal{M}(s,X_1,\ldots,X_i,\overline{p})} (-1)^{(i+1)\mathbb{I}_{0(\mathbb{L}) \neq 0(\partial u)} + \#(\partial u \cap \bigstar_{\mathbb{L}})} P((\partial u)_0)^*,$$

where $P((\partial u)_0) \in \text{Hom}_{\mathbb{C}}(E|_p, E|_s)$ is the parallel transport from $E|_p$ to $E|_s$ along the side of u lying in L.

Notice that it is the same expression with $\mathscr{F}^{\mathbb{L}}(\mathscr{L})[1]$, especially because sign(*u*) is not relevant to the orientation of *L*. This shows that two functors are the same on the object level. It is also straightforward to check that they coincide on the morphism level.

5.2.4. *Orientation-reversing of loops and shift of modules.* The commutativity of diagram (5.14) induces the commutativity of the left square in the following diagram of ordinary categories and functors:

We explain the commutativity of the right square in the general setting:

Definition 5.10. [Buc21] Let (A, \mathfrak{m}) be a Noetherian local ring and $M \in \underline{CM}(A)$ be a maximal Cohen-Macaulay module over A. Choose an injection $i: M \to Q$ of M into a finitely generated projective A-module Q such that its cokernel is still maximal Cohen-Macaulay. We define the **shift** ³⁷ of M as

$$M[1] := \operatorname{coker}(i),$$

which is uniquely determined as an object of $\underline{CM}(A)$ up to isomorphism.

³⁷It was called **translate** in [Buc21], and **AR translation** in [Yos90].

In the case of hypersurface singularities, it can be explicitly given in terms of matrix factorizations under Eisenbud's equivalence:

Proposition 5.11. Let S be the power series ring $\mathbb{C}[[x_1,...,x_m]]$ of m variables, $f \in S$ its nonzero element and A := S/(f) the quotient ring. Then the following diagram is commutative, that is, two compositions of functors are naturally isomorphic to each other:



Proof. Recall that under Eisenbud's equivalence (Theorem A.16), a matrix factorization $P^0 \rightleftharpoons^{\varphi} \Psi^1$ of f corresponds to a maximal Cohen-Macaulay *A*-module $M := \operatorname{coker} \varphi$, which admits a 2-periodic free resolution given by

 $\cdots \longrightarrow P^0 \otimes_S A \xrightarrow{\underline{\varphi}} P^1 \otimes_S A \xrightarrow{\underline{\psi}} P^0 \otimes_S A \xrightarrow{\underline{\varphi}} P^1 \otimes_S A \longrightarrow M \longrightarrow 0.$

From this we have natural isomorphisms

$$M = \operatorname{coker} \underline{\varphi} = (P^1 \otimes_S A) / \operatorname{im} \underline{\varphi} = (P^1 \otimes_S A) / \operatorname{ker} \underline{\psi} \cong \operatorname{im} \underline{\psi} = \operatorname{ker} \underline{\varphi},$$

and hence there is a natural embedding of *M* into a finitely generated free *A*-module $P^0 \otimes_S A$:

$$i: M \cong \ker \varphi \to P^0 \otimes_S A.$$

Taking its cokernel gives

$$M[1] = \operatorname{coker}(i) = (P^0 \otimes_S A) / \ker \varphi \cong (P^0 \otimes_S A) / \operatorname{im} \psi = \operatorname{coker} \psi,$$

which is also maximal Cohen-Macaulay, being the image of the switched matrix factorization (ψ, φ) under Eisenbud's equivalence.

5.2.5. *Correspondence of canonical forms*. Orientation-reversing of a loop with a local system, switching two factors of a matrix factorization, and the shift of a maximal Cohen-Macaulay modules can all be given explicitly in terms of loop/band data.

In Fuk (Σ), let (L, E, ∇) := ℒ(w', η, ρ) be the loop with a local system corresponding to a loop datum (w', η, ρ). Its orientation-reverse is given by (J(L), κ*E, κ*∇). Recall that the free homotopy class of the loop L(w') corresponding to the given normal loop word w' = (l'₁, m'₁, n'₁, ..., l'_τ, m'_τ, n'_τ) ∈ Z^{3τ} is given by

$$[L(w')] = \left[\alpha^{l'_1}\beta^{m'_1}\gamma^{n'_1}\cdots\alpha^{l'_{\tau}}\beta^{m'_{\tau}}\gamma^{n'_{\tau}}\right],$$

where α , β , γ are generators of $\pi_1(\Sigma)$ (Figure 8a). The free homotopy class of the orientation-reversed loop J(L(w')) is

$$[J(L(w'))] = \left[\gamma^{-n'_{\tau}}\beta^{-m'_{\tau}}\alpha^{-l'_{\tau}}\cdots\gamma^{-n'_{1}}\beta^{-m'_{1}}\alpha^{-l'_{1}}\right].$$

= $\left[\alpha^{-l'_{1}}\beta^{0}\gamma^{-n'_{\tau}}\alpha^{0}\beta^{-m'_{\tau}}\gamma^{0}\alpha^{-l'_{\tau}}\beta^{0}\cdots\gamma^{-n'_{1}}\alpha^{0}\beta^{-m'_{1}}\gamma^{0}\right].$

Define a new normal loop word w'[1], the **reverse** of w', as the normal form of the loop word ³⁸

(5.19)

$$(-l'_1, 0, -n'_{\tau}, 0, -m'_{\tau}, 0, -l'_{\tau}, 0, \dots, -n'_1, 0, -m'_1, 0) \in \mathbb{Z}^{6\tau},$$

then two loops J(L(w')) and L(w'[1]) are freely homotopic to each other.

The holonomy of (E, ∇) at some point is represented by a matrix $J_{\rho}(\eta)$ (up to conjugacy). It changes to $J_{\rho}(\eta)^{-1}$ for the pull-back local system $(\kappa^*, \kappa^* \nabla)$.

Thus, the orientation-reversed loop with a local system $(J(L), \kappa^* E, \kappa^* \nabla)$ is isomorphic to the canonical form $\mathscr{L}(w'[1], \eta^{-1}, \rho)$, that is, they give isomorphic matrix factorizations in <u>MF</u>(*xyz*) by Theorem 3.13.

- In <u>MF</u>(*xyz*), it is just easy to switch two factors of canonical form (φ(w', λ, ρ), ψ(w', λ, ρ)) corresponding to a loop datum (w', λ, ρ), but then ψ(w', λ, ρ) no longer appears in the canonical form. Even worse, it is never obvious how to change it into the canonical form φ(w', λ, ρ) for some another loop datum (w', λ, ρ). Example 5.14 shows that the length 3τ of the word w' can be also changed. (In general, we have ¹/₂τ ≤ τ ≤ 2τ, where 3τ is the length of w'.)
- In $\underline{CM}(A)$ or $\underline{Tri}(A)$, there is no easy way or formula to compute the shift of modules.

The above discussions say that so far the only way to compute the canonical form of the shift of maximal Cohen-Macaulay modules is to make a detour to use the geometric operation in the Fukaya category. We summarize the procedure as follows:

Proposition 5.12 (Shift algorithm). Let $M(w, \lambda, \mu) \in \underline{CM}(A)$ be the maximal Cohen-Macaulay module over *A* corresponding to a band datum (w, λ, μ) . Its shift is given by

$$M(w,\lambda,\mu)[1] = M(w[1],\pm\lambda^{-1},\mu)$$

where the band datum $(w[1], \pm \lambda^{-1}, \mu)$ is computed in the following manner:

- (1) Convert the band datum (w, λ, μ) into a loop datum (w', η, ρ) , following Definition 4.11.
- (2) Compute the reverse w'[1] of the loop word w' (i.e. find the normal form of the loop word (5.19)).
- (3) Convert the loop datum $(w'[1], \eta^{-1}, \rho)$ again into a band datum $(w[1], \pm \lambda^{-1}, \mu)$, following Definition 4.10.

We have the following mappings (up to isomorphism) under the diagram (5.18), while two rows are consistent with our main correspondence (1.5):

$$\begin{aligned} \mathscr{L}(w',\eta,\rho) & \longmapsto \mathscr{F}^{\mathbb{L}} & \varphi_{(\deg)}(w',\lambda,\rho) & \longmapsto^{\overset{\operatorname{coker}}{\simeq}} & M(w,\lambda,\mu) \\ & \stackrel{\operatorname{orientation}}{\underset{\operatorname{reversing}}{\longrightarrow}} \int_{J} & \underset{\operatorname{two factors}}{\overset{\operatorname{switching}}{\longrightarrow}} [1] & \underset{\operatorname{coker}}{\overset{\operatorname{coker}}{\longrightarrow}} & M(w,\lambda,\mu) \\ & \mathscr{L}(w'[1],\eta^{-1},\rho) & \stackrel{\mathscr{F}^{\mathbb{L}}}{\longmapsto} & \varphi_{(\deg)}(w'[1],\pm\lambda^{-1},\rho) & \stackrel{\overset{\simeq}{\longrightarrow}}{\longrightarrow} & M(w[1],\pm\lambda^{-1},\mu) \end{aligned}$$

Example 5.13. The following shows the computation of w[1] from the band word w in Example 4.13:

$$w = (6, 0, 2, -1, 0, -3, 0, 0, 5, 0, -2, 1, -1, 3, 4)$$

$$w' = (8, 2, 3, -1, -1, -4, -1, 0, 5, 0, -2, 1, 0, 4, 6)$$

$$w'[1] = (-8, 0, -6, 0, -4, 0, 0, 0, -1, 0, 2, 0, 0, 0, -5, 0, 0, 0, 1, 0, 4, 0, 1, 0, 1, 0, -3, 0, -2, 0)$$

$$\sim (-6, 1, -4, 1, -3, -1, 0, 2, -5, 1, 0, 4, 0, 1, 0, 1, 0, -2, 1, 0, 1)$$

$$w[1] = (-7, 1, -5, 1, -3, 0, 0, 2, -6, 1, -1, 4, -1, 1, -1, 1, 0, -2, 1, -1, 1)$$

)

³⁸It is not normal a priori in general, but Proposition 3.6 ensures that one can deform it to the unique normal loop word by performing five operations in Lemma 3.3 finitely many times.

Example 5.14. The following shows the correspondence of loops with a local system $\mathcal{L}((3, -2, 2), \eta, 1) \leftrightarrow$ $\mathscr{L}((-2,1,-1,0,2,0),\eta^{-1},1)$, matrix factorizations $\varphi((3,-2,2),\lambda,1) \leftrightarrow \varphi((-2,1,-1,0,2,0),-\lambda^{-1},1)$, and maximal Cohen-Macaulay modules $M((2, -3, 1), \lambda, 1) \leftrightarrow M((-2, 1, -1, 0, 2, 0), -\lambda^{-1}, 1)$ (in Tri(A)) up to isomorphism, where $\lambda = -\eta$. (First two in the bottom row are not presented in the canonical forms.)



5.3. Higher rank/multiplicity and twisted complexes. In this subsection, we first recall the concept of *twisted complexes* in an A_{∞} -category and related notions, based on [Sei08, Boc21] (§5.3.1). Then we associate any twisted complex in $MF_{A_{m}}(f)$ an equivalent matrix factorization (§5.3.2). We derive a formula for extending the localized mirror functor to twisted complexes (§5.3.3). Finally, we show that our canonical objects in MF(xyz) as well as Fuk (Σ) of higher rank are isomorphic to twisted complexes of lower rank objects (§5.3.4, §5.3.5).

5.3.1. Twisted complexes and twisted completion.

Definition 5.15. Let \mathscr{A} be a \mathbb{Z}_2 -graded A_∞ -category over a field \Bbbk . An abstract twisted complex ³⁹ in \mathscr{A} is a pair (\mathcal{L}, δ) , which consists of

• a direct sum of shifted objects, which is a formal expression of the form

$$\mathscr{L} := \bigoplus_{i=1}^{N} \mathscr{L}_i [k_i]$$

for some $N \in \mathbb{Z}_{>1}$, $\mathcal{L}_i \in Ob(\mathcal{A})$ and $k_i \in \mathbb{Z}_2$ $(i \in \{1, \ldots, N\})$,

• a collection of morphisms

$$\delta := \left(\delta_{ij} \in \hom^{1}_{\mathscr{A}}\left(\mathscr{L}_{i}, \mathscr{L}_{j}\right) \left[-k_{i} + k_{j}\right]^{40}\right)_{1 \le i < j \le N}$$

satisfying the Maurer-Cartan equation

(5.20)
$$\mathfrak{m}_{0}^{\delta}(\mathscr{L}) := \sum_{n=1}^{\infty} \mathfrak{m}_{n}(\delta, \dots, \delta)^{41} = 0,$$

³⁹The terminology is intended to distinguish it from the *rigid twisted complex* in MF_{A_∞}(*f*) below (5.24). ⁴⁰For a \mathbb{Z}_2 -graded vector space $V := V^0 \oplus V^1$ and $k \in \mathbb{Z}_2$, we denote by V[k] its *k*-shift, i.e., $V[k]^{\bullet} = V^{\bullet+k}$ for $\bullet \in \mathbb{Z}_2$. ⁴¹It is a finite sum because $\mathfrak{m}_n(\delta, ..., \delta)$ vanishes for $n \ge N$.

where $\mathfrak{m}_n(\delta,...,\delta)$ is an element of $\bigoplus_{1\leq i,j\leq N} \hom^2_{\mathscr{A}}(\mathscr{L}_i,\mathscr{L}_j)[-k_i+k_j]$ with components given by

(5.21)
$$(\mathfrak{m}_{n}(\delta,\ldots,\delta))_{ij} := \sum_{\substack{1 \le n \le j-i \\ i < i_{1} < \cdots < i_{n-1} < j}} \mathfrak{m}_{n} \left(\delta_{ii_{1}}, \delta_{i_{1}i_{2}},\ldots,\delta_{i_{n-1}j} \right) \in \hom_{\mathscr{A}}^{2} \left(\mathscr{L}_{i}, \mathscr{L}_{j} \right) \left[-k_{i} + k_{j} \right].$$

Definition 5.16. Given a \mathbb{Z}_2 -graded A_∞ -category \mathscr{A} , its twisted completion $\operatorname{Tw} \mathscr{A}$ is a \mathbb{Z}_2 -graded A_∞ -category defined as follows:

- Its objects are the abstract twisted complexes in \mathcal{A} ,
- The morphism space between $\left(\mathcal{L}_0 = \bigoplus_{i=1}^{N_0} \mathcal{L}_{0i}[k_{0i}], \delta_0\right)$ and $\left(\mathcal{L}_1 = \bigoplus_{i=1}^{N_1} \mathcal{L}_{1i}[k_{1i}], \delta_1\right)$ is

$$\hom_{\mathrm{Tw}\mathscr{A}}^{\bullet}\left(\bigoplus_{i=1}^{N_{0}}\mathscr{L}_{0i}\left[k_{0i}\right],\bigoplus_{j=1}^{N_{1}}\mathscr{L}_{1j}\left[k_{1j}\right]\right):=\bigoplus_{i=1}^{N_{0}}\bigoplus_{j=1}^{N_{1}}\hom_{\mathscr{A}}^{\bullet}\left(\mathscr{L}_{0i},\mathscr{L}_{1j}\right)\left[-k_{0i}+k_{1j}\right] \quad (\bullet \in \mathbb{Z}_{2})$$

• The A_{∞} -operations $\{\mathfrak{m}_{k}^{\mathrm{Tw}\mathscr{A}}\}_{k\geq 1}$ are defined as

$$\mathfrak{m}_{k}^{\mathrm{Tw}\mathscr{A}}:\hom_{\mathrm{Tw}\mathscr{A}}((\mathscr{L}_{0},\delta_{0}),(\mathscr{L}_{1},\delta_{1}))\otimes\cdots\otimes\hom_{\mathrm{Tw}\mathscr{A}}((\mathscr{L}_{k-1},\delta_{k-1}),(\mathscr{L}_{k},\delta_{k}))\to\hom_{\mathrm{Tw}\mathscr{A}}((\mathscr{L}_{0},\delta_{0}),(\mathscr{L}_{k},\delta_{k})),$$

$$(f_{1},\ldots,f_{k})\mapsto\mathfrak{m}_{k}^{\delta_{0},\ldots,\delta_{k}}(f_{1},\ldots,f_{k}):=\sum_{m_{0},\ldots,m_{k}\geq0}\mathfrak{m}_{k+m_{0}+\cdots+m_{k}}^{\mathscr{A}}(\underbrace{\delta_{0},\ldots,\delta_{0}}_{m_{0}},f_{1},\underbrace{\delta_{1},\ldots,\delta_{1}}_{m_{1}},f_{2},\ldots,f_{k},\underbrace{\delta_{k},\ldots,\delta_{k}}_{m_{k}})^{42}.$$

$$for f_{i}\in\hom_{\mathrm{Tw}\mathscr{A}}((\mathscr{L}_{i-1},\delta_{i-1}),(\mathscr{L}_{i},\delta_{i})) \ (i\in\{1,\ldots,k\}).$$

It is a triangulated A_{∞} -category, and its cohomological category $H^0(\text{Tw} \mathscr{A})$ becomes a triangulated category in the classical sense ([Sei08, §I.3]).

Note that there is a natural embedding $\mathscr{A} \hookrightarrow \text{Tw}\mathscr{A}$ of A_{∞} -categories sending each object $\mathscr{L} \in \text{Ob}(\mathscr{A})$ to the trivial abstract twisted complex ($\mathscr{L}[0], 0$).

Taking the twisted completion of A_{∞} -categories is *functorial* in the following sense:

Proposition 5.17. An A_{∞} -functor $\mathscr{F} := \{\mathscr{F}_k\}_{k\geq 0} : \mathscr{A} \to \mathscr{B}$ between A_{∞} -categories induces an A_{∞} -functor $\operatorname{Tw}\mathscr{F} := \{(\operatorname{Tw}\mathscr{F})_k\}_{k\geq 0} : \operatorname{Tw}\mathscr{A} \to \operatorname{Tw}\mathscr{B}$ between their twisted completions defined as follows:

• An abstract twisted complex $\left(\bigoplus_{i=1}^{N} \mathscr{L}_{i}[k_{i}], \delta\right)$ in \mathscr{A} is mapped to the abstract twisted complex in \mathscr{B} given by

$$\left(\bigoplus_{i=1}^{N} \mathscr{F}_{0}\left(\mathscr{L}_{i}\right)\left[k_{i}\right], \sum_{n=1}^{\infty} \mathscr{F}_{n}\left(\delta, \ldots, \delta\right)\right),$$

where $\mathscr{F}_n(\delta,...,\delta)$ is an element of $\bigoplus_{1 \le i,j \le N} \hom^1_{\mathscr{B}} (\mathscr{F}_0(\mathscr{L}_i), \mathscr{F}_0(\mathscr{L}_j)) [-k_i + k_j]$ with components

$$(\mathscr{F}_n(\delta,\ldots,\delta))_{ij} := \sum_{\substack{1 \le n \le j - i\\i < i_1 < \cdots < i_{n-1} < j}} \mathscr{F}_n\left(\delta_{ii_1},\delta_{i_1i_2},\ldots,\delta_{i_{n-1}j}\right) \in \hom^1_{\mathscr{B}}\left(\mathscr{F}_0\left(\mathscr{L}_i\right),\mathscr{F}_0\left(\mathscr{L}_j\right)\right)\left[-k_i + k_j\right].$$

• Higher components $\{(\operatorname{Tw} \mathscr{F})_k\}_{k\geq 1}$ are given by

 $(\mathrm{Tw}\mathscr{F})_k: \hom_{\mathrm{Tw}\mathscr{A}}((\mathscr{L}_0, \delta_0), (\mathscr{L}_1, \delta_1)) \otimes \cdots \otimes \hom_{\mathrm{Tw}\mathscr{A}}((\mathscr{L}_{k-1}, \delta_{k-1}), (\mathscr{L}_k, \delta_k))$

$$\rightarrow \hom_{\mathrm{Tw}\mathscr{B}}((\mathrm{Tw}\mathscr{F})_{0}((\mathscr{L}_{0},\delta_{0})),(\mathrm{Tw}\mathscr{F})_{0}((\mathscr{L}_{k},\delta_{k}))), \\ (f_{1},\ldots,f_{k}) \mapsto \sum_{m_{0},\ldots,m_{k}\geq 0} \mathscr{F}_{k+m_{0}+\cdots+m_{k}}(\underbrace{\delta_{0},\ldots,\delta_{0}}_{m_{0}},f_{1},\underbrace{\delta_{1},\ldots,\delta_{1}}_{m_{1}},f_{2},\ldots,f_{k},\underbrace{\delta_{k},\ldots,\delta_{k}}_{m_{k}}).$$

The induced A_{∞} -functor $\operatorname{Tw} \mathscr{F} : \operatorname{Tw} \mathscr{A} \to \operatorname{Tw} \mathscr{B}$ also boils down to an exact functor $H^0(\operatorname{Tw} \mathscr{F}) : H^0(\operatorname{Tw} \mathscr{A}) \to H^0(\operatorname{Tw} \mathscr{B})$ between classical triangulated categories.

52

 $^{^{42}}$ Each component is defined in the same manner as in (5.21).

5.3.2. *Twisted complexes of matrix factorizations*. Let *S* be the power series ring $\mathbb{C}[[x_1, ..., x_m]]$ of *m* variables, and $f \in S$ its nonzero element. We will demonstrate that the A_{∞} -category of matrix factorizations $MF_{A_{\infty}}(f)$ has an intrinsic notion of twisted complexes, by constructing them as actual objects in it.

In §5.2.2, we already defined the shift functor [1]: $MF_{A_{\infty}}(f) \to MF_{A_{\infty}}(f)$, which simply switches the position of two matrices in a given matrix factorization. More precisely, given a matrix factorization $P^0 \stackrel{\varphi}{\underset{W}{\longrightarrow}} P^1$,

we define its k-shift $(k \in \mathbb{Z}_2)$ as $P[k]^0 \xrightarrow{\varphi[k]} P[k]^1$, where $P[k]^i := P^{i+k}$ and $(\varphi, \psi)[k] := (\varphi[k], \psi[k]) := \begin{cases} (\varphi, \psi) & \text{if } k = 0, \\ (\psi, \varphi) & \text{if } k = 1. \end{cases}$

Now we associate any abstract twisted complex given in Definition 5.15 an object in $MF_{A_{\infty}}(f)$. Suppose that we have finitely many shifted matrix factorizations $P_i[k_i]^0 \xrightarrow{\varphi_i[k_i]} P_i[k_i]^1$ $(N \in \mathbb{Z}_{\geq 1}, i \in \{1, ..., N\})$ and morphisms between them

 $(\gamma_{ij}, \delta_{ij}) \in \text{hom}^1((\varphi_j, \psi_j), (\varphi_i, \psi_i))[-k_j + k_i] = \text{Hom}_S(P_j[k_j]^0, P_i[k_i]^1) \times \text{Hom}_S(P_j[k_j]^1, P_i[k_i]^0)$ for $1 \le i < j \le N$, which form the following left diagram (not necessarily commutative):



We can arrange them into the block matrix form as follows:

$$\begin{aligned} (\varphi,\psi) &:= \left(\bigoplus_{i=1}^{N} \varphi_{i}[k_{i}], \bigoplus_{i=1}^{N} \psi_{i}[k_{i}] \right) \\ &= \left(\begin{array}{cccc} P_{1}[k_{1}]^{0} & P_{2}[k_{2}]^{0} & \cdots & P_{N}[k_{N}]^{0} & P_{1}[k_{1}]^{1} & P_{2}[k_{2}]^{1} & \cdots & P_{N}[k_{N}]^{1} \\ P_{1}[k_{1}]^{1} & \varphi_{1}[k_{1}] & 0 & \cdots & 0 \\ P_{2}[k_{2}]^{1} & 0 & \varphi_{2}[k_{2}] & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ P_{N}[k_{N}]^{1} & 0 & \cdots & \varphi_{N}[k_{N}] \end{array} \right), \begin{array}{c} P_{1}[k_{1}]^{0} & P_{1}[k_{1}] & 0 & \cdots & 0 \\ P_{2}[k_{2}]^{0} & 0 & \cdots & \varphi_{N}[k_{N}] \end{array} \right), \begin{array}{c} P_{1}[k_{1}]^{0} & \psi_{1}[k_{1}] & 0 & \cdots & 0 \\ 0 & \psi_{2}[k_{2}] & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \psi_{N}[k_{N}] \end{array} \right), \end{aligned}$$

$$(\gamma, \delta) := ((\gamma_{ij}, \delta_{ij}) \in \hom^1((\varphi_j, \psi_j), (\varphi_i, \psi_i)) [-k_j + k_i])_{1 \le i < j \le N}$$

δ_{1N}	
÷	
(N-1)N	
₀)	
	$ \sum_{N=1}^{N-1} N $

Observe that (φ, ψ) forms a new matrix factorization $P^0 \stackrel{\varphi}{\underset{\psi}{\longleftrightarrow}} P^1$, where $P^{\bullet} := \bigoplus_{i=1}^N P_i[k_i]^{\bullet}$ (• $\in \mathbb{Z}_2$), and (γ, δ) defines a morphism $(\gamma, \delta) \in \hom^1((\varphi, \psi), (\varphi, \psi))$, which form the right diagram in (5.22) (not necessarily commutative).

The A_{∞} -operations between them are computed as

 $\mathfrak{m}_1((\gamma,\delta)) = (\psi\gamma + \delta\varphi, \varphi\delta + \gamma\psi), \quad \mathfrak{m}_2((\gamma,\delta), (\gamma,\delta)) = (\delta\gamma, \gamma\delta) \quad \text{and} \quad \mathfrak{m}_{k\geq 3}((\delta,\gamma), \dots, (\delta,\gamma)) = 0,$ following (A.10). It is easy to check that their components are also compatible with the abstract definition in (5.21). Therefore, the Maurer-Cartan equation (5.20) is phrased as

(5.23)
$$0 = \mathfrak{m}_1((\gamma, \delta)) + \mathfrak{m}_2((\gamma, \delta), (\gamma, \delta)) = (\psi\gamma + \delta\varphi + \delta\gamma, \varphi\delta + \gamma\psi + \gamma\delta),$$

and an abstract twisted complex in $MF_{A_{\infty}}(f)$ is equivalent to a pair $((\varphi, \psi), (\gamma, \delta))$ satisfying (5.23) ⁴³.

We define the **rigid twisted complex** in $MF_{A_{\infty}}(f)$ associated to such a pair as $P^0 \xleftarrow{\varphi+\gamma}{\psi+\delta} P^1$. More precisely, it is given as

$$\operatorname{Tw}((\varphi,\psi),(\gamma,\delta)) := (\varphi+\gamma,\psi+\delta)$$

(5.24)
$$= \begin{pmatrix} P_{1}[k_{1}]^{0} & P_{2}[k_{2}]^{0} & \cdots & P_{N}[k_{N}]^{0} & P_{1}[k_{1}]^{1} & P_{2}[k_{2}]^{1} & \cdots & P_{N}[k_{N}]^{1} \\ P_{1}[k_{1}]^{1} & \varphi_{1}[k_{1}] & \gamma_{12} & \cdots & \gamma_{1N} \\ P_{2}[k_{2}]^{1} & 0 & \varphi_{2}[k_{2}] & \ddots & \vdots \\ \vdots & \vdots & \ddots & \gamma_{(N-1)N} \\ P_{N}[k_{N}]^{1} & 0 & 0 & \cdots & \varphi_{N}[k_{N}] \end{pmatrix}, \begin{pmatrix} P_{1}[k_{1}]^{0} & P_{1}[k_{1}]^{1} & \delta_{12} & \cdots & \delta_{1N} \\ P_{2}[k_{2}]^{0} & 0 & \psi_{2}[k_{2}] & \ddots & \vdots \\ \vdots & \vdots & \ddots & \delta_{(N-1)N} \\ 0 & 0 & \cdots & \varphi_{N}[k_{N}] \end{pmatrix}, \begin{pmatrix} P_{1}[k_{1}]^{0} & \psi_{1}[k_{1}] & \delta_{12} & \cdots & \delta_{1N} \\ 0 & \psi_{2}[k_{2}] & \ddots & \vdots \\ \vdots & \vdots & \ddots & \delta_{(N-1)N} \\ 0 & 0 & \cdots & \psi_{N}[k_{N}] \end{pmatrix}$$

It is indeed a matrix factorization of f, as a direct consequence of the equation (5.23).

Proposition 5.18. The embedding $i: MF_{A_{\infty}}(f) \hookrightarrow TwMF_{A_{\infty}}(f)$ of A_{∞} -categories is a quasi-equivalence.

Proof. The natural embedding *i* sends each matrix factorization (φ, ψ) to the abstract twisted complex $((\varphi, \psi)[0], 0)$. The morphism space between such two complexes $((\varphi_0, \psi_0)[0], 0)$ and $((\varphi_1, \psi_1)[0], 0)$ is identified with the original hom space hom^{\mathbb{Z}_2} $((\varphi_1, \psi_1), (\varphi_0, \psi_0))$. On the morphism level, the first component *i*₁ is the identity map on that space, and the higher components *i*_{*k*≥2} are defined to be zero.

We define its quasi-inverse π : TwMF_{A_m} $(f) \rightarrow$ MF_{A_m}(f) by sending each abstract twisted complex

$$\left(\left(\varphi,\psi\right):=\left(\bigoplus_{i=1}^{N}\varphi_{i}[k_{i}],\bigoplus_{i=1}^{N}\psi_{i}[k_{i}]\right),\left(\gamma,\delta\right):=\left(\left(\gamma_{ij},\delta_{ij}\right)\in\hom^{1}\left(\left(\varphi_{j},\psi_{j}\right),\left(\varphi_{i},\psi_{i}\right)\right)[-k_{j}+k_{i}]\right)_{1\leq i< j\leq N}\right)$$

to the rigid twisted complex Tw($(\varphi, \psi), (\gamma, \delta)$). Given two abstract twisted complexes ($(\varphi_0, \psi_0), (\gamma_0, \delta_0)$) and ($(\varphi_1, \psi_1), (\gamma_1, \delta_1)$), their hom space

$$\hom_{\text{Tw}MF_{A_{\infty}}(f)}^{\bullet} \left(\left(\left(\varphi_{0}, \psi_{0} \right), \left(\gamma_{0}, \delta_{0} \right) \right), \left(\left(\varphi_{1}, \psi_{1} \right), \left(\gamma_{1}, \delta_{1} \right) \right) \right) = \bigoplus_{i=1}^{N_{0}} \bigoplus_{j=1}^{N_{1}} \hom^{\bullet} \left(\left(\left(\varphi_{1j}, \psi_{1j} \right), \left(\varphi_{0i}, \psi_{0i} \right) \right) \left[-k_{0i} + k_{1j} \right] \right)$$
$$= \bigoplus_{i=1}^{N_{0}} \bigoplus_{j=1}^{N_{1}} \left(\operatorname{Hom}_{S} \left(P_{1j}[k_{1j}]^{0}, P_{0i}[k_{0i}]^{\bullet} \right) \times \operatorname{Hom}_{S} \left(P_{1j}[k_{1j}]^{1}, P_{0i}[k_{0i}]^{1+\bullet} \right) \right) \quad (\bullet \in \mathbb{Z}_{2})$$

is naturally identified with the hom space between rigid twisted complexes

$$\operatorname{hom}_{\operatorname{MF}_{A_{\infty}}(f)}^{\bullet}\left(\operatorname{Tw}\left(\left(\varphi_{0},\psi_{0}\right)\left(\gamma_{0},\delta_{0}\right)\right),\operatorname{Tw}\left(\left(\varphi_{1},\psi_{1}\right),\left(\gamma_{1},\delta_{1}\right)\right)\right)$$

$$=\operatorname{Hom}_{S}\left(\bigoplus_{i=1}^{N_{1}}P_{1j}[k_{1j}]^{0},\bigoplus_{j=1}^{N_{0}}P_{0i}[k_{0i}]^{\bullet}\right)\times\operatorname{Hom}_{S}\left(\bigoplus_{i=1}^{N_{1}}P_{1j}[k_{1j}]^{1},\bigoplus_{j=1}^{N_{0}}P_{0i}[k_{0i}]^{1+\bullet}\right)\quad(\bullet\in\mathbb{Z}_{2})$$

Therefore, we can define the first component π_1 of the A_{∞} -functor π as the identity map on that space, and the higher components $\pi_{k\geq 2}$ as zero. It is straightforward to check that *i* and π are indeed quasi-inverse to each other.

⁴³The pair is also an example of a *bounding cochain*. Compare it with the *weak Maurer-Cartan equation* (2.1).

5.3.3. *Twisted complexes under localized mirror functor.* Combining above discussions, we derive a formula for extending the localized mirror functor to twisted complexes.

Proposition 5.19. The image of an abstract twisted complex $(\mathcal{L} := \bigoplus_{i=1}^{N} \mathcal{L}_i, \delta)$ in W Fuk (Σ) under the induced localized mirror functor Tw $\mathscr{F}^{\mathbb{L}}$: Tw W Fuk $(\Sigma) \to MF_{A_{\infty}}(xyz)$ is the rigid twisted complex

$$\bigoplus_{i=1}^{N} \hom^{0}(\mathscr{L}_{i},\mathbb{L}) \xrightarrow{\Phi^{\mathbb{L}}((\mathscr{L},\delta))=\mathfrak{m}_{1}^{\delta,b}} \bigoplus_{j=1}^{N} \hom^{1}(\mathscr{L}_{j},\mathbb{L}),$$

with components given by (5.25)

$$\left(\mathfrak{m}_{1}^{\delta,b}\right)_{ij} := \sum_{\substack{0 \le n \le j-i \\ i < i_{1} < \cdots < i_{n-1} < j}} \sum_{k=0}^{\infty} \mathfrak{m}_{n+1+k} \left(\delta_{ii_{1}}, \delta_{i_{1}i_{2}}, \dots, \delta_{i_{n-1}j}, -, \underbrace{b, \dots, b}_{k}\right)^{44} : \hom^{\bullet}\left(\mathscr{L}_{j}, \mathbb{L}\right) \to \hom^{\bullet+1}\left(\mathscr{L}_{i}, \mathbb{L}\right)$$

for $1 \le i \le j \le N$ and $\bullet \in \mathbb{Z}_2$.

In particular, its diagonal components are given by the original mirror $(\Phi^{\mathbb{L}}(\mathcal{L}_i), \Psi^{\mathbb{L}}(\mathcal{L}_i))$ (2.3) of \mathcal{L}_i , and its strictly upper triangular components are determined by inserting (j - i)-times of the twisting δ .

Proof. Recall from Proposition 5.17 that the induced functor $\text{Tw}\mathscr{F}^{\mathbb{L}}$: $\text{Tw} W \text{Fuk}(\Sigma) \to \text{Tw} MF_{A_{\infty}}(xyz)$ maps the given abstract twisted complex to the abstract twisted complex in $MF_{A_{\infty}}(xyz)$ given by

(5.26)
$$\left(\bigoplus_{i=1}^{N} \mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{i}), \sum_{n=1}^{\infty} \mathscr{F}_{n}^{\mathbb{L}}(\delta, \dots, \delta)\right),$$

where $\mathscr{F}_{n}^{\mathbb{L}}(\delta,...,\delta)$ is an element of $\bigoplus_{1 \leq i,j \leq N} \hom^{1}_{\operatorname{MF}_{A_{\infty}}(xyz)} \left(\mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{i}), \mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{j}) \right)$ with components

$$\left(\mathscr{F}_{n}^{\mathbb{L}}(\delta,\ldots,\delta)\right)_{ij} := \sum_{\substack{1 \le n \le j-i \\ i < i_{1} < \cdots < i_{n-1} < j}} \mathscr{F}_{n}^{\mathbb{L}}\left(\delta_{ii_{1}},\delta_{i_{1}i_{2}},\ldots,\delta_{i_{n-1}j}\right) \in \hom^{1}_{\mathrm{MF}_{A_{\infty}}(xyz)}\left(\mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{i}),\mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{j})\right).$$

The definition of $\mathscr{F}^{\mathbb{L}}$ in Theorem 2.3 identifies those with

$$\mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{i}) = \mathfrak{m}_{1}^{0,b} : \hom^{\bullet}(\mathscr{L}_{i},\mathbb{L}) \to \hom^{\bullet+1}(\mathscr{L}_{i},\mathbb{L}) \quad \text{and}$$
$$\mathscr{F}_{n}^{\mathbb{L}}(\delta_{ii_{1}},\delta_{i_{1}i_{2}},\ldots,\delta_{i_{n-1}j}) = \mathfrak{m}_{n+1}^{0,\ldots,0,b}(\delta_{ii_{1}},\delta_{i_{1}i_{2}},\ldots,\delta_{i_{n-1}j},-) : \hom^{\bullet}(\mathscr{L}_{j},\mathbb{L}) \to \hom^{\bullet+1}(\mathscr{L}_{i},\mathbb{L}).$$

Finally, under (5.24), the rigid twist complex in $MF_{A_{\infty}}(xyz)$ corresponding to (5.26) is given by

$$\bigoplus_{i=1}^{N} \mathscr{F}_{0}^{\mathbb{L}}(\mathscr{L}_{i}) + \sum_{n=1}^{\infty} \mathscr{F}_{n}^{\mathbb{L}}(\delta, \dots, \delta) = \mathfrak{m}_{1}^{0,b}(-) + \sum_{n=1}^{\infty} \mathfrak{m}_{n+1}^{0,\dots,0,b}(\underbrace{\delta,\dots,\delta}_{n}, -) = \sum_{n=0}^{\infty} \mathfrak{m}_{n+1}^{0,\dots,0,b}(\underbrace{\delta,\dots,\delta}_{n}, -),$$

which is the map $\mathfrak{m}_{1}^{\delta,b}$: $\bigoplus_{i=1}^{N}$ hom[•] ($\mathscr{L}_{i},\mathbb{L}$) $\rightarrow \bigoplus_{i=1}^{N}$ hom^{•+1} ($\mathscr{L}_{i},\mathbb{L}$) with the same components as given in (5.25).

Remark 5.20. One can also directly define the localized mirror functor $\mathscr{F}^{\mathbb{L}}$: Tw W Fuk $(\Sigma) \to MF_{A_{\infty}}(xyz)$ based on the above formula, not passing through Proposition 5.17. For instance, the identity $\left(\mathfrak{m}_{1}^{\delta,b}\right)^{2} = W^{\mathbb{L}} \cdot \mathrm{id}_{\mathrm{hom}(\mathscr{L},\mathbb{L})}$ for any twisted complex (as well as bounding cochain) $(\mathscr{L} := \bigoplus_{i=1}^{N} \mathscr{L}_{i}, \delta)$ in W Fuk (Σ) follows from the same line of proof as in Lemma 2.2. Then one can extend the definition given in Theorem 2.3 by $using \mathfrak{m}_{1}^{\delta,b}$ and $\mathfrak{m}_{k+1}^{\delta,\dots,\delta,b}$ instead of $\mathfrak{m}_{1}^{0,\dots,0,b}$, respectively.

⁴⁴It is just \mathfrak{m}_{1+k} (-, *b*, ..., *b*) if *n* = 0.

5.3.4. Canonical form of matrix factorizations viewed as twisted complexes. Now we show that our canonical form of matrix factorizations of xyz defined in Definition 3.16 with a higher rank is expressed as a twisted complex consisting of several copies of the corresponding object with rank 1.

Proposition 5.21. The canonical form $(\varphi(w', \lambda, \rho), \psi(w', \lambda, \rho))$ of matrix factorizations of xyz corresponding to a non-degenerate loop datum is quasi-isomorphic in MF_{A_∞}(xyz) (and isomorphic in MF(xyz)) to the rigid twisted complex

$$\operatorname{Tw}\left(\left(\varphi\left(w',\lambda,1\right)^{\oplus\rho},\psi\left(w',\lambda,1\right)^{\oplus\rho}\right),(\gamma,\delta)\right)$$

for some morphisms

 $(\gamma, \delta) := \left(\left(\gamma_{ij}, \delta_{ij} \right) \in \hom^1 \left(\left(\varphi\left(w', \lambda, 1 \right), \psi\left(w', \lambda, 1 \right) \right), \left(\varphi\left(w', \lambda, 1 \right), \psi\left(w', \lambda, 1 \right) \right) \right)_{1 \le i < j \le \rho}.$

Proof. Recall from (3.2) that the canonical form of matrix factorizations of xyz corresponding to a nondegenerate loop datum (w', λ, ρ) is written as

$$\begin{split} \varphi\left(w',\lambda,\rho\right) &= \varphi\left(w',0,1\right) \otimes I_{\rho} - x^{l_{1}'-1} K_{3\tau}^{3\tau-1} \otimes J_{\rho}\left(\lambda\right) - x^{-l_{1}'} J_{3\tau}^{3\tau-1} \otimes J_{\rho}\left(\lambda\right)^{-1} \\ &= \varphi\left(w',\lambda,1\right) \otimes I_{\rho} - x^{l_{1}'-1} K_{3\tau}^{3\tau-1} \otimes \left(J_{\rho}\left(\lambda\right) - \lambda I_{\rho}\right) - x^{-l_{1}'} J_{3\tau}^{3\tau-1} \otimes \left(J_{\rho}\left(\lambda\right)^{-1} - \lambda^{-1} I_{\rho}\right) \\ &= \varphi\left(w',\lambda,1\right) \otimes I_{\rho} - x^{l_{1}'-1} K_{3\tau}^{3\tau-1} \otimes J_{\rho} - x^{-l_{1}'} J_{3\tau}^{3\tau-1} \otimes \left(-\lambda^{-2} J_{\rho} + \lambda^{-3} J_{\rho}^{2} - \dots - (-\lambda)^{-\rho} J_{\rho}^{\rho-1}\right). \end{split}$$

Using Lemma 3.20, we know that it is similar to

$$I_{\rho} \otimes \varphi(w',\lambda,1) - x^{l_{1}'-1} J_{\rho} \otimes K_{3\tau}^{3\tau-1} + x^{-l_{1}'} \left(\lambda^{-2} J_{\rho} - \lambda^{-3} J_{\rho}^{2} + \dots + (-\lambda)^{-\rho} J_{\rho}^{\rho-1}\right) \otimes J_{3\tau}^{3\tau-1},$$

which is expressed as

(5.27)

$$\begin{pmatrix} \varphi(w',\lambda,1) - x^{l_1'-1}K_{3\tau}^{3\tau-1} & 0 & \cdots & 0 \\ 0 & \varphi(w',\lambda,1) & -x^{l_1'-1}K_{3\tau}^{3\tau-1} & \cdots & 0 \\ 0 & 0 & \varphi(w',\lambda,1) & -x^{l_1'-1}K_{3\tau}^{3\tau-1} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & -x^{l_1'-1}K_{3\tau}^{3\tau-1} \\ 0 & 0 & 0 & \cdots & \varphi(w',\lambda,1) \end{pmatrix}_{\rho_{3\tau}\times\rho_{3\tau}} \text{ and } \begin{pmatrix} \varphi(w',\lambda,1) \ \lambda^{-2}x^{-l_1'}J_{3\tau}^{3\tau-1} - \lambda^{-3}x^{-l_1'}J_{3\tau}^{3\tau-1} & \cdots & (-\lambda)^{-\rho_{\tau}l_{\tau}'}J_{3\tau}^{3\tau-1} \\ 0 & \varphi(w',\lambda,1) \ \lambda^{-2}x^{-l_{1}'}J_{3\tau}^{3\tau-1} & \cdots & (-\lambda)^{-\rho_{\tau}l_{\tau}'}J_{3\tau}^{3\tau-1} \\ 0 & 0 & \varphi(w',\lambda,1) \ \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \lambda^{-2}x^{-l_{1}'}J_{3\tau}^{3\tau-1} \\ 0 & 0 & 0 & \cdots & \varphi(w',\lambda,1) \end{pmatrix}_{\rho_{3\tau}\times\rho_{3\tau}} \rho_{3\tau}$$

in the case of $l'_1 \ge 1$ and $l'_1 \le 0$, respectively. This observation together with the definition (5.24) of rigid twisted complexes prove the proposition.

5.3.5. *Canonical form of loops with a local system viewed as twisted complexes.* In this subsection, we will realize (5.27) as the image of an abstract twisted complex in Fuk(Σ) under the induced localized mirror functor Tw $\mathscr{F}^{\mathbb{L}}$: Tw *W* Fuk(Σ) \rightarrow MF_{*A*_w}(*xyz*), as a direct consequence of Proposition 5.19.

We first take the loop with a rank 1 local system $\mathcal{L} := (L, E, \nabla) := \mathcal{L}(w', \eta, 1)$, where $(w', \eta, 1)$ is the nondegenerate loop datum corresponding to $(w', \lambda, 1)$ under Theorem 3.17. Its underlying loop L = L(w') has a marked point $o_L \in \chi^1(L, L)$ that we assume is located nearby the point \bigstar as in Figure 20. We denote by $o := \operatorname{id}_{o_L} \in \operatorname{Hom}_{\mathbb{C}}(E|_{o_L}, E|_{o_L}) \subseteq \operatorname{hom}^1(\mathcal{L}, \mathcal{L})$.

Consider the abstract twisted complex

(5.28)
$$\left(\mathscr{L} \left(w', \eta, 1 \right)^{\oplus \rho}, \, \delta = \begin{pmatrix} 0 & o_{12} & 0 & \cdots & 0 \\ 0 & 0 & o_{23} & \cdots & 0 \\ 0 & 0 & 0 & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & o_{(N-1)N} \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \right)$$

that consists of the direct sum of ρ -copies of $\mathcal{L}_i := \mathcal{L} = \mathcal{L}(w', \eta, 1)$ $(i \in \{1, ..., \rho\})$ and a collection of morphisms

$$\delta := \left(\delta_{ij} \in \hom^1(\mathscr{L}_i, \mathscr{L}_j)\right)_{1 \le i < j \le \rho}$$

where δ_{ij} is nontrivial only for j = i + 1, in which case it is $o_{i(i+1)} := o \in \hom^1(\mathcal{L}_i, \mathcal{L}_{i+1}) = \hom^1(\mathcal{L}, \mathcal{L})$.

Proposition 5.22. The mirror image of the abstract twisted complex (5.28) in Fuk (Σ) is quasi-isomorphic in MF_{A_∞}(xyz) (and isomorphic in MF(xyz)) to the rigid twisted complex

$$J_{\rho}(\eta)^{-1} \otimes \varphi_{-1} + I_{\rho} \otimes \varphi_{0} + J_{\rho}(\eta) \otimes \varphi_{1} = \begin{pmatrix} \Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,1)) & \varphi_{1} - \eta^{-2}\varphi_{-1} & \eta^{-3}\varphi_{-1} & \cdots & -(-\eta)^{-\rho}\varphi_{-1} \\ 0 & \Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,1)) & \varphi_{1} - \eta^{-2}\varphi_{-1} & \cdots & -(-\eta)^{-\rho+1}\varphi_{-1} \\ 0 & 0 & \Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,1)) & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \varphi_{1} - \eta^{-2}\varphi_{-1} \\ 0 & 0 & 0 & \cdots & \Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,1)) \end{pmatrix}_{\rho_{3\tau} \times \rho_{3\tau}}$$

where $\Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,1)) = \eta^{-1}\varphi_{-1} + \varphi_0 + \eta\varphi_1$ for $\varphi_{-1}, \varphi_0, \varphi_1 \in \mathbb{C}[[x, y, z]]^{3\tau \times 3\tau}$ (from Proposition 3.15). In particular, it is also isomorphic to the rigid twisted complex (5.27) given in Proposition 5.21.

Proof. According to Proposition 5.19, the mirror image is given in the form

$$\hom^{0}(\mathscr{L},\mathbb{L})^{\oplus\rho} \xrightarrow{\Phi^{\mathbb{L}}((\mathscr{L}^{\oplus\rho},\delta))=\mathfrak{m}_{1}^{\delta,b}} \operatorname{hom}^{1}(\mathscr{L},\mathbb{L})^{\oplus\rho}$$

Each diagonal component of $\Phi^{\mathbb{L}}((\mathscr{L}^{\oplus \rho}, \delta))$ is $\Phi^{\mathbb{L}}(\mathscr{L})$, and strictly upper triangular components are

(5.29)
$$\left(\mathfrak{m}_{1}^{\delta,b}\right)_{ij} \coloneqq \sum_{k=0}^{\infty} \mathfrak{m}_{j-i+1+k}\left(o_{i(i+1)},\ldots,o_{(j-1)j},-,\underbrace{b,\ldots,b}_{k}\right) \colon \hom^{\bullet}\left(\mathscr{L}_{j},\mathbb{L}\right) \to \hom^{\bullet+1}\left(\mathscr{L}_{i},\mathbb{L}\right)$$

for $1 \le i < j \le N$ and $\bullet \in \mathbb{Z}_2$. Note that it depends only on the difference j - i, and hence we assume i = 1 without loss of generality. Substituting $f = f|_p \in \hom^0(\mathcal{L}_j, \mathbb{L})$ for some $p \in \chi^0(L_j, \mathbb{L})$ into it becomes

$$\sum_{\substack{(x_1, X_1), \dots, (x_k, X_k) \\ \in \{(x, X), (y, Y), (z, Z)\}}} x_1 \dots x_k \sum_{s \in \chi^1(L_1, \mathbb{L})} \sum_{u \in \mathcal{M}} (o_{12}, o_{23} \dots, o_{(j-1)j}, p, X_1, \dots, X_k, \overline{s})} \operatorname{sign}(u) \operatorname{hol}_s(\partial u) (o_{12}, o_{23}, \dots, o_{(j-1)j}, f, X_1, \dots, X_k),$$

following the same procedure as in §2.2.

We will associate each element *u* in the moduli space

(5.30)
$$\mathcal{M}(o_{12}, o_{23}, \dots, o_{(j-1)j}, p, X_1, \dots, X_k, \overline{s})$$

with an element u' in $\mathcal{M}(p, X_1, ..., X_k, \overline{s})$, by considering recursive perturbations of ρ -copies of $L_i = L = L(w')$ ($i \in \{1, ..., \rho\}$) as described in Figure 20.



Figure 20. Recursive perturbations of a loop L

Case i) The boundary orientation of u coincides with the orientations of L_i 's. Such a polygon u cannot have angles at more than one of $o_{12}, o_{23}, \ldots, o_{(j-1)j}$ in a consecutive manner because of their arrangement as in Figure 20. Hence it exists only when j = 2, and it has an angle at o_{12} . Ignoring o_{12} and identifying L_1 and L_2 , u has an obvious counterpart u' in $\mathcal{M}(p, X_1, \ldots, X_k, \overline{s})$ that passes through the point \bigstar and has the

same boundary orientation with L_i 's. In other words, such polygons u are identified with the polygons u' in $\mathcal{M}(p, X_1, \ldots, X_k, \overline{s})$ such that the holonomy of $\partial u'$ contains one η -factor.

Under the correspondence, it is easily checked that sign(u) and sign(u') are the same, and the holonomy of ∂u and $\partial u'$ contain zero and one η -factor, respectively, because they pass through the point \bigstar 0 and 1 times, respectively. As those polygons u' contribute to the factor $\eta \varphi_1$ in the decomposition of $\Phi^{\mathbb{L}}(\mathscr{L}(w', \eta, 1))$, their corresponding polygons u contribute φ_1 to the component (5.29) for j = i + 1.

Case ii) The boundary orientation of u differs from the orientations of L_i 's. Such a polygon u now have angles at $o_{12}, o_{23}, \ldots, o_{(j-1)j}$ consecutively as drawn in yellow in Figure 20. Ignoring those angles and all perturbations, u has an obvious counterpart u' in $\mathcal{M}(p, X_1, \ldots, X_k, \overline{s})$ that passes through the point \bigstar and has the opposite boundary orientation with L_i 's. In other words, such polygons u are identified with the polygons u' in $\mathcal{M}(p, X_1, \ldots, X_k, \overline{s})$ such that the holonomy of $\partial u'$ contains one η^{-1} -factor.

Under the correspondence, the quotient of sign(*u*) and sign(*u'*) is given by $(-1)^{j-i}$, because *u* have j-i more angles of odd degree that *u'*, where its boundary orientation is different from loops. The holonomy of ∂u and $\partial u'$ contain one $\eta^{-(j-i+1)}$ and η^{-1} factor, respectively, because they pass through the point \bigstar with the opposite orientation j - i + 1 and 1 times, respectively. As those polygons *u'* contribute to the factor $\eta^{-1}\varphi_{-1}$ in the decomposition of $\Phi^{\mathbb{L}}(\mathscr{L}(w',\eta,1))$, their corresponding polygons *u* contribute $-(-\eta)^{-(j-i+1)}\varphi_{-1}$ to the component (5.29).

To summarize, the polygons in Case i) and Case ii) contribute to $J_{\rho}(\eta) \otimes \varphi_1$ and $J_{\rho}(\eta)^{-1} \otimes \varphi_{-1}$ part of the given matrix factorization, respectively (while the $I_{\rho} \otimes \varphi_0$ part is not relevant to the twisting $o_{i(i+1)}$'s).

The second statement follows from bases change as done in Theorem 3.17. (One can also use (3.2).) \Box

Remark 5.23. Proposition 5.22 also reveals that the canonical form $\mathscr{L}(w',\eta,\rho)$ of loops with a local system corresponding to non-degenerate loop data (w',η,ρ) is quasi-isomorphic in TwFuk (Σ) to the abstract twisted complex (5.28), which is made of ρ -copies of $\mathscr{L}(w',\eta,1)$ and odd-degree morphism o's between them. In fact, it can be also derived from purely Fukaya-categorical discussions, not appealing to homological mirror symmetry. It has been shown in [Bae17, Theorem 5.8] that every higher rank local system over a loop can be realized as an abstract twisted complex of rank 1 local systems, using de Rham version of Fukaya category.

But our specific realization of recursive perturbations as in Figure 20 still suggests how we should perturb underlying loops when we want to work with the perturbation method instead of de Rham version. Especially when inputs of an A_{∞} -operation involve multiple e_L 's and o_L 's, its definition becomes more tricky and unsymmetric in some sense, which was not fully explained in Remark A.9. Nevertheless, some systematic recursive perturbations can be made so that the A_{∞} -relations remain valid.

This also gives an example where the mapping cone of two loops at their intersection is not quasiisomorphic to their surgery at that point, contrary to the usual situation which has been explained in many places in the literature including [Abo08, Lemma 5.4], [OPS18, Theorem 4.1] and [Boc21, Theorem 6.68]. This happens because two perturbed loops cannot satisfy the minimality condition, as also remarked in [AS21, Lemma 2.25]. We hope our explicit construction of twisted complexes involving o_L 's can be extended to realizing mapping cones of arbitrary morphisms in Fukaya category as geometric objects. We will come back to these points in another future work.

APPENDIX A. RELEVANT CATEGORIES

A.1. A_{∞} -category. Let us first recall the definition (and convention) of an A_{∞} -category over a field k and related concepts.

Definition A.1. $A \mathbb{Z}_2$ -graded A_∞ -category \mathscr{A} over \Bbbk consists of a class of objects $Ob(\mathscr{A})$, $a \mathbb{Z}_2$ -graded \Bbbk -vector space hom $(\mathscr{L}_0, \mathscr{L}_1) = hom^0(\mathscr{L}^0, \mathscr{L}^1) \oplus hom^1(\mathscr{L}^0, \mathscr{L}^1)$ for $\mathscr{L}_0, \mathscr{L}_1 \in Ob(\mathscr{A})$, and A_∞ -operations $\{\mathfrak{m}_k\}_{k\geq 1}$ given by \Bbbk -linear maps

$$\mathfrak{m}_k$$
: hom $(\mathscr{L}_0, \mathscr{L}_1) \otimes \cdots \otimes$ hom $(\mathscr{L}_{k-1}, \mathscr{L}_k) \rightarrow$ hom $(\mathscr{L}_0, \mathscr{L}_k)$

of degree $2 - k^{45}$ satisfying A_{∞} -relations

(A.1)
$$\sum_{0 \le i < j \le n} (-1)^{|f_1| + \dots + |f_i| - i} \mathfrak{m}_{n-j+i+1} (f_1, \dots, f_i, \mathfrak{m}_{j-i} (f_{i+1}, \dots, f_j), f_{j+1}, \dots, f_n) = 0$$

for any fixed $n \in \mathbb{Z}_{\geq 1}$ and morphisms $f_i \in \hom^{\bullet} (\mathscr{L}_{i-1}, \mathscr{L}_i)$ $(i \in \{1, ..., n\}, \bullet \in \mathbb{Z}_2)$.

It is called (strictly) **unital** if each object \mathcal{L} has a unit $\mathrm{id}_{\mathcal{L}} \in \mathrm{hom}(\mathcal{L}, \mathcal{L})$ satisfying

(A.2)
$$\mathfrak{m}_2(\mathrm{id}_{\mathscr{L}}, f) = f, \quad \mathfrak{m}_2(g, \mathrm{id}_{\mathscr{L}}) = (-1)^{|g|}g \quad and \quad \mathfrak{m}_k(\dots, \mathrm{id}_{\mathscr{L}}, \dots) = 0 \quad if \ k \neq 2$$

for any $\mathscr{L}' \in \mathrm{Ob}(\mathscr{A}), \ f \in \mathrm{hom}(\mathscr{L}, \mathscr{L}') \text{ and } g \in \mathrm{hom}^{\bullet}(\mathscr{L}', \mathscr{L}) \ (\bullet \in \mathbb{Z}_2).$

Definition A.2. An A_{∞} -functor $\mathscr{F} := \{\mathscr{F}_k\}_{k \ge 0}$ between two A_{∞} -categories \mathscr{A} and \mathscr{B} consists of a mapping $\mathscr{F}_0 : \operatorname{Ob}(\mathscr{A}) \to \operatorname{Ob}(\mathscr{B})$

and \Bbbk -linear maps ($k \ge 1$)

$$\mathscr{F}_k$$
: hom $_{\mathscr{A}}(\mathscr{L}_0, \mathscr{L}_1) \otimes \cdots \otimes hom_{\mathscr{A}}(\mathscr{L}_{k-1}, \mathscr{L}_k) \to hom_{\mathscr{B}}(\mathscr{F}_0(\mathscr{L}_0), \mathscr{F}_0(\mathscr{L}_k))$

of degree 1 - k, satisfying A_{∞} -relations

(A.3)
$$\sum_{1 \le k \le n} \sum_{1 \le i_1 < \dots < i_k = n} \mathfrak{m}_k^{\mathscr{B}} \left(\mathscr{F}_{i_1} \left(f_1, \dots, f_{i_1} \right), \dots, \mathscr{F}_{n-i_{k-1}} \left(f_{i_{k-1}+1}, \dots, f_n \right) \right) \\ = \sum_{0 \le i < j \le n} (-1)^{|f_1| + \dots + |f_i| - i} \mathscr{F}_{n-j+i+1} \left(f_1, \dots, f_i, \mathfrak{m}_{j-i}^{\mathscr{A}} \left(f_{i+1}, \dots, f_j \right), f_{j+1}, \dots, f_n \right)$$

for any fixed $n \in \mathbb{Z}_{\geq 1}$ and morphisms $f_i \in \hom_{\mathscr{A}}^{\bullet}(\mathscr{L}_{i-1}, \mathscr{L}_i)$ $(i \in \{1, ..., n\}, \bullet \in \mathbb{Z}_2)$.

The **composition** $\mathscr{G} \circ \mathscr{F} := \{(\mathscr{G} \circ \mathscr{F})_k\}_{k \ge 0}$ *of two* A_{∞} *-functors* $\mathscr{F} : \mathscr{A} \to \mathscr{B}$ *and* $\mathscr{G} : \mathscr{B} \to \mathscr{C}$ *is given by*

$$(\mathscr{G} \circ \mathscr{F})_n \left(f_1, \dots, f_n \right) := \sum_{1 \le k \le n} \sum_{1 \le i_1 < \dots < i_k = n} \mathscr{G}_k \left(\mathscr{F}_{i_1} \left(f_1, \dots, f_{i_1} \right), \dots, \mathscr{F}_{n-i_{k-1}} \left(f_{i_{k-1}+1}, \dots, f_n \right) \right).$$

An A_{∞} -functor $\mathcal{F} : \mathcal{A} \to \mathcal{B}$ is called **unital** if \mathcal{A} , \mathcal{B} are unital and

 $\mathscr{F}_1(\mathrm{id}_{\mathscr{L}}) = \mathrm{id}_{\mathscr{F}_0(\mathscr{L})} \quad and \quad \mathscr{F}_k(\ldots,\mathrm{id}_{\mathscr{L}},\ldots) = 0 \quad if k \ge 2$

for any $\mathcal{L} \in \mathrm{Ob}(\mathcal{A})$.

The \mathbb{Z}_2 -graded \Bbbk -vector space hom $(\mathscr{L}_0, \mathscr{L}_1) = \text{hom}^0(\mathscr{L}_0, \mathscr{L}_1) \oplus \text{hom}^1(\mathscr{L}_0, \mathscr{L}_1)$ has a degree 1 map \mathfrak{m}_1 : hom $(\mathscr{L}_0, \mathscr{L}_1) \to \text{hom}(\mathscr{L}_0, \mathscr{L}_1)$ satisfying $\mathfrak{m}_1^2 = 0$ by the A_∞ -relation for n = 1. Therefore, it becomes a cochain complex equipped with a **differential** \mathfrak{m}_1 . Taking its cohomology yields an ordinary category:

Definition A.3. For a unital A_{∞} -category \mathcal{A} , its **cohomological category** $H^0(\mathcal{A})$ is an ordinary category whose objects are the same as \mathcal{A} and the morphism space between two objects \mathcal{L}_0 , \mathcal{L}_1 is given by

$$\operatorname{Hom}_{H^{0}(\mathscr{A})}(\mathscr{L}_{0},\mathscr{L}_{1}):=H^{0}(\operatorname{hom}(\mathscr{L}_{0},\mathscr{L}_{1}),\mathfrak{m}_{1})$$

A unital A_{∞} -functor $\mathcal{F} : \mathcal{A} \to \mathcal{B}$ induces an ordinary functor $H^{0}(\mathcal{F}) : H^{0}(\mathcal{A}) \to H^{0}(\mathcal{B})$, whose mapping on objects is \mathcal{F}_{0} and action on morphisms is given by $[f] \mapsto [\mathcal{F}_{1}(f)]$.

⁴⁵Here it is just *k* as we are using the \mathbb{Z}_2 -grading.

A.2. **Compact Fukaya category of a surface.** We establish our geometric setup of the Fukaya category. We refer to [FOOO09, Sei08, AJ10] for its general definitions and properties, [Abo08, Sei11] for Fukaya category of surfaces, and [Aur14, Bae17, Kon17] for higher rank vector bundles in Fukaya category.

A.2.1. *Objects.* Let (Σ, ω) be a 2-dimensional symplectic manifold (possibly with boundary) of finite type. That is, Σ is a connected oriented smooth surface (with boundary) of finite type and ω is an area form on it. Then any smooth curve L in Σ automatically satisfies the Lagrangian conditions $(\omega|_L = 0, \dim L = \frac{1}{2} \dim \Sigma)$ and hence is a Lagrangian submanifold in Σ .

Consider an immersed oriented smooth loop $L: S^1 \to \Sigma \setminus \partial \Sigma$ having only transversal self-intersections. We assign two distinct **marked points** e_L, o_L ⁴⁶ on the image of *L* away from its self-intersections, and call the triple (L, e_L, o_L) a **marked loop** in Σ . When there is no need to specify marked points, we will call a marked loop just a **loop**, and denote it shortly as *L*.

Definition A.4. A set \mathcal{O} consisting of some marked loops in Σ is called **transversal** if it satisfies the following:

- Any two distinct loops in \mathcal{O} meet transversally.
- There are no triple intersections among loops in \mathcal{O} .
- Marked points of each loop in \mathcal{O} do not lie on any intersection of itself or any other loop in \mathcal{O} .

Definition A.5. (1) A loop $L: S^1 \to \Sigma$ is called **obstructed** if it bounds an immersed disk or 'fish-tale'. This means that there is an immersion $i: D^2 \to \Sigma$ which satisfies $i(e^{2\pi i t}) = L(\iota(t))$ for some immersion $\iota: [0, 1] \to S^1$. Otherwise, L is called **unobstructed**. A transversal set \mathcal{O} of marked loops in Σ is called **unobstructed** if all of its elements are unobstructed ⁴⁷.

(2) A transversal set \mathcal{O} of marked loops in Σ is called **full** if it contains at least one element in each primitive free homotopy class other than the null-homotopic one.

Given a full unobstructed set \mathcal{O} of marked loops in Σ , we define a \mathbb{Z}_2 -graded **compact Fukaya category** Fuk $(\Sigma) = \operatorname{Fuk}_{\mathbb{C}}^{\mathbb{Z}_2}(\Sigma; \mathcal{O})$ with respect to \mathcal{O} over \mathbb{C} .

Definition A.6. An object of Fuk (Σ) is given by a triple $\mathcal{L} = (L, E, \nabla)$, which consists of the following:

- a marked loop $L: S^1 \to \Sigma$ in \mathcal{O} ,
- $a \mathbb{C}$ -vector bundle E of finite rank ρ over the domain of L, and
- a flat ⁴⁸ connection ∇ on E.



Figure 21. Objects in Fuk (Σ) where $\Sigma = S^2 \setminus \{3 \text{ points}\}$

⁴⁶Equivalently, we can assign a Morse function $f_L : S^1 \to \mathbb{R}$ on the domain of L which has a minimum at e_L and a maximum at o_L so that they are all critical points of f_L . Graph of its differential df_L induces a C^0 -small Hamiltonian perturbation $\phi_H(L)$ in a neighborhood of L so that they make transversal intersections at e_L and o_L .

⁴⁷We need this condition for the A_{∞} -relations (without \mathfrak{m}_0 -terms) to hold. See [Abo08].

 $^{^{48}}$ In fact, every connection is flat in this case because dim $S^1 = 1$, but we still stick to the terminology to emphasize that it defines a local system.

For a computational purpose, we want to (not globally) trivialize *E* so that the parallel transport between two points is trivial away from a special point on *L*. For that, we choose a point \bigstar on (the domain of) *L* avoiding any intersection points (also with other loops) and denote by

$$\operatorname{hol}_{\bigstar}(E) \in \operatorname{Hom}(E|_{\bigstar}, E|_{\bigstar})$$

the holonomy of (E, ∇) along L at \bigstar ⁴⁹. We choose an identification $E|_{\bigstar} \cong \mathbb{C}^{\rho}$ where ρ is the rank of E, then hol $_{\bigstar}(E)$ is represented by some matrix $H \in \operatorname{GL}_{\rho}(\mathbb{C})$. (We simply say that the object (L, E, ∇) has a holonomy H (at \bigstar).)

Proposition A.7. There is a trivialization

$$E|_{S^1 \setminus \bigstar} \cong (S^1 \setminus \bigstar) \times \mathbb{C}^{\rho}$$

of E over $S^1 \setminus \bigstar$ satisfying the following: For any two points p and q on (the domain of) L other than \bigstar , the parallel transport

$$P(L_{p \to q}) \in \operatorname{Hom}(E|_p, E|_q)$$

from $E|_p$ to $E|_q$ along L (in the shortest way from p to q following the orientation of L) is represented by

(1) *H* if there is \bigstar in the way from *p* to *q*, and

(2) I_{ρ} (the identity matrix) otherwise,

with respect to the induced identifications $E|_p \cong \mathbb{C}^{\rho}$ and $E|_q \cong \mathbb{C}^{\rho}$.

Proof. Note that the linear isomorphisms $P(L_{\bigstar \rightarrow p}) : E_{\bigstar} \xrightarrow{\cong} E_p$ for each $p \in S^1 \setminus \bigstar$ yield the bundle isomorphism

In case (1),

$$P(L_{p \to q}) = P(L_{\bigstar \to p})^{-1} \circ \operatorname{hol}_{\bigstar}(E) \circ P(L_{\bigstar \to q}) : E_p \xrightarrow{\cong} E_{\bigstar} \to E_{\bigstar} \xrightarrow{\cong} E_q$$

coincides with the map $\operatorname{hol}_{\bigstar}(E): E_{\bigstar} \to E_{\bigstar}$ under the identifications of E_p and E_q with E_{\bigstar} .

In case (2),

$$P(L_{p \to q}) = P(L_{\bigstar \to p})^{-1} \circ P(L_{\bigstar \to q}) : E_p \xrightarrow{\cong} E_{\bigstar} \xrightarrow{\cong} E_q$$

is just the identity on E_{\bigstar} under the same identifications.

A.2.2. *Morphisms*. Given two objects $\mathscr{L}_0 = (L_0, E_0, \nabla_0)$ and $\mathscr{L}_1 = (L_1, E_1, \nabla_1)$ in Fuk (Σ) , roughly speaking, their **morphism space** hom $(\mathscr{L}_0, \mathscr{L}_1)$ is defined as a direct sum of \mathbb{Z}_2 -graded \mathbb{C} -vector spaces attached to each (self-)intersection of underlying curves L_0 and L_1 . We will explain the attached vector spaces below by dividing it into two cases:

In the case $L_0 \neq L_1$, as they are compact, they have finitely many intersection points. We define the set

$$\chi(L_0, L_1) := L_0 \cap L_1,$$

which is divided into an *even*-part and an *odd*-part according to orientations of two curves at each element. An element $p \in \chi(L_0, L_1)$ has an *even*-degree if the orientation of $T_pL_1 \oplus T_pL_0$ agrees with that of $T_p\Sigma$, and an *odd*-degree otherwise. Both situations are compared in Figure 22. We use + or – signs to indicate that p is even or odd, respectively, writing |p| = 0 or 1. We denote by $\chi^0(L_0, L_1)$ (the even-part) and $\chi^1(L_0, L_1)$ (the odd-part) the subsets of $\chi(L_0, L_1)$ consisting of even-degree and odd-degree elements, respectively.

⁴⁹It is the only invariant of flat bundles under gauge equivalence. Two gauge equivalent flat bundles (E_1, ∇_1) and (E_2, ∇_2) define quasi-isomorphic objects in the Fukaya category. See, for example, Proposition 4.9 in [Bae17].

Each intersection of L_0 and L_1 contributes one element to each of $\chi(L_0, L_1)$ and $\chi(L_1, L_0)$. We distinguish them by denoting one by p and the other by \overline{p} . Note that $|p| + |\overline{p}| = 1$ always holds. It is convenient to view $p \in \chi(L_0, L_1)$ as a pair of clockwise opposite angles from L_0 to L_1 , and $\overline{p} \in \chi(L_1, L_0)$ as the angles from L_1 to L_0 , as described in Figure 22.



In the case $L := L_0 = L_1$, it has finitely many self-intersections. In a small neighborhood of such a selfintersection p, there are two pieces \tilde{L}_a , \tilde{L}_b of L meeting at p. We may assume that the orientation of $T_p \tilde{L}_b \oplus T_p \tilde{L}_a$ agrees with that of $T_p \Sigma$. As in Figure 23, the pair of clockwise opposite convex angles from \tilde{L}_a to \tilde{L}_b is denoted by p and has an even-degree. Its adjacent pair of clockwise opposite convex angles from \tilde{L}_b to \tilde{L}_a is denoted by \overline{p} and has an odd-degree. Two (pair-of-)angles p and \overline{p} will have different meaning when we count polygons involving them.

We consider the marked points e_L and o_L to have even and odd degrees, respectively. We define the set

$$\chi(L,L) := \{e_L, o_L\} \cup \{p, \overline{p} \mid p : a \text{ self-intersection point of } L\},$$

which is divided into two subsets $\chi^0(L, L)$ and $\chi^1(L, L)$ as before.

In both cases, we define the A_{∞} -morphism spaces as

(A.4)
$$\hom^{\bullet}(\mathscr{L}_{0},\mathscr{L}_{1}) := \bigoplus_{p \in \chi^{\bullet}(L_{0},L_{1})} \operatorname{Hom}_{\mathbb{C}}(E_{0}|_{p},E_{1}|_{p}) \quad (\bullet \in \mathbb{Z}_{2})$$

where $E_0|_p$ and $E_1|_p$ are the fibers of E_0 and E_1 over the preimages (in S^1) of the point $p \in \Sigma$ under L_0 and L_1 (or under different branches \tilde{L}_a and \tilde{L}_b of L in the case $L := L_0 = L_1$), respectively. They yield a \mathbb{Z}_2 -graded morphism space

 $\hom(\mathscr{L}_0,\mathscr{L}_1) := \hom^0(\mathscr{L}_0,\mathscr{L}_1) \oplus \hom^1(\mathscr{L}_0,\mathscr{L}_1).$

An element $f \in \text{Hom}_{\mathbb{C}}(E_0|_p, E_1|_p)$ for each $p \in \chi(L_0, L_1)$ is called a *base morphism* over p, and we denote it by $f|_p$ to specify that fact.

Remark A.8. According to Proposition A.7, we have identifications $E_0|_p \cong \mathbb{C}^{\rho_0}$ and $E_1|_p \cong \mathbb{C}^{\rho_1}$, which yield $\operatorname{Hom}_{\mathbb{C}}(E_0|_p, E_1|_p) \cong \mathbb{C}^{\rho_1 \times \rho_0}$. Denoting by p_{ab} the generator corresponding to the $\rho_1 \times \rho_0$ matrix whose the only nonzero entry is 1 in the (a, b)-th position, we have

$$\hom^{\bullet}(\mathscr{L}_{0},\mathscr{L}_{1}) \cong \bigoplus_{\substack{p \in \chi^{\bullet}(L_{0},L_{1}) \\ 1 \le b \le \rho_{0}}} \operatorname{Span}_{\mathbb{C}} \left\{ p_{ab} \right\} \quad (\bullet \in \mathbb{Z}_{2}).$$

In the simplest case $\rho_0 = \rho_1 = 1$, we just write

$$\hom^{\bullet}(\mathscr{L}_{0},\mathscr{L}_{1}) \cong \bigoplus_{p \in \chi^{\bullet}(L_{0},L_{1})} \operatorname{Span}_{\mathbb{C}} \left\{ p \right\} = \operatorname{Span}_{\mathbb{C}} \left(\chi^{\bullet}(L_{0},L_{1}) \right) \quad (\bullet \in \mathbb{Z}_{2}).$$

A.2.3. A_{∞} -operations. The A_{∞} -operations $\{\mathfrak{m}_k\}_{k\geq 1}$ on base morphisms between objects $\mathscr{L}_i = (L_i, E_i, \nabla_i)$ $(i \in \{0, ..., k\})$ are defined as

(A.5)
$$\mathfrak{m}_{k}: \hom\left(\mathscr{L}_{0}, \mathscr{L}_{1}\right) \otimes \cdots \otimes \hom\left(\mathscr{L}_{k-1}, \mathscr{L}_{k}\right) \to \hom\left(\mathscr{L}_{0}, \mathscr{L}_{k}\right) \\ \left(f_{1}\big|_{p_{1}}, \dots, f_{k}\big|_{p_{k}}\right) \mapsto \sum_{q \in \chi(L_{0}, L_{k})} \sum_{u \in \mathscr{M}(p_{1}, \dots, p_{k}, \overline{q})} \operatorname{sign}\left(u\right) \operatorname{hol}_{q}\left(\partial u\right)\left(f_{1}, \dots, f_{k}\right)$$

for any $p_i \in \chi(L_{i-1}, L_i)$ other than e_L or o_L ⁵⁰, such that $p_{i+1} \neq \overline{p_i}$, and $f_i = f_i|_{p_i} \in \text{Hom}(E_{i-1}|_{p_i}, E_i|_{p_i})$. Then it is linearly extended to other morphisms. We will explain the meaning of each component one by one below.

First, the **moduli space** $\mathcal{M}(p_1, \dots, p_k, \overline{q})$ is the set of immersed (k + 1)-gons bounded by L_0, L_1, \dots, L_k whose angles consist of $p_1, \dots, p_k, \overline{q}$ in counter-clockwise order. To be precise, it means a continuous map $u: D^2 \to \Sigma$ together with k+1 points $z_1, \dots, z_k, z_0 \in \partial D^2$ (in counterclockwise order) such that the segment of ∂D^2 between z_i and z_{i+1} is mapped to L_i ($i \in \mathbb{Z}_{k+1}$), the image of u has a convex corner p_i at $u(z_i) = p_i$ ($p_{k+1} := \overline{q}$), and u is an orientation-preserving immersion on $D^2 \setminus \{z_1, \dots, z_k, z_0\}$. We consider such maps up to automorphisms of the domain D^2 , that is, ($\{z_1, \dots, z_k, z_0\}, u$) and ($\{z'_1, \dots, z'_k, z'_0\}, u'$) define the same element in the moduli space if and only if there is a homeomorphism $\phi: D^2 \to D^2$ such that $\phi(z_i) = z'_i$ ($i \in \mathbb{Z}_{k+1}$), u is a diffeomorphism on $D^2 \setminus \{z_1, \dots, z_k, z_0\}$, and $u' = u \circ \phi$.



Figure 24. An element u of $\mathcal{M}(p_1, \ldots, p_k, \overline{q})$

Second, to determine the **sign** of *u*, we follow the sign rule established and illustrated in [Sei08, Sei11]: Consider the boundary orientation on ∂u as usual, that is, it is given in such a way that *u* lies on the left along it. The orientation of L_0 is irrelevant. For $1 \le i \le k$, whenever the orientation of L_k does not match the orientation of ∂u , $(-1)^{|p_i|}$ is contributed to sign (*u*), which changes the sign only when the angle p_i from L_{i-1} to L_i has odd-degree. In addition, if the orientation of L_k differs from that of ∂u , the sign $(-1)^{|q|}$ of the output angle *q* from L_0 to L_k is also contributed. Summing up, we have

where o(L) denotes the orientation of a curve L and $\mathbb{I}_{statement}$ is 1 if the statement is true and 0 otherwise.

 $[\]overline{}^{50}$ If e_L or o_L 's are involved as inputs, the definition of A_{∞} -operation becomes much complicated. See Remark A.9 for some cases.

Third, the **holonomy operation** of ∂u at *q* is defined as

$$\operatorname{hol}_{q}(\partial u) : \operatorname{Hom}_{\mathbb{C}}\left(E_{0}|_{p_{1}}, E_{1}|_{p_{1}}\right) \otimes \cdots \otimes \operatorname{Hom}_{\mathbb{C}}\left(E_{k-1}|_{p_{k}}, E_{k}|_{p_{k}}\right) \to \operatorname{Hom}_{\mathbb{C}}\left(E_{0}|_{p_{0}}, E_{k}|_{p_{0}}\right)$$
$$\left(f_{1}, \dots, f_{k}\right) \mapsto P\left((\partial u)_{k}\right) \circ f_{k} \circ P\left((\partial u)_{k-1}\right) \circ f_{k-1} \circ \cdots \circ f_{2} \circ P\left((\partial u)_{1}\right) \circ f_{1} \circ P\left((\partial u)_{0}\right)$$

where

$$P((\partial u)_i) \in \operatorname{Hom}_{\mathbb{C}}(E_i|_{p_i}, E_i|_{p_{i+1}})$$

is the parallel transport with respect to ∇_i from $E_i|_{p_i}$ to $E_i|_{p_{i+1}}$ along the side of u (following the boundary orientation) lying in L_i .

Remark A.9. When o_{L_i} or e_{L_i} are involved in the moduli space $\mathcal{M}(p_1, ..., p_k, p_{k+1})$, an appropriate perturbation scheme can be introduced. Here we follow [Sei11] and explain some special cases where we have only one of them in input or output of the A_{∞} -operations, which will be enough for our purpose ⁵¹.

(1) If $\mathcal{L}_{i-1} = \mathcal{L}_i$ ($i \in \{1, ..., k\}$) (and hence $L_{i-1} = L_i$ is a loop) and an input p_i is o_{L_i} from $L_{i-1} = L_i$ to itself, then elements of $\mathcal{M}(p_1, ..., p_{i-1}, o_{L_i}, p_{i+1}, ..., p_k, \overline{q})$ are polygons which have convex corners at $p_1, ..., p_{i-1}$, pass through the point o_{L_i} , and then again have convex corners at $p_{i+1}, ..., p_k, \overline{q}$ in counterclockwise order.

(2) If $\mathcal{L}_{i-1} = \mathcal{L}_i$ $(i \in \{1, ..., k\})$ and p_i is $e_{L_{i-1}} = e_{L_i}$, the set $\mathcal{M}(p_1, ..., e_{L_i}, ..., p_k, \overline{q})$ is empty unless k = 2, which yields

(A.7)
$$\mathfrak{m}_k\left(\ldots,\operatorname{id}_{e_{L_i}},\ldots\right) = 0 \quad if \, k \neq 2.$$

 e_L to any point $p \in \chi(L,L')$ (which doesn't pass through o_L) as an 'infinitesimal triangle' whose angles are e_L , p and \overline{p} . It provides the unique element of $\mathcal{M}(e_L, p, \overline{p})$ and $\mathcal{M}(\overline{p}, e_L, p)$, which yields

(A.8)
$$\mathfrak{m}_{2}\left(\mathrm{id}|_{e_{L}},f|_{p}\right)=f|_{p}\quad and\quad \mathfrak{m}_{2}\left(g|_{\overline{p}},\mathrm{id}|_{e_{L}}\right)=(-1)^{|\overline{p}|}g|_{\overline{p}}$$

for any $f|_p \in \hom(\mathcal{L}, \mathcal{L}')$ and $g|_{\overline{p}} \in \hom(\mathcal{L}', \mathcal{L})$. Equations (A.7) and (A.8) imply that $\operatorname{id}_{\mathcal{L}} := \operatorname{id}|_{e_L}$ is a unit of \mathcal{L} .

(3) If $\mathcal{L}_0 = \mathcal{L}_k =: \mathcal{L}$ and the output q is e_L , elements of $\mathcal{M}(p_1, \dots, p_k, \overline{e_L})$ are polygons which have convex corners at p_1, \dots, p_k and then pass through the point e_L in counterclockwise order.

Remark A.10. In fact, the definition given in (A.5) involves a crucial problem. That is, sometimes there are infinitely many elements in the moduli space $\mathcal{M}(p_1, ..., p_k, \overline{q})$. Therefore, a priori, we must work over the Novikov field

$$\Lambda := \left\{ \sum_{i=0}^{\infty} a_i T^{\lambda_i} \middle| a_i \in \mathbb{C}, \ \lambda_i \in \mathbb{R}, \ \lim_{i \to \infty} \lambda_i = \infty \right\}$$

instead of the base field \mathbb{C} (which makes the hom spaces Λ -vector spaces) and define the A_{∞} -operations as

$$\mathfrak{m}_k$$
: hom $(\mathscr{L}_0, \mathscr{L}_1) \otimes \cdots \otimes$ hom $(\mathscr{L}_{k-1}, \mathscr{L}_k) \rightarrow$ hom $(\mathscr{L}_0, \mathscr{L}_k)$

$$\left(f_1\big|_{p_1},\ldots,f_k\big|_{p_k}\right)\mapsto \sum_{q\in\chi(L_0,L_k)}\sum_{u\in\mathcal{M}(p_1,\ldots,p_k,\overline{q})}\operatorname{sign}(u) T^{\omega(u)}\operatorname{hol}_q(\partial u)\left(f_1,\ldots,f_k\right)$$

instead of (A.5). But we demonstrated in [CJKR22] that we can evaluate T = 1 when we compute the matrix factorizations corresponding to cylinder-free loops under the localized mirror functor (See Definition 3.12 and Theorem 3.13.). In this paper, therefore, we still work over \mathbb{C} and use the definition (A.5) when computing the mirror images of cylinder-free loops.

⁵¹But see also Figure 20 for the case where many o_{L_i} 's are involved.

A.3. **Categories of matrix factorizations.** Let *S* be the formal power series ring $\mathbb{C}[[x_1, ..., x_m]]$ of *m* variables, and $f \in S$ its nonzero element.

Definition A.11. A matrix factorization of f(in S) is a pair (φ, ψ) of S-module homomorphisms $P^0 \xleftarrow{\psi} P^1$ between two finite-rank free S-modules P^0 and P^1 that satisfy

$$\psi \varphi = f \cdot \mathrm{id}_{P^0}$$
 and $\varphi \psi = f \cdot \mathrm{id}_{P^1}^{52}$.

There are several versions of the *categories of matrix factorizations* of f. They all have matrix factorizations of f as objects, but morphism spaces are different, which we will now explain:

Given two matrix factorizations $P_0^0 \xrightarrow{\varphi_0} P_0^1$ and $P_1^0 \xrightarrow{\varphi_1} P_1^1$ of f, an *even-degree* or *odd-degree* **morphism** between them is a pair of *S*-module maps $(\alpha : P_0^0 \to P_1^0, \beta : P_0^1 \to P_1^1)$ or $(\gamma : P_0^0 \to P_1^1, \delta : P_0^1 \to P_1^0)$, respectively. They consist the *even-part* and *odd-part* of the morphism space defined as

 $\hom^{\bullet}\left(\left(\varphi_{0},\psi_{0}\right),\left(\varphi_{1},\psi_{1}\right)\right):=\operatorname{Hom}_{S}\left(P_{0}^{0},P_{1}^{\bullet}\right)\times\operatorname{Hom}_{S}\left(P_{0}^{1},P_{1}^{1+\bullet}\right) \quad (\bullet \in \mathbb{Z}_{2}),$

and form the following diagram but are not required to commute it:

The \mathbb{Z}_2 -graded morphism space is a \mathbb{Z}_2 -graded \mathbb{C} -vector space defined as

$$\hom^{\mathbb{Z}_2}\left(\left(\varphi_0,\psi_0\right),\left(\varphi_1,\psi_1\right)\right) := \hom^0\left(\left(\varphi_0,\psi_0\right),\left(\varphi_1,\psi_1\right)\right) \oplus \hom^1\left(\left(\varphi_0,\psi_0\right),\left(\varphi_1,\psi_1\right)\right).$$

We define a \mathbb{C} -linear map d of degree 1 on it as

(A.9)

$$\hom^{0}((\varphi_{0},\psi_{0}),(\varphi_{1},\psi_{1})) \xrightarrow{d} \hom^{1}((\varphi_{0},\psi_{0}),(\varphi_{1},\psi_{1})) \quad \text{and} \quad \hom^{1}((\varphi_{0},\psi_{0}),(\varphi_{1},\psi_{1})) \xrightarrow{d} \hom^{0}((\varphi_{0},\psi_{0}),(\varphi_{1},\psi_{1})) \xrightarrow{d} (\varphi_{0},\psi_{0}),(\varphi_{1},\psi_{1})) \xrightarrow{d} (\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1})) \xrightarrow{d} (\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1})) \xrightarrow{d} (\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1})) \xrightarrow{d} (\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}),(\varphi_{1},\psi_{1}))$$

It satisfies $d^2 = 0$, making the morphism space a 2-periodic cochain complex with a **differential** *d*.

Definition A.12. We define four different **categories of matrix factorizations** of f. Their objects are matrix factorizations of f. The **morphism space** between two matrix factorizations (φ_0, ψ_0) and (φ_1, ψ_1) will be defined, while the composition of morphisms and the identity morphisms are given in the obvious way.

(1) In differential \mathbb{Z}_2 -graded category $MF_{dg}(f)$, it is given by $(\hom^{\mathbb{Z}_2}((\varphi_0, \psi_0), (\varphi_1, \psi_1)), d)$, which is a \mathbb{Z}_2 -graded \mathbb{C} -vector space equipped with the differential d.

(2) In (ordinary) category MF(f), it is given by $Z^0(\hom^{\mathbb{Z}_2}((\varphi_0, \psi_0), (\varphi_1, \psi_1)), d)$, which consists of evendegree morphisms $(\alpha, \beta) \in \hom^0((\varphi_0, \psi_0), (\varphi_1, \psi_1))$ that commute the left diagram in (A.9).

(3) In homotopy category $\underline{MF}(f)$, it is given by $H^0(\hom^{\mathbb{Z}_2}((\varphi_0, \psi_0), (\varphi_1, \psi_1)), d)$. This category is the same as the stable category $\underline{MF}(f) := \underline{MF}(f)/\{(1, f), (f, 1)\}$ (Definition A.14).

(4) In (\mathbb{Z}_2 -graded) A_∞ -category MF $_{A_\infty}(f)$, it is given by the \mathbb{Z}_2 -graded \mathbb{C} -vector space hom $\mathbb{Z}_2((\varphi_1, \psi_1), (\varphi_0, \psi_0))$. (Note that it is reversed.) The A_∞ -operations $\{\mathfrak{m}_k\}_{k\geq 1}$ are defined as:

(A.10) $\mathfrak{m}_1 := d, \quad \mathfrak{m}_2((\alpha_0, \beta_0), (\alpha_1, \beta_1)) := (-1)^{\bullet - 1}(\alpha_0, \beta_0) \circ (\alpha_1, \beta_1), \quad and \quad \mathfrak{m}_{k \ge 3} := 0$

$$for(\alpha_0,\beta_0) \in \hom^{\bullet}((\varphi_1,\psi_1),(\varphi_0,\psi_0)) (\bullet \in \mathbb{Z}_2) and (\alpha_1,\beta_1) \in \hom^{\mathbb{Z}_2}((\varphi_2,\psi_2),(\varphi_1,\psi_1)).$$

⁵²These conditions imply that P^0 and P^1 have the same rank. Moreover, ψ is completely determined by φ and vice versa.

A.4. Stable categories and Eisenbud's equivalence. ⁵³ Let *S* be the power series ring $\mathbb{C}[[x_1,...,x_m]]$ of *m* variables, $f \in S$ its nonzero element and A := S/(f) the quotient ring. In this case, we can relate the category CM(*A*) of maximal Cohen-Macaulay modules over *A* (Definition 4.1) and the category MF(*f*) of matrix factorizations of *f* (Definition A.12(2)) in a *stable* sense. We need some preparation:

Definition A.13. Let \mathscr{A} be a category whose Hom-sets are abelian groups, and P a set of some objects in \mathscr{A} . For any two objects M, N in \mathscr{A} , we denote by I(M,N) the subgroup of $\operatorname{Hom}_{\mathscr{A}}(M,N)$ generated by all morphisms from M to N that factor through a direct sum of objects in P. We define the quotient of \mathscr{A} by P as the category \mathscr{A} / P whose objects are the same as \mathscr{A} , and the morphism spaces are defined as

$$\operatorname{Hom}_{\mathscr{A}/P}(M,N) := \operatorname{Hom}_{\mathscr{A}}(M,N) / I(M,N).$$

Note that any objects in P are zero objects in the category \mathscr{A}/P , and it is the largest quotient of \mathscr{A} with this property.

Definition A.13 enables us to define the *stable categories* of maximal Cohen-Macaulay modules over A and matrix factorizations of f^{54} :

Definition A.14. (1) The stable category of maximal Cohen-Macaulay modules over A is defined as

$$\underline{\mathrm{CM}}(A) := \mathrm{CM}(A) / \{A\}.$$

(2) The stable category of matrix factorizations of f is defined as

$$\underline{\mathrm{MF}}(f) := \mathrm{MF}(f) / \{(1, f), (f, 1)\}$$

The following justifies our use of the same notation as the homotopy category MF(f) in Definition A.12:

Proposition A.15. The stable category and the homotopy category of matrix factorizations of f are the same.

Proof. A more general statement and proof for the identification of the stable category and the homotopy category of cochain complexes in an additive category can be found in §4.1 in [Kra21]. Here we provide an explicit proof for our setting:

We actually show that their Hom-sets are the same. For that, it is enough to show

$$I((\varphi,\psi),(\varphi',\psi')) = B^0(\hom^{\mathbb{Z}_2}((\varphi,\psi),(\varphi',\psi')),d)$$

for any matrix factorizations $P^0 \xleftarrow{\varphi}{\psi} P^1$ and $P'^0 \xleftarrow{\varphi'}{\psi'} P'^1$ of f.

First let $(\alpha : P^0 \to P'^0, \beta : P^1 \to P'^1)$ be a generator of $I((\varphi, \psi), (\varphi', \psi'))$, i.e., it factors through a direct sum of (1, f)'s and (f, 1)'s, which is written as $S^a \oplus S^b \xrightarrow{\begin{pmatrix} I_a & 0 \\ 0 & fI_b \end{pmatrix}}{\begin{pmatrix} fI_a & 0 \\ 0 & I_b \end{pmatrix}} S^a \oplus S^b$ for some $a, b \in \mathbb{Z}_{\geq 0}$ without loss of generality. That is, (α, β) is a composition $(\alpha'' \circ \alpha', \beta'' \circ \beta')$ of some morphisms that are described in

⁵³All discussions in this section (except for Proposition A.15) can be found in Chapter 7 in [Yos90].

⁵⁴In general, one can take the quotient of a *Frobenius category* by its projective objects to make it a triangulated category, called the *stable category*. See §3.3 in [Kra21].

(and commute the squares in) the following diagram:



Now we know that (α, β) is *null-homotopic* by checking that it is the differential of an odd-degree morphism $(\beta_2'' \circ \alpha_2' : P^0 \to P'^1, \alpha_1'' \circ \beta_1' : P^1 \to P'^0)$. This shows $(\alpha, \beta) \in B^0$ (hom $\mathbb{Z}_2((\varphi, \psi), (\varphi', \psi')), d$).

Conversely, an element of $B^0(\hom^{\mathbb{Z}_2}((\varphi,\psi),(\varphi',\psi')),d)$ is by definition given by the differential of an odd-degree morphism $(\gamma: P^0 \to P'^1, \delta: P^1 \to P'_0)$. As the differential is additive, we may assume $\delta = 0$ without loss of generality. Now the differential of $(\gamma,0)$ is $(\psi' \circ \gamma, \gamma \circ \psi)$, and the following commutative diagram shows that it factors through the direct sum of $P^0 \xleftarrow{\text{id}}{f \cdot \text{id}} P^0$ and $P'^1 \xleftarrow{f \cdot \text{id}}{P'^1} P'^1$, being an element of $I((\varphi,\psi),(\varphi',\psi'))$:

Now we can state Eisenbud's equivalence between two stable categories:

Theorem A.16 (Eisenbud's matrix factorization theorem [Eis80, Yos90]). A matrix factorization $P^0 \xleftarrow{\psi} P^1$ of f defines a 2-periodic acyclic chain complex of A-modules

$$\cdots \longrightarrow P^0 \otimes_S A \xrightarrow{\varphi} P^1 \otimes_S A \xrightarrow{\psi} P^0 \otimes_S A \xrightarrow{\varphi} P^1 \otimes_S A \xrightarrow{\psi} \cdots,$$

where $\underline{\varphi} := \varphi \otimes \operatorname{id}_A$ and $\underline{\psi} := \psi \otimes \operatorname{id}_A$. Taking the cokernel of $\underline{\varphi}$ yields a maximal Cohen-Macaulay A-module $M := \operatorname{coker} \underline{\varphi}^{55}$. It defines a functor coker : MF(f) \rightarrow CM(A), which also induces a functor between stable categories

$$\operatorname{coker}: \underline{\operatorname{MF}}(f) \to \underline{\operatorname{CM}}(A).$$

Conversely, a maximal Cohen-Macaulay A-module M, regarded as an S-module M_S , admits a (not unique) free resolution

 $0 \longrightarrow S^n \xrightarrow{\varphi} S^n \longrightarrow M_S \longrightarrow 0.$

It determines another map $\psi : S^n \to S^n$ such that $\varphi \psi = \psi \varphi = f \cdot id_{S^n}$, yielding a matrix factorization (φ, ψ) of f. This process defines a quasi-inverse to the above induced functor, giving an equivalence of stable categories $\underline{MF}(f)$ and $\underline{CM}(A)$.

⁵⁵Note that *M* admits 2-periodic free resolution
$$\dots \to P^0 \otimes_S A \xrightarrow{\varphi} P^1 \otimes_S A \xrightarrow{\psi} P^0 \otimes_S A \xrightarrow{\varphi} P^1 \otimes_S A \to M = \operatorname{coker} \underline{\varphi} \to 0$$

References

 [Abo08] Mohammed Abouzaid, On the Fukaya categories of higher genus surfaces, Advances in Mathematics 217 (2008), n 1192–1235. [Abo12], On the wrapped Fukaya category and based loops, Journal of Symplectic Geometry 10 (2012), no. 1, 27–75 [AEK21] Denis Auroux, Alexander I Efimov, and Ludmil Katzarkov, Lagrangian floer theory for trivalent graphs and homolog mirror symmetry for curves, arXiv preprint arXiv:2107.01981 (2021). [AJ10] Manabu Akaho and Dominic Joyce, Immersed Lagrangian Floer theory, J. Differential Geom. 86 (2010), no. 3, 381– [AP24] Claire Amiot and Pierre-Guy Plamondon, Skew-group A_∞-categories as Fukaya categories of orbifolds, arXiv preprarXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, A complete derived invariant for gentle algebras via wind 	 b. 3, ical 600. rint 'ling 14 1 of
 [Abo12], On the wrapped Fukaya category and based loops, Journal of Symplectic Geometry 10 (2012), no. 1, 27–79 [AEK21] Denis Auroux, Alexander I Efimov, and Ludmil Katzarkov, Lagrangian floer theory for trivalent graphs and homolog mirror symmetry for curves, arXiv preprint arXiv:2107.01981 (2021). [AJ10] Manabu Akaho and Dominic Joyce, Immersed Lagrangian Floer theory, J. Differential Geom. 86 (2010), no. 3, 381– [AP24] Claire Amiot and Pierre-Guy Plamondon, Skew-group A_∞-categories as Fukaya categories of orbifolds, arXiv preprint arXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, A complete derived invariant for gentle algebras via wind 	ical 500. rint <i>ling</i> 14 1 of
 [AEK21] Denis Auroux, Alexander I Efimov, and Ludmil Katzarkov, Lagrangian floer theory for trivalent graphs and homolog mirror symmetry for curves, arXiv preprint arXiv:2107.01981 (2021). [AJ10] Manabu Akaho and Dominic Joyce, Immersed Lagrangian Floer theory, J. Differential Geom. 86 (2010), no. 3, 381– [AP24] Claire Amiot and Pierre-Guy Plamondon, Skew-group A_∞-categories as Fukaya categories of orbifolds, arXiv preprint arXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, A complete derived invariant for gentle algebras via wind 	<i>ical</i> 500. rint <i>ling</i> 14 1 of
 mirror symmetry for curves, arXiv preprint arXiv:2107.01981 (2021). [AJ10] Manabu Akaho and Dominic Joyce, Immersed Lagrangian Floer theory, J. Differential Geom. 86 (2010), no. 3, 381– [AP24] Claire Amiot and Pierre-Guy Plamondon, Skew-group A_∞-categories as Fukaya categories of orbifolds, arXiv preparXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, A complete derived invariant for gentle algebras via wind 	500. rint <i>ling</i> 14
 [AJ10] Manabu Akaho and Dominic Joyce, <i>Immersed Lagrangian Floer theory</i>, J. Differential Geom. 86 (2010), no. 3, 381– [AP24] Claire Amiot and Pierre-Guy Plamondon, <i>Skew-group A_∞-categories as Fukaya categories of orbifolds</i>, arXiv preparXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, <i>A complete derived invariant for gentle algebras via wind</i> 	500. rint <i>ling</i> 14 1 of
 [AP24] Claire Amiot and Pierre-Guy Plamondon, <i>Skew-group A_∞-categories as Fukaya categories of orbifolds</i>, arXiv preparXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, <i>A complete derived invariant for gentle algebras via wind</i> 	rint <i>ling</i> 14 1 of
arXiv:2405.15466 (2024). [APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, <i>A complete derived invariant for gentle algebras via wind</i>	<i>ing</i> 14 1 of
[APS23] Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll, A complete derived invariant for gentle algebras via wind	<i>111</i>
numbers and Arf invariants, Selecta Mathematica 29 (2023), no. 2, 30.	14 1 of
[AS10] Mohammed Abouzaid and Paul Seidel, <i>An open string analogue of Viterbo functoriality</i> , Geometry & Topolog (2010), no. 2, 627–718.	ı of
[AS21] Denis Auroux and Ivan Smith, Fukaya categories of surfaces, spherical objects and mapping class groups, Forur	.1.4
Mathematics, Sigma, vol. 9, Cambridge University Press, 2021, p. e26.	.1.4
[Aur14] Denis Auroux, <i>A beginner's introduction to Fukaya categories</i> , Contact and symplectic topology, Springer, 2 pp. 85–136.	14,
[Bae17] Hanwool Bae, <i>Higher rank vector bundles in Fukaya category</i> , Ph.D. thesis, Seoul national university, 2017.	
[BD08] Igor Burban and Yuriy Drozd, <i>Maximal Cohen–Macaulay modules over surface singularities</i> , Trends in representa theory of algebras and related topics 1 (2008), 101.	ion
[BD11], Tilting on non-commutative rational projective curves. Mathematische Annalen 351 (2011), 665–709.	
[BD17a] . Maximal Cohen-Macaulay modules over non-isolated surface singularities and matrix problems. vol.	248.
American Mathematical Society. 2017.	,
[BD17b], On the derived categories of gentle and skew-gentle algebras: homological algebra and matrix problems, a	Xiv
preprint arXiv:1706.08358 (2017).	
[BD18], Non-commutative nodal curves and derived tame algebras, arXiv preprint arXiv:1805.05174 (2018).	
[BH98] Winfried Bruns and H. Jürgen Herzog, <i>Cohen-Macaulav rings</i> , 2nd edition, Cambridge Studies in Advanced Ma	he-
matics, Cambridge University Press 39 (1998).	
[Boc16] Raf Bocklandt, Noncommutative mirror symmetry for punctured surfaces, Transactions of the American Mathema	ical
Society 368 (2016), no. 1, 429–469.	
[Boc21], A gentle introduction to homological mirror symmetry, vol. 99, Cambridge University Press, 2021.	
[Buc21] Ragnar-Olaf Buchweitz, Maximal Cohen–Macaulay modules and Tate cohomology, vol. 262, American Mathema	ical
Society, 2021.	
[BZ20] Igor Burban and Alexander Zheglov, Cohen-Macaulay modules over the algebra of planar quasi-invariants	and
Calogero-Moser systems, Proceedings of the London Mathematical Society 121 (2020), 1033.	
[CHL15] Cheol-Hyun Cho, Hansol Hong, and Siu-Cheong Lau, Noncommutative homological mirror functor, Memoirs of	the
American Mathematical Society 271 (2015).	
[CHL17], Localized mirror functor for Lagrangian immersions, and homological mirror symmetry for $\mathbb{P}^1_{a,b,s}$. J. Diffe	en-
tial Geom. 106 (2017), no. 1, 45–126. MR 3640007	
[CHL18], Gluing localized mirror functors, arXiv preprint (2018), arXiv: 1810.02045.	
[CHL19], Localized mirror functor constructed from a Lagrangian torus, Journal of Geometry and Physics 136 (20	19),
284–320.	- , ,
[CHS23] Wen Chang, Fabian Haiden, and Sibylle Schroll, Braid group actions on branched coverings and full exceptiona	se-
<i>quences</i> , arXiv preprint arXiv:2301.04398 (2023).	
[CIKR22] Cheol-Hvun Cho, Wonbo Jeong, Kvoungmo Kim, and Kvungmin Rho, Homological mirror symmetry of indecom	os-
able Cohen-Macaulay modules for some degenerate cusp singularities, arXiv preprint arXiv:2205.02469 (2022).	
[CJS22] Wen Chang, Haibo Jin, and Sibylle Schroll, Recollements of partially wrapped Fukaya categories and surface cuts, a	Xiv
preprint arXiv:2206.11196 (2022).	
[CK24] Cheol-Hyun Cho and Kyoungmo Kim, <i>Topological Fukaya category of tagged arcs</i> , arXiv preprint arXiv:2404.10	294
(2024).	
[Efi12] Alexander I Efimov, Homological mirror symmetry for curves of higher genus, Advances in Mathematics 2 (20	12),
no. 230, 493–530.	
[Eis80] David Eisenbud, <i>Homological algebra on a complete intersection, with an application to group representations</i> , Tractions of the American Mathematical Society 260 (1980), pp. 1, 35–64	.ns-

- [FOOO09] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, Lagrangian intersection Floer theory: anomaly and obstruction. Parts I and II, AMS/IP Studies in Advanced Mathematics, vol. 46, American Mathematical Society, Providence, RI, 2009.
- [HJL24] Hansol Hong, Hyeongjun Jin, and Sangwook Lee, *Orbifold kodaira-spencer maps and closed-string mirror symmetry for punctured riemann surfaces*, arXiv preprint arXiv:2402.14360 (2024).
- [HKK17] Fabian Haiden, Ludmil Katzarkov, and Maxim Kontsevich, *Flat surfaces and stability structures*, Publications mathématiques de l'IHÉS **126** (2017), 247–318.
- [Kon95] Maxim Kontsevich, *Homological algebra of mirror symmetry*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 120–139.
- [Kon98] _____, *Lectures at ens, paris, spring 1998*, notes taken by J. Bellaiche, J.-F. Dat, I. Marin, G. Racinet and H. Randriambololona, unpublished (1998).
- [Kon17] Momchil Konstantinov, Higher rank local systems in Lagrangian Floer theory, arXiv preprint arXiv:1701.03624 (2017).
- [Kra21] Henning Krause, *Homological theory of representations*, vol. 195, Cambridge University Press, 2021.
- [KŠ22] Jakub Kopřiva and Jan Šťovíček, *Semiorthogonal decompositions for bounded derived categories of gentle algebras,* arXiv e-prints (2022), arXiv–2209.
- [Lee16] Heather Lee, *Homological mirror symmetry for open Riemann surfaces from pair-of-pants decompositions*, arXiv preprint arXiv:1608.04473 (2016).
- [LP18] Yanki Lekili and Alexander Polishchuk, *Auslander orders over nodal stacky curves and partially wrapped fukaya categories*, Journal of Topology **11** (2018), no. 3, 615–644.
- [LP20] _____, Derived equivalences of gentle algebras via Fukaya categories, Mathematische Annalen **376** (2020), no. 1, 187–225.
- [Nad16] David Nadler, Wrapped microlocal sheaves on pairs of pants, arXiv preprint arXiv:1604.00114 (2016).
- [Opp19] Sebastian Opper, On auto-equivalences and complete derived invariants of gentle algebras, arXiv preprint arXiv:1904.04859 (2019).
- [OPS18] Sebastian Opper, Pierre-Guy Plamondon, and Sibylle Schroll, *A geometric model for the derived category of gentle algebras*, arXiv preprint arXiv:1801.09659 (2018).
- [Orl03] Dmitri Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, arXiv preprint math/0302304 (2003).
- [Orl12] _____, Matrix factorizations for nonaffine LG–models, Mathematische Annalen 353 (2012), no. 1, 95–108.
- [PPP19] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon, *Non-kissing and non-crossing complexes for locally gentle algebras*, Journal of Combinatorial Algebra **3** (2019), no. 4, 401–438.
- [PS19] James Pascaleff and Nicolò Sibilla, *Topological Fukaya category and mirror symmetry for punctured surfaces*, Compositio Mathematica **155** (2019), no. 3, 599–644.
- [PS21]
 ______, Fukaya categories of higher-genus surfaces and pants decompositions, arXiv preprint arXiv:2103.03366 (2021).

 [PS22]
 ______, Singularity categories of normal crossings surfaces, descent, and mirror symmetry, arXiv preprint
- arXiv:2208.03896 (2022).
- [RAR94] John G Ratcliffe, S Axler, and KA Ribet, Foundations of hyperbolic manifolds, vol. 149, Springer, 1994.
- [Rho23] Kyungmin Rho, *Homological mirror symmetry and geometry of degenerate cusp singularities*, Ph.D. thesis, Seoul national university, 2023.
- [Sei08] Paul Seidel, *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR 2441780
- [Sei11] _____, Homological mirror symmetry for the genus two curve, J. Algebraic Geom. 20 (2011), no. 4, 727–769.
- [SZ03] Jan Schröer and Alexander Zimmermann, *Stable endomorphism algebras of modules over special biserial algebras*, Mathematische Zeitschrift **244** (2003), no. 3, 515–530.
- [Wik24] Wikipedia contributors, Kronecker product Wikipedia, the free encyclopedia, 2024, [Online; accessed 6-April-2024].
- [Yos90] Yuji Yoshino, *Maximal Cohen-Macaulay modules over Cohen-Macaulay rings*, vol. 146, Cambridge University Press, 1990.

DEPARTMENT OF MATHEMATICAL SCIENCES, RESEARCH INSTITUTE IN MATHEMATICS, SEOUL NATIONAL UNIVERSITY, GWANAK-GU, SEOUL, SOUTH KOREA

Email address: chocheol@snu.ac.kr

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT PADERBORN, WARBURGER STR. 100, PADERBORN, GERMANY

Email address: rho@math.uni-paderborn.de