

Hydrodynamically consistent many-body Harada-Sasa relation

Ramin Golestanian^{1,2,*}

¹Max Planck Institute for Dynamics and Self-Organization (MPI-DS), 37077 Göttingen, Germany

²Rudolf Peierls Centre for Theoretical Physics, University of Oxford, Oxford OX1 3PU, United Kingdom
(Dated: June 25, 2024)

The effect of hydrodynamic interactions on the non-equilibrium stochastic dynamics of particles – arising from the conservation of momentum in the fluid medium – is examined in the context of the relationship between fluctuations, response functions, and the entropy production rate. The multiplicative nature of the hydrodynamic interactions is shown to introduce subtleties that preclude a straightforward extension of the Harada-Sasa relation. A generalization of the definitions involved in the framework is used to propose a new form of the relation applicable to systems with hydrodynamic interactions. The resulting framework will enable characterization of the non-equilibrium properties of living and active matter systems, which are predominantly in suspensions.

Since the inception of the Onsager regression hypothesis [1], understanding the violation of the formal correspondence between fluctuations and response functions as observed in a variety of non-equilibrium systems [2–12] has guided many conceptual developments in non-equilibrium statistical physics. A particularly elegant breakthrough has been provided by the establishment of a formal relationship between the spectral summation of the difference between correlation and response, and the rate of energy dissipation, which quantifies how far away from equilibrium a given system operates [6, 7, 11].

With the advent of active matter [13], new horizons have opened in theoretical and experimental studies of non-equilibrium systems where questions pertaining to such relationships, as well as the degree of non-equilibrium activity and entropy production, can be systematically investigated [14–17]. These biological and synthetic active matter systems typically produce mechanical activity in the form of propulsion or delivering work as fuelled by a locally available source of free energy [18–25]. A key question in quantifying the non-equilibrium character of such active matter systems corresponds to the amount of hydrodynamic dissipation, which is inherently related to the faithful implementation of momentum conservation, from the nano-scale [26, 27] to the micro-scale [28–30].

Here, we set out to develop a hydrodynamically consistent extension of the Harada-Sasa relation [6, 7, 11], which can be applied to non-equilibrium suspensions in steady-state conditions. We achieve this by enforcing momentum conservation both on the particles in the suspension and on the medium itself, which is typically an incompressible viscous fluid that satisfies the Stokes equation. After eliminating the dynamics of the fluid medium, the resulting many-body stochastic dynamics constitutes a process with multiplicative noise, and should therefore be handled with care. Our approach is rooted in the correct physical characterization of the fluctuations of the non-equilibrium medium, and as such proves to be free of foundational problems associated with spurious drift terms that otherwise need to be augmented artificially.

We observe that the structure of the many-body dynamics with the multiplicative noise is such that the term responsible for quantifying the breakdown of time-reversal symmetry – that gives the entropy production rate – is independent of the hydrodynamic interactions. This might be construed as suggesting that a straightforward extension of the Harada-Sasa relation for the many-body dynamics could hold independently of the hydrodynamic coupling between the particles. We uncover that this intuition turns out to be untrue, due to the subtle involvement of the hydrodynamic couplings in the relationship between the correlation functions and the response functions. We derive the correct form for the Harada-Sasa relation in the presence of hydrodynamic interactions, which involves appropriately generalized correlation functions and response functions.

We consider the many-body stochastic dynamics of N colloidal particles in d dimensions, with the α th particle ($\alpha \in \{1, \dots, N\}$) being described by position $\mathbf{r}^\alpha(t)$ and force $\mathbf{f}^\alpha(t)$, where the force can depend on the position of the particle itself (arbitrary external force) and the positions of the other particles (arbitrary interactions, including many-body interactions). We can quantify the entropy production rate of the system as the average of the stochastic energy dissipation rate as follows

$$T\dot{\sigma} = \langle \dot{r}_i^\alpha(t) f_i^\alpha(\{\mathbf{r}^\nu(t)\}) \rangle, \quad (1)$$

where summation over repeated indices is assumed ($i \in \{1, \dots, d\}$). The averaging is to be performed using the corresponding path probability distribution weight, which reads

$$\mathbb{P}[\{\mathbf{r}^\alpha(t)\}] = \frac{1}{\mathcal{Z}} \left(\det \mathcal{M}_{ij}^{\alpha\beta} \right)^{-1/2} \exp \left(-\beta \mathcal{S}_{\text{OM}}[\{\mathbf{r}^\alpha(t)\}] \right), \quad (2)$$

with the Onsager-Machlup action given as

$$\mathcal{S}_{\text{OM}} = \frac{1}{4} \int dt \mathcal{Z}_{ij}^{\alpha\beta} \left(\dot{r}_i^\alpha - \mathcal{M}_{ik}^{\alpha\gamma} f_k^\gamma \right) \left(\dot{r}_j^\beta - \mathcal{M}_{jl}^{\beta\delta} f_l^\delta \right). \quad (3)$$

Here, $\mathcal{Z}(\{\mathbf{r}^\alpha(t)\})$ is the friction tensor, and $\mathcal{M}(\{\mathbf{r}^\alpha(t)\})$ is the mobility tensor, with the two being subject to the

relation $\mathcal{Z} = \mathcal{M}^{-1}$. Moreover, Z is a normalization constant, and $\beta \equiv 1/(k_B T)$ (see Appendix for the derivation of Eq. (2)). An inspection of Eq. (3) when re-written as follows

$$\mathcal{S}_{\text{OM}} = \frac{1}{4} \int dt \left[\mathcal{Z}_{ij}^{\alpha\beta} \dot{r}_i^\alpha \dot{r}_j^\beta - 2 \dot{r}_i^\alpha f_i^\alpha + \mathcal{M}_{ij}^{\alpha\beta} f_i^\alpha f_j^\beta \right], \quad (4)$$

reveals that the term in the Onsager-Machlup action that quantifies the breakdown of time-reversal symmetry is independent of the mobility and friction tensors, and that it directly leads to the local definition of entropy production rate as given in Eq. (1). This is related to the fact that the Fokker-Planck equation for the many-particle probability $\mathcal{P}(\{\mathbf{x}^\alpha\}, t)$ associated with the above Onsager-Machlup action, namely $(\partial_i^\alpha \equiv \partial/\partial x_i^\alpha)$

$$\partial_t \mathcal{P}(\{\mathbf{x}^\alpha\}, t) + \partial_i^\alpha \left[\mathcal{M}_{ij}^{\alpha\beta} \left(f_j^\beta \mathcal{P} - k_B T \partial_j^\beta \mathcal{P} \right) \right] = 0, \quad (5)$$

guarantees equilibration without any involvement of the mobility tensor, when $f_j^\beta = -\partial_j^\beta U$ for some scalar potential energy $U(\{\mathbf{x}^\alpha\})$. In this case, the many-body multiplicative coupling of the mobility tensor only affects the transient dynamics through the relaxation to equilibrium as described by $\mathcal{P}_{\text{eq}}(\{\mathbf{x}^\alpha\}) \propto \exp(-\beta U)$. In light of these observations, one naturally wonders if the same fate applies to the Harada-Sasa relation, in the sense that the local definition of entropy production rate given in Eq. (1) still quantifies the spectral sum of the degree of violation of the fluctuation-dissipation relation, as naturally defined in terms of the difference between the correlation function and the response function of the many-body system in the frequency domain. This is what we now investigate.

To develop the framework for the derivation of the Harada-Sasa relation, we start with the average value of the velocity of a given particle

$$\langle \dot{r}_i^\alpha(t) \rangle = \int \prod_\alpha \mathcal{D}\mathbf{r}^\alpha(\tau) \mathcal{P}[\{\mathbf{r}^\alpha(\tau)\}] \dot{r}_i^\alpha(t). \quad (6)$$

We next implement a set of external forces $\mathbf{f}^{\beta, \text{ext}}$ on all the particles, and use them to define the response function

$$\mathcal{R}_{ij}^{\alpha\beta}(t, t') = \mathcal{R}_{ij}^{\alpha\beta}(t - t') \equiv \left. \frac{\delta \langle \dot{r}_i^\alpha(t) \rangle_{\mathbf{f}^{\beta, \text{ext}}}}{\delta f_j^{\beta, \text{ext}}(t')} \right|_{\mathbf{f}^{\beta, \text{ext}} = \mathbf{0}}. \quad (7)$$

The first equality holds because we consider the system to be in steady-state conditions, and as such, expect time-translation invariance to hold. The calculation yields

$$\mathcal{R}_{ij}^{\alpha\beta}(t - t') = \frac{\beta}{2} \left[\bar{V}_i^\alpha \bar{V}_j^\beta + \mathcal{C}_{ij}^{\alpha\beta}(t - t') \right] - \frac{\beta}{2} \left\langle \dot{r}_i^\alpha(t) \mathcal{M}_{jk}^{\beta\gamma}(\{\mathbf{r}^\nu(t')\}) f_k^\gamma(\{\mathbf{r}^\nu(t')\}) \right\rangle, \quad (8)$$

where we have defined the average velocities as $\bar{V}_i^\alpha \equiv \langle \dot{r}_i^\alpha(t) \rangle$, and the correlation functions as

$$\mathcal{C}_{ij}^{\alpha\beta}(t - t') \equiv \left\langle [\dot{r}_i^\alpha(t) - \bar{V}_i^\alpha] [\dot{r}_j^\beta(t') - \bar{V}_j^\beta] \right\rangle. \quad (9)$$

It is straightforward to show that at equilibrium, where the forces are conservative ($f_j^\beta = -\partial_j^\beta U$), the average fluxes vanish ($\bar{V}_i^\alpha = 0$), and the system has both time-reversal and time-translation symmetries, Eq. (8) leads to the celebrated fluctuation-response relation

$$k_B T \mathcal{R}_{\text{eq}, ij}^{\alpha\beta}(t - t') = \Theta(t - t') \mathcal{C}_{\text{eq}, ij}^{\alpha\beta}(t - t'), \quad (10)$$

where $\Theta(t)$ is the Heaviside step function, which enforces causality. In frequency domain, Eq. (10) can be expressed as follows

$$\tilde{\mathcal{C}}_{\text{eq}, ij}^{\alpha\beta}(\omega) = 2k_B T \tilde{\mathcal{R}}_{\text{eq}, ij}^{\alpha\beta}(\omega), \quad (11)$$

where $\tilde{\mathcal{R}}_{\text{eq}, ij}^{\alpha\beta}(\omega)$ corresponds to the real part of the response function.

Back to the the general many-body non-equilibrium steady-state described by Eq. (8), we next construct the symmetrized form of the response functions $\mathcal{R}_{ij}^{\alpha\beta}(t - t') + \mathcal{R}_{ji}^{\beta\alpha}(t' - t)$, set $t = t'$, and take the trace of the tensor, which yields

$$2\mathcal{R}_{ii}^{\alpha\alpha}(0) = \beta [\bar{V}_i^\alpha \bar{V}_i^\alpha + \mathcal{C}_{ii}^{\alpha\alpha}(0)] - \beta \mathcal{J}, \quad (12)$$

where

$$\mathcal{J} \equiv \langle \dot{r}_i^\alpha(t) \mathcal{M}_{ik}^{\alpha\gamma}(\{\mathbf{r}^\nu(t)\}) f_k^\gamma(\{\mathbf{r}^\nu(t)\}) \rangle, \quad (13)$$

plays the role of a generalized flux. We can re-arrange Eq. (12) and express it as follows

$$\mathcal{J} = \bar{V}_i^\alpha \bar{V}_i^\alpha + \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \left[\tilde{\mathcal{C}}_{ii}^{\alpha\alpha}(\omega) - 2k_B T \tilde{\mathcal{R}}_{ii}^{\alpha\alpha}(\omega) \right]. \quad (14)$$

Since \mathcal{J} is not directly related (or proportional) to the entropy production rate as defined in Eq. (1) (as $\langle \dot{r} \mathcal{M} f \rangle \neq \langle \mathcal{M} \dot{r} f \rangle$ in the general case), we observe that the Harada-Sasa relation does not hold in its original form in such a many-body system.

It is, however, possible to construct a generalization of the Harada-Sasa relation for the many-body dissipative dynamics, by using appropriately generalized definitions for the response function and the correlation function. To achieve this, we revisit the starting point of the calculation, namely Eq. (6), and instead start with $\left\langle \mathcal{Z}_{ij}^{\alpha\beta}(\{\mathbf{r}^\nu(t)\}) \dot{r}_j^\beta(t) \right\rangle$ and define the generalized response function

$$\mathcal{R}_{g, ik}^{\alpha\gamma}(t - t') \equiv \left. \frac{\delta \left\langle \mathcal{Z}_{ij}^{\alpha\beta}(\{\mathbf{r}^\nu(t)\}) \dot{r}_j^\beta(t) \right\rangle_{\mathbf{f}^{\gamma, \text{ext}}}}{\delta f_j^{\gamma, \text{ext}}(t')} \right|_{\mathbf{f}^{\gamma, \text{ext}} = \mathbf{0}}. \quad (15)$$

Following the same steps of the calculation and defining the average friction tensor

$$\zeta_{ij}^{\alpha\beta} \equiv \left\langle \mathcal{Z}_{ij}^{\alpha\beta}(\{\mathbf{r}^\nu(t)\}) \right\rangle, \quad (16)$$

and the generalized correlation functions

$$\mathcal{C}_{g,ik}^{\alpha\gamma}(t-t') \equiv \left\langle \mathcal{Z}_{ij}^{\alpha\beta}(\{\mathbf{r}^\nu(t)\}) [\dot{r}_j^\beta(t) - \bar{V}_j^\beta] [\dot{r}_k^\gamma(t') - \bar{V}_k^\gamma] \right\rangle, \quad (17)$$

we obtain the following identity

$$\begin{aligned} \mathcal{R}_{g,ik}^{\alpha\gamma}(t-t') &= \frac{\beta}{2} \left[\zeta_{ij}^{\alpha\beta} \bar{V}_j^\beta \bar{V}_k^\gamma + \mathcal{C}_{g,ik}^{\alpha\gamma}(t-t') \right] \\ &- \frac{\beta}{2} \left\langle \mathcal{Z}_{ij}^{\alpha\beta}(\{\mathbf{r}^\nu(t)\}) \dot{r}_j^\beta(t) \mathcal{M}_{kl}^{\gamma\delta}(\{\mathbf{r}^\nu(t')\}) f_l^\delta(\{\mathbf{r}^\nu(t')\}) \right\rangle. \end{aligned} \quad (18)$$

Constructing the symmetrized form of the generalized response functions $\mathcal{R}_{g,ik}^{\alpha\gamma}(t-t') + \mathcal{R}_{g,ki}^{\gamma\alpha}(t'-t)$, setting $t = t'$, and taking the trace, we obtain

$$\begin{aligned} 2\mathcal{R}_{g,ii}^{\alpha\alpha}(0) &= \beta \left[\zeta_{ij}^{\alpha\beta} \bar{V}_i^\alpha \bar{V}_j^\beta + \mathcal{C}_{g,ii}^{\alpha\alpha}(0) \right] \\ &- \beta \langle \dot{r}_i^\alpha(t) f_i^\alpha(\{\mathbf{r}^\nu(t)\}) \rangle, \end{aligned} \quad (19)$$

which can be re-written as follows

$$T\dot{\sigma} = \zeta_{ij}^{\alpha\beta} \bar{V}_i^\alpha \bar{V}_j^\beta + \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \left[\tilde{\mathcal{C}}_{g,ii}^{\alpha\alpha}(\omega) - 2k_B T \tilde{\mathcal{R}}_{g,ii}^{\alpha\alpha}(\omega) \right]. \quad (20)$$

Note that in this case $\tilde{\mathcal{C}}_{g,ii}^{\alpha\alpha}(\omega)$ can have both real and imaginary parts, because in general $\mathcal{C}_{g,ii}^{\alpha\alpha}(t-t') \neq \mathcal{C}_{g,ii}^{\alpha\alpha}(t'-t)$. The above result (Eq. (20)) is our proposed generalization of the Harada-Sasa relation for a many-body system with hydrodynamic interactions.

In conclusion, we have developed a framework that allows us to relate the rate of entropy production to the difference between appropriately defined observables that characterize correlations and response in many-body stochastic dynamics of particles that are immersed in a background fluid that is subject to the conservation of momentum. The framework can be extended in a variety of ways, including its application to field theories that incorporate a background momentum-conserving fluid, such as model H dynamics [31] and its active generalization [32]. Hydrodynamic consistency is expected to provide the appropriate platform for the characterization and quantification of the non-equilibrium properties of living and active matter systems, as well as driven non-equilibrium processes that have a significant involvement of the background fluid, such as sedimentation.

Appendix: Many-body Onsager-Machlup action for particles with hydrodynamic interactions.—To derive the path probability distribution weight for the many-body stochastic dynamics of the colloidal suspension, we start with the fluctuating hydrodynamics framework of Landau and Lifshitz [33], as formulated in terms of the

velocity field $\mathbf{v}(\mathbf{x}, t)$, which is subject to the incompressibility condition $\nabla \cdot \mathbf{v} = 0$, and the pressure field $p(\mathbf{x}, t)$. This framework has been used to study violation of the fluctuation-response relation in suspensions with spatio-temporally structured temperature field [4], with a related generalization to quantum fluctuations of vacuum developed in Ref. [34]. In the viscous limit (i.e. low Reynolds number and no inertial dynamics), this takes the form of an instantaneous and locally enforced stochastic force balance (i.e. momentum conservation) equation

$$-\eta \nabla^2 \mathbf{v} = -\nabla p + \mathbf{f}(\mathbf{x}, t) + \boldsymbol{\xi}(\mathbf{x}, t), \quad (21)$$

where η is the viscosity of the medium, $\mathbf{f}(\mathbf{x}, t)$ is the body-force density, and $\boldsymbol{\xi}(\mathbf{x}, t)$ is a Gaussian white noise term defined via $\langle \xi_i(\mathbf{x}, t) \rangle = 0$ and $\langle \xi_i(\mathbf{x}, t) \xi_j(\mathbf{x}', t') \rangle = 2\eta k_B T (-\delta_{ij} \nabla^2 + \partial_i \partial_j) \delta^d(\mathbf{x} - \mathbf{x}') \delta(t - t')$. The presence of the colloidal particles necessitates no-slip boundary conditions between the velocities of the particles and the local fluid velocities, namely, $\mathbf{v}(\mathbf{r}^\alpha(t), t) = \dot{\mathbf{r}}^\alpha(t)$, and the identification of the forces experienced by each particle as the body-force exerted on the fluid, namely, $\mathbf{f}(\mathbf{x}, t) = \sum_\alpha \mathbf{f}^\alpha \delta^d(\mathbf{x} - \mathbf{r}^\alpha(t))$.

Using the general path integral formulation of stochastic dynamics through implementation of constraints via Lagrange multiplier fields [35], we can construct the path probability distribution weight for the many-body stochastic dynamics of the colloidal particles with hydrodynamic interactions as follows

$$\begin{aligned} \mathcal{P}_\xi[\{\mathbf{r}^\alpha(t)\}] &= \frac{1}{Z_\xi} \int \prod_\alpha \mathcal{D}\phi^\alpha \mathcal{D}\mathbf{v} \mathcal{D}p \mathcal{D}\bar{\mathbf{v}} \mathcal{D}\bar{p} \\ &\exp \left\{ i \int_{t,\mathbf{x}} \left[\bar{\mathbf{v}} \cdot \left(\eta \nabla^2 \mathbf{v} - \nabla p + \mathbf{f} + \boldsymbol{\xi} \right) + \bar{p} (\nabla \cdot \mathbf{v}) \right] \right. \\ &\left. + i \int_t \phi^\alpha(t) \cdot \left(\mathbf{v}(\mathbf{r}^\alpha(t), t) - \dot{\mathbf{r}}^\alpha(t) \right) \right\}, \end{aligned} \quad (22)$$

where the constant Z_ξ will ensure normalization of the probability distribution, namely, $\int \prod_\alpha \mathcal{D}\mathbf{r}^\alpha(t) \mathcal{P}_\xi[\{\mathbf{r}^\alpha(t)\}] = 1$, and we have used the notation $\int_{t,\mathbf{x}} \equiv \int dt \int d^d \mathbf{x}$, etc. We then define the field $\mathbf{h}(\mathbf{x}, t) = \sum_\alpha \phi^\alpha \delta^d(\mathbf{x} - \mathbf{r}^\alpha(t))$ to put all the relevant terms in the action in Eq. (22) on the same footing, perform noise averaging of the path probability distribution as follows $\mathcal{P}[\{\mathbf{r}^\alpha(t)\}] \equiv \langle \mathcal{P}_\xi[\{\mathbf{r}^\alpha(t)\}] \rangle$, go into the Fourier space, and perform Gaussian integrals over the fields \mathbf{v} , p , $\bar{\mathbf{v}}$, and \bar{p} . After these steps, we obtain

$$\begin{aligned} \mathcal{P}[\{\mathbf{r}^\alpha(t)\}] &= \frac{1}{Z} \int \prod_\alpha \mathcal{D}\phi^\alpha \\ &\exp \left\{ -k_B T \int_{\omega,\mathbf{q}} \frac{(\delta_{ij} - \hat{q}_i \hat{q}_j)}{\eta q^2} h_i(-\mathbf{q}, -\omega) h_j(\mathbf{q}, \omega) \right. \\ &\left. + i \int_{\omega,\mathbf{q}} \frac{(\delta_{ij} - \hat{q}_i \hat{q}_j)}{\eta q^2} h_i(-\mathbf{q}, -\omega) f_j(\mathbf{q}, \omega) - i \int_t \phi_i^\alpha \dot{r}_i^\alpha \right\}, \end{aligned} \quad (23)$$

where $f_j(\mathbf{q}, \omega) = \sum_{\beta} \int_{t'} e^{-i\omega t' + i\mathbf{q} \cdot \mathbf{r}^{\beta}(t')} f_j^{\beta}(t')$, $h_j(\mathbf{q}, \omega) = \sum_{\beta} \int_{t'} e^{-i\omega t' + i\mathbf{q} \cdot \mathbf{r}^{\beta}(t')} \phi_j^{\beta}(t')$, and we have used the shorthand $\int_{\omega, \mathbf{q}} \equiv \int \frac{d\omega}{2\pi} \int \frac{d^d \mathbf{q}}{(2\pi)^d}$. Here again, the constant Z will ensure normalization of the probability distribution, namely, $\int \prod_{\alpha} \mathcal{D}\mathbf{r}^{\alpha}(t) \mathcal{P}[\{\mathbf{r}^{\alpha}(t)\}] = 1$.

Equation (23) can be written explicitly as a Gaussian integral in terms of the Lagrange multiplier fields ϕ^{α} as follows

$$\mathcal{P}[\{\mathbf{r}^{\alpha}(t)\}] = \int \prod_{\alpha} \mathcal{D}\phi^{\alpha} \exp \left\{ -k_{\text{B}}T \int_t \mathcal{M}_{ij}^{\alpha\beta} \phi_i^{\alpha}(t) \phi_j^{\beta}(t) - i \int_t \phi_i^{\alpha}(t) \left[\dot{r}_i^{\alpha}(t) - \mathcal{M}_{ij}^{\alpha\beta} f_j^{\beta}(t) \right] \right\} / Z, \quad (24)$$

where $\mathcal{M}(\{\mathbf{r}^{\alpha}(t)\})$ is the mobility tensor of the colloidal suspension. Considering spherical colloids of radius a in $d = 3$, and in the far field (dilute) limit, the mobility tensor $\mathcal{M}(\{\mathbf{r}^{\alpha}(t)\})$ is obtained as follows

$$\mathcal{M}_{ij}^{\alpha\beta}(\mathbf{r}^{\alpha}, \mathbf{r}^{\beta}) = \begin{cases} \delta_{ij}/(6\pi\eta a), & \alpha = \beta, \\ \mathcal{G}_{ij}(\mathbf{r}^{\alpha} - \mathbf{r}^{\beta}), & \alpha \neq \beta, \end{cases} \quad (25)$$

where the Oseen tensor [36]

$$\mathcal{G}_{ij}(\mathbf{r}) = \frac{1}{8\pi\eta r} \left[\delta_{ij} + \frac{r_i r_j}{r^2} \right], \quad (26)$$

is the Green's function (in free space) for Stokes equation of hydrodynamics, describing the velocity profile of viscous and incompressible fluid flow generated by a point force (at the origin). Note that $\partial_i \mathcal{G}_{ij} = 0$ due to the incompressibility constraint.

Finally, integrating over the ϕ^{α} Lagrange multiplier fields in Eq. (24) yields the closed-form expression as follows

$$\mathcal{P}[\{\mathbf{r}^{\alpha}(t)\}] = \frac{1}{Z} \left(\det \mathcal{M}_{ij}^{\alpha\beta} \right)^{-1/2} \times \exp \left\{ -\frac{\beta}{4} \int_t \mathcal{Z}_{ij}^{\alpha\beta} \left(\dot{r}_i^{\alpha} - \mathcal{M}_{ik}^{\alpha\gamma} f_k^{\gamma} \right) \left(\dot{r}_j^{\beta} - \mathcal{M}_{jl}^{\beta\delta} f_l^{\delta} \right) \right\}, \quad (27)$$

where $\beta = 1/(k_{\text{B}}T)$ and $\mathcal{Z}(\{\mathbf{r}^{\alpha}(t)\}) = \mathcal{M}^{-1}$ is the friction tensor.

I acknowledge support from the Max Planck School Matter to Life and the MaxSynBio Consortium which are jointly funded by the Federal Ministry of Education and Research (BMBF) of Germany and the Max Planck Society.

* ramin.golestanian@ds.mpg.de

- [1] L. Onsager, Reciprocal relations in irreversible processes. ii., *Phys. Rev.* **38**, 2265 (1931).
 [2] L. F. Cugliandolo, J. Kurchan, and L. Peliti, Energy flow, partial equilibration, and effective temperatures in systems with slow dynamics, *Phys. Rev. E* **55**, 3898 (1997).

- [3] L. F. Cugliandolo, D. S. Dean, and J. Kurchan, Fluctuation-dissipation theorems and entropy production in relaxational systems, *Phys. Rev. Lett.* **79**, 2168 (1997).
 [4] R. Golestanian and A. Ajdari, Tracer diffusivity in a time- or space-dependent temperature field, *Europhys. Lett.* **59**, 800 (2002).
 [5] A. Crisanti and F. Ritort, Violation of the fluctuation-dissipation theorem in glassy systems: basic notions and the numerical evidence, *Journal of Physics A: Mathematical and General* **36**, R181–R290 (2003).
 [6] T. Harada and S.-i. Sasa, Equality Connecting Energy Dissipation with a Violation of the Fluctuation-Response Relation, *Phys. Rev. Lett.* **95**, 130602 (2005).
 [7] T. Harada and S.-i. Sasa, Energy dissipation and violation of the fluctuation-response relation in nonequilibrium Langevin systems, *Phys. Rev. E* **73**, 026131 (2006).
 [8] T. Speck and U. Seifert, Restoring a fluctuation-dissipation theorem in a nonequilibrium steady state, *Europhysics Letters (EPL)* **74**, 391–396 (2006).
 [9] M. Baiesi, C. Maes, and B. Wynants, Fluctuations and response of nonequilibrium states, *Phys. Rev. Lett.* **103**, 010602 (2009).
 [10] J. Prost, J.-F. Joanny, and J. M. R. Parrondo, Generalized fluctuation-dissipation theorem for steady-state systems, *Phys. Rev. Lett.* **103**, 090601 (2009).
 [11] T. Harada, Macroscopic expression connecting the rate of energy dissipation with the violation of the fluctuation response relation, *Phys. Rev. E* **79**, 030106 (2009).
 [12] C. Maes, Response theory: A trajectory-based approach, *Frontiers in Physics* **8**, 10.3389/fphy.2020.00229 (2020).
 [13] G. Gompper, R. G. Winkler, T. Speck, A. Solon, C. Nardini, F. Peruani, H. Löwen, R. Golestanian, U. B. Kaupp, L. Alvarez, *et al.*, The 2020 motile active matter roadmap, *J. Phys.: Condens. Matter* **32**, 193001 (2020).
 [14] C. Battle, C. P. Broedersz, N. Fakhri, V. F. Geyer, J. Howard, C. F. Schmidt, and F. C. MacKintosh, Broken detailed balance at mesoscopic scales in active biological systems, *Science* **352**, 604 (2016).
 [15] C. Nardini, E. Fodor, E. Tjhung, F. van Wijland, J. Tailleur, and M. E. Cates, Entropy production in field theories without time-reversal symmetry: Quantifying the non-equilibrium character of active matter, *Phys. Rev. X* **7**, 021007 (2017).
 [16] P. Pietzonka and U. Seifert, Entropy production of active particles and for particles in active baths, *J. Phys. A: Math* **51**, 01LT01 (2017).
 [17] S. Shankar and M. C. Marchetti, Hidden entropy production and work fluctuations in an ideal active gas, *Phys. Rev. E* **98**, 020604 (2018).
 [18] F. Jülicher, A. Ajdari, and J. Prost, Modeling molecular motors, *Rev. Mod. Phys.* **69**, 1269 (1997).
 [19] S. Ramaswamy, The mechanics and statistics of active matter, *Annual Review of Condensed Matter Physics* **1**, 323–345 (2010).
 [20] M. L. Mugnai, C. Hyeon, M. Hinczewski, and D. Thirumalai, Theoretical perspectives on biological machines, *Rev. Mod. Phys.* **92**, 025001 (2020).
 [21] S. Borsley, D. A. Leigh, and B. M. W. Roberts, Chemical fuels for molecular machinery, *Nat. Chem.* **14**, 728 (2022).
 [22] A.-K. Pumm, W. Engelen, E. Kopperger, J. Isensee, M. Vogt, V. Kozina, M. Kube, M. N. Honemann, E. Bertolin, M. Langecker, R. Golestanian, F. C. Sim-

- mel, and H. Dietz, A DNA origami rotary ratchet motor, *Nature* **607**, 492–498 (2022).
- [23] X. Shi, A.-K. Pumm, J. Isensee, W. Zhao, D. Verschueren, A. Martin-Gonzalez, R. Golestanian, H. Dietz, and C. Dekker, Sustained unidirectional rotation of a self-organized DNA rotor on a nanopore, *Nat. Phys.* **18**, 1105–1111 (2022).
- [24] X. Shi, A.-K. Pumm, C. Maffeo, F. Kohler, E. Feigl, W. Zhao, D. Verschueren, R. Golestanian, A. Aksimentiev, H. Dietz, and C. Dekker, A DNA turbine powered by a transmembrane potential across a nanopore, *Nat. Nanotechnol.* (2023).
- [25] R. Golestanian, Phoretic Active Matter, in *Active Matter and Nonequilibrium Statistical Physics: Lectures* (Oxford University Press, 2022).
- [26] R. Golestanian and A. Ajdari, Mechanical response of a small swimmer driven by conformational transitions, *Phys. Rev. Lett.* **100**, 038101 (2008).
- [27] M. Chatzittofi, J. Agudo-Canalejo, and R. Golestanian, Entropy production and thermodynamic inference for stochastic microswimmers, *Phys. Rev. Res.* **6**, L022044 (2024).
- [28] B. Nasouri, A. Vilfan, and R. Golestanian, Minimum dissipation theorem for microswimmers, *Phys. Rev. Lett.* **126**, 034503 (2021).
- [29] A. Daddi-Moussa-Ider, R. Golestanian, and A. Vilfan, Minimum entropy production by microswimmers with internal dissipation, *Nat. Commun.* **14**, 6060 (2023).
- [30] R. Bebon, J. F. Robinson, and T. Speck, *Thermodynamics of active matter: Tracking dissipation across scales* (2024).
- [31] P. C. Hohenberg and B. I. Halperin, Theory of dynamic critical phenomena, *Rev. Mod. Phys.* **49**, 435 (1977).
- [32] A. Tiribocchi, R. Wittkowski, D. Marenduzzo, and M. E. Cates, Active model h: Scalar active matter in a momentum-conserving fluid, *Phys. Rev. Lett.* **115**, 188302 (2015).
- [33] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics*, 11 September 2011, URL: <https://www.cambridge.org/core>.
- [34] M. Krüger, K. Asheichyk, M. Kardar, and R. Golestanian, Scale-dependent heat transport in dissipative media via electromagnetic fluctuations, *Phys. Rev. Lett.* **132**, 106903 (2024).
- [35] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena: Fifth Edition* (Oxford University Press, Oxford, 2021).
- [36] C. W. Oseen, *Neuere Methoden und Ergebnisse in der Hydrodynamik*, Vol. 1 (Akademische Verlagsgesellschaft, 1927).