Concentration Inequalities for (f, Γ) -GANs

A PREPRINT

Jeremiah Birrell

Department of Mathematics Texas State University San Marcos, TX 78666, USA jbirrell@txstate.edu

June 25, 2024

ABSTRACT

Generative adversarial networks (GANs) are unsupervised learning methods for training a generator distribution to produce samples that approximate those drawn from a target distribution. Many such methods can be formulated as minimization of a metric or divergence. Recent works have proven the statistical consistency of GANs that are based on integral probability metrics (IPMs), e.g., WGAN which is based on the 1-Wasserstein metric. IPMs are defined by optimizing a linear functional (difference of expectations) over a space of discriminators. A much larger class of GANs, which allow for the use of nonlinear objective functionals, can be constructed using (f, Γ) -divergences; these generalize and interpolate between IPMs and f-divergences (e.g., KL or α -divergences). Instances of (f,Γ) -GANs have been shown to exhibit improved performance in a number of applications. In this work we study the statistical consistency of (f, Γ) -GANs for general f and Γ . Specifically, we derive finite-sample concentration inequalities. These derivations require novel arguments due to nonlinearity of the objective functional. We demonstrate that our new results reduce to the known results for IPM-GANs in the appropriate limit while also significantly extending the domain of applicability of this theory.

Keywords Generative adversarial networks \cdot (f, Γ) -divergence \cdot statistical consistency \cdot concentration inequalities \cdot error decomposition

1 Introduction

Generative adversarial networks (GANs) are unsupervised learning methods for training a generator distribution to approximate a target distribution by using samples from the target in a minmax game between a generator and a discriminator network Arjovsky et al. [2017], Dziugaite et al. [2015], Goodfellow et al. [2014], Li et al. [2015]. Mathematically, many such methods can be formulated in terms of minimizing an integral probability metric (IPM):

$$\inf_{\Omega \in \Omega} d_{\Gamma}(Q, P_{\theta}), \tag{1}$$

$$\inf_{\theta \in \Theta} d_{\Gamma}(Q, P_{\theta}), \tag{1}$$

$$d_{\Gamma}(Q, P_{\theta}) := \sup_{h \in \Gamma} \{E_{Q}[h] - E_{P_{\theta}}[h]\}. \tag{2}$$
In space Γ (i.e., discriminators) Q is the distribution to be learned (i.e., the

Here d_{Γ} is the IPM with test function space Γ (i.e., discriminators), Q is the distribution to be learned (i.e., the distribution of the data) and P_{θ} is the generator distribution, depending on parameters $\theta \in \Theta$. More specifically, $P_{\theta} = (\Phi_{\theta})_{\#}P_{Z}$ is the pushforward of a noise source P_{Z} under a generator network Φ_{θ} . For instance, Wasserstein GAN Arjovsky et al. [2017] corresponds to Γ being the set of 1-Lipschitz functions. A key distinguishing feature of the IPM-GANs is the linearity in h of the objective functional in (2).

A larger class of GAN methods can be formulated by generalizing the minmax game (1) to use a nonlinear objective functional. Such methods can generally be viewed as minimizing a divergence (a generalized notion of "distance" or discrepancy) between Q and P_{θ} . The (f, Γ) -divergences are a large class of divergences with nonlinear objectives that generalize and interpolate between IPMs and f-divergences (e.g., KL-divergence); see Birrell et al. [2022a,b] for the general theory of these objects. Given a convex function f with f(1) = 0 and a test function space Γ , the (f, Γ) -divergence between probability distributions Q and P is defined by

$$D_f^{\Gamma}(Q||P) := \sup_{h \in \Gamma} \left\{ E_Q[h] - \Lambda_f^P[h] \right\}, \tag{3}$$

where, denoting the Legendre transform of f by f^* , we define

$$\Lambda_f^P[h] := \inf_{\nu \in \mathbb{R}} \{ \nu + E_P[f^*(h-\nu)] \}. \tag{4}$$

We refer to Λ_f^P as the generalized cumulant generating function because in the KL-divergence case $(f_{KL}(z)=z\log(z))$ it is straightforward to show $\Lambda_{f_{KL}}^P[h]=\log E_P[e^g]$, which is the classical cumulant generating function. The relation of Λ_f^P to f-divergences was previously studied in Ben-Tal & Teboulle [2007], Broniatowski & Keziou [2006], Nguyen et al. [2010], Ruderman et al. [2012]. Under appropriate assumptions on f and Γ , $D_f^\Gamma(Q\|P)$ provides a meaningful notion of discrepancy between Q and P due to it satisfying the divergence property, i.e., $D_f^\Gamma(Q\|P) \geq 0$ with equality if and only if Q=P; see Theorem 2.8 in Birrell et al. [2022a]. This property, along with the variational characterization (3), motivates the definition of (f,Γ) -GANs:

$$\inf_{\theta \in \Theta} D_f^{\Gamma}(Q \| P_{\theta}) = \inf_{\theta \in \Theta} \sup_{h \in \Gamma} \left\{ E_Q[h] - \Lambda_f^P[h] \right\}. \tag{5}$$

The family of (f,Γ) -GANs encompasses and generalizes a large number of successful methods in the literature, including Arjovsky et al. [2017], Belghazi et al. [2018], Dupuis, Paul & Mao, Yixiang [2022], Glaser et al. [2021], Goodfellow et al. [2014], Gretton et al. [2012], Gulrajani et al. [2017], Miyato et al. [2018], Nguyen et al. [2010], Nowozin et al. [2016], Song & Ermon [2020]; see Table 2 in Birrell et al. [2022a].

The statistical consistency theory of IPM-GANs has recently been studied in a number of works Chakraborty & Bartlett [2024], Chen et al. [2023b], Huang et al. [2022], Liang [2021]. These derivations take advantage of the special structure of IPMs, namely the linearity of the objective functional. However, the statistical consistency theory for more general GANs with nonlinear objectives is lacking. In this work we develop new tools that allow us to prove statistical consistency of the (f,Γ) -GANs, in the form of finite-sample concentration inequalities. The key technical hurdle is the nonlinearity of the objective functional due to the presence of the generalized cumulant generating function. Section 2 is dedicated to deriving the properties of and bounds on Λ_f^P that will be needed to prove the statistical consistency of (f,Γ) -GANs (5) in Section 3.

1.1 Summary of Results

Key to our results is a decomposition of the (f,Γ) -GAN error into optimization error, approximation error, and various statistical errors. Given an approximate discriminator space $\widetilde{\Gamma}\subset\Gamma$ (e.g., a neural network with spectral normalization as an approximation to the space of 1-Lipschitz functions), and empirical measures Q_n and $P_{\theta,m}:=(\Phi_\theta)_\# P_{Z,m}$ constructed using i.i.d. samples from Q and P_Z respectively, let $\theta_{n,m}^*$ be a solution (with optimization error tolerance $\epsilon_{opt}^{n,m}\geq 0$) to the empirical $(f,\widetilde{\Gamma})$ -GAN problem:

$$D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta_{n,m}^*,m}) \le \inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta,m}) + \epsilon_{opt}^{n,m}.$$
(6)

Given this, we derive the following (f, Γ) -GAN error bound in Lemma 3.5:

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta})$$

$$\leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}]) \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] \right\}$$

$$= \sup_{\text{statistical error from sampling } P_{Z}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\} + \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q_{n}}[h] - E_{Q}[h] \right\}$$

$$= \sup_{\text{statistical error from sampling } Q$$

$$+ \underbrace{\epsilon_{n,m}^{opt}}_{\text{optimization error}} + \underbrace{\left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)\right) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty}}_{\text{discriminator approximation error}}.$$

$$= \inf_{\text{optimization error}} \left\{ \sum_{h \in \Gamma} \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)\right) \right\} = \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty}$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty}$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

$$= \inf_{h \in \Gamma} \left(1 + (h^{*})'_{+}(z_{0} + \beta - \alpha)\right)$$

The distribution $P_{\theta_{n,m}^*} = (\Phi_{\theta_{n,m}^*})_\# P_Z$ is the (f,Γ) -GAN estimator, i.e., the approximation to Q obtained by solving the empirical GAN problem (6). The primary difference between the decomposition (7) and the corresponding result for IPMs in Lemma 9 of Huang et al. [2022] is the presence of the generalized cumulant generating function $\Lambda_f^{P_Z}$ in the statistical error from sampling P_Z . The functional $\Lambda_f^{P_Z}$ is nonlinear in the discriminator and so treating those terms requires new techniques which we develop in Section 2. Using these new results, in the case of a discriminator space Γ such that $\sup_{x,\tilde{x}} |h(x) - h(\tilde{x})| \leq 1$ for all $h \in \Gamma$, we derive concentration inequalities of the following form; see Theorem 3.10 for details:

$$\mathbb{P}\left(D_f^{\Gamma}(Q\|P_{\theta_{n,m}^*}) - \inf_{\theta \in \Theta} D_f^{\Gamma}(Q\|P_{\theta}) \ge \epsilon + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{opt}^{n,m} + 4\mathcal{R}_{\widetilde{\Gamma},Q,n} + 4\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}\right) \le \exp\left(-\frac{\epsilon^2}{\frac{2}{n} + \frac{2}{m}\Delta_{f,m}^2}\right). \tag{8}$$

Here $\epsilon_{approx}^{\Gamma,\widetilde{\Gamma}}$ is the discriminator approximation error, $\mathcal{R}_{\widetilde{\Gamma},Q,n}$ denotes the Rademacher complexity of $\widetilde{\Gamma}$ (see Appendix C), $\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}$ depends on $\mathcal{R}_{\widetilde{\Gamma}\circ\Phi,P_Z,m}$, and $\Delta_{f,m}$ can be bounded uniformly in m. In particular, if the discriminator and optimization errors are zero and assuming that one chooses discriminator and generator classes whose Rademacher complexities decay as $n,m\to\infty$, this result shows that the (f,Γ) -GAN estimator $P_{\theta_{n,m}^*}$ approximately solves the exact (f,Γ) -GAN problem (5) with high probability. If the generator class is sufficiently rich to ensure $\inf_{\theta\in\Theta}D_f^\Gamma(Q\|P_\theta)=0$ then (8) implies the (f,Γ) -GAN estimator is close to the distribution of the data source, Q, with high probability, where closeness is measured by the (f,Γ) -divergence; in this latter scenario we in fact obtain a tighter result than (8), see (85). The exponentially decaying nature of the bounds (8) means that one can use the Borel-Cantelli lemma to also conclude almost-sure convergence of $D_f^\Gamma(Q\|P_{\theta_{n,m}^*})$ to $\inf_{\theta\in\Theta}D_f^\Gamma(Q\|P_\theta)$, provided that the errors and Rademacher complexities approach zero when $n,m\geq k\to\infty$. As part of these derivations we prove bounds on the mean of the error; see Remark 3.13. Due to the asymmetry of D_f^Γ we also provide a concentration inequality where the positions of Q and $P_{\theta_{n,m}^*}$ are reversed; see Theorem 3.11. In addition, our techniques can be used to derive concentration inequalities for the (f,Γ) -divergence estimation problem; see Theorem 3.14. Finally, we note that the quantities $\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}$ and $\Delta_{f,m}$ approach their IPM counterparts in the limit where f^* becomes linear; see Remarks 2.8 and 2.11.

The concentration inequality (8) and other related results presented below reduce the problem of statistical performance guarantees for (f, Γ) -GANs to the problem of bounding the Rademacher complexities of the discriminator and the composition of discriminator and generator function classes. This is a well-studied problem which also arises the case of IPM-GANs and the same techniques and results from that setting can be applied here. We therefore do not focus on this aspect of the problem in the present work, instead referring the reader to Chapter 5 in Wainwright [2019] for background, including the use of Dudley's entropy integral and covering number bounds; see also Lemma 12 and its proof in Huang et al. [2022]. The discussion in Section 3.1 of Huang et al. [2022] and also Chakraborty & Bartlett [2024] shows how covering number bounds imply that, under appropriate assumptions, the IPM-GAN performance depends on the intrinsic dimension, d^* , of the support of Q and not the (possibly much larger) ambient dimension, d. As our new bounds depend on the Rademacher complexity in the same qualitative manner as the IPM results, the aforementioned insights regarding dimension dependence also immediately apply to (f, Γ) -GANs.

2 Properties of Λ_f^P

The fundamental difference between the more commonly studied IPM-GANs (1) and the (f,Γ) -GANs (5) is the nonlinearity of the objective functional in (3), coming from the generalized cumulant generating function (4). A detailed study of the properties of the generalized cumulant generating function, Λ_f^P , is therefore required in order to extend the techniques used for IPM-GANs to obtain statistical guarantees for (f,Γ) -GANs. We undertake that study in this section.

Going forward, we let \mathcal{X} be a measurable space and $\mathcal{M}_b(\mathcal{X})$ be the space of bounded measurable real-valued functions on \mathcal{X} . We let $f:(a,b)\to\mathbb{R},\ 0\le a<1< b\le\infty$, be a convex function that satisfies f(1)=0; we denote the set of such functions by $\mathcal{F}_1(a,b)$. We will make regular use of various properties of such convex functions and their Legendre transforms that are collected in Appendix A. In particular, we note that $a\ge 0$ implies f^* is non-decreasing. The value of the right derivative of f at 1 will play a key role in many of the proofs and so we make the following definition:

$$z_0 \coloneqq f'_+(1) \,. \tag{9}$$

By assumption, 1 is in the interior of the set where f is finite and so this right derivative is guaranteed to exist and be finite.

The first key property we prove is that bounds on a function h translate to bounds on $\Lambda_f^P[h]$ and also allow the minimization over ν in (4) to be restricted to a corresponding compact interval.

Lemma 2.1. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$, P be a probability measure on \mathcal{X} , and $h \in \mathcal{M}_b(\mathcal{X})$ with $\alpha \leq h \leq \beta$. Then:

- 1. $\alpha \leq \Lambda_f^P[h] \leq \beta$
- 2. If f is also strictly convex on a neighborhood of 1 and $z_0 \in \{f^* < \infty\}^o$ then

$$\Lambda_f^P[h] = \inf_{\nu \in [\alpha - z_0, \beta - z_0]} \{ \nu + E_P[f^*(h - \nu)] \}. \tag{10}$$

Remark 2.2. We use A^o to denote the interior of a set A.

Proof. 1. f^* is non-decreasing, therefore

$$\inf_{\nu \in \mathbb{R}} \{ \nu + f^*(\alpha - \nu) \} \le \inf_{\nu \in \mathbb{R}} \{ \nu + E_P[f^*(h - \nu)] \} \le \inf_{\nu \in \mathbb{R}} \{ \nu + f^*(\beta - \nu) \}$$
 (11)

for all ν . For any $c \in \mathbb{R}$ we have

$$\inf_{\nu \in \mathbb{R}} \{ \nu + f^*(c - \nu) \} = \inf_{z} \{ c - z + f^*(c - (c - z)) \}$$

$$= c - \sup_{z \in \mathbb{R}} \{ z - f^*(z) \} = c - (f^*)^*(1) = c - f(1) = c.$$
(12)

This proves the claim.

2. For the following it will be useful to recall that $f^*(z_0)=z_0$ and $(f^*)'_+(z_0)=1$; see Lemma A.1. Suppose $\nu>\beta-z_0$ and first consider the case where $f^*(h-\nu)<\infty$. We have $h-\nu< h-(\beta-z_0)\leq z_0$, therefore $h-\nu+1/n, h-(\beta-z_0)\in\{f^*<\infty\}^o$ for n sufficiently large. f^* is absolutely continuous on compact intervals, therefore the fundamental theorem of calculus holds (see, e.g., Theorem 3.35 and exercise 42 in Folland [2013]) and we have

$$f^*(h - (\beta - z_0)) = f^*(h - \nu + 1/n) + \int_{h - \nu + 1/n}^{h - (\beta - z_0)} (f^*)'_{+}(z)dz.$$
(13)

For $z \le h - (\beta - z_0)$ we have $z \le z_0$, hence $(f^*)'_+(z) \le (f^*)'_+(z_0) = 1$. Therefore

$$f^*(h - (\beta - z_0)) \le f^*(h - \nu + 1/n) + h - (\beta - z_0) - (h - \nu + 1/n)$$

$$= f^*(h - \nu + 1/n) - (\beta - z_0) + \nu - 1/n.$$
(14)

Using the continuity of f^* on $\{f^* < \infty\}$ we can take $n \to \infty$ to get

$$\nu + f^*(h - \nu) > (\beta - z_0) + f^*(h - (\beta - z_0)). \tag{15}$$

This was proven under the assumption that $f^*(h-\nu)<\infty$, but it also trivially holds when this is infinite. Taking the expectation of both sides we therefore find

$$\nu + E_P[f^*(h-\nu)] \ge (\beta - z_0) + E_P[f^*(h-(\beta - z_0))] \ge \inf_{\nu \in [\alpha - z_0, \beta - z_0]} \{\nu + E_P[f^*(h-\nu)]\}$$
 (16)

for all $\nu > \beta - z_0$.

Now suppose $\nu < \alpha - z_0$ and first consider the case where $f^*(h-\nu) < \infty$. We have $h-\nu > h-(\alpha-z_0) \ge z_0$. Therefore $h-\nu-1/n, h-(\alpha-z_0) \in \{f^* < \infty\}^o$ for n sufficiently large and

$$f^*(h-\nu-1/n) = f^*(h-(\alpha-z_0)) + \int_{h-(\alpha-z_0)}^{h-\nu-1/n} (f^*)'_+(z)dz.$$
 (17)

For $z \ge h - (\alpha - z_0)$ we have $z \ge z_0$ and so $(f^*)'_+(z) \ge 1$. Hence

$$f^*(h-\nu-1/n) \ge f^*(h-(\alpha-z_0)) + h-\nu-1/n - (h-(\alpha-z_0))$$

$$= f^*(h-(\alpha-z_0)) - \nu - 1/n + (\alpha-z_0).$$
(18)

Taking $n \to \infty$ gives

$$\nu + f^*(h - \nu) \ge (\alpha - z_0) + f^*(h - (\alpha - z_0)). \tag{19}$$

This was proven under the assumption that $f^*(h-\nu) < \infty$, but it also trivially holds when this is infinite. Taking the expectation of both sides we therefore find

$$\nu + E_P[f^*(h-\nu)] \ge (\alpha - z_0) + E_P[f^*(h-(\alpha - z_0))] \ge \inf_{\nu \in [\alpha - z_0, \beta - z_0]} \{\nu + E_P[f^*(h-\nu)]\}$$
 (20)

for all $\nu < \alpha - z_0$. This completes the proof.

Next we derive a Lipschitz bound for Λ_f^P .

Lemma 2.3. Let P be a probability measure on \mathcal{X} , and $h, \tilde{h} \in \mathcal{M}_b(\mathcal{X})$ with $\alpha \leq h, \tilde{h} \leq \beta$. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$. Then

$$|\Lambda_f^P[\tilde{h}] - \Lambda_f^P[h]| \le (f^*)'_+(z_0 + \beta - \alpha) \|\tilde{h} - h\|_{L^1(P)}. \tag{21}$$

Proof. f^* is non-decreasing, hence $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$ implies $z_0 \in \{f^* < \infty\}^o$. Therefore we can use Lemma 2.1 to write

$$\Lambda_f^P[h] = \inf_{\nu \in [\alpha - z_0, \beta - z_0]} \{ \nu + E_P[f^*(h - \nu)] \}.$$
 (22)

The fact that f^* is non-decreasing also implies that $h-z_0 \le f^*(h-\nu) \le f^*(\beta-\alpha+z_0) < \infty$ for all $\nu \in [\alpha-z_0,\beta-z_0]$ and hence $E_P[f^*(h-\nu)]$ is finite. The same holds for \tilde{h} , therefore

$$|\Lambda_f^P[\tilde{h}] - \Lambda_f^P[h]| \le \sup_{\nu \in [\alpha - z_0, \beta - z_0]} E_P[|f^*(\tilde{h} - \nu) - f^*(h - \nu)|]. \tag{23}$$

Here we made use of the simple bound

$$\pm (\inf_{i \in I} d_i - \inf_{i \in I} c_i) \le \sup_{i \in I} \{ \pm (d_i - c_i) \}$$
 (24)

whenever $c_i, d_i \in \mathbb{R}$ for all $i \in I$ and $\inf_i c_i, \inf_i d_i$ are finite.

 f^* is absolutely continuous on the interval between $h-\nu$ and $\tilde{h}-\nu$, therefore we can use the fundamental theorem of calculus to write

$$f^*(\tilde{h} - \nu) - f^*(h - \nu) = \int_{h - \nu}^{\tilde{h} - \nu} (f^*)'_{+}(z)dz$$
 (25)

and so

$$|f^*(\tilde{h}-\nu)-f^*(h-\nu)| \le |\tilde{h}-h| \sup_{z \in [z_0-(\beta-\alpha), z_0+\beta-\alpha]} |(f^*)'_+(z)|.$$
(26)

The fact that f^* is non-decreasing implies $(f^*)'_+ \ge 0$ on $\{f^* < \infty\}^o$. Combined with the fact that $(f^*)'_+$ is non-decreasing, we can therefore conclude

$$|f^*(\tilde{h} - \nu) - f^*(h - \nu)| \le |\tilde{h} - h|(f^*)'_{+}(z_0 + \beta - \alpha). \tag{27}$$

Taking the expectation of both sides and combining the result with (23) proves the claim.

Next we derive a tight bound on the difference between $\Lambda_f^{P_n}[h]$ and $\Lambda_f^{\widetilde{P}_n}[h]$ when P_n and \widetilde{P}_n are empirical measures that differ by only a single point. For this purpose it will be convenient to make the following definition.

Definition 2.4. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$. Define $\Lambda_f : \mathbb{R}^n \to \mathbb{R}$ by

$$\Lambda_f(x) := \inf_{\nu \in \mathbb{R}} \{ \nu + \frac{1}{n} \sum_{i=1}^n f^*(x_i - \nu) \}.$$
 (28)

The connection to Λ_f^P is given by the following lemma, which is a simple consequence of the respective definitions.

Lemma 2.5. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$. Let $h \in \mathcal{M}_b(\mathcal{X})$ and $P_n = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ be an empirical measure on \mathcal{X} . Then

$$\Lambda_f^{P_n}[h] = \Lambda_f \circ h_n(x) \tag{29}$$

where $h_n(x) := (h(x_1), ..., h(x_n)).$

The Lipschitz bound in Lemma 2.3 translates into the following Lipschitz bound for Λ_f .

Corollary 2.6. Let $f \in \mathcal{F}_1(a,b)$ with $a \ge 0$. Let $\alpha \le \beta$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$. For $x, \tilde{x} \in [\alpha, \beta]^n$ we have

$$|\Lambda_f(\tilde{x}) - \Lambda_f(x)| \le \frac{1}{n} (f^*)'_+(z_0 + \beta - \alpha) \sum_{i=1}^n |\tilde{x}_i - x_i|.$$
(30)

In particular, Λ_f is continuous on $[\alpha, \beta]^n$.

In the following lemma we derive a tight perturbation bounds on Λ_f when the inputs differ in only a single component, which in turn gives the desired tight bound on $\Lambda_f^{P_n}[h] - \Lambda_f^{\widetilde{P}_n}[h]$ when P_n and \widetilde{P}_n are empirical measures that differ by only a single point via Lemma 2.5.

Lemma 2.7. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$. Let $\alpha \leq \beta$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$. Given $j \in \{1, ..., n\}$, let $E = \{(x, \tilde{x}) \in [\alpha, \beta]^n \times [\alpha, \beta]^n : x_i = \tilde{x}_i$ for $i \neq j\}$. Then

$$\sup_{(x,\tilde{x})\in E} \{\Lambda_f(\tilde{x}) - \Lambda_f(x)\} = \inf_{z\in[z_0 - (\beta - \alpha), z_0]} \left\{ -z + \frac{n-1}{n} f^*(z) + \frac{1}{n} f^*(\beta - \alpha + z) \right\}$$
(31)

and we have the simpler loose bounds

$$\frac{\beta - \alpha}{n} \leq \inf_{z \in [z_0 - (\beta - \alpha), z_0]} \left\{ -z + \frac{n - 1}{n} f^*(z) + \frac{1}{n} f^*(\beta - \alpha + z) \right\}
\leq \frac{1}{n} (f^*(\beta - \alpha + z_0) - z_0) \leq (f^*)'_{+} (\beta - \alpha + z_0) \frac{\beta - \alpha}{n}.$$
(32)

Remark 2.8. Note that (32) implies that the bound (31) continuously approaches the result in the linear case (i.e., where $\Lambda_f^P = E_P$, which corresponds to $f^*(z) = z$ on a sufficiently large interval) as $(f^*)'_+(\beta - \alpha + z_0)$ approaches

Proof. First let $x_i = \alpha$ for all i, $\tilde{x}_i = \alpha$ for $i \neq j$ and $\tilde{x}_j = \beta$. We have $(x, \tilde{x}) \in E$ and therefore

$$\sup_{(x,\tilde{x})\in E} \{\Lambda_{f}(\tilde{x}) - \Lambda_{f}(x)\}
\geq \inf_{\nu\in[\alpha-z_{0},\beta-z_{0}]} \left\{\nu + \frac{1}{n}\sum_{i=1}^{n} f^{*}(\tilde{x}_{i}-\nu)\right\} - \inf_{\nu\in\mathbb{R}} \left\{\nu + \frac{1}{n}\sum_{i=1}^{n} f^{*}(x_{i}-\nu)\right\}
= \inf_{\nu\in[\alpha-z_{0},\beta-z_{0}]} \left\{\nu - \alpha + \frac{n-1}{n}f^{*}(\alpha-\nu) + \frac{1}{n}f^{*}(\beta-\nu)\right\} + \sup_{\nu\in\mathbb{R}} \{\alpha - \nu - f^{*}(\alpha-\nu)\}
= \inf_{z\in[z_{0}-(\beta-\alpha),z_{0}]} \left\{-z + \frac{n-1}{n}f^{*}(z) + \frac{1}{n}f^{*}(\beta-\alpha+z)\right\} + (f^{*})^{*}(1)
= \inf_{z\in[z_{0}-(\beta-\alpha),z_{0}]} \left\{-z + \frac{n-1}{n}f^{*}(z) + \frac{1}{n}f^{*}(\beta-\alpha+z)\right\}.$$

To prove the reverse inequality, let $(x, \tilde{x}) \in E$ be arbitrary and compute

$$\Lambda_{f}(\tilde{x}) - \Lambda_{f}(x) = \inf_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ \tilde{\nu} + \frac{1}{n} \sum_{i=1}^{n} f^{*}(\tilde{x}_{i} - \tilde{\nu}) \right\} - \inf_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ \nu + \frac{1}{n} \sum_{i=1}^{n} f^{*}(x_{i} - \nu) \right\}$$

$$= \sup_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \inf_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ \tilde{\nu} - \nu + \frac{1}{n} \sum_{i=1, i \neq j}^{n} f^{*}(x_{i} - \tilde{\nu}) - \frac{1}{n} \sum_{i=1, i \neq j}^{n} f^{*}(x_{i} - \nu) + \frac{1}{n} f^{*}(\tilde{x}_{j} - \tilde{\nu}) - \frac{1}{n} f^{*}(x_{j} - \nu) \right\}$$

$$\leq \sup_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \inf_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ \tilde{\nu} - \nu + \frac{1}{n} \sum_{i=1, i \neq j}^{n} f^{*}(x_{i} - \tilde{\nu}) - \frac{1}{n} \sum_{i=1, i \neq j}^{n} f^{*}(x_{i} - \nu) + \frac{1}{n} f^{*}(\beta - \tilde{\nu}) - \frac{1}{n} f^{*}(\alpha - \nu) \right\},$$

where in the last line we used the fact that f^* is non-decreasing. We have $\tilde{x}_i - \tilde{\nu}, x_i - \nu \in \{f^* < \infty\}^o$ and for each $i \neq j$ the terms involving x_i are absolutely continuous on compact intervals with right derivative

$$\frac{d}{dx_i^+} \left(\frac{1}{n} f^*(x_i - \tilde{\nu}) - \frac{1}{n} f^*(x_i - \nu) \right) = \frac{1}{n} \left((f^*)'_+(x_i - \tilde{\nu}) - (f^*)'_+(x_i - \nu) \right) , \tag{35}$$

which is non-positive when $\tilde{\nu} \geq \nu$ and non-negative when $\tilde{\nu} \leq \nu$ (due to $(f^*)'_+$ being non-decreasing). The fundamental theorem of calculus then implies that $\frac{1}{n}f^*(x_i-\tilde{\nu})-\frac{1}{n}f^*(x_i-\nu)$ is non-increasing in x_i when $\tilde{\nu} \geq \nu$ and non-decreasing in x_i when $\tilde{\nu} \leq \nu$. Therefore

$$\Lambda_{f}(\tilde{x}) - \Lambda_{f}(x) =
\begin{cases}
\sup_{\nu \in [\alpha - z_{0}, \beta - z_{0}]} \inf_{\tilde{\nu} \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ \tilde{\nu} - \nu + f^{*}(\beta - \tilde{\nu}) - \frac{n-1}{n} f^{*}(\beta - \nu) - \frac{1}{n} f^{*}(\alpha - \nu) & \text{if } \tilde{\nu} < \nu \\ \tilde{\nu} - \nu + \frac{n-1}{n} f^{*}(\alpha - \tilde{\nu}) - f^{*}(\alpha - \nu) + \frac{1}{n} f^{*}(\beta - \tilde{\nu}) & \text{if } \tilde{\nu} \ge \nu
\end{cases}$$

$$= \sup_{\nu \in [\alpha - z_{0}, \beta - z_{0}]} \inf_{z \in [z_{0} - (\beta - \alpha), z_{0}]} \left\{ \alpha - z - \nu + f^{*}(\beta - \alpha + z) - \frac{n-1}{n} f^{*}(\beta - \nu) - \frac{1}{n} f^{*}(\alpha - \nu) & \text{if } \alpha - \nu < z \\ \alpha - z - \nu + \frac{n-1}{n} f^{*}(z) - f^{*}(\alpha - \nu) + \frac{1}{n} f^{*}(\beta - \alpha + z) & \text{if } \alpha - \nu \ge z
\end{cases} \right\},$$

where we changed variables to $z = \alpha - \tilde{\nu}$ in the last line. For $z \in (\alpha - \nu, z_0]$, the right derivative of the objective with respect to z is given by

$$\frac{d}{dz^{+}}\left(\alpha - z - \nu + f^{*}(\beta - \alpha + z) - \frac{n-1}{n}f^{*}(\beta - \nu) - \frac{1}{n}f^{*}(\alpha - \nu)\right) = -1 + (f^{*})'_{+}(\beta - \alpha + z). \tag{37}$$

 $(f^*)'_+$ is non-decreasing and $\beta-\alpha+z\geq z_0$, therefore $-1+(f^*)'_+(\beta-\alpha+z)\geq -1+(f^*)'_+(z_0)=0$. This implies that the objective is non-decreasing in $z\in [\alpha-\nu,z_0]$ (the extension to the endpoint $\alpha-\nu$ follows from continuity of the objective). Therefore

$$\Lambda_{f}(\tilde{x}) - \Lambda_{f}(x)
\leq \sup_{\nu \in [\alpha - z_{0}, \beta - z_{0}]} \inf_{z \in [z_{0} - (\beta - \alpha), \alpha - \nu]} \left\{ \alpha - z - \nu + \frac{n - 1}{n} f^{*}(z) - f^{*}(\alpha - \nu) + \frac{1}{n} f^{*}(\beta - \alpha + z) \right\}
= \sup_{\nu \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ -\nu - f^{*}(\alpha - \nu) + \inf_{z \in [z_{0} - (\beta - \alpha), \alpha - \nu]} \left\{ \alpha - z + \frac{n - 1}{n} f^{*}(z) + \frac{1}{n} f^{*}(\beta - \alpha + z) \right\} \right\}.$$
(38)

Let z_* be a minimizer of

$$\inf_{z \in [z_0 - (\beta - \alpha), z_0]} \left\{ \alpha - z + \frac{n - 1}{n} f^*(z) + \frac{1}{n} f^*(\beta - \alpha + z) \right\}, \tag{39}$$

which exists due to compactness of the domain and continuity of the objective. The objective is convex in z, therefore it is non-increasing on $(-\infty, z_*]$ and so

$$\Lambda_{f}(\tilde{x}) - \Lambda_{f}(x)
\leq \sup_{\nu \in [\alpha - z_{0}, \beta - z_{0}]} \left\{ -\nu - f^{*}(\alpha - \nu) + \left\{ \frac{\alpha - z_{*} + \frac{n-1}{n} f^{*}(z_{*}) + \frac{1}{n} f^{*}(\beta - \alpha + z_{*}) & \text{if } \alpha - \nu \geq z_{*} \\ \nu + \frac{n-1}{n} f^{*}(\alpha - \nu) + \frac{1}{n} f^{*}(\beta - \nu) & \text{if } \alpha - \nu < z_{*} \right\}
= \sup_{w \in [z_{0} - (\beta - \alpha), z_{0}]} \left\{ \frac{w - f^{*}(w) - z_{*} + \frac{n-1}{n} f^{*}(z_{*}) + \frac{1}{n} f^{*}(\beta - \alpha + z_{*}) & \text{if } w \geq z_{*} \\ \frac{1}{n} (f^{*}(\beta - \alpha + w) - f^{*}(w)) & \text{if } w < z_{*} \right\},$$
(40)

where we changed variables to $w=\alpha-\nu$. We have $\sup_{w\in\mathbb{R}}\{w-f^*(w)\}=(f^*)^*(1)=0$ and $w\mapsto f^*(\beta-\alpha+w)-f^*(w)$ is non-decreasing on $w\leq z_0$ (this follows from $(f^*)'_+(\beta-\alpha+w)-(f^*)'_+(w)\geq 0$), therefore

$$\Lambda_f(\tilde{x}) - \Lambda_f(x) \le \max \left\{ -z_* + \frac{n-1}{n} f^*(z_*) + \frac{1}{n} f^*(\beta - \alpha + z_*), \frac{1}{n} (f^*(\beta - \alpha + z_*) - f^*(z_*)) \right\}. \tag{41}$$

Using the bound $f^*(z) \ge z$ we can compute

$$-z_* + \frac{n-1}{n}f^*(z_*) + \frac{1}{n}f^*(\beta - \alpha + z_*) - \frac{1}{n}(f^*(\beta - \alpha + z_*) - f^*(z_*))$$

= $-z_* + f^*(z_*) \ge 0$.

Hence

$$\Lambda_{f}(\tilde{x}) - \Lambda_{f}(x) \leq -z_{*} + \frac{n-1}{n} f^{*}(z_{*}) + \frac{1}{n} f^{*}(\beta - \alpha + z_{*})
= \inf_{z \in [z_{0} - (\beta - \alpha), z_{0}]} \left\{ -z + \frac{n-1}{n} f^{*}(z) + \frac{1}{n} f^{*}(\beta - \alpha + z) \right\}.$$
(42)

This holds for all $(x, \tilde{x}) \in E$ and so when combined with (33) we obtain the claimed equality.

To derive the looser bounds, first use $f^*(z) \ge z$ to compute the lower bound

$$\inf_{z \in [z_0 - (\beta - \alpha), z_0]} \left\{ -z + \frac{n-1}{n} f^*(z) + \frac{1}{n} f^*(\beta - \alpha + z) \right\} \ge \frac{\beta - \alpha}{n}.$$
 (43)

For the upper bound we compute

$$\inf_{z \in [z_0 - (\beta - \alpha), z_0]} \left\{ -z + \frac{n-1}{n} f^*(z) + \frac{1}{n} f^*(\beta - \alpha + z) \right\}
\leq -z_0 + \frac{n-1}{n} f^*(z_0) + \frac{1}{n} f^*(\beta - \alpha + z_0) = \frac{1}{n} (f^*(\beta - \alpha + z_0) - z_0)
\leq (f^*)'_+ (\beta - \alpha + z_0) \frac{\beta - \alpha}{n} ,$$
(44)

where to obtain the last line we used the fundamental theorem of calculus together with the facts that $f^*(z_0) = z_0$ and $(f^*)'_+$ is non-decreasing.

Next we derive a pair of uniform law of large numbers (ULLN) results that bound the maximum difference between $\Lambda_f^{P_n}[h]$ and $\Lambda_f^P[h]$ over a set of test function $h \in \Gamma$ in terms of the Rademacher complexity of Γ , where P_n is the empirical measure for n i.i.d. samples from P. These results should be compared with the ULLN for means which provides uniform bounds on the difference between the empirical mean and expectation; see Appendix C for the relevant background. First we obtain a simpler result that uses (10) along with a Lipschitz bound to reduce the problem to the ULLN for means; this can be thought of as a substantially more general version of the estimate in the proof of Lemma 2 in Belghazi et al. [2018], which studied a KL-divergence based mutual information method. However, as we will show, the resulting estimate is overly pessimistic as it does not reduce to the IPM case in the appropriate limit. Following this simpler lemma we will derive a tighter bound that does possess the desired limiting property.

Lemma 2.9. Let Ψ be a nonempty countable collection of measurable functions on \mathcal{Y} and suppose we have $\alpha, \beta \in \mathbb{R}$ such that $\alpha \leq \psi \leq \beta$ for all $\psi \in \Psi$. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$.

Let P be a probability measure on \mathcal{Y} and Y_i , i = 1, ..., n be independent samples from P with P_n the corresponding empirical measure. Then

$$\mathbb{E}\left[\sup_{\psi\in\Psi}\left\{\pm\left(\Lambda_f^P[\psi]-\Lambda_f^{P_n}[\psi]\right)\right\}\right]\leq 2(f^*)'_+(\beta-\alpha+z_0)\left(\mathcal{R}_{\Psi,P,n}+\frac{\beta-\alpha}{2n^{1/2}}\right). \tag{45}$$

Proof. The result is trivial if $\alpha = \beta$ so suppose $\alpha < \beta$. Note that f^* is non-decreasing, and hence $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$ implies $z_0 \in \{f^* < \infty\}^o$. Using Lemma 2.1 along with (24) we obtain

$$\pm \left(\Lambda_f^P[\psi] - \Lambda_f^{P_n}[\psi]\right) \le \sup_{\nu \in [\alpha - z_0, \beta - z_0]} \{ \pm \left(E_P[f^*(\psi - \nu)] - E_{P_n}[f^*(\psi - \nu)]\right) \}. \tag{46}$$

Note that $z_0 - (\beta - \alpha) \le f^*(\psi - \nu) \le f^*(\beta - \alpha + z_0) < \infty$ and so the expectations are finite. Also, the supremum over ν can be restricted to rational ν due to continuity. Maximizing over ψ and taking expectations gives

$$\mathbb{E}\left[\sup_{\psi\in\Psi}\left\{\pm\left(\Lambda_f^P[\psi]-\Lambda_f^{P_n}[\psi]\right)\right\}\right]\leq\mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_0,\beta-z_0]\cap\mathbb{Q}}\left\{\pm\left(E_P[f^*(\psi-\nu)]-E_{P_n}[f^*(\psi-\nu)]\right)\right\}\right]. \tag{47}$$

The ULLN (see Theorem C.4) implies

$$\mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}\left\{\pm\left(E_{P}[f^{*}(\psi-\nu)]-E_{P_{n}}[f^{*}(\psi-\nu))\right]\right\}\right] \\
=\mathbb{E}\left[\sup_{h\in\mathcal{H}}\left\{\pm\left(E_{P}[h]-\frac{1}{n}\sum_{i=1}^{n}h(Y_{i})\right)\right\}\right] \leq 2\mathcal{R}_{\mathcal{H},P,n}, \tag{48}$$

where $\mathcal{H}\coloneqq\{f^*(\psi-\nu):\psi\in\Psi,\nu\in[\alpha-z_0,\beta-z_0]\cap\mathbb{Q}\}$. f^* and $(f^*)'$ are decreasing, therefore f^* is $(f^*)'_+(\beta-\alpha+z_0)$ -Lipschitz on $(-\infty,\beta-\alpha+z_0]$ and so Talagrand's lemma (see Lemma C.3) implies

$$\mathcal{R}_{\mathcal{H},P,n} \le (f^*)'_{+}(\beta - \alpha + z_0)\mathcal{R}_{\Psi - [\alpha - z_0,\beta - z_0] \cap \mathbb{D},P,n}. \tag{49}$$

We can compute

$$\mathcal{R}_{\Psi-[\alpha-z_0,\beta-z_0]\cap\mathbb{Q},P,n} = \mathcal{R}_{\Psi,P,n} + \mathcal{R}_{[\alpha-z_0,\beta-z_0],P,n} \tag{50}$$

and

$$\mathcal{R}_{[\alpha-z_{0},\beta-z_{0}],P,n} = \frac{1}{n} E_{\sigma} \left[\sup_{\nu \in [\alpha-z_{0},\beta-z_{0}]} \nu \sum_{i=1}^{n} \sigma_{i} \right] = \frac{1}{n} E_{\sigma} \left[\sup_{z \in [-(\beta-\alpha)/2,(\beta-\alpha)/2]} (z + (\beta+\alpha)/2 - z_{0}) \sum_{i=1}^{n} \sigma_{i} \right] \\
= \frac{1}{n} E_{\sigma} \left[\sup_{z \in [-(\beta-\alpha)/2,(\beta-\alpha)/2]} z \sum_{i=1}^{n} \sigma_{i} \right] = \frac{\beta-\alpha}{2n} E_{\sigma} \left[\left| \sum_{i=1}^{n} \sigma_{i} \right| \right] \leq \frac{\beta-\alpha}{2n} E_{\sigma} \left[\left| \sum_{i=1}^{n} \sigma_{i} \right|^{2} \right]^{1/2} \\
= \frac{\beta-\alpha}{2n^{1/2}}.$$
(51)

We will now derive the more precise bound that behaves appropriately when f^* becomes linear.

Lemma 2.10. Let Ψ be a countable collection of measurable functions on $\mathcal Y$ that contains at least one constant function and suppose we have $\alpha, \beta \in \mathbb R$ such that $\alpha \leq \psi \leq \beta$ for all $\psi \in \Psi$. Let $f \in \mathcal F_1(a,b)$ with $a \geq 0$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$. Also assume that $(f^*)'_+$ is $L_{\alpha,\beta}$ -Lipschitz on $[z_0 - (\beta - \alpha), z_0 + \beta - \alpha]$.

Let P be a probability measure on \mathcal{Y} and Y_i , i = 1, ..., n be independent samples from P with P_n the corresponding empirical measure. Then

$$\mathbb{E}\left[\sup_{\psi\in\Psi}\left\{\pm\left(\Lambda_{f}^{P}[\psi]-\Lambda_{f}^{P_{n}}[\psi]\right)\right\}\right] \leq 2\min\left\{\left(1+2(\beta-\alpha)L_{\alpha,\beta}\right)\mathcal{R}_{\Psi,P,n}+\frac{(\beta-\alpha)^{2}L_{\alpha,\beta}}{2n^{1/2}},\left(f^{*}\right)'_{+}(\beta-\alpha+z_{0})\left(\mathcal{R}_{\Psi,P,n}+\frac{\beta-\alpha}{2n^{1/2}}\right)\right\}.$$
(52)

Remark 2.11. The result in Lemma 2.9 does not reduce to the ULLN for means in the case where $f^*(z) = z$ (on a sufficiently large interval), i.e., when $\Lambda_f^P[h] = E_P[h]$; specifically the bound (45) differs from the ULLN for means (116) by the term $\frac{\beta-\alpha}{m^{1/2}}$ in that case. However, (52) does reduce to the ULLN for means when $f^*(z) = z$. Moreover, the bound (52) approaches the bound in (116) as the Lipschitz constant for $(f^*)'_+$ on $[z_0 - (\beta - \alpha), z_0 + \beta - \alpha]$ approaches zero.

Proof. As in the proof of Lemma 2.9, the result is trivial when $\alpha = \beta$. For $\alpha < \beta$ we have

$$\mathbb{E}\left[\sup_{\psi\in\Psi}\left\{\pm\left(\Lambda_{f}^{P}[\psi]-\Lambda_{f}^{P_{n}}[\psi]\right)\right\}\right]\leq\mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}\left\{\pm\left(E_{P}[f^{*}(\psi-\nu)]-E_{P_{n}}[f^{*}(\psi-\nu)]\right)\right\}\right]. \quad (53)$$

Now use the fundamental theorem of calculus to compute

$$f^*(\psi - \nu) = f^*(\psi - (\beta - z_0)) + (\beta - z_0 - \nu) \int_0^1 (f^*)'_+(\psi - (1 - t)(\beta - z_0) - t\nu) dt$$
 (54)

for
$$\psi \in \Psi, \nu \in [\alpha - z_{0}, \beta - z_{0}] \cap \mathbb{Q}$$
. Therefore
$$\sup_{\psi \in \Psi, \nu \in [\alpha - z_{0}, \beta - z_{0}] \cap \mathbb{Q}} \left\{ \pm (E_{P}[f^{*}(\psi - \nu)] - E_{P_{n}}[f^{*}(\psi - \nu)]) \right\}$$

$$= \sup_{\psi \in \Psi, \nu \in [\alpha - z_{0}, \beta - z_{0}] \cap \mathbb{Q}} \left\{ \pm (E_{P}[f^{*}(\psi - (\beta - z_{0}))] - E_{P_{n}}[f^{*}(\psi - (\beta - z_{0}))] \right.$$

$$+ (\beta - z_{0} - \nu) \int_{0}^{1} E_{P}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] - E_{P_{n}}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0} - t\nu)]dt \right\}$$

$$\leq \sup_{\psi \in \Psi} \left\{ \pm (E_{P}[f^{*}(\psi - (\beta - z_{0}))] - E_{P_{n}}[f^{*}(\psi - (\beta - z_{0}))] \right\}$$

$$+ \sup_{\psi \in \Psi, \nu \in [\alpha - z_{0}, \beta - z_{0}] \cap \mathbb{Q}} \left\{ (\beta - z_{0} - \nu) \int_{0}^{1} \left(E_{P}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] - E_{P_{n}}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] \right.$$

$$+ \sup_{\psi \in \Psi, \nu \in [\alpha - z_{0}, \beta - z_{0}] \cap \mathbb{Q}} \left\{ (\beta - z_{0} - \nu) \int_{0}^{1} \left(E_{P}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] - E_{P_{n}}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] \right. \right.$$

$$\left. - E_{P_{n}}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] \right.$$

$$\left. - E_{P_{n}}[(f^{*})'_{+}(\psi - (1 - t)(\beta - z_{0}) - t\nu)] \right) dt \right\}.$$

The ULLN (Theorem C.4) implies

$$\mathbb{E}\left[\sup_{h \in f^* \circ (\Psi - (\beta - z_0))} \{ \pm (E_P[h] - E_{P_n}[h]) \} \right] \le 2\mathcal{R}_{f^* \circ (\Psi - (\beta - z_0)), P, n}.$$
(57)

We have $\psi - (\beta - z_0) \le z_0$ and f^* is $(f^*)'_+(z_0)$ -Lipschitz on $(-\infty, z_0]$ where $(f^*)'_+(z_0) = 1$, hence Talagrand's lemma (Lemma C.3) implies

$$\mathcal{R}_{f^* \circ (\Psi - (\beta - z_0)), P, n} \le \mathcal{R}_{\Psi - (\beta - z_0), P, n} = \mathcal{R}_{\Psi, P, n}.$$
(58)

We note that the shift invariance of the Rademacher complexity follows from our use of the definition (113), i.e., without absolute value.

As for the second term in (56), we have

$$\mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}\left\{(\beta-z_{0}-\nu)\int_{0}^{1}\left(E_{P}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]\right)-E_{P_{n}}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]\right]\right] \\
= \left[\sum_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}\left\{\left|E_{P}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]\right|-E_{P_{n}}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0}-t\nu))]\right\}\right]dt.$$
(59)

Using the fact that Ψ contains a constant (which implies that the terms in the following bound are non-negative) and then employing the ULLN (Theorem C.4) we have

$$\mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}|E_{P}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]-E_{P_{n}}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]|\right] (60)$$

$$\leq \mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}\left\{E_{P}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]-E_{P_{n}}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]\right\}\right]$$

$$+\mathbb{E}\left[\sup_{\psi\in\Psi,\nu\in[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}}\left\{-(E_{P}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]-E_{P_{n}}[(f^{*})'_{+}(\psi-(1-t)(\beta-z_{0})-t\nu)]\right)\right\}\right]$$

$$\leq 4\mathcal{R}_{\mathcal{H}_{t},P,n},$$

where $\mathcal{H}_t \coloneqq \{(f^*)'_+(\psi - (1-t)(\beta-z_0) - t\nu) : \psi \in \Psi, \nu \in [\alpha-z_0, \beta-z_0] \cap \mathbb{Q}\}$. Using the Lipschitz assumption on $(f^*)'_+$ together with Talagrand's lemma (Lemma C.3) and the result of the calculation (51) we obtain

$$\mathcal{R}_{\mathcal{H}_{t},P,n} \leq L_{\alpha,\beta} \mathcal{R}_{\Psi-(1-t)(\beta-z_{0})-t[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}),P,n}
= L_{\alpha,\beta} (\mathcal{R}_{\Psi,P,n} + tR_{[\alpha-z_{0},\beta-z_{0}]\cap\mathbb{Q}),P,n}) \leq L_{\alpha,\beta} \left(\mathcal{R}_{\Psi,P,n} + t \frac{\beta-\alpha}{2n^{1/2}} \right).$$
(61)

Putting these bounds together we find

$$\mathbb{E}\left[\sup_{\psi\in\Psi}\left\{\pm\left(\Lambda_{f}^{P}[\psi]-\Lambda_{f}^{P_{n}}[\psi]\right)\right\}\right] \leq 2\mathcal{R}_{\Psi,P,n}+(\beta-\alpha)\int_{0}^{1}4L_{\alpha,\beta}\left(\mathcal{R}_{\Psi,P,n}+t\frac{\beta-\alpha}{2n^{1/2}}\right)dt \qquad (62)$$

$$=2(1+2(\beta-\alpha)L_{\alpha,\beta})\mathcal{R}_{\Psi,P,n}+\frac{(\beta-\alpha)^{2}L_{\alpha,\beta}}{n^{1/2}}.$$

Combining this with the bound from Lemma 2.9 gives the claimed result.

3 Error Bounds for (f, Γ) -GANs

In this section we use the bounds on Λ_f^P derived in Section 2 to derive concentration inequalities for (f,Γ) -GANs.

3.1 (f, Γ) -GAN Error Decomposition

We start by deriving a decomposition of the (f,Γ) -GAN error into statistical, approximation, and optimization error terms. First we consider the error that comes from approximating the idealized discriminator space Γ by a smaller space (e.g., a neural network) $\widetilde{\Gamma}$, a step that is generally required when implementing a GAN method:

Lemma 3.1. Let $\Gamma \subset \Gamma \subset \mathcal{M}_b(\mathcal{X})$ be nonempty and suppose we have $\alpha, \beta \in \mathbb{R}$ such that $\alpha \leq h \leq \beta$ for all $h \in \Gamma$. Let $f \in \mathcal{F}_1(a,b)$ with $a \geq 0$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$. Then for any probability measures Q, P on \mathcal{X} we have

$$0 \le D_f^{\Gamma}(Q||P) - D_f^{\widetilde{\Gamma}}(Q||P) \le (1 + (f^*)'_{+}(z_0 + \beta - \alpha)) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} ||h - \tilde{h}||_{\infty}.$$
 (63)

Proof. For any $h \in \Gamma$, $\tilde{h} \in \widetilde{\Gamma}$ we can use the definition (3) and Lemma 2.3 to compute

$$E_{Q}[h] - \Lambda_{f}^{P}[h] - D_{f}^{\widetilde{\Gamma}}(Q \| P) \leq E_{Q}[h] - E_{Q}[\tilde{h}] + \Lambda_{f}^{P}[\tilde{h}] - \Lambda_{f}^{P}[h]$$

$$\leq \|h - \tilde{h}\|_{L^{1}(Q)} + (f^{*})'_{+}(z_{0} + \beta - \alpha)\|\tilde{h} - h\|_{L^{1}(P)}$$

$$\leq (1 + (f^{*})'_{+}(z_{0} + \beta - \alpha))\|h - \tilde{h}\|_{\infty}.$$

$$(64)$$

Minimizing over $\tilde{h} \in \widetilde{\Gamma}$ gives

$$E_{Q}[h] - \Lambda_{f}^{P}[h] - D_{f}^{\widetilde{\Gamma}}(Q||P) \le (1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)) \inf_{\widetilde{h} \in \widetilde{\Gamma}} ||h - \widetilde{h}||_{\infty}$$
 (65)

for all $h \in \Gamma$. Maximizing over $h \in \Gamma$ then completes the proof.

Next we outline the assumptions we make regarding the discriminator, generator, and the empirical GAN optimization. **Assumption 3.2** $((f, \Gamma)\text{-GAN Assumptions})$. Let $f \in \mathcal{F}_1(a, b)$ with $a \ge 0$ satisfy the following:

- 1. f is strictly convex in a nbhd of 1.
- 2. $z_0 + \beta \alpha \in \{f^* < \infty\}^o$, where z_0 was defined in (9).

Let $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$ be a topological space with the Borel sigma algebra. Suppose $\widetilde{\Gamma} \subset \Gamma \subset C_b(\mathcal{X})$ (the space of bounded continuous functions) are nonempty (the discriminator spaces) and satisfy the following:

- 1. We have $\alpha, \beta \in \mathbb{R}$ such that $\alpha < \beta$ and $\alpha \le h \le \beta$ for all $h \in \Gamma$.
- 2. There exists a countable $\Gamma_0 \subset \Gamma$ such that for all $h \in \Gamma$ there exists a sequence $h_j \in \Gamma_0$ such that $h_j \to h$ pointwise.

3. There exists a countable $\widetilde{\Gamma}_0 \subset \widetilde{\Gamma}$ such that for all $\widetilde{h} \in \widetilde{\Gamma}$ there exists a sequence $\widetilde{h}_j \in \widetilde{\Gamma}_0$ such that $\widetilde{h}_j \to \widetilde{h}$ pointwise.

Let \mathcal{Z} be another measurable space, P_Z a probability measure on \mathcal{Z} , and $\Phi_{\theta}: \mathcal{Z} \to \mathcal{X}$ be measurable for $\theta \in \Theta$, where Θ is a separable metric space. Suppose $\theta \mapsto \Phi_{\theta}(z)$ is continuous for all $z \in \mathcal{Z}$. Define $P_{\theta} := (\Phi_{\theta})_{\#} P_Z$, which is a probability measure on \mathcal{X} .

Let Q be a probability measure on X and X_i , i=1,...,n, Z_i , i=1,...,m be independent and distributed as Q, P_Z respectively. Let Q_n , $P_{Z,m}$. and $P_{\theta,m}$ be the empirical measures corresponding to X_i , Z_i , and $\Phi_{\theta} \circ Z_i$ respectively. Finally, suppose that for each m,n we have an error tolerance $\epsilon_{n,m}^{opt} \geq 0$ and Θ -valued random variables $\theta_{n,m}^*$ that are approximate optimizers to the empirical GAN problem, i.e., that satisfy

$$D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta_{n,m}^*,m}) \le \inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta,m}) + \epsilon_{n,m}^{opt} \quad \mathbb{P}\text{-a.s.}$$
 (66)

Remark 3.3. The assumptions regarding Γ_0 , $\widetilde{\Gamma}_0$, separability of Θ , and continuity of Φ_θ and $h \in \Gamma$ allow us to address the issue of measurability of the various suprema that arise in the derivations below by enabling one to restrict them to countable subsets. These assumptions hold in most cases of interest and can also be replaced by any alternatives that serve the same purpose.

Note that asymmetry of the (f,Γ) -divergences requires us to separately treat the cases where the generator is the first argument and where it is the second; we emphasize that this choice can make a significant difference in practice as shown in the examples in Birrell et al. [2022a]. With this in mind we give a second version of the (f,Γ) -GAN assumptions and will derive concentration inequalities for both cases.

Assumption 3.4 (Reverse (f,Γ) -GAN Assumptions). *Assume that Assumption 3.2 holds, with the sole exception being that* $\theta_{n,m}^*$ *satisfies*

$$D_f^{\widetilde{\Gamma}}(P_{\theta_{n,m}^*,m} \| Q_n) \le \inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(P_{\theta,m} \| Q_n) + \epsilon_{n,m}^{opt} \quad \mathbb{P}\text{-a.s.}$$

$$(67)$$

as opposed to (66).

The concentration inequalities that we derive below rely on the following decompositions of the (f, Γ) -GAN error. These should be compared with the result for IPM-GANs, Lemma 9 of Huang et al. [2022].

Lemma 3.5 $((f, \Gamma)$ -GAN Error Decomposition). *Under the Assumption 3.2 the* (f, Γ) -GAN *error can be decomposed* \mathbb{P} -a.s. as follows:

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta})$$

$$\leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}]) \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\} + \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q_{n}}[h] - E_{Q}[h] \right\}$$

$$+ \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha) \right) \sup_{h \in \Gamma} \inf_{\widetilde{h} \in \widetilde{\Gamma}} \|h - \widetilde{h}\|_{\infty} + \epsilon_{n,m}^{opt}.$$

$$(68)$$

Proof. Using (66) we can compute the \mathbb{P} -a.s. bound

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta})$$

$$= D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta,m}) + \inf_{\theta \in \Theta} D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta,m}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta})$$

$$\leq D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta_{n,m}^{*},m}) + \epsilon_{n,m}^{opt} + \inf_{\theta \in \Theta} D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta,m}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta}) .$$

$$(69)$$

Using the fact that $\widetilde{\Gamma}\subset \Gamma$ implies $D_f^{\widetilde{\Gamma}}\leq D_f^{\Gamma}$ along with Lemma 3.1 we then find

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta})$$

$$\leq D_{f}^{\widetilde{\Gamma}}(Q \| P_{\theta_{n,m}^{*}}) - D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta_{n,m}^{*},m}) + \inf_{\theta \in \Theta} D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta,m}) - \inf_{\theta \in \Theta} D_{f}^{\widetilde{\Gamma}}(Q \| P_{\theta})$$

$$+ \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)\right) \sup_{h \in \Gamma} \inf_{\widetilde{h} \in \widetilde{\Gamma}} \|h - \widetilde{h}\|_{\infty} + \epsilon_{n,m}^{opt}.$$

$$(70)$$

Next we make use of the simple bound

$$\pm (\sup_{i \in I} d_i - \sup_{i \in I} c_i) \le \sup_{i \in I} \{ \pm (d_i - c_i) \}$$
 (71)

whenever $c_i, d_i \in \mathbb{R}$ for all $i \in I$ and $\sup_i c_i, \sup_i d_i$ are finite. Using the definition (3) along with (24) and (71) we can compute

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta}) \leq \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q}[h] - \Lambda_{f}^{P_{\theta_{n,m}^{*}}}[h] - \left(E_{Q_{n}}[h] - \Lambda_{f}^{P_{\theta_{n,m}^{*}},m}[h] \right) \right\}$$

$$+ \sup_{\theta \in \Theta} \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q_{n}}[h] - \Lambda_{f}^{P_{\theta,m}}[h] - \left(E_{Q}[h] - \Lambda_{f}^{P_{\theta}}[h] \right) \right\}$$

$$+ \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha) \right) \sup_{h \in \Gamma} \inf_{h \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty} + \epsilon_{n,m}^{opt}$$

$$\leq \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{\theta,m}}[h] - \Lambda_{f}^{P_{\theta,m}}[h] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q_{n}}[h] - E_{Q}[h] \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{\theta}}[h] - \Lambda_{f}^{P_{\theta,m}}[h] \right\}$$

$$+ \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha) \right) \sup_{h \in \Gamma} \inf_{h \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty} + \epsilon_{n,m}^{opt}.$$

We have $\Lambda_f^{P_{\theta}}[h] = \Lambda_f^{P_Z}[h \circ \Phi_{\theta}]$ and $\Lambda_f^{P_{\theta,m}}[h] = \Lambda_f^{P_{Z,m}}[h \circ \Phi_{\theta}]$ and so this completes the proof.

We also obtain an error decomposition in the case where the arguments of the GAN are reversed. The proof is very similar and so we omit the details.

Lemma 3.6 (Reverse (f, Γ) -GAN Error Decomposition). *Under Assumption 3.4 the* (f, Γ) -GAN *error can be decomposed* \mathbb{P} -a.s. as follows:

$$D_{f}^{\Gamma}(P_{\theta_{n,m}^{*}}\|Q) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(P_{\theta}\|Q)$$

$$\leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ E_{P_{Z}}[h \circ \Phi_{\theta}] - E_{P_{Z,m}}[h \circ \Phi_{\theta}] \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ E_{P_{Z,m}}[h \circ \Phi_{\theta}] - E_{P_{Z}}[h \circ \Phi_{\theta}] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \left\{ \Lambda_{f}^{Q_{n}}[h] - \Lambda_{f}^{Q}[h] \right\} + \sup_{h \in \widetilde{\Gamma}} \left\{ \Lambda_{f}^{Q}[h] - \Lambda_{f}^{Q_{n}}[h] \right\}$$

$$+ \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha) \right) \sup_{h \in \Gamma} \inf_{\widetilde{h} \in \widetilde{\Gamma}} \|h - \widetilde{h}\|_{\infty} + \epsilon_{n,m}^{opt}.$$

$$(73)$$

We note that the setting of Lemmas 3.5 and 3.6 differs in a few ways from that of Huang et al. [2022]. Namely we assume $\widetilde{\Gamma} \subset \Gamma$ and we also do not assume that $\inf_{\theta \in \Theta} D_f^{\Gamma}(Q \| P_{\theta}) = 0$ (resp. $\inf_{\theta \in \Theta} D_f^{\Gamma}(P_{\theta} \| Q) = 0$). Under appropriate assumptions, the latter can be proven to hold for a sufficiently rich class of generators. More specifically, the analogue of Huang et al. [2022] would be to work under the assumption that $\inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(\mu_n \| P_{\theta}) = 0$ (resp. $\inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(P_{\theta} \| \mu_n) = 0$) for all empirical distributions μ_n . As $D_f^{\widetilde{\Gamma}} \leq d_{\widetilde{\Gamma}}$ (see (112) in Appendix B), this zero generator approximation error property holds for the (f, Γ) -divergence whenever it holds for the the corresponding IPM; see Yang et al. [2022] and the discussion in Section 2.2.1 of Huang et al. [2022] for sufficient conditions. Error decompositions that are adapted to the zero-approximation-error setting, and its argument-reversal, are given below.

Lemma 3.7 $((f,\Gamma)\text{-GAN Error Decomposition 2})$. Under Assumption 3.2, and supposing that $\inf_{\theta\in\Theta}D_f^{\widetilde{\Gamma}}(\mu_n\|P_\theta)=0$ for all empirical distributions μ_n , the $(f,\Gamma)\text{-GAN error can be decomposed }\mathbb{P}\text{-}a.s.$ as follows:

$$D_{f}^{\Gamma}(Q||P_{\theta_{n,m}^{*}}) \leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}] \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\} + \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha) \right) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} ||h - \tilde{h}||_{\infty} + \epsilon_{n,m}^{opt}.$$

$$(74)$$

Proof. Using (66) along with Lemma 3.1 we can compute the \mathbb{P} -a.s. bound

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) \leq D_{f}^{\widetilde{\Gamma}}(Q \| P_{\theta_{n,m}^{*}}) + \inf_{\theta \in \Theta} D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta,m}) - D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta_{n,m}^{*},m})$$

$$+ \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)\right) \sup_{h \in \Gamma} \inf_{\widetilde{h} \in \widetilde{\Gamma}} \| h - \widetilde{h} \|_{\infty} + \epsilon_{n,m}^{opt}.$$

$$(75)$$

As a simple consequence of the definition (3), for all $\theta \in \Theta$ we have

$$D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta,m}) \le D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta}) + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \{ \Lambda_f^{P_{\theta}}[h] - \Lambda_f^{P_{\theta,m}}[h] \}. \tag{76}$$

Minimizing over $\theta \in \Theta$ and using the zero-approximation-error property we obtain

$$\inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta,m}) \le \inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(Q_n \| P_{\theta}) + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \{ \Lambda_f^{P_{\theta}}[h] - \Lambda_f^{P_{\theta,m}}[h] \}$$

$$= \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \{ \Lambda_f^{P_{\theta}}[h] - \Lambda_f^{P_{\theta,m}}[h] \}.$$
(77)

Therefore

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) \leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \{ \Lambda_{f}^{P_{\theta}}[h] - \Lambda_{f}^{P_{\theta,m}}[h] \} + (1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)) \sup_{h \in \Gamma} \inf_{\widetilde{h} \in \widetilde{\Gamma}} \| h - \widetilde{h} \|_{\infty} + \epsilon_{n,m}^{opt}$$

$$+ D_{f}^{\widetilde{\Gamma}}(Q \| P_{\theta_{n,m}^{*}}) - D_{f}^{\widetilde{\Gamma}}(Q_{n} \| P_{\theta_{n,m}^{*},m})$$

$$\leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \{ \Lambda_{f}^{P_{\theta}}[h] - \Lambda_{f}^{P_{\theta,m}}[h] \} + (1 + (f^{*})'_{+}(z_{0} + \beta - \alpha)) \sup_{h \in \Gamma} \inf_{\widetilde{h} \in \widetilde{\Gamma}} \| h - \widetilde{h} \|_{\infty} + \epsilon_{n,m}^{opt}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \{ E_{Q}[h] - E_{Q_{n}}[h] \} + \sup_{h \in \widetilde{\Gamma}} \{ \Lambda_{f}^{P_{\theta_{n,m}^{*},m}}[h] - \Lambda_{f}^{P_{\theta_{n,m}^{*},m}}[h] \} .$$

$$(78)$$

Bounding the last term by maximizing over $\theta \in \Theta$ and then using $\Lambda_f^{P_\theta}[h] = \Lambda_f^{P_Z}[h \circ \Phi_\theta]$ and $\Lambda_f^{P_{\theta,m}}[h] = \Lambda_f^{P_{Z,m}}[h \circ \Phi_\theta]$ completes the proof.

Lemma 3.8 (Reverse (f,Γ) -GAN Error Decomposition 2). Under Assumption 3.4, and supposing that $\inf_{\theta\in\Theta}D_f^{\widetilde{\Gamma}}(P_{\theta}\|\mu_n)=0$ for all empirical distributions μ_n , the (f,Γ) -GAN error can be decomposed \mathbb{P} -a.s. as follows:

$$D_{f}^{\Gamma}(P_{\theta_{n,m}^{*}}\|Q) \leq \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ E_{P_{Z}}[h \circ \Phi_{\theta}] - E_{P_{Z,m}}[h \circ \Phi_{\theta}] \right\} + \sup_{h \in \widetilde{\Gamma}, \theta \in \Theta} \left\{ E_{P_{Z,m}}[h \circ \Phi_{\theta}] - E_{P_{Z}}[h \circ \Phi_{\theta}] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}} \left\{ \Lambda_{f}^{Q_{n}}[h] - \Lambda_{f}^{Q}[h] \right\} + \left(1 + (f^{*})'_{+}(z_{0} + \beta - \alpha) \right) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty} + \epsilon_{n,m}^{opt}.$$

$$(79)$$

The proof of (3.8) is similar to that of (3.7) and so we omit the details. Note that the terms in the bounds (74) and (79) reduce to the terms in IPM case, Lemma 9 in Huang et al. [2022], if $f^*(z) = z$ and $-\widetilde{\Gamma} \subset \widetilde{\Gamma}$.

3.2 Concentration Inequalities for (f, Γ) -GANs

We are now ready to derive concentration inequalities for (f,Γ) -GANs. Due to the asymmetry of $D_f^\Gamma(Q\|P)$ in Q and P, the bounds differ based on whether the generator is in the first or second argument of D_f^Γ . We also have variants depending on whether one assumes the zero generator approximation error property for empirical measures. The key ingredients are the ULLN for the generalized cumulant generating function in Lemmas 2.9 and 2.10 along with the perturbation bound from Lemma 2.7, the latter being needed in order to apply McDiarmid's inequality to the terms involving the generalized cumulant generating function. The following definition lists several quantities that will appear in the concentration inequalities below. We note that these quantities approach their counterparts in the linear (i.e., IPM) case in the appropriate limit; see the discussion in Remarks 2.8 and 2.11.

Definition 3.9. With notation as in either Assumption 3.2 or 3.4:

1. Define the discriminator-space approximation error

$$\epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} := \left(1 + (f^*)'_{+}(z_0 + \beta - \alpha)\right) \sup_{h \in \Gamma} \inf_{\tilde{h} \in \widetilde{\Gamma}} \|h - \tilde{h}\|_{\infty}. \tag{80}$$

Note that this vanishes when $\widetilde{\Gamma} = \Gamma$.

2. For $n \in \mathbb{Z}^+$ define

$$\Delta_{f,n} := \inf_{z \in [z_0 - (\beta - \alpha), z_0]} \left\{ n \left(-z + \frac{n-1}{n} f^*(z) + \frac{1}{n} f^*(\beta - \alpha + z) \right) \right\}$$
(81)

and note that $\Delta_{f,n} \in [\beta - \alpha, f^*(\beta - \alpha + z_0) - z_0].$

3. Let $n \in \mathbb{Z}^+$, Ψ be a nonempty family of measurable functions on \mathcal{X} , and P a probability measure on \mathcal{X} . If $(f^*)'_+$ is $L_{\alpha,\beta}$ -Lipschitz on $[z_0 - (\beta - \alpha), z_0 + \beta - \alpha]$ and Ψ contains a constant function then define

$$\mathcal{K}_{f,\Psi,P,n} = \min \left\{ (1 + 2(\beta - \alpha)L_{\alpha,\beta})\mathcal{R}_{\Psi,P,n} + \frac{(\beta - \alpha)^2 L_{\alpha,\beta}}{2n^{1/2}}, (f^*)'_{+}(\beta - \alpha + z_0) \left(\mathcal{R}_{\Psi,P,n} + \frac{\beta - \alpha}{2n^{1/2}}\right) \right\}$$
(82)

and otherwise define

$$\mathcal{K}_{f,\Psi,P,n} := (f^*)'_{+}(\beta - \alpha + z_0) \left(\mathcal{R}_{\Psi,P,n} + \frac{\beta - \alpha}{2n^{1/2}} \right). \tag{83}$$

Theorem 3.10 ((f, Γ) -GAN Concentration Inequalities). *Under Assumption 3.2, and in particular with* $\theta_{n,m}^*$ *the approximate solution to the empirical* (f, Γ) -GAN problem (66), for $\epsilon > 0$ we have

$$\mathbb{P}\left(D_f^{\Gamma}(Q||P_{\theta_{n,m}^*}) - \inf_{\theta \in \Theta} D_f^{\Gamma}(Q||P_{\theta}) \ge \epsilon + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{opt}^{n,m} + 4\mathcal{R}_{\widetilde{\Gamma},Q,n} + 4\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}\right)$$

$$\le \exp\left(-\frac{\epsilon^2}{\frac{2}{n}(\beta - \alpha)^2 + \frac{2}{m}\Delta_{f,m}^2}\right),$$
(84)

where we refer to the quantities in Definition 3.9.

If, in addition, $\inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(\mu_n \| P_{\theta}) = 0$ for all possible empirical distributions μ_n then we obtain the tighter bound

$$\mathbb{P}\left(D_f^{\Gamma}(Q||P_{\theta_{n,m}^*}) \ge \epsilon + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{opt}^{n,m} + 2\mathcal{R}_{\widetilde{\Gamma},Q,n} + 4\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}\right) \le \exp\left(-\frac{\epsilon^2}{\frac{1}{2n}(\beta - \alpha)^2 + \frac{2}{m}\Delta_{f,m}^2}\right). \tag{85}$$

Proof. First note that Lemma 2.3 and Corollary 2.6 together with the dominated convergence theorem imply $\theta \mapsto \Lambda_f^{Pz}[h \circ \Phi_\theta]$ is are continuous and also that $h_j \to h$ pointwise implies $\Lambda_f^{Pz}[h_j \circ \Phi_\theta] \to \Lambda_f^{Pz}[h \circ \Phi_\theta]$, $E_Q[h_j] \to E_Q[h]$, and similarly for the empirical variants. Therefore, letting Θ_0 denote a countable dense subset of Θ , the suprema in Lemma 3.5 can be restricted to the countable subsets $\widetilde{\Gamma}_0$ and Θ_0 , giving the \mathbb{P} -a.s. bound

$$D_{f}^{\Gamma}(Q \| P_{\theta_{n,m}^{*}}) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(Q \| P_{\theta})$$

$$\leq \sup_{h \in \widetilde{\Gamma}_{0}, \theta \in \Theta_{0}} \left\{ \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}]) \right\} + \sup_{h \in \widetilde{\Gamma}_{0}, \theta \in \Theta_{0}} \left\{ \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}_{0}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\} + \sup_{h \in \widetilde{\Gamma}_{0}} \left\{ E_{Q_{n}}[h] - E_{Q}[h] \right\}$$

$$+ \epsilon_{approx}^{\Gamma, \widetilde{\Gamma}} + \epsilon_{n,m}^{opt}.$$

$$(86)$$

Now we will apply McDiarmid's inequality to the right-hand side. The map $H: \mathcal{X}^n \times \mathcal{Z}^m \to \mathbb{R}$ defined by

$$H(x,z) = \sup_{h \in \widetilde{\Gamma}_{0}, \theta \in \Theta_{0}} \left\{ \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}]) \right\} + \sup_{h \in \widetilde{\Gamma}_{0}, \theta \in \Theta_{0}} \left\{ \Lambda_{f}^{P_{Z}}[h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z,m}}[h \circ \Phi_{\theta}] \right\}$$

$$+ \sup_{h \in \widetilde{\Gamma}_{0}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\} + \sup_{h \in \widetilde{\Gamma}_{0}} \left\{ E_{Q_{n}}[h] - E_{Q}[h] \right\}$$

$$(87)$$

is measurable and if $x, \tilde{x} \in \mathcal{X}^n$ differ only in the j'th component then

$$|H(x,z) - H(\tilde{x},z)| \le 2 \sup_{h \in \tilde{\Gamma}_{0}} \left\{ \left| \frac{1}{n} \sum_{i=1}^{n} h(x_{i}) - \frac{1}{n} \sum_{i=1}^{n} h(\tilde{x}_{i}) \right| \right\}$$

$$\le \frac{2}{n} \sup_{h \in \tilde{\Gamma}_{0}} \left\{ |h(x_{j}) - h(x'_{j})| \right\} \le \frac{2}{n} (\beta - \alpha),$$
(88)

while if $z, \tilde{z} \in \mathbb{Z}^m$ differ only in the j'th component then Lemma 2.5 and the perturbation bound from Lemma 2.7 imply

$$|H(x,z) - H(x,\tilde{z})| \le 2 \sup_{h \in \widetilde{\Gamma}_0, \theta \in \Theta_0} \{ |\Lambda_f \circ (h \circ \Phi_\theta)_m(z) - \Lambda_f \circ (h \circ \Phi_\theta)_m(\tilde{z})| \} \le \frac{2}{m} \Delta_{f,m}.$$
 (89)

Therefore we can apply McDiarmid's inequality to H, see, e.g., Theorem D.8 in Mohri et al. [2018], to obtain

$$\mathbb{P}(H(X,Z) \ge \epsilon + \mathbb{E}[H(X,Z)]) \le \exp\left(-\frac{\epsilon^2}{\frac{2}{n}(\beta - \alpha)^2 + \frac{2}{m}\Delta_{f,m}^2}\right). \tag{90}$$

In terms of H, the error bound (86) becomes the a.s. bound

$$D_f^{\Gamma}(Q||P_{\theta_{n,m}^*}) - \inf_{\theta \in \Theta} D_f^{\Gamma}(Q||P_{\theta}) \le H(X,Z) + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{n,m}^{opt}.$$

$$\tag{91}$$

Using the ULLN for means, see Theorem C.4, along with the new ULLN for the generalized cumulant generating function in Lemmas 2.9 and 2.10 we obtain

$$\mathbb{E}[H(X,Z)] \le 4\mathcal{R}_{\widetilde{\Gamma},O,n} + 4\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}.$$
(92)

Combining (90), (91), and (92) we arrive at the claimed result (84).

If we also assume $\inf_{\theta \in \Theta} D_f^{\tilde{\Gamma}}(\mu_n || P_{\theta}) = 0$ for all possible empirical distributions μ_n then we can write the error decomposition 3.7 as

$$D_f^{\Gamma}(Q||P_{\theta_{n,m}^*}) \le H(X,Z) + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{n,m}^{opt}$$
(93)

 \mathbb{P} -a.s., where we now define

$$H(X,Z) := \sup_{h \in \widetilde{\Gamma}_{0}, \theta \in \Theta_{0}} \left\{ \Lambda_{f}^{P_{Z,m}} [h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z}} [h \circ \Phi_{\theta}] \right\} + \sup_{h \in \widetilde{\Gamma}_{0}, \theta \in \Theta_{0}} \left\{ \Lambda_{f}^{P_{Z}} [h \circ \Phi_{\theta}] - \Lambda_{f}^{P_{Z,m}} [h \circ \Phi_{\theta}] \right\}$$
(94)
+
$$\sup_{h \in \widetilde{\Gamma}_{0}} \left\{ E_{Q}[h] - E_{Q_{n}}[h] \right\}.$$

Similar to the above, when x and \tilde{x} differ in only a single component we can bound

$$|H(x,z) - H(\tilde{x},z)| \le \frac{1}{n}(\beta - \alpha) \tag{95}$$

and when z and \tilde{z} differ in only a single component we can bound

$$|H(x,z) - H(x,\tilde{z})| \le \frac{2}{m} \Delta_{f,m}. \tag{96}$$

The mean of H(X, Z) can be bounded via Theorem C.4 and Lemmas 2.9 and 2.10:

$$\mathbb{E}[H(X,Z)] \le 2\mathcal{R}_{\widetilde{\Gamma},Q,n} + 4\mathcal{K}_{f,\widetilde{\Gamma}\circ\Phi,P_Z,m}. \tag{97}$$

Combining (93), and (95) - (97) with McDiarmid's inequality completes the proof of (85).

Similarly, we have concentration inequalities for the reverse (f, Γ) -GANs. The proof is very similar and so we omit the details.

Theorem 3.11 (Reverse (f, Γ) -GAN Concentration Inequalities). *Under Assumption 3.4, and in particular with* $\theta_{n,m}^*$ *the approximate solution to the empirical* (f, Γ) -GAN problem (67), for $\epsilon > 0$ we have

$$\mathbb{P}\left(D_{f}^{\Gamma}(P_{\theta_{n,m}^{*}}\|Q) - \inf_{\theta \in \Theta} D_{f}^{\Gamma}(P_{\theta}\|Q) \ge \epsilon + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{opt}^{n,m} + 4\mathcal{R}_{\widetilde{\Gamma} \circ \Phi, P_{Z}, m} + 4\mathcal{K}_{f,\widetilde{\Gamma}, Q, n}\right)$$

$$\le \exp\left(-\frac{\epsilon^{2}}{\frac{2}{m}(\beta - \alpha)^{2} + \frac{2}{n}\Delta_{f, n}^{2}}\right),$$
(98)

where we refer to the quantities in Definition 3.9.

If, in addition, $\inf_{\theta \in \Theta} D_f^{\widetilde{\Gamma}}(P_{\theta} \| \mu_n) = 0$ for all possible empirical distributions μ_n then we obtain the tighter bound

$$\mathbb{P}\left(D_f^{\Gamma}(P_{\theta_{n,m}^*}\|Q) \ge \epsilon + \epsilon_{approx}^{\Gamma,\widetilde{\Gamma}} + \epsilon_{opt}^{n,m} + 4\mathcal{R}_{\widetilde{\Gamma} \circ \Phi, P_Z, m} + 2\mathcal{K}_{f,\widetilde{\Gamma},Q,n}\right) \le \exp\left(-\frac{\epsilon^2}{\frac{2}{m}(\beta - \alpha)^2 + \frac{1}{2n}\Delta_{f,n}^2}\right). \tag{99}$$

Remark 3.12. As stated, the above results assume a space of discriminators, Γ , that satisfies a uniform bound of the form $\alpha \leq h \leq \beta$ for all $h \in \Gamma$. However, we note that the (f,Γ) -divergence objective functional in (3) is invariant under constant shifts, due to the identity $\Lambda_f^P[h+c] = \Lambda_f^P[h] + c$ for all $c \in \mathbb{R}$. Therefore, if we have discriminators Ψ such that the ranges of $\psi \in \Psi$ have uniformly bounded diameter, i.e., there exists β such that $\sup_{x,\tilde{x}} |\psi(x) - \psi(\tilde{x})| \leq \beta$ for all $\psi \in \Psi$, then we can write

$$D_f^{\Psi}(Q||P) = D_f^{\Gamma}(Q||P), \qquad (100)$$

where $\Gamma \coloneqq \{\psi - \inf \psi : \psi \in \Psi\}$ satisfies $0 \le h \le \beta$ for all $h \in \Gamma$. Thus our theory can be applied to D_f^{Γ} and hence also to D_f^{Ψ} via (100). In this way, our theorems can be applied to (f, Γ) -GANs with discriminators whose ranges have uniformly bounded diameter, e.g., 1-Lipschitz functions on a compact domain.

Remark 3.13 (Bounds on the mean of the (f,Γ) -GAN error). Bounds on the mean of the (f,Γ) -GAN error are implicit in the above derivations. They are obtained by combining the error decompositions in Section 3.1 with the ULLN results in Lemma 2.9, Lemma 2.10, and Theorem C.4. In addition, a standard technique can be used to turn the concentration inequalities into L^q error bounds for any q > 1 as follows: Let Y be a non-negative random variable, $a \in [0, \infty)$, and $K : (0, \infty) \to [0, \infty)$ be measurable such that

$$\mathbb{P}(Y \ge \epsilon + a) \le K(\epsilon) \tag{101}$$

for all $\epsilon > 0$. Then for q > 0 we have

$$\mathbb{E}[Y^q] \le a^q + q \int_0^\infty (a+\epsilon)^{q-1} K(\epsilon) d\epsilon.$$
 (102)

This follows from rewriting

$$\mathbb{E}[Y^q] = \int_0^\infty \mathbb{P}(Y^q \ge r) dr, \qquad (103)$$

breaking the domain of integration into $[0, a^q]$ and $[a^q, \infty)$, then changing variables in the second term and using the bound (101).

3.3 Concentration Inequalities for the Estimation of (f, Γ) -Divergences

The tools developed in Section 2 can also be used to address the simpler problem of concentration inequalities for the estimation of $D_f^{\Gamma}(Q\|P)$ using samples from Q and P. Special cases of this problem were previously considered in Chen et al. [2023a, 2024]. Both considered the case where f corresponded to the family of α -divergences and Γ was the set of L-Lipschitz functions; the former focused on the consequences of group symmetry while the latter focused on heavy-tailed distributions. The techniques developed in this work allow us to derive concentration inequalities for a much wider class of f's and Γ 's; we do not consider the consequences of group symmetry or heavy-tailed distributions in this work.

Theorem 3.14. Let Γ be a nonempty set of measurable functions on $\mathcal X$ and suppose we have $\alpha, \beta \in \mathbb R$ such that $\alpha < \beta$ and $\alpha \le h \le \beta$ for all $h \in \Gamma$. Assume there exists a countable $\Gamma_0 \subset \Gamma$ such that for all $h \in \Gamma$ there exists a sequence $h_j \in \Gamma_0$ such that $h_j \to h$ pointwise. Let $f \in \mathcal F_1(a,b)$ with $a \ge 0$ and assume f is strictly convex in a neighborhood of 1 and $z_0 + \beta - \alpha \in \{f^* < \infty\}^o$.

Let Q, P be probability measures on \mathcal{X} and X_i , i = 1, ..., n, X'_i , i = 1, ..., m be jointly independent samples from Q and P respectively with Q_n , P_m the corresponding empirical measures. Then for $\epsilon > 0$ we have

$$\mathbb{P}\left(D_f^{\Gamma}(Q\|P) - D_f^{\Gamma}(Q_n\|P_m) \ge \epsilon\right) \le \exp\left(-\frac{2\epsilon^2}{\frac{1}{n}(\beta - \alpha)^2 + \frac{1}{m}\Delta_{f,m}^2}\right),\tag{104}$$

$$\mathbb{P}\left(D_f^{\Gamma}(Q_n\|P_m) - D_f^{\Gamma}(Q\|P) \ge \epsilon + 2\mathcal{R}_{\Gamma,Q,n} + 2\mathcal{K}_{f,\Gamma,P,m}\right) \le \exp\left(-\frac{2\epsilon^2}{\frac{1}{n}(\beta - \alpha)^2 + \frac{1}{m}\Delta_{f,m}^2}\right),\tag{105}$$

where $\Delta_{f,m}$ and $\mathcal{K}_{f,\Gamma,P,m}$ were defined in Definition 3.9.

Proof. Similar to the proof of Theorem 3.10, if we define

$$H(X, X') = D_f^{\Gamma}(Q_n || P_m), \qquad (106)$$

then when x and \tilde{x} differ by a single component we have

$$|H(x,x') - H(\tilde{x},x')| \le \frac{1}{n}(\beta - \alpha) \tag{107}$$

and when x' and \tilde{x}' differ by a single component, Lemma 2.7 implies

$$|H(x,x') - H(x,\tilde{x}')| \le \frac{1}{m} \Delta_{f,m}$$
 (108)

Therefore we can use McDiarmid's inequality to conclude that

$$\mathbb{P}\left(\pm(D_f^{\Gamma}(Q_n\|P_m) - \mathbb{E}[D_f^{\Gamma}(Q_n\|P_m)) \ge \epsilon\right) \le \exp\left(-\frac{2\epsilon^2}{\frac{1}{n}(\beta - \alpha)^2 + \frac{1}{m}\Delta_{f,m}^2}\right). \tag{109}$$

Combining (109) (lower sign) with the result of the calculation

$$\mathbb{E}\left[D_{f}^{\Gamma}(Q_{n}\|P_{m})\right] = \mathbb{E}\left[\sup_{h\in\Gamma,\nu\in[\alpha-z_{0},\beta-z_{0}]} \left\{E_{Q_{n}}[h-\nu] - E_{P_{m}}[f^{*}(h-\nu)]\right\}\right]$$

$$\geq \sup_{h\in\Gamma,\nu\in[\alpha-z_{0},\beta-z_{0}]} \mathbb{E}\left[E_{Q_{n}}[h-\nu] - E_{P_{m}}[f^{*}(h-\nu)]\right]$$

$$= \sup_{h\in\Gamma,\nu\in[\alpha-z_{0},\beta-z_{0}]} \left\{E_{Q}[h-\nu] - E_{P}[f^{*}(h-\nu)]\right\} = D_{f}^{\Gamma}(Q\|P)$$
(110)

we obtain (104). Combining (109) (upper sign) with the result of the calculation

$$\mathbb{E}\left[D_f^{\Gamma}(Q_n\|P_m)\right] - D_f^{\Gamma}(Q\|P) \le \mathbb{E}\left[\sup_{h \in \Gamma} \left\{E_{Q_n}[h] - E_Q[h]\right\}\right] + \mathbb{E}\left[\sup_{h \in \Gamma} \left\{\Lambda_f^P[h] - \Lambda_f^{P_m}[h]\right\}\right]$$
(111)

and then using Theorem C.4 and Lemmas 2.9 and 2.10 we obtain (105).

4 Conclusion

We have proven the statistical consistency of (f,Γ) -GANs, a large class of GANs with nonlinear objective functionals. These GANs are based on the (f,Γ) -divergences, which generalize and interpolate between integral probability metrics (IPMs, e.g., 1-Wasserstein) and f-divergences (e.g., KL) and have been show to outperform both IPM and f-divergence-based methods in a number of applications. This paper extends earlier techniques for proving consistency of GANs with linear objective functionals (IPM-GANs) to the nonlinear setting. The key technical results are the tight perturbation bound in Lemma 2.7, uniform law of large numbers bounds for a class of nonlinear functionals in Lemmas 2.9 and 2.10, and the (f,Γ) -GAN error decompositions in Section 3.1. These results allow for the derivation of finite-sample concentration inequalities for (f,Γ) -GANs in Theorems 3.10 and 3.11, thereby proving their statistical consistency. The bounds have the same qualitative form as those previously obtained for IPM-GANs and so we can also conclude that the performance of (f,Γ) -GANs scales with the intrinsic dimension of the data source in a similar fashion to IPM-GANs.

A Properties of f and f^*

In this appendix we collect a number of important properties of the convex function f and its Legendre transform f^* that are needed in the study of (f,Γ) -GANs. For a,b satisfying $-\infty \le a < 1 < b \le \infty$ we define $\mathcal{F}_1(a,b)$ to be the set of convex $f:(a,b)\to\mathbb{R}$ with f(1)=0. For $f\in\mathcal{F}_1(a,b)$, standard convex functions theory, see, e.g., Rockafellar [1970] and Appendix A in Birrell et al. [2022a], implies that f and its Legendre transform $f^*:\mathbb{R}\to(-\infty,\infty]$, $f^*(z)=\sup_{t\in(a,b)}\{zt-f(t)\}$, have the following properties:

- 1. $f^*(z) \ge z$ for all $z \in \mathbb{R}$ (this uses f(1) = 0).
- 2. If $a \ge 0$ then f^* is non-decreasing.
- 3. $(f^*)^* = f$.
- 4. f^* is convex and LSC.
- 5. f^* is continuous on $\overline{\{f^* < \infty\}}$, where \overline{A} denotes the closure of a set A.

- 6. The right derivative $(f^*)'_+$ exists and is finite on $\{f^* < \infty\}^o$, where A^o denotes the interior of a set A. Similarly, f'_+ exists and is finite on (a,b).
- 7. $(f^*)'_+$ is non-decreasing and absolutely continuous on compact intervals; the latter implies that f^* satisfies the fundamental theorem of calculus (see, e.g., Theorem 3.35 and exercise 42 in Folland [2013]).

Of particular relevance will be the value $z_0 := f'_+(1)$, which exists and is finite due to the assumption that $1 \in (a, b)$. The importance of z_0 to the (f, Γ) -divergences was observed in Birrell et al. [2022a], where the following properties were proven (see Lemma A.9):

Lemma A.1. Define $z_0 := f'_+(1)$.

- 1. $f^*(z_0) = z_0$
- 2. If f is strictly convex on a neighborhood of 1 and $z_0 \in \{f^* < \infty\}^o$ then $(f^*)'_+(z_0) = 1$.

These properties will be key to our analysis and so we will generally assume that f is strictly convex on a neighborhood of 1 and $z_0 \in \{f^* < \infty\}^o$.

B Properties of the (f, Γ) -Divergences

In this appendix we collect several important properties of the (f,Γ) -divergences. For \mathcal{X} a measurable space we let $\mathcal{M}_b(\mathcal{X})$ denote the space of bounded measurable real-valued functions on \mathcal{X} and $\mathcal{P}(\mathcal{X})$ be the set of probability measures on \mathcal{X} . The following results are taken from Theorem 2.8 of Birrell et al. [2022a].

Theorem B.1. Let $f \in \mathcal{F}_1(a,b)$, $\Gamma \subset \mathcal{M}_b(\mathcal{X})$ be nonempty, and $Q, P \in \mathcal{P}(\mathcal{X})$.

1.

$$D_f^{\Gamma}(Q||P) \le \inf_{\eta \in \mathcal{P}(\mathcal{X})} \{ D_f(\eta||P) + d_{\Gamma}(Q,\eta) \}. \tag{112}$$

In particular, $D_f^{\Gamma}(Q||P) \leq \min\{D_f(Q||P), d_{\Gamma}(Q,P)\}.$

- 2. The map $(Q, P) \in \mathcal{P}(S) \times \mathcal{P}(S) \mapsto D_f^{\Gamma}(Q||P)$ is convex.
- 3. If there exists $c_0 \in \Gamma \cap \mathbb{R}$ then $D_f^{\Gamma}(Q||P) \geq 0$.
- *4.* Suppose f and Γ satisfy the following:
 - (a) There exist a nonempty set $\Psi \subset \Gamma$ with the following properties: i. Ψ is $\mathcal{P}(\mathcal{X})$ -determining, i.e., for all $Q, P \in \mathcal{P}(\mathcal{X})$, $E_Q[\psi] = E_P[\psi]$ for all $\psi \in \Psi$ implies Q = P. ii. For all $\psi \in \Psi$ there exists $c_0 \in \mathbb{R}$, $\epsilon_0 > 0$ such that $c_0 + \epsilon \psi \in \Gamma$ for all $|\epsilon| < \epsilon_0$.
 - (b) f is strictly convex on a neighborhood of 1.
 - (c) f^* is finite and C^1 on a neighborhood of $f'_+(1)$.

Then D_f^{Γ} has the divergence property, i.e., $D_f^{\Gamma}(Q||P) \geq 0$ with equality if and only if Q = P.

Remark B.2. Under stronger assumptions one can show that (112) is in fact an equality; see Theorem 2.15 in Birrell et al. [2022a].

Remark B.3. Assumptions 4(b) and 4(c) hold, for instance, if f is strictly convex on (a,b) and $f'_{+}(1) \in \{f^* < \infty\}^o$ (see Theorem 26.3 in Rockafellar [1970]).

C Rademacher Complexity and Uniform Law of Large Numbers

In this appendix we recall the definition of Rademacher complexity and its use in proving uniform law of large numbers (ULLN) results. There are two definitions of Rademacher complexity in common use, those being with and without absolute value. In this work we use the version without absolute value as defined below.

Definition C.1. Let Ψ be a collection of functions on \mathcal{Y} and $n \in \mathbb{Z}^+$. The empirical Rademacher complexity of Ψ at $y \in \mathcal{Y}^n$ is defined by

$$\widehat{\mathcal{R}}_{\Psi,n}(y) := E_{\sigma} \left[\sup_{\psi \in \Psi} \left\{ \frac{1}{n} \sigma \cdot \psi_n(y) \right\} \right], \tag{113}$$

where σ_i , i = 1, ..., n are independent uniform random variables taking values in $\{-1, 1\}$, i.e., Rademacher random variables, and $\psi_n(y) := (\psi(y_1), ..., \psi(y_n))$. Given a probability distribution P on \mathcal{Y} , the Rademacher complexity of Ψ relative to P is defined by

$$\mathcal{R}_{\Psi,P,n} := E_{P^n} \left[\widehat{\mathcal{R}}_{\Psi,n} \right] \tag{114}$$

where P^n is the n-fold product of P, i.e., the y_i 's become i.i.d. samples from P.

Remark C.2. Note that in (114) we are implicitly assuming that the empirical Rademacher complexity is measurable. When we use (114) in the main text, this will be guaranteed by other assumptions.

The definition (113) is particularly convenient for our purposes due to the following version of Talagrand's lemma; see Lemma 5.7 in Mohri et al. [2018].

Lemma C.3. Let $\alpha, \beta \in \mathbb{R}$ and Ψ be a collection of real-valued functions on \mathcal{Y} such that $\alpha \leq \psi \leq \beta$ for all $\psi \in \Psi$. If $\phi : [\alpha, \beta] \to \mathbb{R}$ is L-Lipschitz then

$$\widehat{\mathcal{R}}_{\phi \circ \Psi, n}(y) \le L \widehat{\mathcal{R}}_{\Psi, n}(y) \tag{115}$$

for all $n \in \mathbb{Z}^+$, $y \in \mathcal{Y}^n$, where $\phi \circ \Psi := \{\phi \circ \psi : \psi \in \Psi\}$.

Bounds on the Rademacher complexity can be used to prove ULLN bounds; see, e.g., Theorem 3.3, Eq. (3.8) - (3.13) in Mohri et al. [2018] (this reference assumes [0,1]-valued functions but the result can be freely shifted and scaled to apply to a set of uniformly bounded functions):

Theorem C.4 (ULLN). Let P be a probability measure on \mathcal{Y} and Ψ a countable family of real-valued measurable functions on \mathcal{Y} that are uniformly bounded. If Y_i , i=1,...,n are i.i.d. \mathcal{Y} -valued random variables that are distributed as P then for $n \in \mathbb{Z}^+$ we have

$$\mathbb{E}\left[\sup_{\psi\in\Psi}\left\{\pm\left(\frac{1}{n}\sum_{i=1}^{n}\psi(Y_i)-E_P[\psi]\right)\right\}\right]\leq 2\mathcal{R}_{\Psi,P,n}.$$
(116)

Remark C.5. We have stated the ULLN in terms of a countable collection of functions to avoid measurability issues, but under appropriate assumptions one can apply it to an uncountable Ψ , e.g., if there exists a countable $\Psi_0 \subset \Psi$ such that for every $\psi \in \Psi$ there exists a sequence $\psi_j \in \Psi_0$ with $\psi_j \to \psi$ pointwise.

References

Arjovsky, M., Chintala, S., and Bottou, L. Wasserstein generative adversarial networks. In Precup, D. and Teh, Y. W. (eds.), *Proceedings of the 34th International Conference on Machine Learning*, volume 70 of *Proceedings of Machine Learning Research*, pp. 214–223, International Convention Centre, Sydney, Australia, 06–11 Aug 2017. PMLR.

Belghazi, M. I., Baratin, A., Rajeshwar, S., Ozair, S., Bengio, Y., Courville, A., and Hjelm, D. Mutual information neural estimation. In Dy, J. and Krause, A. (eds.), *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pp. 531–540, Stockholmsmässan, Stockholm Sweden, 10–15 Jul 2018. PMLR. URL http://proceedings.mlr.press/v80/belghazi18a.html.

Ben-Tal, A. and Teboulle, M. An old-new concept of convex risk measures: The optimized certainty equivalent. *Mathematical Finance*, 17(3):449–476, 2007. doi: 10.1111/j.1467-9965.2007.00311.x. URL https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-9965.2007.00311.x.

Birrell, J., Dupuis, P., Katsoulakis, M. A., Pantazis, Y., and Rey-Bellet, L. (f, Gamma)-Divergences: Interpolating between f-divergences and integral probability metrics. *Journal of machine learning research*, 23(39):1–70, 2022a.

Birrell, J., Katsoulakis, M., Rey-Bellet, L., and Zhu, W. Structure-preserving GANs. *Proceedings of Machine Learning Research*, 162, 2022b.

Broniatowski, M. and Keziou, A. Minimization of divergences on sets of signed measures. *Studia Scientiarum Mathematicarum Hungarica*, 43(4):403–442, 2006.

Chakraborty, S. and Bartlett, P. L. On the statistical properties of generative adversarial models for low intrinsic data dimension. *arXiv* preprint arXiv:2401.15801, 2024.

Chen, Z., Katsoulakis, M., Rey-Bellet, L., and Zhu, W. Sample complexity of probability divergences under group symmetry. In *International Conference on Machine Learning*, pp. 4713–4734. PMLR, 2023a.

- Chen, Z., Katsoulakis, M. A., Rey-Bellet, L., and Zhu, W. Statistical guarantees of group-invariant GANs. *arXiv* preprint arXiv:2305.13517, 2023b.
- Chen, Z., Gu, H., Katsoulakis, M. A., Rey-Bellet, L., and Zhu, W. Learning heavy-tailed distributions with Wasserstein-proximal-regularized α -divergences. *arXiv* preprint arXiv:2405.13962, 2024.
- Dupuis, Paul and Mao, Yixiang. Formulation and properties of a divergence used to compare probability measures without absolute continuity. *ESAIM: COCV*, 28:10, 2022. doi: 10.1051/cocv/2022002. URL https://doi.org/10.1051/cocv/2022002.
- Dziugaite, G. K., Roy, D. M., and Ghahramani, Z. Training generative neural networks via maximum mean discrepancy optimization. *arXiv preprint arXiv:1505.03906*, 2015.
- Folland, G. Real Analysis: Modern Techniques and Their Applications. Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts. Wiley, 2013. ISBN 9781118626399. URL https://books.google.com/books?id=wI4fAwAAQBAJ.
- Glaser, P., Arbel, M., and Gretton, A. KALE flow: A relaxed KL gradient flow for probabilities with disjoint support. *Advances in Neural Information Processing Systems*, 34:8018–8031, 2021.
- Goodfellow, I. J., Pouget-Abadie, J., Mirza, M., Xu, B., Warde-Farley, D., Ozair, S., Courville, A., and Bengio, Y. Generative adversarial nets. In *Proceedings of the 27th International Conference on Neural Information Processing Systems Volume 2*, NIPS'14, pp. 2672–2680, Cambridge, MA, USA, 2014. MIT Press.
- Gretton, A., Borgwardt, K. M., Rasch, M. J., Schölkopf, B., and Smola, A. A kernel two-sample test. *J. Mach. Learn. Res.*, 13(1):723–773, 2012. ISSN 1532-4435. URL http://dl.acm.org/citation.cfm?id=2503308.2188410.
- Gulrajani, I., Ahmed, F., Arjovsky, M., Dumoulin, V., and Courville, A. Improved Training of Wasserstein GANs. In *Proceedings of the 31st International Conference on Neural Information Processing Systems*, NIPS'17, pp. 5769–5779, Red Hook, NY, USA, 2017. Curran Associates Inc. ISBN 9781510860964.
- Huang, J., Jiao, Y., Li, Z., Liu, S., Wang, Y., and Yang, Y. An error analysis of generative adversarial networks for learning distributions. *Journal of machine learning research*, 23(116):1–43, 2022.
- Li, Y., Swersky, K., and Zemel, R. Generative moment matching networks. In *International conference on machine learning*, pp. 1718–1727. PMLR, 2015.
- Liang, T. How well generative adversarial networks learn distributions. *Journal of Machine Learning Research*, 22 (228):1–41, 2021.
- Miyato, T., Kataoka, T., Koyama, M., and Yoshida, Y. Spectral normalization for generative adversarial networks. In *International Conference on Learning Representations*, 2018. URL https://openreview.net/forum?id=B1QRgziT-.
- Mohri, M., Rostamizadeh, A., and Talwalkar, A. Foundations of Machine Learning, second edition. Adaptive Computation and Machine Learning series. MIT Press, 2018. ISBN 9780262039406. URL https://books.google.com/books?id=V2B9DwAAQBAJ.
- Nguyen, X., Wainwright, M. J., and Jordan, M. I. Estimating divergence functionals and the likelihood ratio by convex risk minimization. *IEEE Transactions on Information Theory*, 56(11):5847–5861, 2010.
- Nowozin, S., Cseke, B., and Tomioka, R. F-GAN: Training Generative Neural Samplers Using Variational Divergence Minimization. In *Proceedings of the 30th International Conference on Neural Information Processing Systems*, NIPS'16, pp. 271–279, Red Hook, NY, USA, 2016. Curran Associates Inc. ISBN 9781510838819.
- Rockafellar, R. *Convex Analysis*. Princeton Landmarks in Mathematics and Physics. Princeton University Press, 1970. ISBN 9780691015866.
- Ruderman, A., Reid, M. D., García-García, D., and Petterson, J. Tighter variational representations of f-divergences via restriction to probability measures. In *Proceedings of the 29th International Coference on International Conference on Machine Learning*, ICML'12, pp. 1155–1162, Madison, WI, USA, 2012. Omnipress. ISBN 9781450312851.
- Song, J. and Ermon, S. Bridging the gap between f-GANs and Wasserstein GANs. In *International Conference on Machine Learning*, pp. 9078–9087. PMLR, 2020.
- Wainwright, M. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2019. ISBN 9781108498029. URL https://books.google.com/books?id=IluHDwAAQBAJ.
- Yang, Y., Li, Z., and Wang, Y. On the capacity of deep generative networks for approximating distributions. *Neural networks*, 145:144–154, 2022.