

## Self-Interacting Dark Matter, Core Collapse and the Galaxy-Galaxy Strong Lensing discrepancy

ISAQUE DUTRA <sup>1</sup>, PRIYAMVADA NATARAJAN <sup>2,1</sup> AND DANIEL GILMAN <sup>3</sup>

<sup>1</sup>*Department of Physics, Yale University, New Haven, CT 06511, USA*

<sup>2</sup>*Department of Astronomy, Yale University, New Haven, CT 06511, USA*

<sup>3</sup>*Department of Astronomy & Astrophysics, University of Chicago, Chicago, IL 60637, USA*

### ABSTRACT

Gravitational lensing by galaxy clusters has emerged as a powerful tool to probe the standard Cold Dark Matter (CDM) paradigm of structure formation in the Universe. Despite the remarkable explanatory power of CDM on large scales, tensions with observations on small scales have been reported. Recent studies find that the observational cross-section for Galaxy-Galaxy Strong Lensing (GGSL) in clusters exceeds the CDM prediction by more than an order of magnitude, and persists even after rigorous examination of various possible systematics. We investigate the impact of modifying the internal structure of cluster dark matter sub-halos on GGSL and report that altering the inner density profile, given by  $r^\gamma$ , to steeper slopes with  $\gamma > 2.5$  can alleviate the GGSL discrepancy. Deviating from the  $\gamma \sim 1.0$  cusps that CDM predicts, these steeper slopes could arise in models of self-interacting dark matter undergoing core collapse. Our results motivate additional study of sub-halo core collapse in dense cluster environments.

*Keywords:* Cosmology (343), Dark Matter(353), Galaxy dark matter halos (1880), Strong gravitational lensing (1643), Galaxy clusters (584)

### 1. INTRODUCTION

The concordance cosmological model of Cold Dark Matter (CDM) posits that dark matter behaves as a collision-less fluid, and forms gravitationally-bound halos with masses  $M \sim 10^{14}M_\odot$ , down to structures comparable to Earth mass  $M \sim 10^{-6}M_\odot$ . CDM halos form with an internal density distribution that is well fit by the Navarro-Frenk-White (NFW) profile (Navarro et al. 1997). Deviation of detected halo properties from these predictions could signal new physics, therefore many cosmic probes

seek to characterize the properties of dark matter halos across a wide range of scales and cosmic environments.

On large cosmological scales, observations agree remarkably well with CDM predictions (Spergel et al. 2007; Planck Collaboration et al. 2020). However, various tensions have emerged on sub-galactic scales, in dark matter halos below masses of  $10^{11}M_{\odot}$  (see Bullock & Boylan-Kolchin 2017, and references therein). On larger cluster-mass scales of  $\sim 10^{14}M_{\odot}$ , the ‘‘arc statistics’’ tension initially reported by (Bartelmann et al. 1998) highlighted the inability of CDM simulations to reproduce the abundance of observed lensed arcs, but several plausible resolutions to resolve this discrepancy have since emerged. Despite the variety of observables and cosmic environments associated with the small-scale and cluster-scale tensions, the common theme among them relates to an apparent failure of simulations to simultaneously reproduce the abundance and internal structure of dark matter halos predicted by CDM. Some of these tensions with CDM predictions were interpreted as a serious crisis for the theory and the underlying assumptions related to dark matter physics. In many if not most instances, however, the implications for dark matter physics became less clear upon further investigation of the implementation of baryonic feedback in galaxy formation simulations. These astrophysical processes can alter the properties of dark matter halos without requiring modifications to the fundamental nature of dark matter <sup>1</sup>.

In this paper, we review and revisit the report of an observational probe of small-scale structure, Galaxy-Galaxy Strong Lensing (GGSL), that reveals a brand-new tension with predictions of the CDM paradigm Meneghetti et al. (2020) (hereafter M20). Comparing lensing observations with the Illustris-TNG cosmological simulation suite, we independently compute the GGSL probability adopting a different methodology from M20 and reaffirm its persistence. We then explore potential explanations to alleviate this discrepancy.

The outline of our paper is as follows: in Section 2 we review the GGSL discrepancy; and outline the methodology including modeling of cluster lenses and the calculation of GGSL from lens mass models in Section 3. The properties of the observed lenses studied here are summarized in Section 4; and the computation of GGSL from the mass-matched simulated analogs from the Illustris-TNG suite is described in Section 5. Alteration of the inner density profile of cluster sub-halos explored with the Lenstronomy package is described in Section 6 and the Results of our analysis and our Conclusions & Discussion are presented in Sections 7 & 8 respectively.

## 2. THE GALAXY-GALAXY STRONG LENSING DISCREPANCY

Gravitational lensing by massive galaxy clusters offers a stringent test of the CDM paradigm of structure formation, both in terms of the number of collapsed halos – the subhalo mass function – and their density profiles (see the recent review by

<sup>1</sup> see review Popolo & Le Delliou 2017 <https://hal.archives-ouvertes.fr/hal-01324388v2/document> and references therein.

Natarajan et al. (2024)). In this work, we focus on the relatively new tension with CDM in galaxy clusters first reported in M20. M20 showed that the measured GGSL cross section of galaxy clusters exceeds the value computed from hydrodynamical CDM simulations. For a fixed total number of cluster sub-halos, the GGSL cross section depends principally on the internal structure of the sub-halos. More centrally concentrated sub-halos are more likely to become super-critical and produce multiple images of background sources. Previously, several works showed that observed clusters contain more high-circular-velocity sub-halos (i.e. sub-halos with maximum circular velocities  $V_{\text{circ}} > 100 \text{ km s}^{-1}$ ) compared to simulations (Grillo et al. 2015; Munari et al. 2016; Bonamigo et al. 2017; Meneghetti et al. 2020), although this feature was not quantified in terms of the strong-lensing cross section at that time.

Mirroring the historical progression of apparent tensions with CDM on smaller scales, several studies highlighted possible solutions to the tension reported by M20 based on astrophysics or systematics in the simulations. These include numerical resolution effects (Robertson 2021) (see also Van Den Bosch et al. (2018)), or choices related to the implementation of baryonic physics (Bahé 2021). Further investigation found that neither of these factors resolves the GGSL tension without creating further new tensions between simulations and observations (Ragagnin et al. 2022; Meneghetti et al. 2022; Tokayer et al. 2024). Resolutions proposed for the other small-scale tensions pertaining to the inner density profiles of low-mass galaxies through baryonic physics effects, in fact exacerbate the GGSL tension, which requires an enhancement, not a suppression, of the central density of cluster sub-halos. Despite the significant advances of the past two decades in understanding the interplay between baryonic physics and dark matter physics inside halos, as well as the numerical advances in the modeling of these processes in clusters, at present no clear resolution to the GGSL discrepancy exists within the framework of CDM (Tokayer et al. 2024).

The tension reported by M20 implies that the concentration of matter in the inner regions of cluster sub-halos significantly exceeds the CDM prediction. As they note, this implies either missing physics in the simulations that would increase the cluster sub-halo lensing efficiency, or a modification to the internal structure of the sub-halos that raises their strong lensing cross section. In this paper, we investigate the GGSL under the assumption that modifications to the internal structure of cluster sub-halos are required to resolve the tension.

### 3. METHODOLOGY

#### 3.1. *Modeling Cluster Lenses*

We briefly outline the standard procedures used to model observed clusters-lenses using parametric mass profiles that are apt and amenable for direct and easy comparison with simulated mass-matched cluster analogs. More details of this procedure can be found in the review by Kneib & Natarajan (2011).

- The total mass distribution of a galaxy cluster is decomposed into parametric mass profile with components to represent larger scale and smaller scale contributions (Natarajan & Kneib 1997):

$$\phi_{tot} = \sum_i \phi_i^{halo} + \sum_j \phi_j^{sub} + \phi_{\kappa,\gamma}; \quad (1)$$

where the first term represents large-scale smooth halos associated with the over-all cluster potential on scales of 100's of kpc; the second term represents smaller-scale perturbations on kpc scales that are associated with the subhalos of cluster member galaxies, and the last term is associated with a possible constant convergence or shear field, referred to as the mass-sheet degeneracy. Observed strong and weak lensing constraints including the positions, brightnesses and shapes of lensed background galaxies are used to constrain the overall mass distribution represented by Eqn. 1. It has been shown that measurements of the shear and convergence for extremely well calibrated lensing clusters with multiple families of strongly lensed images with spectroscopic redshifts such as the ones studied here, can be used to determine the mass sheet degeneracy term above (Bradač et al. 2004; Rexroth et al. 2016; Bergamini et al. 2021). The components in Eqn. 1 are determined using lens inversion algorithms, like for instance, the publicly available software package `Lenstool` that is widely used by the lensing community including by M20 and `lenstronomy`, a new package, that we use for the analysis presented here. Both the larger scale halos and subhalos are parameterized by self-similar *dual Pseudo Isothermal Elliptical mass distributions* (dPIE profiles) in `Lenstool` (Elíasdóttir et al. 2007):

$$\rho_{dPIE}(r) = \frac{\rho_0}{(1 + r^2/r_{core}^2)(1 + r^2/r_{cut}^2)} \quad (2)$$

- The enclosed projected total mass can be found with

$$M_{2D}(R) = 2\pi\Sigma_0 \frac{r_{core}r_{cut}}{r_{cut} - r_{core}} \left( \sqrt{r_{core}^2 + R^2} - r_{core} - \sqrt{r_{cut}^2 + R^2} + r_{cut} \right) \quad (3)$$

- where the projected surface mass density  $\Sigma(R)$  is given by:

$$\Sigma(R) = \Sigma_0 \frac{r_{core}r_{cut}}{r_{cut} - r_{core}} \left( \frac{1}{\sqrt{r_{core}^2 + R^2}} - \frac{1}{\sqrt{r_{cut}^2 + R^2}} \right), \quad (4)$$

- $\rho_0$  and  $\Sigma_0$  can be related to the fiducial velocity dispersion from

$$\sigma_{dPIE}^2 = \frac{4}{3}G\pi\rho_0 \frac{r_{core}^2 r_{cut}^3}{(r_{cut} - r_{core})(r_{cut} + r_{core})^2} \quad (5)$$

$$= \frac{4}{3}G\Sigma_0 \frac{r_{core}r_{cut}^2}{r_{cut}^2 - r_{core}^2} \quad (6)$$

where  $\sigma_{dPIE}$  is found from the lens model optimization obtained by reproducing observed lensed image constraints and is related to the measured central velocity dispersion of cluster galaxies via  $\sigma_{dPIE}^2 = \frac{2}{3}\sigma_0^2$ . The following substitution is done to generalize to the elliptical case,  $R \rightarrow \tilde{R}$ , profiles that are available for use within `Lenstool`:

$$\tilde{R}^2 = \frac{X^2}{(1+\epsilon)^2} + \frac{Y^2}{(1-\epsilon)^2} \quad (7)$$

where the ellipticity  $\epsilon = (A - B)/(A + B)$ , where A and B are the semi-major and semi-minor axis respectively. The ellipticity as defined in `Lenstool`, has a slightly different definition as  $\hat{\epsilon}$  that is given by:

$$\hat{\epsilon} = \frac{A^2 - B^2}{A^2 + B^2}. \quad (8)$$

The dPIE profiles in `Lenstool` are parameterized by projected sky positions  $x$  and  $y$ , fiducial velocity dispersion  $\sigma_{dPIE}$ , ellipticity  $\hat{\epsilon}$ , scale radius  $r_{\text{core}}$ , cut radius  $r_{\text{cut}}$ , and orientation  $\theta_{\hat{\epsilon}}$ . The best-fit cluster lensing model provides constraints on the total mass enclosed within  $r_{\text{cut}}$  of the larger scale halos as well as the integrated mass within  $r_{\text{cut}}$  for the subhalos. How the mass and surface mass density are distributed radially within subhalos is not a quantity that can be currently constrained with the data in hand. Furthermore, to even obtain this integrated constraint on the mass enclosed within the aperture for subhalos, we need to collapse the number of parameters required to describe galaxy-scale subhalos and the associated cluster member galaxies. To do this, typically, an explicit relation between mass and light that is empirically well motivated for cluster galaxies is assumed. The following scaling relations with luminosity  $L$  are adopted involving the characteristic subhalo parameters ( $\sigma_{dPIE}$ ,  $r_{\text{cut}}$ ) (Elíasdóttir et al. 2007, see):

$$\sigma_{dPIE} = \sigma_{dPIE}^{ref} \left( \frac{L}{L_*} \right)^\alpha \quad (9)$$

$$r_{\text{cut}} = r_{\text{cut}}^{ref} \left( \frac{L}{L_*} \right)^\beta. \quad (10)$$


Therefore, for a well constrained lens model, masses are obtained for all constituent subhalos using the above calibrated scaling relation; with these, the overall subhalo mass function can be determined for a given cluster. As noted in previous works, such as Natarajan et al. (2017), the lensing determined subhalo mass function for clusters is in very good agreement with the one computed from mass-matched simulated analogs, as shown in Fig. 6.

### 3.2. Calculation of the GGSL

In this section, we briefly define the GGSL probability and outline its calculation. Starting from the surface mass density maps of the lensing clusters obtained as noted

above, we are interested in computing the probability that a background source could be strongly lensed by foreground substructures, namely, subhalos in the cluster.

All lensing quantities of interest can be derived from the projected Newtonian potential  $\phi$  related to the projected surface mass density  $\Sigma$  as  $4\pi G\Sigma = \nabla^2\phi$ . These can be directly obtained from the derived lensing mass maps. The deflection angle  $\alpha$  is found from the partial derivative of the projected potential while computed second order derivatives characterize induced deformations: the convergence  $\kappa$  quantifies the isotropic deformation and affects the apparent size and brightness of the lensed objects, while the shear  $\gamma$  describes the anisotropic deformation affecting the stretching along the shear direction. The convergence  $\kappa$  and shear  $\gamma$  define the magnification  $\mu^{-1} = (1 - \kappa)^2 - \gamma^2 = (1 - \kappa - \gamma)(1 - \kappa + \gamma)$ , which is infinite when  $\kappa \pm \gamma = 1$ . The set of points of infinite magnification in the image plane define two closed lines, the *critical lines*, with *caustics* being the corresponding lines projected onto the source plane. For a simple mass distribution, the inner critical line can be identified as the *radial* critical line, where background sources are deformed radially, and the outer one as the *tangential* critical line, where background sources are tangentially deformed into arcs and rings. Since the radial critical lines of massive cluster members are expected to be insignificant in terms of area enclosed within (see Meneghetti et al. 2020, Supplementary materials), we consider only the *tangential critical lines* where  $\kappa + \gamma = 1$ . We plot the tangential critical lines and caustics for one of the clusters studied here, MACS J1206, in Fig. 1.



**Figure 1.** The critical lines and caustics for MACS J1206: these are calculated using `lenstronomy` and are shown in the left and right panel, respectively. The primary critical line is identified in orange, while secondaries are shown in white; similarly, primary caustics are identified in blue, while secondary caustics are shown in red. Image credit: The color composite of MACS J1206, NASA, ESA, M. Postman (STScI) and the CLASH Team. We note here that the field of view (FOV) in the left panel is not the same as the FOV in the right panel. The caustics are smaller and harder to see, therefore, we present a zoomed in view in the right panel for visibility.

Critical lines can be assigned an Einstein radius  $\theta_E$  given their average enclosed area  $A$ ,  $\theta_E = \sqrt{A/\pi}$ . Following the same classification criteria adopted in Meneghetti et al. (2022), we identify critical lines with  $\theta_E > 5''$  as *primary*; those with  $0.5'' < \theta_E < 3''$  as *secondary*, and exclude the rest in our analysis. We adopt the same definitions as M20 for consistency for comparison. These selection criteria permit us to filter subhalos with masses in the range of  $\sim 10^{10-12} h^{-1} M_\odot$ . And by definition, GGSL events originate from the secondary critical lines associated with cluster subhalos in this mass range. Substructure associated with Einstein radii less than  $0.5''$  cannot be resolved by simulations, while those with Einstein radii ranging between  $3''$  and  $5''$  are associated with larger, group-scale masses and not individual galaxies (Meneghetti et al. 2022; Ragagnin et al. 2022; Meneghetti et al. 2023). The GGSL cross section  $\sigma_{\text{GGSL}}$  is defined as the total area enclosed by all secondary caustics. Since both the convergence  $\kappa$  and shear  $\gamma$  depend on the source redshift  $z$ ,  $\sigma_{\text{GGSL}} = \sigma_{\text{GGSL}}(z)$  is also a function of redshift. The probability of GGSL events  $P_{\text{GGSL}}$  is defined as the GGSL cross section normalized by the Field of View (FOV) area on the source plane  $A_s$ ,  $P_{\text{GGSL}} = \sigma_{\text{GGSL}}/A_s$ .

In summary, the caustics associated with subhalos mark regions where background galaxies will be strongly lensed and dividing the caustic area by the FOV source plane area defines the probability of GGSL events. It is this derived probability that M20 found to be higher by more than an order of magnitude for observed lensing clusters compared to their simulated counter-parts in  $\Lambda$ CDM.

Here, we adopt a new approach and utilize the recently developed open source gravitational lensing python package `lenstronomy` (Birrer & Amara 2018; Birrer et al. 2021) to perform the lensing calculations. For each cluster, we first load the list of best-fit potentials and their optimized characteristic properties obtained from `Lenstool` representing all cluster members and initialize the appropriate analytical potentials in `lenstronomy`. For a given source redshift, the deflection map and its relevant derivatives are calculated as are the maps for the convergence  $\kappa$  and shear  $\gamma$ . The contours where  $\kappa + \gamma = 1$  are found utilizing the open source package `scikit-image` (Van Der Walt et al. 2014), and by projecting them onto the source plane, the associated caustics are found. Consistent with M20, we delimit the FOV to  $200'' \times 200''$  for all the clusters studied here. Then the areas associated with secondary caustics are summed and normalized by the projected FOV area to compute the  $P_{\text{GGSL}}(z_s)$  for source redshifts  $z_s$  between 0.5 and 7.

We calculate the properties of cluster sub-halos required to reproduce the observed GGSL cross section. This work expands on the earlier analysis by Yang & Yu (2021), who showed that core collapse in Self-Interacting Dark Matter (SIDM) models raises the central density of cluster sub-halos and can hence increase the GGSL cross section. Here, we explore more comprehensively the consequence of core collapse by calculating the strong-lensing cross section with the convergence maps of simulated galaxy clusters. To examine the degree to which feedback processes can impact the computed

GGSL, we analyze a new simulation suite - the Illustris-TNG suite <sup>2</sup> - for comparison with lensing data that implements feedback differently from the DIANOGA suite used for comparison by M20. Unlike in DIANOGA, in the Illustris suite of simulations, feedback from star formation and accreting black holes is modeled very differently. Stellar feedback is driven by galactic-scale outflows and multi-modal feedback from accreting black holes that operate in a thermal quasar mode at high accretion rates and a kinetic wind mode at low accretion rates inform their sub-grid feedback physics models. These implementations of feedback are calibrated to quench gas cooling and star formation that is required to match the properties of observed galaxies. These astrophysical processes are relevant to GGSL as they potentially serve to rearrange mass from the inner regions of halos.

In what follows, we compute the GGSL cross section for sample of simulated clusters that are matched in total mass enclosed within  $R_{200}$  <sup>3</sup> to a set of observed lenses (see Table 1 for the list). We calculate the GGSL from simulated clusters using the software package *lenstronomy* <sup>4</sup> (Birrer & Amara 2018; Birrer et al. 2021), a different code than the `Lenstool` software used by M20<sup>5</sup>. Using *lenstronomy*, we modify the properties of Illustris-TNG CDM cluster sub-halo density profiles to more centrally-concentrated density profiles. In particular, we consider a density profile given by Munoz et al. (2001):

$$\rho_{\text{gNFW}}(r) = \frac{\rho_0}{(r/r_s)^\gamma (1 + (r/r_s)^2)^{(\beta-\gamma)/2}} \quad (11)$$

(hereafter the generalized NFW, or gNFW, profile) where  $\gamma$  and  $\beta$  are the logarithmic inner and outer slopes, respectively. An NFW profile in CDM corresponds to  $\gamma = 1$  and  $\beta = 3$ .<sup>6</sup>

To derive the values of  $\gamma$  and  $\beta$  in the density profile given in Eqn. 11 that are required to match the observed GGSL, we take 3 separate projections of a simulated galaxy cluster analog and replace cluster sub-halos with objects possessing gNFW density profiles. This is implemented in such a way as to conserve the total mass of the cluster, while re-distributing the mass inside the sub-halo. We then compute the GGSL probability with these steepened sub-halo profiles.

#### 4. PROPERTIES OF THE OBSERVED CLUSTER LENS SAMPLE

With the intention of independently verifying the GGSL probability discrepancy reported by M20, we select five observed cluster-lenses to study as our sample. Four of them have well-constrained mass distributions (lensing mass models) derived from the Hubble Frontier Fields (HFF) program data. These four were also selected by

<sup>2</sup> More information on the TNG suite can be found at <https://www.tng-project.org/>

<sup>3</sup> where  $R_{200}$  is the radius within which the density is 200 times the critical density of the Universe at the halo's redshift  $z$

<sup>4</sup> <https://github.com/lenstronomy/lenstronomy>

<sup>5</sup> <https://projets.lam.fr/projects/lenstool/wiki>

<sup>6</sup> The density profile in Eqn. 11 differs from an NFW profile through the  $(1 + (r/r_s)^2)$  in the denominator, whereas the NFW profile has  $(1 + (r/r_s))$ . This detail however is not relevant to our main results.



M20, and are: Abell 2744 (A2744) at  $z = 0.308$  (Merten et al. 2011; Babyk et al. 2012; Jauzac et al. 2016); Abell S1063 (AS1063) at  $z = 0.348$  (Gómez et al. 2012; Gruen et al. 2013); MACS J1206.2-0847 (M1206) at  $z = 0.439$  (Umetsu et al. 2012; Biviano et al. 2013) and MACS J0416.1-2403 (M0416) at  $z = 0.397$  (Jauzac et al. 2014).

The mass models for these four HFF clusters computed using `Lenstool` are publicly available on the MAST Archive <sup>7</sup>. Several groups using independently developed lens inversion algorithms provided magnification and mass maps for the HFF clusters. For our work here, we used the mass models provided by the CATS collaboration. Further details of the `Lenstool` reconstructions from the CATS collaboration can be found in several papers including but not limited to Jauzac et al. (2015, 2016); Mahler et al. (2018); Bergamini et al. (2019, 2021); Meneghetti et al. (2022). The fifth cluster in our sample, PSZ1 G311.65-18.48 (PG311), also included as part of the sample in Meneghetti et al. (2022), is a massive cluster that was selected differently. It was selected not for its lensing properties, but from the ESA Planck catalog of clusters that produce a detectable Sunyaev-Zeldovich (SZ) decrement signal (Planck Collaboration et al. 2014). The properties of our selected cluster sample and their properties are summarized in Table 1.

Cluster	Redshift	$M_{200}$ Estimate [ $10^{15} M_{\odot}$ ]	Subhalos
Abell 2744	0.308	$2.22^{+0.13}_{-0.12}$	246
Abell S1063	0.348	$3.97^{+1.6}_{-0.9}$	222
MACS J0416	0.397	$1.60 \pm 0.01$	191
MACS J1206	0.439	$1.4 \pm 0.2$	258
PSZ1 G311	0.444	0.85	194

**Table 1.** Summary of the observed cluster lenses in our sample.

Abell 2744 (also known as AC118, or RXCJ0014.3-3022) at  $z \sim 0.308$ , is an exceedingly complex system undergoing a complicated merging event, and has been dubbed Pandora’s Cluster. It displays a total of about 34 strongly lensed images identified in 11 multiple-image systems (Merten et al. 2011) and was chosen for the HFF by (Lotz et al. 2017) and more recently, also observed as part of the JWST UNCOVER program (Bezanson et al. 2022). The total enclosed mass within a radius of 1.3 Mpc for Abell 2744 is measured to be  $(1.8 \pm 0.4) \times 10^{15} M_{\odot}$  by Merten et al. (2011) pre-HFF; and with HFF data Jauzac et al. (2015) measure the total cluster mass within 250 kpc as  $(2.765 \pm 0.008) \times 10^{14} M_{\odot}$  and enclosed within 1.3 Mpc as  $(2.3 \pm 0.1) \times 10^{15} M_{\odot}$  using strong and weak lensing analysis in Jauzac et al. (2016). Here, we use the mass estimate of  $M_{200} = 2.22^{+0.13}_{-0.12} \times 10^{15} M_{\odot}$  within  $R_{200} = 2.38^{+0.36}_{-0.31}$  Mpc. The best-fit lensing model for this cluster comprises 258 total mass components, of which six are large scale mass concentrations; six are large scale potentials that lie outside the Hubble

<sup>7</sup> <http://archive.stsci.edu>

Advanced Camera for Surveys FOV but are in the vicinity and need to be included and the rest correspond to confirmed cluster member galaxies.

Abell S1063 at  $z \sim 0.348$ , is a particularly interesting cluster due to its unusually high X-ray luminosity that indicates that it recently underwent a major merging event (Gómez et al. 2012). Part of the HFF, it was also observed previously by HST in the Cluster Lensing And Supernova survey with Hubble (CLASH) program (Postman et al. 2012), and spectroscopically followed up by the CLASH-VLT program (Caminha et al. 2016). The estimated mass is  $M_{200} = 3.31_{-0.68}^{+0.96} \times 10^{15} M_{\odot}$  (Williamson et al. 2011; Gómez et al. 2012; Gruen et al. 2013; Diego et al. 2016). The best-fit lensing mass model for Abell S1063 comprises 227 total mass components, of which two represent large scale central mass concentrations and the rest are all associated with confirmed cluster member galaxies.

MACS J0416.1-2403 at  $z \sim 0.397$  is also a massive cluster in a complex dynamical state: it features a bimodal velocity distribution, showing two central peaks and elongated sub-clusters suggesting that they are being observed in a pre-collisional but strongly interacting phase (Balestra et al. 2016). It was also imaged by the HFF and CLASH programs, and followed up as part of the CLASH-VLT program. Its total projected mass inside a 200 kpc aperture was estimated to be  $(1.60 \pm 0.01) \times 10^{14} M_{\odot}$  by Jauzac et al. (2014), while the CLASH analysis by Umetsu et al. (2016) estimates  $M_{200} = (1.412 \pm 0.366) \times 10^{15}$ , and Balestra et al. (2016) report  $M_{200} = (1.12 \pm 0.26) \times 10^{15} M_{\odot}$  using CLASH-VLT. The best-fit lensing mass model for this cluster contains 197 total mass components, of which six are large scale central mass concentrations and the rest are associated with confirmed cluster member galaxies.

MACS J1206.2-0847 at  $z \sim 0.440$  was previously studied by Ebeling et al. (2009). The CLASH analysis by Umetsu et al. (2016) estimates the cluster mass to be  $M_{200} = (2.596 \pm 0.604) \times 10^{15} M_{\odot}$ . The obtained best-fit lensing mass model contains 264 total mass components, of which three are large scale central potentials; an additional component of external shear and the rest are associated with confirmed cluster member galaxies.

The final selected cluster in our sample is PSZ1 G311.65-18.48 (PSZ1 G311),  $z \sim 0.443$ . Unlike the previous clusters, PSZ1 G311 was not selected via strong-lensing as noted above, but detected in the ESA Planck catalog of Sunyaev-Zeldovich sources (Planck Collaboration et al. 2014). It features a remarkable tangential arc, the highly distorted image of a distant star-forming galaxy at  $z \sim 2.370$  lensed 12 times, dubbed the *Sunburst Arc* by Rivera-Thorsen et al. (2017). Sharon et al. (2022) estimate the projected cluster mass within 250 kpc to be  $2.93_{-0.02}^{+0.01} \times 10^{14} M_{\odot}$  using HST imaging, archival VLT/MUSE spectroscopy, and Chandra X-ray data, while Dahle et al. (2016) estimate its mass using Planck SZ data to be  $M_{500} = 6.6_{-1.0}^{+0.9} \times 10^{14} M_{\odot}$ . Converting to  $M_{200}$  using a typical concentration-mass relation, Meneghetti et al. (2022) estimated  $M_{200} \sim 8.5 \times 10^{14} M_{\odot}$ . The best-fit lensing mass model contains 202

total mass components, of which eight are large scale central potentials and the rest are associated with confirmed cluster member galaxies.

Using the best-fit list of optimized mass components, here we used both `Lenstool` and `lenstronomy` to calculate the GGSL probability for each of these clusters, as outlined in Section 3.2. Our results, computed independently and in a distinct way shown in Fig. 3, are in excellent agreement with the results reported in M20.

## 5. COMPUTING GGSL FROM MASS-MATCHED SIMULATED ANALOGS IN ILLUSTRIS-TNG

We calculate the GGSL probability of subhalos in cluster-scale halos in the Illustris-TNG simulations by modeling them with analytical mass density profiles. The Illustris-TNG simulations are a suite of cosmological magnetohydrodynamical simulations of galaxy formation (Nelson et al. 2019)<sup>8</sup>. In this study, we focus on TNG300, with a periodic box with side length of 302.6 Mpc, particle/cell number of  $2500^3$  for baryons and dark matter; baryonic particle mass of  $1.1 \times 10^7 M_\odot$ ; and dark matter particle mass of  $5.9 \times 10^7 M_\odot$ .

For each cluster in our observational sample, we identify three best total mass-matched halo analogs in TNG300 with comparable  $M_{200}$ 's at the same redshift as the observed sample. For the TNG300 cluster halos,  $M_{200}$  is estimated using the friends-of-friends FoF algorithm. Our simulated analogs have a mean mass  $\sim 10^{15} M_\odot$ .

We model the central halo and subhalos of the selected TNG300 analogs with the NFW profile, which has been shown to be a good fit for DM halos over a wide range of halo masses (Navarro et al. 1997). It is particularly useful since the profile can be specified with two parameters, which can be chosen to be the radius  $R_{200}$  which encloses an average density of  $200\rho_c$  and the scale radius  $R_s$ , where the profile's logarithmic slope steepens from  $-1$  to  $-3$ :

The density profile is given by:

$$\rho_{\text{NFW}}(r) = \frac{\delta_c \rho_c}{(r/r_s)(1+r/r_s)^2}, \quad (12)$$

where  $\rho_c \equiv 3H^2(z)/(8\pi G)$  is the critical density of the Universe at the halo's redshift  $z$ , and  $\delta_c$  is the characteristic over-density of the halo, defined as:

$$\delta_c \equiv \frac{200}{3} \frac{c^3}{\ln(1+c) - c/(1+c)} \quad (13)$$

$c$  is the characteristic concentration of the halo, defined as  $c \equiv R_{200}/R_s$ .

We adopt the same procedure to calculate the GGSL probability for the simulated analogs as done for observed clusters described in Section 4. For each subhalo belonging to the same TNG300 Group, we now initialize an NFW profile in `lenstronomy` with the halo's parameters  $M_{200}$  and concentration  $c$ , wherein the scale radius  $R_s$  is

<sup>8</sup> Publicly available at <https://www.tng-project.org>

found from the relationship  $R_{\max} \simeq 2.163R_s$  (Mo et al. 2010, see Sec 11.1.2), and  $R_{\max}$  is the radius at which the halo’s particles attain maximum velocity.

Furthermore, we project each simulated cluster analog and its subhalos along three independent lines-of-sight corresponding to the simulation’s three spatially orthogonal planes to average over. We assume  $M_{200} \approx M_{\text{sub}}$ , the total mass of all bound member particles. We include only subhalos with virial mass  $M_{200}$  between  $10^{10}h^{-1}M_\odot$  and  $10^{12}h^{-1}M_\odot$  as done in M20, in addition to the massive central large scale halos. The GGSL probability of the TNG300 cluster analogs is then calculated as described in Section 3.2 and the results are shown in Fig. 3. As seen, our results confirm the reported tension with CDM clearly demonstrating the order of magnitude gap.

## 6. ALTERING INNER DENSITY PROFILES TO ALLEVIATE THE GGSL DISCREPANCY

While baryonic physics can, in principle, impact the central density profiles of halos, the structure of a dark matter halo can evolve more dramatically if one relaxes the assumption of collision-less dynamics inherent to CDM. The class of theories with this attribute is commonly referred to as self-interacting dark matter (SIDM). Dark matter self-interactions were initially proposed as a mechanism to generate flat cores inside halos to explain the low central densities (relative to CDM predictions) of low-mass dwarf galaxies (Spergel & Steinhardt 2000). Soon after the initial SIDM proposal, Balberg et al. (2002) pointed out that these dark self-interactions could also result in an opposite effect, due to the ‘gravothermal catastrophe’, a phenomenon initially studied by Lynden-Bell et al. (1968) in the context of dense star clusters. The gravothermal catastrophe, or ‘core collapse’ as it is now more commonly phrased, refers to a runaway contraction of halo density profiles. Core collapse represents the ultimate fate of all halos in SIDM unless an external heat reservoir supplies the central core with sufficient energy to remain intact (e.g. Zeng et al. 2022; Slone et al. 2023). SIDM models have gained traction on small scales because the processes of core formation and eventual core collapse give rise to a diversity of density profile slopes in sub-halos that appears more consistent with the properties of low-mass galaxies (Zavala et al. 2019; Sales et al. 2022; Yang et al. 2023).

The process of core collapse is particularly relevant for lensing because core collapse raises the central density, and therefore the lensing efficiency, of a halo. Recently, Minor et al. (2021) reported the detection of dark object with a central density orders of magnitude higher than the CDM expectation, which could result from core collapse in an SIDM halo (Nadler et al. 2023). Core collapse can also produce a distinct signal in the magnifications among images of quadruply-imaged quasars (Gilman et al. 2021, 2023). These studies focus on strong lensing by isolated individual galaxy-scale lenses, unlike the case of GGSL in clusters. Cosmological simulations of SIDM on cluster scales that permit direct comparison and calculation of the GGSL probability are currently unavailable as there are multiple numerical challenges with resolution and

convergence that are yet to be surmounted (e.g. Mace et al. 2024; Ragagnin et al. 2024).

### 6.1. *Mimicking core-collapse within lenstronomy*

It is clear that more centrally concentrated subhalos likely enhance GGSL and could therefore alleviate the GGSL discrepancy. This however requires steepening the inner density profiles in subhalos beyond  $\rho \propto r^{-1}$  of the NFW profile characteristic of CDM. Cluster galaxies are observed to have a strong concentration of baryons in the inner regions, however as recently shown by Tokayer et al. (2024), the modification to the inner density profiles from these effects cannot account for the GGSL discrepancy.

Such additional steepening is expected to occur in beyond-CDM dark matter models that contain self-interactions, SIDM models. The class of SIDM models in which appreciable steepening occurs are ones in which gravothermal core collapse has commenced in the inner-most regions. In order to mimic this process of core collapse that results in a more centrally concentrated halo, we utilize the generalized NFW (gNFW) profile and alter the inner and outer slopes of the mass density profile for simulated subhalos in the simulated analogs. The larger scale central, smooth halo mass density profiles are kept unaltered and are retained with an NFW fit as they account for the well-constrained large scale lensing features in the observed clusters.

We choose the gNFW profile (Munoz et al. 2001) that is implemented in `lenstronomy` to steepen inner density profiles:

$$\rho_{\text{gNFW}}(r) = \frac{\rho_0}{(r/r_s)^\gamma (1 + (r/r_s)^2)^{(\beta-\gamma)/2}} \quad (14)$$

where  $\gamma$  and  $\beta$  are the logarithmic inner and outer slope, respectively. Compared to standard cusped models such as the NFW profile where 3D density profile  $\rho \propto (r + r_s)^{(\beta-\gamma)}$ , the gNFW profile implemented in `lenstronomy` follows  $\rho \propto (r^2 + r_s^2)^{(\beta-\gamma)/2}$  and possesses convenient analytic properties that facilitate lensing calculations (Munoz et al. 2001).

After loading all TNG300 subhalos into `lenstronomy` as described in the previous section for each observed cluster analog, we modify the mass density profile from the CDM NFW profile into a generalized NFW (gNFW) profile for the subhalo population while conserving mass within  $R_{200}$ . This `lenstronomy` profile conversion procedure is done through the open source python package `pyHalo`<sup>9</sup>. We start by calibrating the gNFW profile by setting  $\gamma = 1$  and finding the logarithmic outer slope  $\beta$  such that the caustic area of the subhalo matches the caustic area produced by an NFW profile; this we find is the case when  $\beta = 2.84$ . The outer slope is now fixed to this value and then we alter the inner slope  $\gamma$  which mimics various stages of core collapse in generic SIDM models. Next, we recalculate the GGSL probability for each simulated cluster analog by fitting a gNFW profile varying  $\gamma = 1.0; 2.0; 2.5; 2.9; 2.99$  for the entire

<sup>9</sup> Publicly available at <https://github.com/dangilman/pyHalo>

subhalo population as a whole. All the subhalos in the simulated analog cluster are assigned the same steepened inner slope. The core-collapse cases with  $\gamma = 2.5; 2.9$  are shown in Fig. 3.

The difference in the computed critical lines of a TNG300 cluster analog fitted with an NFW profile and the gNFW with  $\gamma = 2.9$  can be seen in Fig. 4. As seen, the secondary critical lines for the gNFW profiles representing the core-collapsed subhalos are significantly larger and better resemble the critical lines of observed clusters (see for example the case of MACS J1206 shown in Fig. 1). We also calculate the GGSL probability for the final stage of gravothermal core collapse to a point mass as this represents the limiting case. In `lenstronomy`, this case is initialized by specifying the Einstein radius  $\theta_E$  of the point mass with mass  $M$ ,

$$\theta_E^2 = \frac{4GM}{c^2} \frac{D_{LS}}{D_S D_L} \quad (15)$$

where  $D_{LS}$  is the angular diameter distance between the lens and source plane,  $D_L$  between the observer and the lens plane, and  $D_S$  between the observer and the source plane. The GGSL probability of the core-collapsed cases c1-SIDM and c2-SIDM and that of the point mass are plotted in Fig. 3.

## 7. RESULTS

We compare lensing observations with a new set of cosmological simulations, TNG300 from the Illustris suite and compute GGSL using a new modeling procedure, distinct from M20, to verify that the tension persists independently of simulations and modeling assumptions. We find that the GGSL discrepancy persists for cluster lenses, even with our use of a new sample of mass-matched simulated clusters from the TNG300 simulation suite and computation performed using an independent method utilizing the `lenstronomy` software package instead of `Lenstool`, as done previously by M20. Our observed cluster sample has common candidates with the M20 sample, but also includes a massive cluster PSZ1 G311.65-18.48 that has been selected not for its observed lensing properties, but rather from the Sunyaev-Zeldovich decrement it produced as detected by the Planck satellite (Planck Collaboration et al. 2014). With this new methodology and use of an entirely independently generated Illustris simulation suite for comparison, the computed GGSL still remains discrepant by an order of magnitude, as reported previously. We also quantify the impact of projection effects and the inclusion of sub-halo ellipticity on the computed GGSL for simulated sub-halos, and find no significant increase in the GGSL probability from either factor. Therefore, projection effects and the inclusion of ellipticity for cluster sub-halos cannot bridge the order-of-magnitude GGSL discrepancy.

We now investigate how modifying the central density profile of cluster sub-halos affects the GGSL cross section. We note that explicitly including the contribution of stars in the inner regions also alters CDM density profiles, but this effect results


in  $\gamma < 2.0$ , insufficient to completely alleviate the GGSL discrepancy (Tokayer et al. 2024).

The modifications we make to re-arrange the mass in the inner regions of simulated sub-halo density profiles are intended to replicate the various stages of on-going core-collapse predicted by SIDM theories. We vary the inner logarithmic slopes  $\gamma$  in the profile shown in Eqn. 1, starting initially with the CDM NFW inner slope of 1, and steepening it to 3. Simulations of SIDM sub-halos exhibit a variety of inner profile slopes that fall within this range (Nadler et al. 2023).

We find that the GGSL probability increases as the inner slope steepens, as noted by Yang & Yu (2021). The median GGSL probability depends on the inner slope  $\gamma$ : when  $\gamma = 1.5$  the GGSL probability is roughly 1.5 times higher than for CDM; for  $\gamma = 2$  it increases to roughly 2 times that of CDM; for  $\gamma = 2.5$  it is 3.5 times and for the near ultimate stage of core collapse with  $\gamma = 2.9$  it is found to increase by a factor of 8, bringing the GGSL cross section from simulations into agreement with what is found for observed cluster lenses. We draw attention to the fact that the radial distribution of cluster galaxies and their associated sub-halos is not well reproduced in all CDM simulations including the Illustris-TNG as demonstrated in Natarajan et al. (2017). Therefore, if this additional mismatch is attended to in simulations, it would result in the production of more massive sub-halos closer to the cluster center, which would in turn also increase the GGSL. More accurately reproducing observed cluster galaxy properties within CDM is an on-going challenge and project for all simulators and this involves implementing more physical models for star formation, black hole growth and feedback (Ragagnin et al. 2022).

We conclude that sub-halos possessing internal density profiles given by Eqn. 11 and inner slopes  $\gamma \sim 3$  can resolve the GGSL discrepancy. This enhancement of the GGSL probability occurs due to the large number of sub-halos that become super-critical for lensing in cluster environments. The increased GGSL probability results from the steepened profile slope and the additional boost provided by the larger scale mass components in the cluster (discussed further in the Methods section). Note that NFW halos in field environments with  $\gamma \sim 1$  are sub-critical, and cannot themselves generate multiple images. The super-critical densities on these scales in CDM come from the added contribution of baryonic matter. If one invokes a baryonic solution that increases the central density of a halo by rearranging stellar mass instead of dark matter, our results show that the required outcome must reproduce a mass density profile given by Eqn. 11 with  $\gamma > 2.5$ . However, no known mechanisms cause this level of steepening on the scales required to explain the GGSL discrepancy. The steepening of inner dark matter profiles have been explored arising from (i) accretion of dark matter onto central black holes which achieves  $\gamma \simeq 2.3$  (Gondolo & Silk 2000) on scales  $< 1$  pc; (ii) formation of dark matter spikes around Intermediate Mass Black Holes results in  $\gamma \simeq 2.25$  and (iii) the adiabatic assembly of black holes resulting in dark matter mounds does not steepen NFW profiles on scales of interest for GGSL

(Bertone et al. 2024). Examining the impact of baryons concentrated in the inner regions of dark matter halos on the inner density profiles in the EAGLE suite of cosmological simulations Schaller et al. (2015a,b) report steepening of the NFW to  $\gamma < 2.0$ , once again insufficient to produce the enhancement needed in the GGSL probability.




**Figure 2.** Changes in inner density slope: we show the changes in the density profile shape and the corresponding change in the enclosed projected mass (2D mass) of a fiducial cluster subhalo in the observed lensing cluster M1206 (that is part of our sample) and a subhalo with matching  $M_{200}$  from the simulated cluster analog. Note that steepening the inner logarithmic slope  $\gamma$  from 1 to 2.9 results in an enhancement of the central density of the sub-halo by four orders of magnitude within  $0.1 r_{200}$ .

## 8. CONCLUSIONS & DISCUSSION

Our simulations quantify the central density profiles of dark matter halos required to reproduce the observed GGSL probability of observed cluster lenses. Our results show that the galaxies in dense cluster environments have higher central densities than those produced in large-scale cosmological simulations of these structures. If the discrepancy arises entirely from systematic issues present in the simulations, it must enter at a level that increases the central densities of halos by four orders of magnitude as seen in Fig. 2. Alternatively, the discrepancy between measurements and simulations based on CDM illustrated by Fig. 3 imply that current state-of-the-art cosmological simulations do not accurately predict the internal structure of dark matter halos on scales  $10^{10} - 10^{12} M_{\odot} h^{-1}$ . This scenario implies that these simulations have missing physics associated with the galaxy formation and feedback mechanisms in clusters, or they could point to different properties of the dark matter than predicted by CDM.

The calculation of the GGSL itself does not depend on dark matter physics. However, one can interpret the implied properties of halos within the context of dark matter physics, and in particular SIDM, which predicts an enhanced central densities






**Figure 3.** The GGSL probability as a function of source redshift for observed clusters, simulated mass-matched analogs in CDM and  $c$ -SIDM. The mean GGSL probability of our sample is shown in black and its range are depicted with a gray band. The median GGSL probability for the CDM TNG-300 simulation suite is shown by a purple line; the blue and green lines show the median GGSL probability of the mimicked CDM subhalos undergoing core collapse; we show the 95% percentile with enhanced central density by the blue band; and the GGSL for the final collapsed subhalos in the limit as point masses is plotted with the red line. Pink and orange bands signify the range of GGSL probability for observed clusters and simulations, respectively from M20.

for subhalos through core collapse. To date, this process has mainly been studied in less massive halos outside of galaxy clusters.

The process of core collapse depends on both the internal structure of a dark matter halo and the amplitude of the self-interaction cross section of a halo (e.g. [Yang & Yu 2022](#)). As shown by [Kaplinghat et al. \(2016\)](#), halos with different masses probe the SIDM cross section at different relative velocities. For the cluster subhalos considered in this work, the relevant velocity scales are  $\sim 50 - 100 \text{ kms}^{-1}$ , and thus the GGSL within the context of SIDM requires a self-interaction cross section on these velocity scales high enough to drive core collapse.



**Figure 4.** CDM and c-SIDM critical lines of a  $10^{15}M_{\odot}$  halo found in TNG300 simulation. The left figure shows the critical lines of the halo modeled with CDM profiles, while the right shows the steeper  $\gamma$  profiles. The critical lines of these steepened subhalos increase in area and number as more subhalos become supercritical, while the subhalo critical lines close enough to the center merge with the primary critical line, creating visible distortions on its contours.

A natural question in this context is whether a viable SIDM cross-section exists that can drive core collapse on these scales without violating constraints from other observational probes. At velocities  $\sim 1,000 \text{ km s}^{-1}$  stringent upper limits on the cross-section amplitude  $\sigma \lesssim 1 \text{ cm}^2 \text{ g}^{-1}$  (Peter et al. 2013; Sagunski et al. 2021) preclude values high enough to cause core collapse in cluster sub-halos within the age of the Universe. However, inelastic scattering cross-sections can accelerate the onset of core collapse, and possibly result in core collapse within the age of the Universe on these scales with an interaction cross section  $\mathcal{O}(1 \text{ cm}^2 \text{ g}^{-1})$  (Essig et al. 2019; Huo et al. 2020; O’Neil et al. 2023).

Alternatively, velocity-dependent cross-sections can increase the cross-section amplitude on the velocity scales of interest, especially if a mechanism specific to cluster environments accelerates the onset of core collapse relative to galactic environments. Velocity-dependent cross sections arise naturally in SIDM models in which dark matter particle with mass  $m_{\chi}$  interacts through a light force carrier  $m_{\phi}$  (Vogelsberger et al. 2016; Tulin et al. 2013a; Colquhoun et al. 2021; Yang & Yu 2022). At low speeds ( $v \propto \frac{m_{\phi}}{m_{\chi}}$ ), these cross sections typically scale inversely with relative velocity to a power  $\alpha$  that varies from close to zero for certain models with a weak potential (i.e. those in the Born regime), to values of  $\alpha$  the range  $-1$  to  $-4$  in the semi-classical or classical scattering regimes (Tulin et al. 2013a; Colquhoun et al. 2021). In such models, an immediate consequence of an appreciable cross section on the scales of cluster sub-halos is that the cross-section amplitude on lower mass scales is enhanced

by a factor  $\left(\frac{v}{100 \text{ km s}^{-1}}\right)^{-\alpha}$ , corresponding to a factor 100-1000 increase on the scales of dwarf galaxies  $v \sim 10 - 50 \text{ km s}^{-1}$ , and by an even larger factor on the scales probed by galaxy-galaxy strong lensing (Minor et al. 2021; Gilman et al. 2021, 2023). Alternatively, a cross-section with a pronounced resonance precisely at the velocity scales relevant for cluster sub-halos (e.g. Tulin et al. 2013b; Chu et al. 2019; Gilman et al. 2023) could drive collapse in these structures without violating constraints from other scales, although one could argue that these solutions are disfavored because they require fine tuning of the particle physics model.

To understand whether SIDM presents a viable resolution to the GGSL tension requires high-resolution hydrodynamical simulations of SIDM on galaxy cluster scales with sufficient resolution to resolve the central density profiles of cluster sub-halos as they collapse. Ragagnin et al. (2024) recently simulated SIDM on cluster scales and reported a central density enhancement of  $\sim 20\%$  in cluster subhalos, but their simulations did not have sufficient resolution to study core collapse, and thus this phenomenon on cluster scales remains unexplored. However, if one assumes that the GGSL tension reported by M20 stems entirely from the internal properties of cluster sub-halos, SIDM models offer a viable solution through core collapse. Whether an SIDM solution on the scales of cluster sub-halos can evade stringent constraints from other astrophysical probes remains an open question.

## ACKNOWLEDGMENTS

The authors acknowledge useful discussions with Massimo Meneghetti, Guillaume Mahler, Mathilde Jauzac, Yarone Tokayer, Simon Birrer, Urmila Chadayammuri, and Elena Rasia. PN gratefully acknowledges funding from the Department of Energy grant DE-SC0017660. DG acknowledges support for this work provided by The Brinson Foundation through a Brinson Prize Fellowship grant.


*Software:* astropy (Collaboration et al. 2022), numpy (Harris et al. 2020) scikit-image (Van Der Walt et al. 2014), lenstronomy (Birrer & Amara 2018; Birrer et al. 2021)

## APPENDIX

### A.1 Impact of projection effects on GGSL

We investigate the impact of projection effects on GGSL from the simulated analogs and find it to be negligible. For a simulated analog in our sample, we compute the GGSL probability for each independent projection for both CDM and the steepened version (labeled as c2-SIDM), and show the results in Fig. 5. The highest computed GGSL probability at redshift  $z = 7$  differs from the lowest by less than 20% for CDM and by  $\sim 26\%$  for c2-SIDM, insufficient in and of itself to account on its own


for the reported GGSL discrepancy. Therefore, projection effects cannot alleviate the GGSL discrepancy.



**Figure 5.** Impact of projection effects: change in GGSL probability as a function of source redshift along the three independent projections. Once again, the mean GGSL probability of our sample is shown in black and its range as a gray band. The GGSL probability for each independent projection for the selected CDM TNG300 analog is shown by solid lines in blue, green and red; the dashed lines with corresponding colors show the GGSL probability of the steepened c2-SIDM.

## A.2 Impact of ellipticity on GGSL

We investigate the effect of subhalo ellipticity on our GGSL calculations and also find that it is insufficient to fully account for the GGSL discrepancy. When computing the GGSL probability from simulations as described above, we first initialize a spherical NFW profile in `lenstronomy` for each subhalo in the mass matched simulated cluster analog. We note that there are no observational constraints on the ellipticity of dark matter subhalos. Therefore, all the `Lenstool` parametric model files our observed sample typically adopt spherically averaged profiles. However, if we assume that the ellipticity of the light distribution in an



**Figure 6.** The subhalo mass function of observed clusters and their simulated CDM analogs in the TNG300 box: The solid colored lines are for the observed cluster sample, while gray lines show the subhalo mass distributions of the simulated CDM analogs.

observed cluster member galaxy is a proxy for the ellipticity of its associated dark matter halo, as was done for the case Abell 2744, we can then construct an ellipticity distribution for the subhalo population. The `Lenstool` model for Abell 2744 adopted this ellipticity distribution as a prior, and it is shown in the left panel of Fig. 7.


The ellipticity  $\hat{\epsilon}$  of a projected mass distribution with semi major axis  $A$  and semi minor axis  $B$  are defined by

$$\hat{\epsilon} = \frac{A^2 - B^2}{A^2 + B^2} \quad (1)$$

Further details can be found in (Elíasdóttir et al. 2007, Appendix A.5) and references therein.

We adopt this observationally derived ellipticity distribution for subhalos in Abell 2744 and draw uniformly from it to initialize elliptical NFW profiles in `lenstronomy` for each simulated analog and then compute the GGSL probability for the rest of our observed sample. The result can be seen in the left panel of Fig. 7.

We notice a slight increase in the GGSL probability by a factor of about 1.4 when the effects of subhalo ellipticity are included. However, the effect of ellipticity appears to be insignificant in and of itself to account for the observed GGSL discrepancy between simulations and observed cluster-lenses.



**Figure 7.** Left: Comparison of the GGSL probability for spherical and elliptical subhalos: we compute the GGSL probability for the CDM TNG300 simulated analogs as spherical NFW profiles; and overplot the calculation done with elliptical NFW profiles wherein the ellipticity is assigned to subhalos by drawing uniformly from Abell 2744 ellipticity distribution.

Therefore, we demonstrate that neither projection effects nor ellipticity can fully alleviate the GGSL discrepancy.

## REFERENCES

- Babyk, Iu., Elyiv, A., Melnyk, O., & Krivodubskij, V. N. 2012, *Kinematics and Physics of Celestial Bodies*, 28, 69, doi: [10.3103/S0884591312020031](https://doi.org/10.3103/S0884591312020031)
- Bahé, Y. M. 2021, *Monthly Notices of the Royal Astronomical Society*, 505, 1458, doi: [10.1093/mnras/stab1392](https://doi.org/10.1093/mnras/stab1392)
- Balberg, S., Shapiro, S. L., & Inagaki, S. 2002, *The Astrophysical Journal*, 568, 475, doi: [10.1086/339038](https://doi.org/10.1086/339038)
- Balestra, I., Mercurio, A., Sartoris, B., et al. 2016, *The Astrophysical Journal Supplement Series*, 224, 33, doi: [10.3847/0067-0049/224/2/33](https://doi.org/10.3847/0067-0049/224/2/33)
- Bartelmann, M., Huss, A., Colberg, J. M., Jenkins, A., & Pearce, F. R. 1998, *Astronomy and Astrophysics*, 330, 1, doi: [10.48550/arXiv.astro-ph/9707167](https://doi.org/10.48550/arXiv.astro-ph/9707167)
- Bergamini, P., Rosati, P., Mercurio, A., et al. 2019, *Astronomy & Astrophysics*, 631, A130, doi: [10.1051/0004-6361/201935974](https://doi.org/10.1051/0004-6361/201935974)
- Bergamini, P., Rosati, P., Vanzella, E., et al. 2021, *Astronomy & Astrophysics*, 645, A140, doi: [10.1051/0004-6361/202039564](https://doi.org/10.1051/0004-6361/202039564)
- Bertone, G., Wierda, A. R. A. C., Gaggero, D., et al. 2024, *Dark Matter Mounds: Towards a Realistic Description of Dark Matter Overdensities around Black Holes*, arXiv. <http://ascl.net/2404.08731>

- Bezanson, R., Labbe, I., Whitaker, K. E., et al. 2022, The JWST UNCOVER Treasury Survey: Ultradeep NIRSpec and NIRCам Observations before the Epoch of Reionization, arXiv. <http://ascl.net/2212.04026>
- Birrer, S., & Amara, A. 2018, Physics of the Dark Universe, 22, 189, doi: [10.1016/j.dark.2018.11.002](https://doi.org/10.1016/j.dark.2018.11.002)
- Birrer, S., Shajib, A., Gilman, D., et al. 2021, Journal of Open Source Software, 6, 3283, doi: [10.21105/joss.03283](https://doi.org/10.21105/joss.03283)
- Biviano, A., Rosati, P., Balestra, I., et al. 2013, Astronomy & Astrophysics, 558, A1, doi: [10.1051/0004-6361/201321955](https://doi.org/10.1051/0004-6361/201321955)
- Bonamigo, M., Grillo, C., Ettori, S., et al. 2017, The Astrophysical Journal, 842, 132, doi: [10.3847/1538-4357/aa75cc](https://doi.org/10.3847/1538-4357/aa75cc)
- Bradač, M., Lombardi, M., & Schneider, P. 2004, Astronomy & Astrophysics, 424, 13, doi: [10.1051/0004-6361:20035744](https://doi.org/10.1051/0004-6361:20035744)
- Bullock, J. S., & Boylan-Kolchin, M. 2017, Annual Review of Astronomy and Astrophysics, 55, 343, doi: [10.1146/annurev-astro-091916-055313](https://doi.org/10.1146/annurev-astro-091916-055313)
- Caminha, G. B., Grillo, C., Rosati, P., et al. 2016, Astronomy & Astrophysics, 587, A80, doi: [10.1051/0004-6361/201527670](https://doi.org/10.1051/0004-6361/201527670)
- Chu, X., Garcia-Cely, C., & Murayama, H. 2019, Physical Review Letters, 122, 071103, doi: [10.1103/PhysRevLett.122.071103](https://doi.org/10.1103/PhysRevLett.122.071103)
- Collaboration, T. A., Price-Whelan, A. M., Lim, P. L., et al. 2022, The Astrophysical Journal, 935, 167, doi: [10.3847/1538-4357/ac7c74](https://doi.org/10.3847/1538-4357/ac7c74)
- Colquhoun, B., Heeba, S., Kahlhoefer, F., Sagunski, L., & Tulin, S. 2021, Physical Review D, 103, 035006, doi: [10.1103/PhysRevD.103.035006](https://doi.org/10.1103/PhysRevD.103.035006)
- Dahle, H., Aghanim, N., Guennou, L., et al. 2016, Astronomy & Astrophysics, 590, L4, doi: [10.1051/0004-6361/201628297](https://doi.org/10.1051/0004-6361/201628297)
- Diego, J. M., Broadhurst, T., Wong, J., et al. 2016, Monthly Notices of the Royal Astronomical Society, 459, 3447, doi: [10.1093/mnras/stw865](https://doi.org/10.1093/mnras/stw865)
- Ebeling, H., Ma, C. J., Kneib, J.-P., et al. 2009, Monthly Notices of the Royal Astronomical Society, 395, 1213, doi: [10.1111/j.1365-2966.2009.14502.x](https://doi.org/10.1111/j.1365-2966.2009.14502.x)
- Elíasdóttir, Á., Limousin, M., Richard, J., et al. 2007, Where Is the Matter in the Merging Cluster Abell 2218?, arXiv. <http://ascl.net/0710.5636>
- Essig, R., McDermott, S. D., Yu, H.-B., & Zhong, Y.-M. 2019, Physical Review Letters, 123, 121102, doi: [10.1103/PhysRevLett.123.121102](https://doi.org/10.1103/PhysRevLett.123.121102)
- Gilman, D., Bovy, J., Treu, T., et al. 2021, Monthly Notices of the Royal Astronomical Society, 507, 2432, doi: [10.1093/mnras/stab2335](https://doi.org/10.1093/mnras/stab2335)
- Gilman, D., Zhong, Y.-M., & Bovy, J. 2023, Physical Review D, 107, 103008, doi: [10.1103/PhysRevD.107.103008](https://doi.org/10.1103/PhysRevD.107.103008)
- Gómez, P. L., Valkonen, L. E., Romer, A. K., et al. 2012, The Astronomical Journal, 144, 79, doi: [10.1088/0004-6256/144/3/79](https://doi.org/10.1088/0004-6256/144/3/79)
- Gondolo, P., & Silk, J. 2000, Nuclear Physics B - Proceedings Supplements, 87, 87, doi: [10.1016/S0920-5632\(00\)00640-X](https://doi.org/10.1016/S0920-5632(00)00640-X)
- Grillo, C., Suyu, S. H., Rosati, P., et al. 2015, The Astrophysical Journal, 800, 38, doi: [10.1088/0004-637X/800/1/38](https://doi.org/10.1088/0004-637X/800/1/38)
- Gruen, D., Brimiouille, F., Seitz, S., et al. 2013, Monthly Notices of the Royal Astronomical Society, 432, 1455, doi: [10.1093/mnras/stt566](https://doi.org/10.1093/mnras/stt566)
- Harris, C. R., Millman, K. J., Van Der Walt, S. J., et al. 2020, Nature, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Huo, R., Yu, H.-B., & Zhong, Y.-M. 2020, Journal of Cosmology and Astroparticle Physics, 2020, 051, doi: [10.1088/1475-7516/2020/06/051](https://doi.org/10.1088/1475-7516/2020/06/051)
- Jauzac, M., Clément, B., Limousin, M., et al. 2014, Monthly Notices of the Royal Astronomical Society, 443, 1549, doi: [10.1093/mnras/stu1355](https://doi.org/10.1093/mnras/stu1355)
- Jauzac, M., Richard, J., Jullo, E., et al. 2015, Monthly Notices of the Royal Astronomical Society, 452, 1437, doi: [10.1093/mnras/stv1402](https://doi.org/10.1093/mnras/stv1402)

- Jauzac, M., Eckert, D., Schwinn, J., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 463, 3876, doi: [10.1093/mnras/stw2251](https://doi.org/10.1093/mnras/stw2251)
- Kaplinghat, M., Tulin, S., & Yu, H.-B. 2016, *Physical Review Letters*, 116, 041302, doi: [10.1103/PhysRevLett.116.041302](https://doi.org/10.1103/PhysRevLett.116.041302)
- Kneib, J.-P., & Natarajan, P. 2011, *The Astronomy and Astrophysics Review*, 19, 47, doi: [10.1007/s00159-011-0047-3](https://doi.org/10.1007/s00159-011-0047-3)
- Lotz, J. M., Koekemoer, A., Coe, D., et al. 2017, *The Astrophysical Journal*, 837, 97, doi: [10.3847/1538-4357/837/1/97](https://doi.org/10.3847/1538-4357/837/1/97)
- Lynden-Bell, D., Wood, R., & Royal, A. 1968, *Monthly Notices of the Royal Astronomical Society*, 138, 495, doi: [10.1093/mnras/138.4.495](https://doi.org/10.1093/mnras/138.4.495)
- Mace, C., Zeng, Z. C., Peter, A. H. G., et al. 2024, *Convergence Tests of Self-Interacting Dark Matter Simulations*, arXiv. <http://ascl.net/2402.01604>
- Mahler, G., Richard, J., Clément, B., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 473, 663, doi: [10.1093/mnras/stx1971](https://doi.org/10.1093/mnras/stx1971)
- Meneghetti, M., Davoli, G., Bergamini, P., et al. 2020, *Science*, 369, 1347, doi: [10.1126/science.aax5164](https://doi.org/10.1126/science.aax5164)
- Meneghetti, M., Ragagnin, A., Borgani, S., et al. 2022, *Astronomy & Astrophysics*, 668, A188, doi: [10.1051/0004-6361/202243779](https://doi.org/10.1051/0004-6361/202243779)
- Meneghetti, M., Cui, W., Rasia, E., et al. 2023, *Astronomy & Astrophysics*, 678, L2, doi: [10.1051/0004-6361/202346975](https://doi.org/10.1051/0004-6361/202346975)
- Merten, J., Coe, D., Dupke, R., et al. 2011, *Monthly Notices of the Royal Astronomical Society*, 417, 333, doi: [10.1111/j.1365-2966.2011.19266.x](https://doi.org/10.1111/j.1365-2966.2011.19266.x)
- Minor, Q., Gad-Nasr, S., Kaplinghat, M., & Vegetti, S. 2021, *Monthly Notices of the Royal Astronomical Society*, 507, 1662, doi: [10.1093/mnras/stab2247](https://doi.org/10.1093/mnras/stab2247)
- Mo, H., van den Bosch, F. C., & White, S. 2010, *Galaxy Formation and Evolution* (Cambridge University Press)
- Munari, E., Grillo, C., Lucia, G. D., et al. 2016, *The Astrophysical Journal Letters*, 827, L5, doi: [10.3847/2041-8205/827/1/L5](https://doi.org/10.3847/2041-8205/827/1/L5)
- Munoz, J. A., Kochanek, C. S., & Keeton, C. R. 2001, *The Astrophysical Journal*, 558, 657, doi: [10.1086/322314](https://doi.org/10.1086/322314)
- Nadler, E. O., Yang, D., & Yu, H.-B. 2023, *The Astrophysical Journal Letters*, 958, L39, doi: [10.3847/2041-8213/ad0e09](https://doi.org/10.3847/2041-8213/ad0e09)
- Natarajan, P., & Kneib, J.-P. 1997, *Monthly Notices of the Royal Astronomical Society*, 287, 833, doi: [10.1093/mnras/287.4.833](https://doi.org/10.1093/mnras/287.4.833)
- Natarajan, P., Williams, L. L. R., Bradač, M., et al. 2024, *Space Science Reviews*, 220, 19, doi: [10.1007/s11214-024-01051-8](https://doi.org/10.1007/s11214-024-01051-8)
- Natarajan, P., Chadayammuri, U., Jauzac, M., et al. 2017, *Monthly Notices of the Royal Astronomical Society*, 468, 1962, doi: [10.1093/mnras/stw3385](https://doi.org/10.1093/mnras/stw3385)
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *The Astrophysical Journal*, 490, 493, doi: [10.1086/304888](https://doi.org/10.1086/304888)
- Nelson, D., Springel, V., Pillepich, A., et al. 2019, *Computational Astrophysics and Cosmology*, 6, 2, doi: [10.1186/s40668-019-0028-x](https://doi.org/10.1186/s40668-019-0028-x)
- O'Neil, S., Vogelsberger, M., Heeba, S., et al. 2023, *Monthly Notices of the Royal Astronomical Society*, 524, 288, doi: [10.1093/mnras/stad1850](https://doi.org/10.1093/mnras/stad1850)
- Peter, A. H. G., Rocha, M., Bullock, J. S., & Kaplinghat, M. 2013, *Monthly Notices of the Royal Astronomical Society*, 430, 105, doi: [10.1093/mnras/sts535](https://doi.org/10.1093/mnras/sts535)
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2014, *Astronomy & Astrophysics*, 571, A29, doi: [10.1051/0004-6361/201321523](https://doi.org/10.1051/0004-6361/201321523)
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, *Astronomy & Astrophysics*, 641, A6, doi: [10.1051/0004-6361/201833910](https://doi.org/10.1051/0004-6361/201833910)



- Postman, M., Coe, D., Benítez, N., et al. 2012, *The Astrophysical Journal Supplement Series*, 199, 25, doi: [10.1088/0067-0049/199/2/25](https://doi.org/10.1088/0067-0049/199/2/25)
- Ragagnin, A., Meneghetti, M., Calura, F., et al. 2024, *Dianoga SIDM: Galaxy Cluster Self-Interacting Dark Matter Simulations*, arXiv. <http://ascl.net/2404.01383>
- Ragagnin, A., Meneghetti, M., Bassini, L., et al. 2022, *Astronomy & Astrophysics*, 665, A16, doi: [10.1051/0004-6361/202243651](https://doi.org/10.1051/0004-6361/202243651)
- Rexroth, M., Natarajan, P., & Kneib, J.-P. 2016, *Monthly Notices of the Royal Astronomical Society*, 460, 2505, doi: [10.1093/mnras/stw1017](https://doi.org/10.1093/mnras/stw1017)
- Rivera-Thorsen, T. E., Dahle, H., Gronke, M., et al. 2017, *Astronomy & Astrophysics*, 608, L4, doi: [10.1051/0004-6361/201732173](https://doi.org/10.1051/0004-6361/201732173)
- Robertson, A. 2021, *Monthly Notices of the Royal Astronomical Society: Letters*, 504, L7, doi: [10.1093/mnrasl/slab028](https://doi.org/10.1093/mnrasl/slab028)
- Sagunski, L., Gad-Nasr, S., Colquhoun, B., Robertson, A., & Tulin, S. 2021, *Journal of Cosmology and Astroparticle Physics*, 2021, 024, doi: [10.1088/1475-7516/2021/01/024](https://doi.org/10.1088/1475-7516/2021/01/024)
- Sales, L. V., Wetzel, A., & Fattahi, A. 2022, *Nature Astronomy*, 6, 897, doi: [10.1038/s41550-022-01689-w](https://doi.org/10.1038/s41550-022-01689-w)
- Schaller, M., Frenk, C. S., Bower, R. G., et al. 2015a, *Monthly Notices of the Royal Astronomical Society*, 451, 1247, doi: [10.1093/mnras/stv1067](https://doi.org/10.1093/mnras/stv1067)
- . 2015b, *Monthly Notices of the Royal Astronomical Society*, 452, 343, doi: [10.1093/mnras/stv1341](https://doi.org/10.1093/mnras/stv1341)
- Sharon, K., Mahler, G., Rivera-Thorsen, T. E., et al. 2022, *The Astrophysical Journal*, 941, 203, doi: [10.3847/1538-4357/ac927a](https://doi.org/10.3847/1538-4357/ac927a)
- Slone, O., Jiang, F., Lisanti, M., & Kaplinghat, M. 2023, *Physical Review D*, 107, 043014, doi: [10.1103/PhysRevD.107.043014](https://doi.org/10.1103/PhysRevD.107.043014)
- Spergel, D. N., & Steinhardt, P. J. 2000, *Physical Review Letters*, 84, 3760, doi: [10.1103/PhysRevLett.84.3760](https://doi.org/10.1103/PhysRevLett.84.3760)
- Spergel, D. N., Bean, R., Dore, O., et al. 2007, *The Astrophysical Journal Supplement Series*, 170, 377, doi: [10.1086/513700](https://doi.org/10.1086/513700)
- Tokayer, Y. M., Dutra, I., Natarajan, P., et al. 2024, *The Galaxy-Galaxy Strong Lensing Cross Section and the Internal Distribution of Matter in  $\Lambda$ CDM Substructure*, arXiv. <http://ascl.net/2404.16951>
- Tulin, S., Yu, H.-B., & Zurek, K. M. 2013a, *Physical Review Letters*, 110, 111301, doi: [10.1103/PhysRevLett.110.111301](https://doi.org/10.1103/PhysRevLett.110.111301)
- . 2013b, *Physical Review D*, 87, 115007, doi: [10.1103/PhysRevD.87.115007](https://doi.org/10.1103/PhysRevD.87.115007)
- Umetsu, K., Zitrin, A., Gruen, D., et al. 2016, *The Astrophysical Journal*, 821, 116, doi: [10.3847/0004-637X/821/2/116](https://doi.org/10.3847/0004-637X/821/2/116)
- Umetsu, K., Medezinski, E., Nonino, M., et al. 2012, *The Astrophysical Journal*, 755, 56, doi: [10.1088/0004-637X/755/1/56](https://doi.org/10.1088/0004-637X/755/1/56)
- Van Den Bosch, F. C., Ogiya, G., Hahn, O., & Burkert, A. 2018, *Monthly Notices of the Royal Astronomical Society*, 474, 3043, doi: [10.1093/mnras/stx2956](https://doi.org/10.1093/mnras/stx2956)
- Van Der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., et al. 2014, *PeerJ*, 2, e453, doi: [10.7717/peerj.453](https://doi.org/10.7717/peerj.453)
- Vogelsberger, M., Zavala, J., Cyr-Racine, F.-Y., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 460, 1399, doi: [10.1093/mnras/stw1076](https://doi.org/10.1093/mnras/stw1076)
- Williamson, R., Benson, B. A., High, F. W., et al. 2011, *The Astrophysical Journal*, 738, 139, doi: [10.1088/0004-637X/738/2/139](https://doi.org/10.1088/0004-637X/738/2/139)
- Yang, D., & Yu, H.-B. 2021, *Physical Review D*, 104, 103031, doi: [10.1103/PhysRevD.104.103031](https://doi.org/10.1103/PhysRevD.104.103031)
- . 2022, *Journal of Cosmology and Astroparticle Physics*, 2022, 077, doi: [10.1088/1475-7516/2022/09/077](https://doi.org/10.1088/1475-7516/2022/09/077)

Yang, S., Du, X., Zeng, Z. C., et al. 2023,

The Astrophysical Journal, 946, 47,

doi: [10.3847/1538-4357/acbd49](https://doi.org/10.3847/1538-4357/acbd49)

Zavala, J., Lovell, M. R., Vogelsberger,  
M., & Burger, J. D. 2019, Physical  
Review D, 100, 063007,

doi: [10.1103/PhysRevD.100.063007](https://doi.org/10.1103/PhysRevD.100.063007)

Zeng, Z. C., Peter, A. H. G., Du, X., et al.

2022, Monthly Notices of the Royal  
Astronomical Society, 513, 4845,

doi: [10.1093/mnras/stac1094](https://doi.org/10.1093/mnras/stac1094)