Eigenvalue inequalities and three-term asymptotic formulas of the heat traces for the Lamé operator and Stokes operator

Genqian Liu^{a,*}

^aSchool of Mathematics and Statistics, Beijing Institute of Technology, Beijing 100081, P. R. China

Abstract

This paper is devoted to establish the most essential connections of the eigenvalue problems for the Laplace operator, Lamé operator, Stokes operator, buckling operator and clamped plate operator. We show that the *k*-th Stokes (respectively, Laplace) eigenvalue is the limit of the *k*-th Lamé eigenvalue for the Dirichlet or traction boundary condition as the Lamé coefficient λ tends to $+\infty$ (respectively, to $-\mu$). Furthermore, we establish the eigenvalue inequalities and three-term asymptotic formulas of the heat traces for the Laplace operator, the Lamé operator, the Stokes operator and buckling operator with the Dirichlet and traction boundary conditions.

Keywords: Laplace-type operator; Lamé operator; Stokes operator; Eigenvalues; Spectral asymptotics; Riemannian manifold

Mathematics Subject Classification (2020): 53C21, 58J50, 58C40, 35P20

1. Introduction

Revealing the relationship between the eigenvalues of the most basic physical operators, establishing the heat trace asymptotics for partial differential operators have been the subject of extensive research for over a century. It has attracted the attention of many outstanding mathematicians and physicists. Beyond the wonderful limit laws and beautiful asymptotic formulas that are intimately related to the geometric properties of the domain and its boundary, a sustaining force has been its important role in mathematics, mechanics and theoretical physics (see, for example, [AsB], [BG], [Cha1], [Cha2], [CLN], [CH], [ES], [BG], [Gil2], [Gru], [Ho4], [Iv], [Kac], [Lap], [MS-67], [Pa1], [Ple1], [Ple2], [Po], [Sar], [ShY], [Ste], [Ta2], [Wei], [Wey1], [Wey2], etc).

Let (Ω, g) be a smooth compact Riemannian manifold of dimension $n \ge 2$ with smooth boundary $\partial\Omega$. The Lamé operator $L_{\lambda,\mu}$ of linear elasticity acting on vector fields $\mathbf{u} = (u^1, \dots, u^n)$ can be written as (see [Liu-19], [Liu-23] and [Liu2])

$$L_{\lambda,\mu}\mathbf{u} = 2\mu \operatorname{Def}^* \operatorname{Def} \mathbf{u} - \lambda \operatorname{grad} \operatorname{div} \mathbf{u} \quad \text{in } \Omega, \tag{1.1}$$

*Corresponding author.

Email address: liuswgq@163.com (Genqian Liu)

where $\text{Def } \mathbf{u} = \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T)$ (see, p. 464 of [Ta1]), $(\nabla \mathbf{u})^T$ is the transpose of $\nabla \mathbf{u}$, Def^* is the adjoint of deformation operator Def (see, §2 of [Liu-19], or Exercise 16 on p 153 of [Ta1]), and div and grad are the usual divergence and gradient operators. Here λ and μ are real constants known as Lamé parameters, assumed to satisfy the conditions

$$\mu > 0, \ \lambda + 2\mu > 0,$$
 (1.2)

which guarantee strong ellipticity of $L_{\lambda,\mu}$, see, e.g., [ADN2], [CiMa], [Hah], [MaHu], [McL], [MHNZ], [LiQin] or [Liu-39]. Equivalently, $L_{\lambda,\mu}$ can be written as

$$L_{\lambda,\mu}\mathbf{u} = \mu \left(\nabla^* \nabla \mathbf{u} - \operatorname{Ric}(\mathbf{u})\right) - (\lambda + \mu) \operatorname{grad} \operatorname{div} \mathbf{u}, \tag{1.3}$$

where $(\nabla^* \nabla \mathbf{u})^j := -\nabla_k \nabla^k u^j$ is the Bochner Laplacian (see (2.11) of [Liu2]), $\nabla_k u^j = u^j_{;k} := \frac{\partial u^j}{\partial x_k} + \Gamma^j_{kl} u^l$ are the components of the covariant derivative of the vector field \mathbf{u} , $\nabla^k = g^{kl} \nabla_l$, and

$$\operatorname{Ric}\left(\mathbf{u}\right) = \left(R_{l}^{1}u^{l}, R_{l}^{2}u^{l}, \cdots, R_{l}^{n}u^{l}\right)$$
(1.4)

denotes the action of Ricci tensor $\mathbf{R}_l^{\ j} := \mathbf{R}_{lk}^{k\ j}$ on **u**. Here and further on we adopt the Einstein summation convention over repeated indices. More precisely, $L_{\lambda,\mu}$ can be expressed as

$$(L_{\lambda,\mu}\mathbf{u})^{j} := -\mu \left(\nabla_{k} \nabla^{k} u^{j} + \operatorname{Ric}_{k}^{j} u^{k} \right) - (\lambda + \mu) \nabla^{j} \nabla_{k} u^{k}, \quad j = 1, \cdots, n.$$
(1.5)

For the Lamé operator $L_{\lambda,\mu}$, the Dirichlet boundary condition is $\mathbf{u}|_{\partial\Omega} = 0$, and the traction (i.e., free or Neumann) boundary condition is $\mathcal{T}_{\lambda,\mu}\mathbf{u}|_{\partial\Omega} = 0$, where $\mathcal{T}_{\lambda,\mu}$ is defined by

$$\mathcal{T}_{\lambda,\mu}\mathbf{u} := -2\mu \left(\text{Def }\mathbf{u}\right)^{\#} \boldsymbol{\nu} - \lambda \left(\text{div }\mathbf{u}\right)\boldsymbol{\nu} \quad \text{on } \partial\Omega, \tag{1.6}$$

where # is the sharp operator (for a tensor) by raising index, and $\nu = (\nu^1, \dots, \nu^n)$ is the exterior unit normal vector to the boundary $\partial \Omega$. Clearly, (1.6) is just

$$\left(\mathcal{T}_{\lambda,\mu}\mathbf{u}\right)^{j} = -\lambda v^{j} \nabla_{k} u^{k} - \mu \left(v^{k} \nabla_{k} u^{j} + v_{k} \nabla^{j} u^{k}\right) \quad \text{on} \quad \partial\Omega, \quad j = 1, \cdots, n.$$
(1.7)

In physics, the elastic wave equations $\frac{\partial^2 \mathbf{u}}{\partial t^2} = -L_{\lambda,\mu}\mathbf{u}$ describe the propagations of waves in an isotropic homogeneous elastic medium, and the constants $\sqrt{\lambda + 2\mu}$ and $\sqrt{\mu}$ are the velocities of longitudinal and transverse waves, respectively.

Set

$$\mathcal{V} = \{ \mathbf{u} \in C_0^{\infty}(\Omega, T\Omega) \, \big| \, \operatorname{div} \mathbf{u} = 0 \}.$$
(1.8)

Then

$$J^{k} = \text{closure of } \mathcal{V} \text{ in } H^{k}(\Omega, T\Omega), \quad k = 0, 1.$$
(1.9)

Let *P* be the orthogonal projection of $L^2(\Omega, T\Omega)$ onto J^0 . The operator $S_{\mu}\mathbf{u} := -\mu P\Delta_g \mathbf{u}$ with domain $\mathcal{D}(S_{\mu}) = {\mathbf{u} \in J^0 \cap H^2(\Omega, T\Omega)}$ is called the Stokes operator, where Δ_g is the Beltrami-Laplace operator and $\mu > 0$ is the viscosity (constant) coefficient. The Stokes operator S_{μ} can equivalently be written as (see, Theorem 2.1 of [Liu4]):

$$S_{\mu}\mathbf{u} := -\operatorname{div}\left(\sigma_{\mu}(\mathbf{u}, p)^{\#}, \quad \operatorname{div}\mathbf{u} = 0,\right.$$
(1.10)

where $\mathbf{u} = (u^1, \dots, u^n)$ is the velocity vector field, $\sigma_{\mu}(\mathbf{u}, p)^{\#} = 2\mu (\text{Def}(\mathbf{u}))^{\#} - p\mathbf{I}_n$ is a tensor of field of type (1, 1), and p is the pressure. Precisely, the Stokes operator S_{μ} can be represented as (see [Liu4])

$$S_{\mu}\mathbf{u} = \mu \left(\nabla^* \nabla \mathbf{u} - \operatorname{Ric}\left(\mathbf{u}\right)\right) + \nabla_g p \quad \text{with} \quad \operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega, \tag{1.11}$$

where $-\nabla^* \nabla \mathbf{u}$ and Ric (**u**) are defined as before. In physics, a fluid flow obeying the stationary Stokes equations

$$\begin{cases} -\operatorname{div} \left(\sigma_{\mu}(\mathbf{u}, p)\right)^{\sharp} = 0 & \text{in } \Omega, \\ \operatorname{div} \mathbf{u} = 0 & \text{in } \Omega, \end{cases}$$
(1.12)

is called the Stokes flow (i.e., creeping flow). For the Stokes operator, the Dirichlet boundary condition is $\mathbf{u}\Big|_{\partial\Omega} = 0$ (i.e., $\mathcal{D}(S_{\mu}) = J^0 \cap H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$. Note that in this case, $J^1 = \{\mathbf{u} \in H_0^1(\Omega) | \text{div } \mathbf{u} = 0 \text{ in } \Omega\}$, see Lemma 5.5 on p. 575 of [Ta3]), and the Cauchy force boundary condition is $(\sigma_{\mu}(\mathbf{u}, p))^{\#} v\Big|_{\partial\Omega} = 0$.

We consider the following classical eigenvalue problems:

$$(DL) \qquad \begin{cases} L_{\lambda,\mu} \mathbf{u} = \tau^{(D)} \mathbf{u} & \text{in } \Omega, \\ \mathbf{u} = 0 & \text{on } \partial\Omega; \end{cases}$$
(1.13)

(*TL*)
$$\begin{cases} L_{\lambda,\mu} \mathbf{u} = \tau^{(\mathrm{T})} \mathbf{u} & \text{in } \Omega, \\ \mathcal{T}_{\lambda,\mu} \mathbf{u} = 0 & \text{on } \partial\Omega; \end{cases}$$
(1.14)

$$(DS) \qquad \begin{cases} S_{\mu} \mathbf{v} = \varsigma^{(D)} \mathbf{v} & \text{in } \Omega, \\ \text{div } \mathbf{v} = 0 & \text{in } \Omega, \\ \mathbf{v} = 0 & \text{on } \partial\Omega; \end{cases}$$
(1.15)

(CS)
$$\begin{cases} S_{\mu}\mathbf{v} = \varsigma^{(C)}\mathbf{v} & \text{in } \Omega, \\ \operatorname{div} \mathbf{v} = 0 & \operatorname{in} \Omega, \\ \left(\sigma_{\mu}(\mathbf{v}, p)\right)^{\#} \mathbf{v} = 0 & \text{on } \partial\Omega; \end{cases}$$
(1.16)

$$(DB) \qquad \begin{cases} \mu(\nabla^* \nabla \mathbf{w} - \operatorname{Ric}(\mathbf{w})) = \theta^{(D)} \mathbf{w} & \text{in } \Omega, \\ \mathbf{w} = 0 & \text{on } \partial\Omega; \end{cases}$$
(1.17)

(*TB*)
$$\begin{cases} \mu(\nabla^* \nabla \mathbf{w} - \operatorname{Ric}(\mathbf{w})) = \theta^{(T)} \mathbf{w} & \text{in } \Omega, \\ \mathcal{T}_{-\mu,\mu} \mathbf{w} = 0 & \text{on } \partial\Omega; \end{cases}$$
(1.18)

(1.13) (respectively, (1.14)) is the Lamé eigenvalue problem with Dirichlet (respectively, traction) boundary condition; (1.15) (respectively, (1.16)) is the Stokes eigenvalue problem with Dirichlet (respectively, Cauchy force) boundary condition; (1.17) (respectively, (1.18)) is the Laplace eigenvalue problem with Dirichlet (respectively, traction) boundary condition. In each of these cases, the spectrum is discrete and we arrange the eigenvalues in non-decreasing order (repeated according to multiplicity)

$$\begin{aligned} 0 &< \tau_1^{(\mathrm{D})} \leq \tau_2^{(\mathrm{D})} \leq \cdots \leq \tau_k^{(\mathrm{D})} \leq \cdots ; \\ 0 &\leq \tau_1^{(\mathrm{T})} \leq \tau_2^{(\mathrm{T})} \leq \cdots \leq \tau_k^{(\mathrm{T})} \leq \cdots ; \\ 0 &< \varsigma_1^{(\mathrm{D})} \leq \varsigma_2^{(\mathrm{D})} \leq \cdots \leq \varsigma_k^{(\mathrm{D})} \leq \cdots ; \\ 0 &\leq \varsigma_1^{(\mathrm{C})} \leq \varsigma_2^{(\mathrm{C})} \leq \cdots \leq \varsigma_k^{(\mathrm{C})} \leq \cdots ; \\ 0 &< \theta_1^{(\mathrm{D})} \leq \theta_2^{(\mathrm{D})} \leq \cdots \leq \theta_k^{(\mathrm{D})} \leq \cdots ; \\ 0 &\leq \theta_1^{(\mathrm{T})} \leq \theta_2^{(\mathrm{T})} \leq \cdots \leq \theta_k^{(\mathrm{T})} \leq \cdots . \end{aligned}$$

The corresponding eigenfunctions are expressed as $\mathbf{u}_1^{(D)}, \mathbf{u}_2^{(D)}, \mathbf{u}_3^{(D)}, \cdots; \mathbf{u}_1^{(T)}, \mathbf{u}_2^{(T)}, \mathbf{u}_3^{(T)}, \cdots; \mathbf{v}_1^{(D)}, \mathbf{v}_2^{(D)}, \mathbf{v}_2^{(D)}, \mathbf{v}_3^{(D)}, \cdots; \mathbf{w}_1^{(D)}, \mathbf{w}_2^{(D)}, \mathbf{w}_3^{(D)}, \cdots; \mathbf{w}_1^{(T)}, \mathbf{w}_2^{(T)}, \mathbf{w}_3^{(T)}, \cdots; \mathbf{w}_3^{(T)}, \cdots; \mathbf{w}_4^{(T)}, \mathbf{w}_5^{(T)}, \mathbf{w}_5$

Our main results in this paper are following:

Theorem 1.1. Let (Ω, g) be an n-dimensional smooth compact Riemannian manifold with smooth boundary $\partial\Omega$, $(n \ge 2)$. Then for given Lamé coefficient $\mu > 0$, the spectrum $\sigma_D(L_{\lambda,\mu})$ (respectively, $\sigma_T(L_{\lambda,\mu})$) of $L_{\lambda,\mu}$ with Dirichlet (respectively, traction) boundary condition is discrete for any $\lambda \in (-2\mu, +\infty)$. The eigenvalues $\tau_k^{(D)}(\lambda)$ (respectively, $\tau_k^{(T)}(\lambda)$) of $L_{\lambda,\mu}$ ordered in a non-decreasing sequence, is continuous non-decreasing functions of λ .

Theorem 1.2. Let (Ω, g) be an n-dimensional smooth compact Riemannian manifold with smooth boundary $\partial \Omega$, $(n \ge 2)$. Then for given Lamé coefficient $\mu > 0$ and any $k \ge 1$, the Lamé eigenvalues $\tau_k^{(D)/(T)}(\lambda)$ tend to the Stokes eigenvalues $\varsigma_k^{(D)/(C)}$ (respectively, the Laplace eigenvalues $\theta_k^{(D)/(T)}$) as the Lamé constant $\lambda \to +\infty$ (respectively, $\lambda \to -\tau$), and the corresponding Lamé eigenspaces tend to the corresponding Stokes eigenspaces (respectively, Laplace eigenspaces).

Combining the results of Theorem 1.1 and Theorem 1.2 with the mini-max formulas for the eigenvalues, we have the basic eigenvalue inequalities:

Corollary 1.3. Let (Ω, g) be an n-dimensional smooth compact Riemannian manifold with smooth boundary $\partial\Omega$, $(n \ge 2)$. Let $\tau_k^{(D)/(T)}$, $\varsigma_k^{(D)/(C)}$, $\theta_k^{(D)/(T)}$ be the k-th eigenvalue with repetition according to the multiplicity of the self-adjoint operators $L_{\lambda,\mu}$, S_{μ} and $B_{\mu} := L_{-\mu,\mu}$ with corresponding boundary conditions, respectively. Then for any $\lambda \in (-\mu, +\infty)$ there holds

$$\theta_k^{(D)} \le \tau_k^{(D)}(\lambda) \le \varsigma_k^{(D)}, \quad \theta_k^{(T)} \le \tau_k^{(T)}(\lambda) \le \varsigma_k^{(T)}, \quad k = 1, 2, 3, \cdots.$$
 (1.19)

We introduce the partition function, or the trace of the heat semigroup for the Lamé

operator with Dirichlet (respectively, traction) boundary condition, by $Z^{(D)}(t) := \text{Tr } e^{-tL_{\lambda\mu}^{(D)}} = \sum_{k=1}^{\infty} e^{-t\tau_k^{(D)}}$ (respectively, $Z^{(T)}(t) := \text{Tr } e^{-tL_{\lambda\mu}^{(T)}} = \sum_{k=1}^{\infty} e^{-t\tau_k^{(T)}}$) defined for t > 0 and monotone decreasing in t.

Theorem 1.4. Let (Ω, g) be an n-dimensional compact smooth Riemannian manifold with smooth boundary $\partial\Omega$. Assume that the Lamé coefficients satisfy $\mu > 0$ and $\lambda + 2\mu > 0$. Let $0 < \tau_1^{(D)} < \tau_2^{(D)} \le \tau_3^{(D)} \le \cdots \le \tau_k^{(D)} \le \cdots$ (respectively, $0 \le \tau_1^{(T)} \le \tau_2^{(T)} \le \tau_3^{(T)} \le \cdots \le \tau_k^{(T)} \le$ \cdots) be the eigenvalues of the Lamé operator $L_{\lambda,\mu}$ with respect to the Dirichlet (respectively, traction) boundary condition. Then

where

$$\begin{aligned} a_0 &= \frac{n-1}{(4\pi\mu)^{n/2}} + \frac{1}{(4\pi(\lambda+2\mu))^{n/2}}, \quad a_1^{(D)} &= -\frac{1}{4} \left[\frac{n-1}{(4\pi\mu)^{(n-1)/2}} + \frac{1}{(4\pi(\lambda+2\mu))^{(n-1)/2}} \right], \\ a_1^{(T)} &= \frac{1}{4} \left[\frac{n-1}{(4\pi\mu)^{(n-1)/2}} + \frac{1}{(4\pi(\lambda+2\mu))^{(n-1)/2}} \right], \\ a_2 &= \frac{1}{6(4\pi)^{n/2}} \left[\left(\frac{1}{(\lambda+2\mu)^{(n-2)/2}} + \frac{n-7}{\mu^{(n-2)/2}} + \frac{12\mu}{n} \left(\frac{1}{(\lambda+2\mu)^{n/2}} + \frac{n-1}{\mu^{n/2}} \right) \right) \int_{\Omega} R(x) \, dV \\ &+ 2 \left(\frac{1}{(\lambda+2\mu)^{(n-2)/2}} + \frac{n-7}{\mu^{(n-2)/2}} \right) \int_{\partial\Omega} H(x) \, ds \right]. \end{aligned}$$

Here $Vol_n(\Omega)$ denotes the n-dimensional volume of Ω , $Vol_{n-1}(\partial\Omega)$ is the (n-1)-dimensional volume of $\partial\Omega$, R(x) denotes the scalar curvature at $x \in \Omega$, and H(x) is the mean curvature of the boundary $\partial\Omega$ at $x \in \partial\Omega$.

From Theorem 1.3 and Theorem 1.4 (i.e., as $\lambda \to +\infty$), we immediately obtain:

Corollary 1.5. Let (Ω, g) be an n-dimensional compact smooth Riemannian manifold with smooth boundary $\partial\Omega$. Assume that the viscosity coefficient satisfy $\mu > 0$. Let $0 < \varsigma_1^{(D)} < \varsigma_2^{(D)} \le \varsigma_3^{(D)} \le \cdots \le \varsigma_k^{(D)} \le \cdots$ (respectively, $0 \le \varsigma_1^{(C)} < \varsigma_2^{(C)} \le \varsigma_3^{(C)} \le \cdots \le \varsigma_k^{(C)} \le \cdots$) be the eigenvalues of the Stokes operator S_{μ} with respect to the Dirichlet (respectively, Cauchy force) boundary condition. Then

$$\begin{aligned} \mathcal{Z}^{(D)/(C)}(t) &= \sum_{k=1}^{\infty} e^{-t S_k^{(D)/(C)}} = b_0 t^{-n/2} Vol_n(\Omega) + b_1^{(D)/(C)} t^{(1-n)/2} Vol_{n-1}(\partial \Omega) \\ &+ b_2 t^{(2-n)/2} + O(t^{(3-n)/2}) \quad as \ t \to 0^+, \end{aligned}$$
for $n \ge 2$,

where

$$b_{0} = \frac{n-1}{(4\pi\mu)^{n/2}}, \qquad b_{1}^{(D)} = -\frac{n-1}{4(4\pi\mu)^{(n-1)/2}}, \qquad b_{1}^{(C)} = \frac{n-1}{4(4\pi\mu)^{(n-1)/2}}$$

$$b_{2} = \frac{1}{6(4\pi)^{n/2}} \left[\left(\beta_{n} + \frac{n-7}{\mu^{(n-2)/2}} + \frac{12(n-1)}{n\mu^{(n-2)/2}} \right) \int_{\Omega} R(x) \, dV + 2 \left(\beta_{n} + \frac{n-7}{\mu^{(n-2)/2}} \right) \int_{\partial\Omega} H(x) \, ds \right],$$

and

$$\begin{cases} \beta_n = 0 & \text{when } n \ge 3, \\ \beta_n = 1 & \text{when } n = 2. \end{cases}$$

Remark 1.6. For the Stokes eigenvalues with Dirichlet boundary condition in a bounded domain $\Omega \subset \mathbb{R}^n$, $(n \ge 2)$, the corresponding two-term asymptotic formula have been established in [Liu3] by a completely different method, in which we used the celebrated semigroup theory by Abe and Giga for the Stokes equations (see, [AGi] and [Giga]) and a (pseudodifferential operator) representation of inverse of the Stokes operator by Kozhevnikov [Koz].

Corollary 1.7. Let (Ω, g) be an n-dimensional compact smooth Riemannian manifold with smooth boundary $\partial\Omega$. Let $0 < \theta_1^{(D)} < \theta_2^{(D)} \le \cdots \le \theta_k^{(D)} \le \cdots$ (respectively, $0 \le \theta_1^{(T)} \le \theta_2^{(T)} \le \cdots \le \theta_k^{(T)} \le \cdots$) be the eigenvalues of the Laplace operator $\mu (\nabla^* \nabla \mathbf{u} - Ric(\mathbf{u}))$ with respect to the Dirichlet (respectively, traction) boundary condition. Then

where

$$c_{0} = \frac{n}{(4\pi\mu)^{n/2}}, \quad c_{1}^{(D)} = -\frac{n}{4(4\pi\mu)^{(n-1)/2}}, \quad c_{1}^{(T)} = \frac{n}{4(4\pi\mu)^{(n-1)/2}},$$

$$c_{2} = \frac{1}{6(4\pi)^{n/2}} \left[\frac{n+6}{\mu^{(n-2)/2}} \int_{\Omega} R(x) \, dV + \frac{2(n-6)}{\mu^{(n-2)/2}} \int_{\partial\Omega} H(x) \, ds \right].$$

Let us remark that all the above conclusions hold if Ω is a bounded domain with smooth boundary in the Euclidean space \mathbb{R}^n . Now, let Ω be a bounded domain with smooth boundary in the Euclidean plane \mathbb{R}^2 , and let μ be a positive constant. Denote by $\Lambda^{(D)}$ the buckling eigenvalues with Dirichlet boundary condition

$$\begin{cases} \mu \Delta^2 \psi + \Lambda \Delta \psi = 0 & \text{in } \Omega, \\ u = \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial \Omega, \end{cases}$$
(1.20)

and by $\Lambda^{(C)}$ the buckling eigenvalue with Cauchy force boundary

$$\begin{cases} \mu \Delta^2 \psi + \Lambda \Delta \psi = 0 & \text{in } \Omega, \\ \psi = 0 & \text{on } \partial \Omega, \\ (B\psi) \nu = 0 & \text{on } \partial \Omega, \end{cases}$$
(1.21)

where

$$(B\psi)\mathbf{v} = \left\{ \mu \begin{pmatrix} -2\partial_1\partial_2\psi_k & \partial_1^2\psi_k - \partial_2^2\psi_k \\ \partial_1^2\psi_k - \partial_2^2\psi_k & 2\partial_1\partial_2\psi_k \end{pmatrix} - \mu \sqrt{(\partial_1^2\psi_k - \partial_2^2\psi_k)^2 + (\partial_1\partial_2\psi_k)^2} \mathbf{I}_2 \right\} \mathbf{v} \quad \text{on } \partial\Omega.$$

It is easy to verify that each of spectrum is discrete with

$$0 < \Lambda_1^{(D)} < \Lambda_2^{(D)} \le \dots \le \Lambda_k^{(D)} \le \dots,$$

$$0 < \Lambda_1^{(C)} < \Lambda_2^{(C)} \le \dots \le \Lambda_k^{(C)} \le \dots,$$

each eigenvalue repeated according to its multiplicity.

Theorem 1.8. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with smooth boundary $\partial\Omega$. Let $\Lambda_k^{(D)}$ (respectively, $\Lambda_k^{(C)}$) be the k-th buckling eigenvalue with the Dirichlet (respectively, Cauchy force) boundary condition. Then the set $\{\Lambda_k^{(D)}\}_{k=1}^{\infty}$ (respectively, $\{\Lambda_k^{(C)}\}_{k=1}^{\infty}$) of all the buckling eigenvalues with Dirichlet (respectively, Cauchy force) boundary condition is the same as the set $\{\varsigma_k^{(D)}\}_{k=1}^{\infty}$ (respectively, $\{\varsigma_k^{(C)}\}_{k=1}^{\infty}$) of all the Stokes eigenvalues with Dirichlet (respectively, Cauchy force) boundary condition. That is,

$$\Lambda_k^{(D)} = \varsigma_k^{(D)}, \quad \Lambda_k^{(C)} = \varsigma_{k+1}^{(C)}, \quad k = 1, 2, 3, \cdots.$$
(1.22)

According to Corollary 1.5 and $\varsigma_1 = 0$, we get the following

Corollary 1.9. Let Ω be a bounded domain with smooth boundary in the Euclidean plane \mathbb{R}^2 . Then, for the buckling eigenvalues $\{\Lambda_k^{(D)}\}_{k=1}^{\infty}$ and $\{\Lambda_k^{(C)}\}_{k=1}^{\infty}$,

$$\mathcal{Z}^{(D)/(C)}(t) = \sum_{k=1}^{\infty} e^{-t\Lambda_k^{(D)/(C)}} = \frac{|\Omega|}{4\pi\mu t} + d_1^{(D)/(C)} t^{-1/2} + d_2 + O(1) \quad as \ t \to 0^+,$$

where

$$d_1^{(D)} = -\frac{|\partial\Omega|}{4\sqrt{4\pi\mu}}, \quad d_1^{(C)} = \frac{|\partial\Omega|}{4\sqrt{4\pi\mu}},$$
$$d_2 = -1 - \frac{1}{3\pi} \int_{\partial\Omega} H(x) \, ds.$$

Let Ω be a bounded domain with smooth boundary in the Euclidean space \mathbb{R}^2 , and let μ be a positive constant. Denote by $\{\Gamma_k^2, \phi_k\}_{k=1}^{\infty}$ the clamped plate eigenvalues

$$\begin{cases} \mu^2 \Delta^2 \phi_k - \Gamma_k^2 \phi_k = 0 & \text{in } \Omega, \\ \phi_k = \frac{\partial \phi_k}{\partial \nu} = 0 & \text{on } \partial \Omega. \end{cases}$$
(1.23)

For the Neumann eigenvalues $\{\Theta_k\}_{k=1}^{\infty}$ of the Laplace operator $\mu \Delta$, the Dirichlet eigenvalues $\{\Xi_k\}_{k=1}^{\infty}$ of the Laplace operator $\mu \Delta$, the clamped plate eigenvalues $\{\Gamma_k^2\}_{k=1}^{\infty}$ and the buckling eigenvalues $\{\Lambda_k^{(D)}\}_{k=1}^{\infty}$ on a smooth compact Riemannian manifold (Ω, g) with smooth boundary $\partial\Omega$, the present author in [Liu-11] proved the following basic inequalities:

$$\Theta_k < \Xi_k < \Gamma_k < \Lambda_k^{(D)}, \qquad k = 1, 2, 3, \cdots.$$
(1.24)

The inequalities (1.24) are sharp and they can not been further improved in a Riemannian manifold, see [Liu-11] for the detailed examples. Thus, from the asymptotic formula of

the Dirichlet eigenvalues of the Laplace operator $\mu \Delta$ (see, [MS-67]), Corollary 1.9 and the inequalities

$$\sum_{k=1}^{\infty} e^{-t\Lambda_k^{(D)}} < \sum_{k=1}^{\infty} e^{-t\Gamma_k} < \sum_{k=1}^{\infty} e^{-t\Xi_k},$$

we can obtain the following two-term asymptotic formulas:

Corollary 1.10. Let Ω be a bounded domain with smooth boundary in the Euclidean plane \mathbb{R}^2 . Then, for the Dirichlet eigenvalues $\{\Theta_k\}_{k=1}^{\infty}$ of the Laplace operator $\mu \Delta$, the buckling eigenvalues $\{\Lambda_k^{(D)}\}_{k=1}^{\infty}$ (in (1.20)) and the clamped plate eigenvalues $\{\Gamma_k\}_{k=1}^{\infty}$ (in (1.23)), the following two-term asymptotic formulas holds:

$$\left\{ \begin{array}{c} \sum_{k=1}^{\infty} e^{-t\Xi_k} \\ \sum_{k=1}^{\infty} e^{-t\Gamma_k} \\ \sum_{k=1}^{\infty} e^{-t\Lambda_k^{(D)}} \end{array} \right\} \sim \frac{|\Omega|}{4\pi\mu t} - \frac{|\partial\Omega|}{4\sqrt{4\pi\mu t}} + O(1) \quad as \ t \to 0^+.$$

$$(1.25)$$

The remainder of this paper is organized as follows. In Section 2, we give the proof of Theorem 1.1 by using eigenvalue variational principle. In Section 3, by a construct technique and by applying Gårding inequality we prove Theorem 1.2. Section 4 shows that the elastic Lamé operator is exactly the generalized Ahlfors Laplacian, and provides the third term coefficient of its asymptotic expansion from a result of [BGOP]. In Section 5, we prove Theorem 1.4 by combining double operator (defined on the double closed manifold), pseudodifferential operator technique and the result of Section 4. In Section 6, we discuss the buckling and clamped plate eigenvalue problems and further establish the corresponding eigenvalue inequalities and asymptotic formulas from property of the Stokes operator in a two-dimensional Euclidean bounded domain.

2. Monotonicity of the Lamé eigenvalues with respect to parameter

Proof of Theorem 1.1. By the maximum-minimum property of the Lamé eigenvalues, we have that for any $k \ge 1$,

$$\tau_{k}^{(\mathrm{D})}(\lambda) = \max_{\substack{E \subset H_{0}^{1}(\Omega, T\Omega) \\ \dim E = k-1}} \min_{\boldsymbol{\phi} \in E^{\perp}} \frac{\int_{\Omega} \left[2\mu (\operatorname{Def} \boldsymbol{\phi}, \operatorname{Def} \boldsymbol{\phi}) + \lambda (\operatorname{div} \boldsymbol{\phi})^{2} \right] dV}{\int_{\Omega} \boldsymbol{\phi} \cdot \boldsymbol{\phi} \, dV}$$

and

$$\tau_{k}^{(\mathrm{T})}(\lambda) = \max_{\substack{F \subset H^{1}(\Omega, T\Omega) \\ \dim F = k-1}} \min_{\phi \in F^{\perp}} \frac{\int_{\Omega} \left[2\mu (\operatorname{Def} \phi, \operatorname{Def} \phi) + \lambda (\operatorname{div} \phi)^{2} \right] dV}{\int_{\Omega} \phi \cdot \phi \, dV},$$

where E^{\perp} (respectively, F^{\perp}) is the orthogonal complement space of *E* (respectively, *F*) in $H_0^1(\Omega, T\Omega)$ (respectively, $H^1(\Omega, T\Omega)$) under the L^2 -norm. Or equivalently, there exists a

sequence of eigenpairs $\{(\tau_k^{(D)}(\lambda), \mathbf{u}_k^{(D)}(\lambda))\}_{k=1}^{\infty}$ (respectively, $\{(\tau_k^{(T)}(\lambda), \mathbf{u}_k^{(T)}(\lambda))\}_{k=1}^{\infty}$ such that $\tau_{k}^{(D)}(\lambda) = \frac{\int_{\Omega} \left[2\mu(\operatorname{Def} \mathbf{u}_1^{(D)}(\lambda), \operatorname{Def} \mathbf{u}_1^{(D)}(\lambda)) + \lambda\left(\operatorname{div} \mathbf{u}_1^{(D)}(\lambda)\right)^2\right] dV}{(2.1)}$

.

.....,

$$= \min_{\boldsymbol{\phi} \in H_0^1(\Omega, T\Omega)} \frac{\int_{\Omega} \mathbf{u}_1^{(D)}(\lambda) \cdot \mathbf{u}_1^{(D)}(\lambda) \, dV}{\int_{\Omega} \left[2\mu(\operatorname{Def} \boldsymbol{\phi}, \operatorname{Def} \boldsymbol{\phi}) + \lambda \left(\operatorname{div} \boldsymbol{\phi}\right)^2 \right] dV}{\int_{\Omega} \boldsymbol{\phi} \cdot \boldsymbol{\phi} \, dV},$$
(2.1)

$$\tau_{k}^{(\mathrm{D})}(\lambda) = \frac{\int_{\Omega} \left[2\mu (\operatorname{Def} \mathbf{u}_{k}^{(\mathrm{D})}(\lambda), \operatorname{Def} \mathbf{u}_{k}^{(\mathrm{D})}(\lambda)) + \lambda (\operatorname{div} \mathbf{u}_{k}^{(\mathrm{D})}(\lambda))^{2} \right] dV}{\int_{\Omega} \mathbf{u}_{k}^{(\mathrm{D})}(\lambda) \cdot \mathbf{u}_{k}^{(\mathrm{D})}(\lambda) dV}$$

$$= \min_{\substack{\boldsymbol{\phi} \in H_{0}^{1}(\Omega, T\Omega) \\ \int_{\Omega} \boldsymbol{\phi} \cdot \mathbf{u}_{j}^{(\mathrm{D})}(\lambda) dV = 0, \ j = 1, \cdots, k-1}} \frac{\int_{\Omega} \left[2\mu (\operatorname{Def} \boldsymbol{\phi}, \operatorname{Def} \boldsymbol{\phi}) + \lambda (\operatorname{div} \boldsymbol{\phi})^{2} \right] dV}{\int_{\Omega} \boldsymbol{\phi} \cdot \boldsymbol{\phi} dV},$$

$$(2.2)$$

and

Thus, we have to show that the solutions of the above variational problems are also eigenvectors for the Lamé boundary value problems, and they furnish *all* the eigenvectors. For $\phi \in H_0^1(\Omega, T\Omega)$ or $\phi \in H^1(\Omega, T\Omega)$, we denote the quadratic functionals for the variational eigenvalue problems:

$$A[\boldsymbol{\phi}] = \int_{\Omega} \left(2\mu (\operatorname{Def} \boldsymbol{\phi}, \operatorname{Def} \boldsymbol{\phi}) + \lambda (\operatorname{div} \boldsymbol{\phi})^2 \right) dV$$

and

$$B[\boldsymbol{\phi}] = \int_{\Omega} \boldsymbol{\phi} \cdot \boldsymbol{\phi} \, dV.$$

It is easy to verify that the associated polar forms

$$A[\phi, \psi] = \int_{\Omega} \left(2\mu (\operatorname{Def} \phi, \operatorname{Def} \psi) + \lambda (\operatorname{div} \phi) (\operatorname{div} \psi) \right) dV,$$

$$B[\phi, \psi] = \int_{\Omega} \phi \cdot \psi \, dV$$

satisfy the relations:

$$A[\phi + \psi] = A[\phi] + 2A[\phi, \psi] + A[\psi],$$
$$B[\phi + \psi] = B[\phi] + 2B[\phi, \psi] + B[\psi].$$

Without loss of generality, we only discuss the case of the variational problem with traction boundary condition in $H^1(\Omega, T\Omega)$ (since, for the case of the variational problem with Dirichlet boundary condition in $H^1_0(\Omega, T\Omega)$, it is quite similar and easier). We assume that its solution $\mathbf{u}_1^{(T)}(\lambda)$ satisfies $B[\mathbf{u}_1^{(T)}(\lambda)] = 1$. If **w** is any vector field in $H^1(\Omega, T\Omega)$, and if ϵ is an arbitrary constant, then for every value of ϵ and for $\mathbf{u} = \mathbf{u}_1^{(T)}(\lambda)$, $\tau = \tau_1^{(T)}(\lambda)$, we have

$$A[\mathbf{u} + \boldsymbol{\epsilon} \mathbf{w}] \ge \tau B[\mathbf{u} + \boldsymbol{\epsilon} \mathbf{w}],$$

i.e.,

$$\epsilon \left\{ A[\mathbf{u}, \mathbf{w}] - \tau B[\mathbf{u}, \mathbf{w}] + \frac{\epsilon}{2} (A[\mathbf{w}] - \tau B[\mathbf{w})) \right\} \ge 0$$
(2.5)

in view of the relation $A[\mathbf{u}] = \tau B[\mathbf{u}]$. The inequality (2.5) can be valid for arbitrary values of ϵ only if equality

$$A[\mathbf{u}, \mathbf{w}] - \tau B[\mathbf{u}, \mathbf{w}] = 0 \tag{2.6}$$

holds, i.e., if the expression $A - \tau B$ vanishes (corresponding to the first variational minimum problem (2.3)). By applying Green's formula (see, Section 2 of [Liu-19]), we get

$$A[\mathbf{u},\mathbf{w}] = \int_{\Omega} (L_{\mu,\lambda}\mathbf{u}) \cdot \mathbf{w} \, dV + \int_{\partial\Omega} \left(2\mu \left(\operatorname{Def} \mathbf{u} \right) \mathbf{v} + \lambda \left(\operatorname{div} \mathbf{u} \right) \mathbf{v} \right) \cdot \mathbf{w} \, ds.$$
(2.7)

Since the vector field $\mathbf{w} \in H^1(\Omega, T\Omega)$ is arbitrary, by (2.6) – (2.7) we immediately obtain equations

$$\begin{cases} L_{\lambda,\mu} \mathbf{u} = \tau \mathbf{u} & \text{in } \Omega, \\ 2\mu \left(\text{Def } \mathbf{u} \right) \mathbf{v} + \lambda \left(\text{div } \mathbf{u} \right) \mathbf{v} = 0 & \text{on } \partial \Omega \end{cases}$$
(2.8)

for $\mathbf{u} = \mathbf{u}_1^{(T)}(\lambda)$ and $\tau = \tau_1^{(T)}(\lambda)$. Considering the second minimum problem (in (2.4)) with the additional condition $B[\phi, \mathbf{u}_1^{(T)}(\lambda)] = 0$, where $\phi \in H^1(\Omega, T\Omega)$, we obtain equation (2.6) for $\mathbf{u} = \mathbf{u}_2^{(T)}(\lambda)$ and $\tau = \tau_2^{(T)}(\lambda)$ at first only under the assumption that $\mathbf{w} \in H^1(\Omega, T\Omega)$ satisfies relation

$$B[\mathbf{w}, \mathbf{u}_1^{(\mathrm{T})}(\lambda)] = 0.$$
(2.9)

Now if $\boldsymbol{\eta}$ is any vector field in $H^1(\Omega, T\Omega)$, we can determine a number *t* in such a way that the function $\mathbf{w} = \boldsymbol{\eta} + t \mathbf{u}_1$ satisfies condition (2.9) (this can be realized by setting $t = -B[\mathbf{u}_1^{(T)}(\lambda), \boldsymbol{\eta}]$). Particularly, by substituting the vector field $\mathbf{w} = \mathbf{u}_2^{(T)}(\lambda)$ in equation (2.6) with $\mathbf{u} = \mathbf{u}_1^{(T)}(\lambda)$ and $\tau = \tau_1^{(T)}(\lambda)$, we get

$$A[\mathbf{u}_{2}^{(\mathrm{T})}(\lambda).\mathbf{u}_{1}^{(\mathrm{T})}(\lambda)] = 0$$
(2.10)

since $\mathbf{u}_2^{(\mathrm{T})}(\lambda)$ satisfies the additional condition

$$B[\mathbf{u}_2^{(\mathrm{T})}(\lambda), \mathbf{u}_1^{(\mathrm{T})}(\lambda)] = 0.$$
(2.11)

If we substitute our vector field $\mathbf{w} = \boldsymbol{\eta} + t \mathbf{u}_1^{(T)}(\lambda)$ in equation (2.6), writing $\mathbf{u} = \mathbf{u}_2^{(T)}(\lambda)$, $\tau = \tau_2^{(T)}(\lambda)$, we find

$$A[\mathbf{u}, \boldsymbol{\eta}] - \tau B[\mathbf{u}, \boldsymbol{\eta}] + t \left(A[\mathbf{u}, \mathbf{u}_1^{(\mathrm{T})}(\lambda)] - \tau B[\mathbf{u}, \mathbf{u}_1^{(\mathrm{T})}(\lambda)] \right) = 0$$
(2.12)

or, taking into consideration equations (2.10) and (2.11),

$$A[\mathbf{u},\boldsymbol{\eta}] - \tau B[\mathbf{u},\boldsymbol{\eta}] = 0. \tag{2.13}$$

That is, equation (2.6) holds also for arbitrary vector fields $\boldsymbol{\eta}$ or \mathbf{w} without regard to the auxiliary condition (2.9). From this it follows directly, as above, that equation (2.8) is also valid for $\mathbf{u} = \mathbf{u}_2^{(T)}(\lambda)$ and $\tau = \tau_2^{(T)}(\lambda)$. Continuing in this process, we conclusion that the eigenvalue equation (2.13) holds generally for the solutions $\mathbf{u}_k^{(T)}(\lambda)$ and the minimum values $\tau_k^{(T)}(\lambda)$. For the solutions of the problem, we have the relations

$$A[\mathbf{u}_{k}^{(\mathrm{T})}(\lambda)] = \tau_{k}^{(\mathrm{T})}, \qquad A[\mathbf{u}_{k}^{(\mathrm{T})}(\lambda), \mathbf{u}_{j}^{(\mathrm{T})}(\lambda)] = 0$$

$$B[\mathbf{u}_{k}^{(\mathrm{T})}(\lambda)] = 1, \qquad B[\mathbf{u}_{k}^{(\mathrm{T})}(\lambda), \mathbf{u}_{j}^{(\mathrm{T})}(\lambda)] = 0 \qquad (k \neq j).$$
(2.14)

Our variational problems produce an infinite sequence of eigenpairs of the associated differential equations problem with traction boundary condition. Conversely, it is also true. That is, solutions of the variational eigenvalues problem (in $H^1(\Omega, T\Omega)$) are equivalent to the solutions of Lamé eigenvalue problem with traction boundary condition. Furthermore, the regularity of equations (2.8) implies that all weak solutions { $\mathbf{u}_k^{(T)}(\lambda)$ } belong to $C^{\infty}(\Omega, T\Omega)$.

As pointed out before, the variational eigenvalue problem (2.1)–(2.2) (which is considered in $H_0^1(\Omega, T\Omega)$) produce an infinite sequence of eigenpairs { $(\tau_k^{(D)}(\lambda), \mathbf{u}_k^{(D)}(\lambda))$ } with Dirichlet boundary condition. In addition, it can easily be shown that $\mathbf{u}_k^{(D)}$ belong to $C^{\infty}(\Omega, T\Omega)$. Finally, if $\lambda', \lambda'' \in (-2\mu, +\infty)$ satisfy $\lambda' < \lambda''$, then for any $\boldsymbol{\phi} \in H_0^1(\Omega, T\Omega)$ (in the case of Dirichlet boundary condition) or $\boldsymbol{\phi} \in H^1(\Omega, T\Omega)$ (in the case of traction boundary condition),

$$\frac{\int_{\Omega} \left[2\mu (\operatorname{Def} \boldsymbol{\phi}, \operatorname{Def} \boldsymbol{\phi}) + \lambda' (\operatorname{div} \boldsymbol{\phi})^2 \right] dV}{\int_{\Omega} \boldsymbol{\phi} \cdot \boldsymbol{\phi} \, dV} \le \frac{\int_{\Omega} \left[2\mu \langle \operatorname{Def} \boldsymbol{\phi}, \operatorname{Def} \boldsymbol{\phi}) + \lambda'' (\operatorname{div} \boldsymbol{\phi})^2 \right] dV}{\int_{\Omega} \boldsymbol{\phi} \cdot \boldsymbol{\phi} \, dV}.$$
(2.15)

Combining (2.1) – (2.4) and (2.15), we get that $\tau_k^{(D)}(\lambda)$ (respectively, $\tau_k^{(T)}(\lambda)$) is continuous in λ , and for any $k \ge 1$,

$$\tau_k^{(D)}(\lambda') \le \tau_k^{(D)}(\lambda''), \quad \text{when} \quad -2\mu < \lambda' < \lambda'' < +\infty.$$

$$\tau_k^{(T)}(\lambda') \le \tau_k^{(T)}(\lambda'') \quad \text{when} \quad -2\mu < \lambda' < \lambda'' < +\infty.$$
(2.16)

3. Relationship of eigenvalues for the Lamé operator, Stokes operator and Laplace operator

For the Laplace operator (i.e., the Lamé operator when $\lambda + \mu = 0$) with Dirichlet and traction boundary conditions, we need the following:

Remark 3.1. (*i*) Let (Ω, g) be a smooth compact Riemann manifold with smooth boundary. It is well-known (see, (4.26) and (4.28) on p. 305 of [Ta2]) that

$$Def^*\mathbf{u} = -div \,\mathbf{u},\tag{3.1}$$

where $(\operatorname{div} \mathbf{u})^{j} = u^{jk}_{:k}$, and the following Weitzenbock formula holds on (Ω, g) :

$$2 \operatorname{div} \operatorname{Def} \mathbf{u} = -\nabla^* \nabla \mathbf{u} + \operatorname{grad} \operatorname{div} \mathbf{u} + \operatorname{Ric}(\mathbf{u}). \tag{3.2}$$

Note that (3.2) can also be obtained from (1.1), (1.3), (3.1) and $\mu = -\lambda = 1$. Moreover, from (3.2) and Green's formula for the Lamé operator (see, §2 of [Liu-19]), we have that

$$2\int_{\Omega} \left(\operatorname{Def} \mathbf{u}, \operatorname{Def} \mathbf{u} \right) dV - \left\| \operatorname{div} \mathbf{u} \right\|_{L^2}^2 = \left(\nabla^* \nabla \mathbf{u}, \mathbf{u} \right)_{L^2} - \left(\operatorname{Ric} \left(\mathbf{u} \right), \mathbf{u} \right)_{L^2}$$
(3.3)

for any $\mathbf{u} \in H_0^1(\Omega, T\Omega)$ or $\mathbf{u} \in \{\mathbf{v} \in H^1(\Omega, T\Omega) \mid \mathcal{T}_{-1,1}\mathbf{v} = 0\}$, where $\mathcal{T}_{\lambda,\mu}\mathbf{v} = -2\mu(Def \mathbf{v})^{\#}\mathbf{v} - \lambda$ (div \mathbf{v}) \mathbf{v} . It follows that for all $\mathbf{u} \in H_0^1(\Omega, T\Omega)$ or $\mathbf{u} \in \{\mathbf{v} \in H^1(\Omega, T\Omega) \mid (T_{-\mu,\mu}\mathbf{v}) \mid_{\partial\Omega} = 0\}$,

$$\mu \int_{\Omega} 2(Def\mathbf{u}, Def\mathbf{u}) \, dV - \mu \int_{\Omega} (div \, \mathbf{u}) (div \, \mathbf{u}) \, dV = -\mu \int_{\Omega} (\Delta_B \mathbf{u}) \cdot \mathbf{u} \, dV - \mu (Ric(\mathbf{u}), \mathbf{u})_{L^2}, \quad (3.4)$$

where $\Delta_B := -\nabla^* \nabla$.

(ii) If $\Omega \subset \mathbb{R}^n$ is a bounded domain with smooth boundary $\partial \Omega$, then the corresponding result (3.4) can directly be verified as follows.

Let us recall that $(Def \mathbf{u}, Def \mathbf{w}) = \sum_{j,k=1}^{n} (\frac{1}{2} (\partial_{x_j} u_k + \partial_{x_k} u_j))(\frac{1}{2} (\partial_{x_j} w_k + \partial_{x_k} w_j))$. Similarly, grad \mathbf{u} : grad \mathbf{w} denots $\sum_{j,k=1}^{n} (\partial_{x_j} u_k)(\partial_{x_j} u_k)$ and we write grad \mathbf{u} : $(grad \mathbf{w})^T$ for $\sum_{j,k=1}^{n} (\partial_{x_j} u_k)(\partial_{x_k} w_j)$. Then it is easy to verify that for any vector fields \mathbf{u} and \mathbf{v} in $[H^1(\Omega)]^n$:

$$2 (Def \mathbf{u}, Def \mathbf{w}) = grad \,\mathbf{u} : grad \,\mathbf{w} + grad \,\mathbf{u} : (grad \,\mathbf{w})^T.$$
(3.5)

Applying the integration by parts we have

$$\int_{\Omega} (\operatorname{grad} \mathbf{u} : \operatorname{grad} \mathbf{w}) \, dx = \int_{\Omega} \sum_{j,k=1}^{n} (\partial_{x_j} u_k) (\partial_{x_j} w_k)$$

$$= \int_{\Omega} \left[\sum_{k=1}^{n} \operatorname{div} \left(w_k \partial_{x_1} u_k, \cdots, w_k \partial_{x_n} u_k \right) \, dx - \int_{\Omega} \sum_{k=1}^{n} w_k (\Delta u_k) \, dx \right]$$

$$= \int_{\partial\Omega} \sum_{j,k=1}^{n} (w_k v_j \partial_{x_j} u_k) \, ds - \int_{\Omega} (\Delta \mathbf{u}) \cdot \mathbf{w} \, dx, \quad \forall \, \mathbf{u}, \, \mathbf{w} \in [H^2(\Omega)]^n.$$
(3.6)

Similarly, two integrations by parts we get

$$\int_{\Omega} grad \,\mathbf{u} : (grad \,\mathbf{w})^T dx = -\int_{\Omega} (grad \, div \,\mathbf{u}) \cdot \mathbf{w} \, dx + \int_{\partial\Omega} \sum_{j,k=1}^n w_j v_k \partial_{x_j} u_k \, ds \tag{3.7}$$
$$= \int_{\Omega} (div \,\mathbf{u}) (div \,\mathbf{w}) \, dx - \int_{\partial\Omega} (div \,\mathbf{u}) (\mathbf{v} \cdot \mathbf{w}) \, ds + \int_{\partial\Omega} \sum_{j,k=1}^n w_k v_j \partial_{x_k} u_j \, ds, \quad \forall \mathbf{u}, \mathbf{w} \in [H^2(\Omega)]^n.$$

If follows from (3.5)–(3.8) that for any $\mathbf{u}, \mathbf{w} \in [H^2(\Omega)]^n$,

$$\int_{\Omega} 2 \left(Def \mathbf{u}, Def \mathbf{w} \right) dx = \int_{\Omega} \left[-(\Delta \mathbf{u}) \cdot \mathbf{w} + (div \mathbf{u})(div \mathbf{w}) \right] dx$$

$$+ \int_{\partial \Omega} \left[\sum_{j,k=1}^{n} \left(w_k v_j \partial_{x_j} u_k + w_k v_j \partial_{x_k} u_j \right) + (div \mathbf{u})(\mathbf{v} \cdot \mathbf{w}) \right] ds$$

$$= \int_{\Omega} \left(-(\Delta \mathbf{u}) \cdot \mathbf{w} + (div \mathbf{u})(div \mathbf{w}) \right) dx + \int_{\partial \Omega} \left(2 \left(Def \mathbf{u} \right) \mathbf{v} + (div \mathbf{u}) \mathbf{v} \right) \cdot \mathbf{w} ds.$$
(3.8)

Thus, in the case of $\lambda + \mu = 0$, we have

$$\mu \int_{\Omega} 2 \left(Def \mathbf{u}, Def \mathbf{w} \right) dx - \mu \int_{\Omega} (div \, \mathbf{u}) (div \, \mathbf{w}) \, dx$$

$$= -\mu \int_{\Omega} (\Delta \mathbf{u}) \cdot \mathbf{w} \, dx + \mu \int_{\partial \Omega} \left(2 \left(Def \mathbf{u} \right) \mathbf{v} + (div \, \mathbf{u}) \mathbf{v} \right) \cdot \mathbf{w} \, ds, \quad \forall \, \mathbf{u}, \, \mathbf{w} \in [H^2(\Omega)]^n.$$
(3.9)

This implies that for any $\mathbf{u}, \mathbf{w} \in [H_0^1(\Omega)]^n$ or $\mathbf{u}, \mathbf{w} \in \{\mathbf{v} \in [H^1(\Omega)]^n \mid -2\mu(Def\mathbf{v})^{\#}\mathbf{v} + \mu(div\mathbf{v})\mathbf{v} = 0\},\$

$$\mu \int_{\Omega} 2\left(Def\mathbf{u}, Def\mathbf{w}\right) dx - \mu \int_{\Omega} (div\,\mathbf{u})(div\,\mathbf{w}) \, dx = -\mu \int_{\Omega} (\Delta \mathbf{u}) \cdot \mathbf{w} \, dx.$$
(3.10)

Therefore, for the Dirichlet eigenvalue problem of the Laplacian (i.e., the Lamé operator when $\lambda + \mu = 0$), our definition is the same as the classical one. But, for the Laplacian with traction boundary condition, the eigenvalue problem has a bit difference with classical Neumann Laplacian (the Rayleigh quotient is different). Their relation can be seen in (5.28) of the proof of Theorem 1.4. In physics, the traction Laplacian eigenvalues have significant application, it describes the vibrational case of an elastic body when the velocities of the longitudinal and transverse waves are equal.

The following Lemma plays a key role for the Lamé operator $L_{\lambda,\mu}$:

Lemma 3.2. Let (Ω, g) be a smooth compact Riemannian manifold with smooth boundary $\partial \Omega$, and let the Lamé coefficients satisfy $\mu > 0$ and $\lambda + 2\mu > 0$. Then the following Gårding's inequality holds: there are constants $\tilde{c} > 0$ and $\tilde{d} > 0$ such that for all $\mathbf{u} \in H_{\partial}^{(D)/(T)}$,

$$\int_{\Omega} \left(2\mu \left(\text{Def } \mathbf{u}, \text{Def } \mathbf{u} \right) + \lambda \left(\text{div } \mathbf{u} \right) \left(\text{div } \mathbf{u} \right) \right) dV \ge \tilde{c} \|\mathbf{u}\|_{H^{1}(\Omega, T\Omega)}^{2} - \tilde{d} \|\mathbf{u}\|_{L^{2}(\Omega, T\Omega)}^{2}, \qquad (3.11)$$

where $H^{(D)}_{\partial} := H^1_0(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$ and $H^{(T)}_{\partial} := \{ \mathbf{w} \in H^2(\Omega, T\Omega) \mid \mathcal{T}_{\lambda,\mu} \mathbf{w} = 0 \text{ on } \partial\Omega \}.$

Proof. It is easy to verify that the Lamé operator is strongly elliptic, and the Lamé operator $L_{\lambda,\mu}$ with Dirichlet (respectively, traction) boundary condition is symmetric (i.e., $(L_{\lambda,\mu}^{(D)/(T)}\mathbf{u}, \mathbf{v})_{L^2(\Omega,T\Omega)} = (\mathbf{u}, L_{\lambda,\mu}^{(D)/(T)}\mathbf{v})_{L^2(\Omega,T\Omega)}$ for all $\mathbf{u}, \mathbf{v} \in H_{\partial}^{(D)/(T)}$, see also [Liu-39]). Combing this and Proposition 1.5 on p. 318 of [MaHu], we immediately obtain the desired result (3.11).

Proof of Theorem 1.2. In order to keep the proof as simple as possible, we assume that the Stokes eigenvalues are simple and that each Lamé eigenvalue is simple for λ large enough. The proof is divided into two cases according to the boundary conditions.

i) Let $(\varsigma^{[0]}, \mathbf{u}^{[0]})$ be a Stokes eigenpair with Dirichlet boundary condition (assume that it is simple). Then there exists a unique $p^{[0]} \in L^2(\Omega)$ with $\int_{\Omega} p^{[0]} dV = 0$ such that

| ĺ | $\int \mu (\nabla^* \nabla \mathbf{u}^{[0]} - \operatorname{Ric} (\mathbf{u}^{[0]})) + \nabla_g p^{[0]} = \varsigma^{[0]} \mathbf{u}^{[0]}$ | in s | Ω, |
|---|---|------|---------------------|
| ł | $\operatorname{div} \mathbf{u}^{[0]} = 0$ | in s | Ω, |
| | $\mathbf{u}^{[0]} = 0$ | on | $\partial \Omega$, |

since $p^{[0]}$ is unique up to an additive arbitrary constant. Clearly, by multiplying a constant, it can be assume that $\|\mathbf{u}^{[0]}\|_{H_0^1(\Omega,T\Omega)} < \frac{1}{2}$. We can construct a triplet $(\boldsymbol{\varsigma}^{[1]}, \mathbf{u}^{[1]}, p^{[1]})$ with $\boldsymbol{\varsigma}^{[1]} \in \mathbb{R}$, $\mathbf{u}^{[1]} \in J^0 \cap H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$, and $p^{[1]} \in L^2(\Omega)$, $\int_{\Omega} p^{[1]} dV = 0$ such that

$$\begin{cases} \mu(\nabla^* \nabla \mathbf{u}^{[1]} - \operatorname{Ric}(\mathbf{u}^{[1]})) + \nabla_g p^{[1]} = \varsigma^{[0]} \mathbf{u}^{[1]} + \varsigma^{[1]} \mathbf{u}^{[0]} & \text{in } \Omega, \\ \operatorname{div} \mathbf{u}^{[1]} = -p^{[0]} & \operatorname{in } \Omega \\ \mathbf{u}^{[1]} = 0 & \operatorname{on } \partial\Omega. \end{cases}$$
(3.12)

In fact, there exists a unique solution $\tilde{\mathbf{u}}$ satisfying the following boundary value problem

$$\begin{cases} \operatorname{div} \tilde{\mathbf{u}} = -p^{[0]} & \text{in } \Omega, \\ \tilde{\mathbf{u}} = 0 & \text{on } \partial\Omega. \end{cases}$$
(3.13)

It follows from (3.12) and (3.13) that

$$\begin{pmatrix}
\mu \left(\nabla^* \nabla (\mathbf{u}^{[1]} - \tilde{\mathbf{u}}) - \operatorname{Ric} (\mathbf{u}^{[1]} - \tilde{\mathbf{u}}) \right) + \nabla_g p^{[1]} = \varsigma^{[0]} (\mathbf{u}^{[1]} - \tilde{\mathbf{u}}) + \varsigma^{[1]} \mathbf{u}^{[0]} \\
-\mu (\nabla^* \nabla \tilde{\mathbf{u}} - \operatorname{Ric} \tilde{\mathbf{u}}) + \varsigma^{[0]} \tilde{\mathbf{u}} \quad \text{in } \Omega, \\
\operatorname{div} (\mathbf{u}^{[1]} - \tilde{\mathbf{u}}) = 0 \quad \text{in } \Omega \\
\mathbf{u}^{[1]} - \tilde{\mathbf{u}} = 0 \quad \operatorname{on } \partial\Omega.
\end{cases}$$
(3.14)

Put $\tilde{\tilde{\mathbf{u}}} := \mathbf{u}^{[1]} - \tilde{\mathbf{u}}$ and $\mathbf{f}^{[1]} := -\mu(\nabla^* \nabla \tilde{\mathbf{u}} - \operatorname{Ric}(\tilde{\mathbf{u}})) + \varsigma^{[0]} \tilde{\mathbf{u}}$. Then (3.14) becomes

$$\begin{cases} (S_{\mu} - \varsigma^{[0]}) \tilde{\mathbf{u}} = \mathbf{f}^{[1]} + \varsigma^{[1]} \mathbf{u}^{[0]} & \text{in } \Omega, \\ \operatorname{div} \tilde{\mathbf{u}} = 0 & \operatorname{in } \Omega, \\ \tilde{\mathbf{u}} = 0 & \operatorname{on } \partial\Omega. \end{cases}$$
(3.15)

which is solvable if its right hand side is orthogonal to $\mathbf{u}^{[0]}$: this can be realized by the choice of $\boldsymbol{\varsigma}^{[1]}$:

$$\boldsymbol{\varsigma}^{[1]} = -\frac{(\mathbf{f}^{[1]}, \mathbf{u}^{[0]})_{L^2}}{(\mathbf{u}^{[0]}, \mathbf{u}^{[0]})_{L^2}}.$$

By solving (3.15) (i.e., (3.14)), we get $\tilde{\mathbf{u}}$ and $p^{[1]}$, so that we obtain the required ($\varsigma^{[1]}, \mathbf{u}^{[1]}, p^{[1]}$) (since $\tilde{\mathbf{u}}$ is known by (3.13)). In addition, it can be assume $\|\mathbf{u}^{[1]}\|_{H_0^1(\Omega,T\Omega)} < \frac{1}{2^2}$. Generally, using the same way we can construct a sequence ($\varsigma^{[j]}, \mathbf{u}^{[j]}, p^{[j]}$), $j \ge 1$, with $\varsigma^{[j]} \in \mathbb{R}$, $\mathbf{u}^{[j]} \in J^0 \cap H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$ and $p^{[j]} \in L^2(\Omega)$, $\int_{\Omega} p^{[j]} dV = 0$ such that

$$\begin{cases} \mu(\nabla^* \nabla \mathbf{u}^{[j]} - \operatorname{Ric}(\mathbf{u}^{[j]})) + \nabla_g p^{[j]} = \varsigma^{[0]} \mathbf{u}^{[j]} + \varsigma^{[1]} \mathbf{u}^{[j-1]} + \dots + \varsigma^{[j]} \mathbf{u}^{[0]} & \text{in } \Omega, \\ \operatorname{div} \mathbf{u}^{[j]} = -p^{[j-1]} & \operatorname{in } \Omega, \\ \mathbf{u}^{[j]} = 0 & \operatorname{on } \partial\Omega. \end{cases}$$
(3.16)

Indeed, as the above discussion, after subtraction of a solution $\tilde{\mathbf{w}} \in H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$ of the equation

$$\begin{cases} \operatorname{div} \tilde{\mathbf{w}} = -p^{[j-1]} & \text{in } \Omega, \\ \tilde{\mathbf{w}} = 0 & \text{on } \partial\Omega, \end{cases}$$
(3.17)

(3.16) has the form

$$(S_{\mu} - \boldsymbol{\varsigma}^{[0]}) \,\tilde{\tilde{\mathbf{w}}} = \mathbf{f}^{[j]} + \boldsymbol{\varsigma}^{[j]} \mathbf{u}^{[0]} \tag{3.18}$$

by setting $\tilde{\mathbf{w}} := \mathbf{u}^{[j]} - \tilde{\mathbf{w}}$ and $\mathbf{f}^{[j]} := -\mu(\nabla^*\nabla \tilde{\mathbf{w}} - \operatorname{Ric}(\tilde{\mathbf{w}})) + \varsigma^{[1]}\mathbf{u}^{[j-1]} + \cdots + \varsigma^{[j-1]}\mathbf{u}^{[1]}$, which is solvable if its right hand side is orthogonal to $\mathbf{u}^{[0]}$: this can also be achieved by the choice of $\varsigma^{[j]}$:

$$\boldsymbol{\varsigma}^{[j]} = -\frac{(\mathbf{f}^{[j]}, \mathbf{u}^{[0]})_{L^2}}{(\mathbf{u}^{[0]}, \mathbf{u}^{[0]})_{L^2}}.$$

By solving (3.18) we then get the desired result (i.e., $(\boldsymbol{\varsigma}^{[j]}, \mathbf{u}^{[j]}, p^{[j]})$ satisfying (3.16)) and $\|\mathbf{u}^{[j]}\|_{H_0^1(\Omega,T\Omega)} < \frac{1}{2^{j+1}}$.

With $\epsilon = 1/(\lambda + \mu)$, we see that the power series in ϵ , $(\sum_{j\geq 0} \epsilon^j \varsigma^{[j]}, \sum_{j\geq 0} \epsilon^j \mathbf{u}^{[j]})$ is a formal eigenpair of $L_{\lambda,\mu}$ with Dirichlet boundary condition, since

$$L_{\lambda,\mu}\mathbf{u} = \mu(\nabla^*\nabla \mathbf{u} - \operatorname{Ric}(\mathbf{u})) - (\lambda + \mu) \operatorname{grad} \operatorname{div} \mathbf{u}$$

Let us show now that this also holds in the sense of asymptotic expansions: setting for any $m \ge 1$,

$$\underline{\varsigma}^{(m)} = \sum_{j=0}^{m} \epsilon^{j} \varsigma^{[j]} \text{ and } \underline{\mathbf{u}}^{(m)} = \sum_{j=0}^{m} \epsilon^{j} \mathbf{u}^{[j]}$$

we find by $\lambda + \mu = 1/\epsilon$, div $\mathbf{u}^{[j]} = -p^{[j-1]}$ and (3.16) that

$$\begin{split} & L_{\lambda,\mu} \underline{\mathbf{u}}^{(m)} - \underline{\varsigma}^{(m)} \underline{\mathbf{u}}^{(m)} \\ &= \sum_{j=0}^{m} \epsilon^{j} \Big[\mu (\nabla^{*} \nabla \mathbf{u}^{[j]} - \operatorname{Ric} (\mathbf{u}^{[j]})) \Big] - (\lambda + \mu) \operatorname{grad} \Big(\sum_{j=0}^{m} \epsilon^{j} \operatorname{div} \mathbf{u}^{[j]} \Big) - \Big(\sum_{j=0}^{m} \epsilon^{j} \varsigma^{[j]} \Big) \Big(\sum_{j=0}^{m} \epsilon^{j} \mathbf{u}^{[j]} \Big) \\ &= \epsilon^{m} \Big[\mu \Big(\nabla^{*} \nabla \mathbf{u}^{[m]} - \operatorname{Ric} (\mathbf{u}^{[m]}) \Big) \Big] + \sum_{j=0}^{m-1} \epsilon^{j} \Big[\mu \Big(\nabla^{*} \nabla \mathbf{u}^{[j]} - \operatorname{Ric} (\mathbf{u}^{[j]}) \Big) + \operatorname{grad} p^{[j]} \Big] \end{split}$$

$$-\left[\sum_{j=0}^{m-1} \epsilon^{j} \left(\sum_{k=0}^{j} \varsigma^{[k]} \mathbf{u}^{[j-k]}\right) + \sum_{j=0}^{m} \epsilon^{m+j} \left(\sum_{k=j}^{m} \varsigma^{[k]} \mathbf{u}^{[m+j-k]}\right)\right]$$
$$= \epsilon^{m} \left[\mu \left(\nabla^{*} \nabla \mathbf{u}^{[m]} - \operatorname{Ric}\left(\mathbf{u}^{[m]}\right)\right)\right] + \sum_{j=0}^{m-1} \epsilon^{j} \sum_{k=0}^{j} \varsigma^{[k]} \mathbf{u}^{[j-k]}$$
$$- \left[\sum_{j=0}^{m-1} \epsilon^{j} \left(\sum_{k=0}^{j} \varsigma^{[k]} \mathbf{u}^{[j-k]}\right) + \sum_{j=0}^{m} \epsilon^{m+j} \left(\sum_{k=j}^{m} \varsigma^{[k]} \mathbf{u}^{[m+j-k]}\right)\right]$$
$$= \epsilon^{m} \left[\mu \left(\nabla^{*} \nabla \mathbf{u}^{[m]} - \operatorname{Ric}\left(\mathbf{u}^{[m]}\right)\right)\right] - \sum_{j=0}^{m} \epsilon^{m+j} \left(\sum_{k=j}^{m} \varsigma^{[k]} \mathbf{u}^{[m+j-k]}\right).$$

Whence the uniform estimate for ϵ small enough (i.e., λ large enough)

$$\|L_{\lambda,\mu}\underline{\mathbf{u}}^{(m)}-\underline{\boldsymbol{\varsigma}}^{(m)}\underline{\mathbf{u}}^{(m)}\|_{(H_0^1(\Omega,T\Omega))'} \leq C\boldsymbol{\epsilon}^m \|\underline{\mathbf{u}}^{(m)}\|_{H_0^1(\Omega,T\Omega)},$$

where $(H_0^1(\Omega, T\Omega))'$ is the dual space of $H_0^1(\Omega, T\Omega)$. However, with $\mathbf{u}_k^{(D)}(\lambda)$ the normalized (in $L^2(\Omega, T\Omega)$) eigenvector associated to the eigenvalue $\tau_k^{(D)}(\lambda)$ of the Lamé operator $L_{\lambda,\mu}$ with Dirichlet boundary condition, for any $\varsigma \in \mathbb{R}$ and $\mathbf{u} \in H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$ there holds

$$\|L_{\lambda,\mu}\mathbf{u} - \varsigma \,\mathbf{u}\|_{(H_0^1(\Omega,T\Omega))'}^2 = \sum_{k\geq 1} (\tau_k^{(D)}(\lambda) - \varsigma)^2 (\mathbf{u}, \mathbf{u}_k^{(D)}(\lambda))^2 \|\mathbf{u}_k^{(D)}(\lambda)\|_{(H_0^1(\Omega,T\Omega))'}^2.$$
(3.19)

It follows from Gårding's inequality (3.11) that for any $\mathbf{u} \in H_{\partial}^{(D)}$,

$$\int_{\Omega} \left[2\mu \left(\text{Def } \mathbf{u}, \text{Def } \mathbf{u} \right) + \lambda \left(\text{div } \mathbf{u} \right) (\text{dix } \mathbf{u}) + \tilde{d} \left| \mathbf{u} \right|^2 \right] dV \ge \tilde{c} \left\| \mathbf{u} \right\|_{H^1_0(\Omega, T\Omega)}^2$$

where \tilde{c} and \tilde{d} are the positive constants in Gårding's inequality. That is,

$$\left((L_{\lambda,\mu} + \tilde{d})\mathbf{u}, \mathbf{u} \right)_{L^2(\Omega, T\Omega)} \ge \tilde{c} \|\mathbf{u}\|_{H^1_0(\Omega, T\Omega)}^2, \quad \forall \, \mathbf{u} \in H^{(D)}_{\partial}.$$
(3.20)

Noting that $(L_{\lambda,\mu} + \tilde{d}) \mathbf{u}_k^{(D)}(\lambda) = (\tau_k^{(D)} + \tilde{d}) \mathbf{u}_k^{(D)}(\lambda)$, by (3.20) we have

$$\left(\left(\tau_{k}^{(D)}+\tilde{d}\right)\mathbf{u}_{k}^{(D)}(\lambda),\mathbf{u}_{k}^{(D)}(\lambda)\right)_{L^{2}(\Omega,T\Omega)}\geq\tilde{c}\|\mathbf{u}_{k}^{(D)}(\lambda)\|_{H_{0}^{1}(\Omega,T\Omega)}^{2}.$$
(3.21)

Since

$$\|\mathbf{u}_{k}^{(D)}(\lambda)\|_{(H_{0}^{1}(\Omega,T\Omega))'} = \sup_{\mathbf{w}\in H_{0}^{1}(\Omega,T\Omega)} \frac{\left| (\mathbf{u}_{k}^{(D)}(\lambda), \mathbf{w})_{L^{2}(\Omega,T\Omega)} \right|}{\|\mathbf{w}\|_{H_{0}^{1}(\Omega,T\Omega)}} \ge \frac{\left| (\mathbf{u}_{k}^{(D)}(\lambda), \mathbf{u}_{k}^{(D)}(\lambda))_{L^{2}(\Omega,T\Omega)} \right|}{\|\mathbf{u}_{k}^{(D)}(\lambda)\|_{H_{0}^{1}(\Omega,T\Omega)}}$$

we find from this and (3.21) that

$$(\tau_{k}^{(D)} + \tilde{d}) \|\mathbf{u}_{k}^{(D)}(\lambda)\|_{(H_{0}^{1}(\Omega, T\Omega))'}^{2} \geq \left(\frac{\left((\tau_{k}^{(D)}(\lambda) + \tilde{d})\mathbf{u}_{k}^{(D)}(\lambda), \mathbf{u}_{k}^{(D)}(\lambda)\right)_{L^{2}(\Omega, T\Omega)}}{\|\mathbf{u}_{k}^{(D)}(\lambda)\|_{H_{0}^{1}(\Omega, T\Omega)}^{2}}\right) \|\mathbf{u}_{k}^{(D)}(\lambda)\|_{L^{2}(\Omega, T\Omega)}^{2} \geq \tilde{c},$$

so

$$\|\mathbf{u}_{k}^{(D)}(\lambda)\|_{(H_{0}^{1}(\Omega,T\Omega))'}^{2} \geq \frac{\tilde{c}}{\tau_{k}^{(D)}(\lambda) + \tilde{d}}$$

$$(3.22)$$

(Note that $\tau_k^{(D)}(\lambda) + \tilde{d} \ge \tau_1^{(D)}(\lambda) + \tilde{d} \ge \tau_1^{(D)}(-\mu) + \tilde{d} > 0$ for all $k \ge 1$). Combining this and (3.19), we find that for any $\varsigma \in \mathbb{R}$ and $\mathbf{u} \in H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$,

$$\|L_{\lambda,\mu}\mathbf{u}-\boldsymbol{\varsigma}\,\mathbf{u}\|_{(H_0^1(\Omega,T\Omega))'}^2 \geq C \sum_{k\geq 1} \frac{(\tau_k^{(D)}(\lambda)-\boldsymbol{\varsigma})^2}{\tau_k^{(D)}(\lambda)+\tilde{d}} (\mathbf{u},\mathbf{u}_k^{(D)}(\lambda))^2.$$

We see that

$$\sum_{k\geq 1} \frac{(\tau_k^{(D)}(\lambda) - \underline{\varsigma}^{(m)})^2}{\tau_k^{(D)}(\lambda) + \tilde{d}} (\underline{\mathbf{u}}^{(m)}, \mathbf{u}_k^{(D)}(\lambda))^2 \leq C\epsilon^{2m}.$$
(3.23)

Hence there exists k_0 such that $\lim_{\lambda \to \infty} \tau_{k_0}^{(D)}(\lambda)$ is equal to $\lim_{\epsilon \to 0} \underline{\varsigma}^{(m)} = \varsigma^{[0]}$, and

$$\exists \delta > 0, \quad \forall k \neq k_0, \quad \frac{(\tau_k^{(D)}(\lambda) - \underline{\varsigma}^{(m)})^2}{\tau_k^{(D)}(\lambda) + \tilde{d}} \ge \delta, \tag{3.24}$$

thus

$$\sum_{k\neq k_0} (\underline{\mathbf{u}}^{(m)}, \mathbf{u}_k^{(D)}(\lambda))^2 \leq C \epsilon^{2m}.$$

On the other hand, for $k = k_0$, we must have $|\underline{\varsigma}^{(m)} - \tau_k^{(D)}(\lambda)| \leq C\epsilon^m$. This proves that $\sum_{l\geq 0} \epsilon^l \varsigma^{[l]}$ (respectively, $\sum_{l\geq 0} \epsilon^l \mathbf{u}^{[l]}$) is the asymptotic development of $\tau_{k_0}^{(D)}(\lambda)$ (respectively, $\mathbf{u}_{k_0}^{(D)}(\lambda)$) as $\lambda \to +\infty$.

Conversely, let us fix $\tau^{(D)}(\lambda) = \tau_k^{(D)}(\lambda)$ the *k*-th eigenvalue of $L_{\lambda,\mu}$ with Dirichlet boundary condition. From Theorem 1.1 and the previous proof we deduce that $\tau^{(D)}(\lambda) \leq \varsigma_k^{(D)}$, where $\varsigma_k^{(D)}$ is the *k*-th Stokes eigenvalue with Dirichlet boundary condition. Thus $\tau_k^{(D)}(\lambda)$ is bounded as $\lambda \to \infty$ and has a limit $\varsigma^{[0]}$. The corresponding normalized eigenvectors $\mathbf{u}(\lambda) = \mathbf{u}_k^{(D)}(\lambda)$ are thus bounded in the domain of any power of $L_{\lambda,\mu}$, thus in $H^{1+\delta}(\Omega, T\Omega)$ for $\delta > 0$ small enough. Therefore $\mathbf{u}(\lambda)$ has a limit $\mathbf{u}^{[0]}$ in $J^0 \cap H_0^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$. Going back to the equations satisfied by $\mathbf{u}(\lambda)$ we find that as $\lambda \to +\infty$,

div
$$\mathbf{u}^{[0]} = 0$$
 and $(\lambda + \mu)$ grad div $\mathbf{u}(\lambda) \to \mu(\nabla^* \nabla \mathbf{u}^{[0]} - \operatorname{Ric}(\mathbf{u}^{[0]})) - \varsigma^{[0]} \mathbf{u}^{[0]}$ in $J^0 \cap H^1_0 \cap H^2$.

Setting $p(\lambda) = (\lambda + \mu) \operatorname{div} \mathbf{u}(\lambda)$, we obtain that it converges in $L^2(\Omega)$ to a limit $-p^{[0]}$ as $\lambda \to +\infty$. Thus $(\varsigma^{[0]}, \mathbf{u}^{[0]})$ is a Stokes eigenpair.

ii) For the Stokes operator with Cauchy force boundary condition, the proof is almost the same as in i) only some little modifications are done: from (3.19) on, we have that for any $\varsigma \in \mathbb{R}$ and $\mathbf{u} \in H^1(\Omega, T\Omega) \cap H^2(\Omega, T\Omega)$

$$\|L_{\lambda,\mu}\mathbf{u} - \varsigma \,\mathbf{u}\|_{(H^{1}(\Omega,T\Omega))'}^{2} \ge C \sum_{k\ge 1} (\tau_{k}^{(T)}(\lambda) - \varsigma)^{2} (\mathbf{u}, \mathbf{u}_{k}^{(T)}(\lambda))^{2} \|\mathbf{u}_{k}^{(T)}(\lambda)\|_{(H^{1}(\Omega,T\Omega))'}^{2}, \qquad (3.25)$$

where $\{(\tau_k^{(T)}(\lambda), \mathbf{u}_k^{(T)}(\lambda))\}_{k=1}^{\infty}$ are eigenpairs of $L_{\lambda,\mu}$ with traction boundary condition (note that $\tau_1^{(T)}(\lambda) = 0$ and $\mathbf{u}_1^{(T)}(\lambda) = \text{constant}$). Note that $\tau_k^{(T)}(\lambda) + \tilde{d} \ge \tau_1^{(T)}(\lambda) + \tilde{d} \ge \tau_1^{(T)}(-\mu) + \tilde{d} \ge \tilde{d} > 0$ for all $k \ge 1$. It is similar to the discussion of (i) by applying Gårding's inequality that

$$\|\mathbf{u}_{k}^{(T)}(\lambda)\|_{(H_{0}^{1}(\Omega,T\Omega))'}^{2} \geq \frac{\tilde{c}}{\tau_{k}^{(T)}(\lambda) + \tilde{d}}.$$
(3.26)

From this and (3.25) we get that for any $\varsigma \in \mathbb{R}$ and $\mathbf{u} \in H_{\partial}^{(T)}$,

$$\|L_{\lambda,\mu}\mathbf{u}-\varsigma \mathbf{u}\|_{(H_0^1(\Omega,T\Omega))'}^2 \geq C \sum_{k\geq 1} \frac{(\tau_k^{(T)}(\lambda)-\varsigma)^2}{\tau_k^{(T)}(\lambda)+\tilde{d}} (\mathbf{u},\mathbf{u}_k^{(T)}(\lambda))^2.$$

Therefore

$$\sum_{k\geq 1} \frac{(\tau_k^{(T)}(\lambda) - \underline{\varsigma}^{(m)})^2}{\tau_k^{(T)}(\lambda) + \tilde{d}} (\underline{\mathbf{u}}^{(m)}, \mathbf{u}_k^{(T)}(\lambda))^2 \le C\epsilon^{2m}.$$
(3.27)

Hence there exists k_0 such that $\lim_{\lambda \to \infty} \tau_{k_0}^{(T)}(\lambda)$ is equal to $\lim_{\epsilon \to 0} \underline{\varsigma}^{(m)} = \varsigma^{[0]}$, and

$$\exists \delta > 0, \quad \forall k \neq k_0, \quad \frac{(\tau_k^{(T)}(\lambda) - \underline{\varsigma}^{(m)})^2}{\tau_k^{(T)}(\lambda) + \tilde{d}} \ge \delta, \tag{3.28}$$

thus

$$\sum_{k\neq k_0} (\underline{\mathbf{u}}^{(m)}, \mathbf{u}_k^{(T)}(\lambda))^2 \leq C\epsilon^{2m},$$

and for $k = k_0$ we must have $|\underline{\varsigma}^{(m)} - \tau_k^{(T)}(\lambda)| \le C\epsilon^m$. Others are completely similar to proof of (i). Therefore, the desired result still holds.

Finally, since each of the Lamé eigenvalues is continuous with respect to the Lamé coefficients, by noting Remark 3.1 we immediately obtain the corresponding conclusions for the Laplace-type operator $-\mu(\nabla^*\nabla - \text{Ric})$ with giving boundary conditions as $\lambda \to -\mu$.

4. Generalized Ahlfors Laplacian and the third coefficient of the heat trace asymptotic of the Lamé operator

In order to exactly give the third coefficient of the heat trace asymptotic expansion for the Lamé operator, we need to use a result of T. Branson, P. Gilkey, B. Ørsted, A. Pierzchalski in [BGOP] for the generalized Ahlfors Laplacian. Thus, we must show that the 1-form representation of the Lamé operator $L_{\lambda,\mu}$ (defined on a Riemannian manifold (Ω, g)) is just the generalized Ahlfors Laplacian. Let us remark that the classical Ahlfors Laplacian originated from the conformal geometry and was introduced by Ahlfors in 1974 and 1976 (see [Ahl-74] and [Ahl-76]).

Let $T\Omega$ and $T^*\Omega$ be the tangent and cotangent bundles of the Riemannian *n*-manifold (Ω, g) , respectively. The space of all C^{∞} vector fields will be denoted by \mathscr{X} . Recall that in terms of vector fields, the Lamé operator $L_{\lambda,\mu}$ defined on (Ω, g) can be written as (cf. (1.1)):

$$L_{\lambda,\mu} \mathbf{u} := \mu (\nabla^* \nabla \mathbf{u}) - (\mu + \lambda) \operatorname{grad} \operatorname{div} \mathbf{u} - \mu \operatorname{Ric}(\mathbf{u})$$

$$= \left(-\mu \nabla_k \nabla^k u^j - (\lambda + \mu) \nabla^j \nabla_k u^k - \mu \operatorname{Ric}_k^j u^k \right) \frac{\partial}{\partial x_j}$$

$$(4.1)$$

$$= \left(-\mu \nabla_k \nabla^k u^j - \lambda \nabla^j \nabla_k u^k - \mu \nabla_k \nabla^j u^k\right) \frac{\partial}{\partial x_j}, \quad \mathbf{u} = u^j \frac{\partial}{\partial x_j} \in \mathscr{X},$$

where $\nabla^* \nabla \mathbf{u}$ is given by (1.3), and $\{\frac{\partial}{\partial x_j}\}_{j=1}^n$ are the coordinate basis. On the other hand, the Lamé operator as well as the associated boundary value problems can equivalently be discussed in terms of 1-form. If α is the 1-form dual to the vector field $\mathbf{u} = u^j \frac{\partial}{\partial x_j}$ in the sense that

$$\alpha(\mathbf{X}) = g(\mathbf{u}, \mathbf{X}), \ \mathbf{X} \in \mathscr{X},$$

then

$$\alpha = u_j dx_j \quad \text{and} \quad u_j = g_{jl} u^l. \tag{4.2}$$

Noting that $\nabla g = 0$, one has, in index notation,

$$\nabla_k \nabla^k u^j = \nabla_k \nabla^k g^{jl} u_l = g^{jl} g^{km} \nabla_k \nabla_m u_l = g^{jl} \nabla^m \nabla_m u_l,$$

$$\nabla^j \nabla_k u^k = \nabla^j \nabla_k g^{kl} u_l = \nabla^j \nabla^l u_l = g^{jm} \nabla_m \nabla^l u_l,$$

$$\nabla_k \nabla^j u^k = \nabla_k \nabla^j g^{kl} u_l = g^{kl} \nabla_k g^{jm} \nabla_m u_l = g^{jm} \nabla^l \nabla_m u_l.$$

Combining the above facts and the last line of (4.1) we find that for $\mathbf{u} = u^j \frac{\partial}{\partial x_j} \in \mathscr{X}$,

$$L_{\lambda,\mu}\mathbf{u} = \left(-\mu g^{jl}\nabla^m \nabla_m u_l - \lambda g^{jm}\nabla_m \nabla^l u_l - \mu g^{jm}\nabla^l \nabla_m u_l\right) \frac{\partial}{\partial x_j}$$
(4.3)
$$= g^{jm} \left(-\mu \nabla^l \nabla_l u_m - \lambda \nabla_m \nabla^l u_l - \mu \nabla^l \nabla_m u_l\right) \frac{\partial}{\partial x_j}$$

$$:= \phi^j \frac{\partial}{\partial x_j},$$

where

$$\phi^{j} := g^{jm} \Big(-\mu \nabla^{l} \nabla_{l} u_{m} - \lambda \nabla_{m} \nabla^{l} u_{l} - \mu \nabla^{l} \nabla_{m} u_{l} \Big).$$

Therefore, we have that

$$L^{\flat}_{\lambda,\mu}(\alpha) := \left(L_{\lambda,\mu}\mathbf{u}\right)^{\flat} = \psi_k \, dx_k, \tag{4.4}$$

where

$$\psi_{k} = g_{kj}\phi^{j} = g_{kj}g^{jm} \Big(-\mu\nabla^{l}\nabla_{l}u_{m} - \lambda\nabla_{m}\nabla^{l}u_{l} - \mu\nabla^{l}\nabla_{m}u_{l} \Big)$$

$$= -\mu\nabla^{l}\nabla_{l}u_{k} - \lambda\nabla_{k}\nabla^{l}u_{l} - \mu\nabla^{l}\nabla_{k}u_{l}$$

$$= -\mu\nabla^{l}\nabla_{l}u_{k} - (\lambda + 2\mu)\nabla_{k}\nabla^{l}u_{l} + \mu\nabla^{l}\nabla_{k}u_{l} - 2\mu\operatorname{Ric}_{k}^{l}u_{l},$$

$$(4.5)$$

 α is given by (4.2), and b is the flat operator (for a vector field) by lowering an index. Hence

$$L^{\flat}_{\lambda,\mu}(\alpha) = \left(-(\lambda + 2\mu)\nabla_k \nabla^l u_l + \mu \nabla^l \nabla_k u_l - \mu \nabla^l \nabla_l u_k - 2\mu \operatorname{Ric}^l_k u_l\right) dx_k.$$
(4.6)

Let $d : \Lambda T^*\Omega \to \Lambda T^*\Omega$ be the exterior differential operator, where $\Lambda T^*\Omega = \bigoplus_{p=0}^n \Lambda^p T^*\Omega$. The adjoint operator δ of d acting on a p-form α is defined in terms of d and the Hodge star operator by formula

$$\delta\alpha = (-1)^{np+n+1} * d * \alpha,$$

and the Hodge star operator $*: \Lambda^p T^*\Omega \to \Lambda^{n-p}T^*\Omega$, $p = 0, \dots, n$, is defined by

$$\langle \gamma, \eta \rangle = \gamma \wedge *\eta$$

for any $\gamma, \eta \in \Lambda^p T^*\Omega$. It is well known (see, for example, p. 16–17 of [CLN]) that $\nabla^l u_l = \operatorname{div} \alpha = -\delta \alpha$ for the 1-form $\alpha = u_j dx_j$. Furthermore, from Exercise 5 on p. 561 in [Ta3] we know that $\delta d\alpha = (\nabla^l \nabla_k u_l - \nabla^l \nabla_l u_k) dx_k$ for the above 1-form α . Combining these facts and (4.6) we obtain that

$$L^{\flat}_{\lambda,\mu}(\alpha) = (\lambda + 2\mu)d\delta\alpha + \mu\delta d\alpha - 2\mu\operatorname{Ric}(\alpha).$$
(4.7)

Here Ric (α) denotes the Ricci action on 1-forms α :

$$\operatorname{Ric}(\alpha) = \operatorname{Ric}(\mathbf{u}, \cdot) = (\operatorname{Ric}_{k}^{l} u_{l}) dx_{k}$$

where **u** is the vector field dual to α . (4.7) is the 1-form representation of the Lamé operator.

The generalized Ahlfors Laplacian (see, [BGOP]) is defined as the operator $P = ad\delta + b\delta d - \epsilon \rho$ on $\Lambda^1 T^*\Omega$, where *a* and *b* are positive constants and where $\epsilon \rho$ is an arbitrary constant multiple of the Ricci tensor. Clearly, from (4.7) we see that the 1-form representation $L_{\lambda,\mu}^{b}$ of the elastic Lamé operator $L_{\lambda,\mu}$ is just the generalized Ahlfors Laplacian on $\Lambda^1 T^*\Omega$ with $\lambda + 2\mu = a > 0, \mu = b > 0$ and $2\mu = \epsilon$.

Combining the above discussion and Branson-Gilkey-Ørsted-Pierzchalski's theorem (see, [BGOP]), we get the following:

Lemma 4.1. Let $L_{\lambda,\mu}$ be the Lamé operator defined on an n-dimensional compact Riemannian manifold (Ω, g) with smooth boundary $\partial\Omega$. Then, for the Dirichlet (or traction) boundary conditions, there is an asymptotic expansion of the form:

$$Tr \ e^{-tL_{\lambda,\mu}} \sim \sum_{k=0}^{\infty} a_k(L_{\lambda,\mu}) t^{(k-n)/2} \quad as \ t \to 0^+,$$
 (4.8)

where the first coefficient $a_0(L_{\lambda,\mu})$ and the third coefficient $a_2(L_{\lambda,\mu})$ are given by

$$a_0(L_{\lambda,\mu}) = (4\pi)^{-n/2} ((\lambda + 2\mu)^{-n/2} + (n-1)\mu^{-n/2}) Vol(\Omega),$$
(4.9)

$$a_{2}(L_{\lambda,\mu}) = \frac{1}{6(4\pi)^{n/2}} \Big[\Big(\frac{1}{(\lambda+2\mu)^{(n-2)/2}} + \frac{n-7}{\mu^{(n-2)/2}} + \frac{12\mu}{n} \Big(\frac{1}{(\lambda+2\mu)^{n/2}} + \frac{n-1}{\mu^{n/2}} \Big) \Big]_{\Omega} R(x) \, dV \qquad (4.10)$$
$$+ 2 \Big(\frac{1}{(\lambda+2\mu)^{(n-2)/2}} + \frac{n-7}{\mu^{(n-2)/2}} \Big) \int_{\partial\Omega} H(x) \, ds \Big].$$

Here R(x) *is the scalar curvature at* $x \in \Omega$ *,* $H(x) := L_{\alpha\alpha}(x)$ *be the mean curvature of* Ω *at* $x \in \partial \Omega$ *, and* $L_{\alpha\beta}$ *be the second fundamental form.*

Proof. From the above discussion, we see that the 1-form representation of the Lamé operator $L_{\lambda,\mu}$ is exactly the generalized Ahlfors Laplacian $a d\delta + b \delta d - \epsilon \rho$ with $a = \lambda + 2\mu$, $b = \mu$ and $\epsilon = 2\mu$. It is completely similar to the proof of Theorem 4.2 (or proof of Theorem 4.3) of [BGOP]) (i.e., by calculating the coefficient a_2 on $S^{n-1} \times [0, \pi]$ and by applying the invariance theory of Gilkey [Gil2]) that the first coefficient a_0 and the third coefficient a_2 are obtained. Actually, the desired conclusion is the same as the result in (c) of Theorem 4.2 (or (c) of Theorem 4.3) on p. 5 of [BGOP]) with $a := \lambda + 2\mu$, $b := \mu$ and $\epsilon = 2\mu$.

5. Asymptotics formulas of heat traces for the Lamé operators

Proof of Theorem 1.4. Part proof (particularly, for two-term asymptotic expansions) is similar to [Liu2] (see also [Liu-23]). For the sake of completeness, we shall give a detailed proof up to three-term asymptotics. From the theory of elliptic operators (see [GiTr], [Mo3], [Pa], [Stew]), we see that the Lamé operator $-L_{\lambda,\mu}$ can generate strongly continuous semigroup $(e^{-tL_{\lambda,\mu}^{(D)}})_{t\geq 0}$ (respectively, $(e^{-tL_{\lambda,\mu}^{(T)}})_{t\geq 0}$) with respect to the Dirichlet (respectively, traction boundary) condition, in suitable spaces of vector-valued functions (for example, in $C_0(\Omega, T\Omega)$ (by Stewart [Ste]) or in $L^2(\Omega, T\Omega)$ (by Browder [Bro])), or in $L^p(\Omega, T\Omega)$ (by Friedman [Fri]). Furthermore, there exist matrix-valued functions $\mathbf{K}^{(D)}(t, x, y)$ (respectively, $\mathbf{K}^{(T)}(t, x, y)$), which is called the integral kernel, such that (see [Bro] or p. 4 of [Fri])

$$e^{-tL_{\lambda,\mu}^{(D)/(T)}}\mathbf{w}_0(x) = \int_{\Omega} \mathbf{K}^{(D)/(T)}(t,x,y)\mathbf{w}_0(y) \, dV_y, \quad \mathbf{w}_0 \in L^2(\Omega,T\Omega).$$

Let $\{\mathbf{u}_{k}^{(D)}\}_{k=1}^{\infty}$ (respectively, $\{\mathbf{u}_{k}^{(T)}\}_{k=1}^{\infty}$) be the orthonormal eigenvectors of the elastic operator $L_{\lambda,\mu}^{(D)}$ (respectively, $L_{\lambda,\mu}^{(T)}$) corresponding to the eigenvalues $\{\tau_{k}^{(D)}\}_{k=1}^{\infty}$ (respectively, $L_{\lambda,\mu}^{(T)}$), then the integral kernels $\mathbf{K}^{(D)/(T)}(t, x, y) = e^{-tL_{\lambda,\mu}^{(D)/(T)}} \delta(x - y)$ are given by

$$\mathbf{K}^{(D)/(T)}(t, x, y) = \sum_{k=1}^{\infty} e^{-t\tau_k^{(D)/(T)}} \mathbf{u}_k^{(D)/(T)}(x) \otimes \mathbf{u}_k^{(D)/(T)}(y).$$
(5.1)

This implies that the integrals of the traces of $\mathbf{K}^{(D)/(T)}(t, x, y)$ are actually spectral invariants:

$$\int_{\Omega} \operatorname{Tr} \left(\mathbf{K}^{(D)/(T)}(t, x, x) \right) dV_x = \sum_{k=1}^{\infty} e^{-t\tau_k^{(D)/(T)}}.$$
(5.2)

We will combine calculus of symbols (see [Gru]) and "method of images" to deal with asymptotic expansions for the trace integrals of integral kernels. Let $\mathcal{M} = \Omega \cup (\partial \Omega) \cup \Omega^*$ be the (closed) double of Ω , and $\mathcal{L}_{\lambda,\mu}$ the double to \mathcal{M} of the operator $L_{\lambda,\mu}$ on Ω .

Let us explain the double Riemannian manifold \mathcal{M} and the double differential operator $\mathcal{L}_{\lambda,\mu}$ more precisely, and introduce how to get them from the given Riemannian manifold Ω and the Lamé operator $L_{\lambda,\mu}$. The double of Ω is the manifold $\Omega \cup_{\text{Id}} \Omega$, where $\text{Id} : \partial\Omega \to \partial\Omega$ is the identity map of $\partial\Omega$; it is obtained from $\Omega \sqcup \Omega$ by identifying each boundary point in one copy of Ω with same boundary point in the other. It is a smooth manifold without

boundary, and contains two regular domains diffeomorphic to Ω (see, p. 226 of [Lee]). When considering the double differential system $\mathcal{L}_{\lambda,\mu}$ crossing the boundary, we make use of the coordinates as follows. Let $x' = (x_1, \dots, x_{n-1})$ be any local coordinates for $\partial\Omega$. For each point $(x', 0) \in \partial\Omega$, let x_n denote the parameter along the unit-speed geodesic starting at (x', 0) with initial direction given by the inward boundary normal to $\partial\Omega$ (Clearly, x_n is the geodesic distance from the point (x', 0) to the point (x', x_n)). In such coordinates $x_n > 0$ in Ω , and $\partial\Omega$ is locally characterized by $x_n = 0$ (see, [LU] or [Ta2]). Since the Lamé operator is a linear differential operator defined on Ω , it can be further denoted as (see (1.3)) $L_{\lambda,\mu} := L_{\lambda,\mu}(x, \{g^{jk}(x)\}_{1 \le j,k \le n}, \{\Gamma^j_{kl}(x)\}_{1 \le j,k,l \le n}, \{R^{j'}_{k}(x)\}_{1 \le j,k \le n}, \frac{\partial}{\partial x_n}, \dots, \frac{\partial}{\partial x_{n-1}}, \frac{\partial}{\partial x_n}$. Let $\varpi : (x_1, \dots, x_{n-1}, x_n) \mapsto (x_1, \dots, x_{n-1}, -x_n)$ be the reflection with respect to the boundary $\partial\Omega$ in \mathcal{M} (here we always assume $x_n \ge 0$). Then we can get the Ω^* from the given Ω and ϖ . Now, we discuss the change of the metric g from Ω to Ω^* by ϖ . Recall that the Riemannian metric (g_{ij}) is given in the local coordinates x_1, \dots, x_n , i.e., $g_{ij}(x_1, \dots, x_n)$. In terms of the new coordinates z_1, \dots, z_n , with $x_i = x_i(z_1, \dots, z_n)$, $i = 1, \dots, n$, the same metric is given by the functions $\tilde{g}_{ij} = \tilde{g}_{ij}(z_1, \dots, z_n)$, where

$$\tilde{g}_{ij} = \sum_{k,l=1}^{n} \frac{\partial x_k}{\partial z_i} g_{kl} \frac{\partial x_l}{\partial z_j}.$$
(5.3)

If ϖ is a coordinate change in a neighborhood intersecting with $\partial \Omega$

$$\begin{cases} x_1 = z_1, \\ \dots \\ x_{n-1} = z_{n-1}, \\ x_n = -z_n, \end{cases}$$
(5.4)

then its Jacobian matrix is

$$J := \begin{pmatrix} \frac{\partial x_1}{\partial z_1} & \cdots & \frac{\partial x_1}{\partial z_{n-1}} & \frac{\partial x_1}{\partial z_n} \\ \vdots & \ddots & \vdots & \vdots \\ \frac{\partial x_{n-1}}{\partial z_1} & \cdots & \frac{\partial x_{n-1}}{\partial z_{n-1}} & \frac{\partial x_{n-1}}{\partial z_n} \\ \frac{\partial x_n}{\partial z_1} & \cdots & \frac{\partial x_n}{\partial z_{n-1}} & \frac{\partial x_n}{\partial z_n} \end{pmatrix} = \begin{pmatrix} 1 & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 \\ 0 & \cdots & 0 & -1 \end{pmatrix}.$$
(5.5)

Using this and (5.3), we immediately obtain the corresponding metric on the Ω^* : (see [MS-67], or p. 10169, p. 10183 and p. 10187 of [Liu2])

$$g_{jk}(\overset{*}{x}) = -g_{jk}(x)$$
 for $j < k = n$ or $k < j = n$, (5.6)

$$g(x) = g_{jk}(x)$$
 for $j, k < n$ or $j = k = n$, (5.7)

$$g_{jk}(x) = 0$$
 for $j < k = n$ or $k < j = n$ on $\partial\Omega$, (5.8)

where $x_n(x) = -x_n(x)$. We denote such a new (isometric) metric on Ω^* as g^* . It is easy to verify that

$$\begin{bmatrix} g_{11}(x) & \cdots & g_{1,n-1}(x) & -g_{1n}(x) \\ \vdots & \ddots & \vdots & \vdots \\ g_{n-1,1}(x) & \cdots & g_{n-1,n-1}(x) & -g_{n-1,n}(x) \\ -g_{n1}(x) & \cdots & -g_{n,n-1}(x) & g_{nn}(x) \end{bmatrix}^{-1} = \begin{bmatrix} g^{11}(x) & \cdots & g^{1,n-1}(x) & -g^{1n}(x) \\ \vdots & \ddots & \vdots & \vdots \\ g^{n-1,1}(x) & \cdots & g^{n-1,n-1}(x) & -g^{n-1,n}(x) \\ -g^{n1}(x) & \cdots & -g^{n,n-1}(x) & g^{nn}(x) \end{bmatrix},$$

where $[g^{jk}(x)]_{n\times n}$ is the inverse of $[g_{jk}(x)]_{n\times n}$. In addition, by this reflection ϖ , the differential operators $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_{n-1}}, \frac{\partial}{\partial x_n}$ (defined on Ω) are changed to $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_{n-1}}, -\frac{\partial}{\partial x_n}$ (defined on Ω^*), respectively. It is easy to verify that

$$\Gamma_{kl}^{j}(\overset{*}{x}) = a_{jkl} \Gamma_{kl}^{j}(x), \tag{5.9}$$

where

$$a_{jkl} = \begin{cases} 1 & \text{if there is no } n \text{ among } j, k, l, \\ -1 & \text{if there is an } n \text{ among } j, k, l, \\ 1 & \text{if there are two } n \text{ among } j, k, l, \\ -1 & \text{if there are three } n \text{ among } j, k, l, \end{cases}$$

$$\frac{\partial \Gamma_{jk}^s}{\partial x_l}(x)^* = b_{sjkl} \frac{\partial \Gamma_{jk}^s}{\partial x_l}(x), \tag{5.10}$$

where

$$b_{sjkl} = \begin{cases} 1 & \text{if there is no } n \text{ among } s, j, k, l, \\ -1 & \text{if there is an } n \text{ among } s, j, k, l, \\ 1 & \text{if there are two } n \text{ among } s, j, k, l, \\ -1 & \text{if there are three } n \text{ among } s, j, k, l, \\ 1 & \text{if there are four } n \text{ among } s, j, k, l, \end{cases}$$

and

$$R_k^J(\hat{x}) = c_{jk} R_k^J(x), (5.11)$$

where

$$c_{jk} = \begin{cases} 1 & \text{if there no } n \text{ among } j, k, \\ -1 & \text{if there is an } n \text{ among } j, k, \\ 1 & \text{if there are two } n \text{ among } j, k. \end{cases}$$

In what follows, we will let Greek indices run from 1 to n - 1, Roman indices from 1 to n. We define

$$\mathcal{L}_{\lambda,\mu} = \begin{cases} L_{\lambda,\mu} & \text{on } \Omega\\ L_{\lambda,\mu}^{\star} & \text{on } \Omega^*, \end{cases}$$
(5.12)

where

$$L_{\lambda,\mu}^{\star} := L_{\lambda,\mu} \Big(g^{\alpha\beta}(\overset{*}{x}), -g^{\alpha n}(\overset{*}{x}), -g^{n\beta}(\overset{*}{x}), g^{nn}(\overset{*}{x}), a_{jkl} \Gamma_{kl}^{j}(\overset{*}{x}), b_{sjkl} \frac{\partial \Gamma_{jk}^{s}}{\partial x_{l}}(\overset{*}{x}), \qquad (5.13)$$

$$c_{jk} R_{k}^{j}(\overset{*}{x}), \frac{\partial}{\partial x_{1}}, \cdots, \frac{\partial}{\partial x_{n-1}}, -\frac{\partial}{\partial x_{n}} \Big),$$

and $\overset{*}{x} = (x', -x_n) \in \Omega^*$. Roughly speaking, $L_{\lambda,\mu}^*$ is obtained from the expression of $L_{\lambda,\mu}$ by replacing $\frac{\partial}{\partial x_n}$ by $-\frac{\partial}{\partial x_n}$. But we must rewrite such a $L_{\lambda,\mu}^*$ in the terms of the corresponding metric, Christoffel symbols and Ricci curvatures in Ω^* . That is, $L_{\lambda,\mu}^*$ is got if we replace $g^{\alpha\beta}(x), g^{\alpha n}(x), g^{n\beta}(x), g^{nn}(x), \{\Gamma_{kl}^j(x)\}_{1 \le j,k,l \le n}, \{\frac{\partial \Gamma_{jk}^s}{\partial x_l}(x)\}_{1 \le s,j,k,l \le n}, \{R_k^j(x)\}_{1 \le j,k \le n}, \frac{\partial}{\partial x_n}$ by $g^{\alpha\beta}(\overset{*}{x}), -g^{\alpha n}(\overset{*}{x}), -g^{n\beta}(\overset{*}{x}), g^{nn}(\overset{*}{x}), a_{jkl}\Gamma_{kl}^j(\overset{*}{x}), b_{sjkl}\frac{\partial \Gamma_{jk}^s}{\partial x_l}(\overset{*}{x}), c_{jk}R_k^j(\overset{*}{x}), -\frac{\partial}{\partial x_n}$ in $L_{\lambda,\mu} = L_{\lambda,\mu}(g^{\alpha\beta}(x), g^{\alpha n}(x), g^{n\beta}(x), g^{n\beta}(x), g^{\alpha n}(x), \{\Gamma_{kl}^j(x)\}_{1 \le j,k,l \le n}, \{R_k^j(x)\}_{1 \le j,k \le n}, \frac{\partial}{\partial x_1}, \cdots, \frac{\partial}{\partial x_{n-1}}, \frac{\partial}{\partial x_n})$, respectively. Note that we have used the relations (5.6)–(5.11). In view of the metric matrices g and g* have the same order principal minor determinants, we see that $\mathcal{L}_{\lambda,\mu}$ is still a linear elliptic differential operator on closed Riemannian manifold \mathcal{M} .

Let $\mathbf{K}(t, x, y)$ be the fundamental solution of the parabolic system

$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t} + \mathcal{L}_{\lambda,\mu} \mathbf{u} = 0 & \text{ in } (0, +\infty) \times \mathcal{M}, \\ \mathbf{u} = \boldsymbol{\phi} & \text{ on } \{0\} \times \mathcal{M}. \end{cases}$$

That is, for any $t \ge 0$ and $x, y \in \mathcal{M}$,

$$\begin{cases} \frac{\partial \mathbf{K}(t,x,y)}{\partial t} + \mathcal{L}_{\lambda,\mu} \mathbf{K}(t,x,y) = 0 & \text{for } t > 0, x, y \in \mathcal{M}, \\ \mathbf{K}(0,x,y) = \boldsymbol{\delta}(x-y) & \text{for } x, y \in \mathcal{M}. \end{cases}$$
(5.14)

Here the operator $\mathcal{L}_{\lambda,\mu}$ is acted in the third argument *y* of **K**(*t*, *x*, *y*).

Clearly, the coefficients occurring in $\mathcal{L}_{\lambda,\mu}$ jump as *x* crosses the $\partial\Omega$ (since the extended metric *g* is C^0 -smooth on whole \mathcal{M} and C^∞ -smooth in $\mathcal{M} \setminus \partial\Omega$), but $\frac{\partial \mathbf{u}}{\partial t} + \mathcal{L}_{\lambda,\mu}\mathbf{u} = 0$ with $\mathbf{u}(0, x) = \boldsymbol{\phi}(x)$ still has a nice fundamental solution **K** of class $[C^1((0, +\infty) \times \mathcal{M} \times \mathcal{M})]_{n \times n} \cap [C^\infty((0, +\infty) \times (\mathcal{M} \setminus \partial\Omega) \times (\mathcal{M} \setminus \partial\Omega))]_{n \times n}$, approximable even on $\partial\Omega$ by Levi's sum (see [Liu2], or another proof below). Now, let us restrict $x, y \in \Omega$. Recall that for an (elastic) vector field **u** defined in Ω , the boundary traction operator can also be equivalently written as

$$(\mathcal{T}_{\lambda,\mu}\mathbf{u})^{k} := \mu \sum_{l=1}^{n} \left(\nu^{l} \nabla_{l} u^{k} + \nu_{l} \nabla^{k} u^{l} \right) + \lambda \nu^{k} \sum_{l=1}^{n} \nabla_{l} u^{l} \text{ on } \partial\Omega, \quad k = 1, \cdots, n.$$
(5.15)

Write $\mathbf{K}(t, x, y) = (K^{jk}(t, x, y))_{n \times n}$, $\mathbf{K}_j(t, x, y) = (K^{j1}(t, x, y), \dots, K^{jn}(t, x, y), (j = 1, \dots, n))$, and denote

$$\mathcal{T}_{\lambda,\mu}\mathbf{K}(t,x,y) = \left(\mathcal{T}_{\lambda,\mu}\mathbf{K}_{1}(t,x,y),\cdots,\mathcal{T}_{\lambda,\mu}\mathbf{K}_{n}(t,x,y)\right)$$
(5.16)

for all $t > 0, x \in \Omega, y \in \partial \Omega$. More precisely, for each $j = 1, \dots, n$, the *k*-th component of $\mathcal{T}_{\lambda,\mu}\mathbf{K}_j$ is

$$(\mathcal{T}_{\lambda,\mu}\mathbf{K}_j)^k = \mu \sum_{l=1}^n (\nu^l \nabla_l K^{jk} + \nu_l \nabla^k K^{jl}) + \lambda \nu^k \sum_{l=1}^n \nabla_l K^{jl}.$$
(5.17)

It is easy to see that $\frac{\partial (\mathbf{K}(t,x,y) + \mathbf{K}(t,x,y))}{\partial v_y}\Big|_{\partial \Omega} = 0$ for all $t > 0, x \in \Omega$ and $y \in \partial \Omega$. Changing all terms $\frac{\partial K^{jk}(t,x,y)}{\partial v_y}\Big|_{\partial \Omega}$ into 0 in the expression of $2\mathcal{T}_{\lambda,\mu}\mathbf{K}(t,x,y)$ (i.e., replacing all terms $\frac{\partial K^{jk}(t,x,y)}{\partial v_y}\Big|_{\partial\Omega} = \sum_{m=1}^n \frac{\partial K^{jk}(t,x,y)}{\partial y_m} v_m \text{ by 0 in the expression of } 2\mathcal{T}_{\lambda,\mu}\mathbf{K}(t,x,y)\text{), we obtain a matrix-valued function } \mathbf{\Upsilon}(t,x,y) \text{ for } t > 0, x \in \Omega \text{ and } y \in \partial\Omega. \text{ This implies that } \mathbf{\Upsilon}(t,x,y) \text{ only contains the (boundary) tangent derivatives of } K^{jk}(t,x,y) \text{ with respect to } y \in \partial\Omega \text{ (without normal derivative of } K^{jk}(t,x,y)\text{) in the local expression). That is, in local boundary normal coordinates (the inner normal <math>\nu$ of $\partial\Omega$ is in the direction of x_n -axis),

$$\Upsilon(t, x, y) = (\Upsilon_1(t, x, y), \cdots, \Upsilon_n(t, x, y)),$$

$$\begin{split} \Upsilon_{j}(t,x,y) \\ &= 2\mu \begin{pmatrix} 2\frac{\partial K^{j1}}{\partial x_{1}} + 2\sum_{m=1}^{n} \Gamma_{1m}^{1} K^{jm} & \cdots & \frac{\partial K^{j1}}{\partial x_{n-1}} + \frac{\partial K^{jn-1}}{\partial x_{1}} + \sum_{m=1}^{n} (\Gamma_{n-1,m}^{1} + \Gamma_{1m}^{n-1}) K^{jm} & \frac{\partial K^{jn}}{\partial x_{1}} + \sum_{m=1}^{n} (\Gamma_{nm}^{1} + \Gamma_{1m}^{n}) K^{jm} \\ & \cdots & \cdots & \cdots & \cdots \\ \frac{\partial K^{jn-1}}{\partial x_{1}} + \frac{\partial K^{j1}}{\partial x_{n-1}} + \sum_{m=1}^{n} (\Gamma_{1m}^{n-1} + \Gamma_{n-1,m}^{1}) K^{jm} & \cdots & 2\frac{\partial K^{jn-1}}{\partial x_{n-1}} + 2\sum_{m=1}^{n} \Gamma_{n-1,m}^{n-1} K^{jm} & \frac{\partial K^{jn}}{\partial x_{n-1}} + \sum_{m=1}^{n} (\Gamma_{nm}^{n-1} + \Gamma_{n-1,m}^{n}) K^{jm} \\ \frac{\partial K^{jn}}{\partial x_{1}} + \sum_{m=1}^{n} (\Gamma_{1m}^{n} + \Gamma_{1m}^{1}) K^{jm} & \cdots & \frac{\partial K^{jn}}{\partial x_{n-1}} + (\Gamma_{n-1,m}^{n} + \Gamma_{nm}^{n-1}) K^{jm} & 2\sum_{m=1}^{n} \Gamma_{nm}^{n} K^{jm} \end{pmatrix} \begin{pmatrix} v_{1} \\ \vdots \\ v_{n} \end{pmatrix} \\ &+ 2\lambda \Big(\frac{\partial K^{j1}}{\partial x_{1}} + \sum_{m=1}^{n} \Gamma_{1m}^{1} K^{jm} + \cdots + \frac{\partial K^{j,n-1}}{\partial x_{n-1}} + \sum_{m=1}^{n} \Gamma_{n-1,m}^{n-1} K^{jm} + \sum_{m=1}^{n} \Gamma_{nm}^{n} K^{jm} \Big) \begin{pmatrix} v_{1} \\ \vdots \\ v_{n} \end{pmatrix}, \quad j = 1, \cdots, n \end{split}$$

It is easy to see that $\Upsilon(t, x, y)$ is a continuous (matrix-valued) function for all $t > 0, x \in \Omega$ and $y \in \partial \Omega$. Further, for any fixed $x \in \Omega$ and any $y \in \partial \Omega$, since $x \neq y$ we see that

$$\lim_{t \to 0^+} \mathbf{K}(t, x, y) = 0, \tag{5.18}$$

which implies

$$\lim_{t \to 0^+} \frac{\partial \mathbf{K}(t, x, y)}{\partial T}\Big|_{\partial \Omega} = 0 \quad \text{for} \quad x \in \Omega, \quad y \in \partial \Omega,$$
(5.19)

where $\frac{\partial}{\partial T}$ denotes the tangent derivative along the boundary $\partial \Omega$ in variable y. From (5.18)–(5.19) we get

$$\lim_{t \to 0^+} \nabla_l \mathbf{K}(t, x, y) \Big|_{\partial \Omega} = 0 \quad \text{for all} \ x \in \Omega, \ y \in \partial \Omega, \quad 1 \le l < n,$$

so that

$$\lim_{t \to 0^+} \Upsilon(t, x, y) = 0 \quad \text{for any } x \in \Omega, \ y \in \partial\Omega,$$
(5.20)

where $\nabla_l \mathbf{K} = (\nabla_l \mathbf{K}_1, \dots, \nabla_l \mathbf{K}_n)$ and $\nabla_l K^{jk} := \frac{\partial K^{jk}(t,x,y)}{\partial y_l} + \sum_{m=1}^n \Gamma_{lm}^k K^{jm}(t,x,y), (1 \le j,k,l \le n).$ Let $\mathbf{H}(t,x,y)$ be the solution of

$$\begin{cases} \frac{\partial \mathbf{u}(t,x,y)}{\partial t} = L_{\lambda,\mu} \mathbf{u}(t,x,y) & \text{for } t > 0, \ x, y \in \Omega, \\ 2\mu (\text{Def } \mathbf{u}(t,x,y))^{\#} \mathbf{v} + \lambda (\text{div } (\mathbf{u}(t,x,y))) \mathbf{v} = \mathbf{\Upsilon}(t,x,y) & \text{for } t > 0, \ x \in \Omega, \ y \in \partial\Omega, \\ \mathbf{u}(0,x,y) = \mathbf{0} & \text{for } x, y \in \Omega. \end{cases}$$

From (5.20), we get that the above parabolic system satisfy the compatibility condition. Thus, the matrix-valued solution $\mathbf{H}(t, x, y)$, is smooth in $(0, \infty) \times \Omega \times \Omega$ and continuous on $[0, \infty) \times \overline{\Omega} \times \overline{\Omega}$. Then there exists a constant C > 0 such that

$$|\mathbf{H}(t, x, y)| \le C$$
 for all $0 \le t \le 1$, $x, y \in \overline{\Omega}$,

Actually, we have $\lim_{t\to 0^+} \int_{\Omega} \text{Tr } \mathbf{H}(t, x, x) \, dx = 0$, and hence, for dimensions $n \ge 2$,

$$\int_{\Omega} \operatorname{Tr} \mathbf{H}(t, x, x) \, dx = o(nC(\operatorname{vol}(\Omega))) = o(t^{-\frac{n-2}{2}}) \quad \text{as} \quad t \to 0^+.$$
(5.21)

(5.21) implies that we can precisely get the first three terms asymptotic expansion. It can easily be verified that

$$\mathbf{K}^{(D)}(t, x, y) = \mathbf{K}(t, x, y) - \mathbf{K}(t, x, \dot{y}),$$

$$\mathbf{K}^{(T)}(t, x, y) = \mathbf{K}(t, x, y) + \mathbf{K}(t, x, \dot{y}) - \mathbf{H}(t, x, y),$$

are the Green functions of

$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t} + L_{\lambda,\mu} \mathbf{u} = 0 & \text{ in } (0, +\infty) \times \Omega \\ \mathbf{u} = \boldsymbol{\phi} & \text{ on } \{0\} \times \Omega \end{cases}$$

with zero Dirichlet and zero traction (i.e., free) boundary conditions, respectively, where $y = (y', y_n), y_n \ge 0$, and $\overset{*}{y} := \varsigma(y', y_n) = (y', -y_n)$. In other words,

$$\begin{cases} \frac{\partial \mathbf{K}^{(D)}(t,x,y)}{\partial t} + L_{\lambda,\mu} \mathbf{K}^{(D)}(t,x,y) = 0, \quad t > 0, \ x, \ y \in \Omega, \\ \mathbf{K}^{(D)}(t,x,y) = 0, \quad t > 0, \ x \in \Omega, \ y \in \partial\Omega, \\ \mathbf{K}^{(D)}(0,x,y) = \boldsymbol{\delta}(x-y), \quad x, y \in \Omega \end{cases}$$

and

$$\begin{aligned} \frac{\partial \mathbf{K}^{(T)}(t,x,y)}{\partial t} + L\mathbf{K}^{(T)}(t,x,y) &= 0, \quad t > 0, \ x, \ y \in \Omega, \\ \mathcal{T}_{\lambda,\mu}(\mathbf{K}^{(T)}(t,x,y)) &= 0, \quad t > 0, \ x \in \Omega, \ y \in \partial\Omega, \\ \mathbf{K}^{(T)}(0,x,y) &= \delta(x-y), \quad x, \ y \in \Omega, \end{aligned}$$

where $\mathcal{T}_{\lambda,\mu}(\mathbf{K}^{(T)}(t, x, y))$ is the traction of $\mathbf{K}^{(T)}(t, x, y)$ on $\partial\Omega$. By combining the fact $\frac{\partial(\mathbf{K}(t, x, y) + \mathbf{K}(t, x, y))}{\partial v_y}\Big|_{\partial\Omega} = 0$ and $\mathcal{T}_{\lambda,\mu}(\mathbf{K}(t, x, y) + \mathbf{K}(t, x, y)) = \Upsilon(t, x, y) = \mathcal{T}_{\lambda,\mu}\mathbf{H}(t, x, y)$ for $t > 0, x \in \Omega$ and $y \in \partial\Omega$, we get $\mathcal{T}_{\lambda,\mu}\mathbf{K}^{(T)}(t, x, y) = 0$ for all $t > 0, x \in \Omega$ and $y \in \partial\Omega$. Now, we will show $\mathbf{K}^{(D)}(t, x, y)$ and $\mathbf{K}^{(T)}(t, x, y)$ satisfy the associated parabolic system. In fact, for any $t > 0, x, y \in \Omega$, we have $L_{\lambda,\mu}\mathbf{K}(t, x, y) = \mathcal{L}_{\lambda,\mu}\mathbf{K}(t, x, y)$, so that

$$\begin{cases} \left(\frac{\partial}{\partial t} + L_{\lambda,\mu}\right) \mathbf{K}(t, x, y) = \left(\frac{\partial}{\partial t} + \mathcal{L}_{\lambda,\mu}\right) \mathbf{K}(t, x, y) = 0, \\ \mathbf{K}(0, x, y) = \boldsymbol{\delta}(x - y) \end{cases}$$
(5.22)

by (5.14). Noting that the Jacobian matrix of the reflection ϖ is *J* (see (5.5)), it follows from chain rule that for any fixed t > 0 and $x \in \Omega$, and any $y = (y', y_n) \in \Omega$,

 $\left[L_{\lambda,\mu}(\mathbf{K}(t, x, \overset{*}{y}))\right]$ evaluated at the point y

$$\begin{split} &= \left[L_{\lambda,\mu}(\mathbf{K}(t,x,\varsigma(y',y_n))) \right] \Big|_{\text{evaluated at the point } (y',y_n) \\ &= \left[L_{\lambda,\mu}(\mathbf{K}(t,x,(y',-y_n))) \right] \Big|_{\text{evaluated at the point } (y',y_n) \\ &= \left\{ \left[L_{\lambda,\mu}(g^{\alpha\beta}(y),g^{\alpha n}(y),g^{n\beta}(x),g^{nn}(x),\{\Gamma_{kl}^{j}(x)\}_{1\leq j,k,l\leq n},\left\{\frac{\partial\Gamma_{jk}^{s}}{\partial x_{l}}(x)\right\}_{1\leq s,j,k,l\leq n}, \right. \\ &\left. \left\{ R_{k}^{j}(x) \right\}_{1\leq j,k\leq n}, \frac{\partial}{\partial y_{1}}, \cdots, \frac{\partial}{\partial y_{n-1}}, \frac{\partial}{\partial y_{n}} \right\} \right] \mathbf{K}(t,x,(y',-y_n)) \right\} \Big|_{\text{evaluated at } (y',y_n)} \\ &= \left\{ \left[L_{\lambda,\mu}(g^{\alpha\beta}(\overset{*}{y}), -g^{\alpha n}(\overset{*}{y}), -g^{n\beta}(\overset{*}{y}), g^{nn}(\overset{*}{y}), \{a_{jkl}\Gamma_{kl}^{j}(\overset{*}{x})\}_{1\leq j,k,l\leq n}, \{b_{sjkl}\frac{\partial\Gamma_{jk}^{s}}{\partial x_{l}}(\overset{*}{x})\}_{1\leq s,j,k,l\leq n}, \right. \\ &\left. \left\{ c_{jk}R_{k}^{j}(\overset{*}{x}) \right\}_{1\leq j,k\leq n}, \frac{\partial}{\partial y_{1}}, \cdots, \frac{\partial}{\partial y_{n-1}}, -\frac{\partial}{\partial y_{n}} \right\} \right] \mathbf{K}(t,x,y) \right\} \Big|_{\text{evaluated at } \overset{*}{y} = (y', -y_{n})} \\ &= \left. L_{\lambda,\mu}^{\star}(\mathbf{K}(t,x,\overset{*}{y})) \right|_{\text{evaluated at the point } \overset{*}{y} = (y', -y_{n})}. \end{split}$$

That is, the action of $L_{\lambda,\mu}$ to $\mathbf{K}(t, x, \overset{*}{y})$ (regarded as a vector-valued function of y) at the point $y = (y', y_n)$ is just the action of $L_{\lambda,\mu}^{\star}$ to $\mathbf{K}(t, x, \overset{*}{y})$ (regarded as a vector-valued of $\overset{*}{y}$) at the point $\overset{*}{y} = (y', -y_n)$. Because of $\varsigma(y', y_n) = (y', -y_n) \in \Omega^*$, we see

 $L_{\lambda,\mu}^{\star}(\mathbf{K}(t, x, \overset{*}{y}))\Big|_{\text{evaluated at the point }\overset{*}{y} = (y', -y_n)} = \mathcal{L}_{\lambda,\mu}(\mathbf{K}(t, x, \overset{*}{y}))\Big|_{\text{evaluated at the point }\overset{*}{y} = (y', -y_n)}.$

For any t > 0, $x \in \Omega$ and $(y', -y_n) \in \Omega^*$, we have

$$\left(\frac{\partial}{\partial t} + \mathcal{L}_{\lambda,\mu}\right) (\mathbf{K}(t, x, (y', -y_n))) = 0.$$

In addition, $\mathbf{K}(t, x, (y', -y_n)) = \mathbf{K}(t, x, \varsigma(y))$ for any $t > 0, x, y \in \Omega$. By virtue of $x \neq (y', -y_n)$, this leads to $\mathbf{K}(0, x, (y', -y_n)) = 0$ and

$$\left(\frac{\partial}{\partial t} + L_{\lambda,\mu}\right)$$
 (**K** $(t, x, (y', -y_n))) = 0$ for any $t > 0, x \in \Omega$ and $(y', -y_n) \in \Omega^*$,

i.e.,

$$\begin{cases} \left(\frac{\partial}{\partial t} + L_{\lambda,\mu}\right) \mathbf{K}(t, x, \overset{*}{y}) = 0 & \text{for any } t > 0, x \in \Omega \text{ and } \overset{*}{y} \in \Omega^{*}, \\ \mathbf{K}(0, x, \overset{*}{y}) = 0 & \text{for any } x, y \in \Omega. \end{cases}$$
(5.23)

Combining (5.22) and (5.23), we obtain that

$$\begin{cases} \left(\frac{\partial}{\partial t} + L_{\lambda,\mu}\right) \left(\mathbf{K}(t,x,y) \mp \mathbf{K}(t,x,\overset{*}{y})\right) = 0 & \text{for any } t > 0, x, y \in \Omega, \\ \mathbf{K}(0,x,y) \mp \mathbf{K}(0,x,\overset{*}{y}) = \delta(x-y) & \text{for any } x, y \in \Omega. \end{cases}$$
(5.24)

K(*t*, *x*, *y*) is *C*¹-smooth with respect to *y* in *M* for any fixed *t* > 0 and *x* ∈ Ω (see below), so does it on the hypersurface ∂ Ω. Therefore, we get that **K**^(*D*)(*t*, *x*, *y*) (respectively **K**^(*T*)(*t*, *x*, *y*)) is the Green function in Ω with the Dirichlet (respectively, free) boundary condition on ∂ Ω.

To show C^1 -regularity of the fundamental solution $\mathbf{K}(t, x, y)$, it suffices to prove $C_{loc}^{1,1+\alpha}$ -regularity for a $W_2^{1,2}$ strong solution \mathbf{u} of the parabilic system $(\frac{\partial}{\partial t} + \mathcal{L}_{\lambda,\mu})\mathbf{u} = 0$ in $(0, +\infty) \times \mathcal{M}$, where $W_p^{1,2} := {\mathbf{u} | \mathbf{u}, \frac{\partial \mathbf{u}}{\partial t}, D\mathbf{u}, D^2\mathbf{u} \in L^p}$. When the coefficients of an elliptic system are smooth on both sides of an (n - 1)-dimensional hypersurface (may be discontinuous crossing this hypersurface), the corresponding $C^{1,1+\alpha}$ -regularity for solutions of a parabolic equation system is a special case of Dong's result (see Theorem 4 of p. 141 in [Don]). In fact, Dong in [Don] has given regularity results to the strong solutions for parabolic equation with more general coefficients. This type of system arises from the problems of linearly elastic laminates and composite materials (see, for example, [CKVC], [LiVo], [LiNi], [Don] and [Xi-11]).

Let us discuss the parabolic (elastic) system in more detail. In fact, on p. 10177 in [Liu2], we have expressed the Lamé operator $L_{\lambda,\mu}$ in Ω as the form of components relative to local coordinates:

$$L_{\lambda,\mu}\mathbf{u} = \left\{ -\mu \Big(\sum_{m,l=1}^{n} g^{ml} \frac{\partial^{2}}{\partial x_{m} \partial x_{l}} \Big) \mathbf{I}_{n} - (\mu + \lambda) \begin{bmatrix} \sum_{m=1}^{n} g^{1m} \frac{\partial^{2}}{\partial x_{m} \partial x_{1}} & \cdots & \sum_{m=1}^{n} g^{1m} \frac{\partial^{2}}{\partial x_{m} \partial x_{n}} \\ \vdots & \vdots & \vdots \\ \sum_{m=1}^{n} g^{mn} \frac{\partial^{2}}{\partial x_{m} \partial x_{1}} & \cdots & \sum_{m=1}^{n} g^{mm} \frac{\partial^{2}}{\partial x_{m} \partial x_{n}} \end{bmatrix} \\ +\mu \Big(\sum_{m,l=1}^{n} g^{ml} \Gamma_{ml}^{sl} \frac{\partial}{\partial x_{s}} \Big) \mathbf{I}_{n} - \mu \begin{bmatrix} \sum_{m,l=1}^{n} 2g^{ml} \Gamma_{1m}^{l} \frac{\partial}{\partial x_{l}} & \cdots & \sum_{m=1}^{n} 2g^{ml} \Gamma_{nm}^{l} \frac{\partial}{\partial x_{l}} \\ \vdots & \vdots \\ \sum_{m,l=1}^{n} 2g^{ml} \Gamma_{1m}^{l} \frac{\partial}{\partial x_{m}} & \cdots & \sum_{m,l=1}^{n} 2g^{ml} \Gamma_{nm}^{l} \frac{\partial}{\partial x_{m}} \end{bmatrix} \\ -(\mu + \lambda) \begin{bmatrix} \sum_{m,l=1}^{n} g^{lm} \Gamma_{1l}^{l} \frac{\partial}{\partial x_{m}} & \cdots & \sum_{m,l=1}^{n} g^{lm} \Gamma_{nl}^{l} \frac{\partial}{\partial x_{m}} \\ \vdots & \vdots \\ \sum_{m,l=1}^{n} 2g^{ml} \Gamma_{1m}^{l} \frac{\partial}{\partial x_{m}} & \cdots & \sum_{m,l=1}^{n} g^{mm} \Gamma_{nl}^{l} \frac{\partial}{\partial x_{m}} \end{bmatrix} \\ -\mu \begin{bmatrix} \sum_{l,m=1}^{n} g^{ml} (\frac{\partial\Gamma_{1l}^{l}}{\partial x_{m}} + \Gamma_{hl}^{l} \Gamma_{lm}^{h} - \Gamma_{lh}^{l} \Gamma_{ml}^{h} \Big) & \cdots & \sum_{l,m=1}^{n} g^{ml} (\frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} + \Gamma_{hl}^{l} \Gamma_{nm}^{h} - \Gamma_{nh}^{l} \Gamma_{ml}^{h}) \\ \vdots & \vdots \\ \sum_{m,l=1}^{n} g^{ml} (\frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} + \Gamma_{hl}^{l} \Gamma_{lm}^{h} - \Gamma_{lh}^{l} \Gamma_{ml}^{h} \Big) & \cdots & \sum_{l,m=1}^{n} g^{ml} (\frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} + \Gamma_{ml}^{l} \Gamma_{nm}^{h} - \Gamma_{nh}^{l} \Gamma_{ml}^{h}) \\ -(\mu + \lambda) \begin{bmatrix} \sum_{l,m=1}^{n} g^{ml} \frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} & \cdots & \sum_{l,m=1}^{n} g^{mm} \frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} \\ \vdots & \vdots \\ \sum_{l,m=1}^{n} g^{ml} \frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} & \cdots & \sum_{l,m=1}^{n} g^{mm} \frac{\partial\Gamma_{nl}^{l}}{\partial x_{m}} \end{bmatrix} \\ -\mu \begin{bmatrix} R_{1}^{l} & \cdots & R_{n}^{l} \\ \vdots & \vdots \\ R_{1}^{n} & \cdots & R_{n}^{l} \\ \end{bmatrix} \begin{bmatrix} \mu^{l} \\ \vdots \\ \mu^{l} \end{bmatrix} , \end{bmatrix}$$

where \mathbf{I}_n is the $n \times n$ identity matrix. The above expression $L_{\lambda,\mu}\mathbf{u}$ will played a key role in our proof. From the local expression (5.25) of $L_{\lambda,\mu}$, we see that the top-order coefficients of $\mathcal{L}_{\lambda,\mu}$ are not "too bad" since only the first (n-1) coefficients of the *n*-th column in the

second matrix in $L_{\lambda,\mu}$ defined on Ω are changed their signs in $L_{\lambda,\mu}^{\star}$ at the reflection points of Ω^* . In other words, only $(\sum_{m=1}^n g^{1m}(x) \frac{\partial^2 u^n}{\partial x_m \partial x_n}, \dots, \sum_{m=1}^n g^{n-1,m}(x) \frac{\partial^2 u^n}{\partial x_m \partial x_n})^T$ in Ω is changed into $(-\sum_{m=1}^n g^{1m}(x^*) \frac{\partial^2 u^n}{\partial x_m \partial x_n}, \dots, -\sum_{m=1}^n g^{n-1,m}(x^*) \frac{\partial^2 u^n}{\partial x_m \partial x_n})^T$ in Ω^* in the second term in $\mathcal{L}_{\lambda,\mu}$ (see (1.5) of $L_{\lambda,\mu}$). For any small coordinate chart $V \subset \mathcal{M}$, if $V \subset \mathcal{M} \setminus (\partial \Omega)$, then the solution **u** belongs to $[C^{\infty}((0, +\infty) \times V)]^n \cap [W_2^{1,2}((0, +\infty) \times V)]^n$ since the coefficients of parabolic system

$$\left(\frac{\partial}{\partial t} + \mathcal{L}_{\lambda,\mu}\right)\mathbf{u} = 0 \tag{5.26}$$

are smooth in *V*. If the coordinate chart $V \,\subset \,\mathcal{M}$ and $V \cap \partial\Omega \neq \emptyset$, then we can find a (local) diffeomorphism Ψ such that $\Psi(V) = U \subset \mathbb{R}^n$ and $\partial\Omega$ is mapped onto $U \cap \{x \in \mathbb{R}^n | x_n = 0\}$ (i.e., $V \cap \partial\Omega$ is flatten into hyperplane $\{x \in \mathbb{R}^n | x_n = 0\}$ by Ψ). By this coordinate transformation Ψ , the parabolic system (5.26) is changed into another parabolic system $(\frac{\partial}{\partial t} - L_{\lambda,\mu})\mathbf{v} = 0$, whose coefficients are smooth on both sides of (n - 1)-dimensional hyperplane $\{x \in \mathbb{R}^n | x_n = 0\}$ (may be discontinuous crossing this hyperplane, i.e., the coefficients of $L_{\lambda,\mu}$ have jump only on this hyperplane). It follows from Dong's regularity result (Theorem 4 of [Don]) that $\mathbf{v} \in [C^{1,1+\alpha}((0, +\infty) \times U)]^n$, so that $\mathbf{u} \in [C^{1,1+\alpha}((0, +\infty) \times V)]^n$. Note that \mathcal{M} is a compact closed Riemannian manifold. Thus we find from the above discussion and (global) geometric analysis technique that for any initial value $\phi \in [C^{\infty}(\mathcal{M})]^n$, there exists a (global) strong solution in $[W_2^{1,2}((0, +\infty) \times \mathcal{M})]^n$ which is in $[C^{1,1+\alpha}((0, +\infty) \times \mathcal{M})]^n$. Of course, this result also holds to our case for $C^{1,1+\alpha}$ -regularity of the fundamental solution $\mathbf{K}(t, x, y)$ on $(0, +\infty) \times \mathcal{M} \times \mathcal{M}$ (see [Liu2]).

Therefore, the integral kernels $\mathbf{K}^{(D)/(T)}(t, x, y)$ of $\frac{\partial \mathbf{u}}{\partial t} + L_{\lambda,\mu}^{(D)/(T)}\mathbf{u} = 0$ can be expressed on $(0, \infty) \times \Omega \times \Omega$ as

$$\mathbf{K}^{(D)}(t, x, y) = \mathbf{K}(t, x, y) - \mathbf{K}(t, x, y^{*}),$$
(5.27)

$$\mathbf{K}^{(T)}(t, x, y) = \mathbf{K}(t, x, y) + \mathbf{K}(t, x, y) - \mathbf{H}(t, x, y)$$
(5.28)

^{*} being the double of $y \in \Omega$ (see, p. 53 of [MS-67]). Since the strongly continuous semigroup $(e^{-t\mathcal{L}_{\lambda,\mu}})_{t\geq 0}$ can also be represented as

$$e^{-t\mathcal{L}_{\lambda,\mu}} = \frac{1}{2\pi i} \int_{C} e^{-t\tau} (\tau I - \mathcal{L}_{\lambda,\mu})^{-1} d\tau,$$

where *C* is a suitable curve in the complex plane in the positive direction around the spectrum of $\mathcal{L}_{\lambda,\mu}$ (i.e., a contour around the positive real axis). It follows that

$$\mathbf{K}(t,x,y) = e^{-t\mathcal{L}_{\lambda,\mu}}\delta(x-y) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-y)\cdot\xi} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \iota((\tau I - \mathcal{L}_{\lambda,\mu})^{-1}) d\tau\right) d\xi, \quad \forall t > 0, \ x, y \in \mathcal{M},$$

where $\iota(A)$ denotes the full symbol of a pseudodifferential operator A.

We claim that

$$\frac{1}{2\pi i} \int_{C} (\tau I - \mathcal{L}_{\lambda,\mu})^{-1} e^{-t\tau} \,\delta(x-y) \,d\tau = \frac{1}{2\pi i} \int_{C} \Big(\int_{\mathbb{R}^n} e^{i(x-y)\cdot\xi} \sum_{j\leq -2} \mathbf{q}_j(x,\xi,\tau) \,d\xi \Big) e^{-t\tau} d\tau.$$
(5.29)

In fact, for any smooth vector-valued function ϕ with compact support we have

$$(e^{-t\mathcal{L}_{\lambda,\mu}}\boldsymbol{\phi})(x) = \left(\frac{1}{2\pi i}\int_{C}e^{-t\tau}(\tau I - \mathcal{L}_{\lambda,\mu})^{-1}d\tau\right)\boldsymbol{\phi}(x)$$
$$= \frac{1}{2\pi i}\int_{C}e^{-t\tau}\left(\int_{\mathbb{R}^{n}}e^{ix\cdot\xi}\sum_{j\leq -2}\mathbf{q}_{j}(x,\xi,\tau)\hat{\boldsymbol{\phi}}(\xi)\,d\xi\right)d\tau.$$

On the one hand, from the left-hand side of (5.29), we get

$$\int_{\mathbb{R}^{n}} \left[\left(\frac{1}{2\pi i} \int_{C} (\tau I - \mathcal{L}_{\lambda,\mu})^{-1} e^{-t\tau} d\tau \right) (\delta(x-y)) \right] \phi(y) dy \qquad (5.30)$$
$$= \left(\frac{1}{2\pi i} \int_{C} (\tau - \mathcal{L}_{\lambda,\mu})^{-1} e^{-t\tau} d\tau \right) \phi(x) = e^{-t\mathcal{L}_{\lambda,\mu}} \phi(x).$$

On the other hand, from the right-hand side of (5.29) we obtain

$$\int_{\mathbb{R}^{n}} \left[\frac{1}{2\pi i} \int_{C} \left(\int_{\mathbb{R}^{n}} e^{i(x-y)\cdot\xi} \sum_{j\leq-2} \mathbf{q}_{j}(x,\xi,\tau) d\xi \right) e^{-t\tau} d\tau \right] \boldsymbol{\phi}(y) dy$$

$$= \frac{1}{2\pi i} \int_{C} \left(\int_{\mathbb{R}^{n}} e^{ix\cdot\xi} \sum_{j\leq-2} \mathbf{q}_{j}(x,\xi,\tau) d\xi \right) e^{-t\tau} d\tau \int_{\mathbb{R}^{n}} e^{-y\cdot\xi} \boldsymbol{\phi}(y) dy$$

$$= \frac{1}{2\pi i} \int_{C} \left(\int_{\mathbb{R}^{n}} e^{ix\cdot\xi} \sum_{j\leq-2} \mathbf{q}_{j}(x,\xi,\tau) \hat{\boldsymbol{\phi}}(\xi) d\xi \right) e^{-t\tau} d\tau = e^{-t\mathcal{L}_{\lambda,\mu}} \boldsymbol{\phi}(x).$$
(5.31)

Thus, the desired identity (5.29) is asserted by (5.30) and (5.31).

In particular, for every t > 0 and $x \in \Omega$,

$$\begin{aligned} \mathbf{K}(t,x,x) &= e^{-t\mathcal{L}_{\lambda\mu}} \delta(x-x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \iota((\tau I - \mathcal{L}_{\lambda,\mu})^{-1}) d\tau \right) d\xi \end{aligned} \tag{5.32} \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \iota((\tau I - L_{\lambda,\mu})^{-1}) d\tau \right) d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \sum_{l \ge 0} q_{-2-l}(x,\xi,\tau) d\tau \right) d\xi, \end{aligned}$$
$$\mathbf{K}(t,x,\overset{*}{x}) &= e^{-t\mathcal{L}_{\lambda\mu}} \delta(x-\overset{*}{x}) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-\overset{*}{x})\cdot\xi} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \iota((\tau I - \mathcal{L}_{\lambda,\mu})^{-1}) d\tau \right) d\xi \tag{5.33} \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-\overset{*}{x})\cdot\xi} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \iota((\tau I - \mathcal{L}_{\lambda,\mu})^{-1}) d\tau \right) d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-\overset{*}{x})\cdot\xi} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \iota((\tau I - \mathcal{L}_{\lambda,\mu})^{-1}) d\tau \right) d\xi, \end{aligned}$$

where $\sum_{l\geq 0} \mathbf{q}_{-2-l}(x,\xi,\tau)$ is the full symbol of $(\tau I - L_{\lambda,\mu})^{-1}$.

Firstly, from the discussion on p. 10182 of [Liu2], we know that

$$\mathbf{q}_{-2}(x,\xi,\tau) = \frac{1}{\tau - \mu \sum_{l,m=1}^{n} g^{lm} \xi_l \xi_m} \mathbf{I}_n$$
(5.34)

$$+\frac{\mu+\lambda}{(\tau-\mu\sum_{l,m=1}^{n}g^{lm}\xi_{l}\xi_{m})(\tau-(2\mu+\lambda)\sum_{l,m=1}^{n}g^{lm}\xi_{l}\xi_{m})}\left[\begin{array}{ccc}\sum_{r=1}^{n}g^{1r}\xi_{r}\xi_{1}&\cdots&\sum_{r=1}^{n}g^{1r}\xi_{r}\xi_{n}\\\vdots&\vdots\\\sum_{r=1}^{n}g^{nr}\xi_{r}\xi_{1}&\cdots&\sum_{r=1}^{n}g^{nr}\xi_{r}\xi_{n}\end{array}\right]$$

and

$$\operatorname{Tr}\left(\mathbf{q}_{-2}(x,\xi,\tau)\right) = \frac{n}{(\tau - \mu \sum_{l,m=1}^{n} g^{lm} \xi_{l} \xi_{m})} + \frac{(\mu + \lambda) \sum_{l,m=1}^{n} g^{lm} \xi_{l} \xi_{m}}{(\tau - \mu \sum_{l,m=1}^{n} g^{lm} \xi_{l} \xi_{m})(\tau - (2\mu + \lambda) \sum_{l,m=1}^{n} g^{lm} \xi_{l} \xi_{m})}.$$
(5.35)

For each $x \in \Omega$, we use a geodesic normal coordinate system centered at this *x*. It follows from §11 of Chap.1 in [Ta1] that in such a coordinate system, $g_{jk}(x) = \delta_{jk}$ and $\Gamma_{jk}^{l}(x) = 0$. Then (5.35) reduces to

$$\operatorname{Tr}\left(\mathbf{q}_{-2}(x,\xi,\tau)\right) = \frac{n}{(\tau-\mu|\xi|^2)} + \frac{(\mu+\lambda)|\xi|^2}{(\tau-\mu|\xi|^2)(\tau-(2\mu+\lambda)|\xi|^2)},\tag{5.36}$$

where $|\xi| = \sqrt{\sum_{k=1}^{n} \xi_k^2}$ for any $\xi \in \mathbb{R}^n$. By applying the residue theorem (see, for example, Chap. 4, §5 in [Ahl]) we get

$$\frac{1}{2\pi i} \int_{C} e^{-t\tau} \left(\frac{n}{(\tau - \mu|\xi|^2)} + \frac{(\mu + \lambda)|\xi|^2}{(\tau - \mu|\xi|^2)(\tau - (2\mu + \lambda)|\xi|^2)} \right) d\tau = (n - 1)e^{-t\mu|\xi|^2} + e^{-t(2\mu + \lambda)|\xi|^2}.$$
 (5.37)

It follows that

$$\frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} \left(\frac{1}{2\pi i} \int_{C} e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x,\xi,\tau) \right) d\tau \right) d\xi \qquad (5.38)$$

$$= \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} \left((n-1)e^{-t\mu|\xi|^{2}} + e^{-t(2\mu+\lambda)|\xi|^{2}} \right) d\xi$$

$$= \frac{n-1}{(4\pi\mu t)^{n/2}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}},$$

and hence

$$\int_{\Omega} \left\{ \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} \left(\frac{1}{2\pi i} \int_{C} e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x,\xi,\tau) \right) d\tau \right) d\xi \right\} dV$$

$$= \left(\frac{n-1}{(4\pi\mu t)^{n/2}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} \right) \operatorname{Vol}(\Omega).$$
(5.39)

In the above discussion, if we replace $x \in \Omega$ by $\overset{*}{x} \in \Omega^*$, then (5.34) will become

$$\mathbf{q}_{-2}(\overset{*}{x},\xi,\tau) = \frac{1}{\tau - \mu \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m}} \mathbf{I}_{n} + \frac{\mu + \lambda}{(\tau - \mu \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m})(\tau - (2\mu + \lambda) \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m})}$$

G. Q. Liu / Eigenvalue inequalities and three-term asymptotic formulas of the heat traces

$$\times \begin{bmatrix} \sum_{r=1}^{n} (g^{1r}(\overset{*}{x}))\xi_{r}\xi_{1} & \cdots & \sum_{r=1}^{n} (g^{1r}(\overset{*}{x}))\xi_{r}\xi_{n-1} & \sum_{r=1}^{n} (-g^{1r}(\overset{*}{x}))\xi_{r}\xi_{n} \\ \vdots & \vdots & \vdots \\ \sum_{r=1}^{n} (g^{n-1,r}(\overset{*}{x}))\xi_{r}\xi_{1} & \cdots & \sum_{r=1}^{n} (g^{n-1,r}(\overset{*}{x}))\xi_{r}\xi_{n-1} & \sum_{r=1}^{n} (-g^{n-1,r}(\overset{*}{x}))\xi_{r}\xi_{n} \\ \sum_{r=1}^{n} (-g^{nr}(\overset{*}{x}))\xi_{r}\xi_{1} & \cdots & \sum_{r=1}^{n} (-g^{nr}(\overset{*}{x}))\xi_{r}\xi_{n-1} & \sum_{r=1}^{n} (g^{nr}(\overset{*}{x}))\xi_{r}\xi_{n} \end{bmatrix}$$

and

$$\operatorname{Tr}\left(\mathbf{q}_{-2}(\overset{*}{x},\xi,\tau)\right) = \frac{n}{(\tau - \mu \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m})} + \frac{(\mu + \lambda) \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m}}{(\tau - \mu \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m})(\tau - (2\mu + \lambda) \sum_{l,m=1}^{n} (g^{lm}(\overset{*}{x}))\xi_{l}\xi_{m})}$$

This implies that all expressions (5.35)–(5.39) of the above trace symbols have the same form either in Ω or in Ω^* .

For given (small) $\epsilon > 0$, denote by $U_{\epsilon}(\partial \Omega) = \{z \in \mathcal{M} | \operatorname{dist}(z, \partial \Omega) < \epsilon\}$ the ϵ -neighborhood of $\partial \Omega$ in \mathcal{M} . When $x \in \Omega \setminus U_{\epsilon}(\partial \Omega)$, we see by taking geodesic normal coordinate system at *x* that (5.36) still holds at this *x*. According to (5.37) we have that

$$\operatorname{Tr}\left(\mathbf{q}_{-2}(t,x,\overset{*}{x})\right) = \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} e^{i(x-\overset{*}{x})\cdot\xi} \Big((n-1)e^{-t\mu|\xi|^{2}} + e^{-t(2\mu+\lambda)|\xi|^{2}}\Big)d\xi$$

= $\frac{n-1}{(4\pi\mu t)^{n/2}} e^{-\frac{|x-\overset{*}{x}|^{2}}{4t\mu}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} e^{-\frac{|x-\overset{*}{x}|^{2}}{4t(2\mu+\lambda)}}$ for any $x \in \Omega \setminus U_{\epsilon}(\partial\Omega)$,

which exponentially tends to zero as $t \to 0^+$ because $|x - \overset{*}{x}| \ge \epsilon$. Hence

$$\int_{\Omega \setminus U_{\epsilon}(\partial \Omega)} \left(\operatorname{Tr} \left(\mathbf{q}_{-2}(t, x, \overset{*}{x}) \right) \right) dV = O(t^{1-\frac{n}{2}}) \quad \text{as} \ t \to 0^+.$$
(5.40)

Secondly, for $l \ge 1$, it can be verified that $\text{Tr}(\mathbf{q}_{-2-l}(x,\xi,\tau))$ is a sum of finitely many terms, each of which has the following form:

$$\frac{r_k(x,\xi)}{(\tau-\mu\sum_{l,m=1}^n g^{lm}\xi_l\xi_m)^s(\tau-(2\mu+\lambda)\sum_{l,m=1}^n g^{lm}\xi_l\xi_m)^j}$$

where k - 2s - 2j = -2 - l, and $r_k(x, \xi)$ is the symbol independent of τ and homogeneous of degree *k*. Again we take the geodesic normal coordinate systems center at *x* (i.e., $g_{jk}(x) = \delta_{jk}$ and $\Gamma_{ik}^l(x) = 0$), by applying residue theorem we see that, for $l \ge 1$,

$$\frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr}\left(\mathbf{q}_{-2-l}(x,\xi,\tau)\right) d\tau \right) d\xi = O(t^{l-\frac{n}{2}}) \text{ as } t \to 0^+ \text{ uniformly for } x \in \Omega,$$

and

$$\frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-x)\cdot\xi} \Big(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr}\left(\mathbf{q}_{-2-l}(x,\xi,\tau)\right) d\tau \Big) d\xi = O(t^{l-\frac{n}{2}}) \text{ as } t \to 0^+ \text{ uniformly for } x \in \Omega.$$
(5.41)

Therefore

$$\int_{\Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \sum_{l \ge 1} \operatorname{Tr} \left(\mathbf{q}_{-2-l}(x,\xi,\tau) \right) d\tau \right) d\xi \right\} dV = O(t^{1-\frac{n}{2}}) \text{ as } t \to 0^+, \quad (5.42)$$

and

$$\int_{\Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-x^*)\cdot\xi} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \sum_{l\ge 1} \operatorname{Tr}\left(\mathbf{q}_{-2-l}(x,\xi,\tau)\right) d\tau \right) d\xi \right\} dV = O(t^{1-\frac{n}{2}}) \text{ as } t \to 0^+.$$
(5.43)

Combining (5.32), (5.39) and (5.42), we have

$$\int_{\Omega} \operatorname{Tr} \left(\mathbf{K}(t, x, x) \right) dV = \left[\frac{n-1}{(4\pi\mu t)^{n/2}} + \frac{1}{(4\pi(2\mu + \lambda)t)^{n/2}} \right] \operatorname{Vol}(\Omega) + O(t^{1-\frac{n}{2}}) \text{ as } t \to 0^+.$$
(5.44)

Now, we will consider the case of $\int_{\Omega \cap U_{\epsilon}(\partial \Omega)} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x-x)\cdot\xi} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x,\xi,\tau) \right) d\tau \right) d\xi \right\} dV.$ We pick a self-double patch W of \mathcal{M} (such that $W \subset U_{\epsilon}(\partial \Omega)$) covering a patch $W \cap \partial \Omega$ of $\partial \Omega$ endowed (see the diagram on p. 54 of [MS-67]) with local coordinates x such that



 $\epsilon > x_n > 0$ in $W \cap \Omega$; $x_n = 0$ on $W \cap \partial \Omega$; $x_n(\overset{*}{x}) = -x_n(x)$; and the positive x_n -direction is perpendicular to $\partial \Omega$. This has the effect that (5.6)–(5.8) and

 $\sqrt{|g|/g_{nn}} dx_1 \cdots dx_{n-1}$ = the element of (Riemannian) surface area on $\partial \Omega$. (5.45)

We choose coordinates $x' = (x_1, \dots, x_{n-1})$ on an open set in $\partial\Omega$ and then coordinates (x', x_n) on a neighborhood in $\overline{\Omega}$ such that $x_n = 0$ on $\partial\Omega$ and $|\nabla x_n| = 1$ near $\partial\Omega$ while $x_n > 0$ on Ω and such that x' is constant on each geodesic segment in $\overline{\Omega}$ normal to $\partial\Omega$. Then the metric tensor on $\overline{\Omega}$ has the form (see [LU] or p. 532 of [Ta2])

$$(g_{jk}(x', x_n))_{n \times n} = \begin{pmatrix} (g_{jk}(x', x_n))_{(n-1) \times (n-1)} & 0\\ 0 & 1 \end{pmatrix}.$$
 (5.46)

Furthermore, we can take a geodesic normal coordinate system for $(\partial \Omega, g)$ centered at $x_0 = 0$, with respect to e_1, \dots, e_{n-1} , where e_1, \dots, e_{n-1} are the principal curvature vectors. As Riemann showed, one has (see p. 555 of [Ta2])

$$g_{jk}(x_0) = \delta_{jk}, \quad \frac{\partial g_{jk}}{\partial x_l}(x_0) = 0 \quad \text{for all} \quad 1 \le j, k, l \le n - 1,$$

$$-\frac{1}{2} \frac{\partial g_{jk}}{\partial x_n}(x_0) = \kappa_k \delta_{jk} \quad \text{for all} \quad 1 \le j, k \le n - 1,$$
(5.47)

where $\kappa_1 \cdots , \kappa_{n-1}$ are the principal curvatures of $\partial \Omega$ at point $x_0 = 0$. Due to the special geometric normal coordinate system and (5.47)–(5.46), we see that for any $x \in \{z \in \Omega | \operatorname{dist}(z, \partial \Omega) < \epsilon\}$,

$$x - \hat{x} = (0, \cdots, 0, x_n - (-x_n)) = (0, \cdots, 0, 2x_n).$$
 (5.48)

By (3.17) of [Liu2], (5.47), (5.37) and (5.48), we find that

$$\begin{split} &\int_{W\cap\Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle x-x,\xi\rangle} \Big(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x,\xi,\tau) \right) d\tau \Big) d\xi \right\} dV \\ &= \int_0^{\epsilon} dx_n \int_{W\cap\partial\Omega} \frac{dx'}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle 0,\xi'\rangle + i2x_n\xi_n} \left[\frac{1}{2\pi i} \int_C e^{-t\tau} \Big(\frac{n}{(\tau-\mu|\xi|^2)} + \frac{(\mu+\lambda)|\xi|^2}{(\tau-\mu|\xi|^2)(\tau-(2\mu+\lambda)|\xi|^2)} \Big) d\tau \right] d\xi \\ &= \int_0^{\epsilon} dx_n \int_{W\cap\partial\Omega} \frac{dx'}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i2x_n\xi_n} \Big((n-1)e^{-t\mu|\xi|^2} + e^{-t(2\mu+\lambda)|\xi|^2} d\tau \Big) d\xi \\ &= \int_0^{\epsilon} dx_n \int_{W\cap\partial\Omega} \frac{dx'}{(2\pi)^n} \int_{-\infty}^{\infty} e^{2ix_n\xi_n} \Big[\int_{\mathbb{R}^{n-1}} \Big((n-1)e^{-t\mu(|\xi'|^2+\xi_n^2)} + e^{-t(2\mu+\lambda)(|\xi'|^2+\xi_n^2)} \Big) d\xi' \Big] d\xi_n \\ &= \int_0^{\epsilon} dx_n \int_{W\cap\partial\Omega} \frac{1}{(2\pi)^n} \Big[\int_{-\infty}^{\infty} e^{2ix_n\xi_n} e^{-t\mu\xi_n^2} \Big(\int_{\mathbb{R}^{n-1}} (n-1)e^{-t\mu\sum_{j=1}^{n-1}\xi_j^2} d\xi' \Big) d\xi_n \Big] dx' \\ &+ \int_0^{\epsilon} dx_n \int_{W\cap\partial\Omega} \frac{1}{(2\pi)^n} \Big[\int_{-\infty}^{\infty} e^{2ix_n\xi_n} e^{-t(2\mu+\lambda)\xi_n^2} \Big(\int_{\mathbb{R}^{n-1}} e^{-t(2\mu+\lambda)\sum_{j=1}^{n-1}\xi_j^2} d\xi' \Big) d\xi_n \Big] dx', \end{split}$$

where $\xi = (\xi', \xi_n) \in \mathbb{R}^n, \xi' = (\xi_1, \dots, \xi_{n-1})$. A direct calculation shows that

$$\frac{1}{(2\pi)^{n}} \bigg[\int_{-\infty}^{\infty} e^{2ix_{n}\xi_{n}} e^{-t\mu\xi_{n}^{2}} \bigg(\int_{\mathbb{R}^{n-1}} (n-1)e^{-t\mu\sum_{j=1}^{n-1}\xi_{j}^{2}} d\xi' \bigg) d\xi_{n} \bigg] = \frac{n-1}{(4\pi\mu t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4\mu t}},$$

$$\frac{1}{(2\pi)^{n}} \bigg[\int_{-\infty}^{\infty} e^{2ix_{n}\xi_{n}} e^{-t(2\mu+\lambda)\xi_{n}^{2}} \bigg(\int_{\mathbb{R}^{n-1}} e^{-t(2\mu+\lambda)\sum_{j=1}^{n-1}\xi_{j}^{2}} d\xi' \bigg) d\xi_{n} = \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4(2\mu+\lambda)t}}.$$

Hence

$$(5.49)$$

$$\int_{W\cap\Omega} \left\{ \frac{1}{(2\pi)^{n}} \int_{\mathbb{R}^{n}} e^{i(x-x,\xi)} \left(\frac{1}{2\pi i} \int_{C} e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x,\xi,\tau) \right) d\tau \right) d\xi \right\} dV$$

$$= \int_{0}^{\epsilon} dx_{n} \int_{W\cap\partial\Omega} \left[\frac{n-1}{(4\pi\mu t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4\mu t}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4(2\mu+\lambda)t}} \right] dx'$$

$$= \int_{0}^{\infty} dx_{n} \int_{W\cap\partial\Omega} \left[\frac{n-1}{(4\pi\mu t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4\mu t}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4(2\mu+\lambda)t}} \right] dx'$$

$$- \int_{\epsilon}^{\infty} dx_{n} \int_{W\cap\partial\Omega} \left[\frac{n-1}{(4\pi\mu t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4\mu t}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4(2\mu+\lambda)t}} \right] dx'$$

$$= \frac{n-1}{4} \cdot \frac{\operatorname{Vol}(W\cap\partial\Omega)}{(4\pi\mu t)^{(n-1)/2}} + \frac{1}{4} \cdot \frac{\operatorname{Vol}(W\cap\partial\Omega)}{(4\pi(2\mu+\lambda)t)^{(n-1)/2}}$$

$$- \int_{W\cap\partial\Omega} \left\{ \int_{\epsilon}^{\infty} \left[\frac{n-1}{(4\pi\mu t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4\mu t}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}} e^{-\frac{(2x_{n})^{2}}{4(2\mu+\lambda)t}} \right] dx_{n} \right\} dx'.$$

It is easy to verify that for any fixed $\epsilon > 0$,

$$\int_{\epsilon}^{\infty} \frac{1}{(4\pi\lambda t)^{\frac{n}{2}}} e^{-\frac{(2x_n)^2}{4\mu t}} dx_n = O(t^{1-n/2}) \quad \text{as } t \to 0^+,$$

$$\int_{\epsilon}^{\infty} \frac{1}{(4\pi(2\mu+\lambda)t)^{\frac{n}{2}}} e^{-\frac{(2x_n)^2}{4(2\mu+\lambda)t}} dx_n = O(t^{1-n/2}) \quad \text{as } t \to 0^+.$$
(5.50)

From (5.49) and (5.50), we get that

$$\int_{W\cap\Omega} \left\{ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle x-x,\xi\rangle} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr}\left(\mathbf{q}_{-2}(x,\xi,\tau)\right) d\tau \right) d\xi \right\} dV$$
(5.51)

$$= \frac{n-1}{4} \cdot \frac{\text{Vol}(W \cap \partial \Omega)}{(4\pi\mu t)^{(n-1)/2}} + \frac{1}{4} \cdot \frac{\text{Vol}(W \cap \partial \Omega)}{(4\pi(2\mu + \lambda)t)^{(n-1)/2}} + O(t^{1-n/2}) \quad \text{as } t \to 0^+.$$

For any $x \in \Omega \cap U_{\epsilon}(\partial \Omega)$, we have

$$\operatorname{Tr} \left(K(t, x, \overset{*}{x}) \right) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle x - \overset{*}{x}, \xi \rangle} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x, \xi, \tau) \right) d\tau \right) d\xi$$

$$+ \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle x - \overset{*}{x}, \xi \rangle} \left(\sum_{l \ge 1} \frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2-l}(x, \xi, \tau) \right) d\tau \right) d\xi$$

$$= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle x - \overset{*}{x}, \xi \rangle} \left(\frac{1}{2\pi i} \int_C e^{-t\tau} \operatorname{Tr} \left(\mathbf{q}_{-2}(x, \xi, \tau) \right) d\tau \right) d\xi + O(t^{1+\frac{n}{2}}) \text{ as } t \to 0^+,$$

where the second equality used (5.41). Combining (5.51) and (5.52), we have

$$\int_{W\cap\Omega} \operatorname{Tr}(\mathbf{K}(t,x,\overset{*}{x}))dx = \frac{n-1}{4} \cdot \frac{\operatorname{Vol}(W\cap\partial\Omega)}{(4\pi\mu t)^{(n-1)/2}} + \frac{1}{4} \cdot \frac{\operatorname{Vol}(W\cap\partial\Omega)}{(4\pi(2\mu+\lambda)t)^{(n-1)/2}} + O(t^{1-n/2}) \quad \text{as } t \to 0^+.$$
(5.53)

It follows from (5.27), (5.33), (5.40), (5.43), (5.44) and (5.53) that

$$\int_{W\cap\Omega} \operatorname{Tr}(\mathbf{K}(t,x,x)) dx \neq \int_{W\cap\Omega} \operatorname{Tr}(\mathbf{K}(t,x,x)) dx \qquad (5.54)$$
$$= \left[\frac{n-1}{(4\pi\mu t)^{n/2}} + \frac{1}{(4\pi(2\mu+\lambda)t)^{n/2}}\right] \operatorname{Vol}(W\cap\Omega)$$
$$= \frac{1}{4} \left[(n-1) \frac{\operatorname{Vol}(W\cap\partial\Omega)}{(4\pi\mu t)^{(n-1)/2}} + \frac{\operatorname{Vol}(W\cap\partial\Omega)}{(4\pi(2\mu+\lambda)t)^{(n-1)/2}} \right]$$
$$+ O(t^{1-n/2}) \quad \text{as } t \to 0^+.$$

Note that

$$\sum_{k=1}^{\infty} e^{-t\tau_k^{(D)}} = \int_{\Omega} \operatorname{Tr}\left(\mathbf{K}^{(D)}(t,x,x)\right) dx = \int_{\Omega} \operatorname{Tr}\left(\mathbf{K}(t,x,x) - \mathbf{K}(t,x,x^*)\right) dx$$
(5.55)

and

$$\sum_{k=1}^{\infty} e^{-t\tau_k^{(T)}} = \int_{\Omega} \operatorname{Tr} \left(\mathbf{K}^{(T)}(t, x, x) \right) dx$$

$$= \int_{\Omega} \operatorname{Tr} \left(\mathbf{K}(t, x, x) + \mathbf{K}(t, x, x^*) \right) dx - \int_{\Omega} \operatorname{Tr} \left(\mathbf{H}(t, x, x) \right) dx.$$
(5.56)

Finally, by applying Lemma 2.1, we get the third coefficient a_2 in the asymptotic expansions for the Dirichlet and traction boundary conditions

$$a_{2} = \frac{1}{6(4\pi)^{n/2}} \Big[\Big(\frac{1}{(\lambda + 2\mu)^{(n-2)/2}} + \frac{n-7}{\mu^{(n-2)/2}} + \frac{12\mu}{n} \Big(\frac{1}{(\lambda + 2\mu)^{n/2}} + \frac{n-1}{\mu^{n/2}} \Big) \Big]_{\Omega} R(x) \, dV$$

$$+ 2 \Big(\frac{1}{(\lambda + 2\mu)^{(n-2)/2}} + \frac{n-7}{\mu^{(n-2)/2}} \Big) \int_{\partial\Omega} H(x) \, ds \Big].$$
(5.57)

From (5.54)–(5.56), (5.21) and (5.57), we get the conclusion of Theorem 1.4. \Box

Remark 5.1. (*i*) For the two-term heat trace asymptotic formula for the Lamé operator with Dirichlet or traction boundary conditions, we refer the reader to [Liu2], [Liu-23] or [PiOr], in which an open problem mentioned by Avramidi in [Avr] was affirmatively answered.

6. the Stokes eigenvalues and buckling eigenvalues

The buckling eigenvalue problem with Dirichlet boundary condition has interpretations in physics, that is, it describes the critical buckling load of a clamped plate subjected to a uniform compressive force around its boundary. In 1986, Girault and Raviart (see [GiR] or [ChL]) proved that all the Stokes eigenvalues with Dirichlet boundary condition coincide with all the buckling eigenvalues in the two-dimensional case. In this section, we shall also show that in two dimensions, the eigenvalue problem for the Stokes operator with Cauchy force boundary condition is equivalent to the eigenvalue problem for the buckling plate problem with Cauchy force boundary condition.

Proof of Theorem 1.8. As pointed out above, we shall only prove the case of the Cauchy force boundary condition, i.e., we shall show that

$$\varsigma_{k+1}^{(C)} = \Lambda_k^{(C)} \quad \text{for all } k \ge 1.$$
(6.1)

For a vector field $\mathbf{u} = (u^1, u^2)$ we define $\mathbf{u}^{\perp} = (-u^2, u^1)$ and for a scalar ψ we define $\nabla^{\perp}\psi = (-\partial_2\psi, \partial_1\psi)$. Clearly, $(\mathbf{u}^{\perp})^{\perp} = -\mathbf{u}$ and $(\nabla^{\perp})^{\perp} = -\nabla\psi$. By $\omega(\mathbf{u})$ we mean the vorticity (scalar curl) of \mathbf{u} : $\omega(\mathbf{u}) = \partial_1 u^2 - \partial_2 u^1$. Note that $\omega(\mathbf{u}) = -\operatorname{div} \mathbf{u}^{\perp}$. In particular, if div $\mathbf{u} = 0$, then $\nabla^{\perp}\omega(\mathbf{u}) = \Delta \mathbf{u}$.

Clearly, $(\varsigma_1, \mathbf{u}_1) = (0, c)$ is the first eigenpair of the Lamé operator with the Cauchy force boundary condition. Let $(\varsigma_k, \mathbf{u}_k)$, $(k \ge 2)$, be a Stokes eigenpair with Cauchy force boundary condition. Then there is a scalar function p_k in $H^1(\Omega)$ such that

$$\begin{cases} -\mu \Delta \mathbf{u}_{k} + \nabla p_{k} = \varsigma_{k} \mathbf{u}_{k} & \text{in } \Omega, \\ \text{div } \mathbf{u}_{k} = 0 & \text{in } \Omega, \\ \left(2\mu \left(\text{Def } \mathbf{u}_{k}\right) - p_{k} \mathbf{I}_{2}\right) \mathbf{v} = 0 & \text{on } \partial\Omega. \end{cases}$$
(6.2)

Here the above pressure function p_k is uniquely determined by letting $\inf_{x \in \Omega} p_k(x) = 0$, since p_k is uniquely determined up to an additive arbitrary constant. Note that the Cauchy force boundary condition in eigenvalue problem (6.2) can exactly be expressed as

$$\begin{pmatrix} 2\mu \partial_1 u_k^1 + p_k & \mu(\partial_2 u_k^1 + \partial_1 u_k^2) \\ \mu(\partial_1 u_k^2 + \partial_2 u_k^1) & 2\mu \partial_2 u_k^2 + p_k \end{pmatrix} \begin{pmatrix} v^1 \\ v^2 \end{pmatrix} = 0 \quad \text{on } \partial\Omega.$$
 (6.3)

Since $v = (v^1, v^2)$ is a non-zero solution of linear equation (6.3), we immediately get

$$\det \begin{pmatrix} 2\mu \partial_1 u_k^1 + p_k & \mu(\partial_2 u_k^1 + \partial_1 u_k^2) \\ \mu(\partial_1 u_k^2 + \partial_2 u_k^1) & 2\mu \partial_2 u_k^2 + p_k \end{pmatrix} = 0 \quad \text{on } \partial\Omega,$$
(6.4)

Thus, by (6.4) and div $\mathbf{u}_k = 0$ on Ω we have

$$p_k = \mu \sqrt{\left(\partial_1 u_k^2 + \partial_2 u_k^1\right)^2 + 4\left(\partial_1 u_k^1\right)^2} \quad \text{on} \quad \partial\Omega.$$
(6.5)

Taking the curl of Equation (6.2), we see that the vorticity $\omega_k = \operatorname{curl}(\mathbf{u}_k)$ satisfies

$$\begin{cases} \mu \Delta \omega_k + \varsigma_k \omega_k = 0 & \text{in } \Omega, \\ \left(2\mu \left(\text{Def } \mathbf{u}_k \right) - \mu \sqrt{\left(\partial_1 u_k^2 + \partial_2 u_k^1 \right)^2 + 4\left(\partial_1 u_k^1 \right)^2} \mathbf{I}_2 \right) \mathbf{v} = 0 & \text{on } \partial \Omega, \end{cases}$$
(6.6)

Here we have used the fact that $\omega(\nabla p_k) = 0$ in Ω , and

$$p_k = \mu \sqrt{\left(\partial_1 u_k^2 + \partial_2 u_k^1\right)^2 + 4\left(\partial_1 u_k^1\right)^2}$$
 on $\partial \Omega$.

That is, ω_k is an eigenfunction of $-\mu\Delta$, but with Cauchy force boundary conditions on the velocity \mathbf{u}_k .

Let ψ_k be the stream function for \mathbf{u}_k . That is,

$$\mathbf{u}_k = \nabla^\perp \psi_k \quad \text{in } \ \Omega. \tag{6.7}$$

Then $\omega_k = \Delta \psi_k$ in Ω . For the known function ω_k defined in Ω , let ψ_k is the solution of the following Poisson equation:

$$\begin{cases} \Delta \psi_k = \omega_k & \text{in } \Omega, \\ \psi_k = 0 & \text{on } \partial \Omega. \end{cases}$$
(6.8)

From (6.8), we have

$$\partial_1 u_k^1 = -\partial_1 \partial_2 \psi_k, \quad \partial_1 u_k^2 = \partial_1^2 \psi_k, \\ \partial_2 u_k^1 = -\partial_2^2 \psi_k, \quad \partial_2 u_k^2 = \partial_1 \partial_2 \psi_k$$
 in Ω .

Thus, ψ_k satisfies

$$\begin{cases} \mu \Delta \Delta \psi_k + \varsigma_k \Delta \psi_k = 0 & \text{in } \Omega, \\ \psi_k = 0 & \text{on } \partial \Omega, \\ (B\psi) \nu = 0 & \text{on } \partial \Omega, \end{cases}$$
(6.9)

where

$$B\psi = \mu \begin{pmatrix} -2\partial_1 \partial_2 \psi_k & \partial_1^2 \psi_k - \partial_2^2 \psi_k \\ \partial_1^2 \psi_k - \partial_2^2 \psi_k & 2\partial_1 \partial_2 \psi_k \end{pmatrix} - \mu \sqrt{(\partial_1^2 \psi_k - \partial_2^2 \psi_k)^2 + 4(\partial_1 \partial_2 \psi_k)^2} \mathbf{I}_2 \quad \text{on } \partial\Omega.$$

This is the corresponding eigenvalue problem for the buckling plate with Cauchy force boundary condition.

Conversely, given ψ_k satisfying Equation (6.9), $\omega_k = \Delta \psi_k$ and $\mathbf{u}_k = \nabla^{\perp} \psi_k$ satisfy Equation (6.6) and one can show, at least for sufficiently smooth boundaries, that \mathbf{u}_k satisfies Equation (6.2). Note that (0, c) is not a buckling eigenpair with Cauchy force boundary condition. Thus, except the first Stokes eigenpair, the eigenvalue problem for the Stokes operator and the buckling plate are equivalent for the Cauchy force boundary condition, which implies (6.1).

Remark 6.1. For a bounded domain Ω with smooth boundary in the Euclidean plane \mathbb{R}^2 , the previous formulas (1.25) in Corollary 1.10 only hold upto two-term asymptotics. It is easy to see that the third coefficients of the asymptotic expansions for $\sum_{k=1}^{\infty} e^{-t\Sigma_k}$, $\sum_{k=1}^{\infty} e^{-t\Gamma_k}$ and $\sum_{k=1}^{\infty} e^{-t\Lambda_k^{(D)}}$ are different. Note also that $\mu\Delta$ is the usual Laplacian defined on 0-form and $L_{-\mu,\mu} = \mu\Delta$ is the (vector form) Laplacian defined on 1-form.

Acknowledgments

This research was supported by the NNSF of China (12271031) and NNSF of China (11671033/A010802).

References

- [AGi] K. Abe, Y. Giga, Analyticity of the Stokes semigroup in spaces of bounded functions. Acta Math. 211, 1-46 (2013).
- [Avr] I. G. Avramidi, Non-Laplace type operators on manifolds with boundary, analysis, geometry and topology of elliptic operators, World Sci. Publ, Hackensack, 2006, pp. 107-140.
- [ADN2] S. Agmon, A. Douglis, L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions II, Comm. Pure Appl. Math., 17 (1964), 35–92.
- [Ahl] L. Ahlfors, Complex Analysis, Third Edition, McGraw-Hill, 1979.
- [Ahl-74] L. Ahlfors, *Conditions for quasiconformal deformations in several variables*, in: Contributions to analysis, a collection of papers dedicated to L. Bers, Academic Press, New York, 1974.
- [Ahl-76] L. Ahlfors, Quasiconformal deformations and mappings in Rn, J. Analyse Math. 30(1976), 74–97.
- [AsB] M. Ashbaugh and R. Benguria, A Sharp Bound for the Ratio of the First Two Eigenvalues of Dirichlet Laplacians and Extensions, Annals of Mathematics, no. 3, 135(1992), 601-628.
- [BG] T. P. Branson and P. Gilkey, *The asymptotics of the Laplacian on a manifold with boundary*, Comm. Partial Differential Equations, 15(1990), 245-22.
- [BGOP] T. P. Branson, P. B. Gilkey, B. Ørsted, and A. Pierzchalski, *Heat equation asymptotics of a generalized Ahlfors Laplacian on a manifold with boundary*, Operator Calculus and Spectral Theory, M. Demuth, B. Gramsch and B. Gramsch, and B.Schulze, eds., p. 1–13, Birkhäuser, Boston, 1992.
- [Bro] F. E. Browder, On the spectral theory of elliptic differential operators I, Math. Ann., 142(1960/61), 22–130.
- [CaCa] L. A. Caffarelli and X. Cabré, Fully nonlinear elliptic equations, American Mathematical Society Colloquium Publications, 43. American Mathematical Society, Providence, R.I., 1995.
- [Cha1] I. Chavel, Eigenvalues in Riemannian Geometry, Academic Press, 1984.
- [Cha2] I. Chavel, *Riemannian Geometry* A Modern Introduction, second ed., Cambridge University Press, 2006.
- [CaFrLeVa-23] M. Capoferri, L. Friedlander, M. Levitin, D. Vassiliev, Two-Term Spectral Asymptotics in Linear Elasticity, The Journal of Geometric Analysis (2023) 33:242.
- [ChL] W. Chen and Q. Lin, *Approximation of an eigenvalue problem associated with the Stokes problem by the stream function-vorticity-pressure method*, Appl. Math. 51(1), 73-88 (2006).
- [CiMa] A. Cialdea and V. Maz'ya, Criterion for the functional dissipativity of the Lamé operator, European Journal of Mechanics - A/solids, Vol.100, 2023,104522.
- [CLN] B. Chow, P. Lu, L. Ni, Hamilton's Ricci flow, Graduate Studies in Mathematics, 77, American Mathematical Society, Providence, RI; Science Press, Beijing, (2006).
- [CKVC] M. Chipot, D. Kinderlehrer, G. Vergara-Caffarelli, Smoothness of linear laminates, Arch. Ration. Mech. Anal. 96(1), 81–96 (1986).
- [CH] R. Courant, D. Hilbert, *Methods of Mathematical Physics*, vol. 1, Interscience Publishers, New York, 1953.
- [Don] H. Dong, Gradient Estimates for Parabolic and Elliptic Systems from Linear Laminates, Arch Rational Mech Anal., 205 (2012), 119–149. DOI: 10.1007/s00205-012-0501-z.
- [ES] Y. V. Egorov, M. A. Shubin, Partial Differential Equations II, Springer-Verlag, Berlin, Heidelberg, 1994.
- [Fri] A. Friedman, Partial differential equations, Holt, Rinehart and Winston, New York, 1969.

- [Giga] Y. Giga, Analyticity of the semigroup generated by the Stokes operator in L_r spaces. Math. Z., 178(1981), 297-329.
- [GiTr] D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*, Reprint of the 1998 edition, Classics Math. Springer-Verlag, Berlin, 2001.
- [Gil2] P. Gilkey, Invariance Theory, the Heat Equation and the Atiyah-Singer Index Theorem, CRC Press, Boca Raton, 1995.
- [GiR] V. Girault and P. A. Raviart, Finite Element Methods for Navier-Stokes Equations. Theory and Algorithms, volume 5 of Springer Series in Computational Mathematics. Springer, Berlin (1986).
- [Gru] G. Grubb, Functional Calculus of Pseudo-Differential Boundary Problems, Birkhäuser, Boston, 1986.
- [Gur-72] M. E. Gurtin, The linear theory of elasticity, in Encyclopedia of Physics (ed. S Flügge) Vol.VI a/2, Mechanics of solids II (ed. C. Truesdell), Springer-Verlag, 1972, 1-295.
- [KGBB] V. Kupradze, T. Gegelia, M. Basheleishvili, T. Burchuladze, *Three-dimensional problems of the mathematical theory of elasticity and thermoelasticity*, North-Holland Publishing Company, Amsterdam, New York, Oxford 1979.
- [Hah] P. Hähner, On Acoustic, Electromagnetic, and Elastic Scattering Problems in Inhomogeneous Media, Habilitation thesis, Göttingen, 1998.
- [Ho4] L. Hörmander, *The Analysis of Partial Differential Operators IV*, Springer-Verlag, Berlin, Heidelberg, New York (1985, 2009).
- [Iv] V. Ivrii, Second term of the spectral asymptotic expansion of the Laplace-Beltrami operator on Manifolds with boundary, Funktsional. Anal. i Prilozhen. 14 (2) (1980) 25-34; English transl.: Funct. Anal. Appl. 14 (1980) 98–106.
- [Iv2] V. Ivrii, 100 years of Weyl's law, Bull. Math. Sci. 6 (2016), 379–452.
- [Kac] M. Kac, *Can one hear the shape of a drum?*, Amer. Math. Monthly (Slaught Mem. Papers No. 11) 73 (4) (1966), 1-23.
- [Koz] A. N. Kozhevnikov, On the operator of the linearized steady-state Navier-Stokes problem. Math. USSR Sbornik 53(1), 1-16 (1986).
- [Lap] A. Laptev, Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces, J. Funct. Anal., 151(1997), 531-545.
- [Lee] J. M. Lee, Introduction to smooth manifolds, Springer Science+Business Media, New York, 2013.
- [LU] J. Lee and G. Uhlmann, Determing anisotropic real-analytic conductivities by boundary measurements, Comm. Pure Appl. Math., 42(1989), 1097–1112.
- [LiNi] Y. Y. Li, L. Nirenberg, *Estimates for elliptic systems from composite material*, Comm. Pure Appl. Math. 56(7), 892-925 (2003).
- [LiVo] Y. Y. Li, M. Vogelius, Gradient estimates for solutions to divergence form elliptic equations with discontinuous coefficients, Arch. Ration. Mech. Anal. 153(2), 91-151 (2000).
- [LiQin] T. Li and T. Qin, *Physics and partial differential equations*, Vol.1, Higher Education Press, Beijing, 2013, Translated by Y. Li.
- [Liu1] G. Q. Liu, The Weyl-type asymptotic formula for biharmonic Steklov eigenvalues on Riemannian manifolds. Adv. Math. 228(2011), 2162–2217.
- [Liu2] G. Q. Liu, Geometric Invariants of Spectrum of the Navier–Lamé Operator, J. Geom. Anal. 31 (2021), 10164–10193.
- [Liu3] G. Q. Liu, *The geometric invariants for the spectrum of the Stokes operator*, Math. Ann., 382(3-4),1985-2032(2022), https://doi.org/10.1007/s00208-021-02167-w.
- [Liu4] G. Q. Liu, One can know boundary area and curvatures by hearing the Steklov frequences of a Stokes flow, arXiv: 2207.04018.
- [Liu-11] G. Q. Liu, Some inequalities and asymptotic formulas for eigenvalues on Riemannian manifolds, J. Math. Anal. Appl. 376 (2011), 349–364.
- [Liu-23] G. Q. Liu, Remark on paper "Two-term spectral asymptotic in linear elasticity", arXiv: 2305.14134v5.

- [Liu-19] G. Q. Liu, Determination of isometric real-analytic metric and spectral invariants for elastic Dirichlet-to-Neumann map on Riemannian manifolds, arXiv:1908.05096v3.
- [Liu-39] G. Q. Liu, On an algorithm for two-term spectral asymptotic formulas, arXiv: 2309.07475v3 [math.DG].
- [LiuTan-22] G. Q. Liu, X. Tan, Asymptotic Expansions of the Traces of the Thermoelastic Operators, arXiv:2205.13238.
- [MaHu] J. E. Marsden and T. R. Hughes, *Mathematical Foundations of elasticity, Dover Publications*, Inc., New York, 1983.
- [McL] W. McLean, *Strongly elliptic systems and boundary integral equations*, Cambridge University Press, 2000.
- [MHNZ] M. V. de Hoop, G. Nakammura, J. Zhai, Reconstruction of Lamé Moduli and Density at the Boundary Enabling Directional Elastic Wavefield Decomposition, SIAM Journal Applied Mathematics No.2, 77(2017), 10.1137/16M1087266.
- [Mo3] C. Morrey, Multiple Integrals in the Calculus of Variations, Springer-Verlag, New York, Inc., 1966.
- [MS-67] H. McKean, I. M. Singer, *Curvature and the eigenvalues of the Laplacian*, J. Differential Geometry, 1(1967), 43–69.
- [Pa1] L.E. Payne, Isoperimetric inequalities and their applications, SIAM Rev. 9 (3) (1967) 453–488.
- [Pa] A. Pazy, Semigroups of Linear Operators and Applications to Partial Differential Equations, Springer 1983.
- [PiOr] A. Pierzchalski and B. Ørsted, The Ahlfors Laplacian on a Riemannian manifold with boundary, Michigan Math. J., 4(1996).
- [Ple1] Å. Pleijel, On the eigenvalues and eigenfunctions of elastic plates, Comm. Pure Appl. Math. 3 (1950) 1–10.
- [Ple2] Å. Pleijel, *Green's functions and asymptotic distribution of eigenvalues and eigenfunctions*, in: Proc. of the Symposium on Spectral Theory and Differential Problems, Stillwater, OK, 1951, pp. 439–454.
- [Po] G. Pólya, On the eigenvalues of vibrating membranes, Proc. Lond. Math. Soc. (3) 11 (1961) 419–433.
- [SaV] Yu. Safarov and D. Vassiliev, *The asymptotic distribution of eigenvalues of partial differential operators*, Amer. Math. Soc., Providence, RI, 1997. DOI: 10.1090/mmono/155.
- [Sa] L. Sandgren, A vibration problem, Medd. Lunds Univ. Mat. Sem. 13 (1955), 1–83.
- [Sar] P. Sarnak, Spectra of hyperbolic surfaces, Bull. Amer. Math. Soc. 40 (4) (2003), 441–478.
- [Ste] W. Steklov (V.A. Steklov), *Sur les problèmes fondamentaux de la physique mathématique*, Ann. Sci. Éc. Norm. Super. 19 (1902) 455–490.
- [Stew] H. B. Stewart, *Generation of analytic semigroups by strongly elliptic operators*, Trans. Am. Math. Soc., 199(1974), 141–161.
- [ShY] R. Schoen and S.-T. Yau, Lectures on Differential Geometry, International Press, 2010.
- [Ta1] M. E. Taylor, Partial Differential Equations I, 2nd Edition, Appl. Math. Sci., vol. 115, Springer Science+Business Media, LLC 1996, 2011.
- [Ta2] M. E. Taylor, Partial Differential Equations II, 2nd Edition, Appl. Math. Sci., vol. 116, Springer Science+Business Media, LLC 1996, 2011.
- [Ta3] M. E. Taylor, Partial Differential Equations III, Second Edition, Appl. Math. Sci., vol. 117, Springer-Verlag, New York, 2011.
- [Wei] H. F. Weinberger, An isoperimetric inequality for the n-dimensional free membrane problem, J. Rational Mech. Anal. 5(1956), 633-636.
- [Wey1] H. Weyl, Über die Abh'angigkeit der Eigenschwingungen einer Membran und deren Begrenzung, J. Reine Angew. Math. 141 (1912), 1–11.
- [Wey2] H. Weyl, it Des asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen, Math. Ann. 71 (1912), 441–479.
- [Xi-11] J. G. Xiong, C^{1,1} Estimates for elliptic equations with partial and piecewise continuous coefficients, Methods and Applications of Analysis, no.4, 18(2011), 373–390.