

ARAKELOV GEOMETRY OF CUNTZ-PIMSNER ALGEBRAS

IGOR V. NIKOLAEV¹

ABSTRACT. We use K -theory of the C^* -algebras to study the Arakelov geometry, i.e. a compactification of the arithmetic schemes $V \rightarrow \text{Spec } \mathbf{Z}$. In particular, it is proved that the Picard group of V is isomorphic to the K_0 -group of a Cuntz-Pimsner algebra associated to V . We apply the result to the finiteness problem for the algebraic varieties over number fields.

1. INTRODUCTION

The Arakelov theory deals with a compactification problem for the arithmetic schemes. We refer the reader to the original work [Arakelov 1974] [1] and [Soulé 1992] [9] for an excellent introduction. To outline the idea, consider an arithmetic variety V given by a system of the polynomial equations

$$f_1(x_1, \dots, x_n) = f_2(x_1, \dots, x_n) = \dots = f_k(x_1, \dots, x_n) = 0, \quad (1.1)$$

where $f_1, \dots, f_k \in \mathbf{Z}[x_1, \dots, x_n]$ are homogeneous polynomials with integer coefficients. If (1.1) has a solution in \mathbf{Z} , one can always reduce it modulo prime p thus obtaining a solution in the finite field \mathbf{F}_p . This phenomenon has a remarkable description in terms of commutative algebra and Grothendieck's theory of schemes. Namely, one can think of (1.1) as a fiber bundle over the base $\text{Spec } \mathbf{Z}$ consisting of the prime ideals of the ring \mathbf{Z} taken in the Zariski topology and fibers $V(\mathbf{F}_p)$. The well known analogy between function fields and number fields says that $\text{Spec } \mathbf{Z}$ can be compactified by a prime ideal at the point $p = \infty$. But there is no obvious way to define such an ideal. The problem was settled by Arakelov for $k = 1$ and $n = 2$, i.e. when the fiber is a curve of genus g . Specifically, it was proved that a Riemann surface $X(\mathbf{C})$ endowed with a hermitian metric on the holomorphic vector bundle over $X(\mathbf{C})$ corresponds to a fiber at infinity [Arakelov 1974] [1]. At the heart of Arakelov's construction is the notion of a compactified principal divisor:

$$(\varphi)_c = (\varphi) + v_\infty(\varphi)X(\mathbf{C}), \quad (1.2)$$

where φ is a rational function on V , (φ) is the principal divisor of φ and $v_\infty(\varphi) = -\int_{X(\mathbf{C})} \log |\varphi| d\mu$ so that the measure μ is defined by a Hermitian metric on the holomorphic vector bundle over $X(\mathbf{C})$. In particular, one can define a compactified Picard group $\text{Pic}_c(V)$ by taking the quotient of the abelian group of formal compactified divisors by its subgroup of principal divisors (1.2). The applications of the Arakelov theory so far include a new proof of the Faltings Finiteness Theorem for the algebraic curves of genus $g \geq 2$. Arakelov's theory has been generalized to the higher dimensions in [Soulé 1992] [9].

2010 *Mathematics Subject Classification.* Primary 11M55, 14G40; Secondary 46L85.
Key words and phrases. Arakelov theory, Cuntz-Pimsner algebras.

The Cuntz-Krieger algebra \mathcal{O}_A is a C^* -algebra generated by the partial isometries s_1, \dots, s_n which satisfy the relations $s_i^* s_i = \sum_{j=1}^n a_{ij} s_j s_j^*$, where $A = (a_{ij})$ is a square matrix with $a_{ij} \in \{0, 1, 2, \dots\}$ [Cuntz & Krieger 1980] [3]. Such algebras appear naturally in the study of local factors [6]. The Cuntz-Pimsner algebra is a generalization of \mathcal{O}_A to the countably infinite matrices $A_\infty \in GL_\infty(\mathbf{Z})$ [Pask & Raeburn 1996] [7]. Recall that the matrix A_∞ is called row-finite, if for each $i \in \mathbf{N}$ the number of $j \in \mathbf{N}$ with $a_{ij} \neq 0$ is finite. The matrix A is said to be irreducible, if some power of A is a strictly positive matrix and A is not a permutation matrix. It is known that if A_∞ is row-finite and irreducible, then the Cuntz-Pimsner algebra \mathcal{O}_{A_∞} is a well-defined and simple [Pask & Raeburn 1996] [7, Theorem 1]. The K -theory of the C^* -algebra \mathcal{O}_{A_∞} is determined by an exact sequence $0 \rightarrow K_1(\mathcal{O}_{A_\infty}) \rightarrow \mathbf{Z}^\infty \xrightarrow{1-A_\infty^t} \mathbf{Z}^\infty \xrightarrow{i_*} K_0(\mathcal{O}_{A_\infty}) \rightarrow 0$, so that

$$\begin{cases} K_0(\mathcal{O}_{A_\infty}) \cong \mathbf{Z}^\infty / (1 - A_\infty^t) \mathbf{Z}^\infty, \\ K_1(\mathcal{O}_{A_\infty}) \cong \text{Ker}(1 - A_\infty^t), \end{cases} \quad (1.3)$$

where A_∞^t is the transpose of A_∞ [Pask & Raeburn 1996] [7, Theorem 3]. An interplay between the Cuntz-Pimsner algebras and the local factors at infinity [Serre 1970] [8] is as follows. Let $\text{char } A_\infty$ be the characteristic polynomial of $A_\infty \in GL_\infty(\mathbf{Z})$; such a polynomial is well defined by the row-finiteness of matrix A_∞ [Pask & Raeburn 1996] [7, Note 2.1.11]. For every smooth n -dimensional projective variety V over a number field k there exist the Cuntz-Pimsner algebras $\mathcal{O}_{A_\infty^i}$, such that the Hasse-Weil zeta function of V [Serre 1970] [8] is given by the formula [6, Theorem 1.1]:

$$Z_V(s) = \prod_{i=0}^{2n} (\text{char } A_\infty^i)^{(-1)^{i+1}}. \quad (1.4)$$

The aim of our note is an isomorphism between the K_0 -group of the Cuntz-Pimsner algebra $\mathcal{O}_{A_\infty^1}$ and the Picard group $\text{Pic}_c(V)$ constructed by Arakelov (Theorem 1.1). Such a result is applied to the finiteness problem for the algebraic varieties over number fields (Conjecture 4.1). Namely, let $V \rightarrow \text{Spec } \mathbf{Z}$ be an arithmetic scheme and let $\mathcal{O}_{A_\infty^1}$ be a Cuntz-Pimsner algebra corresponding to V [6, Theorem 1.1]. Our main result can be formulated as follows.

Theorem 1.1. $K_0(\mathcal{O}_{A_\infty^1}) \cong \text{Pic}_c(V)$.

The paper is organized as follows. A brief review of the preliminary facts is given in Section 2. Theorem 1.1 proved in Section 3. An application of Theorem 1.1 is discussed in Section 4.

2. PRELIMINARIES

We briefly review Arakelov theory, Cuntz-Pimsner algebras and local factors at infinity. We refer the reader to [Arakelov 1974] [1], [6], [Pask & Raeburn 1996] [7] and [Serre 1970] [8] for a detailed exposition.

2.1. Arakelov geometry. Let k be a number field and let $\Lambda \subset k$ be its ring of algebraic integers. Denote by $m = \deg(k|\mathbf{Q})$ the degree of the field k over the rationals \mathbf{Q} . Let X be an algebraic variety over k and $V \rightarrow \text{Spec } \Lambda$ the corresponding arithmetic scheme. A finite divisor is the usual divisor of V , i.e. $D_{fin} = \sum k_i C_i$, where $k_i \in \mathbf{Z}$ and $C_i \subset V$ the irreducible closed subsets of V of

codimension 1. By X_∞ one understands a fiber of the bundle $V \rightarrow \text{Spec } \Lambda$ over the infinite point of $\text{Spec } \Lambda$. Along with the finite divisors, one considers divisors of the form:

$$D = D_{fin} + \sum_{i=1}^m \lambda_\infty^{(i)} X_\infty^{(i)}, \quad (2.1)$$

where $\lambda_\infty^{(i)} \in \mathbf{R}$ and summation is taken over all embeddings of the field k . To define a (compactified) principal divisor, it is necessary to fix on each $X_\infty^{(i)}$ a hermitian metric, such that the corresponding volume elements satisfy the condition $\int_{X_\infty^{(i)}} d\mu = 1$. (The construction can be generalized to higher dimensions using hermitian metric on holomorphic vector bundle over $X_\infty^{(i)}$ [Soulé 1992] [9, Section O.2.1].) Let φ be a rational function on V and (φ) be the principal divisor corresponding to φ . One defines $v_\infty(\varphi) = -\int_{X_\infty^{(i)}} \log |\varphi| d\mu$. The compactified principal divisor of φ is given by the formula:

$$(\varphi)_c = (\varphi) + \sum_{i=1}^m v_\infty(\varphi) X_\infty^{(i)}. \quad (2.2)$$

Divisors (2.1) and principal divisors (2.2) make an additive abelian group $Div(V)$ and $P(V)$, respectively. Since $P(V) \subset Div(V)$, one can introduce a compactified Picard group $Pic_c(V) = Div(V)/P(V)$. On the other hand, the well known product formula of valuation theory gives rise to the Picard group $Pic_c(k)$ of the number field k . Roughly speaking, the following result says that the Arakelov geometry is a correct compactification of the arithmetic schemes.

Theorem 2.1. ([1, Section 0]) *Arakelov completion of the fiber map $V \rightarrow \text{Spec } \Lambda$ induces a homomorphism of the abelian groups $Pic_c(k) \rightarrow Pic_c(V)$.*

2.2. Cuntz-Pimsner algebras. The Cuntz-Krieger algebra \mathcal{O}_A is a C^* -algebra generated by the partial isometries s_1, \dots, s_n which satisfy the relations

$$\begin{cases} s_1^* s_1 &= a_{11} s_1 s_1^* + a_{12} s_2 s_2^* + \dots + a_{1n} s_n s_n^* \\ s_2^* s_2 &= a_{21} s_1 s_1^* + a_{22} s_2 s_2^* + \dots + a_{2n} s_n s_n^* \\ \dots & \\ s_n^* s_n &= a_{n1} s_1 s_1^* + a_{n2} s_2 s_2^* + \dots + a_{nn} s_n s_n^* \end{cases} \quad (2.3)$$

where $A = (a_{ij})$ is a square matrix with $a_{ij} \in \{0, 1, 2, \dots\}$. (Note that the original definition of \mathcal{O}_A says that $a_{ij} \in \{0, 1\}$ but is known to be extendable to all non-negative integers [Cuntz & Krieger 1980] [3].) Such algebras appear naturally in the study of local factors [6].

The Cuntz-Pimsner algebra \mathcal{O}_{A_∞} corresponds to the case of the countably infinite matrices $A_\infty \in GL_\infty(\mathbf{Z})$ [Pask & Raeburn 1996] [7]. The matrix A_∞ is called row-finite, if for each $i \in \mathbf{N}$ the number of $j \in \mathbf{N}$ with $a_{ij} \neq 0$ is finite. The matrix A_∞ is said to be irreducible, if some power of A_∞ is a strictly positive matrix and A_∞ is not a permutation matrix. If A_∞ is row-finite and irreducible, then the Cuntz-Pimsner algebra \mathcal{O}_{A_∞} is a well-defined and simple [Pask & Raeburn 1996] [7, Theorem 1].

An AF-core $\mathcal{F} \subset \mathcal{O}_{A_\infty}$ is an Approximately Finite (AF-) C^* -algebra defined by the closure of the infinite union $\cup_{k,j} \cup_{i \in V_k^j} \mathcal{F}_k^j(i)$, where $\mathcal{F}_k^j(i)$ are finite-dimensional C^* -algebras built from matrix A_∞ [Pask & Raeburn 1996] [7, Definition 2.2.1]. Let $\alpha : \mathcal{O}_{A_\infty} \rightarrow \mathcal{O}_{A_\infty}$ be an automorphism acting on the generators s_i of \mathcal{O}_{A_∞} by to

the formula $\alpha_z(s_i) = z s_i$, where z is a complex number $|z| = 1$. One gets an action of the abelian group $\mathbb{T} \cong \mathbf{R}/\mathbf{Z}$ on \mathcal{O}_{A_∞} . The Takai duality [Pask & Raeburn 1996] [7, p. 432] says that:

$$\mathcal{F} \rtimes_{\hat{\alpha}} \mathbb{T} \cong \mathcal{O}_{A_\infty} \otimes \mathcal{K}, \quad (2.4)$$

where $\hat{\alpha}$ is the Takai dual of α and \mathcal{K} is the C^* -algebra of compact operators. Using (2.4) one can calculate the the K -theory of \mathcal{O}_{A_∞} . Namely, the following statement is true.

Theorem 2.2. ([7, Theorem 3]) *If A_∞ is row-finite irreducible matrix, then there exists an exact sequence of the abelian groups:*

$$0 \rightarrow K_1(\mathcal{O}_{A_\infty}) \rightarrow \mathbf{Z}^\infty \xrightarrow{1-A_\infty^t} \mathbf{Z}^\infty \xrightarrow{i_*} K_0(\mathcal{O}_{A_\infty}) \rightarrow 0, \quad (2.5)$$

so that $K_0(\mathcal{O}_{A_\infty}) \cong \mathbf{Z}^\infty / (1 - A_\infty^t) \mathbf{Z}^\infty$ and $K_1(\mathcal{O}_{A_\infty}) \cong \text{Ker}(1 - A_\infty^t)$, where A_∞^t is the transpose of A_∞ and $i : \mathcal{F} \hookrightarrow \mathcal{O}_{A_\infty}$. Moreover, the Grothendieck semigroup $K_0^+(\mathcal{F}) \cong \varinjlim (\mathbf{Z}^\infty, A_\infty^t)$.

2.3. Local factors at infinity. Let V be an n -dimensional smooth projective variety over a number field k and let $V(\mathbf{F}_q)$ be a good reduction of V modulo the prime ideal corresponding to $q = p^r$. The local zeta $Z_q(u) := \exp(\sum_{m=1}^\infty |V(\mathbf{F}_q)| \frac{u^m}{m})$ is a rational function $Z_q(u) = \prod_{i=0}^{2n} (P_i(u))^{(-1)^{i+1}}$, where $P_0(u) = 1 - u$ and $P_{2n}(u) = 1 - q^n u$. Each $P_i(u)$ is the characteristic polynomial of the Frobenius endomorphism $Fr_q^i : (a_1, \dots, a_n) \mapsto (a_1^q, \dots, a_n^q)$ acting on the i -th ℓ -adic cohomology group $H^i(V)$ of variety V . The number of points on $V(\mathbf{F}_q)$ is given by the Lefschetz trace formula $|V(\mathbf{F}_q)| = \sum_{i=0}^{2n} (-1)^i \text{tr}(Fr_q^i)$, where tr is the trace of endomorphism Fr_q^i , see e.g. [Serre 1970] [8]. The Hasse-Weil zeta function of V is an infinite product $Z_V(s) = \prod_p Z_q(p^{-s})$, $s \in \mathbf{C}$, where p runs through all but a finite set of primes. A fundamental analogy between number fields and function fields predicts a prime $p = \infty$ in formula $Z_q(u) = \prod_{i=0}^{2n} (P_i(u))^{(-1)^{i+1}}$. Serre constructed local factors $\Gamma_V^i(s)$ realizing the analogy in terms of the Γ -functions attached to the Hodge structure on V [Serre 1970] [8]. To define $\Gamma_V^i(s)$ in a way similar to finite primes, Deninger introduced an infinite-dimensional cohomology $H_{ar}^i(V)$ and an action of Frobenius endomorphism $Fr_\infty^i : H_{ar}^i(V) \rightarrow H_{ar}^i(V)$, such that $\Gamma_V^i(s) \equiv \text{char}^{-1} Fr_\infty^i$, where $\text{char} Fr_\infty^i$ is the characteristic polynomial of Fr_∞^i [Deninger 1991] [4, Theorem 4.1]. The following result relates the local factors at infinity with the Cuntz-Pimsner algebras.

Theorem 2.3. ([6, Theorem 1.1]) *For every smooth n -dimensional projective variety V over a number field k there exist the Cuntz-Pimsner algebras $\mathcal{O}_{A_\infty^i}$, such that the Hasse-Weil zeta function of V is given by the formula*

$$Z_V(s) = \prod_{i=0}^{2n} (\text{char} A_\infty^i)^{(-1)^{i+1}}. \quad (2.6)$$

3. PROOF

For the sake of clarity, let us outline the main ideas. Let Fr_∞^1 be the Frobenius endomorphism of the Deninger cohomology group $H_{ar}^1(V)$. Recall that $H_{ar}^1(V) \cong \mathbf{Z}[\mathbf{x}^{\pm 1}]$ is the ring of Laurent polynomials in variables $\mathbf{x} = (x_1, \dots, x_{b_1})$, where b_1 is the first Betti number of variety V [Deninger 1991] [4, Section 3]. The commutative ring $A := Fr_\infty^1(\mathbf{Z}[\mathbf{x}^{\pm 1}])$ is a coordinate ring of variety V (Lemma 3.1). One can

consider a divisor class group $C(A)$ of the ring A as the quotient group of divisorial ideals $D(A)$ by the subgroup $F(A)$ consisting of the principal ideals. Since A is a locally factorial ring, the group $C(A)$ is isomorphic to the Picard group of variety V (Lemma 3.2). The rest of the proof follows from Theorem 2.2 and the well known formula $C(A) \cong \mathbf{Z}^\infty / (1 - (Fr_\infty^1)^t) \mathbf{Z}^\infty$ (Lemma 3.3). Let us pass to a detailed argument.

Lemma 3.1. *The commutative ring $A := Fr_\infty^1(\mathbf{Z}[\mathbf{x}^{\pm 1}])$ is a coordinate ring of variety V , i.e. V and V' are isomorphic whenever $A \cong A'$.*

Proof. (i) Let b_1 be the first Betti number of variety $V(\mathbf{C})$. Recall that the first Deninger's cohomology group $H_{ar}^1(V)$ is isomorphic to the additive group of the Laurent polynomials $\mathbf{Z}[x_1^{\pm 1}, \dots, x_{b_1}^{\pm 1}]$ [Deninger 1991] [4, Section 3]. Thus one gets an endomorphism $Fr_\infty^1 : \mathbf{Z}[x_1^{\pm 1}, \dots, x_{b_1}^{\pm 1}] \rightarrow \mathbf{Z}[x_1^{\pm 1}, \dots, x_{b_1}^{\pm 1}]$. Notice that any basis of $\mathbf{Z}[x_1^{\pm 1}, \dots, x_{b_1}^{\pm 1}]$ is countably infinite, and therefore Fr_∞^1 is given by an infinite-dimensional integer matrix; hence the notation.

(ii) Let $V \rightarrow V'$ be an isomorphism between projective varieties V and V' . The cohomology functor induces an isomorphism $\phi : H_{ar}^i(V) \rightarrow H_{ar}^i(V')$ of the corresponding Deninger cohomology groups. Recall that $H_{ar}^i(V) \cong \mathbf{R}[\mathbf{x}^{\pm 1}]$ and since the isomorphism of V is defined over a number field k , one gets an isomorphism $\phi : \mathbf{Z}[\mathbf{x}^{\pm 1}] \rightarrow \mathbf{Z}[\mathbf{y}^{\pm 1}]$. The group isomorphism ϕ extends to a ring isomorphism by choice of a monomial basis in the ring of the Laurent polynomials, and vice versa. It remains to notice that the endomorphism $Fr_\infty^i : \mathbf{Z}[\mathbf{x}^{\pm 1}] \rightarrow \mathbf{Z}[\mathbf{x}^{\pm 1}]$ commutes with ϕ and therefore $\phi(Fr_\infty^i(\mathbf{Z}[\mathbf{x}^{\pm 1}])) = Fr_\infty^i(\mathbf{Z}[\mathbf{y}^{\pm 1}])$. In other words, one gets a ring isomorphism $A \cong A'$. Lemma 3.1 is proved. \square

Lemma 3.2. $C(A) \cong Pic_c(V)$.

Proof. (i) Recall that A is a cluster algebra [6]. Such algebras are Krull domains and, therefore, one can consider the corresponding divisor class group $C(A)$ [Elsener, Lampe & Smertnig 2019] [5].

(ii) Let $Pic(A)$ be the Picard group of the ring A , i.e. a subgroup of $C(A)$ generated by the invertible ideals of A . Since A is a locally factorial ring, one gets a ring isomorphism $Pic(A) \cong C(A)$.

(iii) It is known that $Pic(A) \cong Pic(Spec A)$, where $Spec A \cong V$. On the other hand, we recall that the ring A has been constructed from the action of Frobenius endomorphism Fr_∞^1 on the Deninger cohomology group $H_{ar}^1(V)$ of variety V . In particular, the local factor at the prime $p = \infty$ is the (inverse) of characteristic polynomial of the linear action of Fr_∞^1 on such a cohomology. We conclude therefore that the Picard group of the ring A must encode a compactification of the arithmetic scheme $V \rightarrow Spec A$. In other words, one gets an isomorphism:

$$Pic(Spec A) \cong Pic_c(V). \quad (3.1)$$

(iv) Since $C(A) \cong Pic(A)$ and $Pic(A) \cong Pic(Spec A)$, we conclude from (3.1) that $C(A) \cong Pic_c(V)$. Lemma 3.2 is proved. \square

Lemma 3.3. $K_0(\mathcal{O}_{A_\infty^\perp}) \cong Pic_c(V)$.

Proof. (i) Let us calculate $C(A)$ in terms of the Frobenius endomorphisms $Fr_\infty^i : \mathbf{Z}[\mathbf{x}^{\pm 1}] \rightarrow \mathbf{Z}[\mathbf{x}^{\pm 1}]$. Notice that ideals of A of height i correspond to Frobenius endomorphisms Fr_∞^i for all $0 \leq i \leq 2n$. Since $C(A)$ is generated by the classes of the prime ideals of height 1, we conclude that $C(A)$ depends only on the endomorphism $Fr_\infty^1 : \mathbf{Z}[\mathbf{x}^{\pm 1}] \rightarrow \mathbf{Z}[\mathbf{x}^{\pm 1}]$.

(ii) Consider the additive group $L \cong \mathbf{Z}^\infty$ of the ring $\mathbf{Z}[\mathbf{x}^{\pm 1}]$. The endomorphism Fr_∞^1 defines a sub-lattice $L' := Fr_\infty^1(\mathbf{Z}[\mathbf{x}^{\pm 1}])$ of the lattice L . The index of $L' \subset L$ is given by the formula:

$$\left| \frac{\mathbf{Z}^\infty}{(1 - (Fr_\infty^1))^t \mathbf{Z}^\infty} \right|. \quad (3.2)$$

(iii) On the other hand, $C(A)$ is the quotient of divisorial ideals $D(A)$ by the principal ideals $F(A)$ of the ring A . It is not hard to see that each principal ideal of A can be written as xL for some $x \in L'$. In other words, $F(A) = \{xL \mid x \in L'\}$. Thus

$$C(A) = D(A)/F(A) \cong L/L' \cong \frac{\mathbf{Z}^\infty}{(1 - (Fr_\infty^1))^t \mathbf{Z}^\infty}. \quad (3.3)$$

(iv) Likewise, we set $A_\infty = Fr_\infty^1$ in Theorem 2.2 and we get

$$K_0(\mathcal{O}_{A_\infty}) \cong \frac{\mathbf{Z}^\infty}{(1 - (Fr_\infty^1))^t \mathbf{Z}^\infty}. \quad (3.4)$$

(v) Comparing (3.3) and (3.4) we conclude that $K_0(\mathcal{O}_{A_\infty}) \cong C(A)$. But Lemma 3.2 says that $C(A) \cong Pic_c(V)$. Therefore $K_0(\mathcal{O}_{A_\infty}) \cong Pic_c(V)$. Lemma 3.3 is proved. \square

Theorem 1.1 follows from Lemma 3.3.

4. REMARKS

Recall that the K -theory of a Cuntz-Krieger algebra \mathcal{O}_A is given by the formulas:

$$\begin{cases} K_0(\mathcal{O}_A) \cong \mathbf{Z}^n / (1 - A^t) \mathbf{Z}^n, \\ K_1(\mathcal{O}_A) \cong Ker(1 - A^t). \end{cases} \quad (4.1)$$

It is immediate that $|K_0(\mathcal{O}_A)| < \infty$ if and only if $\det(1 - A^t) \neq 0$. Whenever $\det(1 - A^t) = 0$ both $K_0(\mathcal{O}_A)$ and $K_1(\mathcal{O}_A)$ are infinite groups [Blackadar 1986] [2, Exercise 10.11.9 (c)]. On the other hand, it is known that $|K_0(\mathcal{O}_A)| = |V(\mathbf{F}_p)|$ for a matrix $A \in GL_n(\mathbf{Z})$, see e.g. [6, Section 1].

It is useful to think of $K_0(\mathcal{O}_{A_\infty})$ as the inductive limit of finite abelian groups $K_0(\mathcal{O}_{A_i})$, see definition of the AF-core \mathcal{F} in Section 2.2 and formula (2.4). In other words,

$$\lim_{i \rightarrow \infty} K_0(\mathcal{O}_{A_i}) \cong K_0(\mathcal{O}_{A_\infty}). \quad (4.2)$$

However, if $\det(1 - A_k) = 0$ for some k , one gets $|K_0(\mathcal{O}_{A_k})| = \infty$. In this case the convergent sequence (4.2) must terminate after a finite number of steps. Given that finiteness of the K_0 -groups is related to such of the points of variety V , we have the following conjecture.

Conjecture 4.1.

$$|V(k)| = \begin{cases} \text{finite,} & \text{if } \det(1 - A_\infty^1) = 0, \\ \text{infinite,} & \text{otherwise.} \end{cases} \quad (4.3)$$

REFERENCES

1. S. Yu. Arakelov, *Теория пересечений дивизоров на арифметической поверхности*, Изв. АН СССР **38** (1974), 1179-1192.
2. B. Blackadar, *K-Theory for Operator Algebras*, MSRI Publications, Springer, 1986.
3. J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56** (1980), 251-268.
4. C. Deninger, *On the Γ -factors attached to motives*, Invent. Math. **104** (1991), 245-261.
5. A. G. Elsenner, P. Lampe and D. Smertnig, *Factoriality and class groups of cluster algebras*, Adv. Math. **358** (2019), 106858, 48 pp.
6. I. V. Nikolaev, *Local factors and Cuntz-Pimsner algebras*, arXiv:2404.12179
7. D. Pask and I. Raeburn, *On the K-theory of Cuntz-Krieger algebras*, Publ. RIMS, Kyoto Univ. **32** (1996), 415-443.
8. J.-P. Serre, *Facteurs locaux des fonctions zêta des variétés algébriques (définitions et conjectures)*, Séminaire Delange-Pisot-Poitou (Théorie des nombres) **11** (1969-1970), exp. 19, pp. 1-15.
9. C. Soulé, *Lectures on Arakelov Geometry*, Cambridge Studies in Advanced Mathematics **33**, Cambridge University Press 1992.

¹ DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHN'S UNIVERSITY, 8000 UTOPIA PARKWAY, NEW YORK, NY 11439, UNITED STATES.

Email address: igor.v.nikolaev@gmail.com