

Probabilistic Correlation Functions of the Schwarzian Field Theory

Ilya Losev*

June 26, 2024

Abstract

We study correlation functions of the probabilistic Schwarzian Field Theory. We compute cross-ratio correlation functions exactly in the case when the corresponding Wilson lines do not intersect, confirming predictions made in the physics literature via limit of the conformal bootstrap and the DOZZ formula. Moreover, we prove that these correlation functions characterise the measure uniquely. We use them to define and compute the stress-energy tensor correlation functions, and demonstrate, in particular, that these agree with the results obtained earlier by formal differentiation of the partition function.

1 Introduction and main results

1.1 Introduction

Recently, the Schwarzian Field Theory attracted a lot of attention in the physics literature. It appears in the study of the AdS/CFT correspondence as a holographic dual of Jackiw–Teitelboim (JT) gravity in the disk [Iliesiu et al., 2019; Maldacena et al., 2016; Saad et al., 2019]. The Schwarzian Field Theory also emerges in the low energy limit of the Sachdev–Ye–Kitaev (SYK) random matrix model (e.g. see [Maldacena and Stanford, 2016] and [Kitaev and Suh, 2018]), and is related to Liouville CFT, infinite dimensional symplectic geometry, representation theory of the Virasoro algebra and 2D Yang–Mills.

In our companion paper [Bauerschmidt, Losev, and Wildemann, 2024] we have rigorously constructed a finite measure on $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$, which corresponds to the Schwarzian Field Theory and computed its partition function (i.e. total mass) using methods of stochastic analysis. The obtained result for the partition function agrees with the formula derived in [Stanford and Witten, 2017] using a formal application of the Duistermaat–Heckman theorem on the infinite dimensional symplectic space $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$.

In this paper we continue the study of this measure and rigorously compute a natural class of correlation functions of cross-ratio observables. These correlation functions were originally formally derived in [Mertens et al., 2017] by taking a $c \rightarrow \infty$ limit of two dimensional Liouville CFT and, in particular, a limit of the DOZZ formula. Our results match the formulae obtained in [Mertens et al., 2017]. Moreover, we prove that the computed correlation functions determine the measure uniquely, which further confirms that the measure constructed in [Bauerschmidt, Losev, and Wildemann, 2024]

*DPMMS, University of Cambridge. E-mail: i1320@cam.ac.uk.

corresponds to the Schwarzian Field Theory studied in the physics literature. In addition, we show that using these observables we can make sense of and compute the stress-energy tensor correlation functions, using a procedure similar to one proposed in [Mertens et al., 2017]. We also prove that the stress-energy tensor correlation functions calculated this way agree with values obtained by differentiation of the partition function, as proposed in [Stanford and Witten, 2017] and carried out in [Bauerschmidt, Losev, and Wildemann, 2024]. This, in particular, proves a conjecture from [Mertens et al., 2017].

Formally, the measure corresponding to the Schwarzian Field Theory is supported on the topological space $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$ with density (see [Stanford and Witten, 2017, (1.1)])

$$d\mathcal{M}_{\sigma^2}(\phi) = \exp \left\{ + \frac{1}{\sigma^2} \int_0^1 [\mathcal{S}_\phi(\tau) + 2\pi^2 \phi'^2(\tau)] d\tau \right\} \frac{\prod_{\tau \in [0,1]} \frac{d\phi(\tau)}{\phi'(\tau)}}{\text{PSL}(2, \mathbb{R})}, \quad (1)$$

where $\mathcal{S}_\phi(\tau)$ is the Schwarzian derivative of ϕ given by

$$\mathcal{S}_\phi(\tau) = \mathcal{S}(\phi, \tau) = \left(\frac{\phi''(\tau)}{\phi'(\tau)} \right)' - \frac{1}{2} \left(\frac{\phi''(\tau)}{\phi'(\tau)} \right)^2. \quad (2)$$

Here, $\mathbb{T} = [0, 1]/\{0 \sim 1\}$ is the unit circle, $\text{Diff}^1(\mathbb{T})$ is the space of C^1 orientation preserving diffeomorphisms of \mathbb{T} , and $\text{PSL}(2, \mathbb{R})$ is the group of Möbius transformations of the unit disk (i.e. conformal isomorphisms of the unit disk) restricted to the boundary which is identified with \mathbb{T} . The group $\text{PSL}(2, \mathbb{R})$ acts on $\text{Diff}^1(\mathbb{T})$ by left compositions. Following [Stanford and Witten, 2017] we call it a right action, since in [Stanford and Witten, 2017] it is interpreted as an action on the inverse elements. We denote the quotient of $\text{Diff}^1(\mathbb{T})$ by this action of $\text{PSL}(2, \mathbb{R})$ by $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$. Heuristically, the formal density (1) only depends on the orbit of this action and the quotient by $\text{PSL}(2, \mathbb{R})$ therefore makes sense. In Section 2 we recall the rigorous construction of the measure corresponding to the Schwarzian Field Theory, carried out in [Bauerschmidt, Losev, and Wildemann, 2024].

In our investigation of correlation functions we consider bi-local observables given by cross-ratios

$$\mathcal{O}(\phi; s, t) = \frac{\pi \sqrt{\phi'(t)\phi'(s)}}{\sin(\pi[\phi(t) - \phi(s)])}, \quad s, t \in \mathbb{T}. \quad (3)$$

These observables are defined for $\phi \in \text{Diff}^1(\mathbb{T})$ and are invariant under the $\text{PSL}(2, \mathbb{R})$ action, see Proposition 22. Therefore, these cross-ratios naturally induce well-defined observables on $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$, for which we use the same notation. Correlations of these observables are predicted to correspond to boundary-anchored Wilson line correlators in the JT gravity [Blommaert et al., 2018]. It is also predicted that these correlations describe the boundary-to-boundary propagator of a massive particle in the metric formulation of JT gravity [Iliesiu et al., 2019]. These observables are also known to be related to the Green's function of the SYK model [Bagrets et al., 2016; Saad et al., 2018].

We rigorously compute cross-ratio correlation functions in the case when Wilson lines (lines connecting s and t on the circle) do not intersect, see Figure 2a. In this case we call a set of observables *non-interlaced*, see Section 1.2. Moreover, we prove that the computed correlation functions characterise the measure uniquely. Our methods rely on stochastic analysis and, as far as we know, differ from the methods existing in the physics literature. In particular, we prove our results directly, without relying on the two dimensional Liouville Theory, and the DOZZ formula, which was rigorously proved only recently in [Kupiainen et al., 2017].

Furthermore, these observables are related to the stress-energy tensor of the Schwarzian Field Theory. Formally, the stress-energy tensor is given by the Schwarzian derivative $\mathcal{S}(\tan(\pi\phi), \cdot) = \mathcal{S}_\phi(\tau) + 2\pi^2 \phi'^2(\tau)$, which is connected to observables (3), at least for sufficiently smooth ϕ , by

$$\left(\frac{\pi \sqrt{\phi'(t)\phi'(s)}}{\sin(\pi[\phi(t) - \phi(s)])} \right)^2 = (t - s)^{-2} + \frac{1}{6} \mathcal{S}(\tan(\pi\phi), \tau) + o(1). \quad (4)$$




Figure 1: An example of two observables.

Even though regularity of diffeomorphisms under $d\mathcal{M}_{\sigma^2}$ is not sufficient for the Schwarzian derivative to exist, this relation can be used on the level of correlation functions. In Section 1.3 we show that the corresponding limits of correlation functions exist, and we compute them explicitly. We show that this limit coincides with formulae obtained by differentiating the partition function in [Bauerschmidt, Losev, and Wildemann, 2024], in particular confirming the conjecture from [Mertens et al., 2017], which was checked numerically there.

1.2 Diagrammatic representation on \mathbb{T}

In order to formulate the main results we develop a diagrammatic language. To a given set of observables we associate a diagram in a disk, where every observable is represented with a chord.

Let $\{\mathcal{O}(\phi; s_j, t_j)\}_{j=1}^N$ be a set of observables. Given this set we draw a unit circle and all the points $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ on it. For all $1 \leq j \leq N$ we connect the point s_j with the point t_j with a line segment. These line segments also correspond Wilson lines, see [Blommaert et al., 2018; Iliesiu et al., 2019].

Definition 1. We say that a set of observables $\{\mathcal{O}(\phi; s_j, t_j)\}_{j=1}^N$ is non-interlaced if interiors of all drawn chords (Wilson lines) in the corresponding diagram are pairwise non-intersecting.

Figure 1 gives examples of graphic representations of two observables being interlaced and non-interlaced.


Let $\{\mathcal{O}(\phi; s_j, t_j)\}_{j=1}^N$ be a set of non-interlaced observables. The N drawn chords on the corresponding diagram divide the unit disk into $N+1$ connected domains. We number them with integers from 1 to $N+1$. For each $m \in \{1, \dots, N+1\}$ we associate a Fourier variable k_m to domain number m . We also let τ_m be the total length of all arcs (parts of the initial circle) which form the boundary of m -th domain (see Figure 2a).

For each $j \in \{1, \dots, N\}$ we define $w_1(j)$ and $w_2(j)$ to be the Fourier variables corresponding to the domains that contain the line segment connecting s_j with t_j in their boundaries (see Figure 2b). All formulae will be symmetric in w_1 and w_2 , so the exact order does not matter.

Definition 2. For all $l, k, w \in \mathbb{R}$ we define $\Gamma(l \pm ik \pm iw)$ as

$$\Gamma(l \pm ik \pm iw) := \Gamma(l + ik + iw)\Gamma(l + ik - iw)\Gamma(l - ik + iw)\Gamma(l - ik - iw). \quad (5)$$

Remark 3. For any $l > 0$, the function $\Gamma(l \pm ik \pm iw)$ is non-negative, since $\Gamma(z)\Gamma(\bar{z}) = |\Gamma(z)|^2$.



(a) An example of a diagram. Here,
 $\tau_1 = (t_1 - s_1)$,
 $\tau_2 = (s_1 - s_3) + (s_2 - t_1) + (t_3 - t_2)$,
 $\tau_3 = (t_2 - s_2)$,
 $\tau_4 = (s_3 - t_4)$,
 $\tau_5 = (t_4 - s_4)$.

(b) We use $w_1(j)$ and $w_2(j)$ to denote the Fourier variables k_m corresponding to the domains that lie on both sides of the line segment that connects s_j with t_j . In Figure 2a, for example, $w_1(3) = k_2$ and $w_2(3) = k_4$ (or vice versa).

Figure 2: A diagrammatic representation of Fourier variables.

Remark 4. It is well-known (see e.g. [Gradshteyn and Ryzhik, 2007, 8.332, 8.331.1]) that for all positive integers n and $x \in \mathbb{R}$,

$$|\Gamma(1 + n + ix)|^2 = \frac{\pi x}{\sinh(\pi x)} \prod_{k=1}^n (k^2 + x^2), \quad (6)$$

$$|\Gamma(\frac{1}{2} + n + ix)|^2 = \frac{\pi}{\cosh(\pi x)} \prod_{k=1}^n \left((k - \frac{1}{2})^2 + x^2 \right). \quad (7)$$

Using the diagrammatic language we can now formulate the main result about the correlation functions, confirming predictions from [Mertens et al., 2017].

Theorem 1. For $N \geq 0$ let $\{\mathcal{O}(\phi; s_j, t_j)\}_{j=1}^N$ be a set of non-interlaced observables, and $\{l_j\}_{j=1}^N$ be a set of positive integers. Let also $\{k_m\}_{m=1}^{N+1}$ and $\{\tau_m\}_{m=1}^{N+1}$ be as above. Then

$$\int_{\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})} \prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) d\mathcal{M}_{\sigma^2}(\phi) = \int_{\mathbb{R}_+^{N+1}} \prod_{j=1}^N \frac{\Gamma\left(\frac{l_j}{2} \pm iw_1(j) \pm iw_2(j)\right)}{2\pi^2 \Gamma(l_j)} \cdot \left(\frac{\sigma^2}{2}\right)^{l_j} \times \prod_{m=1}^{N+1} \exp\left(-\frac{\tau_m \sigma^2}{2} \cdot k_m^2\right) \sinh(2\pi k_m) 2k_m dk_m, \quad (8)$$

where the integral in the right-hand side converges absolutely.

Remark 5. Taking $N = 0$ gives a formula for the total mass of the measure $d\mathcal{M}_{\sigma^2}$, which agrees with the result first proposed in [Stanford and Witten, 2017] and obtained rigorously in [Bauerschmidt, Losev, and Wildemann, 2024].

Remark 6. This is also true for $l_j = 0$ if we interpret

$$\frac{\Gamma(\pm ik \pm iw)}{2\pi^2 \Gamma(0)} := \frac{\delta(k-w)}{\sinh(2\pi k) 2k}, \quad (9)$$

where $\delta(\cdot)$ is the delta function.

As an immediate corollary we obtain formulae for moments of the observables.

Corollary 7. For positive integers l the moments of observables are given by

$$\begin{aligned} \int \mathcal{O}^l(\phi; s, t) d\mathcal{M}_{\sigma^2}(\phi) &= \int_{\mathbb{R}_+^2} \frac{\Gamma(\frac{l}{2} \pm ik_1 \pm ik_2)}{2\pi^2 \Gamma(l)} \cdot \left(\frac{\sigma^2}{2}\right)^l \\ &\times \exp\left(-\frac{(t-s)\sigma^2}{2} \cdot k_1^2 - \frac{(1-(t-s))\sigma^2}{2} \cdot k_2^2\right) \sinh(2\pi k_1) 2k_1 \sinh(2\pi k_2) 2k_2 dk_1 dk_2. \end{aligned} \quad (10)$$

As another useful corollary we show that all observables have exponential moments.

Proposition 8. For any $\sigma > 0$ and any $s \neq t \in \mathbb{T}$,

$$\int \exp\left\{\frac{8}{\sigma^2} \mathcal{O}(\phi; s, t)\right\} d\mathcal{M}_{\sigma^2}(\phi) < \infty. \quad (11)$$

Remark 9. The constant $8/\sigma^2$ in the exponential is sharp.

We demonstrate that the computed correlation functions determine the measure uniquely.

Theorem 2. Let \mathcal{P} be a Borel measure on $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$. Suppose that for any $N \geq 0$, any set $\{\mathcal{O}(\phi; s_j, t_j)\}_{j=1}^N$ of non-interlaced observables, and any set $\{l_j\}_{j=1}^N$ of positive integers we have

$$\int \prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) d\mathcal{P}(\phi) = \int \prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) d\mathcal{M}_{\sigma^2}(\phi). \quad (12)$$

Then for any Borel set $A \subset \text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$ we have

$$\mathcal{P}(A) = \mathcal{M}_{\sigma^2}(A). \quad (13)$$

Remark 10. In this work we only consider non-interlaced observables. It would be interesting to compute general correlation functions. It is expected that we get similar formulae, where every crossing contributes an extra factor, called $6j$ -symbol, see [Blommaert et al., 2018; Iliesiu et al., 2019; Mertens et al., 2017]. Theorem 2 tells us that, in principle, it should be possible to deduce this from the correlation functions already computed in Theorem 1.

1.3 Stress-energy tensor

In this section we make sense of the stress-energy tensor correlation functions studied in [Stanford and Witten, 2017]. Formally, in Schwarzian Field Theory the stress-energy tensor is given by the Schwarzian derivative $\mathcal{S}(\tan(\pi\phi), \cdot)$. However, the Schwarzian derivative is a non-linear functional of the field which is a priori not well-defined on the support of $d\mathcal{M}_{\sigma^2}$ due to lack of sufficient regularity (essentially, the measure is supported on diffeomorphisms of regularity $C^{3/2-}$). Nevertheless, we show that it is possible to make sense of its correlation functions.

Our approach is based on the fact that, at least for sufficiently smooth ϕ , we can express the stress-energy tensor $\mathcal{S}(\tan(\pi\phi), \cdot)$ using cross-ratio observables $\mathcal{O}(\phi; s, t)$ by

$$\mathcal{S}(\tan(\pi\phi), \tau) = 6 \lim_{\substack{s \rightarrow \tau^- \\ t \rightarrow \tau^+}} (\mathcal{O}^2(\phi; s, t) - (t - s)^{-2}). \quad (14)$$

Even though the limit in the right-hand side might not exist on the support of $d\mathcal{M}_{\sigma^2}$, we can *define* correlation functions of the stress-energy tensor $\mathcal{S}(\tan(\pi\phi), \cdot)$ by taking the corresponding limit of the correlation functions of the expression appearing in the right-hand side of (14). However, in order to match the resulting correlation functions to formulae obtained by formal differentiation of the partition function in [Stanford and Witten, 2017] and [Bauerschmidt, Losev, and Wildemann, 2024] we need to add an extra shift by a constant. This is often expected when dealing with correlations of classically ill-posed composite fields.

In the following Theorem we compute the joint correlation functions of cross-ratio observables and stress-energy tensors at non-coinciding points.

Theorem 3. *Let $N \geq 0$, $M \geq 0$ be integers, $\{\mathcal{O}(\phi; s_j, t_j)\}_{j=1}^N$ be a set of non-interlaced observables, $\{l_j\}_{j=1}^N$ be non-negative integers, $\{r_p\}_{p=1}^M$ be distinct points on the unit circle \mathbb{T} , which are different from $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$. For $t \in \mathbb{T}$ we write $k(t)$ for the Fourier variable which corresponds to the diagram domain whose boundary contains t . Then for any $\sigma > 0$,*

$$\begin{aligned} & \lim_{\varepsilon_1, \dots, \varepsilon_M \rightarrow 0^+} \int 6^M \prod_{p=1}^M \left(\mathcal{O}^2(\phi; r_p, r_p + \varepsilon_p) - \varepsilon_p^{-2} - \frac{\sigma^4}{240} \right) \prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) d\mathcal{M}_{\sigma^2}(\phi) \\ &= \int_{\mathbb{R}_+^{N+1}} \sigma^{4M} \prod_{p=1}^M k^2(r_p) \prod_{j=1}^N \frac{\Gamma\left(\frac{l_j}{2} \pm iw_1(j) \pm iw_2(j)\right)}{2\pi^2 \Gamma(l_j)} \cdot \left(\frac{\sigma^2}{2}\right)^{l_j} \\ & \quad \times \prod_{m=1}^{N+1} \exp\left(-\frac{\tau_m \sigma^2}{2} \cdot k_m^2\right) \sinh(2\pi k_m) 2k_m dk_m. \quad (15) \end{aligned}$$

Remark 11. *In fact, using integration by parts in τ it is easy to see that the right-hand side depends only on M and does not depend on $\{r_p\}_{p=1}^M$.*

Remark 12. *The Schwarzian derivative also appears in the expression of the stress-energy tensor of Loewner energy, see [Sung and Wang, 2024].*

As an immediate corollary, we verify that non-coinciding stress-energy tensor correlation functions defined this way are given by the moments of spectral density, and thus agree with the expressions obtained in [Stanford and Witten, 2017] and [Bauerschmidt, Losev, and Wildemann, 2024] by differentiating the partition function. This was conjectured and numerically verified in [Mertens et al., 2017] for a one-point function (i.e. for $M = 1$).

Corollary 13. *Let $M > 0$ be an integer, and $\{r_p\}_{p=1}^M$ be distinct points on the unit circle \mathbb{T} . Then for any $\sigma > 0$,*

$$\begin{aligned} & \lim_{\varepsilon_1, \dots, \varepsilon_M \rightarrow 0^+} \int 6^M \prod_{p=1}^M \left(\mathcal{O}^2(\phi; r_p, r_p + \varepsilon_p) - \varepsilon_p^{-2} - \frac{\sigma^4}{240} \right) d\mathcal{M}_{\sigma^2}(\phi) \\ &= \int_{\mathbb{R}_+} \sigma^{4M} \exp\left(-\frac{\sigma k^2}{2}\right) \sinh(2\pi k) 2k^{2M+1} dk. \quad (16) \end{aligned}$$

Remark 14. *We find it interesting that when we turn (14) into the regularisation, proposed in Theorem 3, we do not introduce any new diverging constants, as it often happens when one makes sense of differential operators when there is not enough regularity.*

It is also worth noting that the introduced shift by $\sigma^4/240$ depends on σ . However, if we change the normalisation and, instead of fixing the length of the circle and varying the temperature, we fix the temperature and vary the length of the circle (as, e.g., in [Mertens et al., 2017]), this term becomes an absolute constant.

1.4 Related literature

The finite measure corresponding to the Schwarzian Field Theory was mathematically constructed in the companion paper [Bauerschmidt, Losev, and Wildemann, 2024]. Its partition function (i.e. total mass) was defined and computed in the same paper. For general references about Schwarzian Field Theory and related topics see references therein.

The cross-ratio correlation functions studied in this work were formally derived in the physics literature in [Mertens et al., 2017] by relating Schwarzian Field Theory to a certain degenerate limit of 2D Liouville CFT, and using conformal bootstrap and the DOZZ formula. Alternatively, the same formulae were formally obtained by connecting JT gravity (which is believed to be holographically dual to the Schwarzian Field Theory) to 2D BF theory and weakly coupled limit of 2D Yang-Mills [Blommaert et al., 2018; Iliesiu et al., 2019]. There has been a lot of progress in understanding both 2D Liouville CFT (and conformal bootstrap there) and 2D Yang-Mills from the probabilistic point of view. Below we briefly review some recent progress in these directions. However, as of now there is no rigorous connection between Schwarzian Field Theory and either of these theories from the probabilistic perspective. It would be interesting to rigorously understand relationships between Schwarzian Field Theories and both 2D Yang-Mills and 2D Liouville CFT as predicted in the physics literature.

Two dimensional Liouville CFT was rigorously constructed as a probability measure on a surface in [David et al., 2016; Guillarmou et al., 2019]. For these probability measures the DOZZ formula was rigorously obtained in [Kupiainen et al., 2017], and the conformal bootstrap solution was derived in [Guillarmou et al., 2020, 2021]. See [Guillarmou et al., 2024] and the references therein for the review. In addition, it is possible to make sense of the stress-energy tensor and derive conformal Ward identities for probabilistic Liouville CFT [Kupiainen and Oikarinen, 2019; Kupiainen et al., 2019; Oikarinen, 2022]. The ideas used to make sense of the stress-energy tensor in the context of Liouville CFT are similar to the approaches we take here and in [Bauerschmidt, Losev, and Wildemann, 2024].

Two dimensional Yang-Mills theory, in turn, can also be defined as a probability measure. One of the main approaches is to view it as a probability measure on observables given by Wilson loops [Lévy, 2008; Sengupta, 1997]. Alternatively, one can also define 2D Yang-Mills as a random distribution [Chevyrev, 2018]. Recently, it was proved in [Chandra et al., 2020; Chevyrev and Shen, 2023] that these measures are invariant under the corresponding Langevin dynamics, providing alternative perspective in the context of stochastic quantisation. In addition, Wilson loops in 2D Yang-Mills are known to satisfy the Makeenko-Migdal equation [Driver et al., 2017, 2016; Levy, 2018], which, in particular allows one to study certain limits of 2D Yang-Mills measure [Dahlqvist and Norris, 2017].

Schwarzian Field Theory has also been studied from the algebraic and geometric perspective [Alekseev and Shatashvili, 1989, 2021; Alekseev et al., 2024]. From this point of view, it is naturally associated to a particular coadjoint orbit of the Virasoro algebra. Changing the coadjoint orbit essentially corresponds to changing the regularisation parameter α in (31), where $\alpha = \pi$ corresponds to the Schwarzian Field Theory. In [Alekseev et al., 2024] it was shown that coadjoint orbits corresponding to $\alpha \in i\mathbb{R}$ (which are not $\mathrm{PSL}(2, \mathbb{R})$ invariant) admit global equivariant Darboux charts. Physically this amounts to a bosonisation (diagonalisation) of the theory. As an application, they are

able to obtain certain formulae for lower order correlation functions. We remark that the observables considered in [Alekseev et al., 2024] are, basically, inverses of $\mathcal{O}_\alpha(\phi; s, t)$, as defined in (38).

1.5 Organisation of the paper

In Section 2 we recall the definition of Schwarzian measure, corresponding to the Schwarzian Field Theory, from [Bauerschmidt, Losev, and Wildemann, 2024].

In Section 3 we show how to express correlation functions of $d\mathcal{M}_{\sigma^2}$ as a $\alpha \nearrow \pi$ limit of correlation functions of regularised observables $\mathcal{O}_\alpha(\phi; \cdot, \cdot)$ with respect to regularised measures $d\mathcal{N}_{\sigma^2}^\alpha$ over $\text{Diff}^1(\mathbb{T})$, with regularisation parameter α (see Proposition 24). We also show how the derived expressions can be reduced to computing appropriate expectations for the case $\alpha = 0$ (see Proposition 25). The latter is computed in Section 4.

All the results from Section 1 are proved in Section 5.

Appendices A, and B contain technical results, which are used throughout the proof.

1.6 Preliminaries and notations

Throughout the paper we will be using the following notations.

1. The unit circle is denoted by $\mathbb{T} = [0, 1]/\{0 \sim 1\}$, the non-negative real numbers are denoted by $\mathbb{R}_+ = [0, \infty)$, and for the open disk in the complex plain of radius r we use $\mathbb{D}_r = \{z \in \mathbb{C} : |z| < r\}$. Moreover, for $s, t \in \mathbb{T}$ we write $t - s$ for the length of the interval going from s to t in positive direction. In particular, $t - s \in [0, 1)$.
2. We use $\text{Diff}^k(\mathbb{T})$ for the set of oriented C^k -diffeomorphisms of \mathbb{T} , i.e. $\phi \in \text{Diff}^k(\mathbb{T})$ can be identified with a k -times continuously differentiable function $\phi: \mathbb{R} \rightarrow \mathbb{R}$ satisfying $\phi(\tau + 1) = \phi(\tau) + 1$ and $\phi'(\tau) > 0$ for all $\tau \in \mathbb{R}$. Note that $\text{Diff}^k(\mathbb{T})$ is not a linear space. The topology on $\text{Diff}^k(\mathbb{T})$ is inherited from the natural topology on $C^k(\mathbb{T})$. It turns $\text{Diff}^k(\mathbb{T})$ into a Polish (separable completely metrisable) space as well as a topological group.
It is also useful to consider diffeomorphisms of $[0, 1]$ (or more generally of $[0, T]$) that are not periodic, and we denote by $\text{Diff}^k[0, T]$ the set of oriented C^k -diffeomorphisms of $[0, T]$ satisfying $\phi'(t) > 0$, $\phi(0) = 0$, and $\phi(T) = T$. Thus the derivatives do not have to match at the endpoints. We further set $C_{0, \text{free}}[0, T] = \{f \in C[0, T] \mid f(0) = 0\}$, and $C_0[0, T] = \{f \in C[0, T] \mid f(0) = f(T) = 0\}$.
3. We will abuse the notation and for $\phi \in \text{Diff}^1(\mathbb{T})$ denote its conjugacy class in $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$ by ϕ as well.
4. Throughout the paper we will be encountering expressions of the form $f(\text{arccosh}[z])$, for various even analytic functions f . Even though $\text{arccosh}[z]$ is not analytic at $z = 0$, the composition $f(\text{arccosh}[z])$ still defines an analytic function around $z = 0$. More precisely, we identify $f(\text{arccosh}[z])$ with $\tilde{f}(\text{arccosh}^2[z])$, where \tilde{f} is an analytic function such that $\tilde{f}(\omega) = f(\sqrt{\omega})$, and $\text{arccosh}^2[z]$ is the analytic function described in Statement 40.

2 Measure construction and observables

In this section we recall the rigorous definition and main properties of the measure corresponding to Schwarzian Field Theory from [Bauerschmidt, Losev, and Wildemann, 2024]. For proofs and further details and discussions see [Bauerschmidt, Losev, and Wildemann, 2024].

2.1 Unnormalised Brownian bridge measure

The definition of the Schwarzian measure is based on the appropriate reparametrisation of unnormalised version of the Brownian bridge measure, which we discuss in this subsection. The unnormalised version of the Brownian bridge measure is a finite measure on $C_{0,\text{free}}[0, T] = \{f \in C[0, T] \mid f(0) = 0\}$ formally corresponding to

$$d\mathcal{B}_{\sigma^2}^{a,T}(\xi) = \exp\left\{-\frac{1}{2\sigma^2} \int_0^T \xi'^2(t) dt\right\} \delta(\xi(0)) \delta(\xi(T) - a) \prod_{\tau \in (0, T)} d\xi(\tau). \quad (17)$$

Definition 15. *The unnormalised Brownian bridge measure with variance $\sigma^2 > 0$ is a finite Borel measure $d\mathcal{B}_{\sigma^2}^{a,T}$ on $C_{0,\text{free}}[0, T]$ such that*

$$\sqrt{2\pi T} \sigma \exp\left\{\frac{a^2}{2T\sigma^2}\right\} d\mathcal{B}_{\sigma^2}^{a,T}(\xi) \quad (18)$$

is the distribution of a Brownian bridge $(\xi(t))_{t \in [0, T]}$ with variance σ^2 and $\xi(0) = 0$, $\xi(T) = a$.

For us it will be important that the unnormalised Brownian bridge measure satisfies the following property.

Proposition 16. *For any $T_1, T_2 > 0$, $a \in \mathbb{R}$ and any positive continuous functional F on $C[0, T_1 + T_2]$, we have*

$$\int F(\xi) d\mathcal{B}_{\sigma^2}^{a, T_1 + T_2}(\xi) = \int_{\mathbb{R}} \int F(\xi_1 \sqcup \xi_2) d\mathcal{B}_{\sigma^2}^{b, T_1}(\xi_1) d\mathcal{B}_{\sigma^2}^{a-b, T_2}(\xi_2) db, \quad (19)$$

where for $f \in C_{0,\text{free}}[0, T_1]$ and $g \in C_{0,\text{free}}[0, T_2]$, we denote by $f \sqcup g \in C_{0,\text{free}}[0, T_1 + T_2]$ the function

$$(f \sqcup g)(t) = \begin{cases} f(t) & \text{if } t \in [0, T_1], \\ f(T_1) + g(t - T_1) & \text{if } t \in (T_1, T_1 + T_2]. \end{cases} \quad (20)$$

2.2 Measure construction

In order to define the Schwarzian measure $d\mathcal{M}_{\sigma^2}$, we first need to construct a finite measure μ_{σ^2} on $\text{Diff}^1(\mathbb{T})$ which is similar to what is known as the Malliavin–Shavgulidze measure, see [Bogachev, 2010, Section 11.5]. Formally, this measure corresponds to

$$d\mu_{\sigma^2}(\phi) = \exp\left\{-\frac{1}{2\sigma^2} \int_0^1 \left(\frac{\phi''(\tau)}{\phi'(\tau)}\right)^2 d\tau\right\} \prod_{\tau \in [0, 1)} \frac{d\phi(\tau)}{\phi'(\tau)}. \quad (21)$$

We can make sense of this measure by defining it as a push-forward of an unnormalised Brownian bridge on $[0, 1]$ with respect to a suitable change of variables. We define μ_{σ^2} by

$$d\mu_{\sigma^2}(\phi) := d\mathcal{B}_{\sigma^2}^{0,1}(\xi) \otimes d\Theta, \quad \text{with } \phi(t) = \Theta + P_{\xi}(t) \pmod{1}, \text{ for } \Theta \in [0, 1), \quad (22)$$

where $d\Theta$ is the Lebesgue measure on $[0, 1)$ and

$$P(\xi)(t) := P_{\xi}(t) := \frac{\int_0^t e^{\xi(\tau)} d\tau}{\int_0^1 e^{\xi(\tau)} d\tau}, \quad (23)$$

The variable Θ corresponds to the value of $\phi(0)$. Note that the map $\xi \mapsto P(\xi)$ is a bijection between $C_{0,\text{free}}[0, 1]$ and $\text{Diff}^1[0, 1]$ with inverse map

$$P^{-1} : \text{Diff}^1[0, 1] \rightarrow C_{0,\text{free}}[0, 1] \quad (24)$$

$$\varphi \mapsto \log \varphi'(\cdot) - \log \varphi'(0). \quad (25)$$

In view of (1) and (21), the unquotiented Schwarzian measure is defined by

$$d\widetilde{\mathcal{M}}_{\sigma^2}(\phi) = \exp \left\{ \frac{2\pi^2}{\sigma^2} \int_0^1 \phi'^2(\tau) d\tau \right\} d\mu_{\sigma^2}(\phi). \quad (26)$$

Since μ_{σ^2} is supported on $\text{Diff}^1(\mathbb{T})$, this defines a Borel measure on $\text{Diff}^1(\mathbb{T})$, which is invariant under the right action of $\text{PSL}(2, \mathbb{R})$.

Proposition 17. *The measure $\widetilde{\mathcal{M}}_{\sigma^2}$ is invariant under the right action of $\text{PSL}(2, \mathbb{R})$. In other words, for any $\psi \in \text{PSL}(2, \mathbb{R})$ and Borel $A \subset \text{Diff}^1(\mathbb{T})$ we have*

$$\widetilde{\mathcal{M}}_{\sigma^2}(\psi \circ A) = \widetilde{\mathcal{M}}_{\sigma^2}(A), \quad (27)$$

where $\psi \circ A := \{\psi \circ \phi \mid \phi \in A\}$.

In particular, we remark that $d\widetilde{\mathcal{M}}_{\sigma^2}$ is an infinite measure since $\text{PSL}(2, \mathbb{R})$ has infinite Haar measure. The Schwarzian measure on $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$ with formal density (1) is defined as the quotient of $d\widetilde{\mathcal{M}}_{\sigma^2}$ by $\text{PSL}(2, \mathbb{R})$, see Proposition 18 and Definition 19 below.

Proposition 18. *There exists a unique Borel measure \mathcal{M}_{σ^2} on $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$ such that for any non-negative continuous functional $F: \text{Diff}^1(\mathbb{T}) \rightarrow \mathbb{R}$,*

$$\int_{\text{Diff}^1(\mathbb{T})} d\widetilde{\mathcal{M}}_{\sigma^2}(\phi) F(\phi) = \int_{\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})} d\mathcal{M}_{\sigma^2}([\phi]) \int_{\text{PSL}(2, \mathbb{R})} d\nu_H(\psi) F(\psi \circ \phi), \quad (28)$$

where the right hand side is well-defined, since the second integral only depends on $[\phi]$.

Definition 19. *The Schwarzian measure is given by \mathcal{M}_{σ^2} .*

One of the main tools which will allow us to compute integrals with respect to $d\mu_{\sigma^2}$ exactly is the following analogue of Girsanov's Theorem.

Lemma 20. *For $g \in \text{Diff}^3[0, 1]$ we denote by L_g the left composition operator on $\text{Diff}^1[0, 1]$:*

$$L_g(\phi) = g \circ \phi. \quad (29)$$

Suppose $f \in \text{Diff}^3[0, 1]$. Denote $b = \log f'(1) - \log f'(0)$. Let $f^\# \mathcal{B}_{\sigma^2}^{a,1} = f_\#^{-1} \mathcal{B}_{\sigma^2}^{a,1}$ be the push-forward of $\mathcal{B}_{\sigma^2}^{a,1}$ under $\mathbf{P}^{-1} \circ L_{f^{-1}} \circ \mathbf{P} = (\mathbf{P}^{-1} \circ L_f \circ \mathbf{P})^{-1}$. Then for any $a \in \mathbb{R}$, $f^\# \mathcal{B}_{\sigma^2}^{a,1}$ is absolutely continuous with respect to $\mathcal{B}_{\sigma^2}^{a-b,1}$ and

$$\frac{df^\# \mathcal{B}_{\sigma^2}^{a,1}(\xi)}{d\mathcal{B}_{\sigma^2}^{a-b,1}(\xi)} = \frac{1}{\sqrt{f'(0)f'(1)}} \exp \left\{ \frac{1}{\sigma^2} \left[\frac{f''(0)}{f'(0)} \mathbf{P}'_\xi(0) - \frac{f''(1)}{f'(1)} \mathbf{P}'_\xi(1) \right] + \frac{1}{\sigma^2} \int_0^1 \mathcal{S}_f(\mathbf{P}_\xi(t)) (\mathbf{P}'_\xi(t))^2 dt \right\}. \quad (30)$$

This change of variables formula was used in [Bauerschmidt, Losev, and Wildemann, 2024] to calculate the partition function of $d\mathcal{M}_{\sigma^2}$, as inspired by [Belokurov and Shavgulidze, 2017].

2.3 Measure regularisation

In order to evaluate expectations with respect to the (finite) quotient measure $d\mathcal{M}_{\sigma^2}$, we approximate the (infinite) unquotiented measure $d\widetilde{\mathcal{M}}_{\sigma^2}$ by a particular family of finite measures. Since these measures are finite, they necessarily break the $\text{PSL}(2, \mathbb{R})$ -invariance. The following regularisation was

proposed in [Belokurov and Shavgulidze, 2017] and used in [Bauerschmidt, Losev, and Wildemann, 2024] to calculate the partition function. For $\alpha \in [0, \pi] \cup i\mathbb{R}$ consider the measures given by

$$d\mathcal{N}_{\sigma^2}^\alpha(\phi) = \exp\left\{\frac{2\alpha^2}{\sigma^2} \int_0^1 \phi'^2(t) dt\right\} d\mathcal{B}_{\sigma^2}^{0,1}(\xi), \quad \text{where } \phi = \mathbf{P}(\xi). \quad (31)$$

By definition, this measure is supported on functions ϕ with $\phi(0) = 0$. In particular, $d\widetilde{\mathcal{M}}_{\sigma^2}$ differs from $d\mathcal{N}_{\sigma^2}^\pi$ only by rotation by the random angle Θ , which is chosen independently and uniformly on \mathbb{T} . This feature allows us to use these measures to calculate expectations with respect to the Schwarzian measure $d\mathcal{M}_{\sigma^2}$.

Proposition 21. *Let $F : \text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R}) \rightarrow [0, \infty]$ be a continuous function. Then*

$$\int_{\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})} F(\phi) d\mathcal{M}_{\sigma^2}(\phi) = \lim_{\alpha \rightarrow \pi^-} \frac{4\pi(\pi - \alpha)}{\sigma^2} \int_{\text{Diff}^1(\mathbb{T})} F(\phi) d\mathcal{N}_{\sigma^2}^\alpha(\phi), \quad (32)$$

where, by slight abuse of notation, we denote the lift of F along the quotient map $\text{Diff}^1(\mathbb{T}) \rightarrow \text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$ by F as well.

Another advantage of using the constructed measures $d\mathcal{N}_{\sigma^2}^\alpha$ is that for them integrals of certain functionals can be computed explicitly, see Section 3.3.

3 Observables and their regularisation

3.1 Observables

The main property of cross-ratio observables (3) is that they define observables on the quotient space $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$.

Proposition 22. *Observables $\mathcal{O}(\phi; s, t)$ are invariant under the action of Möbius transformations. In other words, if $\psi \in \text{PSL}(2, \mathbb{R})$, then*

$$\mathcal{O}(\psi \circ \phi; s, t) = \mathcal{O}(\phi; s, t). \quad (33)$$

In particular, they induce well-defined observables on $\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})$, which we, slightly abusing the notation, also denote by $\mathcal{O}(\phi; \cdot, \cdot)$.

Proof. Let $\psi \in \text{PSL}(2, \mathbb{R})$ be Möbius transformation of the unit circle. Fix $\phi \in \text{Diff}^1(\mathbb{T})$ and $s \neq t \in T$. Denote $F(\tau) = \tan(\pi(\tau - a))$, where $a \in \mathbb{T}$ is such that $a \neq \phi(s)$, and $a \neq \phi(t)$. Also denote $f = F \circ \phi$ and $g = F \circ \psi \circ F^{-1}$. Then

$$g(x) = \frac{ax + b}{cx + d}, \quad (34)$$

for some $a, b, c, d \in \mathbb{R}$ with $ad - bc = 1$, and all $x \in \mathbb{R}$.

We have that

$$\mathcal{O}(\phi; s, t) = |\mathcal{O}_0(f; s, t)|, \quad (35)$$

where

$$\mathcal{O}_0(\phi; s, t) = \frac{\sqrt{\phi'(t)\phi'(s)}}{\phi(t) - \phi(s)}. \quad (36)$$

It is well-known that the cross-ratio $\mathcal{O}_0(f; \cdot, \cdot)$ is invariant under transformations of the form (34). Thus,

$$\mathcal{O}(\phi; s, t) = |\mathcal{O}_0(f; s, t)| = |\mathcal{O}_0(g \circ f; s, t)| = \mathcal{O}(\psi \circ \phi; s, t). \quad (37)$$

□

3.2 Regularised observables

It turns out that in order to be able to compute correlation functions for the measures $d\mathcal{N}_{\sigma_2}^\alpha$ we need to modify the observables too. The new observables will depend on α , but they will agree with (3) when $\alpha = \pi$. Thus, we will calculate the expectations of new observables against $d\mathcal{N}_{\sigma_2}^\alpha$, and take $\alpha \nearrow \pi$.

For $\alpha \in [0, \pi]$ and $0 \leq s \neq t \leq 1$ we define

$$\mathcal{O}_\alpha(\phi; s, t) = \begin{cases} \frac{\alpha \sqrt{\phi'(t)\phi'(s)}}{\sin(\alpha[\phi(t) - \phi(s)])} & \text{if } \alpha > 0, \\ \frac{\sqrt{\phi'(t)\phi'(s)}}{\phi(t) - \phi(s)} & \text{if } \alpha = 0. \end{cases} \quad (38)$$

Notice that $\mathcal{O}_\pi(\phi; s, t) = \mathcal{O}(\phi; s, t)$. However, even though it is true that $\lim_{\alpha \rightarrow \pi^-} \mathcal{O}_\alpha(\phi; t, s) = \mathcal{O}_\pi(\phi; s, t)$, this limit is not sufficiently uniform in ϕ , and so it will not allow us to pass to the $\alpha \nearrow \pi$ limit for the integrals of new observables against $d\mathcal{N}_{\sigma_2}^\alpha$ (in fact, doing so leads to wrong answer).

Also observe that, generally, for $\alpha < \pi$ we have $\mathcal{O}_\alpha(\phi; s, t) \neq \mathcal{O}_\alpha(\phi; t, s)$, since, recall, with our notation for \mathbb{T} , $\phi(t) - \phi(s)$ and $\phi(s) - \phi(t)$ sum to 1. This will be important in Lemma 23, when we exploit this property to approximate $\mathcal{O}_\pi(\phi; s, t)$ sufficiently uniformly in $\alpha \nearrow \pi$.

3.3 Observables approximation

Our goal in this subsection is to express correlation functions of $\mathcal{O}(\phi; \cdot, \cdot)$ with respect to $d\mathcal{M}_{\sigma_2}$ as appropriate limits of correlation functions of $\mathcal{O}_\alpha(\phi; \cdot, \cdot)$ with respect to $d\mathcal{N}_{\sigma_2}^\alpha$, see Proposition 24. For this, we first need to approximate products of $\mathcal{O}(\phi; \cdot, \cdot)$ with products of $\mathcal{O}_\alpha(\phi; \cdot, \cdot)$ uniformly in ϕ when $\alpha \rightarrow \pi$, see Lemma 23.

Let $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ be points on \mathbb{T} . Enumerate the union of points $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ counter-clockwise as $r_1 \leq r_2 \leq \dots \leq r_{2N} \leq r_{2N+1} = r_1 + 1$.

For $x \in \mathbb{T}$ define rotation as

$$\left(\mathbb{R}^x \phi\right)(\tau) = \phi(\tau+x) - \phi(x), \quad \mathbb{R}^x s_j = \min\{s_j - x, t_j - x\}, \quad \mathbb{R}^x t_j = \max\{s_j - x, t_j - x\}. \quad (39)$$

Lemma 23. *Let $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ be points on \mathbb{T} , and $\{l_j\}_{j=1}^N$ be positive numbers. Let also $\{r_j\}_{j=1}^N$ be as above. Then*

$$\sum_{m=1}^{2N} \left[\left(\phi(r_{m+1}) - \phi(r_m) \right) \prod_{j=1}^N \mathcal{O}_\alpha^{l_j}(\mathbb{R}^{r_{m+1}} \phi; \mathbb{R}^{r_{m+1}} s_j, \mathbb{R}^{r_{m+1}} t_j) \right] \\ \stackrel{\alpha \rightarrow \pi}{=} \left[\prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) \right] (1 + o(1)), \quad (40)$$

uniformly in $\phi \in \text{Diff}^1(\mathbb{T})$.

Proof. We assume that $\alpha > \pi - \frac{1}{100}$. Denote

$$s_j^m = r_{m+1} + \mathbb{R}^{r_{m+1}} s_j, \quad t_j^m = r_{m+1} + \mathbb{R}^{r_{m+1}} t_j. \quad (41)$$

Note that $\forall m, j$,

$$\frac{\mathcal{O}_\alpha(\mathbb{R}^{r_{m+1}} \phi; \mathbb{R}^{r_{m+1}} s_j, \mathbb{R}^{r_{m+1}} t_j)}{\mathcal{O}(\phi; s_j, t_j)} = \frac{\alpha \sin(\pi[\phi(t_j^m) - \phi(s_j^m)])}{\pi \sin(\alpha[\phi(t_j^m) - \phi(s_j^m)])}. \quad (42)$$

Thus, it is sufficient to prove that

$$\sum_{m=1}^{2N} \left[\left(\phi(r_{m+1}) - \phi(r_m) \right) \prod_{j=1}^N \frac{\sin^{l_j} \left(\pi [\phi(t_j^m) - \phi(s_j^m)] \right)}{\sin^{l_j} \left(\alpha [\phi(t_j^m) - \phi(s_j^m)] \right)} \right] \stackrel{\alpha \rightarrow \pi}{=} 1 + o(1), \quad (43)$$

We take $\delta = \sqrt{\pi - \alpha} \in (0, 1/10)$. Let $X = X(\phi) = \{m \leq 2N : (\phi(r_{m+1}) - \phi(r_m)) > \delta\}$.

Note that since $\sin(y)/y$ is decreasing in y we have

$$\frac{\sin \left(\pi [\phi(t_j^m) - \phi(s_j^m)] \right)}{\sin \left(\alpha [\phi(t_j^m) - \phi(s_j^m)] \right)} \leq \frac{\pi}{\alpha}. \quad (44)$$

Therefore, for any $\phi \in \text{Diff}^1(\mathbb{T})$,

$$\sum_{m \notin X} \left[\left(\phi(r_{m+1}) - \phi(r_m) \right) \prod_{j=1}^N \frac{\sin^{l_j} \left(\pi [\phi(t_j^m) - \phi(s_j^m)] \right)}{\sin^{l_j} \left(\alpha [\phi(t_j^m) - \phi(s_j^m)] \right)} \right] \leq 2N \left(\frac{\pi}{\alpha} \right)^{\sum_{j=1}^N l_j} \delta. \quad (45)$$

Moreover, observe that for $m \in X$ we have $\phi(t_j^m) - \phi(s_j^m) \leq 1 - \delta$, so applying Statement 38 we get

$$\left| \log \prod_{j=1}^N \frac{\sin^{l_j} \left(\pi [\phi(t_j^m) - \phi(s_j^m)] \right)}{\sin^{l_j} \left(\alpha [\phi(t_j^m) - \phi(s_j^m)] \right)} \right| \leq \frac{(\pi - \alpha) \sum_{j=1}^N l_j}{\sin(\pi\delta/2)}, \quad \text{for } m \in X, \quad (46)$$

$$\sum_{m \in X} \left(\phi(r_{m+1}) - \phi(r_m) \right) \geq 1 - 2N\delta, \quad (47)$$

which finishes the proof. \square

Proposition 24. *Let $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ be points on \mathbb{T} , and $\{l_j\}_{j=1}^N$ be positive numbers. Then*

$$\begin{aligned} & \int_{\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})} \left[\prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) \right] d\mathcal{M}_{\sigma^2}(\phi) \\ &= \lim_{\alpha \rightarrow \pi^-} \frac{4\pi(\pi - \alpha)}{\sigma^2} \int_{\mathbb{T}} \left[\int_{\text{Diff}^1(\mathbb{T})} \phi'(0) \prod_{j=1}^N \mathcal{O}_{\alpha}^{l_j}(\phi; \mathbf{R}^x s_j, \mathbf{R}^x t_j) d\mathcal{N}_{\sigma^2}^{\alpha}(\phi) \right] dx. \end{aligned} \quad (48)$$

Proof. From Proposition 21 and Lemma 23 we get

$$\begin{aligned} & \int_{\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})} \left[\prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) \right] d\mathcal{M}_{\sigma^2}(\phi) \\ &= \lim_{\alpha \rightarrow \pi^-} \frac{4\pi(\pi - \alpha)}{\sigma^2} \int_{\text{Diff}^1(\mathbb{T})} \sum_{m=1}^{2N} \left[\left(\phi(r_{m+1}) - \phi(r_m) \right) \prod_{j=1}^N \mathcal{O}_{\alpha}^{l_j}(\mathbf{R}^{r_{m+1}} \phi; \mathbf{R}^{r_{m+1}} s_j, \mathbf{R}^{r_{m+1}} t_j) \right] d\mathcal{N}_{\sigma^2}^{\alpha}(\phi) \end{aligned} \quad (49)$$

Moreover, since

$$\forall x \in (r_m, r_{m+1}) : \quad \mathcal{O}_{\alpha}(\mathbf{R}^x \phi; \mathbf{R}^x s_j, \mathbf{R}^x t_j) = \mathcal{O}_{\alpha}(\mathbf{R}^{r_{m+1}} \phi; \mathbf{R}^{r_{m+1}} s_j, \mathbf{R}^{r_{m+1}} t_j), \quad (50)$$

we get

$$\begin{aligned} & \left(\phi(r_{m+1}) - \phi(r_m) \right) \prod_{j=1}^N \mathcal{O}_\alpha^{l_j}(\mathbf{R}^{r_{m+1}}\phi; \mathbf{R}^{r_{m+1}}s_j, \mathbf{R}^{r_{m+1}}t_j) \\ &= \int_{r_m}^{r_{m+1}} \phi'(x) \prod_{j=1}^N \mathcal{O}_\alpha^{l_j}(\mathbf{R}^x\phi; \mathbf{R}^xs_j, \mathbf{R}^xt_j) dx. \end{aligned} \quad (51)$$

Combining this with (49) and Tonelli's Theorem, we obtain that

$$\begin{aligned} & \int_{\text{Diff}^1(\mathbb{T})/\text{PSL}(2, \mathbb{R})} \left[\prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) \right] d\mathcal{N}_{\sigma^2}(\phi) \\ &= \lim_{\alpha \rightarrow \pi^-} \frac{4\pi(\pi - \alpha)}{\sigma^2} \int_{\mathbb{T}} \left[\int_{\text{Diff}^1(\mathbb{T})} (\mathbf{R}^x\phi)'(0) \prod_{j=1}^N \mathcal{O}_\alpha^{l_j}(\mathbf{R}^x\phi; \mathbf{R}^xs_j, \mathbf{R}^xt_j) d\mathcal{N}_{\sigma^2}^\alpha(\phi) \right] dx, \end{aligned} \quad (52)$$

which together with invariance of $d\mathcal{N}_{\sigma^2}^\alpha$ under \mathbf{R}^x finishes the proof. \square

3.4 Observables and the weight

Now we reduce the integral with respect to $d\mathcal{N}_{\sigma^2}^\alpha$ obtained in Proposition 24 to a calculation of an appropriate integral with respect to $d\mathcal{N}_{\sigma^2}^0$.

Proposition 25. *Let $\left\{ \mathcal{O}_\alpha(\phi; s_j, t_j) \right\}_{j=1}^N$ be a set of observables, and $\{l_j\}_{j=1}^N$ be real numbers. Then*

$$\begin{aligned} & \int_{\text{Diff}^1(\mathbb{T})} \phi'(0) \prod_{j=1}^N \mathcal{O}_\alpha^{l_j}(\phi; s_j, t_j) d\mathcal{N}_{\sigma^2}^\alpha(\phi) \\ &= \int_{\text{Diff}^1(\mathbb{T})} \mathcal{O}_0(\phi; 0, 1) \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \mathcal{O}_0(\phi; 0, 1) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\phi; s_j, t_j) d\mathcal{N}_{\sigma^2}^0(\phi). \end{aligned} \quad (53)$$

Proof. Take

$$f(t) = \frac{1}{2} \left[\frac{1}{\tan \frac{\alpha}{2}} \tan \left(\alpha \left(t - \frac{1}{2} \right) \right) + 1 \right]. \quad (54)$$

It is easy to check that

$$\mathcal{S}_f(t) = 2\alpha^2, \quad f'(0) = f'(1) = \frac{\alpha}{\sin \alpha}, \quad -\frac{f''(0)}{f'(0)} = \frac{f''(1)}{f'(1)} = 2\alpha \tan \frac{\alpha}{2}. \quad (55)$$

Thus, it follows from Lemma 20 that for any non-negative continuous functional F on $\text{Diff}^1(\mathbb{T})$ we have

$$\begin{aligned} & \int_{\text{Diff}^1(\mathbb{T})} F(\phi) d\mathcal{N}_{\sigma^2}^0(\phi) \\ &= \frac{\sin \alpha}{\alpha} \int_{\text{Diff}^1(\mathbb{T})} F(f \circ \phi) \exp \left\{ -\frac{4\alpha}{\sigma^2} \tan \frac{\alpha}{2} \cdot \phi'(0) + \frac{2\alpha^2}{\sigma^2} \int_0^1 \phi'^2(t) dt \right\} d\mathcal{N}_{\sigma^2}^0(\phi). \end{aligned} \quad (56)$$

Now we choose F to be

$$F(\phi) = \phi'(0) \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \cdot \phi'(0) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\phi; s_j, t_j), \quad (57)$$

which guaranties that

$$F(f \circ \phi) = \frac{\alpha}{\sin \alpha} \cdot \phi'(0) \exp \left\{ \frac{4\alpha}{\sigma^2} \tan \frac{\alpha}{2} \cdot \phi'(0) \right\} \prod_{j=1}^N \mathcal{O}_\alpha^{l_j}(\phi; s_j, t_j). \quad (58)$$

Observing that

$$\phi'(0) = \mathcal{O}_0(\phi; 0, 1) \quad (59)$$

gives the desired. \square

4 Main calculations

The main goal of this Section is to calculate the integral involving $\mathcal{O}_0(\phi; s, t)$ with respect to $d\mathcal{N}_\sigma^0$ appearing Proposition 25. An important feature of $\mathcal{O}_0(\phi; s, t)$ that we use here is that it is homogeneous in ϕ . In other words, it does not change if multiply ϕ by a constant. This means that instead of taking $\phi = P(\xi)$, as in (31), we can consider $Q(\xi)$, where we define

$$Q(\xi)(t) := Q_\xi(t) := \int_0^t e^{\xi(s)} ds. \quad (60)$$

An advantage of using Q instead of P is that it is given by a more "local" and simpler expression. We emphasise again that

$$\mathcal{O}_0(P_\xi; s, t) = \mathcal{O}_0(Q_\xi; s, t). \quad (61)$$

Therefore, in order to calculate the integral from the right-hand side of Proposition 25 it is sufficient to study integrals involving $\mathcal{O}_0(Q_\xi; \cdot, \cdot)$ with respect to $d\mathcal{B}_\sigma^{0,1}(\xi)$.

4.1 Key Lemma

The following Lemma is key for the main calculation. Informally, it means that when taking integrals with respect to $d\mathcal{B}_\sigma^{a,T}$ we can trade exponentials of $\mathcal{O}_0(Q_\xi; \cdot, \cdot)$ for the value of parameter a .

Lemma 26. *Fix $T > 0$. Let $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ be points on $[0, T]$, and $\{l_j\}_{j=1}^N$ be positive numbers. Then for any $z < 0$, we have*

$$\int \exp \left\{ \frac{z}{\sigma^2} \mathcal{O}_0(Q_\xi; 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(Q_\xi; s_j, t_j) d\mathcal{B}_\sigma^{a,T}(\xi) = \int \prod_{j=1}^N \mathcal{O}_0^{l_j}(Q_\xi; s_j, t_j) d\mathcal{B}_\sigma^{b,T}(\xi), \quad (62)$$

where $b = 2 \operatorname{arccosh} \left[\cosh \left(\frac{a}{2} \right) - \frac{z}{4} \right]$.

Proof. For $s \in [0, 1]$ define $\tilde{\xi}(s) = \xi(sT)$. If ξ is distributed according to $d\mathcal{B}_\sigma^{p,T}(\xi)$ for some p , then $\tilde{\xi}$ is distributed according to $d\mathcal{B}_\sigma^{p,1}(\tilde{\xi})$.

Notice that

$$\mathcal{O}_0(Q_\xi; s, t) = \frac{1}{T} \mathcal{O}_0(P(\tilde{\xi}); s/T, t/T). \quad (63)$$

Therefore,

$$\begin{aligned} & \int \exp \left\{ \frac{z}{\sigma^2} \mathcal{O}_0(Q_\xi; 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(Q_\xi; s_j, t_j) d\mathcal{B}_\sigma^{a,T}(\xi) \\ &= T^{-\sum_{j=1}^N l_j} \int \exp \left\{ \frac{z}{\sigma^2 T} \mathcal{O}_0(P(\tilde{\xi}); 0, 1) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(P(\tilde{\xi}); s_j/T, t_j/T) d\mathcal{B}_\sigma^{a,1}(\tilde{\xi}) \end{aligned} \quad (64)$$

Consider $f \in \text{Diff}^3[0, 1]$ given by

$$f(t) = \frac{e^\lambda t}{(e^\lambda - 1)t + 1}, \quad (65)$$

with λ to be chosen later. It is easy to see that

$$f'(t) = \frac{e^\lambda}{((e^\lambda - 1)t + 1)^2}, \quad f''(t) = -\frac{2e^\lambda(e^\lambda - 1)}{((e^\lambda - 1)t + 1)^3}, \quad \mathcal{S}_f(t) = 0. \quad (66)$$

Moreover,

$$\frac{f''(0)}{f'(0)} = 2 - 2e^\lambda, \quad \frac{f''(1)}{f'(1)} = 2e^{-\lambda} - 2. \quad (67)$$

Also notice that since f is a fractional linear transformation, we get that observables are invariant under the left composition with f ,

$$\mathcal{O}_0(\phi; s, t) = \mathcal{O}_0(f \circ \phi; s, t), \quad \text{for all } \phi \in \text{Diff}^1(\mathbb{T}), s, t \in \mathbb{T}. \quad (68)$$

Therefore, by Lemma 20 we get

$$\begin{aligned} & \int \exp \left\{ \frac{z}{\sigma^2 T} \mathcal{O}_0(\tilde{\xi}; 0, 1) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\tilde{\xi}; s_j/T, t_j/T) d\mathcal{B}_{\sigma^2 T}^{a,1}(\tilde{\xi}) \\ &= \int \exp \left\{ \frac{z}{\sigma^2 T} \mathcal{O}_0(\tilde{\xi}; 0, 1) + \frac{2 - 2e^\lambda}{\sigma^2 T} \mathbf{P}'(\tilde{\xi})(0) + \frac{2 - 2e^{-\lambda}}{\sigma^2 T} \mathbf{P}'(\tilde{\xi})(1) \right\} \\ & \quad \times \prod_{j=1}^N \mathcal{O}_0^{l_j}(\tilde{\xi}; s_j/T, t_j/T) d\mathcal{B}_{\sigma^2 T}^{a+2\lambda,1}(\tilde{\xi}) \end{aligned} \quad (69)$$

Notice that if $\tilde{\xi} \sim d\mathcal{B}_{\sigma^2 T}^{a+2\lambda,1}(\tilde{\xi})$, then

$$\mathbf{P}'(\tilde{\xi})(0) = e^{-\lambda - \frac{a}{2}} \mathcal{O}_0(\tilde{\xi}; 0, 1), \quad \mathbf{P}'(\tilde{\xi})(1) = e^{\lambda + \frac{a}{2}} \mathcal{O}_0(\tilde{\xi}; 0, 1). \quad (70)$$

Therefore,

$$\frac{2 - 2e^\lambda}{\sigma^2 T} \mathbf{P}'(\tilde{\xi})(0) + \frac{2 - 2e^{-\lambda}}{\sigma^2 T} \mathbf{P}'(\tilde{\xi})(1) = \frac{4[\cosh(\lambda + \frac{a}{2}) - \cosh(\frac{a}{2})]}{\sigma^2 T} \mathcal{O}_0(\tilde{\xi}; 0, 1). \quad (71)$$

Taking

$$\lambda = \text{arccosh} \left[\cosh \left(\frac{a}{2} \right) - \frac{z}{4} \right] - \frac{a}{2} \quad (72)$$

we get the desired. \square

Now we can also prove that observables $\mathcal{O}(\mathbf{Q}_\xi; \cdot, \cdot)$ have exponential moments.

Proposition 27. *Let $T > 0$ and $a \in \mathbb{R}$. For any $0 \leq s < t \leq T$ we have*

$$\int \exp \left\{ \frac{8}{\sigma^2} \mathcal{O}_0(\mathbf{Q}_\xi; s, t) \right\} d\mathcal{B}_{\sigma^2}^{a,T}(\xi) \leq \frac{1}{\sqrt{2\pi T}\sigma} \exp \left(\frac{2000}{(t-s)\sigma^2} - \frac{a^2}{2T\sigma^2} \right). \quad (73)$$

In particular, the expression above is finite.

Proof. We divide the interval $[0, T]$ into three smaller intervals $[0, s]$, $[s, t]$ and $[t, T]$. Let $\xi_1 \in C_{0, \text{free}}[0, s]$, $\xi_2 \in C_{0, \text{free}}[0, t - s]$, and $\xi_3 \in C_{0, \text{free}}[0, T - t]$. Notice that

$$\mathcal{O}_0(\mathbb{Q}(\xi_1 \sqcup \xi_2 \sqcup \xi_3); s, t) = \mathcal{O}_0(\mathbb{Q}(\xi_2); 0, t - s). \quad (74)$$

Thus, using (19) we get

$$\begin{aligned} \int \exp \left\{ \frac{8}{\sigma^2} \mathcal{O}_0(\mathbb{Q}_\xi; s, t) \right\} d\mathcal{B}_{\sigma^2}^{a, T}(\xi) &= \int_{\mathbb{R}^2} \left[\int d\mathcal{B}_{\sigma^2}^{u, s}(\xi_1) \right] \\ &\times \left[\int \exp \left\{ \frac{8}{\sigma^2} \mathcal{O}_0(\mathbb{Q}(\xi_2); 0, t - s) \right\} d\mathcal{B}_{\sigma^2}^{v, t-s}(\xi_2) \right] \\ &\times \left[\int d\mathcal{B}_{\sigma^2}^{a-u-v, T-t}(\xi_3) \right] du dv \quad (75) \end{aligned}$$

Therefore, it is sufficient to prove the Proposition for $s = 0$ and $t = T$.

From Lemma 36 there exists $\varepsilon > 0$ such that

$$\int \exp \{ \varepsilon \mathcal{O}_0(\mathbb{Q}_\xi; 0, T) \} d\mathcal{B}_{\sigma^2}^{a, T}(\xi) < \infty. \quad (76)$$

Moreover, from Lemma 26 for any $z < 0$ we get that

$$\int \exp \left\{ \frac{z}{\sigma^2} \mathcal{O}_0(\mathbb{Q}_\xi; 0, T) \right\} d\mathcal{B}_{\sigma^2}^{a, T}(\xi) = \frac{1}{\sqrt{2\pi T \sigma}} \exp \left(-\frac{2}{T\sigma^2} \operatorname{arccosh}^2 \left[\cosh \left(\frac{a}{2} \right) - \frac{z}{4} \right] \right). \quad (77)$$

Using Lemma 37 and (76) we deduce that the left-hand side is absolutely convergent for all $z \in \mathbb{D}_8$ and the equality above holds for all $z \in \mathbb{D}_8$. Applying Statement 39 we get that for all $z \in [0, 8)$,

$$\int \exp \left\{ \frac{z}{\sigma^2} \mathcal{O}_0(\mathbb{Q}_\xi; 0, T) \right\} d\mathcal{B}_{\sigma^2}^{a, T}(\xi) \leq \frac{1}{\sqrt{2\pi T \sigma}} \exp \left(\frac{2000}{T\sigma^2} - \frac{a^2}{2T\sigma^2} \right). \quad (78)$$

Taking limit $z \rightarrow 8$ we finish the proof. \square

4.2 Diagrammatic representation on an interval

Here we describe a diagrammatic representation for observables $\mathcal{O}_0(\phi; \cdot, \cdot)$ defined on the interval $[0, T]$. This representation is similar to the one described for the circle.

Let $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ be points on $[0, T]$, such that $\forall j : s_j < t_j$. Let also $\{\mathcal{O}_0(\phi; s_j, t_j)\}_{j=1}^N$ be a set of observables. We are going to represent them as a diagram on the interval $[0, T]$.


We draw the interval $[0, T]$ and all the points $\{s_j\}_{j=1}^N$ and $\{t_j\}_{j=1}^N$ on it. For all $1 \leq j \leq N$ we connect the point s_j with the point t_j with a half-circle in the upper half-plane.

Definition 28. *We say that a set of observables $\{\mathcal{O}_0(\phi; s_j, t_j)\}_{j=1}^N$ is non-interlaced if interiors of all drawn half-circles on the corresponding diagram are pairwise non-intersecting. In other words, $\forall p, q \in \{1, \dots, N\}$ we have that one of the following holds: $s_p \leq s_q < t_q \leq t_p$, $s_p < t_p \leq s_q < t_q$, $s_q \leq s_p < t_p \leq t_q$, or $s_q < t_q \leq s_p < t_p$.*

Let $\{\mathcal{O}_0(\phi; s_j, t_j)\}_{j=1}^N$ be a set of non-interlaced observables. The N drawn chords on the corresponding diagram divide the unit disk into $N + 1$ connected domains, one of which is unbounded and N other are bounded. We number the bounded domains with integers from 1 to N , and assign number 0 to the unbounded domain. For each $m \in \{0, \dots, N\}$ we associate a Fourier variable k_m to

domain number m . We also let τ_m be the total length of all line segments (parts of the initial interval $[0, T]$) which form the boundary of m -th domain (see Figure 3a).

For each $j \in \{1, \dots, N\}$ we define $w_1(j)$ and $w_2(j)$ to be the Fourier variables corresponding to the domains that contain the half-circle connecting s_j with t_j in their boundaries (see Figure 3b). All formulae will be symmetric in w_1 and w_2 , so the exact order does not matter.



(a) An example of a diagram. Here,
 $\tau_0 = s_1 + (T - t_1)$,
 $\tau_1 = (s_2 - s_1) + (s_3 - t_2) + (t_1 - t_3)$,
 $\tau_2 = t_2 - t_4$,
 $\tau_3 = t_4 - s_4$,
 $\tau_4 = t_3 - s_3$.

(b) We use $w_1(j)$ and $w_2(j)$ to denote the Fourier variables k_m corresponding to the domains that lie on both sides of the half-circle that connects s_j with t_j . In Figure 3a, for example, $w_1(2) = k_1$ and $w_2(2) = k_2$ (or vice versa).

Figure 3: A diagrammatic representation of Fourier variables.

Proposition 29. Fix $T > 0$, $a \in \mathbb{R}$, and $N \geq 0$. Let $\{\mathcal{O}_0(\phi; s_j, t_j)\}_{j=1}^N$ be a set of non-interlaced observables on $[0, T]$, and $\{l_j\}_{j=1}^N$ be positive integers. Then,

$$\int \prod_{j=1}^N \mathcal{O}_0^{l_j}(\mathbf{Q}_\xi; s_j, t_j) d\mathcal{B}_{\sigma^2}^{a, T}(\xi) = \int_{\mathbb{R}_+^{N+1}} \prod_{j=1}^N \frac{\Gamma\left(\frac{l_j}{2} \pm iw_1(j) \pm iw_2(j)\right)}{2\pi^2 \Gamma(l_j)} \cdot \left(\frac{\sigma^2}{2}\right)^{l_j} \\ \times \exp\left(-\frac{\tau_0 \sigma^2}{2} \cdot k_0^2\right) \frac{\cos(ak_0)}{\pi} dk_0 \cdot \prod_{m=1}^N \exp\left(-\frac{\tau_m \sigma^2}{2} \cdot k_m^2\right) \sinh(2\pi k_m) 2k_m dk_m, \quad (79)$$

where the right-hand side converges absolutely. Moreover, if $\forall j$: either $s_j \neq 0$ or $t_j \neq T$, then the integral in the right-hand side converges absolutely for any $a \in \mathbb{D}_{2\pi}$.

Proof. First, we show that the absolute convergence.

From Remark 4 for every $\varepsilon > 0$ and any positive integer l there exists $C(\varepsilon, l)$ such that

$$\Gamma\left(\frac{l}{2} \pm ik \pm i\omega\right) \leq C(\varepsilon, l) \exp\left(-(\pi - \varepsilon)(|k| + |\omega|)\right), \quad \forall k, \omega \in \mathbb{R}. \quad (80)$$

Furthermore, the boundary of the face number 0 (the unbounded face, corresponding to k_0) contains either at least 1 half-circle or a line segment, and boundary of any other face contains either at

least 3 half-circles or a line segment. Therefore, the integral in the right-hand side of (79) converges absolutely.

Moreover, if $\forall j$: either $s_j \neq 0$ or $t_j \neq T$, then boundary of the face number 0 (the unbounded face) contains either at least 2 half-circles or a line segment. Thus, the integral in the right-hand side of (79) converges absolutely for all $a \in \mathbb{D}_{2\pi}$.

Secondly, we prove (79) by induction on N .

Base case: $N = 0$.

In this case $\tau_0 = T$. Thus, the left-hand side of (79) equals to

$$\int d\mathcal{B}_{\sigma^2}^{a,T}(\xi) = \frac{1}{\sqrt{2\pi T\sigma}} \exp\left(-\frac{a^2}{2T\sigma^2}\right) \quad (81)$$

and the right-hand side is equal to

$$\int_{\mathbb{R}_+^1} \exp\left(-\frac{T\sigma^2}{2} \cdot k_0^2\right) \frac{\cos(ak_0)}{\pi} dk_0 = \frac{1}{\sqrt{2\pi T\sigma}} \exp\left(-\frac{a^2}{2T\sigma^2}\right), \quad (82)$$

which proves the base case.

Induction step: $N - 1 \mapsto N$. By renumbering observables we can assume that $\forall j < N$ we have $s_N \leq s_j$.

There are two possible cases.

Case 1: $s_N = 0$ and $t_N = T$. Renumber Fourier variables so that $w_1(N) = k_0$ and $w_2(N) = k_1$. Using Lemma 26, and induction hypothesis we obtain for all $z < 0$,

$$\begin{aligned} & \int \exp\left\{\frac{z}{\sigma^2} \mathcal{O}_0(\mathbf{Q}_\xi; s_N, t_N)\right\} \prod_{j=1}^{N-1} \mathcal{O}_0^{l_j}(\mathbf{Q}_\xi; s_j, t_j) d\mathcal{B}_{\sigma^2}^{a,T}(\xi) \\ &= \int_{\mathbb{R}_+^{N+1}} \left[\prod_{j=1}^{N-1} \frac{\Gamma\left(\frac{l_j}{2} \pm iw_1(j) \pm iw_2(j)\right)}{2\pi^2 \Gamma(l_j)} \cdot \left(\frac{\sigma^2}{2}\right)^{l_j} \right] \exp\left(-\frac{\tau_1 \sigma^2}{2} \cdot k_1^2\right) \frac{\cos(bk_1)}{\pi} dk_1 \\ & \quad \times \prod_{m=2}^N \exp\left(-\frac{\tau_m \sigma^2}{2} \cdot k_m^2\right) \sinh(2\pi k_m) 2k_m dk_m. \quad (83) \end{aligned}$$

where $b = 2 \operatorname{arccosh} \left[\cosh\left(\frac{a}{2}\right) - \frac{z}{4} \right]$. Notice that it follows from Proposition 27 that the left-hand side is analytic for $z \in \mathbb{D}_8$. Moreover, as we have already proved, the right-hand side is also absolutely convergent whenever $b \in \mathbb{D}_{2\pi}$.

Applying Proposition 32 we obtain

$$\cos(bk_1) = \cos(ak_1) + 2k_1 \sinh(2\pi k_1) \int_0^\infty \sum_{l=1}^\infty \frac{\Gamma\left(\frac{l}{2} \pm ik_1 \pm ik_0\right)}{2\pi^2 \Gamma(l)} \cdot \frac{z^l}{2^l l!} \cos(ak_0) dk_0. \quad (84)$$

Now we substitute this expression into (83). We obtain an absolutely convergent expression, since it is dominated by the same expression with $a = 0$ which, as we have already proved, equals to the left-hand side of (83) with $a = 0$ which, in turn, is finite because of Proposition 27.

Substituting (84) into (83) and taking the coefficient in front of z^{l_0} we get the desired.

Case 2: $t_N < T$. Now we divide the interval $[0, T]$ into three smaller intervals $[0, s_N]$, $[s_N, t_N]$ and $[t_N, T]$. Let $\xi_1 \in C_{0,\text{free}}[0, s_N]$, $\xi_2 \in C_{0,\text{free}}[0, t_N - s_N]$, and $\xi_3 \in C_{0,\text{free}}[0, T - t_N]$. Notice that

$$\mathcal{O}_0(\mathbf{Q}(\xi_1 \sqcup \xi_2 \sqcup \xi_3); s, t) = \mathcal{O}_0(\mathbf{Q}(\xi_2); s - s_N, t - s_N) \quad \text{if } s, t \in [s_N, t_N]; \quad (85)$$

$$\mathcal{O}_0(\mathbf{Q}(\xi_1 \sqcup \xi_2 \sqcup \xi_3); s, t) = \mathcal{O}_0(\mathbf{Q}(\xi_3); s - t_N, t - t_N) \quad \text{if } s, t \in [t_N, T]. \quad (86)$$

Let $X = \{j : s_N \leq t_j \leq t_N\}$ and $Y = \{j : t_N \leq s_j \leq T\}$. Observe that $X \cup Y = \{1, \dots, N\}$, since our observables are non-interlaced. Therefore, we get that

$$\begin{aligned} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\mathbf{Q}(\xi_1 \sqcup \xi_2 \sqcup \xi_3); s_j, t_j) \\ = \prod_{j \in X} \mathcal{O}_0^{l_j}(\mathbf{Q}(\xi_2); s_j - s_N, t_j - s_N) \cdot \prod_{j \in Y} \mathcal{O}_0^{l_j}(\mathbf{Q}(\xi_3); s_j - t_N, t_j - t_N). \end{aligned} \quad (87)$$

Thus, using (19) we get

$$\begin{aligned} \int \prod_{j=1}^N \mathcal{O}_0^{l_j}(\mathbf{Q}_\xi; s_j, t_j) d\mathcal{B}_{\sigma^2}^{a,T}(\xi) &= \int_{\mathbb{R}^2} \left[\int d\mathcal{B}_{\sigma^2}^{b,s_N}(\xi_1) \right] \\ &\times \left[\int \prod_{j \in X} \mathcal{O}_0^{l_j}(\mathbf{Q}(\xi_2); s_j - s_N, t_j - s_N) d\mathcal{B}_{\sigma^2}^{c,t_N-s_N}(\xi_2) \right] \\ &\times \left[\int \prod_{j \in Y} \mathcal{O}_0^{l_j}(\mathbf{Q}(\xi_3); s_j - t_N, t_j - t_N) d\mathcal{B}_{\sigma^2}^{a-b-c,T-t_N}(\xi_3) \right] db dc \end{aligned} \quad (88)$$

Applying induction hypothesis to all three factors in the right-hand side (if $Y = \emptyset$, we apply Case 1 to the second factor) and (114) finishes the proof of Case 2. \square

A similar Proposition holds if we add an exponential of $\mathcal{O}_0(\mathbf{Q}_\xi; 0, T)$ in the expectation.

Proposition 30. *Fix $T > 0$, $a \in \mathbb{R}$, and $N \geq 0$. Let $\{\mathcal{O}_0(\phi; s_j, t_j)\}_{j=1}^N$ be a set of non-interlaced observables on $(0, T)$, and $\{l_j\}_{j=1}^N$ be positive integers. Then,*

$$\begin{aligned} \int \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \mathcal{O}_0(\mathbf{Q}_\xi; 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\mathbf{Q}_\xi; s_j, t_j) d\mathcal{B}_{\sigma^2}^{0,T}(\xi) \\ = \int_{\mathbb{R}_+^{N+1}} \prod_{j=1}^N \frac{\Gamma\left(\frac{l_j}{2} \pm iw_1(j) \pm iw_2(j)\right)}{2\pi^2 \Gamma(l_j)} \cdot \left(\frac{\sigma^2}{2}\right)^{l_j} \\ \times \exp\left(-\frac{\tau_0 \sigma^2}{2} \cdot k_0^2\right) \frac{\cosh(2\alpha k_0)}{\pi} dk_0 \cdot \prod_{m=1}^N \exp\left(-\frac{\tau_m \sigma^2}{2} \cdot k_m^2\right) \sinh(2\pi k_m) 2k_m dk_m. \end{aligned} \quad (89)$$

Proof. Applying Lemma 26 and Proposition 29 we obtain

$$\begin{aligned} \int \exp \left\{ \frac{z}{\sigma^2} \mathcal{O}_0(\mathbf{Q}_\xi; 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\mathbf{Q}_\xi; s_j, t_j) d\mathcal{B}_{\sigma^2}^{0,T}(\xi) \\ = \int_{\mathbb{R}_+^{N+1}} \prod_{j=1}^N \frac{\Gamma\left(\frac{l_j}{2} \pm iw_1(j) \pm iw_2(j)\right)}{2\pi^2 \Gamma(l_j)} \cdot \left(\frac{\sigma^2}{2}\right)^{l_j} \\ \times \exp\left(-\frac{\tau_0 \sigma^2}{2} \cdot k_0^2\right) \frac{\cos(b k_0)}{\pi} dk_0 \cdot \prod_{m=1}^N \exp\left(-\frac{\tau_m \sigma^2}{2} \cdot k_m^2\right) \sinh(2\pi k_m) 2k_m dk_m. \end{aligned} \quad (90)$$

with $b = 2 \operatorname{arccosh}[1 - \frac{z}{4}]$. For $z \in \mathbb{D}_8$ the left-hand side is analytic, and the right-hand side is analytic for $b \in \mathbb{D}_{2\pi}$. Thus, using Statement 40, we can analytically continue the equality above for all $z \in [0, 8)$. Taking $z = 8 \sin^2 \frac{\alpha}{2}$ we get $b = \pm 2i\alpha$, which gives the desired. \square

5 Proofs of the main results

5.1 Proof of Theorem 1

Proof of Theorem 1. Using Proposition 24 and Proposition 25 we get,

$$\int_{\operatorname{Diff}^1(\mathbb{T})/\operatorname{PSL}(2, \mathbb{R})} \left[\prod_{j=1}^N \mathcal{O}^{l_j}(\phi; s_j, t_j) \right] d\mathcal{M}_{\sigma^2}(\phi) = \lim_{\alpha \rightarrow \pi^-} \frac{4\pi(\pi - \alpha)}{\sigma^2} \\ \times \int_{\mathbb{T}} \left[\int_{\operatorname{Diff}^1(\mathbb{T})} \mathcal{O}_0(\phi; 0, T) \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \mathcal{O}_0(\phi; 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\phi; \mathbb{R}^x s_j, \mathbb{R}^x t_j) d\mathcal{N}_{\sigma^2}^0(\phi) \right] dx. \quad (91)$$

Using the fact that

$$d\mathcal{N}_{\sigma^2}^0(\phi) = d\mathcal{B}_{\sigma^2}^{0,T}(\xi), \quad \text{where } \phi = P(\xi), \quad (92)$$

and

$$\mathcal{O}_0(P(\xi); s, t) = \mathcal{O}_0(Q(\xi); s, t) \quad (93)$$

we obtain

$$\int_{\operatorname{Diff}^1(\mathbb{T})} \mathcal{O}_0(\phi; 0, T) \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \mathcal{O}_0(\phi; 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(\phi; \mathbb{R}^x s_j, \mathbb{R}^x t_j) d\mathcal{N}_{\sigma^2}^0(\phi) \\ = \int \mathcal{O}_0(Q(\xi); 0, T) \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \mathcal{O}_0(Q(\xi); 0, T) \right\} \prod_{j=1}^N \mathcal{O}_0^{l_j}(Q(\xi); \mathbb{R}^x s_j, \mathbb{R}^x t_j) d\mathcal{B}_{\sigma^2}^{0,T}(\xi). \quad (94)$$

Differentiating (89) in α gives the desired. \square

5.2 Proof of Proposition 8

Proof of Proposition 8. Using Corollary 7, for

$$C = C(\sigma, s, t) = \sup_{k > 0} \left\{ \exp \left(-\frac{(1 - (t - s))\sigma^2}{2} \cdot k^2 \right) \sinh(2\pi k) 2k \right\} \quad (95)$$

and any positive integer l we have

$$\int \mathcal{O}^l(\phi; s, t) d\mathcal{M}_{\sigma^2}(\phi) \leq C \int_{\mathbb{R}_+^2} \frac{\Gamma(\frac{l}{2} \pm ik_1 \pm ik_2)}{2\pi^2 \Gamma(l)} \cdot \left(\frac{\sigma^2}{2}\right)^l \exp \left(-\frac{(t - s)\sigma^2}{2} \cdot k_1^2 \right) \sinh(2\pi k_1) 2k_1 dk_1 dk_2. \quad (96)$$

Using Tonelli's Theorem and Proposition 32 for $\beta = 0$ we get that for any $z \in [0, 8)$,

$$\int \exp \left\{ \frac{z}{\sigma^2} \mathcal{O}(\phi; s, t) \right\} d\mathcal{M}_{\sigma^2}(\phi) \leq C \int_{\mathbb{R}_+} \exp \left(-\frac{(t - s)\sigma^2}{2} \cdot k^2 \right) \cos \left(2k \operatorname{arccosh} \left[1 - \frac{z}{4} \right] \right) dk. \quad (97)$$

Take $z = 8 \sin^2 \frac{\alpha}{2}$, so that $\cos(2k \operatorname{arccosh}[1 - \frac{z}{4}]) = \cosh(2k\alpha)$. Therefore,

$$\begin{aligned} \int \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \mathcal{O}(\phi; s, t) \right\} d\mathcal{M}_{\sigma^2}(\phi) &\leq C \int_{\mathbb{R}_+} \exp \left(-\frac{(t-s)\sigma^2}{2} \cdot k^2 \right) \cosh(2k\alpha) dk \\ &= C \frac{\sqrt{\pi}}{\sqrt{2(t-s)}\sigma} \exp \left(\frac{2\alpha^2}{(t-s)\sigma^2} \right). \end{aligned} \quad (98)$$

Taking $\alpha \rightarrow \pi$ finishes the proof. \square

5.3 Proof of Theorem 2

Proof of Theorem 2. Denote $\widetilde{\operatorname{Diff}}^1(\mathbb{T}) = \{\phi \in \operatorname{Diff}^1(\mathbb{T}) \mid \phi(0) = 0, \phi'(0) = 1, \phi(1/2) = 1/2\}$. It is easy to see that as a topological space $\widetilde{\operatorname{Diff}}^1(\mathbb{T})$ is isomorphic to $\operatorname{Diff}^1(\mathbb{T})/\operatorname{PSL}(2, \mathbb{R})$. Let \mathcal{P}' and \mathcal{M}'_{σ^2} be the push-forward measures of \mathcal{P} and \mathcal{M}_{σ^2} to $\widetilde{\operatorname{Diff}}^1(\mathbb{T})$ respectively. Then,

$$\int \prod_{j=1}^N \mathcal{O}^{t_j}(\phi; s_j, t_j) d\mathcal{M}'_{\sigma^2}(\phi) = \int \prod_{j=1}^N \mathcal{O}^{t_j}(\phi; s_j, t_j) d\mathcal{P}'(\phi). \quad (99)$$

Now we show that for all Borel $A \subset \widetilde{\operatorname{Diff}}^1(\mathbb{T})$ we have $\mathcal{M}'_{\sigma^2}(A) = \mathcal{P}'(A)$. It is sufficient to prove this for all cylinder sets.

Take any set of points $0 = t_0 < t_1 < \dots < t_{M-1} < t_M = 1/2$ and $1/2 < t_{M+1} < \dots < t_{2M-1} < t_{2M} = 1$. Consider observables $\left\{ \mathcal{O}(\phi; 0, t_j) \right\}_{j=1}^{2M-1}$ and $\left\{ \mathcal{O}(\phi; t_j, t_{j+1}) \right\}_{j=0}^{2M-1}$. This set of observables is non-interlaced.

Observe that for all $\phi \in \widetilde{\operatorname{Diff}}^1(\mathbb{T})$ and $1 < j < 2M-1$, using $\sin(a-b) = \sin(a)\cos(b) - \sin(b)\cos(a)$, we get

$$\frac{\mathcal{O}(\phi; 0, t_j) \mathcal{O}(\phi; 0, t_{j+1})}{\pi \mathcal{O}(\phi; t_j, t_{j+1})} = \frac{\phi'(0) \sin(\pi[\phi(t_{j+1}) - \phi(t_j)])}{\sin(\pi\phi(t_{j+1})) \sin(\pi\phi(t_j))} = \cot(\pi\phi(t_j)) - \cot(\pi\phi(t_{j+1})). \quad (100)$$

Moreover, for any $\phi \in \widetilde{\operatorname{Diff}}^1(\mathbb{T})$ we have $\cot(\pi\phi(t_M)) = 0$. Thus, for all $0 \leq j \leq 2M-1$ it is possible to express $\phi(t_j)$ in terms of observables in question using algebraic operations and trigonometric functions. Since for all $\phi \in \widetilde{\operatorname{Diff}}^1(\mathbb{T})$ and $0 < j \leq 2M-1$,

$$\mathcal{O}(\phi; 0, t_j) = \frac{\pi \sqrt{\phi'(t_j)}}{\sin(\pi\phi(t_j))}, \quad (101)$$

we can express $\phi'(t_j)$ in terms of our observables too.

Moreover, notice that taking $N = 0$ implies that the total mass of both measures is equal to $\mathcal{Z}(\sigma)$.

Now take two random variables $\phi_1 \sim \frac{1}{\mathcal{Z}(\sigma)} \mathcal{P}'$ and $\phi_2 \sim \frac{1}{\mathcal{Z}(\sigma)} \mathcal{M}'_{\sigma^2}$. Notice that joint distribution of observables $\left\{ \mathcal{O}(\phi; 0, t_j) \right\}_{j=1}^{2M-1} \cup \left\{ \mathcal{O}(\phi; t_j, t_{j+1}) \right\}_{j=0}^{2M-1}$ is the same for both ϕ_1 and ϕ_2 , since their correlation functions are equal, and all variables have exponential moments (hence, their characteristic functions are equal). Moreover, we have already shown that we can express both $\{\phi(t_j)\}_{j=0}^{2M-1}$ and $\{\phi'(t_j)\}_{j=0}^{2M-1}$ as continuous functions of observables in question. Therefore, the induced distributions on $\{\phi(t_j), \phi'(t_j)\}_{j=0}^{2M-1}$ are equal. This finishes the proof. \square

5.4 Proof of Theorem 3

Theorem 3 follows immediately from the following Lemma.

Lemma 31. *For $\varepsilon \rightarrow 0$ we have*

$$\begin{aligned} \exp\left(\frac{\varepsilon \sigma^2}{2} \cdot k_2^2\right) \int_{\mathbb{R}_+} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) \sinh(2\pi k_1) \frac{\Gamma(1 \pm ik_1 \pm ik_2)}{2\pi^2 \Gamma(2)} 2k_1 dk_1 \\ = \varepsilon^{-2} + \frac{\sigma^4}{240} + \frac{\sigma^4 k_2^2}{12} + (1 + k_2^5) O(\varepsilon^{1/2}). \end{aligned} \quad (102)$$

Proof. Using the fact that $|\Gamma(1 + ix)|^2 = \pi x / \sinh(\pi x)$, and

$$\frac{\sinh(2\pi k_1)}{\sinh(\pi(k_1 - k_2)) \sinh(\pi(k_1 + k_2))} = \coth(\pi(k_1 - k_2)) + \coth(\pi(k_1 + k_2)), \quad (103)$$

we get

$$\begin{aligned} \int_{\mathbb{R}_+} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) \sinh(2\pi k_1) \frac{\Gamma(1 \pm ik_1 \pm ik_2)}{2\pi^2 \Gamma(2)} 2k_1 dk_1 \\ = \int_{\mathbb{R}} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) k_1 (k_1^2 - k_2^2) \coth(\pi(k_1 - k_2)) dk_1. \end{aligned} \quad (104)$$

It is well-known (e.g. see [Gradshteyn and Ryzhik, 2007, 17.23.21])

$$\mathcal{F}[\coth(\pi x)](w) = -i \coth\left(\frac{w}{2}\right), \quad (105)$$

where by \mathcal{F} we denote the Fourier Transform (in the sense of tempered distribution, e.g. see [Stein and Weiss, 1971, Chapter I.3]) normalized as in (112). Therefore, by Plancherel identity

$$\begin{aligned} \int_{\mathbb{R}} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) k_1^{2n+1} \coth(\pi(k_1 - k_2)) dk_1 \\ = (-1)^n \frac{1}{\sqrt{2\pi}} (\varepsilon \sigma^2)^{-(n+1)} \int_{\mathbb{R}} \exp\left(-\frac{w^2}{2\varepsilon \sigma^2}\right) H_{2n+1}\left(\frac{w}{\sqrt{\varepsilon \sigma}}\right) \coth\left(\frac{w}{2}\right) e^{ik_2 w} dw, \\ = (-1)^n \frac{1}{\sqrt{2\pi}} (\varepsilon \sigma^2)^{-(n+\frac{1}{2})} \int_{\mathbb{R}} \exp\left(-\frac{w^2}{2}\right) H_{2n+1}(w) \coth\left(\frac{\sqrt{\varepsilon \sigma} w}{2}\right) e^{ik_2 \sqrt{\varepsilon \sigma} w} dw, \end{aligned} \quad (106)$$

where $H_k(x) = (-1)^k e^{x^2/2} \frac{d^k}{dx^k} e^{-x^2/2}$ is the k -th Hermite polynomial, and the integrals are understood as Cauchy principal value integrals. It is easy to see that

$$\coth\left(\frac{x}{2}\right) = \frac{2}{x} + \frac{x}{6} - \frac{x^3}{360} + O(|x|^5), \quad (107)$$

$$e^{ik_2 x} = 1 + ik_2 x - \frac{k_2^2 x^2}{2} - \frac{ik_2^3 x^3}{6} + \frac{k_2^4 x^4}{24} + O(k_2^5 x^5), \quad (108)$$

where the remainder terms are uniform over $k_2 \in \mathbb{R}$.

Therefore,

$$\begin{aligned} \int_{\mathbb{R}} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) k_1 \coth(\pi(k_1 - k_2)) dk_1 \\ = \frac{1}{\sqrt{2\pi}} (\varepsilon \sigma^2)^{-1} \int_{\mathbb{R}} \exp\left(-\frac{w^2}{2}\right) \left(2 + i 2\sqrt{\varepsilon \sigma} k_2 w + \varepsilon \sigma^2 \left(\frac{1}{6} - k_2^2\right) w^2 + (1 + k_2^3) O(\varepsilon^{3/2} w^3)\right) dw \\ = 2 (\varepsilon \sigma^2)^{-1} + \frac{1}{6} - k_2^2 + (1 + k_2^3) O(\sqrt{\varepsilon}), \end{aligned} \quad (109)$$

and

$$\begin{aligned}
& \int_{\mathbb{R}} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) k_1^3 \coth(\pi(k_1 - k_2)) dk_1 \\
&= \frac{1}{\sqrt{2\pi}} (\varepsilon \sigma^2)^{-2} \int_{\mathbb{R}} \exp\left(-\frac{w^2}{2}\right) \left(6 - 2w^2 + i \sqrt{\varepsilon} \sigma (6w - 2w^3) k_2 + \varepsilon \sigma^2 (3w^2 - w^4) \left(\frac{1}{6} - k_2^2\right) \right. \\
&\quad \left. - i(\varepsilon \sigma^2)^{3/2} (3w^3 - w^5) \frac{k_2^3}{6} - (\varepsilon \sigma^2)^2 (3w^4 - w^6) \left(\frac{1}{360} + \frac{k_2^2}{12} - \frac{k_2^4}{12}\right) + (1 + k_2^5) O(\varepsilon^{5/2} w^5)\right) dw \\
&= 4 (\varepsilon \sigma^2)^{-2} + \frac{1}{60} + \frac{k_2^2}{2} - \frac{k_2^4}{2} + (1 + k_2^5) O(\varepsilon^{1/2}). \quad (110)
\end{aligned}$$

Hence,

$$\begin{aligned}
& \exp\left(\frac{\varepsilon \sigma^2}{2} \cdot k_2^2\right) \int_{\mathbb{R}_+} \exp\left(-\frac{\varepsilon \sigma^2}{2} \cdot k_1^2\right) \sinh(2\pi k_1) \frac{\Gamma(1 \pm ik_1 \pm ik_2)}{2\pi^2 \Gamma(2)} \left(\frac{\sigma^2}{2}\right)^2 2k_1 dk_1 \\
&= \varepsilon^{-2} + \frac{\sigma^4}{240} + \frac{\sigma^4 k_2^2}{12} + (1 + k_2^5) O(\varepsilon^{1/2}), \quad (111)
\end{aligned}$$

where the remainder terms are uniform over $k_2 \in \mathbb{R}$. \square

A Fourier calculations

We use the following normalization for Fourier transform

$$\hat{f}(\omega) = \int_{\mathbb{R}} f(x) e^{-i\omega x} dx, \quad (112)$$

$$f(x) = \int_{\mathbb{R}} \hat{f}(\omega) \frac{e^{i\omega x}}{2\pi} d\omega. \quad (113)$$

In our normalization

$$\int f(y)g(x-y) dy = \int \hat{f}(\omega)\hat{g}(\omega) \frac{e^{i\omega x}}{2\pi} d\omega. \quad (114)$$

The main goal of this Section is to prove the following proposition.

Proposition 32. *For any $\beta, k \in \mathbb{R}$ and any $z \in \mathbb{D}_2$ we have*

$$\begin{aligned} & \cos\left(2k \cdot \operatorname{arccosh}[\cosh(\beta/2) - z]\right) = \\ & \cos(k\beta) + 2k \sinh(2\pi k) \int_0^\infty \sum_{l=1}^\infty \frac{\Gamma\left(\frac{l}{2} \pm ik \pm iw\right)}{2\pi^2 \Gamma(l)} \cdot \frac{(2z)^l}{l!} \cos(w\beta) dw, \end{aligned} \quad (115)$$

where the right-hand side converges absolutely.

Remark 33. *The right-hand side above can also be written as*

$$2k \sinh(2\pi k) \int_0^\infty \sum_{l=0}^\infty \frac{\Gamma\left(\frac{l}{2} \pm ik \pm iw\right)}{2\pi^2 \Gamma(l)} \cdot \frac{(2z)^l}{l!} \cos(w\beta) dw, \quad (116)$$

if we interpret $l = 0$ term as delta function $\delta(\omega - k)$.

We will deduce it from the following two lemmas.

Lemma 34. *For all $k, w \in \mathbb{R}$ and $l > 0$ we have*

$$\frac{\Gamma(l \pm ik \pm iw)}{\Gamma^2(2l)} = \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} (e^{a/2} + e^{-a/2} + e^{b/2} + e^{-b/2})^{-2l} e^{ika+iwb} da db. \quad (117)$$

Proof. We start by writing

$$\Gamma(l + ix) = \int_0^\infty t^{ix+l-1} e^{-t} dt = \int_{-\infty}^\infty e^{u(l+ix)} e^{-e^u} du. \quad (118)$$

Note that the integrand in the right-hand side decays exponentially as $u \rightarrow \pm\infty$. Applying the equation above for $x \in \{k+w, k-w, -k+w, -k-w\}$ and multiplying them we obtain

$$\begin{aligned} \Gamma(l \pm ik \pm iw) &= \int_{\mathbb{R}^4} \exp\left\{-e^{u_1} - e^{u_2} - e^{u_3} - e^{u_4}\right\} \\ &\times \exp\left\{l(u_1 + u_2 + u_3 + u_4) + ik(u_1 + u_2 - u_3 - u_4) + iw(u_1 - u_2 + u_3 - u_4)\right\} du_1 du_2 du_3 du_4. \end{aligned} \quad (119)$$

Changing the variables

$$v_1 = \frac{1}{2}(u_1 + u_4), \quad v_2 = \frac{1}{2}(u_1 - u_4), \quad (120)$$

$$v_3 = \frac{1}{2}(u_2 + u_3), \quad v_4 = \frac{1}{2}(u_2 - u_3), \quad (121)$$

$$(122)$$

we obtain

$$\begin{aligned} \Gamma(l \pm ik \pm iw) &= 4 \int_{\mathbb{R}^4} \exp \left\{ -e^{v_1+v_2} - e^{v_1-v_2} - e^{v_3+v_4} - e^{v_3-v_4} \right\} \\ &\quad \times \exp \left\{ 2l(v_1 + v_3) + 2ik(v_2 + v_4) + 2iw(v_2 - v_4) \right\} dv_1 dv_2 dv_3 dv_4. \end{aligned} \quad (123)$$

Next, we integrate out v_1 and v_3 using an equality

$$\int_{\mathbb{R}} \exp \{ -y e^x \} e^{\lambda x} dx = \Gamma(\lambda) y^{-\lambda}, \quad (124)$$

and get

$$\begin{aligned} \frac{\Gamma(l \pm ik \pm iw)}{\Gamma^2(2l)} &= 4 \int_{\mathbb{R}^2} (e^{v_2} + e^{-v_2})^{-2l} (e^{v_4} + e^{-v_4})^{-2l} \exp \left\{ 2ik(v_2 + v_4) + 2iw(v_2 - v_4) \right\} dv_2 dv_4 \\ &= 4 \int_{\mathbb{R}^2} (e^{v_2+v_4} + e^{-v_2-v_4} + e^{v_2-v_4} + e^{v_4-v_2})^{-2l} \exp \left\{ 2ik(v_2 + v_4) + 2iw(v_2 - v_4) \right\} dv_2 dv_4. \end{aligned} \quad (125)$$

Changing the variables again

$$a = 2(v_2 + v_4), \quad b = 2(v_2 - v_4), \quad (126)$$

we get the desired. \square

Lemma 35. For all $p, q \in \mathbb{C}$ such that $|p| < \pi, |q| < \pi$ we have

$$\frac{\cos(p\omega) - \cos(q\omega)}{\sinh(\pi\omega)} = \frac{i}{2\pi} \int_{\mathbb{R}} \frac{\sinh(x) (\cosh(p) - \cosh(q))}{(\cosh(q) + \cosh(x)) (\cosh(p) + \cosh(x))} e^{ix\omega} dx. \quad (127)$$

Notice that the integrand in the right-hand side is exponentially decaying as $x \rightarrow \pm\infty$.

Proof. It is well-known (see, e.g. [Gradshteyn and Ryzhik, 2007, 17.23.20]) that for $a, b \in \mathbb{R}$ with $|a| < |b|$ we have

$$\frac{\sinh(a\omega)}{\sinh(b\omega)} = \frac{1}{2} \int_{\mathbb{R}} \frac{\sin(\pi a/b)}{b[\cos(\pi a/b) + \cosh(\pi x/b)]} e^{ix\omega} dx. \quad (128)$$

Notice that the formula above also holds true for $a \in \mathbb{C}$ with $|a| < |b|$, as both sides are analytic in a .

Now we write

$$\frac{\cos(p\omega) - \cos(q\omega)}{\sinh(\pi\omega)} = -\frac{2 \sin\left(\frac{p+q}{2}\omega\right) \sin\left(\frac{p-q}{2}\omega\right)}{\sinh(\pi\omega)}. \quad (129)$$

Thus, combining this with (128) for $a = i(p+q)/2$ and $b = \pi$ we obtain

$$\frac{\cos(p\omega) - \cos(q\omega)}{\sinh(\pi\omega)} = \frac{i}{2\pi} \int_{\mathbb{R}} \left(\frac{\sinh\left(\frac{p+q}{2}\right)}{\cosh\left(\frac{p+q}{2}\right) + \cosh\left(x - \frac{p-q}{2}\right)} - \frac{\sinh\left(\frac{p+q}{2}\right)}{\cosh\left(\frac{p+q}{2}\right) + \cosh\left(x + \frac{p-q}{2}\right)} \right) e^{ix\omega} dx. \quad (130)$$

Moreover,

$$\begin{aligned} &\frac{\sinh\left(\frac{p+q}{2}\right)}{\cosh\left(\frac{p+q}{2}\right) + \cosh\left(x - \frac{p-q}{2}\right)} - \frac{\sinh\left(\frac{p+q}{2}\right)}{\cosh\left(\frac{p+q}{2}\right) + \cosh\left(x + \frac{p-q}{2}\right)} \\ &= \frac{\sinh\left(\frac{p+q}{2}\right) \left(\cosh\left(x + \frac{p-q}{2}\right) - \cosh\left(x - \frac{p-q}{2}\right) \right)}{\left(\cosh\left(\frac{p+q}{2}\right) + \cosh\left(x - \frac{p-q}{2}\right) \right) \left(\cosh\left(\frac{p+q}{2}\right) + \cosh\left(x + \frac{p-q}{2}\right) \right)}. \end{aligned} \quad (131)$$

It is easy to show that the numerator in the right-hand side is equal to $\sinh(x)(\cosh(p) - \cosh(q))$, and the denominator equals to $(\cosh(q) + \cosh(x))(\cosh(p) + \cosh(x))$, which finishes the proof. \square

Proof of Proposition 32. Using Lemma 35 we get

$$\begin{aligned} & \frac{\cos\left(2k \cdot \operatorname{arccosh}[\cosh(\beta/2) - z]\right) - \cos(k\beta)}{\sinh(2\pi k)} \\ &= \frac{-i}{2\pi} \int_{\mathbb{R}} \frac{z \sinh(x)}{(\cosh(x) + \cosh(\beta/2))(\cosh(x) + \cosh(\beta/2) - z)} e^{2ixk} dx. \end{aligned} \quad (132)$$

Now we Taylor expand in z

$$\frac{1}{\cosh(x) + \cosh(\beta/2) - z} = \sum_{l=0}^{\infty} (\cosh(x) + \cosh(\beta/2))^{-l-1} z^l, \quad (133)$$

thus

$$\frac{z \sinh(x)}{(\cosh(x) + \cosh(\beta/2))(\cosh(x) + \cosh(\beta/2) - z)} = \sinh(x) \sum_{l=1}^{\infty} (\cosh(x) + \cosh(\beta/2))^{-l-1} z^l. \quad (134)$$

Differentiating by parts and using Lemma 34 we obtain that for any $l \geq 1$,

$$\begin{aligned} \int_{\mathbb{R}} \sinh(x) (\cosh(x) + \cosh(\beta/2))^{-l-1} e^{2ixk} dx &= \frac{ik2^{l+1}}{l} \int_{\mathbb{R}} (e^x + e^{-x} + e^{\beta/2} + e^{-\beta/2})^{-l} e^{2ixk} dx \\ &= \frac{ik2^l}{\pi l} \int_{\mathbb{R}} \frac{\Gamma(\frac{l}{2} \pm ik \pm iw)}{\Gamma^2(l)} e^{iw\beta} dw. \end{aligned} \quad (135)$$

In particular, taking $\beta = 0$,

$$\frac{k2^l}{\pi l} \int_{\mathbb{R}} \frac{\Gamma(\frac{l}{2} \pm ik \pm iw)}{\Gamma^2(l)} dw \leq \int_{\mathbb{R}} |\sinh(x)| (\cosh(x) + \cosh(\beta/2))^{-l-1} dx, \quad (136)$$

and thus, summing over l we obtain for any $|z| < 2$

$$\frac{k}{\pi} \sum_{l=1}^{\infty} \int_{\mathbb{R}} \frac{\Gamma(\frac{l}{2} \pm ik \pm iw)}{\Gamma(l)} \cdot \frac{(2|z|)^l}{l!} dw \leq \int_{\mathbb{R}} \frac{|z \sinh(x)|}{(\cosh(x) + \cosh(\beta/2))(\cosh(x) + \cosh(\beta/2) - |z|)} dx < \infty. \quad (137)$$

Now using (135) again and summing over l we obtain

$$\begin{aligned} & \frac{-i}{2\pi} \int_{\mathbb{R}} \frac{z \sinh(x)}{(\cosh(x) + \cosh(\beta/2))(\cosh(x) + \cosh(\beta/2) - z)} e^{2ixk} dx \\ &= \frac{k}{2\pi^2} \sum_{l=1}^{\infty} \int_{\mathbb{R}} \frac{\Gamma(\frac{l}{2} \pm ik \pm iw)}{\Gamma(l)} \cdot \frac{(2z)^l}{l!} e^{iw\beta} dw, \end{aligned} \quad (138)$$

where the expression in the right-hand side converges absolutely because of (137). This, together with (132) finishes proof. \square

B Appendix

Lemmas 36 and 37 have already appeared in [Bauerschmidt, Losev, and Wildemann, 2024]. For the reader's convenience we recall them together with their proofs.

Lemma 36 ([Bauerschmidt, Losev, and Wildemann, 2024]). *Fix $\sigma, a, T > 0$. Then there exists $\varepsilon > 0$ such that*

$$\int \exp\left(\frac{\varepsilon}{\int_0^T e^{\tilde{\xi}(\tau)} d\tau}\right) d\mathcal{B}_{\sigma^2}^{a,T}(\xi) < \infty. \quad (139)$$

Proof. Let $\tilde{\xi}$ be a Brownian bridge distributed according to the probability measure $\sqrt{2\pi T}\sigma \exp\left(\frac{a^2}{2T\sigma^2}\right) d\mathcal{B}_{\sigma^2}^{a,T}(\tilde{\xi})$. Then

$$\mathbb{P}\left[\frac{1}{\int_0^T e^{\tilde{\xi}(\tau)} d\tau} > \lambda\right] \leq \mathbb{P}\left[\min_{t \in [0, \varepsilon\lambda^{-1}]} \tilde{\xi}(t) < -1\right] \leq C^{-1}e^{-C\lambda}, \quad (140)$$

for some $C > 0$, independent of $\lambda > 10/T$. \square

Lemma 37 ([Bauerschmidt, Losev, and Wildemann, 2024]). *Let \mathcal{P} be a non-negative measure on \mathbb{R}_+ . Assume that there exists $\varepsilon > 0$ such that the exponential moment generating function $F(z) = \int \exp(zX) d\mathcal{P}(X)$ exists for all $z \in [0, \varepsilon]$. Assume further that for some $R > 0$, $F(z)$ can be analytically continued for all $z \in \mathbb{D}_R$. Then $\int \exp(zX) d\mathcal{P}(X)$ converges absolutely for all $z \in \mathbb{D}_R$, and is equal to the analytic continuation of $F(z)$ to \mathbb{D}_R .*

Proof. Since \mathcal{P} supported on \mathbb{R}_+ , by Fubini's Theorem,

$$F(z) = \sum_{n \geq 0} z^n \frac{\int X^n d\mathcal{P}(X)}{n!}, \quad (141)$$

for $z \in [0, \varepsilon]$. Given that $F(z)$ is analytic in \mathbb{D}_R , we conclude that the right-hand side converges absolutely for $z \in \mathbb{D}_R$. Thus, by Fubini's Theorem again, $\mathbb{E} \exp(zX)$ converges for $z \in [0, R]$ and is equal to $F(z)$. We can continue the equality for the whole disk \mathbb{D}_R , since $|z^n X^n| \leq |z|^n X^n$, and all expressions are absolutely convergent in \mathbb{D}_R . \square

Statement 38. *For any $\delta \in (0, 1/10)$, if $x \in (0, 1 - \delta)$, then for any $\alpha \in (\frac{9\pi}{10}, \pi)$ we have*

$$\left| \log\left(\frac{\sin(\pi x)}{\sin(\alpha x)}\right) \right| \leq \frac{\pi - \alpha}{\sin(\pi\delta/2)}. \quad (142)$$

Proof. Since $\sin(y)/y$ is decreasing for $y \in [0, \pi]$, we get that

$$\left| \frac{d}{d\alpha} \log(\sin(\alpha x)) \right| = \frac{x |\cos(\alpha x)|}{\sin(\alpha x)} \leq \frac{\alpha(1 - \delta)}{\alpha \sin(\alpha(1 - \delta))} \leq \frac{1}{\sin(\pi\delta/2)}. \quad (143)$$

\square

Statement 39. *For any $x \in \mathbb{R}$ we have*

$$\operatorname{arccosh}^2[\cosh(x) - 2] > x^2 - 1000. \quad (144)$$

Proof. Since $\operatorname{arccosh}^2[\cosh(x) - 2]$ is even, it is sufficient to prove the inequality only for $x \geq 0$. For $x \in [0, 10]$ we see that

$$\operatorname{arccosh}^2[\cosh(x) - 2] \geq -\pi^2 > x^2 - 1000. \quad (145)$$

For $x \in (10, \infty)$ we have

$$\cosh(x) - 2 > \cosh\left(x - \frac{3}{x}\right), \quad (146)$$

since

$$\frac{d}{dy} \cosh(y) = \sinh(y) > y \quad (147)$$

for $y > 5$. Therefore,

$$\operatorname{arccosh}^2[\cosh(x) - 2] > \left(x - \frac{3}{x}\right)^2 > x^2 - 10. \quad (148)$$

□

Statement 40. *Function $\operatorname{arccosh}^2(z)$ can be analytically continued from $z \in [1, \infty)$ to $z \in \Omega := \{z \in \mathbb{C} \mid \Re z \geq -1\}$. Moreover, in this continuation $\operatorname{arccosh}^2(z) = -\operatorname{arccos}^2(z)$ for $z \in (-1, 1)$.*

Proof. Let $f(z) = (z + \frac{1}{z})/2$, and $g(z) = e^z$. Notice that $\operatorname{arccosh}^2(z) = [(g^{-1} \circ f^{-1})(z)]^2$.

It is easy to see that f is a double cover from $\mathbb{C} \setminus (-\infty, 0]$ to $\mathbb{C} \setminus (-\infty, -1]$ with only critical point $z = 1$, and g is a bijection between $\{z \in \mathbb{C} \mid -\pi < \Im z < \pi\}$ and $\mathbb{C} \setminus (-\infty, 0]$ with 0 being mapped to 1.

Observation that $(f \circ g)(-z) = (f \circ g)(z)$ finishes the proof of the first part.

The equality $\operatorname{arccosh}^2(z) = -\operatorname{arccos}^2(z)$ for $z \in (-1, 1)$ follows from the fact that $\omega = i \operatorname{arccos}(z)$ and $\omega = -i \operatorname{arccos}(z)$ are the only solutions of $\cosh(\omega) = z$ in the domain $\{\omega \in \mathbb{C} \mid -\pi < \Im \omega < \pi\}$. □

Acknowledgements

The author thanks Roland Bauerschmidt, James Norris and Peter Wildemann for many helpful discussions and careful proofreading.

References

- A. Alekseev and S. Shatashvili. Path integral quantization of the coadjoint orbits of the Virasoro group and 2-d gravity. *Nuclear Phys. B*, 323(3):719–733, 1989. ISSN 0550-3213,1873-1562.
- A. Alekseev and S. L. Shatashvili. Characters, coadjoint orbits and Duistermaat-Heckman integrals. *J. Geom. Phys.*, 170:Paper No. 104386, 20, 2021. ISSN 0393-0440,1879-1662.
- A. Alekseev, O. Chekeres, and D. R. Youmans. Towards Bosonization of Virasoro Coadjoint Orbits. *Annales Henri Poincare*, 25(1):5–34, 2024.
- D. Bagrets, A. Altland, and A. Kamenev. Sachdev–Ye–Kitaev model as Liouville quantum mechanics. *Nuclear Physics B*, 911:191–205, 2016. ISSN 0550-3213.
- R. Bauerschmidt, I. Losev, and P. Wildemann. Probabilistic Definition of the Schwarzian Field Theory. 2024. Preprint.

- V. V. Belokurov and E. T. Shavgulidze. Exact solution of the Schwarzian theory. *Phys. Rev. D*, 96, Nov 2017.
- A. Blommaert, T. G. Mertens, and H. Verschelde. The Schwarzian Theory - A Wilson Line Perspective. *JHEP*, 12:022, 2018.
- V. I. Bogachev. *Differentiable measures and the Malliavin calculus*, volume 164 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010. ISBN 978-0-8218-4993-4.
- A. R. Chandra, I. Chevyrev, M. Hairer, and H. Shen. Langevin dynamic for the 2D Yang–Mills measure. *Publications mathématiques de l’IHÉS*, 136:1 – 147, 2020.
- I. Chevyrev. Yang–Mills measure on the two-dimensional torus as a random distribution. *Communications in Mathematical Physics*, 372:1027 – 1058, 2018.
- I. Chevyrev and H. Shen. Invariant measure and universality of the 2D Yang–Mills Langevin dynamic. 2023. Preprint, arxiv:2302.12160.
- A. Dahlqvist and J. R. Norris. Yang–Mills measure and the master field on the sphere. *Communications in Mathematical Physics*, 377:1163–1226, 2017.
- F. David, A. Kupiainen, R. Rhodes, and V. Vargas. Liouville quantum gravity on the Riemann sphere. *Communications in Mathematical Physics*, 342, 03 2016.
- B. Driver, F. Gabriel, B. Hall, and T. Kemp. The Makeenko–Migdal equation for Yang–Mills theory on compact surfaces. *Communications in Mathematical Physics*, 352, 06 2017.
- B. K. Driver, B. C. Hall, and T. Kemp. Three proofs of the Makeenko–Migdal equation for Yang–Mills theory on the plane. *Communications in Mathematical Physics*, 351:741 – 774, 2016.
- I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Elsevier/Academic Press, Amsterdam, seventh edition, 2007. ISBN 978-0-12-373637-6; 0-12-373637-4. Translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger.
- C. Guillarmou, R. Rhodes, and V. Vargas. Polyakov’s formulation of $2d$ bosonic string theory. *Publications mathématiques de l’IHÉS*, 130, 06 2019.
- C. Guillarmou, A. Kupiainen, R. Rhodes, and V. Vargas. Conformal bootstrap in Liouville Theory. 5 2020. Preprint, arxiv:2005.11530.
- C. Guillarmou, A. Kupiainen, R. Rhodes, and V. Vargas. Segal’s axioms and bootstrap for Liouville theory. 2021. Preprint, arxiv:2112.14859.
- C. Guillarmou, A. Kupiainen, and R. Rhodes. Review on the probabilistic construction and Conformal bootstrap in Liouville Theory. 2024. Preprint, arXiv:2403.12780.
- L. V. Iliesiu, S. S. Pufu, H. Verlinde, and Y. Wang. An exact quantization of Jackiw-Teitelboim gravity. *J. High Energy Phys.*, (11):091, 61, 2019. ISSN 1126-6708,1029-8479.
- A. Kitaev and S. J. Suh. The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual. *J. High Energy Phys.*, (5):183, front matter+66, 2018. ISSN 1126-6708.
- A. Kupiainen and J. Oikarinen. Stress-Energy in Liouville Conformal Field Theory. 11 2019. Preprint, arxiv:1911.05359.

- A. Kupiainen, R. Rhodes, and V. Vargas. Integrability of Liouville theory: proof of the DOZZ formula. *Annals of Mathematics*, 2017.
- A. Kupiainen, R. Rhodes, and V. Vargas. Local conformal structure of Liouville quantum gravity. *Communications in Mathematical Physics*, 371, 11 2019.
- T. Lévy. Two-dimensional Markovian holonomy fields. *Astérisque*, 2008.
- T. Levy. The master field on the plane. *Astérisque*, 2018.
- J. Maldacena and D. Stanford. Remarks on the Sachdev-Ye-Kitaev model. *Phys. Rev. D*, 94(10): 106002, 43, 2016. ISSN 2470-0010,2470-0029.
- J. Maldacena, D. Stanford, and Z. Yang. Conformal symmetry and its breaking in two-dimensional nearly anti-de Sitter space. *Progress of Theoretical and Experimental Physics*, 2016(12):12C104, 11 2016. ISSN 2050-3911.
- T. Mertens, G. Turiaci, and H. Verlinde. Solving the Schwarzian via the Conformal Bootstrap. *Journal of High Energy Physics*, 2017, 05 2017.
- J. Oikarinen. Stress-energy in Liouville Conformal Field Theory on compact riemann surfaces. 2022. Preprint, arxiv:2108.06767.
- P. Saad, S. H. Shenker, and D. Stanford. A semiclassical ramp in SYK and in gravity. 2018. Preprint, arxiv:1806.06840.
- P. Saad, S. H. Shenker, and D. Stanford. JT gravity as a matrix integral. Mar. 2019. Preprint, arXiv:1903.11115.
- A. Sengupta. *Gauge theory on compact surfaces*. 1997. ISBN 978-1-4704-0185-6.
- D. Stanford and E. Witten. Fermionic Localization of the Schwarzian Theory. *JHEP*, 10:008, 2017.
- E. M. Stein and G. Weiss. *Introduction to Fourier Analysis on Euclidean Spaces*. Princeton University Press, 1971. ISBN 9780691080789.
- J. Sung and Y. Wang. Quasiconformal deformation of the chordal Loewner driving function and first variation of the Loewner energy. *Mathematische Annalen*, pages 1–24, 04 2024.