

# Exponential turnpike property for particle systems and mean-field limit

Michael Herty<sup>\*</sup>    Yizhou Zhou<sup>†</sup>

June 26, 2024

## Abstract

This work is concerned with the exponential turnpike property for optimal control problems of particle systems and their mean-field limit. Under the assumption of the strict dissipativity of the cost function, exponential estimates for both optimal states and optimal control are proven. Moreover, we show that all the results for particle systems can be preserved under the limit in the case of infinitely many particles.

**Keywords:** Exponential turnpike property, mean-field limit, optimal control

**AMS subject classification:** 93C20, 35L04, 35Q89, 49N10

## 1 Introduction

For optimal control problems of time-dependent differential equations, the exponential turnpike property states, that the optimal solution remains (exponentially) close to a reference solution. Usually, this reference solution is taken as the optimal solution to the corresponding static problem. The concept of turnpike was first introduced for discrete-time optimal control problems [15, 29]. Since then, many turnpike results have been established and there has been recent interest in the mathematical community [14, 22, 23, 20, 24, 21].

In the present work, we focus on the exponential turnpike phenomenon for optimal control problems of a class of interacting particle systems and their mean-field limit equations. Important applications for these systems occur in the fields of swarm robotics [13], crowd dynamics [3], traffic management [30], or opinion dynamics [5] to name but a few.

The original formulation of the interacting particle system is usually at the so-called microscopic level and given by a coupled system of ODEs. Alternatively, one can also focus on

---

<sup>\*</sup>Chair in Numerical Analysis, IGPM, RWTH Aachen University, Templergraben, 55, D-52062 Aachen, Germany (herty@igpm.rwth-aachen.de)

<sup>†</sup>IGPM, RWTH Aachen University, Templergraben, 55, D-52062 Aachen, Germany (zhou@igpm.rwth-aachen.de)

the collective behavior by considering the probability density distribution of the particles and investigating the corresponding McKean-Vlasov or mean-field equation, see e.g. [8, 9, 1] for results involving control actions. The control of large-scale interacting particle systems has gained recent interest, see e.g., [5, 10, 12]. The control of high-dimensional system is challenging and current approaches resort to e.g. using Riccati-based [27, 2], moment-driven control [4], or model predictive control approaches [6, 5, 31]. Motivated by this, we aim to utilize the turnpike property to control those high-dimensional systems [7, 28, 33]. More precisely, we prove the exponential turnpike estimate for ODE systems with an arbitrary particle number and show that the property also holds in the mean-field limit. Here, we utilize the particular structure of interacting particle systems to derive the turnpike property.

The topic of turnpike property for mean-field optimal control problems has been studied recently in [26]. At this point, we would compare [26] with the present paper and point out our main contributions. (1) In [26], the authors prove the turnpike property with interior decay [25] which is a time integral property [18]. In the present paper, under similar assumptions (with a minor modification), we present a point-wise exponential estimate which is more quantitative. (2) In addition to the estimate of the optimal solution, we also prove the exponential decay for the optimal control.

As in [26], our basic assumption is that the optimal control problems satisfy a strict dissipativity inequality. By considering a feedback control, we obtain the cheap control inequality. Then, we use this inequality iteratively to prove the exponential estimate for the optimal solution. This iteration technique has also been used to prove the turnpike property for other optimal control problems (See i.e [16]). Note that all the estimates for particle systems are independent of the particle number  $N$ . Thus all results are also expected in the mean-field level as  $N \rightarrow \infty$ . By using convergence in the Wasserstein distance and the lower semi-continuity of the cost function, we prove the corresponding exponential decay property for the solution of the mean-field optimal control problem. In order to establish the exponential decay for the optimal control, we design a specific feedback control, see also [17]. In this way, the optimal control can be bounded by the optimal solution. Combining with the estimate for solutions, we also prove the exponential decay property for the optimal control with respect to time  $t$ .

The paper is organized as follows. In Section 2, we state the problem and present some basic assumptions. In Section 3, we prove the cheap control property for the optimal control problem of the particle system. By considering the limit  $N \rightarrow \infty$ , we prove the same property in the mean-field level. Based on these results, we prove the exponential turnpike property for both the particle system and the mean-field problem in Section 4. At last, the auxiliary estimate in the Wasserstein distance is given in Appendix A. The main results are Theorem 4.3 on the exponential turnpike property for the particle system and Theorem 4.4-4.5 for the mean-field problem.

## 2 Preliminaries

Consider the optimal control problem  $\mathcal{Q}(0, T, \mu_0)$ :

$$\begin{aligned} \mathcal{V}(0, T, \mu_0) &= \min_{u \in \mathcal{F}} \int_0^T f(\mu(t, x), u(t, x)) dt, \\ &:= \min_{u \in \mathcal{F}} \int_0^T \int L(x) d\mu(t, x) dt + \int_0^T \int \Psi(u(x, t)) d\mu(t, x) dt. \end{aligned} \quad (2.1)$$

Here  $\mu(t, \cdot) \in P_2(\mathbb{R}^d)$  is a probability measure on  $\mathbb{R}^d$  defined for  $t \in [0, T]$  and it satisfies the following equation in a distributional sense

$$\begin{aligned} \partial_t \mu + \nabla_x \cdot \left( (P * \mu + u) \mu \right) &= 0, \quad 0 < t < T, \quad x \in \mathbb{R}^d, \\ \mu(0, x) &= \mu_0(x). \end{aligned} \quad (2.2)$$

Here,  $P(x) \in \mathbb{R}^d$  is a vector-valued function and

$$(P * \mu)(x, t) = \int_{\mathbb{R}^d} P(x - y) d\mu(t, y).$$

As that in [19], we take the control  $u(t, x) \in \mathcal{F}$  satisfying

**Definition 2.1.** Fix a control bound  $0 < C_B < \infty$ . Then  $u(t, x) \in \mathcal{F}$  if and only if

- (i)  $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  is a Carathéodory function.
- (ii)  $u(t, \cdot) \in W_{loc}^{1, \infty}(\mathbb{R}^d)$  for almost every  $t \in [0, T]$ .
- (iii)  $|u(t, 0)| + \|u(t, \cdot)\|_{Lip} \leq C_B$  for almost every  $t \in [0, T]$ .

**Remark 2.1.** In [19], the control bound can be chosen as an integrable function  $l(t) \in L^q(0, T)$  for  $1 \leq q < \infty$ . For simplicity, we take the bound to be constant.

Next, we show assumptions for the optimal control problem (2.2).

**Assumption 2.1.** The cost function  $f$  satisfies the following assumptions:

- (i) *Strict dissipativity:* there exists a constant  $C_D$  such that, for any  $b \geq a \geq 0$  and any pairs  $(\mu(t, x), u(t, x)) \in P_2(\mathbb{R}^d) \times \mathcal{F}$ , the following inequality holds

$$\int_a^b f(\mu(t, x), u(t, x)) dt \geq C_D \int_a^b \int_{\mathbb{R}^d} \left( |x - \bar{x}|^2 + |u(t, x)|^2 \right) d\mu(t, x) dt.$$

- (ii) There exist constants  $C_\Psi$  and  $C_L$  such that

$$\Psi(u) \leq C_\Psi |u|^2 \quad \text{and} \quad L(x) \leq C_L |x - \bar{x}|^2. \quad (2.3)$$

It holds for all  $x \in B(\bar{x}, R) := \{x \in \mathbb{R}^d : |x - \bar{x}| < R\}$  and  $u \in B(0, R)$ .

(iii) The interaction function  $P(x)$  satisfies  $P(0) = 0$  and the following Lipschitz property:

$$|P(x) - P(y)| \leq C_p |x - y|, \quad \forall x \in \mathbb{R}^d \quad (2.4)$$

with  $C_p > 0$  a constant.

**Remark 2.2.** These assumptions are also used in [26] except for condition (ii). Here, we need to assume that both  $\Psi$  and  $L$  can be bounded by quadratic functions. Note that, this assumption is also satisfied for the example of [26].

For further discussion of the optimal control problem, we consider the empirical measure on  $[0, T] \times \mathbb{R}^d$

$$\mu_N(t, x) = \frac{1}{N} \sum_{i=1}^N \delta(x - x_i(t)). \quad (2.5)$$

Here,  $x_i(t)$  ( $i = 1, 2, \dots, N$ ) is the solution to the optimal control problem  $\mathcal{Q}_N(0, T, x_0)$ :

$$\begin{aligned} \mathcal{V}_N(0, T, x_0) &= \min_{u_N \in \mathcal{F}} \frac{1}{N} \sum_{i=1}^N \int_0^T L(x_i(t)) + \Psi(u_N(t, x_i(t))) dt, \\ \frac{dx_i(t)}{dt} &= \frac{1}{N} \sum_{j=1}^N P(x_i(t) - x_j(t)) + u_N(t, x_i(t)), \end{aligned} \quad (2.6)$$

$$x_i(0) = x_{i0}.$$

Here,  $x(t) = (x_1(t), x_2(t), \dots, x_N(t))$  represents  $N$  particles,  $x_0 = (x_{10}, x_{20}, \dots, x_{N0})$  is the initial data, and  $u_N(t, x_i(t))$  is the control. We use the subscript  $N$  to emphasize the dependence of the optimal control  $u_N$  of (2.6) on the number of particles  $N$ .

**Remark 2.3.** Problem  $\mathcal{Q}_N(0, T, x_0)$  can be formally derived from the original optimal control problem. For any  $N$  we have

$$\begin{aligned} f(\mu_N, u_N) &= \int L(x) d\mu_N(t, x) + \int \Psi(u_N(t, x)) d\mu_N(t, x) \\ &= \frac{1}{N} \sum_{i=1}^N \left[ L(x_i(t)) + \Psi(u_N(t, x_i(t))) \right]. \end{aligned}$$

which implies that the cost function in (2.6) is given by

$$\mathcal{V}_N(0, T, x_0) = \min_{u_N \in \mathcal{F}} \int_0^T f(\mu_N, u_N) dt.$$

As outlined in the remark, the optimal control problem (2.6) and the original problem are intertwined. Under Assumptions 2.1, the existence and uniqueness of the problem (2.2)-(2.1) has been established in [19]. To recall the theorem, the definition of the  $p$ -Wasserstein distance between two probability measures  $\mu$  and  $\nu$  is given:

$$\mathcal{W}_p(\mu, \nu) = \inf_{\gamma \in \Gamma(\mu, \nu)} \left( \int_{\mathbb{R}^{2d}} |x - y|^p d\gamma(x, y) \right)^{1/p}.$$

Here,  $\Gamma(\mu, \nu)$  denotes the set of transport plans, i.e., collection of all probability measures with marginals  $\mu$  and  $\nu$ , see also [32].

**Theorem 2.1.** *Assume that the initial data  $\mu_0$  in (2.2) is compactly supported, i.e., there exists  $R > 0$  such that  $\text{supp } \mu_0 \subset B(0, R)$ . Moreover, the empirical measure  $\mu_N(x, 0)$  converges to  $\mu_0$  in  $\mathcal{W}_1$  distance. Then, there exists an optimal control  $u(t, x)$  and a weak equi-compactly supported solution  $\mu(t, x)$  to the problem (2.1)-(2.2). Namely, for all  $t \in [0, T]$  the distribution  $\mu(t, x) \in C([0, T]; P_1(\mathbb{R}^d))$  satisfies  $\text{supp } \mu(t, \cdot) \subset B(0, R)$  and*

$$\begin{aligned} & \int \phi(t, x) d\mu(t, x) - \int \phi(0, x) d\mu_0(x) \\ &= \int_0^t \int \left[ \partial_t \phi + \nabla_x \phi \cdot (P * \mu + u) \right] d\mu(s, x) ds, \quad \forall \phi \in C_0^\infty([0, T] \times \mathbb{R}^d). \end{aligned} \tag{2.7}$$

The optimal solution satisfies

$$\lim_{k \rightarrow \infty} \mathcal{W}_1(\mu_{N_k}(t, \cdot), \mu(t)) = 0$$

uniformly with respect to  $t \in [0, T]$  and  $u_{N_k}$  converges to  $u$  in  $\mathcal{F}$ . Here  $\mu_{N_k}$  is given by (2.5) and  $(x_i(t), u_{N_k}(t, x))$  is the optimal solution to (2.6) with  $N_k$  particles. Moreover,  $f$  is lower semi-continuous:

$$\int_0^T f(\mu(t, x), u(t, x)) dt \leq \liminf_{k \rightarrow \infty} \int_0^T f(\mu_{N_k}(t, x), u_{N_k}(t, x)) dt. \tag{2.8}$$

For the exponential stability later, we discuss solutions  $\mu(t, x)$  in  $C([0, T]; P_2(\mathbb{R}^d))$  with metric  $\mathcal{W}_2$ . By adapting the method in [11, 19], we have

**Lemma 2.2.** *For fixed control  $u(t, x)$ , if  $\mu(t, x), \nu(t, x)$  are solutions to (2.2) with initial data  $\mu_0$  and  $\nu_0$  satisfying the assumption in Theorem 2.1, then there is a constant  $C > 0$  such that*

$$\mathcal{W}_2(\mu(t, \cdot), \nu(t, \cdot)) \leq e^{Ct} \mathcal{W}_2(\mu_0, \nu_0) \quad \text{for } t \in [0, T].$$

Some remarks are in order. The proof is similar to [11, 19] for the stability in  $\mathcal{W}_1$  and deferred to the Appendix A. Hence, the optimal solution is unique in  $C([0, T]; P_2(\mathbb{R}^d))$  if the initial data  $\mu_0 \in P_2(\mathbb{R}^d)$ . Due to this argument, we assume that the optimal solution  $\mu(t, x)$  also satisfies

$$\lim_{k \rightarrow \infty} \mathcal{W}_2(\mu_{N_k}(t, \cdot), \mu(t, \cdot)) = 0 \tag{2.9}$$

uniformly with respect to  $t \in [0, T]$ . The assumption is justified since we have the convergence in  $\mathcal{W}_1$  and the uniform boundness of the second order moment for  $\mu_N(t, \cdot)$  with respect to  $N$ , see e.g. Theorem 4.3.

### 3 Cheap control property

The cheap control property of the optimal control problem shows that the optimal values are bounded by the distance between the initial state and the desired static state. Combining the cheap control property with the strict dissipativity, we provide a bound on the second-order moments of the probability density. More specifically, for the  $N$ -particles system (2.6), we prove:

**Lemma 3.1.** *Suppose  $u_N$  is an optimal control to the problem  $\mathcal{Q}_N(0, T, x_0)$  and  $x(t)$  is the corresponding solution, then  $u_N|_{t \in [a, T]}$  is also an optimal control to the sub-problem  $\mathcal{Q}_N(a, T, x(a))$  for any  $a \geq 0$ . Moreover, the following inequality holds under Assumption 2.1:*

$$\frac{1}{N} \sum_{i=1}^N \int_a^T |x_i(t) - \bar{x}|^2 + |u_N(t, x_i(t))|^2 dt \leq C_0 \frac{1}{N} \sum_{i=1}^N |x_i(a) - \bar{x}|^2. \quad (3.1)$$

Here,  $C_0$  is a positive constant independent of  $N$ .

*Proof.* Suppose there exists a control  $\tilde{u}_N$ , defined on  $t \in [a, T]$ , such that the corresponding solution  $\tilde{x}(t)$  satisfies  $\tilde{x}(a) = x(a)$  and

$$\int_a^T f(\tilde{\mu}_N, \tilde{u}_N) dt < \int_a^T f_N(\mu_N, u_N) dt.$$

Here  $\tilde{\mu}_N$  is the empirical measure given by

$$\tilde{\mu}_N = \frac{1}{N} \sum_{i=1}^N \delta(x - \tilde{x}_i(t)).$$

Then, we construct a control

$$\hat{u}_N(t, x) = \begin{cases} u_N(t, x), & t \in [0, a) \\ \tilde{u}_N(t, x), & t \in [a, T]. \end{cases}$$

In this case, the cost satisfies

$$\int_0^T f(\hat{\mu}_N, \hat{u}_N) dt = \int_0^a f(\mu_N, u_N) dt + \int_a^T f(\tilde{\mu}_N, \tilde{u}_N) dt < \int_0^T f(\mu_N, u_N) dt.$$

This contradicts to the fact that  $(x(t), u_N(t))$  is an optimal solution on  $[0, T]$ . Therefore,  $u_N|_{t \in [a, T]}$  is an optimal control for the sub-problem  $\mathcal{Q}_N(a, T, x(a))$ .

Thanks to the strict dissipativity, we have

$$\begin{aligned} \int_a^T f(\mu_N, u_N) dt &\geq C_D \int_a^T \int_{\mathbb{R}^d} (|x - \bar{x}|^2 + |u_N(t, x)|^2) d\mu_N(t, x) dt \\ &= C_D \frac{1}{N} \sum_{i=1}^N \int_a^T |x_i(t) - \bar{x}|^2 + |u_N(t, x_i(t))|^2 dt. \end{aligned}$$

By Remark 2.3, we obtain the estimate (3.1) once we prove the following cheap control inequality:

$$\int_a^T f(\mu_N, u_N) dt = \frac{1}{N} \sum_{i=1}^N \int_a^T L(x_i(t)) + \Psi(u_N(t, x_i(t))) dt \leq \tilde{C}_0 \frac{1}{N} \sum_{i=1}^N |x_i(a) - \bar{x}|^2 \quad (3.2)$$

for a constant  $\tilde{C}_0 > 0$  independent of  $N$ .

Next, we focus on the proof of (3.2). To this end, we consider the feedback control for the problem (2.6):

$$\tilde{u}_N(t, \tilde{x}_i(t)) = -\beta(\tilde{x}_i(t) - \bar{x}) - \frac{1}{N} \sum_{j=1}^N P(\tilde{x}_i(t) - \tilde{x}_j(t)), \quad i = 1, 2, \dots, N, \quad t \in [a, T].$$

Note that  $\tilde{u}_N \in \mathcal{F}$  holds. Indeed, due to assumption (2.4), we have that

$$\begin{aligned} |\tilde{u}_N(t, x) - \tilde{u}_N(t, y)| &= \left| \beta(x - y) + \frac{1}{N} \sum_{j=1}^N [P(x - \tilde{x}_j(t)) - P(y - \tilde{x}_j(t))] \right| \\ &\leq \beta|x - y| + C_P \frac{1}{N} \sum_{j=1}^N |x - y| = (\beta + C_P)|x - y|, \end{aligned}$$

which gives a Lipschitz constant for  $\tilde{u}_N(t, \cdot)$ . Based on this feedback control,  $\tilde{x}_i(t)$  satisfies the equation

$$\frac{d\tilde{x}_i(t)}{dt} = -\beta(\tilde{x}_i(t) - \bar{x}), \quad \tilde{x}_i(a) = x_i(a).$$

It follows that

$$|\tilde{x}_i(t) - \bar{x}|^2 = e^{-2\beta(t-a)} |\tilde{x}_i(a) - \bar{x}|^2 = e^{-2\beta(t-a)} |x_i(a) - \bar{x}|^2. \quad (3.3)$$

In the next paragraph, we estimate  $|\tilde{u}_N(t, \tilde{x}_i(t))|^2$ . By definition, we have

$$|\tilde{u}_N(t, \tilde{x}_i(t))|^2 \leq 2\beta^2 |\tilde{x}_i(t) - \bar{x}|^2 + 2 \left| \frac{1}{N} \sum_{j=1}^N P(\tilde{x}_i(t) - \tilde{x}_j(t)) \right|^2.$$

Using Jensen's inequality, we have

$$\left| \frac{1}{N} \sum_{j=1}^N P(\tilde{x}_i(t) - \tilde{x}_j(t)) \right|^2 \leq \frac{1}{N} \sum_{j=1}^N \left| P(\tilde{x}_i(t) - \tilde{x}_j(t)) \right|^2. \quad (3.4)$$

Due to the assumption of  $P(x)$ , we have

$$\begin{aligned} \frac{1}{N} \sum_{j=1}^N \left| P(\tilde{x}_i(t) - \tilde{x}_j(t)) \right|^2 &\leq \frac{C_P^2}{N} \sum_{j=1}^N |\tilde{x}_i(t) - \tilde{x}_j(t)|^2 \\ &\leq 2C_P^2 |\tilde{x}_i(t) - \bar{x}|^2 + \frac{2C_P^2}{N} \sum_{j=1}^N |\tilde{x}_j(t) - \bar{x}|^2. \end{aligned}$$

Then, it follows that

$$|\tilde{u}_N(t, \tilde{x}_i(t))|^2 \leq (2\beta^2 + 4C_P^2) |\tilde{x}_i(t) - \bar{x}|^2 + \frac{4C_P^2}{N} \sum_{j=1}^N |\tilde{x}_j(t) - \bar{x}|^2.$$

We sum  $i$  from 1 to  $N$  and get

$$\frac{1}{N} \sum_{i=1}^N |\tilde{u}_N(t, \tilde{x}_i(t))|^2 \leq C(\beta, C_P) \frac{1}{N} \sum_{i=1}^N |\tilde{x}_i(t) - \bar{x}|^2$$

with  $C(\beta, C_P) = 2\beta^2 + 8C_P^2$ . Since  $u_N$  is optimal in (2.6), we have

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N \int_a^T L(x_i(t)) + \Psi(u_N(t, x_i(t))) dt &\leq \frac{1}{N} \sum_{i=1}^N \int_a^T L(\tilde{x}_i(t)) + \Psi(\tilde{u}_N(t, \tilde{x}_i(t))) dt \\ &\leq (C(\beta, C_P)C_\Psi + C_L) \frac{1}{N} \sum_{i=1}^N \int_a^T |\tilde{x}_i(t) - \bar{x}|^2 dt. \end{aligned}$$

Note that the last inequality is due to Assumption 2.1 (ii). Substituting (3.3) into the last inequality, we have

$$\begin{aligned} &\frac{1}{N} \sum_{i=1}^N \int_a^T L(x_i(t)) + \Psi(u_N(t, x_i(t))) dt \\ &\leq (C(\beta, C_P)C_\Psi + C_L) \left( \int_a^T e^{-2\beta(t-a)} dt \right) \frac{1}{N} \sum_{i=1}^N |x_i(a) - \bar{x}|^2. \end{aligned}$$

It is easy to show that

$$\int_a^T e^{-2\beta(t-a)} dt = \frac{1}{2\beta} e^{-2\beta(t-a)} \Big|_a^T \leq \frac{1}{2\beta}.$$

Then, we conclude

$$\frac{1}{N} \sum_{i=1}^N \int_a^T L(x_i(t)) + \Psi(u_N(t, x_i(t))) dt \leq \tilde{C}_0 \frac{1}{N} \sum_{i=1}^N |x_i(a) - \bar{x}|^2. \quad (3.5)$$

Note that the constant  $\tilde{C}_0 = [C(\beta, C_P)C_\Psi + C_L]/(2\beta)$  is independent of  $N$ .  $\square$

The estimate (3.1) is independent of  $N$ . We consider  $N \rightarrow \infty$  to get the corresponding result for the mean-field problem. To this end, we also need to use the lower semi-continuity of the cost function (2.1). Namely, we prove the following property for the mean-field problem.

**Lemma 3.2.** *Suppose  $(\mu(t, x), u(t, x))$  is the solution to the optimal control problem (2.1)-(2.2), then the following inequality holds under Assumption 2.1:*

$$\int_a^T \int_{\mathbb{R}^d} (|x - \bar{x}|^2 + |u(t, x)|^2) d\mu(t, x) dt \leq C_0 \int |x - \bar{x}|^2 d\mu(a, x). \quad (3.6)$$

*Proof.* Due to lower semi-continuity, we have

$$\begin{aligned} \int_a^T f(\mu(t, x), u(t, x)) dt &\leq \liminf_{k \rightarrow \infty} \int_a^T f(\mu_{N_k}(t, x), u_{N_k}(t, x)) dt \\ &= \liminf_{k \rightarrow \infty} \frac{1}{N_k} \sum_{i=1}^{N_k} \int_a^T L(x_i(t)) + \Psi(u_{N_k}(t, x_i(t))) dt. \end{aligned}$$

On the other hand, since  $u_{N_k}$  is the optimal solution to (2.6), it follows from (3.2) that

$$\begin{aligned} \int_a^T f(\mu(t, x), u(t, x)) dt &\leq \liminf_{k \rightarrow \infty} \frac{1}{N_k} \sum_{i=1}^{N_k} \int_a^T L(x_i(t)) + \Psi(u_{N_k}(t, x_i(t))) dt \\ &\leq \liminf_{k \rightarrow \infty} \tilde{C}_0 \frac{1}{N_k} \sum_{i=1}^{N_k} |x_i(a) - \bar{x}|^2 \\ &= \tilde{C}_0 \int |x - \bar{x}|^2 d\mu(a, x). \end{aligned}$$

Here,  $\tilde{C}_0$  is the constant introduced in Lemma 3.1. Using the strict dissipativity shows that

$$C_D \int_a^T \int_{\mathbb{R}^d} (|x - \bar{x}|^2 + |u(t, x)|^2) d\mu(t, x) dt \leq \int_a^T f(\mu(t, x), u(t, x)) dt \leq \tilde{C}_0 \int |x - \bar{x}|^2 d\mu(a, x).$$

Thus we can take  $C_0 = \tilde{C}_0/C_D$  to conclude the result.  $\square$

We conclude this section with the following remarks:

- The inequality (3.6) is the mean-field limit of relation (3.1).
- The right-hand side of (3.6) is independent of  $T$ . As in other turnpike results, this shows an integral turnpike property. Namely, the second order moments  $\int_{\mathbb{R}^d} (|x - \bar{x}|^2 + |u(t, x)|^2) d\mu(t, x)$  must be small along the largest part of the time-horizon provided that  $T$  is sufficiently large.
- The cheap control idea was also used in [26] to prove the integral turnpike property with interior decay. Different from the results in [26], the present work uses the second-order moment  $\int_{\mathbb{R}^d} |x - \bar{x}|^2 d\mu(a, x)$  as the bound in (3.6) instead of the first-order moment. This is important for the proofs in the next section.

## 4 Exponential turnpike property

In this section, we will prove that the optimal solution to (2.1)-(2.2) converges to the optimal static state exponentially fast. The estimates on the inequalities for the optimal solution  $\mu(t, x)$  and the optimal control  $u(t, x)$  are given separately, see Theorem 4.4 and 4.5 below. To this end, we derive the estimate for the optimal solution  $x_i(t)$  of the  $N$ -particles system. Then, we consider the mean-field limit  $N \rightarrow \infty$  to obtain an estimate for  $\mu(t, x)$ . At last, we prove that the optimal control  $u(t, x)$  can be bounded in terms of the solution  $\mu(t, x)$ .

### 4.1 Estimate for the solution

For the solution  $x_i(t)$  of (2.6), we use Gronwall's inequality to derive

**Lemma 4.1.** *Suppose (3.1) holds, there exists a constant  $C_1 \geq 1$  such that*

$$\frac{1}{N} \sum_{i=1}^N |x_i(t_2) - \bar{x}|^2 \leq C_1 \frac{1}{N} \sum_{i=1}^N |x_i(t_1) - \bar{x}|^2, \quad \forall 0 \leq t_1 \leq t_2 \leq T. \quad (4.1)$$

*Proof.* We estimate  $y_i(t) = x_i(t) - \bar{x}$  by computing:

$$\begin{aligned} \frac{1}{2} \int_{t_1}^{t_2} \frac{d}{dt} \langle y_i(t), y_i(t) \rangle dt &= \int_{t_1}^{t_2} \langle y_i(t), y_i'(t) \rangle dt \\ &= \frac{1}{N} \sum_{j=1}^N \int_{t_1}^{t_2} \langle y_i(t), P(y_i(t) - y_j(t)) \rangle dt + \int_{t_1}^{t_2} \langle y_i(t), u_i(t) \rangle dt. \end{aligned} \quad (4.2)$$

For the second term, we have

$$\int_{t_1}^{t_2} \langle y_i(t), u_i(t) \rangle dt \leq \frac{1}{2} \int_{t_1}^{t_2} |u_i(t)|^2 dt + \frac{1}{2} \int_{t_1}^{t_2} |y_i(t)|^2 dt, \quad (4.3)$$

and for the first term, we have

$$\begin{aligned} \frac{1}{N} \sum_{j=1}^N \int_{t_1}^{t_2} \langle y_i(t), P(y_i(t) - y_j(t)) \rangle dt &\leq \frac{1}{N} \sum_{j=1}^N C_P \int_{t_1}^{t_2} |y_i(t)| |y_i(t) - y_j(t)| dt \\ &\leq \frac{1}{N} \sum_{j=1}^N C_P \int_{t_1}^{t_2} |y_i(t)|^2 + |y_i(t)| |y_j(t)| dt \leq \frac{3C_P}{2} \int_{t_1}^{t_2} |y_i(t)|^2 dt + \frac{C_P}{2N} \sum_{j=1}^N \int_{t_1}^{t_2} |y_j(t)|^2 dt. \end{aligned} \quad (4.4)$$

Combining (4.2)–(4.4) yields

$$\frac{1}{2} \int_{t_1}^{t_2} \frac{d}{dt} \langle y_i(t), y_i(t) \rangle dt \leq \left( \frac{1}{2} + \frac{3C_P}{2} \right) \int_{t_1}^{t_2} |y_i(t)|^2 dt + \frac{C_P}{2N} \sum_{j=1}^N \int_{t_1}^{t_2} |y_j(t)|^2 dt + \frac{1}{2} \int_{t_1}^{t_2} |u_i(t)|^2 dt.$$

We sum  $i$  from 1 to  $N$  and multiply  $1/N$  to obtain

$$\frac{1}{N} \sum_{i=1}^N |y_i(t_2)|^2 \leq \frac{1}{N} \sum_{i=1}^N |y_i(t_1)|^2 + (1 + 4C_P) \frac{1}{N} \sum_{i=1}^N \int_{t_1}^{t_2} |y_i(t)|^2 dt + \frac{1}{N} \sum_{i=1}^N \int_{t_1}^{t_2} |u_i(t)|^2 dt.$$

Combining this with (3.1), we obtain

$$\frac{1}{N} \sum_{i=1}^N |x_i(t_2) - \bar{x}|^2 \leq C_1 \frac{1}{N} \sum_{i=1}^N |x_i(t_1) - \bar{x}|^2, \quad \forall 0 \leq t_1 \leq t_2 \leq T.$$

with  $C_1 = (2 + 4C_P)C_0 + 1$ . □

Combining this lemma with the inequality (3.1), we prove:

**Lemma 4.2.** *Under Assumption 2.1, the following inequality holds for any  $t \in [n\tau, T]$  with a given constant  $\tau > 0$  and an integer  $1 \leq n \leq \frac{T}{\tau}$ :*

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq \left( \frac{C_0 C_1}{\tau} \right)^n \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2.$$

*Proof.* We first prove the case  $n = 1$ . There exists a point  $t_1 \in [0, \tau]$  such that

$$\frac{1}{N} \sum_{i=1}^N |x_i(t_1) - \bar{x}|^2 \leq \frac{1}{\tau} \int_0^\tau \frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 dt \leq \frac{C_0}{\tau} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2.$$

Note that the last inequality follows by (3.1). For any  $t \geq \tau \geq t_1$ , we obtain by Lemma 4.1

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq C_1 \frac{1}{N} \sum_{i=1}^N |x_i(t_1) - \bar{x}|^2 \leq \frac{C_0 C_1}{\tau} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2.$$

Then we suppose the inequality holds for  $n \geq 1$  and prove the result for  $n + 1$ . There exists  $t_n \in [n\tau, (n + 1)\tau]$  such that

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N |x_i(t_n) - \bar{x}|^2 &\leq \frac{1}{\tau} \int_{n\tau}^{(n+1)\tau} \frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 dt \\ &\leq \frac{C_0}{\tau} \frac{1}{N} \sum_{i=1}^N |x_i(n\tau) - \bar{x}|^2 \leq \frac{C_0}{\tau} \left( \frac{C_0 C_1}{\tau} \right)^n \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2. \end{aligned}$$

Thus for any  $t \in [(n + 1)\tau, T]$ , we obtain by Lemma 4.1

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq C_1 \frac{1}{N} \sum_{i=1}^N |x_i(t_n) - \bar{x}|^2 \leq \left( \frac{C_0 C_1}{\tau} \right)^{n+1} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2$$

and this completes the proof.  $\square$

Thanks to the above lemmas, we are in the position to state the main result for the optimal solution  $x_i(t)$  of the particle system (2.6):

**Theorem 4.3.** *Suppose Assumption 2.1 holds. Then there exist constants  $C_2 > 0$  and  $\alpha > 0$  such that the optimal solution for  $\mathcal{Q}_N(0, T, x_0)$  satisfies the exponential turnpike property:*

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq C_2 e^{-\alpha t} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2$$

for any  $t \in (0, T)$ .

*Proof.* In this proof, we need to fix the constant  $\tau$  in Lemma 4.2 such that  $\tau > C_0 C_1$ . Next, we discuss the cases  $t \in (0, \tau)$  and  $t \in [\tau, T)$  separately.

For any  $t \in [\tau, T)$ , we take the integer  $n = \lfloor t/\tau \rfloor$ . Then,  $1 \leq n \leq \frac{T}{\tau}$  and  $t \in [n\tau, T)$  and we obtain by Lemma 4.2:

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq \left( \frac{C_0 C_1}{\tau} \right)^n \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2.$$

Due to the definition of  $n$ , we have  $n > t/\tau - 1$ . Also, the constant  $\tau$  is chosen such that  $\tau > C_0 C_1$ . Thus we have

$$\left( \frac{C_0 C_1}{\tau} \right)^n = \left( \frac{\tau}{C_0 C_1} \right)^{-n} \leq \left( \frac{\tau}{C_0 C_1} \right)^{1-t/\tau}.$$

The exponential estimate is then given by

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq \hat{C}_2 e^{-\alpha t} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2, \quad \forall t \in [\tau, T)$$

with

$$\hat{C}_2 = \frac{\tau}{C_0 C_1}, \quad \alpha = \frac{1}{\tau} \log \left( \frac{\tau}{C_0 C_1} \right) > 0.$$

On the other hand, for  $t \in (0, \tau)$ , we have

$$\hat{C}_2 e^{-\alpha t} \geq \hat{C}_2 e^{-\alpha \tau} = 1.$$

By Lemma 4.1 we have

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq C_1 \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2 \leq C_1 \hat{C}_2 e^{-\alpha t} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2.$$

Recall, that due to the proof of Lemma 4.1,  $C_1 \geq 1$  holds. To combine the results of  $t \in (0, \tau)$  and  $t \in [\tau, T)$ , we take  $C_2 = C_1 \hat{C}_2$  and obtain

$$\frac{1}{N} \sum_{i=1}^N |x_i(t) - \bar{x}|^2 \leq C_2 e^{-\alpha t} \frac{1}{N} \sum_{i=1}^N |x_i(0) - \bar{x}|^2, \quad \forall t \in (0, T). \quad (4.5)$$

□

This theorem implies that the empirical measure has equi-compact support and bounded second-order moments for any number of particles  $N$ . Moreover, we know that the empirical measure  $\mu_N(t, x)$  defined in (2.5) satisfies

$$\mathcal{W}_2(\mu_N(t, \cdot), \delta(x - \bar{x})) \leq \sqrt{C_2} e^{-\alpha t/2} \mathcal{W}_2(\mu_N(0, \cdot), \delta(x - \bar{x})).$$

We established the exponential decay property for the second-order moment of the empirical measures  $\mu_N(t, \cdot)$  with respect to  $t$ :

$$\int |x - \bar{x}|^2 d\mu_N(t, x) \leq C_2 e^{-\alpha t} \int |x - \bar{x}|^2 d\mu_N(0, x).$$

The constant  $C_2$  is independent of  $N$ . Thus, we can use the uniform  $\mathcal{W}_2$  convergence to obtain the exponential turnpike property in the mean field limit. Namely, we have

**Theorem 4.4.** *Suppose Assumption 2.1 holds. For problem  $\mathcal{Q}(0, T, \mu_0)$ , the optimal solution  $\mu(t, x) \in C([0, T]; P_2(\mathbb{R}^d))$  satisfies the exponential turnpike property in the sense that*

$$\int |x - \bar{x}|^2 d\mu(t, x) \leq C_2 e^{-\alpha t} \int |x - \bar{x}|^2 d\mu_0(x)$$

for any  $t \in (0, T)$ . Here the constants  $C_2$  and  $\alpha$  are the same as those in Theorem 4.3.

**Remark 4.1.** *Alternatively, the result on the mean field problem can be also proven by a direct estimate of (2.2). Namely, we may take a test function  $\phi(t, x) = |x - \bar{x}|^2 \chi_R(x)$  with  $\chi_R(x)$  being a mollified characteristic function  $\chi_R(x) = \psi_\delta * \chi_{[-R-\delta, R+\delta]}$ , such that  $\chi_R(x) = 1$  for  $|x| \leq R$ .*

*Then by the same argument as in Lemma 4.1, we have*

$$\int |x - \bar{x}|^2 d\mu(t_2, x) \leq C_1 \int |x - \bar{x}|^2 d\mu(t_1, x), \quad \forall 0 \leq t_1 \leq t_2 \leq T.$$

Similarly, the inequalities analog to those in Lemma 4.2 and Theorem 4.3 can be also obtained.

## 4.2 Estimate on the control

In this subsection, we estimate the optimal control  $u(t, x)$  in the mean field problem. The idea is to construct a novel feedback control and take advantage of the strict dissipativity.

We divide the time interval  $[0, T]$  into three parts:

$$[0, T] = [0, s] \cup [s, s + mh] \cup (s + mh, T].$$

Here  $s \in (0, T)$  is a fixed time point,  $m > 0$  is a scale parameter which will be given later (see (4.22)), and  $h$  is a sufficiently small constant such that  $s + mh \leq T$ . We construct a feedback control  $\hat{u}(t, x)$  by

$$\hat{u}(t, x) = \begin{cases} u(t, x), & t \in [0, s) \\ \frac{1}{m} u\left(s + \frac{t-s}{m}, x\right) - \frac{m-1}{m} (P * \hat{\mu})(t, x) & t \in [s, s + mh] \\ u(t - (m-1)h, x), & t \in (s + mh, T], \end{cases} \quad (4.6)$$

where  $u(t, x)$  is the optimal control to the problem (2.1)-(2.2) on the time interval  $[0, T]$ ,  $\hat{\mu}(t, x)$  is the solution of (2.2) associated to the new control  $\hat{u}(t, x)$ ,

Next, we discuss the solution  $\hat{\mu}(t, x)$  on the different time intervals.

For  $t \in [0, s)$ , we know that  $\hat{u}(t, x) = u(t, x)$  and the initial data satisfies

$$\hat{\mu}(0, \cdot) = \mu_0(\cdot) \quad \text{in } P_2(\mathbb{R}^d).$$

According to the uniqueness of solution to the mean field equation (2.2), it is easy to see that

$$\hat{\mu}(t, \cdot) = \mu(t, \cdot) \quad \text{in } P_2(\mathbb{R}^d), \quad \forall t \in [0, s].$$

Here,  $\mu(t, x)$  is the solution associated to the optimal control  $u(t, x)$ .

On the other hand, for  $t \in [s, s + mh]$ , we use the expression of  $\hat{u}(t, x)$  to compute the equation of  $\hat{\mu}$  (for simplicity in the strong form). A similar computation holds in the weak form.

$$\begin{aligned}
0 &= \partial_t \hat{\mu}(t, x) + \nabla_x \cdot \left( [(P * \hat{\mu})(t, x) + \hat{u}(t, x)] \hat{\mu}(t, x) \right) \\
&= \partial_t \hat{\mu}(t, x) + \nabla_x \cdot \left( \left[ (P * \hat{\mu})(t, x) + \frac{1}{m} u \left( s + \frac{t-s}{m}, x \right) - \frac{m-1}{m} (P * \hat{\mu})(t, x) \right] \hat{\mu}(t, x) \right) \\
&= \partial_t \hat{\mu}(t, x) + \frac{1}{m} \nabla_x \cdot \left( \left[ (P * \hat{\mu})(t, x) + u \left( s + \frac{t-s}{m}, x \right) \right] \hat{\mu}(t, x) \right).
\end{aligned}$$

Moreover, by the first step, we have

$$\hat{\mu}(s, \cdot) = \mu(s, \cdot) \quad \text{in } P_2(\mathbb{R}^d).$$

Thus, the equation for  $\hat{\mu}$  reads (in weak form)

$$\begin{aligned}
&\int \phi(t, x) d\hat{\mu}(t, x) - \int \phi(s, x) d\mu(s, x) \\
&= \int_0^t \int \left[ \partial_t \phi(r, x) + \frac{1}{m} \nabla_x \phi(r, x) \cdot \left( (P * \hat{\mu})(r, x) + u \left( s + \frac{r-s}{m}, x \right) \right) \right] d\hat{\mu}(r, x) dr \quad (4.7) \\
&\quad \forall \phi(t, x) \in C_0^\infty \left( [s, s + mh] \times \mathbb{R}^d \right).
\end{aligned}$$

Since the map

$$t_1 = s + \frac{t-s}{m}$$

is bijective, we consider the test function

$$\phi(t, x) = \hat{\phi}(t_1, x) = \hat{\phi} \left( s + \frac{t-s}{m}, x \right) \quad \text{with } \hat{\phi} \in C_0^\infty \left( [s, s + h] \times \mathbb{R}^d \right)$$

and the formula (4.7) is equivalent to

$$\begin{aligned}
&\int \hat{\phi}(t_1, x) d\hat{\mu}(t, x) - \int \hat{\phi}(s, x) d\mu(s, x) \\
&= \int_0^t \int \left[ \partial_t \hat{\phi}(r_1, x) + \nabla_x \hat{\phi}(r_1, x) \cdot \left( (P * \hat{\mu})(r, x) + u(r_1, x) \right) \right] d\hat{\mu}(r, x) dr_1, \quad (4.8) \\
&\quad \forall \hat{\phi} \in C_0^\infty \left( [s, s + h] \times \mathbb{R}^d \right).
\end{aligned}$$

Here, we use the relation

$$r_1 = s + \frac{r-s}{m}, \quad dr_1 = \frac{1}{m} dr,$$

and obtain that

$$\mu(t_1, x) = \mu \left( s + \frac{t-s}{m}, x \right)$$

is a solution to (4.8). Again,  $\mu(t, x)$  is the solution associated to the optimal control  $u(t, x)$ . Since the solution for (2.2) is unique in  $P_2(\mathbb{R}^d)$ , we have

$$\hat{\mu}(t, \cdot) = \mu(t_1, \cdot) = \mu\left(s + \frac{t-s}{m}, \cdot\right) \quad \text{in } P_2(\mathbb{R}^d), \quad \forall t \in [s, s+mh]. \quad (4.9)$$

In the last interval, for  $t \in (s+mh, T]$ , the control is  $\hat{u}(t, x) = u(t - (m-1)h, x)$  and the equation for  $\hat{\mu}$  reads (in strong form):

$$\begin{aligned} 0 &= \partial_t \hat{\mu}(t, x) + \nabla_x \cdot \left( [(P * \hat{\mu})(t, x) + \hat{u}(t, x)] \hat{\mu}(t, x) \right) \\ &= \partial_t \hat{\mu}(t, x) + \nabla_x \cdot \left( [(P * \hat{\mu})(t, x) + u(t - (m-1)h, x)] \hat{\mu}(t, x) \right). \end{aligned}$$

Considering  $t = s+mh$ , we have

$$\hat{\mu}(s+mh, \cdot) = \mu(s+h, \cdot) \quad \text{in } P_2(\mathbb{R}^d).$$

Thus, the weak form in the time interval  $(s+mh, T]$  reads as

$$\begin{aligned} &\int \phi(t, x) d\hat{\mu}(t, x) - \int \phi(s+mh, x) d\mu(s+h, x) \\ &= \int_0^t \int \left[ \partial_t \phi(r, x) + \nabla_x \phi(r, x) \cdot \left( (P * \hat{\mu})(r, x) + u(t - (m-1)h, x) \right) \right] d\hat{\mu}(r, x) dr \quad (4.10) \\ &\quad \forall \phi(t, x) \in C_0^\infty \left( (s+mh, T] \times \mathbb{R}^d \right). \end{aligned}$$

In the new variable  $t_2 = t - (m-1)h$  and for the test function

$$\phi(t, x) = \hat{\phi}(t_2, x) = \hat{\phi}(t - (m-1)h, x) \quad \text{with} \quad \hat{\phi} \in C_0^\infty \left( (s+h, T - (m-1)h] \times \mathbb{R}^d \right),$$

equation (4.10) reads

$$\begin{aligned} &\int \hat{\phi}(t_2, x) d\hat{\mu}(t, x) - \int \hat{\phi}(s+mh, x) d\mu(s+h, x) \\ &= \int_0^t \int \left[ \partial_t \hat{\phi}(r_2, x) + \nabla_x \hat{\phi}(r_2, x) \cdot \left( (P * \hat{\mu})(r, x) + u(r_2, x) \right) \right] d\hat{\mu}(r, x) dr_2, \quad (4.11) \\ &\quad \forall \hat{\phi} \in C_0^\infty \left( (s+h, T - (m-1)h] \times \mathbb{R}^d \right) \end{aligned}$$

for  $r_2 = r - (m-1)h$  and  $dr_2 = dr$ . It is easy to see that  $\mu(t_2, x) = \mu(t - (m-1)h, x)$  satisfies (4.11). At last, we use the uniqueness of (2.2) in  $P_2(\mathbb{R}^d)$  to conclude that

$$\hat{\mu}(t, \cdot) = \mu(t_2, \cdot) = \mu(t - (m-1)h, \cdot) \quad \text{in } P_2(\mathbb{R}^d), \quad \forall t \in (s+mh, T]. \quad (4.12)$$

Summarizing, we have

$$\hat{\mu}(t, \cdot) = \begin{cases} \mu(t, \cdot), & t \in [0, s), \\ \mu\left(s + \frac{t-s}{m}, \cdot\right), & t \in [s, s+mh], \\ \mu(t - (m-1)h, \cdot), & t \in (s+mh, T]. \end{cases} \quad (4.13)$$

### 4.3 The turnpike estimate

Having the feedback control  $\hat{u}(t, x)$  and its associated solution  $\hat{\mu}(t, x)$ , we proceed to estimate the optimal control  $u(t, x)$ :

**Theorem 4.5.** *Suppose Assumption 2.1 holds. Then there exists a constant  $C_3 > 0$  such that the optimal control  $u(t, x) \in \mathcal{F}$  for  $\mathcal{Q}(0, T, \mu_0)$  satisfies the exponential turnpike property:*

$$\int |u(t, x)|^2 d\mu(t, x) \leq C_3 e^{-\alpha t} \int |x - \bar{x}|^2 d\mu_0(x) \text{ for a.e. } t \in (0, T).$$

*Proof.* Since  $u(t, x)$  is optimal, we have

$$\begin{aligned} & \int_0^T f(\mu(t, x), u(t, x)) dt \leq \int_0^T f(\hat{\mu}(t, x), \hat{u}(t, x)) dt \\ & = \int_0^s f(\hat{\mu}(t, x), \hat{u}(t, x)) dt + \int_s^{s+mh} f(\hat{\mu}(t, x), \hat{u}(t, x)) dt + \int_{s+mh}^T f(\hat{\mu}(t, x), \hat{u}(t, x)) dt. \end{aligned} \quad (4.14)$$

According to (4.6) and (4.13), we have

$$\int_0^s f(\hat{\mu}(t, x), \hat{u}(t, x)) dt = \int_0^s f(\mu(t, x), u(t, x)) dt \quad (4.15)$$

and

$$\begin{aligned} & \int_{s+mh}^T f(\hat{\mu}(t, x), \hat{u}(t, x)) dt = \int_{s+mh}^T f(\mu(t - (m-1)h, x), u(t - (m-1)h, x)) dt \\ & = \int_{s+h}^{T-(m-1)h} f(\mu(t, x), u(t, x)) dt \leq \int_{s+h}^T f(\mu(t, x), u(t, x)) dt. \end{aligned} \quad (4.16)$$

Therefore, it follows by (4.14)-(4.16)

$$\begin{aligned} & \int_s^{s+h} f(\mu(t, x), u(t, x)) dt \leq \int_s^{s+mh} f(\hat{\mu}(t, x), \hat{u}(t, x)) dt \\ & \leq C_4 \int_s^{s+mh} \int |x - \bar{x}|^2 + |\hat{u}(t, x)|^2 d\hat{\mu}(t, x) dt \end{aligned} \quad (4.17)$$

with  $C_4 = \max\{C_\Psi, C_L\}$ . Notice that the last inequality is due to Assumption 2.1. Moreover, we use (4.6) and (4.13) to obtain

$$\begin{aligned} & C_4 \int_s^{s+mh} \int |x - \bar{x}|^2 + |\hat{u}(t, x)|^2 d\hat{\mu}(t, x) dt \\ & = C_4 \int_s^{s+mh} \int |x - \bar{x}|^2 + \left| \frac{1}{m} u(t_1, x) - \frac{m-1}{m} (P * \mu)(t_1, x) \right|^2 d\mu(t_1, x) dt \end{aligned}$$

with  $t_1 = s + \frac{t-s}{m}$ . By change of variables, the above inequality yields

$$\begin{aligned}
& C_4 \int_s^{s+mh} \int |x - \bar{x}|^2 + |\hat{u}(t, x)|^2 d\hat{\mu}(t, x) dt \\
& \leq mC_4 \int_s^{s+h} \int |x - \bar{x}|^2 + \left| \frac{1}{m}u(t, x) - \frac{m-1}{m}(P * \mu)(t, x) \right|^2 d\mu(t, x) dt \\
& \leq mC_4 \int_s^{s+h} \int |x - \bar{x}|^2 + \frac{3}{2} \frac{1}{m^2} |u(t, x)|^2 + 3 \left| \frac{m-1}{m}(P * \mu)(t, x) \right|^2 d\mu(t, x) dt. \tag{4.18}
\end{aligned}$$

Note that the last inequality follows from the basic inequality

$$|a + b|^2 \leq \frac{3}{2}|a|^2 + 3|b|^2.$$

Using Jensen's inequality and Assumption 2.1, we have

$$|(P * \mu)(t, x)|^2 \leq \int |P(x - y)|^2 d\mu(t, y) \leq C_P^2 |x - \bar{x}|^2 + C_P^2 \int |y - \bar{x}|^2 d\mu(t, y). \tag{4.19}$$

By (4.17)-(4.19), there exists a constant  $C_5 > 0$  depending on  $C_P, C_\Psi, C_L$  and  $m$ , such that

$$\int_s^{s+h} f(\mu(t, x), u(t, x)) dt \leq \frac{3}{2} \frac{C_4}{m} \int_s^{s+h} \int |u(t, x)|^2 d\mu(t, x) dt + C_5 \int_s^{s+h} \int |x - \bar{x}|^2 d\mu(t, x) dt. \tag{4.20}$$

On the other hand, by the strict dissipativity we obtain

$$\begin{aligned}
\int_s^{s+h} f(\mu(t, x), u(t, x)) dt & \geq C_D \int_s^{s+h} \int |x - \bar{x}|^2 + |u(t, x)|^2 d\mu(t, x) dt \\
& \geq C_D \int_s^{s+h} \int |u(t, x)|^2 d\mu(t, x) dt. \tag{4.21}
\end{aligned}$$

By equation (4.20)-(4.21) we conclude that

$$\int_s^{s+h} \int |u(t, x)|^2 d\mu(t, x) dt \leq \frac{3}{2} \frac{C_4}{mC_D} \int_s^{s+h} \int |u(t, x)|^2 d\mu(t, x) dt + \frac{C_5}{C_D} \int_s^{s+h} \int |x - \bar{x}|^2 d\mu(t, x) dt.$$

Set

$$m = \max \left\{ 2, \frac{2C_4}{C_D} \right\}, \tag{4.22}$$

and hence,  $\frac{3}{2} \frac{C_4}{mC_D} \leq \frac{3}{4}$ . Therefore,

$$\int_s^{s+h} \int |u(t, x)|^2 d\mu(t, x) dt \leq \frac{3}{4} \int_s^{s+h} \int |u(t, x)|^2 d\mu(t, x) dt + \frac{C_5}{C_D} \int_s^{s+h} \int |x - \bar{x}|^2 d\mu(t, x) dt.$$

Since  $m$  is given, we know that the constant  $C_5 > 0$  depends only on  $C_P, C_\Psi$ , and  $C_L$ , respectively. This holds for any  $h$  satisfying  $s + mh \leq T$ . By Lebesgue's differentiation theorem we obtain  $\int |u(s, x)|^2 d\mu(s, x) \leq \frac{4C_5}{C_D} \int |x - \bar{x}|^2 d\mu(s, x)$  for a.e.  $t \in (0, T)$ . Combining this estimate with the results of Theorem 4.4, the proof is completed for  $C_3 = \frac{4C_2C_5}{C_D}$ .  $\square$

**Remark 4.2.** By Theorem 4.4 and Theorem 4.5, the cost in (2.1) also decreases exponentially since

$$\begin{aligned} \int_0^T f(\mu(t, x), u(t, x)) dt &\leq C_L \int_0^T \int |x - \bar{x}|^2 d\mu(t, x) dt + C_\Psi \int_0^T \int |u(t, x)|^2 d\mu(t, x) dt \\ &\leq (C_L C_2 + C_\Psi C_3) e^{-\alpha t} \int |x - \bar{x}|^2 d\mu_0(x). \end{aligned}$$

**Remark 4.3.** In the proof, we adapt the technique in [17] by considering a new feedback control and introducing an adaptive parameter  $m$  in (4.22). If the cost function in equation (2.1) of quadratic form,

$$f(\mu(t, x), u(t, x)) = \int |x - \bar{x}|^2 d\mu(t, x) + \int |u(t, x)|^2 d\mu(t, x),$$

then, we have  $C_\Psi = 1$ ,  $C_L = 1$  and  $C_D = 1$ . It follows that  $C_4 = 1$  and  $m = 2$ .

**Remark 4.4.** The exponential turnpike property for the optimal control problem of the  $N$ -particles system (2.6) can also be proved by considering the feedback control

$$\tilde{u}_N(t, \tilde{x}_i(t)) = \begin{cases} u_N(t, \tilde{x}_i(t)), & t \in [0, s) \\ \frac{1}{m} u_N(t_1, \tilde{x}_i(t)) - \frac{m-1}{m} \frac{1}{N} \sum_{j=1}^N P(\tilde{x}_i(t) - \tilde{x}_j(t)) & t \in [s, s + mh) \\ u_N(t_2, \tilde{x}_i(t)), & t \in (s + mh, T], \end{cases}$$

where  $t_1$  and  $t_2$  are taken as those in the proof of Theorem 4.5.

## Acknowledgments

The first author thanks the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) for the financial support through 442047500/SFB1481 within the projects B04 (Sparsity fördernde Muster in kinetischen Hierarchien), B05 (Sparsifizierung zeitabhängiger Netzwerkflußprobleme mittels diskreter Optimierung) and B06 (Kinetische Theorie trifft algebraische Systemtheorie) and through SPP 2298 Theoretical Foundations of Deep Learning within the Project(s) HE5386/23-1, Meanfield Theorie zur Analysis von Deep Learning Methoden (462234017). The second author is funded by Alexander von Humboldt Foundation (Humboldt Research Fellowship Programme for Postdocs).

## A Proof of Lemma 2.2

We follow the idea in [19, 11] to prove the estimate in the Wasserstein distance  $\mathcal{W}_2$  of Lemma 2.2:

Let  $\mathcal{T}_t^\mu$  be the flow map associated to the system

$$\frac{dx(t)}{dt} = (P * \mu)(x(t)) + u(t, x(t)) = \int P(x(t) - y) d\mu(t, y) + u(t, x(t)).$$

We know that  $\mu(t) = \mathcal{T}_t^\mu \# \mu_0$  with  $\mathcal{T}_t^\mu \#$  denotes the push-forward of  $\mu_0$ . Then, we have

$$\begin{aligned} \mathcal{W}_2(\mu(t), \nu(t)) &= \mathcal{W}_2(\mathcal{T}_t^\mu \# \mu_0, \mathcal{T}_t^\nu \# \nu_0) \\ &\leq \mathcal{W}_2(\mathcal{T}_t^\mu \# \mu_0, \mathcal{T}_t^\mu \# \nu_0) + \mathcal{W}_2(\mathcal{T}_t^\mu \# \nu_0, \mathcal{T}_t^\nu \# \nu_0). \end{aligned} \quad (\text{A.1})$$

For the first term, we have the following result.

**Lemma A.1.** *Assume that  $P$  satisfies the Lipschitz condition (2.4) and  $u(t, x) \in \mathcal{F}$ . Then, it holds that*

$$\mathcal{W}_2(\mathcal{T}_t^\mu \# \mu_0, \mathcal{T}_t^\mu \# \nu_0) \leq e^{(C_P + C_B)t} \mathcal{W}_2(\mu_0, \nu_0).$$

*Proof.* Set  $\kappa$  to be an optimal transportation between  $\mu_0$  and  $\nu_0$ . One can check that the measure  $\gamma = (\mathcal{T}_t^\mu \times \mathcal{T}_t^\mu) \# \kappa$  has marginals  $\mathcal{T}_t^\mu \# \mu_0$  and  $\mathcal{T}_t^\mu \# \nu_0$ . Then we have

$$\begin{aligned} \mathcal{W}_2(\mathcal{T}_t^\mu \# \mu_0, \mathcal{T}_t^\mu \# \nu_0) &\leq \left( \int_{\mathbb{R}^d \times \mathbb{R}^d} |x_0 - y_0|^2 d\gamma(x_0, y_0) \right)^{1/2} \\ &= \left( \int_{\mathbb{R}^d \times \mathbb{R}^d} |\mathcal{T}_t^\mu(x_0) - \mathcal{T}_t^\mu(y_0)|^2 d\kappa(x_0, y_0) \right)^{1/2}. \end{aligned} \quad (\text{A.2})$$

Denote  $x(t) = \mathcal{T}_t^\mu(x_0)$  and  $y(t) = \mathcal{T}_t^\mu(y_0)$ . We have

$$\begin{aligned} |x(t) - y(t)| &\leq |x_0 - y_0| + \int_0^t |(P * \mu)(x(s)) - (P * \mu)(y(s))| + |u(s, x(s)) - u(s, y(s))| ds \\ &\leq |x_0 - y_0| + C_P \int_0^t |x(s) - y(s)| ds + C_B \int_0^t |x(s) - y(s)| ds. \end{aligned}$$

By Gronwall's inequality, we have

$$|x(t) - y(t)| \leq e^{(C_P + C_B)t} |x_0 - y_0|.$$

Substituting this into (A.2), we have

$$\mathcal{W}_2(\mathcal{T}_t^\mu \# \mu_0, \mathcal{T}_t^\mu \# \nu_0) \leq e^{(C_P + C_B)t} \left( \int_{\mathbb{R}^d \times \mathbb{R}^d} |x_0 - y_0|^2 d\kappa(x_0, y_0) \right)^{1/2} = e^{(C_P + C_B)t} \mathcal{W}_2(\mu_0, \nu_0).$$

□

For the second term in (A.1), we have the following Lemma.

**Lemma A.2.** *Let  $\mathcal{T}_t^\mu$  and  $\mathcal{T}_t^\nu$  be two flow maps associated to  $\mu(t)$  and  $\nu(t)$ . Suppose the initial data  $\nu_0 \in P_2(\mathbb{R}^d)$ . Then,*

$$\mathcal{W}_2(\mathcal{T}_t^\mu \# \nu_0, \mathcal{T}_t^\nu \# \nu_0) \leq \|\mathcal{T}_t^\mu - \mathcal{T}_t^\nu\|_\infty.$$

*Proof.* The proof is similar to that in Lemma 3.11 in [11]. Consider a transportation plan defined by  $\pi := (\mathcal{T}_t^\mu \times \mathcal{T}_t^\nu) \# \nu_0$ . One can check that this measure has marginals  $\mathcal{T}_t^\mu \# \nu_0$  and  $\mathcal{T}_t^\nu \# \nu_0$ . Then, due to the definition of Wasserstein metric, we have

$$\begin{aligned} \mathcal{W}_2(\mathcal{T}_t^\mu \# \nu_0, \mathcal{T}_t^\nu \# \nu_0) &\leq \left( \int_{\mathbb{R}^d \times \mathbb{R}^d} |x_0 - y_0|^2 \pi(x_0, y_0) dx_0 dy_0 \right)^{1/2} \\ &= \left( \int_{\mathbb{R}^d} |\mathcal{T}_t^\mu(x_0) - \mathcal{T}_t^\nu(x_0)|^2 d\nu_0(x_0) \right)^{1/2} \\ &\leq \|\mathcal{T}_t^\mu - \mathcal{T}_t^\nu\|_\infty. \end{aligned}$$

□

Thanks to this, it suffices to estimate  $\|\mathcal{T}_t^\mu - \mathcal{T}_t^\nu\|_\infty$ . To this end, we state

**Lemma A.3.** *Under the assumptions in Lemma A.1, it holds that*

$$\|\mathcal{T}_t^\mu - \mathcal{T}_t^\nu\|_\infty \leq C_P \int_0^t e^{(C_P + C_B)(t-s)} \mathcal{W}_2(\mu(s), \nu(s)) ds.$$

*Proof.* Denote  $x^\mu(t) = \mathcal{T}_t^\mu(x_0)$  and  $x^\nu(t) = \mathcal{T}_t^\nu(x_0)$ . We compute

$$|x^\mu(t) - x^\nu(t)| \leq \int_0^t |(P * \mu)(x^\mu(s)) - (P * \nu)(x^\nu(s))| ds + \int_0^t |u(s, x^\mu(s)) - u(s, x^\nu(s))| ds. \quad (\text{A.3})$$

For the first term on the right hand side, we compute

$$\begin{aligned} &\int_0^t |(P * \mu)(x^\mu(s)) - (P * \nu)(x^\nu(s))| ds \\ &\leq \int_0^t |(P * \mu)(x^\mu(s)) - (P * \mu)(x^\nu(s))| + |(P * \mu)(x^\nu(s)) - (P * \nu)(x^\nu(s))| ds \\ &\leq C_P \int_0^t |x^\mu(s) - x^\nu(s)| ds + \int_0^t \|(P * \mu)(s, \cdot) - (P * \nu)(s, \cdot)\|_\infty ds. \end{aligned} \quad (\text{A.4})$$

Moreover, using the fact that  $u \in \mathcal{F}$ , it follows from (A.3)-(A.4) that

$$|x^\mu(t) - x^\nu(t)| \leq \int_0^t (C_P + C_B) |x^\mu(s) - x^\nu(s)| ds + \int_0^t \|(P * \mu)(s, \cdot) - (P * \nu)(s, \cdot)\|_\infty ds.$$

By Gronwall's inequality, we have

$$|x^\mu(t) - x^\nu(t)| \leq \int_0^t e^{(C_P + C_B)(t-s)} \|(P * \mu)(s, \cdot) - (P * \nu)(s, \cdot)\|_\infty ds.$$

Denote  $\theta(y, z; t)$  the optimal transportation between  $\mu$  and  $\nu$ . Clearly,  $\theta(y, z; t)$  has marginals  $\mu(t, y)$  and  $\nu(t, z)$ . Thus we compute

$$\begin{aligned} (P * \mu - P * \nu)(t, x) &= \int_{\mathbb{R}^d} P(x - y) d\mu(t, y) - \int_{\mathbb{R}^d} P(x - z) d\nu(t, z) \\ &= \int_{\mathbb{R}^{2d}} [P(x - y) - P(x - z)] d\theta(y, z; t). \end{aligned}$$

It follows from Jensen's inequality that

$$\begin{aligned} |(P * \mu - P * \nu)(t, x)| &\leq \left( \int_{\mathbb{R}^{2d}} |P(x - y) - P(x - z)|^2 d\theta(y, z; t) \right)^{1/2} \\ &\leq C_P \left( \int_{\mathbb{R}^{2d}} |y - z|^2 d\theta(y, z; t) \right)^{1/2} = C_P \mathcal{W}_2(\mu(t), \nu(t)). \end{aligned}$$

Note that it holds for arbitrary  $x \in \mathbb{R}^d$ . Thus we know that

$$|x^\mu(t) - x^\nu(t)| \leq C_P \int_0^t e^{(C_P + C_B)(t-s)} \mathcal{W}_2(\mu(s), \nu(s)) ds.$$

□

Combining Lemma A.1-A.3 with the inequality (A.1), we have

$$\begin{aligned} \mathcal{W}_2(\mu(t), \nu(t)) &\leq \mathcal{W}_2(\mathcal{T}_t^\mu \# \mu_0, \mathcal{T}_t^\mu \# \nu_0) + \mathcal{W}_2(\mathcal{T}_t^\mu \# \nu_0, \mathcal{T}_t^\nu \# \nu_0) \\ &\leq e^{(C_P + C_B)t} \mathcal{W}_2(\mu_0, \nu_0) + C_P \int_0^t e^{(C_P + C_B)(t-s)} \mathcal{W}_2(\mu(s), \nu(s)) ds. \end{aligned}$$

Then we have

$$e^{-(C_P + C_B)t} \mathcal{W}_2(\mu(t), \nu(t)) \leq \mathcal{W}_2(\mu_0, \nu_0) + C_P \int_0^t e^{-(C_P + C_B)s} \mathcal{W}_2(\mu(s), \nu(s)) ds.$$

Again, by Gronwall's inequality, we obtain

$$e^{-(C_P + C_B)t} \mathcal{W}_2(\mu(t), \nu(t)) \leq e^{C_P t} \mathcal{W}_2(\mu_0, \nu_0), \quad t \in [0, T].$$

This completes the proof of the stability with respect to the  $\mathcal{W}_2$  distance.

## References

- [1] G. Albi, N. Bellomo, L. Fermo, S.-Y. Ha, J. Kim, L. Pareschi, D. Poyato, and J. Soler. Vehicular traffic, crowds, and swarms: from kinetic theory and multiscale methods to applications and research perspectives. *Math. Models Methods Appl. Sci.*, 29(10):1901–2005, 2019.

- [2] G. Albi, S. Bicego, and D. Kalise. Gradient-augmented supervised learning of optimal feedback laws using state-dependent Riccati equations. *IEEE Control Syst. Lett.*, 6:836–841, 2022.
- [3] G. Albi, M. Bongini, E. Cristiani, and D. Kalise. Invisible control of self-organizing agents leaving unknown environments. *SIAM J. Appl. Math.*, 76(4):1683–1710, 2016.
- [4] G. Albi, M. Herty, D. Kalise, and C. Segala. Moment-driven predictive control of mean-field collective dynamics. *SIAM J. Control Optim.*, 60(2):814–841, 2022.
- [5] G. Albi, M. Herty, and L. Pareschi. Kinetic description of optimal control problems and applications to opinion consensus. *Commun. Math. Sci.*, 13(6):1407–1429, 2015.
- [6] G. Albi and L. Pareschi. Selective model-predictive control for flocking systems. *Commun. Appl. Ind. Math.*, 9(2):4–21, 2018.
- [7] B. D. O. Anderson and P. V. Kokotovic. Optimal control problems over large time intervals. *Autom.*, 23:355–363, 1987.
- [8] N. Bellomo, P. Degond, and E. Tadmor, editors. *Active particles. Vol. 1. Advances in theory, models, and applications*. Modeling and Simulation in Science, Engineering and Technology. Birkhäuser/Springer, Cham, 2017.
- [9] N. Bellomo, P. Degond, and E. Tadmor, editors. *Active particles. Vol. 2. Advances in theory, models, and applications*. Modeling and Simulation in Science, Engineering and Technology. Birkhäuser/Springer, Cham, 2019.
- [10] M. Bongini, M. Fornasier, O. Junge, and B. Scharf. Sparse control of alignment models in high dimension. *Networks and Heterogeneous Media*, 10:647–697, 09 2015.
- [11] J. A. Cañizo, J. A. Carrillo, and J. Rosado. A well-posedness theory in measures for some kinetic models of collective motion. *Math. Models Methods Appl. Sci.*, 21(3):515–539, 2011.
- [12] M. Caponigro, M. Fornasier, B. Piccoli, and E. Trélat. Sparse stabilization and optimal control of the cucker-smale model. *Mathematical Control and Related Fields*, 3:447–466, 2013.
- [13] Y.-P. Choi, D. Kalise, J. Peszek, and A. A. Peters. A collisionless singular Cucker-Smale model with decentralized formation control. *SIAM J. Appl. Dyn. Syst.*, 18(4):1954–1981, 2019.
- [14] T. Damm, L. Grüne, M. Stieler, and K. Worthmann. An exponential turnpike theorem for dissipative discrete time optimal control problems. *SIAM J. Control Optim.*, 52(3):1935–1957, 2014.
- [15] R. Dorfman, P. A. Samuelson, and R. M. Solow. *Linear programming and economic analysis*. McGraw-Hill Book Co., Inc., New York-Toronto-London, 1958. A Rand Corporation Research Study.

- [16] C. Esteve-Yagüe, B. Geshkovski, D. Pighin, and E. Zuazua. Large-time asymptotics in deep learning, 2021.
- [17] C. Esteve-Yagüe, B. Geshkovski, D. Pighin, and E. Zuazua. Turnpike in Lipschitz-nonlinear optimal control. *Nonlinearity*, 35(4):1652–1701, 2022.
- [18] T. Faulwasser, L. Grüne, J.-P. Humaloja, and M. Schaller. The interval turnpike property for adjoints. *Pure Appl. Funct. Anal.*, 7(4):1187–1207, 2022.
- [19] M. Fornasier and F. Solombrino. Mean-field optimal control. *ESAIM Control Optim. Calc. Var.*, 20(4):1123–1152, 2014.
- [20] L. Grüne. Economic receding horizon control without terminal constraints. *Autom.*, 49:725–734, 2013.
- [21] L. Grüne. Dissipativity and optimal control: examining the turnpike phenomenon. *IEEE Control Syst.*, 42(2):74–87, 2022.
- [22] L. Grüne and M. A. Müller. On the relation between strict dissipativity and turnpike properties. *Systems Control Lett.*, 90:45–53, 2016.
- [23] L. Grüne, M. Schaller, and A. Schiela. Exponential sensitivity and turnpike analysis for linear quadratic optimal control of general evolution equations. *J. Differential Equations*, 268(12):7311–7341, 2020.
- [24] L. Grüne and M. Stieler. Asymptotic stability and transient optimality of economic mpc without terminal conditions. *Control. Bd.*, 24(Heft 8):1187–1196, 2014.
- [25] M. Gugat. On the turnpike property with interior decay for optimal control problems. *Mathematics of Control, Signals, and Systems*, 33:1–22, 06 2021.
- [26] M. Gugat, M. Herty, and C. Segala. The turnpike property for mean-field optimal control problems. *European Journal of Applied Mathematics*, page 1–15, 2024.
- [27] M. Herty, L. Pareschi, and S. Steffensen. Mean-field control and Riccati equations. *Netw. Heterog. Media*, 10(3):699–715, 2015.
- [28] A. M. Sahlodin and P. I. Barton. Optimal campaign continuous manufacturing. *Industrial & Engineering Chemistry Research*, 54(45):11344–11359, 2015.
- [29] P. A. Samuelson. A catenary turnpike theorem involving consumption and the golden rule. *The American Economic Review*, 55(3):486–496, 1965.
- [30] A. Tosin and M. Zanella. Kinetic-controlled hydrodynamics for traffic models with driver-assist vehicles. *Multiscale Model. Simul.*, 17(2):716–749, 2019.
- [31] A. Tosin and M. Zanella. Uncertainty damping in kinetic traffic models by driver-assist controls. *Math. Control Relat. Fields*, 11(3):681–713, 2021.
- [32] C. Villani. *Optimal transport*, volume 338 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2009. Old and new.

- [33] A. J. Zaslavski. Necessary and sufficient turnpike conditions. *Pure Appl. Funct. Anal.*, 4(2):463–476, 2019.