

FUSION INEQUALITY FOR QUADRATIC COHOMOLOGY

OLIVER KNILL

ABSTRACT. Classical simplicial cohomology on a simplicial complex G deals with functions on simplices $x \in G$. Quadratic cohomology [6, 7] deals with functions on pairs of simplices $(x, y) \in G \times G$ that intersect. If K, U is a closed-open pair in G , we prove here a quadratic version of the linear fusion inequality [10]. Additional to the quadratic cohomology of G there are five additional interaction cohomology groups. Their Betti numbers are computed from functions on pairs (x, y) of simplices that intersect. Define the Betti vector $b(X)$ computed from pairs $(x, y) \in X \times X$ with $x \cap y \in X$ and $b(X, Y)$ with pairs in $X \times Y$ with $x \cap y \in K$. We prove the fusion inequality $b(G) \leq b(K) + b(U) + b(K, U) + b(U, K) + b(U, U)$ for cohomology groups linking all five possible interaction cases. Counting shows $f(G) = f(K) + f(U) + f(K, U) + f(U, K) + f(U, U)$ for the f -vectors. Super counting gives Euler-Poincaré $\sum_k (-1)^k f_k(X) = \sum_k (-1)^k b_k(X)$ and $\sum_k (-1)^k f_k(X, Y) = \sum_k (-1)^k b_k(X, Y)$ for $X, Y \in \{U, K\}$. As in the linear case, also the proof of the quadratic fusion inequality follows from the fact that the spectra of all the involved Laplacians $L(X), L(X, Y)$ are bounded above by the spectrum of the quadratic Hodge Laplacian $L(G)$ of G .

1. IN A NUTSHELL

1.1. We prove here that if K is a sub-complex of a finite abstract simplicial complex G and U is the open complement $U = G \setminus K$ [1, 12], there are besides the quadratic cohomology of G five quadratic cohomology groups belonging to the five **quadratic Hodge Laplacians** $L(X), L(X, Y)$ with $X, Y \in \{U, K\}$. They all satisfy spectral inequalities:

Theorem 1 (Spectral inequality). $\lambda_k(L) \leq \lambda_k(L(G))$

1.2. The assumption is that all eigenvalues are ordered in an ascending order and that they are **padded left** in comparison with the eigenvalues of G . This result parallels the linear simplicial cohomology case [13], where U and K can not yet interact and L is one of the Hodge Laplacians $L(K), L(U)$ for simplicial cohomology.

1.3. In the linear case, the Betti vectors satisfied the fusion inequality $b(G) \leq b(K) + b(U)$ [10]. This linear fusion inequality had followed from the spectral inequality and the fact that cohomology classes are null-spaces of matrices. Counting gave $f(G) = f(K) + f(U)$ for the f -vectors and the **Euler-Poincaré formula** was $\chi(X) = \sum_k (-1)^k f_k(X) = \sum_k (-1)^k b_k(X)$ for $X \in \{G, K, U\}$, seen directly by heat deformation using the **McKean-Singer symmetry**, rephrasing that D_X is an isomorphism between even and odd parts of image of the Laplacian $L_X = D_X^2$, implying $\text{str}(L^k) = 0$ for $k \geq 1$ so that $\text{str}(\exp(-tL(X))) = \text{str}(1_X) = \chi(X)$.

1.4. In the quadratic cohomology case, where we look at functions on pairs of intersecting simplices, there is the cohomology of G leading to $b(G)$ and five interaction cohomologies. They each lead to **Betti vectors**. We call them $b(K), b(U), b(K, U), b(U, K)$, and $b(U, U)$. Besides pointing out that we have these new cohomologies, we give here a relation between them and

Date: June 24, 2024.

Key words and phrases. Quadratic cohomology.

the cohomology of G . We call it the **quadratic fusion inequality**. We could use the heavier notation $b(X, Y, Z)$ with $X, Y, Z \in \{K, U\}$ dealing with functions on pairs $(x, y) \in X \times Y$ with $x \cap y \in Z$, but we prefer to stick to the simpler notation: one reason is that both $b(U), b(K)$ deal with internal cohomology of U and K and do not involve simplices of the other set, while $b(K, U), b(U, K), b(U, U)$ involve both sets U and K . So, while the quadratic Betti vectors $b(U), b(K)$ are **intrinsic** and only depend on one of the sets U or K , the others are not. We prove:

Theorem 2 (Quadratic fusion inequality). $b(G) \leq b(K) + b(U) + b(K, U) + b(U, K) + b(U, U)$.

1.5. The quadratic Betti vector $b(X)$ belongs to $\{(x, y), x \cap y \in X\}$, the vector $b(K, U)$ belongs to all $(x, y) \in K \times U$ with $x \cap y \in K$, and $b(U, K)$ belongs to all $(x, y) \in U \times K$ with $x \cap y \in K$, and $b(U, U)$ belongs to all $(x, y) \in U \times U$ with $x \cap y \in K$. Note that $x \cap y \in K$ if $x, y \in K$ so that $b(K, K)$ is accounted for in $b(K)$ already. With the heavier notation, from the eight cases $b(X, Y, Z)$ with $X, Y, Z \in \{K, U\}$ only $b(K) = b(K, K, K), b(U) = b(U, U, U), b(K, U) = b(K, U, K), b(U, K) = b(U, K, K), b(U, U) = b(U, U, K)$ can occur.

1.6. The **quadratic characteristics** are defined as $w(X) = \sum_{x, y \in X, x \cap y \in X} w(x)w(y)$ (Wu characteristic) and $w(X, Y) = \sum_{x \in X, y \in Y, x \cap y \in K} w(x)w(y)$ and $w(U, U) = \sum_{x \in U, y \in U, x \cap y \in U} w(x)w(y)$, we have $w(G) = w(U) + w(K) + w(U, K) + w(K, U) + w(U, U)$ which follows from the fact that f -vectors for quadratic cohomology satisfy $f(G) = f(K) + f(U) + f(K, U) + f(U, K) + f(U, U)$. Any of the cohomologies for $X \in \{G, U, K, (U, K), (K, U), (U, U)\}$ satisfy the Euler-Poincaré formula, following heat deformation with L_X or $L_{X, Y}$ using the McKean-Singer symmetry.

1.7. Why is this interesting? If G is finite abstract simplicial complex that is a finite d -manifold, and f is an arbitrary function from G to $P = \{0, \dots, k\}$, then the **discrete Sard theorem** [11] assures that the open set $U = \{x \in G, f(x) = P\}$ is either empty or a $(d - k)$ -manifold in the sense that the graph with vertex set U and edge set $\{(x, y) \text{ for which } x \subset y \text{ or } y \subset x\}$ is a discrete $(d - k)$ -manifold. Since U is open, the complement $K = G \setminus U$ is closed and so a sub-simplicial complex. All these cohomologies are topological invariants.

1.8. For example, if G is a discrete 3-sphere and $f : G \rightarrow \{0, 1, 2\}$ is arbitrary, then U is either empty, a **knot** or a **link**, a finite union of closed disjoint (possibly interlinked) 1-manifolds in the 3-sphere G . The simplicial cohomology of U and the quadratic cohomology of U are not interesting: they are just circles: $b(U) = (l, l)$, where l is the number of connected components of U . The simplicial cohomology of K however can be interesting and leads to knot or link invariants. It is well studied in the continuum as it is a **knot invariant** or a **link invariant**. Additional interaction cohomologies that take into account interaction between U and the complement K are completely unexplored. The inequality shows however that in general, more cohomology classes are created when splitting up G into $U \cup K$. Unlike in the linear case, we have now the possibility of particles (harmonic forms) that are functions of (x, y) with $x \in U, y \in K$ with $x \cap y \in K$ and also of function on (x, y) with $x \in U, y \in U$ with $x \cap y \in K$.

1.9. The quadratic case we look at here would generalize in a straightforward way to **higher characteristics**. One starts with $m = 1$, the linear characteristic which is **Euler characteristic**. The second $m = 2$ is quadratic characteristic or Wu characteristic going back to Wu in 1959. ¹ In general, we would look for m -tuples of points (x_1, \dots, x_m) that do

¹Historically, multi-linear valuations were considered in [14, 3, 6] and its cohomology in [8]. Quadratic cohomology is to Wu characteristic what simplicial cohomology is to Euler characteristic.

simultaneously intersect. There are then much more m -point interaction cohomologies and $b(G)$ is again bounded above by all possible cases of $b(X_1, \dots, X_m)$ with $X_i \in \{U, K\}$ and the intersection in K . The cases $b(U, U, \dots, U)$ has cases $(x_1, \dots, x_m) \in U^m$ with $\bigcap x_i \in U$ which is part of b_U and a new part where $\bigcap x_i \in K$. As in the case $m = 2$, the cohomology of $b(K, K, \dots, K)$ is part of $b(K)$. We have to consider the Betti vector $b(U)$ for functions on $\{(x_1, \dots, x_m) \in U^m, \bigcap_i x_i \in U\}$ and $b(U, \dots, U)$ referring to functions on $\{(x_1, \dots, x_m) \in U^m, \bigcap_i x_i \in K\}$, where $b(K, \dots, K) = b(K)$ in the notation used before. If all x_i are in U , then the intersection can be either in U or K , but if all x_i are in K , then the intersection must be in K , not warranting to distinguish $b(K, \dots, K)$ and $b(K)$. Despite the obvious duality $U \leftrightarrow K$, there is an asymmetry in that $x \in U, y \in U$ allows $x \cap y$ to be in U or K while $x \in K, y \in K$ implies $x \cap y \in K$: technically, a closed set K is a π -system while an open set U is not unless it is \emptyset or G .

Theorem 3. $b(G) \leq b(U) + b(K) + \sum_{X_i \in \{U, K\}} b(X_1, \dots, X_m)$.

There would again be heavier notation $b(X_1, \dots, X_m, X_0)$ dealing with $(x_1, \dots, x_m) \in X_1 \times \dots \times X_m$ with $\bigcap_{k=1}^m x_k \in X_0$ and have $b(G) \leq \sum_{X_i \in \{U, K\}} b(X_1, \dots, X_m, X_0)$ taking into account that some of the cases like (K, \dots, K, U) are empty because $X_0 = U$ only is interesting if all $X_1 = \dots = X_m = K$. Again, like in the case $m = 2$, we have two cases $b(U), b(K)$ which are **intrinsic** while the other cases involve simplices from both U and K .

1.10. We study here higher order chain complexes in finite geometries. Each of the situations is defined by a triple (X, D, R) , where X is a finite set of n elements, $D = d + d^*$ is a finite $n \times n$ matrix such that $d^2 = 0$ and where R is the dimension function compatible with D in the sense that the blocks of $L = D^2$ have constant dimension. We can call this structure an **abstract delta set** because every delta set defines such a structure, but where instead of face maps, we just go directly to the exterior derivative d . The advantage of looking at the Dirac setting is that D can be much more general than coming from face maps. It could be a deformed Dirac matrix for example obtained by isospectral deformation $D' = [D^+ - D^-, D]$ [5, 4], which keeps the spectrum of D invariant but produces $D = d + d^* + B$ leading to new exterior derivatives $d(t)$, a deformation which is invisible to the Hodge Laplacian as $D^2(t) = L$ is not affected. Since $d(t)$ gets smaller, this produces an expansion of space. In general, also in the continuum, there is an inflationary start of expansion.

2. A SMALL EXAMPLE

2.1. Lets illustrate the quadratic fusion inequality in the case K_2 :

2.2. The linear simplicial cohomology is given by $(\begin{bmatrix} \{1\} \\ \{2\} \\ \{1, 2\} \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}, R = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix})$.

Take the open-closed pair $U = \{\{1, 2\}\}$ and $K = \{\{1\}, \{2\}\}$ leading to the abstract delta set structures $(\begin{bmatrix} \{1\} \\ \{2\} \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix})$ ($\{\{1, 2\}\}, [0], [1]$). The Betti vectors are $b(G) = (1, 0), b(U) = (0, 1), b(K) = (2, 0)$ and the f-vectors are $f(G) = (2, 1), f(U) = (0, 1), f(K) = (2, 0)$. The fusion inequality $b(G) < b(K) + b(U)$ is here strict. Merging U and K fuses a harmonic 0 form in K with the 1-form in U . Betti vectors have been considered since Betti and Poincaré. Finite topological spaces were first looked at by Alexandroff [1]. For cohomology of open sets in finite frame works, see [10].

2.3. If we look at **quadratic cohomology** for G , where we have the abstract delta set $(X, D, R) =$

$$\left(\begin{bmatrix} \{2\} & \{2\} \\ \{1\} & \{1\} \\ \{1, 2\} & \{2\} \\ \{1, 2\} & \{1\} \\ \{2\} & \{1, 2\} \\ \{1\} & \{1, 2\} \\ \{1, 2\} & \{1, 2\} \end{bmatrix}, \begin{bmatrix} 0 & 0 & -1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 1 & -1 & 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \\ 1 \\ 1 \\ 2 \end{bmatrix} \right).$$

The Hodge Laplacian $L = D^2 = L_0 \oplus L_1 \oplus L_2$ has the Hodge blocks:

$$L_0 = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}, L_1 = \begin{bmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{bmatrix}, L_2 = [4]$$

with Betti vector $b(G) = (0, 1, 0)$ and f-vector $f(G) = (2, 4, 1)$ and Wu characteristic $w(G) = f_0 - f_1 + f_2 = 2 - 4 + 1 = b_0 - b_1 + b_2 = 0 - 1 + 0 = -1$. The eigenvalues of L_1 are $(0, 2, 2, 4)$, the null-space is spanned by $[1, 1, 1, 1]$. By accident L_1 happens to be a Kirchoff matrix of C_4 . If K_2 is seen as a 1-manifold with boundary δG ² we have $w(G) = \chi(G) - \chi(\delta(G)) = 1 - 2 = -1$, illustrating that in general, for manifolds M with boundary δM , the Wu characteristic is $\chi(M) - \chi(\delta M)$.

2.4. Now to $U = \{\{1, 2\}\}$, where we have the abstract delta set structure $(X, D, R) =$

$$\left(\begin{bmatrix} \{1, 2\} & \{1, 2\} \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \end{bmatrix} \right).$$

with $b(U) = (0, 0, 1)$ and $f(U) = (0, 0, 1)$ and $w(U) = 1$.

For $K = \{\{1\}, \{2\}\}$ we have the abstract delta set structure $(X, D, R) =$

$$\left(\begin{bmatrix} \{2\} & \{2\} \\ \{1\} & \{1\} \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right)$$

with $b(K) = (2, 0, 0)$ and $f(U) = (2, 0, 0)$ and $w(K) = 2$. Obviously the intrinsic cohomologies of U and K are not yet giving a complete picture. The simplices in U and K can interact as we see next.

2.5. Now, we turn to the interactions of K with U

$$\left(\begin{bmatrix} \{2\} & \{1, 2\} \\ \{1\} & \{1, 2\} \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) \cdot$$

$$\left(\begin{bmatrix} \{1, 2\} & \{2\} \\ \{1, 2\} & \{1\} \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right)$$

with $b(K, U) = b(U, K) = (0, 2, 0)$ and $f(U) = (0, 2, 0)$ and $w(U, K) = -2$. There is no pair $(x, y) \in U^2$ such that $x \cap y \in K$ so that $b(U, U) = (0, 0, 0)$.

²We usually assume that manifolds with boundary have an interior. The Barycentric refinement of a complete graph K_{d+1} would be a d -manifold with boundary.

2.6. To summarize, we have

<i>Case</i>	<i>Betti</i>	<i>f - vector</i>	<i>Characteristic</i>
U	(0, 0, 1)	(0, 0, 1)	1
K	(2, 0, 0)	(2, 0, 0)	2
(U, K)	(0, 2, 0)	(0, 2, 0)	-2
(K, U)	(0, 2, 0)	(0, 2, 0)	-2
(U, U)	(0, 0, 0)	(0, 0, 0)	0
G	(0, 1, 0)	(2, 4, 1)	-1

The quadratic fusion inequality $b(U) + b(K) + b(U, K) + b(K, U) + b(U, U) = (2, 3, 1) > b(G) = (0, 1, 0)$ is here strict. The fusion has two 0-form-1-form mergers and one 1-form-2-form merger. The difference in the fusion inequality is $(1, 1, 0) + (1, 1, 0) + (0, 1, 1) = (2, 3, 1)$.

2.7. We see already in this small example, how the closed “laboratory” K and the “observer space” U are no more strictly separated, even so they partition the “world” G . The “tunneling” between K and U is described using algebraic topology, expressed by cohomology groups. Unlike for simplicial cohomology which features homotopy invariance, there is only **topological invariance**. Already the Wu characteristic of contractible balls depends on the dimension. [For a d -ball, the Wu characteristic $w(M)$ is $(-1)^d$ where d is the dimension and illustrates that $w(M) = \chi(M) - \chi(\delta M)$ in general for discrete manifolds M with boundary δM and that for a d -ball, the boundary is a $d - 1$ sphere with Euler characteristic $\chi(\delta M) = 1 - (-1)^d$.] If we take a d -ball in a d -dimensional simplicial complex and replace the interior to get an other d -ball without changing the boundary, then the cohomology does not change because we can for any positive k add add gauge fields (k -forms that are coboundaries) dg to render a cocycle zero in the interior (without changing the equivalence class) and use the heat flow to get back a harmonic form after doing the surgery in the interior. This implements the **chain homotopy** when doing a local homeomorphic deformation: move the field away from the “surgery place”, do the surgery, then use the heat flow to “heal the wound” and get back harmonic forms.

2.8. We have just given the argument for the following result:

Theorem 4. *All quadratic cohomology groups $b(X), b(X, Y)$ are topological invariants.*

2.9. For $G = K_3, K = \{\{1\}\}$, we can look at the complex $G(U, K) = \begin{bmatrix} \{1\} & \{1, 3\} \\ \{1\} & \{1, 2\} \\ \{1\} & \{1, 2, 3\} \end{bmatrix}$ and

$D(K, U) = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 1 \\ -1 & 1 & 0 \end{bmatrix}$ with kernel spanned by $[1, 1, 0]$. For its Barycentric refinement and

still $K = \{1\}$ (on the boundary), and where we look at functions on $X = \begin{bmatrix} \{1\} & \{1, 7\} \\ \{1\} & \{1, 5\} \\ \{1\} & \{1, 4\} \\ \{1\} & \{1, 5, 7\} \\ \{1\} & \{1, 4, 7\} \end{bmatrix}$,

we have $D(K, U) = \begin{bmatrix} 0 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ -1 & 1 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 \end{bmatrix}$ with kernel spanned by $[1, 1, 1, 0, 0]$.

3. QUADRATIC COHOMOLOGY

3.1. **Simplicial cohomology** for a finite abstract simplicial complex G is part of the spectral theory of the **Hodge Laplacian** $L = D^2$ with **Dirac matrix** $D = d + d^*$, where d is the exterior derivative. Note that all these matrices d, D, L are $n \times n$ matrices if G has n elements. The matrix L is a block diagonal matrix $L = \bigoplus_{k=0}^d L_k$. The kernels of the blocks L_k of L are the **k -harmonic forms** or **k -cohomology** vector spaces. In this finite setting, this is linear algebra [2]. The dimensions b_k are the **Betti numbers**, the components of the Betti vector of G .

3.2. If K is a subcomplex of G and U is the open complement, then the **separated system** (K, U) has a Laplacian $L_{K,U} = L_K \oplus L_U$ for which the energies $\lambda_j(L_{K,U})$ are less or equal than $2\lambda_j(L_G)$ [13] implying that the separated system can not have more harmonic forms than G . It can have more: if G is a closed 2-ball for example and K is the boundary 1-sphere then $b_G = (1, 0, 0)$, $b_K = (1, 1, 0)$ and $b_U = (0, 0, 1)$. The closed part K carries a trapped harmonic 1-form. It is fused with the 2-form present on U , if K, U get united to G .

3.3. A complex G defines a delta set $G = \bigcup_{k=0}^d G_k$. The **f -vector** $f(G) = (f_0(G), \dots, f_d(G))$ has components $f_k(G) = |G_k|$, the number of elements in G_k . The **super trace** of an $n \times n$ matrix L ³ is defined as $\text{str}(L) = \sum_{k=0}^d (-1)^k \sum_{x \in G_k} L(x, x)$. Compare with the usual trace $\text{tr}(L) = \sum_{k=0}^d \sum_{x \in G_k} L(x, x) = \sum_{x \in G} L(x, x)$. The Euler characteristic is $\chi(G) = \sum_{x \in G} w(x)$. The **Euler-Poincaré formula** $\chi(G) = \sum_k (-1)^k f_k = \sum_k (-1)^k b_k$ follows directly from the **McKean-Singer identity**, stating that $\text{str}(\exp(-itL)) = \chi(G)$ for all t which in turn follows from the fact that the Dirac matrix D gives an isomorphism between even and odd **non-harmonic forms**. For $t = 0$, the super trace of the heat kernel is the combinatorial Euler characteristic, while for $t = \infty$, it is the cohomological Euler characteristic.

3.4. **Quadratic cohomology** does not build on single simplices $x \in G$ like simplicial cohomology but on **pairs of intersecting simplices** $(x, y) \in G \times G$. Define $w(x) = (-1)^{\dim(x)}$. The quadratic analog of (linear) Euler characteristic $\chi(A) = \sum_{x \in A} w(x)$ is the “Ising type” energy or **Wu characteristic** $w(A) = \sum_{x, y, x \cap y \in A} w(x)w(y)$. It is an example of a multi-linear valuation. We also just call it **quadratic characteristic**, an example of **higher characteristic**. [9].

3.5. The name “quadratic” is chosen because it is multi-linear and for $m = 2$ a quadratic valuation. Similarly as a quadratic form is a multi-linear map, linear in each argument, the quadratic characteristic $w(A, B) = \sum_{x \in A, y \in B, x \cap y \neq \emptyset} w(A)w(B)$ (or variants, where we ask the intersection to be in A or B) satisfies the valuation formula in each of the coordinates, like $w(X, U \cup V) = w(X, U) + w(X, V) - w(X, U \cap V)$.

3.6. Given an open-closed pair (U, K) , one can define quadratic cohomology on **k -forms**. Forms are functions on $\Lambda(X, Y) = \{(x, y) | x \in X, y \in Y, x \cap y \in X\}$ and $\Lambda(X) = \{(x, y) | x \in X, y \in X, x \cap y \in X\}$ and $\Lambda(X, Y) = \{(x, y) | x \in X, y \in Y, x \cap y \neq \emptyset, x \cap y \in K\}$. The k -forms are the forms on functions with $\dim(x) + \dim(y) = k$.

³We write the entries as $L(x, y)$

3.7. In the case of an open-closed pair, we have five different cohomologies $U, K, (U, K), (K, U), (U, U)$. There is no case (K, K) because the intersection of $x \in K, y \in K$ is in K . The case (K, K) is part of K . The case (U, U) looks at pairs such that the intersection is in K while U looks at pairs such that the intersection is in K . We can have a disjoint union

$$\Lambda(G) = \Lambda(U) \cup \Lambda(K) \cup \Lambda(K, U) \cup \Lambda(U, K) \cup \Lambda(U, U) .$$

4

3.8. The **exterior derivative** is inherited from the exterior derivative on products. It is $df(x, y) = d_x f(x, y) + w(x) d_y(f, y)$, where d_x, d_y are the usual simplicial exterior derivatives but with respect to the first or second coordinate. If we would look at this derivative on $X \times Y$, the Hodge Laplacians are the tensor products of the Laplacians on X and Y . Even if the set-theoretical Cartesian product $X \times Y$ is not a simplicial complex any more, we still have a cohomology. But now, we restrict this exterior derivative to pairs (x, y) that intersect. We are not aware of such a construction in the continuum.

4. SPECTRAL MONOTONICITY

4.1. The proof of the quadratic fusion inequality Theorem (2) is analog to the linear case. The key is that in each case, the matrix L is the square $L = D^2$ of a matrix D which has the property that a principal sub-matrix of D has intertwined spectrum so that the left padded spectral functions of L are monotone. This looks like a technical detail but it is important and **at the heart of the entire story**: the matrix L does not have the property that taking away highest or lowest dimensional simplices produces principal sub-matrices which themselves come from a geometry. But the Dirac matrix D does have the property. And since D has symmetric spectrum with respect to the origin and $D^2 = L$, we have also monotonicity for L .

4.2. Let us formulate the Cauchy interlace theorem a bit differently, than usual. The point is that if a principal submatrix B of a self-adjoint matrix A has the eigenvalues padded left when compared to the eigenvalues of A then there is a direct comparison between all eigenvalues. This is very general and allows to talk about monotonicity rather than interlacing.

Lemma 1 (Left Padded Monotonicity). *Let A be a symmetric $n \times n$ matrix and m a principal $m \times m$ submatrix, denote by $\lambda_1 \leq \dots \leq \lambda_n$ the eigenvalues of A and $\mu_{n-m} \leq \dots \leq \mu_n$ the eigenvalues of B . Then $\mu_k \leq \lambda_k$ for all $n - m \leq k \leq n$.*

Proof. This follows directly from the interlace theorem and induction with respect to m . Both induction assumption as well as the induction steps involve the interlace theorem. \square

```
n=300; m=100; B=Table[20*Random[]-10,{n},{n}]; A0=Transpose[B].B;
A=A0; Do[A1=Transpose[Delete[Transpose[Delete[A,1]],1]]; A=A1,{m}];
A=A1; Do[A2=Transpose[Delete[Transpose[Delete[A,1]],1]]; A=A2,{m}];
T=Eigenvalues; S=Sort; {a,b,c}=PadLeft[{S[T[A0]],S[T[A1]],S[T[A2]]}];
ListPlot[{a,b,c},Joined->True,Filling->Bottom,PlotRange->All];
```

⁴In the code we part we identify $\Lambda(U, K)$ with $\Lambda(K, U)$ so that we do not have to probe which of the entries is the closed set.

FUSION INEQUALITY




FIGURE 1. We see the sorted eigenvalues of a random real self-adjoint 300×300 matrix A_0 , then the eigenvalues of a 200×200 principal submatrix A_1 and then the eigenvalues of a 100×100 principal submatrix A_2 of A_1 . The eigenvalues are padded left. The figure illustrates Lemma (1). The code which gave the output is listed below.

4.3. We can now look at the map $K \rightarrow \lambda(K)$ giving for each sub-complex K the spectral function ordered in an ascending way and padded left. The partial order on sub-simplicial complexes and the partial order on spectral functions are compatible:

Corollary 1. *The maps $X \rightarrow \lambda(X)$ and $(X, Y) \rightarrow \lambda(X, Y)$ preserve the partial orders in the sense that if we remove maximal simplices from closed sets, or minimal simplicial from open sets, then the spectral functions can only get smaller.*

4.4. The same holds by iterating the process and take principal $(n - k) \times (n - k)$ sub-matrices. Now, if we look at a Dirac matrix of a closed set K and take a maximal simplex x away, then we get monotonicity. The same happens if we take a minimal simplex x away from an open set. Note that if we look at pairs (x, y) belonging to some pair like (K, U) and we take a maximal element x away, then several pairs (x, y_i) are removed from the complex on (K, U) .

Theorem 5 (Spectral monotonicity). *For all $j \leq n$ we have $\lambda_j(K) \leq \lambda_j(G)$,*

$$\begin{aligned} \lambda_j(K) &\leq \lambda_j(G), \\ \lambda_j(K, U) &\leq \lambda_j(G), \\ \lambda_j(U, K) &\leq \lambda_j(G), \\ \lambda_j(U, U) &\leq \lambda_j(G), \end{aligned}$$

Proof. If we add a locally maximal simplex to a given complex, the spectrum changes monotonically by interlace. For any vector u , $\langle u, Lu \rangle = \langle u, D^2u \rangle = \langle Du, Du \rangle = \|Du\|^2$ Define $\mathcal{S}_k = \{V \subset \mathbb{R}^n, \dim(V) = k\}$

$$\lambda_k(K) = \min_{V \in \mathcal{S}_k} \max_{|u|=1, u \in V} \langle u, L(K)u \rangle \leq \min_{V \in \mathcal{S}_k} \max_{|u|=1, u \in V} \langle u, L(G)u \rangle = \lambda_k(G) .$$

As for the **interlace theorem** applied to D as the Dirac matrix of K is obtained from the Dirac matrix of L by deleting the row and column belonging to the element x which was added. The eigenvalues of the Dirac matrix D_K of K are now interlacing the eigenvalues of the Dirac matrix D_G of G . \square

4.5. In the quadratic case, taking away a largest dimensional simplex (facet) x will affect in general various pairs of simplices (x, y) or (y, x) . The effect is that the quadratic Dirac matrix of $G \setminus x$ is still a principal sub-matrix. We still have spectral monotonicity.

4.6. To conclude the proof of Theorem (2), write down the decoupled Laplacian $L(U) \oplus L(K) \oplus L(K, U) \oplus L(U, K) \oplus L(U, U)$ which is block diagonal and is a $n \times n$ matrix, the same size than $L(G)$. Lets call its eigenvalues μ_k . From the spectral inequalities for each block, we know

$$0 \leq \mu_k \leq 5\lambda_k .$$

where λ_k are the eigenvalues of the quadratic Hodge Laplacian of G . Therefore, there are at least as many 0 eigenvalues for the decoupled system than for G , proving the inequality.

5. AN EXAMPLE

5.1. Here is an example with the **Kite complex** $G = K_{1,2,1}$, where we have a complex with 2 triangles. We will see what happens if we take one of the triangles away. We look at the case (U, U) . The Dirac matrix $D(U, U)$ is a 14×14 matrix.

$$\begin{bmatrix} 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & -1 & 0 & -1 & 0 & 0 & 0 \end{bmatrix} .$$

The Eigenvalues of the Laplacian $L(U, U) = D(U, U)^2$ are $\{4, 4, 4, 4, 2, 2, 2, 2, 2, 2, 2, 2, 0, 0\}$.

5.2. Now lets take away the simplicies which do not involve the triangle $(1, 3, 4)$. We have to select the rows and columns in $\{1, 2, 3, 4, 7, 8, 9, 11\}$. The Dirac matrix is

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} .$$

The eigenvalues of L are now $\{1, 1, 1, 1, 1, 1, 1, 1\}$.

6. CODE

```

Generate[A_]:=If[A=={},{},Sort[Delete[Union[Sort[Flatten[Map[Subsets,A],1]]],1]]];
L=Length; Whitney[s_]:=Generate[FindClique[s,Infinity,All]]; L2[x_]:=L[x[[1]]]+L[x[[2]]];
(* Linear Cohomology *)
sig[x_]:=Signature[x]; nu[A_]:=If[A=={},{},0,L[A]-MatrixRank[A]];
F[G_]:=Module[{l=Map[L,G]},If[G=={},{},Table[Sum[If[l[[j]]==k,1,0],{j,L[l]}],{k,Max[l]}]];
sig[x_,y_]:=If[SubsetQ[x,y]&&(L[x]==L[y]+1),sig[Prepend[y,Complement[x,y][[1]]]*sig[x],0];
Dirac[G_]:=Module[{f=F[G],b,d,n=L[G]},b=Prepend[Table[Sum[f[[1]],{1,k}],{k,L[f]}],0];
d=Table[sig[G[[i]],G[[j]],{i,n},{j,n}];{d+Transpose[d],b}];
Hodge[G_]:=Module[{Q,b,H},{Q,b}=Dirac[G];H=Q.Q;Table[Table[H[[b[[k]]+i,b[[k]]+j]],{i,b[[k+1]]-b[[k]]},{j,b[[k+1]]-b[[k]]}],{k,L[b]-1}];
Betti[s_]:=Module[{G},If[GraphQ[s],G=Whitney[s],G=s];Map[nu,Hodge[G]];
Fvector[A_]:=Delete[BinCounts[Map[Length,A],1];
Euler[A_]:=Sum[(-1)^(Length[A[[k]]]-1),{k,Length[A]}];
(* Quadratic Cohomology *)
F2[G_]:=Module[{},If[G=={},{},Table[Sum[If[L2[G[[j]]]==k,1,0],{j,L[G]}],{k,Max[Map[L2,G]}]];
ev[L_]:=Sort[Eigenvalues[1.0*L]];
WuComplex[A_,B_,opts_]:=Module[{Q={x,y,u},
Do[x=A[[k]];y=B[[1]];u=Intersection[x,y];
If[{(opts=="Open" && Not[x==y] && L[u]>0 && Not[MemberQ[A,u]]) ||
(Not[opts=="Open"] && MemberQ[A,u])},
Q=Append[Q,{x,y}],{k,L[A]},{1,L[B]}];Sort[Q,L2[#1]<L2[#2] &]];
Dirac[G_,H_,opts_]:=Module[{n=L[G],Q,m=L[H],b,d1,d2,h,v,w,l,DD},Q=WuComplex[G,H,opts];
n2=L[Q];f2=F2[Q];b=Prepend[Table[Sum[f2[[1]],{1,k}],{k,L[f2]}],0];
D1[{x_,y_}]:=Table[Sort[Delete[x,k]],y],{k,L[x]}];
D2[{x_,y_}]:=Table[{x,Sort[Delete[y,k]]},{k,L[y]}];
d1=Table[0,{n2},{n2}];Do[v=D1[Q[[m]]];If[L[v]>0,Do[r=Position[Q,v[[k]]];
If[r!={},d1[[m,r[[1,1]]]]=(-1)^k,{k,L[v]}],{m,n2}];
d2=Table[0,{n2},{n2}];Do[v=D2[Q[[m]]];If[L[v]>0,Do[r=Position[Q,v[[k]]];
If[r!={},d2[[m,r[[1,1]]]]=(-1)^(L[Q[[m,1]]]+k),{k,L[v]}],{m,n2}];
d=d1+d2;DD=d+Transpose[d];{DD,b}];
Beltrami[G_,H_,opts_]:=Module[{Q,P,b},{Q,b}=Dirac[G,H,opts];P=Q.Q;
Hodge[G_,H_,opts_]:=Module[{Q,P,b},{Q,b}=Dirac[G,H,opts];P=Q.Q;
Table[Table[P[[b[[k]]+i,b[[k]]+j]],{i,b[[k+1]]-b[[k]]},{j,b[[k+1]]-b[[k]]}],{k,2,L[b]-1}];
Betti[G_,H_,opts_]:=Map[nu,Hodge[G,H,opts]];
Wu[A_,B_,opts_]:=Sum[x=A[[k]];y=B[[1]];u=Intersection[x,y];
If[{(opts=="Open" && Not[x==y] && L[u]>0 && Not[MemberQ[A,u]]) ||
(Not[opts=="Open"] && MemberQ[A,u])},
(-1)^L2[x,y],0],{k,L[A]},{1,L[B]}];
Fvector[A_,B_,opts_]:=Module[{a=F2[WuComplex[A,B,opts]]},Table[a[[k]],{k,2,L[a]}];

s = CompleteGraph[{1,2,1}]; G = Whitney[s]; K = Generate[{{1,4}}]; U=Complement[G,K];
Print["Linear - Cohomology"];
{bU,bK,bG}=PadRight[{Betti[U],Betti[K],Betti[G]}];
{fU,fK,fG}=PadRight[{Fvector[U],Fvector[K],Fvector[G]}];
Print[Grid[{
{"Case","Betti","F-vector","Euler"},{"U",bU,fU,Euler[U]},
{"K",bK,fK,Euler[K]},{"G",bG,fG,Euler[G]},
{"Compare",bU+bK-bG,fU+fK-fG,Euler[U]+Euler[K]-Euler[G]}]];
Print["Quadratic - Cohomology"];
{bU,bK,bKU,bUK,bUU,bG}=PadRight[{Betti[U,U,"Closed"],Betti[K,K,"Closed"],
Betti[K,U,"Closed"],Betti[U,K,"Closed"],Betti[U,U,"Open"],Betti[G,G,"Closed"]}];
{fU,fK,fKU,fUK,fUU,fG}=PadRight[{Fvector[U,U,"Closed"],Fvector[K,K,"Closed"],
Fvector[K,U,"Closed"],Fvector[U,K,"Closed"],Fvector[U,U,"Open"],Fvector[G,G,"Closed"]}];
Print[Grid[{{"Case","Betti","F-vector","Wu"},{"U",bU,fU,Wu[U,U,"Closed"]},
{"K",bK,fK,Wu[K,K,"Closed"]},{"UK",bKU,fKU,Wu[K,U,"Closed"]},{"KU",bKU,fKU,Wu[K,U,"Closed"]},
{"UU",bUU,fUU,Wu[U,U,"Open"]},{"G",bG,fG,Wu[G,G,"Closed"]},
{"Compare",bU+bK+bKU+bUK+bUU-bG,fU+fK+fKU+fUK+fUU-fG,
Wu[U,U,"Closed"]+Wu[K,K,"Closed"]+2Wu[K,U,"Closed"]+Wu[U,U,"Open"]-Wu[G,G,"Closed"]}]];

```

6.1. Here is the output of the above lines for simplicial cohomology

Case	Betti	F-vector	Euler
U	{0, 0, 0}	{2, 4, 2}	0
K	{1, 0, 0}	{2, 1, 0}	1
G	{1, 0, 0}	{4, 5, 2}	1
Compare	{0, 0, 0}	{0, 0, 0}	0

And here the output table for the quadratic cohomology part:

Case	Betti	F-vector	Wu
U	{0, 0, 0, 0, 0}	{2, 8, 12, 8, 2}	0
K	{0, 1, 0, 0, 0}	{2, 4, 1, 0, 0}	-1
UK	{0, 0, 2, 0, 0}	{0, 4, 8, 2, 0}	2
KU	{0, 0, 2, 0, 0}	{0, 4, 8, 2, 0}	2
UU	{0, 0, 0, 2, 0}	{0, 0, 4, 8, 2}	-2
G	{0, 0, 1, 0, 0}	{4, 20, 33, 20, 4}	1
Compare	{0, 1, 3, 2, 0}	{0, 0, 0, 0, 0}	0




FIGURE 2. The Dirac matrix D and the Hodge Laplacian $L = D^2$ in the linear case for the kite graph G . The splittings are given by the f-vector $f(G) = (4, 5, 2)$. There are 4 points, 5 edges and 2 triangles in G .




FIGURE 3. The Dirac matrix D and the Hodge Laplacian $L = D^2$ in the quadratic case for the kite graph. The splittings are given by the f-vector $f(G) = (4, 20, 33, 20, 4)$. The space of 1-forms (intersecting points) is 4-dimensional, the space of 2-forms (intersection of a point with an edge) is 20-dimensional, the space of 3-forms (intersection of two edges or a triangle-point) has dimension 33, the space of 4-forms (intersection of an edge and triangle) is 20-dimensional, the space of 5-forms (intersection of two triangles) is 4-dimensional.

REFERENCES

- [1] P. Alexandroff. Diskrete Räume. *Mat. Sb.* 2, 2, 1937.
- [2] B. Eckmann. Harmonische Funktionen und Randwertaufgaben in einem Komplex. *Comment. Math. Helv.*, 17(1):240–255, 1944.
- [3] B. Grünbaum. Polytopes, graphs, and complexes. *Bull. Amer. Math. Soc.*, 76:1131–1201, 1970.
- [4] O. Knill. An integrable evolution equation in geometry. <http://arxiv.org/abs/1306.0060>, 2013.
- [5] O. Knill. Isospectral deformations of the Dirac operator. <http://arxiv.org/abs/1306.5597>, 2013.
- [6] O. Knill. Gauss-Bonnet for multi-linear valuations. <http://arxiv.org/abs/1601.04533>, 2016.
- [7] O. Knill. The cohomology for Wu characteristics. <http://arxiv.org/abs/1803.06788>, 2017.
- [8] O. Knill. The cohomology for Wu characteristics. <https://arxiv.org/abs/1803.1803.067884>, 2018.
- [9] O. Knill. Characteristic topological invariants. <https://arxiv.org/abs/2302.02510>, 2023.
- [10] O. Knill. Cohomology of Open sets. <https://arxiv.org/abs/2305.12613>, 2023.
- [11] O. Knill. Discrete algebraic sets in discrete manifolds. <https://arxiv.org/abs/2312.14671>, 2023.
- [12] O. Knill. Finite topologies for finite geometries. <https://arxiv.org/abs/2301.03156>, 2023.
- [13] O. Knill. Spectral monotonicity of the Hodge Laplacian. <https://arxiv.org/abs/2304.00901>, 2023.
- [14] Wu W-T. Topologie combinatoire et invariants combinatoires. *Colloq. Math.*, 7:1–8, 1959.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA, 02138