

# Quantum correlation of Hawking radiations for mass-superposed BTZ black holes

Hongbin Zhang and Baocheng Zhang\*

*School of Mathematics and Physics,*

*China University of Geosciences, Wuhan 430074, China*

## Abstract

Recent studies using the Unruh-DeWitt detector model have revealed the quantum signatures of black hole mass superposition. In this paper, we use the equal-time density-density correlation function to reinvestigate the mass superposition of BTZ black holes. Interestingly, the quantum signatures of BTZ black hole mass superposition can be revealed by the equal-time density-density correlation peaks. Additionally, different mass superpositions for the same mass ratio can be distinguished, a phenomenon not previously documented. Specifically, the correlation function method can pinpoint the location where Hawking radiations are generated, even for mass-superposed BTZ black holes. Our findings indicate that this location is approximately at two and a half times the radius of the horizon, where the correlation function reaches its maximum. This supports the concept of a quantum atmosphere as the origin locus of Hawking radiation, even in quantum-superposed spacetimes.

Key words: quantum correlations, black holes, mass superposition, Wightman function.

---

\*Electronic address: [zhangbaocheng@cug.edu.cn](mailto:zhangbaocheng@cug.edu.cn)

## Contents

<b>I. Introduction</b>	3
<b>II. Wightman functions</b>	4
<b>III. Density-density correlation functions</b>	7
<b>IV. Quantum signatures</b>	9
A. Obtaining quantum signals through density-density correlation functions	9
B. Distinguishing quantum signals with same mass ratio	11
<b>V. Creation and evolution of Hawking radiations</b>	13
A. Correlation across the horizon	13
B. Time evolution of correlation	14
<b>VI. Conclusions</b>	16
<b>Acknowledgments</b>	17
<b>References</b>	17

## I. INTRODUCTION

Einstein's general relativity predicts that a sufficiently compact mass can deform spacetime to form a black hole [1]. Black holes provide an arena where interesting effects associated with gravity, both classical and quantum, can manifest. The boundary beyond which nothing can escape is called the event horizon. Quantum field theory in curved spacetime [2] predicts that black holes are not truly black, as they emit thermal radiation known as Hawking radiation [3]. Pairs of correlated particles are created near the black hole event horizon; one particle escapes to infinity as Hawking radiation, while the other is swallowed by the singularity inside the black hole horizon, known as the partner particle. Hawking's discovery is considered a milestone of modern theoretical physics [4].

The exact location where Hawking radiation originates is an interesting question currently being researched. Unruh first discussed the origin of Hawking radiation from gravitational black holes [5]. Subsequent studies analyzed analog black holes [6] and used two-point correlation functions to reanalyze gravitational black holes [7]. Later, Giddings introduced the concept of a quantum atmosphere, displaced from the horizon, as the origin of Hawking radiation [8]. Understanding the origin of Hawking quanta is crucial, as it can address the information loss problem [9–12]. Consequently, research on the origin of Hawking radiation has been extended to various scenarios [13–19].

Recently, the phenomenon of black hole mass superposition has been studied in the context of spacetime superpositions [20, 21]. Examples include quantum superpositions of Minkowski spacetime [22] and the signatures of rotating black holes in quantum superpositions [23]. Researchers are also attempting to apply these concepts to analog black holes [24, 25], building on earlier work [26–29], and testing quantum effects in gravity through tabletop experiments [30–32]. Black hole mass superpositions provide a novel approach to studying quantum black holes and advancing the theory of quantum gravity.

Black holes in mass superpositions also exhibit Hawking radiation, measurable by the Unruh-DeWitt detector [33]. This detector reveals signatures of quantum-gravitational effects when black hole mass superpositions are formed with specific mass ratios [34]. The quantum signatures of black hole mass superpositions support Bekenstein's conjecture [35, 36] that a complete theory of quantum gravity should treat black holes as quantum objects. According to Bekenstein, the black hole horizon area and mass are adiabatic in-

variants with an associated discrete quantization [37]. Bekenstein and Mukhanov proposed that mass quantization should alter the nature of Hawking radiation, resulting in a discrete emission spectrum with evenly spaced energy levels [38]. This idea has drawn significant attention to the study of quantum black holes [39–44]. The quantization of black hole mass implies that black holes can also exist in superpositions. The possibility of black hole mass superpositions has been explored within the framework of loop quantum gravity [45–49].

Since mass-superposed black holes can emit Hawking radiation, investigating the origin of this radiation is crucial. Correlation functions are a valuable tool for studying the origin of Hawking radiation. They have been used to analyze the occurrence position and timing of Hawking radiation in analog black holes [50–54], despite the very low Hawking radiation temperature in such systems. Density-density correlation has become a key piece of evidence for measuring Hawking radiation from analog black holes in laboratory settings [55, 56]. This method has also been applied to determine the origin of Hawking radiation from gravitational black holes [57–62]. In this paper, we will study the density-density correlation functions for mass-superposed BTZ black holes to gain more insight into mass superposition and the original location of Hawking radiation.

This paper is organized as follows. In the second section, we introduce the BTZ black hole metrics and analyze the Wightman functions for both classical BTZ black holes and mass-superposed BTZ black holes. In the third section, based on the Wightman functions, we construct the density-density correlation function for mass-superposed BTZ black holes. Using the density-density correlation function outside the event horizon of mass-superposed BTZ black holes, we demonstrate the quantum signatures of black hole mass superpositions in the fourth section. In the fifth section, we investigate the origin of Hawking radiation using the density-density correlation across the horizon of both classical and mass-superposed BTZ black holes. Finally, we present our conclusions in the sixth section.

## II. WIGHTMAN FUNCTIONS

In (2+1)-dimensions spacetime, Bañados, Teitelboim, and Zanelli found a solution of the Einstein gravitational field equation called BTZ black hole [63]. Geometrically, (2+1)-dimensions AdS space can be obtained from flat space  $\mathbb{R}^{2,2}$ . AdS<sub>3</sub> space is a submanifold

[64]

$$-l^2 = -T_1^2 - T_2^2 + X_1^2 + X_2^2, \quad (1)$$

in flat space  $\mathbb{R}^{2,2}$

$$ds^2 = -dT_1^2 - dT_2^2 + dX_1^2 + dX_2^2. \quad (2)$$

The above embedding coordinates  $(X_1, X_2, T_1, T_2)$  can be chosen as [65, 66]

$$\begin{aligned} T_1 &= \tilde{r} \cosh \tilde{\phi}, \\ X_1 &= \tilde{r} \sinh \tilde{\phi}, \\ T_2 &= \sqrt{\tilde{r}^2 - l^2} \sinh \frac{\tilde{t}}{l}, \\ X_2 &= \sqrt{\tilde{r}^2 - l^2} \cosh \frac{\tilde{t}}{l}, \end{aligned} \quad (3)$$

and then we get the AdS-Rindler metric

$$ds^2 = -\left(\frac{\tilde{r}^2}{l^2} - 1\right)d\tilde{t}^2 + \frac{1}{\left(\frac{\tilde{r}^2}{l^2} - 1\right)}d\tilde{r}^2 + \tilde{r}^2 d\tilde{\phi}^2. \quad (4)$$

The BTZ metric can be derived by the new coordinates  $r = \tilde{r}\sqrt{M}$ ,  $t = \tilde{t}/\sqrt{M}$  and  $\phi = \tilde{\phi}/\sqrt{M}$  [65],

$$ds^2 = -\left(\frac{r^2}{l^2} - M\right)dt^2 + \frac{1}{\left(\frac{r^2}{l^2} - M\right)}dr^2 + r^2 d\phi^2, \quad (5)$$

where  $M$  is the mass of BTZ black hole,  $l$  is the AdS length scale in an asymptotically AdS vacuum, and  $\Lambda = -1/l^2$  is the cosmological constant. In particular, the event horizon is located at  $r_h = \sqrt{M}l$ . Here, we emphasize that in this paper, the discussed BTZ black holes are considered without angular momentum for the sake of convenience in discussing mass-superposed black holes. In these scenarios, the horizon undergoes a superposition characterized by  $r_s = \sqrt{M_1}l$  and  $r_b = \sqrt{M_2}l$ , defined by the two mass values  $M_1, M_2$  ( $M_1 < M_2$ ) of the black hole mass superposition.

To obtain the density-density correlation function we have to know the Wightman function of BTZ black hole spacetime. This Wightman function can be constructed from the well-known vacuum  $AdS_3$  Wightman function [64, 67]

$$W_{AdS}(x, x') = \frac{1}{4\pi l\sqrt{2}} \left( \frac{1}{\sqrt{\sigma(x, x')}} - \frac{\zeta}{\sqrt{\sigma(x, x') + 2}} \right), \quad (6)$$

where the parameter  $\zeta$  is the boundary condition at infinity in  $AdS$  spacetime, taking the values  $\{0, 1, -1\}$  corresponding to transparent, Dirichlet, and Neumann boundary conditions, respectively. The function  $\sigma(x, x')$  is the squared geodesic distance between  $x$  and  $x'$  in the flat  $\mathbb{R}^{2,2}$  space given by

$$\sigma(x, x') = \frac{1}{2l^2}[-(T_1 - T_1')^2 - (T_2 - T_2')^2 + (X_1 - X_1')^2 + (X_2 - X_2')^2]. \quad (7)$$

The Wightman functions for the BTZ black hole spacetime can be derived from AdS vacuum by

$$W_{BTZ}(x, x') = \frac{1}{\sum_p \eta^{2p}} \sum_p \sum_q \eta^p \eta^q W_{AdS}(\Gamma^p x, \Gamma^q x'), \quad (8)$$

where  $\Gamma^p$  is the periodic identification of the angle coordinate  $\tilde{\phi} \rightarrow \tilde{\phi} + 2\pi p\sqrt{M}$  and the parameter  $\eta = +1, -1$ . Inserting the  $AdS_3$  embedding coordinates  $(X_1, X_2, T_1, T_2)$  together with the periodic identification into Eq. (7), we gain the squared geodesic distance in the BTZ black hole spacetime as

$$\begin{aligned} \sigma(\Gamma^p x, \Gamma^q x') &= \frac{\tilde{r}_1 \tilde{r}_2}{l^2} \cosh \left[ 2\pi(p - q)\sqrt{M} \right] - 1 \\ &\quad - \sqrt{\frac{\tilde{r}_1^2}{l^2} - 1} \sqrt{\frac{\tilde{r}_2^2}{l^2} - 1} \cosh \frac{\tilde{t}_1 - \tilde{t}_2}{l}, \end{aligned} \quad (9)$$

and the cross term of squared geodesic distance as

$$\begin{aligned} \sigma(\Gamma_1^p x, \Gamma_2^q x') &= \frac{\tilde{r}_1 \tilde{r}_2}{l^2} \cosh \left[ 2\pi(p\sqrt{M_1} - q\sqrt{M_2}) \right] - 1 \\ &\quad - \sqrt{\frac{\tilde{r}_1^2}{l^2} - 1} \sqrt{\frac{\tilde{r}_2^2}{l^2} - 1} \cosh \frac{\tilde{t}_1 - \tilde{t}_2}{l}, \end{aligned} \quad (10)$$

where  $\Gamma_1^p$  and  $\Gamma_2^q$  are two different periodic isometries with two different mass  $M_1$  and  $M_2$  [34]. For simplicity, we study the Wightman functions with transparent boundary condition  $\zeta = 0$ . In the end, we can get the Wightman function for the BTZ black hole spacetime as

$$W_{BTZ}(x, x') = \frac{1}{\sum_p \eta^{2p}} \frac{1}{4\pi l \sqrt{2}} \sum_p \sum_q \frac{1}{\sqrt{\sigma(\Gamma^p x, \Gamma^q x')}}, \quad (11)$$

and the Wightman function of the mass-superposed BTZ black hole spacetime as

$$W_{BTZ}^{(1,2)}(x, x') = \frac{1}{\sum_p \eta^{2p}} \frac{1}{4\pi l \sqrt{2}} \sum_p \sum_q \frac{1}{\sqrt{\sigma(\Gamma_1^p x, \Gamma_2^q x')}}. \quad (12)$$

Now we consider a massless scalar field  $\hat{\psi}$  in (2+1)-dimension AdS spacetime. The Wightman function of the BTZ black hole with mass  $M$  in Eq. (11) can be rewritten as [68]

$$W_{BTZ}(x, x') = \frac{1}{\mathcal{N}} \sum_{p,q} \eta^p \eta^q \langle 0 | \hat{\psi}(\Gamma^p x) \hat{\psi}(\Gamma^q x') | 0 \rangle, \quad (13)$$

and the Wightman function of the mass-superposed BTZ black hole with superposed mass  $M_1$  and  $M_2$  in Eq. (12) can be rewritten as

$$W_{BTZ}^{(1,2)}(x, x') = \frac{1}{\mathcal{N}} \sum_{p,q} \eta^p \eta^q \langle 0 | \hat{\psi}(\Gamma_1^p x) \hat{\psi}(\Gamma_2^q x') | 0 \rangle, \quad (14)$$

where the parameter  $\eta = +1, -1$  corresponding to the twisted and untwisted fields respectively,  $\mathcal{N} = \sum_p \eta^{2p}$  is normalization factor, and  $|0\rangle$  is the AdS vacuum. The establishment of the connection between the AdS vacuum state and the Whiteman functions will facilitate our construction of the density-density correlation function.

### III. DENSITY-DENSITY CORRELATION FUNCTIONS

In this section, we will utilize Wightman functions to build the density-density correlation functions. One objective of constructing these correlation functions is to gather additional insights into BTZ black holes in mass superpositions. Another aim is to pinpoint where the signal indicating the presence of Hawking radiation emerges in both the classical BTZ black hole and the mass-superposed BTZ black hole.

We choose free-falling observers that are locally inertial observers in the BTZ black hole spacetime. The free-falling observer can measure the energy density [58]

$$\hat{\rho} = \hat{T}_{ab} u^a u^b, \quad (15)$$

where

$$\hat{T}_{ab}(\hat{\phi}) = \partial_a \hat{\phi} \partial_b \hat{\phi} - \frac{g_{ab}}{2} g^{cd} \partial_c \hat{\phi} \partial_d \hat{\phi} \quad (16)$$

is the energy-momentum operator of the massless scalar field  $\hat{\phi}$  that describes an outgoing flux of particles at the Hawking temperature and  $u^a = (1, V)$  is the four-velocity of the free-falling observer. For the convenience to make the physical description from these observers, we transform the BTZ metric in Eq. (5) into *Painlevé* coordinates [69]

$$ds^2 = -f d\tau^2 - 2V d\tau dr + dr^2 + r^2 d\phi^2, \quad (17)$$

where  $f = \frac{r^2}{l^2} - M$  and  $V = -\sqrt{1 - (\frac{r^2}{l^2} - M)}$ . So we gain the vanishing of the trace of  $\hat{T}_{ab}$

$$g^{ab}\hat{T}_{ab} = -\hat{T}_{\tau\tau} - 2V\hat{T}_{\tau r} + f\hat{T}_{rr} = 0, \quad (18)$$

and the energy density

$$\hat{\rho} = \hat{T}_{ab}u^a u^b = \hat{T}_{\tau\tau} + 2V\hat{T}_{\tau r} + V^2\hat{T}_{rr}, \quad (19)$$

when we consider  $\phi$  as a constant. Then we combine Eq. (18) with Eq. (19) to give the energy density as

$$\hat{\rho} = (f + V^2)\hat{T}_{rr} = \hat{T}_{rr}, \quad (20)$$

which is obtained using the *Painlevé* time

$$\tau = t + \int \frac{\sqrt{1 - (r^2/l^2 - M)}}{r^2/l^2 - M} dr = t + r^*. \quad (21)$$

According to massless scalar field in Eq. (16), the energy density can be rewritten as

$$\hat{\rho} = (1 - \frac{f}{2})\partial_r\hat{\phi}\partial_r\hat{\phi} + \frac{1}{2}\partial_\tau\hat{\phi}\partial_\tau\hat{\phi} + V\partial_\tau\hat{\phi}\partial_r\hat{\phi}. \quad (22)$$

Finally, we gain the density-density correlation function from Eq. (22) as

$$\begin{aligned} G(x, x') &= \langle 0|\hat{\rho}(x)\hat{\rho}(x')|0\rangle = \langle 0|[(1 - \frac{f_1}{2})\partial_{r_1}\hat{\phi}\partial_{r_1}\hat{\phi} + \frac{1}{2}\partial_{\tau_1}\hat{\phi}\partial_{\tau_1}\hat{\phi} + V_1\partial_{\tau_1}\hat{\phi}\partial_{r_1}\hat{\phi}] \\ &\quad * [(1 - \frac{f_2}{2})\partial_{r_2}\hat{\phi}\partial_{r_2}\hat{\phi} + \frac{1}{2}\partial_{\tau_2}\hat{\phi}\partial_{\tau_2}\hat{\phi} + V_2\partial_{\tau_2}\hat{\phi}\partial_{r_2}\hat{\phi}]|0\rangle, \end{aligned} \quad (23)$$

where  $f_1 = f(r_1)$ ,  $f_2 = f(r_2)$ , and

$$\hat{\phi} = \frac{1}{\sqrt{\mathcal{N}}} \sum_p \eta^p \hat{\psi}(\Gamma^p x) \quad (24)$$

is an automorphic field of BTZ black hole spacetime [34]. In order to make our paper clear and easy to read, we define four functions that are related to density-density correlation functions by

$$\begin{aligned} G_1 &= \partial_{r_1}\partial_{r_2}\langle 0|\hat{\phi}(x)\hat{\phi}(x')|0\rangle = \partial_{r_1}\partial_{r_2}W_{BTZ}, \\ G_2 &= \partial_{r_1}\partial_{\tau_2}\langle 0|\hat{\phi}(x)\hat{\phi}(x')|0\rangle = \partial_{r_1}\partial_{\tau_2}W_{BTZ}, \\ G_3 &= \partial_{\tau_1}\partial_{r_2}\langle 0|\hat{\phi}(x)\hat{\phi}(x')|0\rangle = \partial_{\tau_1}\partial_{r_2}W_{BTZ}, \\ G_4 &= \partial_{\tau_1}\partial_{\tau_2}\langle 0|\hat{\phi}(x)\hat{\phi}(x')|0\rangle = \partial_{\tau_1}\partial_{\tau_2}W_{BTZ}, \end{aligned} \quad (25)$$



where the Wightman function is just that in Eq. (13) for the classical BTZ black hole or that in Eq. (14) for the mass-superposed BTZ black hole. Finally, we obtain the form of the density-density correlation function as

$$G(x, x') = (1 - \frac{f_1}{2})(1 - \frac{f_2}{2})G_1^2 + \frac{1}{2}(1 - \frac{f_1}{2})G_2^2 + V_2(1 - \frac{f_1}{2})G_1G_2 + \frac{1}{2}(1 - \frac{f_2}{2})G_3^2 + \frac{1}{4}G_4^2 + \frac{V_2}{2}G_3G_4 + V_1(1 - \frac{f_2}{2})G_1G_3 + \frac{V_1}{2}G_2G_4 + V_1V_2G_1G_4. \quad (26)$$

## IV. QUANTUM SIGNATURES

### A. Obtaining quantum signals through density-density correlation functions

We will use the correlation functions obtained in the last section to give the quantum signatures of the black hole mass superposition. Firstly, we express the squared geodesic distance with the radial coordinate  $r = \tilde{r}\sqrt{M}$  and *Painlevé* time  $\tau = t + r^*$  as

$$\sigma(\Gamma^p x, \Gamma^q x') = \frac{r_1 r_2}{M l^2} \cosh \left[ 2\pi(p - q)\sqrt{M} \right] - 1 - \sqrt{\frac{r_1^2}{M l^2} - 1} \sqrt{\frac{r_2^2}{M l^2} - 1} \cosh \frac{\tau_1 - \tau_2 + r_2^* - r_1^*}{l}, \quad (27)$$

and the cross term of squared geodesic distance as

$$\sigma(\Gamma_1^p x, \Gamma_2^q x') = \frac{r_1 r_2}{\sqrt{M_1} \sqrt{M_2} l^2} \cosh \left[ 2\pi(p\sqrt{M_1} - q\sqrt{M_2}) \right] - 1 - \sqrt{\frac{r_1^2}{M_1 l^2} - 1} \sqrt{\frac{r_2^2}{M_2 l^2} - 1} \cosh \frac{\tau_1 - \tau_2 + r_2^* - r_1^*}{l}. \quad (28)$$

Then, we substitute Eq. (27) and Eq. (28) into Eq. (26) to gain the correlation functions of the classical BTZ black hole and the mass-superposed BTZ black hole. In particular, correlations can be calculated from the black hole vacuum outside the event horizon [59, 66, 70]. It is noted that when the superposed mass takes the same value ( $M_1 = M_2 = M$ ) in Eq. (28), the density-density correlation reduces to the same form as in Eq. (27).

Black hole mass superposition induces novel effects that are absent in classical spacetime. As shown in Ref. [34], the detector exhibits resonances at specific mass ratios when interacting with the Hawking radiation from mass-superposed BTZ black holes. These resonances can be understood in light of Bekenstein's conjecture regarding the quantization of black hole mass [35]. Quantum signatures can also be observed through the density-density correlation




FIG. 1: The absolute value of equal-time  $G(x, x')$  outside the horizon of mass-superposed black holes for four different mass pairs:  $(M_1 = 1, M_2 = 9)$ ,  $(M_1 = 1, M_2 = 16)$ ,  $(M_1 = 1, M_2 = 25)$ , and  $(M_1 = 1, M_2 = 36)$ , corresponding to mass ratios,  $M_1/M_2 = 1/n^2$  with  $n = 3, 4, 5, 6$ , respectively. Other parameters are taken as  $r_1 = r_2 = 10$ ,  $l = 1$ ,  $p = q = 20$  and  $\tau_1 = \tau_2$ .

function, as depicted in Fig. 1, which corroborates findings in Ref. [34] for the mass ratio  $M_1/M_2 = 1/n^2$ . Furthermore, it is evident that the value of the equal-time density-density correlation peak gradually decreases with increasing  $n$  in the mass ratio. This indicates that larger differences between the two mass values in the black hole mass superposition lead to reduced density-density correlation peaks.

Regarding the resonance phenomenon observed at the mass ratio  $M_1/M_2 = (n - 1)^2/n^2$  as reported in Ref. [34], we replicate these findings through calculations of the equal-time density-density correlation, illustrated in Fig. 2. In contrast to the scenario depicted in Fig. 1 at the mass ratio  $M_1/M_2 = 1/n^2$ , we observe that the value of the equal-time density-density correlation peak increases gradually with higher values of  $n$  in the mass ratio.

Comparing Fig. 1 with Fig. 2, it becomes apparent that the equal-time density-density correlation value for the mass ratio  $M_1/M_2 = (n - 1)^2/n^2$  is greater than that for  $M_1/M_2 = 1/n^2$  when other parameters remain constant. This difference arises because the disparity between the two mass values in the former mass ratio is smaller than in the latter. Specifically, for each mass ratio, the correlation peak values for black hole mass superpositions increase as the two mass values become closer to each other.




FIG. 2: The absolute value of equal-time  $G(x, x')$  outside the horizon of mass-superposed black holes for four different mass pairs:  $(M_1 = 1, M_2 = 9/4)$ ,  $(M_1 = 1, M_2 = 16/9)$ ,  $(M_1 = 1, M_2 = 25/16)$ , and  $(M_1 = 1, M_2 = 36/25)$ , corresponding to mass ratios  $M_1/M_2 = (n - 1)^2/n^2$  with  $n = 3, 4, 5, 6$ , respectively. Other parameters are kept consistent with those in Fig. (1).

The equal-time density-density correlation also exhibits peaks at certain mass ratios that lie between the special values  $1/n^2$  or  $(n - 1)^2/n^2$ , but these peaks are smaller than those observed at these specific mass ratios. Furthermore, the values of  $p$  and  $q$  in Eqs. (27) and (28) affect the occurrence of these correlation peaks. Higher values of  $p$  and  $q$  result in more peaks in the density-density correlation for black hole mass superpositions. In particular, when the radial coordinates  $r_1 = r_2$ , the equal-time density-density correlation peaks diminish as the radial coordinate moves further away from the horizon  $r_b = \sqrt{Ml}$ , defined by the larger mass of the two black hole mass superpositions.

### B. Distinguishing quantum signals with same mass ratio

It has been observed that resonant peaks occur for  $M_1 < M_2$  or  $M_1 > M_2$  at specific mass ratios, but distinguishing between different mass pairs with the same mass ratio, such as  $(M_1 = 0.5, M_2 = 2)$ ,  $(M_1 = 1, M_2 = 4)$  corresponding to  $M_1/M_2 = 1/n^2$  with  $n = 2$ , has not been clear using earlier methods [34]. However, the density-density correlation method can achieve this distinction.

In Table I, we compute and present four absolute values of  $G(x, x')$  for superposed mass

TABLE I: The absolute value of  $G(x, x')$  outside the horizon of mass-superposed black holes for four different mass pairs:  $(M_1 = 0.5, M_2 = 2)$ ,  $(M_1 = 1, M_2 = 4)$ ,  $(M_1 = 4, M_2 = 16)$ , and  $(M_1 = 4, M_2 = 1)$ . Other parameters are kept consistent with those in Fig. (1).

Mass ratios $(M_1, M_2)$	4, 16	1, 4	4, 1	0.5, 2
$G(x, x')(10^{-6})$	6.048	1.163	1.163	0.480




FIG. 3: The absolute value of equal-time  $G(x, x')$  outside the horizon of mass-superposed black holes for two different mass pairs:  $(M_1 = 1, M_2 = 4)$  (a),  $(M_1 = 4, M_2 = 1)$  (b).

Other parameters are kept consistent with those in Fig. (1).

values:  $(M_1 = 4, M_2 = 16)$ ,  $(M_1 = 1, M_2 = 4)$ ,  $(M_1 = 4, M_2 = 1)$ , and  $(M_1 = 0.5, M_2 = 2)$ . From this table, it is evident that different absolute values are obtained for the density-density correlation even at the same mass ratios in the first, second, and fourth rows. Furthermore, these absolute values increase as the two mass values of the black hole mass superpositions increase.

Moreover, it is noted from the table that the absolute value of  $G(x, x')$  for the mass values  $(M_1 = 1, M_2 = 4)$  is identical to that for  $(M_1 = 4, M_2 = 1)$ . This indicates that these configurations are indistinguishable, demonstrating an exchange symmetry in the density-density correlation functions for the mass-superposed parameters  $M_1$  and  $M_2$ , as illustrated




FIG. 4: Three-dimensional plot of equal-time  $G(x, x')$  across the horizon of the classical BTZ black hole (a) and the mass-superposed BTZ black hole (b). The parameters are taken as  $M = 4$  for the classical BTZ black hole (a) and  $M_1 = 1, M_2 = 4$  for the mass-superposed BTZ black hole (b). Other parameters are taken as  $r_1 = r_2 = 10, l = 1, p = q = 10$  and  $\tau_1 = \tau_2 = 1$ .

in Fig. 3. Specifically, it is seen from Eq. (28) that this symmetry arises when the parameters  $p$  and  $q$  are set to the same value in our calculations.

## V. CREATION AND EVOLUTION OF HAWKING RADIATIONS

### A. Correlation across the horizon

The above analysis indicates that the mass ratio affects the density-density correlation outside the horizon of mass-superposed black holes. In this section, we will study the origin of Hawking radiation through the density-density correlation, as that made for the origin of Hawking radiation from quantum atmosphere [57–61].

To determine the origin of Hawking particles from the quantum atmosphere in both classical BTZ black holes and mass-superposed BTZ black holes, it is essential to examine the density-density correlation function across the horizon [50–54]. This requires rescaling

the time coordinate to  $\tilde{t} = t\sqrt{M}$  and deriving the new squared geodesic distance as

$$\begin{aligned} \sigma(\Gamma^p x, \Gamma^q x') &= \frac{r_1 r_2}{M l^2} \cosh \left[ 2\pi(p - q)\sqrt{M} \right] \\ &- 1 - \sqrt{\frac{r_1^2}{M l^2} - 1} \sqrt{\frac{r_2^2}{M l^2} - 1} \cosh \sqrt{M} \left( \frac{\tau_1 - \tau_2 + r_2^* - r_1^*}{l} \right) \end{aligned} \quad (29)$$

and the new cross term of squared geodesic distance as

$$\begin{aligned} \sigma(\Gamma_1^p x, \Gamma_2^q x') &= \frac{r_1 r_2}{\sqrt{M_1} \sqrt{M_2} l^2} \cosh \left[ 2\pi(p\sqrt{M_1} - q\sqrt{M_2}) \right] \\ &- 1 - \sqrt{\frac{r_1^2}{M_1 l^2} - 1} \sqrt{\frac{r_2^2}{M_2 l^2} - 1} \cosh \frac{\sqrt{M_1} \tau_1 - \sqrt{M_2} \tau_2 + \sqrt{M_2} r_2^* - \sqrt{M_1} r_1^*}{l}. \end{aligned} \quad (30)$$

Then, we substitute Eq. (29) and Eq. (30) into Eq. (26) to get the new density-density correlation function  $G_1(x, x')$  of classical BTZ black holes and mass-superposed BTZ black holes.

In Fig. 4, we depict the density-density correlation functions  $G(x, x')$  across the event horizon of both the classical BTZ black hole and the mass-superposed BTZ black hole. It is evident that a distinct peak appears approximately at a position 2 to 3 times the distance of the event horizon  $r_h = \sqrt{M}l = 2$  for the classical BTZ black hole, and at about 2.5 times the distance of the horizon  $r_b = \sqrt{M}l = 2$ , defined by the larger mass of the two mass values in the black hole mass superposition.

## B. Time evolution of correlation

For classical gravitational black holes, including the BTZ black hole, the equal-time density-density correlation cannot depict time evolution since the time term vanishes at equal times  $\tau_1 = \tau_2$ , as shown in Eq. (27). However, the time term remains present in the density-density correlation for mass-superposed BTZ black holes, as illustrated in Eq. (30). Therefore, we can analyze the evolution of Hawking radiation through the correlation function in mass-superposed BTZ black holes.

In Fig. 5, we compute and present the absolute values of the density-density correlation at four increasing times  $\tau_1 = 5.2$ ,  $\tau_2 = 5.9$ ,  $\tau_3 = 6.4$ , and  $\tau_4 = 9$ . The peak at  $r_s = r_1 = 1$  represents the horizon location defined by the smaller mass of the two mass values in the black hole mass superposition. For correlations where one point is located less than  $r_s$  and the other is located beyond the horizon  $r_b$  defined by the larger mass of the two mass values,




FIG. 5: The absolute value of  $G_1(x, x')$  across the horizon of mass-superposed BTZ black holes at four increasing times:  $\tau_1 = 5.2$  (a),  $\tau_2 = 5.9$  (b),  $\tau_3 = 6.4$  (c), and  $\tau_4 = 9$  (d), with the superposed mass set as  $(M_1 = 1, M_2 = 4)$ . Other parameters are kept consistent with those in Fig. (4).

the density-density correlation peak initially increases to a maximum and then decreases. Moreover, when one point is between  $r_s$  and  $r_b$ , and the other is beyond  $r_b$ , the correlation peaks increase with time. These changes in peak values manifest the evolution of particle creation in the spacetime of mass-superposed BTZ black holes.



Particularly noteworthy is that the density-density correlation peak moves closer to the horizon  $r_s = \sqrt{M}l = 1$  as time progresses. This observation indicates that partner particles of Hawking radiation in mass-superposed BTZ black holes do not fall into the singularity but accumulate near the horizon  $r_s$  over time. This contrasts with findings for gravitational Schwarzschild black holes in Ref. [58].

## VI. CONCLUSIONS

In this paper, we study the equal-time density-density correlation function in the context of classical BTZ black holes and mass-superposed BTZ black holes. We find the quantum signatures of BTZ black hole mass superposition at specific mass ratios,  $\sqrt{M_1/M_2} = 1/n$  and  $\sqrt{M_1/M_2} = (n-1)/n$ , by observing the peaks in the equal-time density-density correlation function outside the event horizon. These peaks diminish when the mass values in the black hole mass superpositions are significantly different. However, the peak values increase as the mass values of the BTZ black hole mass superposition increase. Notably, mass-superposed black holes with the same mass ratio but different mass values can be distinguished by the values of the density-density correlation peaks, which has not been discussed before.

We also investigate the location and evolution of Hawking radiation for mass-superposed BTZ black holes using the equal-time density-density correlation function. Since the location of the peaks in the correlation function represents the creation of Hawking particle pairs, we find that the characteristic peak appears at approximately 2 to 2.5 times the distance of the event horizon  $r_h = \sqrt{M}l = 2$  for classical BTZ black holes, and at about 2.5 times the horizon distance  $r_b = \sqrt{M}l = 2$  for mass-superposed BTZ black holes. As time progresses, the peaks in the density-density correlation of the mass-superposed BTZ black holes gradually increase to a maximum and then decrease. Simultaneously, the peaks move toward the horizon  $r_s$  over time. This movement indicates the evolution of the creation of Hawking particle pairs to some extent.

A clearer description requires further understanding of the superposed horizon in mass-superposed black holes, but it is possible to extend the present research to superpositions of gravitationally collapsed BTZ black holes [71], exotic BTZ black holes [72], or the analog gravity simulation of spacetime superpositions [24, 25].



## Acknowledgments

This work is supported by National Natural Science Foundation of China (NSFC) with Grant No. 12375057, and the Fundamental Research Funds for the Central Universities, China University of Geosciences (Wuhan).

---

- [1] R. M. Wald, Gravitational collapse and cosmic censorship, *Black holes, gravitational radiation and the universe: essays in honor of CV Vishveshwara*, Springer Netherlands (1999) pages 69–86.
- [2] N. D. Birrell and P. Davies, Quantum fields in curved space, *Cambridge University Press, Cambridge, England (1984)*.
- [3] S. W. Hawking, Black hole explosions?, *Nature* **248** (1974) 30.
- [4] S. W. Hawking, Particle creation by black holes, *Commun. Math. Phys.* **43** (1975) 199.
- [5] W. Unruh, Origin of the particles in black-hole evaporation, *Phys. Rev. D* **15** (1977) 365 .
- [6] R. Schützhold and W. Unruh, Origin of the particles in black hole evaporation, *Phys. Rev. D* **78** (2008) 041504 [[gr-qc/0804.1686](#)].
- [7] R. Schützhold and W. Unruh, Quantum correlations across the black hole horizon, *Phys. Rev. D* **81** (2010) 124033 [[gr-qc/1002.1844](#)].
- [8] S. B. Giddings, Hawking radiation, the Stefan–Boltzmann law, and unitarization, *Phys. Lett. B* **754** (2016) 39 [[hep-th/1511.08221](#)].
- [9] S. W. Hawking, Breakdown of predictability in gravitational collapse, *Phys. Rev. D* **14** (1976) 2460.
- [10] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, and A. Tajdini, The entropy of Hawking radiation, *Rev. Mod. Phys.* **93** (2021) 035002 [[hep-th/2006.06872](#)].
- [11] B. Zhang, Q. Y. Cai, L. You, and M. S. Zhan, Hidden messenger revealed in Hawking radiation: a resolution to the paradox of black hole information loss, *Phys. Lett. B* **675** (2009) 98 [[hep-th/0903.08932](#)].
- [12] B. Zhang, Q. Y. Cai, M. S. Zhan, and L. You, Entropy is conserved in Hawking radiation as tunneling: a revisit of the black hole information loss paradox, *Ann. Phys.* **326** (2011) 350 [[hep-th/0906.5033](#)].

- [13] S. Hod, Hawking radiation and the Stefan–Boltzmann law: The effective radius of the black-hole quantum atmosphere, *Phys. Lett. B* **757** (2016) 121 [[gr-qc/1607.02510](#)].
- [14] R. Dey, S. Liberati, and D. Pranzetti, The black hole quantum atmosphere, *Phys. Lett. B* **774** (2017) 308 [[gr-qc/1701.06161](#)].
- [15] C. Buss and M. Casals, Quantum correlator outside a schwarzschild black hole, *Phys. Lett. B* **776** (2018) 168 [[hep-th/1709.05990](#)].
- [16] R. Dey, S. Liberati, Z. Mirzaiyan, and D. Pranzetti, Black hole quantum atmosphere for freely falling observers, *Phys. Lett. B* **797** (2019) 134828 [[gr-qc/1906.02958](#)].
- [17] M. Eune and W. Kim, Test of quantum atmosphere in the dimensionally reduced Schwarzschild black hole, *Phys. Lett. B* **798** (2019) 135020 [[gr-qc/1908.03374](#)].
- [18] A. Z. Kaczmarek and D. Szcześniak, Signatures of the black hole quantum atmosphere in nonlocal correlations, *Phys. Lett. B* **848** (2024) 138364 [[gr-qc/2306.09941](#)].
- [19] Y. C. Ong and M. R. R. Good, Quantum atmosphere of Reissner-Nordström black holes, *Phys. Rev. Res.* **2** (2020) 033322 [[gr-qc/2003.10429](#)].
- [20] M. Zych, F. Costa, I. Pikovski, and Č. Brukner, Bell’s theorem for temporal order, *Nat. Commun.* **10** (2019) 3772 [[quant-ph/1708.00248](#)].
- [21] L. J. Henderson, A. Belenchia, E. Castro-Ruiz, C. Budroni, M. Zych, Č. Brukner, and R.B. Mann, Quantum temporal superposition: the case of quantum field theory, *Phys. Rev. Lett.* **125** (2020) 131602 [[quant-ph/2002.06208](#)].
- [22] J. Foo, C. S. Arabaci, M. Zych, and R. B. Mann, Quantum superpositions of Minkowski spacetime, *Phys. Rev. D* **107** (2023) 045014 [[gr-qc/2208.12083](#)].
- [23] C. Suryaatmadja, C. S. Arabaci, M. P. G. Robbins, J. Foo, M. Zych, and R. B. Mann, Signatures of rotating black holes in quantum superposition, [[gr-qc/2310.10864](#)].
- [24] C. Barceló, L. J. Garay, and G. García-Moreno, Analogue gravity simulation of superpositions of spacetimes, *Eur. Phys. J. C* **82** (2022) 1 [[gr-qc/2104.15078](#)].
- [25] C. Barceló, J. E.Sánchez, G. García-Moreno, and G. Jannes, Chronology protection implementation in analogue gravity, *Eur. Phys. J. C* **82** (2022) 299 [[gr-qc/2201.11072](#)].
- [26] W. Unruh, Experimental black-hole evaporation?, *Phys. Rev. Lett.* **46** (1981) 1351.
- [27] C. Barcelo, S. Liberati, and M. Visser, Analogue gravity, *Living Rev. Relativity* **14** (2011) 1 [[gr-qc/0505065](#)].
- [28] Y. H. Shi, R. Q. Yang, Z. Xiang, Z. Y. Ge, H. Li, Y. Y. Wang, K. Huang, Y. Tian, X.

- Song, D. Zheng et al, Quantum simulation of hawking radiation and curved spacetime with asuperconducting on-chip black hole, *Nat. Commun.* **14** (2023) 3263 [[quant-ph/2111.11092](#)].
- [29] H. Zhang and B. Zhang, Shielding of penrose superradiance in optical black holes, *Eur. Phys. J. C* **83** (2023) 1 [[gr-qc/2303.14418](#)].
- [30] M. Christodoulou and C. Rovelli, On the possibility of laboratory evidence for quantum superposition of geometries, *Phys. Lett. B* **792** (2019) 64 [[gr-qc/1808.05842](#)].
- [31] C. Marletto and V. Vedral, Gravitationally induced entanglement between two massive particles is sufficient evidence of quantum effects in gravity, *Phys. Rev. Lett.* **119** (2017) 240402 [[quant-ph/1707.06036](#)].
- [32] S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Toroš, M. Paternostro, A. A. Geraci, P. F. Barker, M. S. Kim, and G. Milburn, Spin entanglement witness for quantum gravity, *Phys. Rev. Lett.* **119** (2017) 240401 [[quant-ph/1707.06050](#)].
- [33] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, The Unruh effect and its applications, *Rev. Mod. Phys.* **80** (2008) 787 [[gr-qc/0710.5373](#)].
- [34] J. Foo, C. S. Arabaci, M. Zych, and R. B. Mann, Quantum signatures of black hole mass superpositions, *Phys. Rev. Lett.* **129** (2022) 181301 [[gr-qc/2111.13315](#)].
- [35] J.D. Bekenstein, Black holes and entropy, *Phys. Rev. D* **7** (1973) 2333.
- [36] J.D. Bekenstein, The quantum mass spectrum of the kerr black hole, in *JACOB BEKENSTEIN: The Conservative Revolutionary* World Scientific, Singapore, (2020), pp.331–334.
- [37] S. Hod, Bohr’s Correspondence Principle and the Area Spectrum of Quantum Black Holes, *Phys. Rev. Lett.* **81** (1998) 4293 [[gr-qc/9812002](#)].
- [38] J. D. Bekenstein and V. Mukhanov, Spectroscopy of the quantum black hole, *Phys. Lett. B* **360** (1995) 7 [[gr-qc/9505012](#)].
- [39] J. D. Bekenstein, Quantum black holes as atoms, in Proceedings of the 8th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories (MG 8) (World Scientific Publishing Company, Singapore, 1997), pp. 92–111, [[gr-qc/9710076](#)].
- [40] E. C. Vagenas, Area spectrum of rotating black holes via the new interpretation of quasinormal modes, *J. High Energy Phys.* **11** (2008) 073 [[gr-qc/0804.3264](#)].
- [41] R. Banerjee, B. R. Majhi, and E. C. Vagenas, Quantum tunneling and black hole spectroscopy, *Phys. Lett. B* **686** (2010) 279 [[hep-th/0907.4271](#)].

- [42] B. Pourhassan, M. Faizal, Z. Zaz, and A. Bhat, Quantum fluctuations of a BTZ black hole in massive gravity, *Phys. Lett. B* **773** (2017) 325 [[gr-qc/1709.09573](#)].
- [43] K. Lochan and S. Chakraborty, Discrete quantum spectrum of black holes, *Phys. Lett. B* **755** (2016) 37 [[gr-qc/1509.09010](#)].
- [44] Y. Kwon and S. Nam, Area spectra of the rotating BTZ black hole from quasinormal modes, *Classical Quantum Gravity* **27** (2010) 125007 [[hep-th/1001.5106](#)].
- [45] C. Rovelli, Quantum Gravity (Cambridge University Press, Cambridge, England, 2004).
- [46] H. Kastrup and T. Thiemann, Spherically symmetric gravity as a completely integrable system, *Nucl. Phys. B* **425** (1994) 665 [[gr-qc/9401032](#)].
- [47] M. Campiglia, R. Gambini, and J. Pullin, Loop quantization of spherically symmetric midi-superspaces, *Classical Quantum Gravity* **24** (2007) 3649 [[gr-qc/0712.0817](#)].
- [48] R. Gambini, J. Olmedo, and J. Pullin, Quantum black holes in loop quantum gravity, *Classical Quantum Gravity* **31** (2014) 095009 [[gr-qc/1310.5996](#)].
- [49] R. Gambini and J. Pullin, Loop Quantization of the Schwarzschild Black Hole, *Phys. Rev. Lett.* **110** (2013) 211301 [[gr-qc/1302.5265](#)].
- [50] R. Balbinot, A. Fabbri, S. Fagnocchi, A. Recati, and I. Carusotto, Nonlocal density correlations as a signature of Hawking radiation from acoustic black holes, *Phys. Rev. A* **78** (2008) 021603 [[cond-mat/0711.4520](#)].
- [51] I. Carusotto, S. Fagnocchi, A. Recati, R. Balbinot, and A. Fabbri, Numerical observation of Hawking radiation from acoustic black holes in atomic Bose-Einstein condensates, *New J. Phys.* **10** (2008) 103001 [[cond-mat/0803.0507](#)].
- [52] A. Fabbri and R. Balbinot, Ramp-up of Hawking radiation in Bose-Einstein-condensate analog black holes, *Phys. Rev. Lett.* **126** (2021) 111301 [[gr-qc/2005.04443](#)].
- [53] L. Liao, E. C. I. Van Der Wurff, D. Van Oosten, and H. T. C. Stoof, Proposal for an analog Schwarzschild black hole in condensates of light, *Phys. Rev. A* **99** (2019) 023850 [[gr-qc/1806.00023](#)].
- [54] F. Michel, J-F. Coupechoux, and R. Parentani, Phonon spectrum and correlations in a transonic flow of an atomic Bose gas, *Phys. Rev. D* **94** (2016) 084027 [[gr-qc/1605.09752](#)].
- [55] J. Steinhauer, Observation of quantum Hawking radiation and its entanglement in an analogue black hole, *Nat. Phys.* **12** (2016) 959 [[gr-qc/1510.00621](#)].
- [56] J. R. M. de Nova, K. Golubkov, V. I. Kolobov, and J. Steinhauer, Observation of thermal

- Hawking radiation and its temperature in an analogue black hole, *Nature* **569** (2019) 688 [[gr-qc/1809.00913](#)].
- [57] R. Parentani, From vacuum fluctuations across an event horizon to long distance correlations, *Phys. Rev. D* **82** (2010) 025008 [[gr-qc/1003.3625](#)].
- [58] R. Balbinot and A. Fabbri, Quantum correlations across the horizon in acoustic and gravitational black holes, *Phys. Rev. D* **105** (2022) 045010 [[gr-qc/2107.00702](#)].
- [59] P. R. Anderson and J. Traschen, Horizons and correlation functions in 2d Schwarzschild-de Sitter spacetime, *J. High Energy Phys.* **2022** (2022) 192 [[hep-th/2012.08494](#)].
- [60] R. Balbinot and A. Fabbri, Quantum energy momentum tensor and equal time correlations in a Reissner-Nordström black hole, *Phys. Rev. D* **108** (2023) 045004 [[gr-qc/2303.11039](#)].
- [61] M. Fontana and M. Rinaldi, Stress-energy tensor correlations across regular black holes horizons, *Phys. Rev. D* **108** (2023) 125003 [[gr-qc/2302.08804](#)].
- [62] R. Balbinot and A. Fabbri, The Hawking effect in the particles-partners correlations, *Physics* **5** (2023) 968 [[gr-qc/2306.16952](#)].
- [63] M. Banados, C. Teitelboim, and J. Zanelli, Black hole in three-dimensional spacetime, *Phys. Rev. Lett.* **69** (1992) 1849 [[hep-th/9204099](#)].
- [64] S. Carlip, The  $(2+1)$ -dimensional black hole, *Classical Quantum Gravity* **12** (1995) 2853 [[gr-qc/9506079](#)].
- [65] M. Banados, M. Henneaux, C. Teitelboim, and J. Zanelli, Geometry of the  $2+1$  black hole, *Phys. Rev. D* **88** (1993) 069902 [[gr-qc/9302012](#)].
- [66] A. R. H. Smith and R. B. Mann, Looking inside a black hole, *Classical Quantum Gravity* **31** (2014) 082001 [[gr-qc/1309.4125](#)].
- [67] G. Lifschytz and M. Ortiz, Scalar field quantization on the  $(2+1)$ -dimensional black hole background, *Phys. Rev. D* **49** (1994) 1929 [[gr-qc/9310008](#)].
- [68] L. J. Henderson, R. A. Hennigar, R. B. Mann, A. R. H. Smith, and J. Zhang, Harvesting entanglement from the black hole vacuum, *Classical Quantum Gravity* **35** (2018) 21LT02 [[quant-ph/1712.10018](#)].
- [69] P. Painlevé, *C.R.Acad.sci. (paris)* 173 (1921) 677.
- [70] K. Gallock-Yoshimura, E. Tjoa, and R. B. Mann, Harvesting entanglement with detectors freely falling into a black hole, *Phys. Rev. D* **104** (2021) 025001 [[quant-ph/2102.09573](#)].
- [71] A. Saini and D. Stojkovic, Gravitational collapse and Hawking-like radiation of a shell in AdS

spacetime, *Phys. Rev. D* **97** (2018) 025020 [[hep-th/1711.08182](#)].

- [72] P. K. Townsend and B. Zhang, Thermodynamics of “exotic” Banados-Teitelboim-Zanelli black holes, *Phys. Rev. Lett.* **110** (2013) 241302 [[hep-th/1302.3874](#)].