

Attainability and criticality for multipolar Rellich inequality

Yongyang Jin, Shoufeng Shen and Li Tang

Abstract In this paper we obtain optimal multipolar Rellich inequality in \mathbb{R}^N for biharmonic Schrödinger operator with positive multisingular potentials of the form

$$H_n := \Delta^2 - \frac{N^2(N-4)^2}{n^4} \sum_{1 \leq i < j \leq n} \frac{|a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\nu_{k,i,j} \nu_{l,i,j} |a_k - a_l|^2}{|x - a_k|^2 |x - a_l|^2} \right),$$

where a_1, \dots, a_n are n different singular poles, $\nu_{k,i,j} = \frac{N+2n-4}{N}$ when $k = i$ or $k = j$, otherwise $\nu_{k,i,j} = \frac{N-4}{N}$. Moreover, we prove the attainability of the best constant $\frac{N^2(N-4)^2}{n^4}$ in $D^{2,2}(\mathbb{R}^N)$ for $n \geq 3$ and the criticality of the biharmonic Schrödinger operator H_n .

1 Introduction

The classical L^2 -Hardy inequality

$$(1.1) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq \frac{(N-2)^2}{4} \int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx$$

holds for $u \in C_0^\infty(\mathbb{R}^N)$ when $n \geq 3$. Hardy inequality are of fundamental importance in various branches of mathematics such as partial differential equation, spectral theory, analysis, mathematical physics, geometry and quantum mechanics, and have been comprehensively studied, just to name a few, see for example [2, 3, 5, 10, 11, 14, 21, 22, 25, 26, 27] and references therein.

At the 1954 ICM in Amsterdam [24], Rellich presented the following inequality as the second order version of the Hardy inequality for $N \geq 5$:

$$(1.2) \quad \int_{\mathbb{R}^N} |\Delta u|^2 dx \geq \frac{N^2(N-4)^2}{16} \int_{\mathbb{R}^N} \frac{|u|^2}{|x|^2} dx, \quad \forall u \in C_0^\infty(\mathbb{R}^N).$$

The constant $\frac{N^2(N-4)^2}{16}$ is sharp but can not be attained by nontrivial functions. Much effort has been devoted to the study of Rellich type inequalities and their applications. The reader may refer [4, 13, 15, 16, 20, 23] and the references therein for further details.

2020 *Mathematics Subject Classification.* 26D10, 42B37, 35A23, 35B25.

Key words and phrases. Multipolar Rellich inequalities; sharp constant; attainability; criticality.

Funding information: NNSF of China (12071431).

Definition 1.1. (*Sharp/Optimal constant*) Denote the norm $\|\cdot\|_{\sigma,2}$, $\sigma = 1, 2$, as $\|u\|_{1,2} = \|\nabla u\|_2$ and $\|u\|_{2,2} = \|\Delta u\|_2$. Denote Schrödinger/biharmonic Schrödinger operator $H = \Delta^\sigma - \lambda^*V$. We say that the constant λ^* of the following Hardy/Rellich type inequality associated with H

$$(1.3) \quad \|u\|_{\sigma,2}^2 \geq \lambda^* \int_{\mathbb{R}^N} V|u|^2 dx$$

is *sharp* or *optimal* if it is defined through the optimization problem

$$\lambda^* := \inf_{u \in D^{\sigma,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\|u\|_{\sigma,2}^2}{\int_{\mathbb{R}^N} V|u|^2 dx},$$

where

$$D^{\sigma,2}(\mathbb{R}^N) := \overline{C_0^\infty(\mathbb{R}^N)}^{\|\cdot\|_{\sigma,2}}.$$

It is well known that the constants $\frac{(N-2)^2}{4}$ and $\frac{N^2(N-4)^2}{16}$ in inequality (1.1) and (1.2) are both sharp but not attained.

Recently, motivated by problems in quantum mechanics, many authors studied multipolar Hardy inequalities and their applications to Schrödinger operators with multipolar potentials. The readers may find some relevant results in [1, 7, 8, 9, 17, 18, 19] and the reference therein. The two most studied potentials are

$$W_1(x) = \sum_{i=1}^n \frac{1}{|x - a_i|^2}, \quad W_2(x) = \sum_{1 \leq i < j \leq n} \frac{|a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2}.$$

In order to obtain a lower bound of the spectrum of the Schrödinger operators $-\Delta - \mu W_1$, $\mu \in (0, \frac{(N-2)^2}{4}]$, Bosi-Dolbeault-Esteban in [1] proved for $u \in C_0^\infty(\mathbb{R}^N)$ the following multipolar Hardy inequality

$$(1.4) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq \frac{(N-2)^2}{4n} \int_{\mathbb{R}^N} W_1 |u|^2 dx + \frac{(N-2)^2}{4n^2} \int_{\mathbb{R}^N} W_2 |u|^2 dx.$$

Several years later, Cazacu-Zuazua obtained in [9] the following inequality

$$(1.5) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq \frac{(N-2)^2}{n^2} \int_{\mathbb{R}^N} W_2 |u|^2 dx, \quad u \in C_0^\infty(\mathbb{R}^N),$$

where the constant $\frac{(N-2)^2}{n^2}$ is sharp for $N \geq 3$. The inequality (1.5) is an improvement of (1.4) in the sense that: the total mass near a_i of the potential in (1.5) is $\frac{(N-2)^2}{4} \frac{4n-4}{n^2} \frac{1}{|x-a_i|^2}$, which is strictly larger than the total mass $\frac{(N-2)^2}{4} \frac{2n-1}{n^2} \frac{1}{|x-a_i|^2}$ near a_i in inequality (1.4). Another multipolar Hardy inequality proved in [12] by Devyver-Fraas-Pinchover reads as

$$(1.6) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq \left(\frac{N-2}{n+1} \right)^2 \int_{\mathbb{R}^N} (W_1 + W_2) |u|^2 dx, \quad u \in C_0^\infty(\mathbb{R}^N),$$

where the constant $\left(\frac{N-2}{n+1} \right)^2$ is sharp. Furthermore, they gave a stronger concept, say *criticality* of Schrödinger operator, to characterize the *sharpness/optimalty* of an inequality than the "sharp/optimal constant". We recall the original definition in [12].

Definition 1.2. Assume that $H \geq 0$. The operator H is said to be *subcritical* in \mathbb{R}^N if there exists a nonzero nonnegative continuous function W such that

$$\langle Hu, u \rangle \geq \int_{\mathbb{R}^N} W|u|^2 dx, \quad u \in C_0^\infty(\mathbb{R}^N).$$

Otherwise, we say that H is *critical* in \mathbb{R}^N .

Devyver-Fraas-Pinchover proved in [12] that the Schrödinger operators $-\Delta - \frac{(N-2)^2}{n^2}W_2$ and $-\Delta - \left(\frac{N-2}{n+1}\right)^2(W_1 + W_2)$ are both critical, this implies that one cannot add any positive reminders to the right hand side of inequalities (1.5) and (1.6). With this in mind, one may want to know whether the following inequality (1.7) is critical, since these three inequalities are intrinsically related. In fact, assume $2 - N \leq s \leq \frac{2-N}{2}$, we have (from Lemma 2.2 and identity (2.1)), for any $u \in C_0^\infty(\mathbb{R}^N)$, the following inequality

$$(1.7) \quad \int_{\mathbb{R}^N} |\nabla u|^2 dx \geq [s(N-2) - s^2] \sum_{i=1}^n \int_{\mathbb{R}^N} \alpha_i \frac{|u|^2}{|x - a_i|^2} dx \\ + s^2 \sum_{1 \leq i < j \leq n} \int_{\mathbb{R}^N} \alpha_i \alpha_j \frac{|a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2} |u|^2 dx$$

holds, where $\alpha_i \geq 0$, $i = 1, \dots, n$, $\sum_{i=1}^n \alpha_i = 1$. One can also prove that, by the method introduced in [12] of constructing the "optimal Hardy weight", the Schrödinger operator

$$-\Delta - [s(N-2) - s^2] \sum_{i=1}^n \frac{\alpha_i}{|x - a_i|^2} dx - s^2 \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2}$$

is also critical when $0 < \alpha_i \leq \frac{1}{2}$, $i = 1, \dots, n$. Let $\alpha_i = \frac{1}{n}$ for $i = 1, \dots, n$, the inequalities (1.4), (1.5) and (1.6) are all special cases of inequality (1.7) if we let $s = \frac{2-N}{2}$, $2 - N$ and $\frac{n(2-N)}{n+1}$ respectively.

In this paper we always assume $n \geq 2$. We will work in the function space $D^{2,2}(\mathbb{R}^N)$. The first goal of the present article is to obtain the optimal L^2 -version multipolar Rellich inequality. To our shallow knowledge, there has been no result on multipolar Rellich inequality in the existing literatures. A universal approach to obtain Hardy and Rellich inequalities is the so-called method of super-solutions. However, it is not an easy work to compute the fourth-order derivative of a ground state function in the case of Rellich inequality. In this paper, we take an subtle iterative method to solve this difficulty. The main result of the paper is as follows.

Theorem 1.3. Assume $N \geq 5$, $a_1, \dots, a_n \in \mathbb{R}^N$. Then the following inequality holds for any $u \in C_0^\infty(\mathbb{R}^N)$,

$$(1.8) \quad \int_{\mathbb{R}^N} |\Delta u|^2 dx \geq \frac{N^2(N-4)^2}{n^4} \int_{\mathbb{R}^N} V_n(x) |u|^2 dx,$$

where

$$V_n(x) := \sum_{1 \leq i < j \leq n} \frac{|a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\nu_{k,i,j} \nu_{l,i,j} |a_k - a_l|^2}{|x - a_k|^2 |x - a_l|^2} \right),$$

with

$$\nu_{k,i,j} = \begin{cases} \frac{N+2n-4}{N}, & k = i \text{ or } k = j; \\ \frac{N-4}{N}, & \text{otherwise.} \end{cases}$$

The constant $\frac{N^2(N-4)^2}{n^4}$ is sharp. Furthermore, the constant $\frac{N^2(N-4)^2}{n^4}$ is achieved in the space $D^{2,2}(\mathbb{R}^N)$ when $n \geq 3$ by the minimizers (up to a constant)

$$\phi(x) = \prod_{i=1}^n |x - a_i|^{\frac{4-N}{n}},$$

while it can not be attained in the case $n = 2$.

Especially, $\nu_{k,i,j} = 1$ ($\forall i, j, k = 1, 2$) when $n = 2$, we thus have

Corollary 1.4. Assume $N \geq 5$, $a_1, \dots, a_n \in \mathbb{R}^N$. The following inequality holds for any $u \in C_0^\infty(\mathbb{R}^N)$,

$$(1.9) \quad \int_{\mathbb{R}^N} |\Delta u|^2 dx \geq \frac{N^2(N-4)^2}{16} \int_{\mathbb{R}^N} \frac{|a_1 - a_2|^4}{|x - a_1|^4 |x - a_2|^4} |u|^2 dx,$$

where the constant $\frac{N^2(N-4)^2}{16}$ is sharp and it cannot be attained in $D^{2,2}(\mathbb{R}^N)$.

Note that it is a surprising fact that the sharp constant in (1.9) is the same as the classical sharp Rellich constant with single pole.

Next, we consider the criticality of the biharmonic Schrödinger operator

$$H_n := \Delta^2 - \frac{N^2(N-4)^2}{n^4} V_n.$$

We now introduce some definitions for the classification of the critical Schrödinger operator.

Definition 1.5. Assume the Schrödinger/biharmonic Schrödinger operator H is critical in \mathbb{R}^N . H is said to be *positive-critical* in \mathbb{R}^N if the sharp constant λ^* in (1.3) is achieved in $D^{\sigma,2}(\mathbb{R}^N)$. Otherwise, H is called *null-critical* in \mathbb{R}^N .

The attainability of constant $\frac{N^2(N-4)^2}{n^4}$ in (1.8) implies, in fact, the criticality of H_n when $n \geq 3$. More precisely, we have the following criticality result of biharmonic Schrödinger operator H_n for $n \geq 2$.

Proposition 1.6. The biharmonic Schrödinger operator $H_n = \Delta^2 - \frac{N^2(N-4)^2}{n^4} V_n$ is critical in \mathbb{R}^N for $n \geq 2$. Moreover, H_n is positive-critical for $n \geq 3$ and null-critical for $n = 2$.

The plan of this paper is as follows. In section 2 we use the method of super-solutions to obtain the L^2 -case multipolar Rellich inequality (1.8) with sharp constant. In section 3 we prove the attainability or non-attainability of the constant $\frac{N^2(N-4)^2}{n^4}$ and the criticality of the biharmonic Schrödinger operator H_n .

2 L^2 -case multipolar Rellich inequality with sharp constant

In this section we employ the method of super-solutions to prove the sharp multipolar Rellich inequality. Recall that the following Hardy-type identity: Let positive function $\phi \in C^2(\Omega)$. It holds for any $u \in C_0^\infty(\Omega)$ that

$$(2.1) \quad \int_{\Omega} \left| \nabla u - \frac{\nabla \phi}{\phi} u \right|^2 dx = \int_{\Omega} |\nabla u|^2 dx + \int_{\Omega} \frac{\Delta \phi}{\phi} |u|^2 dx.$$

For nonnegative function $W(x) \in L_{loc}^1(\Omega)$, the method of super-solutions shows that if there exists a positive function ϕ which is a super-solution of the Schrödinger operator $-\Delta - W$, i.e. $(-\Delta - W)\phi \geq 0$, then $-\Delta - W \geq 0$, which means that

$$\int_{\Omega} |\nabla u|^2 dx \geq \int_{\Omega} W |u|^2 dx, \quad \forall u \in C_0^\infty(\Omega).$$

Thus, one only need to choose suitable ϕ to obtain Hardy inequalities for any potentials $W \leq \frac{-\Delta \phi}{\phi}$. This method can yield most of the well-known Hardy inequalities, for example, choosing $\phi(x) = |x|^{\frac{2-N}{2}}$ to yield the classical Hardy inequality and $\phi_s(x) = \prod_{i=1}^n |x - a_i|^{s\alpha_i}$ to obtain inequality (1.7) (see also Lemma 2.2).

In order to apply the method of super-solutions to obtain the multipolar Rellich inequality, we need to find an Rellich-type identity analogous to (2.1) for $\Omega = \mathbb{R}^N$, and choose a special auxiliary function ϕ . Assume $0 < \phi \in C^4(\Omega)$ for now, we start from the following identity

$$\int_{\Omega} \left| \Delta u - \frac{\Delta \phi}{\phi} u \right|^2 dx = \int_{\Omega} |\Delta u|^2 dx + \int_{\Omega} \frac{(\Delta \phi)^2}{\phi^2} |u|^2 dx - \int_{\Omega} 2u \Delta u \frac{\Delta \phi}{\phi} dx.$$

Denoting $v = u\phi^{-1}$, then,

$$\begin{aligned} \int_{\Omega} \left| \Delta u - \frac{\Delta \phi}{\phi} u \right|^2 dx &= \int_{\Omega} |\Delta u|^2 dx + \int_{\Omega} (\Delta \phi)^2 |v|^2 dx - \int_{\Omega} 2v \Delta(v\phi) \Delta \phi dx. \\ &= \int_{\Omega} |\Delta u|^2 dx - \int_{\Omega} (\Delta \phi)^2 |v|^2 dx - \int_{\Omega} 2v \Delta \phi (\phi \Delta v + 2\nabla \phi \cdot \nabla v) dx \\ &= \int_{\Omega} |\Delta u|^2 dx - \int_{\Omega} \Delta \phi (\Delta(\phi |v|^2) + 2v \Delta v \phi + 2\nabla \phi \cdot \nabla(|v|^2)) dx \\ &= \int_{\Omega} |\Delta u|^2 dx - \int_{\Omega} \Delta \phi (\Delta(\phi |v|^2) - 2|\nabla v|^2 \phi) dx \\ &= \int_{\Omega} |\Delta u|^2 dx - \int_{\Omega} \frac{\Delta^2 \phi}{\phi} |u|^2 dx + 2 \int_{\Omega} \phi \Delta \phi |\nabla v|^2 dx \end{aligned}$$

Thus, we obtain

$$(2.2) \quad \int_{\Omega} |\Delta u|^2 dx - \int_{\Omega} \frac{\Delta^2 \phi}{\phi} |u|^2 dx = \int_{\Omega} |\phi \Delta v + 2\nabla \phi \cdot \nabla v|^2 dx - 2 \int_{\Omega} \phi \Delta \phi |\nabla v|^2 dx.$$

Lemma 2.1. (See also [6]) *Assume $W(x) \in L^1_{loc}(\Omega)$ is a nonnegative function. If $0 < \phi \in C^4(\Omega)$ and $-\Delta\phi \geq 0$ in Ω , such that the following partial differential inequality holds*

$$(\Delta^2 - W)\phi \geq 0 \text{ in } \Omega,$$

then, we have the following Rellich inequality

$$\int_{\Omega} |\Delta u|^2 dx \geq \int_{\Omega} W|u|^2 dx, \quad \forall u \in C_0^\infty(\Omega).$$

We deduce from Lemma 2.1 that: if there is a positive function ϕ with $\frac{\Delta^2\phi}{\phi} \in L^1_{loc}(\mathbb{R}^N)$, then it holds

$$\int_{\mathbb{R}^N} |\Delta u|^2 dx \geq \int_{\mathbb{R}^N} \frac{\Delta^2\phi}{\phi} |u|^2 dx, \quad \forall u \in C_0^\infty(\mathbb{R}^N).$$

We now establish the following lemma.

Lemma 2.2. *Let $\phi_s(x) = \prod_{i=1}^n |x - a_i|^{s\alpha_i}$, $s \in \mathbb{R}$, $0 \leq \alpha_i \leq 1$ and $\sum_{k=1}^n \alpha_k = 1$. Then*

$$(2.3) \quad \Delta\phi_s = \phi_s \left(s(N + s - 2) \sum_{i=1}^n \frac{\alpha_i}{|x - a_i|^2} - s^2 \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2} \right).$$

Proof. Rewriting ϕ_s with the form

$$\phi_s(x) = \prod_{i=1}^n \phi_i^{\alpha_i},$$

where $\phi_i = |x - a_i|^s$. By direct computation,

$$\Delta\phi_s = \phi_s \left| \sum_{i=1}^n \alpha_i \frac{\nabla\phi_i}{\phi_i} \right|^2 + \phi_s \left(\sum_{i=1}^n \alpha_i \frac{\Delta\phi_i}{\phi_i} - \sum_{i=1}^n \alpha_i \frac{|\nabla\phi_i|^2}{\phi_i^2} \right).$$

Simplifying the expression we obtain

$$(2.4) \quad \Delta\phi_s = \phi_s \left(\sum_{i=1}^n \alpha_i \frac{\Delta\phi_i}{\phi_i} - \sum_{1 \leq i < j \leq n} \alpha_i \alpha_j \left| \frac{\nabla\phi_i}{\phi_i} - \frac{\nabla\phi_j}{\phi_j} \right|^2 \right).$$

Since $\phi_i = |x - a_i|^s$,

$$(2.5) \quad \frac{\Delta\phi_i}{\phi_i} = s(N + s - 2) \frac{1}{|x - a_i|^2}, \quad \frac{|\nabla\phi_i|^2}{\phi_i^2} = s^2 \frac{1}{|x - a_i|^2}.$$

Inserting (2.5) into (2.4) we have

$$\Delta\phi_s = \phi_s \left(s(N + s - 2) \sum_{i=1}^n \alpha_i \frac{1}{|x - a_i|^2} - s^2 \sum_{i < j} \alpha_i \alpha_j \frac{|a_i - a_j|^2}{|x - a_i|^2 |x - a_j|^2} \right).$$

Thus the proof of Lemma 2.2 is complete. \square

In order to compute the $\Delta^2\phi_s$, we write $\Delta\phi_s$ as

$$(2.6) \quad \Delta\phi_s = s(N + s - 2) \sum_{i=1}^n \alpha_i \hat{\phi}_{s,i} - s^2 \sum_{1 \leq i < j \leq n} \alpha_i \alpha_j |a_i - a_j|^2 \hat{\phi}_{s,i,j}.$$

where $\hat{\phi}_{s,i} = \prod_{k=1}^n |x - a_k|^{(s-2)\xi_{k,i}}$, with

$$\xi_{k,i} = \begin{cases} \frac{s\alpha_k - 2}{s-2} & , k = i; \\ \frac{s\alpha_k}{s-2} & , k \neq i, \end{cases}$$

and $\hat{\phi}_{s,i,j} = \prod_{k=1}^n |x - a_k|^{(s-4)\zeta_{k,i,j}}$, with

$$\zeta_{k,i,j} = \begin{cases} \frac{s\alpha_k - 2}{s-4} & , k = i \text{ or } k = j; \\ \frac{s\alpha_k}{s-4} & , \text{otherwise.} \end{cases}$$

It is easy to check that

$$\sum_{k=1}^n \xi_{k,i} = 1, \quad \sum_{k=1}^n \zeta_{k,i,j} = 1.$$

Applying Lemma 2.2 to $\hat{\phi}_i$ and $\hat{\phi}_{i,j}$ we have

$$(2.7) \quad \Delta\hat{\phi}_{s,i} = \hat{\phi}_{s,i} \left((s-2)(N + s - 4) \sum_{k=1}^n \frac{\xi_{k,i}}{|x - a_k|^2} - (s-2)^2 \sum_{1 \leq k < l \leq n} \frac{\xi_{k,i} \xi_{l,i} |a_k - a_l|^2}{|x - a_k|^2 |x - a_l|^2} \right),$$

$$(2.8) \quad \Delta\hat{\phi}_{s,i,j} = \hat{\phi}_{s,i,j} \left((s-4)(N + s - 6) \sum_{k=1}^n \frac{\zeta_{k,i,j}}{|x - a_k|^2} - (s-4)^2 \sum_{1 \leq k < l \leq n} \frac{\zeta_{k,i,j} \zeta_{l,i,j} |a_k - a_l|^2}{|x - a_k|^2 |x - a_l|^2} \right).$$

Since Δ is a linear operator, then

$$(2.9) \quad \Delta^2\phi_s = s(N + s - 2) \sum_{i=1}^n \alpha_i \Delta\hat{\phi}_{s,i} - s^2 \sum_{1 \leq i < j \leq n} \alpha_i \alpha_j |a_i - a_j|^2 \Delta\hat{\phi}_{s,i,j}.$$

Taking (2.7) and (2.8) into (2.9), we get

$$\begin{aligned}
\frac{\Delta^2 \phi_s}{\phi_s} &= s(N+s-2)(s-2)(N+s-4) \sum_{i=1}^n \frac{\alpha_i}{|x-a_i|^2} \left(\sum_{k=1}^n \frac{\xi_{k,i}}{|x-a_k|^2} \right) \\
&\quad - s(N+s-2)(s-2)^2 \sum_{i=1}^n \frac{\alpha_i}{|x-a_i|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\xi_{k,i} \xi_{l,i} |a_k - a_l|^2}{|x-a_k|^2 |x-a_l|^2} \right) \\
&\quad - s^2(s-4)(N+s-6) \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{k=1}^n \frac{\zeta_{k,i,j}}{|x-a_k|^2} \right) \\
&\quad + s^2(s-4)^2 \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\zeta_{k,i,j} \zeta_{k,i,j} |a_k - a_l|^2}{|x-a_k|^2 |x-a_l|^2} \right).
\end{aligned}$$

Thus, we arrive at

$$\begin{aligned}
(2.10) \quad \frac{\Delta^2 \phi_s}{\phi_s} &= s(N+s-2)(s-2)(N+s-4) \sum_{i=1}^n \frac{\alpha_i}{|x-a_i|^2} \left(\sum_{k=1}^n \frac{\xi_{k,i}}{|x-a_k|^2} \right) \\
&\quad + 2s^2(s-4)(4-N-s) \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{k=1}^n \frac{\zeta_{k,i,j}}{|x-a_k|^2} \right) \\
&\quad + s^2(s-4)^2 \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\zeta_{k,i,j} \zeta_{k,i,j} |a_k - a_l|^2}{|x-a_k|^2 |x-a_l|^2} \right).
\end{aligned}$$

The last equality holds because of the the following identity

$$\sum_{i=1}^n \frac{\alpha_i}{|x-a_i|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\xi_{k,i} \xi_{l,i} |a_k - a_l|^2}{|x-a_k|^2 |x-a_l|^2} \right) = \frac{s(s-4)}{(s-2)^2} \sum_{1 \leq i < j \leq n} \frac{\alpha_i \alpha_j |a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{k=1}^n \frac{\zeta_{k,i,j}}{|x-a_k|^2} \right).$$

Theorem 2.3. *Assume $4 - N \leq s < 0$, $N \geq 5$, $a_1, \dots, a_n \in \mathbb{R}^N$. Then the following inequality holds for any $u \in C_0^\infty(\mathbb{R}^N)$,*

$$\begin{aligned}
(2.11) \quad \int_{\mathbb{R}^N} |\Delta u|^2 dx &\geq \frac{s(s-2)(2-N-s)(4-N-s)}{n^2} \int_{\mathbb{R}^N} \sum_{1 \leq i, k \leq n} \frac{\mu_{k,i}}{|x-a_i|^2 |x-a_k|^2} |u|^2 dx \\
&\quad + \frac{2s^3(4-N-s)}{n^3} \int_{\mathbb{R}^N} \sum_{1 \leq i < j \leq n} \frac{|a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{1 \leq k \leq n} \frac{\sigma_{k,i,j}}{|x-a_k|^2} \right) |u|^2 dx \\
&\quad + \frac{s^2(s-4)^2}{n^4} \int_{\mathbb{R}^N} \sum_{1 \leq i < j \leq n} \frac{|a_i - a_j|^2}{|x-a_i|^2 |x-a_j|^2} \left(\sum_{1 \leq k < l \leq n} \frac{\nu_{k,i,j} \nu_{l,i,j} |a_k - a_l|^2}{|x-a_k|^2 |x-a_l|^2} \right) |u|^2 dx,
\end{aligned}$$

where

$$\mu_{k,i} = \begin{cases} \frac{s-2n}{s-2}, & k = i; \\ \frac{s}{s-2}, & k \neq i, \end{cases} \quad \sigma_{k,i,j} = \begin{cases} \frac{s-2n}{s}, & k = i \text{ or } k = j; \\ 1, & \text{otherwise,} \end{cases} \quad \nu_{k,i,j} = \begin{cases} \frac{s-2n}{s-4}, & k = i \text{ or } k = j; \\ \frac{s}{s-4}, & \text{otherwise.} \end{cases}$$

Proof. Taking $\alpha_i = \frac{1}{n}$, $i = 1, \dots, n$, since $N \geq 5$, $\phi_s > 0$, and

$$(2.12) \quad -\Delta\phi_s = \frac{s(2-N-s)}{n}\phi \sum_{i=1}^n \frac{1}{|x-a_i|^2} + \frac{s^2}{n^2}\phi \sum_{1 \leq i < j \leq n} \frac{|a_i - a_j|^2}{|x-a_i|^2|x-a_j|^2} \geq 0,$$

Theorem 2.3 follows from Lemma 2.1. \square

Proof of inequality (1.8): Taking $s = 4 - N$ in (2.11) we obtain inequality (1.8).

3 Attainability and Criticality

In this section we prove the attainability of constant $\frac{N^2(N-4)^2}{n^4}$ in inequality (1.8) and the criticality of the biharmonic schrödinger operator H_n for $N \geq 5$.

3.1 Attainability for constant $\frac{N^2(N-4)^2}{n^4}$

Let $\phi(x) = \prod_{i=1}^n |x - a_i|^{\frac{4-N}{n}}$. In view of (2.10), $\phi(x)$ is the minimizer or virtual minimizer in inequality (1.8). We need to check the integrability of $|\Delta\phi|^2$. Let $s = 4 - N$ in (2.12) we have

$$(3.1) \quad \Delta\phi = -2(N-4)\phi \sum_{i=1}^n \frac{\alpha_i}{|x-a_i|^2} - (N-4)^2\phi \sum_{1 \leq i < j \leq n} \frac{\alpha_i\alpha_j|a_i - a_j|^2}{|x-a_i|^2|x-a_j|^2}.$$

Denote

$$\begin{aligned} a_i(x) &= \frac{\phi(x)^2}{|x-a_i|^4}, \quad b_{i,j}(x) = \frac{\phi(x)^2}{|x-a_i|^2|x-a_j|^2}, \quad c_{i,j}(x) = \frac{\phi(x)^2}{|x-a_i|^4|x-a_j|^2}, \\ d_{i,j,k}(x) &= \frac{\phi(x)^2}{|x-a_i|^2|x-a_j|^2|x-a_k|^2}, \quad e_{i,j}(x) = \frac{\phi(x)^2}{|x-a_i|^4|x-a_j|^4}, \\ f_{i,j,k,l}(x) &= \frac{\phi(x)^2}{|x-a_i|^2|x-a_j|^2|x-a_k|^2|x-a_l|^2}, \quad g_{i,j,k}(x) = \frac{\phi(x)^2}{|x-a_i|^4|x-a_j|^2|x-a_k|^2}. \end{aligned}$$

Expanding $|\Delta\phi|^2$, we know that $|\Delta\phi|^2$ is just a linear combination of these seven classes of functions. The integrability results of these function is as follows.

Proposition 3.1. $a_i(x)$, $b_{i,j}(x)$, $c_{i,j}(x)$, $d_{i,j,k}(x)$, $e_{i,j}(x)$, $f_{i,j,k,l}(x)$, and $g_{i,j,k}(x)$ are all in $L^1(\mathbb{R}^N)$ when $n \geq 3$, while $a_i(x)$, $c_{i,j}(x)$, $e_{i,j}(x)$ and $g_{i,j,k}(x)$ are not in $L^1(\mathbb{R}^N)$ when $n = 2$.

Proof. We only prove the integrability results for $a_i(x)$ and $e_{i,j}(x)$, the argument of others is similar. Assume $\rho > 0$ small enough such that $\{a_1, \dots, a_n\} \subset B_{\frac{1}{\rho}}(0)$ and $B_\rho(a_i) \cap B_\rho(a_j) = \emptyset$ for any $i \neq j$. Since $a_i(x)$ and $e_{i,j}(x)$ are bounded functions in the bounded domain $B_{\frac{1}{\rho}}(0) \setminus \bigcup_{i=1}^n B_\rho(a_i)$, we only need to consider the integrability of them in $B_{\frac{1}{\rho}}(0)^c$ and $B_\rho(a_i)$ for $i = 1, \dots, n$. Note that $|x - a_j| \sim |a_i - a_j|$ for $x \in B_\rho(a_i)$ when $i \neq j$ and $|x - a_i| \sim |x|$ for $x \in B_{\frac{1}{\rho}}(0)^c$, the following asymptotical behavior holds:

1. for $x \in B_\rho(a_i)$,

$$a_i(x) \simeq e_{i,j}(x) \simeq |x - a_i|^{\frac{(n-2)(N-4)}{n} - N};$$

2. for $x \in B_\rho(a_j)$,

$$|x - a_j|^{\frac{(n-2)(N-4)}{n} - N + 4} \simeq a_i(x) \lesssim e_{i,j}(x) \simeq |x - a_j|^{\frac{(n-2)(N-4)}{n} - N};$$

3. for $x \in B_\rho(a_k)$, $k \neq i, j$,

$$a_i(x) \simeq e_{i,j}(x) \simeq |x - a_i|^{\frac{(n-2)(N-4)}{n} - N + 4};$$

4. for $x \in B_{\frac{1}{\rho}}(0)^c$,

$$|x|^{4-2N} \simeq a_i(x) \gtrsim e_{i,j}(x) \simeq |x|^{-2N};$$

Therefore, it shows that $a_i(x), e_{i,j}(x) \in L^1(B_{\frac{1}{\rho}}(0)^c \cup \bigcup_{i=1}^n B_\rho(a_i))$ when $n \geq 3$ and $N \geq 5$, whereas $a_i(x), e_{i,j}(x) \notin L^1(B_\rho(a_i))$ when $n = 2$. Thus, we prove the claim for $a_i(x), e_{i,j}(x)$ in Proposition 3.1 and the proof for other functions is similar. This complete the proof. \square

Proof of the attainability: By (3.1) and Proposition 3.1, we get the integrability of $|\Delta\phi|^2$ when $n \geq 3$, which means $\phi \in D^{2,2}(\mathbb{R}^N)$. So, by using integration by parts, we get the following identity

$$(3.2) \quad \int_{\mathbb{R}^N} |\Delta\phi|^2 dx = \int_{\mathbb{R}^N} \frac{\Delta^2 \phi}{\phi} \phi^2 dx = \frac{N^2(N-4)^2}{n^4} \int_{\mathbb{R}^N} V_n(x) \phi^2 dx.$$

This implies the attainability of $\frac{N^2(N-4)^2}{n^4}$ in $D^{2,2}(\mathbb{R}^N)$ when $n \geq 3$. When $n = 2$, $\phi \notin D^{2,2}(\mathbb{R}^N)$ since $a_i(x)$ is not in $L^1(\mathbb{R}^N)$, which shows that the constant $\frac{N^2(N-4)^2}{16}$ is not attained in $D^{2,2}(\mathbb{R}^N)$. \square

3.2 Criticality for $n = 2$

We know from Theorem 1.3 that H_n is positive-critical for $n \geq 3$, and we study the criticality for $n = 2$ in this subsection. For general subcritical second-order elliptic operator, a convenient method to construct a critical schrödinger operator is given in [12] by constructing the *optimal Hardy-weight*, but we have no such direct approach for biharmonic schrödinger operator. Therefore, we need to find other method to deal with the criticality of nonnegative operators.

Lemma 3.2. *A nonnegative operator H defined in Hilbert space $L^2(\mathbb{R}^N)$ is critical if there exists a function sequence $\{u_\epsilon\}_{\epsilon>0}$ and positive a.e. continuous function $\psi(x)$, such that*

$$\lim_{\epsilon \rightarrow 0} \langle Hu_\epsilon, u_\epsilon \rangle = 0, \text{ and } \lim_{\epsilon \rightarrow 0} u_\epsilon = \psi(x) \text{ a.e. in } \mathbb{R}^N.$$

Proof. We assume that there exists a nonzero nonnegative potential W such that

$$\langle Hu, u \rangle \geq \int_{\mathbb{R}^N} W|u|^2 dx.$$

Then, from Fatou lemma we have

$$0 \leq \int_{\mathbb{R}^N} W|\psi|^2 dx \leq \liminf_{\epsilon \rightarrow 0} \int_{\mathbb{R}^N} W|u_\epsilon|^2 dx \leq \lim_{\epsilon \rightarrow 0} \langle Hu_\epsilon, u_\epsilon \rangle = 0,$$

which forces that $W = 0$ a.e. in \mathbb{R}^N . So, the proof is completed by contradiction. \square

In view of the Rellich-type identity (2.2), we have the following result to determine the criticality of biharmonic schrödinger operator H_2 .

Corollary 3.3. *Let ϕ satisfies the assumptions in Lemma 2.1. The operator $\Delta^2 - \frac{\Delta^2 \phi}{\phi}$ is critical if there exists a function sequence $\{v_\epsilon\}_{\epsilon>0}$, with $0 < v_\epsilon \leq 1$, such that*

$$\lim_{\epsilon \rightarrow 0} \left(\int_{\mathbb{R}^N} |\phi \Delta v_\epsilon + 2 \nabla \phi \cdot \nabla v_\epsilon|^2 dx - \int_{\mathbb{R}^N} \phi \Delta \phi |\nabla v_\epsilon|^2 dx \right) = 0, \text{ and } \lim_{\epsilon \rightarrow 0} v_\epsilon = 1.$$

Proof of Proposition 1.6 for $n = 2$: In this case $\phi(x) = |x - a_1|^{\frac{4-N}{2}} |x - a_2|^{\frac{4-N}{2}}$. Let's select a cut-off function as follows

$$v_\epsilon(x) = \begin{cases} 0 & , x \in B_{\epsilon^2}(a_i), i = 1, 2; \\ \frac{\log(|x-a_i|/\epsilon^2)}{\log 1/\epsilon} & , x \in B_\epsilon(a_i) \setminus B_{\epsilon^2}(a_i), i = 1, 2; \\ 1 & , x \in B_{\frac{1}{\epsilon}}(0) \setminus \bigcup_{i=1}^n B_\epsilon(a_i), i = 1, 2; \\ \frac{\log(\epsilon^2|x|)}{\log \epsilon} & , x \in B_{\frac{1}{\epsilon^2}}(0) \setminus B_{\frac{1}{\epsilon}}(0), i = 1, 2; \\ 0 & , |x| \geq \frac{1}{\epsilon^2}. \end{cases}$$

By direct computation, the gradient and Laplacian of v_ϵ are respectively

$$\nabla v_\epsilon(x) = \begin{cases} \frac{x-a_i}{|x-a_i|^2} (\log \frac{1}{\epsilon})^{-1} & , x \in B_\epsilon(a_i) \setminus B_{\epsilon^2}(a_i), i = 1, 2; \\ \frac{x}{|x|^2} \log \frac{1}{\epsilon} & , x \in B_{\frac{1}{\epsilon^2}}(0) \setminus B_{\frac{1}{\epsilon}}(0), i = 1, 2; \\ 0 & , \text{otherwise.} \end{cases}$$

and

$$\Delta v_\epsilon(x) = \begin{cases} \frac{N-2}{|x-a_i|^2} (\log \frac{1}{\epsilon})^{-1} & , x \in B_\epsilon(a_i) \setminus B_{\epsilon^2}(a_i), i = 1, 2; \\ \frac{N-2}{|x|^2} \log \frac{1}{\epsilon} & , x \in B_{\frac{1}{\epsilon^2}}(0) \setminus B_{\frac{1}{\epsilon}}(0), i = 1, 2; \\ 0 & , \text{otherwise.} \end{cases}$$

On the other hand, we have

$$\begin{aligned}
& \int_{\mathbb{R}^N} |\phi \Delta v_\epsilon + 2\nabla \phi \cdot \nabla v_\epsilon|^2 dx - \int_{\mathbb{R}^N} \phi \Delta \phi |\nabla v_\epsilon|^2 dx \\
& \lesssim \int_{\mathbb{R}^N} |\phi \Delta v_\epsilon|^2 dx + \int_{\mathbb{R}^N} |\nabla \phi \cdot \nabla v_\epsilon|^2 dx + \int_{\mathbb{R}^N} |\phi \Delta \phi| |\nabla v_\epsilon|^2 dx \\
& =: I_1 + I_2 + I_3.
\end{aligned}$$

Using spherical coordinate formula, we get

$$\begin{aligned}
I_1 & \simeq \left(\log \frac{1}{\epsilon} \right)^{-2} |a_1 - a_2|^{4-N} \sum_{i=1}^2 \int_{B_\epsilon(a_i) \setminus B_{\epsilon^2}(a_i)} |x - a_i|^{4-N} \cdot \frac{(N-2)^2}{|x - a_i|^4} dx \\
& \quad + \left(\log \frac{1}{\epsilon} \right)^2 \int_{B_{\frac{1}{\epsilon^2}}(0) \setminus B_{\frac{1}{\epsilon}}(0)} |x|^{2(4-N)} \cdot \frac{(N-2)^2}{|x|^4} dx \\
& \simeq \left(\log \frac{1}{\epsilon} \right)^{-2} \int_{\epsilon^2}^\epsilon r^{-1} dr + \left(\log \frac{1}{\epsilon} \right)^2 \int_{\frac{1}{\epsilon^2}}^{\frac{1}{\epsilon}} r^{4-N-1} dr \\
& \simeq \left(\log \frac{1}{\epsilon} \right)^{-1} + \left(\frac{1}{\epsilon} \right)^{4-N} \left(\log \frac{1}{\epsilon} \right)^2 \rightarrow 0 \text{ as } \epsilon \rightarrow 0.
\end{aligned}$$

By similar computation, we have

$$\begin{aligned}
I_2 & \simeq \left(\log \frac{1}{\epsilon} \right)^{-2} \sum_{i=1}^2 \int_{B_\epsilon(a_i) \setminus B_{\epsilon^2}(a_i)} |x - a_i|^{-N} dx + \left(\log \frac{1}{\epsilon} \right)^2 \int_{B_{\frac{1}{\epsilon^2}}(0) \setminus B_{\frac{1}{\epsilon}}(0)} |x|^{4-2N} dx \\
& \rightarrow 0 \text{ as } \epsilon \rightarrow 0.
\end{aligned}$$

and

$$\begin{aligned}
I_3 & \simeq \left(\log \frac{1}{\epsilon} \right)^{-2} \sum_{i=1}^2 \int_{B_\epsilon(a_i) \setminus B_{\epsilon^2}(a_i)} |x - a_i|^{-N} dx + \left(\log \frac{1}{\epsilon} \right)^2 \int_{B_{\frac{1}{\epsilon^2}}(0) \setminus B_{\frac{1}{\epsilon}}(0)} |x|^{4-2N} dx \\
& \rightarrow 0 \text{ as } \epsilon \rightarrow 0.
\end{aligned}$$

Summing over I_1 , I_2 and I_3 yields

$$\lim_{\epsilon \rightarrow 0} \left(\int_{\mathbb{R}^N} |\phi \Delta v_\epsilon + 2\nabla \phi \cdot \nabla v_\epsilon|^2 dx - \int_{\mathbb{R}^N} \phi \Delta \phi |\nabla v_\epsilon|^2 dx \right) = 0.$$

Thus the operator H_2 is critical by Corollary 3.3, moreover, it is null-critical by Corollary 1.4.

□

References

- [1] Bosi, R., Dolbeault, J., Esteban, M.: Estimates for the optimal constants in multipolar Hardy inequalities for Schrödinger and Dirac operators. *Commun. Pur. Appl. Anal.*, 7(3), 533-562(2008).
- [2] Balinsky, A., Evans, W. D.: Some recent results on Hardy-type inequalities. *Appl. Math. Inf. Sci.*, 4(2), 191-208(2020).
- [3] Barbatis, G., Filippas S., Tertikas A.: A unified approach to improved L^p -Hardy inequalities with best constants. *Trans. Amer. Math. Soc.*, 356, 2169-2196(2004).
- [4] Bez, N., Machihara, S., Ozawa, T.: Revisiting the Rellich inequality. *Math. Z.*, 303(49), (2023).
- [5] Brezis, H., Vázquez, J. L.: Blow-up solutions of some nonlinear elliptic problems. *Rev. Mat. Univ. Complut. Madrid.*, 10, 443-469(1997).
- [6] Cazacu C.: The method of super-solutions in Hardy and Rellich type inequalities in the L^2 setting: an overview of well-known results and short proofs. DOI:10.48550/arXiv.2003.11798(2020).
- [7] Canale, A., Pappalardo, F.: Weighted Hardy inequalities and Ornstein-Uhlenbeck type operators perturbed by multipolar inverse square potentials. *J. Math. Anal. and Appl.*, 463(2), 895-909(2018).
- [8] Canale, A., Pappalardo, F., Tarantino, C.: Weighted multipolar Hardy inequalities and evolution problems with Kolmogorov operators perturbed by singular potentials. *Commun. Pur. Appl. Anal.*, 20(1): 405-425(2021).
- [9] Cazacu, C., Zuazua E.: Improved Multipolar Hardy Inequalities. *Studies in phase space analysis with applications to PDEs*, *Progr. Nonlinear Differential Equations Appl.*, Birkhäuser/Springer, New York, 84, 35-52(2013), MR 3185889.
- [10] Davies, E. B.: The Hardy constant. *Q. J. Math. Oxf.*, 46(2), 417-431(1995).
- [11] Duyckaerts T.: A singular critical potential for the Schrödinger operator. *Canad. Math. Bull.*, 50, 35-47(2007).
- [12] Devyver, B., Fraas, M., Pinchover, Y.: Optimal hardy weight for second-order elliptic operator: An answer to a problem of Agmon. *J. Funct. Anal.*, 266(7), 4422-4489 (2014).
- [13] Davies, E. B., Hinz, A. M.: Explicit constants for Rellich inequalities in $L^p(\Omega)$. *Math. Z.*, 227(3), 511-523(1998).
- [14] Devyver, B., Pinchover, Y.: Optimal L^p Hardy-type inequalities. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 33(1), 93-118(2016).

- [15] Edmunds, D. E., Evans, W. D.: The Rellich inequality. *Rev. Mat. Complut.*, 29(3), 511-530 (2016).
- [16] Evans, W. D., Lewis, R. T.: On the Rellich inequality with magnetic potentials. *Math. Z.*, 251(2), 267-284(2005).
- [17] Faraci, F., Farkas C., Kristály, A.: Multipolar Hardy inequalities on Riemannian manifolds. *Esaim Control Optimisation and Calculus of Variations*, 24(2), 551-567(2016).
- [18] Felli, V., Marchini, E. M., Terracini, S.: On Schrödinger operators with multipolar inverse-square potentials. *J. Funct. Anal.*, 250, 265-316(2007).
- [19] Felli, V., Terracini, S.: Elliptic Equations with Multi-Singular Inverse-Square Potentials and Critical Nonlinearity. *Commun. Part. Diff. Eq.*, 31(3), 469-495(2006).
- [20] Jin, Y., Shen, S.: Weighted Hardy and Rellich inequality on Carnot groups, *Arch. Math.*, 96(3), 263-271(2011).
- [21] Jin, Y., Zhang, G.: Degenerate p -Laplacian operators and Hardy type inequalities on H-type groups. *Canad. J. Math.* 2010, 62(5): 1116-1130.
- [22] Leray, J.: Sur le mouvement d'un liquide visqueux emplissant l'espace (French). *Acta Math.*, 63(1), 193-248(1934).
- [23] Machihara, S., Ozawa, T., Wadade, H.: Remarks on the Rellich inequality. *Math. Z.*, 286, 1367-1373(2017).
- [24] Rellich, F.: Halbbeschränkte Differentialoperatoren höherer Ordnung. In: Gerretsen, J.C.H., et al. (eds) *Proceedings of the International Congress of Mathematicians, Amsterdam (1954), Vol. 3, Groningen, Nordhoff*, pp. 243-250 (1956).
- [25] Tertikas, A.: Critical phenomena in linear elliptic problems. *J. Funct. Anal.*, 154, 42-66(1998).
- [26] Vazquez, J. L., Zuazua, E.: The Hardy Inequality and the Asymptotic Behaviour of the Heat Equation with an Inverse-Square Potential. *J. Funct. Anal.*, 173(1), 103-153(2000).
- [27] Yener, A., Kombe I., Goldstein, J. A.: A unified approach to weighted Hardy type inequalities on Carnot groups. *Discrete and Contin. Dynam. Systems*, 37(4), 2009-2021(2017).