The coherent rank of a graph with three eigenvalues

Gary Greaves * Jose Yip [†]

Abstract

We characterise graphs that have three distinct eigenvalues and coherent ranks 8 and 9, linking the former to certain symmetric 2-designs and the latter to specific quasisymmetric 2-designs. This characterisation leads to the discovery of a new biregular graph with three distinct eigenvalues. Additionally, we demonstrate that the coherent rank of a triregular graph with three distinct eigenvalues is at least 14. Finally, we introduce a conjecturally infinite family of biregular graphs with three distinct eigenvalues, obtained by switching the block graphs of orthogonal arrays.

1 Introduction

Strongly regular graphs are a fundamental object of study in algebraic graph theory. These graphs can be characterised by their spectrum: connected strongly regular graphs are precisely the connected regular graphs that have precisely three distinct eigenvalues. Thus, graphs with exactly three distinct eigenvalues can be viewed as a generalisation of strongly regular graphs.

At the British Combinatorial Conference in 1995, Willem Haemers posed a question: apart from complete bipartite graphs, which non-regular graphs have precisely three distinct eigenvalues? This prompted seminal work by Muzychuk and Klin [23] and Van Dam [12], who each produced infinite families of non-regular graphs with this property. These families include graphs with vertices having exactly two distinct degrees, as well as a finite number of examples with three distinct degrees, such as the multiplicative cones discovered by Bridges and Mena [1] in 1981.

Let \mathscr{G}_3 denote the set of all connected graphs having precisely three distinct eigenvalues. Several attempts have been made to systematically classify the graphs in \mathscr{G}_3 :

- Van Dam [12]: graphs in \mathscr{G}_3 with at most 29 vertices;
- Chuang and Omidi [9]: graphs in \mathscr{G}_3 whose largest eigenvalue is less than 8;
- Van Dam [12] again: graphs in \mathscr{G}_3 whose smallest eigenvalue is at least -2;
- Cheng et al. [8]: graphs in \mathcal{G}_3 whose second largest eigenvalue is at most 1.

These classifications did not yield any graphs in \mathscr{G}_3 with more than three distinct degrees. In fact, the following question by De Caen [4] and [13, Problem 9] remains open:

^{*}School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, Singapore 637371, gary@ntu.edu.sg

[†]School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, Singapore 637371, josezhengho.yip@ntu.edu.sg

Question 1.1. Must each graph in \mathscr{G}_3 have at most three distinct degrees?

Van Dam et al. [14] demonstrated that the condition of having three distinct eigenvalues in Question 1.1 is critical, showing that there exist connected graphs with four or five distinct eigenvalues and arbitrarily many valencies.

Muzychuk and Klin [23] approached the problem of characterising of \mathscr{G}_3 graphs by examining the rank of the Weisfeiler-Leman closure, which we refer to as the coherent rank. Strongly regular graphs can be characterised as graphs whose coherent rank is equal to 3. Loosely speaking, the coherent rank can be thought of as a measure of how close a graph is to being strongly regular. Muzychuk and Klin characterised the graphs in \mathscr{G}_3 that have a coherent rank of at most 6 but claimed without proof [23, Proposition 6.2] that no graph in \mathscr{G}_3 has a coherent rank of 7 or 8. Our first contribution is to prove that no graph in \mathscr{G}_3 has a coherent rank of 7. However, we show that there do exist graphs in \mathscr{G}_3 with a coherent rank of 8, and we characterise such graphs in Theorem 4.3, showing that they correspond to certain symmetric 2-designs.

We continue the characterisation program of Muzychuk and Klin by characterising graphs in \mathscr{G}_3 with coherent rank 9. One consequence is the discovery of a new biregular graph in \mathscr{G}_3 (see Example 5.13). In fact, it turns out that graphs in \mathscr{G}_3 with coherent rank 9 correspond to certain quasi-symmetric designs (see Section 5). Quasi-symmetric designs, particularly affine designs [12], have been used to construct an infinite family of graphs in \mathscr{G}_3 . Rowlinson [26] asked whether graphs in \mathscr{G}_3 can be constructed from quasi-symmetric designs that are not of affine type. We answer Rowlinson's question in the affirmative and characterise the quasi-symmetric designs that correspond to graphs in \mathscr{G}_3 with coherent rank 9.

Our next contribution is a lower bound for the coherent rank of a graph in \mathscr{G}_3 that has three distinct degrees. We show that if a graph in \mathscr{G}_3 has three distinct degrees, then its coherent rank is at least 14 (see Theorem 6.1). This bound has the potential to be sharp contingent on the existence of a certain quasi-symmetric design (see Example 6.6).

All currently known infinite families of graphs in \mathscr{G}_3 have a fixed coherent rank. Our final contribution is a conjecturally infinite construction of biregular graphs in \mathscr{G}_3 with conjecturally arbitrarily large coherent ranks. Each graph in this family corresponds to a prime power in a certain recurrence sequence and is obtained by switching the block graph of an orthogonal array (see Section 7).

All graphs in this paper are finite, undirected, and simple. The **spectrum** and **eigen-values** of a graph are defined to be those of its adjacency matrix. The paper is organised as follows. In Section 2, we develop the foundational theory of graphs with three distinct eigenvalues. In Section 3, we introduce the coherent closure and coherent rank of a graph and present Muzychuk and Klin's characterisation of graphs in \mathscr{G}_3 whose coherent rank is small. In Section 4, we characterise the graphs in \mathscr{G}_3 that have coherent rank 8. In Section 5, we characterise the graphs in \mathscr{G}_3 that have coherent rank 9. In Section 6, we prove a lower bound for the coherent rank of a triregular graph in \mathscr{G}_3 . In Section 7, we exhibit a new family of biregular graphs in \mathscr{G}_3 obtained from switching the block graphs of orthogonal arrays. Finally, in Section 8, we provide a selection of open problems that emerged during our investigations.

2 Graphs with three distinct eigenvalues

In this section, we introduce some basic properties of graphs in \mathscr{G}_3 and tools, which we will use to establish our results, below.

2.1 Fundamentals

We denote by I_n , J_n , O_n , and $\mathbf{1}_n$, the identity matrix, all-ones matrix, zero matrix, and all-ones (column) vector of order n, respectively. We merely write I, J, O, and $\mathbf{1}$ when the order can be determined from context, or in the case of J and O when the matrix is not square. A graph (V, E) is called **empty** or **complete** if $E = \emptyset$ or $E = \binom{V}{2}$, respectively. Denote by K_v the complete graph on v vertices. Suppose that Γ has degree sequence $\{[k_1]^{n_1}, [k_2]^{n_2}, \ldots, [k_d]^{n_d}\}$. If d = 1 then each vertex has k_1 neighbours and Γ is called **regular**. In the cases when d = 2 or d = 3, we call Γ **biregular** or **triregular**, respectively.

For fixed $\theta_0 > \theta_1 > \theta_2$, define $\mathscr{G}_3(\theta_0, \theta_1, \theta_2) \subset \mathscr{G}_3$ as the subset of connected graphs with eigenvalues θ_0 , θ_1 , and θ_2 . Let Γ be a graph with r + 1 distinct eigenvalues $\theta_0 > \theta_1 > \cdots > \theta_r$. The spectrum of Γ is denoted by spec(Γ) and is written as spec(Γ) = $\{[\theta_0]^{m_0}, [\theta_1]^{m_1}, \ldots, [\theta_r]^{m_r}\}$, where m_0, m_1, \ldots, m_r are the multiplicities of $\theta_0, \theta_1, \ldots, \theta_r$ respectively. By the Perron-Frobenius Theorem [2, Theorem 2.2.1], a graph $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ has spectrum $\{[\theta_0]^1, [\theta_1]^{m_1}, [\theta_2]^{m_2}\}$, where m_1 and m_2 are positive integers whose sum is 1 less than the order of Γ . Let $A(\Gamma)$ denote the adjacency matrix of a graph Γ , and $V(\Gamma)$ its vertex set. Again, by the Perron-Frobenius Theorem, if $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$, then

$$A(\Gamma)^2 - (\theta_1 + \theta_2)A(\Gamma) + \theta_1\theta_2 I = \alpha \alpha^{\top}, \qquad (1)$$

where $\boldsymbol{\alpha} = (\alpha_{\mathsf{x}})_{\mathsf{x} \in V(\Gamma)}$ is a positive (column) eigenvector for θ_0 . Conversely, if a connected graph Γ satisfies (1) for some θ_1 , θ_2 , and $\boldsymbol{\alpha}$, then $\Gamma \in \mathscr{G}_3$. The diagonal entries of (1) show that the degree d_{x} of vertex x is $\alpha_{\mathsf{x}}^2 - \theta_1 \theta_2$. Alternatively, for each $\mathsf{x} \in V(\Gamma)$, we can write $\alpha_{\mathsf{x}} = \sqrt{d_{\mathsf{x}} + \theta_1 \theta_2}$. Moreover, (1) implies any graph in \mathscr{G}_3 must have diameter 2.

The following theorem due to Cauchy [2, Theorem 2.5.1], will be used repeatedly, throughout.

Theorem 2.1 (Eigenvalue Interlacing Theorem). Let A be a real symmetric $n \times n$ matrix, and B be an $m \times m$ principal submatrix of A. Suppose A has eigenvalues $\lambda_1 \leq \cdots \leq \lambda_n$, and B has eigenvalues $\mu_1 \leq \cdots \leq \mu_m$. Then

$$\lambda_k \leqslant \mu_k \leqslant \lambda_{n-m+k}.$$

For a square matrix M and U a subset of its rows, denote by M[U] the principal submatrix of M induced on the rows and columns corresponding to U. Let $\Gamma = (V, E)$ be a graph. Similarly, for a subset $U \subset V$, denote by $\Gamma[U]$ the subgraph of Γ induced on the vertices in U. Thus, $A(\Gamma[U]) = A(\Gamma)[U]$.

Suppose $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$. Then Γ is connected and not complete. Hence, Γ must contain the path graph $P_3 = K_{1,2}$ as an induced subgraph. Since $P_3 \in \mathscr{G}_3(\sqrt{2}, 0, -\sqrt{2})$, by Theorem 2.1, we must have

$$\theta_1 \ge 0 \text{ and } \theta_2 \leqslant -\sqrt{2}.$$
 (2)

Regular graphs in \mathscr{G}_3 are known as *strongly regular graphs*. A **strongly regular graph** Γ with parameters (v, k, a, c) is a k-regular graph on v vertices such that each pair of adjacent

vertices in $V(\Gamma)$ has precisely *a* common neighbours and each pair of non-adjacent vertices has precisely *c* common neighbours. Denote by $\operatorname{SRG}(v, k, a, c)$ the set of all strongly regular graphs that have parameters (v, k, a, c). If $A = A(\Gamma)$ then the (x, y)-entry of A^2 is equal to the number of 2-walks from x to y. Thus, if $\Gamma \in \operatorname{SRG}(v, k, a, c)$ then

$$A^{2} = kI + aA + c(J - I - A).$$

Furthermore, if Γ is connected then

$$\Gamma \in \mathscr{G}_3\left(k, \frac{a-c+\sqrt{(a-c)^2+4(k-c)}}{2}, \frac{a-c-\sqrt{(a-c)^2+4(k-c)}}{2}\right)$$

Conversely, it also follows that if $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ is regular with v vertices then

$$\Gamma \in \operatorname{SRG}(v, \theta_0, \theta_0 - 1 + (\theta_1 + 1)(\theta_2 + 1), \theta_0 + \theta_1 \theta_2).$$

2.2 Cones and bipartite graphs

A complete bipartite graph $K_{a,b}$ belongs to the set $\mathscr{G}_3(-\sqrt{ab}, 0, \sqrt{ab})$. Moreover, $K_{a,b}$ has spectrum $\left\{ [-\sqrt{ab}]^1, [0]^{a+b-2}, [\sqrt{ab}]^1 \right\}$. The **cone over** a graph Γ is formed by adjoining a vertex adjacent to every vertex of Γ . Muzychuk and Klin [23] showed that if Γ is a *v*-vertex, *k*-regular strongly regular graph with smallest eigenvalue θ_2 then the cone over Γ is in \mathscr{G}_3 if and only if $v = \theta_2(\theta_2 - k)$.

The **complement** of a graph $\Gamma = (V, E)$ is defined to be the graph $(V, {V \choose 2} - E)$ and is denoted by $\overline{\Gamma}$. The following theorem is a useful characterisation of cones and complete bipartite graphs.

Theorem 2.2 ([7, Theorem 2.2]). Let $\Gamma \in \mathscr{G}_3$ be a non-regular graph whose complement is disconnected. Then Γ is a cone or complete bipartite.

Next, Van Dam [12, Section 4.2] answered Question 1.1 in the affirmative for cones.

Lemma 2.3 ([12, Section 4.2]). Let Γ be a cone in \mathscr{G}_3 . Then either Γ is a cone over a strongly regular graph or Γ is triregular. Furthermore, if $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ is triregular with distinct degrees $k_1 > k_2 > k_3$ then $\sqrt{k_1 + \theta_1 \theta_2} = \sqrt{k_2 + \theta_1 \theta_2} + \sqrt{k_3 + \theta_1 \theta_2}$.

Two examples of triregular cones in \mathscr{G}_3 were discovered by Bridges and Mena [1]. See Table 3, below. In fact, Bridges and Mena [1] classified the triregular cones in \mathscr{G}_3 whose smallest two eigenvalues θ_1 and θ_2 satisfy $\theta_1 + \theta_2 = 0$. There are potentially three such graphs, with the existence of the third possibility still an open problem (see [7, Section 9]).

Van Dam [12, Section 4.2] showed that biregular cones in $\Gamma \in \mathscr{G}_3$ are precisely cones over strongly regular graphs. We record this result as the following lemma.

Lemma 2.4. Let $\Gamma \in \mathscr{G}_3$ be a biregular cone. Then Γ is a cone over a strongly regular graph.

Non-regular complete bipartite graphs can be characterised as the non-regular graphs in \mathscr{G}_3 with second largest eigenvalue 0.

Theorem 2.5 ([7, Corollary 2.3]). Let $\Gamma \in \mathscr{G}_3$ be a non-regular graph. Then the following are equivalent.

- (i) Γ is bipartite;
- (ii) Γ is complete bipartite;

(*iii*) $\theta_1 = 0$.

2.3 Valencies and equitable partitions

Let $\Gamma = (V, E)$ be a graph and let $\pi = \{\pi_1, \ldots, \pi_t\}$ be a partition of the vertex set V. We call π **equitable** if, for each $i, j \in \{1, \ldots, t\}$ there exists n_{ij} such that each vertex in π_i has n_{ij} neighbours in π_j . Denote by $Q_{\pi}(\Gamma)$ the **quotient matrix** $(n_{ij})_{i,j \in \{1,\ldots,t\}}$ that corresponds to the vertex partition π .

Lemma 2.6 ([2, Lemma 2.3.1]). Let Γ be a graph with equitable partition π of $V(\Gamma)$. Then $\operatorname{spec}(Q_{\pi}(\Gamma)) \subset \operatorname{spec}(\Gamma)$.

Suppose Γ has degree sequence $\{[k_1]^{n_1}, [k_2]^{n_2}, \ldots, [k_d]^{n_d}\}$. For each $i \in \{1, \ldots, d\}$, let V_i denote the subset of vertices whose degree is k_i . Clearly, $\{V_1, \ldots, V_d\}$ is a partition of $V(\Gamma)$. We call this partition the **valency partition** of Γ and we denote it by $\mathfrak{D}(\Gamma)$.

Theorem 2.7 ([12, Section 4]). Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ have at most three distinct valencies. Then the valency partition is equitable. Furthermore, θ_0 is an eigenvalue of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$.

Biregular, non-bipartite graphs in \mathscr{G}_3 are subject to stronger conditions, which we list in Theorem 2.8.

Theorem 2.8 ([7, Theorem 4.3]). Let Γ be a non-bipartite biregular graph in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with degree sequence $\{[k_1]^{n_1}, [k_2]^{n_2}\}$ and $\mathfrak{D}(\Gamma) = \{V_1, V_2\}$. Then the following conditions hold:

- (i) All eigenvalues of Γ are integers.
- (ii) The quotient matrix $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$ has eigenvalues θ_0 and θ , where $\theta \in \{\theta_1, \theta_2\}$.

(*iii*)
$$\sqrt{(k_1 + \theta_1 \theta_2)(k_2 + \theta_1 \theta_2)} = -\theta(\theta' + 1)$$
 where $\{\theta, \theta'\} = \{\theta_1, \theta_2\}$.

(iv) If $\Gamma[V_1]$ or $\Gamma[V_2]$ is empty then $\theta = \theta_2$.

3 Coherent configurations and algebras

In this section, we define the coherent rank of a graph.

3.1 The coherent rank

Let \mathfrak{X} be a finite set and let $\mathfrak{R} = \{R_1, \ldots, R_r\}$ be a set of binary relations on \mathfrak{X} . For each R_i let $A_i \in \operatorname{Mat}_{\mathfrak{X}}(\{0,1\})$ be defined such that its (x,y) entry is 1 if $(\mathsf{x},\mathsf{y}) \in R_i$ and 0 otherwise. Suppose that

$$(\text{CC1}) \ \sum_{i=1}^{r} A_i = J;$$

(CC2) For each $i \in \{1, \ldots, r\}$ there exists $j \in \{1, \ldots, r\}$ such that $A_i^{\mathsf{T}} = A_j$;

(CC3) There exists a subset $\Delta \subset \{1, \ldots, r\}$ such that $\sum_{i \in \Delta} A_i = I$;

(CC4)
$$A_i A_j = \sum_{k=1}^r p_{i,j}^k A_k$$
, for each $i, j \in \{1, \dots, r\}$.

Then $(\mathfrak{X}, \mathfrak{R})$ is called a **coherent configuration** of **rank** $r = |\mathfrak{R}|$. The set \mathfrak{X} is called the **point-set** of the coherent configuration.

For each $i \in \Delta$, we call the subset $\mathfrak{X}_i := \{\mathbf{x} \in \mathfrak{X} : (\mathbf{x}, \mathbf{x}) \in R_i\}$ a fibre of the coherent configuration. Clearly, the fibres form a partition of the point-set \mathfrak{X} . When $|\Delta| = 1$, the coherent configuration $(\mathfrak{X}, \mathfrak{R})$ is called an **association scheme**. It follows from (CC4) that, for each $k \in \{1, \ldots, r\}$, there exists *i* and *j* such that $R_k \subset \mathfrak{X}_i \times \mathfrak{X}_j$. Thus, each subset Δ' of Δ induces a coherent configuration with point-set $\bigcup_{i \in \Delta'} \mathfrak{X}_i$. The **type** of $(\mathfrak{X}, \mathfrak{R})$ is defined to be the matrix in $\operatorname{Mat}_{\Delta}(\mathbb{N})$ whose (i, j)-entry t_{ij} is equal to the cardinality $|\{k : R_k \subset \mathfrak{X}_i \times \mathfrak{X}_j\}|$. Note that the sum of the entries of the type matrix is equal to *r*. Furthermore, since the type matrix must be symmetric, we omit the entries below the diagonal. Higman [19] established the following restriction on the type matrix.

Lemma 3.1. For each $i, j \in \Delta$, if $t_{ii} \leq 5$ and $t_{jj} \leq 5$ then $t_{ij} \leq \min(t_{ii}, t_{jj})$.

A coherent algebra is a matrix algebra $\mathcal{A} \subset \operatorname{Mat}_{\mathfrak{X}}(\mathbb{C})$ that satisfies the following axioms.

- (A1) $I, J \in \mathcal{A};$
- (A2) $M^{\mathsf{T}} \in \mathcal{A}$ for each $M \in \mathcal{A}$;
- (A3) $MN \in \mathcal{A}$ and $M \circ N \in \mathcal{A}$ for each $M, N \in \mathcal{A}$, where \circ denotes the entrywise product.

Each coherent algebra \mathcal{A} has a unique basis of $\{0, 1\}$ -matrices $\{A_1, \ldots, A_r\}$ that corresponds to a coherent configuration $(\mathfrak{X}_{\mathcal{A}}, \mathfrak{R}_{\mathcal{A}})$. We denote by $\mathfrak{F}_{\mathcal{A}}$ the set of fibres of the coherent configuration $(\mathfrak{X}_{\mathcal{A}}, \mathfrak{R}_{\mathcal{A}})$ and we define **type** of \mathcal{A} to be that of $(\mathfrak{X}_{\mathcal{A}}, \mathfrak{R}_{\mathcal{A}})$. Clearly, the intersection of any two coherent algebras is itself a coherent algebra. We can thus define the **coherent closure** $\mathcal{W}(\Gamma)$ of Γ to be the minimal coherent algebra that contains the adjacency matrix $\mathcal{A}(\Gamma)$ of Γ . We write $\mathcal{W}(\Gamma) = \langle A_1, \ldots, A_r \rangle$, where $\{A_1, \ldots, A_r\}$ is the unique basis of $\{0, 1\}$ -matrices for $\mathcal{W}(\Gamma)$.

We summarise in the following corollary some straightforward consequences of the above definitions.

Corollary 3.2. Let Γ be a graph with coherent closure $\mathcal{W}(\Gamma) = \langle A_1, \ldots, A_r \rangle$. Then

- (i) For each $\mathfrak{f} \in \mathfrak{F}_{\mathcal{W}(\Gamma)}$, the matrix $A(\Gamma)[\mathfrak{f}]$ belongs to the coherent algebra $\langle A_1[\mathfrak{f}], \ldots, A_r[\mathfrak{f}] \rangle$.
- (ii) If x and y are in the same fibre of $\mathcal{W}(\Gamma)$ then x and y have the same degree.
- (iii) $|\mathfrak{D}(\Gamma)| \leq |\mathfrak{F}_{W(\Gamma)}|$. Furthermore, in the case of equality, we have $\mathfrak{D}(\Gamma) = \mathfrak{F}_{W(\Gamma)}$.

Define the **coherent rank** of Γ to be the rank of its coherent closure $\mathcal{W}(\Gamma)$. The coherent rank of a complete or empty graph on at least 2 vertices is 2 and the coherent rank of a strongly regular graph is 3. Conversely, it is straightforward to verify the following proposition.

Proposition 3.3. Let \mathcal{A} be a coherent algebra of rank r.

- (i) If r = 1 then $\mathcal{A} = \langle I_1 \rangle$.
- (ii) If r = 2 then $\mathcal{A} = \langle I, J I \rangle$.
- (iii) If r = 3 and A contains the adjacency matrix of a graph that is neither empty nor complete then $A = \langle I, A, J I A \rangle$, where A is the adjacency matrix of a strongly regular graph.

3.2 Small coherent rank

Muzychuk and Klin [23] initiated the study of graphs with three eigenvalues that have small coherent rank. We summarise their findings in the following theorem.

Theorem 3.4 ([23, Section 6]). Let $\Gamma \in \mathscr{G}_3$. Then the coherent rank of Γ is

- (i) 3 if and only if Γ is a strongly regular graph;
- (ii) 5 if and only if $\Gamma \cong K_{1,\mathbf{b}}$ where $\mathbf{b} \ge 2$;
- (iii) 6 if and only if

 $- \Gamma \cong K_{\mathsf{a},\mathsf{b}}$ where $2 \leq \mathsf{a} < \mathsf{b}$; or

 $-\Gamma$ is a cone over a strongly regular graph.

Furthermore, the coherent rank of Γ cannot be 4.

Remark 3.5. Note that if we allow disconnected graphs then coherent rank 4 is possible. For example, take the disjoint union of two copies of a strongly regular graph.

In each case of Theorem 3.4 the coherent closure of Γ has type [3], $\begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$, or $\begin{bmatrix} 2 & 1 \\ 2 & 2 \end{bmatrix}$, and $\begin{bmatrix} 1 & 1 \\ 3 & 3 \end{bmatrix}$, respectively. Muzychuk and Klin [23, Proposition 6.2] further claimed without proof that a graph with three distinct eigenvalues cannot have coherent rank 7 or 8. Contrary to their claim, we will show that coherent rank 8 is in fact possible. First, we give a proof that a graph with three distinct eigenvalues cannot have coherent rank 7.

Theorem 3.6. Let $\Gamma \in \mathscr{G}_3$. Then the coherent rank of Γ is not 7.

Proof. Suppose for a contradiction that $\Gamma \in \mathscr{G}_3$ be a graph with coherent rank 7 and let A be its adjacency matrix. By Theorem 3.4, the graph Γ is not regular. Using Lemma 3.1 and Corollary 3.2 (iii), we find that the coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 1 & 1 \\ 4 \end{bmatrix}$ or $\begin{bmatrix} 2 & 1 \\ 3 \end{bmatrix}$. By Corollary 3.2 (iii), the vertices of Γ have two distinct degrees k_1 and k_2 (say). Furthermore, since $A \in \mathcal{W}(\Gamma)$ and Γ is connected, it follows that each vertex of degree k_1 must be adjacent to every vertex of degree k_2 . Whence, the complement of Γ is disconnected. Thus, by Theorem 2.2 and Lemma 2.4 the graph Γ is a cone over a strongly regular graph. But, by Theorem 3.4, such a graph has coherent rank 6, which is a contradiction.

We conclude this section with a corollary of Theorem 3.4 and Theorem 2.5.

Corollary 3.7. Let $\Gamma \in \mathscr{G}_3(\theta_0, 0, \theta_2)$. Then the coherent rank of Γ is at most 6.

4 Coherent rank 8

In this section, we characterise the graphs $\Gamma \in \mathscr{G}_3$ that have coherent rank 8.

4.1 Symmetric 2-designs

A design is a pair $(\mathcal{P}, \mathcal{B})$ consisting of **point-set** \mathcal{P} with cardinality $|\mathcal{P}| = v$ and \mathcal{B} a family of k-sets from $\binom{\mathcal{P}}{k}$ for some $k \in \{1, \ldots, v\}$.¹ Accordingly, the elements of \mathcal{P} and \mathcal{B} are called **points** and **blocks** respectively. If there exists t and λ such that each set in $\binom{\mathcal{P}}{t}$ is contained in precisely λ blocks then $(\mathcal{P}, \mathcal{B})$ is called a t- (v, k, λ) design. We may write t-design in place of t- (v, k, λ) design if the parameters are not required to be specified. Suppose that every pair of blocks intersects in x points. Then the design $(\mathcal{P}, \mathcal{B})$ is called a symmetric design.

Let $(\mathcal{P}, \mathcal{B})$ be a 2- (v, k, λ) design. Write $b = |\mathcal{B}|$ for the number of blocks and r for the number of blocks that contain a particular point. Standard double-counting arguments yield the following equations.

$$vr = bk; (3)$$

$$\lambda(v-1) = r(k-1). \tag{4}$$

We also have the fundamental inequality due to Fisher [15]:

$$b \geqslant v.$$
 (5)

The incidence matrix of $(\mathcal{P}, \mathcal{B})$ is a $v \times b$ matrix whose (i, j)-entry is equal to 1 if $i \in j$ and 0 otherwise. Let M be the incidence matrix for $(\mathcal{P}, \mathcal{B})$. Then M satisfies MJ = rJ, $M^{\mathsf{T}}J = kJ$, and

$$MM^{\mathsf{T}} = rI + \lambda(J - I). \tag{6}$$

Furthermore, if $(\mathcal{P}, \mathcal{B})$ is a symmetric design then

$$M^{\mathsf{T}}M = kI + x(J - I) \tag{7}$$

Note that symmetric designs correspond to equality in Fisher's inequality (5). Moreover, we obtain

$$\lambda(v-1) = k(k-1). \tag{8}$$

Now we state a tool that we will require below.

Lemma 4.1 ([2, Lemma 2.9.2]). Let M be an $n \times m$ real matrix. Then $M^{\mathsf{T}}M$ and MM^{T} have the same nonzero eigenvalues (including multiplicities).

¹We allow the case of repeated blocks, although we will show that only designs without repeated blocks are pertinent in our study. (See Corollary 5.8).

4.2 Graphs with three eigenvalues and coherent rank 8

Next, we show that graphs in \mathscr{G}_3 whose coherent closure has type $\begin{bmatrix} 2 & 2 \\ 2 \end{bmatrix}$ correspond to certain symmetric designs.

Proposition 4.2. Let $\Gamma \in \mathscr{G}_3$ such that $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$. Then the adjacency matrix of Γ has the form $\begin{bmatrix} O & M \\ M^{\intercal} & J-I \end{bmatrix}$ where M is the incidence matrix of a symmetric 2- $(\lambda^3 - \lambda + 1, \lambda^2, \lambda)$ design for some $\lambda \in \mathbb{N}$.

Proof. Suppose $A = A(\Gamma)$ is the adjacency matrix of Γ . By Theorem 3.4 and Corollary 3.2 (iii), the graph Γ must be biregular with degrees k_1 and k_2 (say). Suppose $\mathfrak{D}(\Gamma) = \{V_1, V_2\}$. By Corollary 3.2 (iii), we can write $A = \begin{bmatrix} A(\Gamma[V_1]) & M \\ M^{\intercal} & A(\Gamma[V_2]) \end{bmatrix}$ and

$$A^2 = \begin{bmatrix} A(\Gamma[V_1])^2 + MM^{\mathsf{T}} & A(\Gamma[V_1])M + MA(\Gamma[V_2]) \\ M^{\mathsf{T}}A(\Gamma[V_1]) + A(\Gamma[V_2])M^{\mathsf{T}} & M^{\mathsf{T}}M + A(\Gamma[V_2])^2 \end{bmatrix}$$

Apply (1) to obtain

$$A^{2} - (\theta_{1} + \theta_{2})A + \theta_{1}\theta_{2}I = \begin{bmatrix} (k_{1} + \theta_{1}\theta_{2})J & \sqrt{(k_{1} + \theta_{1}\theta_{2})(k_{2} + \theta_{1}\theta_{2})}J \\ \sqrt{(k_{1} + \theta_{1}\theta_{2})(k_{2} + \theta_{1}\theta_{2})}J & (k_{2} + \theta_{1}\theta_{2})J \end{bmatrix}.$$
 (9)

By Corollary 3.2 (i) and Proposition 3.3, the subgraph $\Gamma[V_1] = K_{n_1}$ or \overline{K}_{n_1} , and the subgraph $\Gamma[V_2] = K_{n_2}$ or \overline{K}_{n_2} , that is, the matrix $A(\Gamma[V_1]) = \varepsilon_{11}(J-I)$ and the matrix $A(\Gamma[V_2]) = \varepsilon_{22}(J-I)$ for some $\varepsilon_{11}, \varepsilon_{22} \in \{0, 1\}$. From the top left block of (9), we obtain

$$MM^{\mathsf{T}} = \begin{cases} (k_{12} + \theta_1 \theta_2) J_{n_1} - \theta_1 \theta_2 I, & \text{if } \Gamma[V_1] = \overline{K}_{n_1}; \\ (k_{12} + (\theta_1 + 1)(\theta_2 + 1)) J_{n_1} - (\theta_1 + 1)(\theta_2 + 1)I, & \text{if } \Gamma[V_1] = K_{n_1}. \end{cases}$$
(10)

Similar to the above, we can use the bottom-right block of (9) to deduce that

$$M^{\mathsf{T}}M = \begin{cases} (k_{21} + \theta_1 \theta_2) J_{n_2} - \theta_1 \theta_2 I, & \text{if } \Gamma[V_2] = \overline{K}_{n_2}; \\ (k_{21} + (\theta_1 + 1)(\theta_2 + 1)) J_{n_2} - (\theta_1 + 1)(\theta_2 + 1)I, & \text{if } \Gamma[V_2] = K_{n_2}. \end{cases}$$
(11)

Next, we show that, without loss of generality, we can assume that $\Gamma[V_1]$ is empty and $\Gamma[V_2]$ is complete. By Theorem 2.5, $\Gamma[V_1]$ and $\Gamma[V_2]$ cannot both be empty. We further claim that $\Gamma[V_1]$ and $\Gamma[V_2]$ cannot both be complete. Suppose (for a contradiction) that both $\Gamma[V_1]$ and $\Gamma[V_2]$ are complete. Using (10) and (11) together with Lemma 4.1, we find that $n_1 = n_2$ and $k_{12} = k_{21}$. This implies that Γ is regular, which gives a contradiction.

Hence, we may assume that $\Gamma[V_1]$ is empty and $\Gamma[V_2]$ is complete. Apply Lemma 4.1 to find that $\theta_1\theta_2 = (\theta_1 + 1)(\theta_2 + 1)$. This implies that $\theta_1 + \theta_2 = -1$. Furthermore, M must be the incidence matrix of a symmetric 2-design \mathcal{D} . Suppose \mathcal{D} has parameters (v, k, λ) . Then $v = b = n_1 = n_2$ and $k = r = \lambda(v - 1)/(k - 1)$.

Now, we can write

$$A^{2} = \begin{bmatrix} (k-\lambda)I + \lambda J & kJ - M \\ kJ - M^{\mathsf{T}} & (k-\lambda+1)I + (v+\lambda-2)J \end{bmatrix}$$

Observe that the off-diagonal entries and the diagonal entries of the diagonal blocks of (9) are equal. Equating the diagonal and off-diagonal entries of the top-left block yields $\theta_1\theta_2 = \lambda - k$. Lastly, we use the equation

$$(A - \theta_0 I)(A - \theta_1 I)(A - \theta_2 I) = O.$$
(12)

The top-left block of A^3 is equal to $(k^2 - \lambda)J + (\lambda - k)I$. Thus, the top-left block of (12) is

$$(k^2 - \lambda)J + (\lambda - k)I - (\theta_0 + \theta_1 + \theta_2)((k - \lambda)I + \lambda J) - \theta_0\theta_1\theta_2I = O.$$

Using $\theta_1 + \theta_2 = -1$ together with the off-diagonal entries of the above yields $\theta_0 = k^2/\lambda$. The top-right block of A^3 is equal to $(k - \lambda + 1)M + k(\lambda + v - 2)J$. Thus, the top-right block of (12) is

$$(k - \lambda + 1)M + k(\lambda + v - 2)J - (\theta_0 + \theta_1 + \theta_2)(kJ - M) + (\theta_0\theta_1 + \theta_0\theta_2 + \theta_1\theta_2)M = O.$$

Combining the coefficients of J yields $k(\lambda+v-2) = k(\theta_0-1)$, which simplifies to the equation $\theta_0 = v + \lambda - 1$. The two expressions for θ_0 combined with the equation $\lambda(v-1) = k(k-1)$ yields $k = \lambda^2$. Furthermore, $v = \lambda^3 - \lambda + 1$.

We can strengthen the above result to show that if a graph in \mathscr{G}_3 has coherent rank 8 then its coherent closure has type $\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$.

Theorem 4.3. Let $\Gamma \in \mathscr{G}_3$ with coherent rank 8. Then the adjacency matrix of Γ has the form $\begin{bmatrix} O & M \\ M^{\mathsf{T}} & J-I \end{bmatrix}$ where M is the incidence matrix of a symmetric $2 \cdot (\lambda^3 - \lambda + 1, \lambda^2, \lambda)$ design for some $\lambda \in \mathbb{N}$.

Proof. By Theorem 3.4, the graph Γ is not regular. If $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & 2 \\ 2 \end{bmatrix}$ then the conclusion follows from Proposition 4.2. Otherwise, by Lemma 3.1 and Corollary 3.2 (iii), the coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 1 & 1 \\ 5 \end{bmatrix}$, $\begin{bmatrix} 2 & 4 \\ 4 \end{bmatrix}$, or $\begin{bmatrix} 3 & 3 \\ 3 \end{bmatrix}$. In each case, the complement of Γ is disconnected. By Theorem 2.2 together with Lemma 2.4 and Theorem 3.4, such a graph has coherent rank at most 6, a contradiction.

Example 4.4. Van Dam [12, Section 2.3] provided an infinite family of graphs corresponding to Theorem 4.3.

The proof of the next proposition can be obtained by the same techniques used in the proof of Proposition 4.2.

Proposition 4.5. Let $\Gamma \in \mathscr{G}_3$ such that $A(\Gamma) = \begin{bmatrix} \varepsilon_1(J-I) & M \\ M^{\mathsf{T}} & \varepsilon_2(J-I) \end{bmatrix}$ for some matrix M with $M\mathbf{1} = r\mathbf{1}, M^{\mathsf{T}}\mathbf{1} = k\mathbf{1}, \text{ and } \varepsilon_1, \varepsilon_2 \in \{0, 1\}$. Then the coherent rank of Γ is at most 8.

Proof. Suppose that M is an $n_1 \times n_2$ matrix. Clearly, Γ has at most two distinct degrees. If Γ is regular then Γ has coherent rank 3. Otherwise, we can assume that Γ is biregular with degrees k_1 and k_2 (say). The top-left block of (1) yields

$$MM^{\mathsf{T}} = (k_1 + \theta_1 \theta_2 - \varepsilon_1 (n_1 - 2 - \theta_1 - \theta_2))J - (\varepsilon_1 (1 + \theta_1 + \theta_2) + \theta_1 \theta_2)I.$$

The bottom-right block of (1) yields

$$M^{\mathsf{T}}M = (k_2 + \theta_1\theta_2 - \varepsilon_2(n_2 - 2 - \theta_1 - \theta_2))J - (\varepsilon_2(1 + \theta_1 + \theta_2) + \theta_1\theta_2)I.$$

Hence, the coherent closure $\mathcal{W}(\Gamma)$ must be a subalgebra of the coherent algebra

 $\left\langle \begin{bmatrix} I & O \\ O & O \end{bmatrix}, \begin{bmatrix} J-I & O \\ O & O \end{bmatrix}, \begin{bmatrix} O & M \\ O & O \end{bmatrix}, \begin{bmatrix} O & J-M \\ O & O \end{bmatrix}, \begin{bmatrix} O & O \\ M^{\mathsf{T}} & O \end{bmatrix}, \begin{bmatrix} O & O \\ J-M^{\mathsf{T}} & O \end{bmatrix}, \begin{bmatrix} O & O \\ O & I \end{bmatrix}, \begin{bmatrix} O & O \\ O & J-I \end{bmatrix} \right\rangle,$

which has rank at most 8.

5 Coherent rank 9

In this section, we characterise the graphs $\Gamma \in \mathscr{G}_3$ that have coherent rank 9.

5.1 The total graph and whole graph of a QSD

Let $(\mathcal{P}, \mathcal{B})$ be a 2- (v, k, λ) design. Suppose that every pair of blocks in \mathcal{B} intersects in either x or y points and both occur. If $x \neq y$ then $(\mathcal{P}, \mathcal{B})$ is called a **quasi-symmetric** design. The numbers x and y are called **intersection numbers**. Let M be the incidence matrix of a quasi-symmetric 2- (v, k, λ) design \mathcal{Q} with intersection numbers x and y. Then

$$M^{\mathsf{T}}M = kI + xX + y(J - I - X), \tag{13}$$

where X is the $\{0, 1\}$ -matrix indexed by \mathcal{B} whose (i, j)-entry is 1 precisely when $|i \cap j| = x$.

The graph whose adjacency matrix is X is called the x-block graph of \mathcal{Q} and is denoted by $B_x(\mathcal{Q})$. Note that $B_y(\mathcal{Q}) = \overline{B_x(\mathcal{Q})}$. For the sake of brevity, we refer to \mathcal{Q} as a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$; recall that r and b can be recovered from v, k, and λ via (3) and (4). The x-total graph of \mathcal{Q} , which we denote by $T_x(\mathcal{Q})$, is defined to be the graph with adjacency matrix

$$\begin{bmatrix} O & M \\ M^{\mathsf{T}} & A(\mathsf{B}_x(\mathcal{Q})) \end{bmatrix}.$$

We define the *x*-whole graph of \mathcal{Q} , which we denote by $W_x(\mathcal{Q})$, to be the graph with adjacency matrix

$$\begin{bmatrix} J-I & M \\ M^{\mathsf{T}} & A(\mathsf{B}_x(\mathcal{Q})) \end{bmatrix}.$$

The block graph of a QSD Q is a strongly regular graph [28, Theorem 48.10] and its eigenvalues can be expressed in terms of the parameters of Q:

Theorem 5.1. The x-block graph $B_x(Q)$ of a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$ has spectrum

$$\operatorname{spec}\left(\mathsf{B}_{x}(\mathcal{Q})\right) = \left\{ \left[\frac{k(r-1) - y(b-1)}{x-y}\right]^{1}, \left[\frac{r-\lambda-k+y}{x-y}\right]^{v-1}, \left[\frac{y-k}{x-y}\right]^{b-v} \right\}.$$

To avoid potential confusion from Theorem 5.1, we make a clarifying remark.

Remark 5.2. Note that Theorem 5.1 does not claim that the x-block graph of a QSD is necessarily connected or that it has precisely three distinct eigenvalues. For example, it is not necessary that

$$\frac{k(r-1) - y(b-1)}{x-y} \neq \frac{r-\lambda - k + y}{x-y}.$$

Indeed, one can obtain a QSD by pooling together $m \ge 2$ copies of the blocks of a symmetric 2- (v, k, λ) design. The resulting QSD Q has parameters $(v, k, m\lambda; mv, mk, \{k, \lambda\})$. Its block graph $\mathsf{B}_k(Q)$ is the disjoint union of v copies of K_m , which is disconnected and has just two distinct eigenvalues.

Next, we show that if the total graph or the whole graph of a QSD has three distinct eigenvalues and is not regular then it must have coherent rank 9.

Lemma 5.3. Let \mathcal{Q} be a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$. Then

- (i) if $T_x(Q) \in \mathscr{G}_3$ is not regular then $T_x(Q)$ has coherent rank 9;
- (ii) if $W_x(\mathcal{Q}) \in \mathscr{G}_3$ is not regular then $W_x(\mathcal{Q})$ has coherent rank 9.

Proof. We give a proof of (i). The proof for (ii) can be obtained mutatis mutandis. Suppose $T_x(\mathcal{Q}) \in \mathscr{G}_3$ is not regular. Let $A = A(\Gamma)$ be the adjacency matrix of Γ , M be the incidence matrix of \mathcal{Q} , and X be the adjacency matrix of $\mathsf{B}_x(\mathcal{Q})$. Since

$$A \in \left\langle \begin{bmatrix} I & O \\ O & O \end{bmatrix}, \begin{bmatrix} J-I & O \\ O & O \end{bmatrix}, \begin{bmatrix} O & M \\ O & O \end{bmatrix}, \begin{bmatrix} O & J-M \\ O & O \end{bmatrix}, \begin{bmatrix} O & O \\ M^{\mathsf{T}} & O \end{bmatrix}, \begin{bmatrix} O & O \\ J-M^{\mathsf{T}} & O \end{bmatrix}, \begin{bmatrix} O & O \\ O & I \end{bmatrix}, \begin{bmatrix} O & O \\ O & X \end{bmatrix}, \begin{bmatrix} O & O \\ O & X \end{bmatrix}, \begin{bmatrix} O & O \\ O & J-X-I \end{bmatrix} \right\rangle,$$

it is clear that the coherent closure $\mathcal{W}(\Gamma)$ has rank at most 9. Observe that Γ is biregular with valency partition $\mathfrak{D}(\Gamma) = \{V_1, V_2\}$ where $\{\Gamma[V_1], \Gamma[V_2]\} = \{\overline{K_v}, \mathsf{B}_x(\mathcal{Q})\}$. By Theorem 4.3, if Γ had coherent rank 8 then $\{\Gamma[V_1], \Gamma[V_2]\} = \{\overline{K_v}, K_b\}$, which is impossible since $\mathsf{B}_x(\mathcal{Q})$ cannot be a complete graph.

Lastly, by Theorem 3.4, if Γ had coherent rank at most 7 then Γ would be either complete bipartite or a cone over a strongly regular graph. If Γ is a complete bipartite graph then $\{\Gamma[V_1], \Gamma[V_2]\} = \{\overline{K_v}, \overline{K_b}\}$, which is impossible since $\mathsf{B}_x(\mathcal{Q})$ cannot be an empty graph. Otherwise, if Γ is a cone over a strongly regular graph then $\{\Gamma[V_1], \Gamma[V_2]\} = \{\overline{K_1}, \mathsf{B}_x(\mathcal{Q})\}$, which is impossible since we must have $v \ge 2$.

The last two results of this subsection are well-known properties of strongly regular graphs and QSDs that will be used below. We call an eigenvalue **restricted** if it has an eigenvector perpendicular to the all-ones vector **1**.

Lemma 5.4 ([3, Section 1.1.1]). Let $\Gamma \in SRG(n, k, a, c)$ have restricted eigenvalues $e_1 > e_2$. Then $a - c = e_1 + e_2$ and $c - k = e_1e_2$.

One more ingredient that we require is a further restriction on the parameters of a QSD. The following necessary condition is due to Calderbank [6].

Theorem 5.5 ([3, Proposition 8.5.4]). Let Q be a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$. Then

$$(v-1)(v-2)(k-x)(k-y) - k(v-k)(v-2)(2k-x-y) + k(v-k)(k(v-k)-1) \ge 0,$$

with equality, if and only if Q is a 3-design.

5.2 Underlying quasi-symmetric designs

First, we show that if a graph in \mathscr{G}_3 has coherent rank 9 then its coherent closure has type $\begin{bmatrix} 2 & 2 \\ 3 \end{bmatrix}$.

Theorem 5.6. Let $\Gamma \in \mathscr{G}_3$ have coherent rank 9. Then $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & 2 \\ 3 \end{bmatrix}$.

Proof. By Theorem 3.4, the graph Γ must not be regular. Using Lemma 3.1 and Corollary 3.2 (iii), we find that the coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 1 & 1 \\ 6 \end{bmatrix}$, $\begin{bmatrix} 2 & 1 \\ 5 \end{bmatrix}$, $\begin{bmatrix} 2 & 2 \\ 3 \end{bmatrix}$, $\begin{bmatrix} 3 & 1 \\ 4 \end{bmatrix}$, or $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 \\ 1 \end{bmatrix}$. Except for $\begin{bmatrix} 2 & 2 \\ 3 \end{bmatrix}$, for each case with two fibres, the complement of Γ is disconnected. The case with three fibres must correspond to a graph with just three vertices. There is just one connected graph on three vertices with three distinct eigenvalues and its coherent rank is 5.

We now show that each graph in \mathscr{G}_3 whose coherent closure has type $\begin{bmatrix} 2 & 2 \\ 3 \end{bmatrix}$ corresponds to a QSD.

Theorem 5.7. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ such that $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & 2 \\ 3 \end{bmatrix}$. Then

- (i) Γ is biregular with degrees $k_1 \neq k_2$ and valency partition $\{V_1, V_2\}$, which is equitable with quotient matrix $\begin{bmatrix} k_{11} & k_{12} \\ k_{21} & k_{22} \end{bmatrix}$ and $|V_i| = n_i$ for $i \in \{1, 2\}$;
- (ii) $A(\Gamma)$ has the form $\begin{bmatrix} \varepsilon(J-I) & M \\ M^{\mathsf{T}} & A(\Gamma[V_2]) \end{bmatrix}$, where $\varepsilon \in \{0,1\}$ and M is the incidence matrix of a QSD \mathcal{Q} with parameters $(n_1, k_{21}, \lambda; n_2, k_{12}, \{x, y\})$ and $\Gamma[V_2] = \mathsf{B}_x(\mathcal{Q})$.

Suppose that $B_x(Q)$ has restricted eigenvalues $e_1 > e_2$. Then either $\theta_1 = e_1 = \frac{y-k_{21}}{x-y}$ or $\theta_2 = e_2 = \frac{y-k_{21}}{x-y}$. Furthermore,

$$(x,y) = (k_{21} + (\theta_1 + 1)(\theta_2 + 1) - (e_1 + 1)(e_2 + 1), k_{21} + \theta_1\theta_2 - e_1e_2);$$
(14)

$$\lambda - k_{12} = \begin{cases} \theta_1 \theta_2, & \text{if } \Gamma[V_1] \text{ is empty;} \\ (\theta_1 + 1)(\theta_2 + 1), & \text{if } \Gamma[V_1] \text{ is complete;} \end{cases}$$
(15)

$$\frac{k_{12} - \lambda + y - k_{21}}{x - y} = \begin{cases} \theta_1 + \theta_2, & \text{if } \Gamma[V_1] \text{ is empty;} \\ \theta_1 + \theta_2 + 1, & \text{if } \Gamma[V_1] \text{ is complete;} \end{cases}$$
(16)

$$\sqrt{(k_1 + \theta_1 \theta_2)(k_2 + \theta_1 \theta_2)} = \begin{cases} \frac{\lambda k_{21} - y k_{12}}{x - y}, & \text{if } \Gamma[V_1] \text{ is empty;} \\ k_{21} + \frac{\lambda k_{21} - y k_{12}}{x - y}, & \text{if } \Gamma[V_1] \text{ is complete.} \end{cases}$$
(17)

Proof. Suppose $A = A(\Gamma)$ is the adjacency matrix of Γ . By Theorem 3.4 and Corollary 3.2 (iii), the graph Γ must be biregular with valencies k_1 and k_2 (say). By Corollary 3.2 (iii), we can write $A = \begin{bmatrix} A(\Gamma[V_1]) & M \\ M^{\mathsf{T}} & A(\Gamma[V_2]) \end{bmatrix}$ and

$$A^2 = \begin{bmatrix} A(\Gamma[V_1])^2 + MM^{\mathsf{T}} & A(\Gamma[V_1])M + MA(\Gamma[V_2]) \\ M^{\mathsf{T}}A(\Gamma[V_1]) + A(\Gamma[V_2])M^{\mathsf{T}} & M^{\mathsf{T}}M + A(\Gamma[V_2])^2 \end{bmatrix}$$

Apply (1) to obtain

$$A^{2} - (\theta_{1} + \theta_{2})A + \theta_{1}\theta_{2}I = \begin{bmatrix} (k_{1} + \theta_{1}\theta_{2})J & \sqrt{(k_{1} + \theta_{1}\theta_{2})(k_{2} + \theta_{1}\theta_{2})}J \\ \sqrt{(k_{1} + \theta_{1}\theta_{2})(k_{2} + \theta_{1}\theta_{2})}J & (k_{2} + \theta_{1}\theta_{2})J \end{bmatrix}.$$
 (18)

By Corollary 3.2 (i) and Proposition 3.3, the subgraph $\Gamma[V_1] = K_{n_1}$ or \overline{K}_{n_1} , i.e., the matrix $A(\Gamma[V_1]) = \varepsilon_{11}(J - I)$ for some $\varepsilon_{11} \in \{0, 1\}$. Moreover, by Proposition 4.5, we have $\Gamma[V_2] \in \text{SRG}(n_2, k_{22}, a, c)$, that is, $A(\Gamma[V_2]) = X$ where

$$X^{2} = k_{22}I + aX + c(J - I - X).$$
(19)

From the top-left block of (18), we obtain

$$MM^{\mathsf{T}} = \begin{cases} (k_{12} + \theta_1 \theta_2) J_{n_1} - \theta_1 \theta_2 I, & \text{if } \Gamma[V_1] = \overline{K}_{n_1}; \\ (k_{12} + (\theta_1 + 1)(\theta_2 + 1)) J_{n_1} - (\theta_1 + 1)(\theta_2 + 1)I, & \text{if } \Gamma[V_1] = K_{n_1}. \end{cases}$$
(20)

Similar to the above, we can use the bottom-right block of (18) together with (19) to deduce that

$$M^{\mathsf{T}}M = (k_2 + \theta_1\theta_2 - c)J_{n_2} - (k_{22} + \theta_1\theta_2 - c)I + (\theta_1 + \theta_2 + c - a)X,$$

which, by Lemma 5.4, becomes

$$M^{\mathsf{T}}M = k_{21}I + (k_{21} + \theta_1\theta_2 - e_1e_2)(J_{n_2} - I - X) + (k_{21} + (\theta_1 + 1)(\theta_2 + 1) - (e_1 + 1)(e_2 + 1))X, \quad (21)$$

where $e_1 > e_2$ are the restricted eigenvalues of X. Observe that M is the incidence matrix of a quasi-symmetric 2-design Q. Suppose Q has parameters $(v, k, \lambda; b, r, \{x, y\})$ then $v = n_1$, $k = k_{21}, b = n_2, r = k_{12}, \text{ and } \lambda(n_1 - 1) = n_2(k_{21} - 1)$. Without loss of generality, we can assume that $X = A(B_x(Q))$.

Using (6), we find that the eigenvalues of MM^{T} are rk and $r - \lambda$ with multiplicities 1 and v - 1 respectively. Using (21), we find the eigenvalues of $M^{\mathsf{T}}M$ are

$$(e_1e_2 - \theta_1\theta_2) + b(k + \theta_1\theta_2 - e_1e_2) + e_0(\theta_1 + \theta_2 - e_1 - e_2),$$

 $(\theta_1 - e_1)(e_1 - \theta_2),$ and
 $(\theta_1 - e_2)(e_2 - \theta_2),$

with multiplicities 1, m_1 , and m_2 respectively, where $\{m_1, m_2\} = \{v - 1, b - v\}$. By Lemma 4.1, we must have

$$0 = \begin{cases} (\theta_1 - e_1)(e_1 - \theta_2), & \text{if } m_1 = b - v; \\ (\theta_1 - e_2)(e_2 - \theta_2), & \text{if } m_2 = b - v. \end{cases}$$

By Corollary 3.7, $\theta_1 > 0$ and $\theta_2 < 0$, whence we have either $\theta_1 = e_1 = \frac{y-k}{x-y}$ or $\theta_2 = e_2 = \frac{y-k}{x-y}$. Comparing coefficients of (21) and (13) yields (14). Comparing coefficients of (6) and

Comparing coefficients of (21) and (13) yields (14). Comparing coefficients of (6) and (20), we obtain (15). Using (13) together with (6), we find that

$$MX = \frac{(\lambda k - yr)J + (r - \lambda + y - k)M}{x - y}.$$
(22)

Combining coefficients for M and J in the top-right block of (18) yields (16) and (17). \Box

Given a graph $\Gamma \in \mathscr{G}_3$ such that $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & 2\\ 3 \end{bmatrix}$, we call the QSD \mathcal{Q} from the conclusion of Theorem 5.7 the **underlying QSD** of Γ .

Corollary 5.8. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ such that $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & 2\\ 3 \end{bmatrix}$. Suppose that \mathcal{Q} is the underlying QSD of Γ . Then \mathcal{Q} cannot have repeated blocks. Furthermore, the block graph $\mathsf{B}_x(\mathcal{Q})$ is connected.

Proof. Suppose that \mathcal{Q} has repeated blocks. Then either k = x or k = y. First, suppose that k = x. Then we must have $\frac{y-k}{x-y} = -1$. By Theorem 5.7, we must have $\theta_2 = -1$, which contradicts (2). Otherwise, if k = y then, in a similar fashion, we find that $\theta_1 = 0$, which contradicts Corollary 3.7. Hence, \mathcal{Q} does not have repeated blocks. Furthermore, by [24, Theorem Q], we can deduce that the block graph $\mathsf{B}_x(\mathcal{Q})$ is connected, as required.

Corollary 5.8 allows us to use the absolute bound (see Theorem A.1) to restrict our search for parametric QSDs in the next section.

Theorem 5.9. Let θ_1 and θ_2 be integers such that $\theta_1 \ge 1$ and $\theta_2 \le -2$. Suppose Q is a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$ where $\mathsf{B}_x(Q) \in \mathscr{G}_3(e_0, e_1, e_2)$ and

$$(x,y) = (k + (\theta_1 + 1)(\theta_2 + 1) - (e_1 + 1)(e_2 + 1), k + \theta_1\theta_2 - e_1e_2).$$

(i) If
$$\lambda = r + \theta_1 \theta_2$$
, $\frac{r - \lambda + y - k}{x - y} = \theta_1 + \theta_2$, and $\frac{\lambda k - yr}{x - y} = \sqrt{\lambda(k + e_0 + \theta_1 \theta_2)}$ then
 $\mathsf{T}_x(\mathcal{Q}) \in \mathscr{G}_3(e_0 - \theta_2, \theta_1, \theta_2)$.

Furthermore, if $r \neq k + e_0$ then $\mathsf{T}_x(\mathcal{Q})$ has coherent rank 9.

(*ii*) If
$$\lambda = r + (\theta_1 + 1)(\theta_2 + 1)$$
, $\frac{r - \lambda + y - k}{x - y} = \theta_1 + \theta_2 + 1$, and

$$k + \frac{\lambda k - yr}{x - y} = \sqrt{(v - 1 + r + \theta_1 \theta_2)(k + e_0 + \theta_1 \theta_2)}$$

then

$$\mathsf{W}_{x}(\mathcal{Q}) \in \mathscr{G}_{3}\left(\frac{v-1+e_{0}+\sqrt{(v-1-e_{0})^{2}+4rk}}{2},\theta_{1},\theta_{2}\right)$$

Furthermore, if $v - 1 + r \neq k + e_0$ then $W_x(Q)$ has coherent rank 9.

Proof. We give a proof for (i). The proof for (ii) can be obtained mutatis mutandis. Let A be the adjacency matrix of $\mathsf{T}_x(\mathcal{Q})$, M be the incidence matrix of \mathcal{Q} , and X be the adjacency matrix of $\mathsf{B}_x(\mathcal{Q})$. Then, we can write

$$A = \begin{bmatrix} O & M \\ M^{\mathsf{T}} & X \end{bmatrix} \quad \text{and} \quad A^2 = \begin{bmatrix} MM^{\mathsf{T}} & MX \\ XM^{\mathsf{T}} & M^{\mathsf{T}}M + X^2 \end{bmatrix}.$$

Next, we find

$$X^{2} = \frac{k(r-1) - y(b-1)}{x - y}I + (e_{1} + e_{2})X + \left(\frac{k(r-1) - y(b-1)}{x - y} + e_{1}e_{2}\right)(J - I).$$

Let $e_0 = \frac{k(r-1)-y(b-1)}{x-y}$. Hence, using (13) and (6), we have

$$A^{2} = \begin{bmatrix} (r-\lambda)I + \lambda J & \frac{(\lambda k - yr)J + (r-\lambda + y - k)M}{x - y} \\ \frac{(\lambda k - yr)J + (r-\lambda + y - k)M^{\mathsf{T}}}{x - y} & (k - y - e_{1}e_{2})I + (x - y + e_{1} + e_{2})X + (e_{0} + e_{1}e_{2} + y)J \end{bmatrix}.$$

Suppose that $\lambda = r + \theta_1 \theta_2$, $\frac{r - \lambda + y - k}{x - y} = \theta_1 + \theta_2$, and $\frac{\lambda k - yr}{x - y} = \sqrt{\lambda(k + e_0 + \theta_1 \theta_2)}$. Then

$$A^{2} - (\theta_{1} + \theta_{2})A + \theta_{1}\theta_{2}I = \begin{bmatrix} \lambda J & \frac{r - \lambda + y - k}{x - y}J\\ \frac{r - \lambda + y - k}{x - y}J & (k + e_{0} + \theta_{1}\theta_{2})J \end{bmatrix}.$$

Let $\sigma = \mathfrak{D}(\mathsf{T}_x(\mathcal{Q}))$. Hence, $\mathsf{T}_x(\mathcal{Q}) \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$, for some θ_0 , which one can determine by finding the eigenvalues of $Q = Q_\sigma(\mathsf{T}_x(\mathcal{Q})) = \begin{bmatrix} 0 & r \\ k & e_0 \end{bmatrix}$. Since $\theta_0 \ge e_0$, the eigenvalues of Qmust be θ_0 and θ_2 . The degree sequence is $\{[r]^v, [k+e_0]^b\}$. Thus, if $r \ne k + e_0$ then, by Lemma 5.3, $\mathsf{T}_x(\mathcal{Q})$ has coherent rank 9.

5.3 Parametric quasi-symmetric designs

Now we apply Theorem 5.7 to find parametrisations for quasi-symmetric designs that are the underlying QSD of a graph in \mathscr{G}_3 of coherent rank 9. Suppose that \mathscr{G}_3 has coherent rank 9. Let \mathcal{Q} be the underlying QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$ and let $e_1 > e_2$ be the restricted values of $\mathsf{B}_x(\mathcal{Q})$. By Theorem 5.7, Γ is biregular with degree sequence $\{[k_1]^{n_1}, [k_2]^{n_2}\}$. According to Theorem 5.7, there are four cases to consider depending on whether $\Gamma[V_1] = \overline{K_{n_1}}$ or $\Gamma[V_1] = K_{n_1}$ and $\theta_1 = e_1 = \frac{y-k}{x-y}$ or $\theta_2 = e_2 = \frac{y-k}{x-y}$. We split our consideration according to each case, which we refer to as *classes* as follows.

- Class 1: $\Gamma[V_1] = \overline{K_{n_1}}$ and $\theta_2 = e_2 = \frac{y-k}{x-y}$;
- Class 2: $\Gamma[V_1] = \overline{K_{n_1}}$ and $\theta_1 = e_1 = \frac{y-k}{x-y}$;
- Class 3: $\Gamma[V_1] = K_{n_1}$ and $\theta_2 = e_2 = \frac{y-k}{x-y}$;
- Class 4: $\Gamma[V_1] = K_{n_1}$ and $\theta_1 = e_1 = \frac{y-k}{x-y}$.

5.3.1 Class 1

Lemma 5.10. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with coherent rank 9. Suppose that the underlying QSD \mathcal{Q} of Γ has parameters $(v, k, \lambda; b, r, \{x, y\})$ and $\mathsf{B}_x(\mathcal{Q})$ has eigenvalues $e_0 > e_1 > e_2$. Suppose that $\Gamma[V_1]$ is empty and $\theta_2 = e_2 = \frac{y-k}{x-y}$. Then $\theta_0 = \theta_2(\theta_1\theta_2 - \theta_1 - 1)$, $e_0 = \theta_1\theta_2(\theta_2 - 1)$, and $e_1 = \theta_1 + \theta_2$. Furthermore, the parameters of \mathcal{Q} can be written in terms of θ_1 and θ_2 according to Table 1. Conversely, suppose that, for some integers θ_1 and θ_2 satisfying

Table 1: Expressions for parameters from Lemma 5.10.

 $\theta_1 \ge 1$ and $\theta_2 \le -2$, there exists a QSD \mathcal{Q} whose parameters $(v, k, \lambda; b, r, \{x, y\})$ can be expressed as above in terms of θ_1 and θ_2 . Then $\mathsf{T}_x(\mathcal{Q}) \in \mathscr{G}_3(\theta_2(\theta_1\theta_2 - \theta_1 - 1), \theta_1, \theta_2)$ and $\mathsf{T}_x(\mathcal{Q})$ has coherent rank 9.

Proof. Since $e_2 = \frac{y-k}{x-y}$, using Theorem 5.1, we must have $e_1 = \frac{r-\lambda-k+y}{x-y}$. We use the equations (3), (4), (14), (15), (16), (17) together with Theorem 5.1 and Theorem 2.8 (iii). Let Q be the quotient matrix of the valency partition of the vertex set of Γ . Since the trace of Q is e_0 , which, by interlacing (Theorem 2.1), is at most θ_0 , we find that θ_2 is an eigenvalue of Q. Hence $e_0 = \theta_0 + \theta_2$ and $\theta_0 \theta_2 = -rk$. The statement of the lemma follows from the above list of equations and the converse follows from Theorem 5.9.

Suppose that Γ satisfies the assumptions of Lemma 5.10. The parameters of the underlying QSD of Γ can each be expressed in terms of θ_1 and θ_2 . First, we consider the subfamily of QSD parameters that satisfies the additional constraint $\theta_1 = -\theta_2$. With this additional constraint, each QSD parameter can be expressed in terms of a single variable θ_1 . Indeed, each such QSD has parameters $(\theta_1^3, \theta_1^2, \theta_1 + 1; \theta_1(\theta_1^2 + \theta_1 + 1), \theta_1^2 + \theta_1 + 1, \{0, \theta_1\})$.

Example 5.11. The biregular graphs $\mathsf{T}_x(\mathcal{Q})$ of Lemma 5.10 whose underlying QSD \mathcal{Q} satisfies the additional constraint $\theta_1 = -\theta_2$ was discovered by Van Dam [12, Section 2.3].

Next, instead, consider the additional constraint $\theta_1 + 1 = \theta_2(\theta_2 - 1)$. The resulting one-parameter family violates Theorem 5.5. Indeed, we have $v = -\theta_2^3 + \theta_2^2 + 2\theta_2 + 1$, $k = \theta_2^2$, $\lambda = \theta_2^2 + \theta_2$, $x = -\theta_2$ and y = 0. Thus, the left-hand side of the inequality of Theorem 5.5 equals $-(\theta_2 + 1)^2 \theta_2^2(1 - \theta_2)$, which is clearly negative.

It remains to consider the QSD parameters of Lemma 5.10 that are not captured by the two infinite parametric families above. By Theorem 2.5, we can assume that $\theta_0 > \theta_1 \ge 1$ and by Theorem 2.1, we can assume that $\theta_2 \le -2$. Furthermore, since $e_1 \ge 0$, we have that $\theta_1 \ge -\theta_2$. In Table 5, we list those for which $\theta_1 \le 100$.

5.3.2 Class 2

Define the function $f_2(w, z)$ by

$$f_2(w,z) := \frac{w(w^2 + w + 1) - 2z(w+1) + w\sqrt{(w^2 + w + 1)^2 - 4wz(w+1)}}{2}$$

Lemma 5.12. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with coherent rank 9. Suppose that the underlying QSD \mathcal{Q} of Γ has parameters $(v, k, \lambda; b, r, \{x, y\})$ and $\mathsf{B}_x(\mathcal{Q})$ has eigenvalues $e_0 > e_1 > e_2$. Suppose that $\Gamma[V_1]$ is empty and $\theta_1 = e_1 = \frac{y-k}{x-y}$. Then $\theta_0 = f_2(\theta_1, \theta_2)$, $e_0 = \theta_0 + \theta_2$, and $e_2 = \theta_1 + \theta_2$. Furthermore, the parameters of \mathcal{Q} can be written in terms of θ_1 and θ_2 according to Table 2. Conversely, suppose that, for some integers θ_1 and θ_2 satisfying $\theta_1 \ge 1$ and $\theta_2 \leqslant -2$, there

$$\begin{array}{c|c|c|c|c|c|c|c|c|}\hline v & k & \lambda & b & r & x & y\\ \hline \hline -\theta_2(\theta_0-\theta_1) & y+\theta_1^2 & \hline -\theta_2(x-1) & r-k+e_0+1 & \lambda-\theta_1\theta_2 & y-\theta_1 & \hline e_0-\theta_1^3+\theta_1\theta_2 & \hline \theta_1 & \hline$$

Table 2: Expressions for parameters from Lemma 5.12.

exists a QSD Q whose parameters $(v, k, \lambda; b, r, \{x, y\})$ can be expressed as above in terms of θ_1 and θ_2 . Then $\mathsf{T}_x(Q) \in \mathscr{G}_3(f_2(\theta_1, \theta_2), \theta_1, \theta_2)$. Furthermore, if $r \neq k + f_2(\theta_1, \theta_2) + \theta_2$ then $\mathsf{T}_x(Q)$ has coherent rank 9.

Proof. Since $e_1 = \frac{y-k}{x-y}$, using Theorem 5.1, we must have $e_2 = \frac{r-\lambda-k+y}{x-y}$. We use the equations (3), (4), (14), (15), (16), (17) together with Theorem 5.1 and Theorem 2.8 (iii). Let Q be the quotient matrix of the valency partition of the vertex set of Γ . Since the trace of Q is e_0 , which, by interlacing (Theorem 2.1), is at most θ_0 , we find that θ_2 is an eigenvalue of Q. Hence $e_0 = \theta_0 + \theta_2$ and $\theta_0 \theta_2 = -rk$. The expressions for the parameters of the underlying QSD Q together with e_0 and e_2 follow from the above equations. We also obtain the equation

$$\theta_0^2 - \theta_0(\theta_1^3 - \theta_1^2 + 2\theta_1\theta_2 - \theta_1 + 2\theta_2) - \theta_1^3\theta_2 + \theta_1^2\theta_2^2 - 2\theta_1^2\theta_2 + 2\theta_1\theta_2^2 - \theta_1\theta_2 + \theta_2^2 = 0.$$

Think of the above equation as a quadratic univariate polynomial equation in θ_0 with coefficients in $\mathbb{Z}[\theta_1, \theta_2]$ and let $r_1(\theta_1, \theta_2) \ge r_2(\theta_1, \theta_2)$ be its roots. Note that we have $r_1(\theta_1, \theta_2) = f_2(\theta_1, \theta_2)$. Using the fact that $x \ge 0$, we can rule out the possibility of $\theta_0 = r_2(\theta_1, \theta_2)$. Indeed, $x \ge 0$ implies that $\theta_0 \ge \theta_1^3 + \theta_1^2 - \theta_1\theta_2 - \theta_2$. On the other hand, $r_2(\theta_1, \theta_2) \le -\theta_2(\theta_1 + 1)$. Hence, $\theta_0 = f_2(\theta_1, \theta_2)$, as required.

The converse follows from Theorem 5.9.

Suppose that Γ satisfies the assumptions of Lemma 5.12. The parameters of the underlying QSD of Γ can each be expressed in terms of θ_1 and θ_2 . First, we consider the subfamily of QSD parameters that satisfies the additional constraint $\theta_2 = -\theta_1^2(\theta_1^2 + 1)$. With this additional constraint, each QSD parameter can be expressed in terms of a single variable θ_1 .

When $\theta_1 = 1$, we find that v = b, which implies the corresponding design is symmetric, and hence not quasi-symmetric. When $\theta_1 = 2$, we obtain a new graph:

Example 5.13. The parameters $(v, k, \lambda; b, r, \{x, y\}) = (22, 15, 80; 176, 120, \{9, 11\})$ correspond to a QSD Q. Indeed, take Q to be the complement of the QSD with parameters $(22, 7, 16; 176, 56, \{1, 3\})$ from [17]. The block graph $\mathsf{B}_9(Q)$ has eigenvalues $e_0 = 70, e_1 = 2$, and $e_2 = -18$ and the total graph $\mathsf{T}_9(Q)$ is a biregular graph in $\mathscr{G}_3(90, 2, -20)$. Moreover,

spec
$$(T_9(\mathcal{Q})) = \left\{ [90]^1, [2]^{175}, [-20]^{22} \right\}.$$

When $\theta_1 \ge 3$, Theorem 5.5 is violated. Indeed, we have

$$v = \theta_1^3 + 2\theta_1^2 + 2\theta_1 + 2; \qquad k = \theta_1^3 + \theta_1^2 + \theta_1 + 1; \qquad x = 1 + \theta_1^3; \qquad y = 1 + \theta_1 + \theta_1^3.$$

Thus, the left-hand side of the inequality of Theorem 5.5 is equal to

$$-(\theta_1 - 2)\theta_1(\theta_1 + 1)^2(\theta_1^2 + \theta_1 + 1),$$

which is clearly negative when $\theta_1 \ge 3$.

Next, instead, consider the additional constraint $\theta_2 = -(\theta_1^4 + 2\theta_1^3 + \theta_1^2 + \theta_1)$. When $\theta_1 = 1$, we obtain the QSD parameters

$$(v, k, \lambda; b, r, \{x, y\}) = (7, 5, 10; 21, 15, \{3, 4\})$$

and the total graph $\mathsf{T}_3(\mathcal{Q}) \in \mathrm{SRG}(28, 15, 6, 10)$. Note that the nonregularity condition $r \neq k + f_2(\theta_1, \theta_2) + \theta_2$ is violated in this case.

When $\theta_1 \ge 2$, Theorem 5.5 is, again, violated. Indeed, we have

$$v = \theta_1^3 + 3\theta_1^2 + 2\theta_1 + 1; \quad k = \theta_1^3 + 2\theta_1^2 + \theta_1 + 1; \quad x = \theta_1^3 + \theta_1^2 + 1; \quad y = (\theta_1 + 1)(\theta_1^2 + 1).$$

Thus, the left hand side of the inequality of Theorem 5.5 equals $-\theta_1(\theta_1+1)^2(\theta_1^3-2\theta_1+1)$, which is clearly negative when $\theta_1 \ge 2$.

It remains to consider QSD parameters of Lemma 5.12 that are not captured by the two infinite parametric families above. By Theorem 2.5, we can assume that $\theta_0 > \theta_1 \ge 1$ and by Theorem 2.1, we can assume that $\theta_2 \le -2$. Since $\theta_1 + \theta_2 = e_2 \le -2$ and $\theta_1 \ge 1$, we find that $1 \le \theta_1 \le -\theta_2 - 2$. In Table 6, we list those for which $\theta_2 \ge -100$.

Remark 5.14. Note that, if the nonregularity condition $r \neq k + f_2(\theta_1, \theta_2) + \theta_2$ is not imposed then Lemma 5.12 can produce QSDs whose total graph is strongly regular. These strongly regular graphs have an *improper strongly regular decomposition* in the sense of Haemers and Higman [18].

5.3.3 Class 3

We need to split into two further subcases according to which of θ_1 or θ_2 is an eigenvalue of the quotient matrix $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$.

Define the functions $f_3(w, z)$ and $g_3(w, z)$ by

$$f_3(w,z) := \frac{wz(w+2)(z+2) + 2z^2 + 4z + 1 + (1-wz)\sqrt{g_3(w,z)}}{-2(z+1)(w+z+2)}$$

$$g_3(w,z) := 5(wz+1)^2 + 4z^3(w+1) + 4wz(w+4z) + 4(z+1)(3z-1).$$

Lemma 5.15. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with coherent rank 9. Suppose that the underlying $QSD \ \mathcal{Q}$ of Γ has parameters $(v, k, \lambda; b, r, \{x, y\})$ and $\mathsf{B}_x(\mathcal{Q})$ has eigenvalues $e_0 > e_1 > e_2$. Suppose that $\Gamma[V_1]$ is complete, $\theta_2 = e_2 = \frac{y-k}{x-y}$, and θ_2 is an eigenvalue of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$. Then $\theta_0 = f_3(\theta_1, \theta_2), e_1 = \theta_1 + \theta_2 + 1$, and

$$e_0 = -\frac{\theta_2(\theta_1 + 1)}{(\theta_2 + 1)} \left(1 + \frac{\theta_2(1 - \theta_1 \theta_2)}{\theta_0 + \theta_1 \theta_2} \right).$$

Furthermore, the parameters of Q can be written in terms of θ_1 and θ_2 as follows.

$$\begin{aligned} v &= \theta_0 + \theta_2 - e_0 + 1; \\ k &= y + \theta_2(\theta_2 + 1); \\ \lambda &= (\theta_1 + 1) \left(1 + \frac{\theta_0 e_1 - \theta_1 e_0}{\theta_0 - \theta_1} \right); \\ y &= \frac{-\theta_2(\lambda - e_1 - 1)}{\theta_1}. \end{aligned}$$

Conversely, suppose that, for some integers θ_1 and θ_2 satisfying $\theta_1 \ge 1$ and $\theta_2 \le -2$, there exists a QSD Q whose parameters $(v, k, \lambda; b, r, \{x, y\})$ can be expressed as above in terms of θ_1 and θ_2 . Then $W_x(Q) \in \mathscr{G}_3(f_3(\theta_1, \theta_2), \theta_1, \theta_2)$ and $W_x(Q)$ has coherent rank 9.

Proof. Since $e_2 = \frac{y-k}{x-y}$, using Theorem 5.1, we must have $e_1 = \frac{r-\lambda-k+y}{x-y}$. We use the equations (3), (4), (14), (15), (16), (17) together with Theorem 5.1 and Theorem 2.8 (iii). Using the trace and determinant of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$, we obtain the equations $v - 1 + e_0 = \theta_0 + \theta_2$ and $\theta_0 \theta_2 = (v - 1)e_0 - rk$. The expressions for the parameters of the underlying QSD \mathcal{Q} together with e_0 and e_1 follow from the above equations. We also obtain the equation

$$\xi_2(\theta_1, \theta_2)\theta_0^2 + \xi_1(\theta_1, \theta_2)\theta_0 + \xi_0(\theta_1, \theta_2) = 0,$$

where

$$\begin{split} \xi_2(\theta_1, \theta_2) &:= \theta_1 \theta_2 + \theta_1 + \theta_2^2 + 3\theta_2 + 2; \\ \xi_1(\theta_1, \theta_2) &:= \theta_1^2 \theta_1^2 + 2\theta_1^2 \theta_2 + 2\theta_1 \theta_2^2 + 4\theta_1 \theta_2 + 2\theta_2^2 + 4\theta_2 + 1; \\ \xi_0(\theta_1, \theta_2) &:= -\theta_1^3 \theta_2^3 + \theta_1^3 \theta_2^2 - \theta_1^2 \theta_2^3 + 4\theta_1^2 \theta_2^2 + 4\theta_1 \theta_2^2 + \theta_2^2. \end{split}$$

Think of the above equation as a quadratic univariate polynomial equation in θ_0 with coefficients in $\mathbb{Z}[\theta_1, \theta_2]$ and let $s_1(\theta_1, \theta_2) \ge s_2(\theta_1, \theta_2)$ be its roots. Note that we have

 $s_1(\theta_1, \theta_2) = f_3(\theta_1, \theta_2)$. Using the fact that $\theta_0 > \theta_1 \ge 0$, we can rule out the possibility of $\theta_0 = s_2(\theta_1, \theta_2)$. Suppose to the contrary that $\theta_0 = s_2(\theta_1, \theta_2)$. Then $\theta_0 > \theta_1 \ge 0$ and $\theta_2 < 0$ implies that

$$\theta_1 \theta_2(\theta_1 + 2)(\theta_2 + 2) + 2\theta_2^2 + 4\theta_2 + 1 - (1 - \theta_1 \theta_2)\sqrt{g_3(\theta_1, \theta_2)} > -2(\theta_2 + 1)(\theta_1 + \theta_2 + 2)\theta_1,$$

which implies

$$(\theta_1\theta_2(\theta_1+2)(\theta_2+2)+2\theta_2^2+4\theta_2+1+2(\theta_2+1)(\theta_1+\theta_2+2)\theta_1)^2-(1-\theta_1\theta_2)^2g_3(\theta_1,\theta_2)$$

= $-4(\theta_2+1)(\theta_1+1)(\theta_1+\theta_2+2)(\theta_2\theta_1^2(\theta_2-3)(\theta_2+1)-\theta_1\theta_2(5\theta_2+4)-\theta_2^2-\theta_1^2-\theta_1)$

is positive. Now, $-4(\theta_2+1)(\theta_1+1)(\theta_1+\theta_2+2)$ is positive since $e_1 = \theta_1+\theta_2+1 \ge 0, \theta_2 \le -2$, and $\theta_1 \ge 0$. However, it also follows that $\theta_2 \theta_1^2(\theta_2-3)(\theta_2+1) - \theta_1 \theta_2(5\theta_2+4) - \theta_2^2 - \theta_1^2 - \theta_1$ is negative, which contradicts our supposition. Hence, $\theta_0 = f_3(\theta_1, \theta_2)$, as required.

The converse follows from Theorem 5.9.

Suppose that Γ satisfies the assumptions of Lemma 5.15. By Theorem 2.5, we can assume that $\theta_0 > \theta_1 \ge 1$ and by Theorem 2.1, we can assume that $\theta_2 \le -2$. Since $\theta_1 + \theta_2 + 1 = e_1 \ge 0$ and $\theta_2 \le -2$, we find that $-\theta_1 - 1 \le \theta_2 \le -2$. In Table 7, we list all the parameters for QSDs corresponding to Lemma 5.15 for $\theta_1 \le 100$.

Example 5.16. When $\theta_1 = 5$ and $\theta_2 = -2$, using the expressions from Lemma 5.15, we obtain the QSD parameters $(8, 6, 15; 28, 21, \{4, 5\})$. A QSD Q having such parameters exists whose blocks are all 6-sets of $\{1, 2, \ldots, 8\}$. The block graph $B_5(Q)$ has eigenvalues $(e_0, e_1, e_2) = (12, 4, -2)$ and the whole graph $W_5(Q)$ is a biregular graph in $\mathscr{G}_3(21, 5, -2)$. This graph was found by Muzychuk and Klin [23].

Define the polynomial h(u, w, z) as

$$h(u, w, z) := C_3(w, z)u^3 + C_2(w, z)u^2 + C_1(w, z)u + C_0(w, z),$$

where

$$\begin{split} C_3(w,z) &= (w+z)(w+1)(z+1);\\ C_2(w,z) &= w^4 + w^3(z+2)^2 + w^2(z^2+7z+5) - w(2z^3+3z^2-z-1) - 2z^2(z+1);\\ C_1(w,z) &= (2w^5+z^3)(z+1) - w^4(z^3+z^2-4z-6) - w^3(z^3-6z-8) \\ &\quad - 3w^2(z^3+3z^2+z+1) + wz(z^3-5z-3);\\ C_0(w,z) &= w(w+1)(w-z)(w^3(z+1)^2+2w^2(z+1) + (w-z)(z+1)(z+2) - wz). \end{split}$$

Lemma 5.17. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with coherent rank 9. Suppose that the underlying QSD \mathcal{Q} has parameters $(v, k, \lambda; b, r, \{x, y\})$ and $\mathsf{B}_x(\mathcal{Q})$ has eigenvalues $e_0 > e_1 > e_2$. Suppose that $\Gamma[V_1]$ is complete, $\theta_2 = e_2 = \frac{y-k}{x-y}$, and θ_1 is an eigenvalue of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$. Then θ_0 is the positive zero of the (univariate) polynomial $p(u) := h(u, \theta_1, \theta_2)$. Furthermore,

$$e_1 = \theta_1 + \theta_2 + 1, \text{ and } e_0 = \frac{(\theta_2 + 1)(\theta_0(\theta_1 e_1 - 1) + \theta_1^2(\theta_2 + 1))}{\theta_0(\theta_1 \theta_2 + e_1) + \theta_1^3(\theta_2 + 1) + 2\theta_1(\theta_1 + 1) - \theta_2(\theta_2 + 2)}$$

The parameters of Q can be written in terms of θ_0 , θ_1 and θ_2 as follows.

$$\begin{split} v &= \theta_0 + \theta_1 - e_0 + 1; \\ k &= y + \theta_2(\theta_2 + 1); \\ \lambda &= \theta_2 + 1 - \frac{\theta_1(y - e_1 - 1)}{\theta_2}; \\ y &= (\theta_2 + 1) \frac{\theta_0(e_1 - 1) - \theta_1(e_0 - t_1 - 1) - \theta_2^2}{\theta_0 - \theta_2}; \end{split}$$

Conversely, suppose that θ_0 is an integer solution to p(u) = 0 for some integers θ_1 and θ_2 satisfying $\theta_0 > \theta_1 \ge 1$ and $\theta_2 \le -2$. If there exists a QSD Q whose parameters $(v, k, \lambda; b, r, \{x, y\})$ can be expressed as above in terms of θ_0 , θ_1 , and θ_2 then $W_x(Q) \in \mathcal{G}_3(\theta_0, \theta_1, \theta_2)$. Suppose $B_x(Q)$ has degree e_0 . If $v - 1 + r \ne k + e_0$ then $W_x(Q)$ has coherent rank 9.

Proof. Since $e_2 = \frac{y-k}{x-y}$, using Theorem 5.1, we must have $e_1 = \frac{r-\lambda-k+y}{x-y}$. We use the equations (3), (4), (14), (15), (16), (17) together with Theorem 5.1 and Theorem 2.8 (iii). Using the trace and determinant of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$, we obtain the equations $v - 1 + e_0 = \theta_0 + \theta_1$ and $\theta_0\theta_1 = (v-1)e_0 - rk$. The expressions for the parameters of the underlying QSD \mathcal{Q} together with e_0 and e_1 follow from the above equations. Now we show that the polynomial p(u) has precisely one real zero. Fix $\theta_1 \ge 1$ and $\theta_2 \leqslant -2$. If $\theta_1 = -\theta_2 - 1$ then it is easy to check that p(u) = 0 has no positive roots. Otherwise, we can assume that $\theta_1 > -\theta_2 - 1$. Since $C_3(\theta_1, \theta_2) < 0$, $C_2(\theta_1, \theta_2) < 0$, $C_1(\theta_1, \theta_2) < 0$, and $C_3(\theta_1, \theta_2) > 0$, by Descartes' rule of signs, we have that p(u) has just one positive zero.

The converse follows from Theorem 5.9.

There do not appear to be any QSD parameters that satisfy Lemma 5.17. However, if we do not impose the nonregularity condition $v - 1 + r \neq k + e_0$ then one can obtain QSDs whose whole graph is strongly regular. These strongly regular graphs correspond to the complement of those of Remark 5.14.

5.3.4 Class 4

Lemma 5.18. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with coherent rank 9. Suppose that the underlying QSD \mathcal{Q} has parameters $(v, k, \lambda; b, r, \{x, y\})$ and $\mathsf{B}_x(\mathcal{Q})$ has eigenvalues $e_0 > e_1 > e_2$. Suppose that $\Gamma[V_1]$ is complete, $\theta_1 = e_1 = \frac{y-k}{x-y}$, and θ_2 is an eigenvalue of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$. Then θ_0 is a zero of the (univariate) polynomial $q(u) := h(u, \theta_2, \theta_1)$. Furthermore,

$$e_2 = \theta_1 + \theta_2 + 1, \text{ and } e_0 = \frac{(\theta_1 + 1)(\theta_0(\theta_2 e_2 - 1) + \theta_2^2(\theta_1 + 1))}{\theta_0(\theta_1 \theta_2 + e_2) + \theta_2^3(\theta_1 + 1) + 2\theta_2(\theta_2 + 1) - \theta_1(\theta_1 + 2)}.$$

The parameters of Q can be written in terms of θ_0 , θ_1 , and θ_2 as follows.

$$\begin{aligned} v &= \theta_0 + \theta_2 - e_0 + 1; \\ k &= y + \theta_1(\theta_1 + 1); \\ \lambda &= \theta_1 + 1 - \frac{\theta_2(y - e_2 - 1)}{\theta_1}; \\ y &= (\theta_1 + 1) \frac{\theta_0(e_2 - 1) - \theta_1(e_0 - t_2 - 1) - \theta_1^2}{\theta_0 - \theta_1}. \end{aligned}$$

Conversely, suppose that θ_0 is an integer solution to q(u) = 0 for some integers θ_1 and θ_2 satisfying $\theta_0 > \theta_1 > 0$ and $\theta_2 \leq -2$. If there exists a QSD Q whose parameters $(v, k, \lambda; b, r, \{x, y\})$ can be expressed as above in terms of θ_0 , θ_1 , and θ_2 then $W_x(Q) \in \mathcal{G}_3(\theta_0, \theta_1, \theta_2)$. Suppose $B_x(Q)$ has degree e_0 . If $v - 1 + r \neq k + e_0$ then $W_x(Q)$ has coherent rank 9.

Proof. Since $e_1 = \frac{y-k}{x-y}$, using Theorem 5.1, we must have $e_2 = \frac{r-\lambda-k+y}{x-y}$. We use the equations (3), (4), (14), (15), (16), (17) together with Theorem 5.1 and Theorem 2.8 (iii). Using the trace and determinant of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$, we obtain the equations $v - 1 + e_0 = \theta_0 + \theta_2$ and $\theta_0 \theta_2 = (v-1)e_0 - rk$. The system of equations can be obtained from that of the proof of Lemma 5.17 by interchanging θ_1 with θ_2 and e_1 with e_2 .

The converse follows from Theorem 5.9.

We are not aware of any QSD parameters that satisfy Lemma 5.18.

Lemma 5.19. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ with coherent rank 9. Suppose that the underlying $QSD \ \mathcal{Q}$ of Γ has parameters $(v, k, \lambda; b, r, \{x, y\})$ and $\mathsf{B}_x(\mathcal{Q})$ has eigenvalues $e_0 > e_1 > e_2$. Suppose that $\Gamma[V_1]$ is complete, $\theta_1 = e_1 = \frac{y-k}{x-y}$, and θ_1 is an eigenvalue of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$. Then $\theta_0 = f_3(\theta_2, \theta_1), e_2 = \theta_1 + \theta_2 + 1$, and

$$e_0 = \frac{\theta_1(\theta_2 + 1)}{(\theta_1 + 1)} \left(1 + \frac{\theta_1(1 - \theta_1\theta_2)}{\theta_0 + \theta_1\theta_2} \right).$$

Furthermore, the parameters of Q can be written in terms of θ_1 and θ_2 as follows.

$$\begin{aligned} v &= \theta_0 + \theta_1 - e_0 + 1; \\ k &= y + \theta_1(\theta_1 + 1); \\ \lambda &= (\theta_2 + 1) \left(1 + \frac{\theta_0 e_2 - \theta_2 e_0}{\theta_0 - \theta_2} \right); \\ y &= \frac{-\theta_1(\lambda - e_2 - 1)}{\theta_2}. \end{aligned}$$

Conversely, suppose that, for some integers θ_1 and θ_2 satisfying $\theta_1 \ge 1$ and $\theta_2 \le -2$, there exists a QSD Q whose parameters $(v, k, \lambda; b, r, \{x, y\})$ can be expressed as above in terms of θ_1 and θ_2 . Then $W_x(Q) \in \mathscr{G}_3(f_3(\theta_2, \theta_1), \theta_1, \theta_2)$ and $W_x(Q)$ has coherent rank 9.

Proof. Since $e_1 = \frac{y-k}{x-y}$, using Theorem 5.1, we must have $e_2 = \frac{r-\lambda-k+y}{x-y}$. We use the equations (3), (4), (14), (15), (16), (17) together with Theorem 5.1 and Theorem 2.8 (iii).

Using the trace and determinant of $Q_{\mathfrak{D}(\Gamma)}(\Gamma)$, we obtain the equations $v - 1 + e_0 = \theta_0 + \theta_1$ and $\theta_0 \theta_1 = (v - 1)e_0 - rk$. The system of equations can be obtained from that of the proof of Lemma 5.15 by interchanging θ_1 with θ_2 and e_1 with e_2 . In particular, we obtain the equation

$$\xi_2(\theta_2, \theta_1)\theta_0^2 + \xi_1(\theta_2, \theta_1)\theta_0 + \xi_0(\theta_2, \theta_1) = 0,$$

where ξ_0 , ξ_1 , and ξ_2 are as defined in the proof of Lemma 5.15. Think of the above equation as a quadratic univariate polynomial equation in θ_0 with coefficients in $\mathbb{Z}[\theta_1, \theta_2]$ and let $s_1(\theta_2, \theta_1) \ge s_2(\theta_2, \theta_1)$ be its roots. Note that $s_1(\theta_2, \theta_1) = f_3(\theta_2, \theta_1)$. Using the fact that $\theta_0 > \theta_1 \ge 0$, we can rule out the possibility of $\theta_0 = s_2(\theta_2, \theta_1)$. Suppose to the contrary that $\theta_0 = s_2(\theta_2, \theta_1)$. Then $\theta_0 > \theta_1 \ge 0$ and $\theta_2 < 0$ implies that

$$\theta_1\theta_2(\theta_1+2)(\theta_2+2) + 2\theta_1^2 + 4\theta_1 + 1 - (1-\theta_1\theta_2)\sqrt{g_3(\theta_2,\theta_1)} > -2(\theta_1+1)(\theta_1+\theta_2+2)\theta_1,$$

which implies

$$(\theta_1\theta_2(\theta_1+2)(\theta_2+2)+2\theta_1^2+4\theta_1+1+2(\theta_1+1)(\theta_1+\theta_2+2)\theta_1)^2-(1-\theta_1\theta_2)^2g_3(\theta_2,\theta_1)$$

= $-4\theta_1(\theta_1+1)(\theta_1+\theta_2+2)(\theta_1\theta_2(\theta_2^2(\theta_1-1)-3(\theta_1+2\theta_2+3))-\theta_1^3-5\theta_1^2-7\theta_1-1)$

is positive. Now, $-4\theta_1(\theta_1+1)(\theta_1+\theta_2+2)$ is positive since $\theta_1 \ge 0$ and $e_2 = \theta_1+\theta_2+1 \le -1$. However, together with the condition that $\theta_2 \le -2$, it also follows that

$$\theta_1\theta_2(\theta_2^2(\theta_1-1) - 3(\theta_1+2\theta_2+3)) - \theta_1^3 - 5\theta_1^2 - 7\theta_1 - 1$$

П

is negative, which contradicts our supposition. Hence, $\theta_0 = f_3(\theta_2, \theta_1)$, as required.

The converse follows from Theorem 5.9.

Suppose that Γ satisfies the assumptions of Lemma 5.19. By Theorem 2.5, we can assume that $\theta_0 > \theta_1 \ge 1$ and by Theorem 2.1, we can assume that $\theta_2 \le -2$. Since $\theta_1 + \theta_2 + 1 = e_2 \le -2$ and $\theta_1 \ge 1$, we find that $1 \le \theta_1 \le -\theta_2 - 3$. In Table 8, we list all the parameters for QSDs corresponding to Lemma 5.15 for $\theta_2 \ge -100$.

6 Triregular graphs with small coherent rank

In this section, we establish a lower bound for the coherent rank of a triregular graph in \mathscr{G}_3 . This bound has the potential to be sharp (see Example 6.6).

Theorem 6.1. Let $\Gamma \in \mathscr{G}_3$ be a triregular graph. Then the coherent closure $\mathcal{W}(\Gamma)$ has rank at least 14.

We will prove Theorem 6.1 after we state and prove a series of three lemmas that restrict the entries of the type matrix.

Lemma 6.2. Let $\Gamma \in \mathscr{G}_3$ be a triregular graph whose coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & t_{12} & t_{13} \\ t_{22} & t_{23} \\ t_{33} \end{bmatrix}$. Then $t_{12}t_{13} > 1$.

Proof. Suppose, for a contradiction, that the coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 2 & t_{12} & t_{23} \\ t_{22} & t_{23} \\ t_{33} \end{bmatrix}$ and suppose that $\mathfrak{D}(\Gamma) = \{V_1, V_2, V_3\}$. Then, by Corollary 3.2 (iii), the adjacency matrix A of Γ has the form

$$A = \begin{bmatrix} A(\Gamma[V_1]) & \varepsilon_2 J & \varepsilon_3 J \\ \varepsilon_2 J & A(\Gamma[V_2]) & \star \\ \varepsilon_3 J & \star & A(\Gamma[V_3]) \end{bmatrix},$$

where $\varepsilon_1, \varepsilon_2, \varepsilon_3 \in \{0, 1\}$. Suppose that $X = A(\Gamma[V_1]) = \varepsilon_1(J - I)$ has order n_1 and $\Gamma[V_2]$ and $\Gamma[V_3]$ have orders n_2 and n_3 , respectively. Then

$$A^{2} = \begin{bmatrix} X^{2} + \varepsilon_{2}n_{2}J + \varepsilon_{3}n_{3}J & \star & \star \\ \star & \star & \star \\ \star & \star & \star \end{bmatrix}.$$

Now, the top-left block of (1) becomes

$$X^{2} + \varepsilon_{2}n_{2}J + \varepsilon_{3}n_{3}J - (\theta_{1} + \theta_{2})X + \theta_{1}\theta_{2}I = (\varepsilon_{1}(n_{1} - 1) + n_{2} + n_{3} + \theta_{1}\theta_{2})J.$$

In the case where X = O, we have $\theta_1 \theta_2 = 0$, but this means that $\theta_1 = 0$ which, by Theorem 2.5, implies that Γ is a complete bipartite graph, which is a contradiction.

In the case where X = J - I, we have

$$(J - I)^{2} - (\theta_{1} + \theta_{2})(J - I) + \theta_{1}\theta_{2}I = O$$
$$J^{2} - 2J - (\theta_{1} + \theta_{2})J + I + (\theta_{1} + \theta_{2})I + \theta_{1}\theta_{2}I = O$$

which means $(\theta_1 + 1)(\theta_2 + 1) = 0$, but this contradicts (2).

Next, we restrict the product of the off-diagonal entries of the type matrix.

Lemma 6.3. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ be a triregular graph whose coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} t_{11} & t_{12} & t_{13} \\ t_{22} & t_{23} \\ t_{33} \end{bmatrix}$. Then $t_{12}t_{13}t_{23} > 1$.

Proof. Suppose, for a contradiction, that $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} t_{11} & 1 & 1 \\ t_{22} & 1 \\ t_{33} \end{bmatrix}$ and suppose that $\mathfrak{D}(\Gamma) = \{V_1, V_2, V_3\}$. Then, by Corollary 3.2 (iii), the adjacency matrix A of Γ has the form

$$A = \begin{bmatrix} A(\Gamma[V_1]) & \varepsilon_{12}J & \varepsilon_{13}J \\ \varepsilon_{12}J & A(\Gamma[V_2]) & \varepsilon_{23}J \\ \varepsilon_{13}J & \varepsilon_{23}J & A(\Gamma[V_3]) \end{bmatrix},$$

where $\varepsilon_{12}, \varepsilon_{13}, \varepsilon_{23} \in \{0, 1\}$ and $\Gamma[V_1], \Gamma[V_2]$, and $\Gamma[V_3]$ are regular graphs orders n_1, n_2 , and n_3 , respectively. First, suppose that $\varepsilon_{12}\varepsilon_{13}\varepsilon_{23} = 0$. Without loss of generality, we can assume that $\varepsilon_{23} = 0$. Since Γ is connected, we must have $\varepsilon_{12} = \varepsilon_{13} = 1$. Thus, the complement of Γ is disconnected. By Theorem 2.2, Γ must be a cone. Hence, $\Gamma[V_1] = K_{n_1}$ and by equating coefficients of I in the (1, 1)-block of (1) yields $(\theta_1 + 1)(\theta_2 + 1) = 0$, which contradicts (2).

Lastly, suppose $\varepsilon_{12} = \varepsilon_{13} = \varepsilon_{23} = 1$. The same argument establishes a contradiction, as required.

Our final lemma rules out the possibility of the type matrix $\begin{vmatrix} 1 & 1 & 1 \\ 2 & 2 \\ 2 \end{vmatrix}$.

Lemma 6.4. Let $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ be a triregular graph whose coherent closure $\mathcal{W}(\Gamma)$ has $type \begin{bmatrix} t_{11} & 1 & 1 \\ & 2 & 2 \\ & 2 \end{bmatrix}$. Then $t_{11} > 1$.

Proof. Suppose, for a contradiction, that the coherent closure $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{2} \\ 2 & 2 \end{bmatrix}$ and suppose that $\mathfrak{D}(\Gamma) = \{V_1, V_2, V_3\}$. Then, by Proposition 3.3 and Corollary 3.2 (iii), the adjacency matrix A of Γ has the form

$$A = \begin{bmatrix} 0 & k_{21} \mathbf{1}^{\mathsf{T}} & k_{31} \mathbf{1}^{\mathsf{T}} \\ k_{21} \mathbf{1} & A(\Gamma[V_2]) & M \\ k_{31} \mathbf{1} & M^{\mathsf{T}} & A(\Gamma[V_3]) \end{bmatrix},$$

where $\Gamma[V_1] = K_1$ and $\Gamma[V_2]$ and $\Gamma[V_3]$ are both empty or complete graphs of orders n_2 and n_3 , respectively. Let $A_2 = A(\Gamma[V_2])$, $A_3 = A(\Gamma[V_3])$, let $k_{12} = k_{21}n_2$, $k_{13} = k_{31}n_3$, and let k_{22} , k_{23} , k_{32} , and k_{33} satisfy $A_2\mathbf{1} = k_{22}\mathbf{1}$, $M\mathbf{1} = k_{23}\mathbf{1}$, $M^{\mathsf{T}}\mathbf{1} = k_{32}\mathbf{1}$, and $A_3\mathbf{1} = k_{33}\mathbf{1}$. Then

$$A^{2} = \begin{bmatrix} k_{12} + k_{13} & (k_{21}k_{22} + k_{31}k_{23})\mathbf{1}^{\mathsf{T}} & (k_{21}k_{32} + k_{31}k_{33})\mathbf{1}^{\mathsf{T}} \\ (k_{21}k_{22} + k_{31}k_{23})\mathbf{1} & k_{21}J + A_{2}^{2} + MM^{\mathsf{T}} & k_{21}k_{31}J + A_{2}M + MA_{3} \\ (k_{21}k_{32} + k_{31}k_{33})\mathbf{1} & k_{21}k_{31}J + M^{\mathsf{T}}A_{2} + A_{3}M^{\mathsf{T}} & k_{31}J + M^{\mathsf{T}}M + A_{3}^{2} \end{bmatrix}.$$

Apply (1) to obtain

$$A^{2} - (\theta_{1} + \theta_{2})A + \theta_{1}\theta_{2}I = \begin{bmatrix} \alpha_{1}^{2}J & \alpha_{1}\alpha_{2}J & \alpha_{1}\alpha_{3}J \\ \alpha_{1}\alpha_{2}J & \alpha_{2}^{2}J & \alpha_{2}\alpha_{3}J \\ \alpha_{1}\alpha_{3}J & \alpha_{2}\alpha_{3}J & \alpha_{3}^{2}J \end{bmatrix}, \text{ where}$$
(23)

 $\alpha_1 = \sqrt{k_{12} + k_{13} + \theta_1 \theta_2}, \ \alpha_2 = \sqrt{k_{21} + k_{22} + k_{23} + \theta_1 \theta_2}, \ \text{and} \ \alpha_3 = \sqrt{k_{31} + k_{32} + k_{33} + \theta_1 \theta_2}.$ From the centre block of (23), we obtain

$$MM^{\mathsf{T}} = \begin{cases} (k_{23} + \theta_1 \theta_2) J_{n_2} - \theta_1 \theta_2 I, & \text{if } \Gamma[V_2] = \overline{K}_{n_2}; \\ (k_{23} + (\theta_1 + 1)(\theta_2 + 1)) J_{n_2} - (\theta_1 + 1)(\theta_2 + 1)I, & \text{if } \Gamma[V_2] = K_{n_2}. \end{cases}$$
(24)

Similarly, we can use the bottom-right block of (23) to deduce that

$$M^{\mathsf{T}}M = \begin{cases} (k_{32} + \theta_1 \theta_2) J_{n_3} - \theta_1 \theta_2 I, & \text{if } \Gamma[V_3] = \overline{K}_{n_3}; \\ (k_{32} + (\theta_1 + 1)(\theta_2 + 1)) J_{n_3} - (\theta_1 + 1)(\theta_2 + 1)I, & \text{if } \Gamma[V_3] = K_{n_3}. \end{cases}$$
(25)

Using (24) and (25) together with Lemma 4.1, we find that $n_2 = n_3$. By double counting edges between V_2 and V_3 , we find that $k_{23} = k_{32}$.

First, we assume that both $\Gamma[V_2]$ and $\Gamma[V_3]$ are empty, that is, $k_{22} = 0$ and $k_{33} = 0$. Since Γ is triregular, we cannot have both $k_{21} = k_{31} = 1$. Without loss of generality, since Γ is connected, we assume that $k_{21} = 1$ and $k_{31} = 0$. Then the valency-partition $\mathfrak{D}(\Gamma)$, which, by Theorem 2.7, is equitable, has quotient matrix

$$Q = Q_{\mathfrak{D}(\Gamma)}(\Gamma) = \begin{bmatrix} 0 & n_1 & 0 \\ 1 & 0 & k_{23} \\ 0 & k_{32} & 0 \end{bmatrix}.$$

However, $\det Q = 0$, which is impossible, by Lemma 2.6 and Corollary 3.7.

Next, we assume that both $\Gamma[V_2]$ and $\Gamma[V_3]$ are complete, that is, $k_{22} = n_2 - 1$ and $k_{33} = n_3 - 1$. Since Γ is triregular, we cannot have both $k_{21} = k_{31} = 1$. Without loss of generality, since Γ is connected, we assume that $k_{21} = 1$ and $k_{31} = 0$. Furthermore, using (24) and (25), we deduce that M is the incidence matrix of a symmetric design with parameters (v, k, λ) , where $v = n_2 = n_3$, $k = k_{23}$, and $\lambda = k + (\theta_1 + 1)(\theta_2 + 1)$. Thus, we obtain

$$(\theta_1 + 1)(\theta_2 + 1) = \lambda - k.$$
(26)

The valency-partition $\mathfrak{D}(\Gamma)$, which, by Theorem 2.7, is equitable, has quotient matrix

$$Q = Q_{\mathfrak{D}(\Gamma)}(\Gamma) = \begin{bmatrix} 0 & v & 0 \\ 1 & v - 1 & k \\ 0 & k & v - 1 \end{bmatrix}.$$

In this case $\det(xI-Q) = x^3 - 2(v-1)x^2 + (v^2 - k^2 - 3v + 1)x + v(v-1)$. Since $\det(Q) < 0$, using Lemma 2.6, it follows that spec $Q = \{[\theta_0]^1, [\theta_1]^1, [\theta_2]^1\}$. Thus, we obtain

$$\theta_0 + \theta_1 + \theta_2 = 2(v-1); \tag{27}$$

$$\theta_0(\theta_1 + \theta_2) + \theta_1\theta_2 = v^2 - k^2 - 3v + 1;$$
(28)

$$\theta_0 \theta_1 \theta_2 = -v(v-1). \tag{29}$$

Putting (26), (27), (28), and (29) together with (8) and the inequalities $\theta_0 > \theta_1 \ge 1$, and $\theta_2 \le -2$ results in no solutions.

Lastly, we assume that $\Gamma[V_2]$ is empty and $\Gamma[V_3]$ is complete, i.e., $k_{22} = 0$ and $k_{33} = n_3 - 1$. Furthermore, again using (24) and (25), we deduce that M is the incidence matrix of a symmetric design with parameters (v, k, λ) , where $v = n_2 = n_3$, $k = k_{23}$, and $\lambda = k + \theta_1 \theta_2 = k + (\theta_1 + 1)(\theta_2 + 1)$. Thus, we obtain

$$\theta_1 + \theta_2 = -1; \tag{30}$$

$$\theta_1 \theta_2 = \lambda - k. \tag{31}$$

The valency-partition $\mathfrak{D}(\Gamma)$, which, by Theorem 2.7, is equitable, with quotient matrix

$$Q = Q_{\mathfrak{D}(\Gamma)}(\Gamma) = \begin{bmatrix} 0 & k_{21}v & k_{31}v \\ k_{21} & 0 & k \\ k_{31} & k & v-1 \end{bmatrix}.$$

Since det $Q = k_{21}v(1 - v + 2k_{31}k)$, by Lemma 2.6 and Theorem 2.5, we must have $k_{21} = 1$. There remain two cases to consider: $k_{31} = 0$ and $k_{31} = 1$.

Consider the case $k_{31} = 0$, i.e.,

$$Q = \begin{bmatrix} 0 & v & 0 \\ 1 & 0 & k \\ 0 & k & v - 1 \end{bmatrix}.$$

Since det Q = -v(v-1) < 0, by Lemma 2.6, we must have that θ_0 , θ_1 , and θ_2 are eigenvalues of Q. Thus, we obtain

$$\theta_0 + \theta_1 + \theta_2 = v - 1;$$
 (32)

$$\theta_0 \theta_1 \theta_2 = -v(v-1). \tag{33}$$

Putting equations (30), (31), (32), and (33) together with equation (8) and the inequalities $\theta_0 > \theta_1 \ge 1$, and $\theta_2 \le -2$ results in no solutions.

Now, consider the case $k_{31} = 1$, i.e.,

$$Q = \begin{bmatrix} 0 & v & v \\ 1 & 0 & k \\ 1 & k & v - 1 \end{bmatrix}.$$

In this case $\det(xI - Q) = x^3 - (v - 1)x^2 - (2v + k^2)x + v(v - 1 - 2k)$. Since 2v + k > 0, using Lemma 2.6, it follows that either spec $Q = \{[\theta_0]^1, [\theta_1]^1, [\theta_2]^1\}$ or spec $Q = \{[\theta_0]^1, [\theta_2]^2\}$. Thus, we obtain

$$\theta_0 + \theta' + \theta_2 = v - 1; \tag{34}$$

$$\theta_0(\theta' + \theta_2) + \theta'\theta_2 = -k^2 - 2v; \tag{35}$$

$$\theta_0 \theta' \theta_2 = v(2k - v + 1) \tag{36}$$

corresponding to the two possible spectra of Q, corresponding to $\theta' \in \{\theta_1, \theta_2\}$. We can check for integer solutions for both of the resulting systems of equations. For $\theta' = \theta_1$, putting (30), (31), (34), (35), and (36) together with (8) and the inequalities $\theta_0 > \theta_1 \ge 1$, and $\theta_2 \le -2$ results in no solutions. Finally, For $\theta' = \theta_2$, putting (30), (31), (34), (35), and (36) together with (8) and the inequalities $\theta_0 > \theta_1 \ge 1$, and $\theta_2 \le -2$ results in no solutions. \Box

Now we are ready to prove Theorem 6.1.

Proof of Theorem 6.1. Suppose the rank of $\mathcal{W}(\Gamma)$ has rank less than 14. By Lemma 3.1 together with Lemma 6.2, Lemma 6.3, and Lemma 6.4, $\mathcal{W}(\Gamma)$ must have type $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$. It is straightforward to check that this is not possible.

Valencies	Spectrum	Coherent rank	Reference	
$\{[45]^1, [25]^{18}, [13]^{27}\}$	$\left\{ [21]^1, [3]^{19}, [-3]^{26} \right\}$	16	[1]	
$\left\{ [15]^4, [10]^{16}, [7]^4 \right\}$	$\{[11]^1, [3]^7, [-2]^{16}\}$	18	[12]	
$\left\{ [96]^1, [61]^{64}, [21]^{32} \right\}$	$\{[56]^1, [4]^{41}, [-4]^{55}\}$	20	[1]	
$\{[24]^{18}, [14]^9, [8]^9\}$	$\left\{ [20]^1, [2]^{17}, [-3]^{18} \right\}$	29	[13]	
$\left\{ [24]^{18}, [14]^9, [8]^9 \right\}$	$\left\{ [20]^1, [2]^{17}, [-3]^{18} \right\}$	240	[7]	
$\left\{ [35]^1, [26]^7, [19]^{35} \right\}$	$\left\{ [21]^1, [\frac{-1\pm\sqrt{41}}{2}]^{21} \right\}$	949	[13]	
$\left\{ [35]^1, [26]^7, [19]^{35} \right\}$	$\left\{ [21]^1, [\frac{-1\pm\sqrt{41}}{2}]^{21} \right\}$	1849	[13]	

Table 3: List of known connected graphs that have three distinct eigenvalues and three distinct valencies.

In Table 3, we list all currently known examples of connected graphs that have precisely three distinct eigenvalues and three distinct valencies. We note that, among these graphs, the smallest coherent rank is 16, which invites the question of whether the lower bound of Theorem 6.1 can be improved.

In view of the following result, we denote by $\widehat{\Gamma}$ the cone over the graph Γ .

Theorem 6.5. Let θ_1 and θ_2 be integers such that $\theta_1 \ge 1$ and $\theta_2 \le -2$. Suppose Q is a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$ where $\mathsf{B}_x(Q) \in \mathscr{G}_3(e_0, e_1, e_2)$ and

$$(x,y) = (k + (\theta_1 + 1)(\theta_2 + 1) - (e_1 + 1)(e_2 + 1), k + \theta_1\theta_2 - e_1e_2).$$

$$\lambda = r + \theta_1\theta_2, \ \frac{r - \lambda + y - k}{x - y} = \theta_1 + \theta_2, \ and$$

$$k + e_0 - \theta_1 - \theta_2 = \sqrt{(v + b + \theta_1 \theta_2)(1 + k + e_0 + \theta_1 \theta_2)}$$
$$r - \theta_1 - \theta_2 = \sqrt{(v + b + \theta_1 \theta_2)(1 + \lambda)}$$
$$1 + k + \frac{\lambda k - yr}{x - y} = \sqrt{(1 + \lambda)(1 + k + e_0 + \theta_1 \theta_2)}$$

then $\widehat{\mathsf{T}_x(\mathcal{Q})} \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ where θ_0 is the largest eigenvalue of $\begin{bmatrix} 0 & v & b \\ 1 & 0 & r \\ 1 & k & e_0 \end{bmatrix}$. Furthermore, if $|\{v+b, 1+r, 1+k+e_0\}| = 3$ then $\widehat{\mathsf{T}_x(\mathcal{Q})}$ has coherent rank 14.

(*ii*) If $\lambda = r + (\theta_1 + 1)(\theta_2 + 1)$, $\frac{r - \lambda + y - k}{x - y} = \theta_1 + \theta_2 + 1$, and

$$k + e_0 - \theta_1 - \theta_2 = \sqrt{(v + b + \theta_1 \theta_2)(1 + k + e_0 + \theta_1 \theta_2)}$$
$$v - 1 + r - \theta_1 - \theta_2 = \sqrt{(v + b + \theta_1 \theta_2)(v + r + \theta_1 \theta_2)}$$
$$1 + k + \frac{\lambda k - yr}{x - y} = \sqrt{(v + r + \theta_1 \theta_2)(1 + k + e_0 + \theta_1 \theta_2)}$$

then $\widehat{W_x(Q)} \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$, where θ_0 is the largest eigenvalue of $\begin{bmatrix} 0 & v & b \\ 1 & v-1 & r \\ 1 & k & e_0 \end{bmatrix}$. Furthermore, if $|\{v+b, v+r, 1+k+e_0\}| = 3$ then $\widehat{W_x(Q)}$ has coherent rank 14.

Proof. The proofs that $\widehat{\mathsf{T}_x(\mathcal{Q})}$ and $\widehat{\mathsf{W}_x(\mathcal{Q})}$ each have precisely three distinct eigenvalues is a modification of the proof of Theorem 5.9, which we leave to the reader. The assumptions also imply that $\widehat{\mathsf{T}_x(\mathcal{Q})}$ and $\widehat{\mathsf{W}_x(\mathcal{Q})}$ are triregular. By Theorem 6.1, the coherent rank of Γ is at least 14. Let M be the incidence matrix of \mathcal{Q} . The upper bound of 14 follows since $\mathcal{W}(\Gamma)$ is a subalgebra of the algebra

which is a coherent algebra of rank 14.

(*i*) If

A cursory search of possible QSD parameters that satisfy Theorem 6.5 resulted in just one possibility.

Example 6.6. Suppose Q is a QSD with parameters (85, 35, 34; 204, 84, {10, 15}). Then Q satisfies Theorem 6.5 with $\theta_1 = 4$ and $\theta_2 = -11$. Furthermore, by Theorem 6.5, we have $\widehat{W_{10}(Q)} \in \mathscr{G}_3(119, 4, -11)$. In particular, $\widehat{W_{10}(Q)}$ has spectrum {[119]¹, [4]²⁰⁴, [-11]⁸⁵}, its degree sequence is {[289]¹, [169]⁸⁵, [64]²⁰⁴} and its coherent rank is 14.

The existence of a QSD corresponding to Example 6.6 is an open problem (see Question 8.6).

7 Large coherent rank

In the previous sections, we have been concerned with graphs in \mathscr{G}_3 that have small coherent rank. We now turn our attention to large coherent rank. Before our work, the largest known coherent rank of a graph in \mathscr{G}_3 has coherent rank at most 19^6 . This graph, which was discovered by Van Dam [12, Section 2.2] was obtained by taking a strongly regular graph on 19^4 vertices and interchanging some of its edges with nonedges (see *switching*, below). Furthermore, all known infinite families of graphs in \mathscr{G}_3 have a fixed coherent rank.

In this section, we exhibit a conjecturally infinite family of biregular graphs in \mathscr{G}_3 (see Remark 7.8) whose coherent ranks can conjecturally become arbitrarily large (see Question 8.1).

7.1 Switching strongly regular graphs

Let $\Gamma = (V, E)$ be a graph and let $\sigma = \{U, V - U\}$ be a partition of the vertex set V. We can write the adjacency matrix of Γ in block form as

$$A(\Gamma) = \begin{bmatrix} A(\Gamma[U]) & M \\ M^{\mathsf{T}} & A(\Gamma[V-U]) \end{bmatrix}.$$

Define the (switched) graph Γ^{σ} via its adjacency matrix as

$$A(\Gamma^{\sigma}) = \begin{bmatrix} A(\Gamma[U]) & J - M \\ J - M^{\mathsf{T}} & A(\Gamma[V - U]) \end{bmatrix}.$$

We say that the graph Γ^{σ} was obtained by **switching** the graph Γ with respect to the vertex-partition σ . We refer to Γ^{σ} as the **switched** graph. Muzychuk and Klin [23] used the following tool to study the spectrum of a graph obtained by switching some other graph.

Proposition 7.1 ([23, Corollary 3.2]). Let σ be an equitable 2-partition of a graph Γ . Then

$$\operatorname{spec}\left(A(\Gamma^{\sigma})\right) = \operatorname{spec}(A(\Gamma)) \cup \operatorname{spec}(Q_{\sigma}(\Gamma^{\sigma})) - \operatorname{spec}(Q_{\sigma}(\Gamma)).$$

In view of Proposition 7.1, to find a graph with three distinct eigenvalues, it suffices to find a strongly regular graph $\Gamma = (V, E)$ and an equitable partition $\sigma = \{U, V - U\}$ such that spec $(Q_{\sigma}(\Gamma^{\sigma}))$ shares an eigenvalue with Γ . This is how we proceed with a focus on the family of block graphs of orthogonal arrays.

7.2 Block graphs of orthogonal arrays

For $m \ge 2$ and $n \ge 1$, an **orthogonal array** OA(m, n) is an $m \times n^2$ matrix M with entries from the set $\{1, \ldots, n\}$ such that the n^2 columns of each $2 \times n^2$ submatrix of M contain all n^2 ordered pairs of elements of $\{1, \ldots, n\}$.

Theorem 7.2 ([30, Theorems 6.39 and 6.40]). Let n be a prime power and $2 \le m \le n+1$. Then there exists an OA(m, n).

We will only consider the construction implicit from Theorem 7.2 (see also [10, Construction 3.29]). However, there exist other constructions of orthogonal arrays [30, Section 6.5]. We denote by OA(m, n) the graph whose vertices are the columns of OA(m, n) where two columns are adjacent if there is a row where they share the same entry. The graph OA(m, n) is commonly known as the **block graph of an orthogonal array** and is known as the *Latin squares graph* in [12], therein denoted by $L_m(n)$.

Theorem 7.3 ([16, Section 5.5]). Let $m \ge 2$ and $n \ge 1$ be integers satisfying $m \le n+1$. Then the graph OA(m, n) has spectrum

$$\left\{ [m(n-1)]^1, [n-m]^{m(n-1)}, [-m]^{(n-1)(n+1-m)} \right\}.$$

Given an orthogonal array OA(m, n) and $i \in \{1, ..., n\}$, let C_i be the set of columns of OA(m, n) that have the entry 1 in the *i*th row. Then C_i is a clique of order n in the graph OA(m, n). Furthermore, the cliques C_i and C_j are vertex disjoint for any $i \neq j$.

7.3 A new family of biregular graphs with three distinct eigenvalues

For $m \ge 2$ and $n \ge 1$, let $\Gamma = \mathsf{OA}(m, n)$ and let $\sigma = \{U, V - U\}$ be a partition of the vertex set $V(\Gamma)$ where $U = C_1 \cup \cdots \cup C_N$. Denote by $\mathsf{S}_N(m, n)$ the switched graph Γ^{σ} .

Lemma 7.4. Let $n \in \mathbb{N}$ and $2 \leq m \leq n+1$. Suppose $1 \leq N \leq n-1$. Then $S_N(m,n)$ has three distinct eigenvalues if and only if

$$N = \frac{n}{2} \pm \frac{\sqrt{(n-2m)(n-2m+2)(n+2)n}}{2(n-2m+2)}.$$

Moreover, $S_N(m,n)$ has spectrum

$$\left\{ [n+m(n-1)]^1, [n-m]^{m(n-1)-1}, [-m]^{(n-1)(n+1-m)+1} \right\}$$

Proof. The quotient matrix

$$Q_{\sigma}(\Gamma) = \begin{bmatrix} n-1+(N-1)(m-1) & (n-N)(m-1) \\ N(m-1) & n-1+(n-N-1)(m-1) \end{bmatrix}$$

has eigenvalues m(n-1) and n-m. Using Theorem 7.3 and Proposition 7.1, for $S_N(m,n)$ to have three distinct eigenvalues, we require -m to be an eigenvalue of the quotient matrix

$$Q_{\sigma}(\Gamma^{\sigma}) = \begin{bmatrix} n - 1 + (N - 1)(m - 1) & (n - N)(n - m + 1) \\ N(n - m + 1) & n - 1 + (n - N - 1)(m - 1) \end{bmatrix}.$$

It is straightforward to check that -m is a eigenvalue of $Q_{\sigma}(\Gamma^{\sigma})$ if and only if

$$N = \frac{n}{2} \pm \frac{\sqrt{(n-2m)(n-2m+2)(n+2)n}}{2(n-2m+2)},$$

For such N, the quotient matrix $Q_{\sigma}(\Gamma^{\sigma})$ has eigenvalues n + m(n-1) and -m.

In view of Lemma 7.4, for $m \ge 2$ and $n \ge 1$ where $N = \frac{n}{2} - \frac{\sqrt{(n-2m)(n-2m+2)(n+2)n}}{2(n-2m+2)}$ is a positive integer less than n, we define the graph $\mathsf{G}(m,n) := \mathsf{S}_N(m,n)$. Clearly, in order for $\mathsf{G}(m,n)$ to exist, we require the existence of an orthogonal array $\mathsf{OA}(m,n)$. We now aim to find integers m and n such that $\mathsf{G}(m,n)$ exists.

Lemma 7.5. Suppose that G(m,n) exists for some $n \ge 2$ and m. Then $\frac{n}{3} < m \le \frac{n}{2}$.

Proof. The proof follows from the condition that $N = \frac{n}{2} - \frac{\sqrt{(n-2m)(n-2m+2)(n+2)n}}{2(n-2m+2)}$ is a positive integer less than n. Suppose n - 2m < -2. Then

$$-n(n-2m+2) < \sqrt{(n-2m)(n-2m+2)(n+2)n},$$

which implies that $N \ge n$, a contradiction. Hence, n - 2m > -2. Furthermore, since $(n-2m)(n-2m+2)(n+2)n \ge 0$, we must have $n-2m \ge 0$. Now, using the requirement that $N \ge 1$, we obtain $3m \ge n + \frac{5n-6}{3n-2}$, as required.

Note that when n = 2m, the graph G(m, n) is regular, and hence, strongly regular. Now, we will consider the special case when n = 2m + 1. In this case, using Theorem 7.2, to show that $G(\frac{n-1}{2}, n)$ exists, it suffices to take n to be a prime power such that $n/2 - \sqrt{3n(n+2)}/6$ is an integer. Let $(a_k)_{k \in \mathbb{N}}$ be the recurrence sequence defined by $a_k = 4a_{k-1} - a_{k-2}$ with initial conditions $a_0 = 1$, $a_1 = 5$.

Lemma 7.6. For each $k \in \mathbb{N}$, we have that $3(a_k^2 + 2) = (a_{k+1} - 2a_k)^2$ is an odd square.

Proof. It is routine to check that $(a_{k+1}-2a_k+a_k\sqrt{3})(a_{k+1}-2a_k-a_k\sqrt{3})=6$, as required. \Box

If $n = a_k^2$ for some $k \in \mathbb{N}$ then, by Lemma 7.6, $\frac{n}{2} - \frac{\sqrt{3n(n+2)}}{6}$ is an integer. Thus, using Theorem 7.2, each prime power in the sequence $(a_k)_{k \in \mathbb{N}}$ can produce a biregular graph with three distinct eigenvalues.

Corollary 7.7. Let $k \in \mathbb{N}$. Suppose that $q = a_k$ is a prime power. Then $G\left(\frac{q^2-1}{2}, q^2\right)$ is biregular and has spectrum

$$\left\{ \left[\frac{q^4+1}{2}\right]^1, \left[\frac{q^2+1}{2}\right]^{\frac{q^4-2q^2-1}{2}}, \left[\frac{1-q^2}{2}\right]^{\frac{q^4+2q^2-1}{2}} \right\}.$$

There are only finitely many terms of the form p^e in the sequence $(a_k)_{k\in\mathbb{N}}$ where e > 1and p is a prime [27]. The (probable) primes in the sequence $(a_k)_{k\in\mathbb{N}}$ are listed in OEIS sequence A299107 [22].

Remark 7.8. It is conjectured [21, Conjecture 47] that $(a_k)_{k\in\mathbb{N}}$ contains infinitely many primes. By Corollary 7.7, the validity of this conjecture implies the existence of a new infinite family of biregular graphs with three distinct eigenvalues: $G\left(\frac{b_k^2-1}{2}, b_k^2\right)$ where $(b_k)_{k\in\mathbb{N}}$ is the subsequence of primes in $(a_k)_{k\in\mathbb{N}}$.

Note that Van Dam [12] already discovered two instances of Corollary 7.7, that is, G(12, 25) and G(180, 361). In the same paper, Van Dam also discovered the graphs G(7, 16) and G(120, 243). One can show, using elementary number theoretic arguments, that G(7, 16) is the only graph of the form $G(2^{k-1} - 1, 2^k)$. We leave this to the reader to verify.

8 Conclusion and open problems

We conclude with a selection of questions that arise naturally from our investigations.

Question 8.1. Can graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ have arbitrarily large coherent rank?

All currently known infinite families in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ have a fixed coherent rank. However, we have empirical evidence that suggests that the coherent rank of the graphs G(m, n)grows with n. In particular, G(7, 16) has coherent rank 2048 and G(12, 25) has coherent rank $25^3 = 15625$. Note that we are assuming the construction implicit in Theorem 7.2 alternative constructions of orthogonal arrays could potentially produce different coherent ranks. We conjecture that if an infinite family of graphs G(m, n) exists then the answer to Question 8.1 is yes.

Question 8.2. Does there exist $\Gamma \in \mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ such that $\mathcal{W}(\Gamma)$ has rank 11?

By Theorem 3.4 together with Example 4.4 and Example 5.11, we observe that there exist infinite families of graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ having coherent ranks 3, 5, 6, 8, and 9. We know that no graph in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ can have coherent rank 4 or 7, by Theorem 3.4 and Theorem 3.6. Van Dam gave a construction of an infinite family of graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ that have coherent rank 10, using a certain combination of symmetric designs and distance regular graphs [12, Section 6.1]. There are currently no known constructions of graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ that have coherent rank 11. In Table 4, we list the known realisable coherent ranks from graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$.

r	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Exists	Y	Ν	Υ	Υ	Ν	Υ	Y	Y	?	Y	Y	?	?	Y	?	Y	?	Y

Table 4: The existence of graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ having coherent rank r for $3 \leq r \leq 20$.

Coherent ranks 16, 18, and 20 correspond to triregular graphs in Table 3. We also have examples of graphs in $\mathscr{G}_3(\theta_0, \theta_1, \theta_2)$ that have coherent ranks 12 and 13.

Example 8.3. Let $\Gamma \in \text{SRG}(40, 12, 2, 4)$. The cardinality of SRG(40, 12, 2, 4) is 28 [29]. Fix a vertex $\mathbf{v} \in V(\Gamma)$ such that $N(\mathbf{v})$ is the disjoint union of four copies of K_3 . Now partition $N(\mathbf{v}) = U \cup W$ such that U and W are disjoint subsets of cardinality 6 and there are no edges between vertices in U and vertices in W. Obtain a new graph on 39 vertices by adding all edges between vertices of U and W and deleting the vertex \mathbf{v} . The resulting graph Δ has degree sequence $\{[12]^{27}, [17]^{12}\}$ and spectrum $\{[14]^1, [2]^{23}, [-4]^{15}\}$. This method can produce 55 pairwise non-isomorphic graphs of which 27 have coherent rank 12.

Moreover, each graph can be switched with respect to its valency partition to produce a graph with degree sequence $\{[16]^{27}, [26]^{12}\}$ and spectrum $\{[20]^1, [2]^{22}, [-4]^{16}\}$. The coherent rank of switched graphs obtained from graphs with coherent rank 12 remains equal to 12. The type of the coherent closure of each of these graphs is $\begin{bmatrix} 4 & 2 \\ 4 & 4 \end{bmatrix}$.

Example 8.4. Let \mathcal{D} be a 2-(45, 12, 3) design that possesses a polarity with 36 absolute points, that is, \mathcal{D} has a symmetric incidence matrix M whose diagonal contains 36 entries equal to 1. Suppose that, in addition, the 9×9 principal submatrix N of M induced on the rows/columns with a 0 on the diagonal is equal to the zero matrix O. Replace all the entries of M in this 9×9 submatrix by 1s. Now, form the adjacency matrix A by setting its off-diagonal entries equal to those of M and its diagonal entries equal to

0. The resulting graph whose adjacency matrix is A has degree sequence $\{[11]^{36}, [20]^9\}$ and spectrum $\{[14]^1, [2]^{27}, [-4]^{17}\}$. Using the list of 2-(45, 12, 3) designs from [11], we can produce nine pairwise non-isomorphic graphs of which four have coherent rank 13.

Moreover, each graph can be switched with respect to its valency partition to produce a graph with degree sequence $\{[14]^{36}, [32]^9\}$ and spectrum $\{[20]^1, [2]^{26}, [-4]^{18}\}$. The coherent rank of switched graphs obtained from graphs with coherent rank 13 remains equal to 13. The type of the coherent closure of each of these graphs is $[2^2 \frac{7}{7}]$.

Question 8.5. Does there exist $\Gamma \in \mathscr{G}_3$ such that $\mathcal{W}(\Gamma)$ has type $\begin{bmatrix} 3 & 2 \\ 3 & 3 \end{bmatrix}$?

Both constructions of Van Dam [12, Section 6.1 and Section 6.2] produce graphs whose coherent closure have type $\begin{bmatrix} 2 & 2 \\ 4 \end{bmatrix}$. Coherent configurations that have type $\begin{bmatrix} 3 & 2 \\ 3 \end{bmatrix}$ are known as *strongly regular designs* [20]. We do not know if any graphs in \mathscr{G}_3 have a coherent closure equal to the adjacency algebra of a strongly regular design.

Question 8.6. Does there exist a QSD with parameters $(85, 35, 34; 204, 84, \{10, 15\})$?

Using Example 6.6, the existence of such a QSD would show that the bound in Theorem 6.1 is sharp.

Question 8.7. Does there exist a graph in $\Gamma \in \mathscr{G}_3$ that has coherent rank 9 and class 4?

We have examples of graphs in \mathscr{G}_3 having coherent rank 9 in classes 1, 2, and 3. However, we are not aware of any example of a graph in class 4.

Question 8.8. Does there exist a graph in $\Gamma \in \mathscr{G}_3$ that satisfies Lemma 5.17?

Example 5.16 is an example of a graph in class 3. This graph is in $\mathscr{G}_3(21, 5, -2)$ and its smallest eigenvalue -2 is an eigenvalue of the quotient matrix of its valency partition. A *priori*, it is possible for the second largest eigenvalue of a graph of class 3 to be an eigenvalue of the quotient matrix of its valency partition. Such graphs must satisfy the assumption of Lemma 5.17. However, we are not aware of any such example.

9 Acknowledgements

The first author is grateful to Saveliy Skressanov for informing us about the stabilization routine due to Sven Reichard [25], to Edwin van Dam, Bill Martin, Misha Klin and Misha Muzychuk for their comments and information about the history of Haemer's question. We have also benefited from insightful conversations with Ian Wanless and Vedran Krčadinac.

The first author was partially supported by the Singapore Ministry of Education Academic Research Fund; grant numbers: RG18/23 (Tier 1) and MOE-T2EP20222-0005 (Tier 2).

References

- W.G. Bridges and R.A. Mena, Multiplicative cones —a family of three eigenvalue graphs, Aequationes Math. 22 (1981), no. 1, 208–214.
- [2] A.E. Brouwer and W.H. Haemers, Spectra of graphs, Springer, 1990.
- [3] A.E. Brouwer and H. Van Maldeghem, *Strongly regular graphs*, Cambridge University Press, 2022.

- [4] D. de Caen, E.R. van Dam, and E. Spence, A nonregular analogue of conference graphs, J. Combin. Theory Ser. A 88 (1999), 194–204.
- [5] A.R. Calderbank, The application of invariant theory to the existence of quasisymmetric designs, J. Combin. Theory Ser. A 44 (1987), no. 1, 94–109.
- [6] _____, Geometric invariants for quasi-symmetric designs, J. Combin. Theory Ser. A 47 (1988), no. 1, 101–110.
- [7] X.-M. Cheng, A.L. Gavrilyuk, G.R.W. Greaves, and J.H. Koolen, *Biregular graphs with three eigenvalues*, European J. Combin. 56 (2016), 57–80.
- [8] X.-M. Cheng, G.R.W. Greaves, and J.H. Koolen, Graphs with three eigenvalues and second largest eigenvalue at most 1, J. Combin. Theory Ser. B 129 (2018), 31–51.
- [9] H. Chuang and G.R. Omidi, Graphs with three distinct eigenvalues and largest eigenvalues less than 8, Linear Algebra Appl. 430 (2009), no. 8–9, 2053–2062.
- [10] C.J. Colbourn and J.H. (Eds.) Dinitz, Handbook of combinatorial designs (2nd ed.), Chapman and Hall/CRC, 2006.
- [11] D. Crnković, D. Dumičić Danilović, and S. Rukavina, Enumeration of symmetric (45,12,3) designs with nontrivial automorphisms, J. Algebra Combin. Discrete Struct. Appl. 3 (2016), 145–154.
- [12] E.R. van Dam, Nonregular graphs with three eigenvalues, J. Combin. Theory Ser. B 73 (1998), no. 2, 101–118.
- [13] _____, The combinatorics of Dom de Caen, Designs, Codes and Cryptography 34 (2005), 137–148.
- [14] E.R. van Dam, J.H. Koolen, and Z.-J. Xia, Graphs with many valencies and few eigenvalues, Electr. J. Linear Algebra 28 (2015), 12–24.
- [15] R. A. Fisher, An examination of the different possible solutions of a problem in incomplete blocks, Annals of Eugenics 10 (1940), 52–75.
- [16] C. Godsil and K. Meagher, Erdős-Ko-Rado theorems: Algebraic approaches, Cambridge University Press, 2015.
- [17] J.-M. Goethals and J.J. Seidel, Strongly regular graphs derived from combinatorial designs, Canad. J. Math. 22 (1970), 587–614.
- [18] W.H. Haemers and D.G. Higman, Strongly regular graphs with strongly regular decomposition, Linear Algebra and its Applications 114-115 (1989), 379–398, Special Issue Dedicated to Alan J. Hoffman.
- [19] D.G. Higman, Coherent algebras, Linear Algebra Appl. 93 (1987), 209–239.
- [20] _____, Strongly regular designs and coherent configurations of type [323], European J. Combin. 9 (1988), no. 4, 411–422.
- [21] A.N.W. Hone, L.E. Jeffery, and R.G. Selcoe, On a family of sequences related to Chebyshev polynomials, J. Integer Sequences 21 (2018), 18.7.2.
- [22] OEIS Foundation Inc., https://oeis.org/A299107, Entry A299107 in The On-Line Encyclopedia of Integer Sequences (2024).
- [23] M. Muzychuk and M. Klin, On graphs with three eigenvalues, Discrete Math. 189 (1998), no. 1, 191–207.
- [24] A. Neumaier, Regular sets and quasi-symmetric 2-designs, Combinatorial Theory (Dieter Jungnickel and Klaus Vedder, eds.), Springer Berlin Heidelberg, 1982, pp. 258–275.
- [25] S. Reichard, https://github.com/sven-reichard/stabilization, Accessed: 2024-05-21.

- [26] P. Rowlinson, More on graphs with just three distinct eigenvalues, Applicable Analysis and Discrete Math. 11 (2017), no. 1, 74–80.
- [27] T.N. Shorey and C.L. Stewart, On the Diophantine equation $ax^{2t} + bx^ty + cy^2 = d$ and pure powers in recurrence sequences, Mathematica Scandinavica **52** (1983), 24–36.
- [28] M.S. Shrikhande, Quasi-symmetric designs, pp. 578–582, Chapman and Hall/CRC, 2006.
- [29] E. Spence, The strongly regular (40, 12, 2, 4) graphs, Electr. J. Combin. (2000), #R22.
- [30] Douglas R. Stinson, Combinatorial designs: Constructions and analysis, Springer New York, NY, 2004.

A Tables of feasible parameters for QSDs

In the appendix we provide tables of QSD parameters that correspond to the lemmas in Section 5.3. Together with Theorem 5.5, we use following necessary condition on the parameters of a QSD.

Theorem A.1 ([28, Theorem 48.13]). Let \mathcal{Q} be a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$ without repeated blocks. Then $b \leq {v \choose 2}$, with equality if and only if $(\mathcal{P}, \mathcal{B})$ is a 4-design.

Calderbank [5, 6] established additional necessary conditions on QSD parameters, which we refer to as C1 and C2 indicated in Theorem A.2. Let \mathbb{F}_p denote the finite field with pelements and let S_p denote the subset of squares in \mathbb{F}_p .

Theorem A.2. Let Q be a QSD with parameters $(v, k, \lambda; b, r, \{x, y\})$ and intersection numbers x and y.

(C1) Suppose $x \equiv y \pmod{2}$ and $r \not\equiv \lambda \pmod{4}$. Then either

 $x \equiv 0 \pmod{2}, \quad k \equiv 0 \pmod{4}, \text{ and } v \equiv \pm 1 \pmod{8}, \text{ or } x \equiv 1 \pmod{2}, \quad k \equiv v \pmod{4}, \text{ and } v \equiv \pm 1 \pmod{8}.$

(C2) Suppose $x \equiv y \pmod{p}$, where p is an odd prime and $r \not\equiv \lambda \pmod{p^2}$. Then either

 $-v \equiv 0 \pmod{2}, \quad v \equiv k \equiv x \equiv 0 \pmod{p}, \text{ and } (-1)^{v/2} \in S_p \text{ or} \\ -v \equiv 1 \pmod{2}, \quad v \equiv k \equiv x \neq 0 \pmod{p}, \text{ and } x(-1)^{(v-1)/2} \in S_p; \\ -\lambda \equiv r \equiv 0 \pmod{p} \text{ and either}$

$$v \equiv 0 \pmod{2}, \quad and \quad v \equiv k \equiv x \not\equiv 0 \pmod{p};$$

$$v \equiv 0 \pmod{2}, \quad k \equiv x \not\equiv 0 \pmod{p}, \quad and \quad v/x \not\in S_p;$$

$$v \equiv 1 \pmod{2p}, \quad r \equiv 0 \pmod{p^2}, \quad and \quad k \equiv x \not\equiv 0 \pmod{p};$$

$$v \equiv p \pmod{2p}, \quad and \quad k \equiv x \equiv 0 \pmod{p};$$

$$v \equiv 1 \pmod{2p}, \quad k \equiv x \equiv 0 \pmod{p}, \quad and \quad v \notin S_p;$$

$$v \equiv 1 \pmod{2p}, \quad k \equiv x \equiv 0 \pmod{p}, v \in S_p \text{ and } (-1)^{(v-1)/2} \in S_p.$$

In the first column of the tables below, we list parameters of QSD corresponding to Class 1, Class 2, Class 3, and Class 4 graphs from Section 5.3. We only list parameters that satisfy both Theorem 5.5 and Theorem A.1. In the second column, we write the three distinct eigenvalues of the resulting total graph or whole graph. The third column indicates the existence of a QSD having the given parameters. If the nonexistence of a QSD with given parameters is due to a violation of C1 or C2 from Theorem A.2 then this is indicated in the fourth column, together with the relevant prime in the case when C2 is violated. If a corresponding QSD exists, further details are given in the fourth column.

$(v, k, \lambda; b, r, \{x, y\})$	$(\theta_0, \theta_1, \theta_2)$	Exists	Remark
(0, 1, 1, 0, 1, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,	(0,0,0,0,0)	2	rtoman
$(225, 30, 10; 400, 64, \{0, 0\})$	(384, 9, -6)	:	
$(232, 36, 15; 638, 99, \{0, 6\})$	(594, 14, -6)	Ν	C2 $(p = 3)$
$(3501, 225, 40; 9725, 625, \{0, 15\})$	(9375, 39, -15)	?	
$(64576, 1600, 65; 105945, 2625, \{0, 40\})$	(105000, 64, -40)	Ν	C2 $(p = 5)$
$(1075, 100, 66; 7697, 716, \{0, 10\})$	(7160, 65, -10)	Ν	C1
$(9549, 441, 70; 32891, 1519, \{0, 21\})$	(31899, 69, -21)	?	
$(2891, 196, 78; 17051, 1156, \{0, 14\})$	(16184, 77, -14)	Ν	C1
$(43561, 1225, 85; 107569, 3025, \{0, 35\})$	(105875, 84, -35)	?	
$(3550, 225, 96; 23998, 1521, \{0, 15\})$	(22815, 95, -15)	Ν	C2 $(p = 3)$

Table 5: List of QSD parameters corresponding to Lemma 5.10 where $1 \leq \theta_1 \leq 100$.

$(v,k,\lambda;b,r,\{x,y\})$	$(heta_0, heta_1, heta_2)$	Exists	Remark
$(121, 46, 69; 484, 184, \{16, 21\})$	(368, 5, -23)	?	
$(172, 64, 112; 817, 304, \{22, 28\})$	(608, 6, -32)	Ν	C2 $(p = 3)$
$(661, 177, 236; 3305, 885, \{45, 56\})$	(2655, 11, -59)	?	
$(379, 136, 340; 2653, 952, \{46, 55\})$	(1904, 9, -68)	?	
$(361, 145, 522; 3249, 1305, \{55, 64\})$	(2175, 9, -87)	?	
$(3627, 540, 330; 14911, 2220, \{78, 99\})$	(13320, 21, -90)	Ν	C2 $(p = 7)$

Table 6: List of QSD parameters corresponding to Lemma 5.12 where $-100 \leq \theta_2 \leq -2$.

$(v,k,\lambda;b,r,\{x,y\})$	$(heta_0, heta_1, heta_2)$	Exists	Remark
$(8, 6, 15; 28, 21, \{4, 5\})$	(21, 5, -2)	Y	Example 5.16
$(120, 50, 35; 204, 85, \{20, 25\})$	(153, 9, -6)	?	
$(76, 40, 52; 190, 100, \{20, 24\})$	(125, 11, -5)	?	
$(120, 75, 370; 952, 595, \{45, 50\})$	(476, 44, -6)	?	
$(169, 105, 585; 1521, 945, \{63, 69\})$	(735, 59, -7)	?	

Table 7: List of QSD parameters corresponding to Lemma 5.15 where $1 \le \theta_1 \le 100$.

$(v,k,\lambda;b,r,\{x,y\})$	$(heta_0, heta_1, heta_2)$	Exists	Remark
$(141, 45, 33; 329, 105, \{9, 15\})$	(175, 5, -13)	?	
$(85, 40, 130; 595, 280, \{15, 20\})$	(224, 4, -31)	?	
$(232, 112, 296; 1276, 616, \{48, 56\})$	(539, 7, -41)	?	
$(5866, 1666, 777; 9637, 2737, \{441, 476\})$	(6647, 34, -57)	Ν	C2 $(p = 5)$
$(3655, 1450, 1380; 8772, 3480, \{550, 580\})$	(5046, 29, -71)	Ν	C2 $(p = 3)$

Table 8: List of QSD parameters corresponding to Lemma 5.19 where $-100 \leq \theta_2 \leq -2$.