

HIGHER GAUSSIAN MAPS ON THE HYPERELLIPTIC LOCUS AND SECOND FUNDAMENTAL FORM

D. FARO, P. FREDIANI, AND A. LACOPO

ABSTRACT. In this paper we study higher even Gaussian maps of the canonical bundle on hyperelliptic curves and we determine their rank, giving explicit descriptions of their kernels. Then we use this descriptions to investigate the hyperelliptic Torelli map j_h and its second fundamental form. We study isotropic subspaces of the tangent space $T_{\mathcal{H}_g, [C]}$ to the moduli space \mathcal{H}_g of hyperelliptic curves of genus g at a point $[C]$, with respect to the second fundamental form ρ_{HE} of j_h . In particular, for any Weierstrass point $p \in C$, we construct a subspace V_p of dimension $\lfloor \frac{g}{2} \rfloor$ of $T_{\mathcal{H}_g, [C]}$ generated by higher Schiffer variations at p , such that the only isotropic tangent direction $\zeta \in V_p$ for the image of ρ_{HE} is the standard Schiffer variation ξ_p at the Weierstrass point $p \in C$.

1. INTRODUCTION

The purpose of this paper is twofold. On the one hand we study higher even Gaussian maps of the canonical bundle on hyperelliptic curves and we determine their rank, giving an explicit description of their kernels. On the other hand, we use this description to investigate the local geometry of the image via the Torelli map of the moduli space \mathcal{H}_g of hyperelliptic curves of genus g in the moduli space \mathcal{A}_g of principally polarised abelian varieties of dimension g .

Let C be a hyperelliptic curve of genus $g \geq 3$. Denote by I_2 the kernel of the multiplication map

$$\mu_0 : S^2 H^0(C, K_C) \rightarrow H^0(C, K_C^{\otimes 2})^+,$$

where $H^0(C, K_C^{\otimes 2})^+$ denotes the invariant subspace of $H^0(C, K_C^{\otimes 2})$ under the action of the hyperelliptic involution. The canonical map is the composition of the $2 : 1$ map $\pi : C \rightarrow \mathbb{P}^1$ with the $(g-1)$ -th Veronese embedding $\nu_{g-1} : \mathbb{P}^1 \rightarrow \mathbb{P}^{g-1}$, hence the space I_2 is identified with the vector space of quadrics containing the rational normal curve $\nu_{g-1}(\mathbb{P}^1)$ in \mathbb{P}^{g-1} . The second Gaussian map μ_2 is a linear map

$$\mu_2 : I_2 = \text{Ker}(\mu_0) \rightarrow H^0(C, K_C^{\otimes 4}),$$

and more generally, for any $k \geq 1$, the $2k$ -th Gaussian map is a linear map

$$\mu_{2k} : \text{Ker}(\mu_{2k-2}) \rightarrow H^0(C, K_C^{\otimes 2k+2}).$$

These maps were introduced by Wahl in [18]. One can also define odd Gaussian maps, each one being defined on the kernel of the previous one. The most important and studied

D. Faro, P. Frediani and A. Lacopo are members of GNSAGA (INdAM) and are partially supported by PRIN project *Moduli spaces and special varieties* (2022).

one is the first Wahl map:

$$\mu_1 : \wedge^2 H^0(C, K_C) \rightarrow H^0(C, K_C^{\otimes 3}).$$

It was proven by Wahl that if C sits on a $K3$ surface, than the first Wahl map μ_1 is not surjective (see [18], see also [1]). On the other hand, Ciliberto, Harris and Miranda proved that for the general curve of genus $g \geq 10$, $g \neq 11$, μ_1 is surjective (see [2], [16] for another proof).

If C is a hyperelliptic curve, it was proven in [17], [3] that the rank of μ_1 is $2g - 3$, while in [4] it was shown that the rank of μ_2 is $2g - 5$.

Our first main result is the following

Theorem 1.1. (See Theorem 3.1) *Let C be a hyperelliptic curve of genus $g \geq 3$. Then for every $0 \leq k \leq \frac{g-1}{2}$*

$$(1) \quad \text{Rank}(\mu_{2k}) = 2g - (4k + 1),$$

$$(2) \quad \dim(\text{Ker}(\mu_{2k})) = \frac{(g-1)(g-2)}{2} - k(2g - 2k - 3).$$

Then, $\text{Rank}(\mu_{2k}) = 0$ for every $k > \lfloor \frac{g-1}{2} \rfloor$.

From this we immediately see (Remark 3.2) that if g is odd we have the following chain of inclusions:

$$0 = \text{Ker}(\mu_{g-1}) \subsetneq \text{Ker}(\mu_{g-3}) \subsetneq \dots \subsetneq \text{Ker}(\mu_2) \subsetneq I_2;$$

while if g is even we have

$$0 = \text{Ker}(\mu_{g-2}) \subsetneq \text{Ker}(\mu_{g-4}) \subsetneq \dots \subsetneq \text{Ker}(\mu_2) \subsetneq I_2.$$

Denote by \mathcal{H}_g the hyperelliptic locus in \mathcal{M}_g and by

$$j_h : \mathcal{H}_g \rightarrow \mathcal{A}_g$$

the restriction of the Torelli map to \mathcal{H}_g .

The map j_h is an orbifold immersion (see [15]). The main tool for studying the local geometry of the hyperelliptic Torelli locus $j_h(\mathcal{H}_g) \subset \mathcal{A}_g$ is the second fundamental form of the map j_h . Recall that the moduli space \mathcal{A}_g is the quotient of the Siegel space $Sp(2g, \mathbb{R})/U(g)$ under the action of $Sp(2g, \mathbb{Z})$. We endow \mathcal{A}_g with the Siegel metric, that is the orbifold metric induced by the symmetric metric on the Siegel space. The second fundamental form of j_h is a map

$$\rho_{HE} : N_{\mathcal{H}_g|\mathcal{A}_g}^* \rightarrow \text{Sym}^2 \Omega_{\mathcal{H}_g}^1,$$

where $N_{\mathcal{H}_g|\mathcal{A}_g}^*$ denotes the conormal bundle of $j_h(\mathcal{H}_g)$ in \mathcal{A}_g . At a point $[C] \in \mathcal{H}_g$, ρ_{HE} is therefore a linear map

$$\rho_{HE} : I_2 \rightarrow \text{Sym}^2(H^0(C, K_C^{\otimes 2})^+).$$

One of the main problems in the study of the local geometry of the image of the hyperelliptic locus in \mathcal{A}_g under the Torelli map is to investigate the existence of totally geodesic subvarieties of \mathcal{A}_g contained in the hyperelliptic locus.

This is related with the hyperelliptic Coleman-Oort conjecture, that says that for $g \geq 8$ there do not exist positive dimensional special subvarieties of \mathcal{A}_g generically contained in the hyperelliptic Torelli locus, namely contained in the closure of $j_h(\mathcal{H}_g)$ and intersecting $j_h(\mathcal{H}_g)$. This conjecture was recently proven by Moonen [14], generalising a previous result of Lu and Zuo [13].

In fact, special subvarieties are totally geodesic, hence the study of the second fundamental form is related with the above problem.

Here we take a different viewpoint, namely, we are interested in the behaviour of the second fundamental form of the map j_h in relation with higher even Gaussian maps.

In [5] it was proven that the map ρ_{HE} can be expressed in terms of the Hodge Gaussian map ρ introduced in [8]. An important property proven in [8] is a formula computing, for any $Q \in I_2$, the Hodge Gaussian map $\rho(Q)$ on Schiffer variations at points p on any curve C in terms of the second Gaussian map $\mu_2(Q)$.

This property is one main tool used in [11, Theorem 6.2] to show that the maximal dimension of a germ of a totally geodesic subvariety of \mathcal{A}_g contained in the hyperelliptic locus is $g + 1$.

In the case of hyperelliptic curves, the result of [8] gives the following expression for $\rho_{HE}(Q)$ on the Schiffer variation $\xi_p \in H^1(C, T_C)^+$ at a Weierstrass point $p \in C$, in terms of the second Gaussian map :

$$\rho_{HE}(Q)(\xi_p \odot \xi_p) = -2\pi i \mu_2(Q)(p).$$

Then we have $\rho_{HE}(Q)(\xi_p \odot \xi_p) = 0$, since p is a Weierstrass point and one can easily see that for all the quadrics $Q \in I_2$, we have $\mu_2(Q)(p) = 0$ (see Remark 6.13).

In [9], a relation between higher even Gaussian maps μ_{2k} of the canonical bundle on a smooth projective curve of genus $g \geq 4$ and the Hodge Gaussian map ρ is given, generalising the above result.

Here we exploit similar techniques to compute $\rho_{HE}(Q)$ for quadrics Q contained in the kernel of higher Gaussian maps on odd higher Schiffer variations $\xi_p^{2k+1} \in H^1(C, T_C)^+$ at a Weierstrass point $p \in C$ (see Section 4 for the definition of higher Schiffer variations). For any $k \leq 2g - 3$, consider the $(k + 1)$ -dimensional subspace

$$V_k := \langle \xi_p^1, \xi_p^3, \dots, \xi_p^{2k+1} \rangle \subset H^1(T_C)^+.$$

We prove the following

Theorem 1.2. (See Theorems 6.6, 6.7).

(1) Let $0 \leq k \leq \lfloor \frac{g-3}{2} \rfloor$ and let $Q \in \text{Ker}(\mu_{2k})$. Then we have

$$\rho_{HE}(Q)(\xi_p^l \odot \xi_p^m) = 0 \quad \forall l + m \leq 4k + 3, \quad l, m \text{ odd.}$$

This implies that the subspace $V_k = \langle \xi_p^1, \xi_p^3, \dots, \xi_p^{2k+1} \rangle \subset H^1(T_C)^+$ is isotropic for $\rho_{HE}(Q)$, $\forall Q \in \text{Ker}(\mu_{2k})$.

(2) There exists $Q \in \text{Ker}(\mu_{2k})$ such that $\rho(Q)(\xi_p^{2k+1} \odot \xi_p^{2k+3}) \neq 0$. Hence the subspace $V_{k+1} = \langle \xi_p^1, \dots, \xi_p^{2k+1}, \xi_p^{2k+3} \rangle \subset H^1(T_C)^+$ is not isotropic for $\rho_{HE}(\text{Ker}(\mu_{2k}))$.

Following [7] we call a nonzero direction $\zeta \in H^1(T_C)^+$ asymptotic if

$$\rho_{HE}(Q)(\zeta \odot \zeta) = 0, \quad \forall Q \in I_2.$$

Clearly tangent directions to totally geodesic subvarieties are asymptotic directions. Notice that saying that a nonzero element $\zeta \in H^1(T_C)^+$ is an asymptotic direction means that the point $[\zeta] \in \mathbb{P}H^1(T_C)^+$ is in the base locus of the linear space of quadrics $\rho_{HE}(I_2) \subset \text{Sym}^2(H^1(T_C)^+)^{\vee}$. In [5] it is shown that ρ_{HE} is injective, hence the image of ρ_{HE} in $\text{Sym}^2(H^1(T_C)^+)^{\vee}$ has dimension $\frac{(g-1)(g-2)}{2}$. So $\rho_{HE}(I_2)$ is a linear system of quadrics in $\mathbb{P}(H^1(C, T_C)^+) \cong \mathbb{P}^{2g-2}$ of dimension $\frac{(g-1)(g-2)}{2}$.

Thus if $g \leq 5$, there are asymptotic directions, in fact there are also examples of special subvarieties of \mathcal{A}_g of positive dimension generically contained in the hyperelliptic locus (see e.g. [10]).

On the other hand, for g sufficiently high one would expect that the intersection of a space of quadrics of dimension $\frac{(g-1)(g-2)}{2}$ in \mathbb{P}^{2g-2} would be empty, hence that asymptotic directions would not exist. This is not true because Schiffer variations at Weierstrass points are asymptotic directions (see Remark 6.13). On the other hand we prove the following

Theorem 1.3. (See Theorem 6.14).

Let C be a hyperelliptic curve of genus $g \geq 3$ and $p \in C$ be a Weierstrass point. Then the asymptotic directions in the space $V_{\lfloor \frac{g-2}{2} \rfloor}$ are exactly the Schiffer variations $\xi_p = \xi_p^1$.

As a consequence we have the following

Remark 1.4. (see Remark 6.15).

In [5, Corollary 3.4] it is shown that ρ_{HE} is injective, so

$$\dim \rho_{HE}(\text{Ker} \mu_{2k}) = \dim \text{Ker} \mu_{2k} = \frac{(g-1)(g-2)}{2} - k(2g-2k-3),$$

for all k . For any hyperelliptic curve C of genus $g \geq 3$ and for any Weierstrass point $p \in C$, we have proven that we have the following chain of subspaces of quadrics in $\mathbb{P}H^1(T_C)^+$ and of corresponding maximal isotropic subspaces in $V_{\frac{g-3}{2}}$ if g is odd and in $V_{\frac{g-2}{2}}$ if g is even.

If g is odd:

$$\rho_{HE}(\text{Ker}(\mu_{g-3})) \subsetneq \rho_{HE}(\text{Ker}(\mu_{g-5})) \dots \subsetneq \rho_{HE}(\text{Ker}(\mu_2)) \subsetneq \rho_{HE}(I_2);$$

$$V_{\frac{g-3}{2}} \supsetneq V_{\frac{g-5}{2}} \supsetneq \dots \supsetneq V_1 \supsetneq V_0 = \langle \xi_p \rangle,$$

while if g is even we have

$$\rho_{HE}(Ker(\mu_{g-4})) \subsetneq \rho_{HE}(Ker(\mu_{g-6})) \dots \subsetneq \rho_{HE}(Ker(\mu_2)) \subsetneq \rho_{HE}(I_2),$$

$$V_{\frac{g-4}{2}} \supsetneq V_{\frac{g-6}{2}} \supsetneq \dots \supsetneq V_1 \supsetneq V_0 = \langle \xi_p \rangle.$$

Finally, we show that Theorem 6.14 also allows us to deduce Corollary 6.16, which gives a bound for the dimension of a germ of a totally geodesic submanifold of \mathcal{A}_g , generically contained in the Hyperelliptic Torelli locus. Nevertheless this is weaker than the one already proven in [11, Theorem 6.2].

The structure of the paper is as follows.

In Section 2 we recall the definition and basic properties of Gaussian maps. In Section 3 we determine the kernel of all even higher Gaussian maps on the hyperelliptic locus and we compute their rank, proving Theorem 1.1. In Section 4 we define and give the basic properties of higher Schiffer variations. In Section 5 we recall the definition and some results on the second fundamental form of the hyperelliptic Torelli map j_h . Finally in Section 6 we prove Theorems 1.2, 1.3 and their consequences (Remark 6.15 and Corollary 6.16).

2. GAUSSIAN MAPS

Let C be a smooth curve of genus g , and let L and M be two line bundles on C . In this section we recall the definition of the (higher) Gaussian maps on C associated with the line bundles L and M .

Set $S := C \times C$ and let $\Delta \subset S$ be the diagonal. Take a non-negative integer k , then the k -th Gaussian (or Wahl map) associated with L and M is the map given by restriction to the diagonal

$$H^0(S, L \boxtimes M(-k\Delta)) \xrightarrow{\Phi_{k,L,M}} H^0(S, L \boxtimes M(-k\Delta)|_{\Delta}) \cong H^0(C, L \otimes M \otimes K_C^{\otimes k}).$$

We will only consider the case $L = M$, and we set $\Phi_{k,L} := \Phi_{k,L,L}$. Since we have the identification $H^0(S, L \boxtimes M) \cong H^0(C, L) \otimes H^0(C, M)$, the map $\Phi_{0,L}$ is given by the multiplication map of global sections

$$H^0(C, L) \otimes H^0(C, L) \rightarrow H^0(C, L^{\otimes 2}),$$

which vanishes identically on $\wedge^2 H^0(C, L)$. Then we have

$$\ker \Phi_{0,L} = H^0(S, L \boxtimes L(-\Delta)) \cong \wedge^2 H^0(C, L) \oplus I_2(L),$$

where $I_2(L)$ is the kernel of the multiplication map $S^2 H^0(C, L) \rightarrow H^0(C, L^{\otimes 2})$. Since $\Phi_{1,L}$ vanishes on symmetric tensors, one writes

$$\mu_{1,L} := \Phi_{1,L}|_{\wedge^2 H^0(C, L)} : \wedge^2 H^0(C, L) \rightarrow H^0(C, L^{\otimes 2}).$$

Take a local frame l for L , local coordinate z and two sections $s_1, s_2 \in H^0(C, L)$ with $s_i = f_i(z)l$. Then we have

$$(3) \quad \mu_{1,L}(s_1 \wedge s_2) = (f_1'f_2 - f_2'f_1)dz \otimes l^2.$$

Consequently, the zero divisor of $\mu_{1,L}(s_1 \wedge s_2)$ is $2B + R$, where B is the base locus of the pencil $\langle s_1, s_2 \rangle$ and R is the ramification divisor of the associated morphism.

Again $H^0(S, L \boxtimes L(-2\Delta))$ decomposes as the sum of $I_2(L)$ and the kernel of $\mu_{1,L}$. Since $\Phi_{2,L}$ vanishes on skew-symmetric tensors, we write

$$\mu_{2,L} := \Phi_{2,L}|_{I_2(L)} : I_2(L) \rightarrow H^0(C, L^{\otimes 2} \otimes K_C^{\otimes 2}).$$

We will denote by μ_2 the second gaussian map of the canonical line bundle K_C on C :

$$\mu_2 := \mu_{2,K_C} : I_2(K_C) \rightarrow H^0(K_C^{\otimes 4}).$$

If $\alpha = \sum l_i \otimes m_i \in I_2(L)$ with $l_i = f_i(z)l$ and $m_i = g_i(z)l$, using that $\mu_{0,L}(\alpha) = 0$, and $\Phi_{1,L}(\alpha) = 0$, one finds the identities

$$(4) \quad \sum f_i g_i \equiv 0, \quad \sum f_i' g_i - f_i g_i' \equiv 0.$$

Taking derivatives of the first equation in (4) and using the second one we obtain the identities:

$$(5) \quad \sum f_i' g_i \equiv \sum f_i g_i' \equiv 0,$$

Derivating the equations in (5) we have:

$$(6) \quad \sum f_i'' g_i = - \sum f_i' g_i' = \sum f_i g_i''.$$

The local expression of $\mu_{2,L}$ is

$$(7) \quad \mu_{2,L}(\alpha) = \sum f_i'' g_i dz^{\otimes 2} \otimes l^{\otimes 2} = \sum f_i g_i'' dz^{\otimes 2} \otimes l^{\otimes 2} = - \sum f_i' g_i' dz^{\otimes 2} \otimes l^{\otimes 2}.$$

More generally the local expression for $\Phi_{2k,L}$ when $k \geq 1$ is given by

$$(8) \quad \Phi_{2k,L}(\alpha) = (-1)^k \sum f_i^{(k)} g_i^{(k)} dz^{\otimes 2k} \otimes l^{\otimes 2},$$

and since it vanishes on antisymmetric tensors one defines

$$(9) \quad \mu_{2k,L} := \Phi_{2k,L}|_{Ker(\mu_{2k-2,L})}.$$

Analogously the local expression for $\Phi_{2k+1,L}$ when $k \geq 0$ is given by

$$(10) \quad \Phi_{2k+1,L}(\alpha) = \sum (f_i^{(k+1)} g_i^{(k)} - f_i^{(k)} g_i^{(k+1)}) dz^{\otimes 2k+1} \otimes l^{\otimes 2},$$

and since it vanishes on symmetric tensors one defines for every $k \geq 2$

$$(11) \quad \mu_{2k+1,L} := \Phi_{2k+1,L}|_{Ker(\mu_{2k-1,L})}.$$

Moreover since $\alpha = \sum l_i \otimes m_i \in Ker(\mu_{2k-2,L})$ if and only if

$$(12) \quad \mu_{0,L}(\alpha) = \Phi_{0,L}(\alpha) = \dots = \mu_{2k-2,L}(\alpha) = \Phi_{2k-1,L}(\alpha) = 0,$$

and these conditions give identities

$$(13) \quad \sum f_i^{(h)} g_i^{(r)} \equiv 0, \forall h, r \text{ such that } h + r \leq 2k - 1,$$

the local expression for $\mu_{2k,L}$ is equivalent to

$$(14) \quad \mu_{2k,L}(\alpha) = (-1)^m \sum f_i^{(2k-m)} g_i^{(m)} dz^{\otimes 2k} \otimes l^{\otimes 2},$$

for every $m = 0, \dots, 2k$. Analogously $\alpha = \sum l_i \otimes m_i \in \text{Ker}(\mu_{2k-1,L})$ if and only if

$$(15) \quad \Phi_{0,L}(\alpha) = \mu_{1,L}(\alpha) = \dots = \mu_{2k-1,L}(\alpha) = \Phi_{2k,L}(\alpha) = 0,$$

and these conditions give the identities

$$(16) \quad \sum f_i^{(h)} g_i^{(r)} \equiv 0, \forall h, r \text{ such that } h + r \leq 2k,$$

and from this, it follows that we can equivalently express (10) as

$$(17) \quad \mu_{2k+1,L}(\alpha) = (-1)^m \sum (f_i^{(2k+1-m)} g_i^{(m)} - f_i^{(m)} g_i^{(2k+1-m)}) dz^{\otimes 2k+1} \otimes l^{\otimes 2}$$

for every $m = 0, \dots, k$.

Remark 2.1. *Observe that for every $k \geq 1$*

$$(18) \quad \text{Ker}(\Phi_{2k,L}) = \text{Ker}(\mu_{2k,L}) \oplus \text{Ker}(\mu_{2k-1,L}),$$

and from the definition of $\mu_{k,L}$ we have the inclusions:

$$(19) \quad \dots \subset \text{Ker}(\mu_{2k,L}) \subset \text{Ker}(\mu_{2(k-1),L}) \subset \dots \subset \text{Ker}(\mu_{2,L}) \subset I_2 = \text{Ker}(\mu_{0,L});$$

$$(20) \quad \dots \subset \text{Ker}(\mu_{2k+1,L}) \subset \text{Ker}(\mu_{2k-1,L}) \subset \dots \subset \text{Ker}(\mu_{1,L}).$$

3. RANK OF HIGHER ORDER GAUSSIAN MAPS ON HYPERELLIPTIC CURVES

Let C be a hyperelliptic curve of genus $g \geq 3$, L be the line bundle giving the g_2^1 . Set $M = K_C \otimes L^\vee$ and $\pi : C \rightarrow \mathbb{P}^1$ the map induced by $|L|$. Denote by $\nu_n : \mathbb{P}^1 \hookrightarrow \mathbb{P}^n$ the n^{th} Veronese embedding. Recall that the canonical map is given by the composition $\nu_{g-1} \circ \pi$, so

$$K_C \cong \pi^*(\mathcal{O}_{\mathbb{P}^1}(g-1)) \cong L^{\otimes(g-1)}$$

and $M \cong L^{\otimes(g-2)}$. Then $H^0(C, M) \cong H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(g-2))$ has dimension $g-1$.

Denote by Φ_k, μ_k , the Gaussian maps Φ_{k,K_C} and μ_{k,K_C} . The main result of this section is the computation of the rank of higher Gaussian maps for any hyperelliptic curve of any genus. More precisely we are going to prove the following:

Theorem 3.1. *Let C be a hyperelliptic curve of genus $g \geq 3$. Then for every $0 \leq k \leq \frac{g-1}{2}$*

$$(21) \quad \text{Rank}(\mu_{2k}) = 2g - (4k + 1),$$

$$(22) \quad \dim(\text{Ker}(\mu_{2k})) = \frac{(g-1)(g-2)}{2} - k(2g - 2k - 3).$$

Then, $\text{Rank}(\mu_{2k}) = 0$ for every $k > \lfloor \frac{g-1}{2} \rfloor$.

Notice that the result was already known for μ_2 (see [4]).

Remark 3.2. *By Theorem 3.1, it is immediate to see that if g is odd we have the following chain of inclusions:*

$$0 = \text{Ker}(\mu_{g-1}) \subsetneq \text{Ker}(\mu_{g-3}) \subsetneq \dots \subsetneq \text{Ker}(\mu_2) \subsetneq I_2(K_C);$$

while if g is even we have

$$0 = \text{Ker}(\mu_{g-2}) \subsetneq \text{Ker}(\mu_{g-4}) \subsetneq \dots \subsetneq \text{Ker}(\mu_2) \subsetneq I_2(K_C).$$

Notice that if g is odd, $\dim(\text{Ker}(\mu_{g-3})) = 1$, and if g is even $\dim(\text{Ker}(\mu_{g-4})) = 3$.

Let us first recall the following well known result (see for example [4]).

Lemma 3.3. *Let C be a hyperelliptic curve of genus $g \geq 3$. Let $|L|$ be the g_2^1 . Set $M = K_C \otimes L^\vee$, let $\omega_1, \dots, \omega_{g-1}$ be a basis for $H^0(M)$ and let $\langle s, t \rangle$ be a basis for $H^0(L)$. Then the map defined by*

$$\begin{aligned} \Lambda^2 H^0(M) &\xrightarrow{\psi} I_2 \\ \omega_i \wedge \omega_j &\rightarrow Q_{ij} := s\omega_i \odot t\omega_j - s\omega_j \odot t\omega_i, \end{aligned}$$

is an isomorphism. In particular, observe that $\{Q_{ij}\}$ gives a basis for I_2 .

In [4, Lemma 4.1] it is shown the following

Lemma 3.4. *For any Q_{ij} as in the previous lemma:*

$$\mu_2(Q_{ij}) = \mu_{1,L}(s \wedge t) \mu_{1,M}(\psi^{-1}(Q_{ij})),$$

where with the expression

$$(23) \quad \mu_{1,L}(s \wedge t) \mu_{1,M}(\psi^{-1}(Q_{ij}))$$

we mean the image of $\mu_{1,L}(s \wedge t) \otimes \mu_{1,M}(\psi^{-1}(Q_{ij}))$ under the multiplication map

$$(24) \quad H^0(K_C \otimes L^{\otimes 2}) \otimes H^0(K_C \otimes M^{\otimes 2}) \rightarrow H^0(K_C^{\otimes 4}).$$

Observe that from Lemma 3.4 it follows that $\text{Rank}(\mu_2) = \text{Rank}(\mu_{1,M})$ since $\mu_{1,L}(s \wedge t)$ is a nonzero section in $H^0(K_C \otimes L^{\otimes 2})$. Indeed the zero locus of $\mu_{1,L}(s \wedge t)$ is given by the base locus of $|L|$ together with the ramification divisor of the induced morphism (recall (3)).

We start generalizing Lemma 3.4 to any higher-order Gaussian maps. We use the same notations as in Lemma 3.3

Lemma 3.5. *Let $k \geq 0$ be an integer and let*

$$Q = \sum_{1 \leq i < j \leq g-1} a_{ij} Q_{ij} \in \text{Ker}(\mu_{2k}).$$

Then

(i) for any $k \geq 1$

$$\sum_{1 \leq i < j \leq g-1} a_{ij}(\omega_i \wedge \omega_j) \in \text{Ker}(\mu_{2k-1, M}),$$

(ii) for any $k \geq 0$

$$\mu_{2k+2}(Q) = (k+1)\mu_{1, L}(s \wedge t)\mu_{2k+1, M}\left(\sum_{1 \leq i < j \leq g-1} a_{ij}(\omega_i \wedge \omega_j)\right) \in H^0(K_C^{\otimes 2k+4}),$$

Proof. Let us proceed by induction. When $k = 0$ (ii) is Lemma 3.4, and when $k = 1$ (i) it is an immediate consequence of the hypothesis $\mu_2(Q) = 0$ together with Lemma 3.4.

Now take $n \geq 2$ and suppose that (ii) holds for every $0 \leq k < n - 1$ and (i) holds for every $1 \leq k < n$. We are going to prove that (ii) holds for $k = n - 1$, which automatically implies that (i) holds for $k = n$. Set $k = n - 1$ and suppose that

$$Q = \sum_{1 \leq i < j \leq g-1} a_{ij}Q_{ij} \in \text{Ker}(\mu_{2k}).$$

On some open sets write

$$\omega_i = f_i dz', \quad t = fT \quad \text{and} \quad s = gT,$$

where f_i is a holomorphic function, dz' is a local generator on $M = K_C \otimes L^\vee$ and T is a local generator on L . From the definition of Q_{ij} in Lemma 3.3, it follows that we can write locally Q as

$$(25) \quad \sum_{1 \leq i < j \leq g-1} a_{ij}Q_{ij} = \sum a_{ij}(ff_i dz \odot gf_j dz - ff_j dz \odot gf_i dz),$$

where we can take z to be a local coordinate on C . Then, by definition

$$\begin{aligned} & \mu_{2k+2}\left(\sum_{1 \leq i < j \leq g-1} a_{ij}Q_{ij}\right) \\ &= \sum_{1 \leq i < j \leq g-1} a_{ij}((ff_i)^{(k+1)}(gf_j)^{(k+1)} - (ff_j)^{(k+1)}(gf_i)^{(k+1)})dz^{\otimes(2k+4)}. \end{aligned}$$

Observe that the latter expression can be written as

$$(26) \quad \begin{aligned} &= \left[\sum_{1 \leq i < j \leq g-1} a_{ij} \sum_{h=0}^{k+1} \binom{k+1}{h} f^{(k+1-h)} f_i^{(h)} \sum_{l=0}^{k+1} \binom{k+1}{l} g^{(k+1-l)} f_j^{(l)} \right] dz^{\otimes(2k+4)} \\ &- \left[\sum_{1 \leq i < j \leq g-1} a_{ij} \sum_{r=0}^{k+1} \binom{k+1}{r} f^{(k+1-r)} f_j^{(r)} \sum_{e=0}^{k+1} \binom{k+1}{e} g^{(k+1-e)} f_i^{(e)} \right] dz^{\otimes(2k+4)} \end{aligned}$$

$$(27) \quad = \left[\sum_{h, l=0}^{k+1} \binom{k+1}{h} \binom{k+1}{l} f^{(k+1-h)} g^{(k+1-l)} \sum_{1 \leq i < j \leq l-1} a_{ij} (f_i^{(h)} f_j^{(l)} - f_j^{(h)} f_i^{(l)}) \right] dz^{\otimes(2k+4)}$$

By hypothesis $Q \in \text{Ker}(\mu_{2k})$ and hence from the inductive hypothesis

$$\mu_{2k-1,M} \left(\sum_{1 \leq i < j \leq g-1} a_{ij}(\omega_i \wedge \omega_j) \right) = 0.$$

Then using 16 it follows that for every $h + l \leq 2k$

$$\sum_{1 \leq i < j \leq g-1} a_{ij}(f_i^{(h)} f_j^{(l)} - f_j^{(h)} f_i^{(l)}) \equiv 0.$$

Then (27) becomes

$$\begin{aligned} & \left[\binom{k+1}{k} f g^{(1)} \left(\sum_{1 \leq i < j \leq l-1} a_{ij}(f_i^{(k+1)} f_j^{(k)} - f_j^{(k+1)} f_i^{(k)}) \right) \right] dz^{\otimes(2k+4)} + \\ & \left[\binom{k+1}{k} f^{(1)} g \left(\sum_{1 \leq i < j \leq l-1} a_{ij}(f_i^{(k)} f_j^{(k+1)} - f_j^{(k)} f_i^{(k+1)}) \right) \right] dz^{\otimes(2k+4)}. \end{aligned}$$

This is just

$$\left[\binom{k+1}{k} (f g^{(1)} - f^{(1)} g) \left(\sum_{1 \leq i < j \leq l-1} a_{ij}(f_i^{(k+1)} f_j^{(k)} - f_j^{(k+1)} f_i^{(k)}) \right) \right] dz^{\otimes(2k+4)},$$

which is equal to

$$(k+1)\mu_{1,L}(t \wedge s)\mu_{2k+1,M} \left(\sum_{1 \leq i < j \leq l-1} a_{ij}(\omega_i \wedge \omega_j) \right),$$

and so (ii) holds. \square

In the following lemma, we are going to describe the equations of the loci

$$\dots \subset \text{Ker}(\mu_{2k}) \subset \text{Ker}(\mu_{2(k-1)}) \subset \dots \subset \text{Ker}(\mu_2) \subset I_2 = \text{Ker}(\mu_0).$$

The equation of $\text{Ker}(\mu_2)$ in I_2 have already been described in the proof of [4, Proposition 4.2]. Indeed we have the following

Lemma 3.6. *Let*

$$Q = \sum_{1 \leq i < j \leq g-1} a_{ij} Q_{ij} \in I_2,$$

Q_{ij} as in 3.3. Then $Q \in \text{Ker}(\mu_2)$ if and only if for all $3 \leq l \leq 2g-3$

$$(28) \quad \sum_{\substack{1 \leq i < j \\ i+j=l}} a_{ij}(j-i) = 0.$$

Now we generalize the approach in [4]. The strategy is to use Lemma 3.5 together the well-known description of a basis of the $H^0(C, K_C)$ for a hyperelliptic curve C . More precisely since we want to describe $\text{Ker}(\mu_{2k})$, we can argue locally and suppose that on some open set with coordinate x , a basis for $H^0(C, K_C)$ is given by

$$(29) \quad \left\{ \frac{dx}{y}, x \frac{dx}{y}, \dots, x^{g-1} \frac{dx}{y} \right\}.$$

and the $2 : 1$ morphism to \mathbb{P}^1 given by $|L|$ is given by $(x, y) \mapsto x$. Hence a local description of a basis for $H^0(C, M)$ is given by

$$(30) \quad \left\{ x \frac{dx}{y}, \dots, x^{g-1} \frac{dx}{y} \right\}.$$

Lemma 3.7. *Let*

$$Q = \sum_{1 \leq i < j \leq g-1} a_{ij} Q_{ij} \in I_2,$$

with Q_{ij} as in 3.3. For any $k \geq 2$, $Q \in \text{Ker}(\mu_{2k})$ if and only if

- $\forall 3 \leq l \leq 2g - 3,$

$$\sum_{\substack{1 \leq i < j \\ i+j=l}} a_{ij} (j - i) = 0,$$

- $\forall 2 \leq m \leq k, \forall 2m - 1 \leq l \leq 2g - 3,$

$$(31) \quad \sum_{\substack{1 \leq i < j \\ i+j=l \\ i \geq m-1, \\ j \geq m-1}} a_{ij} (j - i) i j (i - 1) (j - 1) \dots (i - (m - 2)) (j - (m - 2)) = 0.$$

Proof. We will argue locally. Take the basis of $H^0(C, K_C)$ given in (29) and the one for $H^0(C, M)$ given in (30). We will proceed by induction on k . The thesis holds for $k = 1$ by Lemma 3.6. Now we suppose that the thesis holds for $k - 1$, $k \geq 2$ and take $Q \in \text{Ker}(\mu_{2k-2})$. Since Q also belongs to the previous kernels, proving the lemma is equivalent to prove that $Q \in \text{Ker}(\mu_{2k})$ if and only if

$$(32) \quad \sum_{\substack{1 \leq i < j \\ i+j=l \\ i \geq k-1, \\ j \geq k-1}} a_{ij} (j - i) i j (i - 1) (j - 1) \dots (i - (k - 2)) (j - (k - 2)) = 0,$$

for all $2k - 1 \leq l \leq 2g - 3$. Using lemma 3.5 we have that $\mu_{2k}(Q) = 0$ if and only if $\mu_{2k-1, L}(\sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i \wedge \omega_j) = 0)$, that is if and only if

$$(33) \quad \sum_{1 \leq i < j \leq g-1} a_{ij} ((x^i)^{(k-1)} (x^j)^{(k)} - (x^i)^{(k)} (x^j)^{(k-1)}) \equiv 0,$$

where we are using that dx/y is a local frame. Observe that (33) is equal to

$$(34) \quad \sum_{\substack{1 \leq i < j \leq g-1 \\ i \geq k-1 \\ j \geq k}} a_{ij} i \dots (i - (k - 2)) x^{i-(k-1)} j \dots (j - (k - 1)) x^{j-k}$$

$$(35) \quad - \sum_{\substack{1 \leq i < j \leq g-1 \\ i \geq k \\ j \geq k-1}} a_{ij} i \dots (i - (k - 1)) x^{i-k} j \dots (j - (k - 2)) x^{j-(k-1)},$$

which can be written after some simple algebraic manipulations as

$$(36) \quad \sum_{\substack{1 \leq i < j \leq g-1 \\ i \geq k-1 \\ j \geq k-1}} x^{(i+j-(2k-1))} a_{ij} i \dots (i - (k-2)) j \dots (j - (k-2)) (j - i).$$

So we have shown that for $Q \in \text{Ker} \mu_{2k-2}$, $\mu_{2k}(Q) = 0$ if and only if for every $2k-1 \leq l \leq 2g-3$

$$\sum_{\substack{1 \leq i < j \leq g-1 \\ i+j=l \\ i \geq k-1 \\ j \geq k-1}} a_{ij} (j-i) i \dots (i - (k-2)) j \dots (j - (k-2)) = 0.$$

This concludes the proof. \square

Now we come to the proof Theorem 3.1. The strategy of the proof is clear: we have to determine the number of linearly independent equations in (31).

Proof of Theorem 3.1. We will start introducing some notations. For every i, j such that $1 \leq i < j \leq g-1$, for every $l = 3, \dots, 2g-3$, let $(w_l)^2$ the vector whose coordinates are

$$(w_l)^2_{ij} = \begin{cases} (j-i) & \text{if } i+j=l \\ 0 & \text{if } i+j \neq l \end{cases}$$

which are the coefficients of the l -th equation of the set of equations given in Lemma 3.7, describing $\text{Ker}(\mu_2)$, ordered by increasing values of i . For every $2 \leq r \leq k$ and $l = 2r-1, \dots, 2g-3$, let $w_l^{(2r)}$ be the vector whose coordinates are

$$(37) \quad (w_l)^{(2r)}_{ij} = \begin{cases} (j-i) i j (i-1)(j-1) \dots (i-(r-2))(j-(r-2)) & \text{if } i+j=l, i, j \geq r-1 \\ 0 & \text{if } i+j \neq l, \text{ or } 1 \leq i \leq r-2. \end{cases}$$

which are the coefficients of the l -th equation of the set of equations which describe $\text{Ker}(\mu_{2r})$ inside $(\text{Ker}(\mu_{2(r-1)}))$, ordered by increasing values of i .

For any $3 \leq l \leq 2g-3$ set

$$(38) \quad n_l = \#\{(i, j) : 1 \leq i < j \leq g-1, i+j=l\}.$$

and let $B'_{k,l}$ be the $k \times n_l$ matrix whose r th row, $1 \leq r \leq k$, are the n_l coordinates of $w_l^{(2r)}$ corresponding to the indexes (i, j) such that $i+j=l$.

Observe that $B'_{k,l}$ is just the matrix with rows $w_l^{(2)}, \dots, w_l^{(2k)}$ where we have removed the entries corresponding to $i+j \neq l$. Notice that these entries are all 0 for all the vectors $w_l^{(r)}$.

Define $c_{k,l} = \min\{n_l, k\}$ and let $B_{k,l}$ be the minor of order $c_{k,l}$ given by the first $c_{k,l}$ rows and $c_{k,l}$ columns. We are going to prove that it is not zero for all $3 \leq l \leq 2g-3$ by induction on k .

Consider first the case $k = 2$. Observe that if $l = 3, 4, 2g - 4, 2g - 3$, then $n_l = 1$. In this case $B_{2,l} = j - i > 0$ where (i, j) is the only pair such that $i + j = l$. If $5 \leq l \leq 2g - 5$ then $n_l \geq 2$ and $c_{2,l} = \min(2, n_l) = 2$ and we have

$$B_{2,l} = \begin{pmatrix} j - i & j - 1 - 2 \\ ij(j - i) & (j - i - 2)(i + 1)(j - 1) \end{pmatrix}$$

where i is the minimum such that $i + j = l$. Observe that

$$\det(B_{2,l}) = (j - 1)(j - 1 - 2) \det \begin{pmatrix} 1 & 1 \\ ij & (i + 1)(j - 1) \end{pmatrix}$$

Set

$$A_{2,l} := \begin{pmatrix} 1 & 1 \\ ij & (i + 1)(j - 1) \end{pmatrix}.$$

Subtracting the first column of $A_{2,l}$ from the second we obtain the matrix

$$\begin{pmatrix} 1 & 0 \\ ij & j - i - 1 \end{pmatrix}.$$

whose determinant is $j - i - 1 > 0$. Now assume that for every $2 \leq k_0 \leq k$ and for every $3 \leq l \leq 2g - 3$, $B_{k,l}$ is not zero. We want to prove that the same holds for $k + 1$. If $n_l \leq k$, then $n_l < k + 1$, $c_{k+1,l} = \min\{n_l, k\} = n_l$ and $B_{k+1,l} = B_{k,l}$. In this case, we have nothing to prove. Assume then $n_l > k$. Then $c_{k+1,l} = \min\{n_l, k + 1\} = k + 1$. Then $B_{k+1,l}$ is the following $k + 1 \times k + 1$ matrix

$$\begin{pmatrix} j - i & j - i - 2 & \dots & j - i - 2k \\ (j - i)ji & (j - 2 - i)(i + 1)(j - 1) & \dots & (j - i - 2k)(j - k)(i + k) \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ (j - i)ji \dots (j - (k + 1 - 2))(i - (k + 1 - 2)) & \dots & (j - i - 2k)(j - k)(i + k) \dots (j - 2k + 1)(i + 1) \end{pmatrix}$$

where i is the minimum such that $i + j = l$ and where we mean that every entry in the r -th row, $1 \leq r \leq k + 1$ (corresponding to an index (i, j)) is zero whenever $i \leq r - 1$.

The determinant of the above matrix is equal to the product of $(j - i) \dots (j - i - 2k)$ by the determinant of the following matrix

$$\begin{pmatrix} 1 & 1 & \dots & 1 \\ ji & (i + 1)(j - 1) & \dots & (j - k)(i + k) \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ (j - i)ji \dots (j - (k + 1 - 2))(i - (k + 1 - 2)) & \dots & (j - k)(i + k) \dots (j - 2k + 1)(i + 1) \end{pmatrix}$$

Now consider the matrix above and subtract each column to the previous one. Then one gets a matrix

$$\left(\begin{array}{cccc} & 1 & & 0 \\ & j^i & (j-1)-i & \dots \\ & \cdot & \cdot & \cdot \\ & \cdot & \cdot & \cdot \\ (j-i)j^i \dots (j-(k+1-2))(i-(k+1-2)) & & \dots & (j-1-i-2(k-1))(j-1-(k-1))(i+k-1) \dots (j-2(k-1)+1)(i+1) \end{array} \right)$$

Observe that

$$\left(\begin{array}{ccc} & (j-1)-i & \dots \\ & \cdot & \cdot \\ & \cdot & \cdot \\ (j-1-i)(j-1) \dots ((j-1-(k-2)))(i-((k-2))) & \dots & (j-1-i-2(k-1))(j-1-(k-1))(i+k-1) \dots (j-2(k-1)+1)(i+1) \end{array} \right)$$

is $B_{k,l-1}$ (where as we have already said, we mean that every entry corresponding to a index (i, j) in the r -th row, $1 \leq r \leq k$ is zero whenever $i \leq r - 1$. The matrix has not zero determinant by the inductive hypothesis. Hence we have shown, by induction, that for any $3 \leq l \leq 2g - 3$ and for any $k \geq 2$, $(w_l)^2, \dots, (w_l)^{2k}$ impose exactly $c_{k,l}$ linearly independent conditions.

Now let s_k be the number of the indexes l such that $c_{k,l} > c_{k-1,l}$. Observe that the condition $c_{k,l} > c_{k-1,l}$ is equivalent to say that l -th equation of $Ker(\mu_{2k})$ is independent from the l -th equation of $Ker(\mu_2), \dots, Ker(\mu_{2(k-1)})$. Then we have

$$(39) \quad \dim(Ker(\mu_{2k})) = \dim Ker(\mu_{2(k-1)}) - s_k.$$

Notice that $c_{k,l} > c_{k-1,l}$ if and only if $k - 1 < n_l$. We claim that this happens if and only if $2k + 1 \leq l \leq 2g - (2k + 1)$.

In fact, if $2k + 1 \leq l \leq 2g - (2k + 1)$, we have at least k coordinates (i, j) such that $i + j = l$. Indeed write l as $l = 2k + 1 + m$, $0 \leq m \leq 2g - 2(2k + 1)$ (where we are using the assumption $2g - 2(2k + 1) \geq 0$).

If l is odd (equivalently m is even), these are given by

$$(i, j) = \left(1 + \frac{m}{2}, 2k + \frac{m}{2}\right), \dots, \left(k + \frac{m}{2}, k + 1 + \frac{m}{2}\right)$$

These are easily seen to be admissible indexes (i, j) , that is they satisfy $1 \leq i < j$, $i + j = l$ and $j \leq g - 1$. If l is even (equivalently m is odd), these are given by

$$(i, j) = \left(1 + \lfloor \frac{m}{2} \rfloor, 2k + \lfloor \frac{m}{2} \rfloor\right), \dots, \left(k + \lfloor \frac{m}{2} \rfloor, k + 2 + \lfloor \frac{m}{2} \rfloor\right).$$

This shows that $n_l \geq k$.

On the other hand, if $l \leq 2k$ or $l \geq 2g - 2k$, it is easy to see that the number of admissible indexes (i, j) such that $i + j = l$ is strictly less than k . We then conclude that $s_k = 2g - (2k + 1) - 2k = 2g - (4k + 1)$ and hence $\dim(Ker(\mu_{2k})) = \dim Ker(\mu_{2(k-1)}) - s_k = \dim Ker(\mu_{2(k-1)}) - (2g - (4k + 1))$. So we have proven that $Rank(\mu_{2k}) = 2g - (4k + 1)$.

By the equality: $\dim(Ker(\mu_{2k})) = \dim Ker(\mu_{2(k-1)}) - (2g - (4k + 1))$, and using that $\dim(I_2) = \frac{(g-1)(g-2)}{2}$, we get

$$\dim(Ker(\mu_{2k})) = \frac{(g-1)(g-2)}{2} - \sum_{i=1}^k (2g - (4i + 1)) = \frac{(g-1)(g-2)}{2} + k(2k - 2g + 3),$$

for every $k \leq \frac{g-1}{2}$. So if g is odd and $k = \frac{g-1}{2}$, we have $\dim(Ker(\mu_{2k})) = \dim(Ker(\mu_{g-1})) = 0$, hence for $k > \frac{g-1}{2}$, $\mu_{2k} \equiv 0$. If g is even, and $k = \frac{g-2}{2}$, we have $\dim(Ker(\mu_{2k})) = \dim(Ker(\mu_{g-2})) = 0$, hence for $k > \frac{g-2}{2}$, $\mu_{2k} \equiv 0$.

□

4. SCHIFFER VARIATIONS AND WEIERSTRASS POINTS

In this section we recall the definition and the basic properties of (higher) Schiffer variations.

Let C be a smooth curve of genus g , take a point $p \in C$ and fix a local coordinate z centred in p . For $1 \leq n \leq 3g - 3$, we define the n^{th} Schiffer variation at p to be the element $\xi_p^n \in H^1(C, T_C) \cong H_{\bar{\partial}}^{0,1}(T_C)$ whose Dolbeault representative is given by $\frac{\bar{\partial}\rho_p}{z^n} \frac{\partial}{\partial z}$, where ρ_p is a bump function in p which is equal to one in a small neighborhood U containing p , $\xi_p^n = [\frac{\bar{\partial}\rho_p}{z^n} \frac{\partial}{\partial z}]$. Clearly ξ_p^n depends on the choice of the local coordinate z . Take $1 \leq n \leq 3g - 3$. Consider the exact sequence

$$0 \rightarrow T_C \rightarrow T_C(np) \rightarrow T_C(np)|_{np} \rightarrow 0,$$

and the induced exact sequence in cohomology:

$$0 \rightarrow H^0(T_C(np)) \rightarrow H^0(T_C(np)|_{np}) \xrightarrow{\delta_p^n} H^1(T_C).$$

By Riemann Roch, if $n < 2g - 2$, or $n \leq 3g - 3$ and p is a general point, we have:

$$h^0(T_C(np)) = 0.$$

Hence we have an inclusion

$$\delta_p^n : H^0(T_C(np)|_{np}) \cong \mathbb{C}^n \hookrightarrow H^1(T_C)$$

and the image of δ_p^n in $H^1(C, T_C)$ is the n -dimensional subspace generated by ξ_p^1, \dots, ξ_p^n (see [9]) for more details).

Given an element $\zeta \in H^1(C, T_C)$, consider the map given by cup product:

$$\cup \zeta : H^0(C, K_C) \rightarrow H^1(C, \mathcal{O}_C).$$

The rank of ζ is by definition the rank of the map $\cup \zeta$.

Remark 4.1. Notice that $H^0(K_C(-np)) \subset \text{Ker}(\cup \xi_p^n)$, for $n \leq g$. In fact, $\omega \in \text{ker}(\cup \xi_p^n)$ if and only if for any $\alpha \in H^0(K_C)$, we have $\xi_p^n(\omega\alpha) = 0$. So if z is a local coordinate around p and $\omega = f(z)dz$, $\alpha = g(z)dz$ are local expressions, by [9, Lemma 2.2] we have:

$$\xi_p^n(\omega\alpha) = \frac{2\pi i}{(n-1)!} (fg)^{(n-1)}(p) = 0,$$

if $\omega \in H^0(K_C(-np))$, $\forall \alpha \in H^0(K_C)$. Hence ξ_p^n has rank $\leq n$.

Let \mathcal{H}_g be the moduli space of hyperelliptic curves of genus g . If C is a hyperelliptic curve, then the tangent space of \mathcal{H}_g at $[C]$ is $H^1(T_C)^+$, that is the invariant subspace with respect to the hyperelliptic involution σ .

Assume $p \in C$ is a Weierstrass point. From the Dolbeault representation of the Schiffer variations, one immediately sees that the elements ξ_p^{2k+1} are σ -invariant, hence they belong to $H^1(T_C)^+$. So we have a subspace

$$\langle \xi_p^1, \xi_p^3, \dots, \xi_p^{2g-3} \rangle \subset H^1(T_C)^+$$

of dimension $g - 1$.

Denote as usual by $|L|$ the g_2^1 . We conclude this section giving a basis of $H^0(K_C \otimes L^\vee)$ that will be useful in the next sections.

Lemma 4.2. *There exists a basis $\omega_1, \dots, \omega_{g-1}$ of $H^0(K_C \otimes L^\vee)$ such that $\text{ord}_p \omega_k = 2g - 2k - 2$, for all $k = 1, \dots, g - 1$.*

Proof. Let us set $M := K_C \otimes L^\vee$. By Riemann-Roch we have $h^0(M) = h^0(K_C(-2p)) = g - 1$, $h^0(M(-p)) = h^0(M(-2p)) = g - 2$, since p is a Weierstrass point. Analogously, for every $1 \leq k \leq g - 2$, we get $h^0(M(-(2k - 1)p)) = h^0(M(-2kp)) = g - k - 1$. So we choose a basis of $H^0(M)$ as follows:

- $h^0(M(-(2g - 4)p)) = 1$ and we take $\omega_1 \in H^0(M(-(2g - 4)p))$, $\omega_1 \neq 0$.
- $h^0(M(-(2g - 6)p)) = 2$ and we take $\omega_2 \in H^0(M(-(2g - 6)p)) \setminus H^0(M(-(2g - 4)p))$, so $H^0(M(-(2g - 6)p)) = \langle \omega_1, \omega_2 \rangle$.
- \vdots
- $h^0(M(-2p)) = g - 2$, and we set $H^0(M(-2p)) = \langle \omega_1, \dots, \omega_{g-2} \rangle$.
- $h^0(M) = g - 1$; $\omega_{g-1} \in H^0(M) \setminus H^0(M(-p))$, and we have $H^0(M) = \langle \omega_1, \dots, \omega_{g-1} \rangle$.

□

5. SECOND FUNDAMENTAL FORM OF THE HYPERELLIPTIC TORELLI LOCUS

Let C be a hyperelliptic curve of genus $g \geq 3$, L be the line bundle giving the g_2^1 . Set $M = K_C \otimes L^\vee$ and $\pi : C \rightarrow \mathbb{P}^1$ the map induced by $|L|$.

Denote by σ the hyperelliptic involution and consider the decomposition $H^0(C, K_C^{\otimes 2}) \cong H^0(C, K_C^{\otimes 2})^+ \oplus H^0(C, K_C^{\otimes 2})^-$ of $H^0(C, K_C^{\otimes 2})$ in invariant and anti-invariant subspaces under the action of σ .

Denote by \mathcal{H}_g the hyperelliptic locus in \mathcal{M}_g and by

$$j_h : \mathcal{H}_g \rightarrow \mathcal{A}_g$$

the restriction of the Torelli map to \mathcal{H}_g . We endow \mathcal{A}_g with the Siegel metric, that is the orbifold metric induced by the symmetric metric on the Siegel space $Sp(2g, \mathbb{R})/U(g)$ of which \mathcal{A}_g is the quotient under the action of $Sp(2g, \mathbb{Z})$. The map j_h is an orbifold immersion (see [15]) and we have the following tangent bundle exact sequences

$$(40) \quad \begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ 0 & \longrightarrow & T_{\mathcal{H}_g} & \longrightarrow & T_{\mathcal{A}_g|_{\mathcal{H}_g}} & \longrightarrow & N_{\mathcal{H}_g/\mathcal{A}_g} \longrightarrow 0 \\ & & \downarrow & & \downarrow = & & \downarrow \\ & & T_{\mathcal{M}_g|_{\mathcal{H}_g}} & \longrightarrow & T_{\mathcal{A}_g|_{\mathcal{H}_g}} & \longrightarrow & N_{\mathcal{M}_g/\mathcal{A}_g|_{\mathcal{H}_g}} \longrightarrow 0 \end{array}$$

Denote by

$$(41) \quad \rho_{HE} : N_{\mathcal{H}_g|\mathcal{A}_g}^* \rightarrow \text{Sym}^2 \Omega_{\mathcal{H}_g}^1,$$

the dual of the second fundamental form of j_h .

At a point $[C] \in \mathcal{H}_g$, the dual of (40) is

$$\begin{array}{ccccccc} 0 & \longrightarrow & I_2 & \longrightarrow & S^2 H^0(K_C) & \xrightarrow{m} & H^0(K_C^{\otimes 2}) \\ & & \downarrow & & \downarrow = & & \downarrow \\ 0 & \longrightarrow & I_2 & \longrightarrow & S^2 H^0(K_C) & \xrightarrow{m} & H^0(K_C^{\otimes 2})^+ \longrightarrow 0 \end{array}$$

where m is the multiplication map and I_2 can be identified with the vector space of quadrics containing the rational normal curve.

So at a point $[C]$ the map ρ_{HE} is a linear map

$$I_2 \rightarrow \text{Sym}^2(H^0(C, K_C^{\otimes 2})^+).$$

In [5, Prop. 5.1] it is proven that $\forall Q \in I_2, \forall v, w \in H^1(T_C)^+$ we have

$$(42) \quad \rho_{HE}(Q)(v \odot w) = \rho(Q)(v \odot w),$$

where ρ is the Hodge Gaussian map introduced in [8, Proposition-Definition 1.3].

In [11] the second fundamental form ρ_{HE} of the hyperelliptic locus has been studied using the Hodge Gaussian map ρ and the second gaussian map of the canonical bundle. In [11, Theorem 6.2] it has been proven that if Y is a germ of totally geodesic subvariety of \mathcal{A}_g generically contained in the hyperelliptic Torelli locus, then its dimension is at most $g + 1$.

In [9] a relation between higher even Gaussian maps of the canonical bundle on a smooth projective curve of genus $g \geq 4$ and the second fundamental form of the Torelli map is given. This is a generalisation of a result obtained by Colombo, Pirola and Tortora on the second Gaussian map and the second fundamental form in [8] (see also [6, Theorem 2.2]). In fact, in [6, Theorem 2.2]) it is proven that for any $Q \in I_2$, we have

$$(43) \quad \rho(Q)(\xi_p \odot \xi_p) = -2\pi i \mu_2(Q)(p).$$

More precisely, the computations of $\rho(Q)$ on Schiffer variations given in [9, Proposition 3.1 and Remark 3.2], for quadrics $Q \in \text{Ker}(\mu_{2k})$ will be the main tool in the next section.

6. ISOTROPIC SUBSPACES

In this section, given a hyperelliptic curve C of genus $g \geq 3$ and a Weierstrass point $p \in C$, we will determine for any $k \leq \lfloor \frac{g-3}{2} \rfloor$, the maximum $m \leq \lfloor \frac{g-1}{2} \rfloor$, such that the subspace $V_m = \langle \xi_p^1, \xi_p^2, \dots, \xi_p^{2m+1} \rangle \subset V_{\lfloor \frac{g-1}{2} \rfloor} \subset H^1(T_C)^+$ is isotropic with respect to all the quadrics $\rho(Q), \forall Q \in \text{Ker}(\mu_{2k})$. We will show that $m = k$, namely that V_k is isotropic for $\rho(Q), \forall Q \in \text{Ker}(\mu_{2k})$, while V_{k+1} is not (see Theorems 6.5, 6.7).

In order to do this we recall that, by (42), it suffices to compute $\rho(Q)(\xi_p^r \odot \xi_p^n)$, for odd r, n .

To do this we will use [9, Remark 3.2], that we now recall for the reader's convenience. Assume Q is quadric in I_2 , $Q = \sum_{\alpha, \beta=1}^u b_{\alpha\beta} \gamma_\alpha \gamma_\beta$, where $\gamma_1, \dots, \gamma_u$ are elements of $H^0(K_C)$. Choose a local coordinate z around p and take a local expression of γ_α around p : $\gamma_\alpha = g_\alpha(z)dz$.

Remark 6.1. ([9, Remark 3.2]) *Assume*

$$\sum_{\alpha, \beta} b_{\alpha\beta} g_\alpha^{(h)}(0) g_\beta^{(l)}(0) = 0, \quad \forall h, l \geq 0, \quad h + l \leq m,$$

then

$$\rho_{HE}(Q)(\xi_p^n \odot \xi_p^r) = \rho(Q)(\xi_p^n \odot \xi_p^r) = 0, \quad \text{if } r + n \leq m, \quad \forall \text{ odd } r, n \geq 1,$$

and if $r + n = m + 1$, then

$$\rho_{HE}(Q)(\xi_p^n \odot \xi_p^r) = \rho(Q)(\xi_p^n \odot \xi_p^r) = 2\pi i \left(\sum_{k=0}^{n-1} \left(\sum_{\alpha, \beta=1}^u b_{\alpha\beta} f_i^{(2k+1-l)}(0) f_j^{(l)}(0) \right) \frac{(n-l)}{l!(2k+1-l)!} \right).$$

Recall that a basis of I_2 is given by the following quadrics (see Lemma 3.3):

$$(44) \quad Q_{ij} := s\omega_i \odot t\omega_j - s\omega_j \odot t\omega_i$$

for $1 \leq i < j \leq g-1$, where L is the g_2^1 , $H^0(L) = \langle s, t \rangle$, and $H^0(M) = H^0(K_C \otimes L^\vee) = \langle \omega_1, \dots, \omega_{g-1} \rangle$, where the sections ω_i are as in Lemma 4.2. Let Q be a quadric in I_2 , then

$$Q = \sum_{1 \leq i < j \leq g-1} a_{ij} Q_{ij} = \sum_{\alpha, \beta=1}^u b_{\alpha\beta} \gamma_\alpha \gamma_\beta,$$

where $\gamma_1, \dots, \gamma_u$ are elements of $H^0(K_C)$. Choose a local coordinate z around p and take a local expression of γ_α around p : $\gamma_\alpha = g_\alpha(z)dz$.

Now let $k \geq 0$ and assume $Q \in \text{Ker}(\mu_{2k})$. Recall from Section 2 that

$$Q \in \text{Ker}(\mu_{2k}) \Leftrightarrow \sum_{\alpha, \beta=1}^u b_{\alpha\beta} g_\alpha^{(h)} g_\beta^{(l)} \equiv 0 \quad \forall h + l \leq 2k + 1.$$

Moreover by Lemma 3.5 we have

$$\begin{aligned} \mu_{2k+2}(Q) &= \sum_{\alpha, \beta=1}^u b_{\alpha\beta} g_\alpha^{(h)} g_\beta^{(l)} = \mu_{1,L}(s \wedge t) \mu_{2k+1,M} \left(\sum_{1 \leq i < j \leq g-1} a_{ij} \omega_i \wedge \omega_j \right) = \\ &= (s't - t's) \sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(r)} \omega_j^{(s)} - \omega_i^{(s)} \omega_j^{(r)}) (dz)^{2k+4} \quad \forall h + l = 2k + 2, \quad \forall r + s = 2k + 1. \end{aligned}$$

It follows that

$$\sum_{\alpha, \beta=1}^u b_{\alpha\beta} g_\alpha^{(h)} g_\beta^{(l)}(p) = 0 \quad \forall h + l = 2k + 2,$$

because $(s't - t's)(p) = 0$, since p is a Weierstrass point.

Now we want to determine the least $m \geq 2k + 3$ such that

$$\sum_{\alpha, \beta=1}^u b_{\alpha\beta} g_{\alpha}^{(h)} g_{\beta}^{(l)}(p) \neq 0,$$

for some h, l with $h + l = m$.

Lemma 6.2. *Let $Q = \sum_{1 \leq i < j \leq g-1} a_{ij} Q_{ij} \in I_2$. If $Q \in \text{Ker}(\mu_{2k})$, setting $j = g - s$, we have*

$$a_{i, g-s} = 0 \quad \forall g - (2k + 2 - s) \leq i < g - s$$

Proof. Recall that in Lemma 3.7 we have shown that $Q \in \text{Ker}(\mu_{2k})$, $k \geq 2$, if and only if for every $2 \leq l \leq 2g - 3$,

$$\sum_{\substack{1 \leq i < j \\ i+j=l}} a_{ij}(j-i) = 0,$$

and for every m, l such that $2 \leq m \leq k$, $2m - 1 \leq l \leq 2g - 3$,

$$\sum_{\substack{1 \leq i < j \\ i+j=l \\ i \geq m-1, \\ j \geq m-1}} a_{ij}(j-i)ij(i-1)(j-1)\dots(i-(m-2))(j-(m-2)) = 0.$$

As in the proof of Theorem 3.1), for any $3 \leq l \leq 2g - 3$ set $n_l = \#\{(i, j) : 1 \leq i < j \leq g - 1, i + j = l\}$. In Theorem 3.1 we have shown that when $n_l \geq k$ the equations of μ_2, \dots, μ_{2k} relative to the index l are linearly independent. When $l = 2g - (2k + 1)$, or $l = 2g - (2k + 1) - 1$, we have that $n_l = k$. This indeed follows from the last part of the proof of Theorem 3.1. Hence when l is equal to $2g - (2k + 1)$, or $2g - (2k + 1) - 1$, we have a system of k linearly independent equations in k variables. Applying the same argument for μ_2, \dots, μ_{2k} , this gives $a_{i, g-s} = 0$ for $l = i + g - s \geq 2g - (2k + 1) - 1$, and hence for $i \geq g - (2k + 2 - s)$ (and $j = g - s > i$). □

Remark 6.3. *By Lemma 4.2, with our choice of the basis $\omega_1, \dots, \omega_{g-1}$ of $H^0(M)$, we have*

$$\omega_i^{(h)}(p) = 0 \quad \forall h \leq 2g - (2i + 3).$$

Lemma 6.4. *With the above notation, we have*

$$\sum_{1 \leq i < j \leq g-1} a_{ij}(\omega_i^{(h)}\omega_j^{(l)} - \omega_i^{(l)}\omega_j^{(h)})(p) = 0 \quad \forall h + l \leq 4k + 1.$$

Proof. We may assume $h > l$, since if $h = l$ we have trivially zero. From Remark 6.3, $\omega_j^{(l)}(p) = 0 \quad \forall j$ such that $2j + 3 \leq 2g - l$. Moreover, since $i < j$, this also implies $\omega_i^{(l)}(p) = 0$, so the terms in the sum with $2j + 3 \leq 2g - l$ are identically zero. Then we can assume that the indexes j that appear in the sum are those satisfying $2j + 3 \geq 2g - l + 1$, which gives $j \geq g - \frac{l+2}{2}$. Denote by $\tilde{l} := \frac{l+2}{2}$. Hence we have

$$\sum_{1 \leq i < j \leq g-1} a_{ij}(\omega_i^{(h)}\omega_j^{(l)} - \omega_i^{(l)}\omega_j^{(h)})(p) = \sum_{1 \leq i < j \leq g-1, j \geq g-\tilde{l}} a_{ij}(\omega_i^{(h)}\omega_j^{(l)} - \omega_i^{(l)}\omega_j^{(h)})(p) =$$

$$\begin{aligned}
&= \sum_{1 \leq i < g-1} a_{ig-1}(\omega_i^{(h)}\omega_{g-1}^{(l)} - \omega_i^{(l)}\omega_{g-1}^{(h)})(p) + \sum_{1 \leq i < g-2} a_{ig-2}(\omega_i^{(h)}\omega_{g-2}^{(l)} - \omega_i^{(l)}\omega_{g-2}^{(h)})(p) \\
&\quad + \cdots + \sum_{1 \leq i < g-\tilde{l}} a_{ig-\tilde{l}}(\omega_i^{(h)}\omega_{g-\tilde{l}}^{(l)} - \omega_i^{(l)}\omega_{g-\tilde{l}}^{(h)})(p).
\end{aligned}$$

Let us focus on the first term of the above sum. From Lemma 6.2, $Q \in \text{Ker}(\mu_{2k})$ implies $a_{ig-1} = 0 \quad \forall g - (2k + 1) \leq i \leq g - 2$. Then we have

$$(45) \quad \sum_{1 \leq i < g-1} a_{ig-1}(\omega_i^{(h)}\omega_{g-1}^{(l)} - \omega_i^{(l)}\omega_{g-1}^{(h)})(p) = \sum_{i=1}^{g-2k-2} a_{ig-1}(\omega_i^{(h)}\omega_{g-1}^{(l)} - \omega_i^{(l)}\omega_{g-1}^{(h)})(p).$$

Moreover, again by Remark 6.3, if $2i + 3 \leq 2g - h < 2g - l$, then $\omega_i^{(h)}(p) = \omega_i^{(l)}(p) = 0$. But in the above sum we have $2i + 3 \leq 2(g - 2k - 2) + 3 = 2g - 4k - 1$. Hence, if $h \leq 4k + 1$ (so $l < 4k + 1$), then $2g - 4k - 1 \leq 2g - h < 2g - l$, so $2i + 3 \leq 2g - h < 2g - l$ in all summands in (45), so the sum is identically zero.

Let us now consider the general term, where $j = g - s$. From Lemma 6.2, we have $a_{ig-s} = 0 \quad \forall g - (2k + 2 - s) \leq i \leq g - s$. Hence the i 's that actually appear in the sum are those satisfying $i \leq g - (2k + 2 - s) - 1 = g - 2k - 3 + s$. In the same way as we did for the first term, for $h \leq 4k + 3 - 2s$, we have $2i + 3 \leq 2(g - 2k - 3 + s) \leq 2g - h$, so $\omega_i^{(h)}(p) = \omega_i^{(l)}(p) = 0$ and the term is zero.

It remains to look at the cases when $h \geq 4k + 4 - 2s$. Since $h + l \leq 4k + 1$ by assumption, we must have $4k + 4 - 2s \leq h + l \leq 4k + 1$, which gives $l \leq 2s - 3$. For these values of l , we have $\omega_{g-s}^{(l)}(p) = 0$ by Remark 6.3. Hence the term is zero again, because $i < j$, so $\omega_i^{(l)}(p) = 0$. Thus we have shown that all terms are zero. \square

We are now ready to state and prove our result.

Theorem 6.5. *Let C be an hyperelliptic curve of genus g , $p \in C$ a Weierstrass point. Let $\gamma_1, \dots, \gamma_r \in H^0(K_C)$ where locally $\gamma_\alpha = g_\alpha dz_\alpha$ and let*

$$Q = \sum_{\alpha, \beta=1}^u b_{\alpha\beta} \gamma_\alpha \gamma_\beta = \sum_{1 \leq i < j \leq g-1} a_{ij} Q_{ij} \in \text{Ker}(\mu_{2k}).$$

Then

$$\sum_{\alpha, \beta=1}^u b_{\alpha\beta} g_\alpha^{(h)} g_\beta^{(l)}(p) = 0 \quad \forall h + l \leq 4k + 3.$$

Proof. We have to compute the following expression $\forall h + l \leq 4k + 3$:

$$\begin{aligned}
(46) \quad &\sum_{\alpha, \beta=1}^u b_{\alpha\beta} g_\alpha^{(h)} g_\beta^{(l)}(p) = \\
&= \sum_{1 \leq i < j \leq g-1} a_{ij} [(s\omega_i)^{(h)}(t\omega_j)^{(l)} + (t\omega_j)^{(h)}(s\omega_i)^{(l)} - (s\omega_j)^{(h)}(t\omega_i)^{(l)} - (t\omega_i)^{(h)}(s\omega_j)^{(l)}](p).
\end{aligned}$$

We have

$$(s\omega_i)^{(h)}(t\omega_j)^{(l)} - (s\omega_j)^{(h)}(t\omega_i)^{(l)} = \sum_{n=0}^h \sum_{m=0}^l \binom{h}{n} \binom{l}{m} s^{(h-n)} t^{(l-m)} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(n)} \omega_i^{(m)}).$$

Hence the term inside the square brackets of Equation (46) becomes

$$\sum_{n=0}^h \sum_{m=0}^l \binom{h}{n} \binom{l}{m} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(n)} \omega_i^{(m)}) (s^{(h-n)} t^{(l-m)} - t^{(h-n)} s^{(l-m)}).$$

So equation (46) is

$$(47) \quad \sum_{n=0}^h \sum_{m=0}^l \binom{h}{n} \binom{l}{m} (s^{(h-n)} t^{(l-m)} - t^{(h-n)} s^{(l-m)})(p) \sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(n)} \omega_i^{(m)})(p).$$

From Lemma 6.4 we have that

$$\sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(n)} \omega_i^{(m)})(p) = 0 \quad \forall m + n \leq 4k + 1.$$

Since $n \leq h, m \leq l$ we have $m + n \leq h + l$, so this implies that the expression in (47) is zero for all h, l such that $h + l \leq 4k + 1$.

Now assume $h + l = 4k + 2$. If $n + m \leq 4k + 1$ the expression is zero exactly as before, so the only case to consider is $n + m = 4k + 2$ and we must have $n = h, m = l$. But in this case we have trivially zero, because $(st - ts) = 0$.

We are only left to the case $h + l = 4k + 3$. Now again, if $m + n \leq 4k + 1$ we get zero and if $m + n = 4k + 3$ we get zero as in the previous case (this would imply again $n = h, m = l$). So we can only have $n + m = 4k + 2$. This implies either $n = h - 1, m = l$ or $n = h, m = l - 1$. In both cases, the first factor of (47) is (up to scalar) $(s't - t's)(p) = 0$ since p is a Weierstrass point. Hence we have proven that (47) is zero $\forall h + l \leq 4k + 3$ and this concludes the proof. \square

For any $k \leq 2g - 3$, consider the $(k + 1)$ -dimensional subspace $V_k := \langle \xi_p^1, \xi_p^3, \dots, \xi_p^{2k+1} \rangle \subset H^1(T_C)^+$. We have the following

Theorem 6.6. *Let $0 \leq k \leq \lfloor \frac{g-3}{2} \rfloor$ and let $Q \in \text{Ker}(\mu_{2k})$. Then we have*

$$\rho_{HE}(Q)(\xi_p^l \odot \xi_p^m) = \rho(Q)(\xi_p^l \odot \xi_p^m) = 0 \quad \forall l + m \leq 4k + 3.$$

This implies that the subspace $V_k = \langle \xi_p^1, \xi_p^3, \dots, \xi_p^{2k+1} \rangle \subset H^1(T_C)^+$ is isotropic for $\rho_{HE}(Q)$, $\forall Q \in \text{Ker}(\mu_{2k})$.

Proof. The second assertion follows directly from the first one. The first one follows immediately by Remark 6.1 ([9, Remark 3.2]) and by Theorem 6.5. \square

We have the following theorem.

Theorem 6.7. *Assume $0 \leq k \leq \lfloor \frac{g-3}{2} \rfloor$. Then there exists $Q \in \text{Ker}(\mu_{2k})$ such that $\rho(Q)(\xi_p^{2k+1} \odot \xi_p^{2k+3}) \neq 0$. Hence the subspace $V_{k+1} = \langle \xi_p^1, \dots, \xi_p^{2k+1}, \xi_p^{2k+3} \rangle \subset H^1(T_C)^+$ is not isotropic for $\rho_{HE}(\text{Ker}(\mu_{2k}))$.*

Proof. Let $Q = \sum_{\alpha, \beta=1}^r b_{\alpha, \beta} \gamma_\alpha \otimes \gamma_\beta$, where $\gamma_1, \dots, \gamma_r \in H^0(K_C)$. We will prove that $\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) \neq 0$. By Remark 6.1, we know that

$$(48) \quad \rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) = 2\pi i \sum_{u=0}^{2k} \left[\sum_{\alpha, \beta=1}^r b_{\alpha, \beta} \gamma_\alpha^{(4k+4-u)}(0) \gamma_\beta^{(s)}(0) \right] \frac{(2k+1-u)}{u!(4k+4-u)!}$$

Let us focus our attention on the term inside the square brackets, which is

$$\sum_{\alpha, \beta=1}^r b_{\alpha, \beta} \gamma_\alpha^{(4k+4-u)}(0) \gamma_\beta^{(u)}(0). \quad (*)$$

We see that we must compute the derivatives of order (h, l) of Q where $h + l = 4k + 4$ and $0 \leq l \leq 2k$. In order to do this, we will again use our basis of $H^0(K_C \otimes L^\vee)$ given by the ω_i 's, exactly as in (46) and (47). Hence we must compute

$$\sum_{n=0}^h \sum_{m=0}^l \binom{h}{n} \binom{l}{m} (s^{(h-n)} t^{(l-m)} - t^{(h-n)} s^{(l-m)})(p) \sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(n)} \omega_i^{(m)})(p)$$

where $h + l = 4k + 4$. If $n + m \leq 4k + 1$, we know that it is zero by Lemma 6.4. If $n + m = 4k + 4$, then in the equation above, we have $n = h$, $m = l$, so the first factor is (up to scalar) $(st - ts) = 0$. If $n + m = 4k + 3$, we have either $n = h - 1$ and $m = l$ or $n = h$ and $m = l - 1$. In both cases, the first factor is (up to scalar) $(s't - t's)(p) = 0$, because p is a Weierstrass point. So we can only have $n + m = 4k + 2$, where $h = n$ and $m = l - 2$ or $h = n - 2$ and $m = l$, since if $n = h - 1$ and $m = l - 1$, the first factor is $(s't' - t's') = 0$.

Summarising, in the above equation we only have to consider the following terms:

- (1) $n = h$ and $m = l - 2$, which only works if $l \geq 2$ and gives

$$(st'' - ts'')(p) \sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(h)} \omega_j^{(l-2)} - \omega_j^{(h)} \omega_i^{(l-2)})(p).$$

- (2) $n = h - 2$ and $m = l$, which gives

$$(s''t - t''s)(p) \sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(h-2)} \omega_j^{(l)} - \omega_j^{(h-2)} \omega_i^{(l)})(p).$$

Let us consider the first case. By Remark 6.3, we have $\omega_j^{(l-2)}(p) = 0$ if $2j + 3 \leq 2g - l + 2$, which gives $2j \leq 2g - l - 1$. We have two subcases here: l odd or l even.

- Assume l is odd. We claim that we get zero. In fact, if $j \leq g - \frac{l+1}{2}$, then $\omega_j^{(l-2)}(p) = 0$ by Remark 6.3. Hence we can assume $j \geq g - \frac{l+1}{2} + 1 = g - \frac{l-1}{2}$. Let $j := g - u$, hence $u \in \{1, \dots, \frac{l-1}{2}\}$. From Lemma 6.2 we have $a_{i, g-u} = 0$ for $i \geq g - 2k - 2 + u$, so we may assume $i \leq g - 2k - 3 + u$.

We know that $\omega_i^{(l-2)}(p) = 0$ if $i \leq g - \frac{l+1}{2}$, but $i \leq g - 2k - 3 + u \leq g - \frac{l+1}{2}$ if and only if $u \leq 3 + 2k - \frac{l+1}{2}$. Observe that this is true since $u \leq \frac{l-1}{2}$ and $\frac{l-1}{2} \leq 3 + 2k - \frac{l+1}{2}$, since $l \leq 2k$. Hence $\omega_i^{(l-2)}(p) = 0$.

Now let us consider $\omega_i^{(h)}(p)$. We know that $\omega_i^{(h)}(p) = 0$ if $2i \leq 2g - h - 3$, but $i \leq g - 2k - 3 + u$ implies $2i \leq 2g - 4k - 6 + 2u$. Now, $2g - 4k - 6 + 2u \leq 2g - h - 3$ if and only if $2u \leq 4k + 3 - h$, which gives $l - 1 \leq 4k + 3 - h$, which is true if $l + h = 4k + 4$, that is our assumption. Hence $\omega_i^{(h)}(p) = 0$, for all i and hence every term vanishes.

- Now assume l is even. If $j \leq g - \frac{l+1}{2} = g - \frac{l}{2} - \frac{1}{2} < g - \frac{l}{2}$, we get $\omega_j^{(l-2)}(p) = 0$ by Remark 6.3. So we may assume $j \geq g - \frac{l}{2}$. Setting $j = g - u$, we get $u \leq \frac{l}{2}$. As in the odd case, we may assume $i \leq g - 2k - 3 + u$. Now $\omega_i^{(l-2)}(p) = 0$ if $i \leq g - \frac{l}{2} - 1$ and $g - 2k - 3 + u \leq g - \frac{l}{2} - 1$ if and only if $u \leq 2k + 2 - \frac{l}{2}$. Since $u \leq \frac{l}{2}$, we get that the last inequality is true if and only if $l \leq 2k + 2$, which is the case because $l \leq 2k$. Hence $\omega_i^{(l-2)}(p) = 0$.

We have that $\omega_i^{(h)}(p) = 0$ if $i \leq g - \frac{h}{2} - \frac{3}{2} < g - \frac{h}{2} - 1$, so $\omega_i^{(h)}(p) = 0$ if $i \leq g - \frac{h}{2} - 2$. Hence we can assume $i \geq g - \frac{h}{2} - 1$, but $i \leq g - 2k - 3 + u$ and $g - 2k - 3 + u - (g - \frac{h}{2} - 1) = u - 2 - 2k + \frac{h}{2} \leq \frac{l}{2} + \frac{h}{2} - 2 - 2k = 0$ because $h + l = 4k + 4$. Hence if $u = l/2$, there is a unique nonzero term, which is precisely $a_{g-\frac{h}{2}-1, g-\frac{l}{2}} \omega_{g-\frac{h}{2}-1}^{(h)}(p) \omega_{g-\frac{l}{2}}^{(l-2)}(p)$.

Let us now consider the second case. Again, we should distinguish the cases l odd and l even. However, arguing exactly in the same way as in the previous case, we again get that if l is odd every term is zero.

So let us assume l even, then also h is even. By Remark 6.3, $\omega_j^{(l-2)}(p) = 0$ if $j \leq g - \frac{l}{2} - \frac{3}{2} < g - \frac{l}{2} - 1$, so we may assume $j \geq g - \frac{l}{2} - 1$. As usual we put $j = g - u$ and we have $u \leq \frac{l}{2} + 1$. We have $\omega_i^{(l)}(p) = 0$ if $2i \leq 2g - l - 3$, but, as in the previous case, we can assume $i \leq g - 2k - 3 + u$. Since $l \leq 2k$, and $u \leq \frac{l}{2} + 1$, we have $2g - 6 - 4k + 2u \leq 2g - l - 3$. So $2i \leq 2g - 6 - 4k + 2u \leq 2g - l - 3$, and hence $\omega_i^{(l)}(p) = 0$.

Again, by Remark 6.3, $\omega_i^{(h-2)}(p) = 0$ if $2i \leq 2g - h - 1$. As usual we may assume $2i \leq 2g - 6 - 4k + 2u$. We have $u \leq \frac{l}{2} + 1$. We claim that the only nonzero term occurs for $u = \frac{l}{2} + 1$ and $i = g - \frac{h}{2}$. In fact, if $u \leq \frac{l}{2}$, we have $2g - 6 - 4k + 2u \leq 2g - h - 1$, so $\omega_i^{(h-2)}(p) = 0$. So $u = \frac{l}{2} + 1$, $i = g - \frac{h}{2}$ and the only term is $a_{g-\frac{h}{2}, g-\frac{l}{2}-1} \omega_{g-\frac{h}{2}}^{(h-2)}(p) \omega_{g-\frac{l}{2}-1}^{(l)}(p)$.

We have computed all the terms of (*). These are

- $(st'' - ts'')(p) \left[\binom{l}{2} a_{g-\frac{h}{2}-1, g-\frac{l}{2}} \omega_{g-\frac{h}{2}-1}^{(h)}(p) \omega_{g-\frac{l}{2}}^{(l-2)}(p) - \binom{h}{2} a_{g-\frac{h}{2}, g-\frac{l}{2}-1} \omega_{g-\frac{h}{2}}^{(h-2)}(p) \omega_{g-\frac{l}{2}-1}^{(l)}(p) \right]$ if $l \geq 2$ and h, l even;
- $-(st'' - ts'')(p) \binom{h}{2} a_{g-\frac{h}{2}, g-\frac{l}{2}-1} \omega_{g-\frac{h}{2}}^{(h-2)}(p) \omega_{g-\frac{l}{2}-1}^{(l)}(p)$ if h is even and $l = 0$.

We are now ready to complete the proof. Since $h+l = 4k+4$, this implies $\frac{h}{2} = 2k+2 - \frac{l}{2}$, so that $g - \frac{h}{2} = g - 2 - 2k + \frac{l}{2}$. We have

$$\begin{aligned} \rho(Q)(\xi_p^{2k+3}, \xi_p^{2k+1}) &= 2\pi i \sum_{\substack{2 \leq l \leq 2k \\ l \text{ even}}} \frac{(2k+1-l)}{l!(4k+4-l)!} (st'' - ts'')(p) \cdot \\ &\left[\binom{l}{2} a_{g-3-2k+\frac{l}{2}, g-\frac{l}{2}} \omega_{g-3-2k+\frac{l}{2}}^{(4k+4-l)}(p) \omega_{g-\frac{l}{2}}^{(l-2)}(p) - \binom{h}{2} a_{g-2-2k+\frac{l}{2}, g-\frac{l}{2}-1} \omega_{g-2-2k+\frac{l}{2}}^{(4k+2-l)}(p) \omega_{g-\frac{l}{2}-1}^{(l)}(p) \right] - \\ &- 2\pi i (st'' - ts'')(p) \frac{2k+1}{(4k+4)!} \binom{4k+4}{2} a_{g-2k-2, g-1} \omega_{g-2k-2}^{(4k+2)}(p) \omega_{g-1}(p) \end{aligned}$$

Notice that every term in the equation is a linear combination of $a_{r,u}$ where $r+u = 2g-2k-3$.

Let $\lambda_{k,u}$ be the coefficient of $a_{g-2k-3+u, g-u}$ where $u = 1, \dots, k+1$. We will now see that $\lambda_{k,u} \neq 0$ for every k, u . For $1 \leq u \leq k$, we have

$$\begin{aligned} \lambda_{k,u} &= \omega_{g-3-2k+u}^{(4k+4-2u)}(p) \omega_{g-u}^{(2u-2)}(p) \left[\frac{2k+1-2u}{(4k+4-2u)!} \binom{2u}{2} - \binom{4k+6-2u}{2} \frac{2k+3-2u}{(4k+6-2u)!} \right] = \\ &= \omega_{g-3-2k+u}^{(4k+4-2u)}(p) \omega_{g-u}^{(2u-2)}(p) \frac{-8u^3 + 8u^2(k+1) - 4ku - 2k - 3}{2(4k+4-2u)!} \end{aligned}$$

where the last equality follows after some easy computations. Notice that

$$-8u^3 + 8u^2(k+1) - 4ku - 2k - 3 \neq 0$$

because it is odd for every k, u . Moreover, $\omega_{g-3-2k+u}^{(4k+4-2u)}(p) \omega_{g-u}^{(2u-2)}(p) \neq 0$ because $\omega_{g-3-2k+u}^{(4k+4-2u)}(p)$ has order $4k+4-2u$ at p and $\omega_{g-u}^{(2u-2)}$ has order $2u-2$ at p (see Lemma 4.2).

Finally, if $u = k+1$ we have

$$\lambda_{k,k+1} = -\frac{1}{2(2k+2)!} \omega_{g-2-k}^{(2k+2)}(p) \omega_{g-k-1}^{(2k)}.$$

Hence $\lambda_{k,u} \neq 0$, for all $u = 1, \dots, k+1$.

So assume by contradiction that

$$(49) \quad \rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) = \sum_{u=1}^{k+1} \lambda_{k,u} a_{g-2k-3+u, g-u} = 0$$

for every $Q \in Ker(\mu_{2k})$. Recall that $Q \in Ker(\mu_{2k})$ if and only if the coefficients a_{ij} satisfy the equations given in Lemma 3.7. Moreover observe that from Theorem 3.1 it follows that we have k linearly independent equations in the variables $a_{g-2k-3+u, g-u}$ with $u = 1, \dots, k+1$. We have shown that each coefficient $\lambda_{k,u}$ in equation (49) is different from zero and up to scalar it is the product $\omega_{g-3-2k+u}^{(4k+4-2u)}(p) \omega_{g-u}^{(2u-2)}(p)$. So we can multiply the sections ω_{g-u} by a non zero scalar multiple, hence equation (49) cannot be linearly dependent from the k equations giving $Ker(\mu_{2k})$. Hence there exists a quadric $Q \in ker(\mu_{2k})$ such that $\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) \neq 0$.

□

Remark 6.8. Notice that we have just proven that a quadric $Q \in \text{Ker}(\mu_{2k})$ satisfies $\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) = 0$, if and only if the coefficients of Q satisfy the equations $a_{g-2k+3+u, g-u} = 0, \forall u = 1, \dots, k+1$ in addition to all the other equations in Lemma 3.7 defining $\text{Ker}(\mu_{2k})$. The condition $\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) = 0$ gives a hyperplane in $\text{Ker}(\mu_{2k})$ that we denote by

$$A_{k,0} := \{Q \in \text{Ker}(\mu_{2k}) : \rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+1}) = 0\}.$$

Example 6.9. If $k = 0$ we have a very concrete description of $A_{0,0}$. In fact, we have $\rho(Q_{ij})(\xi_p, \xi_p^3) = 0, \forall (i, j) \neq (g-2, g-1)$ and $\rho(Q_{g-2, g-1})(\xi_p, \xi_p^3) \neq 0$. Hence we get that $A_{0,0} = \langle Q_{i,j} \mid 1 \leq i < j \leq g-1, (i, j) \neq (g-2, g-1) \rangle$.

Theorem 6.10. Let $0 \leq k \leq \lfloor \frac{g-3}{2} \rfloor$ and let us now assume $Q \in A_{k,0}$. Then

$$\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = \sum_{u=1}^{k+1} \alpha_{k,u} a_{g-4-2k+u, g-u} \omega_{g-4-2k+u}^{(4k+6-2u)}(p) \omega_{g-u}^{(2u-2)}(p),$$

where all the coefficients $\alpha_{k,u}$ are different from zero. Moreover,

$$\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = 0 \Leftrightarrow a_{g-4-2k+u, g-u} = 0 \quad \forall u = 1, \dots, k+1.$$

Proof. The proof is very similar to the one given in Theorem 6.7, where by Remark 6.1 we must now compute

$$\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = 2\pi i \left[\sum_{u=0}^{2k+2} \sum_{\alpha, \beta=1}^r b_{\alpha, \beta} \gamma_{\alpha}^{(4k+6-u)}(0) \gamma_{\beta}^{(s)}(0) \right] \frac{(2k+3-u)}{u!(4k+6-u)!}.$$

By looking at the term inside the square brackets, we see that we must compute the derivatives of order (h, l) of Q where $h+l = 4k+6$ and $0 \leq l \leq 2k+2$. Hence we have to compute

$$(50) \quad \sum_{n=0}^h \sum_{m=0}^l \binom{h}{n} \binom{l}{m} (s^{(h-n)} t^{(l-m)} - t^{(h-n)} s^{(l-m)})(p) \sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(n)} \omega_i^{(m)})(p)$$

where $h+l = 4k+6$.

The first step is analogous to Lemma 6.4: if $Q \in A_{k,0}$ we have

$$\sum_{1 \leq i < j \leq g-1} a_{ij} (\omega_i^{(n)} \omega_j^{(m)} - \omega_j^{(m)} \omega_i^{(n)})(p) = 0 \quad \forall m+n \leq 4k+3.$$

Then one immediately sees that in (50) we only have to consider the case $n+m = 4k+4$. Now, after some computations which are analogous to the ones in Theorem 6.7, we get

$$\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = \sum_{u=1}^{k+1} \alpha_{k,u} a_{g-4-2k+u, g-u} \omega_{g-4-2k+u}^{(4k+6-2u)}(p) \omega_{g-u}^{(2u-2)}(p),$$

where $\alpha_{k,u} \neq 0$ for every u . Then we conclude as in Theorem 6.7 that

$$\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = 0 \Leftrightarrow a_{g-4-2k+u, g-u} = 0 \quad \forall u = 1, \dots, k+1.$$

□

Remark 6.11. Let $Q \in A_{k,0}$. By Theorem 6.10, the linear condition $\rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = 0$ gives a hyperplane in $A_{k,0}$. Let us denote by

$$A_{k,0,0} := \{Q \in A_{k,0} : \rho(Q)(\xi_p^{2k+3} \odot \xi_p^{2k+3}) = 0\}.$$

Following [7] we give the following definition

Definition 6.12. A nonzero direction $\zeta \in H^1(T_C)^+$ is said to be asymptotic if $\rho_{HE}(Q)(\zeta \odot \zeta) = \rho(Q)(\zeta \odot \zeta) = 0, \forall Q \in I_2$.

Notice that saying that a nonzero element $\zeta \in H^1(T_C)^+$ is an asymptotic direction means that the point $[\zeta] \in \mathbb{P}H^1(T_C)^+$ is in the base locus of the linear space of quadrics $\rho_{HE}(I_2) \subset \text{Sym}^2(H^1(T_C)^+)^{\vee}$.

Remark 6.13. Recall that by Theorem 6.6 putting $k = 0$, a Schiffer variation at a Weierstrass point ξ_p is asymptotic. Moreover, moving a branch point in \mathbb{P}^1 , one can see that there exist algebraic curves in the hyperelliptic locus having these Schiffer variations as tangent directions (see [12]).

Theorem 6.14. Let C be hyperelliptic of genus $g \geq 3$ and $p \in C$ be a Weierstrass point. Then the asymptotic directions in the space $V_{[\frac{g-2}{2}]}$, which is equal to $\langle \xi_p^1, \xi_p^3, \dots, \xi_p^{g-1} \rangle$ if g is even, and equal to $\langle \xi_p^1, \xi_p^3, \dots, \xi_p^{g-2} \rangle$ if g is odd, are exactly the Schiffer variations $\xi_p = \xi_p^1$.

Proof. By Remark 6.13 we know that the Schiffer variations at the Weierstrass points are asymptotic.

To show that these are the only ones, we will prove the result when g is even, the case g odd been very similar.

Let $v \in \langle \xi_p^1, \xi_p^3, \dots, \xi_p^{g-1} \rangle$, then $v = \sum_{i=0}^{\frac{g-2}{2}} \lambda_{2i+1} \xi_p^{2i+1}$ and let l be the maximum odd integer such $\lambda_l \neq 0$. We may assume $l \geq 3$, since $\rho(Q)(\xi_p \odot \xi_p) = -2\pi i \mu_2(Q)(p) = 0$ for all $Q \in I_2$. Then we have $v \in \langle \xi_p^1, \dots, \xi_p^l \rangle$. Since l is odd, we have $l = 2k + 1$ for some integer $1 \leq k \leq \frac{g-2}{2}$. From Theorem 6.7 we know that there exists $Q \in \text{Ker}(\mu_{2k-2})$ such that $\rho(Q)(\xi_p^{2k+1} \odot \xi_p^{2k-1}) \neq 0$ and $\rho(Q)(\xi_p^i \odot \xi_p^j) = 0$ for every $i, j \leq 2k - 1, i, j$ odd integers. Then we have

$$\begin{aligned} \rho(Q)(v \odot v) &= 2\lambda_{2k-1}\lambda_{2k+1}\rho(Q)(\xi_p^{2k-1} \odot \xi_p^{2k+1}) + \\ &\quad + \lambda_{2k+1}^2 \rho(Q)(\xi_p^{2k+1} \odot \xi_p^{2k+1}). \end{aligned}$$

First, notice that for every $k \leq \frac{g-2}{2}$ we have $A_{k-1,0} \subsetneq \text{Ker}_{\mu_{2k-2}}$ and $A_{k-1,0,0} \subsetneq A_{k-1,0}$. In fact, $\dim \text{Ker}(\mu_{2k-2}) \geq 3$, since $\dim \text{Ker}(\mu_{g-4}) = 3$ by Theorem 3.1 and $2k - 2 \leq g - 4$.

If $\lambda_{2k-1} = 0$, we simply take $Q \in A_{k-1,0} \setminus A_{k-1,0,0}$ and we get $\rho(Q)(v \odot v) \neq 0$ as desired. If $\lambda_{2k-1} \neq 0$, we choose $Q \in \text{Ker}(\mu_{2k-2}) \setminus A_{k-1,0}$ and $Q' \in A_{k-1,0} \setminus A_{k-1,0,0}$. Hence, for a linear combination $Q + aQ'$, we get

$$\rho(Q + aQ')(v \odot v) = 2\lambda_{2k-1}\lambda_{2k+1}\rho(Q)(\xi_p^{2k-1} \odot \xi_p^{2k+1}) +$$

$$+\lambda_{2k+1}^2[\rho(Q)(\xi_p^{2k+1} \odot \xi_p^{2k+1}) + a\rho(Q')(\xi_p^{2k+1} \odot \xi_p^{2k+1})].$$

By the choice of Q' , we have $\rho(Q')(\xi_p^{2k+1} \odot \xi_p^{2k+1}) \neq 0$, so if we take

$$a = -\frac{\rho(Q)(\xi_p^{2k+1} \odot \xi_p^{2k+1})}{\rho(Q')(\xi_p^{2k+1} \odot \xi_p^{2k+1})}$$

we get $\rho(Q + aQ')(v \odot v) = 2\lambda_{2k-1}\lambda_{2k+1}\rho(Q)(\xi_p^{2k-1} \odot \xi_p^{2k+1}) \neq 0$ since $Q \notin A_{k-1,0}$. \square

Remark 6.15. In [5, Corollary 3.4] it is shown that ρ_{HE} is injective, so

$$\dim \rho_{HE}(Ker \mu_{2k}) = \dim Ker \mu_{2k} = \frac{(g-1)(g-2)}{2} - k(2g-2k-3),$$

for all k . For any hyperelliptic curve C of genus $g \geq 3$ and for any Weierstrass point $p \in C$, we have proven in Theorems 6.6, 6.7 that we have the following chain of subspaces of quadrics in $\mathbb{P}H^1(T_C)^+$ and of corresponding maximal isotropic subspaces in $V_{\lfloor \frac{g-1}{2} \rfloor}$.

If g is odd:

$$\rho_{HE}(Ker(\mu_{g-3})) \subsetneq \rho_{HE}(Ker(\mu_{g-5})) \dots \subsetneq \rho_{HE}(Ker(\mu_2)) \subsetneq \rho_{HE}(I_2);$$

$$V_{\frac{g-3}{2}} \supsetneq V_{\frac{g-5}{2}} \supsetneq \dots \supsetneq V_1 \supsetneq V_0 = \langle \xi_p \rangle,$$

while if g is even we have

$$\rho_{HE}(Ker(\mu_{g-4})) \subsetneq \rho_{HE}(Ker(\mu_{g-6})) \dots \subsetneq \rho_{HE}(Ker(\mu_2)) \subsetneq \rho_{HE}(I_2),$$

$$V_{\frac{g-4}{2}} \supsetneq V_{\frac{g-6}{2}} \supsetneq \dots \supsetneq V_1 \supsetneq V_0 = \langle \xi_p \rangle.$$

Theorem 6.14 also allows us to give a bound for the dimension of a germ of a totally geodesic submanifold of \mathcal{A}_g , generically contained in the Hyperelliptic Torelli locus. Notice that, nevertheless this bound is weaker than the one proven in [11, Theorem 6.2].

Corollary 6.16. Let Y be a germ of a totally geodesic submanifold of \mathcal{A}_g generically contained in $j(\mathcal{H}_g)$ passing through $j(C)$, where C is a hyperelliptic curve of genus g . Then we have

$$\dim(Y) \leq \lfloor \frac{3g+1}{2} \rfloor.$$

Proof. We will prove it for g even, the case g odd is analogous.

Set $W := T_{j(C)}Y \subset H_1(T_C)^+$ and $V := V_{\frac{g}{2}-1} = \langle \xi_p^1, \xi_p^3, \dots, \xi_p^{g-1} \rangle$. We have $\dim(V) = \frac{g}{2}$. We claim that $\dim(V \cap W) \leq 1$. Indeed, take $v \in V \cap W$, then v is asymptotic, since Y is totally geodesic. So, by Theorem 6.14 we know that v is a multiple of ξ_p^1 . Hence we have

$$\dim(V) + \dim(W) - 1 \leq \dim(V+W) \leq \dim H^1(T_C)^+ = 2g - 1,$$

so $\dim(Y) = \dim(W) \leq 2g - \frac{g}{2} = \frac{3}{2}g$. \square

REFERENCES

- [1] A. Beauville, J. Merindol. Sections hyperplanes des surfaces K3. *Duke Math. J.* 55 (1987), no. 4, 873-878. [2](#)
- [2] C. Ciliberto, J. Harris, R. Miranda. On the surjectivity of the Wahl map. *Duke Math. Jour.* 57 (1988), 829-858. [2](#)
- [3] C. Ciliberto, R. Miranda. Gaussian maps for certain families of canonical curves. *Complex projective geometry (Trieste, 1989/Bergen, 1989)*, 106- 127. *London Math. Soc. Lecture Note Ser.*, 179 Cambridge University Press, Cambridge, 1992 [2](#)
- [4] Colombo, E., Frediani P. Some results on the second Gaussian map for curves. *Michigan Mathematical Journal*, 58(3) 745-758 December 2009. [2](#), [8](#), [10](#)
- [5] E. Colombo and P. Frediani. Siegel metric and curvature of the moduli space of curves. *Trans. Amer. Math. Soc.*, 362(3):1231-1246, 2010. [3](#), [4](#), [18](#), [28](#)
- [6] E. Colombo, P. Frediani, and A. Ghigi. On totally geodesic submanifolds in the Jacobian locus. *International Journal of Mathematics*, 26 (2015), no. 1, 1550005 (21 pages). [18](#)
- [7] E. Colombo, P. Frediani, G. P. Pirola. Asymptotic directions in the moduli space of curves. arXiv:2405.08641. [4](#), [27](#)
- [8] E. Colombo, G. P. Pirola, and A. Tortora. Hodge-Gaussian maps. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 30(1):125-146, 2001. [3](#), [18](#)
- [9] P. Frediani. Second fundamental form and higher Gaussian maps. arXiv:2208.14794. To appear in *Perspectives on four decades of Algebraic Geometry Volume 1: in Memory of Alberto Collino*, Progress in Mathematics volume 351, Birkhauser. [3](#), [16](#), [18](#), [19](#), [22](#)
- [10] P. Frediani, A. Ghigi and M. Penegini. Shimura varieties in the Torelli locus via Galois coverings. *Int. Math. Res. Not.* 2015(20) (2015) 10595-10623. [4](#)
- [11] P. Frediani, G. P. Pirola. On the geometry of the second fundamental form of the Torelli map. *Proc. Amer. Math. Soc.* 149(3), 1011-1024 (2021). [3](#), [5](#), [18](#), [28](#)
- [12] V. González-Alonso, S. Torelli. Families of curves with Higgs field of arbitrarily large kernel. *Bull. Lond. Math. Soc.* 53 (2021), no.2, 493-506. [27](#)
- [13] X. Lu, K. Zuo, The Oort conjecture on Shimura curves in the Torelli locus of hyperelliptic curves. *J. Math. Pures Appl.* (9) 108 (2017), no. 4, 532-552. [3](#)
- [14] B. Moonen. The Coleman-Oort conjecture: reduction to three key cases. *Bull. Lond. Math. Soc.* 54 (2022), no. 6, 2418-2426. [3](#)
- [15] F. Oort and J. Steenbrink. The local Torelli problem for algebraic curves. In *Journées de Géométrie Algébrique d'Angers, Juillet 1979/Algebraic Geometry*, Angers, 1979, pages 157-204. Sijthoff & Noordhoff, Alphen aan den Rijn, 1980. [2](#), [17](#)
- [16] C. Voisin. Sur l'application de Wahl des courbes satisfaisant la condition de Brill-Noether-Petri. [2](#)
- [17] J. Wahl. Gaussian maps on algebraic curves. *Journal of Differential Geometry*. *J. Differential Geom.* 32(1), 77-98, (1990). [2](#)
- [18] J. Wahl. The Jacobian algebra of a graded Gorenstein singularity. *Duke Math. J.* 55(1987), no.4, 843-871.

1, 2

DARIO FARO, UNIVERSITÀ DEGLI STUDI DI PAVIA, DIPARTIMENTO DI MATEMATICA, VIA FERRATA 1, 27100 PAVIA, ITALY

Email address: `dario.faro@unipv.it`

PAOLA FREDIANI, UNIVERSITÀ DEGLI STUDI DI PAVIA, DIPARTIMENTO DI MATEMATICA, VIA FERRATA 1, 27100 PAVIA, ITALY

Email address: `paola.frediani@unipv.it`

ANTONIO LACOPO, UNIVERSITÀ DEGLI STUDI DI PAVIA, DIPARTIMENTO DI MATEMATICA, VIA FERRATA 1, 27100 PAVIA, ITALY

Email address: `antonio.lacopo01@universitadipavia.it`