

TOTALLY RAMIFIED SUBFIELDS OF p -ALGEBRAS OVER DISCRETE VALUED FIELDS WITH IMPERFECT RESIDUE

S. SRIMATHY

ABSTRACT. Let K be a complete discrete valued field of characteristic p with residue k which is not necessarily perfect. We prove the Conjecture in [CS24] that a p -algebra over K contains a totally ramified cyclic maximal subfield if it contains a totally ramified purely inseparable maximal subfield provided k satisfies some conditions on its p -rank.

1. INTRODUCTION

Let K denote a complete discrete valued field of characteristic p with residue k . The following conjecture appeared in [CS24]:

Conjecture 1.1. ([CS24]) *Let A be a p -algebra over K . Then A contains a totally ramified cyclic maximal subfield if and only if it contains a totally ramified purely inseparable maximal subfield.*

The "only if" part of the conjecture is completely proved in [CS24] and the "if" part is proved for many cases. In this paper, we prove the "if" part of the conjecture for two more cases:

Theorem 1.2. *Let A be a p -algebra of degree p^m , $m > 1$ over K . Suppose A contains a totally ramified purely inseparable maximal subfield. Then it contains a totally ramified cyclic maximal subfield if one of the following conditions holds:*

- (1) $\text{rank}_p(k) \geq 2m$
- (2) $\text{rank}_p(k) \geq m$ and $\dim_{\mathbb{F}_p}(k/\mathcal{P}(k)) \geq 1$

where $\text{rank}_p(k)$ denotes the p -rank of k and \mathcal{P} denotes the Artin-Schreier operator.

In other words, the theorem states that the conjecture is true if k admits at least $2n$ linearly disjoint purely inseparable extensions of degree p or if it admits at least n linearly disjoint purely inseparable extensions of degree p and an Artin-Schreier extension.

The idea of the proof involves first showing the existence of cyclic lifts of purely inseparable extensions of exponent one over the residue. Once we find such cyclic lifts, we construct suitable p -division algebras that contain these cyclic lifts and share the same totally ramified purely inseparable maximal subfield with A . Then we use linkage results of [CFM23] to show the existence of totally ramified cyclic maximal subfields.

2. NOTATIONS

All the fields in this paper have characteristic p . For a field F , the set of non-zero elements of F is denoted by F^\times . Given a field extension L/K , the symbol $N_{L/K}$ denotes the norm function of L over K . The symbol $W_n(F)$ denotes the truncated Witt vector of length n over F .

Fields with discrete valuations are denoted with upper case alphabets and their residue fields are denoted by corresponding lower case alphabets. We denote the valuation ring, its maximal ideal and the value group of a discrete valued field K by \mathcal{O}_K , \mathfrak{m}_K and Γ_K respectively. The set of natural numbers is denoted by \mathbb{N} . The greatest common divisor of $m, n \in \mathbb{N}$ is denoted by (m, n) .

3. PRELIMINARIES

3.1. Albert's theorem. Let F be a field of characteristic p . Let \mathcal{P} denote the Artin-Schreier operator

$$\begin{aligned}\mathcal{P} : F &\rightarrow F \\ x &\mapsto x^p - x\end{aligned}$$

It is well known that cyclic degree p extensions of F upto isomorphism are in bijection with the cyclic subgroups of $F/\mathcal{P}(F)$ of order p ([Lan02, Chapter VI, Theorem 8.3]). Let $W_n(F)$ denote the group of truncated Witt vectors of length n over F . Given any $\omega \in W_n(F)$, one can construct a cyclic extension over F , denoted by F_ω/F . Conversely, given any cyclic extension of degree p^n , one can associate a Witt vector of length n . These are well known ([Jac64, Chapter III], [Tho05]). An explicit way to construct a cyclic extension of degree p^{n+1} containing any given cyclic extension of degree p^n is due to Albert which we recall below.

Theorem 3.1. ([Alb34, Lemma 7], [Jac96, Theorem 4.2.3]) *Let F be a field of characteristic p and let E/F be a cyclic extension of degree p^e , $e \geq 1$ with $\text{Gal}(E/F) = \langle \sigma \rangle$. Then E contains an element β such that $\text{Tr}_{E/F}(\beta) = 1$ and if β is such an element, then there exists an $\alpha \in E$ such that*

$$\mathcal{P}(\beta) = \sigma(\alpha) - \alpha.$$

Then $x^p - x - \alpha$ is irreducible in $E[x]$ and if γ is a root of this polynomial, then $E' = E[\gamma]$ is a cyclic field of degree p^{e+1} over F and any such extension can be obtained this way.

The following remark is obvious:

Remark 3.2. In the above theorem, α can be replaced with $c + \alpha$ for any $c \in F$ and the resulting extension is still cyclic of degree p^{e+1} over F . In other words, the extension $E[\gamma]$ where γ is a root of $x^p - x - (c + \alpha)$ yields a cyclic extension of degree p^{e+1} over F .

3.2. p -rank of a field. Let F be a field of characteristic p . A set of elements $\{x_1, x_2, \dots, x_n\}$ in F is said to be p -independent over F^p if $[F_p(x_1, x_2, \dots, x_n) : F^p] = p^n$. This is equivalent to saying that $[F_p(x_i) : F^p] = p$ and that $F_p(x_i)$ are linearly disjoint over F^p . If moreover, $F_p(x_1, x_2, \dots, x_n) = F$ we say that $\{x_1, x_2, \dots, x_n\}$ is a p -basis for F/F^p . We have that $[F : F^p] = p^n$ where the integer n is called the p -rank (a.k.a the p -dimension) of F , denoted by $\text{rank}_p(F)$. If $[F : F^p]$ is infinite, we set $\text{rank}_p(F) = \infty$.

3.3. A sufficient condition for a cyclic p -algebra to be a division algebra. Let K be a complete discrete valued field with valuation \mathfrak{v} and residue k . We say that an extension L/K is *weakly unramified* if the residue extension satisfies $[l : k] = [L : K]$ (Here we do not assume separability of l/k). Let D be a division algebra over K . The valuation on K extends uniquely to a valuation on D ([Wad02, Corollary 2.2]). Let \overline{D} denote the residue algebra of D and Γ_D denote the value group of D . Denote the degree of D over K by $[D : K]$. We now observe the following:

Lemma 3.3. *Suppose D contains a maximal subfield L that is weakly unramified over K and a maximal subfield that is totally ramified. Then, $\overline{D} = l$ where l is the residue field of L .*

Proof. Since L/K is weakly unramified, $[l : k] = [L : K] = \sqrt{[\overline{D} : K]}$. Since D contains a totally ramified maximal subfield, $[\Gamma_D : \Gamma_K] \geq \sqrt{[\overline{D} : K]}$. By fundamental equality ([Wad02, Equation (2.10)], [Mor95, page 359]),

$$(3.1) \quad [D : K] = [\Gamma_D : \Gamma_K][\overline{D} : k]$$

So we conclude that $[\overline{D} : k] \leq \sqrt{[\overline{D} : K]} = [l : k]$. On the other hand, $l \subseteq \overline{D}$. Therefore, $\overline{D} = l$. \square

Recall that any cyclic p -algebra of degree p^m over K is of the form :

$$\begin{aligned} K \langle x_1, \dots, x_m, y : (x_1^p, \dots, x_m^p) &= (x_1, \dots, x_m) + \omega, \\ y^{p^m} = b, (yx_1y^{-1}, \dots, yx_my^{-1}) &= (x_1, \dots, x_m) + (1, 0, \dots, 0) \rangle \end{aligned}$$

for some $\omega \in W_m(K)$, $b \in K^\times$ and the '+' above denotes addition rule of the Witt vectors. We denote it by the symbol $[\omega, b]_K$.

The following lemma which a simple variant of [CS24, Lemma 3.3], gives a sufficient condition for a p -algebra over K to be a division algebra. The proof is similar to the proof of [CS24, Lemma 3.3], so we skip it.

Lemma 3.4. *Let $b \in K^\times$ with $(\mathfrak{v}(b), p) = 1$. Let $\omega \in W_m(K)$ be such that the corresponding cyclic extension K_ω/K is weakly unramified of degree p^m . Then $[\omega, b]_K$ is a division algebra over K .*

4. CYCLIC EXTENSIONS WITH INSEPARABLE RESIDUE OF EXPONENT ONE

The goal of this section is to prove the following.

Theorem 4.1. *Let K be a complete discrete valued field of characteristic p with valuation \mathfrak{v} and residue k . Suppose $\{\overline{a_1}, \overline{a_2}, \dots, \overline{a_m}\} \subset k^\times$ be elements that are p -independent over k^p . Then there exists a cyclic extension L/K of degree p^m whose residue is $l = k(\sqrt[p]{\overline{a_1}}, \sqrt[p]{\overline{a_2}}, \dots, \sqrt[p]{\overline{a_m}})$.*

Proof. Let t be a uniformizer of K and let $\{a_1, a_2, \dots, a_m\} \subset \mathcal{O}_K^\times$ be arbitrary lifts of $\{\overline{a_1}, \overline{a_2}, \dots, \overline{a_m}\} \subset k^\times$. We will use induction on m to build $L_1, L_2, \dots, L_m = L$ such that L_i/K is cyclic and the residue field of L_i is $l_i = k(\sqrt[p]{\overline{a_1}}, \sqrt[p]{\overline{a_2}}, \dots, \sqrt[p]{\overline{a_i}})$. Let $m = 1$. Consider the Artin-Schreier extension L_1/K given by

$$x_1^p - x_1 = \frac{a_1}{t^p}$$

Let $y_1 = tx_1$. Then y_1 satisfies

$$y_1^p - (t^{p-1})y_1 = a_1$$

Clearly $y_1 \in \mathcal{O}_{L_1}^\times$ and its residue is

$$\overline{y_1} = \sqrt[p]{\overline{a_1}}$$

Therefore the residue field l_1 of L_1 is given by

$$l_1 = k(\sqrt[p]{\overline{a_1}})$$

Now assume by induction hypothesis that we have constructed cyclic L_{i-1}/K with residue field $l_{i-1} = k(\sqrt[p]{\overline{a_1}}, \sqrt[p]{\overline{a_2}}, \dots, \sqrt[p]{\overline{a_{i-1}}})$. By Theorem 3.1 and Remark 3.2, there exists $\alpha_{i-1} \in L_{i-1}$ such that the Artin-Schreier extension L_i/L_{i-1} given by

$$(4.1) \quad x_i^p - x_i = \frac{a_i}{t^{p^{n_i}}} + \alpha_{i-1}$$

is such that L_i/K is cyclic. Choose $n_i \in \mathbb{Z}, n_i \geq 1$ such that $\mathfrak{v}(t^{p^{n_i}}) > -\mathfrak{v}(\alpha_{i-1})$ and let $y_i = t^{p^{n_i-1}}x_i$. Then from (4.1), we get

$$y_i^p - (t^{p^{n_i}-p^{n_i-1}})y_i = a_i + \beta_{i-1}$$

where $\beta_{i-1} = t^{p^{n_i}}\alpha_{i-1} \in \mathfrak{m}_{L_{i-1}}$ by the choice of n_i . From the above equation, we see that $y_i \in \mathcal{O}_{L_i}^\times$ and its residue is given by

$$\overline{y_i} = \sqrt[p]{\overline{a_i}}$$

Therefore the residue field l_i of L_i is given by

$$l_i = l_{i-1}(\sqrt[p]{a_i}) = k(\sqrt[p]{a_1}, \sqrt[p]{a_2}, \dots, \sqrt[p]{a_i})$$

as claimed. \square

5. PROOF OF THEOREM 1.2

Lemma 5.1. *Let $m \in \mathbb{N}$ and let k satisfy either of the following conditions:*

- (1) $\text{rank}_p(k) \geq 2m$
- (2) $\text{rank}_p(k) \geq m$ and $\dim_{\mathbb{F}_p}(k/\mathcal{P}(k)) \geq 1$

Then there exists weakly unramified cyclic extensions L_1/K and L_2/K of degree p^m whose residue satisfy $l_1 \cap l_2 = k$.

Proof. (1) When $\text{rank}_p(k) \geq 2m$: Let $\{\overline{a_1}, \overline{a_2}, \dots, \overline{a_{2m}}\} \subset k^\times$ be elements that are p -independent over k^p . Let $l_1 = k(\sqrt[p]{\overline{a_1}}, \sqrt[p]{\overline{a_2}}, \dots, \sqrt[p]{\overline{a_m}})$ and $l_2 = k(\sqrt[p]{\overline{a_{m+1}}}, \sqrt[p]{\overline{a_{m+2}}}, \dots, \sqrt[p]{\overline{a_{2m}}})$. Then clearly $l_1 \cap l_2 = k$. Let $L_i/K, i = 1, 2$ be the cyclic extensions of degree p^m , whose residue is l_i/k as constructed in Theorem 4.1.

- (2) When $\text{rank}_p(k) \geq m$ and $\dim_{\mathbb{F}_p}(k/\mathcal{P}(k)) \geq 1$: Since $\text{rank}_p(k) \geq m$, there exists $\{\overline{a_1}, \overline{a_2}, \dots, \overline{a_m}\} \subset k^\times$ that are p -independent over k^p . Let $l_1 = k(\sqrt[p]{\overline{a_1}}, \sqrt[p]{\overline{a_2}}, \dots, \sqrt[p]{\overline{a_m}})$. Also since $\dim_{\mathbb{F}_p}(k/\mathcal{P}(k)) \geq 1$, there exists an Artin-Schreier extension of k . By Theorem 3.1, we can extend this to a cyclic extension l_2/k of degree p^m . Clearly, $l_1 \cap l_2 = k$. Let L_1/K be the cyclic extension of degree p^m , whose residue is l_1/k as constructed in Theorem 4.1 and let L_2/K be the inertial lift of l_2/k which is again cyclic of degree p^m .

In both the cases, L_1 and L_2 clearly satisfy the lemma. \square

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2: Let A be a p -algebra of degree p^m over K containing a totally ramified purely inseparable maximal subfield. Then by [CS24, Lemma 5.1, Remark 5.2], $A \simeq [\omega, b)$ for some $\omega \in W_m(F)$, $b \in K^\times$ with $(\mathfrak{v}(b), p) = 1$ and $K(\sqrt[p^m]{b}) \subset A$ is totally ramified purely inseparable over K . Let $L_i/K, i = 1, 2$ be the degree p^m cyclic extension constructed in Lemma 5.1. Let $\omega_i \in W_m(K)$ be Witt vectors corresponding to L_i . Since L_1 and L_2 are weakly unramified, by Lemma 3.4, $D_i = [\omega_i, b)_K, i = 1, 2$ are division algebras over K . Moreover, $\overline{D}_i = l_i$ by Lemma 3.3. Now A, D_1 and D_2 share the same purely inseparable subfield $K(\sqrt[p^m]{b})$ and therefore by [CFM23, Theorem 4.7], share a cyclic maximal subfield L . Now the residue l of L satisfies

$$l \subseteq \overline{D}_1 \cap \overline{D}_2 = l_1 \cap l_2 = k$$

By the fundamental equality [Ser79, Chapter II, §2, Corollary 1], L/K is totally ramified as required. \square

ACKNOWLEDGEMENTS

The author acknowledges the support of the DAE, Government of India, under Project Identification No. RTI4001.

REFERENCES

- [Alb34] A. A. Albert. Cyclic fields of degree p^n over F of characteristic p . *Bull. Amer. Math. Soc.*, 40(8):625–631, 1934.
- [CFM23] Adam Chapman, Mathieu Florence, and Kelly McKinnie. Common splitting fields of symbol algebras. *Manuscripta Math.*, 171(3-4):649–662, 2023.
- [CS24] Adam Chapman and S. Srimathy. Totally ramified subfields of p -algebras, 2024.
- [FJ08] Michael D. Fried and Moshe Jarden. *Field arithmetic*, volume 11 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, third edition, 2008. Revised by Jarden.

- [Jac64] Nathan Jacobson. *Lectures in abstract algebra. Vol III: Theory of fields and Galois theory*. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London-New York, 1964.
- [Jac96] Nathan Jacobson. *Finite-dimensional division algebras over fields*. Springer-Verlag, Berlin, 1996.
- [Lan02] Serge Lang. *Algebra*, volume 211 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2002.
- [Mor95] Patrick J. Morandi. On defective division algebras. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 359–367. Amer. Math. Soc., Providence, RI, 1995.
- [Ser79] Jean-Pierre Serre. *Local fields*, volume 67 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1979. Translated from the French by Marvin Jay Greenberg.
- [Tho05] Lara Thomas. *Arithmétique des extensions d’Artin-Schreier-Witt*. 2005. Thesis (Ph.D.)—Université Toulouse II Le Mirail.
- [Wad02] A. R. Wadsworth. Valuation theory on finite dimensional division algebras. In *Valuation theory and its applications, Vol. I (Saskatoon, SK, 1999)*, volume 32 of *Fields Inst. Commun.*, pages 385–449. Amer. Math. Soc., Providence, RI, 2002.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, MUMBAI, 400005, INDIA

Email address: srimathy@math.tifr.res.in