

Inverse variational problem for equations in the Riccati chain

Pranab Sarkar*

*Department of Chemistry, Visva-Bharati University,
Santiniketan 731235, West Bengal, India*

Pratik Majhi[†] and Madan Mohan Panja[‡]

*Department of Mathematics, Visva-Bharati University,
Santiniketan 731235, West Bengal, India*

Benoy Talukdar[§]

*Department of Physics, Visva-Bharati University,
Santiniketan 731235, West Bengal, India*

Abstract

The nonstandard Lagrangian representations of Riccati and Riccati-type equations that exist in the literature cannot be obtained using Helmholtz solution of the inverse problem. In this work we consider Riccati and higher-order Riccati equations and construct their standard Lagrangian representation by using a simple variant of the Helmholtz theory. We make use of the self-adjoint form of the linear equations corresponding to odd-order equations in the Riccati chain to provide a symmetry-based approach for the solution of inverse problem. Explicit results presented for Lagrangians of the first and third-order Riccati equations show that one cannot Hamiltonize the Riccati family of equations by the traditional method used in classical mechanics.

I. INTRODUCTION

The N th order equation in the Riccati hierarchy can be written as [1]

$$\vartheta^N \omega(x) + \sum_{j=1}^N \alpha_j(x) (\vartheta^{j-1} \omega(x)) + \alpha_0(x) = 0 \quad (1)$$

with the differential operator

$$\vartheta = \frac{d}{dx} + \omega(x). \quad (2)$$

Here α_j , $j = 0, 1, \dots, N$, stand for $N + 1$ well behaved arbitrary functions of x . Clearly, for $N = 1$, the expression in (1) gives the well known Riccati equation

$$\frac{d\omega(x)}{dx} + \omega(x)^2 + \alpha_1(x)\omega(x) + \alpha_0(x) = 0 \quad (3)$$

while for $N > 1$ we get from it higher-order Riccati equations which, together with (3), form the so-called Riccati chain. It is interesting to note that each equation in the Riccati chain obtained from (1) can be linearized by the use of the Cole-Hopf transformation [2, 3]

$$\omega(x) = \frac{d \ln y(x)}{dx} \quad (4)$$

such that use of (4) in (1) gives the linear equation

$$\sum_{j=0}^N \alpha_j(x) \frac{d^j y(x)}{dx^j} + \frac{d^{N+1} y(x)}{dx^{N+1}}. \quad (5)$$

* pranab.sarkar@visva-bharati.ac.in

† iampratikmajhi@gmail.com

‡ madanpanja2005@yahoo.co.in

§ binoy123@bsnl.in

The object of the present work is to construct Lagrangian representations of the linear equations in (5) and then make judicious use of (4) to find the Lagrangians for the associated Riccati and/or higher Riccati equation.

In classical mechanics the standard Lagrangian is represented by $L = T - V$, where T is the kinetic energy of the system modeled by the equation and V , the corresponding potential function. In addition to the traditional Lagrangian L used in classical mechanics, a new type of Lagrangian functions was proposed [4] for the second-order Riccati and some dissipative-like autonomous differential equations related to the generalized Riccati equation. These do involve neither T nor V . Despite that, the sought Lagrangian functions lead to equations of motion via Euler-Lagrange equations. As a result these Lagrangians were qualified as ‘non-standard’. The non-standard Lagrangians do not have a natural space in the theory of inverse problem [5, 6] in the calculus of variation. We are, however, interested here in the inverse variational problem to provide Lagrangian representations of equations in the Riccati chain. For brevity, we shall refer to the associated Lagrangian as standard one. The present study is expected to play a role in several areas of physics since Riccati equation is related to many equations of fundamental physics [7, 8]. In the recent past Faraoni [9] reported a solution of the inverse problem for only the Riccati equation by taking recourse to the use of an analogy with the Friedmann equation [10]. This method seems to be very special and unlikely to be applicable for other equations in the Riccati chain.

From (1) and (5) we see that linear equation corresponding Riccati equation is of order two, while other equations in the Riccati chain are associated with higher-order linear equations. Thus the problem of constructing Lagrangian for higher members in the Riccati chain should be treated within the framework of a generalized mechanics in which the line integral implied in the definition of action depends on derivatives of generalized coordinates higher than the first. Let $L(x, y(x), y'(x), y''(x), \dots, y^{(n)}(x))$, with $y^{(n)}(x) = \frac{d^n y(x)}{dx^n}$ denote the Lagrangian of a one dimensional system of order n . The corresponding Euler-Lagrange equation is given by [11]

$$\sum_{i=0}^n (-1)^i \frac{d^i}{dx^i} \frac{\partial L}{\partial y^{(i)}} = 0. \quad (6)$$

Clearly, the value of n in (6) determines the order of the generalized Euler-Lagrange equation. It is well known that the standard Lagrangian L in (6) for $n = 1$ leading to second-order differential equation is regular or degenerate if the associated Hessian or functional deter-

minant [12] is non-null with the possible exception of a finite number of isolated zeros. Otherwise, L is non-degenerate. Similarly, non-zero and zero values of the functional determinant of the higher-order Lagrangian determine if it is degenerate or non-degenerate. In looking for a generalization of the Ostrogradsky theorem, Motohashi and Suyama [13] have clearly pointed out that Lagrangians for even-order linear differential equations are regular while those of odd-order equations are non-degenerate. The odd-order Riccati and/or Riccati type equations, via Cole-Hopf transformation [2, 3] lead to even order linear equations. Consequently, Lagrangians of these equations will be regular. In the following we restrict ourselves to odd-order Riccati equations only.

The Lagrangian of any mechanical system governed by a second-order linear differential equation can be found by solving the so-called Helmholtz problem [5]. It is, however, not straightforward to adapt the approach to deal with higher-order systems. It appears that investigation in the inverse variational problem for such systems has yet remained largely unexplored [14]. In view of this, we shall make use of certain symmetry considerations [15] to provide a simple recipe to compute Lagrangians of higher-order systems. We postulate that the Lagrangian of a higher-order system is proportional to the self adjoint form of its equation of motion. This simple minded solution of the inverse problem sought by us has an old root in the scientific literature [16].

The theory of self-adjoint differential equation for the second-order equation can be found in any standard text book on mathematical physics [17]. This is, however, not true for higher-order equations. We thus begin Sec. II with a brief introduction to algebraic properties of higher-order self-adjoint systems and then present an ansatz to solve the associated inverse variational problem. In Sec. III we provide some useful checks for the validity of the proposed ansatz in terms of two well-known physically important examples and then demonstrate how our approach could be judiciously employed to find a standard Lagrangian representation of the Riccati equation. Sec. IV is devoted to deal with similar problems for higher-order equations in the Riccati chain. Here we pay special attention to the third-order Riccati equation with an associated fourth-order linear equation and present explicit result for its Lagrangian function. In Appendix we provide a general expression for the Lagrangian of $(2n - 1)$ th order equation, $n \geq 1$, in the Riccati chain. Finally, in Sec. V, we summarize our outlook on the present work and make some concluding remarks.

II. SELF-ADJOINT DIFFERENTIAL EQUATIONS AND INVERSE VARIATIONAL PROBLEM

Let $M(y)$ stand for the differential expression

$$M(y) = r_0(x) \frac{d^n y(x)}{dx^n} + r_1(x) \frac{d^{n-1} y(x)}{dx^{n-1}} + \cdots + r_n(x) y(x), \quad (7)$$

where n is any positive integer. The coefficients r are real or complex function of the real variable x , defined in an interval $a \leq x \leq b$. The relation $M(y) = 0$ defines an n th order non self-adjoint linear differential equation. The general adjoint system associated with the differential operator in (7) is written as

$$N(z) = (-1)^n \frac{d^n (r_0(x) z(x))}{dx^n} + (-1)^{n-1} \frac{d^{n-1} (r_1(x) z(x))}{dx^{n-1}} + \cdots + r_n(x) z(x). \quad (8)$$

The adjoint operator $N(z)$ is connected with $M(y)$ by the Lagrangian identity

$$zM(y) - yN(z) = \frac{dp(y, z)}{dx}, \quad (9)$$

where $p(y, z)$ stands for a bilinear form in $y(x), y'(x), \dots, y^{(n-1)}(x); z(x), z'(x), \dots, z^{(n-1)}(x)$ with coefficients which are functions of x . Ince [18] has carefully demonstrated how to exploit (7), (8) and (9) to construct the self-adjoint equation for a given higher-order linear ordinary differential equation. We shall now present a recipe for the solution of the inverse variational problem for a higher-order equation, where its self-adjoint form plays a crucial role.

The relation between symmetries of a Lagrangian and conserved quantities of the corresponding equation of motion is a very well known result in classical mechanics. But it is less well known that the symmetries of the equation of motion form a larger set than that of the Lagrangian. Hojman [15] supplemented the well-known concept of s-equivalence [19] by a new kind of symmetry such that the resulting set of Lagrangian symmetries coincides with the symmetries of the equations of motions. In this context it was established that a second-order Lagrangian is proportional to a second-order differential equation which follows from the traditional first-order Lagrangian. The constant of proportionality is determined by a gauge term that reduces the expression for the second-order Lagrangian to the first-order one. We postulate that the n th order Lagrangian bears a similar relation with the n th differential equation such that the ansatz

$$L(x, y(x), y'(x), y''(x), \dots, y^{(n)}(x)) = y(x) M_s(y), \quad (10)$$

provides a simple solution for the inverse variational problem for higher-order systems. In equation (10), $M_s(y)$ stands for the self-adjoint differential expression corresponding to $M(y)$.

III. LAGRANGIAN FUNCTION FOR THE RICCATI EQUATION

Before we employ (10) to compute the Lagrangian function of the Riccati equation, we would like to present some useful checks on its validity in terms of simple case studies. To that let us consider the following second- and fourth-order self-adjoint equations.

$$y''(x) + y(x) = 0 \quad (11)$$

and

$$y^{(4)}(x) + y(x) = 0. \quad (12)$$

Because of (10) the Lagrangians for (11) and (12) are given by

$$L(x, y(x), y'(x), y''(x)) = y(x) (y''(x) + y(x)) \quad (13)$$

and

$$L(x, y(x), y'(x), y''(x), y'''(x), y^{(4)}(x)) = y(x) (y^{(4)}(x) + y(x)). \quad (14)$$

When a gauge term $\frac{d}{dx}(y(x)y'(x))$ is subtracted from the second-order Lagrangian in (13) we get the familiar first-order Lagrangian for (11). The gauge term for a fourth-order Lagrangian is $\frac{d}{dx}(y(x)y'''(x))$ [20] which when subtracted from (14) leads to a third-order Lagrangian. It is easy to verify that the constructed lower order Lagrangian via the appropriate Euler-Lagrange equation reproduces the fourth-order equation in (12). These results suggest that application of our solution of the inverse problem to Riccati equation is now in order.

From (5) the linear equation corresponding to the Riccati equation (3) is obtained as

$$y''(x) + \alpha_1(x)y'(x) + \alpha_0(x)y(x) = 0. \quad (15)$$

Equation (15) is not self adjoint. But it can be reduced to the self-adjoint form by multiplying with $e^{\int \alpha_1(x)dx}$ [11] so as to write the Lagrangian for (15) as

$$L = y(x)e^{\int \alpha_1(x)dx} (y''(x) + \alpha_1(x)y'(x) + \alpha_0(x)y(x)). \quad (16)$$

From the Cole-Hopf transformation (4) we write

$$y(x) = ae^{\int \omega(x) dx}, \quad (17)$$

where a is a constant of integration, Equations (16) and (17) give the Lagrangian for the Riccati equation.

$$L = a^2 e^{\int \alpha_1(x) dx + 2 \int \omega(x) dx} (\omega'(x) + \omega^2(x) + \alpha_1(x)\omega(x) + \alpha_0(x)). \quad (18)$$

It is clear from (18) that as with the Lagrangian of the linear equation (15), the Lagrangian for the Riccati equation, which is essentially nonlinear, also involves its own equation of motion. It is easy to verify that L in (18) leads to the Riccati equation via the traditional first-order Euler-Lagrange equation. We shall now extend the above approach to higher-order equations in the Riccati chain.

IV. LAGRANGIAN FUNCTIONS FOR ODD-ORDER EQUATIONS IN THE RICCATI CHAIN

Next to Riccati equation we are interested in the third-order equation in the Riccati chain presumably because the linear equation corresponding to it is of even order. Using $N = 3$ in (1) and (5) we get results for the Riccati-3 and corresponding linear equation in the following form.

$$\begin{aligned} \frac{d^3 \omega}{dx^3} + (\alpha_3(x) + 4\omega(x)) \frac{d^2 \omega}{dx^2} + 3 \left(\frac{d\omega}{dx} \right)^2 + (6\omega(x)^2 + 3\alpha_3(x)\omega(x) + \alpha_2(x)) \frac{d\omega}{dx} + \omega(x)^4 \\ + \alpha_3(x)\omega(x)^3 + \alpha_2(x)\omega(x)^2 + \alpha_1(x)\omega(x) + \alpha_0(x) = 0 \end{aligned} \quad (19)$$

and

$$\frac{d^4 y}{dx^4} + \alpha_3(x) \frac{d^3 y}{dx^3} + \alpha_2(x) \frac{d^2 y}{dx^2} + \alpha_1(x) \frac{dy}{dx} + \alpha_0(x) y = 0. \quad (20)$$

Clearly, Eq. (20) is a special form of the more general equation

$$r_0(x) \frac{d^4 y}{dx^4} + r_1(x) \frac{d^3 y}{dx^3} + r_2(x) \frac{d^2 y}{dx^2} + r_3(x) \frac{dy}{dx} + r_4(x) y = 0. \quad (21)$$

obtained from $M(y) = 0$ for $n = 4$. The self-adjoint form of (21) can be written as [21]

$$(r_0(x)y''(x))'' + (q(x)y'(x))' + r_4(x)y(x) = 0 \quad (22)$$

with the following conditions

$$r_1(x) = 2r_0'(x), \quad q(x) = r_2(x) - r_0''(x) \quad \text{and} \quad q'(x) = r_3(x). \quad (23)$$

In view of (10) we use (22) to write the Lagrangian of the self-adjoint equation in the form

$$L = y(x) \left(r_0(x)y^{(4)}(x) + 2r_0'(x)y^{(3)}(x) + r_0''(x)y''(x) + q(x)y''(x) + q'(x)y'(x) + r_4(x)y(x) \right). \quad (24)$$

It is straightforward to verify that L in (24) via the fourth-order Euler-Lagrange equation obtained from (6) reproduces (22). Making use of (23) we rewrite (24) to read

$$L = y(x) \left(r_0(x)y^{(4)}(x) + r_1(x)y^{(3)}(x) + r_2(x)y''(x) + r_3(x)y'(x) + r_4(x)y(x) \right). \quad (25)$$

Comparing (20) and (21) we have $r_0(x) = 1$, $r_1(x) = \alpha_3(x)$, $r_2(x) = \alpha_2(x)$, $r_3(x) = \alpha_1(x)$ and $r_4(x) = \alpha_0(x)$. When these relations are substituted in (25) we get the Lagrangian for the fourth-order linear equation corresponding third-order equation in the Riccati chain.

$$L = y(x) \left(y^{(4)}(x) + \alpha_3(x)y^{(3)}(x) + \alpha_2(x)y''(x) + \alpha_1(x)y'(x) + \alpha_0(x)y(x) \right). \quad (26)$$

We can now combine (17) and (26) to obtain the Lagrangian of the third-order Riccati equation as

$$L = a^2 e^{2 \int \omega(x) dx} (S_1(x) + S_2(x)) \quad (27)$$

with

$$S_1(x) = \omega^{(3)}(x) + 4\omega(x)\omega''(x) + \alpha_3(x)\omega''(x) + 3\omega'(x)^2 + 6\omega(x)^2\omega'(x) + 3\alpha_3(x)\omega(x)\omega'(x) \quad (28)$$

and

$$S_2(x) = \alpha_2(x)\omega'(x) + \omega(x)^4 + \alpha_3(x)\omega(x)^3 + \alpha_2(x)\omega(x)^2 + \alpha_1(x)\omega(x) + \alpha_0(x). \quad (29)$$

It is straightforward to verify that the result in (29) follows from (A.8) in the Appendix for $n = 2$.

V. CONCLUDING REMARKS

The Newtonian approach to classical mechanics is a force-momentum formulation with cause and effect embedded in it. On the other hand, the Lagrangian approach is developed

in terms of kinetic and potential energies and makes use of a single scalar quantity called the Lagrangian. In the recent past, connections were established between the Newton's laws of motion and Riccati equation [22]. Here we look for Lagrangian representation of equations in the Riccati chain. The explicit result presented for the Riccati equation exhibits that it follows from a first-order Lagrangian depending linearly on $\omega'(x)$ rather than quadratically. This provides an awkward analytic constraint to Hamiltonize the Riccati system because the quantity $\frac{dL}{d\omega'(x)}$ cannot be inverted for $\omega'(x)$. Consequently, we need to take recourse to the use of the Dirac's theory of constraints [23] to obtain the necessary Hamiltonian formulation. We have an identical situation for the third-order Riccati equation because the Lagrangian in (27) is also linear in $\omega^{(3)}$. This is, in general, true for all odd-order equations in the Riccati chain. It will be quite interesting to look for Hamiltonian formulation of Riccati equations because besides classical mechanics Hamiltonian formulation plays important roles in statistical physics and quantum mechanics [24]. Finally, we conclude by noting that in the present study we concentrated on the Lagrangian representation for half of the equations in the Riccati chain. As regards the other half, we noted that each even-order equation in the Riccati chain leads to an odd-order linear equation. This did not permit us to look for a solution of the inverse problem for even-order Riccati-type equations. But we believe that the work of Motohashi and Suyama [13] may be of some help in respect of this.

-
- [1] A. M. Grundland and D. Levi, Higher-order riccati equations as bäcklund transformations, *J. Phys. A: Math-Gen* **32**, 3931 (1999).
 - [2] J. D. Cole, On a quasi-linear parabolic equation occurring in aerodynamics, *Quart. Appl. Math.* **9**, 225 (1951).
 - [3] E. Hopf, The partial differential equation $u_t + uu_x = u_{xx}$, *Comm. Pure Appl. Math.* **3**, 201 (1950).
 - [4] J. F. Cariñena, M. F. Rañada, and M. Santander, Lagrangian formalism for nonlinear second-order riccati systems: one-dimensional integrability and two-dimensional superintegrability, *J. Math. Phys.* **46**, 062703 (2005).
 - [5] H. Helmholtz, Ueber die physikalische bedeutung des principis der kleinsten wirkung, *J. Reine Angew. Math.* **100**, 213 (1887).

- [6] A. Saha and B. Talukdar, Inverse variational problem for nonstandard lagrangians, Rep. Math. Phys. **73**, 299 (2014).
- [7] D. Schuch, Relations between nonlinear riccati equations and other equations in fundamental physics, *J. Phys. Conf. Ser.*, **538**, 012019 (2014).
- [8] D. Schuch, *Quantum theory from a nonlinear perspective* (Springer, New York, 2018).
- [9] V. Faraoni, Helmholtz problem for the riccati equation from an analogous friedmann equation, Eur. Phys. J. C **82**, 13 (2022).
- [10] A. Liddle, *An introduction to modern cosmology* (Wiley, New York, 2015).
- [11] R. Courant and D. Hilbert, *Methods in Mathematical Physics, Vol. 1* (Wiley Eastern Pvt. Ltd., New Delhi, 1975).
- [12] R. M. Santilli, *Foundations of Theoretical Mechanics I: the inverse problem in Newtonian mechanics, Vol. 1* (Springer Verlag, New York, 1978).
- [13] H. Motohashi and T. Suyama, Third order equations of motion and the ostrogradsky instability, Phys. Rev. D **91**, 085009 (2015).
- [14] S. Ghosh, *Inequivalent Lagrangians and their Physical Implications*, Ph.D. thesis, Visva-Bharati University, Santiniketan 731235, West Bengal, India (2010).
- [15] S. Hojman, Symmetries of lagrangians and of their equations of motion, J. Phys. A : Math. Gen. **17**, 2399 (1984).
- [16] R. Hojman, S. Hojman, and J. Sheinbaum, Shortcut for constructing any lagrangian from its equations of motion, Phys. Rev. D **28**, 1333 (1983).
- [17] G. B. Arfken and H. J. Weber, *Mathematical methods for physicists, Sixth ed.* (Elsevier Academic Press, New York, 2012).
- [18] E. L. Ince, *Ordinary differential equations* (Dover Publication, New York, 1958).
- [19] D. G. Currie and E. J. Saletan, q-equivalent particle hamiltonians. i. the classical one-dimensional case, J. Math. Phys. **7**, 967 (1966).
- [20] C. Carathéodory, *Calculus of variations and partial differential equations of the first order, Vol. 2* (Holden-Day Inc., San Fransisco, 1967).
- [21] M. P. Beccar-Varela, M. A. M. Bhuiyan, M. C. Mariani, and O. K. Tweneboah, Analytic methods for solving higher order ordinary differential equations, Mathematics **7**, 826 (2019).
- [22] M. Nowakowski and H. C. Rosu, Newton's laws of motion in the form of a riccati equation, Phys. Rev. E **65**, 047602 (2002).

- [23] P. A. M. Dirac, *Lectures on Quantum Mechanics*, Belfer Graduate School Monograph Series No. 2 (Yeshiva University, NY, 1964).
- [24] E. C. G. Sudarshan and N. Mukunda, *Classical dynamics: a modern perspective* (John-Wiley & Sons, Inc., NY, 1974).

Appendix: General expression for the Lagrangian of any odd-order equation in the Riccati Chain

From Eq. (1) we write

$$\vartheta^{2n-1}\omega(x) + \sum_{j=1}^{2n-1} \alpha_j(x) (\vartheta^{j-1}\omega(x)) + \alpha_0(x) = 0 \quad (\text{A.1})$$

for the odd-order Riccati equations. Here $n \geq 1$. Due to the transformation in Eq. (4) the even-order linear equations corresponding to (A.1) are given by

$$\frac{d^{2n}y}{dx^{2n}} + \sum_{j=0}^{2n-1} \alpha_j(x) \frac{d^j y}{dx^j} = 0. \quad (\text{A.2})$$

The general linear homogeneous ordinary differential equation

$$\sum_{i=0}^{2n} r_i(x) \frac{d^{2n-i}y}{dx^{2n-i}} = 0 \quad (\text{A.3})$$

of order $2n$ is self adjoint if and only if

- (i) the even coefficients $r_0(x), r_2(x), \dots, r_{2n}(x)$ are continuous functions of x and
- (ii) the odd coefficients $r_1(x), r_3(x), \dots, r_{2n-1}(x)$ satisfy the recurrence relation

$$r_{2k+1} = \frac{1}{2} \left\{ \binom{2n}{2n-2k-1} r_0^{(2k+1)} - \binom{2n-1}{2n-2k-1} r_1^{(2k)} + \binom{2n-2}{2n-2k-1} r_2^{(2k-1)} - \dots \right. \\ \left. \dots - \binom{2n-2k+1}{2n-2k-1} r_{2k-1}'' + \binom{2n-2k-1}{2n-2k-1} r_{2k}' \right\}, \quad (\text{A.4})$$

with initial condition $r_1 = n r'_0$ and $k \geq 1$. A solution of the recurrence relation (A.4) can be written as

$$\begin{aligned}
r_{2k+1} = & \left\{ \frac{1}{2} \binom{2n}{2k+1} + \sum_{l=2}^{k+1} \frac{(-1)^{l-1}}{2^l} \sum_{(k_1 k_2 \dots k_l) \in I_l^{2k+1}} \binom{2n}{k_1} \binom{2n-k_1}{k_2} \dots \binom{2n-\sum_{i=1}^{l-1} k_i}{k_l} \right\} r_0^{(2k+1)} \\
+ & \left\{ \frac{1}{2} \binom{2n-2}{2k-1} + \sum_{l=2}^k \frac{(-1)^{l-1}}{2^l} \sum_{(k_1 k_2 \dots k_l) \in I_l^{2k-1}} \binom{2n-2}{k_1} \binom{2n-2-k_1}{k_2} \dots \binom{2n-2-\sum_{i=1}^{l-1} k_i}{k_l} \right\} r_2^{(2k-1)} \\
+ & \left\{ \frac{1}{2} \binom{2n-4}{2k-3} + \sum_{l=2}^{k-1} \frac{(-1)^{l-1}}{2^l} \sum_{(k_1 k_2 \dots k_l) \in I_l^{2k-3}} \binom{2n-4}{k_1} \binom{2n-4-k_1}{k_2} \dots \binom{2n-4-\sum_{i=1}^{l-1} k_i}{k_l} \right\} r_4^{(2k-3)} \\
+ & \dots + \left\{ \frac{1}{2} \binom{2n-2k+2}{3} - \frac{1}{2^2} \binom{2n-2k+2}{1} \binom{2n-2k+1}{2} \right\} r_{2k-2}''' + \frac{1}{2} \binom{2n-2k}{1} r'_{2k}, \tag{A.5}
\end{aligned}$$

where, for a positive odd integer m , the index set I_l^m consists of the elements (k_1, k_2, \dots, k_l) of length $l \geq 2$ defined by

$$(k_1 k_2 \dots k_l) \in I_l^m \Leftrightarrow \begin{cases} k_1 \text{ is odd positive integer } < m, \\ k_1 + k_2 + k_3 + \dots + k_l = m, \\ k_2, k_3, \dots, k_l \text{ all are even positive integers.} \end{cases} \tag{A.6}$$

Thus in analogy with Eq. (24), the Lagrangian for the general equation (A.3) is given by

$$L(x, y, y', y'', \dots, y^{2n}) = y(x) \sum_{i=0}^{2n} r_i(x) \frac{d^{2n-i} y}{dx^{2n-i}}. \tag{A.7}$$

The expression in (A.7) will represent the Lagrangian function for the odd-order Riccati equations in (A.1) and/or (A.2) if $r_0(x) = 1$ and $r_j(x) = \alpha_{2n-j}(x)$, $j = 1, 2, \dots, 2n-1$.

Making use of Eq. (17) in Eq. (A.7) we finally obtain

$$L(\cdot) = a^2 e^{2 \int \omega(x) dx} (\vartheta^{2n-1} \omega + \alpha_{2n-1}(x) \vartheta^{2n-2} \omega + \alpha_{2n-2}(x) \vartheta^{2n-3} \omega \dots + \alpha_2(x) \vartheta \omega + \alpha_1(x) \omega + \alpha_0(x)) \tag{A.8}$$

for the $(2n-1)$ th order Riccati equation.